

Geometric regularisation for optimal transport with strongly p -convex cost

Lukas Koch¹

¹*Max Planck Institute for Mathematics in the Sciences, 04103 Leipzig, Germany*
lkoch@mis.mpg.de

March 21, 2023

Abstract

We prove a geometric regularisation result for minimisers of optimal transport problems where the cost-function is strongly p -convex and of p -growth. Initial and target measures are allowed to be rough, but are assumed to be close to Lebesgue measure.

1 Introduction

The study of the optimal transport problem:

$$\min_{\pi \in \Pi(\lambda, \mu)} \int_{\mathbb{R}^d \times \mathbb{R}^d} c(x - y) d\pi \quad (1.1)$$

is well established. We refer the reader to [18] and [17] for an introduction and overview of the literature. When solutions take the form of a transport map $\pi = (\text{Id} \times T)_{\#}\mu$, under mild assumptions, minimisers are characterised by satisfying the Euler-Lagrange equation

$$\lambda(x) \det(DT(x)) = \mu(T(x)) \quad (1.2)$$

as well as the additional structure condition

$$T(x) = x + \nabla c^*(D\phi), \quad (1.3)$$

where ϕ is a c -concave function and c^* denotes the convex conjugate of c . Assuming $\mu \sim \lambda \sim 1$ and linearising the geometric nonlinearity in (1.2), that is formally expanding $\det(\text{Id} + A) = 1 + \text{tr} A + \dots$, we find that

$$\text{div} \nabla c^*(D\phi) = \mu - \lambda. \quad (1.4)$$

Thus, at least formally, we expect solutions of (1.1) to be well approximated by solutions of (1.4). Note that in general (1.4) is a nonlinear equation. Thus we refer to the process of moving from (1.1) to (1.4) as geometric linearisation. The aim of this paper is to make this connection rigorous. We show the following:

Theorem 1.1. *Let $1 < p < \infty$. Suppose $c: \mathbb{R}^d \rightarrow \mathbb{R}$ is a strongly p -convex cost function of controlled-duality p -growth. Let π be a minimiser of (1.1) for some measures λ, μ . Denote*

$$\begin{aligned} E(R) &:= \frac{1}{|B_R|} \int_{(B_R \times \mathbb{R}^d) \cup (\mathbb{R}^d \times B_R)} c(x - y) \, d\pi \\ D(R) &:= \frac{1}{|B_R|} W_p^p(\lambda \llcorner B_R, \kappa_{\lambda, R} \, dx \llcorner B_R) + \frac{R^p}{\kappa_{\lambda, R}^{p-1}} (\kappa_{\lambda, R} - 1)^p \\ &\quad + \frac{1}{|B_R|} W_p^p(\mu \llcorner B_R, \kappa_{\mu, R} \, dx \llcorner B_R) + \frac{R^p}{\kappa_{\mu, R}^{p-1}} (\kappa_{\mu, R} - 1)^p. \end{aligned}$$

Here $\kappa_{\lambda, R} = \frac{\lambda(B_R)}{|B_R|}$ and $\kappa_{\mu, R} = \frac{\mu(B_R)}{|B_R|}$. Then, for every $\tau > 0$, there exists $\varepsilon(\tau) > 0$ such that if $E(4) + D(4) \leq \varepsilon$, then there exists a radius $R \in (2, 3)$, $c \in \mathbb{R}$ and ϕ satisfying

$$-\operatorname{div} \nabla c^*(D\phi) = c \text{ in } B_R$$

such that

$$\int_{(B_1 \times \mathbb{R}^d) \cup (\mathbb{R}^d \times B_1)} c(x - y - \nabla c^*(D\phi)) \, d\pi \lesssim \tau E(4) + D(4).$$

Moreover

$$\sup_{B_1} |D\phi|^{p'} + \int_{B_R} |D\phi|^{p'} \, dx \lesssim E(4) + D(4).$$

We remark that we explain our assumptions on the cost function in detail in Section 1.1.

Traditionally, (1.1) has been approached via the study of (1.2) using the theory of fully nonlinear elliptic equations developed by CAFFARELLI, see e.g. [3, 2] and the references therein. Recently, an alternative approach using variational techniques has been developed by GOLDMAN AND OTTO in [8]. There, partial $C^{1,\alpha}$ -regularity for solutions to (1.1) in the case of Hölder-regular densities λ, μ and quadratic cost function $c(x - y) = \frac{1}{2}|x - y|^2$ was proven. The key tool in the proof was a version of Theorem 1.1 in this setting. In later papers, continuous densities [5], rougher measures [7], more general cost functions (albeit still close to the quadratic cost functional) [15], as well as almost-minimisers of the quadratic cost functional [15] were considered. The quadratic version of Theorem 1.1 was also used to provide a more refined linearisation result of (1.2) in the quadratic set-up in [7] and of a similar statement in the context of optimal matching in [6]. Finally, quadratic versions of Theorem 1.1 played a key role in disproving the existence of a stationary cyclically monotone Poisson matching in 2-d [11].

We remark that very little information is available about the regularity of minimiser of (1.1) already in the simplest degenerate/singular case $c(x - y) = \frac{|x - y|^p}{p}$. In order to attempt to extend the techniques of [8] to this setting, an essential first step is Theorem 1.1. This result will also play a key role in extending the results of [11] to p -costs with $p \neq 2$.

The strategy of proof is similar to that used in [7], although with a number of simplifications. Further, we point the reader to [12] where a detailed account of the

proof of Theorem 1.1 and the motivations behind the strategy are given in the quadratic case.

Here we only comment on the steps where additional effort is required. An essential part of the proof is to obtain a L^∞ -bound for minimisers of (1.1) in the small-energy regime, see Section 3. In the quadratic case, this relies on the monotonicity (in the classical sense) property of solutions. In the non-quadratic case, c -monotonicity needs to be used directly. Our proof relies on the same intuition but highlights how the L^∞ -bound is a direct consequence of convexity and growth bounds of the cost function. We remark that L^∞ -bounds for p -homogeneous cost functions with $p \geq 2$ in all energy regimes were obtained in [9].

A further key step is the so-called quasi-orthogonality property, see Section 7. In the quadratic case, this relied on expanding squares. Again, if $p \neq 2$, this tool is not available and needs to be replaced by exploiting inequalities expressing the convexity of c . Finally, regularity properties of solutions to (1.4) play a key role in the proof. In the quadratic case, such solutions are harmonic and hence very regular- the proof in [7] requires C^3 -regularity of solutions! Already in the case $c(x-y) = \frac{|x-y|^p}{p}$ with $p \neq 2$, the best regularity that is known for solutions to (1.4) in general is $C^{1,\beta}$ -regularity for some $\beta > 0$. Thus, at various places in the proof, more careful estimates are needed.

Finally, we comment why we restrict our attention to cost functions of the form $c(x-y)$. This is due to the fact that our proof relies on the availability of a dynamical formulation. We want to identify points $(x,y) \in \text{spt } \pi$ with the trajectory $X(t) = tx + (1-t)y$. This is related to the Benamou-Brenier formulation of optimal transport, c.f. [4], which in our case states that (1.1) can be alternatively characterised as

$$\min_{(j,\rho)} \left\{ \int c \left(\frac{dj}{d\rho} \right) d\rho : \partial_t \rho + \text{div } j = 0, \rho(0) = \lambda, \rho(1) = \mu \right\} \quad (1.5)$$

Here $\frac{dj}{d\rho}$ denotes the Radon-Nikodym derivative. This alternative, dynamical formulation of optimal transport is only available for costs of the form $c(x-y)$ where c is convex.

1.1 Assumptions on the cost function and its dual

In this section, we explain our assumptions on the cost function c . The convex theory we quote can be found in [16] and [10].

Let $p \in (1, \infty)$. We consider a C^1 -cost function $c: \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying the following properties: There is $\Lambda \geq 1$ such that

- (i) c is strongly p -convex: for any $x, y \in \mathbb{R}^d$ and $\tau \in [0, 1]$,

$$\Lambda^{-1} \tau(1-\tau) V_p(x, y) + c(\tau x + (1-\tau)y) \leq \tau c(x) + (1-\tau)c(y). \quad (1.6)$$

where

$$V_p(x, y) = \begin{cases} (|x|^2 + |y|^2)^{\frac{p-2}{2}} |x-y|^2 & \text{if } p \leq 2 \\ |x-y|^p & \text{if } p \geq 2. \end{cases}$$

(ii) c has p -growth: for any $x \in \mathbb{R}^d$,

$$\Lambda^{-1}|x|^p \leq c(x) \leq \Lambda|x|^p. \quad (1.7)$$

(iii) for any $x, y \in \mathbb{R}^d$,

$$|c(x) - c(y)| \leq \Lambda U_p(x, y) \quad (1.8)$$

where

$$U_p(x, y) = (|x| + |y|)^{p-1}|x - y|.$$

If the choice of p is clear from the context, we will write $V = V_p$ and $U = U_p$. We further note the following elementary inequality, valid for any $z_1, z_2 \in \mathbb{R}^d$ and with implicit constants depending only on p and d ,

$$|V_p(z_1) - V_p(z_2)| \lesssim (|z_1| + |z_2|)^{p-1}|z_1 - z_2|. \quad (1.9)$$

Note that (1.6) and the fact that c is C^1 imply that for any $x, y \in \mathbb{R}^d$,

$$c(x) \geq c(y) + \langle \nabla c(y), x - y \rangle + \lambda V(x, y) \quad (1.10)$$

$$\langle \nabla c(x) - \nabla c(y), x - y \rangle \geq \lambda V(x, y). \quad (1.11)$$

Introduce the convex conjugate c^* defined on \mathbb{R}^d via

$$c^*(y) = \sup_x \langle y, x \rangle - c(x).$$

Note that due to (1.7), we have

$$|z|^{p'} \lesssim c^*(z) \lesssim |z|^{p'}. \quad (1.12)$$

Due to strict p -convexity of c , c^* satisfies

$$|\nabla c^*(x) - \nabla c^*(y)| \lesssim \begin{cases} (|x| + |y|)^{p'-2}|x - y| & \text{if } p \leq 2 \\ |x - y|^{p'-1} & \text{if } p \geq 2. \end{cases} \quad (1.13)$$

Finally, it follows from (1.12) and (1.13) that for $x, y \in \mathbb{R}^d$,

$$|c^*(x) - c^*(y)| \lesssim U_{p'}(x, y). \quad (1.14)$$

In addition to (1.6)-(1.8), which are standard assumptions quantifying the convexity, smoothness and growth of c , we need to ensure that also c^* is p -convex. This is ensured by requiring a slightly non-standard growth assumption on the Lipschitz constant of c . To be precise, we assume controlled duality p -growth on c , that is

$$|\nabla c(x) - \nabla c(y)| \leq \begin{cases} \Lambda(|\nabla c(x)| + |\nabla c(y)|)^{\frac{p-2}{p-1}}|x - y| & \text{if } p \geq 2 \\ \Lambda|x - y|^{p-1} & \text{if } p \leq 2. \end{cases} \quad (1.15)$$

Assuming (1.15), c^* is p' -convex, that is for some $c = c(p, \Lambda) > 0$,

$$c\tau(1 - \tau)V_{p'}(x, y) + c^*(\tau x + (1 - \tau)y) \leq \tau c^*(x) + (1 - \tau)c^*(y). \quad (1.16)$$

We remark that (1.15) implies the more standard controlled p -growth condition

$$|\nabla c(x) - \nabla c(y)| \lesssim \begin{cases} \Lambda(|x| + |y|)^{p-2}|x - y| & \text{if } p \geq 2 \\ \Lambda|x - y|^{p-1} & \text{if } p \leq 2. \end{cases}$$

Further, we point out that (1.15) is satisfied by polynomial cost functions as well as p -cost [1].

To close this section, we note that $\nabla c^* = (\nabla c)^{-1}$ and recall the Fenchel-Young inequality in the form

$$c(\xi) + c^*(x) \geq \langle \xi, x \rangle \quad \forall \xi, x \in \mathbb{R}^d, \quad (1.17)$$

with equality if and only if $\xi = \nabla c^*(x)$.

1.2 Regularity assumptions on the dual system

In this section, we state the regularity assumptions we make on distributional solutions $\phi \in W^{1,p'}(B)$ of the equation

$$-\operatorname{div} \nabla c^*(D\phi) = c_g \quad \text{on } B \quad (1.18)$$

$$\nabla c^*(D\phi) \cdot \nu = g \quad \text{on } \partial B, \quad (1.19)$$

where $g \in L^p(B)$ and c_g satisfies the compatibility condition $|B|c_g = \int_{\partial B} g$. ν denotes the outward pointing normal vector on ∂B . Note that solutions are only defined up to a constant. Hence, we usually normalise solutions by requiring that $\int_B \phi = 0$.

We assume that solutions to (1.18) exist, are unique up to constant and moreover satisfy the energy estimate

$$\int_{B_R} |D\phi|^{p'} dx \lesssim \int_{\partial B_R} |g|^p. \quad (1.20)$$

Using (1.17), (1.12) and Young's inequality, (1.20) implies that also

$$\int_{B_R} c(\nabla c^*(D\phi)) dx \lesssim \int_{\partial B_R} |g|^p. \quad (1.21)$$

We further assume interior regularity of solutions: for $r < R$,

$$\sup_{x \in B_r} |D\phi|^{p'} \lesssim_{R-r} \int_{\partial B_R} |g|^p. \quad (1.22)$$

Suppose ϕ^r solves (1.18) with data g^r , where g^r denotes convolution with a smooth convolution kernel on ∂B at scale r . Then we require

$$\int_B |D\phi - D\phi^r|^{p'} \lesssim \begin{cases} r \int_{\partial B} |g|^p & \text{if } p \leq 2 \\ r^{\frac{p'}{2}} \int_{\partial B} |g|^p & \text{if } p \geq 2 \end{cases} \quad (1.23)$$

Denote by g^r the convolution of g with a smooth convolution kernel at scale r on ∂B . Denote by ϕ^r the solution of (1.18) with data g^r . Then we require that for some $\beta \in (0, 1)$,

$$r^\beta [D\phi^r]_{C^{0,\beta}(B)}^{p'} + \sup_B |D\phi^r|^{p'} \lesssim \frac{1}{r^{d-1}} \int_{\partial B} |g|^p. \quad (1.24)$$

Moreover, we note by direct calculation using (1.8), (1.14) and (1.13) that

$$[c^*(D\phi) + c(\nabla c^*(D\phi))]_{C^{0,\beta}(B)} \lesssim \|D\phi\|_{L^\infty(B)}^{p'-1} [D\phi]_{C^{0,\beta}(B)}.$$

We note that, if c satisfies controlled-duality p -growth (1.15), (or alternatively, making the slightly weaker assumption that c^* is strictly p' -convex, that is (1.16)) our assumptions are satisfied.

Lemma 1.2. *If c^* satisfies (1.16) and (1.6)-(1.8) hold, then solutions to (1.18) exist and moreover (1.20)-(1.24) are satisfied.*

Proof. As c^* is p' -convex, by the direct method, solutions in $\phi \in W^{1,p'}(B)$ exist and are unique up to constant. (1.20) follows immediately from testing the equation with ϕ and using Young's inequality, the trace estimate and Poincaré's inequality. In order to see (1.23), test the equations for ϕ and ϕ^r against $\phi - \phi^r$ and use the p' -convexity, duality, trace estimate and Poincaré's inequality to see

$$\begin{aligned} \int_B V(D\phi, D\phi^r) &\leq \int_B \langle \nabla c^*(D\phi) - \nabla c^*(D\phi^r), D\phi - D\phi^r \rangle \\ &= \int_{\partial B} (g - g^r)(\phi - \phi^r) \\ &\leq \|g - g^r\|_{W^{-\frac{1}{p},p}(\partial B)} \|\phi - \phi^r\|_{W^{\frac{1}{p},p'}(\partial B)} \\ &\lesssim r^{\frac{1}{p}} \|g\|_{L^p(\partial B)} \|\phi - \phi^r\|_{W^{1,p'}(B)}. \end{aligned}$$

If $p' \geq 2$, this gives the result immediately after re-arranging. If $p' \leq 2$, we note

$$\int_B V(D\phi, D\phi') \geq \left(\int_B |D\phi - D\phi'|^{p'} \right)^{\frac{2}{p'}} \left(\int_B |D\phi|^{p'} + |D\phi'|^{p'} \right)^{\frac{p'-2}{p'}},$$

so that the claimed inequality follows after re-arranging and using (1.20).

(1.22) and (1.24) follows from [14] and [13]. \square

2 Preliminaries

2.1 General notation

Throughout, we let $1 < p < \infty$. $B_r(x)$ will denote a ball of radius $r > 0$ centered at $x \in \mathbb{R}^d$. We further write $B_r = B_r(0)$ and $B = B_1(0)$. c denotes a generic constant that may change from line to line. Relevant dependencies on Λ , say, will be denoted $c(\Lambda)$. We say $a \lesssim b$, if there exists a constant $c > 0$ depending only on d, p and Λ such that $a \leq cb$.

Given $\Omega \subset \mathbb{R}^d$, we denote by $[\cdot]_{C^{0,\alpha}}$, the α -Hölder-seminorm. Given $\alpha \in (0, \infty)$, $L^p(\Omega)$ and $W^{\alpha,p}(\Omega)$ denote the usual Lebesgue and (fractional) Sobolev spaces. If μ is a measure on \mathbb{R}^d , $\mu_\perp \Omega$ denotes its restriction to Ω .

Given $R > 0$, we let $\Pi_R(x) = R \frac{x}{|x|}$ be the projection onto ∂B_R and define for every measure ρ on \mathbb{R}^d the projected measure on ∂B_R , $\hat{\rho} = \Pi_R \# \rho$, i.e.

$$\int \xi \, d\hat{\rho} = \int \xi \left(R \frac{x}{|x|} \right) \, d\rho(x).$$

A set $\Omega \subset \mathbb{R}^d \times \mathbb{R}^d$ is said to be c -cyclically monotone if for any $N \in \mathbb{N}$ and any points $(x_1, y_1), \dots, (x_N, y_N) \in \Omega$, there holds

$$\sum_{i=1}^N c(x_i - y_i) \leq \sum_{i=1}^N c(x_i - y_{i+1}),$$

where we identify $y_{N+1} = y_1$.

A function $f: \mathbb{R}^d \rightarrow \mathbb{R}$ is called c -concave if there exists a function $g: \mathbb{R}^d \rightarrow \mathbb{R}$ such that

$$f(x) = \inf_{y \in \mathbb{R}^d} c(x - y) - g(y)$$

for all $x \in \mathbb{R}^d$.

2.2 Optimal transportation

We recall some definitions and facts about optimal transportation, see [18] for more details. Given a measure π on $\mathbb{R}^d \times \mathbb{R}^d$ we denote its marginals by π_1 and π_2 respectively. The set of measures on $\mathbb{R}^d \times \mathbb{R}^d$ with marginals π_1 and π_2 is denoted $\Pi(\pi_1, \pi_2)$. Given two positive measures with compact support and equal mass λ and μ we define

$$W_c(\lambda, \mu) = \min_{\pi_1=\lambda, \pi_2=\mu} \int c(x - y) d\pi.$$

While our notation is reminiscent of the Wasserstein distance, and in fact gives the Wasserstein p -distance in the case $c(x - y) = |x - y|^p$, in general it is not a distance on measures. Under our hypothesis, an optimal coupling always exists and moreover a coupling π is optimal, if and only if its support is c -cyclical monotone.

Moreover, we note the following triangle-type inequality:

Lemma 2.1. *Let $\varepsilon \in (0, 1)$. There is $c(\varepsilon) > 0$ such that for any admissible measures μ_1, μ_2, μ_3 it holds that*

$$W_c(\mu_1, \mu_3) \leq (1 + \varepsilon)W_c(\mu_1, \mu_2) + c(\varepsilon)W_c(\mu_2, \mu_3).$$

Proof. Due to the gluing lemma, see e.g. [17, Lemma 5.5.], there exists σ , a positive measure on $\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d$ with marginal π_1 on the first two variables and marginal π_2 on the last two variables. Here π_1 and π_2 are the optimal couplings between μ_1 and μ_2 and μ_2 and μ_3 , respectively, with respect to W_c . Set γ to be the marginal of σ with respect to the first and third variable. Then $\gamma \in \Pi(\mu_1, \mu_3)$. It follows using the convexity of c and the triangle inequality in $L^p(\gamma)$ that for any $t \in (0, 1)$,

$$\begin{aligned} W_c(\mu_1, \mu_3) &\leq \left(\int c(x - z) d\gamma \right)^{\frac{1}{p}} \leq \left(\int tc \left(\frac{x - y}{t} \right) + (1 - t) \left(\frac{y - z}{1 - t} \right) d\gamma \right)^{\frac{1}{p}} \\ &\leq \left(\int \left(\left(tc \left(\frac{x - y}{t} \right) \right)^{\frac{1}{p}} + \left((1 - t) \left(\frac{y - z}{1 - t} \right) \right)^{\frac{1}{p}} \right)^p d\gamma \right)^{\frac{1}{p}} \\ &\leq \left(\int tc \left(\frac{x - y}{t} \right) d\gamma \right)^{\frac{1}{p}} + \left(\int (1 - t) c \left(\frac{y - z}{1 - t} \right) d\gamma \right)^{\frac{1}{p}}. \end{aligned}$$

Using (1.7) and recalling the definition of γ , we deduce

$$W_c(\mu_1, \mu_3) \leq \left(\Lambda^2 t^{1-p}\right)^{\frac{1}{p}} W_c(\mu_1, \mu_2) + \left(\Lambda^2 (1-t)^{1-p}\right)^{\frac{1}{p}} W_c(\mu_2, \mu_3).$$

Choosing t sufficiently close to 1, this gives the desired estimate. \square

We require also the following consequence of Lemma 2.1.

Corollary 2.2. *Let μ_1, μ_2 be measures. Then*

$$W_c(\mu_1, \mu_2) \lesssim W_c(\mu_1 + \mu_2, 2\mu_2).$$

Proof. Using Lemma 2.1 and sub-additivity of W_c , we note for any $\delta > 0$,

$$\begin{aligned} W_c(\mu_1, \mu_2) &\leq (1 + \delta) W_c\left(\mu_1, \frac{1}{2}(\mu_1 + \mu_2)\right) + c(\delta) W_c\left(\frac{1}{2}(\mu_1 + \mu_2), \mu_2\right) \\ &= (1 + \delta) W_c\left(\frac{1}{2}\mu_1, \frac{1}{2}\mu_2\right) + c(\delta) W_c\left(\frac{1}{2}(\mu_1 + \mu_2), \mu_2\right) \\ &\leq \frac{1 + \delta}{2} W_c(\mu_1, \mu_2) + c(\delta) W_c(\mu_1 + \mu_2, 2\mu_2). \end{aligned}$$

Re-arranging gives the result. \square

Given $O \subset \mathbb{R}^d$ set $\kappa_{\mu, O}$ to be the generic constant such that $W_c(\mu \llcorner O, \kappa_{\mu, O} dx \llcorner O)$ is well-defined, that is $\kappa_{\mu, O} = \frac{\mu(O)}{|O|}$. If $O = B_R$, we write $\kappa_{\mu, R} = \kappa_{\mu, B_R}$.

It will be convenient to denote $\#_R = (B_R \times \mathbb{R}^d) \cup (\mathbb{R}^d \times B_R)$. We recall the definition of the quantities that we use to measure smallness:

$$\begin{aligned} E(R) &:= \frac{1}{|B_R|} \int_{\#_R} c(x - y) d\pi \\ D(R) &:= \frac{1}{|B_R|} W_p^p(\lambda \llcorner B_R, \kappa_{\lambda, R} dx \llcorner B_R) + \frac{R^p}{\kappa_{\lambda, R}^{p-1}} (\kappa_{\lambda, R} - 1)^p \\ &\quad + \frac{1}{|B_R|} W_p^p(\mu \llcorner B_R, \kappa_{\mu, R} dx \llcorner B_R) + \frac{R^p}{\kappa_{\mu, R}^{p-1}} (\kappa_{\mu, R} - 1)^p. \end{aligned}$$

We will find it convenient to work with trajectories $X(t) = tx + (1-t)y$. In this context, it is useful to work on the domain

$$\Omega_R = \{(x, y) \in \#_3 : \exists t \in [0, 1] \text{ s.t. } X(t) \in \overline{B_R}\}.$$

To every trajectory $X \in \Omega$, we associate entering and exiting times of B_R :

$$\begin{aligned} \sigma_R &:= \min\{t \in [0, 1] : X(t) \in \overline{B_R}\} \\ \tau_R &:= \max\{t \in [0, 1] : X(t) \in \overline{B_R}\}. \end{aligned}$$

Often, we will drop the subscripts and denote $\Omega = \Omega_R$, $\sigma = \sigma_R$ and $\tau = \tau_R$. Further, we will need to track trajectories entering and leaving B_R . This is achieved through the non-negative measures f_R and g_R concentrated on ∂B_R and defined by the relations

$$\int \zeta df_R = \int_{\Omega \cap \{X(\sigma) \in \partial B_R\}} \zeta(X(\sigma)) d\pi,$$

$$\int \zeta dg_R = \int_{\Omega \cap \{X(\tau) \in \partial B_R\}} \zeta(X(\sigma)) d\pi. \quad (2.1)$$

Note that the set of trajectories $\Omega \cap \{X(\sigma) \in \partial B_R\}$ implicitly defines a Borel measurable subset of $\mathbb{R}^d \times \mathbb{R}^d$, namely the pre-image under the mapping $(x, y) \rightarrow X$, which is continuous from $\mathbb{R}^d \times \mathbb{R}^d$ into $C^0([0, 1])$. Thus, the integrals in (2.2) are well-defined. We will often use similar observations without further justification.

2.3 Estimating radial projections

We record a technical estimate concerning radial projections we will require.

Lemma 2.3. *For $R > 0$, there exists $1 \geq \varepsilon(d) > 0$ such that for every $g \geq 0$ with $\text{Spt } g \subset B_{(1+\varepsilon)R} \setminus B_{(1-\varepsilon)R}$ we have*

$$R^{1-d} \left(\int g \right)^p \lesssim \int_{\partial B_R} \hat{g}^p \lesssim \sup g^{p-1} \int |R - |x||^{p-1} g$$

Proof. By scaling we may assume $R = \sup g = 1$. The first inequality is then a direct consequence of Jensen's inequality.

For the second inequality, note that if $\varepsilon \ll 1$, $\sup_{\partial B_1} |\hat{g}| \ll 1$, since we assume $\text{Spt } g \subset B_{1+\varepsilon} \setminus B_{1-\varepsilon}$. Fix $\omega \in \partial B_1$ and set $\psi(r) = r^{d-1} g(r\omega)$ for $r > 0$. Then we have $0 \leq \psi \leq (1 + \varepsilon)^{d-1} \leq 2$ and

$$\int_0^\infty \psi = \hat{g}(\omega).$$

We conclude that for $\omega \in \partial B_1$,

$$\int_0^\infty |1 - r|^{p-1} r^{d-1} g(r\omega) \geq \min_{0 \leq \tilde{\psi} \leq 2, \int \tilde{\psi} = \hat{g}(\omega)} \int_0^\infty |1 - r|^{p-1} \tilde{\psi}(r) \gtrsim \hat{g}(\omega).$$

The last inequality holds, since the minimiser of

$$\min_{0 \leq \tilde{\psi} \leq 2, \int \tilde{\psi} = \hat{g}(\omega)} \int_0^\infty |1 - r|^{p-1} \tilde{\psi}(r)$$

is given by $2I(|r - 1| \leq \frac{1}{4} \hat{g}(\omega))$. □

3 A L^∞ -bound on the displacement

A key point in our proof will be that trajectories do not move very much. Since we assume $E(4) \ll 1$, this is evidently true on average. However, we will require to control the length of trajectories not just on average, but in a pointwise sense. We establish this result in this section. In the quadratic case, the proof in [5] relies on the fact that 2-monotonicity is equivalent to standard monotonicity. In our setting this is not available and we hence provide a different proof.

Lemma 3.1. *Let $1 < p < \infty$. Let π be a coupling between two measures λ and μ . Assume that $\text{Spt } \pi$ is cyclically monotone with respect to c -cost and that $E(4) + D(4) \ll 1$. Then for every $(x, y) \in \text{Spt } \pi \cap \#_3$, we have*

$$|x - y| \lesssim (E(4) + D(4))^{\frac{1}{p+d}}. \quad (3.1)$$

As a consequence, for $(x, y) \in \text{Spt } \pi$ and $t \in [0, 1]$,

$$x \in B_3 \text{ or } y \in B_3 \Rightarrow (1-t)x + ty \in B_4. \quad (3.2)$$

In the proof of Lemma 3.1 we require the following technical result, which we state independently as we will require it again in the future.

Lemma 3.2. *Let $1 < p < \infty$ and $0 < \alpha < 1$. For every $R > 0$, $\xi \in C^{0,\alpha}(B_R)$ and μ supported in B_R with $\mu(B_R) \sim |B_R|$,*

$$\left| \int_{B_R} \xi(d\mu - \kappa_{\mu,R} dx) \right| \leq [\xi]_{C^{0,\alpha}(B_R)} W_c(\mu, \kappa_{\mu,R} dx \llcorner B_R)^{\frac{\alpha}{p}} R^{\frac{2d(p-\alpha)}{p}}. \quad (3.3)$$

In case $\alpha = 1$, (3.3) holds with $C^{0,1}$ replaced by C^1 . Further, if in addition we have $\xi \in C^{[p-1], p-[p-1]}(B_R)$, there is $C > 0$ such that

$$\begin{aligned} \left| \int_{B_R} \xi(d\mu - \kappa_{\mu,R} dx) \right| &\leq C \sum_{i=1}^{[p-1]} \left(\kappa_{\mu,R} \int |D^i \xi|^{\frac{p}{p-i}} dx \right)^{\frac{p-i}{p}} W_c(\mu, \kappa_{\mu,R} dx \llcorner B_R)^{\frac{i}{p}} \\ &\quad + [\xi]_{C^{[p-1], p-[p-1]}(B_R)} W_c(\mu, \kappa_{\mu,R} dx \llcorner B_R). \end{aligned}$$

Proof. Integrate the estimate

$$|\xi(x) - \xi(y)| \leq [\xi]_{C^{0,\alpha}} |x - y|^\alpha$$

against an optimal transport plan π between μ and $\kappa_{\mu,R} dx \llcorner B_R$ to find,

$$\left| \int_{B_R} \xi(d\mu - \kappa_{\mu,R} dx) \right| \leq [\xi]_{C^{0,\alpha}(B_R)} \int_{B_R} |x - y|^\alpha d\pi.$$

Applying Hölder and using (1.7) the result follows.

To obtain the second estimate, we proceed similarly, but start with the estimate

$$|\xi(x) - \xi(y) - \sum_{|\alpha|=1}^{[p-1]} D^\alpha \xi(y) \frac{(x-y)^\alpha}{|\alpha|!}| \leq [\xi]_{C^{[p-1], p-[p-1]}(B_R)} |x - y|^p.$$

The result follows using (1.7) and using Hölder to estimate

$$\int |D^\alpha \xi(y)| |x - y|^{|\alpha|} d\pi \leq \left(\kappa_{\mu,R} \int |D^{|\alpha|} \xi|^{\frac{p}{p-|\alpha|}} dx \right)^{\frac{p-|\alpha|}{p}} W_c(\mu, \kappa_{\mu,R} dx \llcorner B_R)^{\frac{|\alpha|}{p}}.$$

□

We proceed to prove Lemma 3.1.

Proof of Lemma 3.1. Fix $(x, y) \in \text{Spt } \pi \cap \#_3$. Without loss of generality we may assume that $(x, y) \in B_3 \times \mathbb{R}^d$.

Step 1. Barrier points exist in all directions: In this step we show that in all directions we may find points $(x', y') \in \text{Spt } \pi$ with $x' \approx y'$. To be precise, consider an arbitrary

unit vector $\mathbf{n} \in \mathbb{R}^d$ and let $r > 0$. We show that for any \mathbf{n} , and all $r \ll 1$, there is $M = M(p, d, \Lambda) > 0$ and $(x', y') \in \text{Spt } \pi \cap (B_r(x + 2r\mathbf{n}) \times \mathbb{R}^d)$ such that

$$c(x' - y') \leq \frac{ME(4)}{r^d}.$$

Assume, for contradiction, that for any $M > 0$, there is $\mathbf{n} \in \mathbb{R}^d$ and $r > 0$ such that for all $(x', y') \in \text{Spt } \pi \cap (B_r(x + 2r\mathbf{n}) \times \mathbb{R}^d)$, $|x' - y'| \geq \frac{ME(4)}{r^d}$. Let η be a non-negative, smooth cut-off supported in $B_r(x + 2r\mathbf{n})$ satisfying

$$\sum_{i=1}^{\lfloor p-1 \rfloor} r^i \sup |D^i \eta| + r^p [\eta]_{C^{\lfloor p-1 \rfloor, p-\lfloor p-1 \rfloor}} \lesssim 1.$$

Then

$$E(4) \gtrsim \int \int \eta(x) c(x - y) d\pi(x, y) \geq \int \int \frac{ME(4)}{r^d} \eta(x) d\pi(x, y) = \frac{ME(4)}{r^d} \int \eta(x) d\mu(x).$$

However, due to Lemma 3.2 and noting $\kappa_{\mu,4} \sim 1$,

$$\left| \int \eta d\mu(x) - \kappa_{\mu,4} \int \eta dx \right| \lesssim \sum_{i=1}^{\lfloor p-1 \rfloor} r^{\frac{d(p-i)}{p}-i} D(4)^{\frac{i}{p}} + r^{-p} D(4).$$

Normalising η such that $\int_{B_r(x+2r\mathbf{n})} \eta dx \sim r^d$, we can guarantee $\kappa_{\mu,4} \int \eta dx \sim \kappa_{\mu,4} r^d \sim r^d$.

Ensuring $D(4) \ll r^{p+d}$, so that $\sum_{i=1}^{\lfloor p-1 \rfloor} r^{\frac{d(p-i)}{p}-i} D(4)^{\frac{i}{p}} + r^{-p} D(4) \ll r^d$, we may thus conclude

$$E(4) \gtrsim \frac{ME(4)}{r^d} r^d = ME(4).$$

As M was arbitrary, this is a contradiction.

Step 2. Building barriers: In this step, we show that if we are given points

$$(x', y') \in \text{Spt } \pi \cap (B_r(x + 2r\mathbf{n}) \times \mathbb{R}^d)$$

such that $|x' - y'| \leq \frac{ME(4)}{r^d}$ for some $M = M(p, d, \Lambda) > 0$, then there is a cone $C_{x,x'}$ with vertex $x' + r\rho(x' - x)$ for some $\rho = \rho(p, d, \Lambda) > 0$, aperture $\alpha = \alpha(p, d, \Lambda)$ and axis $x' - x$ such that $y \notin C_{x,x'}$.

Without loss of generality, we may assume that $x' - x$ points in the e_n direction. Moreover, considering the cost $c(\cdot) - c(x)$, we may assume that $c(x) = 0$. Suppose for a contradiction that

$$y \in C_{x,x'} = x' + \{a \in \mathbb{R}^{d-1} \times \mathbb{R}^+ : d(a, \Gamma) \leq \alpha(|\bar{a} - x'| - r\rho)\}$$

for some $\alpha, \rho > 0$ to be determined. Here $\Gamma = \{t(x' - x) : t \geq 0\}$ and \bar{a} denotes the orthogonal projection of a point $a \in \mathbb{R}^{d-1} \times \mathbb{R}^+$ onto Γ . We want to show that then

$$c(x - y) \geq c(x' - y) + c(x - y'). \quad (3.4)$$

(3.4) is a contradiction to the c -monotonicity of π and hence proves the stated claim.




Figure 1: Geometric situation in Step 2.

We note that we may assume $x = 0$. Indeed, setting $z = y - x$, $z' = y' - x$ and $\tilde{z} = x' - x$, (3.4) becomes

$$c(-z) \geq c(\tilde{z} - z) + c(-z')$$

with $|\tilde{z}| \leq \frac{ME}{r^d}$ and $z \in C_{0,\tilde{z}}$, which we recognise as precisely the situation we are in if $x = 0$.

Taking $\rho \geq 4$, we then estimate using (1.8) and (1.6)

$$\begin{aligned}
c(-y) &\geq c(-\bar{y}) + c(x' - y') - \Lambda U(-y, -\bar{y}) \\
&\geq \frac{|\bar{y}|}{|-\bar{y} + x'|} c(-\bar{y} + x') + \frac{\lambda|x'|}{|\bar{y}|} V(-\bar{y}, 0) - \Lambda U(-y, -\bar{y}) \\
&\geq c(-\bar{y} + x') + c(x') + \frac{\lambda|x'|}{|\bar{y}|} V(-\bar{y}, 0) + \frac{\lambda|\bar{y} - 2x'|}{|\bar{y} - x'|} V(-\bar{y} + x', 0) - \Lambda U(-y, -\bar{y}) \\
&\geq c(-y + x') + c(-x') - \Lambda U(-y + x', -\bar{y} + x') + \frac{\lambda|x'|}{|\bar{y}|} V(-\bar{y}, 0) \\
&\quad + \frac{\lambda|\bar{y} - 2x'|}{|\bar{y} - x'|} V(-\bar{y} + x', 0) - \Lambda U(-y, -\bar{y}) \\
&\geq c(-y + x') + c(-y') - \Lambda U(-y + x', -\bar{y} + x') + \frac{\lambda|x'|}{|\bar{y}|} V(-\bar{y}, 0) \\
&\quad + \frac{\lambda|\bar{y} - 2x'|}{|\bar{y} - x'|} V(-\bar{y} + x', 0) - \Lambda U(-y, -\bar{y}) - \Lambda U(-x', -y')
\end{aligned}$$

In particular, it suffices to show

$$\begin{aligned}
&c(x' - y') + \frac{\lambda|x'|}{|\bar{y}|} V(-\bar{y}, 0) + \frac{\lambda|\bar{y} - 2x'|}{|\bar{y} - x'|} V(-\bar{y} + x', 0) \\
&\geq \Lambda U(-y, -\bar{y}) + \Lambda U(-y + x', -\bar{y} + x').
\end{aligned} \tag{3.5}$$

We note that, if $\rho \geq 8$, $|\bar{y} - 2x'| \geq \frac{1}{2}|\bar{y}|$. Then we can estimate

$$\begin{aligned} \frac{\lambda|x'|}{|\bar{y}|}V(-\bar{y}, 0) + \frac{\lambda|\bar{y} - 2x'|}{|\bar{y} - x'|}V(-\bar{y} + x', 0) &= \lambda(|x'|\bar{y}|^{p-1} + |\bar{y} - 2x'|\bar{y} - x'|^{p-1}) \\ &\gtrsim |\bar{y}|^p. \end{aligned}$$

Further, if $E(4) \leq \varepsilon r^{d+1}$,

$$\begin{aligned} &\Lambda U(-y, -\bar{y}) + \Lambda U(-y + x', -\bar{y} + x') + \Lambda U(-x', -y') \\ &\leq 2\Lambda(2|y|)^{p-1}|y - \bar{y}| + \Lambda(|x'| + |y'|)^{p-1}|x' - y'| \\ &\lesssim |y|^{p-1}\alpha|\bar{y} - x'| + \frac{ME}{r^d}(2r)^{p-1} \\ &\lesssim \alpha|\bar{y}|^p + \varepsilon|\bar{y}|^p. \end{aligned}$$

Thus choosing $\alpha, \varepsilon > 0$, sufficiently small, we find (3.5) holds, proving our claim.

Step 3. Proving the L^∞ -bounds: Choose $r = c(E(4) + D(4))^{\frac{1}{p+d}}$. For sufficiently large choice of $c > 0$ and selecting $c(d)$ directions n_i , by Step 1 and Step 2 we obtain points

$$(x', y') \in \text{Spt } \pi \cap (B_r(x + 2rn_i) \times \mathbb{R}^d)$$

and cones $(C_{x, x'_i})_{i \leq c(d)}$ with vertices $x'_i + r\rho(x'_i - x)$, aperture α and axis $x'_i - x$ such that for some $c(\alpha) > 0$,

$$y \notin \cup C_{y_i} \text{ and } \mathbb{R}^d \setminus B_{(\rho+c(\alpha))r}(x) \subset \cup C_{y_i}.$$

In particular, we have

$$|y - x| \leq (\rho + c(\alpha))r \lesssim (E + D)^{\frac{1}{p+d}},$$

that is (3.1).

(3.2) is a direct consequence of (3.1), concluding the proof. \square

We record two consequences of Lemma 3.1 we will use later.

Corollary 3.3. *Under the assumptions of Lemma 3.1, it holds that*

$$\begin{aligned} \int_2^3 \int_{\Omega \cap \{\exists t \in [0,1]: X(t) \in \partial B_R\}} c(x - y) \, d\pi \, dR &\lesssim (E(4) + D(4))^{1 + \frac{1}{p+d}}, \\ \int_2^3 \int_{\Omega} I(\{\exists t \in [0,1]: X(t) \in \partial B_R\}) \, d\pi \, dR &\lesssim (E(4) + D(4))^{\frac{1}{p+d}}. \end{aligned}$$

Proof. We use Lemma 3.1 to deduce there is $C > 0$ such that

$$\begin{aligned} &\int_2^3 \int_{\Omega \cap \{\exists t \in [0,1]: X(t) \in \partial B_R\}} c(x - y) \, d\pi \, dR \\ &\leq \int_2^3 \int_{(B_{7/2} \setminus B_{3/2}) \times (B_{7/2} - B_{3/2})} I(\{|x| - R| \leq C(E(4) + D(4))^{\frac{1}{p+d}}\}) c(x - y) \, d\pi \, dR \\ &\lesssim (E(4) + D(4))^{\frac{1}{p+d}} \int_{\#_4} c(x - y) \, d\pi \end{aligned}$$

$$\lesssim (E(4) + D(4))^{1 + \frac{1}{p+d}}.$$

Further, again using Lemma 3.1, there is $C > 0$ such that,

$$\begin{aligned} & \int_2^3 \int_{\Omega} I(\{\exists t \in [0, 1]: X(t) \in \partial B_R\}) \, d\pi \, dR \\ & \leq \int_2^3 \pi(\Omega \cap \{|X(0) - R| \leq C(E(4) + D(4))^{\frac{1}{p+d}}\}) \, dR \\ & = \int_2^3 \mu(\{|x| - R| \leq C(E(4) + D(4))^{\frac{1}{p+d}}\}) \, dR \\ & \leq \int_{B_{7/2} \setminus B_{3/2} \times B_{7/2} \setminus B_{3/2}} I(|x| - R| \leq C(E(4) + D(4))^{\frac{1}{p+d}}) \, dR \, d\mu \\ & \lesssim (E(4) + D(4))^{\frac{1}{p+d}} \mu(B_4) \\ & \lesssim (E(4) + D(4))^{\frac{1}{p+d}}. \end{aligned}$$

□

4 A localisation result

In order to prove the approximation of the optimal transport plan π by a c^* -harmonic map, we will need to use optimality in a localised way. In general, it is not true that the cost of π localised to $\#_R$ is well approximated by the transportation cost between the localised measures $\lambda_{\perp B_R}$ and $\mu_{\perp B_R}$. However, if we take into account the entering points f_R and exiting points g_R , this approximation holds as we show in the next lemma.

Lemma 4.1. *Suppose $\pi \in \Pi(\lambda, \mu)$ minimises (1.1). Let $R \in [2, 3]$. Then for any $\tau, \delta > 0$, there is $\varepsilon > 0$ such that if $E(4) + D(4) \leq \varepsilon$, then*

$$\int_{\Omega} c(x - y) \, d\pi \leq (1 + \delta) W_c(\lambda_{\perp B_R} + f_R, \mu_{\perp B_R} + g_R) + \tau (E(4) + D(4))$$

Proof. Introduce the weakly continuous family of probability measures $\{\lambda_z\}_{z \in \partial B_R}$ such that

$$\int_{\Omega \cap \{X(\sigma) \in \partial B_R\}} \zeta(x, X(\sigma)) \pi(dx \, dy) = \int_{\partial B_R} \int \zeta(x, z) \lambda_z(dx) f_R(dz)$$

for any test function ζ on $\mathbb{R}^d \times \mathbb{R}^d$. Likewise, introduce $\{\mu_w\}_{w \in \partial B_R}$ via

$$\int_{\Omega \cap \{X(\tau) \in \partial B_R\}} \zeta(X(\tau), y) \pi(dx \, dy) = \int_{\partial B_R} \int \zeta(w, y) \mu_w(dy) g_R(dw).$$

Let $\bar{\pi}$ be an optimal plan for $W_c(\lambda_{\perp B_R} + f_R, \mu_{\perp B_R} + g_R)$. Define a competitor $\tilde{\pi}$ for π by requiring the following formula to hold for any test function ζ on $\mathbb{R}^d \times \mathbb{R}^d$,

$$\begin{aligned} & \int \zeta(x, y) \tilde{\pi}(dx \, dy) \\ & = \int_{\Omega^c} \zeta(x, y) \, d\pi(x, y) + \int_{B_R \times B_R} \zeta(x, y) \bar{\pi}(dx \, dy) + \int_{\partial B_R \times B_R} \int \zeta(x, y) \lambda_z(dx) \bar{\pi}(dz \, dy) \end{aligned}$$

$$\begin{aligned}
& + \int_{B_R \times \partial B_R} \zeta(x, y) \mu_w(dy) \bar{\pi}(dx dw) + \int_{\partial B_R \times \partial B_R} \int \int \zeta(x, y) \mu_w(dy) \lambda_z(dx) \bar{\pi}(dz dw) \\
& = I + II + III + IV + V. \tag{4.1}
\end{aligned}$$

In order to see that $\tilde{\pi} \in \Pi(\lambda, \mu)$, by symmetry it suffices to check that the first marginal is λ . Hence test (4.1) against $\zeta(x)$. We begin by noting that due to the definition of μ_w and using that $\bar{\pi}$ is supported in \bar{B}_R ,

$$II + IV = \int_{B_R \times \mathbb{R}^d} \zeta(x) \bar{\pi}(dx dy) = \int_{B_R} \zeta(x) \mu(dx) = \int_{\Omega \cap \{X(\sigma) \in B_R\}} \zeta(x) \pi(dx dy).$$

Similarly, using also the definition of f_R ,

$$\begin{aligned}
III + V & = \int_{\partial B_R \times \mathbb{R}^d} \int \zeta(x) \lambda_z(dx) \bar{\pi}(dz dy) = \int_{\partial B_R} \zeta(z) f_R(dz) \\
& = \int_{\Omega \cap \{X(\sigma) \in \partial B_R\}} \zeta(x) \pi(dx dy).
\end{aligned}$$

In particular, we have shown

$$\int \zeta(x) \tilde{\pi}(dx dy) = \int \zeta(x) \pi(dx dy) = \int \zeta(x) \lambda(dx)$$

as desired.

Using optimality of π in the form

$$\int_{\Omega} c(x-y) d\pi + \int_{\Omega^c} c(x-y) d\pi \leq \int c(x-y) d\tilde{\pi}$$

and testing (4.1) against $\zeta(x, y) = c(x-y)$, we learn

$$\begin{aligned}
& \int_{\Omega} c(x-y) d\pi \\
& \leq \int_{B_R \times B_R} c(x-y) \bar{\pi}(dx dy) + \int_{\partial B_R \times B_R} \int c(x-y) \lambda_z(dx) \bar{\pi}(dz dy) \\
& \quad + \int_{B_R \times \partial B_R} c(x-y) \mu_w(dy) \bar{\pi}(dx dw) \\
& \quad + \int_{\partial B_R \times \partial B_R} \int \int c(x-y) \mu_w(dy) \lambda_z(dx) \bar{\pi}(dz dw) \\
& = \int_{B_R \times B_R} f_1 \bar{\pi}(dx dy) + \int_{\partial B_R \times B_R} f_2 \bar{\pi}(dz dy) + \int_{B_R \times \partial B_R} f_3 \bar{\pi}(dx dw) \\
& \quad + \int_{\partial B_R \times \partial B_R} f_4 \bar{\pi}(dz dw)
\end{aligned}$$

As in the proof of Lemma 2.1, for any $\delta > 0$, there is $C_\delta > 0$ such that for any x, y, z ,

$$c(x-z) \leq (1+\delta)c(x-y) + C_\delta c(y-z).$$

Using this in combination with the fact that λ_z, μ_w are probability measures we deduce

$$f_2 \leq (1+\delta)c(z-y) + C(\delta)\tilde{f}_2, \quad f_3 \leq (1+\delta)c(x-w) + C(\delta)\tilde{f}_3$$

$$f_4 \leq (1 + \delta)c(z - w) + C(\delta)\tilde{f}_4,$$

where

$$\begin{aligned}\tilde{f}_2(z, y) &= \int c(x - z)\lambda_z(\mathrm{d}x), \quad \tilde{f}_3(x, w) = \int c(w - y)\mu_w(\mathrm{d}y), \\ \tilde{f}_4(z, w) &= \tilde{f}_2(z, y) + \tilde{f}_3(x, w).\end{aligned}$$

In particular, we deduce

$$\begin{aligned}& \int_{\Omega} c(x - y) \mathrm{d}\pi \\ & \leq (1 + \delta) \int_{\overline{B}_R \times \overline{B}_R} c(x - y)\overline{\pi}(\mathrm{d}x \mathrm{d}y) + C(\delta) \int_{\partial B_R \times B_R} \tilde{f}_2\overline{\pi}(\mathrm{d}z \mathrm{d}y) \\ & \quad + C(\delta) \int_{B_R \times \partial B_R} \tilde{f}_3\overline{\pi}(\mathrm{d}x \mathrm{d}w) + C(\delta) \int_{\partial B_R \times \partial B_R} \tilde{f}_4\overline{\pi}(\mathrm{d}z \mathrm{d}w) \\ & = (1 + \delta) \int_{\overline{B}_R \times \overline{B}_R} c(x - y)\overline{\pi}(\mathrm{d}x \mathrm{d}y) + 2C(\delta) \int_{\partial B_R \times \mathbb{R}^d} \int c(x - z)\lambda_z(\mathrm{d}x)\overline{\pi}(\mathrm{d}z \mathrm{d}y) \\ & \quad + 2C(\delta) \int_{\mathbb{R}^d \times \partial B_R} c(w - y)\mu_y(\mathrm{d}y)\overline{\pi}(\mathrm{d}x \mathrm{d}w) \\ & = (1 + \delta) \int_{\overline{B}_R \times \overline{B}_R} c(x - y)\overline{\pi}(\mathrm{d}x \mathrm{d}y) + 2C(\delta) \int_{\partial B_R \times \mathbb{R}^d} \int c(x - z)\lambda_z(\mathrm{d}x)f_R(\mathrm{d}z) \\ & \quad + 2C(\delta) \int_{\mathbb{R}^d \times \partial B_R} c(w - y)\mu_y(\mathrm{d}y)g_R(\mathrm{d}w) \\ & = (1 + \delta) \int_{\overline{B}_R \times \overline{B}_R} c(x - y)\overline{\pi}(\mathrm{d}x \mathrm{d}y) + 2C(\delta) \int_{\Omega \cap \{X(\sigma) \in \partial B_R\}} c(x - X(\sigma)) \mathrm{d}\pi \\ & \quad + 2C(\delta) \int_{\Omega \cap \{X(\tau) \in \partial B_R\}} c(X(\tau) - y) \mathrm{d}\pi\end{aligned}$$

In order to obtain the second to last line we used the admissibility of $\overline{\pi}$. Now note on the one hand, that due to optimality of $\overline{\pi}$,

$$\int_{\overline{B}_R \times \overline{B}_R} c(x - y) \mathrm{d}\overline{\pi} = W_c(\lambda_{\perp} B_R + f_R, \mu_{\perp} B_R + g_R).$$

On the other hand, on $\Omega \cap \{X(\sigma) \in \partial B_R\} \cap \{X(\tau) \in \partial B_R\}$, for some $\rho_1, \rho_2 \geq 0$ with $\rho_1 + \rho_2 \leq 1$, due to convexity of c and $c(0) = 0$,

$$c(x - X(\sigma)) + c(X(\tau) - y) = c(\rho_1(x - y)) + c(\rho_2(x - y)) \leq c(x - y).$$

Thus, we have shown

$$\begin{aligned}& \int_{\Omega} c(x - y) \mathrm{d}\pi \\ & \leq (1 + \delta)W_c(\lambda_{\perp} B_R + f_R, \mu_{\perp} B_R + g_R) + C(\delta) \int_{\Omega \cap (\{X(\sigma) \in \partial B_R\} \cup \{X(\tau) \in \partial B_R\})} c(x - y) \mathrm{d}\pi\end{aligned}$$

$$\begin{aligned}
& = (1 + \delta)W_c(\lambda_{\perp}B_R + f_R, \mu_{\perp}B_R + g_R) + C(\delta) \int_{\Omega \cap \{\exists t \in [0,1]: X(t) \in \partial B_R\}} c(x - y) d\pi \\
& \leq (1 + \delta)W_c(\lambda_{\perp}B_R + f_R, \mu_{\perp}B_R + g_R) + C(\delta)(E(4) + D(4))^{1 + \frac{1}{p+d}}.
\end{aligned}$$

To obtain the last line, we used Corollary 3.3. Choosing ε sufficiently small the result follows. \square

5 Approximating the boundary data

Before we can implement the c^* -harmonic approximation, we face another problem. Lemma 4.1 suggests that we should impose Neumann boundary data f_R and g_R . However f_R and g_R are not sufficiently regular for this to make sense. Hence, we need to approximate them.

Lemma 5.1. *Let $\tau > 0$. There is $\varepsilon > 0$ such that if $E(4) + D(4) \leq \varepsilon$, then for every $R \in [2, 3]$ there exist non-negative functions \bar{f}_R, \bar{g}_R such that*

$$\begin{aligned}
W_c(f_R, \bar{f}_R) + W_c(g_R, \bar{g}_R) & \lesssim \tau E(4) + D(4) \\
\int_2^3 \int_{\partial B_R} \bar{g}_R^p + \bar{f}_R^p dR & \lesssim E(4) + D(4)
\end{aligned}$$

Proof. By symmetry it suffices to focus on the terms involving g .

Fix $R \in [2, 3]$. Let $\bar{\pi}$ be optimal for $W_c(\mu_{\perp}B_4, \kappa_{\mu,4} dz_{\perp}B_4)$. Extend $\bar{\pi}$, which is supported on $B_4 \times B_4$ by the identity to $\mathbb{R}^d \times \mathbb{R}^d$. Note that this extension, still denoted $\bar{\pi}$ is admissible for $W_c(\mu, \kappa_{\mu,4} dz_{\perp}B_4 + \mu_{\perp}B_4^c)$. Further due to the definition of D , $W_c(\mu_{\perp}B_4, \kappa_{\mu,4} dz_{\perp}B_4) \leq D(4)$. Introduce the family of probability measures $\{\bar{\pi}(\cdot|y)\}_{y \in \mathbb{R}^d}$ via asking that for every test function ζ

$$\int \zeta(y, z) \bar{\pi}(dz|y) \mu(dy) = \int \zeta(y, z) \bar{\pi}(dy dz).$$

Then define $\tilde{\pi}$ on $\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d$ by the formula

$$\int \zeta(x, y, z) \tilde{\pi}(dx dy dz) = \int \int \zeta(x, y, z) \bar{\pi}(dz|y) \pi(dx dy)$$

valid for any test function ζ . Note that with respect to the (x, y) variables $\tilde{\pi}$ has marginal π , while with respect to the (y, z) variables $\tilde{\pi}$ has marginal $\bar{\pi}$. Extend a trajectory X in Ω in a piecewise affine fashion by setting for $t \in [1, 2]$,

$$X(t) = (t - 1)z + (2 - t)y.$$

Note that the distribution g' of the endpoint of those trajectories that exit \bar{B}_R during the time interval $[0, 1]$ is given by

$$\int \zeta dg' = \int_{\Omega \cap \{X(\tau) \in \partial B_R\}} \zeta(z) \tilde{\pi}(dx dy dz). \tag{5.1}$$

Note that due to Lemma 3.1, $y = X(1) \in B_4$ for any trajectory X that contributes to (5.1). Since $\bar{\pi}(B_4, B_4^c) = 0$, we deduce that also $z = X(2) \in B_4$ and hence that g' is

supported in B_4 . In particular, we may estimate for any $\zeta \geq 0$, using the admissibility of $\bar{\pi}$ for $W_c(\mu, \kappa_{\mu,4} dz \llcorner B_4 + \mu \llcorner B_4^c)$,

$$\int \zeta dg' \leq \int_{\{z \in B_4\}} \zeta(z) \bar{\pi}(dy dz) = \kappa_{\mu,4} \int_{B_4} \zeta.$$

This shows that g' has a density, still denoted g' , satisfying $g' \leq \kappa_{\mu,4}$ and allows us to conclude the construction of \bar{g}_R by defining

$$\int \zeta d\bar{g}_R = \int \zeta \left(R \frac{z}{|z|} \right) g'(dz).$$

We now turn to establishing the claimed estimates for \bar{g}_R . Note that, directly from the definitions of $\tilde{\pi}$, g' and \bar{g}_R , an admissible plan for $W_c(g_R, \bar{g}_R)$ is

$$\int_{\Omega \cap \{X(\tau) \in \partial B_R\}} \zeta \left(X(\tau), R \frac{z}{|z|} \right) d\tilde{\pi}.$$

In particular, using (1.7),

$$W_c(g_R, \bar{g}_R) \lesssim \int_{\Omega \cap \{X(\tau) \in \partial B_R\}} c \left(X(\tau) - R \frac{z}{|z|} \right) d\tilde{\pi} \lesssim \int_{\Omega \cap \{X(\tau) \in \partial B_R\}} \left| X(\tau) - R \frac{z}{|z|} \right|^p d\tilde{\pi}.$$

Noting that $|X(\tau) - R \frac{z}{|z|}| \leq 2|X(\tau) - z|$, we deduce

$$\left| X(\tau) - R \frac{z}{|z|} \right|^p \lesssim |x - y|^p + |y - z|^p.$$

Thus, we deduce using (1.7) and Corollary 3.3,

$$W_c(g_R, \bar{g}_R) \lesssim \int_{\Omega \cap \{\exists t \in [0,1]: X(t) \in \partial B_R\}} c(x - y) d\pi + D(4) \lesssim E(4)^{1 + \frac{1}{p+d}} + D(4).$$

Choosing ε sufficiently small, the first estimate holds.

Noting $\sup g' \leq \kappa_{\mu,4} \lesssim 1$, in order to prove the second inequality, it suffices to prove

$$\int_2^3 \int |R - |x||^{p-1} dg' \lesssim E(4) + D(4)$$

and to apply Lemma 2.3. The condition on the support of g in Lemma 2.3 applies due to Lemma 3.1. Note that by definition of g' ,

$$\begin{aligned} \int |R - |x||^{p-1} dg' &= \int_{\Omega \cap \{X(\tau) \in \partial B_R\}} ||z| - R|^{p-1} \tilde{\pi}(dx dy dz) \\ &\lesssim \int_{\{y \in B_4\} \cap \{\min_{[0,1]} |X| \leq R \leq \max_{[0,1]} |X|\}} |x - y|^{p-1} + |y - z|^{p-1} \tilde{\pi}(dx dy dz). \end{aligned}$$

In order to obtain the second line, we observed that since $|X(\tau)| = R$, it holds that $||z| - R| \leq |x - y| + |y - z|$. In addition we noted that, since $X(\tau) \in \partial B_R$, we have

$\min_{[0,1]} X \leq R \leq \max_{[0,1]} X$ and $X(1) \in B_4$ due to Lemma 3.1. Integrating over R , this gives

$$\begin{aligned}
& \int_2^3 \int ||z| - R|^{p-1} dg' dR \\
& \lesssim \int_{\{y \in B_4\}} (\max_{[0,1]} X - \min_{[0,1]} X) (|x - y|^{p-1} + |y - z|^{p-1}) \tilde{\pi}(dx dy dz) \\
& \leq \int_{\{y \in B_4\}} |x - y| (|x - y|^{p-1} + |y - z|^{p-1}) \tilde{\pi}(dx dy dz) \\
& \lesssim \int_{\{y \in B_4\}} |x - y|^p \pi(dx dy) + \int |y - z|^p \bar{\pi}(dy dz) \\
& \leq E(4) + D(4).
\end{aligned}$$

The second-to last line was obtained applying Young's inequality. This concludes the proof. \square

6 Restricting the data

We also need to control $D(R)$, while at the moment we only control $D(4)$. Unfortunately, this does not follow immediately from the definition but requires a technical proof utilising ideas of the previous sections. The outcome of these considerations is the following lemma:

Lemma 6.1. *For any non-negative measure μ there is $\varepsilon > 0$ such that if $D(4) \leq \varepsilon$, then*

$$\int_2^3 \left(W_c(\mu \llcorner B_R, \kappa_{\mu,R} dx \llcorner B_R) + \frac{1}{\kappa_{\mu,R}} (\kappa_{\mu,R} - 1)^p \right) dR \lesssim D(4).$$

Proof. In this lemma π will denote the optimal transference plan for the problem $W_c(\mu \llcorner B_4, \kappa_{\mu,R} dx \llcorner B_4)$. Define the measures $0 \leq f' \leq \kappa_{\mu,4}$ on \bar{B}_R and $0 \leq g' \leq \kappa_{\mu,4}$ on $\bar{B}_4 \setminus B_R$, which record where exiting and entering trajectories end up:

$$\begin{aligned}
\int \zeta df' &:= \int_{\Omega \cap \{X(0) \notin B_R\} \cap \{X(1) \in B_R\}} \zeta(X(1)) d\pi \\
\int \zeta dg' &:= \int_{\Omega \cap \{X(0) \in B_R\} \cap \{X(1) \notin B_R\}} \zeta(X(1)) d\pi.
\end{aligned}$$

Introduce the mass densities

$$\kappa_f = \frac{f'(\mathbb{R}^D)}{|B_R|} \leq \kappa_{\mu,R} \quad \kappa_g = \frac{g'(\mathbb{R}^d)}{|B_R|}.$$

We use Lemma 2.1 to deduce

$$\begin{aligned}
& W_c(\mu \llcorner B_R, \kappa_{\mu,R} dx \llcorner B_R) \\
& \lesssim W_c(\mu \llcorner B_R, \kappa_{\mu,4} dx \llcorner B_R - f' + g') + W_c(\kappa_{\mu,4} dx \llcorner B_R - f' + g', (\kappa_{\mu,4} - \kappa_f) dx \llcorner B_R + g') \\
& \quad + W_c((\kappa_{\mu,4} - \kappa_f) dx \llcorner B_R + g', \kappa_{\mu,R} dx \llcorner B_R)
\end{aligned}$$

$$=I + II + III.$$

Restricting π to trajectories that start in B_R gives an admissible plan for I , that is,

$$I = W_c(\mu \llcorner B_R, \kappa_{\mu,4} dx \llcorner B_R - f' + g') \leq W_c(\mu \llcorner B_4, \kappa_{\mu,4} dx \llcorner B_4) \leq D(4).$$

Regarding II , we begin by noting that using the sub-additivity of W_c , we have

$$\begin{aligned} & W_c(\kappa_{\mu,4} dx \llcorner B_R - f' + g', (\kappa_{\mu,4} - \kappa_f) dx \llcorner B_R + g') \\ & \lesssim W_c(\kappa_{\mu,4} dx \llcorner B_R - f', (\kappa_{\mu,4} - \kappa_f) dx \llcorner B_R) + W_c(g', g') \\ & = W_c(\kappa_{\mu,4} dx \llcorner B_R - f', (\kappa_{\mu,4} - \kappa_f) dx \llcorner B_R). \end{aligned}$$

Since this term will be estimated in the same way as III but is slightly more tricky, we first turn to III .

In order to estimate III , introduce the projection \bar{g} of g' onto ∂B_R , that is

$$\int \zeta d\bar{g} = \int \zeta \left(\frac{Rx}{|x|} \right) g'(dx).$$

Using Lemma 2.1, we deduce

$$III \lesssim W_c((\kappa_{\mu,4} - \kappa_f) dx \llcorner B_R + \bar{g}, \kappa_{\mu,R} dx \llcorner B_R) + W_c(\bar{g}, g').$$

We claim that

$$W_c((\kappa_{\mu,4} - \kappa_f) dx \llcorner B_R + \bar{g}, \kappa_{\mu,R} dx \llcorner B_R) \lesssim \int_{\partial B_R} \bar{g}^2.$$

Indeed, an admissible density-flux pair (ρ, j) for the Benamou-Brenier formulation is given by

$$\begin{cases} \rho_t = (\kappa_{\mu,4} dx \llcorner - \kappa_f + t\kappa_g) + (1-t)\bar{g} \\ j_t = \nabla c^*(D\phi) dx \llcorner B_R, \end{cases}$$

where ϕ solves

$$\begin{cases} -\operatorname{div} \nabla c^*(D\phi) = \kappa_g & \text{in } B_R \\ \nu \cdot \nabla c^*(D\phi) = \bar{g} & \text{on } \partial B_R. \end{cases}$$

We find, writing $s = (\kappa_{\mu,4} - \kappa_f + t\kappa_g) + (1-t)\bar{g}$, for any ζ supported in B_R ,

$$\begin{aligned} \int \zeta dj_t - \int c^*(\zeta) d\rho_t &= \int_{B_R} \zeta \cdot \nabla c^*(D\phi) - c^*(\zeta)s dx \\ &\leq \int_{B_R} \int_0^1 sc \left(\frac{1}{s} \nabla c^*(D\phi) \right) dt dx \end{aligned}$$

To obtain the second line, we used the Fenchel-Young inequality. Assuming $|s-1| \ll 1$ for now, using (1.7) and (1.8) it is straightforward to see that

$$\int_{B_R} \int_0^1 sc \left(\frac{1}{s} \nabla c^*(D\phi) \right) dt dx \lesssim \int_{B_R} c(\nabla c^*(D\phi)) dx \lesssim \int_{\partial B_R} \bar{g}^p.$$

To obtain the last inequality, we used (1.21). Taking into account Lemma 3.1 and Lemma 2.3, we find

$$\int_{\partial B_R} \bar{g}^p \lesssim \int |R - |x||^{p-1} dg'.$$

Noting $\left| R \frac{x}{|x|} - x \right| = ||x| - R|$ and that g' is supported in $\bar{B}_4 \setminus B_R$, we obtain using (1.7)

$$W_c(\bar{g}, g') \lesssim \int ||x| - R|^{p-1} dg'.$$

Proceeding exactly as in Lemma 6.1 we obtain

$$\int |R - |x||^{p-1} dg' \lesssim D(4).$$

Since $\kappa_{\mu,R} = \kappa_{4,R} - \kappa_f + \kappa_g$ to deduce that $|s - 1| \ll 1$ and to conclude the estimate of *III*, it suffices to show $\kappa_f^p + \kappa_g^p \lesssim D(4)$. By symmetry it suffices to consider κ_g . By definition of \bar{g} and Young's inequality, we find

$$\kappa_g^p = \frac{\bar{g}(\mathbb{R}^d)^p}{|B_R|^p} \leq \frac{|\partial B_R|^{p-1}}{|B_R|^p} \int_{\partial B_R} \bar{g}^p.$$

This concludes the estimate for *III*. It remains to finish the estimate of *II*, by estimating

$$W_c(\kappa_{\mu,4} dx_{\perp} B_R - f', (\kappa_{\mu,4} - \kappa_f) dx_{\perp} B_R).$$

We want to proceed exactly as we did in the estimate for *III*, the only delicate issue being that we do not have $\kappa_{\mu,4} dx_{\perp} B_R - f' \geq c > 0$. However, this can be remedied by using Corollary 2.2 to deduce

$$\begin{aligned} & W_c(\kappa_{\mu,4} dx_{\perp} B_R - f', (\kappa_{\mu,4} - \kappa_f) dx_{\perp} B_R) \\ & \lesssim W_c((2\kappa_{\mu,4} - \kappa_f) dx_{\perp} B_R - f', 2(\kappa_{\mu,4} - \kappa_f) dx_{\perp} B_R). \end{aligned}$$

We can now proceed using the same argument as for *III* to conclude

$$W_c(\kappa_{\mu,4} dx_{\perp} B_R - f', (\kappa_{\mu,4} - \kappa_f) dx_{\perp} B_R) \lesssim D(4).$$

This completes the proof. \square

7 The c^* -harmonic approximation result

The goal of this section is to prove the c^* -harmonic approximation result. In order to define the approximation, note that in light of Lemma 5.1, given $\tau > 0$, we fix $R \in (3/2, 2)$ such that there exist non-negative \bar{f}_R, \bar{g}_R such that

$$\begin{aligned} W_c(f_R, \bar{f}_R) + W_c(g_R, \bar{g}_R) & \lesssim \tau E(4) + D(4) \\ \int_{\partial B_R} \bar{f}_R^p + \bar{g}_R^p & \lesssim E(4) + D(4). \end{aligned}$$

Then let ϕ be a solution with $\int_{B_R} \phi \, dx = 0$ of

$$\begin{cases} -\operatorname{div} \nabla c^*(D\phi) = c_R & \text{in } B_R \\ \nabla c^*(D\phi) \cdot \nu = \bar{g}_R - \bar{f}_R & \text{on } \partial B_R. \end{cases} \quad (7.1)$$

where $c_R = |B_R|^{-1} \left(\int_{\partial B_R} \bar{g}_R - \bar{f}_R \right)$ is the constant so that (7.1) is well-posed. We emphasize that while we do not make the dependence explicit in our notation, ϕ depends on the choice of radius R .

Finally due to Lemma 6.1, we will further assume throughout this section that $D(R) \lesssim D(4)$.

With this notation in place, we state a precise version of our main result.

Theorem 7.1. *For every $0 < \tau$, there exist positive constants $\varepsilon(\tau), C(\tau) > 0$ such that if $E(4) + D(4) \leq \varepsilon(\tau)$, then there exists $R \in (2, 3)$ and ϕ solving (7.1) such that*

$$\int_{\#_1} c(x - y - \nabla c^*(D\phi)) \leq \tau E(4) + C(\tau) D(4).$$

The proof will be a direct consequence of the lemmata we prove in the following subsections.

7.1 Quasi-orthogonality

The key observation in order to prove Theorem 7.1 is contained in the following elementary lemma.

Lemma 7.2. *For any $\pi \in \Pi(\lambda, \mu)$ and ϕ continuously differentiable in \bar{B}_R , there is $c(p, \Lambda)$ such that we have*

$$\begin{aligned} & c(p, \Lambda) \int_{\Omega} \int_{\sigma}^{\tau} V \left(\dot{X}(t) - \nabla c^*(\nabla \phi(X(t))) \right) \, dt \, d\pi \\ & \leq \int_{\Omega} c(x - y) \, d\pi - \int_{B_R} c(\nabla^*(D\phi)) \, dx - \int_{\Omega} \int_{\sigma}^{\tau} \langle \dot{X}(t) - \nabla c^*(D\phi(X(t))), D\phi(X(t)) \rangle \, dt \, d\pi \\ & \quad + \int_{B_R} c(\nabla c^*(D\phi)) \, dx - \int_{\Omega} \int_{\sigma}^{\tau} c(\nabla c^*(D\phi(X(t)))) \, dt \, d\pi. \end{aligned}$$

Proof. We apply (1.10) with $x = \dot{X}$ and $y = \nabla c^*(D\phi(X(t)))$. Noting that we have $\nabla c(\nabla c^*(D\phi)) = D\phi$ and $\int_{\Omega} \int_{\sigma}^{\tau} c(\dot{X}(t)) \, dt \, d\pi \leq \int_{\Omega} c(x - y) \, d\pi$ this gives the desired result. \square

7.2 Error estimates

We would like to apply Lemma 7.2 with ϕ solving (7.1). However note that \bar{g}_R and \bar{f}_R will in general not be sufficiently smooth in order to ensure that ϕ is $C^1(\bar{B}_R)$. Thus, we approximate them using mollification. To be precise, let $0 < r \ll 1$ and denote by \bar{f}_R^r and \bar{g}_R^r , respectively, the convolution with a smooth convolution kernel (on ∂B_R) at scale r of \bar{f}_R and \bar{g}_R . Set ϕ^r to be the solution with $\int_{B_R} \phi^r \, dx = 0$ of

$$\begin{cases} -\operatorname{div} \nabla c^*(D\phi^r) = c^r & \text{in } B_R \\ \nabla c^*(D\phi^r) \cdot \nu = \bar{g}_R^r - \bar{f}_R^r & \text{on } \partial B_R. \end{cases} \quad (7.2)$$

Here $c^r = |B_R|^{-1} \left(\int_{\partial B_R} \overline{g}_R^r - \overline{f}_R^r \right)$ is the constant such that (7.2) is well-posed.

We begin by showing that the left-hand sides of the estimate in Lemma 7.2 are close when evaluated for ϕ and ϕ^r .

Lemma 7.3. *For every $0 < \tau$ there exists $\varepsilon(\tau)$ and $C(\tau), r_0(\tau) > 0$, such that if it holds that $E(4) + D(4) \leq \varepsilon(\tau)$ and $0 < r \leq r_0$, then there exists $R \in [2, 3]$ such that if ϕ solves (7.1) and ϕ^r solves (7.2), then*

$$\int_{\Omega_1} \int_{\sigma_1}^{\tau_1} V \left(\dot{X}(t) - \nabla c^*(D\phi(X(t))) \right) dt d\pi - \int_{\Omega_1} \int_{\sigma_1}^{\tau_1} V \left(\dot{X}(t) - \nabla c^*(D\phi^r(X(t))) \right) dt d\pi \lesssim \tau(E(4) + D(4)).$$

Proof. Write $\xi(x) = \nabla c^*(D\phi(x))$, $\xi^r(x) = \nabla c^*(D\phi^r(x))$. We focus on the case $p \leq 2$. The case $p > 2$ follows by similar arguments, but is easier. In light of (1.9), (1.13) and using Hölder, we find

$$\begin{aligned} & \left| \int_{\Omega_1} \int_{\sigma_1}^{\tau_1} V \left(\dot{X}(t) - \xi(X(t)) \right) dt d\pi - \int_{\Omega} \int_{\sigma}^{\tau} V \left(\dot{X}(t) - \xi^r(X(t)) \right) dt d\pi \right| \\ & \lesssim \int_{\Omega_1} \int_{\sigma_1}^{\tau_1} |\xi(X(t)) - \xi^r(X(t))| \left(|\dot{X}(t)| + |\xi(X(t))| + |\xi^r(X(t))| \right)^{p-1} dt d\pi \\ & \lesssim \int_{\Omega_1} \int_{\sigma_1}^{\tau_1} |D\phi(X(t)) - D\phi^r(X(t))| \left(|D\phi(X(t))| + |D\phi^r(X(t))| \right)^{p'-1} dt d\pi \\ & \leq \left(\int_{\Omega_1} \int_{\sigma_1}^{\tau_1} |D\phi(X(t)) - D\phi^r(X(t))|^{p'} dt d\pi \right)^{\frac{1}{p}} \\ & \quad \times \left(\int_{\Omega_1} \int_{\sigma_1}^{\tau_1} |D\phi(X(t))|^{p'} + |D\phi^r(X(t))|^{p'} dt d\pi \right)^{\frac{1}{p}}. \end{aligned}$$

Note that due to Lemma 3.1, if $X \in \Omega_1$, then $X(0) \in B_{3/2}$. Thus using (1.22) and (1.24),

$$\begin{aligned} & \left| \int_{\Omega_1} \int_{\sigma_1}^{\tau_1} |D\phi(X(t)) - D\phi^r(X(t))|^{p'} dt d\pi - \int_{\Omega_1} \int_{\sigma_1}^{\tau_1} |D\phi(X(0)) - D\phi^r(X(0))|^{p'} dt d\pi \right| \\ & \lesssim \left([D\phi]_{C^{0,\beta}(B_{3/2})} + [D\phi^r]_{C^{0,\beta}(B_{3/2})} \right) \left(\sup_{B_{3/2}} |D\phi| + |D\phi^r| \right)^{p'-1} \int_{\Omega_1} \int_{\sigma_1}^{\tau_1} |X(t) - X(0)|^\beta dt d\pi \\ & \lesssim c(r)(E(4) + D(4))\pi(\Omega_1)(E(4) + D(4))^{\frac{\beta}{p+d}} \lesssim c(r)(E(4) + D(4))^{1+\frac{\beta}{p+d}}. \end{aligned}$$

In order to obtain the last line, we used Lemma 3.1. Moreover, using Lemma 3.2 and (1.23),

$$\begin{aligned} & \int_{\Omega_1} \int_{\sigma_1}^{\tau_1} |D\phi(X(0)) - D\phi^r(X(0))|^{p'} dt d\pi \leq \int_{B_{3/2}} |D\phi(x) - D\phi^r(x)|^{p'} d\mu \\ & \leq \left| \int_{B_{3/2}} |D\phi(x) - D\phi^r(x)|^{p'} (d\mu - \kappa_{\mu,R} dx) \right| + \kappa_{\mu,R} \int_{B_R} |D\phi(x) - D\phi^r(x)|^{p'} dx \\ & \lesssim \left(\sup_{B_{3/2}} |D\phi|^{p'-1} + |D\phi^r|^{p'-1} \right) \left([D\phi]_{C^{0,\beta}(B_{(1+R)/2})} + [D\phi^r]_{C^{0,\beta}(B_{(1+R)/2})} \right) \end{aligned}$$

$$\begin{aligned} & \times W_c(\mu \llcorner B_R, \kappa_{\mu, R} dx \llcorner B_R)^{\frac{\beta}{p}} + r(E(4) + D(4)) \\ & \lesssim c(r)(E(4) + D(4))^{1+\frac{\beta}{p}} + r(E(4) + D(4)). \end{aligned}$$

Arguing similarly, that is first replacing $X(t)$ with $X(0)$, at the cost of making an error of size $c(r)(E(4) + D(4))^{1+\frac{\beta}{p+d}}$, and then replacing $d\mu$ with dx , making an error $c(r)(E(4) + D(4))^{1+\frac{\beta}{p}} + r(E(4) + D(4))$, we find

$$\begin{aligned} & \int_{\Omega_1} \int_{\sigma_1}^{\tau_1} |\nabla \phi(X(t))|^{p'} + |\nabla \phi^r(X(t))|^{p'} dt d\pi \\ & \lesssim r(E(4) + D(4)) + c(r)(E(4) + D(4))^{1+\frac{\beta}{p+d}}. \end{aligned}$$

Collecting estimates and choosing r as well as $\varepsilon(\tau)$ sufficiently small, the desired result follows. \square

We now turn to estimating each of the three terms on the right-hand side of the estimate in Lemma 7.2 in turns. We will see that the second and third term are errors that arise from the approximation of the boundary data and from passing to the perspective of trajectories, respectively. Accordingly, estimating them will be essentially routine. In contrast, estimating the first term requires us to contrast an appropriate competitor to π .

Lemma 7.4. *For every $0 < \tau$ there exists $\varepsilon(\tau), C(\tau), r_0(\tau) > 0$ such that if it holds that $E(4) + D(4) \leq \varepsilon(\tau)$ and $0 < r \leq r_0$, then there exists $R \in [2, 3]$ such that if ϕ^r solves (7.2), then*

$$\int_{\Omega} c(x - y) d\pi - \int_{B_R} c(\nabla c^*(D\phi^r)) dx \lesssim \tau E(4) + D(4).$$

Proof. We note, in the case $p \leq 2$, using (1.8), (1.13) and Hölder,

$$\begin{aligned} & \int_{B_R} c(\nabla c^*(D\phi^r)) - c(\nabla c^*(D\phi)) dx \\ & \lesssim \int_{B_R} |\nabla c^*(D\phi^r) - \nabla c^*(D\phi)| (|\nabla c^*(D\phi^r)| + |\nabla c^*(D\phi)|)^{p-1} dx \\ & \lesssim \int_{B_R} |D\phi - D\phi^r| (|D\phi| + |D\phi^r|)^{p'-1} \\ & \lesssim \|D\phi - D\phi^r\|_{L^{p'}(B_R)} \left(\|D\phi\|_{L^{p'}(B_R)} + \|D\phi^r\|_{L^{p'}(B_R)} \right)^{p'-1} \\ & \lesssim r^{\frac{1}{p'}} (E(4) + D(4)). \end{aligned}$$

To obtain the last line, we used (1.20) and (1.23). In case $p \geq 2$ a similar estimate holds by the same argument. Due to Lemma 4.1 and Corollary 3.3,

$$\begin{aligned} \int_{\Omega} c(x - y) d\pi & \leq W_c(\lambda \llcorner B_R + f_R, \mu \llcorner B_R + g_R) + 2 \int_{\Omega \cap \{\exists t \in [0, 1]: X(t) \in \partial B_R\}} c(x - y) d\pi \\ & \leq W_c(\lambda \llcorner B_R + f_R, \mu \llcorner B_R + g_R) + c(\tau E(4) + D(4)). \end{aligned}$$

In particular, combining the previous two estimates and choosing r sufficiently small, it suffices to prove

$$W_c(\lambda_{\perp} B_R + f_R, \mu_{\perp} B_R + g_R) - \int_{B_R} c(\nabla c^*(D\phi)) dx \lesssim \tau E(4) + D(4).$$

Using Lemma 2.1, we obtain for $\delta \in (0, 1)$ to be fixed

$$\begin{aligned} & W_c(\lambda_{\perp} B_R + f_R, \mu_{\perp} B_R + g_R) \\ & \leq (1 + \delta) W_c(\kappa_{\lambda, R} dx_{\perp} B_R + \bar{f}_R, \kappa_{\mu, R} dx_{\perp} B_R + \bar{g}_R) \\ & \quad + c(\delta) \left(W_c(\lambda_{\perp} B_R, \kappa_{\lambda, R} dx_{\perp} B_R) + W_c(\mu_{\perp} B_R, \kappa_{\mu, R} dx_{\perp} B_R) \right) \\ & \quad + c(\delta) \left(W_c(f_R, \bar{f}_R) + W_c(g_R, \bar{g}_R) \right). \end{aligned}$$

Noting that due to the definition of D and our choice of R ,

$$\begin{aligned} & W_c(\lambda_{\perp} B_R, \kappa_{\lambda, R} dx_{\perp} B_R) + W_c(\mu_{\perp} B_R, \kappa_{\mu, R} dx_{\perp} B_R) + W_c(f_R, \bar{f}_R) + W_c(g_R, \bar{g}_R) \\ & \lesssim \tau E(4) + D(4), \end{aligned}$$

we claim that for some $C > 0$,

$$W_c(\kappa_{\lambda, R} dx_{\perp} B_R + \bar{f}_R, \kappa_{\mu, R} dx_{\perp} B_R + \bar{g}_R) \leq (1 + CD(4)^{\frac{1}{p}}) \int_{B_R} c(\nabla c^*(D\phi)) dx. \quad (7.3)$$

Collecting estimates, choosing first δ and r small, then ε small, once (7.3) is established, the proof is complete.

Establishing (7.3) is easy to do using the Benamou-Brenier formulation. For $t \in [0, 1]$ introduce the non-singular, non-negative measure

$$\rho_t = t(\kappa_{\mu, R} dx_{\perp} B_R + \bar{f}_R) + (1 - t)(\kappa_{\lambda, R} dx_{\perp} B_R + \bar{g}_R),$$

and the vector-valued measure

$$j_t = \nabla c^*(D\phi) dx_{\perp} B_R.$$

Note that (7.2) can be rewritten as

$$\frac{d}{dt} \int \zeta d\rho_t = \int \nabla \zeta \cdot dj_t$$

for all test functions ζ . Set

$$\int c \left(\frac{dj_t}{d\rho_t} \right) d\rho_t = \sup_{\zeta \in C_c^0(\mathbb{R}^d)} \left\{ \int \zeta dj_t - \int c^*(\xi) d\rho_t \right\}$$

The Benamou-Brenier formula gives

$$W_c(\kappa_{\lambda, R} dx_{\perp} B_R + \bar{f}_R, \kappa_{\mu, R} dx_{\perp} B_R + \bar{g}_R) \leq \int_0^1 \int c \left(\frac{dj_t}{d\rho_t} \right) d\rho_t dt.$$

Since j_t is supported in B_R it suffices to consider ζ supported in B_R . Then by definition and the Fenchel-Young inequality for any $s > 0$,

$$\begin{aligned} \int \zeta dj_t - \int c^*(\zeta) d\rho_t &= \int_{B_R} \zeta \cdot \nabla c^*(D\phi) - c^*(\zeta)(t\kappa_{\mu,R} + (1-t)\kappa_{\lambda,R}) dx \\ &\leq \int_{B_R} sc^*(\zeta) + sc\left(\frac{1}{s}\nabla c^*(D\phi)\right) - c^*(\zeta)(t\kappa_{\mu,R} + (1-t)\kappa_{\lambda,R}) dx. \end{aligned}$$

Choosing $s = t\kappa_{\mu,R} + (1-t)\kappa_{\lambda,R}$ and integrating in t , we deduce

$$W_c(\kappa_{\mu,R} dx \llcorner B_R + \bar{f}_R, \kappa_{\lambda,R} dx \llcorner B_R + \bar{g}_R) \leq \int_{B_R} \int_0^1 sc\left(\frac{\nabla c^*(D\phi)}{s}\right) dt dx.$$

Now

$$\begin{aligned} \int_{B_R} \int_0^1 sc\left(\frac{\nabla c^*(D\phi)}{s}\right) dt dx &\leq \int_{B_R} c(\nabla c^*(D\phi)) dx + \int_{B_R} \int_0^1 (s-1)c(\nabla c^*(D\phi)) dx dt \\ &\quad + \int_{B_R} \int_0^1 s\left(c\left(\frac{\nabla c^*(D\phi)}{s}\right) - c(\nabla c^*(D\phi))\right) dt dx. \end{aligned}$$

Note that

$$|s-1| \lesssim D(4)^{\frac{1}{p}}.$$

Further using (1.8) and (1.7),

$$\begin{aligned} &\int_{B_R} \int_0^1 s\left(c\left(\frac{\nabla c^*(D\phi)}{s}\right) - c(\nabla c^*(D\phi))\right) dt dx \\ &\lesssim \int_{B_R} \int_0^1 \left|1 - \frac{1}{s}\right| \left(1 + \frac{1}{s}\right)^{p-1} |\nabla c^*(D\phi)|^p dt dx \lesssim D(4)^{\frac{1}{p}} \int_{B_R} c(\nabla c^*(D\phi)) dx. \end{aligned}$$

Thus the proof of (7.3) is complete. \square

We turn to the second term on the right-hand side of Lemma 7.2. This term will be small due to the definition of ϕ^r .

Lemma 7.5. *For every $0 < \tau$ there exists $\varepsilon(\tau), C(\tau), r_0(\tau) > 0$ such that if it holds that $E(4) + D(4) \leq \varepsilon(\tau)$ and $0 < r \leq r_0$, then there exists $R \in [2, 3]$ such that if ϕ^r solves (7.2), then*

$$\int_{\Omega} \int_{\sigma}^{\tau} \langle \dot{X}(t) - \nabla c^*(D\phi^r(X(t))), D\phi^r(X(t)) \rangle dt d\pi \lesssim \tau E(4) + D(4).$$

Proof. Note that $\frac{d}{dt}\phi^r(X(t)) = \langle \dot{X}(t), \nabla\phi^r(X(t)) \rangle$. Thus, since $\pi \in \Pi(\lambda, \mu)$ as well as using the definition of f_R and g_R ,

$$\int_{\Omega} \int_{\sigma}^{\tau} \langle \dot{X}(t), D\phi^r(X(t)) \rangle dt d\pi = \int_{\Omega} \phi^r(X(\tau)) - \phi^r(X(\sigma)) d\pi$$

$$\begin{aligned}
&= \int_{\Omega \cap \{X(\tau) \in \partial B_R\}} \phi^r(X(\tau)) \, d\pi + \int_{\{x \in B_R\}} \phi^r(x) \, d\pi \\
&\quad - \int_{\Omega \cap \{X(\sigma) \in \partial B_R\}} \phi^r(X(\sigma)) \, d\pi - \int_{\{y \in B_R\}} \phi^r(y) \, d\pi \\
&= \int_{B_R} \phi^r \, d(\mu - \lambda) + \int_{\partial B_R} \phi^r \, d(g_R - f_R).
\end{aligned}$$

On the other hand, as in Lemma 7.3, at cost of an error $c(r)(E(4) + D(4))^{1+\frac{\beta}{p+d}}$, we may replace $D\phi^r(X(t))$ with $D\phi^r(x)$ in the expression

$$\int_{\Omega} \int_{\sigma}^{\tau} \langle \nabla c^*(D\phi^r(X(t)), D\phi^r(X(t))) \rangle \, dt \, d\pi$$

and $\int_{\Omega} \int_{\sigma}^{\tau} dt \, d\pi$ with $\int_{B_R} dx$ at the cost of a further error $c(r)(E(4) + D(4))^{1+\frac{\beta}{p}}$. Thus it suffices to consider

$$\int_{B_R} \langle \nabla c^*(D\phi), D\phi \rangle \, dx = c^r \int_{B_R} \phi^r \, dx + \int_{\partial B_R} \phi^r \, d(\bar{g}_R^r - \bar{f}_R^r).$$

Collecting estimates, we have shown

$$\begin{aligned}
&\int_{\Omega} \int_{\sigma}^{\tau} \langle \dot{X}(t) - \nabla c^*(D\phi^r(X(t)), D\phi^r(X(t))) \rangle \, dt \, d\pi \\
&\lesssim \int_{B_R} \phi^r \, d(\mu - \lambda - c^r) + \int_{\partial B_R} \phi^r \, d(g_R - \bar{g}_R^r + f_R - \bar{f}_R^r) + c(r)(E(4) + D(4))^{1+\frac{\beta}{p+d}} \\
&= I + II + III.
\end{aligned}$$

We find using Lemma 3.2, Young's inequality, (1.24) and (1.20),

$$\begin{aligned}
I &\leq \left| \int_{B_R} \phi^r \, d(\mu - \kappa_{\mu,R} - \lambda + \kappa_{\lambda,R}) \right| + \left| \int_{B_R} \phi^r \, d(\kappa_{\mu,R} - \kappa_{\lambda,R} - c^r) \right| \\
&\lesssim \sup_{B_R} |D\phi^r| \left(W_c(\lambda \llcorner B_R, \lambda_{\mu,R} \, dx \llcorner B_R)^{\frac{1}{p}} + W_c(\mu \llcorner B_R, \kappa_{\mu,R} \, dx \llcorner B_R)^{\frac{1}{p}} \right) \\
&\quad + |\kappa_{\mu,R} - \kappa_{\lambda} - c^r| \|\phi^r\|_{L^p(B_R)} \\
&\lesssim \tau E(4) + c(\tau, r) D(4) + c(\tau) |\kappa_{\mu,R} - \kappa_{\lambda,R} - c^r|^{p'}.
\end{aligned}$$

Noting that as $r \rightarrow 0$,

$$c^r = |B_R|^{-1} (g^r(\partial B_R) - f^r(\partial B_R)) \rightarrow |B_R|^{-1} (g(\partial B_R) - f(\partial B_R)) = \kappa_{\mu,R} - \kappa_{\lambda,R},$$

we deduce, choosing r_0 sufficiently small, that

$$I = \int_{B_R} \phi^r \, d(\mu - \lambda - c^r) \lesssim \tau E(4) + D(4).$$

Finally consider *II*. By symmetry it suffices to consider the terms involving g . We estimate, denoting by $(\phi^r)^r$ convolution of ϕ^r with the convolution kernel used to construct g_R^r ,

$$\int_{\partial B_R} \phi^r \, d(g_R - \bar{g}_R^r) = \int_{\partial B_R} (\phi^r)^r - \phi^r \, d\bar{g}_R + \int_{\partial B_R} \phi^r \, d(\bar{g}_R - g_R).$$

Now note that a standard mollification argument shows

$$\int_{\partial B_R} \phi^r d(g_R - \bar{g}_R^r) \lesssim r^{\frac{1}{p}} \|\mathbf{D}\phi^r\|_{L^{p'}(B_R)} \|\bar{g}_R\|_{L^p(\partial B_R)} \lesssim r^{\frac{1}{p}} (E(4) + D(4)),$$

and that moreover using Lemma 3.2 and Young's inequality,

$$\int_{\partial B_R} \phi^r (d\bar{g}_R - g) \lesssim [\mathbf{D}_{\tan}\phi^r]_{C^{0,\beta}(\partial B_R)} Wc(\bar{g}_R, \bar{g})^{\frac{1}{p}} \lesssim c(r)(\tau E(4) + D(4)).$$

Thus, collecting estimates and first choosing r_0 sufficiently small, then ε small, we conclude the proof. \square

We next estimate the third term on the right-hand side of the estimate in Lemma 7.2.

Lemma 7.6. *For every $0 < \tau$ there exists $\varepsilon(\tau), C(\tau), r_0(\tau) > 0$ such that if it holds that $E(4) + D(4) \leq \varepsilon(\tau)$ and $0 < r \leq r_0$, then there exists $R \in [2, 3]$ such that if ϕ^r solves (7.2), then*

$$\int_{B_R} c(\nabla c^*(\mathbf{D}\phi^r)) dx - \int_{\Omega} \int_{\sigma}^{\tau} c(\nabla c^*(\mathbf{D}\phi^r(X(t)))) dt d\pi \lesssim \tau E(4) + D(4).$$

Proof. Set $\xi = c(\nabla c^*(\mathbf{D}\phi^r))$. Then

$$\begin{aligned} & \int_{B_R} c(\nabla c^*(\mathbf{D}\phi^r)) dx - \int_{\Omega} \int_{\sigma}^{\tau} c(\nabla c^*(\mathbf{D}\phi^r(X(t)))) dt d\pi \\ &= (1 - \kappa_{\mu,R}) \int_{B_R} \xi dx + \left(\kappa_{\mu,R} \int_{B_R} \xi dx - \int_{B_R \times \mathbb{R}^d} \xi d\pi \right) \\ & \quad + \int_{B_R \times \mathbb{R}^d} \xi d\pi - \int_{\Omega} \int_{\sigma}^{\tau} \xi(X(t)) dt d\pi \\ &= I + II + III. \end{aligned}$$

Using (1.21) and Young's inequality, we find

$$I \leq D(4)^{\frac{1}{p}} (E(4) + D(4)) \lesssim \tau E(4) + D(4).$$

Employing Lemma 3.2 we deduce

$$II \lesssim \|\xi\|_{C^{0,\beta}(\bar{B}_R)} Wc(\mu \llcorner B_R, \kappa_{\mu,R} \llcorner B_R)^{\frac{\beta}{p}}.$$

It is straightforward to check that $\|\xi\|_{C^{0,\beta}(\bar{B}_R)} \lesssim \|\mathbf{D}\phi^r\|_{C^{0,\beta}(\bar{B}_R)} \left(\sup_{\bar{B}_R} \mathbf{D}\phi^r \right)^{p'-1}$, so that using (1.24) and Young's inequality,

$$II \lesssim c(r)(E(4) + D(4)) D(4)^{\frac{\beta}{p}} \lesssim \tau E(4) + D(4).$$

In order to estimate III , we first find

$$I(X(0) \in B_R) \xi(X(0)) - I(X(t) \in B_R) \xi(X(t))$$

$$\begin{aligned} &\leq I(\exists s \in [0, 1]: X(s) \in \partial B_R, X(0) \in \overline{B_R})\xi(X(0)) \\ &\quad + I(\forall s \in [0, 1] X(s) \in B_R) (\xi(X(0)) - \xi(X(t))). \end{aligned}$$

Thus, using also (1.24) and Jensen's inequality,

$$\begin{aligned} III &\leq \int_0^1 \int I(X(0) \in B_R)\xi(X(0)) - I(X(t) \in B_R)\xi(X(t)) \, d\pi \, dt \\ &\leq \sup_{\overline{B_R}} |\xi| \int_0^1 \int I(\exists s \in [0, 1]: X(s) \in \partial B_R, X(0) \in \partial B_R) \, d\pi \, dt \\ &\quad + \|D\xi\|_{C^{0,\beta}(\overline{B_R})} \int_0^1 \int I(\forall s \in [0, 1] X(s) \in B_R) |X(t) - X(0)|^\beta \, dt \, d\pi \\ &\lesssim c(r)(E(4) + D(4))\pi(\exists t \in [0, 1]: X(t) \in \partial B_R) + c(r)(E + D) \int_\Omega |x - y|^\beta \, d\pi \\ &\lesssim c(r)(E(4) + D(4))^{1+\frac{1}{p+d}} + c(r)(E(4) + D(4))^{1+\frac{\beta}{p}}. \end{aligned}$$

In order to obtain the last line we used Corollary 3.3. Collecting estimates and choosing first r_0 small, then ε small, we conclude the proof. \square

7.3 Proof of Theorem 7.1

We are now ready to prove Theorem 7.1.

Proof of Theorem 7.1. Applying Lemma 7.2 to ϕ^r and collecting the output of Lemma 7.3, Lemma 7.4, Lemma 7.5 and Lemma 7.6, we have shown that for any $0 < \tau$, there is $\varepsilon, C > 0$ such that if $E(4) + D(4) \leq \varepsilon$, then

$$\int_{\Omega_1} \int_{\sigma_1}^{\tau_1} V(\dot{X}(t) - \nabla c^*(D\phi(X(t)))) \, dt \, d\pi \leq \tau E(4) + CD(4).$$

Arguing as in Lemma 7.3, we may replace $D(\phi(X(t)))$ by $D(\phi(X(0)))$ at the cost of an error of size $(E(4) + D(4))^{1+\frac{\beta}{p+d}}$. Noting that $V(z - \nabla c^*(D\phi(x)))$ is a convex function of z , we employ Jensen's inequality to deduce

$$\int_{\#_1} V(x - y - \nabla c^*(D\phi(x))) \, d\pi \leq \int_{\Omega_1} \int_0^1 V(\dot{X}(t) - \nabla c^*(D\phi(X(0)))) \, dt \, d\pi.$$

Now using (1.14) and (1.22), as well as Corollary 3.3,

$$\begin{aligned} &\int_{\Omega_1 \cap (\{\sigma_1 > 0\} \cup \{\tau_1 < 1\})} \int_{\sigma_1}^{\tau_1} V(\dot{X}(t) - \nabla c^*(D\phi(X(0)))) \, dt \, d\pi \\ &\lesssim \int_{\Omega_1 \cap (\{\sigma_1 > 0\} \cup \{\tau_1 < 1\})} |\dot{X}(t)|^p + |D\phi(X(0))|^{p'} \\ &\lesssim (E(4) + D(4))\pi(\Omega_1 \cap (\{\sigma_1 > 0\} \cup \{\tau_1 < 1\})) \\ &\lesssim (E(4) + D(4))^{1+\frac{1}{p+d}}. \end{aligned}$$

Thus, we conclude

$$\int_{\#_1} V(x - y - \nabla c^*(D\phi(x))) \, d\pi \lesssim \tau E(4) + D(4).$$

This concludes the proof in the case $p \geq 2$. In the case $p \leq 2$, an application of Hölder's inequality combined with (1.7) concludes the proof. \square

References

- [1] C. de Filippis, L. Koch, and J. Kristensen. Regularity in relaxed convex problems. *in Prep.*, 2023.
- [2] G. De Philippis and A. Figalli. Partial regularity for optimal transport maps. *Publ. Math. Inst. Hautes Études Sci.*, 121:81–112, 2015.
- [3] A. Figalli and Y.-H. Kim. Partial regularity of Brenier solutions of the Monge-Ampère equation. *Discret. Contin. Dyn. Syst.*, 28(2):559–565, 2010.
- [4] W. Gangbo and R.J. McCann. The geometry of optimal transportation. *C. R. Acad. Sci. Paris Sér. Mat.*, 321(12):1653–1658, 1995.
- [5] M. Goldman. An ε -regularity result for optimal transport maps between continuous densities. *arXiv Prepr. arXiv2005.06785*, 2020.
- [6] M. Goldman and M. Huesmann. A fluctuation result for the displacement in the optimal matching problem. *Ann. Probab.*, 50(4):144601477, 2022.
- [7] M. Goldman, M. Huesmann, and F. Otto. Quantitative linearization results for the Monge-Ampère equation. *Commun. Pure Appl. Math.*, 74(12):2483–2560, 2021.
- [8] M. Goldman and F. Otto. A variational proof of partial regularity for optimal transportation maps. *Ann. Sci. Éc. Norm. Supér.*, 53(5):1209–1233, 2020.
- [9] E. G. Gutiérrez and A. Montanari. L^∞ -estimates in optimal transport for non quadratic costs. *arXiv Prepr. arXiv2106.02899*, 2021.
- [10] J.B. Hiriart-Urruty and C. Lemar. *Convex Analysis and Minimization Algorithms II*. Springer-Verlag Berlin Heidelberg, 1993.
- [11] M. Huesmann, F. Mattesini, and F. Otto. There is no stationary cyclically monotone Poisson matching in 2d. *arXiv Prepr. arXiv2109.13590*, 2021.
- [12] L. Koch and F. Otto. Lectures notes on the harmonic approximation to quadratic optimal transport. *in Prep.*, 2023.
- [13] O. Ladyzhenskaya and N. Ural'tseva. *Linear and quasilinear elliptic equations*. Academic Press Inc., New York, 1968.
- [14] G.M. Lieberman. Boundary regularity for solutions of degenerate elliptic equations. *Nonlinear Anal. Theory, Methods Appl.*, 12(11):1203–1219, 1988.
- [15] F. Otto, M. Prod'homme, and T. Ried. Variational approach to regularity of optimal transport maps: general cost functions. *Ann. PDE*, 7, 2021.
- [16] R.T. Rockafellar. *Convex analysis*. Princeton University Press, Princeton, N.J., 1970.
- [17] F. Santambrogio. *Optimal Transport for Applied Mathematicians*. Birkhäuser Press, 2015.

- [18] C. Villani. *Optimal Transport. Old and new.* Springer-Verlag Berlin Heidelberg, 2009.