

Non-Abelian homology and homotopy colimit of classifying spaces for a diagram of groups

Ahmet A. Husainov

Abstract

The paper considers non-Abelian homology groups for a diagram of groups introduced as homotopy groups of a simplicial replacement. It is proved that the non-Abelian homology groups of the group diagram are isomorphic to the homotopy groups of the homotopy colimit of the diagram of classifying spaces, with a dimension shift of 1. As an application, a method is developed for finding a nonzero homotopy group of least dimension for a homotopy colimit of classifying spaces. For a group diagram over a free category with a zero colimit, a criterion for the isomorphism of the first non-Abelian and first Abelian homology groups is obtained. It is established that the non-Abelian homology groups are isomorphic to the cotriple derived functors of the colimit functor defined on the category of group diagrams.

2000 Mathematics Subject Classification 55U10, 18C15, 18G10, 55P10, 55R35

Keywords: non-Abelian homology, homotopy groups, simplicial group, group diagrams, cotriple homology, left derived functors, homology of small categories, classifying spaces, homotopy colimit.

Contents

1	Introduction	2
2	Preliminaries	4
2.1	Notation	5
2.2	Diagrams on the category of simplices	6
2.3	Simplicial replacement	8
2.4	A homotopy colimit in the category of pointed simplicial sets	9
3	Classifying space of a simplicial group	11
3.1	Moore complex and its homology groups	11
3.2	Homotopies of a simplicial group and its classifying space . . .	12

4	Non-Abelian homology with coefficients in a group diagram	13
4.1	Simplicial replacement of group diagram	13
4.2	Non-Abelian group diagram homology	14
5	Relationship between non-Abelian homology and homotopy colimits	15
5.1	Homotopy groups of a wedge for classifying spaces	15
5.2	Non-Abelian homology and homotopy groups of the homotopy colimit	16
5.3	Abelian homology of diagrams over a free category	17
5.4	Comparison and application of Abelian and non-Abelian homologies of group diagrams	19
6	Derived functors of the colimit and non-Abelian homology for group diagrams	23
6.1	Left derived with respect to a cotriple	24
6.2	Cotriple for constructing the resolution	24
6.3	Derived functors of colimit	25
7	Conclusion	29

1 Introduction

We introduce non-Abelian homology $\text{colim}_n^{\mathcal{C}} \mathcal{G}$ for an arbitrary group diagram \mathcal{G} over the small category \mathcal{C} (Definition 4.1) using a simplicial replacement of the group diagram in the sense of Bousfield and Kan [3, XII.5.4]. We show that these homology groups can be interpreted as homotopy groups of the homotopy colimit for the diagram of classifying spaces $B\mathcal{G}$ (Theorem 5.2). We prove that these homologies are isomorphic to the cotriple derived functors of the colimit functor for group diagrams (Theorem 6.4).

In homotopy theory, many good methods and models have appeared for the study and calculation of homotopy groups [6]. However, it remains necessary to calculate the homology, and this requires the development of homological methods. This paper is devoted to the theory of non-Abelian homology of group diagrams, which develops the classical theory of homology of small categories. The theory of non-Abelian homology of diagrams is aimed at solving problems of homotopy theory.

The homology of small categories with coefficients in diagrams of abelian groups, shortly abelian homology of diagrams, can be defined in a similar way, using a simplicial replacement. However, the meanings of non-Abelian and Abelian diagram homologies are very different. For example, for the category V consisting of three objects and two morphisms

$$b \leftarrow a \rightarrow c,$$

except for the identical ones, the Abelian homology of any diagram over V is equal to 0 in dimensions $n \geq 2$. Let's compare them with non-Abelian homology. To this aim, we use the following formula, proved in our paper (Theorem 5.2):

$$\operatorname{colim}_n^{\mathcal{C}} \mathcal{G} \cong \pi_{n+1}(\operatorname{hocolim}^{\mathcal{C}} B\mathcal{G}), \quad (1)$$

where $\operatorname{hocolim}^{\mathcal{C}} B\mathcal{G}$ is the homotopy colimit of the diagram in the category of pointed simplicial sets, which consists of the classifying spaces (nerves) of the groups of the diagram \mathcal{G} . It follows from this formula that the n th non-Abelian homology of the group diagram $0 \leftarrow \mathbb{Z} \rightarrow 0$ is isomorphic to the homotopy groups of the two-dimensional sphere $\pi_{n+1}(S^2)$. It was established in [22] that the n th homotopy groups of the two-dimensional sphere are not equal to 0 for all $n \geq 2$. This implies that the n th non-Abelian homology of the diagram $0 \leftarrow \mathbb{Z} \rightarrow 0$ are not equal to its Abelian homology for every $n \geq 1$.

The formula (1) also shows that the non-Abelian homology of the diagram $0 \leftarrow G \rightarrow 0$ is isomorphic to the homotopy groups of the suspension over the classifying space of the group G , which in some cases were calculated in the papers [7] and [26]. In [31], the software for calculation of homotopy groups of such spaces is developed. This points to the possibility of computing non-Abelian homology of group diagrams.

The assertion that we obtained from the formula (1) that for simplicial groups considered as group diagrams (over Δ^{op}), non-Abelian homology is isomorphic to homotopy groups also inspires hope for perspective (Corollary 5.3).

From the formula (1) we also obtain a criterion for the simply connectedness of the homotopy colimit for the diagram of classifying spaces of groups (Corollary 5.4). By the formula (1), the problem of calculating non-Abelian homology of a group diagram

$$\mathcal{G}(b) \leftarrow \mathcal{G}(a) \rightarrow \mathcal{G}(c) \quad (2)$$

is closely related to some problems of describing the homotopy groups of a homotopy pushout, for example [5, Problem 2.2]. Note also that the formulas for the homotopy groups of the homotopy colimit of the diagram of classifying spaces of quotient groups $G/N_{i_1} \cdots N_{i_k}$ over the inclusion-partially ordered set of proper subsets of the set $\{1, 2, \dots, n\}$ proved by Ellis and Mikhailov [12] can be considered as formulas for non-Abelian homology in n and $n - 1$ dimensions.

In our paper, for a group diagram \mathcal{G} over a free category $W\Gamma$ with $\text{colim}^{W\Gamma} \mathcal{G} = 0$, we obtain an isomorphism criterion for the first non-Abelian homology group of this diagram and the first Abelian homology group of its abelianization (Proposition 5.5).

We have introduced the connectivity $\text{con} \mathcal{G}$ of the group diagram \mathcal{G} as the smallest $n \geq 0$ for which the group $\text{colim}_n^{\mathcal{C}} \mathcal{G}$ is non-trivial. Example 5.7 demonstrates a method for finding the connectivity $n = \text{con} \mathcal{G}$ and calculating the group $\text{colim}_n^{\mathcal{C}} \mathcal{G}$. The number $\text{con} \mathcal{G}$ is equal to the largest n for which the space $\text{hocolim}^{\mathcal{C}} B\mathcal{G}$ is n -connected.

The last section is devoted to proving the isomorphism of non-Abelian homology and cotriple derived functors of the colimit functor $\text{colim}^{\mathcal{C}} : \text{Grp}^{\mathcal{C}} \rightarrow \text{Grp}$. The cotriple $T = \Lambda\Omega$ is constructed using adjoint functors $\Lambda : \text{Grp}^{\text{Ob} \mathcal{C}} \rightleftarrows \text{Grp}^{\mathcal{C}} : \Omega$, where Ω assigns to each diagram $\mathcal{G} : \mathcal{C} \rightarrow \text{Grp}$ its restriction to the largest discrete category $\text{Ob} \mathcal{C}$ of the category \mathcal{C} , and Λ is the left Kan extension along the embedding $\text{Ob} \mathcal{C} \subseteq \mathcal{C}$. An investigation of the bar-resolution obtained on the basis of a cotriple shows that applying the colimit functor to this resolution leads to a simplicial group isomorphic to a simplicial replacement of the group diagram. Hence, the derived functors of the colimit are isomorphic to non-Abelian homology.

2 Preliminaries

We work with simplicial sets. We adhere to the notation from Gabriel and Zisman [16] and MacLane [25]. Homotopy colimits will be considered in the category of pointed simplicial sets, the main source is the monograph by Bousfield and Kan [3]. Pointed bisimplicial sets will also be used, from the book by Goerss and Jardine [18] and some information about simplicial groups and their homotopy groups from Wu [35].

2.1 Notation

Let us write out the notation and some definitions. The rest will be given in the course of the presentation.

- $\mathcal{A}^{\mathcal{C}}$ is the category of functors from the small category \mathcal{C} to a category \mathcal{A} . Functors F from a small category \mathcal{C} to an arbitrary \mathcal{A} are called object diagrams of the category \mathcal{A} over \mathcal{C} and can be denoted as the family $\{F(c)\}_{c \in \mathcal{C}}$.
- $\Delta A = \Delta_{\mathcal{C}} A$ is a functor $\mathcal{C} \rightarrow \mathcal{A}$ that takes constant values $A \in \text{Ob} \mathcal{A}$ on objects and 1_A on morphisms.
- \mathbb{N} - set of non-negative integers.
- Cat - category of small categories and functors.
- Set - category of sets and mappings.
- Grp is the category of groups and homomorphisms.
- 0 - a group consisting of one element.
- Δ is the category of finite linearly ordered sets $[n] = \{0, 1, \dots, n\}$, $n \geq 0$, and non-decreasing mappings. The category Δ is generated by morphisms of the following form:
 1. $\partial_n^i : [n-1] \rightarrow [n]$ (for $0 \leq i \leq n$) - increasing mapping whose image does not contain i ,
 2. $\sigma_n^i : [n+1] \rightarrow [n]$ (for $0 \leq i \leq n$) is a non-decreasing surjection that takes the value i twice,
- $\text{colim}_n^{\mathcal{C}} : \text{Grp}^{\mathcal{C}} \rightarrow \text{Grp}$ ($\forall n \geq 0$) are functors assigning to each group diagram the values of the non-Abelian homology of this diagram (the definition will be given in our work),
- $\varinjlim_n^{\mathcal{C}} : \text{Ab}^{\mathcal{C}} \rightarrow \text{Ab}$ ($\forall n \geq 0$) are Abelian homology functors defined in [16, Application II] as left satellites of the colimit functor, and in [21] as homology of small categories with coefficients in object diagrams of the abelian category.

Let $\Phi : \mathcal{C} \rightarrow \mathcal{D}$ be a functor between small categories. For an arbitrary category \mathcal{A} , the inverse image functor $\Phi^* : \mathcal{A}^{\mathcal{D}} \rightarrow \mathcal{A}^{\mathcal{C}}$ is defined, which assigns to each diagram $F \in \mathcal{A}^{\mathcal{D}}$ composition $F\Phi = F \circ \Phi \in \mathcal{A}^{\mathcal{C}}$, and to each natural transformation $\eta : F \rightarrow F'$ the natural transformation $\eta\Phi : F\Phi \rightarrow F'\Phi$ defined by the formula $(\eta\Phi)_c = \eta_{\Phi(c)}$, for all $c \in \text{Ob } \mathcal{C}$. If \mathcal{A} is a cocomplete category, then the functor Φ^* has a left adjoint functor $\text{Lan}^{\Phi} : \mathcal{A}^{\mathcal{C}} \rightarrow \mathcal{A}^{\mathcal{D}}$, which is called the left Kan extensions.

A simplicial set is a functor $X : \Delta^{op} \rightarrow \text{Set}$. The values of this functor on morphisms generating the category Δ are denoted by $d_i^n = X(\partial_n^i)$, $s_i^n = X(\sigma_n^i)$, for all $0 \leq i \leq n$. A simplicial mapping $X \rightarrow Y$ between simplicial sets is a natural transformation.

A simplicial set X is pointed if a vertex $x \in X_0$ is distinguished in it. We call the distinguished vertex x to be base point. A morphism of simplicial sets is said to be pointed if it maps a base point to a base one.

The category of simplicial sets is denoted by $\text{Set}^{\Delta^{op}}$ or sSet . The category of pointed simplicial sets and pointed morphisms is denoted by sSet_* .

The classifying space (nerve) of the small category \mathcal{C} is denoted by $B\mathcal{C}$.

Consider the category of simplices $(\Delta \downarrow X)^{op}$ for the simplicial set X dual to the comma category (in the sense of [25, Chapter 2, §2.6]) over X with respect to the Yoneda embedding functor $\Delta(-) : \Delta \rightarrow \text{Set}^{\Delta^{op}}$. Below we will see (Proposition 2.1) that for an arbitrary category with coproducts \mathcal{A} to every diagram $F : (\Delta \downarrow X)^{op} \rightarrow \mathcal{A}$ there corresponds a simplicial object in \mathcal{A} consisting of coproducts $\coprod_{x \in X_n} F(x)$. This object is used to construct a simplicial replacement of the diagram $F : \mathcal{C} \rightarrow \mathcal{A}$.

We begin our presentation by examining the following question. There are two methods for constructing a simplicial replacement. Bousfield and Kan [3] define it as a simplicial object $\coprod_{c_0 \leftarrow c_1 \leftarrow \dots \leftarrow c_n} F(c_n)$. Quillen [30] uses the notation $\coprod_{c_0 \rightarrow c_1 \rightarrow \dots \rightarrow c_n} F(c_0)$. We will show that these two methods lead to isomorphic simplicial objects.

2.2 Diagrams on the category of simplices

Consider the category of simplices of a simplicial set. Each diagram over this category in the category with coproducts \mathcal{A} corresponds to a simplicial object in \mathcal{A} . We describe its face and degeneration operators.

DEFINITION 2.1 *Let $X : \Delta^{op} \rightarrow \text{Set}$ be a simplicial set. Its category of*

simplices consists of the set of objects $\coprod_{n \geq 0} X_n$. Morphisms of the category of simplices $(m, x) \xrightarrow{\alpha} (n, y)$ are given by nondecreasing mappings $\alpha : [n] \rightarrow [m]$ such that $X(\alpha)(x) = y$. The composition of morphisms is

$$((n, y) \xrightarrow{\beta} (p, z)) \circ ((m, x) \xrightarrow{\alpha} (n, y)) = ((m, x) \xrightarrow{\alpha\beta} (p, z)).$$

The category of simplices is isomorphic to $(\Delta \downarrow X)^{op}$, and we will denote it by the same name.

We need the following statement about diagrams $(\Delta \downarrow X)^{op} \rightarrow \mathcal{A}$ for the category \mathcal{A} with coproducts.

Let $Q_X^{op} : (\Delta \downarrow X)^{op} \rightarrow \Delta^{op}$ denote the forgetful functor defined as $(m, x) \mapsto [m]$ on objects and $(m, x) \xrightarrow{\alpha} (n, y) \mapsto \alpha : [n] \rightarrow [m]$ - on morphisms. Consider the left Kan extension $Lan^{Q_X^{op}} : \mathcal{A}^{(\Delta \downarrow X)^{op}} \rightarrow \mathcal{A}^{\Delta^{op}}$ along the functor Q_X^{op} . We follow the definition of the left Kan extension given in [25, §10.3], with the values of the left Kahn extension being computed pointwise. The following assertion is proved in exactly the same way as the first part of [21, Proposition 3.2] for the case $\mathcal{D} = \Delta$.

Proposition 2.1 *For any diagram $F : (\Delta \downarrow X)^{op} \rightarrow \mathcal{A}$ into the category \mathcal{A} with coproducts, the value of $Lan^{Q_X^{op}} F$ is equal to the simplicial object consisting from objects $Lan^{Q_X^{op}} F[n] = \coprod_{x \in X_n} F(x)$. The boundary operators d_i^n ($n \geq 1$) and degeneracy s_i^n ($n \geq 0$) of this simplicial object are defined as morphisms making the following diagrams commutative for all $x \in X_n$ and $0 \leq i \leq n$:*

$$\begin{array}{ccc} \coprod_{x \in X_n} F(x) & \xrightarrow{d_i^n} & \coprod_{x \in X_{n-1}} F(x) & \quad & \coprod_{x \in X_n} F(x) & \xrightarrow{s_i^n} & \coprod_{x \in X_{n+1}} F(x) \\ \text{in}_x \uparrow & & \uparrow \text{in}_{X(\partial_n^i)x} & & \text{in}_x \uparrow & & \uparrow \text{in}_{X(\sigma_n^i)x} \\ F(x) & \xrightarrow{F(\partial_n^i)} & F(X(\partial_n^i)x) & & F(x) & \xrightarrow{F(\sigma_n^i)} & F(X(\sigma_n^i)x) \\ & & F(x \xrightarrow{\partial_n^i} X(\partial_n^i)x) & & & & F(x \xrightarrow{\sigma_n^i} X(\sigma_n^i)x) \end{array}$$

EXAMPLE 2.2 *If $\mathcal{A} = \text{Grp}$, then for an arbitrary group diagram F over the category of elements $(\Delta \downarrow X)^{op}$ we obtain a simplicial group consisting of free products of the groups $\star_{x \in X_n} F(x)$. The boundary and degeneration operators can be written explicitly*

1. $d_i^n(x \in X_n, f \in F(x)) = (X(\partial_n^i)x, F(x \xrightarrow{\partial_n^i} X(\partial_n^i)x)(f))$,
2. $s_i^n(x \in X_n, f \in F(x)) = (X(\sigma_n^i)x, F(x \xrightarrow{\sigma_n^i} X(\sigma_n^i)x)(f))$.

2.3 Simplicial replacement

The classifying space of a small category \mathcal{C} is its nerve, the simplicial set $B\mathcal{C} = \text{Cat}(-, \mathcal{C})|_{\Delta} : \Delta^{op} \rightarrow \text{Set}$. Its n -dimensional simplices $x \in B_n\mathcal{C}$ can be viewed as functors $x : [n] \rightarrow \mathcal{C}$ or sequences of morphisms $c_0 \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_n} c_n$. The boundary operators $B\mathcal{C}(\partial_n^i) : B_n\mathcal{C} \rightarrow B_{n-1}\mathcal{C}$ map $x \in \text{Cat}([n], \mathcal{C}) \mapsto x \circ \partial_n^i$ or, equivalently, they map to the tuple $(c_0 \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_n} c_n)$, $c_i = x(i)$, one of the sequences

- $(c_0 \xrightarrow{\alpha_1} \dots \rightarrow c_{i-1} \xrightarrow{\alpha_{i+1}\alpha_i} c_{i+1} \rightarrow \dots \xrightarrow{\alpha_n} c_n)$ if $i > 0$,
- $(c_1 \xrightarrow{\alpha_2} \dots \xrightarrow{\alpha_n} c_n)$ if $i = 0$.

The degeneration operators $B\mathcal{C}(\sigma_n^i) : B_n\mathcal{C} \rightarrow B_{n+1}\mathcal{C}$ can be calculated (for all $0 \leq i \leq n$) by the formula $x \mapsto x \circ \sigma_n^i$ or by

$$(c_0 \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_n} c_n) \mapsto (c_0 \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_{i-1}} c_i \xrightarrow{1_{c_i}} c_i \xrightarrow{\alpha_i} \dots \xrightarrow{\alpha_n} c_n).$$

This implies that the categories $\Delta \downarrow \mathcal{C}$ and $\Delta \downarrow B\mathcal{C}$ are naturally isomorphic (with respect to $\mathcal{C} \in \text{Cat}$), and we can consider them as equal categories.

Consider the functors

$$\mathcal{C} \xleftarrow{\partial_0} (\Delta \downarrow B\mathcal{C})^{op} \xrightarrow{Q_{B\mathcal{C}}^{op}} \Delta^{op},$$

the first of which maps each $x : [m] \rightarrow \mathcal{C}$ corresponding to some tuple $(c_0 \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_m} c_m)$ to its object $x(0) = c_0$ and to each morphism $x \rightarrow y$ of the category $(\Delta \downarrow \mathcal{C})$:

$$\begin{array}{ccc} [n] & \xrightarrow{y} & \mathcal{C} \\ & \swarrow g & \nearrow x \\ & [m] & \end{array}$$

morphism $y(0) \xrightarrow{y(0 \leq g(0))} y(g(0)) = x(0)$ of the category \mathcal{C} . This functor ∂_0 is contravariant.

The functor $Q_{B\mathcal{C}}^{op}$ is equal to the functor Q_X^{op} defined above for $X = B\mathcal{C}$.

DEFINITION 2.3 *The simplicial replacement functor $\amalg : \mathcal{A}^{\mathcal{C}} \rightarrow \mathcal{A}^{\Delta^{op}}$ is equal to the composition of the left Kan extension $\text{Lan}^{Q_{B\mathcal{C}}^{op}} : \mathcal{A}^{(\Delta \downarrow B\mathcal{C})^{op}} \rightarrow \mathcal{A}^{\Delta^{op}}$ and the functor $\partial_0^* : \mathcal{A}^{\mathcal{C}} \rightarrow \mathcal{A}^{(\Delta \downarrow B\mathcal{C})^{op}}$ defined by $(F \in \mathcal{A}^{\mathcal{C}}) \mapsto F \circ \partial_0$. In particular, the simplicial change of any diagram F over \mathcal{C} is equal to $\text{Lan}^{Q_{B\mathcal{C}}^{op}}(F\partial_0)$.*

Substituting the composition $F\partial_0$ into the sentence 2.1 instead of F and $X = B\mathcal{C}$, we arrive at the following assertion.

Proposition 2.2 *The simplicial replacement of the diagram $F : \mathcal{C} \rightarrow \mathcal{A}$ consists of the objects $\coprod_n F = \coprod_{c_0 \rightarrow c_1 \rightarrow \dots \rightarrow c_n} F(c_0)$. Boundary operators are defined as morphisms $d_i^n : \coprod_n F \rightarrow \coprod_{n-1} F$, $0 \leq i \leq n$ making commutative the following diagram on the left for every i in the range $0 < i \leq n$, and for $i = 0$ - on the right:*

$$\begin{array}{ccc}
 \coprod_n F & \xrightarrow{d_i^n} & \coprod_{n-1} F \\
 \uparrow \text{in}_{c_0 \rightarrow \dots \rightarrow c_n} & & \uparrow \text{in}_{B\mathcal{C}(\partial_i^n)(c_0 \rightarrow \dots \rightarrow c_n)} \\
 F(c_0) & \xrightarrow{=} & F(c_0)
 \end{array}
 \qquad
 \begin{array}{ccc}
 \coprod_n F & \xrightarrow{d_0^n} & \coprod_{n-1} F \\
 \uparrow \text{in}_{c_0 \rightarrow \dots \rightarrow c_n} & & \uparrow \text{in}_{(c_1 \rightarrow \dots \rightarrow c_n)} \\
 F(c_0) & \xrightarrow{\alpha_1} & F(c_1)
 \end{array}$$

The degeneration operators $s_i^n : \coprod_n F \rightarrow \coprod_{n+1} F$ are defined as morphisms making the following diagrams commutative for all $0 \leq i \leq n$:

$$\begin{array}{ccc}
 \coprod_n F & \xrightarrow{s_i^n} & \coprod_{n+1} F \\
 \uparrow \text{in}_{c_0 \rightarrow \dots \rightarrow c_n} & & \uparrow \text{in}_{B\mathcal{C}(\sigma_i^n)(c_0 \rightarrow \dots \rightarrow c_n)} \\
 F(c_0) & \xrightarrow{=} & F(c_0)
 \end{array}$$

Bousfield and Kan [3] consider the classifying space of the dual category $X = B\mathcal{C}^{op}$ and obtain the simplicial replacement $\coprod' F = Lan_{B\mathcal{C}^{op}}^{Q_B^{op}} F\partial$, where $\partial(c_0 \leftarrow c_1 \leftarrow \dots \leftarrow c_n) = c_n$. Since the simplicial sets $B\mathcal{C}$ and $B\mathcal{C}^{op}$ are isomorphic, the resulting simplicial replacements are isomorphic.

2.4 A homotopy colimit in the category of pointed simplicial sets

A simplicial set X is said to be pointed if a point $\star_0 \in X_0$ is distinguished in it. The distinguished point is called base point. For an arbitrary simplicial set X , we can select any point \star_0 from the set X_0 . Together with the point $\star_0 \in X_0$, for each $n \geq 0$ there will be a single simplex $\star_n = X([n] \rightarrow [0])\star_0 \in X_n$. Since every diagram in the category Δ consisting of morphisms

$$\begin{array}{ccc}
 [0] & \xleftarrow{\quad} & [m] \\
 & \swarrow & \searrow f \\
 & [n] &
 \end{array}$$

is commutative, then faces and degeneracies leave simplices in this sequence, and we can consider a pointed simplicial set as a functor from Δ^{op} to the category of pointed sets whose values are equal to (X_n, \star_n) . The category of pointed simplicial sets is isomorphic to the category of functors $\Delta^{op} \rightarrow \text{Set}_\star$, but homotopy groups are considered with respect to the base point \star_0 .

Let $\text{sSet} = \text{Set}^{\Delta^{op}}$ be the category of simplicial sets, and sSet_\star be the category of pointed simplicial sets and simplicial mappings that preserve the base point.

The homotopy colimit of a diagram of pointed simplicial sets is well described by Bousfield and Kan [3, Chapter XII]. In [3, Lemma 5.2] it is proved that the homotopy colimit functor $\text{hocolim}^{\mathcal{C}} : (\text{sSet}_\star)^{\mathcal{C}} \rightarrow \text{sSet}_\star$ can be defined as functor composition

$$(\text{sSet}_\star)^{\mathcal{C}} \xrightarrow{\amalg} (\text{sSet}_\star)^{\Delta^{op}} = \text{Set}_\star^{\Delta^{op} \times \Delta^{op}} \xrightarrow{\text{diag}} \text{Set}_\star^{\Delta^{op}} = \text{sSet}_\star,$$

where $F \mapsto \amalg F = \text{Lan}^{Q_{\mathcal{C}}^{op}} F \partial_0$ is the simplicial replacement functor for the case $\mathcal{A} = \text{sSet}_\star$ and diag - the diagonal functor [3, Chapter XII], corresponding to each bisimplicial set consisting of the sets

$$\{X([n], [p])\}_{([n],[p]) \in \Delta^{op} \times \Delta^{op}}$$

the simplicial set $\{X([n], [n])\}_{([n]) \in \Delta^{op}}$.

Let $F : \mathcal{C} \rightarrow \text{sSet}_\star$ be an arbitrary diagram of pointed simplicial sets. Its simplicial replacement is equal to a bisimplicial set whose vertical (in the sense of [18, Page 209]) simplicial sets are pointed and equal for each $n \geq 0$ to the coproduct (and hence to the wedge) $\bigvee_{c_0 \rightarrow c_1 \rightarrow \dots \rightarrow c_n} F(c_0)$ over all sequences of morphisms of length n .

Therefore, the homotopy colimit, as the diagonal of the resulting simplicial set, is equal to the pointed simplicial set

$$(\text{hocolim}^{\mathcal{C}} F)_n = \bigvee_{c_0 \rightarrow c_1 \rightarrow \dots \rightarrow c_n} F(c_0)_n, \text{ for all } n \geq 0.$$

EXAMPLE 2.4 *Let $\Delta_{\mathcal{C}} Y : \mathcal{C} \rightarrow \text{sSet}_\star$ be a diagram taking constant values Y . In order to construct $\text{hocolim}^{\mathcal{C}} \Delta_{\mathcal{C}} Y$, we first construct a homotopy colimit of the diagram in which the base points are treated as ordinary. Let $U : \text{sSet}_\star \rightarrow \text{sSet}$ be a functor forgetting base points. Get*

$$(\text{diag} \amalg \Delta_{\mathcal{C}} Y)_n = \amalg_{c_0 \rightarrow c_1 \rightarrow \dots \rightarrow c_n} Y_n = B_n \mathcal{C} \times Y_n.$$

Hence the simplicial set $\text{hocolim}^{\mathcal{C}} U(\Delta_{\mathcal{C}} Y)$ is equal to $B\mathcal{C} \times Y$.

Returning to $\text{hocolim}^{\mathcal{C}} \Delta_{\mathcal{C}} Y$, we find that it consists of pointed sets

$$\bigvee_{c_0 \rightarrow c_1 \rightarrow \dots \rightarrow c_n} Y_n$$

equal to the sets $B_n \mathcal{C} \times Y_n$ with identified simplices from $B_n \mathcal{C} \times \star_n$. Therefore, the homotopy colimit of a diagram on \mathcal{C} taking constant values equal to the pointed simplicial set Y is equal to the pointed simplicial factor set

$$(B\mathcal{C} \times Y) / (B\mathcal{C} \times \star).$$

3 Classifying space of a simplicial group

This section is devoted to the homotopy groups of the simplicial group. To every simplicial group there corresponds a diagram of nerves over Δ^{op} . An auxiliary lemma is considered that the n th homotopy groups of the homotopy colimit of the nerve diagram are isomorphic to the $(n-1)$ th homotopy groups of the simplicial group for all $n \geq 1$.

We adhere to the definition of the homotopy group $\pi_n(X)$ of a pointed simplicial set given in [16, Chapter VI, §3]. In particular, when defining the homotopy groups of a simplicial group its group structure can be ignored. But it is a fibrant simplicial set, and surjective morphisms of simplicial groups are (Kan) fibrations.

3.1 Moore complex and its homology groups

Let $G : \Delta^{op} \rightarrow \text{Grp}$ be a simplicial group.

We describe the Moore complex [28, Ch.II, §3.7] for calculating the homotopy groups $\pi_n(G)$. For this purpose, consider the complex consisting of the groups $N_n G = \bigcap_{i>0} \text{Ker}(d_i^n : G_n \rightarrow G_{n-1})$ for $n > 0$, and $N_0 G = G_0$.

Its differentials $d^n : N_n G \rightarrow N_{n-1} G$ are defined by the restriction of homomorphisms d_0^n for $n > 0$, and $d^0 = 0$.

The homotopy groups $\pi_n(G)$ are isomorphic to the homology groups of this complex (see [35, Chapter 3, Theorem 1.5]):

$$H_n(NG) = \text{Ker}(d^n : N_n G \rightarrow N_{n-1} G) / \text{Im}(d^{n+1} : N_{n+1} G \rightarrow N_n G).$$

For example, for a constant simplicial group $G = \Delta G_0$, this gives isomorphisms $\pi_0(G) \cong G_0$ and $\pi_n(G) = 0$ for all $n > 0$.

Note that the groups $\pi_n(G)$ (see [34, Exercise 8.3.2] or [35, Chapter 3, Theorem 1.5]) are Abelian for $n \geq 1$, while for $n = 0$ they can be non-commutative.

3.2 Homotopies of a simplicial group and its classifying space

Consider an arbitrary bisimplicial set $X(-, =)$. Let $\text{diag } X(-, =)$ denote the diagonal of a double simplicial set. It is a simplicial set defined as $\text{diag } X([n]) = X([n], [n])$. A bisimplicial set $X(-, =)$ is called pointed if for every $m \geq 0$ the simplicial set $X(m, =)$ is pointed, i.e. has a marked vertex $x_m \in X(m, 0)$, and for each morphism $\alpha \in \Delta([m], [n])$ the equality $X(\alpha)(x_m) = x_n$. Any bisimplicial set becomes pointed if we distinguish an element $x_0 \in X(0, 0)$ in it.

Let $G : \Delta^{op} \rightarrow \text{Grp}$ be a simplicial group. Consider the bisimplicial set $G(-)^{[=]}$ formed from the nerves of the groups and consisting of the sets $G(m)^{[n]} = B_n(G(m))$. We will consider it as a pointed simplicial set whose base point is the only element of the set $B_0(G(0))$

Lemma 3.1 *Let $G : \Delta^{op} \rightarrow \text{Grp}$ be a simplicial group. Then for all $n \geq 1$ there are natural isomorphisms*

$$\pi_n(\text{diag } G(-)^{[=]}) \cong \pi_{n-1}G.$$

PROOF. For each $p \geq 0$ the classifying space of the group $G(p)$ is connected. We choose a base point equal to the unique 0-dimensional simplex of the space $BG(p)$ for each $p \geq 0$. One can use the Bousfield-Friedlander spectral sequence [2] or the Zisman spectral sequence (see [7, Appendix A]) for the bisimplicial set $G(-)^{[=]}$:

$$E_{p,q}^2 = \pi_p^h \pi_q^v G(-)^{[=]} \Rightarrow \pi_{p+q}(\text{diag } G(-)^{[=]}).$$

For every $p \geq 0$ the groups $\pi_q G(p)^{[=]}$ are isomorphic to $G(p)$ for $q = 1$, and trivial for $q \neq 1$. This leads to degeneration of the spectral sequence and isomorphisms $E_{p,1}^2 \cong \pi_{p+1} \text{diag } G(-)^{[=]}$. Hence $\pi_p(G) \cong \pi_{p+1}(\text{diag } G(-)^{[=]})$ for all $p \geq 0$. \square

REMARK 3.1 *There are other methods for proving Lemma 3.1, for example, using the exact sequence of homotopy groups for the fibration $G \rightarrow \text{diag } EG \rightarrow \text{diag } BG$ leading to the natural isomorphism $\pi_n \text{diag}(BG) \cong \pi_{n-1}G$ [18, Page 244].*

4 Non-Abelian homology with coefficients in a group diagram

In this section, non-Abelian homology of a small category with coefficients in the group diagram is introduced. They are constructed as homotopy groups of a simplicial replacement of the group diagram. This replacement is a simplicial group, and hence the n th homology for all $n \geq 0$ are groups, commutative for $n \geq 1$.

4.1 Simplicial replacement of group diagram

For an arbitrary small category \mathcal{C} , the following functors are defined

$$\mathcal{C} \xleftarrow{\partial_0} (\Delta \downarrow \mathcal{C})^{op} \xrightarrow{Q_{\mathcal{C}}^{op}} \Delta^{op}.$$

Let \mathcal{A} be a category with coproducts. For each diagram $F : \mathcal{C} \rightarrow \mathcal{A}$ its simplicial replacement $\coprod F \in \mathcal{A}^{\Delta^{op}}$ is defined as $\coprod F = \text{Lan}^{Q_{\mathcal{C}}^{op}}(F \circ \partial_0)$. The object of n -dimensional simplices of the simplicial object $\coprod F$ will be equal to $(\coprod F)_n = \coprod_{c_0 \rightarrow \dots \rightarrow c_n} F(c_0)$. In particular, for the category of groups Grp, it will be equal to the free product $\star_{c_0 \rightarrow \dots \rightarrow c_n} F(c_0)$, for the category of abelian groups Ab it will be equal to the direct sum $\oplus_{c_0 \rightarrow \dots \rightarrow c_n} F(c_0)$, and in the case of the category of pointed simplicial sets sSet_\star it is equal to the wedge $\bigvee_{c_0 \rightarrow \dots \rightarrow c_n} F(c_0)$.

Proposition 4.1 *Let $\mathcal{G} : \mathcal{C} \rightarrow \text{Grp}$ be a group diagram. The simplicial group $\coprod \mathcal{G}$ consists of the groups*

$$C_n(\mathcal{C}, \mathcal{G}) = \star_{c_0 \rightarrow \dots \rightarrow c_n} \mathcal{G}(c_0).$$

Its boundary operators $d_n^i : C_n(\mathcal{C}, \mathcal{G}) \rightarrow C_{n-1}(\mathcal{C}, \mathcal{G})$ and degeneration operators $s_n^i : C_n(\mathcal{C}, \mathcal{G}) \rightarrow C_{n+1}(\mathcal{C}, \mathcal{G})$, for $0 \leq i \leq n$, act on the elements of the

factors of free products by the formulas

$$\begin{aligned}
d_n^i(c_0 \xrightarrow{\alpha_1} c_1 \rightarrow \cdots \rightarrow c_{n-1} \xrightarrow{\alpha_n} c_n, x) &= \\
&= \begin{cases} (c_1 \xrightarrow{\alpha_2} c_2 \rightarrow \cdots \rightarrow c_{n-1} \xrightarrow{\alpha_n} c_n, \mathcal{G}(\alpha_1)(x)), & \text{if } i = 0, \\ (c_0 \xrightarrow{\alpha_1} c_1 \rightarrow \cdots \rightarrow c_{i-1} \xrightarrow{\alpha_{i+1}\alpha_i} c_{i+1} \rightarrow \cdots \xrightarrow{\alpha_n} c_n, x), & \text{if } i > 0, \end{cases} \quad (3)
\end{aligned}$$

$$\begin{aligned}
s_n^i(x, c_0 \xrightarrow{\alpha_1} c_1 \rightarrow \cdots \rightarrow c_{n-1} \xrightarrow{\alpha_n} c_n) &= \\
&= (c_0 \xrightarrow{\alpha_1} c_1 \rightarrow \cdots \rightarrow c_i \xrightarrow{id} c_i \rightarrow \cdots \rightarrow c_{n-1} \xrightarrow{\alpha_n} c_n, x), \quad (4)
\end{aligned}$$

where x denotes an arbitrary element of the group $\mathcal{G}(c_0)$.

4.2 Non-Abelian group diagram homology

According to Proposition 4.1, the simplicial group $C_*(\mathcal{C}, \mathcal{G})$ can be used as a simplicial replacement of a group diagram. We note this in the following definition of non-abelian homology:

DEFINITION 4.1 *For an arbitrary small category \mathcal{C} and a group diagram $\mathcal{G} : \mathcal{C} \rightarrow \text{Grp}$, the n th homologies of the category \mathcal{C} with coefficients in \mathcal{G} are the groups $\text{colim}_n^{\mathcal{C}} \mathcal{G} := \pi_n(\coprod \mathcal{G}) = \pi_n(C_*(\mathcal{C}, \mathcal{G}))$, for all $n \geq 0$.*

For the case when the category \mathcal{C} is a finite poset, one can use the formula for homotopy groups of a simplicial group proved by Castiglioni and Ladra [8, Theorem 25] to calculate non-Abelian homology. We explain below that this can be done.

Let K_0, K_1, \dots, K_m be normal subgroups of an arbitrary group G . The fat commutator subgroup of these subgroups is the subgroup

$$[[K_0, K_1, \dots, K_m]] := \prod_{I \sqcup J = \{0, 1, \dots, m\}} [\cap_{i \in I} K_i, \cap_{j \in J} K_j].$$

For example, $[[K_0, K_1]] = [K_0, K_1]$; $[[K_0, K_1, K_2]] = [K_0, K_1 \cap K_2][K_1, K_0 \cap K_2][K_2, K_0 \cap K_1]$.

Proposition 4.2 *Let $n \geq 1$ be a natural number, and let \mathcal{C} be a small category in which for each sequence morphisms $c_0 \xrightarrow{\alpha_1} c_1 \xrightarrow{\alpha_2} c_2 \rightarrow \dots \rightarrow c_{k-1} \xrightarrow{\alpha_k} c_k$ that are not equal to the identity, the relation $k \leq n$ is true. Then*

for each diagram $\mathcal{G} : \mathcal{C} \rightarrow \text{Grp}$ the non-Abelian homology for all $m \geq n$ is isomorphic to the following quotient groups

$$\text{colim}_m^{\mathcal{C}} \mathcal{G} \cong (K_0 \cap K_1 \cap \cdots \cap K_m) / [[K_0, K_1, \dots, K_m]],$$

where $K_i = \text{Ker } d_i^m$ for all $0 \leq i \leq m$.

PROOF. The simplex $(c_0 \rightarrow c_1 \rightarrow \dots \rightarrow c_k, x \in \mathcal{G}(c_0))$ of the simplicial group $C_*(\mathcal{C}, \mathcal{G})$ is degenerate if and only if among the morphisms $c_i \rightarrow c_{i+1}$, $0 \leq i \leq k-1$, there is at least one identity. Hence, under the given conditions, the group $C_{n+1}(\mathcal{C}, \mathcal{G})$ is generated by degenerate simplices. According to [8], this implies that the image of the $(n+1)$ th differential of the Moore complex, $d^{n+1}(N_{n+1}C_*(\mathcal{C}, F))$, is $[[K_0, K_1, \dots, K_n]]$, and $\text{colim}_n^{\mathcal{C}} \mathcal{G} \cong K_0 \cap K_1 \cap \cdots \cap K_n / [[K_0, K_1, \dots, K_n]]$. If $m \geq n$, then $C_{m+1}(\mathcal{C}, \mathcal{G})$ is all the more generated by degenerate simplices. Hence, the formula being proved is true for all $m \geq n$. \square

5 Relationship between non-Abelian homology and homotopy colimits

In this section, we prove that the n th non-Abelian homology of a group diagram is isomorphic to the $(n+1)$ th homotopy groups of the homotopy colimit for the diagram of the classifying spaces.

5.1 Homotopy groups of a wedge for classifying spaces

We will need a lemma on the permutability of the functor of a classifying space with coproducts up to homotopy.

Coproduct in the category of pointed simplicial sets will be called a wedge. According to [3, Ch.XII, §3.1], the wedge is weakly equivalent to the homotopy colimit of a diagram over a discrete category in the category of pointed simplicial sets.

Lemma 5.1 *For an arbitrary family of groups, there is a weak equivalence $\bigvee_{i \in I} BG_i \approx B(\bigstar_{i \in I} G_i)$.*

PROOF. Note that this assertion follows from a more general theorem due to Whitehead [15, Theorem 4.0]. It will suffice for us to use the natural weak

equivalence $B(G_1 * G_2) \approx BG_1 \vee BG_2$ following from the assertion proved in [9, Lemma 3.8].

In the case of a finite set I this assertion can be proved by induction. If I is infinite, then we use the weak equivalence $\bigvee_{i \in J} BG_i \approx B(\bigstar_{i \in J} G_i)$ for each finite subset of $J \subseteq I$. Passing to a colimit with respect to a set of finite subsets and using the assertion [3, Ch XII, §3.5] that the colimit of pointed simplicial sets with respect to a directed category is weakly equivalent to a homotopy colimit, we obtain the assertion of this lemma. \square

5.2 Non-Abelian homology and homotopy groups of the homotopy colimit

We consider homotopy colimits in the category of pointed simplicial sets. The homotopy colimit of classifying spaces of groups will be considered as a pointed simplicial set having a single vertex.

Theorem 5.2 *For an arbitrary group diagram $\mathcal{G} : \mathcal{C} \rightarrow \text{Grp}$ and for all $n \geq 0$ there are natural isomorphisms*

$$\text{colim}_n^{\mathcal{C}} \mathcal{G} \cong \pi_{n+1}(\text{hocolim}^{\mathcal{C}} B\mathcal{G}). \quad (5)$$

PROOF. By Lemma 3.1, for every simplicial group $G \in \text{Grp}^{\Delta^{op}}$ there exists a natural isomorphism

$$\pi_{n+1} \text{diag } BG \cong \pi_n(G), \text{ for all } n \geq 0, \quad (6)$$

which we want to use.

Using Lemma 5.1, for each $n \geq 0$, we arrive at a weak equivalence

$$(B \coprod \mathcal{G})_n = B \bigstar_{c_0 \rightarrow \dots \rightarrow c_n} \mathcal{G}(c_0) \approx \bigvee_{c_0 \rightarrow \dots \rightarrow c_n} B\mathcal{G}(c_0) = (\coprod B\mathcal{G})_n.$$

This weak equivalence will be natural in $[n] \in \Delta$. Consider pointed bisimplicial sets $B \coprod \mathcal{G}$ and $\coprod B\mathcal{G}$ as diagrams $\Delta^{op} \rightarrow \text{sSet}_*$. The homotopy colimit in $[n] \in \Delta^{op}$ leads to a weak equivalence

$$\text{hocolim}^{\Delta^{op}} B \coprod \mathcal{G} \approx \text{hocolim}^{\Delta^{op}} \coprod B\mathcal{G}.$$

For every pointed bisimplicial set $Y \in \text{sSet}_*^{\Delta^{op}}$ there exists a natural (in Y) weak equivalence $\text{hocolim}^{\Delta^{op}} Y \approx \text{diag } Y$ [3, ch XII, §4.3]. This leads to a weak equivalence of the diagonals $\text{diag } B \coprod \mathcal{G} \approx \text{diag } \coprod B\mathcal{G}$.

Using this weak diagonal equivalence and substituting $G = \coprod \mathcal{G}$ into the formula (6), we arrive at natural isomorphisms (with respect to \mathcal{G}):

$$\pi_{n+1}(\text{diag } \coprod B\mathcal{G}) \cong \pi_{n+1}(\text{diag } B \coprod \mathcal{G}) \cong \pi_n(\coprod \mathcal{G}), \text{ for all } n \geq 0. \quad (7)$$

By the simplicial replacement lemma [3, Ch XII, 5.2], for any diagram of pointed simplicial sets, the homotopy colimit $\text{hocolim}^{\mathcal{C}} Y$ is naturally weakly equivalent to $\text{diag } \coprod Y$. So the left side of the formula (7) is equal to $\pi_{n+1}(\text{hocolim}^{\mathcal{C}} B\mathcal{G})$. By Definition 4.1 the right side of the isomorphism (7) is equal to $\text{colim}_n^{\mathcal{C}} \mathcal{G}$. Formula (5) received. \square

REMARK 5.1 *A similar formula in other terms is given by Toën in [27, Page 136].*

REMARK 5.2 *To construct a pointed homotopy colimit using Thomason's theorem [33, Theorem 1.2], additional steps are needed, since Thomason's theorem has been proved for nonpointed homotopy colimits.*

In the case $\mathcal{C} = \Delta^{op}$ there will be isomorphisms

$$\text{colim}_n^{\Delta^{op}} \mathcal{G} \cong \pi_{n+1} \text{hocolim}^{\Delta^{op}} B\mathcal{G}.$$

Whence, using the natural weak equivalence $\text{hocolim}^{\Delta^{op}} B\mathcal{G} \rightarrow \text{diag } B\mathcal{G}$ [3, Ch XII.3.4] and Lemma 3.1 showing that $\pi_{n+1}(\text{diag } B\mathcal{G}) \cong \pi_n(\mathcal{G})$, $n \geq 0$, we get

Corollary 5.3 *For an arbitrary simplicial group $\mathcal{G} \in \text{Grp}^{\Delta^{op}}$ and a non-negative integer n , there is an isomorphism*

$$\text{colim}_n^{\Delta^{op}} \mathcal{G} \cong \pi_n(\mathcal{G}).$$

For Abelian homology this formula was proved in [3, XII.5.6] when \mathcal{G} is a simplicial Abelian group.

5.3 Abelian homology of diagrams over a free category

Let $\mathcal{A} : \mathcal{C} \rightarrow \text{Ab}$ be a diagram of abelian groups. For the definition of n th Abelian homology $\varinjlim_n^{\mathcal{C}} \mathcal{A}$ one can take the homotopy groups $\pi_n(\coprod \mathcal{A})$ of the simplicial change of the diagram of Abelian groups \mathcal{A} . They will be isomorphic to the homology groups of the simplicial abelian group consisting

of direct sums $C_n(\mathcal{C}, \mathcal{A}) = \bigoplus_{c_0 \rightarrow \dots \rightarrow c_n} \mathcal{A}(c_0)$ with boundary operators d_n^i and degeneracy operators s_n^i acting on the elements of the terms of direct sums according to the formulas (3)-(4) from the proposition 4.1. The Abelian homology groups are isomorphic to the homology groups of the complex of Abelian groups

$$0 \xleftarrow{d_0} C_0(\mathcal{C}, \mathcal{A}) \xleftarrow{d_1} C_1(\mathcal{C}, \mathcal{A}) \xleftarrow{d_2} \dots \xleftarrow{d_n} C_n(\mathcal{C}, \mathcal{A}) \xleftarrow{d_{n+1}} \dots$$

whose differentials are $d_0 = 0$ and $d_n = \sum_{0 \leq i \leq n} (-1)^n d_n^i$, for $n \geq 1$. Thus,

$$\varinjlim_n^{\mathcal{C}} \mathcal{A} = \text{Ker } d_n / \text{Im } d_{n+1}, \text{ for all } n \geq 0.$$

Let $\mathcal{C} = W\Gamma$ be the free category generated by the graph $\Gamma = (A \overset{t}{\rightrightarrows} V)$, where A is the set of arrows (directed edges), V is a set of vertices, $s : A \rightarrow V$ assigns each arrow its beginning, and $t : A \rightarrow V$ its end.

To calculate $\varinjlim_1^{W\Gamma} \mathcal{A}$, [20, Definition 2.1] proposed the following

DEFINITION 5.3 *A flow on the graph $\Gamma = (A \overset{t}{\rightrightarrows} V)$ with coefficients in the diagram $\mathcal{A} : W\Gamma \rightarrow \text{Ab}$ is an element $f = (f_\gamma)_{\gamma \in A} \in \bigoplus_{\gamma \in A} \mathcal{A}(s(\gamma))$ such that for every vertex $v \in V$ there is the following ratio*

$$\sum_{s(\gamma) \in v} f_\gamma = \sum_{t(\gamma) = v} \mathcal{A}(\gamma)(f_\gamma).$$

The set of flows is a subgroup of the Abelian group $\bigoplus_{\gamma \in A} \mathcal{A}(s(\gamma))$. It is proved [20, Theorem 2.5] that this flow subgroup is isomorphic to $\varinjlim_1^{W\Gamma} \mathcal{A}$.

EXAMPLE 5.4 *Consider a graph consisting of two vertices a, b and two arrows $a \xrightarrow{u_0} b$ and $a \xrightarrow{u_1} b$. Denote by $\uparrow\uparrow$ the free category generated by this graph; it is obtained from the graph by adding the identical morphisms 1_a and 1_b . Let $\mathcal{A} : \uparrow\uparrow \rightarrow \text{Ab}$ be a digram of abelian groups. The flow on the graph generating $\uparrow\uparrow$ is given by a pair of elements $f_{u_0}, f_{u_1} \in \mathcal{A}(a)$ such that $f_{u_0} + f_{u_1} = 0$ and $c\mathcal{A}(u_0)(f_{u_0}) + \mathcal{A}(u_1)(f_{u_1}) = 0$. It follows that each flow is given by an arbitrary element $f_{u_0} \in \mathcal{A}(a)$ such that $(\mathcal{A}(u_0) - \mathcal{A}(u_1))(f_{u_0}) = 0$. Hence $\varinjlim_1^{\uparrow\uparrow} \mathcal{A} = \text{Ker}(\mathcal{A}(u_0) - \mathcal{A}(u_1))$.*

5.4 Comparison and application of Abelian and non-Abelian homologies of group diagrams

Let us apply the non-Abelian and Abelian homology of group diagrams to develop a method for finding a non-zero homotopy group of least dimension for the homotopy colimit of classifying group spaces.

For the case when the colimit of a group diagram is equal to the trivial group, for diagrams defined on free categories, we study the conditions under which the first non-Abelian and Abelian homology groups are isomorphic.

The content of this subsection is based on the following statement, which follows from Theorem 5.2.

Corollary 5.4 *For an arbitrary small category \mathcal{C} and a diagram $\mathcal{G} : \mathcal{C} \rightarrow \text{Grp}$ there is an isomorphism*

$$\pi_1(\text{hocolim}^{\mathcal{C}} B\mathcal{G}) \cong \text{colim}^{\mathcal{C}} \mathcal{G}. \quad (8)$$

In particular, a homotopy colimit of a diagram of classifying spaces is simply connected if and only if the colimit of a group diagram is equal to the trivial group.

EXAMPLE 5.5 *It would not be superfluous to check the formula (8) for a diagram that takes the values $\mathcal{G}(c) = 0$, for all $c \in \text{Ob } \mathcal{C}$. Since $B(0) = \star$, then according to Example 2.4, we get $\text{hocolim}^{\mathcal{C}} (\Delta_{\mathcal{C}} \star) = B\mathcal{C} \times \star / B\mathcal{C} \times \star = \star$ and isomorphism (8).*

Consider the problem of finding $n \geq 0$ such that $\text{colim}_k^{\mathcal{C}} \mathcal{G} = 0$ for all $0 \leq k < n$ and $\text{colim}_n^{\mathcal{C}} \mathcal{G} \neq 0$. This number n will be called the connectivity of the group diagram \mathcal{G} and denoted by $\text{con } \mathcal{G}$. If $\text{colim}_n^{\mathcal{C}} \mathcal{G} = 0$ for all $n \geq 0$, then we say that the diagram's connectivity is infinite and write $\text{con } \mathcal{G} = \infty$. The connectivity of the diagram \mathcal{G} is equal to the largest n for which the space $\text{hocolim}^{\mathcal{C}} B\mathcal{G}$ is n -connected.

Let $\uparrow\uparrow$ be the category generated by the graph consisting of a pair of parallel arrows, considered in Example 5.4.

EXAMPLE 5.6 *Let $\mathcal{G} : \uparrow\uparrow \rightarrow \text{Grp}$ be a diagram taking the values $\mathcal{G}(b) = S_3$ be the group of permutations of three elements, $\mathcal{G}(a) = A_3$ be the subgroup of even permutations, $\mathcal{G}(u_0) : A_3 \rightarrow S_3$ is an embedding of an alternating group into a symmetric group, $\mathcal{G}(u_1) : A_3 \rightarrow S_3$ is a homomorphism that takes constant values equal to the neutral element.*

Since $\text{colim}^{\uparrow} \mathcal{G} = \text{Coker } \mathcal{G}(u_0) = S_3/A_3 = \mathbb{Z}_2$, we get that the group $\pi_1(\text{hocolim}^{\mathcal{C}} B\mathcal{G}) = \mathbb{Z}_2$ is not trivial, and $\text{con } \mathcal{G} = 0$.

If in the considered diagram \mathcal{G} we replace the group A_3 by a cyclic subgroup of the second order $\mathbb{Z}_2 \subset S_3$, then we get the equality $\text{colim}^{\uparrow} \mathcal{G} = S_3/\langle \mathbb{Z}_2 \rangle^{S_3} = 0$ and $\text{con } \mathcal{G} \geq 1$.

Before returning to Example 5.6, we examine the conditions under which the values of the non-Abelian homology functor $\text{colim}_1^{\mathcal{C}} : \text{Grp}^{\mathcal{C}} \rightarrow \text{Grp}$ are isomorphic to the values of the Abelian homology functor $\varinjlim_1^{\mathcal{C}} : \text{Ab}^{\mathcal{C}} \rightarrow \text{Ab}$.

For this purpose we need the spectral sequence of Bousfield and Kan [3, Ch. XII, §5.7], which exists for an arbitrary diagram of pointed sets $X(-) : \mathcal{C} \rightarrow \text{sSet}$ and is located in the first quadrant

$$E_{p,q}^2 = \varinjlim_p^{\mathcal{C}} \{ \tilde{H}_q X(c) \}_{c \in \mathcal{C}} \Rightarrow \tilde{H}_{p+q}(\text{hocolim}^{\mathcal{C}} X), \quad (9)$$

where \tilde{H}_n denote the reduced homology groups of pointed simplicial sets [16, Appendix II, §2], $n \geq 0$.

Proposition 5.5 *Let $\mathcal{G} : W\Gamma \rightarrow \text{Grp}$ be a group diagram over the free category $W\Gamma$ generated by an arbitrary graph in the sense of [25, §2.7]. We will assume that $\text{colim}^{W\Gamma} \mathcal{G} = 0$. There is a natural (in $\mathcal{G} \in \text{Grp}^{W\Gamma}$) group homomorphism*

$$f : \text{colim}_1^{W\Gamma} \mathcal{G} \rightarrow \varinjlim_1^{W\Gamma} \mathcal{G}_{ab},$$

where \mathcal{G}_{ab} is the Abelianization of the diagram \mathcal{G} . This homomorphism f is an isomorphism if and only if $\text{colim}^{W\Gamma} H_2(\mathcal{G}) = 0$. In particular, f is an isomorphism if \mathcal{G} is a diagram of free groups.

PROOF. The non-Abelian homology group $\text{colim}_0^{W\Gamma} \mathcal{G}$ in dimension $n = 0$ is isomorphic to $\text{colim}^{W\Gamma} \mathcal{G}$, and hence is equal to 0 by assumption.

Let us calculate the first non-Abelian homology group of the diagram $\mathcal{G} : W\Gamma \rightarrow \text{Grp}$. Since $\text{colim}^{W\Gamma} \mathcal{G} = 0$, then $\pi_1(\text{hocolim}^{W\Gamma} B\mathcal{G}) = 0$ (Corollary 5.4). It follows from the Hurewicz theorem that the group $\pi_2(\text{hocolim}^{W\Gamma} B\mathcal{G})$ is isomorphic to the second integral homology group $H_2(\text{hocolim}^{W\Gamma} B\mathcal{G})$. To calculate this group, we use the spectral sequence (9) for the diagram $X = B\mathcal{G}$ of classifying spaces. In this case, it will degenerate for all $n \geq 2$ into short exact sequences

$$0 \rightarrow \text{colim}^{W\Gamma} H_n(B\mathcal{G}) \rightarrow H_n(\text{hocolim}^{W\Gamma} B\mathcal{G}) \rightarrow \varinjlim_1^{W\Gamma} H_{n-1}(B\mathcal{G}) \rightarrow 0 \quad (10)$$

where $\varinjlim_1^{W\Gamma} H_{n-1}(B\mathcal{G})$ denotes the first Abelian homology group of the category $W\Gamma$ with coefficients in the diagram $H_{n-1}(B\mathcal{G})$.

Since $H_2(\text{hocolim}^{W\Gamma} B\mathcal{G}) \cong \pi_2(\text{hocolim}^{W\Gamma} B\mathcal{G}) \cong \text{colim}_1^{W\Gamma} \mathcal{G}$, and there are isomorphisms $H_1(B\mathcal{G}(c)) \cong \mathcal{G}(c)_{ab}$, $H_2(B(\mathcal{G}(c))) \cong H_2(\mathcal{G}(c))$, then for $n = 2$ the exact sequence (10) leads to the exact sequence natural in \mathcal{G}

$$0 \rightarrow \text{colim}^{W\Gamma} H_2(\mathcal{G}) \rightarrow \text{colim}_1^{W\Gamma} \mathcal{G} \rightarrow \varinjlim_1^{W\Gamma} \mathcal{G}_{ab} \rightarrow 0.$$

It shows that the homomorphism $f : \text{colim}_1^{W\Gamma} \mathcal{G} \rightarrow \varinjlim_1^{W\Gamma} \mathcal{G}_{ab}$ is an isomorphism if and only if $\text{colim}^{W\Gamma} H_2(\mathcal{G}) = 0$.

In particular, if the diagram \mathcal{G} consists of free groups, then it can be given using the presentation $\langle E|R \rangle$, where $R = \{1\}$, whence, by Hopf's formula (see [4, Theorem II.5.3]), the homology groups $H_2(\mathcal{G}(c))$ are equal to 0, which means that in this case the homomorphism f is an isomorphism. \square

The following example continues the group diagram exploration started in Example 5.6.

EXAMPLE 5.7 *Our goal is to find the connectivity of the diagram*

$$\mathcal{G} = \left(\mathbb{Z}_2 \begin{array}{c} \xrightarrow{\leq} \\ \xrightarrow{0} \end{array} S_3 \right),$$

which takes on objects the values $\mathcal{G}(a) = \mathbb{Z}_2$, $\mathcal{G}(b) = S_3$, and on morphisms $\mathcal{G}(u_0) : \mathbb{Z}_2 \rightarrow S_3$ is the embedding of a cyclic subgroup of order 2 into the group S_3 , and $\mathcal{G}(u_1)$ is a homomorphism onto a trivial subgroup. We have established (Example 5.6) that $\text{colim}^{\uparrow\uparrow} \mathcal{G} = 0$, whence the space $\text{hocolim}^{\uparrow\uparrow} B\mathcal{G}$ is simply connected. By the Hurewicz theorem, we get $\pi_2(\text{hocolim}^{\uparrow\uparrow} B\mathcal{G}) \cong H_2(\text{hocolim}^{\uparrow\uparrow} B\mathcal{G})$.

Since $\uparrow\uparrow$ is a free category, then the sequence (10) will be exact, and for $n = 2$ it will lead to the exact sequence

$$0 \rightarrow \text{colim}^{\uparrow\uparrow} H_2(\mathcal{G}) \rightarrow H_2(\text{hocolim}^{\uparrow\uparrow} B\mathcal{G}) \rightarrow \varinjlim_1^{\uparrow\uparrow} H_1(\mathcal{G}) \rightarrow 0 \quad (11)$$

Since $H_2(\mathbb{Z}_2) = 0$ (see, for example, [24, §4.2]) and $H_2(S_3) = 0$ [36], then the diagram $H_2(\mathcal{G})$ is equal to the pair of trivial group homomorphisms $0 \rightrightarrows 0$, whence $\text{colim}^{\uparrow\uparrow} H_2(\mathcal{G}) = 0$. The diagram $H_1(\mathcal{G})$ consists of two homomorphisms $\mathbb{Z}_2 \rightrightarrows (S_3)_{ab} = \mathbb{Z}_2$, one of which is the identity since $\mathcal{G}(u_0)$ is a coretraction, and the other is equal to zero homomorphism $H_1(\mathcal{G}(u_1)) = \mathcal{G}(u_1)_{ab}$, because $\mathcal{G}(u_1)$ is the null homomorphism. Applying the method of calculating

the values of the functor $\varinjlim_1^{\uparrow\uparrow}$ (example 5.4), we obtain $\varinjlim_1^{\uparrow\uparrow} H_1(\mathcal{G}) = 0$. The exact sequence (11) leads to $H_2(\text{hocolim}^{\uparrow\uparrow} B\mathcal{G}) = 0$. Hence $\text{colim}_1^{\uparrow\uparrow} \mathcal{G} = 0$, $\text{con } \mathcal{G} \geq 2$ (the space $\text{hocolim}^{\uparrow\uparrow} B\mathcal{G}$ is 2-connected) and

$$\pi_3(\text{hocolim}^{\uparrow\uparrow} B\mathcal{G}) \cong H_3(\text{hocolim}^{\uparrow\uparrow} B\mathcal{G})$$

by the Hurewicz theorem.

To calculate $H_3(\text{hocolim}^{\uparrow\uparrow} B\mathcal{G})$ we have constructed an exact sequence (10), which for $n = 3$ is equal to the exact sequence

$$0 \rightarrow \text{colim}^{\uparrow\uparrow} H_3(\mathcal{G}) \rightarrow H_3(\text{hocolim}^{\uparrow\uparrow} B\mathcal{G}) \rightarrow \varinjlim_1^{\uparrow\uparrow} H_2(\mathcal{G}) \rightarrow 0 \quad (12)$$

We have shown that the diagram $H_2(\mathcal{G})$ consists of zeros, which implies that $\varinjlim_1^{\uparrow\uparrow} H_2(\mathcal{G}) = 0$ and there is an isomorphism $H_3(\text{hocolim}^{\uparrow\uparrow} B\mathcal{G}) \cong \text{colim}^{\uparrow\uparrow} H_3(\mathcal{G})$. The diagram $H_3(\mathcal{G})$ consists of the Abelian groups $H_3(\mathcal{G}(a)) = H_3(\mathbb{Z}_2) \cong \mathbb{Z}_2$ [24, §4.2] and $H_3(\mathcal{G}(b)) = H_3(S_3) \cong \mathbb{Z}_6$ [36] between which two homomorphisms are given. Since the embedding $\mathcal{G}(u_0) : \mathbb{Z}_2 \rightarrow S_3$ is a coretraction, the homomorphism $H_3(\mathcal{G}(u_0)) : \mathbb{Z}_2 \rightarrow \mathbb{Z}_6$ is a monomorphism. Since the zero homomorphism $\mathcal{G}(u_1) : \mathbb{Z}_2 \rightarrow S_3$ is equal to the composition $\mathbb{Z}_2 \rightarrow 0 \rightarrow S_3$, then the homomorphism $H_3(\mathcal{G}(u_1))$ passes through the group $H_3(0) = 0$ and hence $H_3(\mathcal{G}(u_1))$ will be the zero homomorphism. Hence the colimit $\text{colim}^{\uparrow\uparrow} H_3(\mathcal{G})$ is isomorphic to the cokernel of the monomorphism $\mathbb{Z}_2 \rightarrow \mathbb{Z}_6$. Hence this colimit is isomorphic to the quotient group $\mathbb{Z}_6 / \mathbb{Z}_2 \cong \mathbb{Z}_3$.

We get the following answer: $\text{con } \mathcal{G} = 2$ and

$$\text{colim}_2^{\uparrow\uparrow} \mathcal{G} = \pi_3(\text{hocolim}^{\uparrow\uparrow} B\mathcal{G}) = \mathbb{Z}_3.$$

REMARK 5.8 For the category \mathcal{C} having an initial object, the formula (8) can be obtained from [14, Corollary 5.1].

EXAMPLE 5.9 Let's give an example to illustrate the formula (8). It shows that geometric intuition can be deceiving when constructing pointed homotopy colimits. Let $\mathcal{C} = \uparrow\uparrow$. Consider a diagram in the category sSet consisting of $X(a) = S^1$ is a simplicial circle (coequalizer of the pair $\Delta(\partial^0), \Delta(\partial^1) : \Delta[0] \rightrightarrows \Delta[1]$) [16, II.2.5.2], and let $X(b) = \Delta[0]$. Morphisms $X(u_0) = X(u_1) : X(a) \rightarrow X(b)$ are defined as the only morphisms into a terminal object.

The geometric realization of the homotopy colimit of nonpointed simplicial sets $\text{hocolim}^{\uparrow\uparrow} X$ consists of two cones over a circle whose vertices are identified. Therefore, this homotopy colimit is homotopy equivalent to a sphere

whose two points are identified. Its fundamental group is calculated using the Van Kampen theorem, as shown in [37]. It is equal to $\pi_1(\text{hocolim}^{\uparrow} X) = \mathbb{Z}$.

Now consider the same diagram with only base vertices in S^1 and $\Delta[0]$.

The construction of a homotopy colimit in sSet_* proceeds in steps [3, XII.2]. The first step is to build a wedge of values on objects. In our case, this wedge will be equal to (S^1, \star) (the space $X(b) = \{\star\}$ will be identified with the base point). Then it is necessary to glue the reduced cylinders of mappings $X(u_0), X(u_1) : X(a) \rightarrow X(b)$ to the resulting wedge (and in the general case the process continues further). In our case, these will be circles with a common border equal to the circle. The base points do not need to be connected, they are already identified. Therefore, the geometric realization of the pointed diagram is equal to the sphere. Whence the diagram of pointed simplicial sets satisfies $\pi_1(\text{hocolim}^{\uparrow} X) = 0$.

REMARK 5.10 *The homotopy colimit of nonpointed simplicial sets considered in Example 5.9 is weak equivalent to $S^1 \vee S^2$. To verify this, it suffices to replace the identified points by a segment.*

Farjoun proposed [13] (see also [19, Proposition 18.8.4]) a method for constructing the homotopy colimit of a diagram of pointed simplicial sets from non-pointed ones obtained by forgetting the property of points to be base. This homotopy colimit is equal to the quotient of the space of a nonpointed homotopy colimit with respect to the nerve of the category over which the diagram is given. Example 5.9 shows that this method is understandable.

In our case, the classifying space of the category is a circle, and Farjoun's method yields $\text{hocolim}^{\uparrow} X \approx S^2$.

6 Derived functors of the colimit and non-Abelian homology for group diagrams

Cotriple derived functors were introduced and studied by Barr and Beck [1]. They go back to the work of Godement [17, Appendix. Standard simplicial resolutions]. We adhere to the terminology of Swan's [32]. We prove that the values of the cotriple left derived of the colimit functor on a group diagram over a small category are isomorphic to the non-Abelian homology of the small category with coefficients in this diagram.

6.1 Left derived with respect to a cotriple

Let \mathcal{A} be an arbitrary category. The augmented simplicial object (A_*, ϵ, A) in it is the simplicial object A_* in \mathcal{A} together with the object $A \in \mathcal{A}$ and the morphism $\epsilon : A_0 \rightarrow A$, satisfying the relation $\epsilon d_0^1 = \epsilon d_1^1$.

Let (T, ϵ, μ) be a cotriple over \mathcal{A} . It defines an augmented simplicial functor

$$(T_*, \epsilon, Id_{\mathcal{A}}), \quad (13)$$

where T_* is a simplicial functor consisting of functors $T_n = T^{n+1}$, $n \geq 0$, and natural transformations of the boundary $d_i^n = T^i \epsilon T^{n-i} : T_n \rightarrow T_{n-1}$ (for $n \geq 1$) and degenerations $s_i^n = T^i \mu T^{n-i} : T_n \rightarrow T_{n+1}$ (for $n \geq 0$), for all $0 \leq i \leq n$. The augmentation $\epsilon : T_0 = T \rightarrow Id$ is equal to the counit of the cotriple.

Consider an arbitrary functor $G : \mathcal{A} \rightarrow \text{Grp}$ into the category of groups. Applying the functor G to an augmented simplicial functor (13) results in an augmented simplicial object

$$(GT_*, G\epsilon, G)$$

in the functor category $\mathcal{A} \rightarrow \text{Grp}$.

DEFINITION 6.1 *The left derived functors of the functor $G : \mathcal{A} \rightarrow \text{Grp}$ with respect to the cotriple T over \mathcal{A} are defined by the formula*

$$L_n G(A) := \pi_n(GT_*(A)), \forall A \in \mathcal{A}, \forall n \geq 0.$$

The elements of the group $L_0 G(A) = \pi_0(GT_*(A))$ are the connected components of the simplicial group $GT_*(A)$, and the mapping $G(\epsilon_A)$ is constant on each connected component. Hence we obtain the mapping $L_0 G(A) \rightarrow G(A)$ induced by the augmentation $G(\epsilon_A) : GT_0(A) \rightarrow G(A)$. We will consider the category $\mathcal{A} = \text{Grp}^{\mathcal{C}}$ for some arbitrary small category \mathcal{C} and the colimit functor $\text{colim}^{\mathcal{C}} : \text{Grp}^{\mathcal{C}} \rightarrow \text{Grp}$.

6.2 Cotriple for constructing the resolution

Denote by $O : \text{Ob } \mathcal{C} \xrightarrow{\subseteq} \mathcal{C}$ the embedding of the largest discrete subcategory in \mathcal{C} . The cotriple $T = \Lambda\Omega$ is constructed from a pair of adjoint functors

$$\Lambda : \text{Grp}^{\text{Ob } \mathcal{C}} \xleftrightarrow{\quad} \text{Grp}^{\mathcal{C}} : \Omega.$$

Here the functor $\Omega = O^*$ takes each diagram $F : \mathcal{C} \rightarrow \text{Grp}$ to the composition $F \circ O : \text{Ob}\mathcal{C} \rightarrow \text{Grp}$. Thus $\Omega(F)(c) = F(c)$ for all $c \in \text{Ob}\mathcal{C}$.

The functor $\Lambda = \text{Lan}^O$ is equal to the left Kan extension along $O : \text{Ob}\mathcal{C} \rightarrow \mathcal{C}$. It assigns to each $A \in \text{Grp}^{\text{Ob}\mathcal{C}}$ a diagram taking each $c \in \mathcal{C}$ to the group $(\Lambda A)(c) = \star_{c_0 \rightarrow c} A(c_0)$ equal to the free product of groups by the set of morphisms $c_0 \rightarrow c$ in the category \mathcal{C} . We get the cotriple $(T = \Lambda\Omega, \varepsilon, \Lambda\eta\Omega)$.

Let us describe the unit $\eta : \text{Id} \rightarrow \Omega\Lambda$ and the counit $\varepsilon : \Lambda\Omega \rightarrow \text{Id}$ of the adjunction $\Lambda \dashv \Omega$. Elements from $\star_{c_0 \rightarrow c} A(c_0)$, for $A \in \text{Grp}^{\text{Ob}\mathcal{C}}$, consist of pairs $(c_0 \xrightarrow{\alpha_0} c, x \text{ in } A(c_0))$. The natural transformation η is defined on $A \in \text{Grp}^{\text{Ob}\mathcal{C}}$ by the formula

$$(\eta A)_c(x \in A(c)) = (c \xrightarrow{1_c} c, x \in A(c)) \in \star_{c_0 \rightarrow c} A(c_0).$$

The natural transformation ε has the following property. For each $c \in \mathcal{C}$ the morphism $(\Lambda\Omega F)(c) = \star_{c_0 \rightarrow c} F(c_0) \xrightarrow{(\varepsilon F)_c} F(c)$ makes commutative triangles for all $c_0 \rightarrow c$:

$$\begin{array}{ccc} \star_{c_0 \rightarrow c} F(c_0) & \xrightarrow{\varepsilon F_c} & F(c) \\ & \swarrow \text{in}_{c_0 \rightarrow c} & \nearrow F(c_0 \rightarrow c) \\ & F(c_0)_{c_0 \rightarrow c} & \end{array}$$

Hence, the adjunction counit acts on the elements according to the formula

$$\varepsilon F_c(c_0 \rightarrow c, x \in F(c_0)) = (c, F(c_0 \rightarrow c)(x)).$$

6.3 Derived functors of colimit

DEFINITION 6.2 *The derived functors $L_n \text{colim}^{\mathcal{C}} : \text{Grp}^{\mathcal{C}} \rightarrow \text{Grp}$, $n \geq 0$, are the left derived of the colimit functor with respect to the cotriple $(T, \varepsilon, \mu) = (\Lambda\Omega, \varepsilon, \Lambda\eta\Omega)$.*

Let's construct a resolution. In general, it consists of functors $T_n = T^{n+1}$, $n \geq 0$, boundary natural transformations $d_i^n = T^i \varepsilon T^{n-i} : T^{n+1} \rightarrow T^n$, $0 \leq i \leq n$, and s_i^n degenerations $: T^i \mu T^{n-i} = T^i \Lambda\eta\Omega T^{n-i} : T^{n+1} \rightarrow T^{n+2}$.

Lemma 6.1 *Our resolution consists of functors whose values on $F \in \text{Grp}^{\mathcal{C}}$ and $c \in \text{Ob } \mathcal{C}$ are equal to the groups*

$$(T^{n+1}F)(c) = \star_{c_0 \rightarrow c} \star_{c_1 \rightarrow c_0} \cdots \star_{c_n \rightarrow c_{n-1}} F(c_n).$$

Each element from $T^{n+1}F(c)$ is described by a pair consisting of a tuple and an element of the group

$$(c \xleftarrow{\alpha_0} c_0 \xleftarrow{\alpha_1} c_1 \leftarrow \cdots \leftarrow c_{n-1} \xleftarrow{\alpha_n} c_n, x \in F(c_n))$$

To describe the boundary operators, we need the following assertion.

Lemma 6.2 *The natural homomorphisms $(\varepsilon TF)_c : T^2F(c) \rightarrow TF(c)$ map elements from $\star_{c_0 \rightarrow c} \star_{c_1 \rightarrow c_0} F(c_1)$ to $\star_{c_1 \rightarrow c} F(c_1)$ by formula*

$$(\varepsilon TF)_c(c_1 \xrightarrow{\alpha_1} c_0 \xrightarrow{\alpha_0} c, x \in F(c_1)) = (c_1 \xrightarrow{\alpha_0 \alpha_1} c, x \in F(c_1)).$$

Let us describe the boundary and degeneration operators involved in the resolution:

Proposition 6.3 *The boundary operators $(d_i^n F)_c : T_n F(c) \rightarrow T_{n-1} F(c)$, $0 \leq i \leq n$, act according to the formula*

$$(d_i^n F)_c(c_n \xrightarrow{\alpha_n} c_{n-1} \rightarrow \cdots \rightarrow c_0 \xrightarrow{\alpha_0} c, x \in F(c_n)) = \begin{cases} (c_n \xrightarrow{\alpha_n} c_{n-1} \rightarrow \cdots \rightarrow c_{i+1} \xrightarrow{\alpha_i \alpha_{i+1}} c_{i-1} \rightarrow \cdots \rightarrow c_0 \xrightarrow{\alpha_0} c, x \in F(c_n)), & i < n, \\ (c_{n-1} \xrightarrow{\alpha_{n-1}} c_{n-2} \rightarrow \cdots \rightarrow c_0 \xrightarrow{\alpha_0} c, F(\alpha_n)(x) \in F(c_{n-1})), & i = n. \end{cases}$$

PROOF. In the case of $i < n$, this formula follows from Lemma 6.2, by substituting $T^{n-i}F$ for F , applied to the natural transformation $\varepsilon T^{n-i}F : TT^{n-i}F \rightarrow T^{n-i}F$.

If $i = n$, then $d_i^n F = T^n \varepsilon F$. The homomorphism $(\varepsilon F)_c : TF(c) = \star_{c_0 \rightarrow c} F(c_0) \rightarrow F(c)$ acts like $(\varepsilon F)_c(c_0 \xrightarrow{\alpha_0} c, x \in F(c_0)) = F(\alpha_0)(x) \in F(c)$. Substituting the functor $T^n F$ instead of F , we get $T^n TF(c) \rightarrow T^n F(c)$, whence

$$(T^n \varepsilon F)_c(c \leftarrow c_0 \leftarrow \cdots \leftarrow c_{n-1} \xleftarrow{\alpha_n} c_n, x \in F(c_n)) = (c \leftarrow c_0 \leftarrow \cdots \leftarrow c_{n-1}, F(c_n \xrightarrow{\alpha_n} c_{n-1})(x)).$$

□

Recall that we introduced the n th non-Abelian homology groups $\text{colim}_n^{\mathcal{C}} F$ of the small category \mathcal{C} with coefficients in the group diagram F as homotopy groups of the simplicial replacement $\pi_n(\coprod F)$. Proposition 4.1 contains the definition of the simplicial group $C_*(\mathcal{C}, F)$ and states that this simplicial group is equal to $\coprod F$.

Theorem 6.4 *For an arbitrary small category \mathcal{C} and a group diagram $F : \mathcal{C} \rightarrow \text{Grp}$ the groups $L_n \text{colim}_n^{\mathcal{C}} F$ are isomorphic to the non-Abelian homology $\text{colim}_n^{\mathcal{C}} F$ with coefficients in F for all $n \geq 0$.*

PROOF. It suffices to prove that $\text{colim}^{\mathcal{C}} T_* F \cong C_*(\mathcal{C}, F)$ for all $F : \mathcal{C} \rightarrow \text{Grp}$. Since $\Lambda = \text{Lan}^O$, then for any group diagram $A \in \text{Grp}^{\mathcal{C}}$ there are the relations

$$\text{colim}^{\mathcal{C}} \Lambda A = \text{colim}^{Ob \mathcal{C}} A = \star_{c_0 \in Ob \mathcal{C}} A(c_0).$$

This implies for each $n \geq 0$ the validity of the equalities

$$\begin{aligned} \text{colim}^{\mathcal{C}} (\Lambda \Omega)^{n+1} F &= \star_{c_0 \in Ob \mathcal{C}} (\Lambda \Omega)^n F(c_0) = \\ & \star_{c_0 \in Ob \mathcal{C}} \star_{c_1 \rightarrow c_0} (\Lambda \Omega)^{n-1} F(c_1) = \star_{c_0 \in Ob \mathcal{C}} \star_{c_1 \rightarrow c_0} \star_{c_2 \rightarrow c_1} (\Lambda \Omega)^{n-2} F(c_2) = \dots \\ & \dots = \star_{c_n \rightarrow c_{n-1} \rightarrow \dots \rightarrow c_0} F(c_n) \end{aligned}$$

Changing the symbols in the tuples $c_n \rightarrow c_{n-1} \rightarrow \dots \rightarrow c_0$, c_i to c'_{n-i} for all $0 \leq i \leq n$, we get tuples $c'_0 \rightarrow c'_1 \rightarrow \dots \rightarrow c'_n$ and a simplicial group consisting of groups $\star_{c'_0 \rightarrow c'_1 \rightarrow \dots \rightarrow c'_n} F(c'_0)$ whose boundary operators are $d_i^n = d_{n-i}^n$, and the degeneration operators are equal to $s_i^n = s_{n-i}^n$. We obtain equalities of simplicial groups

$$\star_{c_n \rightarrow c_{n-1} \rightarrow \dots \rightarrow c_0} F(c_n) = \star_{c'_0 \rightarrow c'_1 \rightarrow \dots \rightarrow c'_n} F(c'_0) = \star_{c_0 \rightarrow c_1 \rightarrow \dots \rightarrow c_n} F(c_0)$$

According to the definitions, the derived of the colimit are equal to the homotopy groups of the simplicial group

$$\pi_n(\text{colim}^{\mathcal{C}} T_n F) = \pi_n(\text{colim}^{\mathcal{C}} (\Lambda \Omega)^{n+1} F),$$

and non-Abelian homology groups are equal to $\pi_n(C_*(\mathcal{C}, F))$. So they are isomorphic. □

Theorem 6.4 shows that the values of the derived of the functor $\text{colim}^{\mathcal{C}} : \text{Grp}^{\mathcal{C}} \rightarrow \text{Grp}$ on any diagram F are equal to the homotopy groups of the simplicial change $\pi_n(C_*(\mathcal{C}, F))$. Let us prove that each epimorphism in the category $\text{Grp}^{\mathcal{C}}$ corresponds to a long exact sequence in the category of groups. For this purpose, we prove the following assertion.

Lemma 6.5 *Let $f : F \rightarrow F''$ be an epimorphism in the category $\text{Grp}^{\mathcal{C}}$. Then the corresponding homomorphism $C_*(\mathcal{C}, f) : C_*(\mathcal{C}, F) \rightarrow C_*(\mathcal{C}, F'')$ in the category $\text{Grp}^{\Delta^{op}}$ is an epimorphism.*

PROOF. The simplicial replacement functor that maps the diagrams of their corresponding simplicial groups is equal to the functor composition $\text{Lan}^{\mathcal{Q}_{\mathcal{B}}^{op, \mathcal{C}}} \circ \partial^*$ (Definition 2.3). Each of these functors is left adjoint, and therefore takes an epimorphism to an epimorphism. \square

Let us introduce the notation $\text{colim}_n^{\mathcal{C}}(F \xrightarrow{f} F'') := \pi_n(\text{Ker}(C_*(\mathcal{C}, f)))$. This defines a functor on the category whose objects are epimorphisms in the category $\text{Grp}^{\mathcal{C}}$ and whose morphisms $f \rightarrow g$ are commutative squares

$$\begin{array}{ccc} F & \xrightarrow{f} & F'' \\ \downarrow & & \downarrow \\ G & \xrightarrow{g} & G'' \end{array}$$

Since the epimorphism of simplicial groups is a Kan fibration, Theorem 6.4 and Lemma 6.5 lead to the following assertion.

Corollary 6.6 *For every epimorphism $f : F \rightarrow F''$ in the category $\text{Grp}^{\mathcal{C}}$ there exists a long exact sequence (in Grp)*

$$\begin{aligned} \dots \rightarrow \text{colim}_{n+1}^{\mathcal{C}} F \rightarrow \text{colim}_{n+1}^{\mathcal{C}} F'' \rightarrow \text{colim}_n^{\mathcal{C}}(F \xrightarrow{f} F'') \\ \rightarrow \text{colim}_n^{\mathcal{C}} F \rightarrow \text{colim}_n^{\mathcal{C}} F'' \rightarrow \dots \rightarrow \text{colim}_0^{\mathcal{C}} F \rightarrow \text{colim}_0^{\mathcal{C}} F'' \rightarrow 0, \end{aligned}$$

where $\text{colim}_0^{\mathcal{C}} = \text{colim}^{\mathcal{C}}$.

REMARK 6.3 *In some cases, for homology and cohomology defined by cotriples, long exact sequences are associated with fibrations (see, for example, [11]). This is not necessary in our case, because these fibrations are the morphisms $f : F \rightarrow F''$ for which the homomorphisms f_c are retractions for all $c \in \text{Ob } \mathcal{C}$.*

7 Conclusion

For an arbitrary small category \mathcal{C} , we introduced non-Abelian homology of the category \mathcal{C} with coefficients in group diagrams and proved that they are isomorphic to derived functors of the functor $\text{colim}^{\mathcal{C}} : \text{Grp}^{\mathcal{C}} \rightarrow \text{Grp}$. We have proposed a formula relating non-Abelian homology to the homotopy groups of the homotopy colimit of classifying spaces. This formula provides a method for studying non-Abelian homology, and vice versa, there are applications of non-Abelian homology in homotopy theory. We hope that this will contribute to the development of methods for calculating homotopy groups and generalization of the classical homology theory of small categories to non-Abelian homology of group diagrams.

References

- [1] Barr M., Beck J. *Homology and standard constructions*. Seminar on triples and categorical homology theory (Lecture Notes in Mathematics, V. 80), Berlin-Heidelberg-New York: Springer-Verlag, 1969, P. 245-335.
- [2] Bousfield A.K., Friedlander E.M. *Homotopy theory of Γ -spaces, spectra, and bisimplicial sets*, (Lecture Notes in Mathematics, V.658), Berlin-Heidelberg-New York: Springer-Verlag, 1978, P.80-150.
- [3] Bousfield A. K., Kan D. M. *Homotopy Limits, Completions and Localizations*, (Lecture Notes in Mathematics V. 304), Berlin-Heidelberg-New York: Springer-Verlag, 1972.
- [4] Brown K. S. *Cohomology of groups*, New York-Heidelberg-Berlin: Springer-Verlag, 1982.
- [5] Brown R. *Some problems in non-abelian homotopical and homological algebra*. Homotopy theory and related topics. (Lecture Notes in Mathematics V. 1418), Berlin-Heidelberg-New York: Springer-Verlag, 1990, P.105-129.
- [6] Brown R., Higgins P. J., Sivera R. *Nonabelian Algebraic Topology*. Zürich: European Mathematical Society, 2011.
- [7] Brown R., Loday J.-L. *Van Kampen Theorems for the Diagrams of Spaces*. Topology. 1987. V. 26. P.311-315.

- [8] Castiglioni J. L., Ladra M. *Peiffer elements in simplicial groups and algebras*. Journal of Pure and Applied Algebra. 2008. V.212. P. 2115-2128.
- [9] Dwyer W.G., Kan D.M. *Simplicial localizations of categories*. Journal of Pure and Applied Algebra. 1980. V.17. P. 267-284.
- [10] Dwyer W.G., Kan D.M. *A classification theorem for diagrams of simplicial sets*, Topology, **23**:2 (1984), 139–155.
- [11] Ellis G.J. *Relative derived functors and the homology of groups*. Cahiers de topologie et géométrie différentielle catégoriques. 1990. T. 31, N.2 . P. 121-135
- [12] Ellis G., Mikhailov R. *A colimit of classifying spaces*. Advances in Mathematics. 2010. V. 223. P. 2097-2113.
- [13] Farjoun E. D. *Cellular inequalities*, The Čech Centennial (M. Cenk and H. Miller, eds.), Contemporary Mathematics, vol. 181, American Mathematical Society, 1995, pp. 159-181.
- [14] Farjoun E. D. *Fundamental group of homotopy colimits*. Advances in Mathematics. 2004. V.182. P. 1-27.
- [15] Fiedorowicz Z. *Classifying Spaces of Topological Monoids and Categories*. American Journal of Mathematics. 1984. V. 106, N. 2. P.301-350.
- [16] Gabriel P., Zisman M. *Calculus of fractions and homotopy theory*. Berlin-Heidelberg-New York: Springer-Verlag, 1967.
- [17] Godement R. *Topologie algébrique et théorie des faisceaux*. Paris: Hermann, 1958.
- [18] Goerss P.G., Jardine J.F. *Simplicial Homotopy Theory*, (Modern Birkhäuser Classics), Basel: Birkhäuser, 2009.
- [19] Hirschhorn P.S. *Model categories and their localizations*, Mathematical Surveys and Monographs, vol. 99, American Mathematical Society, Providence, RI, 2003.
- [20] Husainov A. A., Çalışıcı H. *Flows in graphs and the homology of free categories*. Algebra Discrete Math. 2003. Issue 2, P. 36-46.

- [21] Husainov A. A. *Homology and cohomology of cubical sets with coefficients in systems of objects*, Preprint, New York: Cornell University, 2022. <https://arxiv.org/abs/2207.07233>
- [22] Ivanov O.I., Mikhailov R., Wu J. *On Nontriviality of Certain Homotopy Groups of Spheres*. Homology Homotopy Appl. 2016. V. 18, N 2, P.337-344.
- [23] Kan D. M. *A Combinatorial Definition of Homotopy Groups*. Annals of Mathematics, Second Series. 1958. V. 67, N 2. P. 282-312.
- [24] Kuzmin Yu. V. *Homological group theory*. Moscow: “Factorial Press”, 2006 (Advanced Studies in Mathematics and Mechanics 1) (Russian).
- [25] Mac Lane S. *Categories for the Working Mathematician*. New York-Heidelberg: Springer-Verlag, 1998 (Graduate Texts in Mathematics 5).
- [26] Mikhailov R., Wu J. *On homotopy groups of the suspended classifying spaces*. Algebraic and Geometric Topology. 2010. V. 10. P. 565-625.
- [27] Moerdijk I., Toën B. *Simplicial presheaves as stacks*. Simplicial Methods for Operads and Algebraic Geometry. Basel: Springer, 2010. P. 127-141.
- [28] Quillen D. *Homotopical Algebra*. Berlin-Heidelberg: Springer-Verlag. (Lecture Notes in Mathematics V.43), 1967.
- [29] Quillen D.G. *Spectral sequences of a double semi-simplicial group*. Topology. 1966. V.5, N 2. P.155-157.
- [30] Quillen D. G. *Higher algebraic K-theory: I*. Bass H. (eds) Higher K-Theories. Berlin-Heidelberg: Springer-Verlag. (Lecture Notes in Mathematics, V. 341). 1973. P. 85-147.
- [31] Romero A., Rubio J. *Homotopy groups of suspended classifying spaces: an experimental approach*. Mathematics of computation. 2013. V.82, N 284. P. 2237-2244.
- [32] Swan R. G. *Some Relations between Higher K-Functors*. J. Algebra. 1972. V. 21. P. 113-136.
- [33] Thomason R. W. *Homotopy colimits in the category of small categories*. Math. Proc. Camb. Phil. Soc. 1979. V. 85, N. 1. 91-109.

- [34] Weibel C. A. *An Introduction to Homological Algebra*. Cambridge: Cambridge University Press, 1994.
- [35] Wu J. *Simplicial objects and homotopy groups*. Braids. Introductory Lectures on Braids, Configurations and their Applications. Hackensack: World Scientific. 2010, P. 31-181.
- [36] <https://mathoverflow.net/questions/180637/cohomology-of-symmetric-groups>
- [37] <https://math.stackexchange.com/questions/2296538/fundamental-group-of-the-sphere-with-two-points-identified>