

ON AN EXPONENTIAL POWER SUM

NEHA ELIZABETH THOMAS AND K VISHNU NAMBOOTHIRI

ABSTRACT. Using combinatorial techniques, we derive a recurrence identity that expresses an exponential power sum with negative powers in terms of another exponential power sum with positive powers. Consequently, we propose a formula for the power sum of first k natural numbers when the power is odd.

1. INTRODUCTION

A Dirichlet character χ modulo k (where k is a positive integer) is defined to be odd if $\chi(-1) = -1$ and even if $\chi(-1) = 1$. The Dirichlet L -function $L(s, \chi)$ is defined by the infinite series $\sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}$ where $s \in \mathbb{C}$ with $Re(s) > 1$.

The Gauss sum $G(z, \chi)$ for any complex number z is defined as

$$G(z, \chi) := \sum_{m=1}^k \chi(m) e^{\frac{2\pi i m z}{k}}.$$

In [2], E. Alkan established the following identity connecting $L(r, \chi)$ with Gauss sums and Bernoulli numbers.

$$(1) \quad \frac{(-1)^{v+1} k r!}{i^r 2^{r-1} \pi^r} L(r, \chi) = \sum_{q=0}^{2[\frac{r}{2}]} \binom{r}{q} B_q S(r-q, \chi) \quad [2, \text{Theorem 1}]$$

where

$$(2) \quad S(m, \chi) := \sum_{j=1}^k \left(\frac{j}{k}\right)^m G(j, \chi).$$

2020 Mathematics Subject Classification. 05A18, 11B37, 11B75, 11L03.

Key words and phrases. Exponential power sum; binomial theorem; composition; ordered partition; Dirichlet character; Bernoulli numbers.

and B_q are the q^{th} Bernoulli numbers. See [4, Chapter 12] for definition and other properties of Bernoulli numbers and Bernoulli polynomials.

E. Alkan derived formulae [1] for the sums $\sum_{\chi \text{ odd}} |L(1, \chi)|^2$ and $\sum_{\chi \text{ even}} |L(2, \chi)|^2$ using identity (1). His computations involved the simplification of sums of the form $\sum_{s=1}^{k-1} s^p e^{\frac{-2\pi i m s}{k}}$. He mainly tried to express such sums in terms of positive powers of the exponential function. It seems that the complexity of simplifying these exponential power sums prevented him from proceeding further establishing exact formulae for the value of $\sum_{\chi, r \text{ with same parity}} |L(r, \chi)|^2$ in general.

In this paper, we derive an identity that expresses the exponential power sum $\sum_{s=1}^{k-1} s^p e^{\frac{-2\pi i m s}{k}}$ in terms of another exponential power sum with positive powers. As a consequence, we derive a recursive formula for finding the power sum of first k natural numbers $h(p, k) := \sum_{s=1}^k s^p$ when the p is odd.

We note that Schaumberger and Faulhaber gave two different techniques for finding the power sums of natural numbers. In [8], Schaumberger used the exponential generating function

$$\sum_{p=0}^{\infty} h(p, k) \frac{x^k}{k!} = \sum_{r=1}^p e^{rx}$$

and L'Hôpital's rule on it for evaluating $h(p, k)$.

Faulhaber's formula [6, Identity 1] is a non-recursive formula for finding $h(p, k)$. It involves the Bernoulli numbers B_s and is given by

$$(3) \quad h(p, k) = \frac{1}{(p+1)} \left(k^{p+1} + \sum_{j=1}^p (-1)^j \binom{p+1}{j} B_j k^{p-j+1} \right).$$

For non integral power sums, some approximate results only seem to exist in the literature. See, for example theorem 1 in [7].

2. THE MAIN RESULT AND PROOF

An ordered partition or composition of a positive integer n is an m -tuple (j_1, \dots, j_m) with $j_1 + \dots + j_m = n$ where $1 \leq j_l \leq n$. j_l are called parts of the composition and m , the number of parts in it is

called the length of the composition. We use $C(n)$ to denote set of all compositions of n . For $1 \leq m \leq n$, $C(m, n)$ will denote the set of all compositions of n with length m . The number of elements in $C(m, n)$ is $\binom{n-1}{m-1}$ ([3, Theorem 4.1]). More properties and results related to compositions can be found, for example, in [3, Chapter 4].

In this paper, we mainly derive the following identity.

Proposition 2.1. *For k a positive integer, and $k \nmid m$, $p \geq 0$,*

$$(4) \quad \sum_{s=1}^{k-1} s^p e^{-\frac{2\pi i m s}{k}} = -k^p + \sum_{a=0}^{p-1} (-1)^{p-a} \binom{p}{a} k^a \sum_{s=1}^{k-1} s^{p-a} e^{\frac{2\pi i m s}{k}}.$$

We start the proof using binomial expansion.

$$\begin{aligned} \sum_{s=1}^{k-1} s^p e^{\frac{2\pi i m s}{k}} &= \sum_{s=1}^{k-1} (k-s)^p e^{\frac{2\pi i m (k-s)}{k}} \\ &= \sum_{s=1}^{k-1} (k-s)^p e^{-\frac{2\pi i m s}{k}} \\ &= \sum_{s=1}^{k-1} (-1)^p s^p e^{-\frac{2\pi i m s}{k}} + \sum_{a=0}^{p-1} (-1)^a \binom{p}{a} k^{p-a} \sum_{s=1}^{k-1} s^a e^{-\frac{2\pi i m s}{k}}. \end{aligned}$$

Hence

$$(5) \quad \sum_{s=1}^{k-1} s^p e^{-\frac{2\pi i m s}{k}} = (-1)^p \sum_{s=1}^{k-1} s^p e^{\frac{2\pi i m s}{k}} + \sum_{a=0}^{p-1} (-1)^{p+a+1} \binom{p}{a} k^{p-a} \sum_{s=1}^{k-1} s^a e^{-\frac{2\pi i m s}{k}}.$$

For notational convenience, we write $f(p) = \sum_{s=1}^{k-1} s^p e^{-\frac{2\pi i m s}{k}}$ and $g(p) = \sum_{s=1}^{k-1} s^p e^{\frac{2\pi i m s}{k}}$. With this notation, we rewrite identity (5) and expand it

recursively as

$$\begin{aligned}
f(p) &= (-1)^p g(p) + \sum_{a=0}^{p-1} (-1)^{p+a+1} k^{p-a} \binom{p}{a} f(a) \\
&= (-1)^p g(p) + (-1)^{p+1} \binom{p}{p-1} k g(p-1) \\
&\quad + \left[(-1)^{p+1} \binom{p}{p-2} + (-1)^{p+2} \binom{p}{p-1} \binom{p-1}{p-2} \right] k^2 g(p-2) \\
&\quad + \left[(-1)^{p+1} \binom{p}{p-3} + (-1)^{p+2} \binom{p}{p-2} \binom{p-2}{p-3} \right. \\
&\quad \left. + (-1)^{p+2} \binom{p}{p-1} \binom{p-1}{p-3} + (-1)^{p+3} \binom{p}{p-1} \binom{p-1}{p-2} \binom{p-2}{p-3} \right] \\
&\quad \times k^3 g(p-3) + \dots .
\end{aligned}$$

It can be seen by induction that for $a \geq 2$, $g(p-a)$ has coefficient as the sum

$$(6) \quad \sum_{\substack{i_1, i_2, \dots, i_r \in \{p-a+1, \dots, p-1\} \\ i_1 > i_2 > \dots > i_r}} (-1)^{p+r+1} \binom{p}{i_1} \binom{i_1}{i_2} \dots \binom{i_{r-1}}{i_r} \binom{i_r}{p-a} k^a.$$

with the convention that $\{i_1, \dots, i_r\}$ can also be the empty set to allow the binomial coefficient $\binom{p}{p-a}$.

Thus the coefficient of $k^p g(0)$ is

$$(7) \quad \sum_{\substack{i_1, i_2, \dots, i_r \in \{1, \dots, p-1\} \\ i_1 > i_2 > \dots > i_r}} (-1)^{p+r+1} \binom{p}{i_1} \binom{i_1}{i_2} \dots \binom{i_{r-1}}{i_r} \binom{i_r}{0}.$$

Now

$$\begin{aligned}
&\binom{p}{i_1} \binom{i_1}{i_2} \dots \binom{i_{r-1}}{i_r} \binom{i_r}{0} \\
&= \frac{p!}{(p-i_1)! i_1!} \frac{i_1!}{(i_1-i_2)! i_2!} \dots \frac{i_{r-1}!}{(i_{r-1}-i_r)! i_r!} \frac{i_r!}{(i_r-0)! 0!} \\
&= \frac{p!}{(p-i_1)! (i_1-i_2)! \dots (i_r-0)! 0!}.
\end{aligned}$$

Proposition 2.2. *Let $r \geq 1$. The set $\{(i_1, \dots, i_r) : i_1, i_2, \dots, i_r \in \{1, \dots, p-1\}, i_1 > i_2 > \dots > i_r\}$ is in one to one correspondence with the set $C(r+1, p)$*

Proof. Write $p - i_1 = j_1$, $i_1 - i_2 = j_2$, \dots , $i_{r-1} - i_r = j_r$, $i_r - 0 = j_{r+1}$. Since $j_1 + j_2 + \dots + j_{r+1} = p$ and j_l are all positive, $(j_1, \dots, j_{r+1}) \in C(r+1, p)$. Let (i_1, i_2, \dots, i_r) and $(i'_1, i'_2, \dots, i'_r)$ be distinct. Let l be the largest index such that $i_l \neq i'_l$ (with the convention that $i_{l+1} = 0$ if $l = r$). Thus $j_{l+1} = i_l - i_{l+1}$ and $j'_{l+1} = i'_l - i'_{l+1}$ will be distinct. Hence $(j_1, j_2, \dots, j_{r+1})$ and $(j'_1, j'_2, \dots, j'_{r+1})$ will be distinct. On the other hand, if $(j_1, j_2, \dots, j_{r+1}) \in C(r+1, p)$, let $i_1 = p - j_1$, $i_2 = p - (j_1 + j_2)$, \dots so that $i_1, \dots, i_r \in \{1, \dots, p-1\}$ and they satisfy $i_1 > \dots > i_r$, with $r = 0$ giving the empty set. This shows that this correspondence is onto as well. \square

Since $C(r+1, p)$ has $\binom{p-1}{r}$ elements, it follows that

Corollary 2.3. *The set defined in the above proposition has $\binom{p-1}{r}$ elements.*

The binomial product sum (7) consists of the summand $(-1)^{p+1} \binom{p}{0}$. This term corresponds to the empty subset of $\{1, \dots, p-1\}$. Corresponding to this empty set we may take the single available element (p) from $C(1, p)$. Hence, $\binom{p}{i_1} \binom{i_1}{i_2} \dots \binom{i_{r-1}}{i_r} \binom{i_r}{0}$ and $\binom{p}{0}$ can be replaced with the multinomial coefficient $\binom{p}{j_1, j_2, \dots, j_{r+1}}$ and so the coefficient of $k^p g(0)$ can be rewritten as

$$\sum_{r=0}^{p-1} \sum_{(j_1, j_2, \dots, j_{r+1}) \in C(r+1, p)} (-1)^{p+r+1} \binom{p}{j_1, j_2, \dots, j_{r+1}}.$$

Now we use the following lemma to proceed.

Lemma 2.4. [5, Lemma 1] *If u_1, u_2, u_3, \dots is any sequence of numbers, then the coefficient of x^n in $\left(1 - \sum_{i=1}^{\infty} u_i x^i\right)^{-1}$ is $\sum_{(a_1, \dots, a_k) \in C(n)} u_{a_1} u_{a_2} \dots u_{a_k}$.*

Continuing with our computations,

$$\begin{aligned}
& \sum_{r=0}^{p-1} \sum_{(j_1, j_2, \dots, j_{r+1}) \in C(r+1, p)} (-1)^{p+r+1} \binom{p}{j_1, j_2, \dots, j_{r+1}} \\
&= \sum_{r=1}^p \sum_{(j_1, j_2, \dots, j_r) \in C(r, p)} (-1)^{p+r} \binom{p}{j_1, j_2, \dots, j_r} \\
&= (-1)^p p! \sum_{r=1}^p \sum_{(j_1, j_2, \dots, j_r) \in C(r, p)} (-1)^r \frac{1}{j_1! j_2! \dots j_r!} \\
&= (-1)^p p! \times \text{coefficient of } x^p \text{ in } \left(1 + \sum_{i=1}^{\infty} \frac{x^i}{i!} \right)^{-1} \\
&\quad \left(\text{obtained by taking } u_i = \frac{-1}{i!} \text{ in lemma 2.4} \right) \\
&= (-1)^p p! \times \text{coefficient of } x^p \text{ in } (e^x)^{-1} \\
&= 1.
\end{aligned}$$

Now we compute the coefficient of $k^a g(p-a)$. We take the product of binomial coefficients appearing in (6) and simplify it further.

$$\begin{aligned}
& \binom{p}{i_1} \binom{i_1}{i_2} \dots \binom{i_{r-1}}{i_r} \binom{i_r}{p-a} \\
&= \binom{p}{p-i_1} \binom{i_1}{i_1-i_2} \dots \binom{i_{r-1}}{i_{r-1}-i_r} \binom{i_r}{i_r-p+a} \\
&= \frac{p!}{(p-i_1)!(i_1-i_2)! \dots (i_r-p+a)!(p-a)!} \\
&= \frac{a!}{(p-i_1)!(i_1-i_2)! \dots (i_r-p+a)!} \binom{p}{a}.
\end{aligned}$$

Write $p-i_1 = j_1$, $i_1-i_2 = j_2$, \dots , $i_{r-1}-i_r = j_r$, $i_r-(p-a) = j_{r+1}$. So $j_1 + j_2 + \dots + j_{r+1} = a$. As in the above case, we can see that $(j_1, j_2, \dots, j_{r+1})$ is in a one to one correspondence with (i_1, i_2, \dots, i_r) where $i_1 > \dots > i_r$. So we have

$$\binom{p}{i_1} \binom{i_1}{i_2} \dots \binom{i_{r-1}}{i_r} \binom{i_r}{p-a} = \binom{p}{a} \binom{a}{j_1, j_2, \dots, j_{r+1}}.$$

Hence we get the coefficient of $k^a g(p-a)$ as

$$\begin{aligned}
& (-1)^{p-a} \sum_{r=0}^{p-1} \sum_{(j_1, j_2, \dots, j_{r+1}) \in C(a, r+1)} (-1)^{a+r+1} \binom{p}{a} \binom{a}{j_1, j_2, \dots, j_{r+1}} \\
&= (-1)^{p-a} \binom{p}{a} \sum_{r=0}^{p-1} \sum_{(j_1, j_2, \dots, j_{r+1}) \in C(a, r+1)} (-1)^{a+r+1} \binom{a}{j_1, j_2, \dots, j_{r+1}} \\
&= (-1)^{p-a} \binom{p}{a}.
\end{aligned}$$

Thus

$$\begin{aligned}
f(p) &= (-1)^p g(p) + (-1)^{p-1} \binom{p}{p-1} k g(p-1) + \sum_{a=2}^p (-1)^{p-a} \binom{p}{a} k^a g(p-a) \\
&= \sum_{a=0}^{p-1} (-1)^{p-a} \binom{p}{a} k^a g(p-a) + k^p g(0) \\
&= -k^p + \sum_{a=0}^{p-1} (-1)^{p-a} \binom{p}{a} k^a g(p-a) \left(\text{since } g(0) = \sum_{s=1}^{k-1} e^{\frac{2\pi i m s}{k}} = -1 \text{ when } e^{\frac{2\pi i m s}{k}} \neq 1 \right)
\end{aligned}$$

which is what we claimed.

Let us see an application of this identity. When $k|m$, $e^{\frac{2\pi i m s}{k}} = 1$ and $g(0) = k-1$ in the above so that we get

$$(8) \quad \sum_{s=1}^{k-1} s^p = k^p(k-1) + \sum_{a=0}^{p-1} (-1)^{p-a} \binom{p}{a} k^a \sum_{s=1}^{k-1} s^{p-a}.$$

Let $h(p, k) = \sum_{s=1}^k s^p$. If p is odd, from the above, we get a recursive formula for the sum of p th powers of first k natural numbers as

$$h(p, k) = \frac{1}{2} \left((k+1)^p k + \sum_{a=1}^{p-1} (-1)^{p-a} \binom{p}{a} (k+1)^a h(p-a, k) \right).$$

On replacing $p-a$ with j , we get

Corollary 2.5.

$$(9) \quad h(p, k) = \frac{1}{2} \left((k+1)^p k + \sum_{j=1}^{p-1} (-1)^j \binom{p}{j} (k+1)^{p-j} h(j, k) \right).$$

Such a formula cannot be derived when p is even because of the cancellation of $h(p, k)$ taking place from both sides of identity (8).

To illustrate the use of the above identity, let us compute $\sum_{s=1}^k s^3$. Note that the knowledge of $h(1, k) = \sum_{s=1}^k s$ and $h(2, k) = \sum_{s=1}^k s^2$ is necessary for this computation. We have $h(1, k) = \frac{k(k+1)}{2}$ and $h(2, k) = \frac{k(k+1)(2k+1)}{6}$. Thus

$$\begin{aligned} h(3, k) &= \frac{1}{2} \left((k+1)^3 k + \sum_{a=1}^2 (-1)^{3-a} \binom{3}{a} (k+1)^a h(3-a, k) \right) \\ &= \frac{1}{2} \left((k+1)^3 k + (-1)^2 \binom{3}{1} (k+1) h(2, k) + (-1)^1 \binom{3}{2} (k+1)^2 h(1, k) \right) \\ &= \frac{1}{2} \left((k+1)^3 k + 3(k+1) \frac{k(k+1)(2k+1)}{6} - 3(k+1)^2 \frac{k(k+1)}{2} \right) \\ &= \frac{1}{4} (k(k+1)^2(2k+1) - (k+1)^3 k) \\ &= \left[\frac{k(k+1)}{2} \right]^2. \end{aligned}$$

3. ACKNOWLEDGEMENTS

The first author thanks the Kerala State Council for Science, Technology and Environment, Thiruvananthapuram, Kerala, India for providing financial support for carrying out research work.

REFERENCES

- [1] Emre Alkan. On the mean square average of special values of L -functions. *Journal of Number Theory*, 131(8):1470–1485, 2011.
- [2] Emre Alkan. Values of Dirichlet L -functions, Gauss sums and trigonometric sums. *The Ramanujan Journal*, 26(3):375–398, 2011.
- [3] George E Andrews. *The theory of partitions*. Number 2. Cambridge university press, 1998.
- [4] Tom Apostol. *Introduction to analytic number theory*. Springer, 1976.
- [5] Ira M Gessel and Ji Li. Compositions and Fibonacci identities. *Journal of Integer Sequences*, 16(2):3, 2013.
- [6] Kevin J McGown and Harold R Parks. The generalization of Faulhaber’s formula to sums of non-integral powers. *Journal of mathematical analysis and applications*, 330(1):571–575, 2007.

- [7] Harold R Parks. Sums of non-integral powers. *Journal of mathematical analysis and applications*, 297(1):343–349, 2004.
- [8] N Shaumberger. Using L -hospital's rule to sum a series. *PME Journal*, 6(5):281–282, 1976.

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE, THIRUVANANTHAPURAM (RESEARCH CENTRE, UNIVERSITY OF KERALA), KERALA - 695034, INDIA
Email address: nehathomas2009@gmail.com

DEPARTMENT OF MATHEMATICS, GOVERNMENT COLLEGE CHITTUR, KERALA - 678104, INDIA, DEPARTMENT OF COLLEGIATE EDUCATION, GOVERNMENT OF KERALA, INDIA
Email address: kvnamboothiri@gmail.com