

Weak pattern convergence for SLOPE and its robust versions

Ivan Hejný¹, Jonas Wallin¹, and Małgorzata Bogdan^{1,2}

¹Department of Statistics, Lund University

²Institute of Mathematics, University of Wrocław

Abstract

The Sorted L-One Estimator (SLOPE) is a popular regularization method in regression, which induces clustering of the estimated coefficients. That is, the estimator can have coefficients of identical magnitude. In this paper, we derive an asymptotic distribution of SLOPE for the ordinary least squares, Huber, and Quantile loss functions, and use it to study the clustering behavior in the limit. This requires a stronger type of convergence since clustering properties do not follow merely from the classical weak convergence. For this aim, we utilize the Hausdorff distance, which provides a suitable notion of convergence for the penalty subdifferentials and a bridge toward weak convergence of the clustering pattern. We establish asymptotic control of the false discovery rate for the asymptotic orthogonal design of the regressor. We also show how to extend the framework to a broader class of regularizers other than SLOPE.

1 Introduction

Consider the linear model $y = X\beta^0 + \varepsilon$, where $X \in \mathbb{R}^{n \times p}$ is the design matrix, $\beta^0 \in \mathbb{R}^p$ is the vector of regression coefficients and $\varepsilon \in \mathbb{R}^n$ is the random noise vector with independent identically distributed entries $\varepsilon_1, \dots, \varepsilon_n$.

In case when $n > p$ and the columns of X are linearly independent, the vector of regression coefficients β^0 can be estimated using the classical Ordinary Least Squares (OLS) approach. According to the Gauss-Markov Theorem, OLS estimator allows to obtain the minimal mean-squared error among all linear unbiased estimators. However, it is well known that this mean-squared error can be further improved by considering regularized biased estimators, like ridge regression [19] or LASSO [45], which usually have a substantially smaller variance than OLS. This gain in estimation accuracy can be very large when the columns of X are strongly correlated and/or when p approaches n . Moreover, regularized estimators can be applied also when $p > n$.

Ridge regression and LASSO are designed for different data-generating regimes. While ridge regression performs the best when the vector β^0 contains many nonzero elements (i.e. it is dense), LASSO can be much better when β^0 is sparse. This gain results from the non-linear LASSO shrinkage, which allows to obtain sparse estimators of β^0 . The focus of this

article, Sorted L-One Estimator (SLOPE, [8]), is a relatively new extension of LASSO, which allows for further dimensionality reduction and improvement of estimation properties.

Given a vector of penalties $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_p)$ with $\lambda_1 \geq \dots \geq \lambda_p \geq 0$, the SLOPE estimator of β^0 is given by the minimization problem

$$\hat{\boldsymbol{\beta}} = \underset{\beta \in \mathbb{R}^p}{\operatorname{argmin}} \frac{1}{2} \|y - X\beta\|_2^2 + J_{\boldsymbol{\lambda}}(\beta), \quad (1)$$

where $J_{\boldsymbol{\lambda}}(\beta)$ is the SLOPE norm defined as

$$J_{\boldsymbol{\lambda}}(\beta) = \sum_{i=1}^p \lambda_i |\beta|_{(i)},$$

and $|\beta|_{(1)} \geq |\beta|_{(2)} \geq \dots \geq |\beta|_{(p)}$ are the absolute values of the coordinates of β ordered in decreasing order of their magnitude. (see Bogdan et al. [8]).

SLOPE has some very appealing statistical properties. In the seminal works [42, 6], it is proved that SLOPE with a specific predefined sequence of tuning parameters λ adapts to the unknown sparsity of the vector of regression coefficients β^0 and obtains exact minimax estimation and prediction rates. Similar sequences of the tuning parameters yield the False Discovery Rate (FDR) control when the design matrix is orthogonal [9, 8], or the asymptotic FDR control when the explanatory variables are independent [24, 25]. The minimaxity results have been proved also for SLOPE extensions to the binary and multi-class classification [2, 3, 1], identification of group of predictors [11] or for the simultaneous outlier detection and robust regression [46] using the mean-shift model of [17, 32, 39]. SLOPE was also investigated with the asymptotic theory of Approximate Message Passing Algorithms [12, 37, 13]. While in [37] it is shown that in terms of estimation properties SLOPE does not systematically outperform LASSO or ridge regression, the results in [13] illustrate that in terms of feature selection SLOPE has qualitatively different properties from LASSO. Specifically, SLOPE can achieve the full power of signal recovery for the moderately sparse vectors of regression coefficients, when the power of LASSO is restricted. Also, in [13] it is shown that for any problem instance, there exists a choice of the tuning parameters such that SLOPE outperforms LASSO, both with respect to model selection and estimation properties.

These superior model selection properties of SLOPE result from its ability to cluster similar values of estimated regression coefficients. Due to this property, SLOPE can efficiently exploit the situation when some parameters are equal to each other. This allows to reduce the number of estimated parameters and enhances the speed and accuracy of model fitting. A prominent illustration of advantages resulting from such a model reduction is provided by the Convolutional Neural Networks, where the “parameter sharing” has allowed to “dramatically lower the number of unique model parameters and to significantly increase network sizes without requiring a corresponding increase in training data” [18]. In the classical statistics setup identification of the parameter sharing patterns is very beneficial, e.g., in the context of clustering similar levels of multi-valued categorical predictors [31, 41, 33]. Also, clustering properties of SLOPE were crucial for attaining good performance in portfolio optimization and tracking of financial indices [27, 26].

The ability of identification of the parameters tying patterns by SLOPE has been discussed in several articles [10, 15, 26, 38, 40]. Specifically, in [38] the SLOPE clustering

properties are related to the properties of the subdifferential of the SLOPE norm and the SLOPE pattern of the vector $u \in \mathbb{R}^p$ is defined by providing the characteristics which are shared by all vectors in \mathbb{R}^p for which this subdifferential attains the same value as for u . These characteristics can be concisely described in the following definition:

Definition 1.1. A SLOPE pattern is the integer valued function $\mathbf{patt} : \mathbb{R}^p \rightarrow \mathbb{Z}^p$ given by

$$\mathbf{patt}(u)_i := \text{rank}(|u_i|) \text{sgn}(u_i),$$

where $\text{rank}(|u_i|)$ is the rank of the distinct values of $|u|$; (see [40, 7]).

Thus, the SLOPE pattern of the vector u contains the information on the sign of its elements and the ranking of their absolute values, including the information about clusters of coordinates with the same absolute values.

In a recent article [7] a sufficient and necessary condition for the pattern recovery by SLOPE is provided. This condition specifies the patterns which can be recovered by SLOPE for a given design matrix X and when the magnitude of regression coefficients is large enough. The result was further applied to characterize the asymptotic probability of the pattern recovery in the classical asymptotic setup, when the number of features p is fixed and the number of examples n tends to infinity.

In the present paper we substantially expand knowledge on the asymptotic properties of SLOPE and its ability for the pattern recovery and FDR control. Firstly, we extend the asymptotic results of Knight and Fu [16] on Lasso type estimators and derive the limiting distribution of $\sqrt{n}(\hat{\beta}_n - \beta^0)$, as a solution to a regularized random least squares problem. We also observe that, crucially, for a general regularizer, weak convergence of $\sqrt{n}(\hat{\beta}_n - \beta^0)$ does not guarantee weak convergence of $\mathbf{patt}(\sqrt{n}(\hat{\beta}_n - \beta^0))$, as demonstrated in Example A.1 in the Appendix. However, we establish that for the SLOPE estimator this is in fact so. This is achieved by utilizing the Hausdorff distance [36], which provides a suitable notion of convergence for subdifferentials and a bridge toward weak pattern convergence.

Further, building on the ideas of Pollard [34], we investigate the limiting distribution of SLOPE, where we replace the L_2 norm in (1) with a more general loss $\sum_{i=1}^n g(y_i, X_i, \beta)$. We formulate general conditions on g under which we can derive the limiting distribution and establish that these conditions are satisfied for two important loss functions, namely the (robust) Huber loss [20] and the quantile loss [23]. With this result, we are also able to asymptotically control the FDR for the two aforementioned losses under independent regressors by fitting the correct nuisance parameters.

1.1 Overview

In section 2., we give formulas for the limiting distribution of $\hat{\beta}_n$, when $\lambda^n \rightarrow \lambda \geq 0$ (Theorem 2.1) and for the limiting distribution of $\sqrt{n}(\hat{\beta}_n - \beta^0)$, when $\lambda^n/\sqrt{n} \rightarrow \lambda \geq 0$ (Theorem 2.2). In section 3., we study the interplay between the subdifferential and pattern of SLOPE and utilize the Hausdorff distance in Proposition 3.7 on the convergence of SLOPE penalty subdifferentials. Section 4. forms the backbone of the paper. In Theorem 4.1, we prove weak pattern convergence of $\sqrt{n}(\hat{\beta}_n - \beta^0)$ and use this in Corollary 4.3 to provide an alternative proof for the probability of true pattern recovery (Theorem 4.1.i)[7]). Weak pattern convergence leads directly to the asymptotic FDR control under independence in Example 4.7.

We end the section with a discussion 4.4, on how the weak pattern convergence extends to a broader family of regularizers other than SLOPE. In section 5., we prove Theorem 5.4, the main result of the paper, which extends the results from Section 4 to other loss functions, including the Huber loss and the Quantile loss.

2 Asymptotic distribution of SLOPE for the standard loss

Consider the linear model $y = X\beta^0 + \varepsilon$, with $X = (X_1, \dots, X_n)^T$, where X_1, X_2, \dots are i.i.d. centered random vectors in \mathbb{R}^p with covariance matrix C . Further assume $\varepsilon_1, \varepsilon_2, \dots$ are i.i.d. centered random variables with variance σ^2 and $X \perp \varepsilon$. In this section, we want to give exact formulas for the low-dimensional asymptotics with fixed p and $n \rightarrow \infty$, of the SLOPE estimator for the quadratic loss:

$$\hat{\beta}_n = \operatorname{argmin}_{\beta \in \mathbb{R}^p} \frac{1}{2} \|y - X\beta\|_2^2 + J_{\lambda^n}(\beta) \quad (2)$$

$$= \operatorname{argmin}_{\beta \in \mathbb{R}^p} \frac{1}{2} (\beta - \beta^0)^T X^T X (\beta - \beta^0) + (\beta - \beta^0)^T X^T \varepsilon + \|\varepsilon\|_2^2 / 2 + J_{\lambda^n}(\beta) \quad (3)$$

Note that by the LLN and CLT:

$$C_n := \frac{1}{n} X^T X \xrightarrow{a.s.} C \quad \text{and} \quad W_n := \frac{1}{\sqrt{n}} X^T \varepsilon \xrightarrow{d} W \sim \mathcal{N}(0, \sigma^2 C), \quad (4)$$

and $\|\varepsilon\|_2^2/n \xrightarrow{a.s.} \sigma^2$. The following statements and proofs are direct extensions of the results in [16]. For completeness we include the proofs.

Theorem 2.1. *If $\lambda^n/n \rightarrow \lambda \geq 0$ and $C \succ 0$, then $\hat{\beta}_n \xrightarrow{p} \operatorname{argmin}_{\beta} Z(\beta)$, where*

$$Z(\beta) = \frac{1}{2} (\beta - \beta^0)^T C (\beta - \beta^0) + J_{\lambda}(\beta)$$

In particular, in the case $\lambda = 0$, we have consistency $\hat{\beta}_n \xrightarrow{p} \beta^0$.

Proof. Consider

$$Z_n(\beta) := \frac{1}{2n} \|y - X\beta\|_2^2 + \frac{1}{n} J_{\lambda^n}(\beta)$$

Note that $\hat{\beta}_n = \operatorname{argmin} Z_n(\beta)$. From (3) and (4), we get that for every $\beta \in \mathbb{R}^p$:

$$Z_n(\beta) \xrightarrow{p} Z(\beta) + \sigma^2/2.$$

The pointwise convergence together with convexity of Z_n 's implies uniform convergence on compact subsets; i.e., $\sup_{\beta \in K} |Z_n(\beta) - Z(\beta) - \sigma^2/2| \xrightarrow{p} 0$ for every compact K . Moreover, positive definiteness of C guarantees that the minimizer of $Z(\beta)$ is unique. It follows that $\hat{\beta}_n = \operatorname{argmin} Z_n(\beta) \xrightarrow{p} \operatorname{argmin} Z(\beta)$. We refer the reader to [4] and [35] for the general convergence results. \square

Let $J'_{\lambda}(\beta^0; u)$ denote the directional derivative of the SLOPE norm at β^0 in direction u .

Theorem 2.2. *If $\lambda^n/\sqrt{n} \rightarrow \lambda \geq 0$ and $C \succ 0$, then $\hat{\mathbf{u}}_n := \sqrt{n}(\hat{\boldsymbol{\beta}}_n - \beta^0) \xrightarrow{d} \hat{\mathbf{u}}$, where*

$$\begin{aligned} \hat{\mathbf{u}} &:= \operatorname{argmin}_u V(u), \\ V(u) &= \frac{1}{2}u^T C u - u^T W + J'_\lambda(\beta^0; u), \end{aligned} \quad (5)$$

with $W \sim \mathcal{N}(0, \sigma^2 C)$. Moreover, the directional derivative is given by

$$J'_\lambda(\beta^0; u) = \sum_{i=1}^p \lambda_{\pi(i)} [u_i \operatorname{sgn}(\beta_i^0) \mathbb{I}[\beta_i^0 \neq 0] + |u_i| \mathbb{I}[\beta_i^0 = 0]], \quad (6)$$

where π is a permutation which sorts the vector $|(\beta^0 + u/\sqrt{n})|$ for all sufficiently large n ; i.e. $\exists M > 0$ s.t. $\forall n \geq M$ $|(\beta^0 + u/\sqrt{n})|_{\pi^{-1}(1)} \geq \dots \geq |(\beta^0 + u/\sqrt{n})|_{\pi^{-1}(p)}$.

Proof. Consider

$$V_n(u) := \frac{1}{2n}u^T X^T X u - u^T \frac{1}{\sqrt{n}}X^T \varepsilon + J_{\lambda^n}(\beta^0 + u/\sqrt{n}) - J_{\lambda^n}(\beta^0). \quad (7)$$

By substituting $u = \sqrt{n}(\beta - \beta^0)$ it follows that $\operatorname{argmin} V_n(u) = \sqrt{n}(\hat{\boldsymbol{\beta}}_n - \beta^0)$. The stochastic part of $V_n(u)$ converges by (4) and Slutsky:

$$\frac{1}{2n}u^T X^T X u - u^T \frac{1}{\sqrt{n}}X^T \varepsilon \xrightarrow{d} \frac{1}{2}u^T C u - u^T W. \quad (8)$$

Moreover, for a fixed $u \in \mathbb{R}^p$ there exists a permutation π , which sorts $|\beta^0 + u/\sqrt{n}|$ for all sufficiently large n , i.e. $\exists M > 0 : \forall n \geq M : |(\beta^0 + u/\sqrt{n})|_{\pi^{-1}(1)} \geq \dots \geq |(\beta^0 + u/\sqrt{n})|_{\pi^{-1}(p)}$. At the same time we have $|\beta^0|_{\pi^{-1}(1)} \geq \dots \geq |\beta^0|_{\pi^{-1}(p)}$. Consequently

$$\begin{aligned} J_{\lambda^n}(\beta^0 + u/\sqrt{n}) - J_{\lambda^n}(\beta^0) &= \sum_{j=1}^p \lambda_j^n [|\beta^0 + u/\sqrt{n}|_{\pi^{-1}(j)} - |\beta^0|_{\pi^{-1}(j)}] \\ &= \sum_{i=1}^p \lambda_{\pi(i)}^n [|\beta_i^0 + u_i/\sqrt{n}| - |\beta_i^0|] \\ &= \sum_{i=1}^p \lambda_{\pi(i)}^n [(u_i/\sqrt{n}) \operatorname{sgn}(\beta_i^0) \mathbb{I}[\beta_i^0 \neq 0] + (|u_i|/\sqrt{n}) \mathbb{I}[\beta_i^0 = 0]] \\ &\longrightarrow \sum_{i=1}^p \lambda_{\pi(i)} [u_i \operatorname{sgn}(\beta_i^0) \mathbb{I}[\beta_i^0 \neq 0] + |u_i| \mathbb{I}[\beta_i^0 = 0]], \end{aligned} \quad (9)$$

which shows that $V_n(u) \xrightarrow{d} V(u)$. Again, by convexity and uniqueness of the minimizer of $V(u)$, we obtain $\sqrt{n}(\hat{\boldsymbol{\beta}}_n - \beta^0) = \operatorname{argmin} V_n(u) \xrightarrow{d} \operatorname{argmin} V(u)$. \square

Remark 2.3. Observe that the sum appearing in (5), is exactly the directional derivative of J_λ at the point β^0 in direction u , hence the formulation $J'_\lambda(\beta^0, u)$. As such, $u \mapsto J'_\lambda(\beta^0; u)$ is itself convex and satisfies $\partial J'_\lambda(\beta^0; u) \subset \partial J_\lambda(\beta^0)$ for all $u \in \mathbb{R}^p$. Note further that the result

in Theorem 2.2 is valid not only for the SLOPE penalty, but more generally by replacing the sequence of regularizers J_{λ^n} with $n^{1/2}f$ for any convex penalty f . Finally, we remark that we can use proximal methods to solve the optimization problem (5). The proximal operator of the directional SLOPE derivative is described in the Appendix A.2. We also refer the reader to [29], where the directional derivative of the SLOPE penalty is used for a coordinate descent algorithm for SLOPE.

Furthermore, the estimator $\hat{\mathbf{u}}$ minimizes $V(u)$ and satisfies the optimality condition

$$0 \in Cu - W + \partial J'_{\lambda}(\beta^0; u), \quad (10)$$

which yields

$$u \in C^{-1}(W - \partial J'_{\lambda}(\beta^0; u)) \subset C^{-1}(W - \partial J_{\lambda}(\beta^0)).$$

It follows that $\hat{\mathbf{u}} = C^{-1}(W - v)$, for some $v = v_W \in \partial J_{\lambda}(\beta^0)$. Although the exact value of $v = v_W$ depends on W , we can gain some insight into the structure of v merely from the fact that $v \in \partial J_{\lambda}(\beta^0)$. As we shall see, the subdifferential $\partial J_{\lambda}(\beta^0)$ respects the signs of β^0 and we have $\text{sgn}(v_i) = \text{sgn}(\beta_i^0)$, $\forall i$ such that $\beta_i^0 \neq 0$. Consequently, sampling $C\hat{\mathbf{u}}$ amounts to sampling $W \sim \mathcal{N}(0, \sigma^2 C)$ and then pushing each component in the opposite direction of $\text{sgn}(\beta_i^0)$. The total magnitude of the push v satisfies $\sum_{i=1}^p |v_i| \leq \sum_{i=1}^p \lambda_i$. (In fact, the total magnitude of v on the non-zero clusters of β_i^0 is given by a corresponding sum of λ_i 's). We can think of the push by v as introducing a bias, which pushes W in the opposite direction of $\text{sgn}(\beta^0)$. In the special case of LASSO estimator, $\lambda_1 = \dots = \lambda_p = c > 0$ and on the non-zero entries of β^0 , the bias is given by $|v_i| = c$. In case $\lambda = 0$, the subdifferential vanishes, hence $v = 0$ and $\hat{\mathbf{u}} \sim \mathcal{N}(0, \sigma^2 C^{-1})$.

3 Subdifferential and Pattern

In this section we investigate a “duality” between the SLOPE pattern and its subdifferential. The subdifferential of the SLOPE norm has already been explored in [12, 14, 28, 44]. For clarity of exposition, we provide a concise derivation of the subdifferential formula from [12] in terms of symmetries of the pattern (14). In subsection 3.1, we make use of the Hausdorff distance, a suitable notion for the convergence of SLOPE subdifferentials.

Note that pattern induces an equivalence relation on \mathbb{R}^p ; $u \sim v \iff \mathbf{patt}(u) = \mathbf{patt}(v)$. The information contained in a pattern $\mathbf{p} = \mathbf{patt}(u) \in \mathfrak{P}$ is threefold: First, it captures the information about the partition of $\{1, \dots, p\}$ into clusters $\mathcal{I}(\mathbf{p}) = \mathcal{I}(u) = \{I_0, I_1, \dots, I_m\}$. Second, it preserves the ranking of the respective clusters, with I_0 corresponding to the 0-cluster and I_m corresponding to the cluster of the greatest magnitude. And third, it preserves the signs of u_i . The full information of a pattern \mathbf{p} is captured by the ordered partition $(I_0, I_1^+, I_1^-, \dots, I_m^+, I_m^-)$, where $I_j^+ = \{i \in I_j : u_i > 0\}$ and $I_j^- = \{i \in I_j : u_i < 0\}$.

Remark 3.1. A crucial property of pattern is its close connection to the subdifferential, which was extensively studied in [38]. In fact,

$$\mathbf{patt}(u) = \mathbf{patt}(v) \implies \partial J_{\lambda}(u) = \partial J_{\lambda}(v) \quad \forall u, v \in \mathbb{R}^p. \quad (11)$$

Moreover, if $\lambda_1 > \dots > \lambda_p > 0$, the reverse implication also holds:

$$\mathbf{patt}(u) = \mathbf{patt}(v) \iff \partial J_\lambda(u) = \partial J_\lambda(v) \quad \forall u, v \in \mathbb{R}^p;$$

see [7, 38].

In particular, (11) asserts that $\partial J_\lambda(u)$ is constant on $\mathbf{patt}^{-1}(\mathbf{p})$, which allows for a well-defined notion of a subdifferential at a pattern.

Definition 3.2. Let $\mathbf{p} \in \mathfrak{P}$ be any pattern. We define the SLOPE-subdifferential at \mathbf{p} as $\partial J_\lambda(\mathbf{p}) := \partial J_\lambda(u)$, where $u \in \mathbb{R}^p$ with $\mathbf{patt}(u) = \mathbf{p}$.

We introduce some helpful notation to illuminate how the subdifferential arises naturally from the symmetries of the pattern. Abbreviate the diagonal matrices with entries $+1$ or -1 with $\{\pm 1\}^p$ and for $\mathbf{p} \in \mathfrak{P}$ with $\mathcal{I}(\mathbf{p}) = \{I_0, I_1, \dots, I_m\}$, define the sign matrix $S_{\mathbf{p}} \in \{\pm 1\}^p$ by $(S_{\mathbf{p}})_{ii} := 1$ for $i \in I_0$ and $(S_{\mathbf{p}})_{ii} := \text{sgn}(\mathbf{p}_i)$ else. The chain rule yields:

$$\begin{aligned} \text{i)} \quad & \partial J_\lambda(S\mathbf{p}) = S\partial J_\lambda(\mathbf{p}) \quad \forall S \in \{\pm 1\}^p \\ \text{ii)} \quad & \partial J_\lambda(|\mathbf{p}|) = S_{\mathbf{p}}\partial J_\lambda(\mathbf{p}) \end{aligned}$$

Further, denote by \mathcal{S}_p the set of permutation matrices on \mathbb{R}^p and \mathcal{S}_I the subset of all $p \times p$ matrices, where the $I \times I$ submatrix is a permutation matrix and all other entries are zero. Also, consider the direct product $G = \{\pm 1\}^p \cdot \mathcal{S}_p$. In particular, let $\mathcal{S}(\mathbf{p})$ denote the group of symmetries of $|\mathbf{p}|$ in G :

$$\begin{aligned} \mathcal{S}(\mathbf{p}) &:= \{\Sigma \in G : \Sigma|\mathbf{p}| = |\mathbf{p}|\} \\ &= \mathcal{S}_{I_0}^{+/-} \oplus \mathcal{S}_{I_1} \oplus \dots \oplus \mathcal{S}_{I_m}. \end{aligned}$$

These are the matrices, which act as permutations on the non-zero clusters I_1, \dots, I_m ; and allow additional swapping of signs on I_0 .

Moreover, for a given $v \in \mathbb{R}^p$, consider a permutation matrix Π_v^T , which sorts $|v|$, meaning that $(\Pi_v^T|v|)_1 \geq \dots \geq (\Pi_v^T|v|)_p$. In particular, we have $J_\lambda(v) = \langle \boldsymbol{\lambda}, \Pi_v^T|v| \rangle = \langle \Pi_v \boldsymbol{\lambda}, |v| \rangle$. Analogously, for any pattern $\mathbf{p} \in \mathfrak{P}$ we define the set

$$\Pi(\mathbf{p}) := \{\Pi \in \mathcal{S}_p : (\Pi^T|\mathbf{p}|)_1 \geq \dots \geq (\Pi^T|\mathbf{p}|)_p\},$$

and shall write $\Pi_{\mathbf{p}}$ for an arbitrary, but fixed element in $\Pi(\mathbf{p})$.

Example 3.3. Let $\mathbf{p} = (0, 2, -2, 1, 2, 1)^T$, $\mathcal{I}(\mathbf{p}) = \{\{1\}, \{4, 6\}, \{2, 3, 5\}\}$.

$$\begin{aligned} \Sigma = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix} \in \mathcal{S}(\mathbf{p}) = \mathcal{S}_{I_0}^{+/-} \oplus \mathcal{S}_{I_1} \oplus \mathcal{S}_{I_2} & \quad \Pi_{\mathbf{p}}^T = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \in \Pi(\mathbf{p}) \\ \Sigma\mathbf{p} = (0, -2, 2, 1, 2, 1)^T & \quad \Pi_{\mathbf{p}}^T\mathbf{p} = (2, -2, 2, 1, 1, 0)^T \end{aligned}$$

Now consider a fixed $u \in \mathbb{R}^p$, $\mathbf{p} = \mathbf{patt}(u)$, and $\Pi_{\mathbf{p}} \in \Pi(\mathbf{p})$. We can write

$$\begin{aligned} J_{\lambda}(u) &= \max_{\Pi \in \mathcal{S}_{\mathbf{p}}} \langle \Pi \boldsymbol{\lambda}, |u| \rangle \\ &= \langle \Pi_{\mathbf{p}} \boldsymbol{\lambda}, |u| \rangle. \end{aligned}$$

More importantly, for a sufficiently small¹ $\varepsilon > 0$, we have for all $v \in B_{\varepsilon}(u)$:

$$\begin{aligned} J_{\lambda}(v) &= \max \{ \langle \Sigma \Pi_{\mathbf{p}} \boldsymbol{\lambda}, |v| \rangle : \Sigma \in \mathcal{S}(\mathbf{p}) \} \\ &= \max \{ \langle S_{\mathbf{p}} \Sigma \Pi_{\mathbf{p}} \boldsymbol{\lambda}, v \rangle : \Sigma \in \mathcal{S}(\mathbf{p}) \} \stackrel{\text{not.}}{=} J_{\lambda}(\mathbf{p}, v). \end{aligned} \quad (12)$$

In general, for $f = \max\{f_1, \dots, f_N\}$, where f_i are smooth functions, the subdifferential of f at a point x is given by the convex hull:

$$\partial f(x) = \text{con} \{ \nabla f_i(x) : i \in I(x) \}, \quad (13)$$

where $I(x) := \{i : f_i(x) = f(x)\}$; (see [36] Exercise 8.31). Combining (12) and (13), one obtains an explicit representation of the subdifferential of SLOPE as a convex hull:

$$\begin{aligned} \partial J_{\lambda}(\mathbf{p}) &= \text{con} \{ S_{\mathbf{p}} \Sigma \Pi_{\mathbf{p}} \boldsymbol{\lambda} : \Sigma \in \mathcal{S}(\mathbf{p}) \} \\ &= S_{\mathbf{p}} \text{con} \{ \mathcal{S}(\mathbf{p}) \} \Pi_{\mathbf{p}} \boldsymbol{\lambda}. \end{aligned} \quad (14)$$

Notice how the subdifferential is fully captured by the group of symmetries $\mathcal{S}(\mathbf{p})$ of the pattern. We can also view the subdifferential as a direct sum of convex hulls on each of the clusters (see [12]):

$$\begin{aligned} \partial J_{\lambda}(\mathbf{p}) &= \left\{ v \in \mathbb{R}^p \left| \begin{array}{l} v_{I_j} \in S_{\mathbf{p}} \text{con} \{ \mathcal{S}_{I_j} \} \Pi_{\mathbf{p}} \boldsymbol{\lambda}, \quad j = 1, \dots, m \\ v_{I_0} \in S_{\mathbf{p}} \text{con} \{ \mathcal{S}_{I_0}^{+/-} \} \Pi_{\mathbf{p}} \boldsymbol{\lambda} \end{array} \right. \right\} \\ &= \text{con} \{ S_{\mathbf{p}} \mathcal{S}_{I_0}^{+/-} \Pi_{\mathbf{p}} \boldsymbol{\lambda} \} \oplus \text{con} \{ S_{\mathbf{p}} \mathcal{S}_{I_1} \Pi_{\mathbf{p}} \boldsymbol{\lambda} \} \oplus \dots \oplus \text{con} \{ S_{\mathbf{p}} \mathcal{S}_{I_m} \Pi_{\mathbf{p}} \boldsymbol{\lambda} \}. \end{aligned} \quad (15)$$

In other words, the subdifferential is obtained by assigning $\boldsymbol{\lambda}'$ s to each cluster according to $\Pi_{\mathbf{p}}$ and then taking the convex hull of all possible permutations within each cluster, whilst keeping track of the signs. The convex hull $\text{con} \mathcal{S}_{I_j}$ is given by the Birkhoff polytopes on I_j (also known as the doubly stochastic matrices), which are the $I_j \times I_j$ matrices, where each row and column sums to one. We refer the reader to [12] for further details about the representation of the SLOPE subdifferential in terms of Birkhoff polytopes and to [38] and [28] for different derivations of the SLOPE subdifferential.

3.1 Hausdorff distance and directional SLOPE derivative

We shall now consider the directional derivative in (6):

$$J'_{\lambda}(\beta^0; u) = \sum_{i=1}^p \lambda_{\pi(i)} [u_i \text{sgn}(\beta_i^0) \mathbb{I}[\beta_i^0 \neq 0] + |u_i| \mathbb{I}[\beta_i^0 = 0]].$$

¹We want ε to be smaller than one half of the smallest gap between the clusters.

Recall from (9) that for every u ; $J_{\lambda^n}(\beta^0 + u/\sqrt{n}) - J_{\lambda^n}(\beta^0) \rightarrow J_{\lambda}(\beta^0; u)$ as $\lambda^n/\sqrt{n} \rightarrow \lambda$. We will argue, that the convergence is carried over to the convergence of the subdifferentials $\partial_u(J_{\lambda^n}(\beta^0 + u/\sqrt{n})) \xrightarrow{d_H} \partial J_{\lambda}(\beta^0; u)$ in the Hausdorff metric.

To that end let $B^\delta := \{x \in \mathbb{R}^d : d(x, B) \leq \delta\} = B + \overline{B_\delta(0)}$, where $\overline{B_\delta(x)}$ is the closed δ -ball around x , and denote $B^{-\delta} := \{x \in \mathbb{R}^d : \overline{B_\delta(x)} \subset B\}$. For non-empty sets $A, B \subset \mathbb{R}^p$, the Hausdorff distance (also Pompeiu-Hausdorff distance) is defined as

$$d_H(A, B) := \inf\{\delta \geq 0 \mid A \subset B^\delta, B \subset A^\delta\};$$

see [36]. The Hausdorff distance defines a pseudo-metric, and yields a metric on the space of all closed, bounded, non-empty subsets of \mathbb{R}^p . For a sequence of sets A_n , we write $A_n \xrightarrow{d_H} A$, if $d_H(A_n, A) \rightarrow 0$ as $n \rightarrow \infty$. We collect several basic properties of the Hausdorff distance. First, if $A_n \xrightarrow{d_H} A, A'_n \xrightarrow{d_H} A'$, then also $A_n + A'_n \xrightarrow{d_H} A + A'$. It follows that if $B_n \xrightarrow{d_H} B$, then $B_n^\delta \xrightarrow{d_H} B^\delta$ and $B_n^{-\delta} \xrightarrow{d_H} B^{-\delta}$, because $d_H(B_n^{-\delta}, B^{-\delta}) \leq d_H(B_n, B)$. Finally, one can check that if $x_n^i \rightarrow x^i$ as $n \rightarrow \infty$ are N convergent sequences of points in \mathbb{R}^p , then

$$\text{con}\{x_n^1, \dots, x_n^N\} \xrightarrow{d_H} \text{con}\{x^1, \dots, x^N\}. \quad (16)$$

In particular, convergence of convex sets in Hausdorff distance is compatible with convergence in distribution of random vectors.

Lemma 3.4. *Suppose $B_n \xrightarrow{d_H} B$, where B_n and B are convex sets in \mathbb{R}^p . If $W_n \xrightarrow{d} W$, for some W with a continuous bounded density w.r.t. the Lebesgue measure on \mathbb{R}^p . Then $\mathbb{P}[W_n \in B_n] \rightarrow \mathbb{P}[W \in B]$.*

Proof. Let $\delta > 0$ be arbitrary. Since $B_n \xrightarrow{d_H} B$, it follows that $B_n \subset B^\delta$ eventually. Similarly, for all sufficiently large n we have $B \subset B_n^\delta$, thus $B^{-\delta} \subset (B_n^\delta)^{-\delta} = B_n$, where last equality follows from convexity² and a hyperplane separation argument. As a result, for any $\delta > 0$ $B^{-\delta} \subset B_n \subset B^\delta$ eventually. Moreover, since B is convex, one can show that for every $\varepsilon > 0$ there exists³ a $\delta > 0$ such that

$$\mathbb{P}[W \in B^\delta] - \varepsilon \leq \mathbb{P}[W \in B] \leq \mathbb{P}[W \in B^{-\delta}] + \varepsilon. \quad (17)$$

Consequently, for any $\varepsilon > 0$ we can choose $\delta > 0$ sufficiently small such that:

$$\begin{aligned} \limsup_{n \rightarrow \infty} \mathbb{P}[W_n \in B_n] &\leq \limsup_{n \rightarrow \infty} \mathbb{P}[W_n \in B^{\delta/2}] \leq \mathbb{P}[W \in B^\delta] \leq \mathbb{P}[W \in B] + \varepsilon \\ \liminf_{n \rightarrow \infty} \mathbb{P}[W_n \in B_n] &\geq \liminf_{n \rightarrow \infty} \mathbb{P}[W_n \in B^{-\delta/2}] \geq \mathbb{P}[W \in B^{-\delta}] \geq \mathbb{P}[W \in B] - \varepsilon, \end{aligned}$$

where we have used the Portmanteau Lemma and that $\overline{B^{\delta/2}} \subset B^\delta$, $(B^{-\delta/2})^\circ \supset B^{-\delta}$. This shows the desired convergence $\mathbb{P}[W_n \in B_n] \rightarrow \mathbb{P}[W \in B]$. \square

²Convexity is necessary: The annuli $B_n = \overline{B_1(0)} \setminus B_{1/n}(0) \xrightarrow{d_H} \overline{B_1(0)}$, but $\overline{B_1(0)}^{-\delta} \not\subset B_n$.

³In fact, the bounds with tubular sets hold uniformly over all convex sets; i.e., for each $\varepsilon > 0$ there even exists a $\delta > 0$ such that (17) holds for every convex set B .

An immediate consequence of (14) and (16) is that $\partial J_{\lambda^n/\sqrt{n}}(\mathbf{p}) \xrightarrow{d_H} \partial J_{\lambda}(\mathbf{p}) \forall \mathbf{p} \in \mathfrak{P}$. To get an analogous statement for $J'_{\lambda}(\beta^0; u)$, observe that for any fixed $u \in \mathbb{R}^p$, the pattern $\mathbf{patt}(\beta^0 + u/\sqrt{n})$ eventually stabilizes⁴ as n increases. This simple, but key idea of a limiting pattern lies at the core of the directional SLOPE derivative $J'_{\lambda}(\beta^0; u)$, and deserves its own definition.

Definition 3.5. Let $u \in \mathbb{R}^p$, we define the *limiting pattern of u with respect to β^0* , denoted by $\mathbf{patt}_{\beta^0}(u)$, as

$$\mathbf{patt}_{\beta^0}(u) := \lim_{\varepsilon \searrow 0} \mathbf{patt}(\beta^0 + \varepsilon u).$$

For a pattern $\mathbf{p} \in \mathfrak{P}$ we define its limiting pattern by $\mathbf{patt}_{\beta^0}(\mathbf{p}) = \mathbf{patt}_{\beta^0}(u)$, where $u \in \mathbb{R}^p$ is any vector with $\mathbf{patt}(u) = \mathbf{p}$.

Note that $\mathbf{patt}_{\beta^0}(\mathbf{p})$ is well defined, since if $\mathbf{patt}(u) = \mathbf{patt}(u')$ then also $\mathbf{patt}_{\beta^0}(u) = \mathbf{patt}_{\beta^0}(u')$.

Example 3.6. An example of such a triple $\beta^0, \mathbf{p}, \mathbf{patt}_{\beta^0}(\mathbf{p})$:

$$\begin{aligned} \beta^0 &= (0 & 0 & | 3 & 3 & -3 & | 7 & 7 & 7 & 7 & 7 &) \\ \mathbf{p} &= (0 & | 1 & 1 & 1 & -1 & -1 & -1 & | 2 & -2 & -2 &) \\ \mathbf{patt}_{\beta^0}(\mathbf{p}) &= (0 & | 1 & 2 & 2 & -2 & | 4 & 4 & | 5 & 3 & 3 &) \end{aligned}$$

Recall the notation in (12): $J_{\lambda}(\mathbf{p}; v) = \max \{ \langle S_{\mathbf{p}} \Sigma \Pi_{\mathbf{p}} \boldsymbol{\lambda}, v \rangle : \Sigma \in \mathcal{S}(\mathbf{p}) \}$ and the key insight that locally around some $u \in \mathbb{R}^p$ with $\mathbf{patt}(u) = \mathbf{p}$, we can represent $J_{\lambda}(v) = J_{\lambda}(\mathbf{p}, v)$. The following result asserts that the directional derivative $J'_{\lambda}(\beta^0, v)$ has a similar local representation in a neighborhood of $u \in \mathbb{R}^p$: $J'_{\lambda}(\beta^0, v) = J_{\lambda}(\mathbf{p}_0; v)$, where \mathbf{p}_0 is the limiting pattern of u w.r.t. β^0 .

Proposition 3.7. *Given $u \in \mathbb{R}^p$, $\mathbf{p} = \mathbf{patt}(u)$, $\mathbf{p}_0 = \mathbf{patt}_{\beta^0}(\mathbf{p})$, the following holds:*

$$i) \exists \varepsilon > 0 \forall v \in B_{\varepsilon}(u) : J'_{\lambda}(\beta^0; v) = J_{\lambda}(\mathbf{p}_0; v), \quad (18)$$

$$ii) \partial J'_{\lambda}(\beta^0; \mathbf{p}) = \partial J_{\lambda}(\mathbf{p}_0), \quad (19)$$

$$iii) \partial_u J_{\lambda^n}(\beta^0 + u/\sqrt{n}) \xrightarrow{d_H} \partial J'_{\lambda}(\beta^0; \mathbf{p}). \quad (20)$$

Proof. Take $M > 0$ s.t. $\mathbf{patt}(\beta^0 + u/\sqrt{n}) = \mathbf{p}_0$ for all $n \geq M$. By (12), for all v in a sufficiently small neighborhood $B_{\varepsilon}(u)$ around u ; $J_{\lambda^n}(\beta^0 + v/\sqrt{n}) = J_{\lambda^n}(\mathbf{p}_0; \beta^0 + v/\sqrt{n}) \forall n \geq M$. Moreover, observe that $J_{\lambda^n}(\beta^0) = \langle \Sigma \Pi_{\beta^0} \boldsymbol{\lambda}^n, |\beta^0| \rangle = \langle \Sigma \Pi_{\mathbf{p}_0} \boldsymbol{\lambda}^n, S_{\mathbf{p}_0} \beta^0 \rangle$ for every $\Sigma \in \mathcal{S}(\mathbf{p}_0)$, since $|\beta^0| = S_{\mathbf{p}_0} \beta^0$, $\Pi_{\mathbf{p}_0}^T$ orders $|\beta^0|$ and $\mathcal{S}(\mathbf{p}_0) \subset \mathcal{S}(\mathbf{patt}(\beta^0))$. Hence $\forall v \in B_{\varepsilon}(u)$:

$$\begin{aligned} J_{\lambda^n}(\beta^0 + v/\sqrt{n}) - J_{\lambda^n}(\beta^0) &= J_{\lambda^n}(\mathbf{p}_0; \beta^0 + v/\sqrt{n}) - J_{\lambda^n}(\beta^0) \\ &= \max_{\Sigma \in \mathcal{S}(\mathbf{p}_0)} \langle S_{\mathbf{p}_0} \Sigma \Pi_{\mathbf{p}_0} \boldsymbol{\lambda}^n, \beta^0 + v/\sqrt{n} \rangle - J_{\lambda^n}(\beta^0) \\ &= \max_{\Sigma \in \mathcal{S}(\mathbf{p}_0)} \langle S_{\mathbf{p}_0} \Sigma \Pi_{\mathbf{p}_0} \boldsymbol{\lambda}^n / \sqrt{n}, v \rangle \\ &\longrightarrow \max_{\Sigma \in \mathcal{S}(\mathbf{p}_0)} \langle S_{\mathbf{p}_0} \Sigma \Pi_{\mathbf{p}_0} \boldsymbol{\lambda}, v \rangle = J_{\lambda}(\mathbf{p}_0; v). \end{aligned}$$

⁴This means there exists $M > 0$ such that $\forall n \geq M \mathbf{patt}(\beta^0 + u/\sqrt{n}) = \mathbf{patt}_{\beta^0}(u)$.

From (9), we also know that $J_{\lambda^n}(\beta^0 + v/\sqrt{n}) - J_{\lambda^n}(\beta^0) \rightarrow J'_\lambda(\beta^0; v)$, which yields claim *i*). As a result of *i*); $\partial J'_\lambda(\beta^0; u) = \partial J_\lambda(\mathbf{p}_0; u) = \text{con} \{S_{\mathbf{p}_0} \Sigma \Pi_{\mathbf{p}_0} \boldsymbol{\lambda} : \Sigma \in \mathcal{S}(\mathbf{p}_0)\} = \partial J_\lambda(\mathbf{p}_0)$, where the second equality follows from (13) by observing that the maximum of $\langle S_{\mathbf{p}_0} \Sigma \Pi_{\mathbf{p}_0} \boldsymbol{\lambda}, u \rangle$ is attained for all $\Sigma \in \mathcal{S}(\mathbf{p}_0)$. This shows *ii*). Finally, for $n \geq M$; $\partial_u J_{\lambda^n}(\beta^0 + u/\sqrt{n}) = \partial J_{\lambda^n/\sqrt{n}}(\mathbf{p}_0)$, which converges in Hausdorff distance to $\partial J_\lambda(\mathbf{p}_0)$ by (14) and (16), finishing the proof. \square

4 Weak pattern convergence

In this section we show that on top of the weak convergence established in Theorem 2.2 the sequence of patterns $\mathbf{patt}(\hat{\mathbf{u}}_n)$ converges weakly to $\mathbf{patt}(\hat{\mathbf{u}})$. We remark that the weak pattern convergence does not follow merely from the weak convergence of $\hat{\mathbf{u}}_n$ to $\hat{\mathbf{u}}$, since the pattern function $\mathbf{patt} : \mathbb{R}^d \rightarrow \mathbb{Z}^d$ is not continuous. (See Example A.1.)

Theorem 4.1. *For any pattern $\mathbf{p} \in \mathfrak{P}$ we have:*

$$\mathbb{P}[\mathbf{patt}(\hat{\mathbf{u}}_n) = \mathbf{p}] \xrightarrow[n \rightarrow \infty]{} \mathbb{P}[\mathbf{patt}(\hat{\mathbf{u}}) = \mathbf{p}].$$

Before proving the theorem, we make a few preparatory observations. Using the notation from (4), $C_n := n^{-1} X^T X$, $W_n = n^{-1/2} X^T \varepsilon$ and $W \sim \mathcal{N}(0, \sigma^2 C)$, the optimality conditions arising from (5) and (7) in terms of subdifferentials read

$$\begin{aligned} W &\in C\hat{\mathbf{u}} + \partial J'_\lambda(\beta^0; \hat{\mathbf{u}}), \\ W_n &\in C_n \hat{\mathbf{u}}_n + \partial J_{\lambda^n/\sqrt{n}}(\beta^0 + \hat{\mathbf{u}}_n/\sqrt{n}). \end{aligned}$$

Given $\mathbf{p} \in \mathfrak{P}$ and any bounded set $D^{\mathbf{p}} \subset \mathbf{patt}^{-1}(\mathbf{p})$, we have for sufficiently large n :

$$\begin{aligned} \hat{\mathbf{u}} \in D^{\mathbf{p}} &\iff W \in CD^{\mathbf{p}} + \partial J'_\lambda(\beta^0; \mathbf{p}) =: B, \\ \hat{\mathbf{u}}_n \in D^{\mathbf{p}} &\iff W_n \in C_n D^{\mathbf{p}} + \partial J_{\lambda^n/\sqrt{n}}(\mathbf{p}_0) =: B_n, \end{aligned} \tag{21}$$

where $\mathbf{p}_0 = \mathbf{patt}_{\beta^0}(\mathbf{p})$, and B_n is a random set.

Proof. (Theorem 4.1) Let $\varepsilon > 0$ be arbitrary. By tightness of $\hat{\mathbf{u}}_n$ there exists $M > 0$ such that $\mathbb{P}[\hat{\mathbf{u}}_n \notin B_M(0)] < \varepsilon \forall n$. For each pattern $\mathbf{p} \in \mathfrak{P}$ define $D^{\mathbf{p}} := \mathbf{patt}^{-1}(\mathbf{p}) \cap B_M(0)$. Further, let $B = CD^{\mathbf{p}} + \partial J'_\lambda(\beta^0; \mathbf{p})$ and $B_n = C_n D^{\mathbf{p}} + \partial J_{\lambda^n/\sqrt{n}}(\mathbf{p}_0)$ with $\mathbf{p}_0 = \mathbf{patt}_{\beta^0}(\mathbf{p})$, as in (21). We have already established in Proposition 3.7 that $d_H(\partial J_{\lambda^n/\sqrt{n}}(\mathbf{p}_0), \partial J'_\lambda(\beta^0; \mathbf{p})) \rightarrow 0$. Moreover, since $C_n \xrightarrow{a.s.} C$, we have $d_H(C_n D^{\mathbf{p}}, CD^{\mathbf{p}}) \xrightarrow{a.s.} 0$.⁵ It follows that $d_H(B_n, B) \xrightarrow{a.s.} 0$ and Lemma 3.4 yields

$$\mathbb{P}[\hat{\mathbf{u}}_n \in D^{\mathbf{p}}] = \mathbb{P}[W_n \in B_n] \xrightarrow[n \rightarrow \infty]{} \mathbb{P}[W \in B] = \mathbb{P}[\hat{\mathbf{u}} \in D^{\mathbf{p}}].$$

Hence

$$\begin{aligned} \limsup_{n \rightarrow \infty} \mathbb{P}[\hat{\mathbf{u}}_n \in \mathbf{patt}^{-1}(\mathbf{p})] &\leq \limsup_{n \rightarrow \infty} \mathbb{P}[\hat{\mathbf{u}}_n \in D^{\mathbf{p}}] + \varepsilon = \mathbb{P}[\hat{\mathbf{u}} \in D^{\mathbf{p}}] + \varepsilon, \\ \liminf_{n \rightarrow \infty} \mathbb{P}[\hat{\mathbf{u}}_n \in \mathbf{patt}^{-1}(\mathbf{p})] &\geq \liminf_{n \rightarrow \infty} \mathbb{P}[\hat{\mathbf{u}}_n \in D^{\mathbf{p}}] = \mathbb{P}[\hat{\mathbf{u}} \in D^{\mathbf{p}}], \end{aligned}$$

and letting $\varepsilon \downarrow 0$ yields the desired pattern convergence. \square

⁵Boundedness of $D^{\mathbf{p}}$ grants continuity of the map $A \mapsto d_H(AD^{\mathbf{p}}, CD^{\mathbf{p}})$ on $M_n(\mathbb{R})$ and thus measurability of $\omega \mapsto d_H(C_n(\omega)D^{\mathbf{p}}, CD^{\mathbf{p}})$.

4.1 Weak pattern recovery

As a consequence of Theorem 4.1, we can characterize the asymptotic distribution of the SLOPE pattern $\mathbf{patt}(\hat{\beta}_n)$ in terms of $\hat{\mathbf{u}}$.

Corollary 4.2. *For any pattern $\mathbf{p} \in \mathfrak{P}$ we have:*

$$\mathbb{P}[\mathbf{patt}(\hat{\beta}_n) = \mathbf{p}] \xrightarrow[n \rightarrow \infty]{} \mathbb{P}[\mathbf{patt}_{\beta^0}(\hat{\mathbf{u}}) = \mathbf{p}].$$

Proof. By Theorem 4.1, sampling a pattern $\mathbf{p} \in \mathfrak{P}$ according to $\mathbf{p} \sim \mathbf{patt}(\hat{\mathbf{u}}_n)$ is asymptotically equivalent to sampling $\mathbf{p} \sim \mathbf{patt}(\hat{\mathbf{u}})$. Therefore, sampling a pattern $\mathbf{p} \in \mathfrak{P}$ according to $\mathbf{p} \sim \mathbf{patt}(\hat{\beta}_n) = \mathbf{patt}(\beta^0 + \hat{\mathbf{u}}_n/\sqrt{n})$ is asymptotically equivalent to sampling according to $\mathbf{p} \sim \mathbf{patt}(\beta^0 + \hat{\mathbf{u}}/\sqrt{n})$, which is in turn asymptotically equivalent to sampling $\mathbf{p} \sim \mathbf{patt}_{\beta^0}(\hat{\mathbf{u}})$. \square

A natural question concerns the limiting probability of pattern recovery of the true signal vector β^0 . More precisely, we're interested in $\mathbb{P}[\mathbf{patt}(\hat{\beta}_n) = \mathbf{patt}(\beta^0)]$ as $n \rightarrow \infty$. Using properties of $\hat{\mathbf{u}}$, we find a new way of proving the result given in Theorem 4.2 i) [7]. Let $\mathcal{I}(\beta^0) = \{I_0, I_1, \dots, I_m\}$ and write the basis vectors $\mathbf{1}_{I_1}, \dots, \mathbf{1}_{I_m}$ as columns in a ‘‘pattern’’ matrix $U_{\beta^0} = S_{\beta^0}(\mathbf{1}_{I_m} | \dots | \mathbf{1}_{I_1})$ and let $\Lambda_0 = S_{\beta^0} \Pi_{\beta^0} \boldsymbol{\lambda}$. (In the notation in [7] $U_{\beta^0}^T \Lambda_0 = \tilde{\Lambda}_{b_0}$ denotes the clustered parameter.) Also, by Proposition 2.1. in [7], the dual unit ball coincides with the subdifferential at zero $\{v \in \mathbb{R}^p : J_{\boldsymbol{\lambda}}^*(v) \leq 1\} = \partial J_{\boldsymbol{\lambda}}(0)$ and the subdifferential satisfies

$$\begin{aligned} \partial J_{\boldsymbol{\lambda}}(\beta^0) &= \{v \in \mathbb{R}^p : v \in \partial J_{\boldsymbol{\lambda}}(0) \text{ and } U_{\beta^0}^T(v - \Lambda_0) = 0\} \\ &= \partial J_{\boldsymbol{\lambda}}(0) \cap (\Lambda_0 + \langle U_{\beta^0} \rangle^\perp), \end{aligned} \quad (22)$$

where $\langle U_{\beta^0} \rangle$ denotes the column span of U_{β^0} and $\langle U_{\beta^0} \rangle^\perp$ its orthogonal complement. Equivalently, we have $\partial J_{\boldsymbol{\lambda}}(\beta^0) - \Lambda_0 = (\partial J_{\boldsymbol{\lambda}}(0) - \Lambda_0) \cap \langle U_{\beta^0} \rangle^\perp$ and in particular $\partial J_{\boldsymbol{\lambda}}(\beta^0) - \Lambda_0 \perp \langle U_{\beta^0} \rangle$. Finally, notice the elegant relation between the limiting pattern $\mathbf{patt}_{\beta^0}(\cdot)$ and the pattern matrix $\langle U_{\beta^0} \rangle$:

$$\{u \in \mathbb{R}^p : \mathbf{patt}_{\beta^0}(u) = \mathbf{patt}(\beta^0)\} = \langle U_{\beta^0} \rangle. \quad (23)$$

Corollary 4.3 (Theorem 4.2 i) in [7]). *The limiting probability of pattern recovery satisfies*

$$\begin{aligned} \mathbb{P}[\mathbf{patt}(\hat{\beta}_n) = \mathbf{patt}(\beta^0)] &\xrightarrow[n \rightarrow \infty]{} \mathbb{P}[\hat{\mathbf{u}} \in \langle U_{\beta^0} \rangle] = \mathbb{P}[Z \in \partial J_{\boldsymbol{\lambda}}(0)], \\ Z &\sim \mathcal{N}(C^{1/2} P C^{-1/2} \Lambda_0, \sigma^2 C^{1/2} (I - P) C^{1/2}), \end{aligned}$$

where $P = C^{1/2} U_{\beta^0} (U_{\beta^0}^T C U_{\beta^0})^{-1} U_{\beta^0}^T C^{1/2}$ is the projection matrix onto $C^{1/2} \langle U_{\beta^0} \rangle$.

Proof. By Corollary 4.2 and (23), we obtain:

$$\mathbb{P}[\mathbf{patt}(\hat{\beta}_n) = \mathbf{patt}(\beta^0)] \longrightarrow \mathbb{P}[\mathbf{patt}_{\beta^0}(\hat{\mathbf{u}}) = \mathbf{patt}(\beta^0)] = \mathbb{P}[\hat{\mathbf{u}} \in \langle U_{\beta^0} \rangle].$$

Moreover, by Proposition 3.7 and (23), $\forall u \in \langle U_{\beta^0} \rangle; \partial J_{\boldsymbol{\lambda}}'(\beta^0; u) = \partial J_{\boldsymbol{\lambda}}(\mathbf{patt}_{\beta^0}(u)) = \partial J_{\boldsymbol{\lambda}}(\beta^0)$. Consequently, the optimality condition (10) $W \in C u + \partial J_{\boldsymbol{\lambda}}'(\beta^0; u)$ yields:

$$\begin{aligned} \hat{\mathbf{u}} \in \langle U_{\beta^0} \rangle &\iff W \in C \langle U_{\beta^0} \rangle + \partial J_{\boldsymbol{\lambda}}(\beta^0) \\ &\iff C^{-1/2} W \in C^{1/2} \langle U_{\beta^0} \rangle + C^{-1/2} \partial J_{\boldsymbol{\lambda}}(\beta^0) \\ &\iff \underbrace{-C^{-1/2} \Lambda_0 + C^{-1/2} W}_{=: Y} \in C^{1/2} \langle U_{\beta^0} \rangle + C^{-1/2} (\partial J_{\boldsymbol{\lambda}}(\beta^0) - \Lambda_0). \end{aligned} \quad (24)$$

Furthermore, (22) implies $C^{1/2}\langle U_{\beta^0} \rangle \perp C^{-1/2}(\partial J_{\lambda}(\beta^0) - \Lambda_0)$ and because $C^{1/2}\langle U_{\beta^0} \rangle = \langle P \rangle$, we have $C^{-1/2}(\partial J_{\lambda}(\beta^0) - \Lambda_0) \subset \langle P \rangle^{\perp} = \langle I - P \rangle$. Decomposing $Y = PY + (I - P)Y$, (24) splits into two conditions: $PY \in C^{1/2}\langle U_{\beta^0} \rangle$ and $(I - P)Y \in C^{-1/2}(\partial J_{\lambda}(\beta^0) - \Lambda_0)$. The first condition is always satisfied and the second is equivalent to $(I - P)Y \in C^{-1/2}(\partial J_{\lambda}(0) - \Lambda_0)$ by (22). Thus (24) yields

$$\begin{aligned} \hat{\mathbf{u}} \in \langle U_{\beta^0} \rangle &\iff (I - P)Y \in C^{-1/2}(\partial J_{\lambda}(0) - \Lambda_0) \\ &\iff \Lambda_0 + C^{1/2}(I - P)(-C^{-1/2}\Lambda_0 + C^{-1/2}W) \in \partial J_{\lambda}(0) \\ &\iff C^{1/2}PC^{-1/2}\Lambda_0 + C^{1/2}(I - P)C^{-1/2}W \in \partial J_{\lambda}(0), \end{aligned}$$

and using that $W \sim \mathcal{N}(0, \sigma^2 C)$, the above Gaussian vector has expectation $C^{1/2}PC^{-1/2}\Lambda_0$ and covariance matrix $\sigma^2 C^{1/2}(I - P)C^{1/2}$, which finishes the proof. \square

4.2 Pattern attainability

In the following section, we want to characterize all patterns that $\hat{\mathbf{u}}$ can attain. This will depend on the signal vector β^0 and the penalty vector λ .

Definition 4.4. We shall say a pattern $\mathbf{p} \in \mathfrak{P}$ is *attainable* by $\hat{\mathbf{u}}$, if $\mathbb{P}[\mathbf{patt}(\hat{\mathbf{u}}) = \mathbf{p}] > 0$.

First, consider a partial ordering on the partitions of $\{1, \dots, p\}$, where $\mathcal{I}^1 \preceq \mathcal{I}^2$ if every element (cluster) in \mathcal{I}^1 is a subset of some element in \mathcal{I}^2 ; in that case we will say that \mathcal{I}^1 is a *refinement* of \mathcal{I}^2 , for example $\{\{1, 2\}, \{3\}, \{5\}, \{4, 6\}\} \preceq \{\{1, 2, 3\}, \{4, 5, 6\}\}$. For $u, v \in \mathbb{R}^p$ such that $\max_i |u_i| < \min\{|v_i - v_j| : v_i \neq v_j\} / 2$, we have $\mathcal{I}(v + u) \preceq \mathcal{I}(v)$, and $\mathcal{I}(v + u) \preceq \mathcal{I}(u)$. In particular, for $\mathbf{p} \in \mathfrak{P}$ and $\mathbf{p}_0 = \mathbf{patt}_{\beta^0}(\mathbf{p})$ we have both $\mathcal{I}(\mathbf{p}_0) \preceq \mathcal{I}(\mathbf{p})$ and $\mathcal{I}(\mathbf{p}_0) \preceq \mathcal{I}(\beta^0)$.

In order to determine, which patterns are attainable by $\hat{\mathbf{u}}$, we need to consider the dimension of $\partial J_{\lambda}(\mathbf{p})$. To that end, using (15) we rewrite

$$\begin{aligned} \partial J_{\lambda}(\mathbf{p}) &= (\partial J_{\lambda}(\mathbf{p}) - S_{\mathbf{p}}\Pi_{\mathbf{p}}\lambda) + S_{\mathbf{p}}\Pi_{\mathbf{p}}\lambda \\ &= \text{con}\{S_{\mathbf{p}}(\mathcal{S}(\mathbf{p}) - \mathbb{I})\Pi_{\mathbf{p}}\lambda\} + S_{\mathbf{p}}\Pi_{\mathbf{p}}\lambda \\ &= \text{con}\{S_{\mathbf{p}}(\mathcal{S}_{I_0}^{+/-} - \mathbb{I}_{I_0})\Pi_{\mathbf{p}}\lambda\} \oplus \dots \oplus \text{con}\{S_{\mathbf{p}}(\mathcal{S}_{I_m} - \mathbb{I}_{I_m})\Pi_{\mathbf{p}}\lambda\} + S_{\mathbf{p}}\Pi_{\mathbf{p}}\lambda. \end{aligned}$$

For $j \in \{1, \dots, m\}$, we have $\dim\{\text{con}\{S_{\mathbf{p}}(\mathcal{S}_{I_j} - \mathbb{I}_{I_j})\Pi_{\mathbf{p}}\lambda\}\} = |I_j| - 1$, unless λ is constant on I_j . If λ is constant on I_j , the convex hull equals 0 and is zero-dimensional. For the zero cluster $\dim\{\text{con}\{S_{\mathbf{p}}(\mathcal{S}_{I_0}^{+/-} - \mathbb{I}_{I_0})\Pi_{\mathbf{p}}\lambda\}\} = |I_0|$, whenever λ is not constant zero on I_0 , otherwise the convex hull is 0. Consequently, for any $\mathbf{p} \in \mathfrak{P}$:

$$\dim \partial J_{\lambda}(\mathbf{p}) \leq p - |\mathcal{I}(\mathbf{p})| + 1, \quad (25)$$

with equality if and only if λ is non-constant on $I_j \in \{I_1, \dots, I_m\}$ and not equal to zero everywhere on I_0 . Also, consider a set of the form:

$$D^{\mathbf{p}} = \{u \in \mathbf{patt}^{-1}(\mathbf{p}) : |u_i| \in K_j \text{ for } i \in I_j, j = 1, \dots, m_{\mathbf{p}}\}, \quad (26)$$

where $K_1, \dots, K_{m_{\mathbf{p}}}$ are ordered compact disjoint positive intervals corresponding to the non-zero clusters in $\mathcal{I}(\mathbf{p}) = \{I_0, I_1, \dots, I_{m_{\mathbf{p}}}\}$ and $K_0 = \{0\}$. We denote the family of sets of the

form (26) by $\mathcal{D}^{\mathbf{p}}$. Using the refined partition $\{I_0, I_1^+, I_1^-, \dots, I_{m_{\mathbf{p}}}^+, I_{m_{\mathbf{p}}}^-\}$ we can also represent any $D^{\mathbf{p}} \in \mathcal{D}^{\mathbf{p}}$ explicitly as

$$\begin{aligned} D^{\mathbf{p}} &= \left\{ 0_{I_0} \oplus \alpha_1(\mathbf{1}_{I_1^+} - \mathbf{1}_{I_1^-}) \oplus \dots \oplus \alpha_m(\mathbf{1}_{I_{m_{\mathbf{p}}}^+} - \mathbf{1}_{I_{m_{\mathbf{p}}}^-}) : \alpha_i \in K_i, i = 1, \dots, m_{\mathbf{p}} \right\} \\ &= S_{\mathbf{p}} \{ \alpha_1 \mathbf{1}_{I_1} \oplus \dots \oplus \alpha_m \mathbf{1}_{I_m} : \alpha_i \in K_i, i = 1, \dots, m_{\mathbf{p}} \}, \end{aligned} \quad (27)$$

where $\mathbf{1}_I$ denotes the vector in \mathbb{R}^p with 1's on I and zeros everywhere else. We can characterize all attainable patterns by $\hat{\mathbf{u}}$:

Theorem 4.5. *A pattern $\mathbf{p} \in \mathfrak{P}$ with $\mathcal{I}(\mathbf{p}) = \{I_0, I_1, \dots, I_m\}$ is attainable by $\hat{\mathbf{u}}$ if and only if*

- i) $\mathcal{I}(\mathbf{p}) \preceq \mathcal{I}(\beta^0)$,*
- ii) $\text{sgn}(\mathbf{p}) \odot \text{sgn}(\beta^0)$ is constant on $I_j \in \mathcal{I}(\mathbf{p}) \ \forall j \neq 0$,*
- iii) $\boldsymbol{\lambda}$ is not constant on $I_j \in \mathcal{I}(\mathbf{p}) \ \forall j \neq 0$,*
- iv) $\boldsymbol{\lambda}$ is not constant zero on $I_0 \in \mathcal{I}(\mathbf{p})$.*

Remark 4.6. Condition *i)* asserts that, with probability one, the clusters of $\hat{\mathbf{u}}$ are contained within the clusters of β^0 . The first two conditions *i)* and *ii)* together are equivalent to $\mathcal{I}(\mathbf{p}) = \mathcal{I}(\mathbf{p}_0)$, where $\mathbf{p}_0 = \mathbf{patt}_{\beta^0}(\mathbf{p})$ is the limiting pattern of \mathbf{p} w.r.t. β^0 . Indeed, if either *i)* or *ii)* is violated then $\mathcal{I}(\mathbf{p}_0) \prec \mathcal{I}(\mathbf{p})$ is a proper refinement, see Example 3.6.

Proof. Let $\mathbf{p}_0 = \mathbf{patt}_{\beta^0}(\mathbf{p})$ be the limiting pattern of \mathbf{p} w.r.t. β^0 . Following the notation in (21) and (26), a pattern $\mathbf{p} \in \mathfrak{P}$ is attainable by $\hat{\mathbf{u}}$ if and only if $0 < \mathbb{P}[\hat{\mathbf{u}} \in D^{\mathbf{p}}] = \mathbb{P}[W \in B]$, where $W \sim \mathcal{N}(0, \sigma^2 C)$, $D^{\mathbf{p}} \in \mathcal{D}^{\mathbf{p}}$ and

$$\begin{aligned} B &= CD^{\mathbf{p}} + \partial J'_{\boldsymbol{\lambda}}(\beta^0; \mathbf{p}) \\ &= CD^{\mathbf{p}} + \partial J_{\boldsymbol{\lambda}}(\mathbf{p}_0), \end{aligned}$$

which is a consequence of (19). Therefore, a pattern \mathbf{p} is attainable if and only if B is a full-dimensional subset in \mathbb{R}^p . From (27), we have $\dim CD^{\mathbf{p}} = \dim D^{\mathbf{p}} = |\mathcal{I}(\mathbf{p})| - 1$ and from (25) we get $\dim \partial J_{\boldsymbol{\lambda}}(\mathbf{p}_0) \leq p - |\mathcal{I}(\mathbf{p}_0)| + 1$. It follows

$$\begin{aligned} \dim B &\leq \dim CD^{\mathbf{p}} + \dim \partial J_{\boldsymbol{\lambda}}(\mathbf{p}_0) \\ &\leq p - (|\mathcal{I}(\mathbf{p}_0)| - |\mathcal{I}(\mathbf{p})|) \end{aligned} \quad (28)$$

$$\leq p. \quad (29)$$

If *i)* or *ii)* is violated, then by Remark 4.6 we have $\mathcal{I}(\mathbf{p}_0) \prec \mathcal{I}(\mathbf{p})$, thus the inequality in (29) is strict and $\dim B < p$. If *iii)* or *iv)* are violated, the inequality in (28) is strict. Hence *i) - iv)* are necessary for attainability.

If *i) - iv)* are satisfied, then (29) and (28) are equalities. Therefore, to see that *i) - iv)* are sufficient for attainability, it remains to show that

$$\dim B = \dim CD^{\mathbf{p}} + \dim \partial J_{\boldsymbol{\lambda}}(\mathbf{p}_0).$$

This is a completely general fact, which is valid without any assumptions on \mathbf{p} , β^0 or $\boldsymbol{\lambda}$. It suffices to argue that $\text{span}CD^{\mathbf{p}} \cap \text{span}(\partial J_{\boldsymbol{\lambda}}(\mathbf{p}_0) - S_{\mathbf{p}_0}\Pi_{\mathbf{p}_0}\boldsymbol{\lambda}) = 0$, which will guarantee linear independence of the spans and additivity of their dimensions⁶. We set

$$\begin{aligned} V(\mathbf{p}_0) &= \text{span}(\partial J_{\boldsymbol{\lambda}}(\mathbf{p}_0) - S_{\mathbf{p}_0}\Pi_{\mathbf{p}_0}\boldsymbol{\lambda}) \\ &= \text{span}\{S_{\mathbf{p}_0}(\mathcal{S}(\mathbf{p}_0) - \mathbb{I})\Pi_{\mathbf{p}_0}\boldsymbol{\lambda}\}. \end{aligned}$$

Assume that the above intersection is non-trivial, meaning that there exists a non-zero vector $\boldsymbol{\alpha} = S_{\mathbf{p}}(\alpha_1\mathbf{1}_{I_1} \oplus \dots \oplus \alpha_m\mathbf{1}_{I_m})$ in $D^{\mathbf{p}}$ (representation (27)), such that $C\boldsymbol{\alpha}$ lies in $V(\mathbf{p}_0)$. Then there are coefficients $\{\beta_{\Sigma} : \Sigma \in \mathcal{S}(\mathbf{p}_0)\}$ such that

$$C\boldsymbol{\alpha} = \sum_{\Sigma \in \mathcal{S}(\mathbf{p}_0)} \beta_{\Sigma} S_{\mathbf{p}_0}(\Sigma - \mathbb{I})\Pi_{\mathbf{p}_0}\boldsymbol{\lambda}.$$

Also, one can check that $S_{\mathbf{p}_0}S_{\mathbf{p}}$ has a constant sign on each of the non zero clusters of \mathbf{p}_0 . It follows that $S_{\mathbf{p}_0}\boldsymbol{\alpha}$ is constant on each cluster of \mathbf{p}_0 . Therefore, $(S_{\mathbf{p}_0}\boldsymbol{\alpha})^T(\Sigma - \mathbb{I}) = 0$ for any $\Sigma \in \mathcal{S}(\mathbf{p}_0)$. This leads to a contradiction:

$$0 < \langle \boldsymbol{\alpha}, C\boldsymbol{\alpha} \rangle = \sum_{\Sigma \in \mathcal{S}(\mathbf{p}_0)} \beta_{\Sigma} \langle S_{\mathbf{p}_0}\boldsymbol{\alpha}, (\Sigma - \mathbb{I})\Pi_{\mathbf{p}_0}\boldsymbol{\lambda} \rangle = 0.$$

□

4.3 Asymptotic FDR control

We can harness the weak pattern convergence from Theorem 4.1 to characterize the asymptotic false discovery rate of the SLOPE estimator $\hat{\boldsymbol{\beta}}_n$ as n goes to infinity. Recall the notation from Theorem 2.2, where $\hat{\mathbf{u}}_n$ and $\hat{\mathbf{u}}$ are the minimizers of (7) and (5) respectively and $\hat{\boldsymbol{\beta}}_n = \beta^0 + \hat{\mathbf{u}}_n/\sqrt{n}$. Moreover, under the scaling $\boldsymbol{\lambda}^n/\sqrt{n} \rightarrow \boldsymbol{\lambda} \geq 0$, $\hat{\mathbf{u}}_n$ converges to $\hat{\mathbf{u}}$ both weakly and weakly in pattern. By Corollary 4.2, sampling a random pattern according to $\text{patt}(\hat{\boldsymbol{\beta}}_n)$ is asymptotically equivalent to sampling a limiting pattern $\hat{\mathbf{b}} := \text{patt}_{\beta^0}(\hat{\mathbf{u}})$.

Given the true signal β^0 , we formulate p null-hypotheses $H_{0i} : \beta_i^0 = 0$; and reject H_{0i} whenever $\hat{\boldsymbol{\beta}}_{n,i} \neq 0$. Let $R^n = |\{i : \hat{\boldsymbol{\beta}}_{n,i} \neq 0\}|$ denote the number of rejections and $V^n = |\{i : \hat{\boldsymbol{\beta}}_{n,i} \neq 0, \beta_i^0 = 0\}|$ the number of false rejections. Similarly, let $R = |\{i : \hat{\mathbf{b}}_i \neq 0\}|$ and $V = |\{i : \hat{\mathbf{b}}_i \neq 0, \beta_i^0 = 0\}|$ be the corresponding numbers of rejections and false rejections in the limiting experiment $\hat{\mathbf{b}}$. Asymptotic FDR of $\hat{\boldsymbol{\beta}}_n$ is matched by the FDR of $\hat{\mathbf{b}}$:

$$\text{FDR}(\hat{\boldsymbol{\beta}}_n) = \mathbb{E} \left[\frac{V^n}{1 \vee R^n} \right] \xrightarrow{n \rightarrow \infty} \mathbb{E} \left[\frac{V}{1 \vee R} \right] = \text{FDR}(\hat{\mathbf{b}}).$$

Moreover, note that by consistency $\hat{\mathbf{b}}_i \neq 0$, whenever $\beta_i^0 \neq 0$, and $\hat{\mathbf{b}}$ has full power.

Example 4.7. Suppose $C = \mathbb{I}_p$, and let $p_0 = |I_0|$. The objective function (5) simplifies to

$$\begin{aligned} V(u) &= \frac{1}{2}u^T C u - u^T W + J'_{\boldsymbol{\lambda}}(\beta^0; u) \\ &= \frac{1}{2}\|W - u\|_2^2 - \frac{1}{2}\|W\|_2^2 + J'_{\boldsymbol{\lambda}}(\beta^0; u), \end{aligned} \tag{30}$$

⁶The affine shift of $\partial J_{\boldsymbol{\lambda}}(\mathbf{p}_0)$ by $S_{\mathbf{p}_0}\Pi_{\mathbf{p}_0}\boldsymbol{\lambda}$ does not affect the dimension.

where $W \sim \mathcal{N}(0, \sigma^2 \mathbb{I}_p)$. The penalty $J'_\lambda(\beta^0; u)$ is separable (see Appendix A.2), and it suffices to consider the minimization problem on I_0 : $\|W_{I_0} - u_{I_0}\|^2/2 + J'_\lambda(u)$, with the smallest p_0 lambdas appearing in the penalty. On I_0 , the rejection procedure $\hat{\mathbf{b}}$ is determined by the restricted optimization and outside of I_0 , it rejects with probability one.

Furthermore, it is shown in [8] that minimization of

$$\frac{1}{2} \|\tilde{W} - \beta\|_2^2 + J_\lambda(\beta), \quad (31)$$

with $\tilde{W} \sim \mathcal{N}(\beta^0, \sigma^2 \mathbb{I}_p)$, and λ_i given by the BHq coefficients:

$$\lambda_i = \sigma \lambda_i^{BH(q)} = \sigma \Phi^{-1} \left(1 - \frac{iq}{2p} \right),$$

controls the FDR at level $q(p_0/p)$. The FDR control holds for any magnitudes of β^0 . Letting $|\beta_i^0| \rightarrow \infty \forall i \notin I_0$, the rejection procedure (31) separates into I_0 and I_0^c and is asymptotically equivalent to $\hat{\mathbf{b}}$, whence it follows that $\text{FDR}(\hat{\mathbf{b}}) \leq q(p_0/p)$.

In particular, $\text{FDR}(\hat{\beta}_n)$ is asymptotically controlled at level $q(p_0/p)$:

$$\text{FDR}(\hat{\beta}_n) \xrightarrow{n \rightarrow \infty} \text{FDR}(\hat{\mathbf{b}}) \leq q \frac{p_0}{p}.$$

Remark 4.8. Note that the covariance structure among the true signals does not matter for the asymptotic FDR control. To see this, consider a block diagonal covariance $C = \mathbb{I}_{I_0} \oplus \Sigma$, with identity matrix \mathbb{I}_{I_0} on I_0 and an arbitrary positive definite Σ on I_0^c . The optimization (5) separates into I_0 and I_0^c and the rejection procedure by $\hat{\mathbf{b}}_\Sigma = \mathbf{patt}_{\beta^0}(\hat{\mathbf{u}})$ does not depend on Σ , as it rejects with probability one on I_0^c . Hence $\text{FDR}(\hat{\beta}_n) \rightarrow \text{FDR}(\hat{\mathbf{b}}_\Sigma) = \text{FDR}(\hat{\mathbf{b}}) \leq q(p_0/p)$.

4.4 General penalty and pattern

The arguments so far, have predominantly relied on the fact that the SLOPE norm is a maximum over finitely many linear functions $J_\lambda(\beta) = \max\{\langle \Sigma \lambda, \beta \rangle : \Sigma \in \{\pm 1\}^n \cdot \mathcal{S}_p\}$. This is an example of a polyhedral gauge, which takes the form $f(x) := \max\{0, v_1^T x, \dots, v_p^T x\}$ for some $v_1, \dots, v_p \in \mathbb{R}^p$. Penalties given by polyhedral norms/gauges and the related notion of a model pattern is explored in [43]. In what follows, we show how our results on weak pattern convergence extend to more general penalties, including polyhedral norms.

To that end, let $f = \max\{f_1, \dots, f_N\}$ with convex and differentiable functions f_1, \dots, f_N . We have seen in (13) that $\partial f(x) = \text{con}\{\nabla f_i(x) : i \in I(x)\}$, where $I(x) := \{i : f_i(x) = f(x)\}$. This motivates the following definition of a pattern for a function f at a point x .

Definition 4.9. Given $f = \max\{f_1, \dots, f_N\}$, we define a *pattern* of f as an equivalence relation on \mathbb{R}^p given by $x \sim_f y \iff I(x) = I(y)$, where $I(x) = \{i : f_i(x) = f(x)\}$. We also say that $I(x) \subset \{1, \dots, N\}$ is the pattern of f at x . The set of all patterns \mathfrak{P}_f is the powerset of $\{1, \dots, N\}$.

Further, the directional derivative of f at x in direction u satisfies

$$f'(x; u) = \sup_{g \in \partial f(x)} \langle g, u \rangle = \max_{i \in I(x)} \langle \nabla f_i(x), u \rangle, \quad (32)$$

thus $\partial_u f'(x; u) = \text{con}\{\nabla f_i(x) : i \in I_x(u)\}$, where $I_x(u) := \{i \in I(x) : \langle \nabla f_i(x), u \rangle = f'(x; u)\}$. We call $I_x(u)$ the *limiting pattern* of u with respect to x .

As in Theorem 2.2, the minimizer

$$\widehat{\mathbf{u}}_{f,n} = \operatorname{argmin}_u u^T C_n u / 2 - u^T W_n + n^{1/2} [f(\beta^0 + u\sqrt{n}) - f(\beta^0)],$$

converges in distribution to

$$\widehat{\mathbf{u}}_f = \operatorname{argmin}_u u^T C u / 2 - u^T W + f'(\beta^0; u).$$

It is typical for a penalty f with non-linear f_i 's that the general pattern $I(\widehat{\mathbf{u}}_{f,n})$ converges weakly to $I(\widehat{\mathbf{u}}_f)$, but the SLOPE pattern $\mathbf{patt}(\widehat{\mathbf{u}}_{f,n})$ does not converge weakly to $\mathbf{patt}(\widehat{\mathbf{u}}_f)$, see Example A.1 in the Appendix.

To ensure convergence of the SLOPE pattern, we assume $f = \max\{f_1, \dots, f_N\}$, where $f_i(x) = \langle v_i, x \rangle + g(x)$, with $v_i \in \mathbb{R}^p$ and g smooth and convex. This includes all penalties of the type $J_\lambda(x) + \sum_{i=1}^p \alpha_i |x_i| + \gamma_i x_i^2$, with $\alpha_i, \gamma_i \in \mathbb{R} \forall i$, which encompasses for instance the LASSO, SLOPE, elastic net, quadratic loss, and any combination thereof. For such f , there is a one-to-one correspondence between patterns and sets of constant subdifferential: $\partial f(x) = \partial f(y) \iff I(x) = I(y)$, and a subdifferential at a pattern $\partial f(I)$ is well defined. Also, if $I(u) = I(v)$, then also $I_x(u) = I_x(v)$ and we can write $I_x(I)$ for the limiting pattern of $I \in \mathfrak{P}_f$ w.r.t. x . A direct consequence of (32) is that $\partial_u f'(x; u) = \partial f(I_x(u))$, generalizing (19). Moreover, for a sequence of functions $f_n = \max\{f_{1,n}, \dots, f_{N,n}\}$, under suitable conditions on f_n (such as $f_n = n^{1/2}f$), we have $\partial_u f_n(x + u/\sqrt{n}) \xrightarrow{d_H} \partial_u f'(x; u)$, generalizing (20). Crucially, linearity assumptions on f grant that $I(x + u/\sqrt{n})$ stabilizes at $I_x(u)$, which allows for the Hausdorff convergence of subdifferentials.

Theorem 4.10. *For every convex set $\mathcal{K} \subset \mathbb{R}^p$: $\mathbb{P}[\widehat{\mathbf{u}}_{f,n} \in \mathcal{K}] \longrightarrow \mathbb{P}[\widehat{\mathbf{u}}_f \in \mathcal{K}]$ as $n \rightarrow \infty$. In particular, $I(\widehat{\mathbf{u}}_{f,n})$ converges weakly to $I(\widehat{\mathbf{u}}_f)$ and $\mathbf{patt}(\widehat{\mathbf{u}}_{f,n})$ converges weakly to $\mathbf{patt}(\widehat{\mathbf{u}}_f)$.*

Proof. The sets $I^{-1}(\mathbf{p}_f)$, where $\mathbf{p}_f \in \mathfrak{P}_f$, are convex cones, which partition \mathbb{R}^p into sets of constant pattern. For $\varepsilon > 0$ and a convex \mathcal{K} , there exists $M > 0$ s.t. $\mathbb{P}[\widehat{\mathbf{u}}_{f,n} \in \mathcal{K} \setminus B_M(0)] < \varepsilon \forall n$. For each $\mathbf{p}_f \in \mathfrak{P}_f$ consider the bounded convex set $\mathcal{K}^{\mathbf{p}_f} = \mathcal{K} \cap I^{-1}(\mathbf{p}_f) \cap B_M(0)$. We get

$$\mathbb{P}[\widehat{\mathbf{u}}_{f,n} \in \mathcal{K}^{\mathbf{p}_f}] = \mathbb{P}[W_n \in C_n \mathcal{K}^{\mathbf{p}_f} + \partial f(\beta^0 + \mathcal{K}^{\mathbf{p}_f}/\sqrt{n})] \quad (33)$$

$$\rightarrow \mathbb{P}[W \in C \mathcal{K}^{\mathbf{p}_f} + \partial f'(\beta^0; \mathbf{p}_f)] = \mathbb{P}[\widehat{\mathbf{u}}_f \in \mathcal{K}^{\mathbf{p}_f}], \quad (34)$$

where the convergence follows by Lemma 3.4, since $I(\beta^0 + u/\sqrt{n})$ stabilizes for every $u \in \mathcal{K}^{\mathbf{p}_f}$ at the limiting pattern $I_{\beta^0}(\mathbf{p}_f)$ and $\partial f'(\beta^0; \mathbf{p}_f) = \partial f(I_{\beta^0}(\mathbf{p}_f))$.

$$\limsup_{n \rightarrow \infty} \mathbb{P}[\widehat{\mathbf{u}}_{f,n} \in \mathcal{K}] \leq \limsup_{n \rightarrow \infty} \sum_{\mathbf{p}_f \in \mathfrak{P}_f} \mathbb{P}[\widehat{\mathbf{u}}_{f,n} \in \mathcal{K}^{\mathbf{p}_f}] + \varepsilon = \sum_{\mathbf{p}_f \in \mathfrak{P}_f} \mathbb{P}[\widehat{\mathbf{u}}_f \in \mathcal{K}^{\mathbf{p}_f}] + \varepsilon \leq \mathbb{P}[\widehat{\mathbf{u}}_f \in \mathcal{K}] + \varepsilon,$$

$$\liminf_{n \rightarrow \infty} \mathbb{P}[\widehat{\mathbf{u}}_{f,n} \in \mathcal{K}] \geq \liminf_{n \rightarrow \infty} \sum_{\mathbf{p}_f \in \mathfrak{P}_f} \mathbb{P}[\widehat{\mathbf{u}}_{f,n} \in \mathcal{K}^{\mathbf{p}_f}] = \sum_{\mathbf{p}_f \in \mathfrak{P}_f} \mathbb{P}[\widehat{\mathbf{u}}_f \in \mathcal{K}^{\mathbf{p}_f}] \geq \mathbb{P}[\widehat{\mathbf{u}}_f \in \mathcal{K}],$$

which shows that $\widehat{\mathbf{u}}_{f,n}$ converges to $\widehat{\mathbf{u}}_f$ on all convex sets. The remaining assertions of the Theorem on weak pattern convergence follow from setting $\mathcal{K} = I^{-1}(\mathbf{p}_f)$ for some $\mathbf{p}_f \in \mathfrak{P}_f$ or $\mathcal{K} = \mathbf{patt}^{-1}(\mathbf{p})$ for $\mathbf{p} \in \mathfrak{P}$, respectively. \square

In this framework, the results on weak SLOPE pattern convergence in Section 4. and 5. extend to a broader family of regularizers f . We decide not to pursue this line of inquiry in this paper, and formulate and prove the subsequent results in the context of SLOPE.

5 General Loss function

Recall the assumption of the linear model $y = X\beta^0 + \varepsilon$, where $X = (X_1, \dots, X_n)^T$, $X_i \sim_{i.i.d} \mathbb{P}$, with covariance C , $\varepsilon_1, \dots, \varepsilon_n$ i.i.d. and $X \perp \varepsilon$. We can rewrite (2) as

$$\hat{\beta}_n = \operatorname{argmin}_{b \in \mathbb{R}^p} \frac{1}{n} \sum_{i=1}^n g(y_i, X_i, \beta) + J_{\lambda^n/n}(\beta), \quad (35)$$

where we have sofar assumed $g(y_i, X_i, \beta) = (y_i - X_i^T \beta)^2/2 = (\varepsilon_i - X_i^T(\beta - \beta^0))^2/2$. In this section, we show how the arguments from previous sections can be extended to a broader class of loss functions, including the Quantile and the Huber loss:

$$g(y_i, X_i, \beta) = \begin{cases} (y_i - X_i^T \beta)^2/2 & \text{Quadratic loss,} \\ H(y_i - X_i^T \beta) & \text{Huber loss,} \\ |y_i - X_i^T \beta|_\alpha & \text{Quantile loss,} \end{cases}$$

where $|\cdot|_\alpha : \mathbb{R} \rightarrow \mathbb{R}$, given by

$$|x|_\alpha := \begin{cases} (1 - \alpha)(-x) & x \leq 0, \\ \alpha x & x > 0, \end{cases}$$

and $\alpha \in (0, 1)$. Note that $|x|_{1/2} = \frac{1}{2}|x|$. For the Quadratic loss and the Huber loss, we shall assume that ε_i is centered with variance σ^2 . For the Quantile loss, we specify the assumptions in the example section. We apply the results from Pollard's paper [34], harnessing the notion of stochastic differentiability. We consider

$$\begin{aligned} G_n(\beta) &:= \frac{1}{n} \sum_{i=1}^n g(y_i, X_i, \beta), \\ G(\beta) &:= \mathbb{E}[g(\cdot, \beta)], \\ \nu_n g(\cdot, \beta) &:= n^{-1/2} \sum_{i=1}^n (g(\cdot, \beta) - G(\beta)). \end{aligned} \quad (36)$$

Observe that $G(\beta) = G_n(\beta) + n^{-1/2} \nu_n g(\cdot, \beta)$. Throughout, we shall express $g(\cdot, \beta)$ in the form

$$g(\cdot, \beta) = g(\cdot, \beta^0) + (\beta - \beta^0)^T \Delta(\cdot, \beta^0) + \|\beta - \beta^0\| r(\cdot, \beta), \quad (37)$$

where $\Delta(\cdot, \beta^0)$ is a random vector, which does not depend on β , and $r(\cdot, \beta)$ a rest term. Pollard [34] shows how the rather weak assumptions of twice differentiability of $G(\beta)$ together with a stochastic differentiability condition on $g(\cdot, \beta)$ are sufficient for asymptotic normality of the minimizer in (36), centered around β^0 .

These conditions lend themselves well to describe the asymptotics of the SLOPE estimator $\hat{\beta}_n$, which minimizes $G_n(\beta) + J_{\lambda^n/n}(\beta)$. We refer the reader to [34] for a more thorough discussion, but include the definition of stochastic differentiability.

Definition 5.1. A function $g(\cdot, \beta)$ of the form (37) is called stochastically differentiable at β^0 if for any shrinking sequence of neighborhoods U_n around β^0 :

$$\sup_{\beta \in U_n} \frac{|\nu_n r(\cdot, \beta)|}{1 + n^{1/2} \|\beta - \beta^0\|} \xrightarrow[n \rightarrow \infty]{p} 0.$$

Remark 5.2. Stochastic differentiability is a relatively weak condition on $g(\cdot, \beta)$. If $g(\cdot, \beta)$ is differentiable at β^0 , and $\Delta(\cdot, \beta^0) = \nabla_{\beta} g(\cdot, \beta^0)$ has a finite second moment $\mathbb{E}[\Delta(\cdot, \beta^0) \Delta(\cdot, \beta^0)^T] < \infty$, then $G(\beta)$ has a non-singular Hessian at β^0 and $g(\cdot, \beta)$ is stochastically differentiable at β^0 (Section 4. in [34]). However, differentiability of $g(\cdot, \beta)$ is not necessary for stochastic differentiability. More general ways of verifying stochastic differentiability, including the Huber loss or Median loss, are discussed in Section 5. [34].

The following Lemma is a direct corollary the proof of Theorem 2 in [34].

Lemma 5.3. Assume $G(\beta)$ has a non-singular Hessian $\tilde{C} = \nabla^2|_{\beta=\beta^0} G(\beta)$ at it's minimizing value β^0 and $g(\cdot, \beta)$ is stochastically differentiable. Then

$$n \left(G_n \left(\beta^0 + \frac{u}{\sqrt{n}} \right) - G_n(\beta^0) \right) = \frac{1}{2} u^T \tilde{C} u + u^T [\nu_n \Delta(\cdot, \beta^0) + R_n(u)],$$

where $\sup_{u \in K} \|R_n(u)\| \rightarrow 0$ in probability as $n \rightarrow \infty$, for any compact set K .

Proof. Abbreviate $\beta_n = \beta^0 + u/\sqrt{n}$. We have

$$G_n(\beta_n) - G_n(\beta^0) = G(\beta_n) - G(\beta^0) + n^{-1/2} \nu_n [g(\cdot, \beta_n) - g(\cdot, \beta^0)].$$

Taylor expansion of $G(\beta)$ around β^0 yields:

$$\begin{aligned} n(G(\beta_n) - G(\beta^0)) &= n[(\beta_n - \beta^0)^T \tilde{C} (\beta_n - \beta^0)/2 + o(\|\beta_n - \beta^0\|^2)] \\ &= u^T \tilde{C} u/2 + n \cdot o(\|u\|^2/n). \end{aligned}$$

At the same time

$$\begin{aligned} n \cdot n^{-1/2} \nu_n [g(\cdot, \beta_n) - g(\cdot, \beta^0)] &= n^{1/2} (\beta_n - \beta^0)^T \nu_n \Delta(\cdot, \beta^0) + n^{1/2} \|(\beta_n - \beta^0)\| \nu_n r(\cdot, \beta_n) \\ &= u^T \nu_n \Delta(\cdot, \beta^0) + \|u\| \nu_n r(\cdot, \beta^0 + u/\sqrt{n}). \end{aligned}$$

Combining both expressions yields the desired representation with a rest term

$$R_n(u) = u \cdot \nu_n r(\cdot, \beta^0 + u/\sqrt{n}) + u \cdot \frac{o(\|u\|^2/n)}{(\|u\|^2/n)}.$$

Stochastic differentiability guarantees that for any compact set K , $\sup_{u \in K} \|R_n(u)\| \rightarrow 0$ in probability. \square

The following statement unifies Theorem 2 in [34] and Theorem 2 in [16] into one result.

Theorem 5.4. Let $g(\cdot, \beta)$ be a convex loss function and $\lambda^n/\sqrt{n} \rightarrow \lambda \geq 0$. Suppose that

1. $G(\beta)$ has a non-singular Hessian $\tilde{C} = \nabla^2|_{\beta=\beta^0}G(\beta)$ at its minimizing value β^0 ,
2. $\Delta(\cdot, \beta^0)$ in (37) satisfies $\mathbb{E}(\Delta(\cdot, \beta^0)) = 0$ and $C_\Delta := \mathbb{E}(\Delta(\cdot, \beta^0)\Delta(\cdot, \beta^0)^T) < \infty$,
3. $g(\cdot, \beta)$ is stochastically differentiable at β^0 .

Then $\hat{\mathbf{u}}_n = \sqrt{n}(\hat{\beta}_n - \beta^0)$ converges in distribution to the random vector $\hat{\mathbf{u}} := \operatorname{argmin}_u V(u)$, where

$$V(u) = \frac{1}{2}u^T \tilde{C}u - u^T \tilde{W} + J'_\lambda(\beta^0; u), \quad (38)$$

and $\tilde{W} \sim \mathcal{N}(0, C_\Delta)$. Moreover, $\mathbf{patt}(\hat{\mathbf{u}}_n)$ converges in distribution to $\mathbf{patt}(\hat{\mathbf{u}})$, hence $\mathbf{patt}(\hat{\beta}_n)$ converges in distribution to $\mathbf{patt}_{\beta^0}(\hat{\mathbf{u}})$ and $FDR(\hat{\beta}_n) \rightarrow FDR(\mathbf{patt}_{\beta^0}(\hat{\mathbf{u}}))$.

Remark 5.5. In the case $\lambda = 0$, the minimum is attained at $u = \tilde{C}^{-1}\tilde{W} \sim \mathcal{N}(0, \tilde{C}^{-1}C_\Delta\tilde{C}^{-1})$, which is exactly the assertion of Theorem 2 in [34].

Proof. By Lemma 5.3:

$$\begin{aligned} \sqrt{n}(\hat{\beta}_n - \beta^0) &= \operatorname{argmin}_u G_n \left(\beta^0 + \frac{u}{\sqrt{n}} \right) + J_{\lambda^n/n} \left(\beta^0 + \frac{u}{\sqrt{n}} \right) \\ &= \operatorname{argmin}_u n \left(G_n \left(\beta^0 + \frac{u}{\sqrt{n}} \right) - G_n(\beta^0) \right) + J_{\lambda^n} \left(\beta^0 + \frac{u}{\sqrt{n}} \right) - J_{\lambda^n}(\beta^0) \\ &= \operatorname{argmin}_u \frac{1}{2}u^T \tilde{C}u + u^T [\nu_n \Delta(\cdot, \beta^0) + R_n(u)] + J_{\lambda^n} \left(\beta^0 + \frac{u}{\sqrt{n}} \right) - J_{\lambda^n}(\beta^0), \end{aligned} \quad (39)$$

For every fixed u , the convex objective in (39) converges in distribution and it follows that

$$\sqrt{n}(\hat{\beta}_n - \beta^0) \xrightarrow{d} \operatorname{argmin}_u \left\{ \frac{1}{2}u^T \tilde{C}u + u^T \tilde{W} + J'_\lambda(\beta^0; u) \right\} = \hat{\mathbf{u}}.$$

This shows weak convergence, which is not sufficient to conclude pattern convergence. The key to pattern convergence is to absorb the rest term $R_n(u)$ into the $O_p(1)$ stochastic term $W_n := \nu_n \Delta(\cdot, \beta^0)$. We abbreviate the objective in (39) by $F_n(W_n + R_n(u), u)$. The goal is to get rid of the dependency on u in the first argument of F_n . To that end, let us fix $\varepsilon > 0$. By uniform tightness of $\hat{\mathbf{u}}_n$, there exists a compact set K such that $\mathbb{P}[\hat{\mathbf{u}}_n \in K^\circ] > 1 - \varepsilon$ for all $n \in \mathbb{N}$, where K° denotes the interior of K . Moreover, by Lemma 5.3, there exists a sequence $\delta_M \downarrow 0$ as $M \rightarrow \infty$, such that $\mathbb{P}[\sup_{u \in K} \|R_n(u)\| < \delta_M] > 1 - \varepsilon$ for every $n \geq M$. Let $\mathfrak{p} \in \mathfrak{P}$ be a fixed pattern and consider $p_n := \mathbb{P}[\mathbf{patt}(\hat{\mathbf{u}}_n) = \mathfrak{p}]$.

$$\begin{aligned} p_n &= \mathbb{P} \left[\mathbf{patt}(\operatorname{argmin}_{u \in \mathbb{R}^p} F_n(W_n + R_n(u), u)) = \mathfrak{p} \right] \\ &\leq \mathbb{P} \left[\mathbf{patt}(\operatorname{argmin}_{u \in K^\circ} F_n(W_n + R_n(u), u)) = \mathfrak{p}, \hat{\mathbf{u}}_n \in K^\circ, \sup_{u \in K} \|R_n(u)\| < \delta_M \right] + 2\varepsilon \\ &\leq \mathbb{P} \left[\exists \delta \in B_{\delta_M}(0) : \mathbf{patt}(\operatorname{argmin}_{u \in K^\circ} F_n(W_n + \delta, u)) = \mathfrak{p}, \hat{\mathbf{u}}_n \in K^\circ \right] + 2\varepsilon \\ &\leq \mathbb{P} \left[\exists \delta \in B_{\delta_M}(0) : W_n + \delta \in \tilde{C}D^{\mathfrak{p}} + \partial J_{\lambda^n/\sqrt{n}}(\mathfrak{p}_0) \right] + 2\varepsilon \\ &\leq \mathbb{P} \left[W_n \in \left(\tilde{C}D^{\mathfrak{p}} + \partial J_{\lambda^n/\sqrt{n}}(\mathfrak{p}_0) \right)^{\delta_M} \right] + 2\varepsilon, \end{aligned} \quad (40)$$

where in (40) we have used optimality condition $0 \in \partial F_n(W_n + \delta, u)$ and denoted by $D^{\mathfrak{p}}$ the intersection of $\mathbf{patt}^{-1}(\mathfrak{p})$ with K and $\mathfrak{p}_0 = \mathbf{patt}_{\beta^0}(\mathfrak{p})$. Setting $B_n := \tilde{C}D^{\mathfrak{p}} + \partial J_{\lambda^n/\sqrt{n}}(\mathfrak{p}_0)$, we have $\tilde{C}D^{\mathfrak{p}} + \partial J_{\lambda/\sqrt{n}}(\mathfrak{p}_0) \xrightarrow{d_H} \tilde{C}D^{\mathfrak{p}} + \partial J_{\lambda}(\mathfrak{p}_0) =: B$ converging in the Hausdorff distance. Consequently, also $B_n^{\delta_M} \xrightarrow{d_H} B^{\delta_M}$ and applying Lemma 3.4:

$$\limsup_{n \rightarrow \infty} p_n \leq \limsup_{n \rightarrow \infty} \mathbb{P}[W_n \in B_n^{\delta_M}] + 2\varepsilon = \mathbb{P}[\tilde{W} \in B^{\delta_M}] + 2\varepsilon \leq \mathbb{P}[\tilde{W} \in B] + 3\varepsilon,$$

where in the last inequality we have taken δ_M small enough and used (17). Similarly, we can bound p_n from below

$$\begin{aligned} p_n &\geq \mathbb{P} \left[\mathbf{patt}(\operatorname{argmin}_{u \in K^\circ} F_n(W_n + R_n(u), u)) = \mathfrak{p}, \hat{\mathbf{u}}_n \in K^\circ, \sup_{u \in K} \|R_n(u)\| < \delta_M \right] \\ &\geq \mathbb{P} \left[\forall \delta \in B_{\delta_M}(0) : \mathbf{patt}(\operatorname{argmin}_{u \in K^\circ} F_n(W_n + \delta, u)) = \mathfrak{p}, \hat{\mathbf{u}}_n \in K^\circ, \sup_{u \in K} \|R_n(u)\| < \delta_M \right] \\ &\geq \mathbb{P} \left[\forall \delta \in B_{\delta_M}(0) : W_n + \delta \in \tilde{C}D^{\mathfrak{p}} + \partial J_{\lambda^n/\sqrt{n}}(\mathfrak{p}_0) \right] - 2\varepsilon \\ &= \mathbb{P} \left[W_n \in \left(\tilde{C}D^{\mathfrak{p}} + \partial J_{\lambda^n/\sqrt{n}}(\mathfrak{p}_0) \right)^{-\delta_M} \right] - 2\varepsilon, \end{aligned}$$

Since $B_n^{-\delta_M} \xrightarrow{d_H} B^{-\delta_M}$, by the same argument applying Lemma 3.4 and (17):

$$\liminf_{n \rightarrow \infty} p_n \geq \liminf_{n \rightarrow \infty} \mathbb{P}[W_n \in B_n^{-\delta_M}] - 2\varepsilon = \mathbb{P}[\tilde{W} \in B^{-\delta_M}] - 2\varepsilon \geq \mathbb{P}[\tilde{W} \in B] - 3\varepsilon,$$

As a result $\mathbf{patt}(\hat{\mathbf{u}}_n)$ converges in distribution to $\mathbf{patt}(\hat{\mathbf{u}})$. The remaining assertions are direct consequences of the weak pattern convergence of $\hat{\mathbf{u}}_n$. \square

Example 5.6. (Quadratic loss) Set $g(y_i, X_i, \beta) = (\varepsilon_i - t(\beta))^2/2$, with $t(\beta) = X_i^T(\beta - \beta^0)$. We have $\nabla_\beta g(y_i, X_i, \beta) = X_i(t(\beta) - \varepsilon_i)$, hence the first order term in (37) reads $\Delta(\cdot, \beta^0) = -\varepsilon_i X_i$ and by independence $C_\Delta = \mathbb{E}(\Delta(\cdot, \beta^0)\Delta(\cdot, \beta^0)^T) = \mathbb{E}[\varepsilon_i^2] \mathbb{E}[X_i X_i^T] = \sigma^2 C$. Moreover, $\nabla_\beta^2 g(y_i, X_i, \beta) = X_i X_i^T$, which gives $\tilde{C} = \nabla^2|_{\beta=\beta^0} G(\beta) = \mathbb{E}[\nabla^2|_{\beta=\beta^0} g(y_i, X_i, \beta)] = C$. This coincides with Theorem 2.2.

Example 5.7. (Huber loss [21]) Consider $g(y_i, X_i, \beta) = H(\varepsilon_i - t(\beta))$, with $t(\beta) := X_i^T(\beta - \beta^0)$ and the Huber loss $H : \mathbb{R} \rightarrow \mathbb{R}$, given by

$$H(x) := k \left(-x - \frac{k}{2} \right) \mathbf{1}_{\{x < -k\}} + \frac{1}{2} x^2 \mathbf{1}_{\{|x| \leq k\}} + k \left(x - \frac{k}{2} \right) \mathbf{1}_{\{x > k\}}.$$

We consider

$$\begin{aligned} D(x, t) &:= \frac{\partial}{\partial t} H(x - t) = k \mathbf{1}_{\{x-t < -k\}} - (x-t) \mathbf{1}_{\{|x-t| \leq k\}} - k \mathbf{1}_{\{x-t > k\}} \\ \frac{\partial^2}{\partial t^2} H(x-t) &= \mathbf{1}_{\{|x-t| \leq k\}} \end{aligned}$$

By chain rule $\nabla_\beta g(y_i, X_i, \beta) = X_i D(\varepsilon_i, t(\beta))$, hence the $\Delta(\cdot, \beta^0)$ term in expansion (37) reads $\Delta(\varepsilon_i, X_i, \beta^0) = X_i D(\varepsilon_i, 0)$. The covariance matrix of $\Delta(\cdot, \beta^0)$ equals $C_\Delta = \delta C$, where C is the covariance matrix of X_i and

$$\delta = \mathbb{E}[D(\varepsilon_i, 0)^2] = \mathbb{E}[\varepsilon_i^2 \mathbf{1}_{|\varepsilon_i| < k}] + k^2 \mathbb{P}[|\varepsilon_i| > k]. \quad (41)$$

Moreover, $\nabla_{\beta}^2 g(y_i, X_i, \beta) = X_i X_i^T \mathbf{1}_{\{|\varepsilon_i - t(\beta)| \leq k\}}$, and we get $\tilde{C} = \nabla^2|_{\beta=\beta^0} G(\beta) = \gamma C$, where $\gamma = \mathbb{P}[|\varepsilon_i| < k]$. In particular, letting $k \rightarrow \infty$, we get $\delta \rightarrow \mathbb{E}[\varepsilon_i^2] = \sigma^2$ and $\gamma \rightarrow 1$, and recover the standard quadratic case with $\tilde{C} = C$ and $C_{\Delta} = \sigma^2 C$. Further, for $C = \mathbb{I}_p$ we have $\tilde{C} = \gamma \mathbb{I}_p$, $\tilde{W} \sim \mathcal{N}(0, \delta \mathbb{I}_p)$ and for $\lambda_i := \sqrt{\delta} \lambda_i^{BH(q)} = \sqrt{\delta} \Phi^{-1}(1 - iq/2p)$, $\text{FDR}(\hat{\beta}_n)$ is asymptotically controlled at level $q(|I_0|/p)$. (This can be verified by substituting $\tilde{u} = \sqrt{\gamma}u$ in (38) and using Example 4.7; see Appendix A.3) Note that λ_i do not depend on γ , but only on δ . Because $\delta = \delta_k \leq \sigma^2$ for any k , it suffices to penalize with smaller penalties to obtain FDR control for the Huber loss than for the standard Quadratic loss. In particular, $\delta \rightarrow 0$ as $k \rightarrow 0$, so that λ_i are significantly smaller for small k . Similar to Example 4.7, the FDR control holds more generally for a block diagonal $C = \mathbb{I}_{I_0} \oplus \Sigma$.

Quantile regression [23] is a popular method where one assumes that there is a linear relation between the fixed effect and a fixed quantile of y . It has been studied earlier with L_1 regularization in [30].

Example 5.8. (Quantile) For the Quantile regression we drop the assumption of finite variance and zero mean of the noise ε . Instead we assume that the CDF of ε has a continuous density f and CDF F , which satisfies $F^{-1}(\alpha) = 0$, and $f(0) > 0$. Consider $g(y_i, X_i, \beta) = |\varepsilon_i - t(\beta)|_{\alpha}$, with $t(\beta) := X_i^T(\beta - \beta^0)$. The loss function $|\cdot|_{\alpha}$ is everywhere differentiable except at 0. The weak derivative is given by $|x|'_{\alpha} = -(1 - \alpha)\mathbf{1}_{\{x \leq 0\}} + \alpha\mathbf{1}_{\{x > 0\}}$. Note that $t \mapsto \mathbb{E}[|\varepsilon_i - t|_{\alpha}]$ attains its minimum at the solution to $0 = -(1 - \alpha)\mathbb{P}(\varepsilon_i \leq t) + \alpha\mathbb{P}(\varepsilon_i > t)$, which is the α quantile of ε_i . Now since this minimum is obtained by setting $t = 0$ it is clear that G has a minimum value at β^0 .

By chain rule $\nabla_{\beta} g(y_i, X_i, \beta) = -X_i |\varepsilon_i - t(\beta)|'_{\alpha} = X_i((1 - \alpha)\mathbf{1}_{\{\varepsilon_i \leq t(\beta)\}} - \alpha\mathbf{1}_{\{\varepsilon_i > t(\beta)\}})$ and the $\Delta(\cdot, \beta^0)$ term in expansion (37) reads $\Delta(\varepsilon_i, X_i, \beta^0) = -X_i |\varepsilon_i|_{\alpha}$. Consequently, the covariance matrix of $\Delta(\cdot, \beta^0)$ equals $C_{\Delta} = \delta C$, where

$$\delta = \mathbb{E}[(|\varepsilon_i|'_{\alpha})^2] = (1 - \alpha)^2 + \alpha^2$$

Moreover,

$$\begin{aligned} \nabla_{\beta} \mathbb{E}[g(y_i, X_i, \beta)] &= \mathbb{E}[\nabla_{\beta} g(y_i, X_i, \beta)] \\ &= \mathbb{E}\left[X_i \mathbb{E}[(1 - \alpha)\mathbf{1}_{\{\varepsilon_i \leq t(\beta)\}} - \alpha\mathbf{1}_{\{\varepsilon_i > t(\beta)\}} | X_i]\right] \\ &= \mathbb{E}\left[X_i((1 - \alpha)\mathbb{P}(\varepsilon_i \leq t(\beta) | X_i) - \alpha(1 - \mathbb{P}(\varepsilon_i \leq t(\beta) | X_i)))\right] \\ &= \mathbb{E}\left[X_i(\mathbb{P}(\varepsilon_i \leq t(\beta) | X_i) - \alpha)\right], \end{aligned}$$

Thus $\tilde{C} = \nabla^2|_{\beta=\beta^0} G(\beta) = f(0)C$. Also, for $C = \mathbb{I}_p$, and $\lambda_i := \sqrt{\delta} \Phi^{-1}(1 - iq/2p)$, $\text{FDR}(\hat{\beta}_n)$ is asymptotically controlled at level $q(|I_0|/p)$. For the Median loss $\alpha = 1/2$, we obtain $\lambda_i = (1/\sqrt{2}) \Phi^{-1}(1 - iq/2p)$. Again, the result remains valid for a block diagonal $C = \mathbb{I}_{I_0} \oplus \Sigma$.

6 Discussion

In this article we proposed a general theoretical framework for the asymptotic analysis of the pattern recovery and FDR control by SLOPE and its extensions. We mainly concentrated

on the extensions to robust loss of functions, which are very important from the practical perspective. However, our proof techniques can be generalized to obtain similar results for applications of SLOPE to other important Generalized Linear Models, like the logistic or multinomial regressions, or the Gaussian Graphical models. Another point for future interest is the analysis of algorithms combining the SLOPE penalty with L_1 or L_2 penalties. Our preliminary investigation suggests that combining the SLOPE and L_2 penalty allows to obtain FDR control under a wider set of scenarios than for SLOPE itself and we consider the investigation of the properties of this method as a very promising direction of research. Finally, our techniques can be extended for the analysis of adaptive versions of SLOPE. In [22] SLOBE, the first version of the adaptive SLOPE, was constructed. It was illustrated that SLOBE can control FDR when predictors are strongly correlated. We expect that our techniques can be useful for the analysis of SLOBE properties as well as for construction of new adaptive versions of SLOPE, which apart from FDR control will also enhance the pattern recovery properties of SLOPE.

The asymptotic results presented in this paper are concerned with the classical asymptotics, where the model dimension p is fixed and n diverges to infinity. Our analysis shows that even this classical setup deriving the results on the pattern convergence requires the development of new tools and substantially more care as compared to the analysis of the convergence of the vector of parameter estimates. We believe that the our analysis framework based on the set Hausdorff distance can be extended for the analysis of the SLOPE properties in the high dimensional setup and consider our work as an important first step in this direction.

7 Acknowledgments

The authors acknowledge the support of the Swedish Research Council, grant no. 2020-05081. We would like to thank Elvezio Ronchetti and Michał Kos for the discussions on the robust versions of SLOPE and further Piotr Graczyk, Bartosz Kołodziejek, Tomasz Skalski, Patrick Tardivel and Ulrike Schneider for their helpful comments and discussions.

References

- [1] F. Abramovich. Classification by sparse additive models. *arXiv preprint, arXiv:2212.01792*, 2022.
- [2] F. Abramovich and V. Grinshtein. High-dimensional classification by sparse logistic regression. *IEEE Transactions on Information Theory*, PP, 06 2017.
- [3] F. Abramovich, V. Grinshtein, and T. Levy. Multiclass classification by sparse multinomial logistic regression. *IEEE Transactions on Information Theory*, 65(5):3068–3079, 2018.
- [4] Per Kragh Andersen and Richard D Gill. Cox’s regression model for counting processes: a large sample study. *The annals of statistics*, pages 1100–1120, 1982.
- [5] R. E. Barlow and H. D. Brunk. The isotonic regression problem and its dual. *Journal of the American Statistical Association*, 67(337):140–147, 1972.
- [6] P.C. Bellec, G. Lecué, and A.B. Tsybakov. Slope meets lasso: improved oracle bounds and optimality. *Annals of Statistics*, 46(6B):3603–3642, 2018.
- [7] Małgorzata Bogdan, Xavier Dupuis, Piotr Graczyk, Bartosz Kołodziejek, Tomasz Skalski, Patrick Tardivel, and Maciej Wilczyński. Pattern recovery by slope. *arXiv preprint arXiv:2203.12086*, 2022.
- [8] Małgorzata Bogdan, Ewout Van Den Berg, Chiara Sabatti, Weijie Su, and Emmanuel J Candès. Slope—adaptive variable selection via convex optimization. *The annals of applied statistics*, 9(3):1103, 2015.
- [9] Małgorzata Bogdan, Ewout Van Den Berg, Weijie Su, and Emmanuel J Candès. Statistical estimation and testing via the sorted l1 norm. *arXiv preprint arXiv:1310.1969*, 2013.
- [10] Howard D Bondell and Brian J Reich. Simultaneous regression shrinkage, variable selection, and supervised clustering of predictors with oscar. *Biometrics*, 64(1):115–123, 2008.
- [11] D. Brzyski, A. Gossmann, W. Su, and M. Bogdan. Group slope - adaptive selection of groups of predictors. *Journal of the American Statistical Association*, 2018. DOI: 10.1080/01621459.2017.1411269.
- [12] Zhiqi Bu, Jason Klusowski, Cynthia Rush, and Weijie Su. Algorithmic analysis and statistical estimation of slope via approximate message passing. *Advances in Neural Information Processing Systems*, 32, 2019.
- [13] Zhiqi Bu, Jason Klusowski, Cynthia Rush, and Weijie Su. Characterizing the slope trade-off: A variational perspective and the donoho-tanner limit. *Annals of Statistics*, to appear, available at arXiv:2105.13302, 2023.

- [14] Zhiqi Bu, Jason M Klusowski, Cynthia Rush, and Weijie J Su. Algorithmic analysis and statistical estimation of slope via approximate message passing. *IEEE Transactions on Information Theory*, 67(1):506–537, 2020.
- [15] Mario Figueiredo and Robert Nowak. Ordered weighted l_1 regularized regression with strongly correlated covariates: Theoretical aspects. In *Artificial Intelligence and Statistics*, pages 930–938. PMLR, 2016.
- [16] Wenjiang Fu and Keith Knight. Asymptotics for lasso-type estimators. *The Annals of statistics*, 28(5):1356–1378, 2000.
- [17] I. Gannaz. Robust estimation and wavelet thresholding in partial linear models. *Technical report, University Joseph Fourier, Grenoble, France*, 2006.
- [18] I. Goodfellow, Y. Bengio, and A. Courville. *Deep Learning*. MIT Press, 2016.
- [19] Arthur E. Hoerl and Robert W. Kennard. Ridge regression: Biased estimation for nonorthogonal problems. *Technometrics*, 12(1):55–67, 1970.
- [20] Peter J. Huber. Robust Estimation of a Location Parameter. *The Annals of Mathematical Statistics*, 35(1):73 – 101, 1964.
- [21] Peter J. Huber. The behavior of maximum likelihood estimates under nonstandard conditions. In *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability*, volume 1: Statistics, pages 221–233. University of California Press, Berkeley, CA, 1967.
- [22] Wei Jiang, Małgorzata Bogdan, Julie Josse, Szymon Majewski, Błażej Miasojedow, Veronika Rockova, and TraumaBase Group. Adaptive Bayesian SLOPE: Model Selection With Incomplete Data. *J. Comput. Graph. Statist.*, 31(1):113–137, 2022.
- [23] Roger Koenker and Gilbert Bassett Jr. Regression quantiles. *Econometrica: journal of the Econometric Society*, pages 33–50, 1978.
- [24] Michał Kos. Identification of statistically important predictors in high-dimensional data. theoretical properties and practical applications. *PhD Thesis, University of Wrocław*, 2019.
- [25] Michał Kos and Małgorzata Bogdan. On the asymptotic properties of slope. *Sankhya A*, 82(2):499–532, 2020.
- [26] Philipp J. Kremer, Damian Brzyski, Małgorzata Bogdan, and Sandra Paterlini. Sparse index clones via the sorted l_1 -norm. *Quantitative Finance*, 22(2):349–366, 2022.
- [27] Philipp J Kremer, Sangkyun Lee, Małgorzata Bogdan, and Sandra Paterlini. Sparse portfolio selection via the sorted l^1 -norm. *Journal of Banking & Finance*, 110:105687, 2020.
- [28] Johan Larsson, Małgorzata Bogdan, and Jonas Wallin. The strong screening rule for slope. *Advances in neural information processing systems*, 33:14592–14603, 2020.

- [29] Johan Larsson, Quentin Klopfenstein, Mathurin Massias, and Jonas Wallin. Coordinate descent for slope, 2022.
- [30] Youjuan Li and Ji Zhu. L 1-norm quantile regression. *Journal of Computational and Graphical Statistics*, 17(1):163–185, 2008.
- [31] Aleksandra Maj-Kańska, Piotr Pokarowski, and Agnieszka Prochenka. Delete or merge regressors for linear model selection. *Electron. J. Stat.*, 9(2):1749–1778, 2015.
- [32] L. McCann and R. E. Welsch. Robust variable selection using least angle regression and elemental set sampling. *Computational Statistics and Data Analysis*, 52(1):249–257, 2007.
- [33] S. Nowakowski, P. Pokarowski, and W. Rejchel. Group lasso merger for sparse prediction with high-dimensional categorical data. *arXiv:2112.11114*, 2021.
- [34] David Pollard. New ways to prove central limit theorems. *Econometric Theory*, 1(3):295–313, 1985.
- [35] David Pollard. Asymptotics for least absolute deviation regression estimators. *Econometric Theory*, 7(2):186–199, 1991.
- [36] R.T. Rockafellar, M. Wets, and R.J.B. Wets. *Variational Analysis*. Grundlehren der mathematischen Wissenschaften. Springer Berlin Heidelberg, 2009.
- [37] A. Maleki S. Wang, H. Weng. Does slope outperform bridge regression? *Information and Inference: A Journal of the IMA*, 11(1):1–54, 2022.
- [38] Ulrike Schneider and Patrick Tardivel. The geometry of uniqueness, sparsity and clustering in penalized estimation. *Journal of Machine Learning Research*, 23(331):1–36, 2022.
- [39] Yiyuan She and Art B. Owen. Outlier detection using nonconvex penalized regression. *Journal of the American Statistical Association*, 106(494):626–639, 2011.
- [40] T. Skalski, P. Graczyk, B. Kołodziejek, and M. Wilczyński. Pattern recovery and signal denoising by slope when the design matrix is orthogonal. *Probability and Mathematical Statistics*, 42:283–302, 2022.
- [41] Benjamin G. Stokell, Rajen D. Shah, and Ryan J. Tibshirani. Modelling high-dimensional categorical data using nonconvex fusion penalties. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 83(3):579–611, 2021.
- [42] W. Su and E. Candès. Slope is adaptive to unknown sparsity and asymptotically minimax. *Ann. Statist.*, 44(3):1038–1068, 06 2016.
- [43] Patrick J C Tardivel, Tomasz Skalski, Piotr Graczyk, and Ulrike Schneider. The Geometry of Model Recovery by Penalized and Thresholded Estimators. working paper or preprint, June 2021.

- [44] Patrick JC Tardivel, Rémi Servien, and Didier Concordet. Simple expressions of the lasso and slope estimators in low-dimension. *Statistics*, 54(2):340–352, 2020.
- [45] Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B (Methodological)*, 58(1):267–288, 1996.
- [46] A. Virouleau, A. Guilloux, S. Gaïffas, and M. Bogdan. High-dimensional robust regression and outliers detection with slope. *arXiv preprint, arXiv:1712.02640*, 2017.

A Appendix

A.1 Weak pattern convergence is not trivial

Consider the penalty $f(x) = \max\{x_1^2, x_2\}$ on \mathbb{R}^2 , and let $\widehat{\beta}_f = \operatorname{argmin}_{\beta} G_n(\beta) + f_n(\beta)$, with $f_n = n^{1/2}f$ and G_n as in (36) for the standard quadratic loss. Replicating Theorem 2.2,

$$\begin{aligned} \widehat{\mathbf{u}}_{f,n} &= \sqrt{n}(\widehat{\beta}_f - \beta^0) = \operatorname{argmin} u^T C_n u / 2 - u^T W_n + n^{1/2}[f(\beta^0 + u/\sqrt{n}) - f(\beta^0)] \\ &\xrightarrow{d} \operatorname{argmin} u^T C u / 2 - u^T W + f'(\beta^0; u) =: \widehat{\mathbf{u}}_f. \end{aligned}$$

For $\beta^0 = 0$, we have $n^{1/2}[f(\beta^0 + u/\sqrt{n}) - f(\beta^0)] = \max\{n^{-1/2}u_1^2, u_2\} =: g_n(u)$, and on the half line $\mathcal{K} = \{u_1 > 0, u_2 = 0\}$; the subdifferential $\partial g_n(u) = (2u_1/\sqrt{n}, 0)^T$ is zero-dimensional. We obtain

$$\mathbb{P}[\widehat{\mathbf{u}}_{f,n} \in \mathcal{K}] = \mathbb{P}\left[W_n \in \left\{C_n \begin{pmatrix} u_1 \\ 0 \end{pmatrix} + \begin{pmatrix} 2u_1/\sqrt{n} \\ 0 \end{pmatrix} : u_1 > 0\right\}\right] = 0 \quad \forall n,$$

provided W_n is absolutely continuous w.r.t. the Lebesgue measure. Furthermore, from (32) we get $f'(0; u) = \max\{\langle(0, 0), (u_1, u_2)\rangle, \langle(0, 1), (u_1, u_2)\rangle\} = \max\{0, u_2\}$, hence on \mathcal{K} the subdifferential $\partial f'(0; u) = \operatorname{con}\{(0, 0)^T, (0, 1)^T\}$ is one-dimensional. We get

$$\mathbb{P}[\widehat{\mathbf{u}}_f \in \mathcal{K}] = \mathbb{P}\left[W \in \left\{C \begin{pmatrix} u_1 \\ 0 \end{pmatrix} + \partial f'(0; u) : u_1 > 0\right\}\right] > 0,$$

since $C_{11} > 0$. In particular, the SLOPE pattern $\mathbf{patt}(\widehat{\mathbf{u}}_{f,n})$ does not converge weakly to $\mathbf{patt}(\widehat{\mathbf{u}}_f)$, despite the weak convergence of $\widehat{\mathbf{u}}_{f,n}$ to $\widehat{\mathbf{u}}_f$.

Observe that $\widehat{\mathbf{u}}_{f,n}$ puts positive mass on the parabola $\{u_2 = n^{-1/2}u_1\}$, where $\partial g_n(u)$ is one-dimensional, whereas $\widehat{\mathbf{u}}_f$ puts positive mass on the tangential space of the parabola at 0 given by $\{u_2 = 0\}$. More precisely, the Lebesgue decompositions of $\widehat{\mathbf{u}}_{f,n}$ and $\widehat{\mathbf{u}}_f$ w.r.t. the Lebesgue measure yield different singular sets; the (flattening) parabola and the x-axis respectively. This gives some intuition for why linearity of the functions in the penalty $f = \max\{f_1, \dots, f_N\}$ is essential for convergence on convex cones. By applying the Portmanteau Lemma, we get weak convergence of $I(\widehat{\mathbf{u}}_{f,n})$ to $I(\widehat{\mathbf{u}}_f)$ w.r.t. the general pattern as described in 4.4.

A.2 Proximal operator

Similar to the SLOPE optimization problem, the minimization of (5) and (38) can be solved using proximal methods. The proximal operator of $J'_{\lambda}(\beta^0; u)$ is given by:

$$\operatorname{prox}_{J'_{\lambda, \beta^0}}(y) := \operatorname{argmin}_{u \in \mathbb{R}^p} (1/2)\|u - y\|_2^2 + J'_{\lambda}(\beta^0; u)$$

Let $\mathcal{I}(\beta^0) = \{I_0, I_1, \dots, I_m\}$ be the partition of β^0 into the clusters of the same magnitude. The directional SLOPE derivative $J'_{\lambda}(\beta^0; u)$ is separable:

$$J'_{\lambda}(\beta^0; u) = J_{\lambda}^{I_0}(u) + J_{\lambda, \beta^0}^{I_1}(u) + \dots + J_{\lambda, \beta^0}^{I_m}(u),$$

with

$$J_{\lambda}^{I_0}(u) = \sum_{i \in I_0} \lambda_{\pi(i)} |u_i|,$$

$$J_{\lambda, \beta^0}^{I_j}(u) = \sum_{i \in I_j} \lambda_{\pi(i)} u_i \operatorname{sgn}(\beta_i^0),$$

where the permutation π in $J_{\lambda}'(\beta^0; u)$ sorts the limiting pattern of u w.r.t. β^0 , i.e; $|\mathbf{p}_0|_{\pi^{-1}(1)} \geq \dots \geq |\mathbf{p}_0|_{\pi^{-1}(p)}$, with $\mathbf{p}_0 = \mathbf{patt}_{\beta^0}(u)$.

Hence

$$\operatorname{prox}_{J_{\lambda, \beta^0}}(y) = \operatorname{prox}_{J_{\lambda, \beta^0}^{I_0}}(y) \oplus \operatorname{prox}_{J_{\lambda, \beta^0}^{I_1}}(y) \oplus \dots \oplus \operatorname{prox}_{J_{\lambda, \beta^0}^{I_m}}(y)$$

Since we can treat each cluster separately, we can w.l.o.g. assume that β^0 consists of one cluster only. There are only two possible cases:

In the first case, $\beta^0 = 0$ and the proximal operator is described in [8]:

$$\begin{aligned} \operatorname{prox}_{J_{\lambda}}(y) &= \operatorname{argmin}_{u \in \mathbb{R}^p} (1/2) \|u - y\|_2^2 + J_{\lambda}(u) \\ &= S_y \Pi_y \operatorname{argmin}_{\tilde{u}_1 \geq \dots \geq \tilde{u}_p \geq 0} (1/2) \|\tilde{u} - |y|_{(\cdot)}\|_2^2 + \sum_{i=1}^p \lambda_i \tilde{u}_i, \end{aligned} \quad (42)$$

where $|y|_{(\cdot)} = \Pi_y^T S_y y$ arises by sorting the absolute values of y . (See Proposition 2.2 in [8] and notation in the section Subdifferential and Pattern.)

In the second case, β^0 consists of a single non zero cluster. In this case the penalty becomes $J_{\lambda}'(\beta^0; u) = \sum_{i=1}^p \lambda_{\pi(i)} (S_{\beta^0} u)_i = \sum_{i=1}^p \lambda_i (S_{\beta^0} u)_{\pi^{-1}(i)}$, where $(S_{\beta^0} u)_{\pi^{-1}(1)} \geq \dots \geq (S_{\beta^0} u)_{\pi^{-1}(p)}$. In particular, $J_{\lambda, \beta^0}(S_{\beta^0} u) = \sum_{i=1}^p \lambda_i u_{\pi^{-1}(i)} = \sum_{i=1}^p \lambda_i u_{(i)}$, with $u_{(1)} \geq \dots \geq u_{(p)}$.

$$\begin{aligned} \operatorname{prox}_{J_{\lambda, \beta^0}}(y) &= \operatorname{argmin}_{u \in \mathbb{R}^p} (1/2) \|u - y\|_2^2 + J_{\lambda}'(\beta^0; u) \\ &= S_{\beta^0} \operatorname{argmin}_{\tilde{u} \in \mathbb{R}^p} (1/2) \|\tilde{u} - S_{\beta^0} y\|_2^2 + \sum_{i=1}^p \lambda_i \tilde{u}_{(i)} \\ &= S_{\beta^0} \Pi \operatorname{argmin}_{\tilde{u}_1 \geq \dots \geq \tilde{u}_p} (1/2) \|\tilde{u} - (S_{\beta^0} y)_{(\cdot)}\|_2^2 + \sum_{i=1}^p \lambda_i \tilde{u}_i, \end{aligned} \quad (43)$$

where $(S_{\beta^0} y)_{(\cdot)} = \Pi^T (S_{\beta^0} y)$ is the sorted ⁷ $(S_{\beta^0} y)$ vector. The optimization problem in (43) is very similar to the optimization problem in (42). The only difference is in the relaxed constraint, where the set of feasible solutions in (43) allows for negative values. The optimization (43) is a special case of the isotonic regression problem [5]:

$$\begin{aligned} &\text{minimize} && \|x - z\|_2^2 \\ &\text{subject to} && x_1 \geq \dots \geq x_p, \end{aligned}$$

where we set $z = (S_{\beta^0} y)_{(\cdot)} - \lambda$.

⁷Note that the permutation matrix Π depends both on y and β^0 .

A.3 FDR constant

Let $Z \sim \mathcal{N}(0, \mathbb{I}_p)$ and $\gamma > 0$. Consider the rejection procedure $\hat{\mathbf{b}} = \mathbf{patt}_{\beta^0}(\hat{\mathbf{u}})$, with

$$\begin{aligned}\hat{\mathbf{u}} &= \operatorname{argmin}_u u^T \gamma \mathbb{I}_p u / 2 - u^T \sigma Z + J'_\lambda(\beta^0; u) \\ &= \sqrt{\gamma} \operatorname{argmin}_{\tilde{u}} \tilde{u}^T \mathbb{I}_p \tilde{u} / 2 - \tilde{u}^T (\sigma / \sqrt{\gamma}) Z + J'_{(\lambda/\sqrt{\gamma})}(\beta^0; \tilde{u}),\end{aligned}$$

where $\tilde{u} = \sqrt{\gamma}u$. Therefore, if $\lambda_i = \sigma \lambda_i^{BH(q)} = \sigma \Phi^{-1}(1 - iq/2p)$, we have $\text{FDR}(\hat{\mathbf{b}}) \leq q(|I_0|/p)$, by Example 4.7. In particular, the penalties λ_i do not depend on the constant γ .