

AN ANTICIPATING TANGENT TO A SEMIMARTINGALE

A. S. ÜSTÜNEL

Abstract: We construct a perturbation of identity type mapping on an abstract Wiener space where the Cameron-Martin space has an orthonormal basis indexed by the jumps a one dimensional semimartingale. We then derive a change of variables formula and a degree type result for this map.

Keywords: Semimartingale, entropy, anticipating perturbation of identity, Wiener measure, invertibility, degree, measure transportation.

1. INTRODUCTION

Denote by $(\Omega, \mathcal{F}, (\mathcal{F}_t), P)$ a general probability space satisfying the usual conditions of the theory of stochastic processes. Let $Z = (Z_t, t \geq 0)$ be a real-valued semimartingale, denote its Doléans-Dade exponential as $\mathcal{E}_t(Z)$, i.e.,

$$\mathcal{E}_t(Z) = \exp\left(Z_t - \frac{1}{2}\langle Z^c, Z^c \rangle_t\right) \prod_{s \leq t} (1 + \Delta Z_s) e^{-\Delta Z_s},$$

where Z^c denotes the continuous local martingale part, $\langle Z^c, Z^c \rangle$ is the associated Doob-Meyer process and ΔZ denotes the jumps of Z . Since P -almost surely

$$(1.1) \quad \sum_{s \leq t} (\Delta Z_s)^2 < \infty$$

for any $t \geq 0$, the product term is well-defined. Consequently, for P -almost all $z \in \Omega$, the sequence $\{\Delta Z_s(z) : s \in [0, t], \Delta Z_s(z) \neq 0\}$ can be regarded as the spectrum of a Hilbert-Schmidt operator and then the term

$$D_t = \prod_{s \leq t} (1 + \Delta Z_s(z)) e^{-\Delta Z_s(z)}$$

becomes its modified Carleman-Fredholm determinant. Starting from this observation, we construct, on a Wiener space an anticipating perturbation of identity whose Gaussian Jacobian is the product of D_t with Wick-Girsanov exponential of this perturbation of identity. This paper is devoted to studying the properties of this Jacobian.

2. PRELIMINARIES

Let (W, H, μ) be any Wiener space, denote by $(e_i, i \geq 1)$ a CONB in the Cameron-Martin space H , constructed from the elements of W^* , i.e., the continuous dual of the Fréchet space W . Let $(T_n, n \geq 1)$ be a sequence of stopping times absorbing the jumps of the semimartingale Z with

disjoint graphs, cf. [2]. We shall construct a Hilbert-Schmidt operator ∂u_t , whose eigenvalues will consist of the jumps of Z : define

$$\partial u_t = \sum_{T_n \leq t} \Delta Z_{T_n} e_n \otimes e_n.$$

Because of the property (1.1), ∂u_t is a measurable map defined on $\mathbb{R}_+ \times \Omega$ with values in $H \tilde{\otimes}_2 H$, where the latter denotes the completed Hilbert-Schmidt tensor product. In fact

$$\|\partial u_t(z)\|_2^2 = \sum_{T_n(z) \leq t} (\Delta Z_{T_n}(z))^2 \leq \sum_{s \leq t} (\Delta Z_s(z))^2 < \infty$$

P -almost surely. Since this operator is also symmetric, for fixed z , we can construct an H -valued map, whose Sobolev derivative on the Wiener space will be equal to ∂u_t explicitly. In fact, it suffices to put

$$u_t(w, z) = \sum_{T_n(z) \leq t} \Delta Z_{T_n}(z) \delta e_n(w) e_n,$$

where δe_n denotes the Gaussian divergence of $e_n \in H$. Note that, defining

$$u_t^\varepsilon(w, z) = \sum_{T_n \leq t, |\Delta Z_{T_n}| > \varepsilon} \Delta Z_{T_n}(z) \delta e_n(w) e_n$$

we get well-defined, H -valued, measurable maps. For $\varepsilon, \eta > 0$, we have

$$|u_t^\varepsilon - u_t^\eta|_H^2 = |u_t^\varepsilon|_H^2 + |u_t^\eta|_H^2 - 2 \sum_{T_n(z) \leq t, \varepsilon \wedge \eta < |\Delta Z_{T_n}| \leq \varepsilon \vee \eta} (\Delta Z_{T_n}(z))^2 (\delta e_n(w))^2$$

since the graphs of (T_n) are disjoint. Consequently, for P -almost all z , $\lim_{\varepsilon \rightarrow 0} u_t^\varepsilon(\cdot, z) = u_t(\cdot, z)$ in $L^2(\mu, H)$, since $\lim \nabla u_t^\varepsilon(\cdot, z)$ exists also, $u_t(\cdot, z) \in \mathbb{D}_{p,1}(H)$ for any $p > 1$. We have also a more delicate result:

Theorem 1. *The mapping $U_t(w, z) = w + u_t(w, z)$ defined on $W \times \Omega$ with values in W , where u_t is given as*

$$u_t(w, z) = \sum_{T_n(z) \leq t} \Delta Z_{T_n} \delta e_n(w) e_n,$$

is $\mu \times P$ -almost surely well-defined perturbation of identity on the Wiener space W , besides, for P -almost all z , the partial map

$$w \rightarrow u_t(w, z)$$

is an $H - C^1$ -map on W , i.e., the map $h \rightarrow u_t(w + h, z)$ is a C^1 -map on H outside a set of capacity zero (even an H -invariant set).

Proof: For any $h \in H$, we have

$$u_t^\varepsilon(w + h, z) = u_t^\varepsilon(w, z) + \nabla u_t^\varepsilon(z) h$$

P -a.s. for any $w \in W$ (recall that $e_n \in W^*$). Hence

$$\begin{aligned} |u_t^\varepsilon(w + h, z) - u_t^\eta(w + h, z)|_H^2 &\leq 2|u_t^\varepsilon(w, z) - u_t^\eta(w, z)|_H^2 \\ &\quad + 2|h|_H^2 \|\nabla u_t^\varepsilon(z) - \nabla u_t^\eta(z)\|_2^2. \end{aligned}$$

Hence $(h \rightarrow u_t^\varepsilon(w + h, z), \varepsilon > 0)$ converges to $h \rightarrow u_t(w + h, z)$ uniformly w.r.to $h \in H$ and this implies that $h \rightarrow u_t(w + h, z)$ is an $H - C$ -map for almost all $z \in \Omega$. Moreover, $\nabla u_t(w, z)$ does not depend on w , hence $h \rightarrow u_t(w, z)$ is even $H - C^\infty$. \square

In the sequel $\Lambda_t = \Lambda_t(w, z)$ will represent the Gaussian Jacobian associated to the perturbation of identity $U_t(w, z)$, namely

$$\begin{aligned} \Lambda_t(w, z) &= \det_2(I_H + \nabla u_t(w, z)) \exp(-\delta u_t(w, z) - \frac{1}{2} |u_t(w, z)|_H^2) \\ &= \prod_{T_n(z) \leq t} (1 + \Delta Z_{T_n}) e^{-\Delta Z_{T_n}} \exp\left(-\sum_{T_n \leq t} \Delta Z_{T_n} ((\delta e_n)^2 - 1) - \frac{1}{2} \sum_{T_n \leq t} (\Delta Z_{T_n})^2 (\delta e_n)^2\right) \end{aligned}$$

We have the following degree-type result (cf. [1] or [7]):

Theorem 2. *Let us denote by M_t the set of nondegeneracy of Λ_t , namely*

$$M_t = \bigcap_{T_n \leq t} \{z \in \Omega : \Delta Z_{T_n} \neq -1\}.$$

Then we have

$$\mu(U_t(M_t^c)) = 0$$

and

$$(2.2) \quad \int f(z, U_t(w, z)) |\Lambda_t(w, z)| g(w, z) d\mu dP = \int f(w, z) \left(\sum_{y \in U_t^{-1}(\cdot, z)\{w\}} g(y, z) \right) d\mu dP$$

for any positive, measurable functions on $W \times \Omega$. Consequently, we have

$$(2.3) \quad \int_W f \circ U_t(w, z) \Lambda_t(w, z) d\mu = \int_W f d\mu \int_W \Lambda_t d\mu$$

P -almost surely and hence

$$(2.4) \quad \int_W \Lambda_t d\mu = \sum_{y \in U_t^{-1}(\cdot, z)\{w\}} \text{sign} \Lambda_t(y, z) = |\{U_t^{-1}(\cdot, z)\{w\}\}| \text{sign}(D_t(z))$$

$P \times \mu$ -almost surely, where $|\{\dots\}|$ denotes the cardinality of the set inside the brackets. Moreover, in the case of classical Wiener space, the mapping $w \rightarrow U_t(w, z)$ is μ -almost surely invertible on the set $\{z \in \Omega : E_\mu[|\Lambda_t|] = 1\}$, and its (measurable) inverse, denoted as $(\tau, w) \rightarrow V_t(w, z)(\tau)$, is the strong solution of the following functional stochastic differential equation

$$V_t(w, z)(\tau) = W_\tau - \int_0^\tau \dot{u}_t(V_t(w, z), z)(s) ds.$$

Proof: Since U_t is an $H - C^1$ -map P -almost surely, the Sard inequality implies that the image under itself of its set of degeneracy has zero μ -measure and the change of variables formula (2.2) holds cf. [7], Chapter 4, Section 4.4. The relation (2.3) follows from Theorem 9.2.3 of [7] and finally the equality (2.4) is a sequence of the relations (2.3) and (2.4). As explained in [8], the equivalence of the measures $U_t \mu$ and μ , combined with the fact that the cardinality of $U_t^{-1}(\cdot, z)\{w\}$ being equal to one μ -almost everywhere implies the almost sure invertibility the map $w \rightarrow U_t(w, z)$ P -almost everywhere and the functional stochastic differential equation is a direct consequence of this fact. \square

Remark: Note that the delicate point in the above result comes from the lack of information about the instants of the distorted noise posterior to $t \in \mathbb{R}_+$.

We can view $(\Lambda_t \in [0, 1])$ as a semimartingale with respect to the filtration (\mathcal{F}_t) . First let us define the process $(e(t), t \in [0, 1])$ as

$$e(t) = \sum_n e_n 1_{[T_n, T_{n+1}[}(t).$$

We have

Theorem 3. *Assume that Z is a pure jump semimartingale, then the process $(t, z) \rightarrow \Lambda_t(w, z)$ satisfies the following stochastic differential equation:*

$$\begin{aligned} \Lambda_t &= \Lambda_0 - \int_0^t \Lambda_{s-} \left[(\delta e_s)^2 \left(1 + \frac{1}{2} \Delta Z_s \right) - 1 \right] dZ_s \\ &\quad + \sum_{s \leq t} \Lambda_{s-} e^{\Delta Z_s [(\delta e_s)^2 (1 - \frac{1}{2} \Delta Z_s) - 1]} (e^{-\Delta Z_s} (1 + \Delta Z_s) - 1) \end{aligned}$$

REFERENCES

- [1] A. Al-Hussein and K. D. Elworthy: Infinite-dimensional degree theory and stochastic analysis. *J. Fixed Point Theory Appl.* 7 (2010), no. 1, 33–65.
- [2] C. Dellacherie: *Capacités et Processus Stochastiques*. Ergebnisse de Math. No. 67, Springer, 1972.
- [3] D. Feyel, A.S. Üstünel: Monge-Kantorovitch measure transportation and Monge-Ampère equation on Wiener space. *Probab. Theor. Relat. Fields*, **128**, no. 3, pp. 347–385, 2004.
- [4] A. S. Üstünel: *Introduction to Analysis on Wiener Space*. Lecture Notes in Math. Vol. 1610. Springer, 1995.
- [5] A. S. Üstünel: *Analysis on Wiener Space and Applications*. <http://arxiv.org/abs/1003.1649>, 2010.
- [6] A. S. Üstünel: “Entropy, invertibility and variational calculus of adapted shifts on Wiener space”. *J. Funct. Anal.* 257 (2009), no. 11, 3655–3689.
- [7] A. S. Üstünel and M. Zakai: *Transformation of Measure on Wiener Space*. Springer Verlag, 1999.
- [8] A. S. Üstünel and M. Zakai: “Sufficient conditions for the invertibility of adapted perturbations of identity on the Wiener space”. *Probab. Theory Relat. Fields*, **139**, p. 207-234, 2007.

A. S. Üstünel,
 Mathematics Departement, Bilkent University, Ankara, Turkey,
 ustunel@fen.bilkent.edu.tr