

FUNDAMENTAL GROUPS OF PROPER VARIETIES ARE FINITELY PRESENTED

MARCIN LARA, VASUDEVAN SRINIVAS, AND JAKOB STIX

ABSTRACT. It was proven in [ESS22], that the étale fundamental group of a connected smooth projective variety over an algebraically closed field k is topologically finitely presented. In this note, we extend this result to all connected proper schemes over k .

1. INTRODUCTION

For a connected algebraic variety X over an algebraically closed field k of characteristic 0, the étale fundamental group $\pi_1^{\text{ét}}(X, \bar{x})$ of X is a topologically finitely presented profinite group. This is proven by first reducing to the case of $k = \mathbb{C}$ and then applying the Riemann Existence Theorem [SGA1, Exp. XII, Thm. 5.1] together with the fact that the topological fundamental group $\pi_1^{\text{top}}(X(\mathbb{C}), x)$ is of finite presentation as a discrete group (see e.g. [Loj64] or [Hir75]).

In characteristic $p > 0$, the picture is much more subtle due to the existence of Artin-Schreier covers of affine schemes, which makes $\pi_1^{\text{ét}}(\text{Spec}(A), \bar{x})$ typically not even topologically finitely generated. Remark 5.7 of [SGA1, Exp. IX] raised doubts whether $\pi_1^{\text{ét}}(X, \bar{x})$ is topologically finitely presented for proper varieties, even for proper smooth curves. In recent work, however, Shusterman (in the case of curves [Shu22]), and Esnault, Shusterman and the second named author [ESS22] have shown that for smooth projective varieties $\pi_1^{\text{ét}}(X, \bar{x})$ is still topologically finitely presented. Both results are based on a criterion for finite presentation of profinite groups due to Lubotzky [Lub01].

From now on, we will omit the base points.

Theorem 1.1 (part of Thm. 1.1 of [ESS22]). *Let X be a connected smooth projective variety over an algebraically closed field k . Then the étale fundamental group $\pi_1^{\text{ét}}(X)$ is topologically finitely presented.*

Our goal is to generalize Thm. 1.1 to all connected schemes that are proper over $\text{Spec}(k)$ (which is new only if k has characteristic $p > 0$). Such a generalization responds affirmatively to a question raised by Esnault.

Theorem 1.2. *Let X be a connected scheme that is proper over $\text{Spec}(k)$ for an algebraically closed field k . Then $\pi_1^{\text{ét}}(X)$ is topologically finitely presented.*

To prove the theorem, we use descent along an alteration map to X and the van Kampen presentation of $\pi_1^{\text{ét}}(X)$ arising in this way. More precisely, we use this trick twice.

The authors acknowledge support by Deutsche Forschungsgemeinschaft (DFG) through the Collaborative Research Centre TRR 326 "Geometry and Arithmetic of Uniformized Structures", project number 444845124. The second author (VS) was supported during part of the preparation of the article by a J. C. Bose Fellowship of the Department of Science and Technology, India. He also acknowledges support of the Department of Atomic Energy, India under project number RTI4001. This material is partly based upon work supported by the NSF Grant No. DMS-1928930 while the third named author (JS) was in residence at MSRI in Berkeley (Spring 2023).

2. THE PROOF

For a scheme T , let $\mathbf{F}\acute{\text{E}}\text{t}_T$ denote the category of finite étale covers of T . This gives rise to a category fibred over schemes. Recall that a morphism $g : T' \rightarrow T$ of schemes is said to be of effective descent for $\mathbf{F}\acute{\text{E}}\text{t}$, if g^* induces an equivalence of categories between $\mathbf{F}\acute{\text{E}}\text{t}_T$ and the category of descent data in $\mathbf{F}\acute{\text{E}}\text{t}$ along g .

Proposition 2.1 (Exp. IX, Thm. 4.12 of [SGA1]). *Let $f : X' \rightarrow X$ be a proper surjective morphism of finite presentation. Then f is of effective descent for $\mathbf{F}\acute{\text{E}}\text{t}$.*

Morphisms of effective descent $f : X' \rightarrow X$ for $\mathbf{F}\acute{\text{E}}\text{t}$ give rise to a van Kampen-like presentation of $\pi_1^{\acute{\text{e}}\text{t}}(X)$ as the profinite completion of a quotient of the free topological product of the étale fundamental groups of the connected components of X' and the usual topological fundamental group of a suitably defined “dual graph”. This goes back to [SGA1, Exp. IX, § 5] and has been worked out in detail in [Sti06, Cor. 5.3].

The existence of such a presentation allows one to “descend” finite generation/presentation of the fundamental groups involved, as made precise in the following proposition. Every statement below is about *topological* finite generation/presentation.

Proposition 2.2 (Exp. IX, Cor. 5.2 + Cor. 5.3 of [SGA1]). *Let $f : X' \rightarrow X$ be a morphism of effective descent for $\mathbf{F}\acute{\text{E}}\text{t}$. We denote $X' \times_X X'$ by X'' and $X' \times_X X' \times_X X'$ by X''' .*

- (a) *Assume that X', X'' have finite π_0 's and that $\pi_1^{\acute{\text{e}}\text{t}}$'s of the connected components of X' are finitely generated. Then $\pi_1^{\acute{\text{e}}\text{t}}(X)$ is finitely generated.*
- (b) *Assume that X', X'', X''' have finite π_0 's, that $\pi_1^{\acute{\text{e}}\text{t}}$'s of the connected components of X' are of finite presentation and that $\pi_1^{\acute{\text{e}}\text{t}}$'s of the connected components of X'' are finitely generated. Then $\pi_1^{\acute{\text{e}}\text{t}}(X)$ is of finite presentation.*

Let us now recall a result of de Jong specialized to our setting.

Proposition 2.3 (see Thm. 4.1 of [dJ96]). *Let X be a scheme that is proper over $\text{Spec}(k)$ for an algebraically closed field k . Then there exists a proper surjective morphism (of finite presentation) $f : X' \rightarrow X$ from a smooth projective variety X' over $\text{Spec}(k)$.*

Proof. Let $\nu : X^\nu \rightarrow X$ be the normalization of X . The map ν is finite, and thus the scheme X^ν is still proper over $\text{Spec}(k)$.

Let then $f_1 : X' \rightarrow X^\nu$ be the alteration map of [dJ96, Thm. 4.1] applied to each connected component of X^ν . The map f_1 is proper, dominant, and thus surjective. Moreover, *loc. cit.* guarantees that X' is regular and projective (and not merely proper!) over $\text{Spec}(k)$. Now, as k is algebraically closed, X' is smooth over $\text{Spec}(k)$. The composition $f = \nu \circ f_1 : X' \rightarrow X$ has all the requested properties. \square

We are now ready to finish the proof of the main result.

Proof of Thm. 1.2. Take $f : X' \rightarrow X$ as in Prop. 2.3. By Thm. 1.1 and Prop. 2.1 the map f satisfies the assumptions for Prop. 2.2(a). This shows that $\pi_1^{\acute{\text{e}}\text{t}}(X)$ is finitely generated, for any scheme X that is proper over $\text{Spec}(k)$. In fact, finite generation was already proven in [SGA1, Exp. X, Thm. 2.9], and we included the argument here for the convenience of the reader.

We are going to apply finite generation to the connected components of $X'' = X' \times_X X'$, which are connected proper schemes over $\text{Spec}(k)$. Indeed, using Thm. 1.1 (this time crucially!) and Prop. 2.1 again, the map f now satisfies the assumptions of Prop. 2.2(b). This shows that $\pi_1^{\acute{\text{e}}\text{t}}(X)$ is finitely presented. \square

3. MORE GENERAL BASE FIELDS

Similarly to [ESS22, §5], our main result extends to more arithmetic settings. We thank Peter Haine for essentially suggesting the following corollary.

Corollary 3.1. *Let X be a connected scheme that is proper over $\mathrm{Spec}(k)$ for a field k . Then $\pi_1^{\text{ét}}(X)$ is finitely presented if and only if the absolute Galois group Gal_k is finitely presented.*

Proof. We may assume X is reduced and thus $k' = H^0(X, \mathcal{O}_X)$ is a finite field extension of k . Let \bar{k} be an algebraic closure of k containing k' . Then $X \rightarrow \mathrm{Spec}(k')$ being the Stein factorization of $X \rightarrow \mathrm{Spec}(k)$ implies that $\bar{X} = X \times_{k'} \bar{k}$ is connected. By Thm. 1.2, the group $\pi_1(\bar{X})$ is finitely presented. The fundamental exact sequence [SGA1, Exp. IX, Thm. 6.1]

$$1 \rightarrow \pi_1(\bar{X}) \rightarrow \pi_1(X) \rightarrow \mathrm{Gal}_{k'} \rightarrow 1$$

shows that $\pi_1(X)$ is finitely presented if and only if $\mathrm{Gal}_{k'}$ is finitely presented. The latter is equivalent to Gal_k being finitely presented, see for example [ESS23, Prop. 2.3]. \square

Remark 3.2. Examples of fields k with finitely presented Gal_k include: fields algebraic over a finite field, local p -adic fields, \mathbb{R} and more generally real closed fields, $K((t))$ for a field of characteristic 0 with Gal_K of finite presentation, and by [Jar74, Thm. 5.1] for a hilbertian field k , probabilistically almost always (for the Haar measure on Gal_k) the fixed field k^Σ in the separable closure \bar{k} of a finite subset $\Sigma \subset \mathrm{Gal}_k$.

REFERENCES

- [dJ96] Aise Johan de Jong. Smoothness, semi-stability and alterations. *IHES Publ. Math.*, 83:51–93, 1996.
- [ESS22] Hélène Esnault, Mark Shusterman, and Vasudevan Srinivas. Finite presentation of the tame fundamental group. *Selecta Math. (N.S.)*, 28(2):Paper No. 37, 19, 2022.
- [ESS23] Hélène Esnault, Vasudevan Srinivas, and Jakob Stix. An obstruction to lifting to characteristic 0. *Algebraic Geometry*, 10(3):327–347, 2023.
- [Hir75] Heisuke Hironaka. Triangulations of algebraic sets. In *Algebraic geometry (Proc. Sympos. Pure Math., Vol. 29, Arcata, Calif., 1974)*, pages 165–185. Amer. Math. Soc., Providence, R.I., 1975.
- [Jar74] Moshe Jarden. Algebraic extensions of finite corank of Hilbertian fields. *Israel J. Math.*, 18:279–307, 1974.
- [Łoj64] Stanisław Łojasiewicz. Triangulation of semi-analytic sets. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)*, 18:449–474, 1964.
- [Lub01] Alexander Lubotzky. Pro-finite presentations. *J. Algebra*, 242(2):672–690, 2001.
- [SGA1] *Revêtements étales et groupe fondamental*. Lecture Notes in Mathematics, Vol. 224. Springer-Verlag, Berlin-New York, 1971. Séminaire de Géométrie Algébrique du Bois Marie 1960–1961 (SGA 1), Dirigé par Alexandre Grothendieck. Augmenté de deux exposés de Michèle Raynaud.
- [Shu22] Mark Shusterman. Balanced presentations for fundamental groups of curves over finite fields. *Mathematical Research Letters*, 29(4):1251–1259, 2022.
- [Sti06] Jakob Stix. A general Seifert-Van Kampen theorem for algebraic fundamental groups. *Publ. Res. Inst. Math. Sci.*, 42(3):763–786, 2006.

INSTITUTE OF MATHEMATICS, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, JAGIELLONIAN UNIVERSITY, ŁOJASIEWICZA 6, 30-348 KRAKÓW, POLAND

Email address: marcin.lara@uj.edu.pl

TIFR, SCHOOL OF MATHEMATICS, HOMI BHABHA ROAD, COLABA, 400005 MUMBAI, INDIA

Email address: srinivas@math.tifr.res.in

INSTITUT FÜR MATHEMATIK, GOETHE-UNIVERSITÄT FRANKFURT, ROBERT-MAYER-STRASSE 6–8, 60325 FRANKFURT AM MAIN, GERMANY

Email address: stix@math.uni-frankfurt.de