

BUBBLING OF KÄHLER-EINSTEIN METRICS

SONG SUN

In Memory of Professor Jean-Pierre Demailly

ABSTRACT. We prove the finite step termination of bubble trees for singularity formation of polarized Kähler-Einstein metrics in the non-collapsing situation. We also raise several questions and conjectures in connection with algebraic geometry and Riemannian geometry.

1. INTRODUCTION

The study of Kähler-Einstein metrics has been a central topic in geometry for the past few decades. The main existence results on compact Kähler manifolds are due to Aubin [2]-Yau [36] (in the case $c_1 < 0$), Yau [36] (in the case $c_1 = 0$) and Chen-Donaldson-Sun [7] (in the case $c_1 > 0$, in terms of the K-stability criterion). One interesting research direction is lead by the interplay between geometric analysis and complex algebraic geometry of singularities. On the one hand, bridges with algebraic geometry allow one to prove deep results concerning the structure of singularities of Kähler-Einstein metrics that are not yet accessible using traditional techniques in geometric analysis. On the other hand, geometric analytic results motivate the right questions concerning singularities in algebraic geometry that would not have been envisioned otherwise and enable the application of sophisticated algebro-geometric techniques to answer these questions and form algebro-geometric generalizations.

In this paper we will make some further progress along these lines. More precisely, we aim toward understanding the algebro-geometric meaning of singularity formation process of Kähler-Einstein metrics, building upon previous results and discussion in [17, 18, 16, 30]. To set the stage we consider a sequence of polarized Kähler-Einstein metrics $(X_j, L_j, \omega_j, p_j)$ with $\omega_j \in 2\pi c_1(L_j)$ and $Ric(\omega_j) = c_j \omega_j$ ($c_j \in \{1, 0, -1\}$). When $c_j = 0$ we also assume K_{X_j} is holomorphically trivial. We assume that $\text{diam}(X_j, \omega_j) \leq D$ for some $D \geq 0$. The latter in particular implies a uniform *volume non-collapsing* condition, i.e., there exists a $\kappa > 0$ such that

$$(1.1) \quad \text{Vol}(B(p_j, 1)) \geq \kappa$$

for all j . It is proved in [17] that after passing to a subsequence we obtain a pointed Gromov-Hausdorff limit $(X_\infty, d_\infty, p_\infty)$ which has the structure of a normal projective variety with log terminal singularities and d_∞ is induced by a weak Kähler-Einstein metric ω_∞ on X_∞ . It is proved in [18] that there is a unique metric tangent cone at p_∞ , which we denote by $\mathcal{C}_0(X_\infty)$. The latter is naturally a normal affine variety endowed with a weak Calabi-Yau (i.e., Ricci-flat Kähler) cone metric. Furthermore, the underlying

Date: March 21, 2023.

algebraic structure on $\mathcal{C}_0(X_\infty)$ can be obtained in terms of a 2-step degeneration from the local ring $\mathcal{O}_{X_\infty, p_\infty}$ defined by the metric. The cone $\mathcal{C}_0(X_\infty)$ and an intermediate K-semistable Fano cone W are conjectured in [18] to be algebro-geometric invariants of the germ of X_∞ at p_∞ , thus leading to a local notion of stability. This conjecture was confirmed by Li-Wang-Xu [23]. There are also many follow-up work on related topics in algebraic geometry; see for example [35] and the references therein for more information.

From the viewpoint of geometric analysis, tangent cones can be regarded as only the *first order* approximation of the singular behavior of solutions to an elliptic non-linear PDE. In our case the tangent cone $\mathcal{C}_0(X_\infty)$ only depends on the limit space X_∞ , as such it does not encode the dynamic information on how the singularity is formed. To investigate the latter we consider the limits of the rescaled spaces $(X_j, \lambda_j \omega_j, p_j)$ for *all* possible sequences of integers $\lambda_j \geq 1$. Clearly if λ_j stays uniformly bounded then the limit is simply a rescaling of X_∞ ; if $\lambda_j \rightarrow \infty$, then passing to a subsequence by [18] we obtain a non-compact pointed Gromov-Hausdorff limit (Z, d_Z, p_Z) ; the latter is naturally an affine variety with a weak Calabi-Yau metric ω_Z which is asymptotic to a unique metric cone $\mathcal{C}_\infty(Z)$ at infinity. Moreover, there is also a unique metric tangent cone $\mathcal{C}_0(Z)$ at p_Z .

From now on we fix the given convergent sequence $(X_j, L_j, \omega_j, p_j)$. For our interest in this article we will always assume p_∞ is a singular point.

Definition 1.1. A *bubble limit* (Z, ω_Z, p_Z) associated to the above sequence is a subsequential pointed Gromov-Hausdorff limit of the sequence $(X_j, \lambda_j \omega_j, p_j)$ for some $\lambda_j \geq 1$.

Remark 1.2. *For notational simplicity we will sometimes drop ω_Z and p_Z when discussing bubble limits, but it should be understood that Z always carries the data (ω_Z, p_Z) implicitly.*

Remark 1.3. *The notion of (pointed) Gromov-Hausdorff convergence we are referring to in this paper is stronger than the standard definition in terms of metric spaces, since we also include the information on the convergence of complex structures. For example, a priori the same metric space Z may be seen as an affine variety in different ways when passing to different further subsequences. See [18], Section 2.2 for more precise definition of the convergence.*

By definition a bubble limit is endowed with both an algebraic structure and a metric structure. We denote by \mathfrak{R} the class of all bubble limits modulo the equivalence relation given by holomorphic isometries keeping the base points. Obviously X_∞ is the only compact bubble limit. By a diagonal sequence argument we know $\mathcal{C}_0(X_\infty) \in \mathfrak{R}$. Also for any non-compact $Z \in \mathfrak{R}$, both the tangent cone $\mathcal{C}_0(Z)$ and the asymptotic cone $\mathcal{C}_\infty(Z)$ are in \mathfrak{R} . Notice that \mathfrak{R} depends not only on the limit point p_∞ , but also on the sequence of points p_j converging to p_∞ .

Definition 1.4. For any $Z \in \mathfrak{R}$, its volume density at p_Z is defined to be

$$\text{Vol}_0(Z) \equiv \lim_{r \rightarrow 0} \frac{\text{Vol}(B(p_Z, r))}{\text{Vol}(B_0(r))};$$

when Z is non-compact, its volume density at infinity is defined to be

$$\text{Vol}_\infty(Z) \equiv \lim_{r \rightarrow \infty} \frac{\text{Vol}(B(p_Z, r))}{\text{Vol}(B_0(r))}.$$

Here $B_0(r)$ denotes the radius r ball in the Euclidean space \mathbb{C}^n .

The Bishop-Gromov inequality implies that these are well-defined positive numbers and we have $\text{Vol}_0(Z) \geq \text{Vol}_\infty(Z) \geq \text{Vol}_0(X_\infty)$. Moreover, the first equality holds if and only if Z is a metric cone with p_Z as a vertex. If this occurs we also denote

$$\text{Vol}(Z) \equiv \text{Vol}_\infty(Z).$$

In particular, $\text{Vol}_0(Z) = \text{Vol}(\mathcal{C}_0(Z))$ and $\text{Vol}_\infty(Z) = \text{Vol}(\mathcal{C}_\infty(Z))$.

Definition 1.5. A bubble limit (\mathcal{C}, O) is a *cone limit* if \mathcal{C} is a metric cone with O as a vertex.

Remark 1.6. As above we will often simply denote by \mathcal{C} a cone limit.

Our first result is that

Theorem 1.7 (Finiteness of volumes). *There is a constant $\mathfrak{V} = \mathfrak{V}(n, D)$ such that the set*

$$\mathcal{V}(\mathfrak{R}) = \{\text{Vol}_0(Z) \mid Z \in \mathfrak{R}\}$$

is a finite set with cardinality bounded by \mathfrak{V} .

Remark 1.8. *It is proved in [27, 18] that $\text{Vol}_0(Z)$ is always an algebraic number, as a consequence of the volume minimization principle discovered by Martelli-Sparks-Yau [27]. This was used in [18] to show the rigidity of the holomorphic spectrum (and the volume density) in a continuous family of cone limits. The latter plays an indispensable role in the study of tangent cones and asymptotic cones for Kähler-Einstein metrics. However the algebraicity of volume does not seem to directly yield Theorem 1.7.*

Our proof of Theorem 1.7 uses connections with algebraic geometry. It is natural to ask if the same statement holds in the more general Riemannian setting, see Section 5.3 for more discussion.

Definition 1.9. A *minimal bubble* associated to the above sequence is a bubble limit which is not a cone limit but is asymptotic to $\mathcal{C}_0(X_\infty)$.

Next we prove

Theorem 1.10 (Existence of minimal bubbles). *There is a sequence of scales $\lambda_j \rightarrow \infty$ such that the bubble limits arising from subsequential limits of $(X_j, \lambda_j \omega_j, p_j)$ are minimal bubbles. Furthermore, for any sequence of scales $\lambda'_j \rightarrow \infty$ with $\lambda'_j \lambda_j^{-1} \rightarrow 0$, $(X_j, \lambda'_j \omega_j, p_j)$ converges to $\mathcal{C}_0(X_\infty)$ in the pointed Gromov-Hausdorff topology.*

Remark 1.11. *The second statement and the Bishop-Gromov inequality imply that a minimal bubble has the minimum volume density at infinity among all non-compact bubble limits and it is the unique such minimizer beside the tangent cone $\mathcal{C}_0(X_\infty)$. This justifies our choice of the terminology “minimal bubbles”.*

To deduce Theorem 1.10 from Theorem 1.7 the key step is to rule out the possible appearance of a continuous family of cone limits containing $\mathcal{C}_0(X_\infty)$. Our argument again makes crucial use of algebraic geometry. It can be viewed as an extension of the proof for the uniqueness of tangent cones in [18].

Corollary 1.12 (Finite step termination of bubble tree). *Given any subsequence of $\{j\}$, passing to a further subsequence the set of bubble limits of (X_j, ω_j, p_j) modulo metric scalings is a finite ordered set $\{(Z_\alpha, q_\alpha)\}_{\alpha=1}^{2k}$ with $\text{Vol}_0(Z_\alpha)$ strictly increasing, $(Z_1, q_1) = (X_\infty, p_\infty)$, $(Z_{2k}, q_{2k}) = (\mathbb{C}^n, 0)$, and such that for all α , $(Z_{2\alpha}, q_{2\alpha})$ is a cone limit, and $\mathcal{C}_0(Z_{2\alpha-1}) = Z_{2\alpha} = \mathcal{C}_\infty(Z_{2\alpha+1})$. Here k is uniformly bounded by an integer depending on n and D .*

It is a natural intriguing question to ask for algebro-geometric interpretations of these minimal bubbles. See Section 5.1 for discussion.

Acknowledgements. The author is grateful to Professor Simon Donaldson and Cristiano Spotti for helpful conversations on the topics of this article some years ago at Stony Brook. He would like to thank Xuemiao Chen and Junsheng Zhang for relevant discussions in related context. He also thanks Martin de Borbon, Cristiano Spotti and Junsheng Zhang for valuable comments that improved the exposition. The author is partially supported by NSF Grant DMS-2004261 and the Simons Collaboration Grant in special holonomy.

2. PRELIMINARIES

2.1. Projective and affine algebraic structures. Let $(X_j, L_j, \omega_j, p_j)$ be given as in the introduction. Let $\lambda_j \geq 1$ be a sequence of integers and consider the rescaled sequence $(X_j, L_j^{\lambda_j}, \lambda_j \omega_j, p_j)$. From [17, 18], after passing to a subsequence we have the following

(1). There is a pointed Gromov-Hausdorff limit (Z, d_Z, p_Z) . The latter is a complete length space which is equipped with a regular-singular decomposition $Z = Z^{reg} \cup Z^{sing}$ in the sense of Cheeger-Colding. The regular set Z^{reg} is an open subset which is a smooth complex manifold and the limit metric d_Z is induced by a smooth Kähler-Einstein metric ω_Z on Z^{reg} . The convergence over Z^{reg} is in the smooth Cheeger-Gromov sense.

(2). Z is a normal complex-analytic space with structure sheaf given by $\iota_* \mathcal{O}_{Z^{reg}}$, where $\iota : Z^{reg} \rightarrow Z$ is the natural inclusion map. Moreover, ω_Z extends to a global closed positive $(1, 1)$ current with locally continuous potentials. The metric singular set Z^{sing} coincides with the complex-analytic singular set of Z .

(3). Z has only log terminal singularities and ω_Z satisfies the weak Kähler-Einstein equation globally on Z .

(4). Fix a sequence of pointed Gromov-Hausdorff approximations realizing the convergence (as in [18], see also the discussion at the end of this subsection). Then there are $0 < \lambda_2 < \lambda_1 \leq 1$ (depending only on n and D) such that given any holomorphic function f on $B(q, \lambda_1) \subset Z$ and $q_j \in X_j$ converging to q , for j large there is a holomorphic function f_j on $B(q_j, \lambda_2) \subset X_j$ that naturally converges to f .

Remark 2.1. *Item (4) says that the convergence is complex-analytic in a weak sense. It implies that locally one can define a holomorphic map $F_j : B(q_j, \lambda) \rightarrow \mathbb{C}^N$ for all j large, which is generically one-to-one (so is a normalization map) and converges naturally to a holomorphic embedding $F_\infty : B(q_\infty, \lambda) \rightarrow \mathbb{C}^N$. A non-trivial question is whether F_j can be made an embedding for j large. Moreover, as remarked in [18], it is not yet known how to describe the convergence of the images of F_j in a certain “flat” family. For example,*

are locally the defining equations for the image of F_j given by small perturbations of the defining equations of Z ? This is a technical point that will be addressed in our special setting in this paper; in general it seems to be a missing link between geometric analysis and complex analytic geometry, which is worth further investigation.

If Z is compact then it is simply given by a rescaling of the original limit space X_∞ . In this case by [17] we know that (possibly passing to a further subsequence) there is a limit hermitian holomorphic line bundle L_∞ over X_∞^{reg} , which extends as an ample \mathbb{Q} -line bundle to the entire X_∞ . The latter makes X_∞ a projective variety. Moreover, the convergence $X_j \rightarrow X_\infty$ can be realized algebro-geometrically as the convergence in a fixed Hilbert scheme of projective varieties in some $\mathbb{C}\mathbb{P}^M$. These results have applications to the study of moduli of Fano manifolds, see [29, 22, 28].

If Z is non-compact, then by [18] we have

- Z is a normal affine variety whose coordinate ring $R(Z)$ is given by the space of holomorphic functions with polynomial growth at infinity.
- For any non-zero $f \in R(Z)$, it has a well-defined degree at infinity

$$(2.1) \quad \deg(f) \equiv \lim_{r \rightarrow \infty} \frac{\log \sup_{B(p_Z, r)} |f|}{\log r} \geq 0.$$

For later purpose we recall slightly more details on the enhanced (pointed) Gromov-Hausdorff topology we use in this paper, which incorporates the convergence of complex structures (see [18] for more details). Given metric balls $B \subset Z, B' \subset Z'$ for $Z, Z' \in \mathfrak{R}$, we define $d_{EGH}(B, B')$ to be the smallest $\epsilon > 0$ such that the following holds: there is a metric d on the disjoint union $B \sqcup B'$ extending the metrics on B and B' , such that there are open subsets $U \subset B^{reg}, U' \subset B'^{reg}$ which are both ϵ dense in $B \sqcup B'$ and there exists a diffeomorphism $\chi : U \rightarrow U'$ with $d(x, \chi(x)) \leq \epsilon$ and satisfying

$$\|\chi^* g' - g\|_{C^2(U)} + \|\chi^* J' - J\|_{C^2(U)} \leq \epsilon,$$

where (g, J) and (g', J') denote the Kähler structure on U, U' respectively and $C^2(U)$ is the C^2 norm on U defined by the metric g . When no such $\epsilon > 0$ exists we simply define $d_{EGH}(B, B') = 1$. Notice d_{EGH} is not necessarily a metric on \mathfrak{R} but we do not need this property. It is clear from the definition that $d_{EGH}(B, B') = 0$ if and only if there is a holomorphic isometry between B and B' . We say a sequence B_j converges to B_∞ if $d_{EGH}(B_j, B_\infty) \rightarrow 0$ as $j \rightarrow \infty$. More generally, we say that a sequence $(Z_j, p_{Z_j}) \in \mathfrak{R}$ converges to a limit (Z_∞, p_{Z_∞}) if for any $R > 0$, $d_{EGH}(B(p_{Z_j}, R), B(p_{Z_\infty}, R)) \rightarrow 0$ as $j \rightarrow \infty$.

2.2. Cone limits. Recall that we denoted by \mathfrak{R} the class of all bubble limits associated to the sequence $(X_j, L_j, \omega_j, p_j)$, and an element (\mathcal{C}, O) in \mathfrak{R} is a *cone limit* if \mathcal{C} is a metric cone with O as the vertex. We denote by \mathfrak{C} the subclass of \mathfrak{R} consisting of all cone limits. Given a metric cone \mathcal{C} we denote by $B_{\mathcal{C}}$ the unit ball in \mathcal{C} centered at O . Due to the cone structure one sees that the enhanced pointed Gromov-Hausdorff topology on \mathfrak{C} is induced by the enhanced Gromov-Hausdorff topology on the unit balls $B_{\mathcal{C}}$. Namely, a sequence of cone limits \mathcal{C}_j converge to \mathcal{C} if and only if $d_{EGH}(B_{\mathcal{C}_j}, B_{\mathcal{C}}) \rightarrow 0$.

Differential geometrically the cone structure is reflected by the dilation invariance. Over the regular set of a cone limit \mathcal{C} there is a dilation vector field given by $r\partial_r$ (where

r is the distance function to O). The *Reeb vector field* $\xi \equiv Jr\partial_r$ is a holomorphic Killing field, which generates a holomorphic isometric action of a compact torus \mathbb{T} on \mathcal{C} . In [18] we discussed the algebro-geometric interpretation of the cone structure. In particular we have

(1). \mathcal{C} is a polarized normal affine cone. By this we mean that there is a positive \mathbb{R} -grading on the coordinate ring

$$(2.2) \quad R(\mathcal{C}) = \bigoplus_{\mu \in \mathcal{S}(\mathcal{C})} R_\mu(\mathcal{C}),$$

where $\mathcal{S}(\mathcal{C}) \subset \mathbb{R}_{\geq 0} \cap \{\langle \alpha, \xi \rangle \mid \alpha \in \Gamma^*\}$ for $\Gamma^* \subset \text{Lie}(\mathbb{T})^*$ the weight lattice of \mathbb{T} . As in [18], we call $\mathcal{S}(\mathcal{C})$ the *holomorphic spectrum* of \mathcal{C} . In this paper we will denote

$$R_I(\mathcal{C}) = \bigoplus_{\mu \in I} R_\mu(\mathcal{C})$$

for any interval I .

(2). $f \in R_\mu(\mathcal{C})$ if and only if f is homogeneous of degree μ , i.e., $\mathcal{L}_\xi f = \sqrt{-1}\mu f$. For $f \in R(\mathcal{C})$, $\deg(f) = \mu$ if and only if the highest degree term in the expansion of f with respect to (2.2) has degree μ .

(3). The \mathbb{R} -grading is induced by a positive multi-grading

$$R(\mathcal{C}) = \bigoplus_{\alpha \in \Gamma^*} \mathcal{H}_\alpha(\mathcal{C}),$$

where $\mathcal{H}_\alpha(\mathcal{C}) = R_\mu(\mathcal{C})$ for $\mu = \langle \alpha, \xi \rangle$.

(4). A choice of a homogeneous generating set of $R(\mathcal{C})$ gives rise to an algebraic embedding of \mathcal{C} as an affine variety in some \mathbb{C}^N and the Reeb vector field ξ extends to a linear diagonal vector field $\xi = \text{Re}(\sqrt{-1} \sum_{\alpha} \zeta_\alpha z_\alpha \partial_{z_\alpha})$ on \mathbb{C}^N such that each $\zeta_\alpha = \deg(z_\alpha) > 0$.

Obviously the decomposition (2.2) is orthogonal with respect to the L^2 inner product on $R(\mathcal{C})$ defined by

$$(2.3) \quad \langle f_1, f_2 \rangle \equiv \int_{\mathcal{C}} f_1 \bar{f}_2 e^{-\frac{r^2}{2}} d\mu.$$

We define the Hilbert function of \mathcal{C} to be $h_{\mathcal{C}} : \mathbb{R}_{\geq 0} \rightarrow \mathbb{Z}_{\geq 0}; \mu \mapsto \dim R_\mu(\mathcal{C})$.

Now we fix an embedding of \mathcal{C} in some \mathbb{C}^N as given in item (4) above. Denote by \mathbb{S} the coordinate ring of \mathbb{C}^N . We will make use of the theory of multi-graded Hilbert schemes developed by Haiman-Sturmfels [20]. There is a projective scheme **Hilb** parameterizing affine schemes of the form $\text{Spec}(\mathbb{S}/\mathcal{I})$, where \mathcal{I} is a ξ -homogeneous ideal with Hilbert function $h_{\mathcal{I}}(d) \equiv \dim(\mathbb{S}/\mathcal{I})_d = h_{\mathcal{C}}(d)$ for all $d \geq 0$. In particular \mathcal{C} corresponds to a point $[\mathcal{C}] \in \mathbf{Hilb}$. By the construction in [20], there is a flat universal family over **Hilb** and a boundedness result is proved using the theory of Gröbner basis. Namely, there exists a $\mu_0 = \mu_0(\mathcal{C})$ depending only on \mathcal{C} (more precisely, on the function $h_{\mathcal{C}}$) such that if a ξ -homogeneous ideal \mathcal{I} satisfies that $h_{\mathcal{I}}(d) = h_{\mathcal{C}}(d)$ for $d \leq \mu_0$, then $h_{\mathcal{I}}(d) = h_{\mathcal{C}}(d)$ for all d . Let $GL_\xi(N)$ be the subgroup of $GL(N; \mathbb{C})$ that preserves ξ . It naturally acts on **Hilb**. The stabilizer group of $[\mathcal{C}]$ agrees with $\text{Aut}(\mathcal{C})$, the group of holomorphic transformations of \mathcal{C} fixing the vertex O .

The above discussion does not make use of the weak Calabi-Yau metric on \mathcal{C} . A crucial consequence of the latter is the volume minimization property of Martelli-Sparks-Yau (see [27, 18]). In our setting we know \mathcal{C} has \mathbb{Q} -Gorenstein singularities, thus it is a *Fano cone* in the sense of [11]. This means that there exists a non-vanishing holomorphic section Ω of $K_{\mathcal{C}}^l$ (for some positive integer l) in a neighborhood \mathcal{U} of O . We may assume \mathcal{U} is \mathbb{T} invariant. Now $\mathcal{L}_{\xi}\Omega = f\Omega$ for some holomorphic function f on \mathcal{U} . As is explained in [18] we may perform a weight decomposition $f = \sum_{\alpha} f_{\alpha}$, where $\mathcal{L}_{\xi}f_{\alpha} = \sqrt{-1}\langle \alpha, \xi \rangle f_{\alpha}$.

Set

$$F = \exp\left(\sum_{\langle \alpha, \xi \rangle \neq 0} -\frac{\sqrt{-1}}{\langle \alpha, \xi \rangle} f_{\alpha}\right),$$

then replacing Ω by $F\Omega$ we may assume f is a constant. Since \mathbb{T} is compact it follows that f is an imaginary constant.

Given a compact torus \mathbb{T}' acting on \mathcal{C} holomorphically such that $\xi \in \text{Lie}(\mathbb{T}')$. For any $\xi' \in \text{Lie}(\mathbb{T}')$, we may write $\mathcal{L}_{\xi'}\Omega = f'\Omega$ for a holomorphic function f' . Using the fact that $[\xi, \xi'] = 0$ it follows that $\xi(f') = 0$. So $r\partial_r(f') = 0$. Then f' is globally bounded hence must be a constant. Then we get a linear map

$$c : \text{Lie}(\mathbb{T}') \rightarrow \mathbb{R}$$

such that $\mathcal{L}_{\xi'}\Omega = \sqrt{-1}c(\xi')\Omega$.

Now the Calabi-Yau cone metric $\omega_{\mathcal{C}}$ on \mathcal{C} satisfies the equation

$$\omega_{\mathcal{C}}^n = e^U (\Omega \wedge \bar{\Omega})^{1/l},$$

where U is pluri-harmonic. Since $\mathcal{L}_{\xi}\omega_{\mathcal{C}} = 2\omega_{\mathcal{C}}$, it follows that

$$\mathcal{L}_{\xi}U = \sqrt{-1}(2n - 2c(\xi)l^{-1}).$$

Hence U must be a constant and then

$$(2.4) \quad c(\xi) = nl.$$

Denote by

$$(2.5) \quad \mathbb{H} = \{\xi' \in \text{Lie}(\mathbb{T}') \mid c(\xi') = nl\},$$

and by $\text{Lie}(\mathbb{T}')^+$ the set of ξ' such that $\langle \alpha, \xi' \rangle \geq 0$ whenever $\mathcal{H}_{\alpha} \neq 0$. Then we have the volume minimization principle [27, 18]

- There is a volume function

$$\mathbb{V} : \text{Lie}(\mathbb{T}')^+ \cap \mathbb{H} \rightarrow \mathbb{R}_{>0}$$

which is determined by the function $h_{\mathcal{C}}$ in terms of the index character. It is a rational function with rational coefficients and $\mathbb{V}(\xi)$ coincides with the Riemannian volume of the cross section of the cone \mathcal{C} .

- \mathbb{V} is strictly convex.
- ξ is a critical point of \mathbb{V} .

As a consequence we know that ξ and \mathbb{V} are both algebraic. We will need two results proved in [18]:

- (Matsushima theorem) For any cone limit \mathcal{C} , the automorphism group $\text{Aut}(\mathcal{C})$ is reductive. More precisely, it is given by the complexification of the holomorphic isometry group of \mathcal{C} fixing the vertex.
- (Bando-Mabuchi theorem) The weak Calabi-Yau metric on a cone limit is uniquely determined by the underlying Fano cone structure, up to holomorphic isometries that preserve the Reeb vector field.

2.3. Algebraic geometry of tangent cones and asymptotic cones. Given any bubble limit $(Z, \omega_Z, p_Z) \in \mathfrak{R}$, we consider the rescaled spaces $(Z, \lambda\omega_Z, p_Z)$ for $\lambda \in (0, \infty)$. As $\lambda \rightarrow \infty$, passing to subsequences we obtain bubble limits which are *tangent cones* at p_Z . It is proved in [18] that in our setting the tangent cone is unique, which we denote by $(\mathcal{C}_0(Z), O)$.

More interestingly, in [18] it is established a 2-step degeneration theory, i.e., $\mathcal{C}_0(Z)$ can be described algebro-geometrically in terms of a degeneration from the local germ at p_Z in two steps. There is a valuation on the local ring \mathcal{O}_{Z, p_Z} defined by the degree function

$$\deg_{p_Z}(f) \equiv \lim_{r \rightarrow 0} \frac{\log \sup_{B(p_Z, r)} |f|}{\log r} \in \mathcal{S}(\mathcal{C}_0(Z)).$$

Denote by $R(p_Z)$ the associated graded ring. Then $W = \text{Spec}(R(p_Z))$ defines a Fano cone and there is then a further equivariant degeneration from W to $\mathcal{C}_0(Z)$.

There is also a dual picture for $\lambda \rightarrow 0$ when Z is non-compact. In this case it follows from [18] that $(Z, \lambda\omega_Z, p_Z)$ converges to a unique asymptotic cone $\mathcal{C}_\infty(Z)$. There is a negative valuation on $R(Z)$ (in the sense of [31]) which is related to the degree function by

$$\nu(f) = -\deg(f) = -\lim_{r \rightarrow \infty} \frac{\log \sup_{B(p_Z, r)} |f|}{\log r} \in -\mathcal{S}(\mathcal{C}_\infty(Z)).$$

Denote by $R_\infty(Z)$ the graded ring associated to the filtration defined by the negative valuation. Then $W_\infty = \text{Spec}(R_\infty(Z))$ defines a Fano cone and there is an equivariant degeneration from W_∞ to $\mathcal{C}_\infty(Z)$.

In both cases $\mathcal{C}_0(Z)$ and $\mathcal{C}_\infty(Z)$ can be viewed as K-polystable Fano cones, whereas W and W_∞ can be viewed as intermediate K-semistable Fano cones. For the definitions of stabilities used here one is referred to for example [11, 23]. However, there is a sharp contrast between the local and the asymptotic situation. In the local case, it is conjectured in [18] that both W and $\mathcal{C}_0(Z)$ are invariants of the algebraic singularity of Z at p_Z , which should lead to a local notion of stability for general singularities. This conjecture is confirmed in [24] and [23], based on the earlier work of Li [21] which further reformulated and generalized the original conjecture in [18] using more algebro-geometric language and the volume minimization principle of Martelli-Sparks-Yau [27]. As a result, one can compute both W and $\mathcal{C}_0(Z)$ in certain explicit examples; in particular, there are examples where W and $\mathcal{C}_0(Z)$ are genuinely distinct. In the asymptotic setting, it is the case that neither W_∞ nor $\mathcal{C}_\infty(Z)$ is an invariant of the affine variety Z only. For example, even \mathbb{C}^3 admits a complete Calabi-Yau metric with non-flat asymptotic cone (see [25, 12, 33]). Moreover, there is a new phenomenon discovered in [31], which is referred to as “no semistability at infinity”. The latter says that in this case W_∞ must be equal to $\mathcal{C}_\infty(Z)$ if $\mathcal{C}_\infty(Z)$ has a smooth cross section; this is conjectured to be always true in general.

3. DISCRETENESS OF VOLUMES

In this section we prove Theorem 1.7. We start with

Lemma 3.1. *Let $\mathcal{C} \in \mathfrak{C}$ be a cone limit and $\mu \in \mathcal{S}(\mathcal{C})$. For any $\epsilon > 0$ small, there exists a neighborhood \mathcal{U} of \mathcal{C} in \mathfrak{C} such that for any cone limit \mathcal{C}' in \mathcal{U} , it holds that*

$$\dim R_{(\mu-\epsilon, \mu+\epsilon)}(\mathcal{C}') = \dim R_\mu(\mathcal{C}).$$

Proof. Suppose otherwise, then we may find $\epsilon > 0$ with $[\mu - \epsilon, \mu + \epsilon] \cap \mathcal{S}(\mathcal{C}) = \{\mu\}$ and a sequence $(\mathcal{C}_i, O_i) \in \mathfrak{S}$ converging to \mathcal{C} such that $\dim R_{(\mu-\epsilon, \mu+\epsilon)}(\mathcal{C}_i) \neq \dim R_\mu(\mathcal{C})$ for all i .

We first claim this implies that for i large $\dim R_{(\mu-\epsilon, \mu+\epsilon)}(\mathcal{C}_i) < \dim R_\mu(\mathcal{C})$. Suppose otherwise, then we can choose a homogeneous orthonormal basis of $R_{(\mu-\epsilon, \mu+\epsilon)}(\mathcal{C}_i)$ with respect to the L^2 inner product defined by (2.3). Passing to a subsequence we may take limits of the basis elements to obtain an L^2 orthonormal set of elements in $R_\mu(\mathcal{C})$ whose cardinality is strictly bigger than $\dim R_{[\mu-\epsilon, \mu+\epsilon]}(\mathcal{C})$. This is a contradiction.

Given a homogeneous holomorphic function $f \in R_\mu(\mathcal{C})$, by the discussion in Section 2.1 for i large we may find a holomorphic function f_i on the unit ball $B_{\mathcal{C}_i}$ that naturally converges to f . Now using the cone structure on \mathcal{C}_i we may write $f_i = \sum_{\alpha} f_{i,\alpha}$, where

each $f_{i,\alpha}$ is homogeneous with respect to ξ_i with weight α . The decomposition is L^2 orthogonal so in particular each term $f_{i,\alpha}$ has uniformly bounded L^2 norm on B_i . We may write $f_i = u_i + v_i$, where u_i consists of homogeneous terms of degree in $[\mu - \frac{\epsilon}{2}, \mu + \frac{\epsilon}{2}]$, and v_i consists of homogeneous terms of degree not in $[\mu - \frac{\epsilon}{2}, \mu + \frac{\epsilon}{2}]$. It follows that $\|v_i\|_{L^2(B(O_i, \frac{3}{4}))}$ converges to 0 as $i \rightarrow \infty$. So we know u_i converges to f uniformly in the half size ball. Applying this construction to a basis of $R_\mu(\mathcal{C})$ would imply that for i large $\dim R_{(\mu-\epsilon, \mu+\epsilon)}(\mathcal{C}_i) \geq \dim R_\mu(\mathcal{C})$. Contradiction. \square

Now we suppose a sequence of cone limits $\mathcal{C}_i \in \mathfrak{C}$ converge to a cone limit \mathcal{C} . Our next goal is to realize this convergence algebraically. First of all, we choose $\underline{\mu} \notin \mathcal{S}(\mathcal{C})$ large so that $R(\mathcal{C})$ is generated by $R_{[0, \underline{\mu}]}(\mathcal{C})$. By Lemma 3.1 we may choose $\epsilon > 0$ small such that $\dim R_{[0, \underline{\mu}+\epsilon]}(\mathcal{C}_i) = \dim R_{[0, \underline{\mu}]}(\mathcal{C})$ for i large. Now we choose an L^2 orthonormal basis of $R_{[0, \underline{\mu}+\epsilon]}(\mathcal{C}_i)$. Passing to a subsequence we may assume that they converge to an L^2 orthonormal basis of $R_{[0, \underline{\mu}]}(\mathcal{C})$. Using these functions we obtain holomorphic maps $F_i : \mathcal{C}_i \rightarrow \mathbb{C}^N$ which converge naturally to $F : \mathcal{C} \rightarrow \mathbb{C}^N$ under the Gromov-Hausdorff convergence. By the choice of $\underline{\mu}$ we know F is an algebraic embedding onto a normal affine variety $W \subset \mathbb{C}^N$. It follows similar to [18] that for i large F_i is generically one-to-one onto a reduced affine variety $W_i \subset \mathbb{C}^N$. Hence F_i is the normalization map. In particular, it is an isomorphism if we know W_i is normal. Clearly W_i converges in the (local) Hausdorff sense to W .

By construction W_i is invariant under a linear vector field ξ_i on \mathbb{C}^N which converges to ξ as $i \rightarrow \infty$. It is easy to see that by suitable linear transformations on \mathbb{C}^N (i.e., by choosing a basis of $R_{[0, \underline{\mu}+\epsilon]}(\mathcal{C}_i)$ consisting of ξ_i -homogeneous functions) we may assume ξ and each ξ_i are in fact diagonalized with respect to the standard coordinates on \mathbb{C}^N .

Fix a Euclidean metric ω_0 on \mathbb{C}^N so that it induces a weighted L^2 inner product on the space of all polynomial functions:

$$(3.1) \quad \langle F_1, F_2 \rangle = \int_{\mathbb{C}^N} F_1 \overline{F_2} e^{-\frac{|z|^2}{2}} \omega_0^n.$$

Now we set some notations. We say a holomorphic polynomial function $f \in \mathbb{S}$ is ξ_i (reps. ξ)-homogeneous if $\mathcal{L}_{\xi_i} f = \sqrt{-1}\mu f$ (resp. $\mathcal{L}_\xi f = \sqrt{-1}\mu f$) for some $\mu \geq 0$; in this case we denote $\deg_{\xi_i}(f) = \mu$ (resp. $\deg_\xi(f) = \mu$). Clearly a monomial is both ξ -homogeneous and ξ_i -homogeneous. Given a polynomial f , its weighted degree with respect to ξ_i (resp. ξ) is defined in the obvious way using the weighted degree on monomials in terms of \deg_{ξ_i} (resp. \deg_ξ). We denote by $\mathbb{S}_{i,\mu}$ (resp. \mathbb{S}_μ) the space of polynomial functions on \mathbb{C}^N whose weighted degree with respect to ξ_i (resp. ξ) is at most μ , by \mathcal{I}_i (resp. \mathcal{I}) the ideal of \mathbb{S} consisting of polynomial functions on \mathbb{C}^N which vanish on W_i (resp. W) and by $\mathcal{I}_{i,\mu}$ (resp. \mathcal{I}_μ) the finite dimensional subspace of \mathcal{I}_i (resp. \mathcal{I}) consisting of those polynomials f with weighted degree at most μ with respect to ξ_i (resp. ξ).

Let $\mu_0 = \mu_0(\mathcal{C})$ be given in Section 2.2. Choose $\mu_1 > \max(\mu, \mu_0)$ sufficiently large and $\epsilon > 0$ small so that \mathcal{I}_{μ_1} generates \mathcal{I} , $\mathcal{I}_{\mu_1} = \mathcal{I}_{\mu_1+\epsilon}$ and $\dim \mathbb{S}_{i,\mu_1+\epsilon} = \dim \mathbb{S}_{\mu_1+\epsilon} = \dim \mathbb{S}_{\mu_1}$.

Lemma 3.2. *For i large we have $\dim \mathcal{I}_{i,\mu_1+\epsilon} = \dim \mathcal{I}_{\mu_1+\epsilon}$.*

Proof. To see this, on the one hand, for each i we can choose an L^2 orthonormal basis of $\mathcal{I}_{i,\mu_1+\epsilon}$ with respect to the inner product defined in (3.1). By passing to a subsequence we can take limits of the basis elements, which then implies that $\dim \mathcal{I}_{\mu_1+\epsilon} \geq \dim \mathcal{I}_{i,\mu_1+\epsilon}$ for i large. On the other hand, for each i we choose an L^2 orthonormal basis of $R_{[0,\mu_1+\epsilon]}(\mathcal{C}_i)$. Then by passing to a subsequence we can take limits under the Gromov-Hausdorff convergence and obtain an L^2 orthonormal set of holomorphic functions on \mathcal{C} . This implies that for i large $\dim R_{[0,\mu_1+\epsilon]}(\mathcal{C}) \geq \dim R_{[0,\mu_1+\epsilon]}(\mathcal{C}_i)$. But we have

$$\dim R_{[0,\mu_1+\epsilon]}(\mathcal{C}_i) \geq \dim \mathbb{S}_{i,\mu_1+\epsilon} - \dim \mathcal{I}_{i,\mu_1+\epsilon},$$

and

$$\dim R_{[0,\mu_1+\epsilon]}(\mathcal{C}) = \dim \mathbb{S}_{\mu_1+\epsilon} - \dim \mathcal{I}_{\mu_1+\epsilon}.$$

So we see the desired identity must hold. □

As a consequence we can choose a basis $\{f_{i,a}\}$ of $\mathcal{I}_{i,\mu_1+\epsilon}$ such that each $f_{i,a}$ is ξ_i -homogeneous and as $i \rightarrow \infty$, they converge to a basis $\{f_a\}$ of \mathcal{I}_{μ_1} such that each f_a is ξ -homogeneous. By the choice of μ_1 we know the latter generates the ideal \mathcal{I} .

Denote by \mathcal{I}'_i the ideal of \mathbb{S} generated by $\mathcal{I}_{i,\mu_1+\epsilon}$. Let W'_i be the affine scheme in \mathbb{C}^N defined by \mathcal{I}'_i . Since W_i is ξ_i invariant we see that $\mathcal{I}_{i+\epsilon}$ is spanned by ξ_i -homogeneous elements, so \mathcal{I}'_i is ξ_i -homogeneous. In particular W'_i is also ξ_i invariant. A priori W'_i could be strictly bigger than W_i as a scheme, i.e., \mathcal{I}'_i might be strictly contained in \mathcal{I}_i .

Lemma 3.3. *For i large W'_i is ξ invariant and W is ξ_i invariant.*

Proof. Since each $f_{i,a}$ is ξ_i -homogeneous and $f_{i,a}$ converges to f_a as $i \rightarrow \infty$, f_a must also be ξ_i -homogeneous for i large. Thus W is ξ_i invariant.

To show W_i is ξ invariant for i is large we argue by contradiction. Suppose not, without loss of generality we may assume that for some fixed a and for all i large, $f_{i,a}$ is not ξ -homogeneous. Passing to a subsequence we may further assume that there is a fixed monomial $h \in \mathbb{S}_{\mu_1}$ which appears in the monomial expansion of $f_{i,a}$ for all i and such that $\deg_{\xi}(h) \neq \deg_{\xi}(f_a)$. Since $\xi_i \rightarrow \xi$, we see that for i large, $\deg_{\xi_i}(h) \neq \deg_{\xi_i}(f_a)$. This contradicts the fact that $f_{i,a}$ is ξ_i -homogeneous. \square

Denote by \mathbb{T} the compact torus in $GL(N; \mathbb{C})$ generated by ξ . Denote by **Hilb** the multi-graded Hilbert scheme defined by the Hilbert function $h_{\mathcal{C}}$ with respect to the \mathbb{T} action. Lemma 3.2 implies that for i large, $h_{\mathcal{T}'_i}(d) = h_{\mathcal{C}}(d)$ for all $d \leq \mu_1$. Then by the result of Haiman-Sturmfels [20] (as reviewed in Section 2.2) we see W'_i is also represented by a point $[W'_i]$ in **Hilb**. In particular, $h_{\mathcal{T}'_i}(d) = h_{\mathcal{C}}(d)$ for all $d \geq 0$. Furthermore, $[W'_i] \rightarrow [\mathcal{C}]$ as $i \rightarrow \infty$. Since the universal family over **Hilb** is flat and normality is an open condition it follows that for i large W'_i is normal. In particular it is reduced and irreducible, but we have $\dim W_i = \dim W'_i$, so we conclude that $W'_i = W_i$. It follows that F_i must be an isomorphism.

Proposition 3.4. *For i large we have $\xi_i = \xi$ and $\text{Vol}(\mathcal{C}_i) = \text{Vol}(\mathcal{C})$.*

Proof. It suffices to show that given any subsequence of $\{i\}$, by passing to a further subsequence for i large it holds that $\xi_i = \xi$ and $\text{Vol}(\mathcal{C}_i) = \text{Vol}(\mathcal{C})$. Without loss of generality we may assume the subsequence is the original sequence. First by the openness of rational singularities [19] we may assume W_i has rational singularities for $i \geq i_0$. We take the set of points $P = \{[W_i] | i \geq i_0\} \in \mathbf{Hilb}$ and take its Zariski closure \mathcal{Z} . Denote by $\pi : \mathcal{W} \rightarrow \mathcal{Z}$ the restriction of the flat universal family. Since each affine cone by definition contains the origin, we have an obvious section σ of π . By [14] we can find an integer $l > 0$ and a Zariski open $\mathcal{O} \subset \mathcal{Z}$ such that for any $t \in \mathcal{O}$, the corresponding affine cone $\mathcal{W}_t = \pi^{-1}(t)$ is normal with rational singularities and $K^l_{\mathcal{W}_t}$ is trivial. Moreover, we may take a holomorphic section Ω of $K^l_{\mathcal{W}/\mathcal{O}}$ over a neighborhood of σ . Using the \mathbb{T} action we can as in Section 2.2 modify Ω_t simultaneously for all $t \in \mathcal{O}$ so that $\mathcal{L}_{\xi} \Omega_t = \sqrt{-1} c_t \Omega_t$, where c_t is a real-valued function which depends holomorphically on t . It follows that c_t must be a constant in t . So we have $c_t = c_0 = nl$.

This implies that passing to a subsequence for i large we know both ξ and ξ_i belong to the same hyperplane $\mathbb{H} = \mathbb{H}_i \subset \text{Lie}(\mathbb{T})$ (as defined in (2.5)). On the other hand by the properties of the volume function listed in Section 2.2 we see $\mathbb{V}_i = \mathbb{V}$. Since \mathbb{V} is strictly convex and both ξ and ξ_i are critical points of $\mathbb{V}|_{\mathbb{H}}$, we conclude that $\xi = \xi_i$. Consequently, we also have $\text{Vol}(\mathcal{C}_i) = \text{Vol}(\mathcal{C})$. \square

By a simple contradiction argument the above discussion also gives

Corollary 3.5. *There exists $\underline{\epsilon} > 0$ depending only on n and D such that for a given cone limit $\mathcal{C} \in \mathfrak{C}$, all the cone limits $\mathcal{C}' \in \mathfrak{C}$ with $d_{EGH}(B_{\mathcal{C}}, B_{\mathcal{C}'}) < \underline{\epsilon}$ can be parametrized by a connected component in a multi-graded Hilbert scheme **Hilb** of \mathbb{C}^N such that the Reeb vector field of \mathcal{C}' is given by the restriction of a common diagonal linear vector field ξ on \mathbb{C}^N .*

Proof of Theorem 1.7. Suppose otherwise, then we can find a sequence $Z_i \in \mathfrak{R}$ such that $\text{Vol}_{\infty}(Z_i)$ are all distinct. Let \mathcal{C}_i be the unique asymptotic cone of Z_i . Then

$\text{Vol}(\mathcal{C}_i) = \text{Vol}_\infty(Z_i)$. Passing to a subsequence we may assume \mathcal{C}_i converges to a limit \mathcal{C} . Then we get a contradiction with Proposition 3.4.

From the argument we can also see that the bound only depends on D , which guarantees the non-collapsing property and Gromov-Hausdorff compactness of the relevant cones. □

Remark 3.6. *The above arguments work as long as [18] can be applied so may be useful in more general setting. For example, Theorem 1.7 holds for general Ricci-flat Kähler cones with isolated singularities or with orbifold singularities along rays, without assuming that they arise as bubble limits. We leave this direction for interested readers to explore further.*

4. EXISTENCE OF MINIMAL BUBBLES

We adopt the notation in the introduction. Let $(X_j, L_j, \omega_j, p_j)$ be a sequence of polarized Kähler-Einstein manifolds converging to $(X_\infty, \omega_\infty, p_\infty)$. By [17] the convergence can be understood algebro-geometrically in the flat universal family over a fixed Hilbert scheme of \mathbb{P}^M . As a consequence there is a neighborhood U of p_∞ in \mathbb{P}^M such that $X_\infty \cap U$ is cut out by finitely many holomorphic functions $\{h^a\}_{a=1}^m$ on U and for j large $X_j \cap U$ is cut out by $\{h_j^a\}_{a=1}^m$. Moreover, for each a , h_j^a converges uniformly to h^a over U as $j \rightarrow \infty$.

Let $\mathcal{C}_0(X_\infty)$ be the unique tangent cone at p_∞ , and denote $V \equiv \text{Vol}_0(X_\infty)$, $\mathcal{S} \equiv \mathcal{S}(\mathcal{C}_0(X_\infty))$. Let $\mu_1 = \mu_1(\mathcal{C}_0(X_\infty))$ be given in Lemma 3.2 and fix $\epsilon > 0$ such that

$$[V - \epsilon, V + \epsilon] \cap \mathcal{V}(\mathfrak{R}) = \{V\}.$$

Choose $\mu_2 \geq \mu_1$ such that the graded ring $R(p_\infty)$ is generated by functions in degrees at most μ_2 . For $\mu \geq \mu_2$, we choose a finite set $\{f^\alpha\}_{\alpha=0}^N \subset \mathcal{O}(p_\infty)$ which descends to a basis of $R_{[0, \mu]}(p_\infty)$ and such that $\deg(f^\alpha) \leq \deg(f^\beta)$ whenever $\alpha \leq \beta$. Notice f^0 is a constant. Choose an extension of f^α to a holomorphic function \hat{f}^α on a neighborhood of p_∞ in \mathbb{P}^M . We may fix $\mu_* > \mu_2$ large so that $\{\hat{f}^\alpha\}_{\alpha \geq 1}$ define a holomorphic embedding of a neighborhood $U \subset \mathbb{P}^M$. For j large we denote by f_j^α the restriction of \hat{f}^α to $U \cap X_j$. Clearly, as $j \rightarrow \infty$, f_j^α converges naturally to f^α under the convergence $X_j \rightarrow X_\infty$.

Denote

$$\delta = \inf\{|\mu - \nu| \mid \mu, \nu \in \mathcal{S} \cap [0, \mu_*], \mu \neq \nu\}$$

For $\lambda > 0$ denote by $X_{j, \lambda}$ the rescaled space $(X_j, p_j, \lambda^2 \omega_j)$ and by $B_{j, \lambda}$ the unit ball in $X_{j, \lambda}$ centered at p_j . We denote

$$\underline{\text{Vol}}(B_{j, \lambda}) = \frac{\text{Vol}(B_{j, \lambda})}{\text{Vol}(B_0(1))},$$

where $B_0(r)$ denotes the radius r ball in the Euclidean space \mathbb{C}^n .

From the Bishop-Gromov monotonicity and Colding's volume convergence theorem we see that there exists a $\underline{\lambda} > 0$ such that $\underline{\text{Vol}}(B_{j, \underline{\lambda}}) \geq V - \epsilon/4$ for all j large. Denote by $\bar{\lambda}_j$ the largest λ such that $\underline{\text{Vol}}(B_{j, \lambda}) - V \leq \epsilon/4$. Notice $\bar{\lambda}_j \rightarrow \infty$ as $j \rightarrow \infty$. We may assume $\underline{\lambda}$ is large so that $B_{j, \underline{\lambda}} \subset U$ for all j large.

Lemma 4.1. *Given a sequence $\lambda_j \in [\underline{\lambda}, \bar{\lambda}_j]$ with $\lambda_j \rightarrow \infty$ and $\lambda_j^{-1}\bar{\lambda}_j \rightarrow \infty$, any subsequential limit of X_{j,λ_j} is a cone limit.*

Proof. Passing to a subsequence we may take a pointed Gromov-Hausdorff limit Z . The assumptions imply that the asymptotic cone $\mathcal{C}_\infty(Z)$ and the tangent cone $\mathcal{C}_0(Z)$ at p_Z both have volume ϵ -close to V . It follows from our choice of ϵ that $\text{Vol}(\mathcal{C}_\infty(Z)) = \text{Vol}(\mathcal{C}_0(Z)) = V$. Hence Z itself is a cone limit. \square

Denote by \mathfrak{C}' the class of all cone limits arising from the above lemma. Notice we have $\mathcal{C}_0(X_\infty) \in \mathfrak{C}'$.

Lemma 4.2. *All the cone limits in \mathfrak{C}' can be parametrized by a fixed multi-graded Hilbert scheme \mathbf{Hilb} of \mathbb{C}^N such that the Reeb vector fields are given by the restriction of a common diagonal linear vector field ξ on \mathbb{C}^N . In particular, all the cone limits in \mathfrak{C}' have the same holomorphic spectrum which coincides with \mathcal{S} and have volume density V .*

Proof. We suppose otherwise, then by Lemma 4.1 after passing to a subsequence we may find scales $\underline{\lambda} < \lambda'_j < \lambda''_j < \bar{\lambda}_j$ such that $X_{j,\lambda'_j} \rightarrow \mathcal{C}'$ and $X_{j,\lambda''_j} \rightarrow \mathcal{C}''$ but \mathcal{C}' and \mathcal{C}'' do not belong to the same multi-graded Hilbert scheme. Now by a simple contradiction argument we can see that for all j large and all $\lambda \in [\lambda'_j, \lambda''_j]$ there is a $\mathcal{C}_{j,\lambda} \in \mathfrak{C}'$ with $d_{EGH}(B_{j,\lambda}, B_{\mathcal{C}_{j,\lambda}}) \leq \epsilon/10$ such that $\mathcal{C}_{j,\lambda'_j} = \mathcal{C}'$ and $\mathcal{C}_{j,\lambda''_j} = \mathcal{C}''$, where ϵ is given in Corollary 3.5. Then it follows easily that for all $\lambda \in [\lambda'_j, \lambda''_j]$ we can find $\delta_\lambda > 0$ such that $d_{EGH}(B_{\mathcal{C}_{j,\beta}}, B_{\mathcal{C}_{j,\lambda}}) < \epsilon$ for all $\beta \in [\lambda - \delta_\lambda, \lambda + \delta_\lambda]$. So by Corollary 3.5 we see that for all $\beta \in [\lambda - \delta_\lambda, \lambda + \delta_\lambda]$, $\mathcal{C}_{j,\beta}$ is parametrized in a fixed multi-graded Hilbert scheme of \mathbb{C}^N with the same Reeb vector field. Since $[\lambda'_j, \lambda''_j]$ can be covered by finitely many such intervals, we obtain a contradiction. \square

The following is the main result of this section.

Theorem 4.3. $\mathcal{C}_0(X_\infty)$ is an isolated point in \mathfrak{C}' .

Proof of Theorem 1.10. We first prove that the subsequential limits of $X_{j,\bar{\lambda}_j}$ are minimal bubbles. Let Z be such a subsequential limit. By our choice of $\bar{\lambda}_j$ and Bishop-Gromov monotonicity we know $\text{Vol}_\infty(Z) < \text{Vol}_0(Z)$, so Z is not a cone limit, so it suffices to show $\mathcal{C}_\infty(Z) = \mathcal{C}_0(X_\infty)$. Suppose otherwise, notice $\mathcal{C}_0(X_\infty)$ is the bubble limit associated to a sequence of scales $\lambda_j \in (\underline{\lambda}, \bar{\lambda}_j)$, so for each integer $k \gg 0$ and j large we may find $\lambda_{j,k} \in [\underline{\lambda}, \bar{\lambda}_j]$ such that $d_{EGH}(B_{j,\lambda_{j,k}}, B_{\mathcal{C}_0(X_\infty)}) = k^{-1}$. Taking a subsequential limit we obtain a cone limit \mathcal{C}_k with $d_{EGH}(B_{\mathcal{C}_k}, B_{\mathcal{C}_0(X_\infty)}) = k^{-1}$. This contradicts Theorem 4.3.

The second statement can be proved similarly. \square

The rest of this section is devoted to the proof of Theorem 4.3. First we need to slightly extend the convexity result in [18]. Define

$$\underline{k} \equiv \log_2(\sigma \underline{\lambda}), \quad \bar{k}_j \equiv \log_2(\sigma^{-1} \bar{\lambda}_j).$$

The proof of the lemma below follows from exactly the same contradiction argument as in [18], Proposition 3.7. So we omit the proof.

Lemma 4.4. *Given any fixed $\mu \in (0, \infty) \setminus \mathcal{S}$, there exist a $\underline{j} \geq 0$ and $\sigma > 0$ such that for all $j \geq \underline{j}$ and $k \in [\underline{k}, \bar{k}_j]$, if a holomorphic function f defined on $B_{j,2^k}$ satisfies $\|f\|_{B_{j,2^{k+1}}} \geq 2^{-\mu} \|f\|_{B_{j,2^k}}$, then $\|f\|_{B_{j,2^{k+2}}} > 2^{-\mu} \|f\|_{B_{j,2^{k+1}}}$, where $\|\cdot\|$ denotes the natural L^2 norm.*

From the uniform convergence of f_j^α to f^α that we can choose $\underline{k} > 0$ such that

$$\|f_j^\alpha\|_{B_{j,2^{\underline{k}+1}}} \geq 2^{-\deg(f^\alpha) - \frac{\delta}{2}} \|f_j^\alpha\|_{B_{j,2^{\underline{k}}}}$$

for all α and for all j large. It follows from the lemma that

$$\|f_j^\alpha\|_{B_{j,2^{k+1}}} \geq 2^{-\deg(f^\alpha) - \frac{\delta}{2}} \|f_j^\alpha\|_{B_{j,2^k}}$$

for all $k \in [\underline{k}, \bar{k}_j]$ when j is large. On $B_{j,k}$ we define

$$\tilde{f}_{j,k}^\alpha = \frac{f_j^\alpha}{\|f_j^\alpha\|_{B_{j,2^k}}}.$$

Using Lemma 4.4 and the arguments in [18] we see that when $j, k \rightarrow \infty$, passing to a subsequence each $\tilde{f}_{j,k}^\alpha$ converges to a non-trivial limit holomorphic function f_∞^α on some limit $Z \in \mathfrak{R}$. Moreover, we have $\deg(f_\infty^\alpha) \leq \deg(f^\alpha)$.

Now for a holomorphic function on $B_{j,2^k}$ we denote by $\Pi_{j,k}(f)$ the L^2 orthogonal projection of f to the orthogonal complement of f^1 , then normalized so that $\|\Pi_{j,k}f\|_{B_{j,2^k}} = 1$. The following Lemma can be proved similarly to [18], Proposition 3.11. Again we omit the proof here.

Lemma 4.5. *Given any $\mu \in (0, \infty) \setminus \mathcal{S}$, by possibly enlarging \underline{j} and σ , we have that for all $j \geq \underline{j}$ and $k \in [\underline{k}, \bar{k}_j]$, if f is a holomorphic function f on $B_{j,2^k}$ such that $\|\Pi_{j,k+1}f\|_{B_{j,2^{k+1}}} \geq 2^{-\mu} \|\Pi_{j,k}f\|_{B_{j,2^k}}$, then $\|\Pi_{j,k+2}f\|_{B_{j,2^{k+2}}} > 2^{-\mu} \|\Pi_{j,k+1}f\|_{B_{j,2^{k+1}}}$.*

Given this Lemma, as in [18, 8] arguing by induction and by possibly enlarging \underline{j} and \underline{k} again, we may find functions $\{\tilde{f}_{j,k}^\alpha\}$ ($0 \leq \alpha \leq N$, $j \geq \underline{j}$ and $k \in [\underline{k}, \bar{k}_j]$) such that

- $\tilde{f}_{j,k}^\alpha = \text{const}$ when $\alpha = 0$;
- $\tilde{f}_{j,k}^\alpha$ belongs to the linear span of f_j^β for $\beta \leq \alpha$;
- $\{\tilde{f}_{j,k}^\alpha\}$ is L^2 orthonormal over $B_{j,2^k}$;
- as $j, k \rightarrow \infty$, passing to a subsequence $\tilde{f}_{j,k}^\alpha$ converges uniformly to a limit f_∞^α on a limit $Z \in \mathfrak{R}$ with $\deg(f_\infty^\alpha) \leq \deg(f^\alpha)$.

In particular, $\{f_\infty^\alpha\}$ are L^2 orthonormal on the unit ball $B(p_Z, 1)$ in Z . We denote by $R_{\leq \mu}$ the space of holomorphic function on Z with degree at most μ .

Lemma 4.6. *For all α we have $\deg(f_\infty^\alpha) = \deg(f^\alpha)$ and $R_{\leq \mu_*}(Z)$ is precisely spanned by $\{f_\infty^\alpha\}$. Moreover, if Z is a cone limit, then $\{f_\infty^\alpha\}$ is a homogeneous L^2 orthonormal basis.*

Proof. Since $\deg(f_\infty^\alpha) \leq \deg(f^\alpha)$ for all α , it follows that $\dim R_{\leq \mu}(Z) \geq \dim R_{[0, \mu]}(p_\infty)$ for all $\mu \leq \mu_*$. But since the asymptotic cone $\mathcal{C}_\infty(Z)$ belongs to **Hilb** we must have

$h_{\mathcal{C}_\infty(Z)} = h_{\mathcal{C}_0(X_\infty)}$, so $\dim R_{\leq \mu}(Z) = \dim R_{[0, \mu]}(\mathcal{C}_\infty(Z)) = \dim R_{[0, \mu]}(p_\infty)$. It then follows from an easy induction argument that $\deg(f_\infty^\alpha) = \deg(f^\alpha)$ for all α , hence $R_{\leq \mu_*}(Z)$ is spanned by $\{f_\infty^\alpha\}$.

Now we assume Z is a cone limit. Lemma 4.4 and Lemma 4.5 imply by induction that for all $k \in [\underline{k}, \overline{k}_j]$,

$$\log_2 \frac{\|\tilde{f}_{j,k+1}^\alpha\|_{B_{j,2^{k+1}}}}{\|\tilde{f}_{j,k}^\alpha\|_{B_{j,2^k}}} \leq -\deg(f^\alpha).$$

But we also have that for given k ,

$$\lim_{j \rightarrow \infty} \log_2 \frac{\|\tilde{f}_{j,k+1}^\alpha\|_{B_{j,2^{k+1}}}}{\|\tilde{f}_{j,k}^\alpha\|_{B_{j,2^k}}} = \lim_{j \rightarrow \infty} \log_2 \frac{\|\tilde{f}_{k+1}^\alpha\|_{B_{j,2^{k+1}}}}{\|\tilde{f}_k^\alpha\|_{B_{j,2^k}}} \geq -\deg(f^\alpha) - \epsilon_k$$

for $\epsilon_k \rightarrow 0$. It follows that if $k_j \rightarrow \infty$, then

$$\lim_{j \rightarrow \infty} \log_2 \frac{\|\tilde{f}_{j,k_j+2}^\alpha\|_{B_{j,2^{k_j+2}}}}{\|\tilde{f}_{j,k_j+1}^\alpha\|_{B_{j,2^{k_j+1}}}} = \lim_{j \rightarrow \infty} \log_2 \frac{\|\tilde{f}_{j,k_j+1}^\alpha\|_{B_{j,2^{k_j+1}}}}{\|\tilde{f}_{j,k_j}^\alpha\|_{B_{j,2^{k_j}}}} = -\deg(f^\alpha).$$

This easily implies that the limit is homogeneous. \square

Now as in [18] we can do a further perturbation to the above set $\tilde{f}_{j,k}^\alpha$. In particular we will replace $\tilde{f}_{j,k}^\alpha$ by $\tilde{f}_{j,k}^\alpha + \sum_{\beta} \epsilon_{\alpha\beta,jk} \tilde{f}_{j,k}^\beta$, where $\epsilon_{\alpha\beta,jk} \rightarrow 0$ as $j, k \rightarrow \infty, k^{-1}\overline{k}_j \rightarrow \infty$ and $\epsilon_{\alpha\beta,jk} = 0$ unless $\beta < \alpha$ and $\deg(f^\beta) = \deg(f^\alpha)$, such that

$$\Lambda_{j,k} \tilde{f}_{j,k}^\alpha = 2^{-\mu_{j,k}} \tilde{f}_{j,k+1}^\alpha + \sum_{\beta < \alpha, \deg(f^\beta) = \deg(f^\alpha)} \tau_{\alpha\beta,jk} \tilde{f}_{j,k+1}^\beta,$$

where $\mu_{j,k} \rightarrow \deg(f^\alpha)$ and $\tau_{\alpha\beta,jk} \rightarrow 0$ as $j, k \rightarrow \infty$ and $k^{-1}\overline{k}_j \rightarrow \infty$. The proof is similar to that of Proposition 3.12 and Lemma 3.13 in [18]. We omit the details here.

Using the functions $\{\tilde{f}_{j,k}^\alpha\}_{\alpha=1}^N$ we obtain holomorphic embeddings $\Phi_{j,k} : B_{j,2^k} \rightarrow \mathbb{C}^N$. Notice there is a natural inclusion $B_{j,2^k} \hookrightarrow B_{j,2^{k'}}$ whenever $k \geq k'$. By construction we have $\Phi_{j,k+1} = \Lambda_{j,k} \circ \Phi_{j,k}|_{B_{j,2^{k+1}}}$, where $\Lambda_{j,k}$ is a linear transformation of \mathbb{C}^N that preserves the Reeb vector field ξ (viewed as a linear vector field on \mathbb{C}^N) and satisfies $\Lambda_{j,k} = \Lambda(\text{Id} + \tau_{j,k})$ for a diagonal linear transformation $\Lambda = \text{diag}(2^{\zeta_1}, \dots, 2^{\zeta_N})$ with $\tau_{j,k} \rightarrow 0$ as $j, k \rightarrow \infty$ and $k^{-1}\overline{k}_j \rightarrow \infty$. Furthermore, as $j, k \rightarrow \infty$ and $k^{-1}\overline{k}_j \rightarrow \infty$, passing to subsequences $\Phi_{j,k}$ converges to holomorphic embeddings of the cone limits into \mathbb{C}^N .

Let W be the weighted tangent cone in \mathbb{C}^N of X_∞ at p_∞ with respect to ξ . From [18] and the discussion above we know that W defines a point $[W]$ in **Hilb**. So its defining ideal \mathcal{I}_W is generated by finitely many homogeneous functions $\{G^b\}_{b=1}^l$. We list them in degree increasing order, so we have $\deg_\xi(G^b) = d_s$ for $b_{s-1} < b \leq b_s$, $b_0 = 0$ and $d_1 \leq d_2 \leq \dots \leq \mu_*$. It follows that X_∞ is cut out by $\{F^b\}_{b=1}^l$, where $F^b = G^b + H^b$ such that H^b is a local holomorphic function on a neighborhood of

$0 \in \mathbb{C}^N$ with $\deg_\xi(H^b) > \deg_\xi(G^b)$. Here we use \deg_ξ to denote the weighted vanishing order at 0 of a local holomorphic function on \mathbb{C}^N .

Now by the discussion at the beginning of this section and the construction of $\Phi_{j,k}$ it is not hard to see that the image of $\Phi_{j,k}$ is cut out by finitely many local holomorphic functions $\{F_j^b\}_{b=1}^l$ such that for each b , F_j^b converges locally uniformly to F^b . Write

$$F_j^b = F^b + S_j^b + H_j^b + L_j^b,$$

where $\deg_\xi(L_j^b) < \deg_\xi(S_j^b) = \deg_\xi(F^b) < \deg_\xi(H_j^b)$ and S_j^b, H_j^b, L_j^b all converge uniformly to 0 as $j \rightarrow \infty$. Denote $G_j^b = G^b + S_j^b$ and $Q_j^b = G_j^b + L_j^b$. For $k \in [\underline{k}, \bar{k}_j]$, the image of $\Phi_{j,k}$ is cut out by $\{F_{j,k}^b = 0\}$, where $F_{j,k}^b = \Gamma_{j,k} \cdot F_j^b$ for $\Gamma_{j,k} = \Lambda_{j,k} \cdots \Lambda_{j,\underline{k}+1}$. Denote $G_{j,k}^b = \Gamma_{j,k} \cdot G_j^b$.

Now we try to take limits of the defining functions $\{F_{j,k}^b\}_{b=1}^l$ as $j, k \rightarrow \infty$. We denote by \mathbb{S}_μ the linear space of all ξ -homogeneous polynomials on \mathbb{C}^N with ξ -degree equal to μ . We have a natural inner product on \mathbb{S}_μ such that an orthonormal basis is given by the set of monomials with coefficient 1. Denote by $\|\cdot\|_\mu$ the corresponding norm on \mathbb{S}_μ . By abusing notation we also denote by $\|\cdot\|_\mu$ the induced operator norm on the space of linear transformations on \mathbb{S}_μ . We also have an induced inner product on the direct sum $\mathbb{S}_{\leq \mu_*} \equiv \bigoplus_{\mu \leq \mu_*} \mathbb{S}_\mu$. We denote by $\|\cdot\|_\bullet$ the induced norm. By a linear transformation we

may assume $\{G_j^b\}$ is orthonormal in $\mathbb{S}_{\leq \mu_*}$ with respect to this inner product.

Notice Λ and $\tau_{j,k}$ naturally act on \mathbb{S}_μ for all μ . It is also easy to check that there exists $C > 0$ such that for all μ , $\|\Lambda\|_\mu \leq C2^{-\mu}$, and

$$(4.1) \quad \|\tau_{j,k}\|_\mu \leq C(1 + \|\tau_{j,k}\|_1)^\mu.$$

Since as $j, k \rightarrow \infty$, $k^{-1}\bar{k}_j \rightarrow \infty$ we have $\|\tau_{j,k}\| \rightarrow 0$, it is easy to see that for any b , the subsequential limit of $F_{j,k}^b / \|Q_{j,k}^b\|_\bullet$ coincides with the limit of $Q_{j,k}^b / \|Q_{j,k}^b\|_\bullet$, which is a polynomial of degree at most $\deg_\xi(G^b)$.

We first consider the case $b \leq b_1$. It is clear that the subsequential limits of each $F_{j,k}^b / \|Q_{j,k}^b\|_\bullet$ must be homogeneous of degree d_1 . But the limits for different b 's may coincide. As usual we need to perform the Gram-Schmidt transform \mathcal{L}_j to make an orthonormal basis $\{\tilde{Q}_{j,k}^b\}$ out of the linear span of $\{Q_{j,k}^b\}$. Applying the same \mathcal{L}_j to $\{F_{j,k}^b\}$ and $\{G_{j,k}^b\}$ we get $\{\tilde{F}_{j,k}^b\}$ and $\{\tilde{G}_{j,k}^b\}$. Notice in fact $G_{j,k}^b = Q_{j,k}^b$ by the minimality of d_1 .

Lemma 4.7. *As $j, k \rightarrow \infty$, subsequential limits of $\{\tilde{F}_{j,k}^b\}_{b=1}^{b_1}$ are polynomials on Z with degree d_1 . If Z is a cone limit, then the limits are homogeneous.*

Proof. It is easy to check using (4.1) that in the Taylor expansion of $\tilde{F}_{j,k}^b$, the degree higher than d_1 terms will not survive as $j, k \rightarrow \infty$. So $\tilde{F}_{j,k}^b$ converges to a polynomial \tilde{Q}_∞^b of degree at most d_1 , which is the same as the limit of $\tilde{Q}_{j,k}^b$. When Z is a cone limit, then by the minimality of d_1 we know $\tilde{Q}_{j,k}^b$ must be homogeneous. \square

In particular, since $\mathcal{C}_\infty(Z) \in \mathbf{Hilb}$, these limit functions $\{\tilde{Q}_\infty^b\}_{b=1}^{b_1}$ precisely span $\mathcal{I}_{\leq d_1}(Z)$.

Next we consider the case $b \leq b_2$. We argue as above to extend \mathcal{L}_j to the linear span of $\{Q_{j,k}^b\}_{b=1}^{b_2}$. This time we can only say that the subsequential limits of $\{\tilde{F}_{j,k}^b\}_{b=b_1+1}^{b_2}$ have degree at most d_2 . Also these limits agree with the limits of $\{\tilde{Q}_{j,k}^b\}_{b=b_1+1}^{b_2}$. But from the above discussion we know that for $b_1 < b \leq b_2$, the limit \tilde{Q}_∞^b is orthonormal to $\mathcal{I}_{\leq d_1}(Z)$ (with respect to the L^2 inner product over the unit ball $B(p_Z, 1)$), so must have degree exactly d_2 by dimension counting. Again if Z is a cone limit the limit must be homogeneous; in this case \tilde{Q}_∞^b agrees with the limit \tilde{G}_∞^b of $\tilde{G}_{j,k}^b$ and all such limits for $1 \leq b \leq b_2$ precisely span $\mathcal{I}_{\leq d_2}(Z)$.

By a straightforward induction argument, we then obtain linear transformations \mathcal{L}_j on the linear span of $\{Q_{j,k}^b\}_{b=1}^l$, such that $\{\mathcal{L}_j \cdot F_{j,k}^b\}_{b=1}^l$ and $\{\mathcal{L}_j \cdot Q_{j,k}^b\}_{b=1}^l$ both converge to an orthonormal basis $\{\tilde{Q}_\infty^b\}_{b=1}^l$ of $\mathcal{I}_{\leq \mu^*}(Z)$ (with respect to the L^2 inner product over the unit ball $B(p_Z, 1)$), and $\{\mathcal{L}_j \cdot G_{j,k}^b\}_{b=1}^l$ converge to a linear independent set $\{\tilde{G}_\infty^b\}_{b=1}^l$ of \mathbb{S} . If Z is a cone limit, then each \tilde{Q}_∞^b is homogeneous and coincides with \tilde{G}_∞^b .

Now we define a sequence of affine schemes $\{\mathcal{W}_j\}$ in \mathbb{C}^N using the ideal generated by $\{G_j^b\}_{b=1}^l$. Then the above discussion implies that $h_{\mathcal{W}_j} = h_{\mathcal{C}_0(X_\infty)}$. In particular \mathcal{W}_j defines a point $[\mathcal{W}_j] \in \mathbf{Hilb}$. For any cone limit $\mathcal{C} \in \mathfrak{C}'$ which corresponds to a subsequence $k_j \rightarrow \infty$, it then follows that $\Gamma_{j,k_j} \cdot [\mathcal{W}_j]$ converges to $[\mathcal{C}]$ as $j \rightarrow \infty$.

Remark 4.8. *When \mathcal{C} is the tangent cone $\mathcal{C}_0(X_\infty)$ at p_∞ , the arguments above reduce to that of [18]. In that case \mathcal{W}_j is isomorphic to W for all j .*

As reviewed in Section 2.2, we have the Matsushima theorem which implies that for each $\mathcal{C} \in \mathfrak{C}'$, the corresponding element $[\mathcal{C}] \in \mathbf{Hilb}$ has a reductive stabilizer group in $GL_\xi(N; \mathbb{C})$. We also have the Bando-Mabuchi uniqueness theorem, therefore Theorem 4.3 is a consequence of the following

Proposition 4.9. *Let V be a finite dimensional complex representation of a reductive group G and let $\{x_j\}$ be a sequence of points in $\mathbb{P}(V)$ converging to a limit x_∞ . Let $\{I_j\}$ be a sequence of closed intervals in \mathbb{R} and $g_j : I_j \rightarrow G$ be a sequence of continuous maps. Denote by \mathcal{Y} the set of all subsequential limits of $g_j(t_j) \cdot x_j$ ($g_j \in G, t_j \in I_j$) as $j \rightarrow \infty$. Suppose that for all $y \in \mathcal{Y}$, the stabilizer group of y is reductive, then \mathcal{Y} is contained in a single G orbit.*

Proof. It is not hard to see we only need to prove that for any given subsequence of $\{j\}$ there is a further subsequence such that all the subsequential limits arising from this subsequence is contained in a single G orbit. Without loss of generality we may assume the given subsequence is the original sequence.

Clearly \mathcal{Y} is compact. Choose $y_0 \in \mathcal{Y}$ such that its stabilizer group has minimal dimension among all points in \mathcal{Y} . Without loss of generality we may assume $0 \in I_j$, $g_j(0) = \text{Id}$ and $y_0 = \lim_{j \rightarrow \infty} x_j$. Denote by G_0 the stabilizer group of y_0 . By assumption G_0 is reductive.

Applying the slice theorem at y (see for example [15]) we may find a G_0 -invariant projective subspace $\mathbb{P}(V') \subset \mathbb{P}(V)$ containing y , which is transversal to $G \cdot y$ at y and has the property that for any $y' \in \mathbb{P}(V')$ close to y , locally the intersection of $G \cdot y'$ with $\mathbb{P}(V')$ coincides with the G_0 orbit of y' and furthermore, the identity component

of the stabilizer group of y' in G is contained in G_0 . As a consequence, we can find a neighborhood \mathcal{N} of y_0 in $\mathbb{P}(V)$ such that the following hold

- any $y \in \mathcal{Y} \cap \mathbb{P}(V') \cap \mathcal{N}$ is the limit of points in $G.x_j \cap \mathcal{N} \cap \mathbb{P}(V')$;
- for any $z \in U$, $G.z \cap \mathcal{N} \cap \mathbb{P}(V') = \bigsqcup_{\alpha=1}^{m_z} G_0.z_\alpha$ for $z_\alpha \in \mathbb{P}(V') \cap \mathcal{N}$.

Since the degrees of all the G orbits in $\mathbb{P}(V)$ are uniformly bounded, one can find $m > 0$

such that $m_z \leq m$ for all $z \in \mathcal{N}$. For j large we then have $G.x_j \cap \mathcal{N} \cap \mathbb{P}(V') = \bigsqcup_{\alpha=1}^{m_{x_j}} G_0.x_{j\alpha}$

for $x_{j\alpha} \in \mathbb{P}(V') \cap \mathcal{N}$. Passing to a subsequence (noticing this changes \mathcal{Y}) we may assume each $x_{j\alpha}$ converges to a limit $z_\alpha \in \mathbb{P}(V')$ as $j \rightarrow \infty$. Shrinking \mathcal{N} we can ensure that there is exactly one $z_\alpha \in \mathbb{P}(V') \cap \mathcal{N}$, which is given by y_0 .

Now we consider the decomposition as G_0 representations $V' = V_0 \oplus V_1$, where V_0 is the invariant subspace of G_0 . For j large we write $x_{j\alpha} = [v_{j\alpha}, \hat{v}_{j\alpha}]$, where $v_{j\alpha} \in V_0$ and $\hat{v}_{j\alpha} \in V_1$. By the choice of y_0 we see that any $y = [v, \hat{v}] \in \mathcal{Y} \cap \mathbb{P}(V') \cap \mathcal{N}$ must satisfy $\hat{v} = 0$. But then it follows that $y = z_\alpha$ since the action of G_0 on V_0 is trivial. Hence by the choice of \mathcal{N} we conclude that $y = y_0$.

The above argument shows that y_0 is an isolated point in \mathcal{Y} . Then using the fact that I_j is connected a simple continuity argument then implies that \mathcal{Y} is contained in a single G orbit, which must be the G orbit of x_∞ . □

Proof of Corollary 1.12. From Theorem 1.10 we can define $(Z_1, q_1) = (X_\infty, p_\infty)$, $(Z_2, q_2) = (\mathcal{C}_0(X_\infty), O)$. Let (Z_3, q_3) be the minimal bubble after passing to subsequence, so it is asymptotic to $\mathcal{C}_0(X_\infty)$. It is then given by the limit corresponding to the subsequence $\{\bar{\lambda}_j\}$. Denote by \tilde{Q}_∞^b the subsequential limits of $\mathcal{L}_j.F_{j,\bar{\lambda}_j}^b$. Then it follows that the ideal of Z in \mathbb{C}^N is exactly generated by $\{\tilde{Q}_\infty^b\}_{b=1}^l$. Then we can repeat the above discussion in a neighborhood of q_3 to obtain the next minimal bubble. By Theorem 1.7 we know this process stops in finitely many steps. The last bubble limit (Z_{2k}, q_{2k}) must then be flat. □

5. DISCUSSION

5.1. Algebro-geometric meaning. It is a natural question to study the algebro-geometric meaning of the minimal bubbles constructed in this paper. For our purposes we will call these *analytic minimal bubbles*. They are endowed with both algebraic and metric information.

Recall that given an affine variety Y , in [31] we defined the notion of a negative valuation ν on Y .

Definition 5.1. A polarized Calabi-Yau affine variety is an affine variety Y with log terminal singularities, which is endowed with a negative valuation ν whose associated graded ring $Gr_\nu(Y)$ is finitely generated and defines a Fano cone \mathcal{C}_ν (in the sense of [11]). The cone \mathcal{C}_ν is called the *algebraic asymptotic cone* of Y .

Remark 5.2. Notice that a Fano cone itself is by definition a polarized Calabi-Yau affine variety. From the discussion in [31] the notion of a polarized Calabi-Yau variety is naturally designed to study complete Calabi-Yau metrics with maximal volume growth.

Definition 5.3. A polarized affine Calabi-Yau variety (Y, ν) is K-semistable (K-polystable) if the associated Fano cone \mathcal{C}_ν is K-semistable (K-polystable).

From the discussion in Section 2.3 we know that an analytic minimal bubble Z is always a K-semistable polarized affine Calabi-Yau variety. Moreover, one expects it to be K-polystable: this is discovered and proved in [31] under the quadratic curvature decaying condition, and it is conjectured in [31] to be always the case in general. Notice that the weak Calabi-Yau metric on Z is not uniquely determined by the negative valuation; an explicit example is shown by Chiu [9] for complete Calabi-Yau metrics on \mathbb{C}^3 asymptotic to the flat cone $\mathbb{C}^2/\mathbb{Z}_3 \times \mathbb{C}$. Nevertheless one can still formulate a generalized uniqueness, see Conjecture 6.8 in [31], which ensures only a finite dimensional freedom.

One can ask whether the analytic minimal bubble depends only on the local complex-analytic/algebraic-geometric information of the sequence (X_j, L_j, p_j) . One can try to ask a uniqueness question in this general context but there is a complication related to the fact that analytic minimal bubbles are sensitive to the sequence $\{p_j\}$ converging to p_∞ . Instead below we will make an attempt formulating it only for algebraic families. This is meant to be a generalization of the 2-step degeneration theory (see [18]) for tangent cones of singularities. Here we explain a strategy and the difficulty involved. Suppose we are given a flat \mathbb{Q} -Gorenstein family of local germs $\pi : \mathcal{X} \rightarrow \Delta$ and a holomorphic section $\sigma : \Delta \rightarrow \mathcal{X}$. Denote $p_t = \sigma(t)$ and $X_t = \pi^{-1}(t)$. We suppose X_t is smooth for $t \neq 0$ and X_0 has only log terminal singularities. Locally in a neighborhood of p_0 , we may embed the family holomorphically into the family $\pi : \mathbb{C}^N \times \mathbb{C} \rightarrow \mathbb{C}$ such that p_t is mapped to $(0, t)$. Given positive weights $(\zeta_1, \dots, \zeta_N)$ on \mathbb{C}^N and $\zeta_{N+1} = 1$ on \mathbb{C} , we can take a weighted tangent cone \mathcal{W} of \mathcal{X} at $(0, 0)$. The latter is also equipped with a projection map to \mathbb{C} . Denote by W_t the fiber over t . We are only interested in the case when W_1 is of pure dimension n and has mild singularities. Geometrically W_1 is given by the limit in \mathbb{C}^N of the rescaled varieties $\lambda(t) \cdot X_t$, where $\lambda(t) \cdot (z_1, \dots, z_N) = (|t|^{-\zeta_1} z_1, \dots, |t|^{-\zeta_N} z_N)$.

In general W_0 may not agree with the weighted tangent cone \mathcal{W} of X_0 at 0. When \mathcal{X} comes from degeneration of polarized Kähler-Einstein varieties, one can choose the embedding above and the weights (ζ_j) such that \mathcal{W} is the intermediate K-semistable Fano cone discovered in [18], and there is a further equivariant degeneration to the metric tangent cone of X_0 at p_0 . More generally, by Xu-Zhuang [35] we can always make \mathcal{W} a K-semistable Fano cone in an essentially unique way. Denote such a \mathcal{W} by W^{ss} . By Li-Wang-Xu [23] there is an equivariant degeneration from W^{ss} to a unique K-polystable Fano cone \mathcal{C}^{ps} .

Now we choose $\lambda > 0$ and adjust the weights (ζ_j) to $(\lambda\zeta_j)$. It seems possible to show that a suitable chosen λ would give rise to a limit W_1 whose weighted asymptotic cone at infinity (with respect to the inverse weights $(-\zeta_j)$) is given by W^{ss} . Moreover, one expects such a W_1 to be essentially unique. Suppose all these work out we call W_1 an *algebraic minimal bubble*. It is a K-semistable polarized affine Calabi-Yau variety.

It is tempting to relate the algebraic minimal bubble with the analytic minimal bubble. However, the situation is more subtle than one would naively think. First as mentioned above an analytic minimal bubble Z is expected to be K-polystable (rather than only

K -semistable). This means that in general we need a further degeneration from W_1 to Z . To speculate what this should be one can go back to the proof in Section 4, where we have the error terms $\tau_{j,k}$ which goes to zero as $j, k, k^{-1}\bar{k}_j \rightarrow \infty$. These may accumulate and diverge when we compose from \underline{k} to \bar{k}_j . So naturally one expects to embed both W_1 and Z into \mathbb{C}^N and a sequence of elements $\Phi_j \in GL_\xi(N)$ (where ξ is the diagonal linear vector field associated to the weights (ζ_j)) such that $\Phi_j.W_1$ converges to Z in an appropriate sense. The subtlety here, according to the author, is that if one only takes purely algebro-geometric consideration this way, then such a limit Z may not be unique in general. This is related to the discussion in [31] that the class of all K -polystable polarized affine Calabi-Yau varieties with a fixed algebraic asymptotic cone modulo isomorphisms, may not be Hausdorff. Also to determine the metric on the analytic minimal bubble from the algebraic information is another issue, since as mentioned above, the Calabi-Yau metric may not be uniquely determined by the underlying polarized affine Calabi-Yau variety. Nevertheless, we make a tentative conjecture here, which may be subject to modification with further progress.

Conjecture 5.4 (Algebraicity of minimal bubbles). *Given a family \mathcal{X} as above, then*

- (1) *There exists a unique algebraic minimal bubble W_1 ;*
- (2) *There exists a unique K -polystable polarized affine Calabi-Yau variety Z such that the following properties hold*
 - (a) *one can find affine embeddings of W_1 and Z into a common \mathbb{C}^N such that both negative valuations are induced by a weight vector $(\zeta_1, \dots, \zeta_N)$ and a one parameter subgroup $\lambda(u) \in GL_\xi(N)$ such that $Z = \lim_{u \rightarrow 0} \lambda(u).W_1$;*
 - (b) *The algebraic asymptotic cone of Z is given by \mathcal{C}^{ps} ;*
 - (c) *Suppose the family comes from a polarized Kähler-Einstein degeneration, then the analytic minimal bubble is isomorphic to Z as polarized affine Calabi-Yau varieties.*
- (3) *There exists a unique complete (weak) Calabi-Yau metric on Z up to holomorphic isometry and metric scaling, which gives the metric on the analytic minimal bubble associated to a polarized Kähler-Einstein degeneration.*

Remark 5.5. *A version of this conjecture was formulated in a discussion with Cristiano Spotti in 2016. For low dimensional examples, progress has been made by him and Martin de Borbon [13].*

Remark 5.6. *From the above discussion we know that in item (2), the uniqueness does not hold if we only assume (a) and (b) but it seems to be only a finite dimensional freedom. Also it is possible to use the more intrinsic language of valuations instead of affine embeddings in \mathbb{C}^N but here we will not delve into it. The main purpose of this conjecture is to motivate further study.*

Remark 5.7. *One may consider even more general \mathbb{Q} -Gorenstein families of log terminal singularities where the general fibers are not necessarily smooth. In this case one needs to assume the volume density at p_0 (in the sense of [21]) is strictly smaller than that of p_t so that bubbling actually occurs. It is straightforward to extend the above discussion in this generality, which we omit here.*

Remark 5.8. *One may also ask about the dependence of both W_1 and Z on the section σ and the point p_0 . For example, is it generically the same?*

To show this is of interest, we consider an example here. Let \mathcal{X} be the family defined by $x_1^2 + x_2^2 + x_3^2 + x_4^4 + tx_4 = 0$ in $\mathbb{C}^4 \times \mathbb{C}$ and $p_t = (0, 0, 0, 0, t)$. Notice the central fiber X_0 is itself a K-semistable Fano cone with weights $\zeta_1 = \zeta_2 = \zeta_3 = 2, \zeta_4 = 1$ (see for example the discussion in [18] on A_k singularities). The corresponding K-polystable Fano cone \mathcal{C}^{ps} is given by $\mathbb{C}^2/\mathbb{Z}_2 \times \mathbb{C}$, which admits the flat orbifold metric. The obvious algebraic minimal bubble is given by $W_1 = X_1$ if we choose $\zeta_1 = \zeta_2 = \zeta_3 = \frac{2}{3}, \zeta_4 = \frac{1}{3}$. Suppose \mathcal{X} arises from degeneration of polarized Kähler-Einstein metrics (for example, of polarized Calabi-Yau varieties), then the main results in this paper seem to imply one of the following two possibilities must occur

- (1) The analytic minimal bubble is W_1 itself equipped with a complete Calabi-Yau metric which is asymptotic to the above flat orbifold cone. But we do not expect this to be the case since it would violate the general no-semistability at infinity conjecture in [31].
- (2) There is a degeneration of W_1 to another polarized affine Calabi-Yau variety Z with algebraic asymptotic cone \mathcal{C}^{ps} and the analytic minimal bubble is given by Z . Algebraically, one way is to use the one-parameter subgroup $\lambda(u) = (u^2, u^2, u^2, u^4)$ and Z is given by $x_1^2 + x_2^2 + x_3^2 + x_4 = 0$, which is simply \mathbb{C}^3 with a non-standard negative valuation. We know the latter admits an essentially unique complete Ricci-flat metric asymptotic to the flat orbifold cone, by [25, 12, 33, 34].

We leave these for interested readers to explore further.

5.2. More general situation. It is desirable to relax the assumptions used in this paper. Our arguments depend crucially on the existence of a polarization, which allows the application of the Hörmander L^2 technique. Without this the following is an important open question. Let (X_j, ω_j, p_j) be a sequence of n dimensional Kähler-Einstein manifolds with $Ric(\omega_j) = c_j \omega_j$ ($|c_j| \leq 1$) such that $\text{Vol}(B(p_j, 1)) \geq \kappa > 0$ and suppose $(X_\infty, d_\infty, p_\infty)$ is a Gromov-Hausdorff limit. Denote by $X_\infty = X_\infty^{reg} \cup X_\infty^{sing}$ the regular-singular decomposition in the sense of Cheeger-Colding. Notice that X_∞^{reg} is an open Kähler manifold. Denote by \mathcal{O} the push-forward to X_∞ of the structure sheaf of X_∞^{reg} .

Conjecture 5.9 (Complex analyticity). *(X_∞, \mathcal{O}) is a normal complex-analytic space and the singular set in the complex-analytic sense coincides with X_∞^{sing} .*

Noticing that by Liu-Szekelyhidi [26] cone limits in this general case are still affine varieties, and it is natural to conjecture that the singularities to be log terminal. The latter seems to be an necessary ingredient in order to extend Theorem 1.7.

Moving further, one can also study the bubbling phenomenon in this more general situation. Notice that here we do not expect the bubble limits to be affine varieties since compact holomorphic cycles may get contracted in the limiting process. A model case is to consider the convergence to the asymptotic cone of a complete Calabi-Yau metric with maximal volume growth. It is also an open question to classify the later in terms of algebro-geometric data, see [31] for more details.

Finally, the volume collapsing situation is substantially much more complicated. The only known general result is in the case of 2 dimensional Calabi-Yau metrics with uniformly bounded L^2 curvature energy studied in [32]. In particular, in this more restrictive case (which corresponds to degenerations of K3 surfaces) we know the bubble tree is also finite, but the algebro-geometric meaning is not yet well-understood. In general one expects that non-Archimedean geometry may be involved to study the bubble tree structure. Without the bounded energy hypothesis, even in complex dimension 2, if one allows collapsing then in the local setting the bubble tree may not even terminate in finite steps.

5.3. The Riemannian setting. The notion of a *bubble tree* is commonly studied in many problems in geometric analysis, such as minimal surfaces, harmonic maps, Yang-Mills connections and Einstein metrics etc. Often one works with an energy functional which is conformally invariant and studies the singularity formation for a sequence of solutions with uniformly bounded energy. Specific to the case of Einstein metrics in real m dimensions, assuming a uniform bound on the curvature energy $\int |Rm|^{m/2}$ and non-collapsing volume, bubble tree convergence was studied in [1, 3, 4]; in dimension 4, by the result of Cheeger-Naber [6] one gets the same results under the volume non-collapsing assumption alone. In higher dimensions, very little is known in general regarding the bubble tree structure if we only assume non-collapsing volume. We formulate some conjectures and questions related to generalizations of our main results in this paper.

First we will allow Einstein metrics with mild singularities. For simplicity by *singular* Einstein metrics we mean a non-collapsed Gromov-Hausdorff limit of smooth Einstein manifolds. Given $m \geq 3$, we denote by $\mathfrak{C}(m)$ the isometry classes of m dimensional singular Ricci-flat cones, and by $\mathcal{E}(m)$ the isometry classes of $m - 1$ dimensional compact singular Einstein metrics with Einstein constant $m - 2$. For $X \in \mathcal{E}(m)$ we denote by $\text{Vol}(X)$ the Riemannian volume of X and for $\mathcal{C} \in \mathfrak{C}(m)$ we denote by $\text{Vol}(\mathcal{C})$ the volume density of \mathcal{C} . A natural generalization of Theorem 1.7 is

Conjecture 5.10 (Volume rigidity). *For any $m \geq 3$, the set*

$$\mathcal{V}_1(m) = \{\text{Vol}(\mathcal{C}) | \mathcal{C} \in \mathfrak{C}(m)\}$$

and

$$\mathcal{V}_2(m) = \{\text{Vol}(X) | X \in \mathcal{E}(m)\}$$

are both discrete subsets of $(0, \infty)$.

Remark 5.11. *One might also want to allow even more general singularities using more synthetic formulations, without referring to Gromov-Hausdorff limits. However for the conjecture to hold one has to exclude for example codimension 2 cone singularities; otherwise one notices that the volume of flat 2 dimensional cones can deform continuously.*

This is known to be true if we restrict to cones with isolated singularities and smooth X . This follows from the local analytic structure of the moduli space of smooth Einstein metrics (see for example Besse [5], Chapter 12). A strategy to tackle this question in general would require more precise understanding of the singularity formations in one dimension lower, which suggests a possible (but possibly complicated) inductive approach.

Now we consider a sequence of Einstein metrics (X_j, g_j, p_j) converging to a singular Einstein metric $(X_\infty, g_\infty, p_\infty)$ with non-collapsing volume. Denote by $\text{Vol}(p_\infty)$ the volume density at p_∞ . A folklore question is the uniqueness of tangent cones at p_∞ . This is known if one a priori assumes the curvature has only quadratic blow-up at p_∞ (c.f. [10]). As a generalization of Theorem 1.10, one needs to study

Conjecture 5.12 (Cone rigidity). *The set of bubble limits with volume density equal to $\text{Vol}(p_\infty)$ consists of a single point.*

This can be viewed as an upgrade of the uniqueness of tangent cones. It seems natural to apply the strategy in [10] to prove this when the tangent cone at p_∞ is smooth. Notice Conjecture 5.10 and 5.12 together imply the existence of minimal bubbles and finite step termination of bubble trees, in exactly the same way as the Kähler-Einstein case.

Finally, a key reason why one can prove far stronger results in the (polarized) Kähler setting is due to the rigidity of holomorphic spectrum (as discussed in [18]). In the general Riemannian setting it is desirable to find a substitute. A naive candidate is the Laplacian spectrum. We propose the following questions for future study

Question 5.13. Are there special arithmetic properties of the Laplacian spectrum on compact Einstein manifolds with Einstein constant $m - 1$?

Question 5.14 (Spectrum rigidity). Suppose g_t is a smooth family of Einstein metrics on a compact manifold with Einstein constant $m - 1$, is it true that the Laplacian spectrum must be constant in t ?

Of course, one can formulate analogous conjectures/questions to the above for other geometric PDEs.

REFERENCES

- [1] Michael T. Anderson and Jeff Cheeger. Diffeomorphism finiteness for manifolds with Ricci curvature and $L^{n/2}$ -norm of curvature bounded. *Geom. Funct. Anal.*, 1(3):231–252, 1991.
- [2] Thierry Aubin. Équations du type Monge-Ampère sur les variétés kähleriennes compactes. *C. R. Acad. Sci. Paris Sér. A-B*, 283(3):Aiii, A119–A121, 1976.
- [3] Shigetoshi Bando. Bubbling out of Einstein manifolds. *Tohoku Math. J. (2)*, 42(2):205–216, 1990.
- [4] Shigetoshi Bando. Correction and addition: “Bubbling out of Einstein manifolds”. *Tohoku Math. J. (2)*, 42(4):587–588, 1990.
- [5] Arthur L. Besse. *Einstein manifolds*. Classics in Mathematics. Springer-Verlag, Berlin, 2008. Reprint of the 1987 edition.
- [6] Jeff Cheeger and Aaron Naber. Regularity of Einstein manifolds and the codimension 4 conjecture. *Ann. of Math. (2)*, 182(3):1093–1165, 2015.
- [7] Xiuxiong Chen, Simon Donaldson, and Song Sun. Kähler-Einstein metrics on Fano manifolds. I, II, III. *J. Amer. Math. Soc.*, 28(1):183–278, 2015.
- [8] Xuemiao Chen and Song Sun. Singularities of Hermitian-Yang-Mills connections and Harder-Narasimhan-Seshadri filtrations. *Duke Math. J.*, 169(14):2629–2695, 2020.
- [9] Shih-Kai Chiu. Nonuniqueness of calabi-yau metrics with maximal volume growth. *arXiv preprint arXiv:2206.08210*, 2022.
- [10] Tobias Holck Colding and William P Minicozzi. On uniqueness of tangent cones for einstein manifolds. *Inventiones mathematicae*, 196(3):515–588, 2014.
- [11] Tristan Collins and Gábor Székelyhidi. Sasaki-einstein metrics and k-stability. *Geometry & Topology*, 23(3):1339–1413, 2019.

- [12] Ronan J Conlon and Frédéric Rochon. New examples of complete Calabi-Yau metrics on \mathbb{C}^n for $n \geq 3$. *arXiv preprint arXiv:1705.08788*, 2017.
- [13] Martin de Borboon and Cristiano Spotti. Some models for bubbling of (log) kähler-einstein metrics. *in preparation*.
- [14] Tommaso de Fernex, Lawrence Ein, and Mircea Mustață. Log canonical thresholds on varieties with bounded singularities. In *Classification of algebraic varieties*, EMS Ser. Congr. Rep., pages 221–257. Eur. Math. Soc., Zürich, 2011.
- [15] Simon Donaldson. Stability, birational transformations and the Kahler-Einstein problem. *Surveys in differential geometry. Vol. XVII*, 17:203–228, 2012.
- [16] Simon Donaldson. Stability of algebraic varieties and Kähler geometry. *arXiv 1702.05745*, 2017.
- [17] Simon Donaldson and Song Sun. Gromov–Hausdorff limits of Kähler manifolds and algebraic geometry. *Acta Mathematica*, 213(1):63–106, 2014.
- [18] Simon Donaldson and Song Sun. Gromov–Hausdorff limits of Kähler manifolds and algebraic geometry, II. *Journal of Differential Geometry*, 107(2):327–371, 2017.
- [19] Renée Elkik. Singularités rationnelles et déformations. *Invent. Math.*, 47(2):139–147, 1978.
- [20] Mark Haiman and Bernd Sturmfels. Multigraded Hilbert schemes. *J. Algebraic Geom.*, 13(4):725–769, 2004.
- [21] Chi Li. Minimizing normalized volumes of valuations. *Math. Z.*, 289(1-2):491–513, 2018.
- [22] Chi Li, Xiaowei Wang, and Chenyang Xu. On the proper moduli spaces of smoothable Kähler-Einstein Fano varieties. *Duke Math. J.*, 168(8):1387–1459, 2019.
- [23] Chi Li, Xiaowei Wang, and Chenyang Xu. Algebraicity of the metric tangent cones and equivariant K-stability. *Journal of the American Mathematical Society*, 34(4):1175–1214, 2021.
- [24] Chi Li and Chenyang Xu. Stability of Valuations: Higher Rational Rank. *Peking Math. J.*, 1(1):1–79, 2021.
- [25] Yang Li. A new complete Calabi–Yau metric on \mathbb{C}^3 . *Inventiones mathematicae*, 217(1):1–34, 2019.
- [26] Gang Liu and Gábor Székelyhidi. Gromov-Hausdorff limits of Kähler manifolds with Ricci curvature bounded below II. *Comm. Pure Appl. Math.*, 74(5):909–931, 2021.
- [27] Dario Martelli, James Sparks, and Shing-Tung Yau. Sasaki-Einstein manifolds and volume minimisation. *Comm. Math. Phys.*, 280(3):611–673, 2008.
- [28] Yuji Odaka. Compact moduli spaces of Kähler-Einstein Fano varieties. *Publ. Res. Inst. Math. Sci.*, 51(3):549–565, 2015.
- [29] Cristiano Spotti, Song Sun, and Chengjian Yao. Existence and deformations of Kähler-Einstein metrics on smoothable \mathbb{Q} -Fano varieties. *Duke Math. J.*, 165(16):3043–3083, 2016.
- [30] Song Sun. Degenerations and moduli spaces in Kähler geometry. In *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. II. Invited lectures*, pages 993–1012. World Sci. Publ., Hackensack, NJ, 2018.
- [31] Song Sun and Junsheng Zhang. No semistability at infinity for calabi-yau metrics asymptotic to cones. *arXiv preprint*, 2022.
- [32] Song Sun and Ruobing Zhang. Collapsing geometry of hyperkaehler 4-manifolds and applications. *arXiv preprint arXiv:2108.12991*, 2021.
- [33] Gábor Székelyhidi. Degenerations of \mathbb{C}^n and calabi–yau metrics. *Duke Mathematical Journal*, 168(14):2651–2700, 2019.
- [34] Gábor Székelyhidi. Uniqueness of some Calabi–Yau metrics on \mathbb{C}^n . *Geometric and Functional Analysis*, 30(4):1152–1182, 2020.
- [35] Chenyang Xu and Ziquan Zhuang. Stable degenerations of singularities. *arXiv preprint*, 2022.
- [36] Shing Tung Yau. On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I. *Comm. Pure Appl. Math.*, 31(3):339–411, 1978.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720

Email address: sosun@berkeley.edu