

ALEXANDER AND JONES POLYNOMIALS OF WEAVING 3-BRAID LINKS AND WHITNEY RANK POLYNOMIALS OF LUCAS LATTICE

MARK E. ALSUKAITI AND NAFAA CHBILI

ABSTRACT. We establish a relationship between the Jones polynomial of generalized weaving knots of type $W(3, n, m)$ and the Chebyshev polynomial of the first kind. Consequently, we prove that the coefficients of the Jones polynomial of weaving knots are basically the Whitney numbers of Lucas lattices. Furthermore, we give an explicit formula for the Alexander polynomial of weaving knots $W(3, n)$ and we prove that it satisfies Fox's trapezoidal conjecture.

1. INTRODUCTION

Let $f(x) = \sum_{k=0}^n \alpha_k x^k$ be a polynomial with real positive coefficients. Then, f is said to be logarithmically concave, or log-concave for short, if its coefficients satisfy the condition $\alpha_k^2 \geq \alpha_{k-1} \alpha_{k+1}$ for all $0 < k < n$. If there exists an integer $r \leq n/2$ such that: $\alpha_0 < \alpha_1 < \dots < \alpha_{\frac{n}{2}-r} = \dots = \alpha_{\frac{n}{2}+r} > \dots > \alpha_{n-1} > \alpha_n$, then f is said to be trapezoidal. It is well known that every log-concave polynomial is trapezoidal. Polynomials and sequences with such properties have been subject to extensive studies. The importance of these polynomials in combinatorics, algebra and geometry is highlighted in [16], [34] and the references therein.

An n -component link is an embedding of n disjoint circles $\coprod S^1$ into the 3-dimensional sphere S^3 . A knot is a one-component link. A link diagram is a regular planar projection of the link together with hight information at each double point showing which of the two strands passes over the other. A link is said to be alternating if it admits an alternating diagram; a diagram where the overpass and the underpass alternate as one travels along any strand of the diagram. Let $n \geq 2$ be an integer and B_n be the group of braids on n strands. This group is generated by the elementary braids $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ subject to the following relations:

$$\begin{aligned} \sigma_i \sigma_j &= \sigma_j \sigma_i \text{ if } |i - j| \geq 2 \\ \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1}, \quad \forall 1 \leq i \leq n - 2. \end{aligned}$$

By Alexander's theorem [1], any link L in S^3 can be represented as the closure of an n -braid b . We write $L = \hat{b}$. This representation is not unique and the minimum integer n such that L admits an n -braid representation is called the braid index of L . Recall that two braids represent the same link if and only if they are related by a finite sequence of Markov moves [28]. The group B_2 is infinite cyclic. Consequently, a link of braid index two is a $(2, k)$ -torus link. A complete classification of 3-braids up to conjugacy can be found in [27]. This classification has been very useful in the study of links of braid index 3. In particular, Stoimenow [35] determined which 3-braid links are alternating. Baldwin [4] characterized quasi-alternating links of braid index 3. More recently, Lee and Vafaee [22] proved that twisted torus knots are the only L -space knots of braid index 3. As for knot invariants, signatures of 3-braid links are computed in [11] and explicit formulas for their determinants are given in [31]. The structure of the Jones polynomials of 3-braid links is studied in [8, 9]. Colored HOMFLY-PT invariants of some families of 3-braid links have been computed in [32, 33]. A list of open questions related

Key words and phrases. Weaving knots, knot polynomials, Whitney numbers, Trapezoidal Conjecture.

to the study of 3-braids is given by Birman and Menasco [6].

The Alexander polynomial [2] is a topological invariant of oriented links which associates with each link L a Laurent polynomial with integral coefficients $\Delta_L(t) \in \mathbb{Z}[t^{\pm 1/2}]$. Conway proved that this polynomial can be defined in a simple recursive way. This polynomial can be also recovered from the Burau representation of the braid group [7]. More recently, Ozsváth and Szabó [30] proved that this polynomial, up to the multiplication by some factor, is obtained as the Euler characteristic of link Floer homology. The Alexander polynomial of a knot K is known to be symmetric; it satisfies $\Delta_K(t) = \Delta_K(t^{-1})$. Furthermore, up to the multiplication by a power of t it is always possible to write $\Delta_K(t) = \sum_{i=0}^{2n} \alpha_i t^i$ with $\alpha_0 \neq 0$ and $\alpha_{2n} \neq 0$.

Murasugi, studied the Alexander polynomial of alternating knots and proved that the coefficients of this polynomial alternate in sign and that the polynomial has no gaps, equivalently $\alpha_i \alpha_{i+1} < 0$ for all $0 \leq i < 2n$. Moreover, the degree of $\Delta_K(t)$ is equal to twice the genus of the knot [25, 26]. In [12], R. Fox conjectured that the coefficients of the Alexander polynomial of an alternating knot are trapezoidal. Stoimenow, later conjectured that these coefficients are indeed log-concave [36].

Conjecture 1.1. [12] *Let K be an alternating knot and $\Delta_K(t) = \pm \sum_{i=0}^{2n} \alpha_i (-t)^i$, with $\alpha_i > 0$, its Alexander polynomial. Then there exists an integer $r \leq n$ such that:*

$$\alpha_0 < \alpha_1 < \cdots < \alpha_{n-r} = \cdots = \alpha_{n+r} > \cdots > \alpha_{2n-1} > \alpha_{2n}.$$

This conjecture, known as Fox's trapezoidal conjecture, was listed by J. Huh [16], as one of the most interesting open problems about log-concavity. The conjecture has been confirmed for several classes of alternating knots. Hartley proved the conjecture for 2-bridge knots [13]. Murasugi, showed that the conjecture holds for a large family of alternating algebraic knots [28]. Using knot Floer homology, Ozsváth and Szabó [30] confirmed the conjecture for knots of genus 2. The same result has been obtained by Jong using combinatorial methods [19, 20]. Hirasawa and Murasugi [14] showed that Conjecture 1.1 holds for stable alternating knots and suggested a refinement of this conjecture by stating that the length of the stable part is less than the signature of the knot $\sigma(K)$, which means $r \leq |\sigma(K)|/2$. In [10], Chen proved that this refinement holds for two-bridge knots. Alrfai and the second author, showed that certain families of 3-braid knots satisfy Fox trapezoidal conjecture in its refined form [3].

Let $m \geq 1$ be an integer. The closure of the 3-braid $(\sigma_1^m \sigma_2^{-m})^n$ is called the generalized weaving link $W(3, n, m)$, see [32, 33]. Notice that if $m = 1$, then we get the classical weaving link often denoted as $W(3, n)$. In this paper, we establish a relationship between the trace of the Burau representation of the braids of type $(\sigma_1^m \sigma_2^{-m})^n$ and the Chebyshev polynomials of the first kind. As an application, we introduce explicit formulas for the coefficients of Jones and Alexander polynomials of weaving links. Furthermore, we verify that the Alexander polynomial of weaving knots satisfy Fox's trapezoidal conjecture, see Figure 1. We also compute the zeroes of the Alexander polynomials of weaving knots and we verify that the real part of any such zero is greater than -1 , hence these zeroes satisfy Hoste's condition, see [23] and [15]. By setting $s = -t$, we have:

Theorem 1.2. (1) *The Alexander polynomial of the weaving link $W(3, n)$ is given by the following formula: $\Delta_{W(3, n)}(s) = \sum_{k=0}^{2n-2} \alpha_{n, k} s^k$, with:*

$$\alpha_{n, k} = \sum_{i=0}^{n-1} (-1)^{k-n+i+1} \frac{2n(n+i)!}{(n-i-1)!(2i+2)!} \binom{i; 3}{k+i-n+1}$$

where $\binom{n; 3}{k}$ is the coefficient of q^k in the expansion of $(1 + q + q^2)^n$.
 (2) For any $n \geq 1$, the coefficients of $\Delta_{W(3,n)}(s)$ are trapezoidal.

1																	
1	3	1															
1	4	6	4	1													
1	5	10	13	10	5	1											
1	6	15	24	29	24	15	6	1									
1	7	21	40	58	66	58	40	21	7	1							
1	8	28	62	104	140	155	140	104	62	28	8	1					
1	9	36	91	173	266	341	371	341	266	173	91	36	9	1			
1	10	45	128	272	468	676	838	900	838	676	468	272	128	45	10	1	

FIGURE 1. The coefficients of the Alexander polynomials of $\Delta_{W(3,n)}(s)$, for $n = 1$ to 10.

Here is an outline of this paper. In Section 2, we briefly recall some basic definitions and notations needed in the sequel. In Section 3, we study the Jones polynomial of generalized weaving links. Section 4 is devoted to prove our main theorem. Finally, the zeroes of the Alexander polynomial of weaving knots are computed in Section 5.

2. PRELIMINARIES

In 1928, Alexander [2] introduced a topological invariant of oriented links. This invariant is a one-variable Laurent polynomial with integral coefficients $\Delta_L(t)$. It can be defined in several different, but equivalent, ways. In particular, it is uniquely determined recursively using Conway skein relations:

$$\Delta_U(t) = 1,$$

$$\Delta_{L_+}(t) - \Delta_{L_-}(t) = (\sqrt{t} - \frac{1}{\sqrt{t}})\Delta_{L_0}(t),$$

where U is the unknot and L_+ , L_- and L_0 are three oriented link diagrams which are identical except in a small region where they are as pictured in Figure 2.




FIGURE 2. The three oriented links L_+ , L_- and L_0 , respectively.

This polynomial is known to be symmetric, in the sense that it satisfies $\Delta_L(t) = \Delta_L(t^{-1})$. The reduced Burau representation of the braid group permits another way to define the Alexander polynomial. Since we are only interested in 3-braid links, we shall restrict ourselves to the Burau representation of B_3 . This representation is faithful and can be described as follows. Let b be a given 3-braid and e_b the exponent sum of b as a word in the elementary braids σ_1 and σ_2 . Let $\psi_t : B_3 \rightarrow GL(2, \mathbb{Z}[t, t^{-1}])$ be the Burau representation defined on the generators

of B_3 by $\psi_t(\sigma_1) = \begin{bmatrix} -t & 1 \\ 0 & 1 \end{bmatrix}$ and $\psi_t(\sigma_2) = \begin{bmatrix} 1 & 0 \\ t & -t \end{bmatrix}$. The Alexander polynomial, up to the multiplication by a unit of $\mathbb{Z}[t, t^{-1}]$, of the link $L = \hat{b}$ is given by:

$$\Delta_L(t) = (-1/\sqrt{t})^{e_b-2} \frac{1-t}{1-t^3} \det(I_2 - \psi_t(b)),$$

where I_2 is the identity matrix. Given that this representation is 2-dimensional, then $\det(I_2 - \psi_t(b)) = 1 - \text{tr}(\psi_t(b)) + \det(\psi_t(b))$, where tr denotes the usual matrix-trace function defined as the sum of entries on the main diagonal of the matrix.

In 1984, V. F. Jones introduced a new polynomial invariant of oriented links. This one-variable Laurent polynomial, denoted hereafter by $V_L(t)$, was originally defined through the study of representations of the braid group via certain von Neumann algebras [18]. The Jones polynomial is uniquely determined by the following skein relations:

$$\begin{aligned} V_U(t) &= 1, \\ tV_{L_+}(t) - t^{-1}V_{L_-}(t) &= (\sqrt{t} + \frac{1}{\sqrt{t}})V_{L_0}(t). \end{aligned}$$

In the case of 3-braid links, Birman [5] showed that the Jones polynomial can be defined using the Burau representation of B_3 by the following formula:

$$V_{\hat{\alpha}}(t) = (-\sqrt{t})^{e_{\alpha}}(t + t^{-1} + \text{tr}(\psi_t(\alpha))).$$

Throughout the rest of this paper, we set $s = -t$. Moreover, for any positive integer n we set $[n] = \frac{1-s^n}{1-s}$. Let $T_n(x)$ be the Chebyshev polynomial of the first kind. This polynomial is determined recursively by the following relations:

$$\begin{aligned} T_0(x) &= 1 \\ T_1(x) &= x \\ T_{n+1}(x) &= 2xT_n(x) - T_{n-1}(x). \end{aligned}$$

Let $F_n(q) = \sum_{k=0}^n f_{n,k}q^k$ and $C_n(q) = \sum_{k=0}^{2n} c_{n,k}q^k$ be the rank polynomials of the Fibonacci lattice and the Lucas lattice, respectively. Explicit formulas for the Whitney numbers $f_{n,k}$ and $c_{n,k}$ have been obtained by Munarini and Zagaglia Salvi [24]. Another formula for $c_{n,k}$ is given in [29]:

$$c_{n,k} = \begin{cases} \sum_{j=0}^{\lfloor k/2 \rfloor} \frac{n}{n-j} \binom{n-j}{n-k+j} \binom{k-j-1}{j} & k < 2n \\ 1 & k = 2n. \end{cases}$$

The relation between the rank polynomial of Lucas lattice and the Chebyshev polynomial is given by the following proposition [24].

Proposition 2.1. *For any $n \geq 0$, we have: $C_n(q) = 2q^n T_n(\frac{1+q+q^2}{2q})$.*

3. BURAU REPRESENTATION OF 3-BRAIDS

In this section, we shall study the Burau representation of the braids $b_n = (\sigma_1 \sigma_2^{-1})^n$ as well as $b_{n,m} = (\sigma_1^m \sigma_2^{-m})^n$, where n and m are positive integers. Recall that these braids close to the weaving link $W(3, n)$ and the generalized weaving link $W(3, n, m)$, respectively. In particular, we will give an explicit formula for the coefficients of Jones polynomials of weaving links.

Proposition 3.1. *The trace of the Burau representation of the generalized weaving braid is given by:*

$$\text{tr}(\psi_s(b_{n,m})) = 2^{-n} s^{-nm} (\lambda_{1,m}^n + \lambda_{2,m}^n)$$

where

$$\begin{aligned}\lambda_{1,m} &= (1 + [m]^2 s + s^{2m}) + \sqrt{(1 + [m]^2 s + s^{2m})^2 - 4s^{2m}} \text{ and} \\ \lambda_{2,m} &= (1 + [m]^2 s + s^{2m}) - \sqrt{(1 + [m]^2 s + s^{2m})^2 - 4s^{2m}}.\end{aligned}$$

Proof. Notice first that one can easily show that for any $m > 0$, we have:

$$\psi_s(\sigma_1^m) = \begin{bmatrix} s^m & [m] \\ 0 & 1 \end{bmatrix}, \text{ and } \psi_s(\sigma_2^{-m}) = s^{-m} \begin{bmatrix} s^m & 0 \\ s[m] & 1 \end{bmatrix}.$$

Thus,

$$\psi_s(\sigma_1^m \sigma_2^{-m}) = s^{-m} \begin{bmatrix} s^{2m} + s[m]^2 & [m] \\ s[m] & 1 \end{bmatrix}.$$

Now, assume that $\psi_s(b_{n,m}) = s^{-nm} \begin{bmatrix} A_n(s) & B_n(s) \\ C_n(s) & D_n(s) \end{bmatrix}$, where all entries are polynomials on the variable s . Since $b_{n+1,m} = b_{n,m} \sigma_1^m \sigma_2^{-m}$, we get the following system of recurrence relations:

$$\begin{aligned}A_{n+1}(s) &= (s^{2m} + s[m]^2)A_n(s) + s[m]B_n(s) \\ B_{n+1}(s) &= [m]A_n(s) + B_n(s) \\ C_{n+1}(s) &= (s^{2m} + s[m]^2)C_n(s) + s[m]D_n(s) \\ D_{n+1}(s) &= [m]C_n(s) + D_n(s)\end{aligned}$$

with $A_0(s) = D_0(s) = 1$ and $B_0(s) = C_0(s) = 0$. Hence:

$$\begin{aligned}\begin{bmatrix} A_{n+1}(s) \\ B_{n+1}(s) \end{bmatrix} &= \begin{bmatrix} (s^{2m} + s[m]^2) & s[m] \\ [m] & 1 \end{bmatrix} \begin{bmatrix} A_n(s) \\ B_n(s) \end{bmatrix} \\ &= \begin{bmatrix} (s^{2m} + s[m]^2) & s[m] \\ [m] & 1 \end{bmatrix}^{n+1} \begin{bmatrix} A_0(s) \\ B_0(s) \end{bmatrix} \\ &= \begin{bmatrix} (s^{2m} + s[m]^2) & s[m] \\ [m] & 1 \end{bmatrix}^{n+1} \begin{bmatrix} 1 \\ 0 \end{bmatrix}.\end{aligned}$$

Similarly, we also get for $C_n(s)$ and $D_n(s)$:

$$\begin{bmatrix} C_{n+1}(s) \\ D_{n+1}(s) \end{bmatrix} = \begin{bmatrix} (s^{2m} + s[m]^2) & s[m] \\ [m] & 1 \end{bmatrix}^{n+1} \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

Solving for the eigenvalues of $\begin{bmatrix} (s^{2m} + s[m]^2) & s[m] \\ [m] & 1 \end{bmatrix}^n$ then diagonalizing the matrix gives us:

$$\begin{aligned}A_n(s) + D_n(s) &= 2^{-n}(\lambda_{1,m}^n + \lambda_{2,m}^n). \text{ Thus,} \\ \text{tr}(\psi_s(b_{n,m})) &= s^{-n}(A_n(s) + D_n(s)) \\ &= 2^{-n} s^{-n}(\lambda_{1,m}^n + \lambda_{2,m}^n).\end{aligned}$$

□

Corollary 3.2. *We have: $\text{tr}(b_n) = 2^{-n} s^{-n}(\lambda_1^n + \lambda_2^n)$, where*

$$\begin{aligned}\lambda_1 &= (1 + s + s^2) + \sqrt{(1 + s + s^2)^2 - 4s^2} \text{ and} \\ \lambda_2 &= (1 + s + s^2) - \sqrt{(1 + s + s^2)^2 - 4s^2}.\end{aligned}$$

Theorem 3.3. *The Jones polynomial of the generalized weaving knot $W(3, n, m)$ is given by:*

$$V_{W(3,n,m)}(s) = -s - s^{-1} + 2T_n\left(\frac{1 + [m]^2s + s^{2m}}{2s^m}\right).$$

Proof. The Jones polynomial can be obtained through the Burau representation:

$$V_{W(3,n,m)}(s) = -s - s^{-1} + \text{tr}(\psi_s(\sigma_1^m \sigma_2^{-m})^n).$$

It is known that $T_n(x) = \frac{1}{2}((x - \sqrt{x^2 - 1})^n + (x + \sqrt{x^2 - 1})^n)$, hence it becomes clear that:

$$T_n\left(\frac{1 + [m]^2s + s^{2m}}{2s^m}\right) = 2^{-n-1} s^{-nm} (\lambda_1^n + \lambda_2^n) = \frac{\text{tr}(\psi_s(\sigma_1^m \sigma_2^{-m})^n)}{2}.$$

Consequently,

$$V_{W(3,n,m)}(s) = -s - s^{-1} + 2T_n\left(\frac{1 + [m]^2s + s^{2m}}{2s^m}\right).$$

□

Corollary 3.4. *The Jones polynomial of the weaving knot $W(3, n)$ is given by:*

$$V_{W(3,n)}(s) = -s - s^{-1} + s^{-n} C_n(s) = \sum_{k=0}^{2n} a_k s^{k-n},$$

where

$$a_k = \begin{cases} c_{n,k} & |k - n| \neq 1 \\ c_{n,k} - 1 & |k - n| = 1 \end{cases}$$

Proof. By Theorem 3.3, we know that $\text{tr}(\psi_s((\sigma_1 \sigma_2^{-1})^n)) = -s - s^{-1} + 2T_n\left(\frac{1+s+s^2}{2s}\right)$, and by Proposition 2.1 we have that: $2T_n\left(\frac{1+s+s^2}{2s}\right) = s^{-n} C_n(s) = \sum_{k=0}^{2n} c_{n,k} s^{k-n}$. □

The determinant of an oriented link L is a classical numerical invariant of links, $\det(L)$, which can be obtained by evaluation of the Jones polynomial at $t = -1$; $\det(L) = |V_L(-1)|$. The Lucas numbers L_k are defined recursively by the relations: $L_0 = 2, L_1 = 1$ and $L_k = L_{k-1} + L_{k-2}$ for any $k \geq 2$. The generalized Lucas numbers $L_{m,k}$ are defined by the relation $L_0 = 2, L_1 = 1$ and $L_{m,k} = mL_{m,k-1} + L_{m,k-2}$. The relationship between the determinants of weaving links and the Lucas numbers has been observed first in [33]. Then this relation was extended to the wider class of generalized weaving links in [21]. These results can be obtained as a direct consequence of Theorem 3.3 above.

Corollary 3.5. *For any $n \geq 1$ and $m \geq 1$, we have:*

$$\begin{aligned} \det(W(3, n)) &= L_{2n} - 2 \\ \det(W(3, n, m)) &= L_{m,2n} - 2. \end{aligned}$$

Proof. Using the recurrence relation for $L_{m,k}$ we obtain

$$L_{m,k} = \left(\frac{m + \sqrt{m^2 + 4}}{2}\right)^k + \left(\frac{m - \sqrt{m^2 + 4}}{2}\right)^k.$$

Then notice that:

$$\begin{aligned}
 \det(W(3, n, m)) &= |V_{W(3, n, m)}(1)| \\
 &= -2 + 2T_n\left(\frac{2+m^2}{2}\right) \\
 &= -2 + \left(\frac{2+m^2}{2} + \sqrt{\left(\frac{2+m^2}{2}\right)^2 - 1}\right)^n + \left(\frac{2+m^2}{2} - \sqrt{\left(\frac{2+m^2}{2}\right)^2 - 1}\right)^n \\
 &= -2 + \left(\frac{m + \sqrt{m^2 + 4}}{2}\right)^{2n} + \left(\frac{m - \sqrt{m^2 + 4}}{2}\right)^{2n} \\
 &= -2 + L_{m, 2n}.
 \end{aligned}$$

Similarly, $\det(W(3, n)) = L_n - 2$. □

We close this section by the following proposition which will be used later to prove that the Alexander polynomial of weaving knots is trapezoidal.

Proposition 3.6. *The Whitney numbers of the Lucas lattice $c_{n,k}$ satisfy the following recurrence relation:*

$$\begin{aligned}
 (1) \quad &c_{n,0} = 1 \\
 &c_{n,1} = n = c_{n-1,1} + c_{n-1,0} \\
 &c_{n,k} = c_{n-1,k} + c_{n-1,k-1} + c_{n-1,k-2} - c_{n-2,k-2}.
 \end{aligned}$$

Proof. For $k = 0$ and $k = 1$ this follows straightforward from the explicit formula for $c_{n,k}$. For $k \geq 0$, we have the following identity relating the sequences $c_{n,k}$ and $f_{n,k}$, [24]:

$$(2) \quad c_{n+2,k+2} = f_{2n+4,k+2} + f_{2n,k},$$

where

$$(3) \quad f_{n+4,k+2} = f_{n+2,k+2} + f_{n+2,k+1} + f_{n+2,k} - f_{n,k}.$$

Substituing 3 in 2 we get:

$$\begin{aligned}
 c_{n+2,k+2} &= f_{2n+2,k+2} + f_{2n+2,k+1} + f_{2n+2,k} - f_{2n,k} + f_{2n-2,k} \\
 &\quad + f_{2n-2,k-1} + f_{2n-2,k-2} - f_{2n-4,k-2} \\
 &= c_{n+1,k+2} + c_{n+1,k+1} + c_{n+1,k} - c_{n,k}.
 \end{aligned}$$

Giving us $c_{n,k} = c_{n-1,k} + c_{n-1,k-1} + c_{n-1,k-2} - c_{n-2,k-2}$, for $k \geq 2$. □

4. PROOF OF THE MAIN THEOREM

This section is devoted to the proof of Theorem 1.2. First, we shall give explicit formulas for the coefficients of the Alexander polynomial of weaving links $W(3, n)$. More precisely, we will prove that: $\Delta_{W(3,n)}(s) = (\sum_{k=0}^{2n-2} \alpha_{n,k} s^k)$ where

$$\alpha_{n,k} = \sum_{i=0}^{n-1} (-1)^{k-n+i} \frac{2n(n+i)!}{(n-i-1)!(2i+2)!} \binom{i; 3}{k+i-n+1}.$$

We know that $T_n(x) = n \sum_{k=0}^n (-2)^k \frac{(n+k-1)!}{(n-k)!(2k)!} (1-x)^k$, then:

$$\begin{aligned}
2 - 2T_n\left(\frac{1+s+s^2}{2s}\right) &= 2 - 2n \sum_{k=0}^n (-2)^k \frac{(n+k-1)!}{(n-k)!(2k)!} \left(1 - \frac{1+s+s^2}{2s}\right)^k \\
&= 2 - 2n \sum_{k=0}^n (2)^k \frac{(n+k-1)!}{(n-k)!(2k)!} \left(\frac{1-s+s^2}{2s}\right)^k \\
&= -2n \sum_{k=1}^n (2)^k \frac{(n+k-1)!}{(n-k)!(2k)!} \left(\frac{1-s+s^2}{2s}\right)^k \\
&= -2\left(\frac{1-s+s^2}{2s}\right)n \sum_{k=1}^n (2)^k \frac{(n+k-1)!}{(n-k)!(2k)!} \left(\frac{1-s+s^2}{2s}\right)^{k-1} \\
&= \frac{-(1-s+s^2)}{s} n \sum_{k=1}^n (2)^k \frac{(n+k-1)!}{(n-k)!(2k)!} \left(\frac{1-s+s^2}{2s}\right)^{k-1} \\
&= \frac{-(1-s+s^2)}{s} n \sum_{k=0}^{n-1} (2)^{k+1} \frac{(n+k)!}{(n-k-1)!(2k+2)!} \left(\frac{1-s+s^2}{2s}\right)^k \\
&= \frac{-(1-s+s^2)}{s} n \sum_{k=0}^{n-1} \frac{2(n+k)!}{(n-k-1)!(2k+2)!} \sum_{l=0}^{2k} (-1)^l \binom{k; 3}{l} s^{l-k} \\
&= \frac{-(1-s+s^2)}{s} n \sum_{k=0}^{n-1} \frac{2(n+k)!}{(n-k-1)!(2k+2)!} \sum_{l=-k}^k (-1)^{l+k} \binom{k; 3}{l+k} s^l \\
&= -\frac{1-s+s^2}{s} \sum_{l=-n+1}^{n-1} \sum_{k=0}^{n-1} (-1)^{l+k} \frac{2n(n+k)!}{(n-k-1)!(2k+2)!} \binom{k; 3}{l} s^l \\
&= -\frac{1-s+s^2}{s} \sum_{l=-n+1}^{n-1} \sum_{k=0}^{n-1} (-1)^{l+k} \frac{2n(n+k)!}{(n-k-1)!(2k+2)!} \binom{k; 3}{l+k} s^l \\
&= -\frac{1-s+s^2}{s} \sum_{l=-n+1}^{n-1} \sum_{k=0}^{n-1} (-1)^{l+k} \frac{2n(n+k)!}{(n-k-1)!(2k+2)!} \binom{k; 3}{l+k} s^l.
\end{aligned}$$

Therefore, up to some multiplication of a unit in $\mathbb{Z}[s, s^{-1}]$, we have:

$$\Delta_{W(3,n)}(s) = \frac{2 - 2T_n\left(\frac{1+s+s^2}{2s}\right)}{1-s+s^2} = - \sum_{l=-n+1}^{n-1} \sum_{k=0}^{n-1} (-1)^{l+k} \frac{2n(n+k)!}{(n-k-1)!(2k+2)!} \binom{k; 3}{l+k} s^l.$$

We could then shift the polynomial by s^{n-1} to give us

$$\Delta_{W(3,n)}(s) = \sum_{l=0}^{2n-2} \sum_{k=0}^{n-1} (-1)^{l-n+k+1} \frac{2n(n+k)!}{(n-k-1)!(2k+2)!} \binom{k; 3}{l+k-n+1} s^l.$$

Now, in order to prove the second part of Theorem 1.2 we need the following Lemma.

Lemma 4.1. *Let $\Delta_{W(3,n)}(s) = (\sum_{k=0}^{2n-2} \alpha_{n,k} s^k)$, then $\alpha_{k,n}$ satisfies the following recurrence relations:*

$$\begin{aligned}
\alpha_{n,0} &= 1 \\
\alpha_{n,1} &= \alpha_{n-1,1} + \alpha_{n-1,0} \\
\alpha_{n,k} &= \alpha_{n-1,k} + \alpha_{n-1,k-1} + \alpha_{n-1,k-2} - \alpha_{n-2,k-2} \quad \text{for } 1 < k < n-1 \\
\alpha_{n,n-1} &= \alpha_{n-1,n-1} + \alpha_{n-1,n-2} + \alpha_{n-1,n-3} - \alpha_{n-2,n-3} + 2.
\end{aligned}$$

Proof. Recall that the Burau representation defines the Alexander polynomial up to the multiplication by a unit. In the following calculations, we shall use the symbol $f \equiv g$ to indicate that the polynomials f and $g \in \mathbb{Z}[s, s^{-1}]$ satisfy $f = u.g$ for some unit $u \in \mathbb{Z}[s, s^{-1}]$. Since $\Delta_{W(3,n)}(s) \equiv \frac{1+s}{1+s^3} \det(I - \psi_s((\sigma_1\sigma_2^{-1})^n))$ and $\det(I - \psi_s((\sigma_1\sigma_2^{-1})^n))$ is the characteristic polynomial evaluated at $\lambda = 1$ of the 2×2 matrix $\psi_s((\sigma_1\sigma_2^{-1})^n)$ then:

$$\Delta_{W(3,n)}(s) \equiv \frac{1+s}{1+s^3} (1 - \text{tr}(\psi_s((\sigma_1\sigma_2^{-1})^n)) + \det(\psi_s((\sigma_1\sigma_2^{-1})^n))).$$

Notice that $\det(\psi_s((\sigma_1\sigma_2^{-1})^n)) = \det(s^{-1} \begin{bmatrix} s^2 + s & 1 \\ s & 1 \end{bmatrix})^n = 1$.

Therefore:

$$\begin{aligned} \Delta_{W(3,n)}(s) &\equiv \frac{1+s}{1+s^3} (2 - \text{tr}(\psi_s((\sigma_1\sigma_2^{-1})^n))) \\ &\equiv \frac{1+s}{1+s^3} (2s^n - \sum_{k=0}^{2n} c_{n,k} s^k) \\ &\equiv (1+s) (2s^n - \sum_{k=0}^{2n} c_{n,k} s^k) (\sum_{i=0}^{2n} (-s)^{3i}). \end{aligned}$$

Now if we define $c_{n,k}^*$ to be:

$$\begin{aligned} c_{n,0}^* &= c_{n,0} = 1 \\ c_{n,k}^* &= c_{n,k} + c_{n,k-1} \quad \text{for } 0 < k < n \\ c_{n,n}^* &= c_{n,n} + c_{n,n-1} - 2 \\ c_{n,k}^* &= c_{n,2n-k+1}^* \quad \text{for } k > n \end{aligned}$$

and $\alpha_{n,k} = c_{n,k}^* - c_{n,k-3}^* + c_{n,k-6}^* - \dots = \sum_{i=0}^{k/3} (-1)^i c_{n,k-3i}^*$, then we get:

$$\begin{aligned} \Delta_{W(3,n)}(s) &\equiv \left(\sum_{k=0}^{2n+1} c_{n,k}^* s^k \right) \left(\sum_{i=0}^{2n} (-s)^{3i} \right) \\ &\equiv \sum_{k=0}^{2n-2} \alpha_{n,k} s^k. \end{aligned}$$

Notice that by Proposition 3.6 we have:

If $k = 1$, then

$$\begin{aligned} c_{n,1}^* &= c_{n,1} + c_{n,0} \\ &= c_{n-1,1} + c_{n-1,0} + c_{n-1,0} \\ &= c_{n-1,1}^* + c_{n-1,0}^*. \end{aligned}$$

If $2 \leq k < n-1$, then

$$\begin{aligned} c_{n,k}^* &= c_{n,k} + c_{n,k-1} \\ &= c_{n-1,k} + c_{n-1,k-1} + c_{n-1,k-2} - c_{n-2,k-2} \\ &\quad c_{n-1,k-1} + c_{n-1,k-2} + c_{n-1,k-3} - c_{n-2,k-3} \\ &= c_{n-1,k}^* + c_{n-1,k-1}^* + c_{n-1,k-2}^* - c_{n-2,k-2}^*. \end{aligned}$$

If $k = n - 1$, then

$$\begin{aligned}
c_{n,n-1}^* &= c_{n,n-1} + c_{n,n-2} \\
&= c_{n-1,n-1} + c_{n-1,n-2} + c_{n-1,n-3} - c_{n-2,n-3} \\
&\quad c_{n-1,n-2} + c_{n-1,n-3} + c_{n-1,n-4} - c_{n-2,n-4} \\
&= c_{n-1,k}^* + c_{n-1,k-1}^* + c_{n-1,k-2}^* - c_{n-2,k-2}^* + 2.
\end{aligned}$$

Hence it follows from the definition of $\alpha_{n,k}$ and the recurrence relation above that:

$$\begin{aligned}
(4) \quad &\alpha_{n,0} = 1 \\
&\alpha_{n,1} = \alpha_{n-1,1} + \alpha_{n-1,0} \\
&\alpha_{n,k} = \alpha_{n-1,k} + \alpha_{n-1,k-1} + \alpha_{n-1,k-2} - \alpha_{n-2,k-2} \quad \text{for } 1 < k < n - 1 \\
&\alpha_{n,n-1} = \alpha_{n-1,n-1} + \alpha_{n-1,n-2} + \alpha_{n-1,n-3} - \alpha_{n-2,n-3} + 2.
\end{aligned}$$

□

Notice that this immediately gives us $\alpha_{n,0} < \alpha_{n,1} < \alpha_{n,2}$. For $k > 2$, we use the above system to show that for $2 < k < n - 1$ we have:

$$\alpha_{n,k} = \alpha_{n-1,k} + \alpha_{n-1,k-1} + \alpha_{n-2,k-3} + \alpha_{n-3,k-5} + \dots = \alpha_{n-1,k} + \sum_{i=1}^{\frac{k-1}{2}} \alpha_{n-i,k-2i+1}$$

and for $k = n - 1$ we have:

$$\alpha_{n,k} = 2 + \alpha_{n-1,k} + \sum_{i=1}^{\frac{k-1}{2}} \alpha_{n-i,k-2i+1}.$$

This directly comes from the recurrence relation 4. Since $\alpha_{n-1,k-2}$ is related to $\alpha_{n-2,k-2}$ we can expand down to obtain the above relation. Notice that $\Delta_{W(3,2)}(s) = s^{-1} + 3 + s$ is trapezoidal, thus we can induct on n to show that $\Delta_{W(3,n)}(s)$ is trapezoidal. Indeed, if we assume that $\Delta_{W(n-1,3)}(s)$ is trapezoidal we obtain:

$$\begin{aligned}
\alpha_{n,k} &\geq \alpha_{n-1,k} + \sum_{i=1}^{\frac{k-1}{2}} \alpha_{n-i,k-2i+1} \\
&> \alpha_{n-1,k-1} + \sum_{i=1}^{\frac{k-1}{2}} \alpha_{n-i,k-2i} \\
&= \alpha_{n,k-1}.
\end{aligned}$$

Hence completing our proof since $\alpha_{n,k-1} < \alpha_{n,k}$ for $k \leq n - 1$. Moreover, $\Delta_{W(3,n)}(s)$ is a symmetric polynomial, thus it has its center at $n - 1$. Therefore, it satisfies Fox's trapezoidal conjecture and the length of the satble part is $r = 0$ which is less than or equal to $\frac{|\sigma(W(3,n))|}{2}$.

5. ZEROES OF THE ALEXANDER POLYNOMIAL OF WEAVING KNOTS

In 2022, J. Hoste conjectured that if z is a complex zero of the Alexander polynomial of an alternating knot, then z satisfies the condition $\Re(z) > -1$. Lyubich and Murasugi proved that this conjecture holds for a large class of two-bridge knots, see [23]. Stoimenow [37] proved it is true for knots of genus less than or equal 4. Ishikawa [17] proved that Hoste's conjecture holds for any 2-bridge knot. A thorough discussions of the distribution of the zeros of Alexander polynomial of alternating knots can be found in [14]. More recently, Hirasawa, Ishikawa and Suzuki [15] found counter-examples to Hoste's conjecture. It is worth mentioning here that in all these counterexamples, the zeroes have real parts which are very close to -1. Thus, the existence of a lower bound to the real part of these zeros remains still an open question. In this

section, we shall compute the zeroes of $\Delta_{W(3,n)}(t)$ and prove that for every such zero the real part is greater than -1.

Proposition 5.1. *The zeroes of $\Delta_{W(3,n)}(t)$ are of the form:*

$$z = -\frac{1}{2} \left(2\cos\left(\frac{2k}{n}\pi\right) - 1 \pm \sqrt{(2\cos\left(\frac{2k}{n}\pi\right) - 1)^2 - 4} \right),$$

with $1 \leq k \leq \lfloor n/2 \rfloor$.

Proof. The zeroes of $\Delta_{W(3,n)}(s)$ correspond to the zeroes of $2 - 2T_n(\frac{1+s+s^2}{2s})$ with the zeroes of $1 - s + s^2$ ignored. If $2 - 2T_n(\frac{1+s+s^2}{2s}) = 0$ then $T_n(\frac{1+s+s^2}{2s}) = 1$. These values correspond to the peaks of $T_n(x)$ for $-1 \leq x \leq 1$ which are indeed given by $x_k = \cos(\frac{2k}{n}\pi)$ for $0 \leq k \leq \lfloor n/2 \rfloor$. Hence, the values of the zeroes of the Alexander polynomial are the solutions of $\frac{1+s+s^2}{2s} = \cos(\frac{2k}{n}\pi)$. Solving for s and taking $t = -s$ gives us the desired form. \square

Corollary 5.2. *If z is a zero of $\Delta_{W(3,n)}(t)$, then $\Re(z) > -1$. Moreover, if z is a non-real root then $|z| = 1$.*

Proof. If $(2\cos(\frac{2k}{n}\pi) - 1)^2 - 4 \geq 0$, then we have $\cos(\frac{2k}{n}\pi) \leq \frac{-1}{2}$. Since $\cos(\frac{2k}{n}\pi) \geq -1$ we get that $(2\cos(\frac{2k}{n}\pi) - 1)^2 - 4 \leq 5$. If $(2\cos(\frac{2k}{n}\pi) - 1)^2 - 4 \geq 0$, then z is real and we have

$$\begin{aligned} -3 - \sqrt{5} &\leq 2\cos\left(\frac{2k}{n}\pi\right) - 1 \pm \sqrt{(2\cos\left(\frac{2k}{n}\pi\right) - 1)^2 - 4} \leq -2 + \sqrt{5} \\ \implies \frac{-3 - \sqrt{5}}{2} &\leq -z \leq \frac{-2 + \sqrt{5}}{2} \\ \implies z &> -1. \end{aligned}$$

Otherwise, if $(2\cos(\frac{2k}{n}\pi) - 1)^2 - 4 < 0$ then z is a non-real root and we have

$$\begin{aligned} \Re(z) &= \frac{1}{2} \left(1 - 2\cos\left(\frac{2k}{n}\pi\right) \right) \\ \Im(z) &= \pm \frac{1}{2} \sqrt{4 - (1 - 2\cos\left(\frac{2k}{n}\pi\right))^2}. \end{aligned}$$

One can easily see that $\Re(z) > -1$ and $|z| = 1$. \square

ACKNOWLEDGMENTS

This research was funded by United Arab Emirates University, UPAR grant #G00004167.

REFERENCES

- [1] J. W. Alexander, A lemma on a system of knotted curves, *Proc. Nat. Acad. Sci. USA.* **9** (1923), 93-95.
- [2] J. W. Alexander, Topological invariants of knots and links, *Trans. Amer. Math. Soc.*, **30** (1928) 275-306.
- [3] M. Alrefai and N. Chbili, On Fox's Trapezoidal Conjecture for Closed 3-braids, *International Journal of Mathematics*, **32** (01), (2021) 2150002.
- [4] J. Baldwin, Heegaard Floer homology and genus one, one boundary component open books, *J. Topol.* **1**(4) (2008) 963-992.
- [5] J. Birman, On the Jones polynomial of closed 3-braids, *Inventiones Mathematicae* **81** (1985) 287-294.
- [6] J. Birman and W. Menasco, A note on closed 3-braids, *Commun. Contemp. Math.* **10**(1) (2008) 1033-1047.
- [7] W. Burau, Über Zopfgruppen und gleichsinnig verdrillte Verkettungen, *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg* **11** (1935) 179-186.
- [8] N. Chbili and K. Qazaqzeh, On the Jones Polynomial of quasi-atlernating links, *Topology and its Applications*, **264** (2019) 1-11.

- [9] N. Chbili, A Note on the Jones Polynomial of 3-braid links, *Siberian Mathematical Journal*, **63** (2022) 983–994.
- [10] W. Chen, On two-bridge knots and a conjecture of Hirasawa–Murasugi, *Journal of Knot Theory and Its Ramifications*, **30**(2) 2021, 2150007.
- [11] D. Erle, Calculation of the signature of a 3-braid link, *Kobe J. Math.* **16**(2) (1999) 161–175.
- [12] R. H. Fox, Some problems in knot theory, in *Topology of 3-manifolds and related topics, Proc. The Univ. of Georgia Institute, 1961*, (Prentice-Hall, Englewood Cliffs, N.J. 1962), pp. 168–176.
- [13] R. I. Hartley, On two-bridged knot polynomials, *J. Austral. Math. Soc. Ser. A* **28**(2) (1979) 241–249.
- [14] M. Hirasawa and K. Murasugi, Various stabilities of the Alexander polynomials of knots and links, *arXiv preprint arXiv:1307.1578* (2013).
- [15] M. Hirasawa, K. Ishikawa and M. Suzuki, Alternating knots with Alexander polynomials having unexpected zeros, *Topology and its Applications*, **253**, (2019), 48–56.
- [16] J. Huh, Combinatorial applications of the Hodge–Riemann relations, *Proceedings of the International Congress of Mathematicians* **3** (2018), 3079–3098.
- [17] K. Ishikawa, Hoste’s conjecture for the 2-bridge knots, *Proc. Amer. Math. Soc.* **147** (2019), 2245–2254.
- [18] V. F. R. Jones, Polynomial invariant for knots via von Neumann algebras, *Bull. Amer. Math. Soc. (N.S.)* **12**(1) (1985) 103–111.
- [19] I. D. Jong, Alexander polynomials of alternating knots of genus two, *Osaka J. Math.* **46**(2) (2009) 353–371.
- [20] I. D. Jong, Alexander polynomials of alternating knots of genus two II, *J. Knot Theory Ramifications* **19** (8) (2010) 1075–1092.
- [21] S. Joshi and M. Prabhakar, Determinants of twisted generalized hybrid weaving knots, *Journal of Knot Theory and Its Ramifications*, **31** (14) (2022), 2250104.
- [22] C. Lee and F. Vafaee, On 3-braids and L-space Knots, *Geom. Dedicata* **213** (1) (2021), 513–521.
- [23] L. Lyubich and K. Murasugi, On zeros of the Alexander polynomial of an alternating knot, *Topology and its Applications*, **159** (1) (2012), 290–303.
- [24] E. Munarini and N. Zagaglia Salvi, On the rank polynomial of the lattice of order ideals of fences and crowns, *Discrete Mathematics*, **259**, 1–3, (2002) 163–177.
- [25] K. Murasugi, On the genus of the alternating knots I, *J. Math. Soc. Japan* **10** (1958) 94–105.
- [26] K. Murasugi, On the Alexander polynomial of the alternating knot, *Osaka Math. J.* **10** (1958) 181–189.
- [27] K. Murasugi, On closed 3-braids, *Memoirs of the American Mathematical Society*. 151 (AMS, Providence, RI 1974).
- [28] K. Murasugi, On the Alexander polynomial of alternating algebraic knots, *J. Austral. Math. Soc. Ser. A* **39**(3) (1985) 317–333.
- [29] OEIS Foundation Inc. (2023), Entry A205810 in The On-Line Encyclopedia of Integer Sequences, <https://oeis.org/A205810>
- [30] P. Ozsváth and Z. Szabó, Heegaard Floer homology and alternating knots, *Geom. Topol.* **7** (2003) 225–254.
- [31] K. Qazaqzeh and N. Chbili, A new obstruction of quasi-alternating links, *Algebr. Geom. Topol.* **15** (2015) 1847–1862.
- [32] Singh, V.K., Mishra, R. and Ramadevi, P. Colored HOMFLY-PT for hybrid weaving knot $W^3(m, n)$. *J. High Energ. Phys.* **63** (2021).
- [33] V. K. Singh and N. Chbili, Colored HOMFLY-PT polynomials of quasi-alternating 3-braid knots, *Nuclear Physics B*, **980**, (2022), 115800,
- [34] R. P. Stanley, Log-Concave and Unimodal Sequences in Algebra, Combinatorics, and Geometry, *Annals of the New York Academy of Sciences*, **576** (1989), 500–535.
- [35] A. Stoimenow, The skein polynomial of closed 3-braids *J. Reine Angew. Math.* **564** (2003), 167–180.
- [36] A. Stoimenow, Newton-like polynomials of links. *Enseign. Math. (2)* **51** 3–4 (2005), 211–230.
- [37] A. Stoimenow, Diagram Genus, Generators, and Applications Monographs and Research Notes in Mathematics, CRC Press, Boca Raton (2016)

DEPARTMENT OF MATHEMATICAL SCIENCES, COLLEGE OF SCIENCE, UNITED ARAB EMIRATES UNIVERSITY, 15551 AL AIN, U.A.E.

Email address: 700035655@uaeu.ac.ae

DEPARTMENT OF MATHEMATICAL SCIENCES, COLLEGE OF SCIENCE, UNITED ARAB EMIRATES UNIVERSITY, 15551 AL AIN, U.A.E.

Email address: nafaachbili@uaeu.ac.ae

URL: <http://faculty.uaeu.ac.ae/nafaachbili>