

# Propagation of chaos for point processes induced by particle systems with mean-field drift interaction\*

Nikolaos Kolliopoulos<sup>†</sup>      Martin Larsson<sup>‡</sup>      Zeyu Zhang<sup>§</sup>

March 22, 2023

## Abstract

We study the asymptotics of the point process induced by an interacting particle system with mean-field drift interaction. Under suitable assumptions, we establish propagation of chaos for this point process: it has the same weak limit as the point process induced by i.i.d. copies of the solution of a limiting McKean–Vlasov equation. This weak limit is a Poisson point process whose intensity measure is related to classical extreme value distributions. In particular, this yields the limiting distribution of the normalized upper order statistics.

## 1 Introduction

This paper extends our earlier work [5], which establishes propagation of chaos for the largest particle in a system of real-valued diffusive particles  $X^{1,N}, \dots, X^{N,N}$  with mean-field interaction through the drifts. In the present paper we do not only consider the maximum particle, but we study the point process  $\tilde{U}_N$  on  $[0, 1] \times \mathbb{R}$  defined by

$$\tilde{U}_N(I) = \sum_{i=1}^N \mathbf{1}_{\{(\frac{i}{N}, (X_T^{i,N} - b_T^N)/a_T^N) \in I\}} \quad (1.1)$$

for any Borel subset  $I \subset [0, 1] \times \mathbb{R}$ , where  $a_T^N$  and  $b_T^N$  are suitable normalizing constants. In particular, this allows us to derive the large  $N$  asymptotics for any fixed number  $k$  of upper order statistics. Our result can be viewed as a propagation of chaos property for the point process  $U_N$ . Alternatively, it can be viewed as an extension of classical extreme value theory for i.i.d. random variables to the case of mean-field interaction.

---

\*This work has been partially supported by the National Science Foundation under grant NSF DMS-2206062.

<sup>†</sup>Department of Mathematical Sciences, Carnegie Mellon University, [nkolliop@andrew.cmu.edu](mailto:nkolliop@andrew.cmu.edu).

<sup>‡</sup>Department of Mathematical Sciences, Carnegie Mellon University, [larsson@cmu.edu](mailto:larsson@cmu.edu).

<sup>§</sup>Department of Mathematical Sciences, Carnegie Mellon University, [zeyuzhan@andrew.cmu.edu](mailto:zeyuzhan@andrew.cmu.edu).

We work with the same particle dynamics as in [5]. Specifically, for each  $N \in \mathbb{N}$  the  $N$ -particle system evolves according to a stochastic differential equation of the form

$$dX_t^{i,N} = A(t, X_{[0,t]}^{i,N}) \left( B \left( t, X_{[0,t]}^{i,N}, \int g(t, X_{[0,t]}^{i,N}, \mathbf{y}_{[0,t]}) \mu_t^N(d\mathbf{y}) \right) dt + dW_t^i \right) + C(t, X_{[0,t]}^{i,N}) dt \quad (1.2)$$

for  $i = 1, \dots, N$ , with i.i.d. initial conditions  $X_0^{i,N} \sim \nu_0$ , where  $\nu_0$  is a given probability measure on  $\mathbb{R}$ . Moreover,  $W^i$ ,  $i \in \mathbb{N}$ , is family of independent standard Brownian motions, and we write  $\mathbf{x}_{[0,t]} = (x(s))_{s \in [0,t]}$  for any continuous function  $\mathbf{x}$ . For each  $t \in \mathbb{R}_+$  we defined the following empirical measure on trajectory up to time  $t$ ,

$$\mu_t^N = \frac{1}{N} \sum_{i=1}^N \delta_{X_{[0,t]}^{i,N}}. \quad (1.3)$$

The coefficients  $A(t, \mathbf{x}_{[0,t]})$ ,  $B(t, \mathbf{x}_{[0,t]}, r)$ ,  $C(t, \mathbf{x}_{[0,t]})$  and interaction function  $g(t, \mathbf{x}_{[0,t]}, \mathbf{y}_{[0,t]})$  are defined for all  $t \in \mathbb{R}_+$ ,  $\mathbf{x}, \mathbf{y} \in C(\mathbb{R}_+)$ , and  $r \in \mathbb{R}$ . Note that here we only allow interaction in the drift  $B$  but not in  $A$  or  $C$ . This restriction is crucial for the methods used later, which rely on the ability to use equivalent changes of probability measure to remove the interaction.

Classical propagation of chaos [8, 10, 3, 6] states, under suitable assumption, that for any fixed number  $k \in \mathbb{N}$ , the first  $k$  particles  $(X^{1,N}, \dots, X^{k,N})$  converge jointly as  $N \rightarrow \infty$  to  $k$  independent copies  $(X^1, \dots, X^k)$  of the solution to the McKean–Vlasov equation

$$dX_t = A(t, X_{[0,t]}) \left( B \left( t, X_{[0,t]}, \int g(t, X_{[0,t]}, \mathbf{y}_{[0,t]}) \mu_t(d\mathbf{y}) \right) dt + dW_t \right) + C(t, X_{[0,t]}) dt$$

$$\mu_t = \text{Law}(X_{[0,t]}) \quad (1.4)$$

with initial condition  $\mu_0 = \nu_0$ . See in particular [4, Theorem 2.1]. This leads to the heuristic that the interacting  $N$ -particle system behaves asymptotically like a system of i.i.d. copies of the solution of the limiting McKean–Vlasov equation. This suggests that the large- $N$  asymptotics of the point process  $\tilde{U}_N$  in (1.1) should match the asymptotics of the point process  $U_N$  induced by the independent copies  $X^i$  of the solution of (1.4), namely

$$U_N(I) = \sum_{i=1}^N \mathbf{1}_{\left\{ \left( \frac{i}{N}, (X_T^i - b_T^N)/a_T^N \right) \in I \right\}}. \quad (1.5)$$

Due to the i.i.d. property, the latter falls within the framework of classical extreme value theory, and its asymptotic behavior is therefore well-known. Specifically, under suitable assumptions,  $U_N$  converges weakly to a Poisson random measure whose intensity measure depends on the so-called extreme value index of the distribution of  $X_T^1$ . See [2, Theorem 2.1.2] for a precise statement and, more generally, [2, 9] for an introduction about extreme value theory.

Our main result, Theorem 2.3, states (under assumptions) that the above intuition is correct:  $\widetilde{U}_N$  converges to the same Poisson point process as  $U_N$ . However, to prove this, existing results on standard propagation of chaos, such as the rate of convergence statements in [4, 7], appear to be insufficient. Instead, we rely on a method developed in our earlier paper [5]. As in [1, 6, 4], our method starts with a Girsanov transformation. We then depart from earlier methods by performing an iterated integral expansion of the Radon–Nikodym derivative. By carefully controlling the terms in this expansion, we are able to complete the proof.

The rest of the paper organized as follow: we discuss the assumptions, notation, and state our main result precisely in Section 2. We then discuss some examples and consequences of our main result in Section 3. In Section 5, we give the proof of the main result, Theorem 2.3. We will frequently use the notation

$$[n] = \{1, \dots, n\} \text{ for any } n \in \mathbb{N} = \{1, 2, \dots\},$$

and  $\mathbb{R}_+ = [0, \infty)$ . We will allow generic constants  $C$  to vary from line to line, and occasionally indicate the dependence on parameters by writing  $C(n)$ ,  $C(p, n)$ , etc. We also use the notation

$$\mathcal{F}_t^V = \sigma(X_s^i, W_s^i : s \leq t, i \in V) \text{ for any index set } V \subset \mathbb{N},$$

and write  $\mathbb{L}$  for the space of all progressively measurable processes  $Y$  with locally integrable moments.

## 2 Assumptions and main result

We work in the same setup as in our previous work, and under the same assumption; see [5, Assumption 2.1, Assumption 2.3]. For the convenience of the reader, we recall these here.

**Assumption 2.1.** *The coefficient functions  $(t, \mathbf{x}) \mapsto A(t, \mathbf{x}_{[0,t]})$ ,  $(t, \mathbf{x}, r) \mapsto B(t, \mathbf{x}_{[0,t]}, r)$ ,  $(t, \mathbf{x}) \mapsto C(t, \mathbf{x}_{[0,t]})$  and the interaction function  $(t, \mathbf{x}, \mathbf{y}) \mapsto g(t, \mathbf{x}_{[0,t]}, \mathbf{y}_{[0,t]})$  are real-valued measurable functions on  $\mathbb{R}_+ \times C(\mathbb{R}_+)$ ,  $\mathbb{R}_+ \times C(\mathbb{R}_+) \times \mathbb{R}$ ,  $\mathbb{R}_+ \times C(\mathbb{R}_+)$ , and  $\mathbb{R}_+ \times C(\mathbb{R}_+) \times C(\mathbb{R}_+)$ , respectively. They satisfy the following conditions:*

- *A and C are uniformly bounded,*
- *for every  $t \in \mathbb{R}_+$  and  $\mathbf{x} \in C(\mathbb{R}_+)$ , the function  $r \mapsto B(t, \mathbf{x}_{[0,t]}, r)$  is twice continuously differentiable, and its first and second derivatives are bounded uniformly in  $(t, \mathbf{x})$ .*

As in [5], we avoid imposing specific conditions for well-posedness of the limiting equation, but rather assume existence directly (uniqueness is not actually required, so we do not assume it.) We also assume certain moment bounds. Two examples where these assumptions can be verified are discussed below; see Examples 3.1 and 3.2. See also [5, Remark 2.7 and Section 3].

**Assumption 2.2.** Fix a probability measure  $\nu_0$  on  $\mathbb{R}$  and assume that the McKean–Vlasov equation (1.4) admits a weak solution  $(X, W)$  with  $X_0 \sim \nu_0$ . Construct (for instance as a countable product) a filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$  with a countable sequence  $(X^i, W^i)$ ,  $i \in \mathbb{N}$  of independent copies of  $(X, W)$ . Then, assume that there is a continuous function  $K(t)$  such that for all  $p \in \mathbb{N}$ ,  $t \in \mathbb{R}_+$ ,  $N \in \mathbb{N}$ , and  $i, j \in [N]$ , one has the moment bounds

$$\mathbb{E} \left[ g(t, X_{[0,t]}^i, X_{[0,t]}^j)^{2p} \right] \leq p! K(t)^p \quad (2.1)$$

and

$$\mathbb{E} \left[ \left( \int g(t, X_{[0,t]}^i, \mathbf{y}_{[0,t]}) (\mu_t^N - \mu_t)(d\mathbf{y}) \right)^{2p} \right] \leq \frac{1}{N^p} p! K(t)^p. \quad (2.2)$$

We used the same notation as in [9]. In general a point process  $U$  is a random distribution of points in a state space  $E$  which is locally compact with a countable basis. Let  $\varepsilon$  be the Borel  $\sigma$ -algebra of subsets of  $E$ . More precisely letting  $M_p(E)$  denote the space of all point measures on  $E$  and define a  $\sigma$ -algebra  $\mathcal{M}_p(E)$  of subsets of  $M_p(E)$  to be the smallest  $\sigma$ -algebra making all evaluation maps  $m \rightarrow m(F)$  measurable for all  $F \in \varepsilon$ . Then  $U$  is a measurable map from some probability space  $(\Omega, \mathcal{F})$  into  $(M_p(E), \mathcal{M}_p(E))$ . Readers may refer to [9, Section 3] for the background of point process. In our situation, the state space  $E$  for  $U_N, \tilde{U}_N$  is  $[0, 1] \times \mathbb{R}$ .

In preparation for the Theorem 2.3, we now introduce some notation and definitions. Suppose Assumptions 2.1 and 2.2 are satisfied. For each  $N \in \mathbb{N}$  we use the processes  $(X^i, W^i)$  to construct the  $N$ -particle systems through a change of probability measure. The  $N$ -particle empirical measure is defined by

$$\mu_t^N = \frac{1}{N} \sum_{i=1}^N \delta_{X_{[0,t]}^i}. \quad (2.3)$$

We then define a Radon–Nikodym density process by

$$Z^N = \exp(M^N - \frac{1}{2} \langle M^N \rangle) \quad (2.4)$$

where

$$M_t^N = \sum_{i=1}^N \int_0^t \Delta B_s^{i,N} dW_s^i \quad (2.5)$$

and

$$\begin{aligned} \Delta B_t^{i,N} = & B \left( t, X_{[0,t]}^i, \int g(t, X_{[0,t]}^i, \mathbf{y}_{[0,t]}) \mu_t^N(d\mathbf{y}) \right) \\ & - B \left( t, X_{[0,t]}^i, \int g(t, X_{[0,t]}^i, \mathbf{y}_{[0,t]}) \mu_t(d\mathbf{y}) \right). \end{aligned} \quad (2.6)$$

Under Assumptions 2.1 and 2.2,  $M^N$  is well-defined and a positive martingale. Moreover,  $\mathbb{E}[Z_T^N] = 1$  for all  $T \in (0, \infty)$ , which implies that the positive local martingale  $Z^N$  is in fact a true martingale. These facts are argued in [5]; see the discussion leading up to Theorem 2.4 in that paper.

Since  $Z^N$  is a true martingale, it induces a locally equivalent probability measure  $\mathbb{Q}^N \sim_{\text{loc}} \mathbb{P}$  under which the processes defined by

$$W_t^{i,N} = W_t^i - \int_0^t \Delta B_s^{i,N} ds$$

are mutually independent standard Brownian motions. Thus under  $\mathbb{Q}^N$  we find that  $X^1, \dots, X^N$  follow the  $N$ -particle dynamics (1.2),

$$dX_t^i = A(t, X_{[0,t]}^i) \left( B \left( t, X_{[0,t]}^i, \int g(t, X_{[0,t]}^i, \mathbf{y}_{[0,t]}) \mu_t^N(d\mathbf{y}) \right) dt + dW_t^{i,N} \right) + C(t, X_{[0,t]}^i) dt. \quad (2.7)$$

We thus model the point process  $\tilde{U}_N$  in (1.1) by  $U_N$  (1.5) viewed under  $\mathbb{Q}^N$ , and our main result establishes convergence of the laws  $\mathbb{Q}^N \circ U_N^{-1}$ . The following is our main result.

**Theorem 2.3.** *Suppose Assumptions 2.1 and 2.2 are satisfied. Fix  $T \in (0, \infty)$  and suppose that for some normalizing constants  $a_T^N, b_T^N$  the normalized maxima of the i.i.d. system (1.4) converge weakly to a nondegenerate distribution function  $\Gamma_T$  on  $\mathbb{R}$ :*

$$\mathbb{P} \left( \max_{i \leq N} \frac{X_T^i - b_T^N}{a_T^N} \leq x \right) \rightarrow \Gamma_T(x) \text{ as } N \rightarrow \infty, \quad x \in \mathbb{R}. \quad (2.8)$$

*Then, the sequence  $(\mathbb{Q}^N \circ U_N^{-1})_{N \in \mathbb{N}}$  of probability laws on  $M_p(E)$  converges to the law of a Poisson random measure on  $(0, 1] \times (x_*, x^*]$  with intensity measure  $\nu$ , where*

$$\nu((a, b] \times (c, d]) = (b - a)((1 + \gamma c)^{-1/\gamma} - (1 + \gamma d)^{-1/\gamma}), \quad a < b, \quad c < d.$$

*Here  $x_*, x^*$  are the left and right endpoints of the support of the limiting distribution  $\Gamma_T$ , and  $\gamma$  is the extreme value index.*

**Remark 2.4.** The condition (2.8) is equivalent to the weak convergence of  $\mathbb{P} \circ U_N^{-1}$  to a Poisson random measure on  $(0, 1] \times (x_*, x^*]$  with the same intensity measure  $\nu$  as in Theorem 2.3. One may check this by looking at [2, Theorem 2.1.2].

**Remark 2.5.** Readers who are not familiar with the extreme value index may refer to [2, Definition 1.1.4]. In the case  $\gamma = 0$ , the expression  $(1 + \gamma x)^{-1/\gamma}$  is interpreted as  $\exp(-x)$ . Note that the assumptions of Theorem 2.3 are precisely those of the main theorem in [5], even though the conclusion is stronger in that it applies to the full point process rather than to the maximum particle only.

We now introduce some additional notation. This will prepare us to state an auxiliary result that simplifies the proof of Theorem 2.3. We write  $D_3B(t, \mathbf{x}_{[0,t]}, r)$  for the derivative with respect to  $r$  and define the process

$$D_3B_t^i = D_3B \left( t, X_{[0,t]}^i, \int g(t, X_{[0,t]}^i, \mathbf{y}_{[0,t]}) \mu_t(d\mathbf{y}) \right). \quad (2.9)$$

Note that  $D_3B^i$  is adapted to the filtration  $(\mathcal{F}_t^{\{i\}})_{t \geq 0}$  generated by  $(X^i, W^i)$ . We also write

$$H_t^i(\mu) = \int g(t, X_{[0,t]}^i, \mathbf{y}_{[0,t]}) \mu(d\mathbf{y}) \quad (2.10)$$

for any signed measure  $\mu$  on  $C(\mathbb{R}_+)$ . We then have the Taylor formula

$$\Delta B_t^{i,N} = D_3B_t^i H_t^i(\mu_t^N - \mu_t) + R_t^{i,N} (H_t^i(\mu_t^N - \mu_t))^2,$$

where  $R^{i,N}$  is a process which is uniformly bounded in terms of the bound on the second derivative of  $r \mapsto B(t, \mathbf{x}_{[0,t]}, r)$  given by Assumption 2.1. Furthermore, we define the local martingales

$$\widetilde{M}_t^N = \sum_{i=1}^N \int_0^t D_3B_s^i H_s^i(\mu_s^N - \mu_s) dW_s^i. \quad (2.11)$$

With all these preparations, we are ready to introduce an auxiliary theorem which is a crucial intermediate step toward establishing Theorem (2.3). Recall that  $X^1, \dots, X^N$  are i.i.d. copies of the solution to (1.4), and that  $Z_T^N$  is defined in (2.4). For each  $m \in \mathbb{N}$  and  $0 \leq s \leq t$ , we define the  $m$ -fold iterated integral  $\widetilde{I}_m^N(s, t)$  by

$$\widetilde{I}_m^N(s, t) = \int_s^t \int_s^{t_1} \dots \int_s^{t_{m-1}} d\widetilde{M}_{t_m}^N \dots d\widetilde{M}_{t_1}^N. \quad (2.12)$$

**Theorem 2.6.** *Let Assumptions 2.1 and 2.2 be in force and fix  $T \in (0, \infty)$ . Let  $\phi_N: \mathbb{R}^N \rightarrow \mathbb{R}$ ,  $N \in \mathbb{N}$ , be a sequence of measurable functions such that*

$$\sup_{N \in \mathbb{N}, (x_1, \dots, x_N) \in \mathbb{R}^N} |\phi_N(x_1, \dots, x_N)| < \infty.$$

*To show that*

$$\lim_{N \rightarrow \infty} \mathbb{E} [\phi_N(X_T^1, \dots, X_T^N) (Z_T^N - 1)] = 0, \quad (2.13)$$

*it suffices to show that*

$$\lim_{N \rightarrow \infty} \left| \mathbb{E} \left[ \phi_N(X_T^1, \dots, X_T^N) \widetilde{I}_{k_1}^N(T_0, T_1) \dots \widetilde{I}_{k_\alpha}^N(T_{\alpha-1}, T_\alpha) \right] \right| = 0 \quad (2.14)$$

*for all  $\alpha \in [n]$  and all  $\alpha$ -tuples  $(k_1, \dots, k_\alpha)$  such that  $k_\beta \in [m_\beta] \cup \{0\}$  for  $\beta < \alpha$  and  $k_\alpha \in [m_\alpha]$ , where  $m_\beta$  ( $\beta = 1, \dots, \alpha$ ) are some positive integers not depending on  $N$ , and  $T_i = iT/n$ .*

### 3 Consequences of the main result and examples

We now discuss some examples and applications of our main result.

#### 3.1 Marginal point process

The point process  $U_N$  in (1.5) is defined over the product space  $(0, 1] \times (x_*, x^*]$ . The first coordinate, which accounts for the index of the particle, is included for technical reasons: it ensures that no two points overlap. Using Theorem 2.3 it is straightforward to derive a result for the marginal point process which excludes this coordinate. Indeed, by setting  $I = (0, 1] \times I'$  where  $I'$  is a subset of  $(x_*, x^*]$ , we obtain that the point process defined on  $(x_*, x^*]$  by

$$H_N(I') = \sum_{i=1}^N \mathbf{1}_{\{(X_T^i - b_T^N)/a_T^N \in I'\}}.$$

According to Theorem 2.3, the laws  $\mathbb{Q}^N \circ H_N^{-1}$  converges weakly to a Poisson point process  $H_\infty$  on  $(x_*, x^*]$  with intensity measure  $\mu((a, b]) = (1 + a\gamma)^{-1/\gamma} - (1 + b\gamma)^{-1/\gamma}$ . The laws  $\mathbb{P} \circ H_N^{-1}$  converges weakly to the same Poisson point process  $H_\infty$  with the same intensity measure  $\mu$  according to [2, Theorem 2.1.2].

#### 3.2 Propagation of chaos for order statistics

For any  $N$  particles in (1.2), we use the following notation to denote the upper order statistics:

$$X_T^{(1),N} \geq X_T^{(2),N} \geq X_T^{(3),N} \geq \dots \geq X_T^{(N),N}.$$

As discussed in Section 2, we model the  $N$  populated mean-field interacting particle system (1.2) by the  $N$  independent copies of the solution of the McKean–Vlasov equation (1.4) under  $\mathbb{Q}^N$ . Hence, we model the order statistics of the mean-field interacting particles

$$X_T^{(1),N} \geq X_T^{(2),N} \geq X_T^{(3),N} \geq \dots \geq X_T^{(N),N}$$

via the order statistics of the  $N$  independent copies of the solution of the McKean–Vlasov equation

$$X_T^{(1)} \geq X_T^{(2)} \geq X_T^{(3)} \geq \dots \geq X_T^{(N)}$$

under  $\mathbb{Q}^N$ . The time- $T$  marginal distribution of the rank- $k$  particle  $X_T^{(k)}$  has the representation

$$\mathbb{Q}^N \left( \frac{X_T^{(k)} - b_T^N}{a_T^N} \geq x \right) = \mathbb{Q}^N (H_N([x, \infty)) \geq k).$$

Thus, in view of Subsection 3.1 above, we have

$$\lim_{N \rightarrow \infty} \mathbb{Q}^N \left( \frac{X_T^{(k)} - b_T^N}{a_T^N} \geq x \right) = \lim_{N \rightarrow \infty} \mathbb{Q}^N (H_N([x, \infty)) \geq k) = \mathbb{P}(H_\infty([x, \infty)) \geq k),$$

where  $H_\infty([x, \infty))$  has a Poisson distribution with parameter  $(1 + x\gamma)^{-1/\gamma}$  under  $\mathbb{P}$ . The same reasoning applies to the system of independent copies, resulting in

$$\mathbb{P}\left(\frac{X_T^{(k)} - b_T^N}{a_T^N} \geq x\right) = \mathbb{P}(H_N([x, \infty)) \geq k),$$

and hence the propagation of chaos property holds for the  $k$ -th upper order statistic,

$$\lim_{N \rightarrow \infty} \mathbb{Q}^N \left( \frac{X_T^{(k)} - b_T^N}{a_T^N} \geq x \right) = \lim_{N \rightarrow \infty} \mathbb{P} \left( \frac{X_T^{(k)} - b_T^N}{a_T^N} \geq x \right),$$

where

$$X_T^{(1)} \geq X_T^{(2)} \geq \dots \geq X_T^{(N)}. \quad (3.1)$$

Thus, the  $k$ th upper order statistic from the interacting particle system (1.2) has the same weak limit as the  $k$ th upper order statistics from the independent copies of the limiting McKean–Vlasov system.

### 3.3 Propagation of chaos for the joint distribution of the top $k$ order statistics

We now discuss the asymptotic joint distribution of the fixed time top- $k$  ranked particles of the mean–field interacting particle system,

$$\lim_{N \rightarrow \infty} \mathbb{Q}^N \left( \frac{X_T^{(1)} - b_T^N}{a_T^N} \geq x_1, \dots, \frac{X_T^{(k)} - b_T^N}{a_T^N} \geq x_k \right), \quad x_1 \geq x_2 \geq \dots \geq x_k.$$

Note that

$$\begin{aligned} & \mathbb{Q}^N \left( \frac{X_T^{(1)} - b_T^N}{a_T^N} \geq x_1, \dots, \frac{X_T^{(k)} - b_T^N}{a_T^N} \geq x_k \right) \\ &= \mathbb{Q}^N (H_N([x_1, \infty)) \geq 1, H_N([x_2, \infty)) \geq 2, \dots, H_N([x_k, \infty)) \geq k) \\ &= \sum_{\{i_1, \dots, i_k\} \in S_k} \mathbb{Q}^N (H_N([x_1, \infty)) = i_1, H_N([x_2, x_1]) = i_2, \dots, H_N([x_k, x_{k-1}]) = i_k) \end{aligned}$$

where the set  $S_k$  is defined as

$$S_k = \{(i_1, \dots, i_k) : i_1 + \dots + i_j \geq j \quad \forall j \in [k]\}.$$

Moreover, if we consider the order statistics (3.1) of the  $N$  copies of the solution to the limiting McKean–Vlasov equation, we have the following equation which establish the prop-



agation of chaos property for the top  $k$  order statistics.

$$\begin{aligned}
& \lim_{N \rightarrow \infty} \mathbb{Q}^N \left( \frac{X_T^{(1)} - b_T^N}{a_T^N} \geq x_1, \dots, \frac{X_T^{(k)} - b_T^N}{a_T^N} \geq x_k \right) \\
&= \lim_{N \rightarrow \infty} \sum_{\{i_1, \dots, i_k\} \in S_k} \mathbb{Q}^N (H_N([x_1, \infty)) = i_1, H_N([x_2, x_1]) = i_2, \dots, H_N([x_k, x_{k-1}]) = i_k) \\
&= \sum_{\{i_1, \dots, i_k\} \in S_k} \mathbb{P} (H_\infty([x_1, \infty)) = i_1, H_\infty([x_2, x_1]) = i_2, \dots, H_\infty([x_k, x_{k-1}]) = i_k) \\
&= \lim_{N \rightarrow \infty} \mathbb{P} \left( \frac{X_T^{(1)} - b_T^N}{a_T^N} \geq x_1, \dots, \frac{X_T^{(k)} - b_T^N}{a_T^N} \geq x_k \right).
\end{aligned}$$

On the other hand, for each fixed tuple  $(i_1, \dots, i_k)$ , as discussed in Section 3.1, we know that the limiting distribution of  $H_N$  is a Poisson point process. Thus for any non-overlapping intervals  $I_1, I_2$ ,  $H_\infty(I_1)$  is independent from  $H_\infty(I_2)$ . Hence we have

$$\begin{aligned}
& \lim_{N \rightarrow \infty} \mathbb{Q}^N (H_N([x_1, \infty)) = i_1, H_N([x_2, x_1]) = i_2, \dots, H_N([x_k, x_{k-1}]) = i_k) \\
&= \mathbb{P} (H_\infty([x_1, \infty)) = i_1, H_\infty([x_2, x_1]) = i_2, \dots, H_\infty([x_k, x_{k-1}]) = i_k) \\
&= \prod_{j=1}^k \mathbb{P} (H_\infty([x_j, x_{j-1}]) = i_j) \\
&= \prod_{j=1}^k \frac{\lambda_j^{i_j} e^{-\lambda_j}}{i_j!}
\end{aligned}$$

where  $\lambda_j = (1 + \gamma x_j)^{-1/\gamma} - (1 + \gamma x_{j-1})^{-1/\gamma}$ . Thus we have:

$$\lim_{N \rightarrow \infty} \mathbb{Q}^N \left( \frac{X_T^{(1)} - b_T^N}{a_T^N} \geq x_1, \dots, \frac{X_T^{(k)} - b_T^N}{a_T^N} \geq x_k \right) = \sum_{\{i_1, \dots, i_k\} \in S_k} \prod_{j=1}^k \frac{\lambda_j^{i_j} e^{-\lambda_j}}{i_j!}. \quad (3.2)$$

There is another way to characterize this limit. Due to [2, Theorem 2.1.1], the normalized top  $k$  ranked particles of the independent system,

$$\left( \frac{X_T^{(1)} - b_T^N}{a_T^N}, \frac{X_T^{(2)} - b_T^N}{a_T^N}, \dots, \frac{X_T^{(k)} - b_T^N}{a_T^N} \right),$$

converge in distribution to

$$\left( \frac{(E_1)^{-\gamma} - 1}{\gamma}, \frac{(E_1 + E_2)^{-\gamma} - 1}{\gamma}, \dots, \frac{(E_1 + E_2 + \dots + E_k)^{-\gamma} - 1}{\gamma} \right),$$

where  $\gamma$  is the extreme value index, and  $E_i$  are i.i.d exponential distribution. Theorem 2.3 implies that this limit coincides with the one in (3.2).

### 3.4 Propagation of chaos when $k \rightarrow \infty$

Let us comment briefly on the *intermediate order statistics* of the interacting particle system (1.2). The intermediate order statistics are particles  $X_t^{k(N),N}$  where the rank  $k(N)$  increases with  $N$  in such a way that  $k(N)/N \rightarrow 0$  as  $N \rightarrow \infty$ . Let us consider the general case where  $k := k(N) \rightarrow \infty$ . Our previous computations show that

$$\begin{aligned} \lim_{N \rightarrow \infty} \mathbb{Q}^N \left( \frac{X_T^{k(N)} - b_T^N}{a_T^N} \geq x \right) &= \lim_{N \rightarrow \infty} \mathbb{Q}^N (H_N([x, \infty)) \geq k(N)) \\ &= \lim_{N \rightarrow \infty} \mathbb{Q}^N \left( \frac{H_N([x, \infty))}{k(N)} \geq 1 \right) \\ &= 0. \end{aligned}$$

This indicates that our results do not yield any information about the asymptotic behavior of the  $k(N)$ -th order statistics. In particular, our results do not yield any information about the asymptotic behavior of intermediate order statistics as well. In fact, it is well-known in the classical extreme value literature that certain second-order conditions are required to establish convergence of suitably normalized intermediate order statistics. An interesting open question is whether those conditions are also sufficient to obtain the analogous result for the interacting systems that we consider here, but this is outside the scope of the current paper.

### 3.5 Two examples

We now discuss two examples that illustrate the main result as well as its applications.

**Example 3.1** (Gaussian particles). We build on [5, Example 3.1]. Consider the following Gaussian particle system:

$$X_t^{i,N} = X_0^i - \kappa \int_0^t \left( X_s^{i,N} - \frac{1}{N} \sum_{j=1}^N X_s^{j,N} \right) ds + \sigma W_t^i, \quad i = 1, \dots, N,$$

with i.i.d.  $N(m_0, \sigma_0^2)$  initial conditions. Here  $\kappa, m_0 \in \mathbb{R}$  and  $\sigma, \sigma_0 \in (0, \infty)$  are parameters. In our setting this example arises by taking  $A(t, \mathbf{x}_{[0,t]}) = \sigma$ ,  $B(t, \mathbf{x}_{[0,t]}, r) = -\kappa(x_t - r)/\sigma$ ,  $C(t, \mathbf{x}_{[0,t]}) = 0$ , and  $g(t, \mathbf{x}_{[0,t]}, \mathbf{y}_{[0,t]}) = y_t$ . Clearly Assumption 2.1 is satisfied. The McKean–Vlasov equation (1.4) reduces to

$$dX_t = -\kappa (X_t - \mathbb{E}[X_t]) dt + \sigma dW_t.$$

From the discussion in [5, Example 3.1] we know that for suitable parameters  $a_T^N, b_T^N$ ,

$$\mathbb{P} \left( \max_{i \leq N} \frac{X_T^i - b_T^N}{a_T^N} \leq x \right) \rightarrow \exp(-e^{-x}) \text{ as } N \rightarrow \infty.$$

That is,  $X_T$  belongs to the Gumbel domain of attraction. Our main result Theorem 2.3 applies, and by the discussion in Subsection 3.1, we know that the marginal point process converges weakly to a Poisson random measure on the real line  $\mathbb{R}$  with intensity measure  $\mu((a, b]) = \exp(-a) - \exp(-b)$ . Thanks to (3.2) we have the following asymptotics:

$$\lim_{N \rightarrow \infty} \mathbb{P} \left( X_T^{(1),N} \geq x_1, \dots, X_T^{(k),N} \geq x_k \right) = \sum_{\{i_1, \dots, i_k\} \in S_k} \prod_{j=1}^k \frac{\lambda_j^{i_j} e^{-\lambda_j}}{i_j!},$$

where  $\lambda_j = \exp(-x_j) - \exp(-x_{j-1})$  and the set  $S_k$  is defined as

$$S_k = \{(i_1, \dots, i_k) : i_1 + \dots + i_k \geq j \quad \forall j \in [k]\}.$$

**Example 3.2** (Rank-based diffusions). This is another example from [5]. Consider the  $N$ -particle system evolving according to

$$dX_t^{i,N} = B \left( F_t^N(X_t^{i,N}) \right) dt + \sqrt{2} dW_t^i$$

for  $i \in [N]$ , where  $B(r)$  is a twice continuously differentiable function on  $[0, 1]$  and

$$F_t^N(x) = \frac{1}{N} \sum_{j=1}^N \mathbf{1}_{\{X_t^{j,N} \leq x\}}$$

is the empirical distribution function. If  $X$  is stationary, an sufficient condition for the stationary distribution of the limiting McKean–Vlasov equation was given in Example 3.2 in [5]. Under this condition, the limiting stationary distribution belongs to the Gumbel domain of attraction, which is exactly the same as in the previous example. Hence we may apply the main Theorem 2.3 to derive the weak limit of the marginal point process as well as the asymptotic joint distribution of the top  $k$  order statistics.

## 4 Proof of Theorem 2.6

The proof of the theorem is exactly the same word by word repeating step 1-3 and the first part of step 4, leading up to (6.13) in [5] with

$$\prod_{i=1}^N \mathbf{1}_{\{X_T^i < X_N\}}$$

replaced with

$$\phi_N(X_T^1, \dots, X_T^N).$$

Showing that there exists  $n$  such that for all  $\epsilon > 0$ , we have

$$\mathbb{E} [\phi_N(X_T^1, \dots, X_T^N)(Z_T^N - 1)] = \sum_{\alpha=1}^n A_\alpha^N,$$

where

$$|A_\alpha^N| \leq \alpha\epsilon + \frac{C}{\sqrt{N}} + \sum_{k_1, \dots, k_\alpha} \left| \mathbb{E} \left[ \phi_N(X_T^1, \dots, X_T^N) \tilde{I}_{k_1}^N(T_0, T_1) \cdots \tilde{I}_{k_\alpha}^N(T_{\alpha-1}, T_\alpha) \right] \right|. \quad (4.1)$$

The sum in (4.1) ranges over all  $(k_1, \dots, k_\alpha)$  such that  $k_\beta \in [m_\beta] \cup \{0\}$  for  $\beta < \alpha$  and  $k_\alpha \in [m_\alpha]$ . Note that  $\epsilon$  is arbitrary,  $n$  and constant  $C$  here do not depend on  $N$ , hence to show (2.13) holds, it suffices to show (2.14) holds for all  $\alpha \in [n]$  and all  $\alpha$ -tuples  $(k_1, \dots, k_\alpha)$  such that  $k_\beta \in [m_\beta] \cup \{0\}$  for  $\beta < \alpha$  and  $k_\alpha \in [m_\alpha]$ , where  $m_\beta$  ( $\beta = 1, \dots, \alpha$ ) are some positive integers not depending on  $N$ , and  $T_i = iT/n$ .

## 5 Proof of Theorem 2.3

To prove Theorem 2.3, we employ a variation of [9, Proposition 3.22] which provides conditions under which a sequence  $\{\xi_n\}_{n \in \mathbb{N}}$  of point processes converges weakly to a candidate limit  $\xi$ . Observe that the proof of that result uses [9, Equation (3.24)] only for showing that the sequence  $\{\xi_n\}_{n \in \mathbb{N}}$  is tight and for obtaining the inequality

$$\mathbb{E}[\eta(I)] \leq \mathbb{E}[\xi(I)] \quad (5.1)$$

for any weak limit  $\eta$  of  $\{\xi_n\}_{n \in \mathbb{N}}$  (through some subsequence) and any set  $I$  taken from a certain collection  $\mathcal{T}$  of subsets of the state space  $E$ . The inequality (5.1) for all  $I \in \mathcal{T}$  is established at the end of the proof as part of showing that the arbitrary weak limit  $\eta$  is a simple point process. Therefore, we are going to replace [9, Equation (3.24)] with another condition that implies both the tightness of  $\{\xi_n\}_{n \in \mathbb{N}}$  and (5.1).

We start by noting that we only need (5.1) to hold as an equality, which is equivalent to

$$\mathbb{P}(\eta(I) = t) = \mathbb{P}(\xi(I) = t)$$

for each  $t \in \mathbb{N}$  and all  $I \in \mathcal{T}$ . By the continuity argument immediately after [9, Display (3.28)], the latter reduces to having

$$\lim_{n \rightarrow \infty} \mathbb{P}(\xi_n(I) = t) = \mathbb{P}(\xi(I) = t) \quad (5.2)$$

for all  $I \in \mathcal{T}$  and  $t \in \mathbb{N}$ . We claim now that (5.2) is also sufficient for tightness. Indeed, by [9, Exercise 3.5.2], we only need to show that

$$\lim_{t \rightarrow \infty} \limsup_{n \rightarrow \infty} \mathbb{P}(\xi_n(K) > t) = 0 \quad (5.3)$$

for any compact subset  $K$  of the state space  $E$ . Thanks to the properties of  $\mathcal{T}$ , one can pick a set  $I \in \mathcal{T}$  such that  $K \subset I$ , so we can bound

$$\begin{aligned} \limsup_{N \rightarrow \infty} \mathbb{P}(\xi_n(K) > t) &\leq \limsup_{N \rightarrow \infty} \mathbb{P}(\xi_n(I) > t) \\ &= \mathbb{P}(\xi(I) > t), \end{aligned}$$

where the last equality follows from (5.2). Sending  $t \rightarrow \infty$  in the above estimate we get (5.3) so the required tightness follows.

Observe now that [9, Equation (3.23)] is also obtained from (5.2) by simply setting  $t = 0$ . Therefore, one can deduce the weak convergence  $\xi_n \rightarrow \xi$  as  $n \rightarrow \infty$  by verifying the following two conditions:

- (a)  $\mathbb{P}(\xi(\partial I) = 0) = 1$  for all  $I \in \mathcal{T}$ .
- (b)  $\lim_{n \rightarrow \infty} \mathbb{P}(\xi_n(I) = t) = \mathbb{P}(\xi(I) = t)$  for all  $I \in \mathcal{T}$  and  $t \in \mathbb{N}$ .

In our case,  $\xi = U_\infty$  is a Poisson point process with intensity  $\nu$  so that  $\mathbb{P} \circ U_N^{-1} \rightarrow \text{Law}(U_\infty)$  as  $N \rightarrow \infty$ ,  $\{\xi_n\}_{n \in \mathbb{N}}$  is a sequence of point processes  $\{\tilde{U}_N\}_{N \in \mathbb{N}}$  with  $\mathbb{Q}^N \circ U_N^{-1}$  being the law of  $\tilde{U}_N$  for all  $N \in \mathbb{N}$ ,  $E = (0, 1] \times (x_*, x^*]$ , and finally  $\mathcal{T} = \mathcal{I}$  where  $\mathcal{I}$  is the family of finite unions of rectangles of the form:

$$I = \bigcup_{j=1}^m (a_j, b_j] \times (c_j, d_j],$$

where  $m \in \mathbb{N}$ ,  $0 < a_j < b_j \leq 1$  and  $x_* < c_j < d_j \leq x^*$  for  $j \in [m]$ . Without loss of generality, we assume that all the rectangles that constitute each  $I$  are pairwise disjoint. Since  $\xi = U_\infty$  is a Poisson point process with intensity measure that is absolutely continuous with respect to the Lebesgue measure, (a) follows immediately. Then, since  $\mathbb{P} \circ U_N^{-1} \rightarrow \text{Law}(U_\infty)$  as  $N \rightarrow \infty$ , condition (b) translates to

$$\lim_{N \rightarrow \infty} \mathbb{Q}^N(U_N(I) = t) = \lim_{N \rightarrow \infty} \mathbb{P}(U_N(I) = t), \quad (5.4)$$

for any set  $I \in \mathcal{I}$  and any  $t \in \mathbb{N}$ . Therefore, the proof of our result reduces to showing that (5.4) is true for any set  $I \in \mathcal{I}$  and any  $t \in \mathbb{N}$ . This will be done by employing ideas from [5].

**Proof of condition (5.4):** Observe that what we want to show now is equivalent to

$$\lim_{N \rightarrow \infty} \mathbb{E} [\mathbf{1}_{\{U_N(I)=t\}} (Z_T^N - 1)] = 0, \quad (5.5)$$

for any  $I \in \mathcal{I}$  and any integer  $t \geq 0$ . Recalling Theorem 2.6 with  $\phi_N(X_T^1, \dots, X_T^N)$  taken to be the indicator  $\mathbf{1}_{\{U_N(I)=t\}}$  which is a bounded and measurable function of  $X_T^1, \dots, X_T^N$ , it suffices to show that

$$\lim_{N \rightarrow \infty} \left| \mathbb{E} \left[ \mathbf{1}_{\{U_N(I)=t\}} \tilde{I}_{k_1}^N(T_0, T_1) \cdots \tilde{I}_{k_\alpha}^N(T_{\alpha-1}, T_\alpha) \right] \right| = 0. \quad (5.6)$$

Under the convention

$$I_N = \bigcup_{j=1}^m (Na_j, Nb_j) \times (a_T^N c_j + b_T^N, a_T^N d_j + b_T^N)$$

and also

$$\mathbf{X}_T^i = (i, X_T^i),$$

we can re-write the indicator  $\mathbf{1}_{\{U_N(I)=t\}}$  as follows:

$$\begin{aligned} \mathbf{1}_{\{U_N(I)=t\}} &= \sum_{\{i_1, \dots, i_t\} \subset [N]} \prod_{l=1}^t \mathbf{1}_{\{\mathbf{X}_T^{i_l} \in I_N\}} \prod_{i \in [N] / \{i_1, \dots, i_t\}} \mathbf{1}_{\{\mathbf{X}_T^i \notin I_N\}} \\ &= \sum_{\{i_1, \dots, i_t\} \subset [N]} \prod_{l=1}^t \mathbf{1}_{\{\mathbf{X}_T^{i_l} \in I_N\}} \sum_{\kappa=0}^{N-t} (-1)^\kappa \sum_{\{i'_1, \dots, i'_\kappa\} \subset [N] / \{i_1, \dots, i_t\}} \prod_{l=1}^{\kappa} \mathbf{1}_{\{\mathbf{X}_T^{i'_l} \in I_N\}} \\ &= \sum_{\{i_1, \dots, i_t\} \subset [N]} \sum_{\kappa=0}^{N-t} (-1)^\kappa \sum_{\{i'_1, \dots, i'_\kappa\} \subset [N] / \{i_1, \dots, i_t\}} \prod_{l=1}^t \mathbf{1}_{\{\mathbf{X}_T^{i_l} \in I_N\}} \prod_{l=1}^{\kappa} \mathbf{1}_{\{\mathbf{X}_T^{i'_l} \in I_N\}}, \end{aligned} \quad (5.7)$$

and we proceed now to the expansion of the iterated integrals  $\tilde{I}_{k_\beta}^N(T_{\beta-1}, T_\beta)$  in equation (5.6). In view of (2.11) and the definitions of  $H_s^i(\mu_s^N - \mu_s)$  and  $\mu_s^N$  in (2.10) and (1.3) respectively, we can write

$$\tilde{M}_t^N = \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^N \int_0^t G_s^{ij} dW_s^i, \quad (5.8)$$

where

$$G_t^{ij} = D_3 B_t^i \left( g \left( t, X_{[0,t]}^i, X_{[0,t]}^j \right) - \int g \left( t, X_{[0,t]}^i, \mathbf{y}_{[0,t]} \right) \mu_t(d\mathbf{y}) \right) \quad (5.9)$$

for all  $i, j \in [N]$ . Recall that  $D_3 B_t^i$  is the partial derivative of drift function  $B(t, \mathbf{x}_{[0,t]}, r)$  with respect to its third argument as defined in (2.9). Plugging (5.8) and (5.9) into the definition of  $\tilde{I}_{k_\beta}^N(T_{\beta-1}, T_\beta)$ , see (2.12), gives

$$\tilde{I}_{k_\beta}^N(T_{\beta-1}, T_\beta) = \frac{1}{N^{k_\beta}} \sum_{\mathbf{i} \in [N]^{k_\beta}} \sum_{\mathbf{j} \in [N]^{k_\beta}} I_{\mathbf{i}, \mathbf{j}}^N(T_{\beta-1}, T_\beta), \quad (5.10)$$

where we use the iterated integral notation from [5, equation (4.1)], namely:

$$I_{\mathbf{i}, \mathbf{j}}^N(s, t) = \int_s^t G_{t_1}^{i_1, j_1} \int_s^{t_1} G_{t_2}^{i_2, j_2} \dots \int_s^{t_{k-1}} G_{t_k}^{i_k, j_k} dW_{t_k}^{i_k} \dots dW_{t_1}^{i_1} \quad (5.11)$$

for any  $k \in \mathbb{N}$  and any multiindices  $\mathbf{i} = (i_1, \dots, i_k) \in [N]^k$  and  $\mathbf{j} = (j_1, \dots, j_k) \in [N]^k$ . Using the representation of  $\tilde{I}_{k_\beta}^N(T_{\beta-1}, T_\beta)$  in (5.10), the product of iterated integrals appearing in (5.6) can be written as

$$\tilde{I}_{k_1}^N(T_0, T_1) \dots \tilde{I}_{k_\alpha}^N(T_{\alpha-1}, T_\alpha) = \frac{1}{N^{k_1 + \dots + k_\alpha}} \sum_{(\mathbf{i}_1, \mathbf{j}_1), \dots, (\mathbf{i}_\alpha, \mathbf{j}_\alpha)} \prod_{\beta=1}^{\alpha} I_{\mathbf{i}_\beta, \mathbf{j}_\beta}^N(T_{\beta-1}, T_\beta). \quad (5.12)$$

Now we can combine the identities (5.12) and (5.7) and apply the triangle inequality to get

$$\begin{aligned} & \left| \mathbb{E} \left[ \mathbf{1}_{\{U_N(I)=t\}} \tilde{I}_{k_1}^N(T_0, T_1) \dots \tilde{I}_{k_\alpha}^N(T_{\alpha-1}, T_\alpha) \right] \right| \\ & \leq \frac{1}{N^s} \sum_{\{i_1, \dots, i_t\} \subset [N]} \sum_{\kappa=0}^{N-t} \sum_{\{i'_1, \dots, i'_\kappa\} \subset [N] / \{i_1, \dots, i_t\}} \sum_{(\mathbf{i}_1, \mathbf{j}_1), \dots, (\mathbf{i}_\alpha, \mathbf{j}_\alpha)} \\ & \quad \left| \mathbb{E} \left[ \prod_{l=1}^t \mathbf{1}_{\{\mathbf{X}_T^{i_l} \in I_N\}} \prod_{l=1}^{\kappa} \mathbf{1}_{\{\mathbf{X}_T^{i'_l} \in I_N\}} \prod_{\beta=1}^{\alpha} I_{\mathbf{i}_\beta, \mathbf{j}_\beta}^N(T_{\beta-1}, T_\beta) \right] \right| \\ & = \frac{1}{N^s} \sum_{\kappa=0}^{N-t} \sum_{\{i_1, \dots, i_t\} \subset [N]} \sum_{\{i'_1, \dots, i'_\kappa\} \subset [N] / \{i_1, \dots, i_t\}} \sum_{(\mathbf{i}_1, \mathbf{j}_1), \dots, (\mathbf{i}_\alpha, \mathbf{j}_\alpha)} \\ & \quad \left| \mathbb{E} \left[ \prod_{l=1}^t \mathbf{1}_{\{\mathbf{X}_T^{i_l} \in I_N\}} \prod_{l=1}^{\kappa} \mathbf{1}_{\{\mathbf{X}_T^{i'_l} \in I_N\}} \prod_{\beta=1}^{\alpha} I_{\mathbf{i}_\beta, \mathbf{j}_\beta}^N(T_{\beta-1}, T_\beta) \right] \right|, \quad (5.13) \end{aligned}$$

where  $s = k_1 + \dots + k_\alpha$ . Therefore, to show (5.6), it suffices to show the last sum in the above goes to zero as  $N \rightarrow \infty$ .

We count now the number of non-zero terms in the last sum in (5.13). Note that each term in that sum has the form

$$\mathbb{E} \left[ \Psi \prod_{\beta=1}^{\alpha} I_{\mathbf{i}_\beta, \mathbf{j}_\beta}^N(T_{\beta-1}, T_\beta) \right] \quad (5.14)$$

where

$$\Psi = \prod_{l=1}^t \mathbf{1}_{\{\mathbf{X}_T^{i_l} \in A_N\}} \prod_{l=1}^{\kappa} \mathbf{1}_{\{\mathbf{X}_T^{i'_l} \in A_N\}},$$

is a bounded  $\mathcal{F}_T^K$ -measurable random variable for  $K := \{i_1, \dots, i_t, i'_1, \dots, i'_\kappa\}$ , each iterated integral  $I_{i_\beta, j_\beta}^N(T_{\beta-1}, T_\beta)$  is given by equation (5.11) in terms of the processes  $G^{ij}$ , and we can verify that for all  $V \subset [N]$ ,  $i \in V$ ,  $j \notin V$  we have:

$$\begin{aligned} \mathbb{E} [G_s^{ij} | \mathcal{F}_t^V] &= \mathbb{E} \left[ D_3 B_t^i \left( g \left( t, X_{[0,t]}^i, X_{[0,t]}^j \right) - \int g \left( t, X_{[0,t]}^i, \mathbf{y}_{[0,t]} \right) \mu_t(d\mathbf{y}) \right) | \mathcal{F}_t^V \right] \\ &= D_3 B_t^i \left( \mathbb{E} \left[ g \left( t, X_{[0,t]}^i, X_{[0,t]}^j \right) | \mathcal{F}_t^V \right] - \mathbb{E} \left[ g \left( t, X_{[0,t]}^i, X_{[0,t]}^j \right) | \mathcal{F}_t^V \right] \right) \\ &= 0 \end{aligned}$$

Thus, to give an upper bound for the number of terms that survive in the last sum in (5.13), we need to count the terms in that sum for which  $K$  violates both Conditions 1 and 2 of [5, Lemma 4.1]. By [5, Lemma 4.2], there are at most

$$\binom{N}{\kappa+t} (\kappa+t)(\kappa+t+1) \cdots (\kappa+t+s) N^{s-1},$$

choices of  $K$  with  $|K| = \kappa+t$  for which (5.14) is not zero. Moreover, the last sum in (5.13) consists of terms of the form (5.14) for  $|K| \geq t$ , where each term with  $|K| = \kappa+t$  for  $\kappa \geq 0$  appears precisely

$$\binom{t+\kappa}{t} = \binom{t+\kappa}{\kappa}$$

times. The latter follows from the fact that the term in (5.14) for that particular  $K$  emerges exactly once for each choice of the  $\kappa$  indices  $i'_1, \dots, i'_\kappa$  for which  $\{i'_1, \dots, i'_\kappa\} \subset K$ , by picking then the remaining  $t$  indices  $\{i_1, \dots, i_t\}$ . Hence, there are at most

$$\binom{t+\kappa}{t} \binom{N}{\kappa+t} (\kappa+t)(\kappa+t+1) \cdots (\kappa+t+s) N^{s-1}, \quad (5.15)$$

surviving terms of the form (5.14) in the last sum in (5.13) with  $|K| = t+\kappa$

Next, we are going to derive a common bound for all surviving terms in (5.13). Note that for any  $(i, x) \in I_N$ , one has  $x > a_T^N c + b_T^N$  where  $c = \min\{c_j : 1 \leq j \leq m\}$ . Moreover, from our assumptions we have

$$\begin{aligned} (1 - \mathbb{P}(X_T^1 > a_T^N c + b_T^N))^N &= \mathbb{P} \left( \max_{i \leq N} X_T^i < a_T^N c + b_T^N \right) \\ &= \mathbb{P} \left( \frac{\max_{i \leq N} X_T^i - b_T^N}{a_T^N} < c \right) \\ &\rightarrow \Gamma_T(c) > 0 \end{aligned}$$

as  $N \rightarrow \infty$ , which implies

$$N \mathbb{P}(X_T^1 > a_T^N c + b_T^N) \leq -N \log(1 - \mathbb{P}(X_T^1 > a_T^N c + b_T^N)) \leq C$$



for all  $N$  and some  $C > 0$ . Hence, we can combine the above to obtain

$$\begin{aligned}\mathbb{P}(\mathbf{X}_T^i \in I_N) &\leq P(X_T^i > a_T^N c + b_T^N) \\ &= P(X_T^1 > a_T^N c + b_T^N) \\ &\leq \frac{C}{N}.\end{aligned}\tag{5.16}$$

Furthermore, the estimations in [5, Page 30] give

$$\left\| I_{i_\beta, j_\beta}^N(T_{\beta-1}, T_\beta) \right\|_p \leq C(4\sqrt{p})^{k_\beta} \prod_{\ell=1}^{k_\beta} \sqrt{p} = Cp^{k_\beta}.\tag{5.17}$$

for some arbitrary constant  $C$ . Therefore, we can apply Hölder's inequality with some exponents  $p_N$  and  $q_N = p_N/(p_N - 1)$  and use also (5.16) to obtain

$$\begin{aligned}&\left| \mathbb{E} \left[ \prod_{l=1}^t \mathbf{1}_{\{\mathbf{X}_T^{i_l} \in I_N\}} \prod_{l=1}^{\kappa} \mathbf{1}_{\{\mathbf{X}_T^{i'_l} \in I_N\}} \prod_{\beta=1}^{\alpha} I_{i_\beta, j_\beta}^N(T_{\beta-1}, T_\beta) \right] \right| \\ &\leq \mathbb{E} \left[ \prod_{l=1}^t \mathbf{1}_{\{\mathbf{X}_T^{i_l} \in I_N\}} \prod_{l=1}^{\kappa} \mathbf{1}_{\{\mathbf{X}_T^{i'_l} \in I_N\}} \right]^{\frac{1}{q_N}} \prod_{\beta=1}^{\alpha} \left\| I_{i_\beta, j_\beta}^N(T_{\beta-1}, T_\beta) \right\|_{\alpha p_N} \\ &= \left( \prod_{l=1}^t \mathbb{P}(\mathbf{X}_T^{i_l} \in I_N) \prod_{l=1}^{\kappa} \mathbb{P}(\mathbf{X}_T^{i'_l} \in I_N) \right)^{\frac{1}{q_N}} \prod_{\beta=1}^{\alpha} \left\| I_{i_\beta, j_\beta}^N(T_{\beta-1}, T_\beta) \right\|_{\alpha p_N} \\ &\leq C \left( \frac{C}{N} \right)^{\frac{\kappa+t}{q_N}} p_N^s.\end{aligned}\tag{5.18}$$

which is the required common bound.

Finally, we apply on the last sum in (5.13) the upper bounds (5.15) and (5.18) we have derived for the numbers and the sizes of non-zero terms respectively, so taking also  $p_N = \ln N$  we can bound that sum by

$$\begin{aligned}&\frac{1}{N^s} \sum_{\kappa=0}^{N-t} \binom{t+\kappa}{t} \binom{N}{\kappa+t} N^{s-1} (\kappa+t) \cdots (\kappa+t+s) C [\ln N]^s \left( \frac{C}{N} \right)^{(\kappa+t)(1-\frac{1}{\ln N})} \\ &= \sum_{\kappa=t}^N \frac{1}{N} \binom{N}{\kappa} \binom{\kappa}{t} \kappa \cdots (\kappa+s) C [\ln N]^s \left( \frac{C}{N} \right)^{\kappa(1-\frac{1}{\ln N})} \\ &\leq \frac{C [\ln N]^s}{N} \sum_{\kappa=1}^N \binom{N}{\kappa} \kappa \cdots (\kappa+s+t) \left( \frac{C}{N} \right)^{\kappa(1-\frac{1}{\ln N})} \\ &\leq \frac{C [\ln N]^s}{N} (s+t+2)(s+t+1)^{2(s+t+1)} e^{Ce} (ce)^{s+t+1}\end{aligned}\tag{5.19}$$

In the above display, the last inequality came from applying [5, Lemma 4.6] with  $S = s + t$ , and we have also used the bound

$$\binom{\kappa}{t} \kappa \cdots (\kappa + s) \leq \kappa \cdots (\kappa + s + t).$$

Since  $s = k_1 + \cdots + k_\alpha$  and  $t$  do not depend on  $N$ , the right hand side of (5.19) vanishes as  $N \rightarrow \infty$ , which means that the last sum in (5.13) goes also to 0 as  $N \rightarrow \infty$ . This implies (5.6) and thus (5.5) which is equivalent to (5.4), so the proof of the theorem is now complete.

## References

- [1] D. A. Dawson and J. Gärtner. Large deviations from the McKean-Vlasov limit for weakly interacting diffusions. *Stochastics: An International Journal of Probability and Stochastic Processes*, 20(4):247–308, 1987. 3
- [2] L. de Haan and A. Ferreira. *Extreme Value Theory: An Introduction*. Springer Series in Operations Research and Financial Engineering. Springer New York, 2007. 2, 5, 7, 9
- [3] J. Gärtner. On the McKean-Vlasov limit for interacting diffusions. *Math. Nachr.*, 137:197–248, 1988. 2
- [4] J.-F. Jabir. Rate of propagation of chaos for diffusive stochastic particle systems via girsanov transformation. *arXiv: Probability*, 2019. <https://arxiv.org/abs/1907.09096>. 2, 3
- [5] N. Kolliopoulos, M. Larsson, and Z. Zhang. Propagation of chaos for maxima of particle systems with mean-field drift interaction. *arXiv preprint arXiv:2206.10018*, 2022. 1, 2, 3, 5, 10, 11, 13, 15, 16, 17, 18
- [6] D. Lacker. On a strong form of propagation of chaos for McKean-Vlasov equations. *Electron. Commun. Probab.*, 23:Paper No. 45, 11, 2018. 2, 3
- [7] D. Lacker. Hierarchies, entropy, and quantitative propagation of chaos for mean field diffusions. *arXiv: Probability*, 2021. <https://arxiv.org/abs/2105.02983>. 3
- [8] H. P. McKean, Jr. A class of Markov processes associated with nonlinear parabolic equations. *Proc. Nat. Acad. Sci. U.S.A.*, 56:1907–1911, 1966. 2
- [9] S. I. Resnick. *Extreme values, regular variation and point processes*. Springer Series in Operations Research and Financial Engineering. Springer, New York, 2008. Reprint of the 1987 original. 2, 4, 12, 13

- [10] A.-S. Sznitman. Topics in propagation of chaos. In *École d'Été de Probabilités de Saint-Flour XIX—1989*, volume 1464 of *Lecture Notes in Math.*, pages 165–251. Springer, Berlin, 1991. [2](#)