

Unitarity of the SoV transform for $SL(2, \mathbb{C})$ spin chains.

A.N. MANASHOV

Max-Planck-Institut für Physik, Werner-Heisenberg-Institut, 80805 München, Germany.

Abstract:

We prove the unitarity of the separation of variables transform for $SL(2, \mathbb{C})$ spin chains by a method based on the use of Gustafson integrals.

1 Introduction

Theory of quantum integrable models is an important part of modern theoretical physics. The solution of such models relies on the Quantum Inverse Scattering Method (QISM) which includes such techniques as the Algebraic Bethe Ansatz (ABA) [1] and Separation of Variables (SoV) [2,3]. The ABA allows one to effectively calculate energies and eigenstates of integrable models and to address more complicated problems such as calculating norms [4], scalar products [5] and correlation functions [6,7]. Models with infinite dimensional Hilbert spaces, the Toda chain [8] being the most famous example, are, however, beyond ABA's grasp. The solution of such models relies on the SoV method proposed by Sklyanin [2,3]. The method consists in constructing a map between the original Hilbert space, \mathbb{H}_{org} , in which the model is formulated, and an auxiliary Hilbert space, \mathbb{H}_{SoV} . This map is constructed in such a way that a multidimensional spectral problem associated with the original Hamiltonian is reduced to a one-dimensional problem on an auxiliary Hilbert space which usually takes the form of the Baxter $T - Q$ relation. Technically constructing the SoV representation is equivalent to finding the eigenfunctions of an element of the monodromy matrix associated with the model. For the Toda chain it was done by Kharchev and Lebedev [9,10]. Later, a regular method for obtaining eigenfunctions for models with an R-matrix of the rank one * was developed in [16], and at present the SoV representation is known for a number of models [17–21].

In order to be sure that the spectral problems in the original and auxiliary Hilbert spaces are equivalent it is necessary to show that the corresponding map, $\mathbb{H}_{\text{SoV}} \mapsto \mathbb{H}_{\text{org}}$, is unitary (or that the eigenfunctions form a complete set in \mathbb{H}_{org}). If $\dim \mathbb{H}_{\text{org}} < \infty$ the problem can be solved, at least in principle, by counting the dimensions of the Hilbert spaces. For the models with infinite dimensional Hilbert space, such as the Toda chain, the noncompact $\text{SL}(2, \mathbb{C})$ spin chain, etc., the task becomes more difficult. For the Toda chain unitarity was first established by using harmonic analysis of Lie groups techniques [22,23]. However this method is quite sophisticated and can hardly be generalized to more complicated cases. The rigorous proof of the unitarity of the SoV transform for the Toda chain based on the use of natural objects for the QISM was given by Kozłowski [24]. This technique was later applied to the modular XXZ magnet [25]. Later it was realized [26] that there exists a close relation between $\text{SL}(2, \mathbb{R})$ symmetric spin chains and the multidimensional Mellin-Barnes integrals studied by Gustafson [27,28] that allowed to greatly simplify the proof of the unitarity of the SoV transform for $\text{SL}(2, \mathbb{R})$ symmetric spin chains [29].

In the present paper we apply this technique to the analysis of the noncompact spin chains with the $\text{SL}(2, \mathbb{C})$ symmetry group. Such models appear in the studies of the Regge limit of scattering amplitudes in gauge theories, in QCD in particular [30–34], see also [35–37] for recent developments. The SoV representation for the $\text{SL}(2, \mathbb{C})$ spin chains was constructed in [16]. The generalization of Gustafson integrals relevant for the $\text{SL}(2, \mathbb{C})$ spin chains was obtained recently in [38]. Based on these results, we present below a proof of unitarity of the SoV transform for a generic $\text{SL}(2, \mathbb{C})$ spin chain.

The paper is organized as follows: in sect. 2 we recall elements of the QISM relevant for further analysis. The eigenfunctions of the elements of the monodromy matrix are constructed in sect. 3. In sect. 4 we calculate several scalar products of the eigenfunctions and discuss their properties. Sect. 5 contains the proof of unitarity of the SoV transform. Sect. 6 is reserved for a summary and several appendices contain a discussion of technical details.

*In recent years, significant progress has been made in constructing SoV representations for higher rank finite-dimensional models, see refs. [11–15]

2 $SL(2, \mathbb{C})$ spin chains

Spin chains are quantum mechanical systems whose dynamical variables are spin generators. We consider models with spin generators belonging to the unitary continuous principle series representation, $\mathbb{T}^{(s_k, \bar{s}_k)}$, of the unimodular group of complex two by two matrices. Namely, each site of the chain is equipped with two sets of generators, holomorphic (S^α) and anti-holomorphic ones (\bar{S}^α),

$$S_k^- = -\partial_{z_k}, \quad S_k^0 = z_k \partial_{z_k} + s_k, \quad S_k^+ = z_k^2 \partial_{z_k} + 2s_k z_k, \quad (2.1a)$$

$$\bar{S}_k^- = -\partial_{\bar{z}_k}, \quad \bar{S}_k^0 = \bar{z}_k \partial_{\bar{z}_k} + \bar{s}_k, \quad \bar{S}_k^+ = \bar{z}_k^2 \partial_{\bar{z}_k} + 2\bar{s}_k \bar{z}_k. \quad (2.1b)$$

The generators $S_k^\alpha (\bar{S}_k^\alpha)$ satisfy the standard $\mathfrak{sl}(2)$ commutation relations, while the generators at different sites and holomorphic and anti-holomorphic generators commute, $[S_k^\alpha, \bar{S}_k^{\alpha'}] = 0$. The parameters s_k, \bar{s}_k specifying the representation take the form [39]

$$s_k = \frac{1 + n_k}{2} + i\rho_k, \quad \bar{s}_k = \frac{1 - n_k}{2} + i\rho_k, \quad (2.2)$$

where n_k is an integer or half-integer number and ρ_k is real, so that

$$s_k + \bar{s}_k^* = 1 \quad \text{and} \quad s_k - \bar{s}_k = n_k \in \mathbb{Z}/2. \quad (2.3)$$

The later condition comes from the requirement for the finite group transformations to be well defined while the former one guarantees the unitary character of transformations and anti-hermiticity of the generators, $(S_k^\alpha)^\dagger = -\bar{S}_k^\alpha$.

The Hilbert space of the model is given by the direct product of the Hilbert spaces at each node. For a chain of length N , $\mathbb{H}_N = \bigotimes_{k=1}^N \mathcal{H}_k$, where $\mathcal{H}_k = L_2(\mathbb{C})$.

In the QISM [1, 3, 40, 41] the dynamics of the model is determined by a family of mutually commuting operators. Namely, one defines the so-called L -operators,

$$L_k(u) = u + i \begin{pmatrix} S_k^0 & S_k^- \\ S_k^+ & -S_k^0 \end{pmatrix}, \quad \bar{L}_k(\bar{u}) = \bar{u} + i \begin{pmatrix} \bar{S}_k^0 & \bar{S}_k^- \\ \bar{S}_k^+ & -\bar{S}_k^0 \end{pmatrix}, \quad (2.4)$$

which are the basic building blocks in the QISM. The complex variables u, \bar{u} are called spectral parameters. The next important object – a monodromy matrix – is given by the product of L operators

$$\begin{aligned} T_N(u) &= L_1(u + \xi_1) L_2(u + \xi_2) \dots L_N(u + \xi_N), \\ \bar{T}_N(\bar{u}) &= \bar{L}_1(\bar{u} + \bar{\xi}_1) \bar{L}_2(\bar{u} + \bar{\xi}_2) \dots \bar{L}_N(\bar{u} + \bar{\xi}_N), \end{aligned} \quad (2.5)$$

where $\xi_k, \bar{\xi}_k$ are the so-called impurity parameters [†]. The entries of the monodromy matrix,

$$T_N(u) = \begin{pmatrix} A_N(u) & B_N(u) \\ C_N(u) & D_N(u) \end{pmatrix}, \quad (2.6)$$

are polynomials in u with the operator valued coefficients, e.g.

$$A_N(u) = u^N + u^{N-1} (iS^0 + \Xi) + \sum_{k=2}^N u^{N-k} a_k, \quad B_N(u) = u^{N-1} iS^- + \sum_{k=2}^N u^{N-k} b_k. \quad (2.7)$$

[†]As it can already be noticed any formula in the holomorphic sector has its exact copy in the anti-holomorphic one. Therefore, from now on we write explicitly only holomorphic formulae tacitly implying its anti-holomorphic counterparts.

where $\Xi = \sum_{k=1}^N \xi_k$ and S^0, S^- are the total generators:

$$S^\alpha = S_1^\alpha + \dots + S_N^\alpha. \quad (2.8)$$

The entries of the monodromy matrix form commuting operator families [1, 42]

$$[A_N(u), A_N(v)] = [B_N(u), B_N(v)] = [C_N(u), C_N(v)] = [D_N(u), D_N(v)] = 0. \quad (2.9)$$

In particular, each entry commutes with the corresponding total generator, S^α ,

$$[S^0, A_N(u)] = [S^0, D_N(u)] = 0 \quad \text{and} \quad [S^-, B_N(u)] = [S^+, C_N(u)] = 0. \quad (2.10)$$

The same equations hold for the anti-holomorphic operators $\bar{A}_N, \bar{B}_N, \bar{C}_N, \bar{D}_N$ and, of course, the holomorphic and anti-holomorphic operators commute. Moreover it can be checked that if the impurity parameters satisfy the constraint $\bar{\xi}_k = \xi_k^*$ for all k , the following relations between holomorphic and anti-holomorphic operators hold

$$(A_N(u))^\dagger = \bar{A}_N(u^*), \quad (B_N(u))^\dagger = \bar{B}_N(u^*), \quad (2.11)$$

etc. This ensures that the operators a_k and \bar{a}_k in the expansion of $A_N(u)$, Eq. (2.7), and $\bar{A}_N(u)$, are adjoint to each other $a_k^\dagger = \bar{a}_k$ ($b_k^\dagger = \bar{b}_k$ etc.)

The commutativity of the operators $A_N(u), B_N(u), C_N(u), D_N(u)$ implies that the following families of self-adjoint operators

$$\begin{aligned} \mathfrak{A}_N &= \{iS^0, i\bar{S}^0, a_k + \bar{a}_k, i(a_k - \bar{a}_k), k = 2, \dots, N\}, \\ \mathfrak{B}_N &= \{iS^-, i\bar{S}^-, b_k + \bar{b}_k, i(b_k - \bar{b}_k), k = 2, \dots, N\}, \end{aligned}$$

(and similarly for others) are commutative and can be diagonalized simultaneously[‡]. The corresponding eigenfunctions provide a convenient basis – Sklyanin’s representation of Separated Variables (SoV) – for the analysis of spin chain models [3].

The operators B_N and C_N , (A_N and D_N) are related to each other by the inversion transformation, see ref. [21] for detail, so it is sufficient to construct eigenfunctions for the operators B_N and A_N . The eigenfunctions of B_N for the homogeneous chain were constructed in ref. [16] and later on for the operator A_N , [21]. Extending this approach to the inhomogeneous case is rather straightforward.

3 Eigenfunctions

In this section we present explicit expressions for the eigenfunctions of the operators B_N and A_N for a generic inhomogeneous spin chain with impurities. We start with the operator B_N where the construction follows the lines of ref. [16] with minimal modifications.

3.1 B_N operator

Let Λ_n be an integral (layer) operator which maps functions of $n - 1$ variables into functions of n variables and depends on the spectral parameters x, \bar{x} and the complex vectors $\gamma, \bar{\gamma}$ of dimension $2n - 2$

$$[\Lambda_n(x|\gamma)f](z_1, \dots, z_n) = \int \dots \int \Lambda_n(x|\gamma)(z_1, \dots, z_n|w_1, \dots, w_{n-1})f(w_1, \dots, w_{n-1}) \prod_{k=1}^{n-1} d^2 w_k. \quad (3.1)$$

[‡]The impurity parameters must also satisfy the condition $i(\xi_k - \bar{\xi}_k) = r_k$, where r_k are (half)integers.

The kernel is given by the following expression

$$\Lambda_n(x|\gamma)(z_1, \dots, z_n | w_1, \dots, w_{n-1}) = \prod_{k=1}^{n-1} D_{\gamma_{2k-1}-ix}(z_k - w_k) D_{\gamma_{2k}+ix}(z_{k+1} - w_k), \quad (3.2)$$

where the function $D_\alpha(z)$ (propagator) is defined as follows

$$D_\alpha(z) \equiv D_{\alpha, \bar{\alpha}}(z, \bar{z}) = z^{-\alpha} \bar{z}^{-\bar{\alpha}}. \quad (3.3)$$

We will assume that the indices $\alpha, \bar{\alpha}$ satisfy the condition $[\alpha] \equiv \alpha - \bar{\alpha} \in \mathbb{Z}$ so that the propagator is a single-valued function on the complex plane. It implies that the parameters γ_k and x have the form

$$\gamma_k = \frac{1}{2} + \frac{r_k}{2} + i\sigma_k, \quad \bar{\gamma}_k = \frac{1}{2} - \frac{r_k}{2} + i\sigma_k, \quad x = \frac{im}{2} + \nu, \quad \bar{x} = -\frac{im}{2} + \nu. \quad (3.4)$$

The numbers $\{m, r_1, \dots, r_{2N-2}\}$ are either integer or half-integer and depending on this we call the corresponding variables integer (half-integer). The continuous parameters σ_k and ν are subject to the constraints

$$\text{Im}(\sigma_{2k+1} - \nu) > -1/2 \quad \text{and} \quad \text{Im}(\sigma_{2k} + \nu) > -1/2 \quad (3.5)$$

which guarantee the convergence of the integral (3.1) for a smooth function f with finite support. In the case we are most interested in, $\gamma_k + \bar{\gamma}_k = 1$, the parameters $\sigma_k \in \mathbb{R}$, and the variable ν lies in the strip $-1/2 < \text{Im}\nu < 1/2$.

The operators Λ_n possess two important properties:

- (i) Let ρ be a map which takes M -dimensional vectors

$$\gamma = (\gamma_1, \dots, \gamma_M), \quad \bar{\gamma} = (\bar{\gamma}_1, \dots, \bar{\gamma}_M) \quad (3.6)$$

to vectors of dimension $M - 2$ as follows

$$\rho\gamma = (\gamma'_2, \gamma'_3, \dots, \gamma'_{M-1}) \quad \rho\bar{\gamma} = (\bar{\gamma}'_2, \bar{\gamma}'_3, \dots, \bar{\gamma}'_{M-1}), \quad (3.7)$$

where $a' \equiv 1 - a$. It can be shown that the operators Λ_n and Λ_{n-1} obey the following exchange relation

$$\Lambda_n(u|\gamma) \Lambda_{n-1}(v|\rho\gamma) = \omega_n(\gamma, u, v) \Lambda_n(v|\gamma) \Lambda_{n-1}(u|\rho\gamma). \quad (3.8)$$

Here $\gamma(\bar{\gamma})$ is $2n - 2$ dimensional vector and the factor ω_n is given by the following expression

$$\omega_n(\gamma, u, v) = \prod_{m=1}^{n-1} \Gamma \left[\frac{\gamma_{2m-1} - iv, \bar{\gamma}_{2m} + i\bar{v}}{\gamma_{2m-1} - iu, \bar{\gamma}_{2m} + i\bar{u}} \right] = \prod_{m=1}^{n-1} \Gamma \left[\frac{\bar{\gamma}_{2m-1} - i\bar{v}, \gamma_{2m} + iv}{\bar{\gamma}_{2m-1} - i\bar{u}, \gamma_{2m} + iu} \right], \quad (3.9)$$

where

$$\Gamma \left[\frac{a_1, a_2, \dots, a_n}{b_1, b_2, \dots, b_m} \right] \equiv \frac{\prod_{k=1}^n \Gamma[a_k]}{\prod_{k=1}^m \Gamma[b_k]} \quad (3.10)$$

and Γ is the Gamma function of the complex field \mathbb{C} [43]

$$\Gamma[u] \equiv \Gamma[u, \bar{u}] = \Gamma(u)/\Gamma(1 - \bar{u}). \quad (3.11)$$

The relation (3.8) is a direct consequence of the exchange relation for the propagators, see (A.4). Its proof is exactly the same as for the homogeneous spin chain. For more details see refs. [16, 21].

(ii) Let us choose the vector γ as follows

$$\begin{aligned}\gamma &= (s_1 - i\xi_1, s_2 + i\xi_2, s_2 - i\xi_2, \dots, s_{N-1} + i\xi_{N-1}, s_{N-1} - i\xi_{N-1}, s_N + i\xi_N), \\ \bar{\gamma} &= (\bar{s}_1 - i\bar{\xi}_1, \bar{s}_2 + i\bar{\xi}_2, \bar{s}_2 - i\bar{\xi}_2, \dots, \bar{s}_{N-1} + i\bar{\xi}_{N-1}, \bar{s}_{N-1} - i\bar{\xi}_{N-1}, \bar{s}_N + i\bar{\xi}_N),\end{aligned}\quad (3.12)$$

where s_k and ξ_k are the spins and impurity parameters of the spin chain, respectively. For such a choice of the vector γ the operator $B_N(x)$ annihilates $\Lambda_N(x|\gamma)$ [16, 44]

$$B_N(x)\Lambda_N(x|\gamma) = 0. \quad (3.13)$$

Let us define a function

$$\Psi_{p,x}^{(N)}(z) \equiv \Psi_{p,x_1,\dots,x_{N-1}}^{(N)}(z_1, \dots, z_N) = \pi^{-N^2/2} |p|^{N-1} \int d^2z U_{x_1,\dots,x_{N-1}}(z_1, \dots, z_N|z) e^{i(pz + \bar{p}\bar{z})}, \quad (3.14)$$

where the kernel $U_{x_1,\dots,x_{N-1}}$ is given by the product of the layer operators,

$$U_{x_1,\dots,x_{N-1}} = \varpi(x|\gamma)\Lambda_N(x_1|\gamma)\Lambda_{N-1}(x_2|\rho\gamma)\Lambda_{N-2}(x_3|\rho^2\gamma)\dots\Lambda_2(x_{N-1}|\rho^{N-2}\gamma), \quad (3.15)$$

and γ is given by Eq. (3.12). Equation (3.8) guarantees that $U_{x_1,\dots,x_{N-1}} \sim U_{x_{i_1},\dots,x_{i_{N-1}}}$ for any permutation of x_1, \dots, x_{N-1} . The kernel U_x becomes totally symmetric for the following choice of the prefactor $\varpi(x|\gamma)$:

$$\varpi(x|\gamma) = \varpi(x_1, \dots, x_{N-1}|\gamma) = \prod_{m=1}^{N-1} \prod_{k=1}^m \varpi(x_k|\rho^{m-1}\gamma), \quad (3.16)$$

where

$$\varpi(x|\gamma) = \varpi(x|\gamma_1, \dots, \gamma_{2n}) = \prod_{m=1}^n \Gamma[\gamma_{2m-1} - ix, \bar{\gamma}_{2m} + i\bar{x}]. \quad (3.17)$$

Thus the function $\Psi_{p,x_1,\dots,x_{N-1}}^{(N)}$ is a symmetric function of the variables x_1, \dots, x_{N-1} . Together with Eq. (3.13) it implies that

$$B_N(x_k)\Psi_{p,x_1,\dots,x_{N-1}}^{(N)} = 0 \quad \text{for} \quad k = 1, \dots, N-1. \quad (3.18)$$

Invariance of the kernel $U_{x_1,\dots,x_{N-1}}(z_1, \dots, z_N|z)$ is under shifts

$$U_{x_1,\dots,x_{N-1}}(z_1 + w, \dots, z_N + w|z + w) = U_{x_1,\dots,x_{N-1}}(z_1, \dots, z_N|z) \quad (3.19)$$

results in

$$iS^- \Psi_{p,x_1,\dots,x_{N-1}}^{(N)} = p \Psi_{p,x_1,\dots,x_{N-1}}^{(N)}, \quad i\bar{S}^- \Psi_{p,x_1,\dots,x_{N-1}}^{(N)} = \bar{p} \Psi_{p,x_1,\dots,x_{N-1}}^{(N)}. \quad (3.20)$$

It follows then from Eqs. (2.7), (3.13) and (3.20) that [§]

$$B_N(u)\Psi_{p,x}^{(N)}(z) = p \prod_{k=1}^{N-1} (u - x_k)\Psi_{p,x}^{(N)}(z), \quad \bar{B}_N(\bar{u})\Psi_{p,x}^{(N)}(z) = \bar{p} \prod_{k=1}^{N-1} (\bar{u} - \bar{x}_k)\Psi_{p,x}^{(N)}(z). \quad (3.21)$$

[§]We recall that the variables $x_k, \bar{x}_k, k = 1, \dots, N-1$ take the form $x_k = in_k/2 + \nu_k, \bar{x}_k = -in_k/2 + \nu_k$, where, depending on the spin and impurities parameters, all n_k are either integer or half-integer numbers.

For $N = 1$ the functions $\Psi_p^{(1)}(z, \bar{z}) = \pi^{-1/2} e^{i(pz + \bar{p}\bar{z})}$ form the complete orthonormal system in $\mathbb{H}_1 = L_2(\mathbb{C})$. The aim of this paper is to extend this statement to $N > 1$. Namely, we will show in sect. 5 that if the spins and impurities parameters of the spin chain obey the ‘‘unitarity’’ condition,

$$\gamma_k + \bar{\gamma}_k^* = 1, \quad (3.22)$$

for all k (γ_k has the form (3.4) with $\sigma_k \in \mathbb{R}$) then the set of functions $\{\Psi_{p,x}^{(N)}, x_k = \bar{x}_k^* (\nu_k \in \mathbb{R}), k = 1, \dots, N-1\}$ is complete in $\mathbb{H}_N = (\otimes L_2(\mathbb{C}))^N$.

Note that the functions $\Psi_{p,x}^{(N)}$ are well defined for the complex parameters ν_k in the vicinity of the real line. For further analysis, it will be useful to consider regularized functions, $\Psi_{p,x}^{(N),\epsilon}$, by relaxing the last of the conditions (3.22) to $\gamma_{2N-2} + \bar{\gamma}_{2N-2}^* = 1 + 2\epsilon$. This can be achieved by shifting the impurity parameter $\xi_N \rightarrow \xi_N - i\epsilon$ [¶], i.e.

$$\Psi_{p,x}^{(N),\epsilon}(z) \stackrel{\text{def}}{=} \Psi_{p,x}^{(N)}(z) \Big|_{\xi_N \rightarrow \xi_N - i\epsilon}. \quad (3.23)$$

3.2 A_N operator

Construction of the eigenfunctions of the operator A_N follows the scheme described in the previous subsection. We define a layer operator Λ'_n which maps functions of $n-1$ variables into functions of n variables

$$[\Lambda'_n(x|\gamma)f](z_1, \dots, z_n) = \int \cdots \int \Lambda'_n(x|\gamma)(z_1, \dots, z_n | w_1, \dots, w_{n-1}) f(w_1, \dots, w_{n-1}) \prod_{k=1}^{n-1} d^2 w_k, \quad (3.24)$$

where the kernel is given by the following expression

$$\Lambda'_n(x|\gamma)(z_1, \dots, z_n | w_1, \dots, w_{n-1}) = D_{\gamma_{2N-1}-ix}(z_N) \prod_{k=1}^{n-1} D_{\gamma_{2k-1}-ix}(z_k - w_k) D_{\gamma_{2k}+ix}(z_{k+1} - w_k). \quad (3.25)$$

The layer operator Λ'_n depends on the spectral parameters $x(\bar{x})$ and the vector $\gamma(\bar{\gamma})$ of dimension $2n-1$ which have the form Eq. (3.4).

These operators satisfy the exchange relation

$$\Lambda'_n(u|\gamma)\Lambda'_{n-1}(v|\rho\gamma) = \omega_n(\gamma, u, v)\Lambda'_n(v|\gamma)\Lambda'_{n-1}(u|\rho\gamma), \quad (3.26)$$

and the factor ω_n is defined in Eq. (3.9).

Let $\Phi_x^{(N)}(z)$ be the following function

$$\begin{aligned} \Phi_x^{(N)}(z) &\equiv \Phi_{x_1, \dots, x_N}^{(N)}(z_1, \dots, z_N) \\ &= \pi^{-N^2/2} \varpi(x|\gamma) [\Lambda'_N(x_1|\gamma)\Lambda'_{N-1}(x_2|\rho\gamma) \cdots \Lambda'_1(x_N|\rho^{N-1}\gamma)](z_1, \dots, z_N), \end{aligned} \quad (3.27)$$

where γ is $2N-1$ dimensional vector and the prefactor ϖ is given by Eq. (3.16). For such a choice of ϖ the function $\Phi_x^{(N)}$ is a symmetric function of the variables x_1, \dots, x_N .

It can be shown that the operator $A_N(x)$ annihilates the layer operator $\Lambda'_N(x|\gamma)$,

$$A_N(x)\Lambda'_N(x|\gamma) = 0, \quad (3.28)$$

[¶]Of course, one also can regularize the function by shifting the parameter γ_1 instead of γ_{2N-2} , $\gamma_1 + \bar{\gamma}_1^* = 1 + 2\epsilon$.




Figure 1: The diagrammatic representation for the function Ψ (left) and Φ (right) for $N = 3$. The arrow from z to w with an index α stands for the propagator $D_\alpha(z - w)$, Eq. (3.3).

for the following choice of the vector γ

$$\begin{aligned}\gamma &= (s_1 - i\xi_1, s_2 + i\xi_2, s_2 - i\xi_2, \dots, s_N + i\xi_N, s_N - i\xi_N), \\ \bar{\gamma} &= (\bar{s}_1 - i\bar{\xi}_1, \bar{s}_2 + i\bar{\xi}_2, \bar{s}_2 - i\bar{\xi}_2, \dots, \bar{s}_N + i\bar{\xi}_N, \bar{s}_N - i\bar{\xi}_N).\end{aligned}\quad (3.29)$$

Taking into account polynomiality of $A_N(u)$, see Eq. (2.7), one obtains

$$A_N(u)\Phi_x^{(N)}(z) = \prod_{k=1}^N (u - x_k)\Phi_x^{(N)}(z), \quad \bar{A}_N(\bar{u})\Phi_x^{(N)}(z) = \prod_{k=1}^N (\bar{u} - \bar{x}_k)\Phi_x^{(N)}(z). \quad (3.30)$$

Again, the variables x_k, \bar{x}_k are integers (half-integers) for all k . We will show that these functions, $\{\Phi_x^{(N)}(z), x_k = \bar{x}_k^*, k = 1, \dots, N\}$, form a complete set in the Hilbert space \mathbb{H}_N .

4 Scalar products, momentum representation, etc.

The functions constructed in the previous section are given by multi-dimensional integrals. In this section we show that these integrals converge for the parameters ν_k in the vicinity of real axis. To this end it will be quite helpful, as was advocated in ref. [16], to visualize the integrals as Feynman diagrams. The examples for $N = 3$ are shown in Fig. 1. It will be convenient to convert diagrams (functions) to momentum space

$$\Psi(z_1, \dots, z_N) = \pi^{-N} \int \dots \int \tilde{\Psi}(p_1, \dots, p_N) e^{i \sum_{k=1}^N (p_k z_k + \bar{p}_k \bar{z}_k)} d^2 p_1 \dots d^2 p_N. \quad (4.1)$$

In momentum space the function $\Psi_{p,x}^{(N),\epsilon}$, Eq. (3.23), takes the form

$$\tilde{\Psi}_{p,x}^{(N),\epsilon}(p_1, \dots, p_N) = \delta^{(2)}\left(p - \sum_{k=1}^N p_k\right) \Psi_x^{(N),\epsilon}(p_1, \dots, p_N). \quad (4.2)$$

Let us remark here that the "ε" regularization is reduced to a multiplication by the factor $(p_N \bar{p}_N)^\epsilon$

$$\Psi_x^{(N),\epsilon}(p_1, \dots, p_N) = (p_N \bar{p}_N)^\epsilon \Psi_x^{(N)}(p_1, \dots, p_N). \quad (4.3)$$

The function $\Psi_x^{(N),\epsilon}$ can be read from the Feynman diagram in Fig.1 as follows

$$\Psi_x^{(N),\epsilon}(p_1, \dots, p_N) = \int \cdots \int \mathcal{J}_x^\epsilon(\{p_k\}, \{\ell_{ij}\}) \prod_{1 \leq j \leq i \leq N-2} d^2 \ell_{ij}, \quad (4.4)$$

with the integrand $\mathcal{J}_x^\epsilon(\{p_k\}, \{\ell_{ij}\})$ given by the product of the propagators, $D_\alpha(k)$. Up to a momentum independent factor

$$\mathcal{J}_x^\epsilon(\{p_k\}, \{\ell_{ij}\}) \simeq \prod_{k=1}^{N-1} \prod_{j=1}^k D_{\alpha_{kj}}(\ell_{k,j} - \ell_{k-1,j-1}) D_{\beta_{kj}}(\ell_{k-1,j} - \ell_{k,j}), \quad (4.5)$$

where $\ell_{k0} \equiv 0$, $\ell_{k-1,k} \equiv p$ and $\ell_{N-1,j} = (p_1 + \dots + p_j)$. The indices α_{kj} , β_{kj} take the following values

$$\alpha_{kj} = \gamma_{2j-1}^{(N-k)} + ix_{N-k}, \quad \beta_{kj} = \gamma_{2j}^{(N-k)} - ix_{N-k}, \quad (4.6)$$

where we introduced the notations:

$$a^{(1)} = a' = 1 - a \quad \text{and} \quad a^{(k+1)} = 1 - a^{(k)}. \quad (4.7)$$

In many cases, Feynman diagrams can be evaluated diagrammatically. In particular, the computation of diagrams for the scalar product of Ψ (Φ) functions is based on the successive application of the exchange relation (A.4) to the diagram.

Let us consider the scalar product of two functions $\Psi_{p,x}^{(N),\epsilon}$ and $\Psi_{q,y}^{(N),\epsilon'}$

$$\left(\Psi_{q,y}^{(N),\epsilon'}, \Psi_{p,x}^{(N),\epsilon} \right) = \pi \delta^2(p - q) (p\bar{p})^{\epsilon+\epsilon'} I^{\epsilon,\epsilon'}(x, y), \quad (4.8)$$

where

$$I^{\epsilon,\epsilon'}(x, y) = \frac{1}{\pi} (p\bar{p})^{-\epsilon-\epsilon'} \int \cdots \int \delta^{(2)}\left(p - \sum_k p_k\right) \Psi_x^{(N),\epsilon}(\vec{p}) \left(\Psi_y^{(N),\epsilon'}(\vec{p}) \right)^\dagger \prod_{j=1}^N d^2 p_j. \quad (4.9)$$

The function $I_p^{\epsilon,\epsilon'}(x, y)$ is given by the Feynman diagram shown in Fig. 2 (left panel), which is a multidimensional integral

$$I_p^{\epsilon,\epsilon'}(x, y) = \int \cdots \int \mathcal{I}_{x,y}^{\epsilon\epsilon'}(p, \{\ell_{pr}\}|\gamma) \prod_{p,r=1}^{N-1} d^2 \ell_{pr} \quad (4.10)$$

with the integrand given by the product of the propagators. The diagram can be evaluated in a closed form by successively applying the exchange relation (A.4), that is equivalent to calculating the loop integrals in a certain order. The answer takes the form

$$\begin{aligned} I^{\epsilon,\epsilon'}(x, y) &= C_N(\gamma) \Gamma \left[\frac{\epsilon + \epsilon' + iX - i\bar{Y}^*}{\epsilon + \epsilon'} \right] \frac{\prod_{k,j=1}^{N-1} \Gamma[i(y_k^* - \bar{x}_j)]}{\prod_{k=1}^{N-1} \bar{\phi}_N(\bar{x}_k) (\phi_N(y_k))^*} \\ &= C_N(\gamma) \Gamma \left[\frac{\epsilon + \epsilon' + i\bar{X} - iY^*}{\epsilon + \epsilon'} \right] \frac{\prod_{k,j=1}^{N-1} \Gamma[i(\bar{y}_k^* - x_j)]}{\prod_{k=1}^{N-1} \phi_N(x_k) (\bar{\phi}_N(\bar{y}_k))^*}, \end{aligned} \quad (4.11)$$

where $X = \sum_{k=1}^{N-1} x_k$, $Y = \sum_{k=1}^{N-1} Y_k$ and

$$\begin{aligned}\phi_N(x) &= \mathbf{\Gamma} \left[\gamma_{2N-3} - ix, \gamma_{2N-4}^{(1)} - ix, \gamma_{2N-5} - ix, \dots, \gamma_N^{(N-3)} - ix \right], \\ \bar{\phi}_N(\bar{x}) &= \mathbf{\Gamma} \left[\bar{\gamma}_{2N-3} - i\bar{x}, \bar{\gamma}_{2N-4}^{(1)} - i\bar{x}, \bar{\gamma}_{2N-5} - i\bar{x}, \dots, \bar{\gamma}_N^{(N-3)} - i\bar{x} \right].\end{aligned}\quad (4.12)$$

For the sign factor $C_N(\gamma)$ we get :

$$C_N(\gamma_1, \gamma_2, \dots, \gamma_{2N-2}) = \begin{cases} 1 & \text{odd } N \\ (-1)^{\sum_{k=1}^{N-3} [\gamma_{2N-2-k}^{(k-1)} - \gamma_N^{(N-3)}]} & \text{even } N. \end{cases}\quad (4.13)$$

Here $[a] \equiv a - \bar{a}$. Details of the calculation can be found in [Appendix B](#).

Let us show now that integrations in (4.10) can be done in an arbitrary order. The integrand in (4.10), $\mathcal{I}_{x,y}^{\epsilon\epsilon'}(p, \{\ell_{pr}\}|\gamma)$, is given by the product of the propagators $D_\alpha(k)$, with each index being of the form $\alpha = \frac{1}{2} + \frac{n}{2} + i\sigma$, momentum k being a linear combination of loop momenta, ℓ_{ij} , and the external momentum p . Since

$$|D_\alpha(k)| = |k^{-\alpha} \bar{k}^{-\bar{\alpha}}| = |k|^{-1+2\text{Im}\sigma} = D_{1/2-\text{Im}\sigma}(k)\quad (4.14)$$

then for the parameters γ satisfying the unitarity condition (3.22), and x_k, y_k having the form

$$x_k = in_k/2 + \nu_k, \quad y_k = im_k/2 + \mu_k,\quad (4.15)$$

one obtains for the modulus of the integrand

$$\left| \mathcal{I}_{x,y}^{\epsilon\epsilon'}(p, \{\ell_{pr}\}|\gamma) \right| = \mathcal{I}_{\underline{x},\underline{y}}^{\epsilon\epsilon'}(p, \{\ell_{pr}\}|\underline{\gamma}) > 0,\quad (4.16)$$

where the underlined variables are: $\underline{\gamma} = (1/2, \dots, 1/2)$,

$$(\underline{x})_k = \text{Im}(\nu_k) = \epsilon_k, \quad (\underline{y})_k = \text{Im}(\mu_k) = \epsilon'_k.$$

Thus the integral of $|\mathcal{I}_{x,y}^{\epsilon\epsilon'}(p, \{\ell_{pr}\}|\gamma)|$ is a particular case of the integral (4.10) which was calculated by performing loop integrations in a certain order. Since all integrals converge under the conditions

$$\epsilon_{kj} \equiv \epsilon_k + \epsilon'_j > 0 \quad \text{for} \quad k, j = 1, \dots, N-1 \quad \text{and} \quad \epsilon + \epsilon' > \sum_{k=1}^{N-1} (\epsilon_k + \epsilon'_k),\quad (4.17)$$

by Fubini theorem, the integral (4.10) exists and the integrations can be done in an arbitrary order.

The following statements can immediately be deduced from this result:

- For any bounded function $\varphi(p, x)$ with a finite support the function

$$\Psi_\varphi^\epsilon = \int \dots \int \varphi(p, x) \Psi_{p,x^\epsilon}^{(N),\epsilon} d^2p \mathcal{D}x_1 \dots \mathcal{D}x_{N-1},\quad (4.18)$$

where $x^\epsilon = (x_1 + i\epsilon_1, \dots, x_{N-1} + i\epsilon_{N-1})$, $x_k = in_k/2 + \nu_k$, $\epsilon_k > 0$, $\epsilon > \sum_{k=1}^{N-1} \epsilon_k$ and

$$\int \mathcal{D}x_k \equiv \sum_{n_k=-\infty}^{\infty} \int_{-\infty}^{\infty} d\nu_k,$$

belongs to the Hilbert space \mathbb{H}_N , $\|\Psi_\varphi^\epsilon\|^2 < \infty$, for sufficiently small ϵ .

- It follows from the finiteness of the integral $I_p^{\epsilon, \epsilon'}(x, y)$, Eq. (4.9), that the function $\Psi_x^{(N), \epsilon}(\vec{p})$, Eq. (4.4), exists almost for all \vec{p} for the separated variables x_k close to the real axis:

$$\text{Im } \nu_k = \frac{1}{2} \text{Im}(x_k + \bar{x}_k) \sim 0 \quad \text{for all } k$$

and $\Psi_x^{(N), \epsilon}(\vec{p})$ is a continuous function of ν_k in this region. Indeed, let us fix $m < N$ and put $u_m = \text{Re} \nu_m$ and $v_m = \text{Im} \nu_m$, $|v_m| < \delta$. One gets the following estimate for the integrand (4.10)

$$|\mathcal{J}_x^\epsilon(\{p_k\}, \{\ell_{ij}\})| < \left| \mathcal{J}_{x_+}^\epsilon(\{p_k\}, \{\ell_{ij}\}) \right| + \left| \mathcal{J}_{x_-}^\epsilon(\{p_k\}, \{\ell_{ij}\}) \right|, \quad (4.19)$$

where x_\pm are defined as follows: for $k \neq m$ $(x_\pm)_k = x_k$ and for $k = m$ $(x_\pm)_m = u_m \pm i\delta$. The integrals of the functions on the rhs of (4.19) are finite for sufficiently small δ . It follows then from the Lebesgue theorem that the function $\Psi_x^{(N), \epsilon}(\vec{p})$ is continuous in the variable ν_m [‡].

The scalar product of the functions $\Psi_{p,y}^{(N)}$ and $\Phi_x^{(N)}$ constructed in sect. 3.2 can be calculated in a similar way. Note that there is no need to introduce “ ϵ ” regulator here. The corresponding integral is absolutely convergent when $\text{Im}(\nu_k + \mu_j) > 0$ for all k, j , (x_k, y_j given by (4.15)). The scalar product takes the form

$$\left(\Psi_{p,y}^{(N)} | \Phi_x^{(N)} \right) = C_N^{AB}(\gamma) |p|^{N-1} (-ip)^{-A_N - iX} (i\bar{p})^{-\bar{A}_N - i\bar{X}} \frac{\prod_{k=1}^N \prod_{j=1}^{N-1} \Gamma[i(\bar{y}_j^* - x_k)]}{\left(\prod_{j=1}^N \vartheta_N(x_j) \right) \left(\prod_{j=1}^{N-1} \bar{\vartheta}_N(\bar{y}_j) \right)^\dagger}, \quad (4.20)$$

where

$$\vartheta_N(x) = \prod_{k=1}^N \Gamma \left[\gamma_{2N-k}^{(k-1)} - ix_j \right], \quad \bar{\vartheta}_N(\bar{x}) = \prod_{k=1}^N \Gamma \left[\bar{\gamma}_{2N-k}^{(k-1)} - i\bar{x}_j \right], \quad (4.21)$$

$A_N = \sum_{k=N}^{2N-1} \gamma_k^{(k)}$, $X = \sum_{k=1}^N x_k$ and

$$C_N^{AB}(\gamma_1, \dots, \gamma_{2N-1}) = \begin{cases} 1 & \text{odd } N \\ (-1)^{\sum_{k=1}^N [\gamma_{2N-k}^{(k-1)} - \gamma_N^{(N-1)}]} & \text{even } N. \end{cases} \quad (4.22)$$

Similar to the previous case one can argue that $\Phi_x^{(N)}$ is a continuous function of ν_k in the vicinity of the real axis.

Finally, the scalar product of the functions $\Psi_{p,x}^{(N+1)}(z_1, \dots, z_{N+1})$ and $\Psi_{q_1,y}^{(N)}(z_1, \dots, z_N) \otimes \Psi_{q_2}^{(1)}(z_{N+1})$ which we need in the proof of Theorem 1, takes the form

$$\begin{aligned} \left(\Psi_{q_1,y}^N \otimes \Psi_{q_2}^{(1)}, \Psi_{p,x}^{(N+1)} \right) &= C_{NN+1}(\gamma) \pi \delta^{(2)}(p - q_1 - q_2) |p|^N |q_1|^{N-1} \times \\ &\quad (ip)^{-\bar{A}_{N+1}^*} (-i\bar{p})^{-A_{N+1}^*} (iq_2)^{-\gamma'_{2N}} (-i\bar{q}_2)^{-\bar{\gamma}'_{2N}} (-iq_1)^{-A_N} (i\bar{q}_1)^{-\bar{A}_N} \times \\ &\quad \left(1 + \frac{q_1}{q_2} \right)^{i\bar{Y}^*} \left(1 + \frac{\bar{q}_1}{\bar{q}_2} \right)^{iY^*} \left(-\frac{q_2}{q_1} \right)^{iX} \left(-\frac{\bar{q}_2}{\bar{q}_1} \right)^{i\bar{X}} \times \\ &\quad \frac{\prod_{k=1}^{N-1} \prod_{j=1}^N \Gamma[i(\bar{y}_k^* - x_j)]}{\left(\prod_{j=1}^N \prod_{k=1}^{N-1} \Gamma \left[\gamma_{2N-k}^{(k-1)} - ix_j \right] \right) \left(\prod_{k=1}^N \prod_{j=1}^{N-1} \Gamma \left[\bar{\gamma}_{2N-k}^{(k-1)} - i\bar{y}_j \right] \right)^\dagger}, \end{aligned} \quad (4.23)$$

[‡]Since the integrand is analytic function of ν_k $\Psi_x^{(N), \epsilon}(\vec{p})$ is an analytic function of ν_k in the vicinity of the real axis.

where

$$A_N = \sum_{m=N}^{2N-1} \gamma_m^{(m)}, \quad A_{N+1} = A_N - \gamma_N^{(N)} = \sum_{m=N+1}^{2N-1} \gamma_m^{(m)} \quad (4.24)$$

and

$$C_{NN+1}(\gamma_1, \dots, \gamma_{2N}) = \begin{cases} 1 & \text{for odd } N \\ (-1)^{\sum_{k=1}^{N-1} [\gamma_{2N-k}^{(k-1)} - \gamma_N^{(N-1)}]} & \text{for even } N. \end{cases} \quad (4.25)$$

The calculation is almost the same as in the previous cases so we omit the details.

5 SoV representation

In the previous section we constructed the functions $\Psi_{p,x}^{(N)}$ and $\Phi_x^{(N)}$ associated with the entries B_N and A_N of the monodromy matrix (2.5). For a given vector $\Psi \in \mathbb{H}_N$ we define two functions by projecting it on $\Psi_{p,x}^{(N)}$ and $\Phi_x^{(N)}$:

$$\varphi(p, x_1, \dots, x_{N-1}) = \left(\Psi_{p,x}^{(N)}, \Psi \right), \quad \chi(x_1, \dots, x_N) = \left(\Phi_x^{(N)}, \Psi \right). \quad (5.1)$$

These functions are symmetric functions of the variables x . It was shown by Sklyanin [3] that the transformation $\Psi \mapsto \varphi$ ($\Psi \mapsto \chi$) reduces the original multi-dimensional spectral problem for the transfer matrix to the set of one-dimensional spectral problems that greatly simplifies the analysis. We want to show that the maps $\Psi \mapsto \varphi$ and $\Psi \mapsto \chi$ can be extended to the isomorphism between the Hilbert spaces, $\mathbb{H}_N \mapsto \mathbb{H}_{\text{SoV}}$.

Let us define

$$\begin{aligned} (\varphi_1, \varphi_2)_{B_N} &= \int_{\mathbb{R} \times \mathbb{R}} \int_{\mathcal{D}_{N-1}^\sigma} (\varphi_1(p, x))^\dagger \varphi_2(p, x) \mu_{N-1}(x) d^2p d\mu_{N-1}^B(x), \\ (\chi_1, \chi_2)_{A_N} &= \int_{\mathcal{D}_N^\sigma} (\chi_1(x))^\dagger \chi_2(x) d\mu_N^A(x). \end{aligned} \quad (5.2)$$

The variables x_k, \bar{x}_k take the form $x_k = in_k/2 + \nu_k$, $\bar{x}_k = -in_k/2 + \nu_k$, where all n_k are either integers or half-integers,

$$n_k \in \mathbb{Z}^\sigma \equiv \mathbb{Z} + \frac{\sigma}{2}, \quad \sigma = 0, 1 \quad (5.3)$$

and

$$\mathcal{D}_N^\sigma \equiv (\mathbb{R} \times \mathbb{Z}^\sigma)^N. \quad (5.4)$$

The measures are defined as follows

$$d\mu_N^{B(A)}(x) = \mu_N^{B(A)}(x) \prod_{k=1}^N \mathcal{D}x_k, \quad \mu_N^{B(A)}(x) = c_N^{B(A)} \mu_N(x). \quad (5.5)$$

The symbol $\mathcal{D}x$ stands for

$$\int \mathcal{D}x \equiv \sum_{n \in \mathbb{Z}^\sigma} \int_{-\infty}^{\infty} d\nu. \quad (5.6)$$

The weight function $\mu_N(x)$ is given by the following expression

$$\mu_N(x_1, \dots, x_N) = \prod_{1 \leq k < j \leq N} x_{kj} \bar{x}_{kj} = \prod_{1 \leq k < j \leq N} \left(\nu_{kj}^2 + \frac{1}{4} n_{kj}^2 \right), \quad (5.7)$$

where $x_{kj} = x_k - x_j$, $\nu_{kj} = \nu_k - \nu_j$, $n_{kj} = n_k - n_j$ while the coefficients $c_N^{B(A)}$ take the form

$$(c_N^B)^{-1} = \frac{1}{2} (2\pi)^{N+1} N!, \quad (c_N^A)^{-1} = (2\pi)^N N!. \quad (5.8)$$

Let $\mathbb{H}_N^{B,\sigma}$, $\mathbb{H}_N^{A,\sigma}$ be the Hilbert spaces of symmetric functions corresponding to the scalar products (5.2):

$$\mathbb{H}_N^{B,\sigma} = L^2(\mathbb{R} \times \mathbb{R}) \otimes L_{\text{sym}}^2(\mathcal{D}_{N-1}^\sigma, d\mu_{N-1}^B(x)), \quad (5.9a)$$

$$\mathbb{H}_N^{A,\sigma} = L_{\text{sym}}^2(\mathcal{D}_N^\sigma, d\mu_N^A(x)). \quad (5.9b)$$

Given that $\varphi(p, x)$ and $\chi(x)$ are smooth and compactly supported functions on $\mathbb{R}^2 \times \mathcal{D}_{N-1}^\sigma$ and \mathcal{D}_N^σ , respectively, we introduce transforms $\mathbb{T}_N^B : \varphi \mapsto \Psi_\varphi$ and $\mathbb{T}_N^A : \chi \mapsto \Phi_\chi$,

$$\Psi_\varphi(z) \equiv [\mathbb{T}_N^B \varphi](z) = \int_{\mathbb{R}^2} \int_{\mathcal{D}_{N-1}^\sigma} \varphi(p, x) \Psi_{p,x}^{(N)}(z) d^2 p d\mu_{N-1}^B(x), \quad (5.10a)$$

$$\Phi_\chi(z) \equiv [\mathbb{T}_N^A \chi](z) = \int_{\mathcal{D}_N^\sigma} \chi(x) \Phi_{p,x}^{(N)}(z) d\mu_N^A(x). \quad (5.10b)$$

Note that the function Ψ_φ depends on the vector γ , Eq. (3.12), which appears in the definition of the function $\Psi_{p,x}^{(N)}$. That is $\mathbb{T}_N^B \equiv \mathbb{T}_N^B(\gamma)$ and the same applies to the operator \mathbb{T}_N^A . In order to not overload the notation we do not display this dependence explicitly.

5.1 B system

We begin the proof of the unitarity of the transform \mathbb{T}_N^B with the following lemma:

Lemma 1. *For any smooth fast decreasing function φ on $\mathbb{R}^2 \times \mathcal{D}_{N-1}^\sigma$ the function $\mathbb{T}_N^B \varphi$ belongs to the Hilbert space \mathbb{H}_N and it holds*

$$\|\mathbb{T}_N^B \varphi\|_{\mathbb{H}_N}^2 = \|\varphi\|_{\mathfrak{D}_{N-1}^{B,\pm}}^2 = \int_{\mathfrak{D}_{N-1}^\pm} |\varphi(p, x)|^2 d^2 p d\mu_{N-1}^B(x). \quad (5.11)$$

Proof: Let Ψ_φ^ϵ be a function defined by Eq. (5.10a) with $\Psi_{p,x}^{(N)}$ replaced by $\Psi_{p,x^\epsilon}^{(N),\epsilon}$, see Eqs. (4.3) and (4.18). It can be shown that $\Psi_\varphi^\epsilon(\vec{p}) \xrightarrow{\epsilon \rightarrow 0} \Psi_\varphi(\vec{p})$ almost everywhere. Next, taking into account Eq. (4.8) one gets

$$\left(\Psi_\varphi^\epsilon, \Psi_{\varphi'}^{\epsilon'} \right)_{\mathbb{H}_N} = \pi \int d^2 p \int d\mu_{N-1}^B(x) \int d\mu_{N-1}^B(x') (p\bar{p})^{\epsilon+\epsilon'} \varphi(p, x) (\varphi'(p, x'))^\dagger I^{\epsilon,\epsilon'}(x, x'), \quad (5.12)$$

with $I^{\epsilon,\epsilon'}(x, x')$ given by Eq. (4.11). Let us assume that the function $\varphi(\varphi')$ has the form

$$\varphi(p, x_1, \dots, x_{N-1}) = \kappa(p) \phi(x_1, \dots, x_{N-1}), \quad (5.13)$$

where $\phi(x_1, \dots, x_{N-1})$ is a symmetric function

$$\phi(x_1, \dots, x_{N-1}) = \sum_{S_{N-1}} \phi_1(x_{i_1}) \dots \phi_{N-1}(x_{i_{N-1}}) \quad (5.14)$$

and the sum goes over all permutations. We also assume that the functions $\phi_k(x_k) = \phi_k(n_k, \nu_k)$ are local in n_k , $\phi(n_k, \nu_k) = \delta_{n_k, m_k} \phi_k(\nu_k)$ and $\phi_k(\nu_k)$ is an analytic function of ν_k in some strip $|\text{Im}\nu_k| < \delta_k$ which vanishes sufficiently fast at $\nu_k \rightarrow \pm\infty$. Such functions form a dense subspace in the Hilbert space $\mathbb{H}_N^{B, \sigma}$. Since the momentum integral in (5.12) factorizes one has to consider the integrals over $x_k = (n_k, \nu_k)$, $x'_k = (n'_k, \nu'_k)$ which have the form

$$\int d\mu_{N-1}^B(x) \int d\mu_{N-1}^B(x') \dots \equiv \prod_{j=1}^{N-1} \sum_{n_j \in \mathbb{Z} + \frac{\sigma}{2}} \sum_{n'_j \in \mathbb{Z} + \frac{\sigma}{2}} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \mu_{N-1}^B(\vec{n}, \vec{\nu}) \mu_{N-1}^B(\vec{n}', \vec{\nu}') \prod_{k=1}^{N-1} d\nu_k d\nu'_k \dots \quad (5.15)$$

According to our assumptions only finite number of terms contribute to the sum in (5.15). Let us study behaviour of a particular term in the sum in the limit $\epsilon, \epsilon' \mapsto 0$. The functions ϕ, ϕ' are smooth and fast decreasing functions of ν, ν' . The function $I^{\epsilon, \epsilon'}(x, x')$ contains the factor $\Gamma[\epsilon + \epsilon' + i\bar{X} - i(X')^*] / \Gamma[\epsilon + \epsilon']$ and the product of the Γ -functions

$$\Gamma [i((\bar{x}'_k)^* - x_j)] = \Gamma \left[\frac{n'_k}{2} - \frac{n_j}{2} + i(\nu'_k - \nu_j) + \epsilon_{jk} \right] = \frac{\Gamma \left(\frac{n'_k}{2} - \frac{n_j}{2} + i(\nu'_k - \nu_j) + \epsilon_{jk} \right)}{\Gamma \left(1 + \frac{n'_k}{2} - \frac{n_j}{2} - i(\nu'_k - \nu_j) - \epsilon_{jk} \right)}, \quad (5.16)$$

where $\epsilon_{jk} \equiv \epsilon_j + \epsilon'_k$. In the $\epsilon'_k, \epsilon_j \rightarrow 0$ this function becomes singular at $\nu'_k = \nu_j$ if $n'_k = n_j$. Let us shift the contours of integrations over ν'_k variables to the upper half-plane, $\text{Im}\nu'_k = \delta > \epsilon_{jk}$, and pick up the residues at the corresponding poles. After this we can send $\epsilon'_k, \epsilon_j \mapsto 0$. Let us consider a generic contribution arising after this rearrangement. It has the form

$$\int_{C_\delta} \dots \int_{C_\delta} \prod_{k=1}^M d\nu'_k f(x_1, \dots, x_{N-1}, S(x'_1), \dots, S(x'_{N-1})), \quad (5.17)$$

where $S(x'_k) = x'_k$ if $k \in (i_1, \dots, i_M)$ and $S(x'_k) = x_{p_k}$ if k does not belong to this set. The integrand f is given by the product of the functions ϕ_k, ϕ'_k, Γ -functions (5.16) and the factor $A = \Gamma[\epsilon + \epsilon' + i\bar{X} - i(X')^*] / \Gamma[\epsilon + \epsilon']$. All these factors are regular on the contours of integration. Moreover, if $M \geq 1$ the last factor, A , tends to zero at $\epsilon, \epsilon' \mapsto 0$. Thus the only non-vanishing contribution comes from the term with $M = 0$, i.e. when all $x'_k \mapsto x_{i_k}$ for $k = 1, \dots, N-1$. It takes the form

$$\left(\Psi_\varphi^\epsilon, \Psi_{\varphi'}^{\epsilon'} \right)_{\mathbb{H}_N} = \int d^2p \int d\mu_{N-1}^B(x) \varphi(p, x) (\varphi'(p, x))^\dagger + O(\epsilon + \epsilon') \quad (5.18)$$

that results in the following estimate for the norm of the function Ψ_φ^ϵ ,

$$\|\Psi_\varphi^\epsilon\|_{\mathbb{H}_N}^2 = K + O(\epsilon), \quad (5.19)$$

where

$$K = \|\varphi\|_{\mathbb{H}_N^{B, \sigma}}^2 \equiv \int_{\mathbb{R}^2} \int_{\mathcal{D}_{N-1}^\sigma} |\varphi(p, x)|^2 d^2p d\mu_{N-1}^B(x). \quad (5.20)$$

Since $\Psi_\varphi^\epsilon(\vec{p}) \mapsto \Psi_\varphi(\vec{p})$ at $\epsilon \rightarrow 0$ it follows from Fatou's theorem that $\|\Psi_\varphi\|_{\mathbb{H}_N}^2 < K$. At the same time the inequality

$$\|\Psi_\varphi - \Psi_\varphi^\epsilon\|_{\mathbb{H}_N}^2 \geq 0 \quad (5.21)$$

implies $\|\Psi_\varphi\|_{\mathbb{H}_N}^2 \geq K$ that results in $\|\Psi_\varphi\|_{\mathbb{H}_N}^2 = K$.

Since the set of functions (5.13), (5.14) is dense in the Hilbert spaces $\mathbb{H}_N^{B,\pm}$ the transformation T_N^B can be extended to the entire Hilbert space $\mathbb{H}_N^{B,\pm}$ and (5.22a) holds for any function $\varphi \in \mathbb{H}_N^{B,\pm}$. ■

Taking this result into account we formulate the following theorem.

Theorem 1. *The map T_N^B defined in Eq. (5.10a) can be extended to the linear bijective isometry of the Hilbert spaces, $\mathbb{H}_N^{B,\sigma} \mapsto \mathbb{H}_N$, i.e.*

$$\|T_N^B \varphi\|_{\mathbb{H}_N}^2 = \|\varphi\|_{\mathbb{H}_N^{B,\sigma}}^2 \quad (5.22a)$$

and

$$\mathcal{R}(T_N^B) = \mathbb{H}_N. \quad (5.22b)$$

Proof: Eq. (5.22a) is a direct consequence of Lemma 1. It implies that $\|T_N^B\| = 1$, hence $\mathcal{R}(T_N^B)$ is a closed subspace in \mathbb{H}_N and $\mathbb{H}_N = \mathcal{R}(T_N^B) \oplus \mathcal{R}(T_N^B)^\perp$. Since $\mathcal{R}(T_N^B)^\perp = \ker(T_N^B)^*$ in order to prove (5.22b) it is enough to show that $\ker(T_N^B)^* = 0$.

We prove this statement using induction on N . For $N = 1$ the map $T_{N=1}^B$ is a two-dimensional Fourier transform, hence Eq. (5.22b) is true. Let us now assume that $\mathcal{R}(T_N^B) = \mathbb{H}_N$ and prove that it implies $\mathcal{R}(T_{N+1}^B) = \mathbb{H}_{N+1}$. As was stated above it is sufficient to prove that $\ker(T_{N+1}^B)^* = 0$. To this end let us consider the map

$$S_N = (T_{N+1}^B)^* (T_N^B \otimes T_1^B), \quad \mathbb{H}_N^{B,\sigma} \otimes L^2(\mathbb{R}^2) \xrightarrow{T_N^B \otimes T_1^B} \mathbb{H}_{N+1} \xrightarrow{(T_{N+1}^B)^*} \mathbb{H}_{N+1}^{B,\sigma}. \quad (5.23)$$

Since by the assumption $T_N^B \otimes T_1^B$ is a bijective isometry $\ker S_N = 0$ if and only if $\ker(T_{N+1}^B)^* = 0$.

The adjoint operator $(T_{N+1}^B)^*$ is a bounded operator which acts on a vector $\Psi \in \mathbb{H}_{N+1}$ by projecting it on the eigenfunction $\Psi_{p,x}^{(N+1)}$,

$$(T_{N+1}^B)^* \Psi = \left(\Psi_{p,x}^{(N+1)}, \Psi \right)_{\mathbb{H}_{N+1}} = \left(\Psi_{p,x}^{(N+1)}, P_{N+1} \Psi \right)_{\mathbb{H}_{N+1}} \equiv \varphi(p, x), \quad (5.24)$$

where P_{N+1} is the projector on $\mathcal{R}(T_{N+1}^B)$. It follows from (5.24) that

$$\|\varphi\|_{\mathbb{H}_{N+1}^{B,\sigma}}^2 = \int_{\mathbb{R}^2} \int_{\mathcal{D}_N^\sigma} |\varphi(p, x)|^2 d^2 p d\mu_N^B(x) = \|P_{N+1} \Psi\|_{\mathbb{H}_{N+1}}^2 \leq \|\Psi\|_{\mathbb{H}_{N+1}}^2. \quad (5.25)$$

For $\phi \in \mathbb{H}_N^{B,\sigma} \otimes L^2(\mathbb{R}^2)$ the function $\Psi_\phi = (T_N^B \otimes T_1^B) \phi$ reads

$$\Psi_\phi(z) = \int_{\mathbb{R}^2 \otimes \mathbb{R}^2} \int_{\mathcal{D}_{N-1}^\sigma} \Psi_{q_1, x}^{(N)}(z_1, \dots, z_N) \Psi_{q_2}^{(1)}(z_{N+1}) \phi(q_1, q_2, x) d^2 q_1 d^2 q_2 d\mu_{N-1}^B(x). \quad (5.26)$$

Replacing $\Psi_{q_1, x}^{(N)} \mapsto \Psi_{q_1, x}^{(N), \epsilon}$ in (5.26) we define a new function, Ψ_ϕ^ϵ . Since by Lemma 1 $\Psi_\phi^\epsilon \xrightarrow{\epsilon \rightarrow 0^+} \Psi_\phi$ in \mathbb{H}_{N+1} for smooth rapidly decreasing functions we obtain

$$\varphi(p, x) = [S_N \phi](p, x) = \left(\Psi_{p,x}^{(N+1)}, \Psi_\phi \right)_{\mathbb{H}_{N+1}} = \lim_{\epsilon \rightarrow 0^+} \left(\Psi_{p,x}^{(N+1)}, \Psi_\phi^\epsilon \right)_{\mathbb{H}_{N+1}} \equiv \lim_{\epsilon \rightarrow 0^+} \varphi_\epsilon(p, x), \quad (5.27)$$

where

$$\varphi_\epsilon(p, x) = \int_{\mathbb{R}^2 \times \mathbb{R}^2} \int_{\mathcal{D}_{N-1}^\sigma} S_N^\epsilon(p, x|q_1, q_2, x') \phi(q_1, q_2, x') d^2 q_2 d^2 q_1 d\mu_{N-1}^B(x'). \quad (5.28)$$

The kernel S_N^ϵ reads

$$S_N^\epsilon(p, x|q_1, q_2, x') = \left(\Psi_{p,x}^{(N+1)}, \Psi_{q_1, x'_\epsilon}^{(N)} \otimes \Psi_{q_2}^{(1)} \right), \quad (5.29)$$

see Eq. (4.23), and $x'_\epsilon = (x'_1 + i\epsilon_1, \dots, x'_{N-1} + i\epsilon_{N-1})$. We assume that function ϕ takes the form

$$\phi(q_1, q_2, x_1, \dots, x_{N-1}) = \kappa_1(q_1) \kappa_2(q_2) \sum_{S_{N-1}} \phi_1(x_{i_1}) \dots \phi_{N-1}(x_{i_{N-1}}), \quad (5.30)$$

where the sum goes over all permutations and that the functions ϕ_k are local in “ n ” variable, that is $\phi_k(x_k) = \phi_k(n_k, \nu_k) = \delta_{n_k m_k} \phi_{n_k}(\nu_k)$ and ϕ_{n_k} are compactly supported.

The function $\varphi(p, y)$ does not decrease sufficiently fast for large y_k in order to justify changing the order of integration after substituting $\varphi_\epsilon(p, y)$ in the form (5.27), (5.28) into (5.25). To overcome this difficulty we, following the lines of ref. [29], consider the integral

$$I_Z(\varphi) = \int_{\mathbb{R}^2} \int_{\mathcal{D}_N^\sigma} |\varphi(p, y)|^2 \Omega_Z(y) d^2 p d\mu_N^B(y), \quad (5.31)$$

where

$$\Omega_Z(y) = \prod_{k=1}^N \frac{\Gamma[Z + iy_k, Z - iy_k]}{\Gamma[Z, Z]}, \quad Z = \bar{Z} = \frac{1}{2} + iM. \quad (5.32)$$

For $y_k^* = \bar{y}_k$ the factor Ω is a pure phase, $|\Omega_Z(y)| = 1$ and $\Omega_Z(y) \mapsto 1$ when $M \rightarrow \infty$, y is fixed. Since the integral (5.25) is convergent

$$\|\varphi\|_{\mathbb{H}_{N+1}^{B,\sigma}}^2 = \lim_{M \rightarrow \infty} \int_{\mathbb{R}^2} \int_{\mathcal{D}_N^\sigma} |\varphi(p, y)|^2 \Omega_Z(y) d^2 p d\mu_N^B(y). \quad (5.33)$$

It follows from Eqs. (5.28), (5.29) and (4.23) that for compactly supported functions ϕ_k the function $f(\nu) = |\varphi_\epsilon(p, y)|^2$ is an analytic function of ν_k in the vicinity of the real axis for sufficiently large ν_k . Thus we can write

$$I_Z(\varphi) = \lim_{\omega \rightarrow 0} I_Z^\omega(\varphi) = \lim_{\omega \rightarrow 0} \int_{\mathbb{R}^2} \int_{\mathcal{D}_N^{\sigma,\omega}} |\varphi(p, y)|^2 \Omega_{Z-\omega}(y) d^2 p d\mu_N^B(y), \quad (5.34)$$

where the integration contours over ν_k are deformed in order to separate the poles due to the Gamma functions, $\Gamma[Z - \omega \pm iy_k]$, in the factor Ω . The integral $I_Z^\omega(\varphi)$ is an analytic function of ω . Substituting $\varphi(p, y)$ in (5.34) in the form (5.28) one can show that for $\text{Re } \omega > 1$ the integrals over y decay fast enough to allow the change of the order of integration over x, x' and y . Thus we obtain

$$\begin{aligned} I_Z^\omega(\varphi) &= \lim_{\epsilon, \epsilon' \rightarrow 0^+} \int_{\mathbb{R}^2 \times \mathbb{R}^2} \int_{\mathcal{D}_{N-1}^\sigma \times \mathcal{D}_{N-1}^\sigma} \delta^{(2)}(q_1 + q_2 - q'_1 - q'_2) \phi(q_1, q_2, x) (\phi(q'_1, q'_2, x'))^\dagger \left| \frac{q_1 + q_2}{q_1 q'_2} \right|^2 \left| \frac{q'_1}{q_1} \right|^{N-1} \times \\ &\quad \left(1 + \frac{q'_1}{q_2} \right)^{iX'} \left(1 + \frac{\bar{q}'_1}{\bar{q}'_2} \right)^{i\bar{X}'} \left(1 + \frac{q_1}{q_2} \right)^{-iX} \left(1 + \frac{\bar{q}_1}{\bar{q}_2} \right)^{-i\bar{X}} \left(\frac{q_1}{q'_1} \right)^{A_N} \left(\frac{\bar{q}_1}{\bar{q}'_1} \right)^{\bar{A}_N} \left(\frac{q'_2}{q_2} \right)^{\gamma_{2N}} \left(\frac{\bar{q}'_2}{\bar{q}_2} \right)^{\bar{\gamma}_{2N}} \times \\ &\quad R(x, x') J_\omega^{(\epsilon)}(Z, \zeta, x, x') d^2 q_1 d^2 q_2 d^2 q'_1 d^2 q'_2 d\mu_{N-1}^B(x) d\mu_{N-1}^B(x'), \end{aligned} \quad (5.35)$$

where $\zeta = \frac{q_1 q_2'}{q_2 q_1'}$, A_N is defined in Eq. (4.24),

$$R(x, x') = \prod_{k=1}^N \prod_{j=1}^{N-1} \Gamma \left[\bar{\gamma}_{2N-k}^{(k-1)} - ix'_j \right] / \Gamma \left[\bar{\gamma}_{2N-k}^{(k-1)} - ix_j \right] \quad (5.36)$$

and

$$J_{\omega}^{(\epsilon, \epsilon')} (Z, \zeta, x, x') = \pi^2 \int_{\mathcal{D}_N^{\omega, \sigma}} \zeta^{iY} \bar{\zeta}^{i\bar{Y}} \prod_{j=1}^N \frac{\Gamma[Z - \omega \pm iy_j]}{\Gamma^2(Z)} \prod_{k=1}^{N-1} \Gamma[i(\bar{x}_k - \bar{y}_j)] \Gamma[i(y_j - x'_k)] d\mu_N^B(y). \quad (5.37)$$

We recall that the variables ν_k, ν'_k , ($x_k = in_k/2 + \nu_k$, $x'_k = in'_k/2 + \nu'_k$) have small negative (positive) imaginary parts, $\text{Im}\nu_k = -\epsilon_k$, $\text{Im}\nu'_k = \epsilon'_k$, which must be sent to zero at the end of the calculation.

The integral (5.37) can be obtained in the closed form with the help of Eq. (C.4). Indeed,

$$\begin{aligned} \prod_{1 \leq j \neq k \leq N} \frac{1}{\Gamma[i(y_k - y_j)]} &= \mu_N(y) (-1)^{\sum_{k < j} [i(y_k - y_j)]}, \\ \prod_{j=1}^N \prod_{k=1}^{N-1} \Gamma[i(\bar{x}_k - \bar{y}_j)] &= \prod_{j=1}^N \prod_{k=1}^{N-1} \Gamma[i(x_k - y_j)] (-1)^{\sum_{j=1}^N \sum_{k=1}^{N-1} [i(y_j - x_k)]}, \end{aligned} \quad (5.38)$$

where $y_k = im_k/2 + \nu_k$, $\bar{y}_k = -im_k/2 + \nu_k$ and we recall that $[iy_k] = i(y_k - \bar{y}_k) = -m_k$. Taking into account that

$$(-1)^{\sum_{k < j} [i(y_k - y_j)]} (-1)^{\sum_{j=1}^N \sum_{k=1}^{N-1} [i(y_j - x_k)]} = (-1)^{\sum_{1 \leq k < j \leq N-1} [i(x_k - x_j)]} \quad (5.39)$$

one finds that the integral (5.37) is nothing else as Gustafson's integral (C.4) [$u_k \rightarrow iy_k$ for all k , $\{z_1, \dots, z_N\} \mapsto \{ix_1, \dots, ix_{N-1}, Z - \omega\}$ and $\{w_1, \dots, w_N\} \mapsto \{-ix'_1, \dots, -ix'_{N-1}, Z - \omega\}$]. Thus we obtain for $J_{\omega}^{(\epsilon, \epsilon')}$:

$$\begin{aligned} J_{\omega}^{(\epsilon, \epsilon')} (Z, \zeta, x, x') &= \pi (-1)^{\sum_{k < j} [i(x_k - x_j)]} \frac{\Gamma[2Z - 2\omega]}{\Gamma^2[Z]} \frac{\zeta^{Z - \omega + iX}}{(1 + \zeta)^{2(Z - \omega) + i(X - X')}} \frac{\bar{\zeta}^{\bar{Z} - \omega + i\bar{X}}}{(1 + \bar{\zeta})^{2(\bar{Z} - \omega) + i(\bar{X} - \bar{X}')}} \times \\ &\quad \prod_{k=1}^{N-1} \frac{\Gamma[Z - \omega + ix_k, Z - \omega - ix'_k]}{\Gamma[Z, Z]} \prod_{k, j=1}^{N-1} \Gamma[i(x_k - x'_j)]. \end{aligned} \quad (5.40)$$

Let us substitute this expression into (5.35) and calculate the corresponding limits. First of all, since all factors containing ω are regular at $\omega, \epsilon_k, \epsilon'_k \rightarrow 0$ one can interchange the limits and first send $\omega \rightarrow 0$.

At $M \rightarrow \infty$ the integral over q, q' is dominated by the contribution from the stationary point at $\zeta = 1$,

$$\frac{\Gamma[1 + 2iM]}{\Gamma^2[\frac{1}{2} + iM]} \int d^2\zeta \frac{(\zeta \bar{\zeta})^{iM + \frac{1}{2}}}{((1 + \zeta)(1 + \bar{\zeta}))^{1 + 2iM}} \varphi(\zeta) \stackrel{M \rightarrow \infty}{=} \pi \varphi(1) \left(1 + O\left(\frac{1}{M^{1/2}}\right) \right). \quad (5.41)$$

Taking this into account and expanding the first factor in the second line in (5.40) one gets for (5.35)

$$\begin{aligned} I_{\omega=0}(Z) &= \lim_{\epsilon, \epsilon' \rightarrow 0^+} \int_{\mathcal{D}_{N-1}^{\sigma} \times \mathcal{D}_{N-1}^{\sigma}} \phi(q_1, q_2, x) (\phi(q_1, q_2, x'))^{\dagger} \times \\ &\quad \pi (-1)^{\sum_{k < j} [i(x_k - x_j)]} i^{N-N'} R(x, x') \left(1 + \frac{q_1}{q_2} \right)^{i(X' - X)} \left(1 + \frac{\bar{q}_1}{\bar{q}_2} \right)^{i(\bar{X}' - \bar{X})} \times \\ &\quad \left(\frac{M}{2} \right)^{2i(\mathcal{V} - \mathcal{V}')} \prod_{k, j=1}^{N-1} \Gamma[i(x_k - x'_j) + \epsilon_{kj}] d^2q_1 d^2q_2 d\mu_{N-1}^B(x) d\mu_{N-1}^B(x') + \dots, \end{aligned} \quad (5.42)$$

where ellipses stand for terms vanishing at $M \rightarrow \infty$ and

$$x_k = \frac{in_k}{2} + \nu_k, \quad x'_k = \frac{in'_k}{2} + \nu'_k, \quad \epsilon_{kj} = \epsilon_k + \epsilon'_j, \quad X = \sum_{k=1}^{N-1} x_k, \quad \mathcal{V} = \sum_{k=1}^{N-1} \nu_k, \quad N = \sum_{k=1}^{N-1} n_k,$$

etc. The analysis of this integral is similar to the analysis of the integral (5.12) **. In the limit $\epsilon, \epsilon' \rightarrow 0$ the poles of the Gamma functions, $x_k = x'_j$, approach the integration contour, while all other factors remain regular. Let us shift the integration contour in x_k to the upper complex half-plane picking up the residues at the poles at $x_k = x'_j$. We recall that the Gamma functions develop poles only when $n_k = n'_j$, otherwise they are regular at $\nu_k = \nu'_j$. Afterwards we can send $\epsilon, \epsilon' \rightarrow 0$. The answer is given by the sum of terms

$$\int \dots \int M^{i \sum_{k=1}^m (\nu_{i_k} - \nu'_{j_k})} \times f_m(x, x') d\nu_{i_1} \dots d\nu_{i_m} d\nu'_{j_1} \dots d\nu'_{j_m}, \quad (5.43)$$

where $f_m(x, x')$ is a smooth function. Note, the contours of integration over ν variables lay in the upper half-plane, so that $|M^{i \sum_{k=1}^m (\nu_{i_k} - \nu'_{j_k})}| < 1$ in the integration region. Since the functions $f_m(x, x')$ are smooth functions all such terms with $m > 0$ vanish after integration in the limit $M \rightarrow \infty$. Thus the only contribution with $m = 0$, i.e. when $x_k = x'_{k_j}$, survives in this limit. Then one obtains after some algebra

$$\|\varphi\|_{\mathbb{H}_{N+1}^{B,\sigma}}^2 = \|S_N \phi\|_{\mathbb{H}_{N+1}^{B,\sigma}}^2 = \int_{\mathbb{R}^2 \times \mathbb{R}^2} \int_{\mathcal{D}_{N-1}^\sigma} |\phi(q_1, q_2, x)|^2 d^2 q_1 d^2 q_2 d\mu_{N-1}^B(x) = \|\phi\|_{\mathbb{H}_N^{B,\sigma} \otimes L^2(\mathbb{R}^2)}^2. \quad (5.44)$$

Since the space of functions (5.30) dense in $\mathbb{H}_N^{B,\sigma} \otimes L^2(\mathbb{R}^2)$ this relation can be extended to the whole Hilbert space. Thus one concludes that $\ker S_N = 0$, and, hence, $\ker (\mathbb{T}_{N+1}^B)^* = 0$. ■

5.2 A system

Using the results of the previous section it becomes quite easy to prove the unitarity of \mathbb{T}_N^A transform. First, we prove an analogue of the Lemma 1

Lemma 2. *For any smooth fast decreasing function χ on \mathcal{D}_N^σ the function $\mathbb{T}_N^A \chi$, Eq. (5.10b), belongs to the Hilbert space \mathbb{H}_N and it holds*

$$\|\mathbb{T}_N^A \chi\|_{\mathbb{H}_N}^2 = \|\chi\|_{\mathbb{H}_N^{A,\sigma}}^2 = \int_{\mathcal{D}_N^\sigma} |\chi(x)|^2 d\mu_N^A(x). \quad (5.45)$$

Proof: The proof is similar to the proof of the lemma 1. It suffices to prove (5.45) for functions of the form

$$\chi(x_1, \dots, x_N) = \sum_{S_N} \chi_1(x_{i_1}) \dots \chi_N(x_{i_N}), \quad \chi_k(x_k) = \chi_k(n_k, \nu_k) = \delta_{n_k m_k} \chi_k(\nu_k). \quad (5.46)$$

We assume that the functions $\chi_k(\nu)$ are analytic in some strip near the real axis. Let us calculate the projection

$$\varphi_\chi(p, y) = \left(\Psi_{p,y}^{(N)}, \Phi_\chi \right) = \lim_{\epsilon \rightarrow 0} \int_{\mathcal{D}_N^\sigma} \left(\Psi_{p,y}^{(N)}, \Phi_{x+i\epsilon}^{(N)} \right) \chi(x) d\mu_N^A(x). \quad (5.47)$$

**We do it assuming that the functions $\phi_k(x_k)$ have the properties discussed around Eq. (5.13).

Here we have given the variables $x_k \rightarrow x_k + i\epsilon_k$, $\epsilon_k = \bar{\epsilon}_k > 0$ small imaginary parts which allows us to change the order of integration. In order to show that $\|\varphi_\chi\|_{\mathbb{H}_N^{B,\sigma}} = \|\chi\|_{\mathbb{H}_N^{A,\sigma}}$ we write

$$\|\varphi_\chi\|_{\mathbb{H}_N^{B,\sigma}}^2 = \int_{\mathbb{R}^2} \int_{\mathcal{D}_{N-1}^\sigma} |\varphi(p, y)|^2 d^2 p d\mu_{N-1}^B(y) = \lim_{\sigma \rightarrow 0} \int_{\mathbb{R}^2} e^{-\sigma|p|^2} \left(\int_{\mathcal{D}_{N-1}^\sigma} |\varphi(p, y)|^2 d\mu_{N-1}^B(y) \right) d^2 p. \quad (5.48)$$

Using the representation (5.47) for $\varphi_\chi(p, y)$ we first evaluate the y -integral $\dagger\dagger$. This integral coincides with the so-called SL(2, \mathbb{C}) Gustafson integral and can be evaluated in a closed form (C.1) resulting in

$$\|\varphi_\chi\|_{\mathbb{H}_N^{B,\sigma}}^2 = \frac{1}{\pi} \lim_{\sigma \rightarrow 0} \lim_{\epsilon, \epsilon' \rightarrow 0^+} \int_{\mathbb{R}^2} \int_{\mathcal{D}_N^\sigma \times \mathcal{D}_N^\sigma} e^{-\sigma|p|^2} i^{N-N'} p^{i(X'-X)-1+\mathcal{E}+\mathcal{E}'} \bar{p}^{i(\bar{X}'-\bar{X})-1+\mathcal{E}+\mathcal{E}'} (-1)^{\sum_{k<j} [i(x'_k-x'_j)]} \frac{\chi(x)}{\left(\prod_{j=1}^N \vartheta_N(x_j)\right)} \left(\frac{\chi(x')}{\left(\prod_{j=1}^N \vartheta_N(x'_j)\right)}\right)^\dagger \frac{\prod_{k,j=1}^N \Gamma[i(x'_k-x_j)+\epsilon_{jk}]}{\Gamma[i(X'-X)+\mathcal{E}+\mathcal{E}']} d\mu_N^A(x) d\mu_N^A(x') d^2 p, \quad (5.49)$$

where $X = \sum_{k=1}^N x_k$, $N = \sum_{k=1}^N n_k$, $\mathcal{E} = \sum_{k=1}^N \epsilon_k$, $\epsilon_{jk} = \epsilon_j + \epsilon'_k$, etc. For the momentum integral one gets

$$\pi \delta_{NN'} \sigma^{i(\mathcal{V}-\mathcal{V}')-\mathcal{E}-\mathcal{E}'} \Gamma(i(\mathcal{V}'-\mathcal{V})+\mathcal{E}+\mathcal{E}'), \quad (5.50)$$

where Γ is Euler's gamma function. Thus

$$\|\varphi_\chi\|_{\mathbb{H}_N^{B,\sigma}}^2 = \lim_{\sigma \rightarrow 0} \lim_{\epsilon, \epsilon' \rightarrow 0^+} \int_{\mathcal{D}_N^\sigma \times \mathcal{D}_N^\sigma} (-1)^{\sum_{k<j} [i(x'_k-x'_j)]} \delta_{NN'} \sigma^{i(\mathcal{V}-\mathcal{V}')} \prod_{k,j=1}^N \Gamma[i(x'_k-x_j)+\epsilon_{jk}] \Gamma(1+i(\mathcal{V}-\mathcal{V}')) \frac{\chi(x)}{\left(\prod_{j=1}^N \vartheta_N(x_j)\right)} \left(\frac{\chi(x')}{\left(\prod_{j=1}^N \vartheta_N(x'_j)\right)}\right)^\dagger d\mu_N^A(x) d\mu_N^A(x'), \quad (5.51)$$

where we put $\epsilon_k, \epsilon'_k = 0$ in all nonsingular factors. The analysis of this integral in the $\sigma, \epsilon, \epsilon' \rightarrow 0$ limit is exactly the same as in Theorem 1, see discussion around Eq. (5.42), and results in

$$\|\Phi_\chi\|_{\mathbb{H}_N}^2 = \|\varphi_\chi\|_{\mathbb{H}_N^{B,\sigma}}^2 = \int_{\mathcal{D}_N^\sigma} |\varphi(x)|^2 d\mu_N^A(x). \quad (5.52)$$

Since the space of the functions (5.46) is dense in $\mathbb{H}_N^{A,\sigma}$ the relation (5.52) extends to the whole Hilbert space. \blacksquare

Finally, we formulate the analog of Theorem 1 for the map \mathbb{T}_N^A .

Theorem 2. *The map \mathbb{T}_N^A defined in Eq. (5.10b) can be extended to the linear bijective isometry of the Hilbert spaces, $\mathbb{H}_N^{A,\sigma} \mapsto \mathbb{H}_N$, i.e.*

$$\|\mathbb{T}_N^A \chi\|_{\mathbb{H}_N}^2 = \|\varphi\|_{\mathbb{H}_N^{A,\sigma}}^2 \quad (5.53a)$$

and

$$\mathcal{R}(\mathbb{T}_N^A) = \mathbb{H}_N. \quad (5.53b)$$

Proof: As in the Theorem 1 we only need to prove Eq. (5.53b). As was discussed in earlier Eq. (5.53b) is equivalent to the statement that $\ker(\mathbb{T}_N^A)^* = 0$ or to the assertion $\ker \mathbf{S}_N = 0$, where $\mathbf{S}_N = (\mathbb{T}_N^A)^* \mathbb{T}_N^B$. In order to prove this it suffices to show that $\|\mathbf{S}_N \varphi\|_{\mathbb{H}_N^{A,\sigma}} = \|\varphi\|_{\mathbb{H}_N^{B,\sigma}}$. The proof of this statement repeats step by step the proof given in the Theorem 1, and on the technical level is reduced to the evaluation of the integral (5.37). \blacksquare

$\dagger\dagger$ The x, x', y integral can be interchanged since the integral of modulus is convergent.

6 Summary

In this work we consider a generic inhomogeneous $SL(2, \mathbb{C})$ spin chain with impurities and construct the eigenfunctions of the B and A entries of the monodromy matrix. We prove the unitarity of the SoV transform associated with these systems or, equivalently, the completeness of the corresponding systems in the Hilbert space of the model. Namely, the following identities hold in the sense of distributions,

$$\begin{aligned} \int_{\mathbb{R}^2} \int_{\mathcal{D}_{N-1}^\sigma} \Psi_{p,x}^{(N)}(z) (\Psi_{p,x}^{(N)}(z'))^\dagger d^2 p d\mu_N^B(x) &= \prod_{k=1}^N \delta^2(z_k - z'_k), \\ \int_{\mathcal{D}_N^\sigma} \Phi_x^{(N)}(z) (\Phi_x^{(N)}(z'))^\dagger d\mu_N^A(x) &= \prod_{k=1}^N \delta^2(z_k - z'_k), \end{aligned} \quad (6.1)$$

and

$$\begin{aligned} \int_{\mathbb{C}^N} \Psi_{p,x}^{(N)}(z) (\Psi_{p',x'}^{(N)}(z))^\dagger \prod_{k=1}^N d^2 z_k &= (\mu_N^B(x))^{-1} \delta^2(p - p') \delta^{N-1}(x, x'), \\ \int_{\mathbb{C}^N} \Phi_x^{(N)}(z) (\Phi_{x'}^{(N)}(z))^\dagger \prod_{k=1}^N d^2 z_k &= (\mu_N^A(x))^{-1} \delta^N(x, x'), \end{aligned} \quad (6.2)$$

where

$$\delta^N(x, x') = \frac{1}{N!} \sum_{w \in S_N} \delta^N(x' - wx) \quad , \quad wx = (x_{w_1}, \dots, x_{w_N}) \quad (6.3)$$

and $\delta^N(x' - x) = \prod_{k=1}^N \delta^2(x'_k - x_k)$, $\delta^2(x' - x) = \delta_{nn'} \delta(\nu - \nu')$.

The method relies heavily on the use of multidimensional Mellin-Barnes integrals which generalize integrals calculated by R. A. Gustafson [28]. The attractive feature of our approach is that it does not depend on the details of the spin chain such as spins and inhomogeneity parameters. We believe that this technique can also be used to prove the unitarity of the SoV transform for the open $SL(2, \mathbb{C})$ spin chain.

Acknowledgements

The author is grateful to S. É. Derkachov for fruitful discussions and T. A. Sinkevich for critical remarks.

Appendix A The diagram technique

Throughout this paper we used a diagrammatic representation for the functions under consideration. The calculation of relevant scalar products is, most conveniently, performed diagrammatically with the help of a few simple identities. Below we give some of these rules (see also ref. [16]).

- (i) An arrow with the index α directed from w to z stands for a propagator $D_\alpha(z - w) = [z - w]^{-\alpha}$:

$$w \xrightarrow{\alpha} z = [z - w]^{-\alpha}$$

- (ii) The Fourier transform reads

$$\int d^2 z e^{i(pz + \bar{p}\bar{z})} D_\alpha(z) = \pi i^{\alpha - \bar{\alpha}} a(\alpha) D_{1-\alpha}(p), \quad (A.1)$$

where the function $a(\alpha) \equiv 1/\mathbf{\Gamma}[\alpha] = \Gamma(1 - \bar{\alpha})/\Gamma(\alpha)$.

(iii) Chain rule

$$\int \frac{d^2 w}{[z_1 - w]^\alpha [w - z_2]^\beta} = \pi \frac{a(\alpha, \beta)}{a(\gamma)} \frac{1}{[z_1 - z_2]^\gamma}, \quad (\text{A.2})$$

where $\gamma = \alpha + \beta - 1$. Its diagrammatic form is

$$\begin{array}{c} \alpha \quad \beta \\ \longrightarrow \bullet \longrightarrow \end{array} = \pi \frac{a(\alpha)a(\beta)}{a(\gamma)} \begin{array}{c} \gamma \\ \longrightarrow \end{array}$$

(iv) Star-triangle relation

$$\begin{array}{c} \alpha \\ \uparrow \\ \gamma \swarrow \bullet \searrow \beta \end{array} = \pi a(\alpha, \beta, \gamma) \begin{array}{c} 1-\beta \quad 1-\gamma \\ \swarrow \quad \searrow \\ \triangle \\ \swarrow \quad \searrow \\ 1-\alpha \end{array} \quad (\text{A.3})$$

(v) Exchange relation

$$\begin{array}{c} a(\alpha', \bar{\beta}') \\ \uparrow \\ \alpha' - \alpha \\ \uparrow \\ \alpha \quad \beta \\ \swarrow \bullet \searrow \\ 1-\alpha' \quad 1-\beta' \end{array} = \begin{array}{c} \alpha' \quad \beta' \\ \swarrow \quad \searrow \\ \triangle \\ \swarrow \quad \searrow \\ 1-\alpha \quad 1-\beta \\ \uparrow \\ \beta - \beta' \\ \uparrow \\ a(\alpha, \bar{\beta}) \end{array} \quad (\text{A.4})$$

where $\alpha + \beta = \alpha' + \beta'$.

Appendix B Scalar products

Here we discuss the calculation of scalar products of $\Psi_{p,x}^{(N),\epsilon}$ and $\Phi_x^{(N)}$ functions. The diagrams for the scalar products (4.8), (4.20) are shown in Fig. 2. The leftmost vertex on both diagrams has only two propagators attached to it. We call such a vertex – free vertex. On the first step one integrates over the free vertex (on both diagrams) using the chain relation for propagators and move the resulting line to the right with the help of the exchange relation. After that two new free vertices appear and one repeat the same procedure again. In this way one can integrate over all vertices on the left edge of both diagrams (they are shown by black blobs). Keeping trace of all factors arising in the process one represent the initial diagram D as

$$D_N(\{x_1, x_2, \dots\}, \{y_1, y_2, \dots\}, \{\gamma_1, \gamma_2, \dots\}) = f(x_1, y_1, \gamma) D'_N(\{x_2, \dots\}, \{y_2, \dots\}, \{\gamma_3, \dots\}). \quad (\text{B.1})$$

Taking into account that the function $\Psi_{p,x}^{(N)}$ and $\Phi_x^{(N)}$ are symmetric functions of the separated variables it follows from (B.1) that

$$D_N(\{x_1, x_2, \dots\}, \{y_1, y_2, \dots\}, \{\gamma_1, \gamma_2, \dots\}) = C_N(\gamma) \prod_{k,j} f(x_k, y_j, \gamma). \quad (\text{B.2})$$

The factor $C_N(\gamma)$ does not depend on x, y variables. The easiest way to fix it is to evaluate both sides of (B.2) for special values of x, y . For example, one can take $x_k \rightarrow x$ and $y_k \rightarrow \bar{x}^*$. Both sides, in this limits, contain divergent factors, $\Gamma[i(\bar{y}_j^* - x_k)]$ which cancel out. It is easy check that the result of the




Figure 2: Examples of diagrams for scalar products, Eqs. (4.8), (4.20) for $N = 4$.

integration over any free vertex in this limit (after removing this singular factor) gives one. Therefore the equation on $C_N(\gamma)$ for the scalar product (4.11) takes the form

$$1 = C_N(\gamma)(\chi(x)(\bar{\chi}(\bar{x}^*))^*)^{N-1} = C_N(\gamma)(-1)^{(N-1)\sum_{k=0}^{N-3}[\gamma_{2N-3-k}^{(k)} - ix]}. \quad (\text{B.3})$$

Since $[\gamma_m^{(k)} - ix]$ is an integer number one gets that $C_N = 1$ for odd N , while for even N

$$\begin{aligned} \sum_{k=0}^{N-3} [\gamma_{2N-3-k}^{(k)} - ix] &= \sum_{k=0}^{N-3} ([\gamma_{2N-3-k}^{(k)} - \gamma_N^{(N-3)}] + [\gamma_N^{(N-3)} - ix]) \\ &= \sum_{k=0}^{N-3} [\gamma_{2N-3-k}^{(k)} - \gamma_N^{(N-3)}] + (N-2)[\gamma_N^{(N-3)} - ix]. \end{aligned} \quad (\text{B.4})$$

Taking into account that the last term in the above equation is an even number one gets that $C_N(\gamma)$ is given by the expression (4.13). For the second diagram the analysis follows exactly the same lines.

Appendix C Gustafson's integral reduction

The extension of the first Gustafson integral [28, Theorem 5.1] to the complex case was obtained in [38]. It takes the form

$$\prod_{j=1}^N \sum_{n_j \in \mathbb{Z} + \frac{\sigma}{2}} \int_{-i\infty}^{i\infty} \frac{\prod_{m=1}^{N+1} \prod_{k=1}^N \Gamma(z_m - u_k) \Gamma(u_k + w_m)}{\prod_{m < j} \Gamma(u_m - u_j) \Gamma(u_j - u_m)} \prod_{p=1}^N \frac{dv_p}{2\pi i} = \frac{N! \prod_{k,j=1}^{N+1} \Gamma(z_k + w_j)}{\Gamma\left(\sum_{k=1}^{N+1} (z_k + w_k)\right)}, \quad (\text{C.1})$$

where Γ is the Gamma function of the complex field \mathbb{C} [43]

$$\Gamma(u) \equiv \Gamma(u, \bar{u}) = \frac{\Gamma(u)}{\Gamma(1 - \bar{u})} = \frac{1}{a(u)}. \quad (\text{C.2})$$

The variables u_k, w_m, z_m have the form

$$\begin{aligned} u_k &= \frac{n_k}{2} + \nu_k, & z_m &= \frac{n_m}{2} + x_m, & w_m &= \frac{\ell_m}{2} + y_m, \\ \bar{u}_k &= -\frac{n_k}{2} + \nu_k, & \bar{z}_m &= -\frac{n_m}{2} + x_m, & \bar{w}_m &= -\frac{\ell_m}{2} + y_m. \end{aligned}$$

and the integration contours over ν_k separate the series of poles associated with the Γ -functions: $\Gamma(z_m - u_k)$ and $\Gamma(u_k + w_m)$, see ref. [38] for more detail. The integral converges for $\sum_{m=1}^{N+1} \text{Re}(z_m + w_m) < 1$.

Let us put

$$\begin{aligned} z_{N+1} &= M \left(\frac{1}{2} + ix \right), & \bar{z}_{N+1} &= M \left(-\frac{1}{2} + ix \right), \\ w_{N+1} &= M' \left(\frac{1}{2} + ix' \right), & \bar{w}_{N+1} &= M' \left(-\frac{1}{2} + ix' \right) \end{aligned} \quad (\text{C.3})$$

and send $M, M' \rightarrow \infty$ keeping $M/M' = \xi$ fixed, so that $w_{N+1}/z_{N+1} \mapsto \zeta$ and $\bar{w}_{N+1}/\bar{z}_{N+1} \mapsto \bar{\zeta}$.

Dividing both sides of (C.1) by $(\Gamma(z_{N+1})\Gamma(w_{N+1}))^N$ we get in this limit

$$\frac{1}{N!} \prod_{j=1}^N \sum_{n_j \in \mathbb{Z} + \frac{\sigma}{2}} \int_{-i\infty}^{i\infty} [\zeta]^U \frac{\prod_{m,k=1}^N \Gamma(z_m - u_k) \Gamma(u_k + w_m)}{\prod_{m < j} \Gamma(u_m - u_j) \Gamma(u_j - u_m)} \prod_{p=1}^N \frac{d\nu_p}{2\pi i} = \frac{[\zeta]^Z}{[1 + \zeta]^{Z+W}} \prod_{k,j=1}^N \Gamma(z_k + w_j), \quad (\text{C.4})$$

where $|\arg \zeta| < \pi$, $Z = \sum_{k=1}^N z_k$, $W = \sum_{k=1}^N w_k$ and we recall that $[\zeta]^U \equiv \zeta^U \bar{\zeta}^{\bar{U}}$.

References

- [1] E. K. Sklyanin, L. A. Tahtadzjan, and L. D. Faddeev, “Quantum inverse problem method. I,” *Teoret. Mat. Fiz.*, vol. 40, no. 2, pp. 194–220, 1979.
- [2] E. K. Sklyanin, “The quantum Toda chain,” in *Nonlinear equations in classical and quantum field theory (Meudon/Paris, 1983/1984)*, vol. 226 of *Lecture Notes in Phys.*, pp. 196–233, Springer, Berlin, 1985.
- [3] E. K. Sklyanin, “Quantum inverse scattering method. Selected topics,” in *Quantum group and quantum integrable systems*, Nankai Lectures Math. Phys., pp. 63–97, World Sci. Publ., River Edge, NJ, 1992.
- [4] V. E. Korepin, “Calculation of norms of Bethe wave functions,” *Comm. Math. Phys.*, vol. 86, no. 3, pp. 391–418, 1982.
- [5] N. A. Slavnov, “Calculation of scalar products of wave functions and form-factors in the framework of the algebraic Bethe ansatz,” *Teoret. Mat. Fiz.*, vol. 79, no. 2, pp. 232–240, 1989.
- [6] A. G. Izergin and V. E. Korepin, “The quantum inverse scattering method approach to correlation functions,” *Comm. Math. Phys.*, vol. 94, no. 1, pp. 67–92, 1984.
- [7] N. Kitanine, J. M. Maillet, and V. Terras, “Correlation functions of the XXZ Heisenberg spin- $\frac{1}{2}$ chain in a magnetic field,” *Nuclear Phys. B*, vol. 567, no. 3, pp. 554–582, 2000.
- [8] M. C. Gutzwiller, “The quantum mechanical Toda lattice. II,” *Ann. Physics*, vol. 133, no. 2, pp. 304–331, 1981.

- [9] S. Kharchev and D. Lebedev, “Integral representation for the eigenfunctions of a quantum periodic Toda chain,” *Lett. Math. Phys.*, vol. 50, no. 1, pp. 53–77, 1999.
- [10] S. Kharchev and D. Lebedev, “Integral representations for the eigenfunctions of quantum open and periodic Toda chains from the QISM formalism,” vol. 34, pp. 2247–2258, 2001. Kowalevski Workshop on Mathematical Methods of Regular Dynamics (Leeds, 2000).
- [11] J. M. Maillet and G. Niccoli, “On quantum separation of variables,” *J. Math. Phys.*, vol. 59, no. 9, pp. 091417, 47, 2018.
- [12] J. M. Maillet and G. Niccoli, “On quantum separation of variables beyond fundamental representations,” *SciPost Phys.*, vol. 10, no. 2, pp. Paper No. 026, 38, 2021.
- [13] P. Ryan and D. Volin, “Separated variables and wave functions for rational $gl(N)$ spin chains in the companion twist frame,” *J. Math. Phys.*, vol. 60, no. 3, pp. 032701, 23, 2019.
- [14] N. Gromov, F. Levkovich-Maslyuk, and G. Sizov, “New construction of eigenstates and separation of variables for $SU(N)$ quantum spin chains,” *J. High Energy Phys.*, no. 9, pp. 111, front matter+39, 2017.
- [15] N. Gromov, F. Levkovich-Maslyuk, and P. Ryan, “Determinant form of correlators in high rank integrable spin chains via separation of variables,” *J. High Energy Phys.*, no. 5, pp. Paper No. 169, 79, 2021.
- [16] S. E. Derkachov, G. P. Korchemsky, and A. N. Manashov, “Noncompact Heisenberg spin magnets from high-energy QCD. I. Baxter Q -operator and separation of variables,” *Nuclear Phys. B*, vol. 617, no. 1-3, pp. 375–440, 2001.
- [17] S. E. Derkachov, G. P. Korchemsky, and A. N. Manashov, “Separation of variables for the quantum $SL(2, \mathbb{R})$ spin chain,” *J. High Energy Phys.*, no. 7, pp. 047, 30, 2003.
- [18] S. E. Derkachov, G. P. Korchemsky, and A. N. Manashov, “Baxter \mathbb{Q} -operator and separation of variables for the open $SL(2, \mathbb{R})$ spin chain,” *J. High Energy Phys.*, no. 10, pp. 053, 31, 2003.
- [19] A. G. Bytsko and J. Teschner, “Quantization of models with non-compact quantum group symmetry: modular XXZ magnet and lattice sinh-Gordon model,” *J. Phys. A*, vol. 39, no. 41, pp. 12927–12981, 2006.
- [20] A. V. Silant’ev, “Transition function for the Toda chain,” *Teoret. Mat. Fiz.*, vol. 150, no. 3, pp. 371–390, 2007.
- [21] S. E. Derkachov and A. N. Manashov, “Iterative construction of eigenfunctions of the monodromy matrix for $SL(2, \mathbb{C})$ magnet,” *J. Phys. A*, vol. 47, no. 30, pp. 305204, 25, 2014.
- [22] M. A. Semenov-Tian-Shansky, *Quantization of Open Toda Lattices*, pp. 226–259. Berlin, Heidelberg: Springer Berlin Heidelberg, 1994.
- [23] N. R. Wallach, *Real reductive groups. II*, vol. 132 of *Pure and Applied Mathematics*. Academic Press, Inc., Boston, MA, 1992.
- [24] K. K. Kozłowski, “Unitarity of the SoV transform for the Toda chain,” *Comm. Math. Phys.*, vol. 334, no. 1, pp. 223–273, 2015.

- [25] S. E. Derkachov, K. K. Kozlowski, and A. N. Manashov, “On the separation of variables for the modular XXZ magnet and the lattice sinh-Gordon models,” *Ann. Henri Poincaré*, vol. 20, no. 8, pp. 2623–2670, 2019.
- [26] S. E. Derkachov and A. N. Manashov, “Spin chains and Gustafson’s integrals,” *J. Phys. A*, vol. 50, no. 29, pp. 294006, 20, 2017.
- [27] R. A. Gustafson, “Some q -beta and Mellin-Barnes integrals with many parameters associated to the classical groups,” *SIAM J. Math. Anal.*, vol. 23, no. 2, pp. 525–551, 1992.
- [28] R. A. Gustafson, “Some q -beta and Mellin-Barnes integrals on compact Lie groups and Lie algebras,” vol. 341, no. 1, pp. 69–119, 1994.
- [29] S. E. Derkachov, K. K. Kozlowski, and A. N. Manashov, “Completeness of SoV representation for $SL(2, \mathbb{R})$ spin chains,” *SIGMA Symmetry Integrability Geom. Methods Appl.*, vol. 17, pp. Paper No. 063, 26, 2021.
- [30] L. N. Lipatov, “Asymptotic behavior of multicolor QCD at high energies in connection with exactly solvable spin models,” *JETP Lett.*, vol. 59, pp. 596–599, 1994.
- [31] L. N. Lipatov, “High-energy asymptotics of multicolor QCD and two-dimensional conformal field theories,” *Phys. Lett. B*, vol. 309, pp. 394–396, 1993.
- [32] L. D. Faddeev and G. P. Korchemsky, “High-energy QCD as a completely integrable model,” *Phys. Lett. B*, vol. 342, pp. 311–322, 1995.
- [33] L. N. Lipatov, “Integrability of scattering amplitudes in N=4 SUSY,” *J. Phys. A*, vol. 42, p. 304020, 2009.
- [34] J. Bartels, L. N. Lipatov, and A. Prygarin, “Integrable spin chains and scattering amplitudes,” *J. Phys. A*, vol. 44, p. 454013, 2011.
- [35] S. Derkachov, V. Kazakov, and E. Olivucci, “Basso-Dixon correlators in two-dimensional fishnet CFT,” *J. High Energy Phys.*, no. 4, pp. 032, 31, 2019.
- [36] S. Derkachov and E. Olivucci, “Exactly solvable magnet of conformal spins in four dimensions,” *Phys. Rev. Lett.*, vol. 125, no. 3, pp. 031603, 7, 2020.
- [37] S. Derkachov and E. Olivucci, “Exactly solvable single-trace four point correlators in χCFT_4 ,” *J. High Energy Phys.*, no. 2, pp. Paper No. 146, 84, 2021.
- [38] S. E. Derkachov and A. N. Manashov, “On complex gamma-function integrals,” *SIGMA Symmetry Integrability Geom. Methods Appl.*, vol. 16, pp. Paper No. 003, 20, 2020.
- [39] I. M. Gel’fand, M. I. Graev, and N. Y. Vilenkin, *Generalized functions. Vol. 5: Integral geometry and representation theory*. Translated from the Russian by Eugene Saletan, Academic Press, New York-London, 1966.
- [40] L. A. Tahtadžjan and L. D. Faddeev, “The quantum method for the inverse problem and the XYZ Heisenberg model,” *Uspekhi Mat. Nauk*, vol. 34, no. 5(209), pp. 13–63, 256, 1979.
- [41] P. P. Kulish and E. K. Sklyanin, “Quantum spectral transform method. Recent developments,” in *Integrable quantum field theories (Tvärminne, 1981)*, vol. 151 of *Lecture Notes in Phys.*, pp. 61–119, Springer, Berlin-New York, 1982.

- [42] L. D. Faddeev, “How the algebraic Bethe ansatz works for integrable models,” in *Symétries quantiques (Les Houches, 1995)*, pp. 149–219, North-Holland, Amsterdam, 1998.
- [43] I. M. Gel’fand, M. I. Graev, and V. S. Retakh, “Hypergeometric functions over an arbitrary field,” *Uspekhi Mat. Nauk*, vol. 59, no. 5(359), pp. 29–100, 2004.
- [44] S. E. Derkachov, “Baxter’s Q -operator for the homogeneous XXX spin chain,” *J. Phys. A*, vol. 32, no. 28, pp. 5299–5316, 1999.