

# SOME LINKS BETWEEN $F$ -HARMONICITY, SUBMERSION AND COHOMOLOGY

BANG-YEN CHEN AND SHIHSHU WALTER WEI

ABSTRACT. By studying cohomology classes that are related with  $p$ -harmonic morphisms,  $F$ -harmonic maps, and  $f$ -harmonic maps, we extend several of our previous results on Riemannian submersions and  $p$ -harmonic morphisms to  $F$ -harmonic maps, and  $f$ -harmonic maps which are submersions.

## 1. INTRODUCTION

We note in the study of  $p$ -harmonic maps between Riemannian manifolds that the following duality occurs between (A) conformal immersions and (B) horizontally conformal submersions:

(A) **Theorem** ([24]) *Let  $u : (M, g_M) \rightarrow (N, g_N)$  be a conformal immersion with  $\dim M = p$ . Then  $u$  is a  $p$ -harmonic map if and only if  $u(M)$  is a (parametric) minimal submanifold in  $N$ .*

(B) **Theorem** ([3]) *Let  $u : (M, g_M) \rightarrow (N, g_N)$  be a horizontally conformal submersion with  $\dim N = p$ . Then  $u$  is a  $p$ -harmonic map if and only if all the fibers of  $u$  are (parametric) minimal submanifolds in  $M$ .*

This duality between (A) and (B) links  $p$ -harmonic maps and (parametric) minimal submanifolds.

On the other hand, there also exists a duality between (C) (non-parametric) minimal hypersurface in the Euclidean space  $\mathbb{R}^{n+1}$  and (D) maximal spacelike (non-parametric) hypersurface in the Minkowski space  $\mathbb{R}^{n,1}$  equipped with the Lorentzian metric  $ds^2 = \sum_{i=1}^n (dx^i)^2 - dt^2$ , where

(C) The minimal hypersurface in the Euclidean space  $\mathbb{R}^{n+1}$  is given by the graph of a function  $u$  on a Euclidean domain satisfying the differential equation

$$(1.1) \quad \operatorname{div} \left( \frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right) = 0;$$

and

---

2000 *Mathematics Subject Classification.* Primary 58E20; Secondary 31B35, 53C40.

*Key words and phrases.*  $F$ -harmonic maps,  $p$ -harmonic morphism, Cohomology class, minimal submanifold, submersion.

(D) The maximal spacelike hypersurface in the Minkowski space  $\mathbb{R}^{n,1}$  is furnished by the graph of a function  $v$  on a Euclidean domain satisfying a dual differential equation

$$(1.2) \quad \operatorname{div} \left( \frac{\nabla v}{\sqrt{1 - |\nabla v|^2}} \right) = 0, \quad |\nabla v|^2 < 1.$$

E. Calabi showed in [5] that equations (1.1) and (1.2) are equivalent over any simply-connected domain in  $\mathbb{R}^2$ . Furthermore, the notion of  $F$ -harmonic maps unifies minimal hypersurfaces and maximal spacelike hypersurfaces; namely, the solutions  $u$  and  $v$  are  $F$ -harmonic maps from a domain in  $\mathbb{R}^{n+1}$  to  $\mathbb{R}$  with  $F(t) = \sqrt{1+2t}-1$  and  $F(t) = 1-\sqrt{1-2t}$ , respectively (see [27, 12]). Hence, one can unify harmonic maps,  $p$ -harmonic maps,  $\alpha$ -harmonic maps, exponential harmonic maps, (nonparametric) minimal hypersurfaces in Euclidean space, and (nonparametric) maximal spacelike hypersurface in Minkowski space  $\mathbb{R}^{n,1}$  by  $F$ -harmonic maps (see [1, 27]). Also, the “ $F$ ”-idea can be extended to Gauge Theory (see [12]). Here

$$(1.3) \quad F : [0, \infty) \rightarrow [0, \infty) \quad \text{with} \quad F(0) = 0 \quad \text{is } C^2 \text{ strictly increasing.}$$

We recall in the study of topology on a compact Riemannian manifold  $M$ , it is well-known that nontrivial fundamental groups  $\pi_1(M)$ , homology groups, and cohomology classes can be represented by stable closed geodesics, stable minimal rectifiable currents, and harmonic forms on  $M$ , respectively, by Cartan’s Theorem ([6]), Federer-Fleming’s Theorem ([14]), and Hodge Theorem ([15]). In an analogous spirit, it was shown in [26] that homotopy classes can be represented by  $p$ -harmonic maps (For definition and examples of  $p$ -harmonic maps, see e.g. [27]):

**Theorem A.** *If  $N$  is a compact Riemannian manifold, then for any positive integer  $i$ , each class in the  $i$ -th homotopy group  $\pi_i(N)$  can be represented by a  $C^{1,\alpha}$   $p$ -harmonic map  $u_0$  from an  $i$ -dimensional sphere  $S^i$  into  $N$  minimizing  $p$ -energy in its homotopy class for any  $p > i$ .*

Further applications and homotopically vanishing theorems were explored in [26, 28], using stable  $p$ -harmonic maps as catalysts, whereas homologically vanishing theorem was given in [19, 25, 16], using stable rectifiable currents as catalysts.

On the other hand, B.-Y. Chen established in [8] the following result involving Riemannian submersion, minimal immersion, and cohomology class.

**Theorem B.** ([8]) *Let  $b = \dim B$  and let  $\pi : (M, g_M) \rightarrow (B, g_B)$  be a Riemannian submersion with minimal fibers and orientable base manifold  $B$ . If  $M$  is a closed manifold with cohomology class  $H^b(M, \mathbf{R}) = 0$ , then the horizontal distribution  $\mathcal{H}$  of the Riemannian submersion is never integrable. Thus the submersion  $\pi$  is never non-trivial.*

Whereas  $p$ -harmonic maps represent homotopy classes, B.-Y. Chen and S. W. Wei connected the two seemingly unrelated areas of  $p$ -harmonic morphisms and cohomology classes in the following.

**Theorem C.** ([10, 11]) *Let  $u : (M, g_M) \rightarrow (N, g_N)$  be an  $n$ -harmonic morphism which is a submersion with  $\dim N = n$ . If  $N$  is orientable and  $M$  is a closed manifold with  $n$ -th cohomology class  $H^n(M, \mathbf{R}) = 0$ , then the horizontal distribution  $\mathcal{H}$  of  $u$  is never integrable. Hence the submersion  $u$  is always non-trivial.*

This recaptures Theorem B when  $\pi : M \rightarrow B$  is a Riemannian submersion with minimal fibers and orientable base manifold  $B$ . While a horizontally weak conformal  $p$ -harmonic map is a  $p$ -harmonic morphism (see Theorem 2.5),  $p$ -harmonic morphism is also linked to cohomology class as follows.

**Theorem D.** ([10, 11]) *Let  $u : (M, g_M) \rightarrow (N, g_N)$  be an  $n$ -harmonic morphism with  $n = \dim N$  which is a submersion. Then the pull back of the volume element of  $N$  is a harmonic  $n$ -form if and only if the horizontal distribution  $\mathcal{H}$  of  $u$  is completely integrable.*

Following the proofs given in [10, 11], and by applying a characterization theorem of a  $p$ -harmonic morphism (see Theorem 2.5) and Theorem 2.6, we obtain a dual version of Theorem D. In particular, we have  $p$ -harmonic maps and cohomology classes are interrelated in [27] as follows.

**Theorem E.** *Let  $M$  be a closed manifold, and  $u : M \rightarrow N$  be an  $n$ -harmonic map with  $n = \dim N$  which is a submersion. Assume that the horizontal distribution  $\mathcal{H}$  of  $u$  is integrable and  $u$  is an  $n$ -harmonic morphism. Then we have  $H^n(M, \mathbf{R}) \neq 0$ .*

**Theorem F.** ([27]) *Let  $u : M \rightarrow N$  be an  $n$ -harmonic map with  $n = \dim N$  which is a submersion. Let the horizontal distribution  $\mathcal{H}$  of  $u$  be integrable. If  $M$  is a closed manifold with cohomology class  $H^n(M, \mathbf{R}) = 0$ . Then  $u$  is not an  $n$ -harmonic morphism. Thus the submersion  $u$  is always nontrivial.*

## 2. PRELIMINARIES

We recall some basic facts, notations, definitions, and formulas for minimal submanifolds, submersions,  $p$ -harmonic morphisms,  $F$ -harmonic maps, and  $f$ -harmonic maps for later use (see, e.g., [7, 8, 9, 10] for details).

**2.1. Basic formulas and equations.** Let  $\widetilde{M}$  be a Riemannian manifold with Levi-Civita connection  $\widetilde{\nabla}$ . The tangent bundle of  $\widetilde{M}$  is denoted by  $T\widetilde{M}$ , and the (infinite dimensional) vector space of smooth sections of a smooth vector bundle  $E$  is denoted by  $\Gamma(E)$ . Let  $M$  be a submanifold of dimension  $n \geq 2$  in  $\widetilde{M}$ . Denote by  $\nabla$  and  $D$ , the Levi-Civita connection and the normal connection of  $M$ , respectively. For each normal vector  $\xi \in T_x^\perp M$ ,  $x \in M$ , the shape operator  $A_\xi$  is a symmetric

endomorphism of the tangent space  $T_x M$  at  $x$ . Then the shape operator and the second fundamental form  $\mathbf{h}$  are related by

$$(2.1) \quad \langle \mathbf{h}(X, Y), \xi \rangle = \langle A_\xi X, Y \rangle$$

for  $X, Y$  tangent to  $M$  and  $\xi$  normal to  $M$ .

The formulas of Gauss and Weingarten are given respectively by (cf. [7])

$$(2.2) \quad \tilde{\nabla}_X Y = \nabla_X Y + \mathbf{h}(X, Y),$$

$$(2.3) \quad \tilde{\nabla}_X \xi = -A_\xi X + D_X \xi$$

for tangent vector fields  $X, Y$  and normal vector field  $\xi$  on  $M$ .

The *mean curvature vector field* of a submanifold  $M$  is defined by  $H = \frac{1}{n} \text{trace } \mathbf{h}$ . A submanifold  $M$  in  $\tilde{M}$  is called *totally geodesic* (respectively, *minimal*) if its second fundamental form  $\mathbf{h}$  (respectively, its mean curvature vector field  $H$ ) vanishes identically.

**2.2. Energy and harmonic maps.** Let  $u : (M, g_M) \rightarrow (N, g_N)$  be a differential map between two Riemannian manifolds  $M$  and  $N$ . Denote  $e_u$  the *energy density* of  $u$ , which is given by

$$(2.4) \quad e_u = \frac{1}{2} \sum_{i=1}^m g_N(du(e_i), du(e_i)) = \frac{1}{2} |du|^2,$$

where  $\{e_1, \dots, e_n\}$  is a local orthonormal frame field on  $M$  and  $|du|$  is the Hilbert-Schmidt norm of  $du$ , determined by the metric  $g_M$  of  $M$  and the metric  $g_N$  of  $N$ . The *energy of  $u$* , denoted by  $E(u)$ , is defined to be

$$E(u) = \int_M e_u dv_g.$$

A smooth map  $u : M \rightarrow N$  is called *harmonic* if  $u$  is a critical point of the energy functional  $E$  with respect to any compactly supported variation.

**2.3. Submersions.** A differential map  $u : (M, g_M) \rightarrow (N, g_N)$  between two Riemannian manifolds is called a submersion at a point  $x \in M$  if its differential  $du_x : T_x(M) \rightarrow T_{u(x)}(N)$  is a surjective linear map. A differentiable map  $u$  that is a submersion at each point  $x \in M$  is called a *submersion*. For each point  $x \in N$ ,  $u^{-1}(x)$  is called a *fiber*. For a submersion  $u : M \rightarrow N$ , let  $\mathcal{H}_x$  denote the orthogonal complement of Kernel( $du_x : T_x(M) \rightarrow T_{u(x)}(N)$ ) in  $T_x(M)$  at  $x \in M$ . Let  $\mathcal{H} = \{\mathcal{H}_x : x \in M\}$  denote the horizontal distribution of  $u$ . A submersion  $u : (M, g_M) \rightarrow (N, g_N)$  is called *horizontally weakly conformal* if the restriction of  $du_x$  to  $\mathcal{H}_x$  is conformal, i.e., there exists a smooth function  $\lambda$  on  $M$  such that

$$(2.5) \quad u^* g_N = \lambda^2 g_M|_{\mathcal{H}} \quad \text{or} \quad g_N(du_x(X), du_x(Y)) = \lambda^2(x) g_M(X, Y)$$

for all  $X, Y \in \mathcal{H}_x$  and  $x \in M$ . If the function  $\lambda$  in (2.5) is positive, then  $u$  is called *horizontally conformal* and  $\lambda$  is called the *dilation* of  $u$ . For a horizontally conformal

submersion  $u$  with dilation  $\lambda$ , the *energy density* is  $e_u = \frac{1}{2}n\lambda^2$ . A horizontally conformal submersion with dilation  $\lambda \equiv 1$  is called a *Riemannian submersion*. For Riemannian submersions, we have the following results.

**Theorem 2.1** ([8], Theorem 2). *Let  $\pi : (M, g_M) \rightarrow (B, g_B)$  be a Riemannian submersion from a closed manifold  $M$  onto an orientable base manifold  $B$ . Then the pullback of the volume element of  $B$  is harmonic if and only if the horizontal distribution  $\mathcal{H}$  is integrable and fibers are minimal.*

**Theorem 2.2** ([27], Theorem 2.5). *Let  $\pi : (M, g_M) \rightarrow (B, g_B)$  be a Riemannian submersion. Then  $\pi$  is a  $p$ -harmonic map for every  $p > 1$  if and only if all fibers  $\pi^{-1}(y), y \in B$ , are minimal submanifolds in  $M$ .*

**2.4. Harmonic morphisms.** A  $C^2$ -map  $u : (M, g_M) \rightarrow (N, g_N)$  is said to be a *harmonic morphism* if for any harmonic function  $f$  defined on an open set  $V$  of  $N$ , the composition  $f \circ u$  is harmonic on  $u^{-1}(V)$ .

P. Baird and J. Eells used the stress-energy tensor to establish the following.

**Theorem 2.3** ([2]). *Let  $u : (M, g_M) \rightarrow (N, g_N)$  be a harmonic morphism which is a submersion everywhere on  $M$ . Then (setting  $n = \dim N$ , and  $e_u$  as in (2.4))*

- (a) *if  $n = 2$ , the fibres are minimal submanifolds ;*
- (b) *if  $n > 2$ , then the following properties are equivalent :*
  - (i) *the fibres are minimal submanifolds;*
  - (ii)  *$\text{grad}(e_u)$  is a vertical field;*
  - (iii) *the horizontal distribution has mean curvature vector  $\frac{\text{grad}(e_u)}{2e_u}$ .*

**2.5.  $p$ -harmonic morphisms.** A  $C^2$ -map  $u : (M, g_M) \rightarrow (N, g_N)$  is said to be a  *$p$ -harmonic morphism* if for any  $p$ -harmonic function  $f$  defined on an open set  $V$  of  $N$ , the composition  $f \circ u$  is  $p$ -harmonic on  $u^{-1}(V)$ . We have the following link between  $p$ -harmonic morphisms for every  $p > 1$  and minimal fibers:

**Theorem 2.4** ([27], Proposition 2.4). *If  $\pi : (M, g_M) \rightarrow (N, g_N)$  is a Riemannian submersion, then  $\pi$  is a  $p$ -harmonic morphism for every  $p > 1$  if and only if all fibers  $\pi^{-1}(y), y \in N$ , are minimal submanifolds of  $M$ .*

E. Loubeau [18] and J. M. Burel and E. Loubeau [4] obtained the following characterization of  $p$ -harmonic morphisms.

**Theorem 2.5** ([18, 4]). *A  $C^2$ -map  $u : (M, g_M) \rightarrow (N, g_N)$  is a  $p$ -harmonic morphism with  $p \in (1, \infty)$  if and only if it is a  $p$ -harmonic, horizontally conformal map.*

Theorem 2.4 ([27], Proposition 2.4) links  $p$ -harmonic maps for every  $p > 1$  with minimal fibers in the presence of Riemannian submersion. This is dual to a minimal submanifold occurs from a  $p$ -harmonic map for every  $p > 1$  in the presence of

isometric immersion (cf. [26]). In [3], P. Baird and S. Gudmundsson linked  $n$ -harmonic morphisms with minimal fibers as follows.

**Theorem 2.6.** ([3]) *If  $u : (M, g_M) \rightarrow (N, g_N)$  is a horizontally conformal submersion from a manifold  $M$  onto a manifold  $N$  with  $n = \dim N$ , then  $u$  is  $n$ -harmonic if and only if the fibers of  $u$  are minimal in  $M$ .*

**2.6.  $F$ -harmonic maps.** Let  $F$  be as in (1.3) and let  $u : (M, g_M) \rightarrow (N, g_N)$  be a smooth map between two compact Riemannian manifolds. Then the map  $u : M \rightarrow N$  is called  $F$ -harmonic if it is a critical point of the  $F$ -energy functional:  $E_F(u) = \int_M F\left(\frac{|du|^2}{2}\right) dv_g$ . In particular, when

$$F(t) = t, \quad \frac{1}{p}(2t)^{\frac{p}{2}}, \quad (1+2t)^\alpha - 1 \quad (\alpha > 1, \dim M = 2),$$

$$e^t - 1, \quad \sqrt{1+2t} - 1, \quad 1 - \sqrt{1-2t}, \quad \text{respectively,}$$

the  $F$ -energy  $E_F(u)$  becomes energy,  $p$ -energy, (normalized)  $\alpha$ -energy, (normalized) exponential energy, (normalized) area functional in Euclidean space, and (normalized) area functional in Minkowski space. Hence, its critical point  $u$  or its graph is harmonic,  $p$ -harmonic,  $\alpha$ -harmonic, exponential harmonic, minimal hypersurface in Euclidean space  $\mathbb{R}^{n+1}$ , maximal spacelike hypersurface in Minkowski space  $\mathbb{R}_1^{n+1}$ , respectively.

For horizontally conformal  $F$ -harmonic maps, M. Ara proved the following.

**Theorem 2.7** (cf. [1]). *Let  $u : (M, g_M) \rightarrow (N, g_N)$ ,  $m > n$ , be an  $F$ -harmonic map, which is horizontally conformal with dilation  $\lambda$ . Assume that the zeros of  $(n-2)F'(t) - 2tF''(t)$  are isolated. Then the following three properties are equivalent:*

- (1) *The fibers of  $u$  are minimal submanifolds.*
- (2)  *$\text{grad}(\lambda^2)$  is vertical.*
- (3) *The horizontally distribution of  $u$  has mean curvature vector  $\frac{\text{grad}(\lambda^2)}{2\lambda^2}$ .*

**2.7.  $f$ -harmonic maps.** Let  $f : (M, g) \rightarrow (0, \infty)$  be a smooth function. The notion of  $f$ -harmonic maps was first introduced and studied by A. Lichnerowicz in 1970 (see [17]). A map  $u : (M, g_M) \rightarrow (N, g_N)$  between two Riemannian manifolds is said to be  $f$ -harmonic, if  $u$  is a critical point of the  $f$ -energy functional  $E_f$  with respect to any compactly supported variation (cf. [13, 17]), where

$$(2.6) \quad E_f(u) = \frac{1}{2} \int_M f |du|^2 dv_g.$$

Examples of  $f$ -harmonic maps include harmonic maps with  $f \equiv C$ , a positive constant, and *submersive  $p$ -harmonic maps* with  $f = |du|^{p-2}$ .

The next result is due to Y.-L. Ou.

**Theorem 2.8** ([21]). *Let  $u : (M, g_M) \rightarrow (N, g_N)$  be a horizontally weakly conformal map with dilation  $\lambda$ . Then, any two of the following conditions imply the other one:*

- (a)  $u$  is an  $f$ -harmonic map;
- (b)  $\text{grad}(f\lambda^{2-n})$  is vertical;
- (c)  $u$  has minimal fibers.

### 3. STATEMENTS OF THEOREMS

In the following we shall assume that  $\dim M = m > n = \dim N$ . The purpose of this article is to extend the ideas in [10, 27] and to connect the seemingly unrelated areas of  $F$ -harmonic maps,  $f$ -harmonic maps, and cohomology classes. More precisely we prove the following

**Theorem 1.** *Let  $u : M \rightarrow N$  be an  $F$ -harmonic map which is a horizontally conformal submersion with dilation  $\lambda$ . And let  $\text{grad } \lambda^2$  be vertical, and the zeros of  $(n-2)F'(t) - 2tF''(t)$  be isolated. If  $H^n(M, \mathbf{R}) = 0$ , then the horizontal distribution  $\mathcal{H}$  of  $u$  is not integrable. Thus the submersion is always non-trivial.*

*Remark 3.1.* (i) Theorem 1 generalizes Theorem B, in which by assumption  $u$  is a horizontal conformal submersion with dilation  $\lambda = 1$ , and  $u$  is a  $p$ -harmonic map for every  $p > 1$  according to Theorem 2.4. If we choose  $p \neq n$ , then  $\text{grad } \lambda^2 = 0$ ,  $F(t) = \frac{1}{p}(2t)^{\frac{p}{2}}$  so that  $F$ -harmonic map becomes  $p$ -harmonic map,  $p \neq n$ , and the zeros of  $(n-2)F'(t) - 2tF''(t)$  are isolated.

(ii) Theorem 1 augments Theorem C but does not generalize Theorem C, since when  $F(t) = \frac{1}{n}(2t)^{\frac{n}{2}}$ ,  $F$ -harmonic map  $u$  becomes  $n$ -harmonic map, and the zeros of  $(n-2)F'(t) - 2tF''(t)$  are not isolated. Thus  $n$ -harmonic maps are not applicable to Theorem 1.

(iii) Theorem 1 is equivalent to the following Theorem 2, due to the equivalence in Theorem 2.7.

**Theorem 2.** *Let  $u : M \rightarrow N$  be an  $F$ -harmonic map which is a horizontally conformal submersion with dilation  $\lambda$ . Assume that the horizontal distribution has mean curvature vector  $\frac{\text{grad}(\lambda^2)}{2\lambda^2}$ , and the zeros of  $(n-2)F'(t) - 2tF''(t)$  are isolated. If  $H^n(M, \mathbf{R}) = 0$ , then the horizontal distribution  $\mathcal{H}$  of  $u$  is not integrable. Thus the submersion is always non-trivial.*

**Theorem 3.** *Let  $u : M \rightarrow N$  be an  $F$ -harmonic map which is a horizontally conformal submersion with dilation  $\lambda$ . Assume that the horizontal distribution has mean curvature vector  $\frac{\text{grad}(\lambda^2)}{2\lambda^2}$ , and the zeros of  $(n-2)F'(t) - 2tF''(t)$  are isolated. Then the pullback of the volume element of  $N$  is a harmonic  $n$ -form if and only if the horizontal distribution  $\mathcal{H}$  of  $u$  is completely integrable.*

*Remark 3.2.* By similar arguments as in Remark 3.1, (i) Theorem 2 generalizes Theorem 2.1; (ii) Theorem 2 augments Theorem D but does not generalize Theorem D; and (iii) Theorem 2 is equivalent to the following Theorem 4.

**Theorem 4.** *Let  $u : M \rightarrow N$  be an  $F$ -harmonic map which is a horizontally conformal submersion with dilation  $\lambda$ . Assume that  $\text{grad } \lambda^2$  be vertical, and the zeros of  $(n-2)F'(t) - 2tF''(t)$  are isolated. Then the pullback of the volume element of  $N$  is a harmonic  $n$ -form if and only if the horizontal distribution  $\mathcal{H}$  of  $u$  is completely integrable.*

We then study  $f$ -harmonic maps which was first introduced and studied by A. Lichnerowicz in 1970 (see [17]), where  $f : (M, g) \rightarrow (0, \infty)$  is a smooth function.. Examples of  $f$ -harmonic maps include harmonic maps with  $f \equiv C$ , a positive constant, and *submersive  $p$ -harmonic maps* with  $f = |du|^{p-2}$  (cf. §4). We have the following results.

**Theorem 5.** *Let  $u : M \rightarrow N$  be an  $f$ -harmonic map which is a horizontally conformal submersion with dilation  $\lambda$  such that  $\text{grad}(f\lambda^{2-n})$  is vertical. If  $M$  is a closed manifold with cohomology class  $H^n(M, \mathbf{R}) = 0$ , then the horizontal distribution  $\mathcal{H}$  of  $u$  is never integrable. Thus the submersion  $u$  is always non-trivial.*

This recaptures Theorem B, when  $f$ -harmonic map is an  $n$ -harmonic map, then  $f = |du|^{n-2}$ .

**Theorem 6.** *Let  $u : M \rightarrow N$  be an  $f$ -harmonic map which is a horizontally conformal submersion and with dilation  $\lambda$  such that  $\text{grad}(f\lambda^{2-n})$  is vertical. Then the pullback of the volume element of  $N$  is a harmonic  $n$ -form if and only if horizontal distribution is completely integrable.*

This recaptures Theorems D, when  $f$ -harmonic map is an  $n$ -harmonic map.

#### 4. PROOFS OF THEOREMS 1-6

*Proof. Proof of Theorem 3:* Let  $u : (M, g_M) \rightarrow (N, g_N)$  be an  $F$ -harmonic map which is a horizontally conformal submersion with dilation  $\lambda$  and  $n = \dim N$ . Assume that the horizontal distribution has mean curvature vector  $\frac{\text{grad}(\lambda^2)}{2\lambda^2}$ , and the zeros of  $(n-2)F'(t) - 2tF''(t)$  are isolated. Then the submersion  $u$  has minimal fibers according to Theorem 2.7.

Let  $\{\bar{e}_1, \dots, \bar{e}_n\}$  be an oriented local orthonormal frame of the base manifold  $(N, g_N)$  and let  $\bar{\omega}^1, \dots, \bar{\omega}^n$  denote the dual 1-forms of  $\{\bar{e}_1, \dots, \bar{e}_n\}$  on  $N$ . Then

$$\bar{\omega} = \bar{\omega}^1 \wedge \dots \wedge \bar{\omega}^n$$

is the volume form of  $(N, g_N)$ , which is a closed  $n$ -form on  $N$ .

Consider the pull back of the volume form  $\bar{\omega}$  of  $N$  via  $u$ , which is denoted by  $u^*(\bar{\omega})$ . Then  $u^*(\bar{\omega})$  is a simple  $n$ -form on  $M$  satisfying

$$(4.1) \quad d(u^*(\bar{\omega})) = u^*(d\bar{\omega}) = 0,$$

since the exterior differentiation  $d$  and the pullback  $u^*$  commute.

Assume that  $\dim M = m = n + k$  and let  $e_1, \dots, e_{n+k}$  be a local orthonormal frame field with  $\omega^1, \dots, \omega^{n+k}$  being its dual coframe fields on  $M$  such that



- (i)  $e_1, \dots, e_n$  are basic horizontal vector fields satisfying  $du(e_i) = \lambda \bar{e}_i$ ,  $i = 1, \dots, n$ , and  $du(e_1), \dots, du(e_n)$  give a positive orientation of  $N$ ; and  
 (ii)  $e_{n+1}, \dots, e_{n+k}$  are vertical vector fields.

Then we have

$$(4.2) \quad \omega^j(e_s) = 0, \quad \omega^i(e_j) = \delta_{ij}, \quad 1 \leq i, j \leq n; \quad n+1 \leq s \leq n+k.$$

Also, it follows from (i) that

$$(4.3) \quad u^* \bar{\omega}^i = \frac{1}{\lambda} \omega^i, \quad i = 1, \dots, n.$$

If we put

$$(4.4) \quad \omega = \omega^1 \wedge \dots \wedge \omega^n \quad \text{and} \quad \omega^\perp = \omega^{n+1} \wedge \dots \wedge \omega^{n+k},$$

then

$$(4.5) \quad d\omega^\perp = \sum_{i=1}^k (-1)^i \omega^{n+1} \wedge \dots \wedge d\omega^{n+i} \wedge \dots \wedge \omega^{n+k}.$$

It follows from (4.2) and (4.5) that  $d\omega^\perp = 0$  holds identically if and only if the following two conditions are satisfied:

$$(4.6) \quad d\omega^\perp(e_i, e_{n+1}, \dots, e_{n+k}) = 0$$

for  $i = 1, \dots, n$ ; and

$$(4.7) \quad d\omega^\perp(X, Y, V_1, \dots, V_{k-1}) = 0.$$

for any horizontal vector fields  $X, Y$  and vertical vector fields  $V_1, \dots, V_{k-1}$ .

Since the fibers of  $u$  are minimal in  $M$ , for each  $1 \leq i \leq n$ , we find

$$(4.8) \quad \begin{aligned} & d\omega^\perp(e_i, e_{n+1}, \dots, e_{n+k}) \\ &= \sum_{j=1}^k (-1)^{j+1} \omega^\perp([e_i, e_{n+j}], e_{n+1}, \dots, \hat{e}_{n+j}, \dots, e_{n+k}) \\ &= \sum_{j=1}^k (-1)^{j+1} (\omega^{n+j}(\nabla_{e_i} e_{n+j}) - \omega^{n+j}(\nabla_{e_{n+j}} e_i)) \\ &= \sum_{j=1}^k -\langle \nabla_{e_{n+j}} e_i, e_{n+j} \rangle \\ &= \sum_{j=1}^k \langle h(e_{n+j}, e_{n+j}), e_i \rangle \\ &= 0, \end{aligned}$$

where “ $\hat{\cdot}$ ” denotes the missing term, which proves that condition (4.6) holds.

Next, we suppose that the horizontal distribution  $\mathcal{H}$  is integrable. If  $X, Y$  are horizontal vector fields, then  $[X, Y]$  is also horizontal by Frobenius theorem. So,

for vertical vector fields  $V_1, \dots, V_k$ , we find (cf. [8, formula (6.7)] or [27, formula (3.5)])

$$(4.9) \quad d\omega^\perp(X, Y, V_1, \dots, V_{k-1}) = \omega^\perp([X, Y], V_1, \dots, V_{k-1}) = 0.$$

Consequently, from (4.8) and (4.9) we get

$$(4.10) \quad d\omega^\perp = 0.$$

Next, we show that if  $\mathcal{H}$  is integrable, then we have  $d(u^*\bar{\omega})^\perp = 0$ . Since  $u$  is a horizontally conformal submersion with dilation  $\lambda$ , it preserves orthogonality, which is crucial to horizontal and vertical distributions, and the pullback  $u^*$  expands the length of 1-form by  $\frac{1}{\lambda}$  in every direction. This, via (4.3) and (4.10), leads to

$$(4.11) \quad \begin{aligned} d(u^*\bar{\omega})^\perp &= d(u^*\omega^1 \wedge \dots \wedge u^*\omega^n)^\perp \\ &= d\left(\frac{1}{\lambda}\omega^1 \wedge \dots \wedge \frac{1}{\lambda}\omega^n\right)^\perp \\ &= d\left(\frac{1}{\lambda}\omega^{n+1} \wedge \dots \wedge \frac{1}{\lambda}\omega^{n+k}\right)^\perp \\ &= d\left(\frac{1}{\lambda^k}\omega^\perp\right)^\perp \\ &= -\frac{k}{2}\lambda^{-k-2}d(\lambda^2) \wedge \omega^\perp + \frac{1}{\lambda^k}d\omega^\perp \\ &= -\frac{k}{2}\lambda^{-k-2}d(\lambda^2) \wedge \omega^\perp \\ &= 0. \end{aligned}$$

The last equality of (4.11) is due to the fact that  $\text{grad}(\lambda^2)$  is vertical and Theorem 2.7. Therefore,  $\text{grad}(\lambda^2), e_{n+1}, \dots, e_{n+k}$  are linearly dependent. Hence, we obtain  $d(\lambda^2) \wedge \omega^\perp = 0$ . Because  $d(u^*\omega)^\perp = 0$  is equivalent to  $u^*\omega$  being co-closed, it follows from  $d(u^*\omega) = 0$  and  $d(u^*\omega)^\perp = 0$  that the pullback of the volume form,  $u^*\omega$ , is a harmonic  $n$ -form on  $M$  whenever  $\mathcal{H}$  is completely integrable.

Conversely, it follows from the proof given above that if  $u^*\omega$  is a harmonic  $n$ -form on  $M$ , then the horizontal distribution  $\mathcal{H}$  is completely integrable. This proves Theorem 3.

**Proof of Theorem 1:** Since each nonzero harmonic form represents a nontrivial cohomology class by Hodge theorem ([15]), and since on a closed manifold a differential form is harmonic if and only if it is closed and co-closed, Theorem 1 follows from Theorem 3.

**Proof of Theorem 2 and Theorem 4:** Theorem 2 and Theorem 4 follow from Theorem 1 together with Remarks 3.1 and 3.2 respectively.

**Proof of Theorem 5 and Theorem 6:** By virtue of Theorem 2.8, the submersion  $u$  in Theorem 5 and Theorem 6 has minimal fibers, we proceed as before,  $u$  is an  $n$ -harmonic map by virtue of Theorem 2.6, and hence  $u$  is an  $n$ -harmonic morphism

according to Theorem 2.5. We are then ready to apply Theorem C and Theorem D to complete the proof of Theorem 5 and Theorem 6 , respectively.  $\square$

## REFERENCES

- [1] M. Ara, *Geometry of  $F$ -harmonic maps*. Kodai Math. J. **22**(2) (1999), 243–263.
- [2] P. Baird and J. Eells, *A conservation law for harmonic maps*. Geometry Symposium, Utrecht 1980 (Utrecht, 1980), 1–25, Lecture Notes in Math. **894** Springer, Berlin-New York, 1981.
- [3] P. Baird and S. Gudmundsson,  *$p$ -harmonic maps and minimal submanifolds*. Math. Ann. **294**(4) (1992), 611–624.
- [4] J. M. Burel and E. Loubeau,  *$p$ -harmonic morphisms: the  $1 < p < 2$  case and a non-trivial example*. Contemp. Math. **308** (2002), 21–37.
- [5] E. Calabi, *Examples of Bernstein problems for some nonlinear equations*. Global Analysis, Proc. Sympos. Pure Math. **15** (1970), 223–230.
- [6] E. Cartan, *Lecons sur la Géométrie des Espaces de Riemann*, 2nd edition. Gauthier-Villars, Paris, 1951.
- [7] B.-Y. Chen, *Geometry of Submanifolds*. Marcer Dekker, New York, NY, 1973.
- [8] B.-Y. Chen, *Riemannian submersions, minimal immersions and cohomology class*. Proc. Japan Acad. Ser. A Math. Sci. **81**(10) (2005), 162–167.
- [9] B.-Y. Chen, *Pseudo-Riemannian Geometry,  $\delta$ -invariants and Applications*. World Scientific, Hackensack, NJ, 2011.
- [10] B.-Y. Chen and S. W. Wei,  *$p$ -harmonic morphisms, cohomology classes and submersions*. Tamkang J. Math. **40**(4) (2009), 377–382.
- [11] B.-Y. Chen and S. W. Wei,  *$n$ -Harmonicity, minimality, conformality and cohomology*, <https://doi.org/10.48550/arXiv.2302.14019> (2023).
- [12] Y. X. Dong and S. W. Wei, *On vanishing theorems for vector bundle valued  $p$ -forms and their applications*. Comm. Math. Phys. **304** (2011), 329–368.
- [13] J. Eells and L. Lemaire, *A report on harmonic maps*. Bull. London Math. Soc. **10** (1978), 1–68.
- [14] H. Federer and W. H. Fleming, *Normal and integral currents*. Ann. Math. **72** (1960), 458–520.
- [15] W.V.D. Hodge, *The theory and application of harmonic intergrals*. Cambridge University Press, New York, NY, 1941 (2nd ed. 1952)
- [16] R. Howard and S. W. Wei, *On the existence and nonexistence of stable submanifolds and currents in positively curved manifolds and the topology of submanifolds in Euclidean spaces*. Geometry and Topology of Submanifolds and Currents. 127–167, Contemp. Math. **646** Amer. Math. Soc., Providence, RI, 2015.
- [17] A. Lichnerowicz, *Applications harmoniques et variétés Kähleriennes*. Rend. Sem. Mat. Fis. Milano **39** (1969), 186–195.
- [18] E. Loubeau, *On  $p$ -harmonic morphisms*. Differential Geom. Appl. **12** (2000), 219–229.
- [19] H. B. Lawson and J. Simons, *On stable currents and their application to global problems in real and complex geometry*. Ann. of Math. **98** (1973), 427–450.
- [20] B. O’Neill, *Semi-Riemannian Geometry with Applications to Relativity*. Academic Press, New York, NY, 1982.
- [21] Y.-L. Ou,  *$f$ -harmonic morphisms between Riemannian manifolds*. Chinese Ann. Math. Ser. B **35**(2) (2014), 225–236.
- [22] A. V. Pogorelov, *On the stability of minimal surfaces*. Dokl. Akad. Nauk SSSR **260**, 293–295 (1981, Zbl. 495.53005), English transl.: Soviet Math. Dokl. **24**, (1981), 274–276.
- [23] S.-I. Tachibana, *On harmonic simple forms*, Tensor (N.S.) **27** (1973), 123–130.

- [24] H. Takeuchi, *Some conformal properties of  $p$ -harmonic maps and a regularity for sphere-valued  $p$ -harmonic maps*. J. Math. Soc. Japan **46**(2) (1994), 217–234.
- [25] S. W. Wei, *On topological vanishing theorems and the stability of Yang-Mills fields*. Indiana Univ. Math. J. **33**(4) (1984), 511–529.
- [26] S. W. Wei, *Representing Homotopy Groups and Spaces of Maps by  $p$ -harmonic maps*. Indiana Univ. Math. J. **47**(2) (1998), 625–670.
- [27] S. W. Wei, *The unity of  $p$ -harmonic geometry*. Recent developments in geometry and analysis, 439–483, Adv. Lect. Math. (ALM) **23**, Int. Press, Somerville, MA, 2012.
- [28] S. W. Wei and L. Wu, *Homotopy groups and  $p$ -harmonic maps*. Commun. Math. Anal., Conference **1** (2008), 40–45.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN 48824–1027, U.S.A.

*Email address:* `chenb@msu.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OKLAHOMA 73019-0315, U.S.A.

*Email address:* `wwei@ou.edu`