

QUASI MAXIMUM LIKELIHOOD ESTIMATION OF HIGH-DIMENSIONAL FACTOR MODELS:

A CRITICAL REVIEW

Matteo Barigozzi[†]

This version: May 12, 2023

Abstract

We review Quasi Maximum Likelihood estimation of factor models for high-dimensional panels of time series. We consider two cases: (1) estimation when no dynamic model for the factors is specified (Bai and Li, 2012, 2016); (2) estimation based on the Kalman smoother and the Expectation Maximization algorithm thus allowing to model explicitly the factor dynamics (Doz et al., 2012; Barigozzi and Luciani, 2019). Our interest is in approximate factor models, i.e., when we allow for the idiosyncratic components to be mildly cross-sectionally, as well as serially, correlated. Although such setting apparently makes estimation harder, we show, in fact, that factor models do not suffer of the *curse of dimensionality* problem, but instead they enjoy a *blessing of dimensionality* property. In particular, given an approximate factor structure, if the cross-sectional dimension of the data, N , grows to infinity, we show that: (i) identification of the model is still possible, (ii) the misspecification error due to the use of an exact factor model log-likelihood vanishes. Moreover, if we let also the sample size, T , grow to infinity, we can also consistently estimate all parameters of the model and make inference. The same is true for estimation of the latent factors which can be carried out by weighted least-squares, linear projection, or Kalman filtering/smoothing. We also compare the approaches presented with: Principal Component analysis and the classical, fixed N , exact Maximum Likelihood approach. We conclude with a discussion on efficiency of the considered estimators.

Keywords: Approximate Dynamic Factor Model; Maximum Likelihood Estimation; Weighted Least Squares; Expectation Maximization Algorithm; Kalman smoother.

[†]Department of Economics - Università di Bologna. Email: matteo.barigozzi@unibo.it

Section 6 heavily draws on Barigozzi and Luciani (2019) to which the reader is referred for the original results. I thank Matteo Luciani and Rolf Sundberg for helpful comments.

Contents

1	Introduction	3
2	The approximate factor model	4
2.1	Structural assumptions	5
2.2	Identification conditions	8
2.3	Factor dynamics	9
3	A review of the Principal Component estimators	11
4	The log-likelihood function	13
4.1	Exact log-likelihood	14
4.2	Mis-specified log-likelihoods	14
4.3	Identification of the QML estimator	15
5	QML estimation of an approximate static factor model	16
5.1	Loadings	16
5.2	Factors	19
6	QML estimation of an approximate dynamic factor model	21
6.1	Loadings	23
6.2	Factors	26
7	Classical ML estimation	29
7.1	The incidental parameter problem	31
8	On the mis-specification error and the search for efficient estimators	33
9	Concluding remarks	37
A	Kalman filter and Kalman smoother	37
	References	38

1 Introduction

Factor analysis is one of the earliest proposed multivariate statistical techniques: it dates back to the studies of Spearman (1904) in experimental psychology. The idea is to decompose a vector of N random variables into two components: (i) a common component driven by $r < N$ latent factors, and (ii) a component which is idiosyncratic, in the sense that it can be due either to measurement errors or to specific features of a single, or a sub-group of variables. As such we can retrospectively consider factor analysis as a pioneering technique in the field of unsupervised statistical learning.

Over the years, two main strands of applications have emerged: first, in psychometrics (Thomson, 1936; Bartlett, 1937, 1938; Thomson, 1951; Lawley and Maxwell, 1971; Bartholomew et al., 2011), second, in econometrics, for the analysis of financial markets (Connor et al., 2006; Ait-Sahalia and Xiu, 2017; Kim and Fan, 2019; Barigozzi and Hallin, 2020), the measurement and prediction of macroeconomic aggregates (Stock and Watson, 2002b; Bernanke et al., 2005; De Mol et al., 2008; Giannone et al., 2008; Barigozzi and Luciani, 2021), and the analysis of demand systems (Stone, 1945; Barigozzi and Moneta, 2016).

The former strand of literature deals with random samples of data, i.e., independent observations for each variable, and estimation is almost exclusively by means of Maximum Likelihood (Anderson and Rubin, 1956; Amemiya et al., 1987; Anderson and Amemiya, 1988; Tipping and Bishop, 1999; Bai and Li, 2012). The latter strand of literature deals with time series data, i.e., dependent observations for each variable, and estimation is either by means of Principal Component analysis (Stock and Watson, 2002a; Bai, 2003; Fan et al., 2013) or by means of Quasi Maximum Likelihood (Quah and Sargent, 1993; Doz et al., 2012; Bai and Li, 2016).

Our focus is on high-dimensional factor models for time series. Specifically, we allow for the dimension of the vector, N , to grow to infinity. As a consequence, it is realistic to consider an *approximate* factor model where we allow for the idiosyncratic components to be cross-sectionally correlated, rather than an *exact* factor model where the idiosyncratic components are not correlated. Furthermore, since we deal with time series, both the factors and the idiosyncratic components are allowed to be autocorrelated, and, if we specify also a model describing the time evolution of the factors, e.g., a VAR, we speak of a *dynamic* factor model, while, if we do not, we speak of a *static* factor model.

The idea of an approximate factor structure dates back to Chamberlain and Rothschild (1983) who showed that we if let the number of time series, N , to grow we are able to identify the common components even when allowing for idiosyncratic correlations. Intuitively, since estimation is based on cross-sectional aggregation, the more series we use, i.e., the more we increase N , the more the signal, factor-driven, component emerges from the noisy idiosyncratic component because of the cross-sectional weak Law of Large Numbers. Technically, this is seen by noticing that, under a factor model, the gap between the largest r eigenvalues of the covariance matrix of the data and the remaining $N - r$ widens as N grows. As a consequence of this phenomenon, an approximate factor model does not suffer of the *curse of dimensionality* problem, which typically affects high-dimensional parametric models, but it enjoys instead a *blessing of dimensionality* property.

The idea of a dynamic factor model goes back to Sargent and Sims (1977), while the setting considered here is the same as the one considered by Forni et al. (2009) and Doz et al. (2011, 2012) and it consists in writing the model as a linear state-space system, having as latent states the factors. This, in turn, is a restricted version of the more general model proposed by Forni et al. (2000) and recently

surveyed by Lippi (2018, 2019).

In this paper, we review likelihood-based methods for estimating high-dimensional approximate static and dynamic factor models, as considered by Bai and Li (2016) and Doz et al. (2012), respectively. On the one hand, Bai and Li (2016) treat the factors as non-stochastic and follow the classical approach based on first estimating the loadings and the idiosyncratic variances by Quasi Maximum Likelihood and then estimating the factors by Weighted Least Squares. On the other hand, Doz et al. (2012) explicitly model the factors as an autocorrelated stochastic process and propose to use the Expectation Maximization (EM) algorithm together with the Kalman smoother to obtain joint estimates of the loadings and the idiosyncratic variances, which approximate the Quasi Maximum Likelihood estimators, and of the factors. Both approaches are based on the maximization of a mis-specified log-likelihood where we do not take into account any of the idiosyncratic cross- and serial correlations, nevertheless, we show that as $N \rightarrow \infty$, the mis-specification error vanishes: yet another instance of the blessing of dimensionality.

2 The approximate factor model

Let us consider the $T \times N$ double-array \mathbf{X} having elements $\{x_{it} \in, i = 1, \dots, N, t = 1, \dots, T\}$ of N observed times series over T time periods. We say that \mathbf{X} follows an r -factor model if

$$x_{it} = \alpha_i + \boldsymbol{\lambda}'_i \mathbf{F}_t + \xi_{it}, \quad i = 1, \dots, N, \quad t = 1, \dots, T. \quad (1)$$

where $\boldsymbol{\lambda}_i := (\lambda_{i1} \cdots \lambda_{ir})'$ and $\mathbf{F}_t := (F_{1t} \cdots F_{rt})'$ are the r -dimensional latent vectors of loadings for series i and factors, respectively. We call ξ_{it} the idiosyncratic component and $\chi_{it} := \boldsymbol{\lambda}'_i \mathbf{F}_t$ the common component.

It is convenient to write model (1), which is written in scalar notation, also in other equivalent ways which we list below.

Let $\mathbf{x}_t := (x_{1t} \cdots x_{Nt})'$ be the N -dimensional vector of observables at time t . Then, in vector notation (1) reads

$$\mathbf{x}_t = \boldsymbol{\alpha} + \mathbf{\Lambda} \mathbf{F}_t + \boldsymbol{\xi}_t, \quad t = 1, \dots, T,$$

where $\boldsymbol{\xi}_t := (\xi_{1t} \cdots \xi_{Nt})'$ is the corresponding N -dimensional vector of idiosyncratic components, $\mathbf{\Lambda} := (\boldsymbol{\lambda}_1 \cdots \boldsymbol{\lambda}_N)' = (\boldsymbol{\ell}_1 \cdots \boldsymbol{\ell}_r)$ is the $n \times r$ matrix of factor loadings with rows $\boldsymbol{\lambda}'_i$, $i = 1, \dots, N$ and columns $\boldsymbol{\ell}_j$, $j = 1, \dots, r$, and $\boldsymbol{\alpha} := (\alpha_1 \cdots \alpha_N)'$. We call $\boldsymbol{\chi}_t := \mathbf{\Lambda} \mathbf{F}_t$ the vector of common components.

Equivalently, let $\mathbf{x}_i := (x_{i1} \cdots x_{iT})'$ be the T -dimensional time series vector of the i th observable, in vector notation (1) also reads

$$\mathbf{x}_i = \alpha_i \boldsymbol{\nu}_T + \mathbf{F} \boldsymbol{\lambda}_i + \boldsymbol{\zeta}_i, \quad i = 1, \dots, N,$$

where $\boldsymbol{\zeta}_i := (\xi_{i1} \cdots \xi_{iT})'$ is the corresponding T -dimensional time series vector of the i th idiosyncratic component, $\mathbf{F} := (\mathbf{F}_1 \cdots \mathbf{F}_T)'$ is the $T \times r$ matrix of factors, and $\boldsymbol{\nu}_T$ is a T -dimensional column vector of ones.

Finally, given that $\mathbf{X} = (\mathbf{x}_1 \cdots \mathbf{x}_T)'$, in matrix notation (1) reads

$$\mathbf{X} = \boldsymbol{\nu}_T \boldsymbol{\alpha}' + \mathbf{F} \mathbf{\Lambda}' + \boldsymbol{\Xi}, \quad (2)$$

where $\Xi := (\xi_1 \cdots \xi_T)'$ is the $T \times N$ matrix of idiosyncratic components.

2.1 Structural assumptions

In this section we provide the minimal set of assumptions required to derive the results given in the rest of the paper. Notice that, although these assumptions do not coincide with those made by Bai (2003), Bai and Li (2012), and Bai and Li (2016), they nest all assumptions made in those papers. More precisely, they nest all necessary assumptions in those papers, since, in fact, some of the assumptions made therein turn out to be redundant. In the following, we compare our assumptions only with Bai and Li (2012, 2016), while for a comparison with Bai (2003) we refer to Barigozzi (2022).

We characterize the common component by means of the following assumption.

Assumption 1 (COMMON COMPONENT).

- (a) $\lim_{N \rightarrow \infty} \|N^{-1} \mathbf{A}' \mathbf{A} - \Sigma_\Lambda\| = 0$, where Σ_Λ is $r \times r$ positive definite, and, for all $i = 1, \dots, N$ and all $N \in \mathbb{N}$, $\|\boldsymbol{\lambda}_i\| \leq M_\Lambda$ for some finite positive real M_Λ independent of i .
- (b) For all $t = 1, \dots, T$ and all $T \in \mathbb{N}$, $\mathbb{E}[\mathbf{F}_t] = \mathbf{0}_r$, $\mathbf{\Gamma}^F := \mathbb{E}[\mathbf{F}_t \mathbf{F}_t']$ is $r \times r$ positive definite, $\|\mathbf{\Gamma}^F\| \leq M_F$ and $\mathbb{E}[\|\mathbf{F}_t\|^4] \leq K_F$, for some finite positive reals M_F and K_F independent of t .
- (c) $\text{p-lim}_{T \rightarrow \infty} \|T^{-1} \mathbf{F}' \mathbf{F} - \mathbf{\Gamma}^F\| = 0$.
- (d) There exists a positive integer N_0 such that for all $N \in \mathbb{N}$ with $N > N_0$, r is a finite positive integer, independent of N .
- (e) For all $i = 1, \dots, N$ and all $N \in \mathbb{N}$, $\alpha_i \in \mathbb{R}$.

Parts (a) and (b), are also assumed in Bai and Li (2016, Assumptions A and B), and are similar to what assumed by Bai and Li (2012, Assumptions A and C). They imply pervasiveness of the factors, i.e., all factors have a non-negligible effect on all observables. Indeed, the loadings matrix has asymptotically maximum column rank r (part (a)), the factors have a finite full-rank covariance matrix (part (b)), and, because of the upper bound on $\|\boldsymbol{\lambda}_i\|$ in part (a), for any given $N \in \mathbb{N}$, the contribution of each factor to every series is finite.

Part (c) is analogous to what assumed by Bai and Li (2012, 2016, Assumption A) in the case of deterministic factors. It is very general, it simply says that the sample covariance matrix of the factors is a consistent estimator of its population counterpart $\mathbf{\Gamma}^F$. It is satisfied by linear processes having innovations with finite 4th order moments, or strong mixing processes with finite 4th order moments, or, more in general, by processes having summable 4th order cross-cumulants, which is a necessary and sufficient condition for consistency of the sample covariance matrix (Hannan, 1970, pp. 209-211).

Part (d) implies the existence of a finite of factors for all $N, T \in \mathbb{N}$. In particular, the numbers of common factors, r , is identified only for $N \rightarrow \infty$. Here N_0 is the minimum number of series we need to be able to identify r so that it must be such that $N_0 \geq r$. Without loss of generality hereafter when we say “for all $N \in \mathbb{N}$ ” we always mean that $N > N_0$ so that r can be identified. In practice, we must always work with N such that $r < \min(N, T)$.

Last, part (e), assumes a α_i to be a fixed idiosyncratic effect. Sometimes (1) is also written as (Bai and Li, 2012, see the remarks to Assumption A) $x_{it} = \alpha_i + \boldsymbol{\lambda}_i' \bar{\mathbf{F}} + \boldsymbol{\lambda}_i' (\mathbf{F}_t - \bar{\mathbf{F}}) + \xi_{it} = \alpha_i^* + \boldsymbol{\lambda}_i' \mathbf{F}_t^* + \xi_{it}$, with $\bar{\mathbf{F}} := T^{-1} \sum_{t=1}^T \mathbf{F}_t$, $\mathbf{F}_t^* := \mathbf{F}_t - \bar{\mathbf{F}}$, and $\alpha_i^* := \alpha_i + \boldsymbol{\lambda}_i' \bar{\mathbf{F}}$ which is then a random idiosyncratic effect.

Now, because of Assumption 1(a), we have that the eigenvalues of Σ_Λ , denoted as $\mu_j(\Sigma_\Lambda)$, $j = 1, \dots, r$ are such that $m_\Lambda^2 \leq \mu_j(\Sigma_\Lambda) \leq M_\Lambda^2$ (see also Lemma 3(iv) in Barigozzi, 2022). Similarly, because

of Assumptions 1(b) and 1(c), we have that the eigenvalues of $\mathbf{\Gamma}^F$, denoted as $\mu_j(\mathbf{\Gamma}^F)$, $j = 1, \dots, r$ are such that $m_F \leq \mu_j(\mathbf{\Gamma}^F) \leq M_F$ (see also Lemma 3(v) in Barigozzi, 2022).

Hereafter, denote the covariance matrix of the common component as

$$\mathbf{\Gamma}^\chi := \mathbb{E}[\boldsymbol{\chi}_t \boldsymbol{\chi}_t'] = \mathbf{\Lambda} \mathbf{\Gamma}^F \mathbf{\Lambda}',$$

with j th largest eigenvalue μ_j^χ , $j = 1, \dots, r$. Then, it follows that (see also Lemma 1(iv) in Barigozzi, 2022) for all $j = 1, \dots, r$,

$$\underline{C}_j := m_\Lambda^2 m_F \leq \liminf_{N \rightarrow \infty} \frac{\mu_j^\chi}{N} \leq \limsup_{N \rightarrow \infty} \frac{\mu_j^\chi}{N} \leq M_\Lambda^2 M_F =: \overline{C}_j. \quad (3)$$

This means we consider only strong factors. For weak factors see, e.g., Onatski (2012), Uematsu and Yamagata (2021), Bai and Ng (2021), and Freyaldenhoven (2022), among many others.

To characterize the idiosyncratic component, we make the following assumptions.

Assumption 2 (IDIOSYNCRATIC COMPONENT).

- (a) For all $i = 1, \dots, N$, all $t = 1, \dots, T$, and all $N, T \in \mathbb{N}$, $\mathbb{E}[\xi_{it}] = 0$ and $\sigma_i^2 := \mathbb{E}[\xi_{it}^2] \geq C_\xi$ for some finite positive real C_ξ independent of i , t , N , and T .
- (b) For all $i, j = 1, \dots, N$, all $t = 1, \dots, T$, all $N, T \in \mathbb{N}$, and all $k \in \mathbb{Z}$, $|\mathbb{E}[\xi_{it} \xi_{j, t-k}]| \leq \rho^{|k|} M_{ij}$ for some finite positive reals ρ and M_{ij} independent of t and such that $0 \leq \rho < 1$, $\sum_{j=1}^N M_{ij} \leq M_\xi$, and $\sum_{i=1}^N M_{ij} \leq M_\xi$ for some finite positive real M_ξ independent of i , j , and N .
- (c) For all $i = 1, \dots, N$, all $t = 1, \dots, T$, and all $N, T \in \mathbb{N}$, $\mathbb{E}[|\xi_{it}|^4] \leq K_\xi$ for some finite positive real K_ξ independent of i and t .
- (d) For all $j = 1, \dots, N$, all $s = 1, \dots, T$, and all $N, T \in \mathbb{N}$,

$$\max \left\{ \mathbb{E} \left[\left| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^T \{\xi_{it} \xi_{jt} - \mathbb{E}[\xi_{it} \xi_{jt}]\} \right|^2 \right], \mathbb{E} \left[\left| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^T \{\xi_{it} \xi_{is} - \mathbb{E}[\xi_{it} \xi_{is}]\} \right|^2 \right] \right\} \leq K_\xi$$

for some finite positive real K_ξ independent of j , s , N , and T .

Part (a), is a minimum requirement for having non-degenerate idiosyncratic components. Although, in principle, this assumption could be removed (Bartholomew et al., 2011, Chapter 3.4), here it is maintained for simplicity, so that the log-likelihood is always well-defined. Moreover, from part (a), jointly with Assumption 1(b) by which $\mathbb{E}[\chi_{it}] = 0$, we see that we are implicitly assuming that the observables are such that $\mathbb{E}[x_{it}] = \alpha_i$, $i = 1, \dots, N$.

Part (b) has a twofold purpose. First, it limits the degree of serial correlation of the idiosyncratic components. Second, it also limits the degree of cross-sectional correlation between idiosyncratic components. In particular, part (b) implies the conditions required by Bai and Li (2016, Assumption C3,

C4, and E1) (see Lemma 1(i)-1(iii) in Barigozzi, 2022), i.e.,

$$\begin{aligned}
\sup_{N,T \in \mathbb{N}} \frac{1}{NT} \sum_{i,j=1}^N \sum_{t,s=1}^T |\mathbb{E}[\xi_{it}\xi_{js}]| &\leq \frac{M_\xi(1+\rho)}{1-\rho}, \\
\sup_{N,T \in \mathbb{N}} \max_{t=1,\dots,T} \frac{1}{N} \sum_{i,j=1}^N |\mathbb{E}[\xi_{it}\xi_{jt}]| &\leq M_\xi, \\
\sup_{N,T \in \mathbb{N}} \max_{i=1,\dots,N} \frac{1}{T} \sum_{t,s=1}^T |\mathbb{E}[\xi_{it}\xi_{is}]| &\leq \frac{M_\xi(1+\rho)}{1-\rho},
\end{aligned} \tag{4}$$

where ρ and M_ξ are defined in Assumption 2(b).

Define the $N \times N$ idiosyncratic covariance matrix and the diagonal matrix of idiosyncratic variances as:

$$\mathbf{\Gamma}^\xi := \mathbb{E}[\boldsymbol{\xi}_t \boldsymbol{\xi}_t'] \quad \text{and} \quad \boldsymbol{\Sigma}^\xi := \text{diag}(\sigma_1^2, \dots, \sigma_N^2),$$

respectively. Then, letting μ_1^ξ be the largest eigenvalue of $\mathbf{\Gamma}^\xi$, it follows that (see Lemma 1(v) in Barigozzi, 2022)

$$\sup_{N \in \mathbb{N}} \mu_1^\xi \leq M_\xi, \tag{5}$$

where M_ξ is defined in Assumption 2(b). Condition (5) was originally assumed by Chamberlain and Rothschild (1983) to control the amount of idiosyncratic correlation that we can allow for. Moreover, by setting $k = 0$ in Assumption 2(b), it follows also that for all $i = 1, \dots, N$ and all $N \in \mathbb{N}$, $\sigma_i^2 \leq M_\xi$. Thus, all idiosyncratic components have finite variance.

Parts (c) and (d) require finite 4th order moments as in Bai and Li (2012, Assumption B) and finite sums of 4th order cross-moments. Notice that this is weaker than the requirement of finite 8th order cross-moments assumed by Bai and Li (2016, Assumption C). Part (d) also implies that the sample covariance between $\{\xi_{it}\}$ and $\{\xi_{jt}\}$ is a consistent estimator of $\mathbb{E}[\xi_{it}\xi_{jt}]$ (Hannan, 1970, pp. 209-211).

The basic set of assumption is completed by the following requirement.

Assumption 3 (INDEPENDENCE). *For all $N, T \in \mathbb{N}$, the sequences $\{\xi_{it}, i = 1, \dots, N, t = 1, \dots, T\}$ and $\{F_{jt}, j = 1, \dots, r, t = 1, \dots, T\}$ are mutually independent.*

This assumption obviously implies that the common components are independent of the idiosyncratic components at all leads and lags and across all units. Notice that Bai and Li (2012, 2016) treat the factors as constant parameters, therefore, by construction, they have no relation with the idiosyncratic components. Finally, Assumption 3 is compatible with a structural macroeconomic interpretation of factor models, according to which the factors driving the common component are independent of the idiosyncratic components representing measurement errors or local dynamics.

We conclude with an assumption of two Central Limit Theorems which are required to allow us to conduct inference on the loadings and the factors. It coincides with the assumption made by Bai and Li (2016, Assumption F).

Assumption 4 (CENTRAL LIMIT THEOREMS).

(a) For all $i = 1, \dots, N$ and all $N \in \mathbb{N}$, as $T \rightarrow \infty$,

$$\frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{F}_t \xi_{it} \rightarrow_d \mathcal{N} \left(\mathbf{0}_r, \lim_{T \rightarrow \infty} \frac{\mathbb{E}[\mathbf{F}' \zeta_i \zeta_i' \mathbf{F}]}{T} \right).$$

(b) For all $t = 1, \dots, T$ and all $T \in \mathbb{N}$, as $N \rightarrow \infty$,

$$\frac{1}{\sqrt{N}} \sum_{i=1}^N \lambda_i \xi_{it} \rightarrow_d \mathcal{N} \left(\mathbf{0}_r, \lim_{N \rightarrow \infty} \frac{\mathbb{E}[\boldsymbol{\Lambda}' \boldsymbol{\xi}_t \boldsymbol{\xi}_t' \boldsymbol{\Lambda}]}{N} \right).$$

Remark 1. Assumption 4(a) can be derived from more primitive assumptions, for example it follows from Ibragimov (1962, Theorem 1.4) under the assumption of strong mixing factors and idiosyncratic components (see also Section 3 in Barigozzi, 2022). Assumption 4(b) of course holds if we assumed cross-sectionally independent idiosyncratic components. In general, to derive it from primitive assumptions we should introduce some notion of ordering of the N cross-sectional units, as, e.g., the notion of mixing random fields (Bolthausen, 1982). Since a notion of ordering might be unnatural in many contexts, we could alternatively resource to the notion of exchangeability, which is instead independent of the ordering (see, e.g., Bolthausen, 1984).

2.2 Identification conditions

Let μ_i^x , $i = 1, \dots, N$, be the i th largest eigenvalue of $\boldsymbol{\Gamma}^x := \mathbb{E}[(\mathbf{x}_t - \boldsymbol{\alpha})(\mathbf{x}_t - \boldsymbol{\alpha})']$. Then, we have the following result.

Lemma 1. Under Assumptions 1 through 3:

- (i) for all $j = 1, \dots, r$, $\underline{C}_j \leq \liminf_{N \rightarrow \infty} N^{-1} \mu_j^x \leq \limsup_{n \rightarrow \infty} N^{-1} \mu_j^x \leq \overline{C}_j$;
- (ii) $\sup_{N \in \mathbb{N}} \mu_{r+1}^x \leq M_\xi$;

where \underline{C}_j and \overline{C}_j are defined in (3), and M_ξ is defined in Assumption 2(b).

The proof is omitted since it is a direct consequence of Weyl's inequality, combined with (3) which follows from Assumption 1, (5) which follows from Assumption 2, and since $\boldsymbol{\Gamma}^x = \boldsymbol{\Gamma}^\lambda + \boldsymbol{\Gamma}^\xi$ which follows from Assumption 3.

Lemma 1 shows that the spectrum of the population covariance matrix of $\{\mathbf{x}_t\}$ has an eigen-gap which widens as N grows. This is a necessary condition for identification of the model, and, indeed, virtually all existing methods for estimating the number of factors, r , are based on such property (Bai and Ng, 2002, Alessi et al., 2010, Onatski, 2010, Ahn and Horenstein, 2013, Trapani, 2018). Notice, however, that even if an eigen-gap is displayed by an observed data set, this is not sufficient to say that the underlying data generating process is a factor model. Of course, in practice, requiring $N \rightarrow \infty$ is not sufficient for identification, since, as shown later, in order to consistently estimate all quantities we also need $T \rightarrow \infty$.

For QML estimation it is crucial that the parameters of the model and the factors are identified. However, it is well known that in general the factor loadings, as well as the factors, are not identified. Indeed, all the structures equivalent to (1) can be obtained through an $r \times r$ invertible matrix \mathbf{R} , which defines the transformed quantities: $\mathbf{F}_t^o := \mathbf{R}^{-1} \mathbf{F}_t$, $\boldsymbol{\Lambda}^o := \boldsymbol{\Lambda} \mathbf{R}$, and $\boldsymbol{\Gamma}^{\xi o} := \boldsymbol{\Gamma}^\xi$. Under such relationships,

using only first- and second-moment information, as it is the case with gaussian QML estimation, the model specified by \mathbf{F}_t^o , $\mathbf{\Lambda}^o$, and $\mathbf{\Gamma}^{\xi o}$, is indistinguishable from the one given by \mathbf{F}_t , $\mathbf{\Lambda}$, and $\mathbf{\Gamma}^\xi$.

To obtain an identified model, we then need enough a priori structure to preclude any but the trivial transformation $\mathbf{R} = \mathbf{I}_r$. This can be achieved by imposing additional r^2 identifying constraints. In particular, we impose the following standard identifying conditions.

Assumption 5 (LOADINGS AND FACTORS IDENTIFICATION).

- (a) For all $N \in \mathbb{N}$, $\frac{\mathbf{\Lambda}'\mathbf{\Lambda}}{N}$ diagonal with distinct elements.
- (b) For all $T \in \mathbb{N}$ $\frac{\mathbf{F}'\mathbf{F}}{T} = \mathbf{I}_r$.
- (c) For all $j = 1, \dots, r$, one of the two following conditions hold: (i) $\lambda_{1j} \geq 0$; (ii) $F_{j1} \geq 0$.

Under parts (a) and (b) the loadings and the factors are identified up to a sign (see conditions PC1 in Bai and Ng, 2013), hence, they are only locally identified. To achieve global identification, in part (c) we fix the sign indeterminacy.

Let now \mathbf{V} be the $r \times r$ diagonal matrix of non-zero eigenvalues of $(\mathbf{\Sigma}_\Lambda)^{1/2}\mathbf{\Gamma}^F(\mathbf{\Sigma}_\Lambda)^{1/2}$ or equivalently of $(\mathbf{\Gamma}^F)^{1/2}\mathbf{\Sigma}_\Lambda(\mathbf{\Gamma}^F)^{1/2}$. Then, because of Assumption 5: (A) $\mathbf{\Sigma}_\Lambda = \mathbf{V}$; (B) $\mathbf{\Gamma}^F = \mathbf{I}_r$. Moreover, from parts (a) and (b) it follows also that for all $N \in \mathbb{N}$, the $r \times r$ matrix $N^{-1}\mathbf{\Lambda}'\mathbf{\Lambda}$ is the diagonal matrix having as diagonal entries $N^{-1}\mu_1^\chi, \dots, N^{-1}\mu_r^\chi$, i.e., the eigenvalues of $\mathbf{\Gamma}^\chi$ scaled by N , which therefore are assumed to be distinct. Note that this implies that $\overline{C}_j < \underline{C}_{j-1}$, $j = 2, \dots, r$, in (3) and that also the diagonal elements of \mathbf{V} are distinct. Requiring distinct eigenvalues is useful for identifying the eigenvectors of $\mathbf{\Gamma}^\chi$. Although, in principle, this requirement is needed only for PC estimation (Bai, 2003, Assumption G), it is also needed to ensure consistency of the numerical maximization algorithms presented below and used for QML estimation, which are initialized with the PC estimators.

The identifying conditions in Assumption 5 are similar to the set of constraints IC3 considered in Bai and Li (2012, 2016), where, however, part (a) is replaced by the requiring $N^{-1}\mathbf{\Lambda}'(\mathbf{\Sigma}^\xi)^{-1}\mathbf{\Lambda}$ to be diagonal.

Remark 2. Clearly, Assumption 5 does not provide any economic meaning to the factors; hence, they can be used only for an exploratory factor analysis. For alternative identification conditions, which allow to interpret the factors, hence to conduct confirmatory factor analysis, see, e.g., the classical approach by Jöreskog (1969), then partially adopted also by Bai and Li (2012, 2016) and Li et al. (2018). In this paper we do not deal with confirmatory factor analysis since in most macroeconomic applications the focus is only the estimated common component, which is always identified, so no interpretation of the factors is required.

2.3 Factor dynamics

Under Assumptions 1(b)-1(d) the factors can be seen either as an rT -dimensional sequence of constant parameters (in which case in 1(b) the expectations would be replaced by averages and in 1(c) we would have an ordinary limit), or as a realization of an r -dimensional stochastic process $\{\mathbf{F}_t, t \in \mathbb{Z}\}$. The former case is commonly considered in classical factor analysis where observations are collected from random samples. The latter case is especially relevant in time series analysis. Nevertheless, if we believe the factors to be truly autocorrelated, then it would be desirable to explicitly model the dynamic evolution of the factors. This is the approach originally adopted by Sargent and Sims (1977)

and Quah and Sargent (1993) for dynamic factor analysis and then re-introduced by Forni et al. (2009) and Doz et al. (2011, 2012).

In order to keep things simple, but without loss of generality, here we assume the VAR(1) specification:

$$\mathbf{F}_t = \mathbf{A}\mathbf{F}_{t-1} + \mathbf{H}\mathbf{u}_t. \quad (6)$$

We make the following assumption.

Assumption 6.

- (a) The roots of $\det(\mathbf{I}_r - \mathbf{A}z) = 0$ are $z_j^* \in \mathbb{C}$, $j = 1, \dots, r$ such that $|z_j^*| \geq M_A$ for some finite positive real $M_A > 1$ independent of j .
- (b) \mathbf{H} is lower-triangular, $\text{rk}(\mathbf{H}) = r$, and $\|\mathbf{H}\| \leq M_H$ for some finite positive real M_H .
- (c) For all $t = 1, \dots, T$ and all $T \in \mathbb{N}$, $\mathbb{E}[\mathbf{u}_t] = \mathbf{0}_r$ and $\mathbb{E}[\mathbf{u}_t \mathbf{u}_t'] = \mathbf{I}_r$.
- (d) For all $t = 1, \dots, T$, all $T \in \mathbb{N}$, and all $k \in \mathbb{Z}$ with $k \neq 0$, $\mathbb{E}[\mathbf{u}_t \mathbf{u}_{t-k}'] = \mathbf{0}_{r \times r}$.
- (e) For all $j_1, j_2, j_3, j_4 = 1, \dots, r$, all $t_1, t_2, t_3, t_4 = 1, \dots, T$ and all $T \in \mathbb{N}$,

$$\mathbb{E}[|u_{j_1 t_1} u_{j_2 t_2} u_{j_3 t_3} u_{j_4 t_4}|] \leq K_u$$

for some finite positive real K_u independent of $j_1, j_2, j_3, j_4, t_1, t_2, t_3, t_4$, and T .

- (f) For all $j_1, j_2, j_3, j_4 = 1, \dots, r$, all $k \in \mathbb{Z}^+$, all $t, s = 1, \dots, T$ and all $T \in \mathbb{N}$

$$\frac{1}{T} \sum_{t,s=1}^T |\mathbb{E}[u_{j_1 t} u_{j_2 t-k} u_{j_3 s} u_{j_4 s-k}]| \leq C_u$$

for some finite positive real K_u independent of $j_1, j_2, j_3, j_4, k_1, k_2, t_1, t_2$, and T .

- (g) For all $t \leq 0$, $\mathbf{u}_t = \mathbf{0}_r$.

Assumption 6 is standard in VAR literature. Part (a), requires the factors to have a causal autoregressive representation. If we let $\mathbf{v}_t := \mathbf{H}\mathbf{u}_t$ and $\mathbf{\Gamma}^v := \mathbb{E}[\mathbf{v}_t \mathbf{v}_t']$, then, under parts (b) and (c), \mathbf{H} is the Choleski factor of $\mathbf{\Gamma}^v$, i.e., $\mathbf{H} = (\mathbf{\Gamma}^v)^{1/2}$. In part (e), we fix the initial conditions.

As a consequence, $\{\mathbf{F}_t\}$ admits the Wold representation $\mathbf{F}_t = \sum_{k=0}^{t-1} \mathbf{C}_k \mathbf{v}_{t-k}$, with coefficients $\mathbf{C}_k := \mathbf{A}^k$ such that $\sum_{k=0}^{\infty} \|\mathbf{C}_k\|^2 \leq M_C$ for some finite positive real M_C and with innovations process $\{\mathbf{v}_t\}$ which is a zero-mean white noise with finite and summable 4th order cumulants, and with covariance matrix $\mathbf{\Gamma}^v$ finite and positive definite. In fact, parts (a) through (e) imply Assumption 1(b) and, jointly with part (f), they also imply Assumption 1(c), and both become redundant in this setting.¹

Remark 3. Under Assumptions 5(a), 6(a), and 6(b), the linear system described by (1) and (6) is controllable and observable, as well as stabilizable and detectable.² These properties have two consequences.

¹Specifically, it follows that $\mathbb{E}[\mathbf{F}_t] = \sum_{k=0}^{t-1} \mathbf{C}_k \mathbb{E}[\mathbf{v}_{t-k}] = \mathbf{0}_r$, $\mathbf{\Gamma}^F = \sum_{k=0}^{t-1} \mathbf{C}_k \mathbf{H} \mathbf{H}' \mathbf{C}_k' \leq M_C^2 M_H^2$ and it is positive definite since $\mu_r(\mathbf{\Gamma}^F) \geq \mu_r(\mathbf{H} \mathbf{H}') > 0$, $\mathbb{E}[\|\mathbf{F}_t\|^4] \leq r^{10} M_C^4 M_H^4 K_u$, and, last, $\mathbb{E}[|T^{-1/2} \sum_{t=1}^T F_{it} F_{jt} - \mathbb{E}[F_{it} F_{jt}]|^2] \leq r^{10} M_C^4 M_H^4 K_u$, which implies Assumption 1(c) by Chebychev's inequality.

²The couple (\mathbf{A}, \mathbf{H}) is controllable if and only if $\text{rk}[\mathbf{H}(\mathbf{A}\mathbf{H}) \dots (\mathbf{A}^{(r-1)}\mathbf{H})] = r$ and the couple $(\mathbf{A}, \mathbf{\Lambda})$ is observable if and only if $\text{rk}[\mathbf{\Lambda}'(\mathbf{\Lambda}\mathbf{A})' \dots (\mathbf{\Lambda}\mathbf{A}^{(r-1)})'] = r$, moreover, the linear system is stabilizable if its unstable states are controllable and all uncontrollable states are stable, and it is detectable if its unstable states are observable and all unobservable states are stable (Anderson and Moore, 1979, Appendix C, pp. 341-342).

First, the standard mini-phase condition:

$$\text{rk} \begin{pmatrix} \mathbf{I}_r - \mathbf{A}z & \mathbf{H} \\ \mathbf{\Lambda} & \mathbf{0}_{r \times r} \end{pmatrix} = 2r,$$

is satisfied, and (1)-(6) is the minimal state space representation of a dynamic approximate factor model, having as McMillan degree the number of factors r (Anderson and Deistler, 2008, Section II). This result, combined with Assumption 5, makes the linear system fully identified. Second, the one-step-ahead covariance matrix of the linear prediction error, denoted as $\mathbf{P}_{t|t-1}$ has a steady-state \mathbf{P} which is such that $\|\mathbf{P}_{t|t-1} - \mathbf{P}\| = O(N^{-1})$ for all $t \geq \bar{t}$ with $\bar{t} \geq 2$ and finite for all T (see Barigozzi and Luciani, 2019, Lemma 7).

Remark 4. The factor model defined in (1) is called a static factor model, in that the factors are loaded only contemporaneously by the data. If we include also a dynamic model for the factors, such as (6), we could also write (1) as

$$x_{it} = \alpha_i + \boldsymbol{\lambda}'_i \sum_{k=0}^{\infty} \mathbf{A}^k \mathbf{H} \mathbf{u}_{t-k} + \xi_{it}, \quad i = 1, \dots, N, \quad t = 1, \dots, T, \quad (7)$$

where $\{\mathbf{u}_t\}$ is an orthonormal r -dimensional white noise process of so-called dynamic factors, in that they are loaded by the data together with their lags. For this reason the model described by (1)-(6) is called a dynamic factor model, and (7) is a special case of the Generalized Dynamic Factor model introduced by Forni et al. (2000). The GDFM is actually more general and it is defined as

$$x_{it} = \alpha_i + \sum_{k=0}^{\infty} \mathbf{b}'_{ik} \mathbf{e}_{t-k} + \xi_{it}, \quad i = 1, \dots, N, \quad t = 1, \dots, T, \quad (8)$$

for some square-summable sequence of $N \times q$ matrix of coefficients $\{\mathbf{b}_{ik}\}$ and an orthonormal q -dimensional white noise process $\{\mathbf{e}_t\}$ with $q < r$. As shown in Stock and Watson (2016, Section 2.1), the GDFM admits a state-space representation as (1)-(6) only if in (8) we allow for just finite number of lags, and we impose specific zero restrictions on \mathbf{A} and \mathbf{H} .

3 A review of the Principal Component estimators

In this section we briefly review Principal Components (PC) estimation of the static approximate factor model (1). This is arguably the most popular estimation approach and it is a reference estimator also for QML estimation. The PC estimators are non-parametric and, in particular, their implementation does not require to specify any second order structure of the idiosyncratic components as long as Assumption 2(b) is satisfied. There are a few different, although asymptotically equivalent, definitions of PCs. We refer to Barigozzi (2022) for a discussion of the various approaches.

In particular, here we consider the classical definition of PC estimators, where the factors are estimated as the orthonormal PCs of \mathbf{X} (see, e.g., Mardia et al., 1979, Chapter 9.2). This is also the definition adopted also by Forni et al. (2009). In particular, the PC estimator of the loadings is defined as:

$$\widehat{\boldsymbol{\lambda}}_i^{\text{PC}} := \left(\widehat{\mathbf{M}}^x\right)^{1/2} \widehat{\mathbf{v}}_i^x, \quad i = 1, \dots, N, \quad (9)$$

where $\widehat{\mathbf{v}}_i^{x'}$ is the r -dimensional i th row of the $N \times r$ matrix $\widehat{\mathbf{V}}^x$ having as columns the r normalized eigenvectors of the $N \times N$ sample covariance matrix $T^{-1}(\mathbf{X} - \boldsymbol{\nu}_T \bar{\mathbf{x}})'(\mathbf{X} - \boldsymbol{\nu}_T \bar{\mathbf{x}})$ corresponding to the r largest eigenvalues collected in the diagonal matrix $\widehat{\mathbf{M}}^x$. Notice that, by construction, the $N \times r$ matrix of PC estimated loadings $\widehat{\boldsymbol{\Lambda}}^{\text{PC}} := (\widehat{\boldsymbol{\lambda}}_1^{\text{PC}} \cdots \widehat{\boldsymbol{\lambda}}_N^{\text{PC}})'$ is such that $N^{-1} \widehat{\boldsymbol{\Lambda}}^{\text{PC}'} \widehat{\boldsymbol{\Lambda}}^{\text{PC}} = N^{-1} \widehat{\mathbf{M}}^x$, which is diagonal. Hence, $\widehat{\boldsymbol{\Lambda}}^{\text{PC}}$ satisfies Assumption 5(a).

Under the identifying Assumptions 5, we have that the PC estimator is asymptotically equivalent to the unfeasible Ordinary Least Squares (OLS) estimator we would get if the factors were observed:

$$\boldsymbol{\lambda}_i^{\text{OLS}} := (\mathbf{F}' \mathbf{F})^{-1} \mathbf{F}'(\mathbf{x}_i - \bar{x}_i \boldsymbol{\nu}_T), \quad i = 1, \dots, N. \quad (10)$$

In particular, it can be shown that (Barigozzi, 2022, Proposition 2 and Theorem 1)

$$\|\widehat{\boldsymbol{\lambda}}_i^{\text{PC}} - \boldsymbol{\lambda}_i^{\text{OLS}}\| = O_P\left(\frac{1}{\sqrt{NT}}\right) + O_P\left(\frac{1}{N}\right), \quad (11)$$

from which we prove the following result.

Proposition 1. *Under Assumptions 1 through 5, for any $i = 1, \dots, N$, as $N, T \rightarrow \infty$, if $\sqrt{T}/N \rightarrow 0$,*

$$\sqrt{T}(\widehat{\boldsymbol{\lambda}}_i^{\text{PC}} - \boldsymbol{\lambda}_i) \rightarrow_d \mathcal{N}(\mathbf{0}_r, \boldsymbol{\nu}_i^{\text{OLS}}),$$

where $\boldsymbol{\nu}_i^{\text{OLS}} := (\boldsymbol{\Gamma}^F)^{-1} \left\{ \lim_{T \rightarrow \infty} \frac{\mathbb{E}[\mathbf{F}' \boldsymbol{\zeta}_i \boldsymbol{\zeta}_i' \mathbf{F}]}{T} \right\} (\boldsymbol{\Gamma}^F)^{-1} = \lim_{T \rightarrow \infty} \frac{\mathbb{E}[\mathbf{F}' \mathbb{E}[\boldsymbol{\zeta}_i \boldsymbol{\zeta}_i' | \mathbf{F}]]}{T}$ (because of Assumption 5(b)).

Since the PC estimator is non-parametric the asymptotic covariance matrix $\boldsymbol{\nu}_i^{\text{OLS}}$ has a sandwich form reflecting the fact that we do not take into account possible idiosyncratic serial correlations. The final expression of $\boldsymbol{\nu}_i^{\text{OLS}}$ comes from the law of iterated expectations, and since factors and idiosyncratic components are independent by Assumption 3.

The PC estimator of the factors is then obtained by linear projection of \mathbf{X} onto $\widehat{\boldsymbol{\Lambda}}^{\text{PC}}$:

$$\widehat{\mathbf{F}}_t^{\text{PC}} := \left(\widehat{\boldsymbol{\Lambda}}^{\text{PC}'} \widehat{\boldsymbol{\Lambda}}^{\text{PC}} \right)^{-1} \widehat{\boldsymbol{\Lambda}}^{\text{PC}'} (\mathbf{x}_t - \bar{\mathbf{x}}) = (\widehat{\mathbf{M}}^x)^{-1/2} \widehat{\mathbf{V}}^x (\mathbf{x}_t - \bar{\mathbf{x}}), \quad t = 1, \dots, T. \quad (12)$$

It follows that the $T \times r$ matrix of PC estimated factors $\widehat{\mathbf{F}}^{\text{PC}} := (\widehat{\mathbf{F}}_1^{\text{PC}} \cdots \widehat{\mathbf{F}}_T^{\text{PC}})'$ is such that $T^{-1} \widehat{\mathbf{F}}^{\text{PC}'} \widehat{\mathbf{F}}^{\text{PC}} = \mathbf{I}_r$. Hence $\widehat{\mathbf{F}}^{\text{PC}}$ satisfies Assumption 5(b).

Under the identifying Assumptions 5, we have that the PC estimator is asymptotically equivalent to the unfeasible OLS estimator we would get if the loadings were observed:

$$\mathbf{F}_t^{\text{OLS}} := (\boldsymbol{\Lambda}' \boldsymbol{\Lambda})^{-1} \boldsymbol{\Lambda}' (\mathbf{x}_t - \bar{\mathbf{x}}_t), \quad t = 1, \dots, T. \quad (13)$$

In particular, it can be shown that (Barigozzi, 2022, Proposition 5 and Theorem 2)

$$\|\widehat{\mathbf{F}}_t^{\text{PC}} - \mathbf{F}_t^{\text{OLS}}\| = O_P\left(\frac{1}{T}\right) + O_P\left(\frac{1}{\sqrt{NT}}\right), \quad (14)$$

from which we prove the following result.

Proposition 2. Under Assumptions 1 through 5, for any $t = 1, \dots, T$, as $N, T \rightarrow \infty$, if $\sqrt{N}/T \rightarrow 0$,

$$\sqrt{N}(\widehat{\mathbf{F}}_t^{\text{PC}} - \mathbf{F}_t) \rightarrow_d \mathcal{N}(\mathbf{0}_r, \mathbf{W}_t^{\text{OLS}}),$$

where $\mathbf{W}_t^{\text{OLS}} := (\boldsymbol{\Sigma}_\Lambda)^{-1} \left\{ \lim_{N \rightarrow \infty} \frac{\Lambda' \mathbb{E}[\boldsymbol{\xi}_t \boldsymbol{\xi}_t' \Lambda]}{N} \right\} (\boldsymbol{\Sigma}_\Lambda)^{-1}$.

Since the PC estimator is non-parametric the asymptotic covariance matrix $\mathbf{W}_t^{\text{OLS}}$ has a sandwich form reflecting the fact that we do not take into account possible idiosyncratic cross-sectional correlations. Notice that the asymptotic covariance could be time dependent if we had serially heteroskedastic idiosyncratic components, since it would then depend on $\boldsymbol{\Gamma}_t^\xi := \mathbb{E}[\boldsymbol{\xi}_t \boldsymbol{\xi}_t']$.

Remark 5. An equivalent set of PC estimators is adopted by Bai (2003), where, however, first the factors are estimated as \sqrt{T} times the eigenvectors of the $T \times T$ sample covariance $N^{-1}(\mathbf{X} - \boldsymbol{\iota}_T \bar{\mathbf{x}})(\mathbf{X} - \boldsymbol{\iota}_T \bar{\mathbf{x}})'$, hence they are the orthonormal PCs of \mathbf{X}' rather than of \mathbf{X} . The estimated loadings are then obtained by linear projection of \mathbf{X}' onto the estimated factors. Such estimators also satisfy the identifying Assumptions 5(a) and 5(b). This approach is considered also in Fan et al. (2013) and Bai and Ng (2020).

Finally, Stock and Watson (2002a) define the PC estimator of the loadings as \sqrt{N} times the eigenvectors of the $N \times N$ sample covariance matrix. However, such definition does not satisfy Assumption 5(a) and 5(b), since now the estimated loadings would have all the same scale, while the corresponding estimated factors would have different scales.

4 The log-likelihood function

First, let us introduce the following notation. Define $\boldsymbol{\mathcal{X}} := \text{vec}(\mathbf{X}') = (\mathbf{x}'_1 \cdots \mathbf{x}'_T)'$ and $\boldsymbol{\mathcal{Z}} := \text{vec}(\boldsymbol{\Xi}') = (\boldsymbol{\xi}'_1 \cdots \boldsymbol{\xi}'_T)'$ as the NT -dimensional vectors of observations and idiosyncratic components, $\boldsymbol{\mathcal{A}} := \text{vec}(\boldsymbol{\alpha} \boldsymbol{\iota}'_T)$ as the NT -dimensional vector of constants, $\boldsymbol{\mathcal{F}} := \text{vec}(\mathbf{F}') = (\mathbf{F}'_1 \cdots \mathbf{F}'_T)'$ as the rT -dimensional vector of factors, and $\boldsymbol{\mathcal{L}} := \mathbf{I}_T \otimes \boldsymbol{\Lambda}$ as the $NT \times rT$ block-diagonal matrix containing the loadings matrix replicated T times.

Then, by taking the vectorized version of the transposed of (2), we can write model (1) as:³

$$\boldsymbol{\mathcal{X}} = \boldsymbol{\mathcal{A}} + \boldsymbol{\mathcal{L}}\boldsymbol{\mathcal{F}} + \boldsymbol{\mathcal{Z}}. \quad (15)$$

Let $\boldsymbol{\Omega}^x := \mathbb{E}[(\boldsymbol{\mathcal{X}} - \boldsymbol{\mathcal{A}})(\boldsymbol{\mathcal{X}} - \boldsymbol{\mathcal{A}})']$ and $\boldsymbol{\Omega}^\xi := \mathbb{E}[\boldsymbol{\mathcal{Z}}\boldsymbol{\mathcal{Z}}']$, be the $NT \times NT$ covariance matrices of the NT -dimensional vectors of data and idiosyncratic components, respectively. Let also $\boldsymbol{\Omega}^F := \mathbb{E}[\boldsymbol{\mathcal{F}}\boldsymbol{\mathcal{F}}']$ be the $rT \times rT$ covariance matrix of the rT -dimensional factor vector. Then, because of Assumption 3,

$$\boldsymbol{\Omega}^x = \boldsymbol{\mathcal{L}} \boldsymbol{\Omega}^F \boldsymbol{\mathcal{L}}' + \boldsymbol{\Omega}^\xi.$$

The parameters of the model that need to be estimated are then $\boldsymbol{\alpha}$ and

$$\boldsymbol{\varphi} := (\text{vec}(\boldsymbol{\Lambda})', \text{vech}(\boldsymbol{\Omega}^\xi)', \text{vech}(\boldsymbol{\Omega}^F)')'.$$

³Recall that $\text{vec}(\mathbf{ABC}) = (\mathbf{C}' \otimes \mathbf{A})\text{vec}(\mathbf{B})$.

4.1 Exact log-likelihood

The gaussian quasi-log-likelihood computed in a generic value of the parameters, denoted as $\underline{\alpha}$ and $\underline{\varphi}$, is given by

$$\ell_*(\mathcal{X}; \underline{\alpha}, \underline{\varphi}) := -\frac{NT}{2} \log(2\pi) - \frac{1}{2} \log \det(\underline{\Omega}^x) - \frac{1}{2} [(\mathcal{X} - \underline{\mathbf{A}})' (\underline{\Omega}^x)^{-1} (\mathcal{X} - \underline{\mathbf{A}})]. \quad (16)$$

Let $\bar{\mathcal{X}} = \text{vec}(\bar{\mathbf{x}}\mathbf{1}'_T)$ with $\bar{\mathbf{x}} = T^{-1} \sum_{t=1}^T \mathbf{x}_t$. Then, it holds that

$$(\mathcal{X} - \underline{\mathbf{A}})' (\underline{\Omega}^x)^{-1} (\mathcal{X} - \underline{\mathbf{A}}) = (\mathcal{X} - \bar{\mathcal{X}})' (\underline{\Omega}^x)^{-1} (\mathcal{X} - \bar{\mathcal{X}}) + (\bar{\mathcal{X}} - \underline{\mathbf{A}})' (\underline{\Omega}^x)^{-1} (\bar{\mathcal{X}} - \underline{\mathbf{A}}),$$

which shows that (16) is maximized when $\underline{\mathbf{A}} = \bar{\mathcal{X}}$, i.e., the QML estimator of α is $\hat{\alpha}^{\text{QML}} := \bar{\mathbf{x}}$.

The concentrated quasi-log-likelihood computed in a generic value of the parameters $\underline{\varphi}$ is then given by:

$$\begin{aligned} \ell(\mathcal{X}; \underline{\varphi}) &:= -\frac{NT}{2} \log(2\pi) - \frac{1}{2} \log \det(\underline{\Omega}^x) - \frac{1}{2} [(\mathcal{X} - \bar{\mathcal{X}})' (\underline{\Omega}^x)^{-1} (\mathcal{X} - \bar{\mathcal{X}})] \\ &= -\frac{NT}{2} \log(2\pi) - \frac{1}{2} \log \det(\underline{\underline{\Sigma}} \underline{\Omega}^F \underline{\underline{\Sigma}}' + \underline{\Omega}^\xi) - \frac{1}{2} [(\mathcal{X} - \bar{\mathcal{X}})' (\underline{\underline{\Sigma}} \underline{\Omega}^F \underline{\underline{\Sigma}}' + \underline{\Omega}^\xi)^{-1} (\mathcal{X} - \bar{\mathcal{X}})]. \end{aligned} \quad (17)$$

In general, it might be very hard, if not impossible, to find the QML estimator of φ , i.e., the maximizer of (17). This is due to the very large number of parameters that need to be estimated. The QML estimator might not even exist unless other restrictions are imposed on the model. The main problem here is that, under our assumptions, we should estimate both Ω^ξ and Ω^F , which are in general full block Toeplitz matrices containing autocovariance matrices up to lag $(T-1)$. Moreover, each of the T blocks of Ω^ξ is in general an $N \times N$ full matrix, so, without imposing further restrictions, Ω^ξ alone contains already $NT(NT+1)/2$ parameters to be estimated, which is more than the NT available data points.

4.2 Mis-specified log-likelihoods

Henceforth, we will be working with a mis-specified log-likelihood where we treat the idiosyncratic components as serially and cross-sectionally uncorrelated. Thus, we consider a mis-specified model for the idiosyncratic 2nd order structure described by: $\Omega^\xi = \mathbf{I}_T \otimes \Sigma^\xi$, which reduces the parameters in Ω^ξ to be estimated to just the N elements of Σ^ξ . The log-likelihood (17) is then replaced by:

$$\begin{aligned} \ell_0(\mathcal{X}; \underline{\varphi}) &= -\frac{1}{2} \log \det \left(\{\mathbf{I}_T \otimes \underline{\mathbf{A}}\} \underline{\Omega}^F \{\mathbf{I}_T \otimes \underline{\mathbf{A}}'\} + \{\mathbf{I}_T \otimes \underline{\Sigma}^\xi\} \right) \\ &\quad - \frac{1}{2} \left[(\mathcal{X} - \bar{\mathcal{X}})' \left(\{\mathbf{I}_T \otimes \underline{\mathbf{A}}\} \underline{\Omega}^F \{\mathbf{I}_T \otimes \underline{\mathbf{A}}'\} + \{\mathbf{I}_T \otimes \underline{\Sigma}^\xi\} \right)^{-1} (\mathcal{X} - \bar{\mathcal{X}}) \right]. \end{aligned} \quad (18)$$

In general, maximization of (18) is still unfeasible because it requires estimating Ω^F which contains other $rT(rT+1)/2$ parameters. In this paper we consider two approaches. First, we consider a further mis-specification by imposing zero autocorrelation in the factors, i.e., by setting $\Omega^F = \mathbf{I}_T \otimes \Gamma^F = \mathbf{I}_{rT}$, where we used also Assumption 5(b). This is the approach reviewed in Section 5, which defines the QML estimator $\hat{\varphi}^{\text{QML,S}}$. Second, we consider a parametric linear model describing the factor dynamics, so that Ω^F depends on the parameters characterizing such model. In particular, in the case of the

VAR(1) specification given in (6), the (t, s) -block of $\boldsymbol{\Omega}^F$ is the $r \times r$ lag- $(t - s)$ autocovariance matrix: $\boldsymbol{\Gamma}_{t-s}^F := \mathbb{E}[\mathbf{F}_t \mathbf{F}_s']$ such that $\text{vec}(\boldsymbol{\Gamma}_{t-s}^F) = (\mathbf{I}_r \otimes \mathbf{A})^{t-s} (\mathbf{I}_{r^2} - \mathbf{A} \otimes \mathbf{A})^{-1} \text{vec}(\mathbf{H}\mathbf{H}')$. This reduces the number of parameters in $\boldsymbol{\Omega}^F$ to be estimated to $r^2 + r(r + 1)/2$. This is the approach reviewed in Section 6, which defines the QML estimator $\hat{\boldsymbol{\varphi}}^{\text{QML,D}}$.

Remark 6. An alternative QML estimation approach, not covered in this paper, consists in replacing the log-likelihood (18) by its Whittle frequency domain asymptotic approximation which is then maximized to estimate the parameters (Sargent and Sims, 1977; Dunsmuir, 1979; Geweke and Singleton, 1981). This approach is appealing since, in principle, it does not require specifying a parametric model for the dynamics of the factors. However, its implementation requires to first estimate a large dimensional spectral density matrix, which will produce an additional estimation error depending also on the chosen bandwidth (Zhang and Wu, 2021).

Remark 7. So far, we implicitly considered the idiosyncratic components as stationary, and, in particular, as serially homoskedastic. Nevertheless, under Assumption 2 we could in principle allow also for serial heteroskedasticity. In this case, we would have $\boldsymbol{\Gamma}_t^\xi := \mathbb{E}[\boldsymbol{\xi}_t \boldsymbol{\xi}_t']$ to be a time-dependent matrix with diagonal elements σ_{it}^2 collected into the diagonal matrix $\boldsymbol{\Sigma}_t^\xi$. Then, letting $\boldsymbol{\Sigma}^\xi := T^{-1} \sum_{t=1}^T \boldsymbol{\Sigma}_t^\xi$, with diagonal entries $\sigma_i^2 := T^{-1} \sum_{t=1}^T \sigma_{it}^2$, we could still consider the same log-likelihood (18), but where now we introduced an additional mis-specification in that we treat the idiosyncratic components as if they were serially homoskedastic. In this case, the QML estimators of the diagonal entries of $\boldsymbol{\Sigma}^\xi$, i.e., of σ_i^2 , $i = 1, \dots, N$, have to be viewed as estimators of the idiosyncratic variances averaged over time. This is the approach suggested by Bai and Li (2016). Given that in this context we cannot properly address the presence of serially heteroskedastic idiosyncratic components, hereafter, we rule out such possibility.

4.3 Identification of the QML estimator

Hereafter, we define the parameter space as the set $\mathcal{O} \in \mathbb{R}^Q$ where Q is the number of parameters to be estimated. It is intended that any generic parameter vector $\boldsymbol{\varphi} \in \mathcal{O}$ has elements satisfying all the assumptions given in Section 2. In particular, notice that, under those assumptions, the dimension of \mathcal{O} grows with N .

Usually in QML estimation it is required to first prove the existence of the QML estimator. However, to this end we cannot rely on compactness of \mathcal{O} since its dimension increases to infinity. Nevertheless, as shown in the next sections, in the present setting the parameter estimates maximizing the log-likelihood have an asymptotic linear representation, so we can easily bound the estimation errors directly by the distance between the estimated and the true parameter vector without the need of using any Taylor expansion for asymptotic analysis. This allows us to bypass the need to work with a compact parameter space. The only condition we need is for each element of the parameter vector $\boldsymbol{\varphi}$ to be in a bounded set, and this is guaranteed by Assumptions 1(a), 2(a), 2(b), 6(a), and 6(b).

Finally, in order to have a well defined log-likelihood in correspondence of the QML estimator of $\boldsymbol{\varphi}$, it is common to make the following assumption.

Assumption 7. For all $i = 1, \dots, N$ and all $N \in \mathbb{N}$, $\hat{\sigma}_i^{2\text{QML,S}} \in [C'_\xi, M'_\xi]$ and $\hat{\sigma}_i^{2\text{QML,D}} \in [C'_\xi, M'_\xi]$, for some finite positive reals C'_ξ and M'_ξ independent of i .

This assumption is identical to the requirement imposed by Bai and Li (2016, Assumption D). In

fact, it is a redundant condition which we assume only for the sake of simplicity. Indeed, it can be easily proved that in the present context Assumption 7 is always satisfied (Gao et al., 2021, Theorem 3.1).

5 QML estimation of an approximate static factor model

In this section we review the case in which the dynamics of the factors is not explicitly modeled, i.e., when we set $\mathbf{\Omega}^F = \mathbf{I}_{rT}$. So we consider an approximate static factor model described by the equation:

$$x_{it} = \alpha_i + \boldsymbol{\lambda}'_i \mathbf{F}_t + \xi_{it}, \quad i = 1, \dots, N, \quad t = 1, \dots, T,$$

together with Assumptions 1 through 5. Then, either $\{\mathbf{F}_t\}$ is effectively an uncorrelated process, so that it can also be considered as a deterministic sequence, or we are introducing a further mis-specification on top of the mis-specifications related to the idiosyncratic 2nd order structure introduced in the previous section.

For a static factor model the log-likelihood (18) simplifies to

$$\ell_{0,s}(\boldsymbol{\mathcal{X}}; \underline{\boldsymbol{\varphi}}) \simeq -\frac{T}{2} \log \det \left(\mathbf{\Lambda} \mathbf{\Lambda}' + \underline{\boldsymbol{\Sigma}}^\xi \right) - \frac{1}{2} \sum_{t=1}^T \left[(\mathbf{x}_t - \bar{\mathbf{x}})' \left(\mathbf{\Lambda} \mathbf{\Lambda}' + \underline{\boldsymbol{\Sigma}}^\xi \right)^{-1} (\mathbf{x}_t - \bar{\mathbf{x}}) \right]. \quad (19)$$

Notice that, up to a scaling by T , this is also known in the literature as the Wishart log-likelihood for the sample covariance $T^{-1}(\mathbf{x}_t - \bar{\mathbf{x}})(\mathbf{x}_t - \bar{\mathbf{x}})'$ (Anderson and Rubin, 1956). The parameters to be estimated are: $\underline{\boldsymbol{\varphi}} = (\text{vec}(\mathbf{\Lambda})', \sigma_1^2, \dots, \sigma_N^2)'$, which is a $Q = (r+1)N$ -dimensional vector. We shall denote the maximizer of $\ell_{0,s}(\boldsymbol{\mathcal{X}}; \underline{\boldsymbol{\varphi}})$ as $\widehat{\boldsymbol{\varphi}}^{\text{QML,S}}$.

5.1 Loadings

Despite all introduced simplifications, no closed form solution exists for the elements of the vector of QML estimators, but there exist few numerical ways to compute it. Classical iterative approaches, dealing with cross-sections of small dimension N , were proposed by, e.g., Jöreskog (1969) and Lawley and Maxwell (1971, Chapter 2), and extended to the large N case by Sundberg and Feldmann (2016). Similarly, Ng et al. (2015) adapted the Newton-Raphson maximization algorithm to the large N case by exploiting the properties of tri-diagonal blocked matrices. However, all these algorithms are heuristic and their relation to proper QML estimation is unclear. In this respect, an EM algorithm, thus in principle accomplishing QML estimation, was introduced by Rubin and Thayer (1982), and considered in the large N setting by Bai and Li (2012, 2016).

Under our assumption of an approximate factor model, Bai and Li (2016, Theorem 1) prove that

$$\left\| \widehat{\boldsymbol{\lambda}}_i^{\text{QML,S}} - \boldsymbol{\lambda}_i^{\text{OLS}} \right\| = O_P \left(\frac{1}{N} \right) + O_P \left(\frac{1}{\sqrt{NT}} \right). \quad (20)$$

Although this result is derived under the alternative identifying condition $N^{-1} \mathbf{\Lambda}' (\boldsymbol{\Sigma}^\xi)^{-1} \mathbf{\Lambda}$ being diagonal (IC3 in the original papers), it is still quite straightforward to see that it is applicable also under our identifying conditions. Moreover, in Barigozzi (2022, Theorem 4) it is shown that, under the assumptions

of this paper, the following holds:

$$\left\| \widehat{\boldsymbol{\lambda}}_i^{\text{QML,S}} - \widehat{\boldsymbol{\lambda}}_i^{\text{PC}} \right\| = O_{\text{P}} \left(\frac{1}{N} \right). \quad (21)$$

The proof of this result is based on studying the easier problem of QML estimation when using a projection of the data on an r -dimensional subspace (see also Rao, 1973, Chapter 8a, pp. 527-528).

Therefore, (21), together with (11), implies (20), and we immediately see that the QML, PC, and the unfeasible OLS estimators of the loadings are asymptotically equivalent, as long as $N \rightarrow \infty$. This is formalized in the following result, of which part (a) is an immediate consequence of (21) and Proposition 1, and part (b), which holds regardless of the imposed identification conditions, is proved by Bai and Li (2016, Theorem S.2).

Proposition 3. *For any $i = 1, \dots, N$, under Assumptions 1 through 5:*

(a) *if $\sqrt{T}/N \rightarrow 0$, as $N, T \rightarrow \infty$,*

$$\sqrt{T}(\widehat{\boldsymbol{\lambda}}_i^{\text{QML,S}} - \boldsymbol{\lambda}_i) \rightarrow_d \mathcal{N}(\mathbf{0}_r, \boldsymbol{\nu}_i^{\text{OLS}}),$$

where $\boldsymbol{\nu}_i^{\text{OLS}} := (\boldsymbol{\Gamma}^F)^{-1} \left\{ \lim_{T \rightarrow \infty} \frac{\mathbb{E}[\mathbf{F}'\boldsymbol{\zeta}_i\boldsymbol{\zeta}_i'\mathbf{F}]}{T} \right\} (\boldsymbol{\Gamma}^F)^{-1} = \lim_{T \rightarrow \infty} \frac{\mathbb{E}[\mathbf{F}'\mathbb{E}[\boldsymbol{\zeta}_i\boldsymbol{\zeta}_i'\mathbf{F}]]}{T}$ (because of Assumption 5(b));

(b) *if also Assumption 7 holds, as $N, T \rightarrow \infty$, $\min(\sqrt{T}, N) \|\widehat{\sigma}_i^{2\text{QML,S}} - \sigma_i^2\| = O_{\text{P}}(1)$.*

There are three mis-specifications that we introduced in the log-likelihood (19). First, we treat the factors as serially uncorrelated. This is, however, an innocuous mis-specification that has no effect on the asymptotic properties of the loadings. Indeed, the OLS estimator, which is asymptotically equivalent to the QML estimator, is not affected by the autocorrelation of the regressors, i.e., of the factors, as long as their covariance matrix is well defined as required by Assumptions 1(b) and 1(c).

Second, the QML and PC estimators of the loadings are not the most efficient estimators, since they both neglect the possible idiosyncratic serial correlations, hence the sandwich form of the asymptotic covariance matrix, which for the QML estimator is due to the use of a mis-specified log-likelihood, while for the PC estimator is due to its non-parametric nature. Clearly, if the idiosyncratic components were truly serially uncorrelated, then, we would have $\mathbb{E}[\boldsymbol{\zeta}_i\boldsymbol{\zeta}_i'] = \sigma_i^2 \mathbf{I}_T$, and the asymptotic covariance matrix would reduce to $\sigma_i^2 (\boldsymbol{\Gamma}^F)^{-1} = \sigma_i^2 \mathbf{I}_r$ (because of Assumption 5(b)), which is the Gauss-Markov lower bound under serial homoskedasticity.

Third, as a consequence of the fact that we are estimating an approximate factor model using the mis-specified log-likelihood (19) of an exact factor model, we would not get consistent estimators of the loadings if N were fixed. Indeed, only in the limit $N \rightarrow \infty$ we can disentangle the common components, i.e., the loadings and the factors, from the idiosyncratic components (see Lemma 1). This is reflected in the asymptotic expansion (20), where it is clear that if N is fixed then the estimation error is non-vanishing. In other words, the mis-specification error, which we introduce by using a mis-specified log-likelihood, vanishes asymptotically only if $N \rightarrow \infty$. So, despite being fully parametric, the QML estimator does not suffer of the curse of dimensionality, but, in fact, it produces consistent estimates only in a high-dimensional setting, i.e., it enjoys a blessing of dimensionality.

Remark 8. An alternative intuitive argument for the equivalence between QML and OLS is the following. Consider the decomposition of the log-likelihood (19):

$$\ell_{0,s}(\mathcal{X}; \underline{\varphi}) = \ell_{0,s}(\mathcal{X}|\mathcal{F}; \underline{\varphi}) + \ell_{0,s}(\mathcal{F}; \underline{\varphi}) - \ell_{0,s}(\mathcal{F}|\mathcal{X}; \underline{\varphi}).$$

Then, since $\|\mathcal{F}\| = O_P(\sqrt{T})$, $\|\mathbf{A}\| = O(\sqrt{N})$, and $\|\mathcal{X}\| = O_P(\sqrt{NT})$ (Barigozzi, 2022, Lemma 2), intuitively, we have $\sup_{\underline{\varphi} \in \mathcal{O}} |\ell(\mathcal{F}; \underline{\varphi})| = O_P(T)$ and $\sup_{\underline{\varphi} \in \mathcal{O}} |\ell(\mathcal{F}|\mathcal{X}; \underline{\varphi})| = O_P(T)$, while $\sup_{\underline{\varphi} \in \mathcal{O}} |\ell(\mathcal{X}; \underline{\varphi})| = O_P(NT)$ and $\sup_{\underline{\varphi} \in \mathcal{O}} |\ell(\mathcal{X}|\mathcal{F}; \underline{\varphi})| = O_P(NT)$, so that

$$\sup_{\underline{\varphi} \in \mathcal{O}} \frac{1}{NT} |\ell(\mathcal{X}; \underline{\varphi}) - \ell(\mathcal{X}|\mathcal{F}; \underline{\varphi})| \leq \frac{1}{NT} \sup_{\underline{\varphi} \in \mathcal{O}} |\ell(\mathcal{F}; \underline{\varphi})| + \frac{1}{NT} \sup_{\underline{\varphi} \in \mathcal{O}} |\ell(\mathcal{F}|\mathcal{X}; \underline{\varphi})| = O_P\left(\frac{1}{N}\right).$$

This implies that, for any given $i = 1, \dots, N$, as $N \rightarrow \infty$, the value of the loadings maximizing $\ell_{0,s}(\mathcal{X}; \underline{\varphi})$, i.e., $\hat{\lambda}_i^{\text{QML,S}}$ coincides asymptotically with the value of the loadings maximizing $\ell_{0,s}(\mathcal{X}|\mathcal{F}; \underline{\varphi})$, which is λ_i^{OLS} . This reasoning was also suggested, but not formally proved, by Breitung and Tenhofen (2011), and used by Doz et al. (2012) in their proofs.

Remark 9. Not only the full log-likelihood $\ell_{0,s}(\mathcal{X}; \underline{\varphi})$ coincides asymptotically with the conditional log-likelihood $\ell_{0,s}(\mathcal{X}|\mathcal{F}; \underline{\varphi})$, but also their derivatives with respect to the loadings coincide. Namely, Barigozzi (2022, Proposition 10) proves that:

$$\frac{1}{\sqrt{T}} \left\| \frac{\partial \ell_{0,s}(\mathcal{X}; \underline{\varphi})}{\partial \lambda_i'} \Big|_{\underline{\varphi}=\underline{\varphi}} - \frac{\partial \ell_{0,s}(\mathcal{X}|\mathcal{F}; \underline{\varphi})}{\partial \lambda_i'} \Big|_{\underline{\varphi}=\underline{\varphi}} \right\| = O_P\left(\max\left(\frac{1}{\sqrt{N}}, \frac{\sqrt{T}}{N}\right)\right);$$

$$\frac{1}{T} \left\| \frac{\partial^2 \ell_{0,s}(\mathcal{X}; \underline{\varphi})}{\partial \lambda_i' \partial \lambda_i} \Big|_{\underline{\varphi}=\underline{\varphi}} - \frac{\partial^2 \ell_{0,s}(\mathcal{X}|\mathcal{F}; \underline{\varphi})}{\partial \lambda_i' \partial \lambda_i} \Big|_{\underline{\varphi}=\underline{\varphi}} \right\| = O_P\left(\max\left(\frac{1}{N}, \frac{1}{\sqrt{NT}}\right)\right).$$

By denoting the Fisher information and the population Hessian matrices for λ_i derived from the full log-likelihood as:

$$\mathcal{I}_i(\mathcal{X}; \varphi) = \lim_{T \rightarrow \infty} \mathbb{E} \left[\left(\frac{1}{\sqrt{T}} \frac{\partial \ell_{0,s}(\mathcal{X}; \underline{\varphi})}{\partial \lambda_i'} \Big|_{\underline{\varphi}=\underline{\varphi}} \right) \left(\frac{1}{\sqrt{T}} \frac{\partial \ell_{0,s}(\mathcal{X}; \underline{\varphi})}{\partial \lambda_i} \Big|_{\underline{\varphi}=\underline{\varphi}} \right) \right],$$

$$\mathcal{H}_i(\mathcal{X}; \varphi) = \lim_{T \rightarrow \infty} \mathbb{E} \left[\frac{1}{T} \frac{\partial^2 \ell_{0,s}(\mathcal{X}; \underline{\varphi})}{\partial \lambda_i' \partial \lambda_i} \Big|_{\underline{\varphi}=\underline{\varphi}} \right],$$

respectively, we have that, as $N, T \rightarrow \infty$:

$$\mathcal{V}_i^{\text{OLS}} = (\mathcal{H}_i(\mathcal{X}; \varphi))^{-1} \mathcal{I}_i(\mathcal{X}; \varphi) (\mathcal{H}_i(\mathcal{X}; \varphi))^{-1} + o_P(1).$$

As expected, the asymptotic covariance matrix of the QML estimator of the loadings coincides asymptotically with the classical QML sandwich form which we have for a linear regression with autocorrelated and heteroskedastic residuals (White, 1980; Newey and West, 1987; Andrews, 1991).

Remark 10. It is possible to consider QML estimation based on an even simpler and further misspecified log-likelihood, where we also avoid to model idiosyncratic cross-sectional heteroskedasticities. In this case, the log-likelihood (19) is simplified by imposing $\Sigma^\xi = \sigma^2 \mathbf{I}_N$ for some $\sigma^2 > 0$. This is the

approach proposed by [Tipping and Bishop \(1999\)](#). In this case the QML estimator of the loadings has a closed form solution given by

$$\widehat{\boldsymbol{\lambda}}_i^{\text{QML},S_0} := \left(\widehat{\mathbf{M}}^x - \widehat{\sigma}^{2\text{QML},S_0} \mathbf{I}_r \right)^{1/2} \widehat{\mathbf{v}}_i^x, \quad i = 1, \dots, N, \quad (22)$$

with $\widehat{\sigma}^{2\text{QML},S_0} := (N - r)^{-1} \sum_{j=r+1}^N \widehat{\mu}_j^x$, which can be considered as an estimator of $N^{-1} \sum_{i=1}^N \sigma_i^2$. By comparing (22) with the PC estimator defined in (9), it is easy to show that (see [Barigozzi, 2022](#), Proposition 9):

$$\left\| \widehat{\boldsymbol{\lambda}}_i^{\text{QML},S_0} - \widehat{\boldsymbol{\lambda}}_i^{\text{PC}} \right\| = O_{\text{P}} \left(\frac{1}{N} \right), \quad (23)$$

which is a special case of (21). So, once again, by virtue of (11) and (23), the QML estimator coincides asymptotically with the PC and the unfeasible OLS estimators, and so it still satisfies Proposition 3.

Remark 11. The EM algorithm by [Rubin and Thayer \(1982\)](#) is the method used by [Bai and Li \(2012, 2016\)](#) compute the QML estimator of the loadings maximizing the log-likelihood (19). However, no investigation about its convergence to a (global) maximum of the log-likelihood is reported. Nor any indication is given on how to initialize such algorithm.

5.2 Factors

In this section we consider estimation of the factors when these are treated as a sequence of constant parameters. It is natural to start again from the mis-specified log-likelihood (19), where we now treat the factors as additional parameters to be estimated. Then, the log-likelihood (19) can be rewritten as (see, e.g., [Anderson and Rubin, 1956](#) and [Bai and Li, 2012](#)):

$$\ell_{0,s}(\boldsymbol{\mathcal{X}}; \boldsymbol{\mathcal{F}}, \boldsymbol{\varphi}) \simeq -\frac{T}{2} \log \det(\boldsymbol{\Sigma}^\xi) - \frac{1}{2} \sum_{t=1}^T (\mathbf{x}_t - \bar{\mathbf{x}} - \boldsymbol{\Lambda} \mathbf{F}_t)' (\boldsymbol{\Sigma}^\xi)^{-1} (\mathbf{x}_t - \bar{\mathbf{x}} - \boldsymbol{\Lambda} \mathbf{F}_t). \quad (24)$$

It is immediate to see that for known parameters $\boldsymbol{\varphi}$, the estimator of the factors maximizing (24), is the unfeasible Weighted Least Squares (WLS):

$$\mathbf{F}_t^{\text{WLS}} := \left(\boldsymbol{\Lambda}' (\boldsymbol{\Sigma}^\xi)^{-1} \boldsymbol{\Lambda} \right)^{-1} \boldsymbol{\Lambda}' (\boldsymbol{\Sigma}^\xi)^{-1} (\mathbf{x}_t - \bar{\mathbf{x}}), \quad t = 1, \dots, T. \quad (25)$$

Once we have also the QML estimator of the parameters $\widehat{\boldsymbol{\varphi}}^{\text{QML},S}$, maximizing the log-likelihood (19), we can compute (25), and obtain the feasible WLS:

$$\widehat{\mathbf{F}}_t^{\text{WLS}} := \left(\widehat{\boldsymbol{\Lambda}}^{\text{QML}'} (\widehat{\boldsymbol{\Sigma}}^{\xi\text{QML}})^{-1} \widehat{\boldsymbol{\Lambda}}^{\text{QML}} \right)^{-1} \widehat{\boldsymbol{\Lambda}}^{\text{QML}'} (\widehat{\boldsymbol{\Sigma}}^{\xi\text{QML}})^{-1} (\mathbf{x}_t - \bar{\mathbf{x}}), \quad t = 1, \dots, T, \quad (26)$$

which is nothing else but the classical “least-squares” estimator of the factors originally proposed by [Bartlett \(1937\)](#) (see also [Thomson, 1936](#), [Bartlett, 1938](#), and [Lawley and Maxwell, 1971](#), Chapter 7.2). In the large N case, [Bai and Li \(2012, Theorem 6.1\)](#) and [Bai and Li \(2016, Theorem 2\)](#) prove that:

$$\left\| \widehat{\mathbf{F}}_t^{\text{WLS}} - \mathbf{F}_t^{\text{WLS}} \right\| = O_{\text{P}} \left(\frac{1}{T} \right) + O_{\text{P}} \left(\frac{1}{\sqrt{NT}} \right). \quad (27)$$

The following result follows.

Proposition 4. For any $t = 1, \dots, T$, under Assumptions 1 through 7 if $\sqrt{N}/T \rightarrow 0$, as $N, T \rightarrow \infty$,

$$\sqrt{N}(\widehat{\mathbf{F}}_t^{\text{WLS}} - \mathbf{F}_t) \rightarrow_d \mathcal{N}(\mathbf{0}_r, \mathbf{W}_t^{\text{WLS}}),$$

where $\mathbf{W}_t^{\text{WLS}} := (\boldsymbol{\Sigma}_{\Lambda\xi\Lambda})^{-1} \left\{ \lim_{N \rightarrow \infty} \frac{\boldsymbol{\Lambda}'(\boldsymbol{\Sigma}^\xi)^{-1} \mathbb{E}[\boldsymbol{\xi}_t \boldsymbol{\xi}_t'] (\boldsymbol{\Sigma}^\xi)^{-1} \boldsymbol{\Lambda}}{N} \right\} (\boldsymbol{\Sigma}_{\Lambda\xi\Lambda})^{-1}$, with $\boldsymbol{\Sigma}_{\Lambda\xi\Lambda} := \lim_{N \rightarrow \infty} \frac{\boldsymbol{\Lambda}'(\boldsymbol{\Sigma}^\xi)^{-1} \boldsymbol{\Lambda}}{N}$.

If instead we treat the factors as random variables, but we do not model their dynamics, then their optimal (in mean-squared sense) linear estimator is the linear projection of the true factors onto the observed data. Given that we are considering a mis-specified exact static factor model, such linear projection can be approximated by:

$$\mathbf{F}_t^{\text{LP}} := \left(\boldsymbol{\Lambda}'(\boldsymbol{\Sigma}^\xi)^{-1} \boldsymbol{\Lambda} + \mathbf{I}_r \right)^{-1} \boldsymbol{\Lambda}'(\boldsymbol{\Sigma}^\xi)^{-1} (\mathbf{x}_t - \bar{\mathbf{x}}), \quad t = 1, \dots, T. \quad (28)$$

Once we have also the QML estimator of the parameters $\widehat{\boldsymbol{\varphi}}^{\text{QML},S}$ maximizing the log-likelihood (19) of a static factor model. In this way, we obtain the feasible Linear Projection (LP) estimator

$$\widehat{\mathbf{F}}_t^{\text{LP}} := \left(\widehat{\boldsymbol{\Lambda}}^{\text{QML},S'} (\widehat{\boldsymbol{\Sigma}}^{\xi\text{QML},S})^{-1} \widehat{\boldsymbol{\Lambda}}^{\text{QML},S} + \mathbf{I}_r \right)^{-1} \widehat{\boldsymbol{\Lambda}}^{\text{QML},S'} (\widehat{\boldsymbol{\Sigma}}^{\xi\text{QML},S})^{-1} (\mathbf{x}_t - \bar{\mathbf{x}}), \quad t = 1, \dots, T, \quad (29)$$

which is nothing else but the classical ‘‘regression’’ estimator of the factors proposed by Thomson (1951) (see also Scott, 1966, and Lawley and Maxwell, 1971, Chapter 7.4). Clearly, the unfeasible LP estimator (28) has always a smaller MSE than the unfeasible WLS estimator (25). Nevertheless, since because of Proposition 3, and Assumptions 1(a) and 2(a), $\|\widehat{\boldsymbol{\Lambda}}^{\text{QML},S}\| = O_P(\sqrt{N})$ and $\|(\widehat{\boldsymbol{\Sigma}}^{\xi\text{QML},S})^{-1}\| = O_P(1)$, then, the feasible WLS and LP estimators coincide asymptotically (see also Bai and Li, 2012, Proposition 6.1):

$$\left\| \widehat{\mathbf{F}}_t^{\text{WLS}} - \widehat{\mathbf{F}}_t^{\text{LP}} \right\| = O_P\left(\frac{1}{N}\right). \quad (30)$$

The following result follows.

Proposition 5. For any $t = 1, \dots, T$, under Assumptions 1 through 7 if $\sqrt{N}/T \rightarrow 0$, as $N, T \rightarrow \infty$,

$$\sqrt{N}(\widehat{\mathbf{F}}_t^{\text{LP}} - \mathbf{F}_t) \rightarrow_d \mathcal{N}(\mathbf{0}_r, \mathbf{W}_t^{\text{WLS}}),$$

where $\mathbf{W}_t^{\text{WLS}} := (\boldsymbol{\Sigma}_{\Lambda\xi\Lambda})^{-1} \left\{ \lim_{N \rightarrow \infty} \frac{\boldsymbol{\Lambda}'(\boldsymbol{\Sigma}^\xi)^{-1} \mathbb{E}[\boldsymbol{\xi}_t \boldsymbol{\xi}_t'] (\boldsymbol{\Sigma}^\xi)^{-1} \boldsymbol{\Lambda}}{N} \right\} (\boldsymbol{\Sigma}_{\Lambda\xi\Lambda})^{-1}$, with $\boldsymbol{\Sigma}_{\Lambda\xi\Lambda} := \lim_{N \rightarrow \infty} \frac{\boldsymbol{\Lambda}'(\boldsymbol{\Sigma}^\xi)^{-1} \boldsymbol{\Lambda}}{N}$.

The asymptotic covariance matrix of the estimated factors is the same for both the WLS and LP estimator. It has a sandwich form reflecting the mis-specification of the cross-sectional correlation structure of the idiosyncratic components. It differs from the asymptotic covariance of the PC estimator given in Proposition 2, which is an OLS-type estimator. Notice that, in general, under our assumptions of an approximate factor model, we cannot say if the WLS/LP estimators are more or less efficient than the PC estimator, i.e., the matrix $\mathbf{W}_t^{\text{OLS}} - \mathbf{W}_t^{\text{WLS}}$ is neither positive nor negative definite. The only thing we can say is that neither the WLS/LP nor the PC estimator are the most efficient, the efficiency loss of the former being due to the use of a mis-specified log-likelihood, and the efficiency loss of the latter being due to its non-parametric nature.

Clearly, if the true model were an exact factor model then we would have $\mathbb{E}[\boldsymbol{\xi}_t \boldsymbol{\xi}_t'] = \boldsymbol{\Sigma}^\xi$, and $\mathbf{W}_t^{\text{WLS}}$ would reduce to $(\boldsymbol{\Sigma}_{\Lambda\xi\Lambda})^{-1}$, which is the Gauss-Markov lower bound under cross-sectional heteroskedas-

ticity. In this case the WLS/LP estimators are always more efficient than the PC estimator, since the latter, being non-parametric does not even account explicitly for idiosyncratic cross-sectional heteroskedasticity.

It is clear from Propositions 4 and 5, that in the fixed N case no consistency can be proved for the estimated factors. Indeed, for fixed N there is no hope of estimating the whole rT -dimensional vector of factors \mathbf{F} consistently, simply due to a lack of degrees of freedom. Moreover, for any given $t = 1, \dots, T$, the WLS/LP estimators of the factors in (25) are the result of the aggregation of data along the cross-sectional dimension, i.e., they are the result of a cross-sectional regression. Hence, for consistency we must require $N \rightarrow \infty$.⁴ This is another manifestation of the blessing of dimensionality.

In fact, from (27) it is clear that we must have also $T \rightarrow \infty$ in order to consistently estimate the factors. This is because the feasible WLS/LP estimators in (26) and (29) require the use of the QML estimator of the loadings and the idiosyncratic variances, which are consistent if and only if $T \rightarrow \infty$ (see Proposition 3).

Remark 12. Two other estimators of the factors are worth recalling. First, if we compute the WLS estimator in (25) using the PC estimators of the loadings in (9) and of the idiosyncratic variances⁵, we get the Generalized PC estimator of the factors studied by [Breitung and Tenhofen \(2011\)](#) and [Choi \(2012\)](#), which, by virtue of (21), is asymptotically equivalent to the feasible WLS estimator in (26). Second, if we further simplify the log-likelihood (24) by treating the idiosyncratic components as cross-sectionally homoskedastic (see Remark 10), and we use the QML estimator of the loadings in (22) to compute the WLS estimator in (25), we obtain a feasible OLS estimator of the factors which, by (23), is asymptotically equivalent to the PC estimator defined in (12).

6 QML estimation of an approximate dynamic factor model

In this section we review the case in which the dynamics of the factors is explicitly modeled, so we consider an approximate dynamic factor model described by the linear system:

$$x_{it} = \alpha_i + \lambda_i' \mathbf{F}_t + \xi_{it}, \quad i = 1, \dots, N, \quad t = 1, \dots, T, \quad (31)$$

$$\mathbf{F}_t = \mathbf{A}\mathbf{F}_{t-1} + \mathbf{H}\mathbf{u}_t, \quad (32)$$

together with Assumptions 1 through 6. In this case, $\{\mathbf{F}_t\}$ is an autocorrelated stochastic process and Ω^F is a function of \mathbf{A} and \mathbf{H} .

For a dynamic factor model the log-likelihood (18) can be decomposed as:

$$\ell_{0,D}(\mathcal{X}; \underline{\varphi}) = \ell_{0,D}(\mathcal{X}|\mathcal{F}; \underline{\varphi}) + \ell_{0,D}(\mathcal{F}; \underline{\varphi}) - \ell_{0,D}(\mathcal{F}|\mathcal{X}; \underline{\varphi}), \quad (33)$$

⁴Notice that although the PC estimator of the factors proposed by [Bai \(2003\)](#) is just an eigenvector so no cross-sectional aggregation is required, we still require $N \rightarrow \infty$ to prove its consistency. Indeed, in this case we need to compute eigenvectors of a $T \times T$ covariance matrix, which is consistently estimating the true covariance only if $N \rightarrow \infty$.

⁵The PC estimator of the idiosyncratic variances is $T^{-1} \sum_{t=1}^T (x_{it} - \hat{\lambda}_i^{\text{PC}'} \hat{\mathbf{F}}_t^{\text{PC}})^2$.

where (recall that $\mathbf{F}_0 = \mathbf{0}_r$ because of Assumption 6(g))

$$\begin{aligned}\ell_{0,\text{D}}(\mathcal{X}|\mathcal{F}; \underline{\varphi}) &\simeq -\frac{T}{2} \log \det(\underline{\Sigma}^\xi) - \frac{1}{2} \sum_{t=1}^T (\mathbf{x}_t - \bar{\mathbf{x}} - \underline{\mathbf{A}}\mathbf{F}_t)' (\underline{\Sigma}^\xi)^{-1} (\mathbf{x}_t - \bar{\mathbf{x}} - \underline{\mathbf{A}}\mathbf{F}_t), \\ \ell_{0,\text{D}}(\mathcal{F}; \underline{\varphi}) &\simeq -\frac{T}{2} \log \det(\underline{\mathbf{H}}\underline{\mathbf{H}}') - \frac{1}{2} \sum_{t=1}^T (\mathbf{F}_t - \underline{\mathbf{A}}\mathbf{F}_{t-1})' (\underline{\mathbf{H}}\underline{\mathbf{H}}')^{-1} (\mathbf{F}_t - \underline{\mathbf{A}}\mathbf{F}_{t-1}), \\ \ell_{0,\text{D}}(\mathcal{F}|\mathcal{X}; \underline{\varphi}) &\simeq -\frac{1}{2} \sum_{t=1}^T \log \det \left(\text{Var}_{\underline{\varphi}}(\mathbf{F}_t|\mathcal{X}) \right) - \frac{1}{2} \sum_{t=1}^T (\mathbf{F}_t - \mathbb{E}_{\underline{\varphi}}[\mathbf{F}_t|\mathcal{X}])' \left(\text{Var}_{\underline{\varphi}}(\mathbf{F}_t|\mathcal{X}) \right)^{-1} (\mathbf{F}_t - \mathbb{E}_{\underline{\varphi}}[\mathbf{F}_t|\mathcal{X}]),\end{aligned}\tag{34}$$

with $\text{Var}_{\underline{\varphi}}(\mathbf{F}_t|\mathcal{X}) = \mathbb{E}_{\underline{\varphi}}[(\mathbf{F}_t - \mathbb{E}_{\underline{\varphi}}[\mathbf{F}_t|\mathcal{X}])(\mathbf{F}_t - \mathbb{E}_{\underline{\varphi}}[\mathbf{F}_t|\mathcal{X}])'|\mathcal{X}]$. The parameters to be estimated are $\underline{\varphi} = (\underline{\phi}', \underline{\theta}')$ with $\underline{\phi} = (\text{vec}(\underline{\mathbf{A}})', \sigma_1^2, \dots, \sigma_N^2)'$ and $\underline{\theta} = (\text{vec}(\underline{\mathbf{A}})', \text{vec}(\underline{\mathbf{H}})')$, so $\underline{\varphi}$ is a $Q = (r+1)N + 2r^2$ -dimensional vector. We shall denote the maximizer of $\ell_{0,\text{D}}(\mathcal{X}; \underline{\varphi})$ as $\widehat{\underline{\varphi}}^{\text{QML,D}}$.

We study QML estimation of the approximate dynamic factor model under the following additional assumption.

Assumption 8.

- (a) For all $s > 0$, all $t = 1, \dots, T$, all $T \in \mathbb{N}$, and all $j = 1, \dots, r$, $\text{P}(|F_{jt}| > s) \leq \exp\{-K_F s^{\delta_F}\}$ for some finite positive reals K_F and $\delta_F \leq 2$ independent of t and j .
- (b) For all $s > 0$, all $t = 1, \dots, T$, all $T \in \mathbb{N}$, all $i = 1, \dots, N$, and all $N \in \mathbb{N}$, $\text{P}(|\xi_{it}| > s) \leq \exp\{-K_\xi s^{\delta_\xi}\}$ for some finite positive reals K_ξ and $\delta_\xi \leq 2$ independent of t and i .
- (c) For all $s > 0$, all $t = 1, \dots, T$, all $T \in \mathbb{N}$, and all $N \in \mathbb{N}$,

$$\text{P} \left(\left\| \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{\lambda_i \xi_{it}}{[\Gamma^\xi]_{ii}} \right\| > s \right) \leq r \exp\{-\kappa_1 s^2\} + rN \exp\left\{-\kappa_2 \left(s\sqrt{N}\right)^\alpha\right\},$$

for some finite positive reals κ_1, κ_2 , and $\alpha \leq 2$ independent of t and N .

- (d) For all $s > 0$, all $i = 1, \dots, N$, all $N \in \mathbb{N}$, and all $T \in \mathbb{N}$,

$$\text{P} \left(\left\| \frac{1}{\sqrt{T}} \sum_{t=1}^T \mathbf{F}_t \xi_{it} \right\| > s \right) \leq r \exp\{-\kappa_3 s^2\} + rT \exp\left\{-\kappa_4 \left(s\sqrt{T}\right)^\beta\right\},$$

for some finite positive reals κ_3, κ_4 , and $\beta \leq 2$ independent of i and T .

Assumption 8 is a high-level one and it deserves few comments. Parts (a) and (b) are also assumed by Fan et al. (2013) for PC estimation. Although they imply that all moments exist, depending on the value of δ_F and δ_ξ , we are able to consider not only distributions with sub-Gaussian ($\delta_F = \delta_\xi = 2$) or sub-exponential tails ($\delta_F = \delta_\xi = 1$), which include the Laplace and the Generalized Error distribution, but also distributions with sub-Weibull tails ($\delta_F < 1, \delta_\xi < 1$), which, in finite samples, can generate a heavy tail behavior comparable to the power-law behavior, see, e.g., Mikosch and Nagaev (1998) and Kuchibhotla and Chakraborty (2022).

Part (c) is a cross-sectional Bernstein type inequality and it can be shown to hold if $\{\xi_{it}\}$ is independent across i , see, e.g., Bakhshizadeh et al. (2020, Corollary 4). More in general, it could be derived if we introduced some notion of dependence for random fields (see the comments after Assumption 4). Part (d) is a time series Bernstein type inequality and it could be derived for example if we assumed

$\{\mathbf{F}_t\}$ and $\{\xi_{it}\}$ to be strong mixing processes with exponentially decaying mixing coefficients, see, e.g., Merlevède et al. (2011, Theorem 1).

6.1 Loadings

The classical approach to estimate a dynamic factor model by QML is based on the maximization of the prediction error log-likelihood which depends on the linear prediction of the factors, usually computed via the Kalman filter, which is defined in the next section. This approach is equivalent to maximizing the log-likelihood (33) (see, e.g., Hannan and Deistler, 2012, Chapter 4). Although there is no analytic solution for the QML estimator of the parameters obtained in this way, numerical maximization is standard practice at least for low dimensional exact dynamic factor models, see, e.g., Stock and Watson (1989, 1991) and Harvey (1990, Chapter 3.4). In high-dimensions Jungbacker and Koopman (2015) propose to maximize the log-likelihood (33) subject to a preliminary step in which the data are projected onto a lower dimensional space. It is not clear, however, what are the effects of this first step on the asymptotic properties of the final estimators.

In high-dimensions maximization of the log-likelihood (33) is usually achieved by means of EM algorithm, which is an iterative procedure proposed by Dempster et al. (1977) as a general way to achieve QML estimation when dealing with incomplete information (see also Sundberg, 1974, 1976, and Sundberg, 2019, Chapter 8, for EM and incomplete data in exponential families). The use of the EM algorithm for estimating small dimensional state space models dates back to Shumway and Stoffer (1982), Watson and Engle (1983), Harvey and Peters (1990), and Ghahramani and Hinton (1996), while its use to estimate a high-dimensional approximate dynamic factor model was first suggested by Quah and Sargent (1993) and then formalized by Doz et al. (2012), while its theoretical properties are studied by Barigozzi and Luciani (2019).

In the present context, the EM algorithm has two main features: (i) it gives closed form solutions for the estimated parameters, and (ii) it runs in few iterations. In a nutshell, it works as follows. Consider a given iteration $k \geq 0$ of the EM algorithm and let us assume to have an estimate of the parameters $\hat{\varphi}^{(k)}$. Then, for each k repeat two steps.

E-step Compute the expected full-information log-likelihood with respect to the conditional distribution of the factors given the data and computed using the estimated parameters $\hat{\varphi}^{(k)}$:

$$Q(\underline{\varphi}, \hat{\varphi}^{(k)}) := \mathbb{E}_{\hat{\varphi}^{(k)}}[\ell_{0,D}(\mathcal{X}, \mathcal{F}; \underline{\varphi}) | \mathcal{X}];$$

M-step Compute a new estimate of the parameters maximizing the expected full-information log-likelihood:

$$\hat{\varphi}^{(k+1)} = \arg \max_{\underline{\varphi}} Q(\underline{\varphi}, \hat{\varphi}^{(k)}).$$

The algorithm is initialized and terminated as follows.

Initialization The loadings are initialized with their PC estimator defined in (9), then, given also the PC estimates of the factors and of the idiosyncratic components, we estimate: (i) the VAR parameters, and (ii) the idiosyncratic variances.

Stopping The iterations are stopped at iteration k^* such that the log-likelihoods $\ell_{0,D}(\mathcal{X}; \hat{\varphi}^{(k^*+1)})$ and $\ell_{0,D}(\mathcal{X}; \hat{\varphi}^{(k^*)})$ do not differ more than a pre-specified threshold. We define the EM estimator of the parameters as $\hat{\varphi}^{\text{EM}} := \hat{\varphi}^{(k^*+1)}$.

The rationale for the EM algorithm is the following. By taking the conditional expectations of the log-likelihood (33) and computed using $\hat{\varphi}^{(k)}$, we get:

$$\begin{aligned} \ell_{0,D}(\mathcal{X}; \underline{\varphi}) &= \mathbb{E}_{\hat{\varphi}^{(k)}}[\ell_{0,D}(\mathcal{X}, \mathcal{F}; \underline{\varphi}) | \mathcal{X}] - \mathbb{E}_{\hat{\varphi}^{(k)}}[\ell_{0,D}(\mathcal{F} | \mathcal{X}; \underline{\varphi}) | \mathcal{X}] \\ &=: \mathcal{Q}(\underline{\varphi}, \hat{\varphi}^{(k)}) - \mathcal{H}(\underline{\varphi}, \hat{\varphi}^{(k)}), \text{ say.} \end{aligned} \quad (35)$$

The QML estimator of φ is then a maximum of the right hand side of (35). In fact, since, by definition of Kullback-Leibler divergence, $\mathcal{H}(\underline{\varphi}, \hat{\varphi}^{(k)}) \leq \mathcal{H}(\hat{\varphi}^{(k)}, \hat{\varphi}^{(k)})$ for any pair $(\underline{\varphi}, \hat{\varphi}^{(k)})$ and any $k \geq 0$ (see Dempster et al., 1977, Lemma 1, and Wu, 1983, Theorem 3), it is enough to maximize $\mathcal{Q}(\underline{\varphi}, \hat{\varphi}^{(k)})$. It follows that the EM algorithm defines a continuous path in the parameter space from the starting point to the stopping point along which the log-likelihood monotonically increases without leaping over valleys.

In general, the implementation of the EM steps might be non-trivial. However, things simplify considerably in the present setup. First, the full-information expected log-likelihood $\mathcal{Q}(\underline{\varphi}, \hat{\varphi}^{(k)})$ can be further decomposed as:

$$\mathcal{Q}(\underline{\varphi}, \hat{\varphi}^{(k)}) = \mathbb{E}_{\hat{\varphi}^{(k)}}[\ell_{0,D}(\mathcal{X} | \mathcal{F}; \underline{\varphi}) | \mathcal{X}] + \mathbb{E}_{\hat{\varphi}^{(k)}}[\ell_{0,D}(\mathcal{F}; \underline{\varphi}) | \mathcal{X}]. \quad (36)$$

Then, it is clear from (34) that, in order to compute the expected full information log-likelihood in the E-step, we just need to compute the conditional moments of the factors: $\mathbb{E}_{\hat{\varphi}^{(k)}}[\mathbf{F}_t | \mathcal{X}]$, $\mathbb{E}_{\hat{\varphi}^{(k)}}[\mathbf{F}_t \mathbf{F}'_t | \mathcal{X}]$, and $\mathbb{E}_{\hat{\varphi}^{(k)}}[\mathbf{F}_t \mathbf{F}'_{t-1} | \mathcal{X}]$. And, in turn, these moments can be obtained by means of the Kalman smoother implemented using the estimated parameters $\hat{\varphi}^k$, as detailed in the next section.

The M-step has then a closed form solution. Indeed, from (34) we immediately see that the final EM estimator of the parameters, ϕ , in the measurement equation (31) is:

$$\begin{aligned} \hat{\lambda}_i^{\text{EM}} &= \left(\sum_{t=1}^T \mathbb{E}_{\hat{\varphi}^{(k^*)}}[\mathbf{F}_t \mathbf{F}'_t | \mathcal{X}] \right)^{-1} \left(\sum_{t=1}^T \mathbb{E}_{\hat{\varphi}^{(k^*)}}[\mathbf{F}_t | \mathcal{X}] (x_{it} - \bar{x}_i) \right), \quad i = 1, \dots, N, \quad (37) \\ \hat{\sigma}_i^{2\text{EM}} &= \frac{1}{T} \sum_{t=1}^T \left\{ (x_{it} - \bar{x}_i)^2 + \hat{\lambda}_i^{\text{EM}'} \left(\mathbb{E}_{\hat{\varphi}^{(k^*)}}[\mathbf{F}_t \mathbf{F}'_t | \mathcal{X}] \right) \hat{\lambda}_i^{\text{EM}} \right. \\ &\quad \left. - 2(x_{it} - \bar{x}_i) \hat{\lambda}_i^{\text{EM}'} \mathbb{E}_{\hat{\varphi}^{(k^*)}}[\mathbf{F}_t | \mathcal{X}] \right\}, \quad i = 1, \dots, N, \quad (38) \end{aligned}$$

such that $\hat{\Lambda}^{\text{EM}} := (\hat{\lambda}_1^{\text{EM}} \dots \hat{\lambda}_N^{\text{EM}})'$ is the estimated matrix of all loadings and $\hat{\Sigma}^{\xi\text{EM}} := \text{diag}(\hat{\sigma}_1^{2\text{EM}}, \dots, \hat{\sigma}_N^{2\text{EM}})$ is the estimated diagonal matrix of idiosyncratic variances.

Notice that, since to compute the estimators in (37) and (38) we use the mis-specified expected conditional log-likelihood (36) of an exact factor model, we reduce a potentially hard maximizations problem of estimating a high-dimensional parameter vector ϕ into the maximizations of N expected log-likelihoods, each depending on a finite $r + 1$ -dimensional parameter vector $(\text{vec}(\lambda_i)', \sigma_i^2)'$. This can be seen as an extreme form of regularization of the idiosyncratic covariance matrix which makes estimation of the large dimensional parameter vector ϕ straightforward.

Closed form expressions are immediately derived also for the final estimators of the parameters, $\boldsymbol{\theta}$, in the state equation (32):

$$\widehat{\mathbf{A}}^{\text{EM}} = \left(\sum_{t=1}^T \mathbb{E}_{\widehat{\boldsymbol{\varphi}}^{(k^*)}} [\mathbf{F}_t \mathbf{F}'_{t-1} | \boldsymbol{\mathcal{X}}] \right) \left(\sum_{t=2}^T \mathbb{E}_{\widehat{\boldsymbol{\varphi}}^{(k^*)}} [\mathbf{F}_{t-1} \mathbf{F}'_{t-1} | \boldsymbol{\mathcal{X}}] \right)^{-1}, \quad (39)$$

$$\widehat{\mathbf{H}}^{\text{EM}} = \left[\frac{1}{T} \sum_{t=1}^T \left\{ \mathbb{E}_{\widehat{\boldsymbol{\varphi}}^{(k^*)}} [\mathbf{F}_t \mathbf{F}'_t | \boldsymbol{\mathcal{X}}] - \left(\mathbb{E}_{\widehat{\boldsymbol{\varphi}}^{(k^*)}} [\mathbf{F}_{t-1} \mathbf{F}'_t | \boldsymbol{\mathcal{X}}] \right) \widehat{\mathbf{A}}^{\text{EM}} \right\} \right]^{1/2}. \quad (40)$$

For a log-likelihood belonging to the exponential family, as the Gaussian one, it is possible to prove that the EM algorithm produces a sequence of estimators $\{\widehat{\boldsymbol{\lambda}}_i^{(k)}\}$ which converges to a local maximum of the log-likelihood (33) as $k \rightarrow \infty$, say $\widehat{\boldsymbol{\lambda}}_i^{(\infty)}$ (Wu, 1983, Theorem 6). It is then common practice to consider the output of the EM algorithm $\widehat{\boldsymbol{\lambda}}_i^{\text{EM}}$ as equivalent to the QML estimator $\widehat{\boldsymbol{\lambda}}_i^{\text{QML,D}}$. This, however, is not always true, because there are two additional sources of errors: (i) the local maximum might not be the global maximum, and (ii) the EM algorithm runs for a finite number of iterations k^* .

The two aforementioned sources of error are addressed by Barigozzi and Luciani (2019, Proposition 2) who prove that $\widehat{\boldsymbol{\lambda}}_i^{\text{EM}}$ is asymptotically equivalent to the QML estimator $\widehat{\boldsymbol{\lambda}}_i^{\text{QML,D}}$ maximizing the full log-likelihood (33), which, in turn, is asymptotically equivalent to the QML estimator $\widehat{\boldsymbol{\lambda}}_i^{\text{QML,S}}$ obtained by maximizing the log-likelihood of a static factor model (19). Namely,

$$\left\| \widehat{\boldsymbol{\lambda}}_i^{(\infty)} - \widehat{\boldsymbol{\lambda}}_i^{\text{QML,D}} \right\| = O_{\text{P}} \left(\frac{1}{N} \right) + O_{\text{P}} \left(\frac{1}{T} \right) + O_{\text{P}} \left(\frac{1}{\sqrt{NT}} \right), \quad (41)$$

and (up to logarithmic terms)

$$\left\| \widehat{\boldsymbol{\lambda}}_i^{\text{EM}} - \widehat{\boldsymbol{\lambda}}_i^{(\infty)} \right\| = O_{\text{P}} \left(\frac{1}{N} \right) + O_{\text{P}} \left(\frac{1}{T} \right) + O_{\text{P}} \left(\frac{1}{\sqrt{NT}} \right), \quad (42)$$

$$\left\| \widehat{\boldsymbol{\lambda}}_i^{\text{QML,D}} - \widehat{\boldsymbol{\lambda}}_i^{\text{QML,S}} \right\| = O_{\text{P}} \left(\frac{1}{N} \right). \quad (43)$$

In particular, (41) is a consequence of the fact that we initialize the EM algorithm with the PC estimator which is fully identified under our Assumption 5(b), so the sequence $\{\widehat{\boldsymbol{\lambda}}_i^{(k)}\}$ actually converges to the global identified maximum $\widehat{\boldsymbol{\lambda}}_i^{\text{QML,D}}$ (see also Ruud, 1991, Section 4, for a similar result in the case of one-to-one mapping from the factors to the data, corresponding to the case of no idiosyncratic component). Moreover, (42) follows from the fact that the numerical error is proportional to the information loss between using the expected joint log-likelihood in (36) instead of the full log-likelihood (33), i.e., it depends on the fraction of missing information due to the fact that the factors are not observed (Sundberg, 1976, Meng and Rubin, 1994, and McLachlan and Krishnan, 2007, Chapter 3.9). Last, (43) follows from the fact that the exact factor model log-likelihood (19) and the approximate factor model log-likelihood (33) are both asymptotically equivalent to the log-likelihood of conditional on the factors which is the same in both cases, i.e., $\ell_{0,\text{S}}(\boldsymbol{\mathcal{X}} | \boldsymbol{\mathcal{F}}; \underline{\boldsymbol{\varphi}}) = \ell_{0,\text{D}}(\boldsymbol{\mathcal{X}} | \boldsymbol{\mathcal{F}}; \underline{\boldsymbol{\varphi}})$, so asymptotically their maxima should coincide and they both coincide with the unfeasible OLS estimator (see also Remark 8).

From (41), (42), and (43), jointly with Proposition 3, it follows that.

Proposition 6. *For any $i = 1, \dots, N$, under Assumptions 1 through 8:*

(a) if $\sqrt{T} \log^\kappa T/N \rightarrow 0$ with $\kappa := \max(\delta_F^{-1}, \delta_\xi^{-1}) \geq 1/2$, as $N, T \rightarrow \infty$,

$$\sqrt{T}(\widehat{\boldsymbol{\lambda}}_i^{\text{EM}} - \boldsymbol{\lambda}_i) \rightarrow_d \mathcal{N}(\mathbf{0}_r, \boldsymbol{\nu}_i^{\text{OLS}}),$$

where $\boldsymbol{\nu}_i^{\text{OLS}} = (\boldsymbol{\Gamma}^F)^{-1} \left\{ \lim_{T \rightarrow \infty} \frac{\mathbb{E}[\mathbf{F}'_i \boldsymbol{\zeta}_i \boldsymbol{\zeta}'_i \mathbf{F}_i]}{T} \right\} (\boldsymbol{\Gamma}^F)^{-1} = \lim_{T \rightarrow \infty} \frac{\mathbb{E}[\mathbf{F}'_i \boldsymbol{\zeta}_i \boldsymbol{\zeta}'_i \mathbf{F}_i]}{T}$ (because of Assumption 5(b));

(b) as $N, T \rightarrow \infty$, $\min(\sqrt{T/\log N}, \sqrt{N}) \|\widehat{\sigma}_i^{2EM} - \sigma_i^2\| = O_P(1)$.

Up to logarithmic terms, the EM estimator of the loadings has the same asymptotic properties of the QML estimator of a static factor model. Therefore, the same comments made after Proposition 3 about efficiency apply also in this case. Moreover, the EM estimator of the loadings has also the same asymptotic properties of the PC estimator. This is not surprising since the EM algorithm is initialized with the PC estimator, and, therefore, $\widehat{\boldsymbol{\lambda}}_i^{\text{EM}}$ can be seen as a one-step estimator, which, by construction, must also be consistent and as efficient as the initial estimator (see, e.g., van der Vaart, 2000, Section 5.7, and Lehmann and Casella, 2006, Theorem 4.3).

6.2 Factors

In a dynamic factor models, the factors are explicitly treated as autocorrelated random variables, thus, their optimal linear predictor is the linear projection of the true factors onto the all available data which is collected into the NT -dimensional vector $\boldsymbol{\mathcal{X}}$. For generic values of the parameters and any given $t = 1, \dots, T$, such prediction is given by the linear projection: $\text{Proj}_{\boldsymbol{\varphi}}(\mathbf{F}_t | \boldsymbol{\mathcal{X}}) =: \boldsymbol{\mathcal{P}}'_{\boldsymbol{\varphi}, t} (\boldsymbol{\mathcal{X}} - \bar{\boldsymbol{\mathcal{X}}})$, where $\boldsymbol{\mathcal{P}}_{\boldsymbol{\varphi}, t}$ is an $NT \times r$ matrix, which depends on $\boldsymbol{\varphi}$, and such that it minimizes the mean-squared error (MSE):

$$\boldsymbol{\mathcal{P}}_{\boldsymbol{\varphi}, t} = \underset{\boldsymbol{\mathcal{P}}_t \in \mathbb{R}^{r \times NT}}{\text{argmin}} \mathbb{E}_{\boldsymbol{\varphi}} \left[(\mathbf{F}_t - \boldsymbol{\mathcal{P}}'_t (\boldsymbol{\mathcal{X}} - \bar{\boldsymbol{\mathcal{X}}})) (\mathbf{F}_t - \boldsymbol{\mathcal{P}}'_t (\boldsymbol{\mathcal{X}} - \bar{\boldsymbol{\mathcal{X}}}))' \right], \quad t = 1, \dots, T. \quad (44)$$

By solving (44) for known parameters $\boldsymbol{\varphi}$, and treating again the idiosyncratic components as if they were serially and cross-sectionally uncorrelated, we get:

$$\begin{aligned} \mathbf{F}_t^{\text{WK}} &:= (\boldsymbol{\iota}'_t \otimes \mathbf{I}_r) \left(\mathbf{I}_T \otimes \left(\boldsymbol{\Lambda}' (\boldsymbol{\Sigma}^\xi)^{-1} \boldsymbol{\Lambda} \right) + (\boldsymbol{\Omega}^F)^{-1} \right)^{-1} \left(\mathbf{I}_T \otimes \boldsymbol{\Lambda}' (\boldsymbol{\Sigma}^\xi)^{-1} \right) (\boldsymbol{\mathcal{X}} - \bar{\boldsymbol{\mathcal{X}}}) \\ &= (\boldsymbol{\iota}'_t \otimes \mathbf{I}_r) \left(\mathbf{I}_T \otimes \left(\boldsymbol{\Lambda}' (\boldsymbol{\Sigma}^\xi)^{-1} \boldsymbol{\Lambda} \right) + (\boldsymbol{\Omega}^F)^{-1} \right)^{-1} \left(\mathbf{I}_T \otimes \boldsymbol{\Lambda}' (\boldsymbol{\Sigma}^\xi)^{-1} \right) \begin{pmatrix} (\mathbf{x}_1 - \bar{\mathbf{x}})' \\ \vdots \\ (\mathbf{x}_T - \bar{\mathbf{x}})' \end{pmatrix}, \quad t = 1, \dots, T, \end{aligned} \quad (45)$$

which is nothing else but the unfeasible estimator obtained by taking the inverse Fourier transform of the smoother originally proposed by Wiener (1949) and Kolmogorov (1941).

Now since $\{\mathbf{F}_t\}$ is an autocorrelated process, $\boldsymbol{\Omega}^F$ is not diagonal and at each point in time the estimator of the factors (45) is a weighted average of all T present, past, and future values of all N time series. Hence, differently from the PC, WLS, and LP estimators, this estimator is obtained not only by cross-sectional aggregation, but also by temporal aggregation of the data, thus effectively taking into account all its 2nd order dependencies. Given that (45) uses all available history of $\{\mathbf{x}_t\}$, it is also called a smoother estimator, rather than a filter which, at a given point in time t , would use only past and present information.

Direct implementation of (45) is still unfeasible due to the presence of the $rT \times rT$ full matrix, $\mathbf{\Omega}^F$ which needs to be estimated and inverted.⁶ There are two main solutions to this problem. First, we could ignore the autocorrelation of the factors and impose $\mathbf{\Omega}^F = \mathbf{I}_{rT}$, because of Assumption 5(b), and then (45) would become the LP estimator defined in (29).

Second, in the spirit of this section on estimation of a dynamic factor model, we can use a parametric model for the dynamics of the factors, so that $\mathbf{\Omega}^F$ is parametrized accordingly. In this case, we can estimate the factors by means of the well-known iterative procedures proposed by Kálmán (1960), producing the Kalman filter and smoother estimators, together with their conditional second moments. For known parameters $\boldsymbol{\varphi}$, these are nothing else but the linear projections: $\mathbf{F}_{t|t} := \text{Proj}_{\boldsymbol{\varphi}}(\mathbf{F}_t | \mathbf{x}_t, \dots, \mathbf{x}_1)$ and $\mathbf{F}_{t|T} := \text{Proj}_{\boldsymbol{\varphi}}(\mathbf{F}_t | \mathcal{X})$, respectively (see also Rauch, 1963, and De Jong, 1989). Hence, under the chosen parametrization of $\mathbf{\Omega}^F$, the Kalman smoother coincides with the estimator given in (45).

It is now sensible to compute the Kalman smoother by using the EM estimator of the parameters $\hat{\boldsymbol{\varphi}}^{\text{EM}}$, which approximate the QML estimators, defined in (37) through (40). However, those estimators of the parameters depend again on the factors. To break this mutual dependence, we can iterate between the Kalman smoother and the EM algorithm to obtain joint estimates of the parameters and the factors. In particular, at each iteration $k \geq 0$ of the EM algorithm we can implement the Kalman smoother using the estimated parameters $\boldsymbol{\varphi}^{(k)}$, estimated at the previous iteration, thus giving estimated factors, $\mathbf{F}_{t|T}^{(k)}$, their conditional MSE, $\mathbf{P}_{t|T}^{(k)}$, and lag-1 conditional MSE, $\mathbf{C}_{t,t-1|T}^{(k)}$ (see below for the explicit definitions), necessary to compute the expected log-likelihood in the E-step. By means of these moments we can compute new estimates of the parameters $\boldsymbol{\varphi}^{(k+1)}$ in the M-step.

In practice, the sufficient statistics (moments of 1st and 2nd order) needed to compute the final estimators of the parameters given in (37) through (40), are given by:

$$\begin{aligned} \mathbb{E}_{\hat{\boldsymbol{\varphi}}^{(k^*)}}[\mathbf{F}_t | \mathcal{X}] &= \mathbf{F}_{t|T}^{(k^*)}, \\ \mathbb{E}_{\hat{\boldsymbol{\varphi}}^{(k^*)}}[\mathbf{F}_t \mathbf{F}_t' | \mathcal{X}] &= \mathbf{F}_{t|T}^{(k^*)} \mathbf{F}_{t|T}^{(k^*)'} + \mathbf{P}_{t|T}^{(k^*)}, \quad \mathbb{E}_{\hat{\boldsymbol{\varphi}}^{(k^*)}}[\mathbf{F}_t \mathbf{F}_{t-1}' | \mathcal{X}] = \mathbf{F}_{t|T}^{(k^*)} \mathbf{F}_{t-1|T}^{(k^*)'} + \mathbf{C}_{t,t-1|T}^{(k^*)}. \end{aligned} \quad (46)$$

Notice, that, unless the factors and the idiosyncratic components are Gaussian, these expressions are correct only asymptotically since, as $N, T \rightarrow \infty$, the Kalman smoother $\mathbf{F}_{t|T}^{(k)}$ is a consistent estimator of the true factors at any given iteration $k \geq 0$, and any non-linearity is in $\mathbb{E}[\boldsymbol{\xi}_t | \mathcal{X}]$.

The final Kalman filter and Kalman smoother estimators are obtained at the end of the EM algorithm, i.e., they are given by $\hat{\mathbf{F}}_t^{\text{KF}} := \mathbf{F}_{t|t}^{(k^*+1)}$ and $\hat{\mathbf{F}}_t^{\text{KS}} := \mathbf{F}_{t|T}^{(k^*+1)}$. In particular, under the VAR(1) model in (32), these are defined by the following iterations:

$$\begin{aligned} \hat{\mathbf{F}}_t^{\text{KF}} &:= \hat{\mathbf{A}}^{\text{EM}} \hat{\mathbf{F}}_{t-1}^{\text{KF}} \\ &+ \left(\hat{\boldsymbol{\Lambda}}^{\text{EM}'} (\hat{\boldsymbol{\Sigma}}^{\xi \text{EM}})^{-1} \hat{\boldsymbol{\Lambda}}^{\text{EM}} + \hat{\mathbf{P}}_{t|t-1}^{-1} \right)^{-1} \hat{\boldsymbol{\Lambda}}^{\text{EM}'} (\hat{\boldsymbol{\Sigma}}^{\xi \text{EM}})^{-1} \left(\mathbf{x}_t - \bar{\mathbf{x}} - \hat{\boldsymbol{\Lambda}}^{\text{EM}} \hat{\mathbf{A}}^{\text{EM}} \hat{\mathbf{F}}_{t-1}^{\text{KF}} \right), \end{aligned} \quad t = t, \dots, T, \quad (47)$$

$$\hat{\mathbf{F}}_t^{\text{KS}} := \hat{\mathbf{F}}_t^{\text{KF}} + \hat{\mathbf{P}}_{t|t} \hat{\mathbf{A}}^{\text{EM}'} \hat{\mathbf{P}}_{t+1|t}^{-1} \left(\hat{\mathbf{F}}_{t+1}^{\text{KS}} - \hat{\mathbf{A}}^{\text{EM}} \hat{\mathbf{F}}_{t-1}^{\text{KF}} \right), \quad t = T, \dots, 1, \quad (48)$$

where $\hat{\mathbf{P}}_{t|t-1} = \hat{\mathbf{A}}^{\text{EM}} \hat{\mathbf{P}}_{t-1|t-1} \hat{\mathbf{A}}^{\text{EM}'} + \hat{\mathbf{H}}^{\text{EM}} \hat{\mathbf{H}}^{\text{EM}'}$, with $\hat{\mathbf{P}}_{t|t}$, being the filtered conditional MSEs, which

⁶Although in this paper we do not consider such possibility, it is also worth noticing that in the case of macroeconomic data we often have factors with a singular spectral density, i.e., of rank $q < r$ (see, e.g., the empirical evidence in D'Agostino and Giannone, 2012). This implies $\text{rk}(\mathbf{\Omega}^F) < rT$, so that, in such case, the smoother estimator (45) is not even defined.

evolves according to the iterations given in Appendix A. These iterations are initialized by $\widehat{\mathbf{F}}_0^{\text{KF}} = \mathbf{0}_r$, $\widehat{\mathbf{F}}_{T+1}^{\text{KS}} = \widehat{\mathbf{A}}^{\text{EM}} \widehat{\mathbf{F}}_T^{\text{KF}}$, and $\widehat{\mathbf{P}}_{0|0} = c\mathbf{I}_r$, for some finite positive real c .

Under the present setting, Barigozzi and Luciani (2019, Proposition 3) show that the Kalman smoother and the Kalman filter are asymptotically equivalent

$$\left\| \widehat{\mathbf{F}}_t^{\text{KS}} - \widehat{\mathbf{F}}_t^{\text{KF}} \right\| = O_P \left(\frac{1}{N} \right), \quad (49)$$

and also (up to logarithmic terms):

$$\left\| \widehat{\mathbf{F}}_t^{\text{KF}} - \widehat{\mathbf{F}}_t^{\text{WLS}} \right\| = O_P \left(\frac{1}{N} \right), \quad \left\| \widehat{\mathbf{F}}_t^{\text{KF}} - \widehat{\mathbf{F}}_t^{\text{LP}} \right\| = O_P \left(\frac{1}{N} \right), \quad (50)$$

that is, the Kalman filter and the feasible WLS/LP estimators are asymptotically equivalent (see also Bai and Li, 2016, Theorem 3 where however it is also required $T/N^3 \rightarrow 0$). From (49), (50), and Propositions 4 or 5, we get the following result.

Proposition 7. *For any $t = 1, \dots, T$, under Assumptions 1 through 8, if $\sqrt{N} \log^\tau N/T \rightarrow 0$ with $\tau := \max(\delta_F^{-1}, \delta_\xi^{-1}) + 1/2 \geq 3/2$, as $N, T \rightarrow \infty$,*

$$\sqrt{N}(\widehat{\mathbf{F}}_t^{\text{KS}} - \mathbf{F}_t) \rightarrow_d \mathcal{N}(\mathbf{0}_r, \mathbf{W}_t^{\text{WLS}}),$$

where $\mathbf{W}_t^{\text{WLS}} := (\boldsymbol{\Sigma}_{\Lambda\xi\Lambda})^{-1} \left\{ \lim_{N \rightarrow \infty} \frac{\boldsymbol{\Lambda}'(\boldsymbol{\Sigma}^\xi)^{-1} \mathbb{E}[\boldsymbol{\xi}_t \boldsymbol{\xi}_t'] (\boldsymbol{\Sigma}^\xi)^{-1} \boldsymbol{\Lambda}}{N} \right\} (\boldsymbol{\Sigma}_{\Lambda\xi\Lambda})^{-1}$, with $\boldsymbol{\Sigma}_{\Lambda\xi\Lambda} := \lim_{N \rightarrow \infty} \frac{\boldsymbol{\Lambda}'(\boldsymbol{\Sigma}^\xi)^{-1} \boldsymbol{\Lambda}}{N}$.

Up to logarithmic terms, the Kalman smoother estimator, computed using the EM estimator of the parameters, has the same asymptotic properties of the feasible WLS/LP estimators. Therefore, the same comments made after Propositions 4 and 5 about efficiency apply also in this case.

Remark 13. The iterations given in (47) and (48) might be computationally challenging because at each point in time they require inverting $\widehat{\mathbf{P}}_{t|t-1}$, which has a complex form. To avoid incurring in such numerical problems, Durbin and Koopman (2012, Chapter 4.4, pp. 70-73) provide a definition of the Kalman smoother, equivalent to (48), but which does not require the inversions of $\widehat{\mathbf{P}}_{t|t-1}$. An alternative numerically efficient approach is represented by the matrix formulation, instead of the recursive formulation, of the Kalman filter and smoother (see, e.g., Delle Monache and Petrella, 2019).

Remark 14. The previous results might lead to think that the Kalman filter and the Kalman smoother are useless in a high-dimensional setting, since they are equivalent to the WLS/LP estimator. But there is at least one good reason for preferring these estimators over the WLS/LP ones. Consider for simplicity the case of one factor, $r = 1$, and known parameters. Then, following Ruiz and Poncela (2022, Section 2.3), and denoting $B := \boldsymbol{\Lambda}'(\boldsymbol{\Sigma}^\xi)^{-1} \boldsymbol{\Lambda}$, we get:

$$F_t^{\text{KF}} = \frac{A}{1 + BP} F_{t-1}^{\text{KF}} + \frac{BP}{1 + BP} F_t^{\text{WLS}} = \frac{A}{1 + BP} F_{t-1}^{\text{KF}} + \frac{(B+1)P}{1 + BP} F_t^{\text{LP}}, \quad (51)$$

where A is the scalar autoregressive coefficient in (32), and P is the steady-state solution of the Riccati equation describing the dynamics of the one-step-ahead MSE $P_{t|t-1}$ (see Remark 3).⁷ Clearly, $B = O(N)$

⁷The steady-state P is given by (Poncela and Ruiz, 2015, Section 1.3):

$$P = \left\{ H^2 B - 1 + A^2 \left[1 + \sqrt{1 + \frac{4H^2 B}{(H^2 B - 1 + A^2)^2}} \right] \right\} \frac{1}{2B}.$$

because of Assumptions 1(a) and 2(a), so, as predicted from (50), the Kalman filter is asymptotically equivalent to the WLS/LP estimators. However, if $\{F_t\}$ is a very persistent process which does not fluctuate much, i.e., $A \lesssim 1$ and $H \gtrsim 0$ so that $P \simeq 0$, the first term on the right-hand-side of (51) might be non-negligible.

Notice that, because of (49) the same argument holds also for the Kalman smoother, which, in fact, has to be preferred over the Kalman filter, since, by being computed by conditioning on a larger set, has always a smaller variance.

Remark 15. By means of the Kalman filter and smoother we could treat any autocorrelated idiosyncratic component as additional latent states, thus introducing a dynamic model for each of them and by adding in the measurement equation (31) an error term with small variance (see, e.g., Bańbura and Modugno, 2014). This is crucial when the idiosyncratic components are very persistent (Barigozzi and Luciani, 2021). In this case, it can be shown that, in order for the presented asymptotic results to still hold, the number of additional latent states should be limited, and they should share some common driving force.

Remark 16. The factors can also be estimated by means of the Wiener-Kolmogorov filter, which, as mentioned above, is the frequency domain counterpart of the smoother estimator in (45) (Hannan, 1970, Chapter III.7). By means of such estimator an EM algorithm can be implemented in the spectral domain by first running the Wiener-Kolmogorov filter in the E-step, and then maximizing the expected Whittle log-likelihood in the M-step (Quah and Sargent, 1993; Fiorentini et al., 2018). As for all spectral methods, in order to implement a spectral EM we must first estimate a large dimensional spectral density matrix.

Remark 17. A last approach to estimate the factors by accounting of their dynamics is proposed by Bai and Li (2016), and it is based on the three following steps. Step 1: compute estimators of $\mathbf{\Lambda}$ and $\mathbf{\Sigma}^\xi$ via QML, and of the factors via WLS, as described in Section 5. Step 2: use the estimated factors to compute estimators of \mathbf{A} and \mathbf{H} in (32). Step 3: use those estimated parameters to compute the Kalman smoother estimator of the factors. Similar multi-step approaches have been proposed by Doz et al. (2011), who replace step 1 with the PC estimators of loadings and factors, and Ng et al. (2015), who replace step 1 with an estimator of the loadings obtained by maximizing the log-likelihood (19) using the Newton-Raphson algorithm. However, none of those approaches exploits the mutual feedback from the loadings to the factors and viceversa; hence, at least in finite samples, they might incur in efficiency losses.

7 Classical ML estimation

In this section, we review classical factor analysis for cross-sectional data, which is characterized by: (i) fixed N , (ii) uncorrelated Gaussian, i.e., independent, idiosyncratic components $\mathbf{\Omega}^\xi = \mathbf{I}_T \otimes \mathbf{\Sigma}^\xi$, and (iii) non-stochastic factors, i.e., $\mathbf{\Omega}^F = \mathbf{I}_T \otimes \mathbf{\Gamma}^F$. This is the setting considered mainly in psychometric applications (see Lawley and Maxwell, 1971 and references therein).

Classical Maximum Likelihood (ML) estimation of the factor model can be carried out in two ways.

- (A) Maximize the, now correctly specified, log-likelihood (19) subject to some given identifying constraint, in order to obtain an ML estimator of the loadings and of the idiosyncratic variances using, e.g., the EM algorithm by Rubin and Thayer (1982). And, then, use these estimates to compute an estimator of the factors, either by least squares (see Bartlett, 1937, and (26) in Section 5.2) or by linear projection (see Thomson, 1951, and (29) in Section 5.2).
- (B) Use the formulation of the log-likelihood in (24) and jointly estimate loadings and factors, which, in this case, are always estimated via least squares.

In principle, approach A is considered to be the appropriate one in the case stochastic factors, while approach B is considered to be the appropriate one in the case non-stochastic factors (Anderson, 2003, p. 587). However, approach A can be used also with non-stochastic factors. Indeed, the log-likelihood (19) does not depend on the factors, nor on their second moments if we also assume $\mathbf{\Gamma}^F = \mathbf{I}_r$. Hence, approach A is commonly considered the most sensible one.

Another reason to prefer approach A is that the log-likelihood (24), used in approach B, might diverge to infinity under certain choices of the parameters. To be more precise, this issue is related to just one specific configuration, namely the case in which at least one idiosyncratic component has zero variance (Anderson and Rubin, 1956, Section 7.7, Anderson, 2003, p. 587, Breitung and Tenhofen, 2011). This implies that the ML estimator of the idiosyncratic variances might not be defined. To solve this problem, Lawley (1942) suggested to look for solutions satisfying just the first-order conditions, but, as shown by Solari (1969), this does not lead to discovering a maximum, but just a stationary point of the log-likelihood. Notice that, if we instead follow approach A, then the log-likelihood (19) can easily accommodate some idiosyncratic components with zero variance, see, e.g., Bartholomew et al. (2011, Chapter 3.4).

Nevertheless, approach B has the nice feature of allowing us to exploit the mutual dependence of factors and loadings, and, moreover, the log-likelihood (24) has a more tractable form than the log-likelihood (19).

Regardless of the approach considered, when N is fixed ML estimation of factor models suffers of two main problems: (I) inference for the loadings is computationally challenging, and (II) the factors cannot be estimated consistently and their estimation might even pose an incidental parameter problem for the estimation of the other parameters of the model. We now consider both issues in detail and show how letting $N \rightarrow \infty$ helps solving both problems and would allow us to use the estimation approach in B.

Consider first, the estimation of the loadings. Under the assumption of an exact factor model, Bai and Li (2012, Theorems 5.2 and 5.4) prove that the ML estimator of the loadings, maximizing (19), is such that:

$$\left\| \widehat{\boldsymbol{\lambda}}_i^{\text{ML}} - \boldsymbol{\lambda}_i^{\text{OLS}} \right\| = O_{\text{P}} \left(\frac{1}{\sqrt{NT}} \right). \quad (52)$$

From this result it is clear that since now the factor model is truly exact, then, the ML estimator of the loadings is \sqrt{T} -consistent, regardless of N . However, while if $N \rightarrow \infty$, the asymptotic covariance is the one given in Proposition 3, when N is fixed the expression is much more complex. Indeed, from the asymptotic expansion (52) it is clear that if N is fixed we still have \sqrt{T} consistency, but an additional term on top of the OLS estimation error, must be included in the asymptotic distribution. The specific expression of the asymptotic covariance in the fixed N case is given, for example, in Anderson and Rubin

(1956, Theorem 12.3), Amemiya et al. (1987, Theorem 2F), and Anderson and Amemiya (1988, Theorems 1, 2, and 3) (see also Barigozzi, 2022, Proposition 10). Since for fixed N defining a consistent estimator of such covariance matrix is not easy, inference might become a computationally hard problem. Notice that the same problem exists even if we assumed the simplest possible case of an exact factor model with the unrealistic assumption of cross-sectionally homoskedastic idiosyncratic components, as in Young (1940) and Whittle (1952).

Second, the factors represent an additional rT -dimensional vector of parameters that need to be estimated. They can either be estimated after we have an ML estimator of the loadings and the idiosyncratic variances (approach A) or they can be estimated jointly with the other parameters (approach B). Now, when N is fixed none of the two approaches allows to retrieve the factors consistently. The impossibility to retrieve the factors consistently when N is fixed is precisely the reason why in classical factor analysis the factors are usually identified only indirectly through their associated loadings which, as we have seen above are \sqrt{T} -consistent. This is the spirit of confirmatory analysis (Jöreskog, 1969), an example of which is in Mardia et al. (1979, Chapter 9.6), where the factors are identified through a VARIMAX orthogonal rotation of the loadings, providing rotated loadings with a few large entries and as many near-zero entries as possible. Another example is in Terada (2014), where the loadings are identified by the reduced K -means clustering method.

Moreover, besides the aforementioned problem of zero idiosyncratic variances, approach B, although appealing, poses also an incidental parameter problem. Indeed, we need to simultaneously estimate $rT + (r + 1)N$ parameters using NT observations, and we might easily run out of degrees-of-freedom. This is especially true if N is small and T is large, while the larger N gets, the less binding the issue is (see Remark 7.1 for details).

Now, if we let $N \rightarrow \infty$, then we can prove that under approach A the factors are consistently estimated (see Propositions 4 and 5). Furthermore, as shown in Section 7.1 below, when $N \rightarrow \infty$ also the incidental parameter problem vanishes asymptotically and we could use the estimation approach B by just iterating between the two sets of estimates. Say we start at iteration $k = 0$ with an estimate of the loadings and the idiosyncratic variances, e.g., given by the PC estimator. Then, at any iteration $k \geq 0$ we would have the feasible WLS and OLS estimators:

$$\widehat{\mathbf{F}}_t^{(k)} = (\widehat{\mathbf{\Lambda}}^{(k)'} (\widehat{\mathbf{\Sigma}}^{\xi(k)})^{-1} \widehat{\mathbf{\Lambda}}^{(k)})^{-1} \widehat{\mathbf{\Lambda}}^{(k)'} (\widehat{\mathbf{\Sigma}}^{\xi(k)})^{-1} (\mathbf{x}_t - \bar{\mathbf{x}}), \quad t = 1, \dots, T, \quad (53)$$

$$\widehat{\boldsymbol{\lambda}}_i^{(k+1)} = (\widehat{\mathbf{F}}^{(k)'} \widehat{\mathbf{F}}^{(k)})^{-1} \widehat{\mathbf{F}}^{(k)'} (\mathbf{x}_i - \bar{x}_i \boldsymbol{\nu}_T), \quad i = 1, \dots, N. \quad (54)$$

Clearly since we initialize the algorithm with a consistent estimator of the parameters, then at each iteration $k \geq 0$ both estimators can be shown to be consistent too as $N, T \rightarrow \infty$. Approach B would then become feasible and, possibly, more preferable than approach A. Indeed, approach B would allow us to explicitly take into account the mutual dependence of factors and loadings and we would also have closed form expressions for all estimators.

7.1 The incidental parameter problem

In this section we formally show that if N is fixed, then the score of the log-likelihood (24) with respect to the loadings does not satisfy the orthogonality condition by Neyman (1979). Hence, any estimator the loadings using an estimator of the factors, is going to be biased. More precisely, for any specific

value of the factors, $\tilde{\mathbf{F}}_t$, of the loadings, $\tilde{\mathbf{\Lambda}}$, and of the idiosyncratic variances, $\tilde{\mathbf{\Sigma}}^\xi$, consider the score with respect to the loadings $\underline{\lambda}_i$, $i = 1, \dots, N$,

$$\begin{aligned} s_{it}(\mathbf{x}_t; \tilde{\mathbf{F}}_t, \tilde{\mathbf{\Lambda}}, \tilde{\mathbf{\Sigma}}^\xi) &:= \frac{\partial}{\partial \underline{\lambda}_i} \left\{ -\frac{1}{2} \log \det(\underline{\mathbf{\Sigma}}^\xi) - \frac{1}{2} (\mathbf{x}_t - \bar{\mathbf{x}} - \underline{\mathbf{\Lambda}} \mathbf{F}_t)' (\underline{\mathbf{\Sigma}}^\xi)^{-1} (\mathbf{x}_t - \bar{\mathbf{x}} - \underline{\mathbf{\Lambda}} \mathbf{F}_t) \right\} \Big|_{\substack{\mathbf{F}_t = \tilde{\mathbf{F}}_t \\ \underline{\mathbf{\Lambda}} = \tilde{\mathbf{\Lambda}}, \underline{\mathbf{\Sigma}}^\xi = \tilde{\mathbf{\Sigma}}^\xi}} \\ &= \frac{\tilde{\mathbf{F}}_t(x_{it} - \tilde{\mathbf{F}}_t' \tilde{\lambda}_i)}{\tilde{\sigma}_i^2}. \end{aligned}$$

Clearly, in the true values of factors and parameters the first-order conditions are satisfied:

$$\mathbb{E}[s_{it}(\mathbf{x}_t; \mathbf{F}_t, \mathbf{\Lambda}, \mathbf{\Sigma}^\xi)] = \frac{\mathbb{E}[\mathbf{F}_t \xi_{it}]}{\sigma_i^2} = \frac{\mathbf{F}_t \mathbb{E}[\xi_{it}]}{\sigma_i^2} = \mathbf{0}_r, \quad (55)$$

since we treat the factors as constant parameters (but this would hold even for stochastic factors uncorrelated with the idiosyncratic components).

We now ask ourselves what happens if we replace the true factors with an estimator, and let us choose the best possible estimator, which is the unfeasible WLS in (25). To this end, consider the curve $\gamma : [0, 1] \rightarrow \mathbb{R}^r$ connecting \mathbf{F}_t and $\mathbf{F}_t^{\text{WLS}}$, i.e., such that $\gamma(\eta) = \mathbf{F}_t + \eta(\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t)$ for $\eta \in [0, 1]$, and consider the r -dimensional function $\mathbf{D}_\eta : \mathbb{R}^r \rightarrow \mathbb{R}^r$ describing the variation of the score along the curve γ , so that for any $\eta \in [0, 1]$ we have the $r \times 1$ differential vector

$$\mathbf{D}_\eta(\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t) := \partial_\eta \left\{ \mathbb{E} \left[s_i \left(\mathbf{x}_t; \mathbf{F}_t + \eta(\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t), \mathbf{\Lambda}, \mathbf{\Sigma}^\xi \right) \right] \right\}.$$

In order, to have unbiased estimators of the loadings the first-order conditions should be locally insensitive to the value of the factors, which here play the role of nuisance parameters. A necessary condition for this to happen is that the Gateaux derivative along γ should be such that (see also Chernozhukov et al., 2018, Definition 2.1):

$$\mathbf{D}_\eta[\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t]|_{\eta=0} = \lim_{\eta \rightarrow 0} \frac{\mathbb{E} [s_i(\mathbf{x}_t; \mathbf{F}_t + \eta(\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t), \mathbf{\Lambda}, \mathbf{\Sigma}^\xi)] - \mathbb{E} [s_i(\mathbf{x}_t; \mathbf{F}_t, \mathbf{\Lambda}, \mathbf{\Sigma}^\xi)]}{\eta} = \mathbf{0}_r, \quad (56)$$

If condition (56) is satisfied, then we could estimate the loadings by plugging into the log-likelihood (24) noisy estimates of the factors, and then maximize the concentrated log-likelihood $\ell(\mathcal{X}; \mathcal{F}^{\text{WLS}}, \underline{\varphi})$, without strongly violating the moment conditions (55). However, it is easy to see that for fixed N condition (56) is not satisfied. Indeed, by using (55) and the expression of the WLS estimator (25) into (56), we get:

$$\begin{aligned} \mathbf{D}_\eta[\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t]|_{\eta=0} &= \lim_{\eta \rightarrow 0} \frac{\mathbb{E} [\mathbf{F}_t \xi_{it} - \eta \mathbf{F}_t (\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t)' \lambda_i + \eta (\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t) \xi_{it} - \eta^2 (\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t) (\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t)' \lambda_i]}{\eta \sigma_i^2} \\ &= \frac{1}{\sigma_i^2} \mathbb{E} [(\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t) \xi_{it}] - \frac{1}{\sigma_i^2} \mathbb{E} [\mathbf{F}_t (\mathbf{F}_t^{\text{WLS}} - \mathbf{F}_t) \lambda_i] \\ &= \frac{1}{\sigma_i^2} (\mathbf{\Lambda}' (\mathbf{\Sigma}^\xi)^{-1} \mathbf{\Lambda})^{-1} \mathbf{\Lambda}' (\mathbf{\Sigma}^\xi)^{-1} \mathbb{E} [\xi_t \xi_{it}] - \frac{1}{\sigma_i^2} \mathbb{E} [\mathbf{F}_t \xi_t] (\mathbf{\Sigma}^\xi)^{-1} \mathbf{\Lambda} (\mathbf{\Lambda}' (\mathbf{\Sigma}^\xi)^{-1} \mathbf{\Lambda})^{-1} \\ &= \frac{1}{\sigma_i^2} \left(\frac{\mathbf{\Lambda}' (\mathbf{\Sigma}^\xi)^{-1} \mathbf{\Lambda}}{N} \right)^{-1} \frac{1}{N} \sum_{j=1}^N \frac{\lambda_j \mathbb{E} [\xi_{jt} \xi_{it}]}{\sigma_j^2}, \end{aligned} \quad (57)$$

which for fixed N is never zero, not even if the model is exact, in which case the summation will just made of one term when $j = i$. So even if we use the ideal unfeasible WLS estimator of the factors, we would get biased estimates for the loadings. However, we clearly see that in the limit $N \rightarrow \infty$, the expression in (57) is $O_P(N^{-1})$, because of Assumptions 1(a), 2(a), and 2(b), and condition (56) is satisfied. So, as expected, the problem of incidental parameters vanishes asymptotically.

8 On the mis-specification error and the search for efficient estimators

Consider the two equivalent ways of writing the factor model (1) when stacking all NT observations in one NT dimensional vector, i.e., by vectorizing the matrix representation (2) or its transposed:

$$\mathbf{x} := \text{vec}(\mathbf{X}) = \text{vec}(\boldsymbol{\nu}_T \boldsymbol{\alpha}') + (\mathbf{I}_N \otimes \mathbf{F}) \text{vec}(\boldsymbol{\Lambda}') + \text{vec}(\boldsymbol{\Xi}) =: \boldsymbol{\alpha} + \boldsymbol{\mathfrak{F}} \boldsymbol{\mathcal{L}} + \boldsymbol{\mathcal{E}}, \quad (58)$$

$$\boldsymbol{\mathcal{X}} := \text{vec}(\mathbf{X}') = \text{vec}(\boldsymbol{\alpha} \boldsymbol{\nu}_T') + (\mathbf{I}_T \otimes \boldsymbol{\Lambda}) \text{vec}(\mathbf{F}') + \text{vec}(\boldsymbol{\Xi}') =: \boldsymbol{\mathcal{A}} + \boldsymbol{\mathcal{L}} \boldsymbol{\mathcal{F}} + \boldsymbol{\mathcal{E}}. \quad (59)$$

Let $\boldsymbol{\Theta}^\xi := \mathbb{E}[\text{vec}(\boldsymbol{\Xi}) \text{vec}(\boldsymbol{\Xi})'] = \mathbb{E}[\boldsymbol{\mathcal{E}} \boldsymbol{\mathcal{E}}']$ and $\boldsymbol{\Omega}^\xi := \mathbb{E}[\text{vec}(\boldsymbol{\Xi}') \text{vec}(\boldsymbol{\Xi}')'] = \mathbb{E}[\boldsymbol{\mathcal{E}} \boldsymbol{\mathcal{E}}']$, which are $NT \times NT$ matrices having the same elements just ordered differently.

If the factors were observed, the true, i.e., not mis-specified, log-likelihood (17) could be written also as:

$$\ell(\boldsymbol{\mathfrak{X}}, \boldsymbol{\mathfrak{F}}; \boldsymbol{\mathcal{L}}, \text{vech}(\boldsymbol{\Theta}^\xi)) \simeq -\frac{1}{2} \log \det(\boldsymbol{\Theta}^\xi) - \frac{1}{2} \left[(\boldsymbol{\mathfrak{X}} - \bar{\boldsymbol{\mathfrak{X}}} - \boldsymbol{\mathfrak{F}} \boldsymbol{\mathcal{L}})' (\boldsymbol{\Theta}^\xi)^{-1} (\boldsymbol{\mathfrak{X}} - \bar{\boldsymbol{\mathfrak{X}}} - \boldsymbol{\mathfrak{F}} \boldsymbol{\mathcal{L}}) \right]. \quad (60)$$

Then, for known $\boldsymbol{\Theta}^\xi$ the QML estimator of the loadings, maximizing (60), would be the unfeasible GLS estimator:

$$\boldsymbol{\mathcal{L}}^{\text{GLS}} = \text{vec}(\boldsymbol{\Lambda}^{\text{GLS}'}) := \left\{ \boldsymbol{\mathfrak{F}}' (\boldsymbol{\Theta}^\xi)^{-1} \boldsymbol{\mathfrak{F}} \right\}^{-1} \left\{ \boldsymbol{\mathfrak{F}}' (\boldsymbol{\Theta}^\xi)^{-1} (\boldsymbol{\mathfrak{X}} - \bar{\boldsymbol{\mathfrak{X}}}) \right\}. \quad (61)$$

This would be the QML estimator of the loadings for an approximate factor model with known factors.

Likewise, for known loadings the true, i.e., not mis-specified, log-likelihood (17) could be written also as:

$$\ell(\boldsymbol{\mathcal{X}}, \boldsymbol{\mathcal{L}}; \boldsymbol{\mathcal{F}}, \text{vech}(\boldsymbol{\Omega}^\xi)) \simeq -\frac{1}{2} \log \det(\boldsymbol{\Omega}^\xi) - \frac{1}{2} \left[(\boldsymbol{\mathcal{X}} - \bar{\boldsymbol{\mathcal{X}}} - \boldsymbol{\mathcal{L}} \boldsymbol{\mathcal{F}})' (\boldsymbol{\Omega}^\xi)^{-1} (\boldsymbol{\mathcal{X}} - \bar{\boldsymbol{\mathcal{X}}} - \boldsymbol{\mathcal{L}} \boldsymbol{\mathcal{F}}) \right]. \quad (62)$$

Then, for known $\boldsymbol{\Omega}^\xi$ the QML estimator of the factors, maximizing (62), would be the unfeasible GLS estimator:

$$\boldsymbol{\mathcal{F}}^{\text{GLS}} = \text{vec}(\mathbf{F}^{\text{GLS}'}) := \left\{ \boldsymbol{\mathcal{L}}' (\boldsymbol{\Omega}^\xi)^{-1} \boldsymbol{\mathcal{L}} \right\}^{-1} \left\{ \boldsymbol{\mathcal{L}}' (\boldsymbol{\Omega}^\xi)^{-1} (\boldsymbol{\mathcal{X}} - \bar{\boldsymbol{\mathcal{X}}}) \right\}. \quad (63)$$

This would be the QML estimator of the factors for an approximate factor model with known loadings.

Now, even if the factors or the loadings were known, the GLS estimators (61) and (63) would still require the unfeasible task to estimate and invert $\boldsymbol{\Theta}^\xi$ or $\boldsymbol{\Omega}^\xi$. As done in Section 5, we can then consider a mis-specified version of the log-likelihoods (60) or (62), where we do not account for idiosyncratic correlations and we set $\boldsymbol{\Theta}^\xi = \boldsymbol{\Sigma}^\xi \otimes \mathbf{I}_T$ and $\boldsymbol{\Omega}^\xi = \mathbf{I}_T \otimes \boldsymbol{\Sigma}^\xi$. Then, it is easy to see that the maximizers of such mis-specified log-likelihoods are the unfeasible OLS estimator of the loadings and the unfeasible

WLS estimator of the factors. Indeed, in this case (61) would simplify as follows:

$$\begin{aligned}
\mathcal{L}^{\text{OLS}} = \text{vec}(\mathbf{\Lambda}^{\text{OLS}'}) &:= \left\{ \mathfrak{F}'(\mathbf{\Sigma}^\xi \otimes \mathbf{I}_T)^{-1} \mathfrak{F} \right\}^{-1} \left\{ \mathfrak{F}'(\mathbf{\Sigma}^\xi \otimes \mathbf{I}_T)^{-1} (\mathbf{x} - \bar{\mathbf{x}}) \right\} \\
&= \left\{ (\mathbf{\Sigma}^\xi)^{-1} \otimes \mathbf{F}'\mathbf{F} \right\}^{-1} \left\{ \left((\mathbf{\Sigma}^\xi)^{-1} \otimes \mathbf{F}' \right) (\mathbf{x} - \bar{\mathbf{x}}) \right\} \\
&= \left\{ \mathbf{\Sigma}^\xi \otimes (\mathbf{F}'\mathbf{F})^{-1} \right\} \left\{ \text{vec} \left(\mathbf{F}'(\mathbf{X} - \bar{\mathbf{X}})(\mathbf{\Sigma}^\xi)^{-1} \right) \right\} \\
&= \text{vec} \left((\mathbf{F}'\mathbf{F})^{-1} \mathbf{F}'(\mathbf{X} - \bar{\mathbf{X}}) \right), \tag{64}
\end{aligned}$$

with components λ_i^{OLS} given in (10). And, similarly, (63) would simplify as follows:

$$\begin{aligned}
\mathcal{F}^{\text{WLS}} = \text{vec}(\mathbf{F}^{\text{WLS}'}) &:= \left\{ \mathfrak{L}'(\mathbf{I}_T \otimes \mathbf{\Sigma}^\xi)^{-1} \mathfrak{L} \right\}^{-1} \left\{ \mathfrak{L}'(\mathbf{I}_T \otimes \mathbf{\Sigma}^\xi)^{-1} (\mathbf{x} - \bar{\mathbf{x}}) \right\} \\
&= \left\{ \mathbf{I}_T \otimes \mathbf{\Lambda}'(\mathbf{\Sigma}^\xi)^{-1} \mathbf{\Lambda} \right\}^{-1} \left\{ \left(\mathbf{I}_T \otimes \mathbf{\Lambda}'(\mathbf{\Sigma}^\xi)^{-1} \right) (\mathbf{x} - \bar{\mathbf{x}}) \right\} \\
&= \left\{ \mathbf{I}_T \otimes \left(\mathbf{\Lambda}'(\mathbf{\Sigma}^\xi)^{-1} \mathbf{\Lambda} \right)^{-1} \mathbf{\Lambda}'(\mathbf{\Sigma}^\xi)^{-1} \right\} (\mathbf{x} - \bar{\mathbf{x}}) \\
&= \text{vec} \left(\left(\mathbf{\Lambda}'(\mathbf{\Sigma}^\xi)^{-1} \mathbf{\Lambda} \right)^{-1} \mathbf{\Lambda}'(\mathbf{\Sigma}^\xi)^{-1} (\mathbf{X} - \bar{\mathbf{X}})' \right), \tag{65}
\end{aligned}$$

with components $\mathbf{F}_t^{\text{WLS}}$ given in (25).

As shown in Section 5, these unfeasible OLS and WLS estimators are asymptotically equivalent to the QML and feasible WLS estimators studied in Propositions 3 and 4, respectively. So to understand what is the effect of the mis-specifications introduced in the log-likelihood (19), it is enough to study how different the unfeasible OLS and WLS estimators in (64) and (65) are from the most efficient estimators which are the unfeasible GLS estimators in (61) and (63), respectively.

On average the full and the mis-specified idiosyncratic covariances are asymptotically equivalent since, for all $N, T \in \mathbb{N}$,

$$\begin{aligned}
\frac{1}{NT} \left\| \mathbf{\Theta}^\xi - (\mathbf{\Sigma}^\xi \otimes \mathbf{I}_T) \right\| &= \frac{1}{NT} \left\| \mathbf{\Omega}^\xi - (\mathbf{I}_T \otimes \mathbf{\Sigma}^\xi) \right\| \\
&\leq \frac{1}{NT} \max_{i=1, \dots, N} \max_{t=1, \dots, T} \sum_{j=1}^N \sum_{s=1}^T |\mathbb{E}[\xi_{it}\xi_{js}]| \leq \frac{1}{NT} \max_{i=1, \dots, N} \max_{t=1, \dots, T} \sum_{j=1}^N \sum_{s=1}^T M_{ij} \rho^{|t-s|} \\
&\leq \frac{M_\xi}{(1-\rho)NT}, \tag{66}
\end{aligned}$$

because of Assumption 2(b) and since M_ξ and ρ do not depend on i, j, t, s . Nevertheless, the difference between the two sets of estimators depends on the inverse idiosyncratic covariances, i.e., on

$$\begin{aligned}
\left\| (\mathbf{\Theta}^\xi)^{-1} - (\mathbf{\Sigma}^\xi \otimes \mathbf{I}_T)^{-1} \right\| &\leq \left\| (\mathbf{\Theta}^\xi)^{-1} \right\| \left\| \mathbf{\Theta}^\xi - (\mathbf{\Sigma}^\xi \otimes \mathbf{I}_T) \right\| \left\| (\mathbf{\Sigma}^\xi \otimes \mathbf{I}_T)^{-1} \right\|, \\
\left\| (\mathbf{\Omega}^\xi)^{-1} - (\mathbf{I}_T \otimes \mathbf{\Sigma}^\xi)^{-1} \right\| &\leq \left\| (\mathbf{\Omega}^\xi)^{-1} \right\| \left\| \mathbf{\Omega}^\xi - (\mathbf{I}_T \otimes \mathbf{\Sigma}^\xi) \right\| \left\| (\mathbf{I}_T \otimes \mathbf{\Sigma}^\xi)^{-1} \right\|,
\end{aligned}$$

which, even by assuming that $\mathbf{\Theta}^\xi$ and $\mathbf{\Omega}^\xi$ are positive definite, are still finite for all N and T , but do not vanish. Therefore, as expected, the GLS estimators are in principle better than the unfeasible QML and WLS estimators described in the previous section.

In light of this discussion, it is natural to consider extensions of QML estimation of the static

approximate factor model where we model explicitly some of the serial and cross-sectional idiosyncratic correlations. This would allow us to compute new estimators of the loadings and the factors which are closer to the unfeasible GLS estimators and thus are more efficient than the OLS and WLS.

In particular, the second best that we can hope for are the following pseudo-GLS estimators. Consider, first, estimation of the loadings when we do not model either the idiosyncratic cross- or cross-autocorrelations, i.e., when imposing $\mathbb{E}[\zeta_i \zeta_j'] = \mathbf{0}_{T \times T}$ for $i \neq j$. So we maximize the log-likelihood (60) when Θ^ξ is block-diagonal with N blocks $\Delta_i^\xi := \mathbb{E}[\zeta_i \zeta_i']$, $i = 1, \dots, N$, each being a $T \times T$ matrix, with entries $\mathbb{E}[\xi_{it} \xi_{is}]$, $t, s = 1, \dots, T$, hence, under serial homoskedasticity, Δ_i^ξ is a Toeplitz matrix. The QML estimator of the loadings is then the pseudo-GLS estimator:

$$\lambda_i^{\text{GLS}_0} := \left(\mathbf{F}'(\Delta_i^\xi)^{-1} \mathbf{F} \right)^{-1} \mathbf{F}'(\Delta_i^\xi)^{-1} (\mathbf{x}_i - \bar{x}_i \mathbf{1}_T), \quad i = 1, \dots, N, \quad (67)$$

This estimator has an asymptotic covariance matrix which attains the Gauss-Markov lower bound: $\lim_{T \rightarrow \infty} T(\mathbb{E}[\mathbf{F}'(\Delta_i^\xi)^{-1} \mathbf{F}])^{-1}$.

However, even for known factors, computing (67) is in general unfeasible, since it requires estimation and inversion of a large matrix Δ_i^ξ . A common approach is to assume a parametric expression for Δ_i^ξ . For example, let $\xi_{it} = \alpha_i \xi_{i,t-1} + \nu_{it}$ with $|\alpha_i| < 1$ and $\nu_{it} \sim (0, \omega_i^2)$ and uncorrelated across i and t . Then, both Δ_i^ξ and its inverse depend only on α_i and ω_i^2 .⁸ In practice, we could first estimate the loadings by QML, then the factors by WLS, and last the idiosyncratic components, from which, by fitting an AR(1), we could get estimates of α_i and ω_i^2 and thus of Δ_i^ξ and its inverse. By using all these estimates we can compute a feasible version of the pseudo-GLS estimator of the loadings in (67). This is similar to the approach used in PC estimation by [Breitung and Tenhofen \(2011\)](#). and, for known factors, it coincides with the classical feasible GLS estimator of the loadings, which is asymptotically equivalent to their QML estimator when considering the mis-specified log-likelihood (24), but for the filtered data $(1 - \alpha_i L)x_{it}$, $i = 1, \dots, N$, and, thus, with Σ^ξ replaced by a diagonal matrix with entries ω_i^2 , $i = 1, \dots, N$ ([Cochrane and Orcutt, 1949](#)).

Similarly, in order to explicitly model the autocorrelation of the idiosyncratic components, the EM algorithm presented in Section 6.1 can also be modified. Namely, in the M-step of any iteration $k \geq 0$, we could iterate between estimates of the loadings conditional on α_i and ω_i , and estimates of α_i and ω_i conditional on the loadings, until convergence, and then move to iteration $k + 1$ of the EM algorithm. This is the procedure adopted by [Reis and Watson \(2010\)](#) and it is an instance of an Expectation Conditional Maximization (ECM) algorithm ([Meng and Rubin, 1993](#)).

Second, consider estimation of the factors when we do not model either idiosyncratic auto- and cross-autocorrelations, i.e., when we impose $\mathbb{E}[\xi_t \xi_s'] = \mathbf{0}_{N \times N}$ for $t \neq s$. So we maximize the log-likelihood (62) when $\Omega^\xi = \mathbf{I}_T \otimes \Gamma^\xi$, where $\Gamma^\xi := \mathbb{E}[\xi_t \xi_t']$ is a $N \times N$ matrix, with entries $\mathbb{E}[\xi_{it} \xi_{jt}]$, $i, j = 1, \dots, N$,

⁸Indeed, we have that the (t, s) entry of Δ_i^ξ is $\frac{\omega_i^2 \alpha_i^{|t-s|}}{1 - \alpha_i^2}$ and

$$\left(\Delta_i^\xi \right)^{-1} = \frac{\omega_i^2}{(1 - \alpha_i^2)^2} \begin{pmatrix} 1 & -\alpha_i & 0 & \dots & 0 & 0 & 0 \\ -\alpha_i & 1 + \alpha_i^2 & -\alpha_i & \dots & 0 & 0 & 0 \\ 0 & -\alpha_i & 1 + \alpha_i^2 & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & -\alpha_i & 1 + \alpha_i^2 & -\alpha_i \\ 0 & 0 & 0 & \dots & 0 & -\alpha_i & 1 \end{pmatrix}.$$

hence, under serial homoskedasticity, it is independent of t . The QML estimator of the factors is then the pseudo-GLS estimator:

$$\mathbf{F}_t^{\text{GLS}_0} := \left(\mathbf{\Lambda}'(\mathbf{\Gamma}^\xi)^{-1}\mathbf{\Lambda} \right)^{-1} \mathbf{\Lambda}'(\mathbf{\Gamma}^\xi)^{-1}(\mathbf{x}_t - \bar{\mathbf{x}}), \quad t = 1, \dots, T. \quad (68)$$

This estimator has an asymptotic covariance matrix which attains the Gauss-Markov lower bound: $\lim_{N \rightarrow \infty} N(\mathbf{\Lambda}'(\mathbf{\Gamma}^\xi)^{-1}\mathbf{\Lambda})^{-1}$.

However, even for known loadings computing (68) is in general unfeasible, since it requires estimation and inversion of a large matrix $\mathbf{\Gamma}^\xi$. To this end, Bai and Liao (2016) propose an EM algorithm to maximize the log-likelihood (19) when replacing $\mathbf{\Sigma}^\xi$ with the full matrix $\mathbf{\Gamma}^\xi$ but subject to an ℓ_1 penalty imposed on its off-diagonal entries. They then use the QML estimators of the loadings and the idiosyncratic covariance obtained in this way to compute a feasible version of the pseudo-GLS estimator of the factors in (68). Similar approaches are also in Wang et al., 2019, and Pognard and Terada, 2020. Generalizations of the EM algorithm, e.g., by considering penalized M-steps in order to account for cross-sectional idiosyncratic correlations, are in principle possible too. This is the spirit of the approach by Lin and Michailidis (2020) who assume a sparse VAR model for the idiosyncratic components, thus accounting also for cross-autocorrelation, and they embed into the EM algorithm a penalized M-step.

Nevertheless, in general, the pseudo-GLS estimators (67) and (68) are not necessarily more efficient than the QML and WLS estimators studied in Proposition 3 and 4, since, they do not address possible cross-autocorrelations, so they are still maximizers of mis-specified log-likelihoods. We might have gains in efficiency only if the cross-autocorrelations are small enough, i.e., if the $T \times T$ off-diagonal blocks of $\mathbf{\Theta}^\xi$ or the $N \times N$ off-diagonal blocks of $\mathbf{\Omega}^\xi$ are sparse enough. Moreover, the feasible pseudo-GLS estimators discussed above may suffer if we do not correctly specify the autoregressive order of the idiosyncratic components or we do not select correctly the degree of penalization when thresholding the idiosyncratic covariance matrix. For these reasons such estimators are hardly considered in empirical econometric applications.

Remark 18. Lets us briefly consider the case in which we had serial idiosyncratic heteroskedasticity. Then, regarding estimation of the loadings, if we do not model any idiosyncratic cross- or cross-autocorrelations, the Gauss-Markov lower bound would be $\lim_{T \rightarrow \infty} T(\mathbb{E}[\mathbf{F}'(\mathbf{H}_i^\xi)^{-1}\mathbf{F}])^{-1}$, where $\mathbf{H}_i^\xi := \mathbb{E}[\zeta_i \zeta_i']$ is a $T \times T$ matrix with entries $\mathbb{E}[\xi_{it}\xi_{is}]$, $t, s = 1, \dots, T$, which now depend on t and s , so it is no more a Toeplitz matrix. Regarding the estimated factors, if we do not model any idiosyncratic auto- or cross-autocorrelations, the Gauss-Markov lower bound would become $\lim_{N \rightarrow \infty} N(\mathbf{\Lambda}'(\mathbf{\Gamma}_t^\xi)^{-1}\mathbf{\Lambda})^{-1}$, where $\mathbf{\Gamma}_t^\xi := \mathbb{E}[\xi_t \xi_t']$ is a $N \times N$ matrix with entries $\mathbb{E}[\xi_{it}\xi_{jt}]$, $t, s = 1, \dots, T$, which now depend on t . As a consequence, the pseudo-GLS estimators (67) and (68) become even less efficient since they do not account also for serial heteroskedasticity. Notice, however, that estimating and inverting \mathbf{H}_i^ξ and $\mathbf{\Gamma}_t^\xi$ is a very complex task since both matrices have entries which are time dependent, hence, generalizations of the approaches by Breitung and Tenhofen (2011) or Bai and Liao (2016), described above, are not straightforward.

9 Concluding remarks

In this paper, we have shown two main results. First, when estimating an approximate static or dynamic factor model by QML, possibly via the EM algorithm, and using a mis-specified log-likelihood for an exact factor model, the estimated loadings are asymptotically equivalent to the loadings estimated via PC analysis. Why should then we be using the QML estimator rather than the PC estimator of the loadings? The main reason put forward by the literature seems to be the possibility of easily imposing constraints on the loadings, e.g., incorporating prior beliefs.

Second, we have also shown that the factors estimated via WLS or via the Kalman filter/smoothing, are all asymptotically equivalent, and these estimators should be always preferred over the PC estimators as they address the heteroskedasticity of the idiosyncratic components so they might even be more efficient (depending on the amount of neglected cross-sectional correlations). Furthermore, in a time series setting the Kalman filter/smoothing should be preferred as it allows to easily deal with missing values and given its dynamic aggregation property it can handle very persistent or even non-stationary data without any modification.

Summing up, although the QML plus WLS estimation approach, proposed by Bai and Li (2016), is asymptotically equivalent to the EM plus Kalman smoother approach, proposed by Doz et al. (2012), the latter is likely to produce estimated factors which enjoy better final sample properties and is a more flexible approach for dealing with complex economic datasets. Relevant examples of the use of this estimation technique are: (i) counterfactual analysis (Giannone et al., 2019); (ii) conditional forecasts (Bańbura et al., 2015); (iii) nowcasting (Giannone et al., 2008); (iv) dealing with data irregularly spaced in time (Bańbura and Modugno, 2014); (v) imposing a block-specific factor structure on the loadings (Coroneo et al., 2016); (vi) building low and high frequency indicators of the economic activity (Barigozzi and Luciani, 2021; Ng and Scanlan, 2023); (vii) impulse response analysis (Luciani, 2015); (viii) risk management (Linton et al., 2022); (ix) firm-level or household-level repeated cross-sections of data (Barigozzi and Pellegrino, 2023).

A Kalman filter and Kalman smoother

The following iterations are stated assuming that the true value of the parameters φ is given and for given initial conditions $\mathbf{F}_{0|0}$ and $\mathbf{P}_{0|0}$.

The Kalman filter is based on the forward iterations for $t = 1, \dots, T$:

$$\begin{aligned}\mathbf{F}_{t|t-1} &= \mathbf{A}\mathbf{F}_{t-1|t-1}, \\ \mathbf{P}_{t|t-1} &= \mathbf{A}\mathbf{P}_{t-1|t-1}\mathbf{A}' + \mathbf{H}\mathbf{H}', \\ \mathbf{F}_{t|t} &= \mathbf{F}_{t|t-1} + \mathbf{P}_{t|t-1}\mathbf{\Lambda}'(\mathbf{\Lambda}\mathbf{P}_{t|t-1}\mathbf{\Lambda}' + \mathbf{\Sigma}^\xi)^{-1}(\mathbf{x}_t - \bar{\mathbf{x}} - \mathbf{\Lambda}\mathbf{F}_{t|t-1}), \\ \mathbf{P}_{t|t} &= \mathbf{P}_{t|t-1} - \mathbf{P}_{t|t-1}\mathbf{\Lambda}'(\mathbf{\Lambda}\mathbf{P}_{t|t-1}\mathbf{\Lambda}' + \mathbf{\Sigma}^\xi)^{-1}\mathbf{\Lambda}\mathbf{P}_{t|t-1}.\end{aligned}$$

Alternatively for $t = 1, \dots, T$ (by Woodbury formula)

$$\begin{aligned}\mathbf{F}_{t|t} &= \mathbf{F}_{t|t-1} + (\mathbf{\Lambda}'(\mathbf{\Sigma}^\xi)^{-1}\mathbf{\Lambda} + \mathbf{P}_{t|t-1}^{-1})^{-1}\mathbf{\Lambda}'(\mathbf{\Sigma}^\xi)^{-1}(\mathbf{x}_t - \bar{\mathbf{x}} - \mathbf{\Lambda}\mathbf{F}_{t|t-1}), \\ \mathbf{P}_{t|t} &= \mathbf{P}_{t|t-1} - (\mathbf{\Lambda}'(\mathbf{\Sigma}^\xi)^{-1}\mathbf{\Lambda} + \mathbf{P}_{t|t-1}^{-1})^{-1}\mathbf{\Lambda}'(\mathbf{\Sigma}^\xi)^{-1}\mathbf{\Lambda}\mathbf{P}_{t|t-1}.\end{aligned}$$

The Kalman Smoother is then based on the backward iterations for $t = T, \dots, 1$:

$$\begin{aligned}\mathbf{F}_{t|T} &= \mathbf{F}_{t|t} + \mathbf{P}_{t|t} \mathbf{A}' \mathbf{P}_{t+1|t}^{-1} (\mathbf{F}_{t+1|T} - \mathbf{F}_{t+1|t}), \\ \mathbf{P}_{t|T} &= \mathbf{P}_{t|t} + \mathbf{P}_{t|t} \mathbf{A}' \mathbf{P}_{t+1|t}^{-1} (\mathbf{P}_{t+1|T} - \mathbf{P}_{t+1|t}) \mathbf{P}_{t+1|t}^{-1} \mathbf{A} \mathbf{P}_{t|t}.\end{aligned}$$

Alternatively for $t = T, \dots, 1$ (see, e.g., [Durbin and Koopman, 2012](#), Chapter 4.4, pp.87-91)

$$\begin{aligned}\mathbf{F}_{t|T} &= \mathbf{F}_{t|t-1} + \mathbf{P}_{t|t-1} \mathbf{r}_{t-1}, \\ \mathbf{r}_{t-1} &= \mathbf{\Lambda}' (\mathbf{\Lambda} \mathbf{P}_{t|t-1} \mathbf{\Lambda}' + \mathbf{\Sigma}^\xi)^{-1} (\mathbf{x}_t - \bar{\mathbf{x}} - \mathbf{\Lambda} \mathbf{F}_{t|t-1}) + \mathbf{L}'_t \mathbf{r}_t, \\ \mathbf{P}_{t|T} &= \mathbf{P}_{t|t-1} (\mathbf{I}_r - \mathbf{N}_{t-1} \mathbf{P}_{t|t-1}), \\ \mathbf{N}_{t-1} &= \mathbf{\Lambda}'_n (\mathbf{\Lambda} \mathbf{P}_{t|t-1} \mathbf{\Lambda}' + \mathbf{\Sigma}^\xi)^{-1} \mathbf{\Lambda} + \mathbf{L}'_t \mathbf{N}_t \mathbf{L}_t, \\ \mathbf{L}_t &= \mathbf{A} - \mathbf{A} \mathbf{P}_{t|t-1} \mathbf{\Lambda}' (\mathbf{\Lambda} \mathbf{P}_{t|t-1} \mathbf{\Lambda}' + \mathbf{\Sigma}^\xi)^{-1} \mathbf{\Lambda}, \\ \mathbf{C}_{t,t+1|T} &= \mathbf{P}_{t|t-1} \mathbf{L}'_t (\mathbf{I}_r - \mathbf{N}_t \mathbf{P}_{t+1|t}), \quad \mathbf{C}_{t,t-1|T} = \mathbf{C}'_{t,t+1|T},\end{aligned}$$

where $\mathbf{r}_T = \mathbf{0}_r$, $\mathbf{N}_T = \mathbf{0}_r$.

References

- Ahn, S. C. and A. R. Horenstein (2013). Eigenvalue ratio test for the number of factors. *Econometrica* 81, 1203–1227.
- Aït-Sahalia, Y. and D. Xiu (2017). Using principal component analysis to estimate a high dimensional factor model with high-frequency data. *Journal of Econometrics* 201, 384–399.
- Alessi, L., M. Barigozzi, and M. Capasso (2010). Improved penalization for determining the number of factors in approximate static factor models. *Statistics and Probability Letters* 80, 1806–1813.
- Amemiya, Y., W. A. Fuller, and S. G. Pantula (1987). The asymptotic distributions of some estimators for a factor analysis model. *Journal of Multivariate Analysis* 22, 51–64.
- Anderson, B. D. O. and M. Deistler (2008). Generalized linear dynamic factor models-A structure theory. In *Proceedings of the 47th IEEE Conference on Decision and Control*, pp. 1980–1985.
- Anderson, B. D. O. and J. B. Moore (1979). *Optimal Filtering*. Dover Publications, Inc.
- Anderson, T. W. (2003). *An Introduction to Multivariate Statistical Analysis*. Wiley series in Probability and Mathematical Statistics.
- Anderson, T. W. and Y. Amemiya (1988). The asymptotic normal distribution of estimators in factor analysis under general conditions. *The Annals of Statistics* 16, 759–771.
- Anderson, T. W. and H. Rubin (1956). Statistical inference in factor analysis. In *Proceedings of the third Berkeley symposium on mathematical statistics and probability*, Volume 5, pp. 111–150.
- Andrews, D. W. (1991). Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica* 59, 817–858.
- Bai, J. (2003). Inferential theory for factor models of large dimensions. *Econometrica* 71, 135–171.
- Bai, J. and K. Li (2012). Statistical analysis of factor models of high dimension. *The Annals of Statistics* 40, 436–465.
- Bai, J. and K. Li (2016). Maximum likelihood estimation and inference for approximate factor models of high dimension. *The Review of Economics and Statistics* 98, 298–309.
- Bai, J. and Y. Liao (2016). Efficient estimation of approximate factor models via penalized maximum likelihood. *Journal of Econometrics* 191, 1–18.
- Bai, J. and S. Ng (2002). Determining the number of factors in approximate factor models. *Econometrica* 70,

191–221.

- Bai, J. and S. Ng (2013). Principal components estimation and identification of static factors. *Journal of Econometrics* 176, 18–29.
- Bai, J. and S. Ng (2020). Simpler proofs for approximate factor models of large dimensions. Technical Report arXiv:2008.00254.
- Bai, J. and S. Ng (2021). Approximate factor models with weaker loadings. Technical Report arXiv:2109.03773.
- Bakhshizadeh, M., A. Maleki, and V. H. de la Pena (2020). Sharp concentration results for heavy-tailed distributions. Technical Report arXiv:2003.13819.
- Bañbura, M., D. Giannone, and M. Lenza (2015). Conditional forecasts and scenario analysis with vector autoregressions for large cross-sections. *International Journal of Forecasting* 31, 739–756.
- Bañbura, M. and M. Modugno (2014). Maximum likelihood estimation of factor models on datasets with arbitrary pattern of missing data. *Journal of Applied Econometrics* 29, 133–160.
- Barigozzi, M. (2022). On estimation and inference of large approximate dynamic factor models via the principal component analysis. Technical report. arXiv:2211.01921.
- Barigozzi, M. and M. Hallin (2020). Generalized dynamic factor models and volatilities: Consistency, rates, and prediction intervals. *Journal of Econometrics* 216, 4–34.
- Barigozzi, M. and M. Luciani (2019). Quasi maximum likelihood estimation and inference of large approximate dynamic factor models via the EM algorithm. Technical Report arXiv:1910.03821.
- Barigozzi, M. and M. Luciani (2021). Measuring the output gap using large datasets. *The Review of Economics and Statistics*. forthcoming.
- Barigozzi, M. and A. Moneta (2016). Identifying the independent sources of consumption variation. *Journal of Applied Econometrics* 31, 420–449.
- Barigozzi, M. and F. Pellegrino (2023). Multidimensional dynamic factor models. Technical report. arXiv:2301.12499.
- Bartholomew, D. J., M. Knott, and I. Moustaki (2011). *Latent variable models and factor analysis: A unified approach*. John Wiley & Sons.
- Bartlett, M. S. (1937). The statistical conception of mental factors. *British Journal of Psychology* 28, 97–104.
- Bartlett, M. S. (1938). Methods of estimating mental factors. *Nature* 141(3570), 609–610.
- Bernanke, B. S., J. Boivin, and P. S. Elias (2005). Measuring the effects of monetary policy: A Factor-Augmented Vector Autoregressive (FAVAR) approach. *The Quarterly Journal of Economics* 120, 387–422.
- Bolthausen, E. (1982). On the central limit theorem for stationary mixing random fields. *The Annals of Probability* 10, 1047–1050.
- Bolthausen, E. (1984). An estimate of the remainder in a combinatorial central limit theorem. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* 66, 379–386.
- Breitung, J. and J. Tenhofen (2011). GLS estimation of dynamic factor models. *Journal of the American Statistical Association* 106, 1150–1166.
- Chamberlain, G. and M. Rothschild (1983). Arbitrage, factor structure, and mean-variance analysis on large asset markets. *Econometrica* 51, 1281–1304.
- Chernozhukov, V., D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, W. Newey, and J. Robins (2018). Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal* 21, C1–C68.
- Choi, I. (2012). Efficient estimation of factor models. *Econometric Theory* 28, 274–308.
- Cochrane, D. and G. H. Orcutt (1949). Application of least squares regression to relationships containing autocorrelated error terms. *Journal of the American statistical association* 44, 32–61.
- Connor, G., R. A. Korajczyk, and O. Linton (2006). The common and specific components of dynamic volatility. *Journal of Econometrics* 132, 231–255.

- Coroneo, L., D. Giannone, and M. Modugno (2016). Unspanned macroeconomic factors in the yield curve. *Journal of Business and Economic Statistics* 34, 472–485.
- D’Agostino, A. and D. Giannone (2012). Comparing alternative predictors based on large-panel factor models. *Oxford Bulletin of Economics and Statistics* 74, 306–326.
- De Jong, P. (1989). Smoothing and interpolation with the state-space model. *Journal of the American Statistical Association* 84, 1085–1088.
- De Mol, C., D. Giannone, and L. Reichlin (2008). Forecasting using a large number of predictors: Is bayesian shrinkage a valid alternative to principal components? *Journal of Econometrics* 146, 318–328.
- Delle Monache, D. and I. Petrella (2019). Efficient matrix approach for classical inference in state space models. *Economics Letters* 181, 22–27.
- Dempster, A. P., N. M. Laird, and D. B. Rubin (1977). Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 39, 1–38.
- Doz, C., D. Giannone, and L. Reichlin (2011). A two-step estimator for large approximate dynamic factor models based on Kalman filtering. *Journal of Econometrics* 164, 188–205.
- Doz, C., D. Giannone, and L. Reichlin (2012). A quasi maximum likelihood approach for large approximate dynamic factor models. *The Review of Economics and Statistics* 94(4), 1014–1024.
- Dunsmuir, W. (1979). A central limit theorem for parameter estimation in stationary vector time series and its application to models for a signal observed with noise. *The Annals of Statistics* 7, 490–506.
- Durbin, J. and S. J. Koopman (2012). *Time Series Analysis by State Space Methods*. Oxford University Press.
- Fan, J., Y. Liao, and M. Mincheva (2013). Large covariance estimation by thresholding principal orthogonal complements. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 75, 603–680.
- Fiorentini, G., A. Galesi, and E. Sentana (2018). A spectral EM algorithm for dynamic factor models. *Journal of Econometrics* 205, 249–279.
- Forni, M., D. Giannone, M. Lippi, and L. Reichlin (2009). Opening the black box: Structural factor models versus structural VARs. *Econometric Theory* 25, 1319–1347.
- Forni, M., M. Hallin, M. Lippi, and L. Reichlin (2000). The Generalized Dynamic Factor Model: Identification and estimation. *The Review of Economics and Statistics* 82, 540–554.
- Freyaldenhoven, S. (2022). Factor models with local factors – determining the number of relevant factors. *Journal of Econometrics* 229, 80–102.
- Gao, Z., J. Guo, and Y. Ma (2021). A note on statistical analysis of factor models of high dimension. *Science China Mathematics* 64, 1905–1916.
- Geweke, J. F. and K. J. Singleton (1981). Latent variable models for time series: A frequency domain approach with an application to the permanent income hypothesis. *Journal of Econometrics* 17, 287–304.
- Ghahramani, Z. and G. Hinton (1996). Parameter estimation for linear dynamical systems. Technical report, Cambridge University. mimeo.
- Giannone, D., M. Lenza, and L. Reichlin (2019). Money, credit, monetary policy and the business cycle in the euro area: what has changed since the crisis? *International Journal of Central Banking* 15, 137–173.
- Giannone, D., L. Reichlin, and D. Small (2008). Nowcasting: The real-time informational content of macroeconomic data. *Journal of Monetary Economics* 55, 665–676.
- Hannan, E. J. (1970). *Multiple time series*. John Wiley & Sons.
- Hannan, E. J. and M. Deistler (2012). *The statistical theory of linear systems*. SIAM.
- Harvey, A. C. (1990). *Forecasting, structural time series models and the Kalman filter*. Cambridge University Press.
- Harvey, A. C. and S. Peters (1990). Estimation procedures for structural time series models. *Journal of Forecasting* 9, 89–108.
- Ibragimov, I. A. (1962). Some limit theorems for stationary processes. *Theory of Probability and its Applications* 7, 349–382.

- Jöreskog, K. G. (1969). A general approach to confirmatory maximum likelihood factor analysis. *Psychometrika* 34, 183–202.
- Jungbacker, B. and S. J. Koopman (2015). Likelihood-based dynamic factor analysis for measurement and forecasting. *Econometrics Journal* 18, C1–C21.
- Kálmán, R. E. (1960). A new approach to linear filtering and prediction problems. *Journal of Basic Engineering* 82, 35–45.
- Kim, D. and J. Fan (2019). Factor GARCH-Itô models for high-frequency data with application to large volatility matrix prediction. *Journal of Econometrics* 208, 395–417.
- Kolmogorov, A. N. (1941). Interpolation und Extrapolation von stationären zufälligen Folgen. *Izvestiya Akademii Nauk. Seriya Matematicheskaya* 5, 3–14. Translated in “Selected Works of A. N. Kolmogorov”, (1992), A. N. Shirayev (ed.), Springer.
- Kuchibhotla, A. K. and A. Chakraborty (2022). Moving beyond sub-gaussianity in high-dimensional statistics: Applications in covariance estimation and linear regression. *Information and Inference: A Journal of the IMA* 11, 1389–1456.
- Lawley, D. N. (1942). Further investigations in factor estimation. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics* 61, 176–185.
- Lawley, D. N. and A. E. Maxwell (1971). *Factor Analysis as a Statistical Method*. Butterworths, London.
- Lehmann, E. L. and G. Casella (2006). *Theory of point estimation*. Springer Science & Business Media.
- Li, K., Q. Li, and L. Lu (2018). Quasi maximum likelihood analysis of high dimensional constrained factor models. *Journal of Econometrics* 206, 574–612.
- Lin, J. and G. Michailidis (2020). System identification of high-dimensional linear dynamical systems with serially correlated output noise components. *IEEE Transactions on Signal Processing* 68, 5573–5587.
- Linton, O. B., H. Tang, and J. Wu (2022). A model for daily global stock market returns. Technical Report arXiv:2202.03638.
- Lippi, M. (2018). Frequency-domain approach in high-dimensional dynamic factor models. In *Oxford Research Encyclopedia of Economics and Finance*. Oxford University Press.
- Lippi, M. (2019). Time-domain approach in high-dimensional dynamic factor models. In *Oxford Research Encyclopedia of Economics and Finance*. Oxford University Press.
- Luciani, M. (2015). Monetary policy and the housing market: A structural factor analysis. *Journal of Applied Econometrics* 30, 199–218.
- Mardia, K., J. Kent, and J. Bibby (1979). *Multivariate analysis*.
- McLachlan, G. and T. Krishnan (2007). *The EM algorithm and extensions*, Volume 382. John Wiley & Sons.
- Meng, X.-L. and D. B. Rubin (1993). Maximum likelihood estimation via the ECM algorithm: A general framework. *Biometrika* 80, 267–278.
- Meng, X.-L. and D. B. Rubin (1994). On the global and componentwise rates of convergence of the EM algorithm. *Linear Algebra and its Applications* 199, 413–425.
- Merlevède, F., M. Peligrad, and E. Rio (2011). A Bernstein type inequality and moderate deviations for weakly dependent sequences. *Probability Theory and Related Fields* 151, 435–474.
- Mikosch, T. and A. V. Nagaev (1998). Large deviations of heavy-tailed sums with applications in insurance. *Extremes* 1, 81–110.
- Newey, W. K. and K. D. West (1987). A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica* 55, 703–708.
- Neyman, J. (1979). C (α) tests and their use. *Sankhyā: The Indian Journal of Statistics, Series A* 41, 1–21.
- Ng, C. T., C. Y. Yau, and N. H. Chan (2015). Likelihood inferences for high-dimensional factor analysis of time series with applications in finance. *Journal of Computational and Graphical Statistics* 24(3), 866–884.
- Ng, S. and S. Scanlan (2023). Constructing high frequency economic indicators by imputation. Technical report. arXiv:2303.01863.

- Onatski, A. (2010). Determining the number of factors from empirical distribution of eigenvalues. *The Review of Economics and Statistics* 92, 1004–1016.
- Onatski, A. (2012). Asymptotics of the principal components estimator of large factor models with weakly influential factors. *Journal of Econometrics* 168, 244–258.
- Poignard, B. and Y. Terada (2020). Statistical analysis of sparse approximate factor models. *Electronic Journal of Statistics* 14, 3315–3365.
- Poncela, P. and E. Ruiz (2015). More is not always better: Kalman filtering in dynamic factor models. In S. J. Koopman and N. Shephard (Eds.), *Unobserved Components and Time Series Econometrics*. Oxford Scholarship Online.
- Quah, D. and T. J. Sargent (1993). A dynamic index model for large cross sections. In *Business cycles, indicators and forecasting*. University of Chicago Press.
- Rao, C. R. (1973). *Linear statistical inference and its applications*. Wiley New York.
- Rauch, H. (1963). Solutions to the linear smoothing problem. *IEEE Transactions on Automatic Control* 8, 371–372.
- Reis, R. and M. W. Watson (2010). Relative goods’ prices, pure inflation, and the Phillips correlation. *American Economic Journal Macroeconomics* 2, 128–157.
- Rubin, D. B. and D. T. Thayer (1982). EM algorithms for ML factor analysis. *Psychometrika* 47, 69–76.
- Ruiz, E. and P. Poncela (2022). Factor extraction in Dynamic Factor Models: Using Kalman Filter and Principal Components in practice. *Foundations and Trends in Econometrics*. forthcoming.
- Ruud, P. A. (1991). Extensions of estimation methods using the EM algorithm. *Journal of Econometrics* 49(3), 305–341.
- Sargent, T. J. and C. A. Sims (1977). Business cycle modeling without pretending to have too much a priori economic theory. In *New methods in business cycle research*. Federal Reserve Bank of Minneapolis.
- Scott, J. T. (1966). Factor analysis and regression. *Econometrica* 34, 552–562.
- Shumway, R. H. and D. S. Stoffer (1982). An approach to time series smoothing and forecasting using the EM algorithm. *Journal of Time Series Analysis* 3, 253–264.
- Solari, M. E. (1969). The “maximum likelihood solution” of the problem of estimating a linear functional relationship. *Journal of the Royal Statistical Society: Series B (Methodological)* 31, 372–375.
- Spearman, C. (1904). General intelligence objectively determined and measured. *American Journal of Psychology* 15, 201–293.
- Stock, J. H. and M. W. Watson (1989). New indexes of coincident and leading economic indicators. In O. J. Blanchard and S. Fischer (Eds.), *NBER Macroeconomics Annual 1989*. MIT press.
- Stock, J. H. and M. W. Watson (1991). A probability model of the coincident economic indicators. In K. Lahiri and G. Moore (Eds.), *Leading Economic Indicators: New Approaches and Forecasting Records*, pp. 63–90. Cambridge University Press.
- Stock, J. H. and M. W. Watson (2002a). Forecasting using principal components from a large number of predictors. *Journal of the American Statistical Association* 97, 1167–1179.
- Stock, J. H. and M. W. Watson (2002b). Macroeconomic forecasting using diffusion indexes. *Journal of Business and Economic Statistics* 20, 147–162.
- Stock, J. H. and M. W. Watson (2016). Dynamic Factor Models, Factor-Augmented Vector Autoregressions, and Structural Vector Autoregressions in Macroeconomics. In J. B. Taylor and H. Uhlig (Eds.), *Handbook of Macroeconomics*, Volume 2, pp. 415–525. Elsevier.
- Stone, R. (1945). The analysis of market demand. *Journal of the Royal Statistical Society* 108, 286–391.
- Sundberg, R. (1974). Maximum likelihood theory for incomplete data from an exponential family. *Scandinavian Journal of Statistics* 1, 49–58.
- Sundberg, R. (1976). An iterative method for solution of the likelihood equations for incomplete data from exponential families. *Communication in Statistics-Simulation and Computation* 5, 55–64.

- Sundberg, R. (2019). *Statistical modelling by exponential families*. Cambridge University Press.
- Sundberg, R. and U. Feldmann (2016). Exploratory factor analysis—Parameter estimation and scores prediction with high-dimensional data. *Journal of Multivariate Analysis* 148, 49–59.
- Terada, Y. (2014). Strong consistency of reduced k-means clustering. *Scandinavian Journal of Statistics* 41, 913–931.
- Thomson, G. H. (1936). Some points of mathematical technique in the factorial analysis of ability. *Journal of Educational Psychology* 27, 37–54.
- Thomson, G. H. (1951). *The factorial analysis of human ability*. London University Press.
- Tipping, M. E. and C. M. Bishop (1999). Probabilistic principal component analysis. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 61, 611–622.
- Trapani, L. (2018). A randomized sequential procedure to determine the number of factors. *Journal of the American Statistical Association* 113, 1341–1349.
- Uematsu, Y. and T. Yamagata (2021). Inference in sparsity-induced weak factor models. *Journal of Business & Economic Statistics*, 1–14.
- van der Vaart, A. W. (2000). *Asymptotic Statistics*. Cambridge University Press.
- Wang, S., H. Yang, and C. Yao (2019). On the penalized maximum likelihood estimation of high-dimensional approximate factor model. *Computational Statistics* 34, 819–846.
- Watson, M. W. and R. F. Engle (1983). Alternative algorithms for the estimation of dynamic factor, mimic and varying coefficients regression models. *Journal of Econometrics* 23, 385–400.
- White, H. (1980). A heteroskedasticity-consistent covariance matrix estimator and a direct test for heteroskedasticity. *Econometrica* 48, 817–838.
- Whittle, P. (1952). On principal components and least square methods of factor analysis. *Scandinavian Actuarial Journal* 1952, 223–239.
- Wiener, N. (1949). *Extrapolation, Interpolation, and Smoothing of Stationary Time Series*. MIT press.
- Wu, J. C. F. (1983). On the convergence properties of the EM algorithm. *The Annals of Statistics* 11, 95–103.
- Young, G. (1940). Maximum likelihood estimation and factor analysis. *Psychometrika* 6, 49–53.
- Zhang, D. and W. B. Wu (2021). Convergence of covariance and spectral density estimates for high-dimensional locally stationary processes. *The Annals of Statistics* 49(1), 233–254.