

CONES OF DIVISORS ON \mathbb{P}^3 BLOWN UP AT EIGHT VERY GENERAL POINTS

ISABEL STENGER AND ZHIXIN XIE

ABSTRACT. Let X be the blowup of \mathbb{P}^3 at eight very general points. We give a complete description of its nef and effective cones. Moreover, we show that there exists a rational polyhedral fundamental domain for the action of a certain Weyl group on the effective movable cone of X .

CONTENTS

1. Introduction	1
2. Notation and preliminaries	4
3. Threefolds with nef anticanonical class	6
4. Blowup of \mathbb{P}^3 at eight points	9
5. The cone of curves and the nef cone	14
6. The effective cone	17
7. The movable cone	26
8. Other examples	32
References	34

1. INTRODUCTION

Due to Mori's fundamental insights from the 1980s, the minimal model program of a projective variety X is closely intertwined with the structure of the nef and the movable cones of X . When X is a Fano variety, or more generally a Mori dream space, Hu and Keel proved in [HK00] that the nef cone is rational polyhedral, and the movable cone is the union of nef cones of finitely many small modifications of X .

When X is a Calabi-Yau variety, these cones have much more complicated structure. Nevertheless, the Morrison-Kawamata cone conjecture ([Mor93], [Kaw97]) predicts that the groups of automorphisms and pseudo-automorphisms act respectively with rational polyhedral fundamental domain on the nef and the movable cones of the variety. Despite of being verified in several instances, the cone conjecture for Calabi-Yau varieties remains widely open; we refer the reader to [LOP18] for a detailed survey on

2020 *Mathematics Subject Classification*:

Keywords: 14E07, 14E30.

this topic. In dimension three, Kawamata proved in [Kaw97] that the movable effective cone of X decomposes into a union of nef cones of its marked minimal models. However, in contrast to the case of Mori dream spaces, this decomposition is usually not finite.

In general, it is difficult to describe the nef and the movable cones of a variety, even under weaker assumptions on the anticanonical class $-K_X$. For instance, varieties with nef anticanonical class interpolate between Fano varieties and varieties with numerically trivial canonical class. We may thus expect a similar description of the nef and the movable cones of this class of varieties. We note that when $-K_X$ is semi-ample, there are several examples where the nef and the movable cones are fully described, see Subsection 1.2.

1.1. Main results. Considering a smooth projective threefold with nef anticanonical class, we obtain a similar decomposition of its effective movable cone as in the Calabi-Yau case:

Proposition 1.1. *Let X be a smooth projective threefold with $-K_X$ nef. Then*

$$\overline{\text{Mov}}^+(X) \supset \overline{\text{Mov}}^e(X) = \bigcup_{\varphi: X \dashrightarrow X' \text{ a finite sequence of flops}} \varphi^*(\text{Nef}^e(X')).$$

We refer to Theorem 3.5 for a decomposition of the effective movable cone for a more general class of varieties, formulated in the setting of generalised pairs.

We further study a specific example of a threefold whose anticanonical class is nef but **not** semi-ample. For this example, we give a complete description of its nef and effective cones, and we show that a certain Weyl group acts with a rational polyhedral fundamental domain on the effective movable cone.

Let X be the blowup of \mathbb{P}^3 at eight very general points. Then $-K_X$ is nef and non semi-ample, as proven in [LO16]. Moreover, it is shown in [CLO16] that the cone of curves of X is rational polyhedral. As the dual cone, the nef cone of X is also rational polyhedral and we obtain the following:

Theorem 1.2. *Let X be the blowup of \mathbb{P}^3 at eight very general points. Then the nef cone $\text{Nef}(X)$ is rational polyhedral. Furthermore, we have $\text{Nef}^e(X) = \text{Nef}(X) = \text{Nef}^+(X)$ and there exists a rational polyhedral fundamental domain for the action of $\text{Aut}(X)$ on $\text{Nef}^e(X)$.*

We construct an explicit set of extremal rays generating the nef cone $\text{Nef}(X)$ in Lemma 5.3.

For the structure of the movable and the effective cones, we study the action of a Weyl group \mathcal{W} on the Néron-Severi group $N^1(X)$ introduced by Dolgachev ([Dol83]); see Subsection 4.1 for the definition of the group \mathcal{W} and its action. We obtain the following description for the movable cone:

Theorem 1.3. *Let X be the blowup of \mathbb{P}^3 at eight very general points and let \mathcal{W} be the Weyl group of the Dynkin diagram $T_{2,4,4}$ acting on $N^1(X)$. Then:*

- (i) *We have $\overline{\text{Mov}}^e(X) = \overline{\text{Mov}}^+(X) = \overline{\text{Mov}}(X)$.*
- (ii) *The effective movable cone $\overline{\text{Mov}}^e(X)$ is not rational polyhedral.*
- (iii) *There exists a rational polyhedral cone which is a fundamental domain for the action of \mathcal{W} on $\overline{\text{Mov}}^e(X)$.*

As for the effective cone, we note that X is an example of a blowup of a projective space at general points such that the corresponding Weyl group \mathcal{W} is infinite (see Subsection 1.3). This implies that the effective cone $\text{Eff}(X)$ is **not** rational polyhedral. However, we are able to determine an explicit set of infinitely many extremal rays generating $\text{Eff}(X)$ as convex cone and to show that $\text{Eff}(X)$ is closed:

Theorem 1.4. *Let X and \mathcal{W} be as in Theorem 1.3. Then $\text{Eff}(X) = \overline{\text{Eff}}(X)$, and $\text{Eff}(X)$ is the convex cone generated by the classes in $\mathcal{W} \cdot E_8$ and the class $-\frac{1}{2}K_X$, where E_8 is an exceptional divisor.*

1.2. Cone conjecture on Calabi-Yau pairs. Totaro introduced a generalisation of the cone conjecture to klt Calabi-Yau pairs (see [Tot08]) and verified the conjecture in dimension two (see [Tot10]). Moreover, using the example of \mathbb{P}^2 blown up at nine very general points, Totaro showed that the conjecture is in general not true for log canonical Calabi-Yau pairs. However, using a different group action on the nef effective cone, Li verified in [Li22] a variant of the cone conjecture on the nef effective cone for a class of log canonical Calabi-Yau surface pairs.

When X has a semi-ample anticanonical class, one can relate the description of the nef and the movable cones of X to the cone conjecture for klt Calabi-Yau pairs: since $-K_X$ is semi-ample, choosing m large enough so that $-mK_X$ is basepoint-free, a general divisor $D \in |-mK_X|$ is smooth. Thus putting $\Delta = \frac{1}{m}D$, the pair (X, Δ) is a klt Calabi-Yau pair. For many examples of threefolds with semi-ample anticanonical class, the cone conjecture for klt Calabi-Yau pairs has been verified; see [PS12, PS15] and [CPS13, CPS19].

Now let X be \mathbb{P}^3 blown up at eight very general points. Then $-K_X$ is nef, not semi-ample and choosing a general divisor $\Delta \in |-K_X|$, the pair (X, Δ) is a log canonical Calabi-Yau pair. Theorem 1.2 shows that the pair (X, Δ) verifies the statement of the cone conjecture on the nef cone. For the movable cone, one major difficulty is to describe the pseudo-automorphisms of X . We remark that the pseudo-automorphism group of the blowup of \mathbb{P}^3 at very general points is not well understood; see [Dol11, Remark 1] and [BH15, Section 8]. Our strategy to overcome this problem is to study the cone conjecture with respect to a different subgroup of $\text{GL}(N^1(X))$ preserving $\overline{\text{Mov}}^e(X)$, which is related to the geometry of X . In this sense,

Theorem 1.3 can be seen as a variant of the cone conjecture for the effective movable cone.

1.3. Blowups of \mathbb{P}^n and the Weyl group. If $X_{n,k}$ denotes the blowup of \mathbb{P}^n in $k > n + 1$ points in general position, Dolgachev [Dol83] realised a root system in the group $N^1(X_{n,k})$ and interpreted the action of the Weyl group $\mathcal{W}_{n,k}$ corresponding to this root system by means of Cremona transformations. Mukai [Muk01], Castravet and Tevelev [CT06] showed that the semi-group of effective divisor classes in $\text{Pic}(X_{n,k})$ is finitely generated if and only if

$$\frac{1}{n+1} + \frac{1}{k-n-1} > \frac{1}{2}.$$

When the above inequality holds, the corresponding Weyl group $\mathcal{W}_{n,k}$ is finite and the effective cone $\text{Eff}(X_{n,k})$ is spanned as a convex cone by the finitely many classes in $\mathcal{W}_{n,k} \cdot E_k$, where E_k is an exceptional divisor of the blowup $X_{n,k}$ of \mathbb{P}^n . Thus $\text{Eff}(X_{n,k})$ is a rational polyhedral cone.

In the case of an infinite Weyl group, the effective cone $\text{Eff}(X_{n,k})$ is only poorly understood: Mukai showed in [Muk01] that the classes in the orbit $\mathcal{W}_{n,k} \cdot E_k$ are contained in any generating system of the effective cone. If $n = 3$, then $k = 8$ is the smallest number which yields an infinite Weyl group $\mathcal{W} = \mathcal{W}_{3,8}$, and hence an effective cone which is not rational polyhedral. Theorem 1.4 shows that the infinitely many classes in the \mathcal{W} -orbit of an exceptional divisor are not enough to span the effective cone, but together with the class $-\frac{1}{2}K_X$ we obtain a generating system of the effective cone.

Acknowledgments. We would like to thank Vladimir Lazić for his helpful comments and discussions leading to Theorem 3.5. We are grateful to Artie Prendergast-Smith for his inspiring ideas for the proof of Proposition 6.11. Moreover, we would like to thank Cécile Gachet, Andreas Höring, John Lesieutre and Susanna Zimmermann for useful discussions. The first author is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Project-ID 286237555 - TRR 195.

2. NOTATION AND PRELIMINARIES

In this paper we work over the field \mathbb{C} .

2.1. Notation. First we fix some notation.

Let X be a normal projective variety. Let $N^1(X)$ be the Néron-Severi group generated by the classes of Cartier divisors modulo numerical equivalence; let $N^1(X)_{\mathbb{R}} = N^1(X) \otimes \mathbb{R}$. Inside $N^1(X)_{\mathbb{R}}$, we have:

- the effective cone $\overline{\text{Eff}}(X)$ and the pseudoeffective cone $\overline{\text{Eff}}(X)$;
- the movable cone $\overline{\text{Mov}}(X)$, which is the closure of the cone generated by classes of effective Cartier divisors D such that the base locus of $|D|$ has codimension at least 2; and its interior $\text{Mov}(X)$;
- the effective movable cone $\overline{\text{Mov}}^e(X) := \overline{\text{Mov}}(X) \cap \overline{\text{Eff}}(X)$;
- the convex hull $\overline{\text{Mov}}^+(X)$ of all rational classes in $\overline{\text{Mov}}(X)$;

- the nef cone $\text{Nef}(X)$; and its interior $\text{Amp}(X)$;
- the effective nef cone $\text{Nef}^e(X) := \text{Nef}(X) \cap \text{Eff}(X)$;
- the convex hull $\text{Nef}^+(X)$ of all rational classes in $\text{Nef}(X)$.

We call an \mathbb{R} -divisor D *movable* if its class is in $\overline{\text{Mov}}(X)$.

We denote by $N_1(X)$ the group of 1-cycles modulo numerical equivalence and $N_1(X)_{\mathbb{R}} = N_1(X) \otimes \mathbb{R}$. Inside $N_1(X)_{\mathbb{R}}$, we consider the convex cone $\text{NE}(X)$ generated by effective 1-cycles and its closure, the cone of curves, $\overline{\text{NE}}(X)$.

We denote by $\text{Aut}(X)$ and $\text{PsAut}(X)$ the group of automorphisms and pseudo-automorphisms of X , that means birational maps $X \dashrightarrow X$ which are isomorphisms in codimension one. Any automorphism induces an isomorphism on $N^1(X)_{\mathbb{R}}$ which preserves the nef cone, whereas any pseudo-automorphism induces an isomorphism on $N^1(X)_{\mathbb{R}}$ preserving the effective and the movable cones. We denote by $\text{Aut}(X)^*$ and $\text{PsAut}(X)^*$ the image of $\text{Aut}(X)$ and $\text{PsAut}(X)$ in $\text{GL}(N^1(X)_{\mathbb{R}})$, respectively.

Finally, we say that an \mathbb{R} -divisor on X is an *NQC divisor* if it is a non-negative linear combination of nef \mathbb{Q} -divisors on X . The acronym NQC stands for *nef \mathbb{Q} -Cartier combinations*.

2.2. The Morrison-Kawamata cone conjecture. In this subsection we state a generalisation of the Morrison-Kawamata cone conjecture for klt Calabi-Yau pairs.

A *pair* consists of a normal variety X and an effective \mathbb{R} -divisor Δ on X such that the divisor $K_X + \Delta$ is \mathbb{R} -Cartier. Given a \mathbb{Q} -factorial projective klt (respectively log canonical) pair (X, Δ) , we say that the pair is *Calabi-Yau* if $K_X + \Delta$ is numerically trivial.

Conjecture 2.1. [Tot08] *Let (X, Δ) be a klt Calabi-Yau pair. Then:*

- There exists a rational polyhedral cone Π which is a fundamental domain for the action of $\text{Aut}(X)$ on $\text{Nef}^e(X) = \text{Nef}(X) \cap \text{Eff}(X)$, in the sense that*

$$\text{Nef}^e(X) = \bigcup_{g \in \text{Aut}(X)} g^* \Pi$$

with $\text{Int } \Pi \cap \text{Int } g^ \Pi = \emptyset$ unless $g^* = \text{id}$.*

- There exists a rational polyhedral cone Π' which is a fundamental domain for the action of $\text{PsAut}(X)$ on $\overline{\text{Mov}}^e(X) = \overline{\text{Mov}}(X) \cap \text{Eff}(X)$.*

Remark 2.2. We remark that in [Tot10] there exists a different formulation of the Morrison-Kawamata cone conjecture for klt Calabi-Yau pairs using the group $\text{Aut}(X, \Delta)$ (respectively $\text{PsAut}(X, \Delta)$) of automorphisms (respectively pseudo-automorphisms) of X preserving the boundary divisor Δ . As $\text{Aut}(X, \Delta) \subset \text{Aut}(X)$ and analogously for the pseudo-automorphisms, the version in [Tot10] is stronger than the one stated above.

A useful tool to show the existence of a fundamental domain for a group action on a convex cone is the following result of Looijenga:

Theorem 2.3. [Loo14, Theorem 4.1 and Application 4.15] *Let V be a real vector space with rational structure $V(\mathbb{Q})$, C an open strictly convex cone in V , and let G be a subgroup of $\mathrm{GL}(V)$ stabilising the cone C . Let C_+ denote the convex hull of $\overline{C} \cap V(\mathbb{Q})$. Then the following are equivalent:*

- (i) *There exists a rational polyhedral cone Π in C_+ with $C_+ = G \cdot \Pi$.*
- (ii) *There exists a rational polyhedral cone Π in C_+ with $C \subset G \cdot \Pi$.*

Moreover, if these conditions are satisfied, then there exists a rational polyhedral fundamental domain for the action of G on C_+ .

3. THREEFOLDS WITH NEF ANTICANONICAL CLASS

In this subsection we prove some general results for the movable cone of threefolds with nef anticanonical class.

Definition 3.1. Let (X, D) be a projective \mathbb{Q} -factorial pair. A *flopping contraction* is an extremal birational contraction $f: X \rightarrow Y$ to a normal variety Y such that the exceptional locus of f has codimension at least two in X and K_X is numerically f -trivial. If additionally $-(K_X + D)$ is f -ample, then the $(K_X + D)$ -flip of f is called the *D -flop*.

Proposition 3.2. *Let X be a smooth projective threefold with $-K_X$ nef and let $D \in \overline{\mathrm{Mov}}^e(X)$. Then there exists a finite sequence of D -flops $X \dashrightarrow X'$ such that the strict transform of D on X' is nef, where X' is a smooth projective threefold with $-K_{X'}$ nef.*

Proof. Let $D \in \overline{\mathrm{Mov}}^e(X)$ be a divisor on X which is not nef.

We first assume that D is a \mathbb{Q} -divisor. Since X is smooth, for sufficiently small $\epsilon > 0$, the pair $(X, \epsilon D)$ is klt. Since D is not nef, by [Xie20, Lemma 2.5] there exists a $(K_X + \epsilon D)$ -negative extremal ray Γ such that $D \cdot \Gamma < 0$. Let c_Γ be the contraction of the extremal ray Γ and let ℓ be a contracted curve.

Since D is movable, c_Γ does not contract any component of D , see for example [Dru11, Lemma 2.5]. Hence c_Γ is a small contraction, which implies $K_X \cdot \ell = 0$, as there is no flipping contraction for smooth threefolds. Hence there exists a D -flop $X \dashrightarrow X^+$ of the K_X -trivial small contraction c_Γ and the flopped threefold X^+ is again smooth by [Kol89, Theorem 2.4] with $-K_{X^+}$ nef.

Now if D is not a \mathbb{Q} -divisor, we can choose a small ample \mathbb{R} -divisor $D' \subset X$ such that $D + D' \in \overline{\mathrm{Mov}}^e(X)$ is a \mathbb{Q} -divisor which is not nef. By the same argument as in the previous paragraph, there exists a $(D + D')$ -flop $X \dashrightarrow X^+$, where X^+ is smooth with $-K_{X^+}$ nef. Since D' is ample, this is also a D -flop.

By repeating the above argument and by the termination of flops in dimension three ([KM98, Corollary 6.19]), we deduce that there exists a sequence of flops $\varphi: X \dashrightarrow X'$ such that $D' := \varphi_* D$ is nef and X' is smooth with $-K_{X'}$ nef. \square

As corollaries, we obtain the following two results. The proof of the first result uses the notion of generalised pairs which are introduced below.

Corollary 3.3. *Let X be a smooth projective threefold with $-K_X$ nef. Then*

$$\overline{\text{Mov}}^e(X) \subset \overline{\text{Mov}}^+(X).$$

Proof. Let $D \in \overline{\text{Mov}}^e(X)$. By Proposition 3.2, there exists a finite sequence of flops $\varphi: X \dashrightarrow X'$ such that the strict transform $D' \subset X'$ of D is nef and X' is smooth.

Since X' is smooth, by replacing D' with $\epsilon D'$ for sufficiently small $\epsilon > 0$, the pair (X', D') is klt. Moreover, $-K_{X'}$ is an NQC divisor. Thus $(X', D' + (-K_{X'}))$ is an NQC klt generalised pair (see [HL22, Definition 2.15]) and we can apply [HL22, Proposition 3.20] to $K_{X'} + D' + (-K_{X'})$ and we obtain $D' \in \text{Nef}^+(X')$. Hence, we have $D = \varphi^* D' \in \overline{\text{Mov}}^+(X)$. \square

Corollary 3.4. *Let X be a smooth projective threefold with $-K_X$ nef. Then*

$$\overline{\text{Mov}}^e(X) = \bigcup_{\varphi: X \dashrightarrow X' \text{ a finite sequence of flops}} \varphi^*(\text{Nef}^e(X')).$$

Proof. Let $D \in \overline{\text{Mov}}^e(X)$. By Proposition 3.2 there exist a smooth threefold X' and a finite sequence of flops $\varphi: X \dashrightarrow X'$ such that the strict transform $D' \subset X'$ of D is nef and thus $D' \in \text{Nef}^e(X')$.

The other inclusion follows from the fact that the pullback of an effective nef divisor by an isomorphism in codimension one is an effective movable divisor. \square

We can now prove our first main result.

Proof of Proposition 1.1. The statement follows from Corollaries 3.3 and 3.4. \square

Proposition 1.1 is an important case which fits into a more general setting related to generalised pairs. These objects are introduced in [BH14, BZ16]: very roughly speaking, generalised pairs are couples $(X, B + M)$, where (X, B) is a usual pair and M is the pushforward of a nef divisor from a higher birational model of X . We refer to [HL22, LT22] for a precise definition and a more detailed discussion. When (X, B) is a projective \mathbb{Q} -factorial klt pair such that $-(K_X + B)$ is nef, one may take $M = -(K_X + B)$. Then M is an NQC divisor on X by [HL21, Corollary 3.6], and $(X, B + M)$ is an NQC klt generalised pair. We will now give a more general result formulated in the language of generalised pairs.

The following theorem is obtained in discussions with Vladimir Lazić.

Theorem 3.5. *Let $(X, B + M)$ be a projective \mathbb{Q} -factorial klt NQC generalised pair of dimension n such that $K_X + B + M \equiv 0$.*

Assume the existence of minimal models for smooth varieties in dimension n . Then

$$\overline{\text{Mov}}^+(X) \supset \overline{\text{Mov}}^e(X) = \bigcup_{\varphi: X \dashrightarrow X' \text{ an isomorphism in codimension 1}} \varphi^*(\text{Nef}^e(X')).$$

The conclusion holds unconditionally if:

- (i) $n \leq 4$, or
- (ii) $n = 5$ and X is uniruled.¹

Proof. Step 1. Since the pullback of an effective nef divisor by an isomorphism in codimension one is an effective movable divisor, we obtain

$$\bigcup_{\varphi: X \dashrightarrow X' \text{ an isomorphism in codimension 1}} \varphi^*(\text{Nef}^e(X')) \subset \overline{\text{Mov}}^e(X).$$

Step 2. In this step we show the following:

$$(1) \quad \overline{\text{Mov}}^e(X) \subset \bigcup_{\varphi: X \dashrightarrow X' \text{ an isomorphism in codimension 1}} \varphi^*(\text{Nef}^e(X')),$$

$$(2) \quad \overline{\text{Mov}}^e(X) \subset \overline{\text{Mov}}^+(X).$$

Let $D \in \overline{\text{Mov}}^e(X)$. By replacing D with ϵD for small enough $\epsilon > 0$, we have that $(X, (B + D) + M)$ is an NQC log canonical generalised pair, and

$$(3) \quad D \equiv K_X + (B + D) + M$$

as $K_X + B + M \equiv 0$. By [LT22, Theorem 1.2 and Corollary 1.3], we may run a $(K_X + (B + D) + M)$ -MMP with scaling of an ample divisor

$$\varphi: (X, (B + D) + M) \dashrightarrow (X', (B' + D') + M'),$$

where $B' = \varphi_* B$, $D' = \varphi_* D$, and the divisors M and M' are pushforwards of the same nef \mathbb{R} -divisors on a common birational model of X and X' . Since D is movable and by (3), we infer that this MMP does not contract any divisor. Moreover, $(X', (B' + D') + M')$ is a \mathbb{Q} -factorial NQC log canonical generalised pair and together with (3), we obtain that D' is nef.

Note that D' is also an effective divisor, since D is effective and φ is an isomorphism in codimension one. Thus we have $D = \varphi^* D' \in \varphi^*(\text{Nef}^e(X'))$, which proves (1).

Applying [HL22, Proposition 3.20] to $K_{X'} + (B' + D') + M'$, we have $D' \in \text{Nef}^+(X')$. Thus we obtain $D = \varphi^* D' \in \overline{\text{Mov}}^+(X)$, which proves (2).

Step 3. For $n \leq 4$, or for $n = 5$ and X is uniruled, we repeat verbatim *Step 2*, except that we invoke [LT22, Corollary 1.4] instead of [LT22, Theorem 1.2 and Corollary 1.3]. \square

¹By [LMP⁺22, Lemma 3.18], X is not uniruled if and only if X is canonical, $B = 0$ and $K_X \sim_{\mathbb{Q}} 0$.

4. BLOWUP OF \mathbb{P}^3 AT EIGHT POINTS

Let p_1, \dots, p_8 be eight very general points in \mathbb{P}^3 and let $\pi: X \rightarrow \mathbb{P}^3$ be the blowup of \mathbb{P}^3 at the points p_1, \dots, p_8 .

From this point on, we use the following notation for the blowup X of \mathbb{P}^3 at eight points:

- $E_i \subset X$ is the exceptional divisor over $p_i \in \mathbb{P}^3$, for $i = 1, \dots, 8$;
- $H \in \text{Pic}(X)$ is the pullback $\pi^* \mathcal{O}_{\mathbb{P}^3}(1)$;
- e_i is the class of line in E_i , for $i = 1, \dots, 8$;
- $h = H^2$ is the class of the transform of a general line in \mathbb{P}^3 ;
- ℓ_{ij} is the class of the transform of the line through the two points p_i and p_j , for $1 \leq i < j \leq 8$.

We have $N^1(X) = \mathbb{Z}H \oplus \mathbb{Z}E_1 \oplus \dots \oplus \mathbb{Z}E_8$ and $N_1(X) = \mathbb{Z}h \oplus \mathbb{Z}e_1 \oplus \dots \oplus \mathbb{Z}e_8$, and we will call these generators the *standard bases* for $N^1(X)$ and $N_1(X)$, respectively. In this standard basis, the anticanonical divisor class $-K_X$ is given by

$$-K_X = 4H - 2 \sum_{i=1}^8 E_i.$$

As shown in [LO16, Section 2], $-K_X$ is nef and not semiample with $(-K_X)^3 = 0$. Moreover, a general member Q in the linear system $|-\frac{1}{2}K_X|$ is smooth and isomorphic to \mathbb{P}^2 blown up at nine very general points. Two general members in $|-\frac{1}{2}K_X|$ intersect transversally along a smooth elliptic curve which is the unique member in $| -K_Q |$. Furthermore, a general member $\Delta \in | -K_X |$ is reducible and (X, Δ) forms a log canonical Calabi-Yau pair.

4.1. The Weyl group. In this subsection, we introduce a Weyl group acting on $N^1(X)_{\mathbb{R}}$. For more detail we refer to [Dol83] and [Muk04]. Let us consider a symmetric bilinear form (\cdot, \cdot) on $N^1(X)$:

$$(4) \quad (H, H) = 2, \quad (E_i, E_j) = -\delta_{i,j}, \quad (H, E_j) = 0$$

for $i, j \in \{1, \dots, 8\}$.

Lemma 4.1 ([Muk04]). *The classes $\alpha_0, \dots, \alpha_7, E_8$, where*

$$\begin{aligned} \alpha_0 &= H - E_1 - \dots - E_4, \\ \alpha_1 &= E_1 - E_2, \dots, \alpha_7 = E_7 - E_8, \end{aligned}$$

form another basis of $N^1(X)$. Moreover, $\alpha_0, \dots, \alpha_7$ is a \mathbb{Z} -basis of the orthogonal complement K_X^\perp of K_X under (\cdot, \cdot) and a system of simple roots of an affine root system with Dynkin diagram $T_{2,4,4}$.

Consider the real vector space $N^1(X)_{\mathbb{R}}$. Then $\{\alpha_0, \dots, \alpha_7\}$ is a set of linearly independent vectors in $N^1(X)_{\mathbb{R}}$. Let \mathcal{W} be the Weyl group generated by the orthogonal reflections with respect to $\alpha_0, \dots, \alpha_7$, that means,

$$(5) \quad \mathcal{W} = \langle s_0, \dots, s_7 \rangle,$$

where $s_i(D) = D + (\alpha_i, D) \cdot \alpha_i$ for any $D \in N^1(X)_{\mathbb{R}}$.

The *fundamental Weyl chamber* is defined as

$$C = \{D \in N^1(X)_{\mathbb{R}} \mid (D, \alpha_i) > 0 \text{ for all } i \in \{0, \dots, 7\}\},$$

where the α_i are the simple roots of Lemma 4.1 and (\cdot, \cdot) the symmetric bilinear form from (4). The closure \overline{C} of C is again a convex cone and characterized in the following way. Given any divisor class $D = dH - \sum_{i=1}^8 m_i E_i$ in $N^1(X)_{\mathbb{R}}$, we have:

$$(6) \quad \begin{aligned} D \in \overline{C} & \quad \text{if and only if } (D, \alpha_i) \geq 0 \text{ for all } i \in \{0, \dots, 7\} \\ & \quad \text{if and only if } 2d \geq \sum_{i=1}^4 m_i \text{ and } m_i \geq m_{i+1} \text{ for any } i \in \{0, \dots, 7\}. \end{aligned}$$

Note that $(K_X, \alpha_i) = 0$ for all $i \in \{0, \dots, 7\}$, hence any \mathbb{R} -multiple of K_X is in \overline{C} . The set \overline{C} is thus determined modulo K_X by the *fundamental weights* f_0, \dots, f_7 satisfying

$$(f_i, \alpha_j) = \delta_{i,j}.$$

More precisely, setting

- (i) $f_0 = \frac{1}{2}H$,
- (ii) $f_1 = \frac{1}{2}H - E_1$,
- (iii) $f_2 = H - E_1 - E_2$,
- (iv) $f_3 = \frac{3}{2}H - E_1 - E_2 - E_3$,
- (v) $f_4 = 2H - \sum_{i=1}^4 E_i$,
- (vi) $f_5 = 2H - \sum_{i=1}^5 E_i$,
- (vii) $f_6 = 2H - \sum_{i=1}^6 E_i$,
- (viii) $f_7 = 2H - \sum_{i=1}^7 E_i$,

then any element in \overline{C} is of the form

$$\sum_i \lambda_i f_i + \mu K_X$$

with $\lambda_i \in \mathbb{R}_{\geq 0}$ and $\mu \in \mathbb{R}$.

The union

$$T = \bigcup_{w \in \mathcal{W}} w \cdot \overline{C}$$

is called the *Tits cone*. It is a convex cone in $N^1(X)_{\mathbb{R}}$.

Now we analyse the action of the generators of \mathcal{W} on the standard basis $\{H, E_1, \dots, E_8\}$ of $N^1(X)_{\mathbb{R}}$. It is easy to see that for each $i \in \{1, \dots, 8\}$, the simple reflection s_i permutes the elements E_i and E_{i+1} and leaves E_j for $j \neq i, i+1$ and H invariant. Moreover, for the reflection s_0 we obtain

- (i) $s_0(H) = 3H - 2 \sum_{i=1}^4 E_i$,
- (ii) $s_0(E_j) = H - \sum_{i=1, i \neq j}^4 E_i$, for $j = 1, \dots, 4$,
- (iii) $s_0(E_j) = E_j$, else.

In the next subsection we see that the simple reflection s_0 is exactly the automorphism $\phi_{\{1, \dots, 4\}}: N^1(X) \rightarrow N^1(X)$ induced by the standard Cremona transformation centered at the first four points p_1, \dots, p_4 . More generally,

given an element $w \in \mathcal{W}$, there exists a corresponding isomorphism in codimension one for X :

Theorem 4.2 ([Muk04]). *For each element $w: N^1(X) \rightarrow N^1(X)$ of the Weyl group \mathcal{W} there exists an isomorphism in codimension one $\varphi_w: X \dashrightarrow X_w$, where X_w is also \mathbb{P}^3 blown up at 8 points in general position and the pullback of the standard basis of X_w coincides with the transformation of that of X by w .*

Remark 4.3. Let $\mathcal{V} = ((\mathbb{P}^3)^8)/\mathrm{PGL}(4)$, and consider $\mathbf{p} = (p_1, \dots, p_8) \in \mathcal{V}$. Then for an element $w \in \mathcal{W}$, the variety X_w from Theorem 4.2 is the blowup of \mathbb{P}^3 at $\mathbf{q} = \mathrm{cr}_{3,8}(w)(\mathbf{p})$, where

$$\mathrm{cr}_{3,8}: \mathcal{W} \rightarrow \mathrm{Bir}(\mathcal{V})$$


is the so called *Coble presentation* of \mathcal{W} (see [Muk04]). Then, using [Les14, Lemma 2.2], there exists a set $U \subset \mathcal{V}$, the complement of a countable union of subvarieties of \mathcal{V} such that if $\mathbf{p} \in U$, then for X being the blowup of \mathbb{P}^3 at \mathbf{p} , the action of \mathcal{W} on $N^1(X)_{\mathbb{R}}$ preserves the cones $\mathrm{Eff}(X)$ and $\overline{\mathrm{Mov}}(X)$, respectively.

4.2. The standard Cremona transformations. In this subsection, we summarise the action of a standard Cremona transformation of \mathbb{P}^3 centered at four points. We refer the reader to [DM22, Section 3] and [LO16, Section 3] for more detail.

The standard Cremona transformation of \mathbb{P}^3 (centered at the four coordinate points) is defined as

$$\mathrm{Cr}: \mathbb{P}^3 \dashrightarrow \mathbb{P}^3, \quad [x_0 : x_1 : x_2 : x_3] \mapsto \left[\frac{1}{x_0} : \frac{1}{x_1} : \frac{1}{x_2} : \frac{1}{x_3} \right].$$

Geometrically, it is realised by first blowing up the four coordinate points $\pi_1: Y_1 \rightarrow \mathbb{P}^3$, and then the strict transforms of the six lines through pairs of coordinate points $p: Y \rightarrow Y_1$, giving rise to six exceptional divisors isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$; then one blows down the “other ruling” of each of the six exceptional divisors $q: Y \rightarrow Y_2$, and then contracts the strict transforms of the four planes through three of the four coordinate points $\pi_2: Y_2 \rightarrow \mathbb{P}^3$.



Note that the induced map $\overline{\mathrm{Cr}}: Y_1 \dashrightarrow Y_2$ enjoys the following properties:

- (i) $\overline{\mathrm{Cr}}$ is an isomorphism in codimension 1;
- (ii) $\overline{\mathrm{Cr}}^*(K_{Y_2}) = K_{Y_1}$;
- (iii) $\overline{\mathrm{Cr}}$ induces an isomorphism $N^1(Y_1) \rightarrow N^1(Y_2)$.

After the first stage of performing the blowups, the lifted morphism $\widetilde{\text{Cr}} = \pi_2 \circ q: Y \rightarrow \mathbb{P}^3$ resolves the indeterminacies of the Cremona transformation Cr . The subspace in $N^1(Y)_{\mathbb{R}}$ generated by the six exceptional divisors of $p: Y \rightarrow Y_1$ is invariant under the lifted Cremona transformation $\widetilde{\text{Cr}}$, and the quotient space inherits the action induced by $\widetilde{\text{Cr}}$. This quotient space is naturally isomorphic to $N^1(Y_1)_{\mathbb{R}}$, and this action extends trivially to an action on $N^1(\text{Bl}_{\mathbf{p}} \mathbb{P}^3)_{\mathbb{R}}$, where \mathbf{p} is a very general configuration of eight points in \mathbb{P}^3 containing the four coordinate points.

Now consider the threefold X . Since the eight points p_1, \dots, p_8 are very general, any set of four of them can be the centered points of a corresponding Cremona transformation. For any subset $I \subset \{1, \dots, 8\}$ with $|I| = 4$, we denote by ϕ_I the corresponding Cremona transformation, which induces an action on $N^1(X)$. The action on $N^1(X)$ is explicitly described as follows.

Proposition 4.4. ([DM22, Proposition 3.1], [BDP16, Section 3.3])

Fix any subset $I \subset \{1, \dots, 8\}$ with $|I| = 4$.

(i) The action of ϕ_I on $N^1(X)$ is given by sending $dH - \sum_{i=1}^8 m_i E_i$ to $d'H - \sum_{i=1}^8 m'_i E_i$, where

$$d' = 3d - \sum_{j \in I} m_j = d + t,$$

$$m'_i = 2d - \sum_{j \in I, j \neq i} m_j = m_i + t \text{ for } i \in I \text{ and } m'_i = m_i \text{ for } i \notin I,$$

and $t := 2d - \sum_{i \in I} m_i$. In particular, we have $\phi_I^2 = \text{Id}$ and

$$\phi_I(H) = 3H - 2 \sum_{i \in I} E_i,$$

$$\phi_I(E_i) = H - \sum_{j \in I, j \neq i} E_j \text{ for } i \in I \text{ and } \phi_I(E_i) = E_i \text{ for } i \notin I.$$

(ii) Let $D \subset X$ be an effective Cartier divisor. Then

$$h^0(X, \mathcal{O}_X(D)) = h^0(X, \mathcal{O}_X(\phi_I(D))).$$

Consider a divisor $D \subset X$ which is the strict transform of a hypersurface in \mathbb{P}^3 . The divisor class of D can be written as $dH - \sum_{i=1}^8 m_i E_i$ with $d \in \mathbb{N}^*$ and $m_i \in \mathbb{N}$. We call d the *degree* of D . With notation from Proposition 4.4, if $t = 2d - \sum_{i \in I} m_i < 0$ for some subset $I \subset \{1, \dots, 8\}$ with $|I| = 4$, then the degree of D can be lowered by ϕ_I . In view of this property, we give the following definition.

Definition 4.5. ([DM22]) A divisor class $dH - \sum_{i=1}^8 m_i E_i$ in $N^1(X)$ with $d \in \mathbb{N}^*$ and $m_i \in \mathbb{N}$ is *Cremona-reduced* if $2d \geq \sum_{i \in I} m_i$ for every subset of indices $I \subset \{1, \dots, 8\}$ with $|I| = 4$.

In particular, the degree of a Cremona-reduced divisor class cannot be lowered by any of the ϕ_I 's.

Remark 4.6. Note that not any divisor class can be transformed into a Cremona-reduced one. For example, the effective class

$$D = 3H - 2 \sum_{i=1}^4 E_i - \sum_{i=5}^7 E_i$$

is not Cremona-reduced. Applying one Cremona transformation for $I = \{1, \dots, 4\}$ we obtain

$$\phi_I(D) = H - E_5 - E_6 - E_7$$

which is again not Cremona-reduced. Applying another Cremona transformation with $I' = \{5, 6, 7, 8\}$ we obtain $\phi_{I'}(\phi_I(D)) = E_8$.

This remark motivates the following definition.

4.3. Divisor classes in standard form. In this subsection we introduce the notion of a divisor class in standard form due to DeVolder and Laface (see [DL07]). The existence of this standard form (see Remark 4.8) together with the decomposition of a linear system in standard form (see Proposition 4.10) are one of the main tools to prove Theorems 1.3 and 1.4.

Definition 4.7. ([DL07]) Fix $d, m_i \in \mathbb{Z}$ for $i \in \{1, \dots, 8\}$. Let us denote by $(d; m_1, \dots, m_8)$ the divisor class $dH - \sum_{i=1}^8 m_i E_i$. We say that the class $(d; m_1, \dots, m_8)$ is in *standard form* if $m_1 \geq m_2 \geq \dots \geq m_8$ and $2d \geq \sum_{i=1}^4 m_i$. We denote by $\mathcal{L}(d; m_1, \dots, m_8)$ the linear system $|dH - \sum_{i=1}^8 m_i E_i|$ and we say that $\mathcal{L}(d; m_1, \dots, m_8)$ is in *standard form* if the class $(d; m_1, \dots, m_8)$ is in standard form.

Remark 4.8. Note that by applying Proposition 4.4 and permuting the order of the points, every divisor class can be transformed into standard form. Recall that the Weyl group \mathcal{W} from (5) is generated by s_0 (corresponding to the standard Cremona transformation) and the elements s_1, \dots, s_7 (corresponding to generators of the symmetric group S_8 permuting the points p_1, \dots, p_8). In particular, we have that for any effective Cartier divisor class D , there exists a class $(d; m_1, \dots, m_8)$ in standard form and an element $w \in W$ such that

$$w \cdot D = dH - \sum_{i=1}^8 m_i E_i,$$

that means $m_1 \geq m_2 \geq \dots \geq m_8$ and $2d \geq \sum_{i=1}^4 m_i$. In particular,

$$\dim |D| = \dim \mathcal{L}(d; m_1, \dots, m_8).$$

Notation 4.9. For a divisor class $(d; m_1, \dots, m_8)$ respectively a linear system $\mathcal{L}(d; m_1, \dots, m_8)$ in standard form we usually omit those m_i which are zero. For example, we write $\mathcal{L}(1; 1^2)$ for the system $|H - E_1 - E_2|$.

Proposition 4.10. ([DL07, Proposition 2.2]) *Let $(d; m_1, \dots, m_8)$ be a divisor class in standard form with $m_8 \geq 0$. Then*

$$(7) \quad (d; m_1, \dots, m_8) = D' + \sum_{i=4}^{a-1} (m_i - m_{i+1}) \cdot (2; 1^i) + m_a \cdot (2; 1^a),$$

where $D' = (d - 2m_4; m_1 - m_4, m_2 - m_4, m_3 - m_4)$ and $a = \max\{i \mid m_i \neq 0\}$.

Remark 4.11. Note that every divisor class in (7) is in standard form and all the coefficients are non-negative. The divisor class D' is effective if and only if $d - 2m_4 \geq m_1 - m_4$ (see [DL07, Lemma 5.1]).

Writing a linear system in standard form allows us to determine its base locus, see [DL06, Theorem 6.2]. For our purpose, we only need the following result, see [DL06, Theorem 6.2] and [BDP16, Corollary 4.4].

Proposition 4.12. *Let $\mathcal{L}(d; m_1, \dots, m_8)$ be a linear system in standard form. Assume that $m_8 \geq 0$. Then $\mathcal{L}(d; m_1, \dots, m_8)$ has no fixed divisor.*

5. THE CONE OF CURVES AND THE NEF CONE

Let X be \mathbb{P}^3 blown up at eight very general points p_1, \dots, p_8 . We use the notation introduced at the beginning of Section 4. In this section we determine a set of generators for the cone of curves $\overline{\text{NE}}(X)$ and respectively for its dual cone $\text{Nef}(X)$, and we prove Theorem 1.2.

The fact that the cone $\text{NE}(X)$ is a rational polyhedral cone is a particular case of [CLO16, Lemma 2.6] (applied for $r = 8$ and $n = 2$). We include the proof for readers' convenience.

Lemma 5.1. *Let $c = ah - \sum_{i=1}^8 b_i e_i$ be a class in $N_1(X)$ satisfying*

- (i) $a \in \mathbb{N}^*$ and $b_i \in \mathbb{N}$,
- (ii) $a \geq b_i$ for every i ,
- (iii) $2a \geq \sum_{i=1}^8 b_i$.

Then c is a non-negative linear combination of the classes e_i for $1 \leq i \leq 8$, and the classes $\ell_{ij} = h - e_i - e_j$ for $1 \leq i < j \leq 8$.

Proof. We proceed by induction on $a \in \mathbb{N}^*$. The case $a = 1$ is clear: by (ii), we have $a \geq b_i$ for each i and thus each b_i is either 0 or 1. By (iii), there are at most two non-zero b_i and the class c is of the form claimed.

Suppose that $a > 1$. Let

$$J := \{1 \leq i \leq 8 \mid b_i > 0\}$$

and let $j = \min(2, |J|)$. Let I be a set of j indices $\{i_1, \dots, i_j\}$ such that $b_{i_1} \geq \dots \geq b_{i_j} \geq b_i$ for any $i \in \{1, \dots, 8\} \setminus I$.

If $j = 2$, we have $I = \{i_1, i_2\}$. Set

$$c' = c - \ell_{i_1 i_2} =: a'h - \sum_{i=1}^8 b'_i e_i,$$

where $a' = a - 1$, $b'_i = b_i - 1$ if $i \in I$ and $b'_i = b_i$ if $i \notin I$. We check that the class c' satisfies the inequalities in question, as follows.

- By assumption, we have $a - 1 > 0$ and $b_{i_1} - 1 \geq b_{i_2} - 1 \geq 0$.
- Clearly, $a - 1 \geq b_{i_1} - 1 \geq b_{i_2} - 1$. We also have $a - 1 \geq b_i$ for all $i \notin I$: otherwise by (ii) we have $b_i = a$ for some $i \notin I$, which implies $b_{i_1} = b_{i_2} = b_i = a$. But by (iii) we have $2a \geq b_{i_1} + b_{i_2} + b_i$, which implies $a = 0$ and gives a contradiction.
- By (iii) we have $2(a - 1) \geq (b_{i_1} - 1) + (b_{i_2} - 1) + \sum_{i \notin I} b_i$.

Now if $j = 1$, then $c = ah - b_{i_1}e_{i_1}$ with $a \geq b_{i_1}$. Let $k \in \{1, \dots, 8\} \setminus I$. Then we can write

$$c = (a - b_{i_1})e_{i_1} + ae_k + a(h - e_{i_1} - e_k).$$

If $j = 0$, then $c = ah$ and we can write

$$c = ae_1 + ae_2 + a(h - e_1 - e_2).$$

This completes the proof by induction on a . \square

Now let C be an irreducible curve in X . If C is contained in some exceptional divisor E_i for $i \in \{1, \dots, 8\}$, then the class of C is proportional to e_i . If C is not contained in any exceptional divisor, then C is the strict transform of a curve in \mathbb{P}^3 with positive degree a and non-negative multiplicity $b_i \leq a$ at the point p_i for $i \in \{1, \dots, 8\}$. Since $H - E_i$ is nef for any $i \in \{1, \dots, 8\}$, and $2H - \sum_{i=1}^8 E_i$ is nef, the class of C satisfies the assumptions of Lemma 5.1 and thus is a non-negative linear combination of e_i for $1 \leq i \leq 8$, and ℓ_{ij} for $1 \leq i < j \leq 8$. Therefore, we have shown the following result:

Proposition 5.2. ([CLO16, Proposition 4.1]) *The cone $\text{NE}(X)$ is a rational polyhedral cone spanned by the classes e_i , for $1 \leq i \leq 8$, and the classes ℓ_{ij} , for $1 \leq i < j \leq 8$.*

In particular, Proposition 5.2 implies that $\text{NE}(X)$ is closed. We can now describe the nef cone $\text{Nef}(X)$ which is dual to $\overline{\text{NE}}(X) = \text{NE}(X)$.

Lemma 5.3. *The cone $\text{Nef}(X)$ is a rational polyhedral cone spanned by the classes*

- H ,
- $H - E_i$ for $i = 1, \dots, 8$,
- $2H - \sum_{j \in J \subset \{1, \dots, 8\}, |J|=m} E_j$ for $3 \leq m \leq 8$.

Proof. Consider a nef Cartier divisor class $D = dH - \sum_{i=1}^8 m_i E_i$. Then by Proposition 5.2, we have

$$m_i \geq 0 \text{ for any } i = 1, \dots, 8 \text{ and } d \geq m_i + m_j \text{ for any } 1 \leq i < j \leq 8.$$

By reordering, we may assume that $m_1 \geq m_2 \geq \dots \geq m_8$. This implies that D is in standard form. Similarly as in Proposition 4.10, we can write

$$D = (d - 2m_2; m_1 - m_2) + \sum_{i=2}^{a-1} (m_i - m_{i+1}) \cdot (2; 1^i) + m_a \cdot (2; 1^a),$$

where $a = \max\{i \mid m_i \neq 0\}$. Any class of the form $(2; 1^i)$ for $i \in \{2, \dots, a\}$ is an \mathbb{N} -linear combination of the classes in the lemma. Furthermore,

$$d - 2m_2 \geq m_1 - m_2 \geq 0,$$

as $d \geq m_1 + m_2$ and $m_1 \geq m_2$. Thus,

$$(d - 2m_2; m_1 - m_2) = (d - m_1 - m_2)H + (m_1 - m_2)(H - E_1),$$

which finishes the proof. \square

As an immediate consequence, we obtain the result below, which is a special case of a more general result from [BC13, Corollary 3.2]. Recall that $\text{Aut}(X)$ acts on $N^1(X)_{\mathbb{R}}$ by pulling back divisors and we denote by $\text{Aut}(X)^*$ the image of $\text{Aut}(X)$ in $\text{GL}(N^1(X)_{\mathbb{R}})$.

Corollary 5.4. *The group $\text{Aut}(X)^*$ is finite.*

Proof. Any element $g \in \text{Aut}(X)$ induces an action $g^*: N^1(X)_{\mathbb{R}} \rightarrow N^1(X)_{\mathbb{R}}$ which preserves the cone $\text{Nef}(X)$. By Lemma 5.3, $\text{Nef}(X)$ is a rational polyhedral cone spanned by finitely many extremal rays. Any $g^* \in \text{Aut}(X)^*$ permutes the finitely many primitive integral elements on the extremal rays and is determined by the action on these elements. Hence, we can consider $\text{Aut}(X)^*$ as a subgroup of a symmetric group. \square

Corollary 5.5. *The action of $\text{Aut}(X)$ on the effective nef cone $\text{Nef}^e(X)$ admits a fundamental domain and we have*

$$\text{Nef}(X) = \text{Nef}^e(X) = \text{Nef}^+(X).$$

Proof. By Lemma 5.3, we see that $\text{Nef}(X)$ is a polyhedral cone and that every extremal ray of $\text{Nef}(X)$ contains a rational effective class. Hence, we have $\text{Nef}(X) \subset \text{Nef}^+(X)$, $\text{Nef}(X) \subset \text{Eff}(X)$ and the cone

$$\text{Nef}^e(X) = \text{Nef}(X)$$

is rational polyhedral. As $\text{Aut}(X)^*$ is finite and

$$\text{Nef}^e(X) = \text{Aut}(X)^* \cdot \text{Nef}^e(X),$$

this implies that the action of $\text{Aut}(X)$ on $\text{Nef}^e(X)$ admits a fundamental domain.

For the equality $\text{Nef}(X) = \text{Nef}^+(X)$, Theorem 2.3 applied to $C = \text{Amp}(X)$, $G = \text{Aut}(X)^*$ and $\Pi = \text{Nef}(X) \subset \text{Nef}^+(X)$ implies that there exists a rational polyhedral fundamental domain for the action of G on $\text{Nef}^+(X)$ and

$$\text{Nef}(X) = G \cdot \Pi = \text{Nef}^+(X). \quad \square$$

Proof of Theorem 1.2. The first statement follows from Lemma 5.3 and the second statement follows from Corollary 5.5. \square

6. THE EFFECTIVE CONE

As before, X denotes the blowup of \mathbb{P}^3 at eight very general points. In this section, we explicitly describe a set of extremal rays for the convex cone $\text{Eff}(X)$ and we prove Theorem 1.4.

Definition 6.1. ([Muk01]) A divisor $D \subset X$ is a (-1) -divisor if there is an birational map $f: X \dashrightarrow X'$ and a morphism $\pi': X' \rightarrow Y$ such that f is an isomorphism in codimension one and π' is the blowup of a projective variety Y at a smooth point and D is the strict transform of the exceptional divisor of π' .

In [Muk01, Lemma 3], Mukai proved that the class of a (-1) -divisor is contained in any set of generators of the semi-group of effective divisor classes in $\text{Pic}(X)$. Thus (-1) -divisors play an important role in understanding the cone $\text{Eff}(X)$. Furthermore, Mukai proved in [Muk04, Theorem 1] that any divisor in the orbit of the Weyl group of an exceptional divisor of $\pi: X \rightarrow \mathbb{P}^3$ is a (-1) -divisor. To summarise, we have the following inclusions:

$$\begin{aligned} \{\mathcal{W} \cdot E_8\} &\subset \{\text{classes of } (-1)\text{-divisors}\} \\ &\subset \{\text{generators of the semi-group of the effective divisor classes}\} \end{aligned}$$

We first describe all the prime divisors on X in the following lemma.

Lemma 6.2. *Let $D \subset X$ be a prime divisor on X . Assume that D is non-nef. Then D is either a (-1) -divisor or a movable divisor.*

Proof. Since X is smooth, for sufficiently small $\epsilon > 0$, the pair $(X, \epsilon D)$ is log-canonical. Since D is not nef, by [Xie20, Lemma 2.5] there exists a $(K_X + \epsilon D)$ -negative extremal ray Γ such that $D \cdot \Gamma < 0$. Let c_Γ be the contraction of the extremal ray Γ and let ℓ be a contracted curve. Then $D \cdot \ell < 0$, which implies $\ell \subset \text{Bs}|D|$.

If c_Γ is small, then $K_X \cdot \ell = 0$, as there is no flipping contraction for smooth threefolds. Hence there exists a flop of c_Γ and the flopped threefold X^+ is again smooth by [Kol89, Theorem 2.4]. By repeating the above argument and by the termination of three-dimensional flops [KM98, Corollary 6.19], we deduce that there exists a sequence of flops $\varphi: X \dashrightarrow X'$ with $D' := \varphi_* D$, such that we are in one of the following cases.

- (i) If D is movable, then D' is nef.
- (ii) If D is not movable, i.e. $\text{Bs}|D| = D$, then there exists a divisorial contraction $c_{\Gamma'}$ corresponding to a $(K_{X'} + \epsilon D')$ -negative extremal ray Γ' such that $D' \cdot \Gamma' < 0$. Thus $c_{\Gamma'}$ contracts the divisor D' .

Now we show that in above case (ii), the divisor D' is contracted to a smooth point by $c_{\Gamma'}$, which implies that D is a (-1) -divisor.

Indeed, if Γ' is $K_{X'}$ -negative, then by the classification of Mori-Mukai (see [MM83, Section 3]), we obtain that $D' \simeq \mathbb{P}^2$ and $c_{\Gamma'}$ contracts D' to a smooth point as $-K_{X'}$ is divisible by two in $\text{Pic}(X')$. If Γ' is $K_{X'}$ -trivial,

then D' is covered by $K_{X'}$ -trivial curves. Since ψ is a sequence of flops, we obtain that D is covered by K_X -trivial curves. This implies that the set of K_X -trivial is uncountable, which contradicts [LO16, Lemma 10] and finishes the proof. \square

Now, the following lemma implies that the set of all (-1) -divisors on X is equal to the set $\mathcal{W} \cdot E_8$.

Lemma 6.3. *Let $D \subset X$ be a (-1) -divisor. Then D is in the orbit of the Weyl group of an exceptional divisor of $\pi: X \rightarrow \mathbb{P}^3$.*

Proof. Since $D \subset X$ is a prime divisor, either D is one of the exceptional divisors E_i of $\pi: X \rightarrow \mathbb{P}^3$, or we can write

$$(8) \quad D \sim dH - \sum_{i=1}^8 m_i E_i$$

with $d \in \mathbb{N}^*$ and $m_1, \dots, m_8 \in \mathbb{N}$.

Assume that D is not one of the exceptional divisors E_i . Since the divisor D is not movable, by Proposition 4.12 we obtain that D is not Cremona-reduced, i.e. the degree d of D can be lowered by a Cremona transformation ϕ_I .

By Proposition 4.4(ii), we have $h^0(X, \mathcal{O}_X(\phi_I(D))) = h^0(X, \mathcal{O}_X(D)) = 1$. Thus the divisor $\phi_I(D)$ is also a (-1) -divisor by Lemma 6.2. Therefore, we may assume $d = 1$ in (8) and thus $\pi(D)$ is a plane. Together with $h^0(X, \mathcal{O}_X(D)) = 1$, we obtain

$$D = H - (E_{i_1} + E_{i_2} + E_{i_3})$$

for some pairwise distinct indices $i_1, i_2, i_3 \in \{1, \dots, 8\}$.

Now, choose $i_4 \in \{1, \dots, 8\} \setminus \{i_1, i_2, i_3\}$ and fix $I = \{i_1, i_2, i_3, i_4\}$, we have $\phi_I(D) = E_{i_4}$ by Proposition 4.4(i). \square

Revisiting the chain of inclusion stated above, we have now

$$\begin{aligned} \{\mathcal{W} \cdot E_8\} &= \{\text{classes of } (-1)\text{-divisors}\} \\ &\subset \{\text{generators of the semi-group of the effective divisor classes}\}. \end{aligned}$$

We recall from the introduction that if $X_{n,k}$ denotes the blowup of \mathbb{P}^n in $k > n + 1$ points in general position, then by [CT06, Theorem 1.3] the effective cone is rational polyhedral if and only if

$$(9) \quad \frac{1}{n+1} + \frac{1}{k-n-1} > \frac{1}{2}.$$

If (9) is satisfied, then $\text{Eff}(X_{n,k})$ is spanned by the finitely many elements in the orbit $\mathcal{W}_{n,k} \cdot E_k$, where $\mathcal{W}_{n,k}$ denotes the associated Weyl group. If $n = 3$ and $k = 8$, the Weyl group $\mathcal{W} = \mathcal{W}_{3,8}$ is infinite, $\text{Eff}(X)$ is not rational polyhedral, and we show that in this case the infinitely many divisors in the \mathcal{W} -orbit of E_8 span only a proper subcone of $\text{Eff}(X)$:

Lemma 6.4. *The class $-\frac{1}{2}K_X \in \text{Eff}(X)$ is not contained in the convex cone generated by the classes of $\mathcal{W} \cdot E_8$.*

Proof. To decide whether $-\frac{1}{2}K_X = 2H - \sum_{i=1}^8 E_i$ is a non-negative combination of elements in the orbit of E_8 it is enough to consider all elements in $\mathcal{W} \cdot E_8$ up to degree 2. We state these elements *up to permutation*: for a given element written in the table below, all elements obtained by permutations of the last eight entries are also in the orbit.

H	E_1	E_2	E_3	E_4	E_5	E_6	E_7	E_8
0	1	0	0	0	0	0	0	0
1	-1	-1	-1	0	0	0	0	0
2	-2	-1	-1	-1	-1	-1	0	0

TABLE 1. (-1) -divisors up to degree 2

The entries in the table denote the coefficient of the corresponding divisor. For example, the second row represents the class $H - E_1 - E_2 - E_3$.

We deduce that $2H - \sum_{i=1}^8 E_i$ is not a non-negative combination of the elements in table which shows the claim. \square

We can now describe the extremal rays generating the cone $\text{Eff}(X)$ as a convex cone.

Theorem 6.5. *The effective cone $\text{Eff}(X)$ is the convex cone generated by the classes in $\mathcal{W} \cdot E_8$ and the class of $-\frac{1}{2}K_X$.*

Proof. Let us denote by \mathcal{C} the convex cone generated by the classes in $\mathcal{W} \cdot E_8$ and the class $-\frac{1}{2}K_X$. We remark that $w \cdot (-\frac{1}{2}K_X) = -\frac{1}{2}K_X$ for any $w \in \mathcal{W}$. Hence the cone \mathcal{C} is preserved by \mathcal{W} . In particular, we have that $D \in \mathcal{C}$ if and only if there exists $w \in \mathcal{W}$ such that $w \cdot D \in \mathcal{C}$. Recall that $\text{Eff}(X)$ is the convex cone spanned by all the effective Cartier divisor classes. Therefore, to conclude $\text{Eff}(X) = \mathcal{C}$, it is sufficient to show that for any effective Cartier divisor class D , there exists $w \in \mathcal{W}$ such that $w \cdot D \in \mathcal{C}$.

Let D be an effective Cartier divisor class. By Remark 4.8, there exists an element $w \in \mathcal{W}$ such that

$$w \cdot D = dH - \sum_{i=1}^8 m_i E_i$$

is an effective divisor class, where

$$(10) \quad 2d \geq \sum_{i=1}^4 m_i$$

and

$$(11) \quad m_1 \geq \dots \geq m_8,$$

i.e. $w \cdot D$ is in standard form. Assume that $w \cdot D$ is not one of the exceptional divisor class E_i for $i \in \{1, \dots, 8\}$, otherwise $w \cdot D \in \mathcal{W} \cdot E_8$ and we are finished. We may further assume that there exists a prime divisor in the linear system $\mathcal{L}(d; m_1, \dots, m_8)$, since otherwise $w \cdot D$ can be written as a sum of effective divisors. Thus

$$(12) \quad d \geq m_i \geq 0, \quad \forall i \in \{1, \dots, 8\}.$$

In the remainder of the proof, we will show that $w \cdot D \in \mathcal{C}$.

Step 1. We first consider the case when $d \geq m_1 + m_4$.

Applying Proposition 4.10 we have

$$(d; m_1, \dots, m_8) = D' + \sum_{i=4}^8 c_i \cdot (2; 1^i),$$

where $D' = (d - 2m_4; m_1 - m_4, m_2 - m_4, m_3 - m_4)$ and $c_i \geq 0$.

First we remark that any $2H - \sum_{i=1}^k E_i$ for $4 \leq k \leq 8$ is contained in \mathcal{C} . Indeed using the list of elements in $\mathcal{W} \cdot E_8$ up to degree 2 in Table 1, we see that for any $4 \leq k \leq 6$, the class of $2H - \sum_{i=1}^k E_i$ is a non-negative combination of classes in $\mathcal{W} \cdot E_8$, whereas $2H - \sum_{i=1}^8 E_i$ is the class of $-\frac{1}{2}K_X$ and $2H - \sum_{i=1}^7 E_i = (2H - \sum_{i=1}^8 E_i) + E_8 \in \mathcal{C}$.

Consequently, it remains to show that the class D' is contained in \mathcal{C} . We write $D' = d'H - \sum_{i=1}^3 m'_i E_i$. Then $2d' \geq m'_1 + m'_2 + m'_3$ by (10) and $m'_1 \geq m'_2 \geq m'_3 \geq 0$ by (11). Since $d \geq m_1 + m_4$, we also have $d' \geq m'_i$ for all $i \in \{1, 2, 3\}$. Then we have

$$d'H - \sum_{i=1}^3 m'_i E_i = d'(H - \sum_{i=1}^3 E_i) + \sum_{i=1}^3 (d' - m'_i) E_i,$$

where the right hand side is contained in \mathcal{C} . Thus, we have shown that

$$w \cdot D \in \mathcal{C}.$$

Step 2. Now we consider the case when $d < m_1 + m_4$. The proof of this step is inspired by the proof of [DL06, Theorem 6.2]. Let $t_{i,j} := m_i + m_j - d$ for $1 \leq i < j \leq 8$. Then $d < m_1 + m_4$ is equivalent to

$$(13) \quad t_{1,4} > 0.$$

Therefore,

$$(14) \quad m_1 \geq m_2 + 2 \geq 3.$$

Indeed, we have

$$\begin{aligned} m_1 &\geq m_1 + \left(\sum_{i=1}^4 m_i - 2d \right) && \text{by (10)} \\ &= m_2 + (m_3 - m_4) + 2t_{1,4} \\ &\geq m_2 + 2 && \text{by (11)(13)}. \end{aligned}$$

Let $a := \max\{i \in \{1, \dots, 8\} \mid m_i \neq 0\}$. Since $m_1 \leq d < m_1 + m_4$, we have $m_4 > 0$ and thus $a \geq 4$.

Note that using the list of elements in $\mathcal{W} \cdot E_8$ up to degree 2 in Table 1, we have

$$(3; 3, 1^7) = (2H - 2E_1 - \sum_{i=2}^6 E_i) + (H - E_1 - E_7 - E_8) \in \mathcal{C},$$

and $(3; 3, 1^{a-1}) = (3; 3, 1^7) + \sum_{i=a+1}^8 E_i \in \mathcal{C}$ for $4 \leq a < 8$.

Now consider

$$(d'; m'_1, \dots, m'_{a'}) := (d; m_1, \dots, m_a) - (3; 3, 1^{a-1}).$$

Note that by (12) and (14), the case $d < m_1 + m_4$ can only occur if $d \geq m_1 \geq 3$. Then

$$\begin{aligned} d' &= d - 3, \\ m'_1 &= m_1 - 3, \\ m'_i &= m_i - 1, \quad \forall i \in \{2, \dots, a\}, \\ a' &:= \max\{i \in \{1, \dots, 8\} \mid m'_i \neq 0\} \leq a. \end{aligned}$$

We obtain that $(d'; m'_1, \dots, m'_{a'})$ is in standard form with $d' \geq m'_i \geq 0$ for any $i \in \{1, \dots, a'\}$. Indeed, by construction we have $m'_i = 0$ for $a' < i \leq 8$, and $d' \geq m'_1 \geq m'_2 \geq \dots \geq m'_{a'} > 0$ by (11) and (14). Moreover, by (10) we have

$$2d' = 2d - 6 \geq \sum_{i=1}^4 m_i - 6 = \sum_{i=1}^4 m'_i.$$

Furthermore, set $t'_{i,j} := m'_i + m'_j - d'$ for $1 \leq i < j \leq 8$. Then we have $t'_{1,4} = t_{1,4} - 1$. If $t'_{1,4} > 0$, since $(d'; m'_1, \dots, m'_{a'})$ is in standard form, we can start our above procedure in *Step 2* again and do this until $t'_{1,4} = 0$ for some $(d'; m'_1, \dots, m'_{a'})$. We are now under the assumption in *Step 1* and thus $(d'; m'_1, \dots, m'_{a'}) \in \mathcal{C}$. \square

In the remainder of the section, we will show that the cone $\text{Eff}(X)$ is closed. One crucial ingredient for the proof of this result is that the cone of curves of \mathbb{P}^2 blown up at nine very general points is closed. We thank Artie Prendergast-Smith for communicating the following result to us:

Proposition 6.6. ([Bor91, Proposition 6.2]) *Let S be a complex projective smooth surface such that $-K_S$ is nef and non-torsion. Then $\text{NE}(S)$ is a closed convex cone generated by the classes of rational curves and possibly the class $-K_S$, accumulating at most to $\mathbb{R}_{>0}[-K_S]$.*

In particular, $\text{NE}(S) \cap K_S^\perp$ is a rational polyhedral subcone of $\text{NE}(S)$ generated by the classes of (-2) -curves, and possibly the class $-K_S$ if $(-K_S)^2 = 0$.

Remark 6.7. In Proposition 6.6, if S is isomorphic to the blowup $X_{2,9}$ of \mathbb{P}^2 at nine very general points, then the unique element in $|-K_S|$ is a smooth elliptic curve and there is no (-2) -curve on the surface S . Moreover, by

[Dol83, Corollary 1] there is a bijection between the set of (-1) -curves on S and the orbit $\mathcal{W}_{2,9} \cdot e$, where e is an exceptional curve of the blowup and $\mathcal{W}_{2,9}$ is the Weyl group of the Dynkin diagram $T_{2,6,3}$ acting on $N^1(X_{2,9})$. Hence $\text{Eff}(S)$ is a closed convex cone generated by the classes in $\mathcal{W}_{2,9} \cdot e$ and the class $-K_S$.

We need the following lemma to show that any (-1) -divisor on X restricts to a (-1) -curve on a general member of $|-\frac{1}{2}K_X|$.

Lemma 6.8. *Let D be a (-1) -divisor on X and let Q be a general member in the linear system $|-\frac{1}{2}K_X|$. Then we have $D^2 \cdot Q = -1$ and $D \cdot Q^2 = 1$.*

Proof. Recall that $Q \sim 2H - \sum_{i=1}^8 E_i$ and we may write

$$D \sim dH - \sum_{i=1}^8 m_i E_i$$

with $d, m_1, \dots, m_8 \in \mathbb{Z}$.

Since $H^3 = 1, H^2 \cdot E_i = H \cdot E_i^2 = 0, E_i^3 = 1$ and $E_i \cdot E_j = 0$ for $1 \leq i < j \leq 8$, we obtain

$$(15) \quad D^2 \cdot Q = (dH - \sum_{i=1}^8 m_i E_i)^2 \cdot (2H - \sum_{i=1}^8 E_i) = 2d^2 - \sum_{i=1}^8 m_i^2,$$

$$(16) \quad D \cdot Q^2 = (dH - \sum_{i=1}^8 m_i E_i) \cdot (2H - \sum_{i=1}^8 E_i)^2 = 4d - \sum_{i=1}^8 m_i.$$

Consider the symmetric bilinear form (\cdot, \cdot) on $N^1(X)$ defined in (4) and we have

$$(17) \quad (D, D) = (dH - \sum_{i=1}^8 m_i E_i, dH - \sum_{i=1}^8 m_i E_i) = 2d^2 - \sum_{i=1}^8 m_i^2 = D^2 \cdot Q$$

by (15), and

$$(18) \quad (D, Q) = (dH - \sum_{i=1}^8 m_i E_i, 2H - \sum_{i=1}^8 E_i) = 4d - \sum_{i=1}^8 m_i = D \cdot Q^2$$

by (16). Now it follows from [DP19, Theorem 2.10(1)] that the form (\cdot, \cdot) is preserved under the action of the Weyl group \mathcal{W} . Hence, by Lemma 6.3 we obtain

$$(D, D) = (E_8, E_8) = -1 \text{ and } (D, Q) = (E_8, Q) = 1.$$

Together with (17) and (18), we conclude $D^2 \cdot Q = -1$ and $D \cdot Q^2 = 1$. \square

Lemma 6.9. *Let $Q \in |-\frac{1}{2}K_X|$ be a general member. Then we have an injection:*

$$\begin{aligned} \{(-1)\text{-divisors on } X\} &\hookrightarrow \{(-1)\text{-curves on } Q\} \\ D &\mapsto D|_Q \end{aligned}$$

Proof. By the adjunction formula, we have

$$-K_Q \sim (-K_X - Q)|_Q \sim Q|_Q.$$

Let D be a (-1) -divisor on X and denote by Γ the curve $D \cap Q$. On the surface Q , we have

$$(19) \quad -K_Q \cdot \Gamma = Q|_Q \cdot D|_Q = 1 \text{ and } \Gamma^2 = (D|_Q)^2 = -1,$$

since $D \cdot Q^2 = 1$ and $D^2 \cdot Q = -1$ by Lemma 6.8.

Since D is a (-1) -divisor, by the proof of Lemma 6.2 there exists a finite sequence of flops $X \dashrightarrow X'$ such that the strict transform D' of D on X' is isomorphic to \mathbb{P}^2 . Denote by Q' the strict transform of Q on X' . As $\mathcal{O}_{D'}(D') = \mathcal{O}_{\mathbb{P}^2}(-1)$, we have that $|Q'|_{D'} \subset |Q'|_{D'}$ is a pencil of smooth rational curves on $D' \simeq \mathbb{P}^2$. Hence $|Q|_D$ is a pencil of integral rational curves; in particular, Γ is an integral rational curve. Together with (19), we obtain that Γ is a (-1) -curve on the surface Q ; in particular, Γ is a smooth rational curve. Thus $D|_Q$ is a (-1) -curve on the surface Q and we obtain a well-defined map:

$$\begin{aligned} \{(-1)\text{-divisors on } X\} &\rightarrow \{(-1)\text{-curves on } Q\} \\ D &\mapsto D|_Q \end{aligned}$$

By the short exact sequence

$$0 \rightarrow \mathcal{O}_X(D - Q) \rightarrow \mathcal{O}_X(D) \rightarrow \mathcal{O}_Q(D) \rightarrow 0$$

and since $H^0(X, \mathcal{O}_X(D - Q)) = 0$, the map defined above is injective. \square

In the following, we will work with the real projective space $\mathbb{R}\mathbb{P}^8 = (N^1(X)_{\mathbb{R}} \setminus \{0\})/\mathbb{R}^*$. For any class $x \in \overline{\text{Eff}}(X)$ seen as a non-zero vector in $N^1(X)_{\mathbb{R}} \simeq \mathbb{R}^9$,

the ray spanned by the class x intersects the unit sphere $\mathbb{S}^8 \subset \mathbb{R}^9$ in a unique point. Thus, we can identify each ray with a point in the projective space $\mathbb{R}\mathbb{P}^8$. We denote by $x = (d; m_1, \dots, m_8)$ an element in $\overline{\text{Eff}}(X) \setminus \{0\}$ and by $\mathbf{x} = [d : m_1 : \dots : m_8]$ its corresponding point in $\mathbb{R}\mathbb{P}^8$.

Proposition 6.10. *Let $A = \{w \cdot E_8 \mid w \in \mathcal{W}\}$. Then the only accumulation point of the set $\{\mathbf{x}\}_{x \in A} \subset \mathbb{R}\mathbb{P}^8$ is the point corresponding to the class of $-\frac{1}{2}K_X$.*

Proof. Fix a very general member $Q \in |-\frac{1}{2}K_X|$. Now since Q is the blowup of a smooth quadric surface in \mathbb{P}^3 at eight very general points, we can write

$$N^1(Q) = \mathbb{Z}\ell_1 \oplus \mathbb{Z}\ell_2 \oplus \mathbb{Z}e_1 \oplus \dots \oplus \mathbb{Z}e_8 \cong \mathbb{Z}^{10},$$

where ℓ_1 and ℓ_2 are respectively the pullbacks of the two rulings of the quadric surface in \mathbb{P}^3 . Now, analogously as for X , we identify the rays in $N^1(Q)_{\mathbb{R}}$ spanned by (-1) -curves and $-K_Q$ with their images in $\mathbb{R}\mathbb{P}^9$.

We have

$$(20) \quad H|_Q \sim \ell_1 + \ell_2 \text{ and } E_i|_Q \sim e_i \text{ for } i \in \{1, \dots, 8\}.$$

By compactness of \mathbb{RP}^8 , there exists an accumulation point of the set $\{\mathbf{x}\}_{x \in A} \subset \mathbb{RP}^8$. In the following, we show that there is exactly one accumulation point.

Let $\{\mathbf{x}_k\}_{k \in \mathbb{N}}$ be a converging subsequence of $\{\mathbf{x}\}_{x \in A}$ with limit point $\mathbf{y} \in \mathbb{RP}^8$, let D_k be a (-1) -divisor on X corresponding to the class x_k for each $k \in \mathbb{N}$ and let B be an \mathbb{R} -divisor on X corresponding to the class y . We can write

$$(21) \quad B \sim aH - \sum_{i=1}^8 b_i E_i,$$

where $a, b_i \in \mathbb{R}$ for any $i \in \{1, \dots, 8\}$.

By (20), the map in Lemma 6.9 can be written as:

$$\begin{aligned} \{(-1)\text{-divisors on } X\} &\leftrightarrow \{(-1)\text{-curves on } Q\} \\ D \sim dH - \sum_{i=1}^8 m_i E_i &\mapsto D|_Q \sim d(\ell_1 + \ell_2) - \sum_{i=1}^8 m_i e_i, \end{aligned}$$

which gives the corresponding injection:

$$\begin{aligned} \mathbb{RP}^8 &\hookrightarrow \mathbb{RP}^9 \\ [d : m_1 : \dots : m_8] &\mapsto [d : d : m_1 : \dots : m_8]. \end{aligned}$$

Since the sequence $\{\mathbf{x}_k\}_{k \in \mathbb{N}}$ converges to the point \mathbf{y} , we obtain that the sequence $\{D_k|_Q\}_{k \in \mathbb{N}}$ converges to $B|_Q$ in \mathbb{RP}^9 . Since Q is isomorphic to \mathbb{P}^2 blown up at nine very general points, by Proposition 6.6 and Remark 6.7 we obtain $[B|_Q] \in \mathbb{R}_{>0}[-K_Q]$, and thus there exists $\lambda \in \mathbb{R}_{>0}$ such that

$$B|_Q \sim \lambda(2\ell_1 + 2\ell_2 - \sum_{i=1}^8 e_i).$$

Hence, by (21) we have $a = 2\lambda$ and $b_i = \lambda$ for any $i \in \{1, \dots, 8\}$ and thus the limit point \mathbf{y} corresponds to the ray spanned by the class $-\frac{1}{2}K_X$. \square

We would like to thank Artie Prendergast-Smith for communicating the proof of the following statement to us:

Proposition 6.11. *The effective cone $\text{Eff}(X)$ is closed.*

Proof. Recall from Theorem 6.5 that $\text{Eff}(X)$ is generated by the classes in $\mathcal{W} \cdot E_8$ and the class of $-\frac{1}{2}K_X$. Consider the symmetric bilinear pairing (\cdot, \cdot) introduced in Section 4. Then for $D = dH - \sum_{i=1}^8 m_i E_i$ we have

$$(D, -\frac{1}{2}K_X) = 4d - \sum_{i=1}^8 m_i \text{ and } (D, D) = 2d^2 - \sum_{i=1}^8 m_i^2.$$

Furthermore, by [DP19, Theorem 2.10(1)] the pairing (\cdot, \cdot) is invariant under the Weyl group operation, hence

$$(22) \quad (D, -\frac{1}{2}K_X) = 1 \text{ for any } D \in \mathcal{W} \cdot E_8 \text{ and } (-\frac{1}{2}K_X, -\frac{1}{2}K_X) = 0.$$

Let us denote by $K_X^\perp = \{x \in N^1(X)_\mathbb{R} \mid (x, -\frac{1}{2}K_X) = 0\}$ and by $K_X^{\leq 0} = \{x \in N^1(X)_\mathbb{R} \mid (x, -\frac{1}{2}K_X) \leq 0\}$. As before, we can identify any ray spanned by an element in $\overline{\text{Eff}}(X)$ with its image in the projective space $\mathbb{R}\mathbb{P}^8$. In the following, we denote by $x = (d; m_1, \dots, m_8)$ an element in $\overline{\text{Eff}}(X)$ and by $\mathbf{x} = [d : m_1 : \dots : m_8]$ its corresponding point in \mathbb{P}^8 .

Claim 1. $\overline{\text{Eff}}(X) \subset K_X^{\leq 0}$.

Indeed, by Theorem 6.5 any element x in $\text{Eff}(X)$ is a finite non-negative combination of elements in $\mathcal{W} \cdot E_8$ and the class of $-\frac{1}{2}K_X$. Thus, by (22) we have $(x, -\frac{1}{2}K_X) \geq 0$ which implies $(x, K_X) \leq 0$. Now, as $K_X^{\leq 0}$ is closed we obtain the desired claim.

Claim 2. All extremal rays of $\overline{\text{Eff}}(X) \cap K_X^{\leq 0}$ are given by the classes in $\mathcal{W} \cdot E_8$.

We know that any element x in $\mathcal{W} \cdot E_8$ is a (-1) -divisor on X . Hence, $(x, K_X) < 0$ and x spans an extremal ray of $\overline{\text{Eff}}(X)$. The claim follows then by Proposition 6.10, as the rays spanned by (-1) -divisors can only accumulate towards $\overline{\text{Eff}}(X) \cap K_X^\perp$.

Claim 3. $\overline{\text{Eff}}(X) \cap K_X^\perp = \{-\frac{1}{2}K_X\}$.

As introduced above, we will identify $(\overline{\text{Eff}}(X) \cap K_X^\perp) \setminus \{0\}$ with its image, denoted by S , in $\mathbb{R}\mathbb{P}^8$. Then S is a compact set with $\mathbf{y} := [2 : 1 : \dots : 1] \in S$. Now let $x = (d, m_1, \dots, m_8) \in \overline{\text{Eff}}(X) \cap K_X^\perp$. By Lemma 4.1 the pairing (\cdot, \cdot) is negative semi-definite on K_X^\perp with a 1-dimensional kernel spanned by K_X . Thus, we have

$$(x, x) = 2d^2 - \sum_{i=1}^8 m_i^2 < 0$$

unless x is contained in the ray spanned by K_X . Consider the function $g: S \rightarrow \mathbb{R}$ defined by

$$[d : m_1 \dots m_8] \mapsto \frac{2d^2 - \sum_{i=1}^8 m_i^2}{2d^2 + \sum_{i=1}^8 m_i^2}.$$

Now as g is continuous it takes a minimum $\mathbf{x}_{\min} = [d : m_1 : \dots : m_8]$ on the compact set S , represented by a class $x_{\min} \in \overline{\text{Eff}}(X) \cap K_X^\perp$. Now assume that S contains a point different from \mathbf{y} . Then

$$g(\mathbf{x}_{\min}) < g(\mathbf{y}) = 0.$$

As $x_{\min} \in K_X^\perp$, we have $4d = \sum_{i=1}^8 m_i$. Together with $g(\mathbf{x}_{\min}) < 0$, this implies that there exists a set $I \subset \{1, \dots, 8\}$ of length 4 such that $2d < \sum_{i \in I} m_i$. So, we can perform a degree-decreasing Cremona transformation ϕ_I as defined in Proposition 4.4 to obtain an element $(d', m'_1, \dots, m'_8) = x' := \phi_I(x_{\min})$ with

$$d' < d, m'_i < m_i \text{ for } i \in I, m'_i = m_i \text{ for } i \notin I.$$

Moreover, as $(x', x') = (x_{\min}, x_{\min}) < 0$, we have

$$g(\mathbf{x}') < g(\mathbf{x}_{\min}),$$

which is a contradiction to \mathbf{x}_{\min} minimising g . Consequently, $\overline{\text{Eff}}(X)$ intersects K_X^\perp in a single ray which is therefore extremal. \square

Proof of Theorem 1.4. It follows from Theorem 6.5 and Proposition 6.11. \square

7. THE MOVABLE CONE

We recall that X denotes the blowup of \mathbb{P}^3 at eight very general points. In this section we prove Theorem 1.3. The first part of Theorem 1.3 is a direct consequence of the fact that the infinite Weyl group preserves the effective movable cone $\overline{\text{Mov}}^e(X) = \overline{\text{Mov}}(X) \cap \text{Eff}(X)$:

Lemma 7.1. *The cone $\overline{\text{Mov}}^e(X)$ is not rational polyhedral.*

Proof. Suppose by contradiction that $\overline{\text{Mov}}^e(X)$ is a rational polyhedral cone spanned by finitely many extremal rays and denote by D_1, \dots, D_k the primitive classes generating these extremal rays, i.e. each D_i is the smallest Cartier divisor class generating an extremal ray. Since the Weyl group \mathcal{W} preserves the cone $\overline{\text{Mov}}^e(X)$, any $w \in \mathcal{W}$ permutes the classes D_1, \dots, D_k . Hence the class $\sum_{i=1}^k D_i$ is fixed by the group \mathcal{W} . Moreover, $\sum_{i=1}^k D_i$ is contained in the interior of $\overline{\text{Mov}}^e(X)$ and thus is big. But any class fixed by \mathcal{W} is proportional to the class of $-\frac{1}{2}K_X$ which is not big, we obtain a contradiction. \square

The proof of the second and third part of Theorem 1.3 is more involved and we split it in several steps. First, we show that we can cover the rational points of $\text{Mov}(X)$ by \mathcal{W} -translates of a rational polyhedral cone Π , see Theorem 7.3. Then, to deduce that the whole cone $\text{Mov}(X)$ is contained in $\mathcal{W} \cdot \Pi := \bigcup_{w \in \mathcal{W}} w \cdot \Pi$, we prove that $\mathcal{W} \cdot \Pi$ is convex, see Theorem 7.9. Finally, the existence of a fundamental domain for the action of \mathcal{W} on $\overline{\text{Mov}}^e(X)$ follows from Theorem 2.3.

Lemma 7.2. *Let Π be the rational polyhedral cone in $N^1(X)_{\mathbb{R}}$ spanned by the classes of*

- H ,
- $H - E_i$ for $1 \leq i \leq 3$,
- $H - E_i - E_j$ for $1 \leq i < j \leq 3$,
- $L_k = 2H - \sum_{i=1}^k E_i$ for $4 \leq k \leq 8$,
- $M_\ell = 3H - 3E_1 - \sum_{i=2}^\ell E_i$ for $4 \leq \ell \leq 8$.

Let $D = (d; m_1, \dots, m_8)$ be a Cartier divisor class in standard form such that $d \geq m_i \geq 0$ for all $i \in \{1, \dots, 8\}$. Then $D \in \Pi$.

Proof. Let $D = (d; m_1, \dots, m_8)$ be a Cartier divisor class in standard form such that $d \geq m_i \geq 0$ for all $i \in \{1, \dots, 8\}$. Then $2d \geq \sum_{i=1}^4 m_i$ and $d \geq m_1 \geq \dots \geq m_8 \geq 0$.

Step 1. We first consider the case when $d \geq m_1 + m_4$.

By Proposition 4.10, we can write

$$D = (d - 2m_4; m_1 - m_4, m_2 - m_4, m_3 - m_4) + \sum_{i=4}^8 c_i \cdot (2; 1^i)$$

with $c_i \geq 0$.

Since every divisor class $(2; 1^i)$ for $i \geq 4$ is contained in Π , it is enough to show that for any divisor class D of the form:

$$D = (d; m_1, m_2, m_3),$$

where $2d \geq m_1 + m_2 + m_3$ and $d \geq m_1 \geq m_2 \geq m_3 \geq 0$, we have $D \in \Pi$. Note that we can assume $d > 0$, as $d = 0$ implies that $D = 0$.

We show the following stronger statement by induction on d :

Claim. Any class

$$D = (d; m_1, m_2, m_3)$$

with $d > 0$, $2d \geq m_1 + m_2 + m_3$ and $0 \leq m_i \leq d$ is contained in the cone Π .

For $d = 1$, the only possible classes are exactly the generators of Π of degree 1. Now let $d > 1$. We distinguish the following cases.

- If all m_i are zero, then D is a multiple of H which is in Π .
- If there exists exactly one $j \in \{1, 2, 3\}$ so that m_j is non-zero, then D is a non-negative combination of H and $H - E_j$ which are both in Π .
- Now let us consider the case where at least two m_j are non-zero. We assume first that $m_3 = \min\{m_i \mid i = 1, 2, 3\}$. Then $m_1 > 0$, $m_2 > 0$ and $m_3 < d$ since $2d \geq m_1 + m_2 + m_3$. Hence the class

$$(d'; m'_1, m'_2, m'_3) = (d - 1; m_1 - 1, m_2 - 1, m_3)$$

satisfies $2d' \geq \sum_{i=1}^3 m'_i$ and $0 \leq m'_i \leq d'$.

We remark that $m_1 - 1$ or $m_2 - 1$ can be strictly smaller than m_3 which explains the use of the stronger statement here. Consequently,

$$dH - \sum_{i=1}^3 m_i E_i = \underbrace{((d-1)H - \sum_{i=1}^2 (m_i - 1)E_i - E_3)}_{\in \Pi \text{ by induction}} + \underbrace{(H - E_1 - E_2)}_{\in \Pi}.$$

is contained in Π . For the case that $m_1 = \min\{m_i \mid i = 1, 2, 3\}$ (resp. $m_2 = \min\{m_i \mid i = 1, 2, 3\}$), we subtract $H - E_2 - E_3$ (resp. $H - E_1 - E_3$) from the divisor $dH - \sum_{i=1}^3 m_i E_i$ and obtain the same result.

This proves the claim.

Step 2. Now let us consider the case $d < m_1 + m_4$. We proceed as in the proof of Theorem 1.4. Let $t_{i,j} := m_i + m_j - d$ for $1 \leq i < j \leq 8$. Then $d < m_1 + m_4$ is equivalent to $t_{1,4} > 0$. This implies that $d \geq 3$ and $m_1 \geq m_2 + 2 \geq 3$. Let $a = \max\{i \mid m_i \neq 0\}$. As $d < m_1 + m_4$, we have $a \geq 4$. We consider

$$(d'; m'_1, \dots, m'_a) := (d; m_1, \dots, m_a) - (3; 3, 1^{a-1}),$$

where $a' = \max\{i \mid m'_i \neq 0\}$. For any $4 \leq a \leq 8$, the divisor class $(3; 3, 1^{a-1})$ is contained in Π . So it is enough to show that $(d'; m'_1, \dots, m'_8) \in \Pi$.

We obtain that $(d'; m'_1, \dots, m'_{a'})$ is in standard form and $d' \geq m'_i > 0$ for any $i \in \{1, \dots, a'\}$. Furthermore, setting $t'_{1,4} := m'_1 + m'_4 - d'$, we have $t'_{1,4} = t_{1,4} - 1$. If $t'_{1,4} > 0$, since $(d'; m'_1, \dots, m'_{a'})$ is in standard form, we can start our above procedure in *Step 2* again and do this until $t'_{1,4} = 0$ for some $(d'; m'_1, \dots, m'_{a'})$. We are now under the assumption in *Step 1* and thus $(d'; m'_1, \dots, m'_{a'}) \in \Pi$. \square

Theorem 7.3. *Let Π be the rational polyhedral cone in $N^1(X)_{\mathbb{R}}$ defined as in Lemma 7.2. Then*

$$\{D \in \text{Mov}(X) \mid D \text{ is a rational class}\} \subset \mathcal{W} \cdot \Pi = \bigcup_{w \in \mathcal{W}} w \cdot \Pi \subset \overline{\text{Mov}}^e(X).$$

Before proving the theorem, we make the following remark.

Remark 7.4. Consider an effective Cartier divisor class D such that $\text{Bs} |D|$ has codimension at least 2. By Remark 4.8 there exists an element $w \in \mathcal{W}$ such that $w \cdot D = (d; m_1, \dots, m_8)$ is in standard form, and we have $d \geq m_i$ for all $i \in \{1, \dots, 8\}$ as $w \cdot D$ is again effective by Remark 4.3. Now assume that the class $(d; m_1; \dots, m_8)$ has a negative coefficient m_k for some $k \in \{1, \dots, 8\}$, then the linear system $\mathcal{L}(d; m_1, \dots, m_8)$ contains E_k with multiplicity $-m_k$ in its base locus. Note that if this is the case, then there exists a fixed component with multiplicity $-m_k$ in the base locus of the starting linear system $|D|$; it is the image of E_k by w^{-1} . This gives a contradiction.

In particular, if we start with a Cartier divisor class $D \in \text{Mov}(X)$ in Remark 4.8, then we obtain a class $(d; m_1, \dots, m_8)$ in standard form with $d \geq m_i \geq 0$ for all $i \in \{1, \dots, 8\}$.

Proof. We first show the second inclusion. Clearly, $\Pi \subset \overline{\text{Mov}}^e(X)$ as we can easily verify that every generator of Π is a movable effective divisor. Moreover, for a class $D \in \overline{\text{Mov}}^e(X)$ we have $w \cdot D \in \overline{\text{Mov}}^e(X)$ for any $w \in \mathcal{W}$ by Theorem 4.2 and Remark 4.3.

For the first inclusion, let us consider a Cartier divisor class $D \in \text{Mov}(X)$. Then by Remark 4.8 there exists a divisor class $D_s = (d; m_1, \dots, m_8)$ in standard form and an element $w \in \mathcal{W}$ such that $D_s = w \cdot D$. Moreover, by Remark 7.4 we have $d \geq m_i \geq 0$ for all $i \in \{1, \dots, 8\}$. Thus by Lemma 7.2, we obtain

$$D = w^{-1} \cdot D_s \in \mathcal{W} \cdot \Pi.$$

Now let $D \in \text{Mov}(X)$ be a \mathbb{Q} -divisor class. Then for sufficiently large and divisible $m \geq 1$, we have that mD is an effective Cartier class in $\text{Mov}(X)$. Thus $D \in \mathcal{W} \cdot \Pi$. \square

To cover the whole cone $\text{Mov}(X)$ by \mathcal{W} -translates of the rational polyhedral cone Π , we need to show that $\mathcal{W} \cdot \Pi$ is convex. To do this, we use the fact that the Tits cone $T = \bigcup_{w \in \mathcal{W}} w \cdot \overline{C}$ is convex (see Subsection 4.1) and

we show that $\mathcal{W} \cdot \Pi$ is precisely the intersection of the Tits cone T with a convex closed set.

Lemma 7.5. *We have $\mathcal{W} \cdot \Pi \subset T$.*

Proof. Let us first show that $\Pi \subset T$. Since Π is a rational polyhedral cone and T is convex, it is enough to show that any generator of Π is in T . Recall from the characterization of \overline{C} in (6) that all generators of Π which are in standard form are contained in \overline{C} , and thus clearly in T . For each generator D of Π which is not in standard form, there exists an element of $w \in \mathcal{W}$ such that $w \cdot D$ is in standard form, hence $D \in w^{-1} \cdot \overline{C} \subset T$.

Since T is invariant under the action of \mathcal{W} , we have $w \cdot \Pi \subset T$ for any $w \in \mathcal{W}$ and this shows the statement. \square

Lemma 7.6. *We have $\text{Mov}(X) \subset T$.*

Proof. Let $D \in \text{Mov}(X)$. Since $\text{Mov}(X)$ is open, we can write

$$D = \sum_{i=1}^k \lambda_i D_i,$$

where $\lambda_i \in \mathbb{R}_{\geq 0}$ and $D_i \in \text{Mov}(X)$ is a Cartier divisor class for each $i \in \{1, \dots, k\}$.

Now for each $i \in \{1, \dots, k\}$, since $D_i \in \text{Mov}(X)$ is Cartier, we have $D_i \in \bigcup_{w \in \mathcal{W}} w \cdot \Pi$ by Theorem 7.3 and thus $D_i \in T$ by Lemma 7.5. The statement follows from the fact that T is convex. \square

Remark 7.7. We remark that T strictly contains $\text{Mov}(X)$. For example, $E_8 \in \overline{C} \subset T$ is effective but not movable. Moreover, since $\mathcal{W} \cdot E_8$ and $-\frac{1}{2}K_X$ are contained in T , the Tits cone T contains the whole effective cone $\text{Eff}(X)$ by Theorem 1.4.

Now we are ready to show that there exists a closed convex set \mathcal{B} such that

$$T \cap \mathcal{B} = \bigcup_{w \in \mathcal{W}} w \cdot \Pi.$$

As before we identify $N^1(X)_{\mathbb{R}}$ with \mathbb{R}^9 : an element $(d, m_1, \dots, m_8) \in \mathbb{R}^9$ corresponds to a class $dH - \sum_{i=1}^8 m_i E_i \in N^1(X)_{\mathbb{R}}$. Let

$$(23) \quad B = \{(d; m_1, \dots, m_8) \in N^1(X)_{\mathbb{R}} \mid d - m_i \geq 0 \text{ and } m_i \geq 0 \text{ for all } i = 1, \dots, 8\}.$$

Then B is the intersection of closed half-spaces in \mathbb{R}^9 , hence it is closed and convex. Furthermore, we have $\Pi \subset B$.

Fix an element $w \in \mathcal{W}$. As \mathcal{W} is generated by the orthogonal reflections in (5), we can consider w as an invertible linear map on \mathbb{R}^9 . We define

$$B_w := w^{-1} \cdot B.$$

Then B_w is closed and convex. Moreover, $D \in B_w$ if and only if $w \cdot D \in B$.

We set

$$\mathcal{B} = \bigcap_{w \in \mathcal{W}} B_w.$$

Then \mathcal{B} is the intersection of infinitely many closed convex sets, thus it is closed and convex. In particular, $D \in \mathcal{B}$ if and only if $w \cdot D \in B$ for any $w \in \mathcal{W}$.

Lemma 7.8. *We have $\mathcal{W} \cdot \Pi \subset \mathcal{B}$.*

Proof. Let $D \in \Pi$ be a generator as in Lemma 7.2. Since the class D contains a movable prime divisor P which is the strict transform of a hypersurface in \mathbb{P}^3 , we have $D \in B$. By Theorem 4.2, for any $w \in \mathcal{W}$, the class $w \cdot D$ also contains a movable prime divisor P' which is the strict transform of a hypersurface in \mathbb{P}^3 . Hence $w \cdot D \in B$ for any $w \in \mathcal{W}$. Therefore, $D \in \mathcal{B}$ and $w \cdot D \in \mathcal{B}$ for any $w \in \mathcal{W}$.

Now let $D = \sum_i \lambda_i D_i \in \Pi$, where $\lambda_i \in \mathbb{R}_{\geq 0}$ and D_i is a generator of Π for each i . For any $w \in \mathcal{W}$, we have $w \cdot D = \sum_i \lambda_i (w \cdot D_i)$ and $w \cdot D_i \in B$, thus $w \cdot D \in B$ by convexity of B . Therefore, we obtain $D \in \mathcal{B}$ and thus $w \cdot D \in \mathcal{B}$ for any $w \in \mathcal{W}$. \square

Theorem 7.9. *We have*

$$T \cap \mathcal{B} = \bigcup_{w \in \mathcal{W}} w \cdot \Pi.$$

In particular, $\mathcal{W} \cdot \Pi$ is convex.

Proof. By Lemmas 7.5 and 7.8, we have $\bigcup_{w \in \mathcal{W}} w \cdot \Pi \subset T \cap \mathcal{B}$. It remains to show the other inclusion.

Step 1. In this step we show that $\overline{\mathcal{C}} \cap B \subset \Pi$.

Let $D \in \overline{\mathcal{C}} \cap B$. We first assume that D is a Cartier divisor class and we can write it as $(d; m_1, \dots, m_8)$. By (6) and (23), we have

- $d \geq m_i \geq 0$ for all $i \in \{1, \dots, 8\}$,
- $2d \geq \sum_{i=1}^4 m_i$,
- $m_i \geq m_{i+1}$ for all $i \in \{1, \dots, 7\}$.

Hence D is in standard form with $d \geq m_i \geq 0$ for all $i \in \{1, \dots, 8\}$ and by Lemma 7.2, we obtain $D \in \Pi$. For a \mathbb{Q} -divisor $D \in \overline{\mathcal{C}} \cap B$, we have that mD is Cartier for sufficiently large and divisible m , thus $D \in \Pi$. Now for an \mathbb{R} -divisor $D \in \overline{\mathcal{C}} \cap B$, there exists a sequence of \mathbb{Q} -divisors $(D_n)_{n \in \mathbb{N}}$ in $\overline{\mathcal{C}} \cap B$ converging to D . Since $D_n \in \Pi$ for any $n \in \mathbb{N}$ and Π is closed, this implies $D \in \Pi$.

Step 2. Let $D \in T \cap \mathcal{B}$ be an \mathbb{R} -divisor. Since $D \in T = \bigcup_{w \in \mathcal{W}} w \cdot \overline{\mathcal{C}}$, there exists an element $w \in \mathcal{W}$ and an \mathbb{R} -divisor $D' \in \overline{\mathcal{C}}$ such that $w \cdot D' = D$. But $D \in \mathcal{B}$ implies in particular $w^{-1} \cdot D \in B$. Consequently,

$$D' = w^{-1} \cdot D \in \overline{\mathcal{C}} \cap B \subset \Pi$$

by Step 1, and thus $D = w \cdot D' \in w \cdot \Pi$. \square

As an immediate consequence, we obtain the following result.

Corollary 7.10. *Let Π be the rational polyhedral cone in $N^1(X)_{\mathbb{R}}$ defined as in Lemma 7.2. Then*

$$\text{Mov}(X) \subset \mathcal{W} \cdot \Pi = \bigcup_{w \in \mathcal{W}} w \cdot \Pi \subset \overline{\text{Mov}}^e(X).$$

Proof. Let $D \in \text{Mov}(X)$. Then we can write

$$D = \sum_{i=1}^k \lambda_i D_i$$

with $\lambda_i \in \mathbb{R}_{\geq 0}$ and $D_i \in \text{Mov}(X)$ is a Cartier divisor. By Theorem 7.3 we have that each D_i is contained in $\mathcal{W} \cdot \Pi$. Now, as $\mathcal{W} \cdot \Pi$ is convex by Theorem 7.9, we have $D \in \mathcal{W} \cdot \Pi$. \square

Finally, we have:

Proof of Theorem 1.3. The statement (ii) follows from Lemma 7.1.

To show that the action of \mathcal{W} on $\overline{\text{Mov}}^e(X)$ admits a fundamental domain, we apply Theorem 2.3 to $V = N^1(X)_{\mathbb{R}}$, $C = \text{Mov}(X)$ and $G = \mathcal{W} \subset \text{GL}(N^1(X)_{\mathbb{R}})$. Then $\overline{C} = \overline{\text{Mov}}(X)$ and $C_+ = \overline{\text{Mov}}^+(X)$.

By Corollary 7.10 there exists a rational polyhedral cone $\Pi \subset C_+$ with $C \subset G \cdot \Pi$. Indeed, $\Pi \subset \overline{\text{Mov}}^+(X)$ as every generator of Π is movable and rational. Thus, the second condition of Theorem 2.3 is satisfied and there exists a rational polyhedral cone $\tilde{\Pi}$ which is a fundamental domain for the action of \mathcal{W} on $\overline{\text{Mov}}^+(X)$:

$$\overline{\text{Mov}}^+(X) = \mathcal{W} \cdot \tilde{\Pi}.$$

Then, combining Corollary 7.10 and the first condition in Theorem 2.3, we obtain

$$\overline{\text{Mov}}^e(X) \supset \mathcal{W} \cdot \Pi = \overline{\text{Mov}}^+(X) = \mathcal{W} \cdot \tilde{\Pi}.$$

Furthermore, by Corollary 3.3 we obtain $\overline{\text{Mov}}^e(X) = \overline{\text{Mov}}^+(X)$. Hence, $\tilde{\Pi}$ is a fundamental domain for the action of \mathcal{W} on $\overline{\text{Mov}}^e(X)$ and this proves statement (iii). Since the cone $\text{Eff}(X)$ is closed by Proposition 6.11, we have $\overline{\text{Mov}}^e(X) = \overline{\text{Mov}}(X)$ and thus statement (i) follows. \square

Remark 7.11. We recall that for a general element $\Delta \in |-K_X|$, the pair (X, Δ) is log canonical. The question remains whether there exists a fundamental domain for the action of $\text{PsAut}(X)$ on $\overline{\text{Mov}}^e(X)$ or, even stronger, for the action of $\text{PsAut}(X, \Delta)$, the subgroup of pseudo-automorphisms preserving the boundary divisor Δ . Up to our knowledge, the group $\text{PsAut}(X)$ is not described and we expect it to be finite. If this is indeed true, then (X, Δ) is an example of a log canonical Calabi-Yau pair of dimension 3 which does not satisfy the movable cone conjecture, as $\overline{\text{Mov}}^e(X)$ is not rational polyhedral. This can be compared to the counterexample to the cone

conjecture for log canonical Calabi-Yau pairs in dimension 2 constructed as \mathbb{P}^2 blown up at 9 very general points ([Tot10]).

8. OTHER EXAMPLES

In this section, we give some further examples of log canonical Calabi-Yau pairs and study their cones of effective curves.

Our first example is a smooth threefold which is isomorphic in codimension one to \mathbb{P}^3 blown up at eight very general points. More precisely, the threefold in the example below is obtained by flopping three of the six rational curves that are flopped by the standard Cremona transformation described in Subsection 4.2.

Example 8.1. ([LO16, Subsection 5.1]) Let $Y \rightarrow \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ be the blow up of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ at 6 very general points q_1, \dots, q_6 . Let H_i be the pullback of the hyperplane class of \mathbb{P}^1 under i -th projection $p_i: \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$ for $i = 1, 2, 3$, and let $E_j \subset Y$ be the exceptional divisor over the point q_j for $j = 1, \dots, 6$. Then the anticanonical class $-K_Y$ is given by

$$-K_Y = 2\left(\sum_{i=1}^3 H_i - \sum_{j=1}^6 E_j\right).$$

We have that $-K_Y$ is nef and not semi-ample, and by taking a general member $D \in |-K_Y|$ we obtain a log canonical Calabi-Yau pair (Y, D) .

Proposition 8.2. *In Example 8.1 the cone $\overline{\text{NE}}(Y)$ is a rational polyhedral cone.*

Proof. We follow the same strategy as in the proof of [CPS13, Theorem 3.1(6)]. Although [CPS13] deals with the blowup of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ at 6 special points such that the obtained threefold has nef and semi-ample anticanonical divisor, the same argument works here. We indicate below the part of their proof which is needed in our case.

Let ℓ_i be the class dual to H_i for $i = 1, 2, 3$ and let e_j be the class of line in E_j for $1 \leq j \leq 6$. Consider the subcone \mathcal{C} of the closed cone of curves $\overline{\text{NE}}(Y)$ spanned by the curve classes:

$$\ell_i, e_j, \ell_i - e_j, \text{ for } i = 1, 2, 3 \text{ and } 1 \leq j \leq 6.$$

We can easily verify that the following divisor classes span the dual cone \mathcal{C}^\vee of \mathcal{C} :

$$(24) \quad H_1, H_2, H_3, H_1 + H_2 + H_3 - \sum_{j \in J} E_j, \text{ for } J \subset \{1, \dots, 6\} \text{ with } 1 \leq |J| \leq 6.$$

Thus we have $\text{Nef}(Y) \subset \mathcal{C}^\vee$.

It remains to show that every divisor class in (24) is nef, so that $\mathcal{C}^\vee \subset \text{Nef}(Y)$, and thus we conclude that $\text{Nef}(Y) = \mathcal{C}^\vee$ and $\overline{\text{NE}}(Y) = \mathcal{C}$ are rational polyhedral cones.

Note that H_1, H_2, H_3 are nef, being pullbacks of ample divisors by morphisms. Now consider a subset $J \subset \{1, \dots, 6\}$ with $1 \leq |J| \leq 6$ and a divisor

$$(25) \quad D_J := H_1 + H_2 + H_3 - \sum_{j \in J} E_j = -\frac{1}{2}K_Y + \sum_{k \notin J} E_k.$$

Assume by contradiction that D_J is not nef. Then there exists an irreducible curve $C \subset Y$ such that $D_J \cdot C < 0$. Since $-K_Y$ is nef, by (25) there exists an exceptional divisor E_{k_0} with $k_0 \notin J$ such that $E_{k_0} \cdot C < 0$. Hence $C \subset E_{k_0}$ and we obtain that C is proportional to e_{k_0} . But $E_{k_0} \cdot e_{k_0} = -1$, $E_k \cdot e_{k_0} = 0$ for $k \neq k_0$ and $-\frac{1}{2}K_Y \cdot e_{k_0} = 1$, and thus we obtain $D_J \cdot C = 0$ by (25). This gives a contradiction and finishes the proof. \square

Now we give an example of a log canonical Calabi-Yau pair whose cone of effective curves has infinitely many extremal rays. Note that this example is not isomorphic in codimension one to \mathbb{P}^3 blown up at eight very general points, since they have different Picard numbers.

Example 8.3. ([Xie20, Example 3.6]) Let S be \mathbb{P}^2 blown up at 9 very general points such that $-K_S$ is nef, not semi-ample and the unique member in $|-K_S|$ is a smooth elliptic curve. Now define $\mathcal{E} := \mathcal{O}_S \oplus \mathcal{O}_S(-K_S)$ and $p: Y := \mathbb{P}(\mathcal{E}) \rightarrow S$. Then $-K_Y \sim 2\xi$ is nef and not semi-ample, where ξ is a tautological divisor associated to $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$. Moreover, a general member in $|\frac{1}{2}K_Y|$ is a smooth surface isomorphic to S , and two general members in $|\frac{1}{2}K_Y|$ intersect transversally along a smooth elliptic curve. Furthermore, a general member $D \in |-K_Y|$ is reducible, and (Y, D) forms a log canonical Calabi-Yau pair.

Proposition 8.4. *In Example 8.3 the cone $\overline{\text{NE}}(Y)$ has infinitely many K_Y -trivial extremal rays.*

Proof. We follow closely the proof of [CJR08, Lemma 2.6]. Let C be a (-1) -curve on the surface S . Twisting by some line bundle \mathcal{L} on C , we may write

$$\mathcal{V} := \mathcal{E}|_C \otimes \mathcal{L} = \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(a)$$

for some integer $a \geq 0$.

Let $S_C := p^{-1}(C) \simeq \mathbb{P}(\mathcal{V})$ and let $\tilde{C} \rightarrow C$ be the section in S_C over C corresponding to the quotient $\mathcal{V} \rightarrow \mathcal{O}_{\mathbb{P}^1} \rightarrow 0$. Since C is a (-1) -curve on S , we have $S_C \cdot \tilde{C} = p^*C \cdot \tilde{C} = C \cdot p_*\tilde{C} = -1$ and thus

$$(26) \quad N_{S_C/Y}|_{\tilde{C}} \simeq \mathcal{O}_{\mathbb{P}^1}(-1).$$

Let C_0 be a tautological divisor associated to the line bundle $\mathcal{O}_{\mathbb{P}(\mathcal{V})}(1)$ and let ℓ be a fibre of $p|_{S_C}: S_C \rightarrow C \simeq \mathbb{P}^1$. Then $\tilde{C} \in |C_0 - a\ell|$. Moreover, by the Grothendieck relation we have

$$(27) \quad C_0^2 - C_0 \cdot (p|_S)^*(c_1(\mathcal{V})) = 0, \text{ i.e. } C_0 \cdot \tilde{C} = 0.$$

Hence, by the adjunction formula and (26)(27), we obtain

$$\begin{aligned}
-K_Y \cdot \tilde{C} &= (-K_{S_C} + S_C)|_{S_C} \cdot \tilde{C} \\
&= (2C_0 - (p|_{S_C})^*(K_C + \det \mathcal{V})) \cdot \tilde{C} - 1 \\
&= 2C_0 \cdot \tilde{C} - (-2 + a) - 1 \\
&= 1 - a.
\end{aligned}$$

Since $-K_Y$ is nef and divisible by two in $\text{Pic}(Y)$, we have $a = 1$. Thus $-K_Y \cdot \tilde{C} = 0$, and we obtain

$$S_C \simeq \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(1)) \simeq \mathbb{F}_1$$

with $-K_{S_C} \cdot \tilde{C} = 1$, i.e. \tilde{C} is the minimal section of the ruled surface \mathbb{F}_1 .

Therefore, we have $N_{\tilde{C}/S_C} \simeq \mathcal{O}_{\mathbb{P}^1}(-1)$. Together with (26), this implies the splitting of the normal bundle sequence

$$0 \rightarrow N_{\tilde{C}/S_C} \rightarrow N_{\tilde{C}/Y} \rightarrow N_{S_C/Y}|_{\tilde{C}} \rightarrow 0,$$

and thus $N_{\tilde{C}/Y} \simeq \mathcal{O}_{\mathbb{P}^1}(-1)^{\oplus 2}$. Blowing up the curve \tilde{C} in Y , the exceptional divisor is isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$ and we may blow down in the other direction onto some smooth threefold Y^+ :

$$\begin{array}{ccc}
& \text{Bl}_{\tilde{C}}(Y) & \\
& \swarrow \quad \searrow & \\
Y & \overset{\chi}{\dashrightarrow} & Y^+,
\end{array}$$

where the birational map χ is a flop. Note that the K_Y -trivial ray generated by the class $[\tilde{C}]$ is extremal by [Xie20, Lemma 2.5] applied to the pair (Y, S_C) . Therefore, every (-1) -curve on S corresponds to a K_Y -trivial extremal ray. Now since there are infinitely many (-1) -curves on the surface S (see [Dol83, Corollary 1]), we obtain infinitely many K_Y -trivial extremal rays. \square

In view of these examples and the decomposition of the effective movable cone

$$\overline{\text{Mov}}^e(X) = \bigcup_{\varphi: X \dashrightarrow X' \text{ a finite sequence of flops}} \varphi^*((\text{Nef}^e(X'))).$$

from Corollary 3.4, it is natural to ask the following question:

Question. Let X be the blowup of \mathbb{P}^3 at 8 very general points and let $X \dashrightarrow X'$ be a finite sequence of flops to another smooth projective threefold X' . Is the nef cone $\text{Nef}(X')$ a rational polyhedral cone?

REFERENCES

- [BC13] T. Bayraktar and S. Cantat. Constraints on automorphism groups of higher dimensional manifolds. *J. Math. Anal. Appl.*, 405(1):209–213, 2013.
- [BDP16] M. C. Brambilla, O. Dumitrescu, and E. Postinghel. On the effective cone of \mathbb{P}^n blown-up at $n + 3$ points. *Exp. Math.*, 25(4):452–465, 2016.

- [BH14] C. Birkar and Z. Hu. Polarized pairs, log minimal models, and Zariski decompositions. *Nagoya Math. J.*, 215:203–224, 2014.
- [BH15] J. Blanc and I. Hedén. The group of Cremona transformations generated by linear maps and the standard involution. *Ann. Inst. Fourier (Grenoble)*, 65(6):2641–2680, 2015.
- [Bor91] C. Borcea. On desingularized Horrocks-Mumford quintics. *J. Reine Angew. Math.*, 421:23–41, 1991.
- [BZ16] C. Birkar and D.-Q. Zhang. Effectivity of Iitaka fibrations and pluricanonical systems of polarized pairs. *Publ. Math. Inst. Hautes Études Sci.*, 123:283–331, 2016.
- [CJR08] C. Casagrande, P. Jahnke, and I. Radloff. On the Picard number of almost Fano threefolds with pseudo-index > 1 . *Internat. J. Math.*, 19(2):173–191, 2008.
- [CLO16] I. Coskun, J. Lesieutre, and J. C. Ottem. Effective cones of cycles on blowups of projective space. *Algebra Number Theory*, 10(9):1983–2014, 2016.
- [CPS13] I. Coskun and A. Prendergast-Smith. Fano manifolds of index $n - 1$ and the cone conjecture. *Int. Math. Res. Not.*, 2014(9):2401–2439, 01 2013.
- [CPS19] I. Coskun and A. Prendergast-Smith. Fano manifolds of index $n - 2$ and the cone conjecture. *Math. Proc. Cambridge Philos. Soc.*, 166(1):1–31, 2019.
- [CT06] A.-M. Castravet and J. Tevelev. Hilbert’s 14th problem and Cox rings. *Compositio Mathematica*, 142(6):1479–1498, 2006.
- [DL06] C. De Volder and A. Laface. Base locus of linear systems on the blowing-up of \mathbb{P}^3 along at most 8 general points. *Pacific J. Math.*, 223(1):17–34, 2006.
- [DL07] C. De Volder and A. Laface. On linear systems of \mathbb{P}^3 through multiple points. *Journal of Algebra*, 310(1):207–217, 2007.
- [DM22] O. Dumitrescu and R. Miranda. Coxeter theory for curves on blowups of \mathbb{P}^r . [arXiv:2205.13605](https://arxiv.org/abs/2205.13605), 2022.
- [Dol83] I. V. Dolgachev. Weyl groups and Cremona transformations. In *Singularities, Part 1 (Arcata, Calif., 1981)*, volume 40 of *Proc. Sympos. Pure Math.*, pages 283–294. Amer. Math. Soc., Providence, RI, 1983.
- [Dol11] I. V. Dolgachev. Cremona special sets of points in products of projective spaces. In *Complex and differential geometry*, volume 8 of *Springer Proc. Math.*, pages 115–134. Springer, Heidelberg, 2011.
- [DP19] O. Dumitrescu and N. Priddis. On divisorial (i) classes. [arXiv:1905.00074v3](https://arxiv.org/abs/1905.00074v3), 2019.
- [Dru11] S. Druel. Quelques remarques sur la décomposition de Zariski divisorielle sur les variétés dont la première classe de Chern est nulle. *Math. Z.*, 267(1-2):413–423, 2011.
- [HK00] Y. Hu and S. Keel. Mori dream spaces and GIT. volume 48, pages 331–348. 2000.
- [HL21] J. Han and W. Liu. On a generalized canonical bundle formula for generically finite morphisms. *Ann. Inst. Fourier (Grenoble)*, 71(5):2047–2077, 2021.
- [HL22] J. Han and Z. Li. Weak Zariski decompositions and log terminal models for generalized pairs. *Math. Z.*, 302(2):707–741, 2022.
- [Kaw97] Y. Kawamata. On the cone of divisors of Calabi-Yau fiber spaces. *Int. J. Math.*, 8(5):665–687, 1997.
- [KM98] J. Kollár and S. Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [Kol89] J. Kollár. Flops. *Nagoya Math. J.*, 113:15–36, 1989.
- [Les14] John Lesieutre. The diminished base locus is not always closed. *Compositio Mathematica*, 150(10):1729–1741, 2014.

- [Li22] J. Li. A cone conjecture for log Calabi-Yau surfaces. [arXiv:2207.12483](#), 2022.
- [LMP⁺22] V. Lazić, S. Matsumura, Th. Peternell, N. Tsakanikas, and Z. Xie. The nonvanishing problem for varieties with nef anticanonical bundle. [arXiv:2202.13814v3](#), 2022.
- [LO16] J. Lesieutre and J. C. Ottem. Curves disjoint from a nef divisor. *Mich. Math. J.*, 65(2):321–332, 2016.
- [Loo14] E. Looijenga. Discrete automorphism groups of convex cones of finite type. *Compositio Mathematica*, 150(11):1939–1962, 2014.
- [LOP18] V. Lazić, K. Oguiso, and Th. Peternell. The Morrison-Kawamata Cone Conjecture and Abundance on Ricci flat manifolds. In *Uniformization, Riemann-Hilbert correspondence, Calabi-Yau manifolds & Picard-Fuchs equations*, volume 42 of *Advanced Lectures in Mathematics*, pages 157–186. International Press, 2018.
- [LT22] V. Lazić and N. Tsakanikas. Special MMP for log canonical generalised pairs (with an appendix joint with Xiaowei Jiang). *Selecta Math. (N.S.)*, 28(5):Paper No. 89, 2022.
- [MM83] S. Mori and S. Mukai. On Fano 3-folds with $B_2 \geq 2$. In *Algebraic varieties and analytic varieties (Tokyo, 1981)*, volume 1 of *Adv. Stud. Pure Math.*, pages 101–129. North-Holland, Amsterdam, 1983.
- [Mor93] D. R. Morrison. Compactifications of moduli spaces inspired by mirror symmetry. In *Journées de géométrie algébrique d’Orsay - Juillet 1992*, number 218 in Astérisque. Société mathématique de France, 1993.
- [Muk01] S. Mukai. Counterexample to hilbert’s fourteenth problem for three dimensional additive groups. RIMS preprint no. 1343, Kyoto, 2001.
- [Muk04] S. Mukai. Geometric realization of T -shaped root systems and counterexamples to Hilbert’s fourteenth problem. In *Algebraic transformation groups and algebraic varieties*, volume 132 of *Encyclopaedia Math. Sci.*, pages 123–129. Springer, Berlin, 2004.
- [PS12] A. Prendergast-Smith. The cone conjecture for some rational elliptic threefolds. *Math. Z.*, 272(1-2):589–605, 2012.
- [PS15] A. Prendergast-Smith. Finiteness results for 3-folds with semiample anticanonical bundle. *Math. Res. Lett.*, 22(2):549–578, 2015.
- [Tot08] B. Totaro. Hilbert’s 14th problem over finite fields and a conjecture on the cone of curves. *Compositio Mathematica*, 144(5):1176–1198, 2008.
- [Tot10] B. Totaro. The cone conjecture for Calabi-Yau pairs in dimension 2. *Duke Math. J.*, 154(2):241–263, 2010.
- [Xie20] Z. Xie. Rationally connected threefolds with nef and bad anticanonical divisor. [arXiv:2006.08234v2](#).

FACHRICHTUNG MATHEMATIK, CAMPUS, GEBÄUDE E2.4, UNIVERSITÄT DES SAARLANDES, 66123 SAARBRÜCKEN, GERMANY

Email address: stenger@math.uni-sb.de

FACHRICHTUNG MATHEMATIK, CAMPUS, GEBÄUDE E2.4, UNIVERSITÄT DES SAARLANDES, 66123 SAARBRÜCKEN, GERMANY

Email address: xie@math.uni-sb.de