

Aron–Berner–type extension in complex Banach manifolds*

László Lempert
 Department of Mathematics
 Purdue University
 West Lafayette, IN 47907-2067, USA

Abstract

Let S be a compact Hausdorff space and X a complex manifold. We consider the space $C(S, X)$ of continuous maps $S \rightarrow X$, and prove that any bounded holomorphic function on this space can be continued to a holomorphic function, possibly multivalued, on a larger space $B(S, X)$ of Borel maps. As an application we prove two theorems about bounded holomorphic functions on $C(S, X)$, one reminiscent of the Monodromy Theorem, the other of Liouville’s Theorem.

Introduction

One version of the Aron–Berner extension theorem considers a complex Banach space \mathfrak{X} and for every bounded holomorphic function on its unit ball provides a holomorphic extension to the unit ball of the bidual (second dual) $\mathfrak{X}'' \supset \mathfrak{X}$. After the initial, somewhat weaker result of Aron and Berner, the sharp form above was proved by Davie and Gamelin. Aron and Berner also proved a similar theorem with the ball in \mathfrak{X} replaced by a general open subset of \mathfrak{X} , that was sharpened subsequently by Zalduendo [AB78, DG89, Z90].—It is well understood in infinite dimensional complex analysis that the result is tight in many ways. While linear forms on \mathfrak{X} can be extended to an arbitrary superspace \mathfrak{Y} , this is no longer generally true for holomorphic functions, not even for quadratic functions. Work of Dineen and Josefson gives that extension even to \mathfrak{X}'' may fail without the boundedness assumption. For example, a holomorphic function f on the space $\mathfrak{X} = c_0$ of complex sequences tending to 0 extends to a holomorphic function on $\mathfrak{X}'' \approx l^\infty$ if and only if f is bounded on every bounded subset of \mathfrak{X} [D71, J78].

A natural question is whether similar extension is possible for complex Banach manifolds \mathfrak{X} rather than open subsets of Banach spaces. Of course, before talking

*2020 Mathematics subject classification 32K05, 46G20, 58D10

about extensions of holomorphic functions, one should first construct a manifold \mathfrak{X}'' to play the role of the bidual, apparently a hard problem. Dineen and Venkova addressed it when \mathfrak{X} is a Riemann domain over a certain type of Banach space, and extended holomorphic functions to a Riemann domain lying above the bidual [DV04]. But in this paper we do not attempt a corresponding construction of \mathfrak{X}'' in general, and instead produce extensions in a more specialized framework, of Banach manifolds that arise as mapping spaces.

Let us start with a finite dimensional connected complex manifold X and a compact Hausdorff space S . The space $C(S, X)$ of continuous maps $\mathfrak{r} : S \rightarrow X$ has the natural structure of a complex Banach manifold, see [L04, LS07] and section 2 here. (When working with X , we will use x, y , etc. to denote points of X , and the corresponding fraktur fonts $\mathfrak{r}, \mathfrak{r}, \dots$ to denote mappings into X .) Whatever the bidual of $C(S, X)$ should be, it must have something to do with the bidual of the Banach space $C(S)$ of continuous functions $S \rightarrow \mathbb{C}$. This bidual has been described, in slightly different ways, first by Kakutani, then by Gordon, as the space of continuous functions on a related space T [G66, K41]; the book [D⁺16] gives yet another variant of the construction of T and of the isomorphism $C(S)'' \approx C(T)$. This suggests that the bidual of a mapping space $\mathfrak{X} = C(S, X)$ should be $\mathfrak{X}'' = C(T, X)$. It turns out that the associations $(S, X) \mapsto C(S, X) = \mathfrak{X}$ and $(S, X) \mapsto C(T, X) = \mathfrak{X}''$ are parts of (bi)functors between which there is a natural transformation that embeds \mathfrak{X} as a complex submanifold into \mathfrak{X}'' . The question then would be whether bounded holomorphic functions on \mathfrak{X} extend to \mathfrak{X}'' .

This is still not the question we will answer in this paper, though. With an application in mind, that we will discuss shortly, we rather consider extension to a smaller manifold

$$(0.1) \quad B(S, X) = \{\mathfrak{r} : S \rightarrow X \text{ is Borel} \mid \mathfrak{r}(S) \subset X \text{ has compact closure}\}.$$

(A map $\mathfrak{r} : S \rightarrow X$ is Borel if preimages of Borel subsets of X are Borel.) The space of Borel maps in (0.1) is also a complex Banach manifold. This is the space to which holomorphic functions on $C(S, X) \subset B(S, X)$ are to be extended.

It turns out that unlike in the Aron–Berners extension theorem, instead of holomorphic functions it is better to work with possibly multivalued holomorphic functions, that is, holomorphic germs that admit analytic continuations along paths. We will define the relevant notions in section 7. For the moment we first submit that the Aron–Berners extension provides for any holomorphic function germ \mathbf{f} on $C(S, X)$ at some \mathfrak{r}_1 a holomorphic germ \mathbf{f}^{ab} on $B(S, X)$ (see section 5). We will denote the germ of a complex manifold M at some $p \in M$ by (M, p) and, accordingly, a germ of a function on M at $p \in M$ as $\mathbf{f} : (M, p) \rightarrow \mathbb{C}$.

Theorem 0.1. *Let $\mathfrak{r}_1 \in C(S, X)$. If a holomorphic germ $\mathbf{f} : (C(S, X), \mathfrak{r}_1) \rightarrow \mathbb{C}$ can be analytically continued along any path in $C(S, X)$ starting at \mathfrak{r}_1 , and these continuations are uniformly bounded, then \mathbf{f}^{ab} can be analytically continued along any path in $B(S, X)$ starting at \mathfrak{r}_1 .*

Without the boundedness assumption the theorem would be false, already if $X = \mathbb{C}$ and S is either $[0, 1]$, or it consists of a convergent sequence and its limit. Interestingly,

in the theorem even if the analytic continuations of \mathbf{f} define a single valued function on $C(S, X)$, the continuations of \mathbf{f}^{ab} may define a multi-valued function on $B(S, X)$. We will discuss this and related examples in section 11.

Since $B(S, X)$ is connected, Theorem 0.1 has the paradoxical consequence that if the germ \mathbf{f} has uniformly bounded analytic continuations along paths in $C(S, X)$ starting at \mathfrak{r}_1 , then it has *canonical* continuations to all of the possibly disconnected $C(S, X)$.

Spaces of continuous maps are, of course, central objects in topology, while Borel maps occur rarely, if ever, and for good reason: The topology of $B(S, X)$ is too directly related to the topology of X , see Lemma 10.2 (and (7.1), Lemma 9.4a). But this is precisely the reason why the connection between the two mapping spaces, described in Theorem 0.1, has implications about the continuous mapping spaces themselves:

Theorem 0.2 (Monodromy Theorem). *Suppose X is simply connected and $\mathfrak{r}_1 \in C(S, X)$. Assume a holomorphic germ $\mathbf{f} : (C(S, X), \mathfrak{r}_1) \rightarrow \mathbb{C}$ has analytic continuation along any path in $C(S, X)$ starting at \mathfrak{r}_1 . If these continuations are uniformly bounded, then*

- (a) *they define a single valued holomorphic function on the component of \mathfrak{r}_1 in $C(S, X)$.*
- (b) *If, in addition, X is compact, then this single valued function is in fact constant.*

We emphasize that X , and not as in the traditional Monodromy theorem the source manifold $C(S, X)$, is assumed to be simply connected.—Both theorems will be proved in greater generality in sections 9 and 10.

That holomorphic functions (bounded or not) on certain mapping spaces must be constant for deeper reasons was first discovered by Dineen and Mellon [DM98]. Subsequently with Szabó we generalized to spaces of ‘based’ maps when X is rationally connected, e.g., a Grassmannian; if S is a manifold, constancy follows even for spaces of C^k maps [L04, LS07]. Nonetheless, similarity with Theorem 0.2b is formal only, and the underlying mechanisms are different. That this must be so can be seen upon comparing what is covered here and there. On the one hand, the manifolds X of Theorem 0.2 are vastly more general than the rationally connected manifolds of [LS07]; on the other, the requirement of boundedness and that we work with spaces of continuous, rather than C^k maps, severely restricts the scope of Theorem 0.2b, say. But this is inevitable. For general compact simply connected X and a compact smooth manifold S the spaces $C^k(S, X)$ of C^k maps may support nonconstant holomorphic functions when $k \geq 1$. For example, a Borel measure μ on the circle S^1 and a holomorphic section α of $(T^*X)^{\otimes m}$, not identically vanishing on multivectors of form $\xi \otimes \xi \otimes \cdots \in (TX)^{\otimes m}$, induces a holomorphic function $a : C^1(S^1, X) \rightarrow \mathbb{C}$,

$$a(\mathfrak{r}) = \int_{S^1} \alpha(\mathfrak{r} \otimes \mathfrak{r} \otimes \cdots) d\mu,$$

typically not even locally constant.

Related results have been proved by Aron, Galindo, Pinasco, and Zalduendo. In [A⁺16] they consider, for a compact space S , holomorphic functions on $C(S)$ that are invariant under the group of linear maps $C(S) \rightarrow C(S)$ induced by homeomorphisms $S \rightarrow S$. Among other things they find that when S is a circle or a closed disc, such

functions must be constant. While their assumptions are very different from ours: here boundedness, there invariance, both proofs rely on the idea of Aron–Berner extension.

Contents. Section 1 discusses the notion of Borel vector bundles. These are needed to endow the space $B(S, X)$ with the structure of a complex Banach manifold. Sections 2 and 3 review and discuss basic properties of mapping spaces and of the Aron–Berner extension in Banach spaces. Section 4 is about extension in local models of mapping spaces: from spaces of continuous sections of certain topological vector bundles to spaces of Borel sections. In particular, we prove a naturality property of this extension, Corollary 4.6. Section 5 then uses this naturality to extend holomorphic germs \mathbf{f} on $C(S, X)$ to holomorphic germs \mathbf{f}^{ab} on $B(S, X)$. All of this is rather straightforward. Sections 6 and 7 discuss notions of holomorphic Banach bundles, flat norms, and analytic continuation. Theorem 7.5 generalizes Theorem 0.1 to sections of certain Banach bundles. Section 8 features a key result: The construction of a family of holomorphic maps into X of balls in finite dimensional vector spaces. These will be used in section 9 to reduce analytic continuation in $B(S, X)$ to analytic continuation in Banach spaces. The proofs of Theorems 0.1, 0.2, generalized, are given in sections 9 and 10. Section 11 contains examples that show that the main theorems are optimal in certain ways.

During the preparation of this paper I have profited from discussion with Pablo Galindo.

Our basic reference to complex analysis in Banach spaces is Mujica’s book [M86], and to infinite dimensional complex manifolds, [L98, Section 2]. The infinite dimensional complex manifolds that we will encounter are what [L98] calls rectifiable, and modeled on complex Banach spaces; all this means is that they are locally biholomorphic to open subsets of Banach spaces. We denote the set of holomorphic maps between complex manifolds M, N by $\mathcal{O}(M, N)$, and by $\mathcal{O}(M)$ when $N = \mathbb{C}$.

1 Borel vector bundles

While continuous mapping spaces $C(S, X)$ are Banach manifolds modeled on spaces of continuous sections of topological vector bundles, the local models of spaces $B(S, X)$ of Borel maps are spaces of sections of what we call Borel vector bundles. In this section we develop the notion in the generality we will need. Related but different notions, such as measurable fields and bundles of Banach, especially Hilbert, spaces have been introduced before.

Let S be a topological space and $B(S)$ the algebra of bounded Borel functions $S \rightarrow \mathbb{C}$. Consider a map $\pi : V \rightarrow S$ of sets, with each $V_s = \pi^{-1}s$, $s \in S$, endowed with the structure of a finite dimensional complex vector space. The space of all sections of π , with pointwise addition and multiplication by bounded Borel functions, forms a $B(S)$ –module. By a Borel field of (finite dimensional) vector spaces we mean the datum of such $\pi : V \rightarrow S$ plus a $B(S)$ –submodule Ξ of the module of all sections. As is customary, instead of $(\pi : V \rightarrow S, \Xi)$, often we just write $\pi : V \rightarrow S$, or even just V , for a Borel field of vector spaces, and Ξ is only implied. If $\dim V_s$ is the same for all $s \in S$, we call this number the rank of V .

Suppose $T \subset S$ is Borel. We write $V|T$ first for $\pi^{-1}T$, but also for the Borel field $(\pi^{-1}T \rightarrow T, \Xi|T)$, where $\Xi|T = \{\xi|T : \xi \in \Xi\}$. This is the restriction of V to T .

A homomorphism of Borel fields of vector spaces $(V \rightarrow S, \Xi)$, $(W \rightarrow S, H)$ is a map $F : V \rightarrow W$, linear between V_s and W_s , $s \in S$, such that $F \circ \xi \in H$ for all $\xi \in \Xi$. A homomorphism is an isomorphism if it has an inverse, which is also a homomorphism.

Any finite dimensional complex vector space A defines a trivial Borel field of vector spaces $S \times A \rightarrow S$ and Ξ consisting of sections that correspond to bounded Borel functions $S \rightarrow A$. Denote the space of linear maps between vector spaces A, B by $\text{Hom}(A, B)$. Any bounded Borel map $f : S \rightarrow \text{Hom}(A, B)$ induces a homomorphism

$$F : S \times A \ni (s, a) \mapsto (s, f(s)a) \in S \times B$$

of trivial Borel fields of vector spaces, and any homomorphism arises in this way.

Definition 1.1. *A Borel vector bundle is a Borel field of vector spaces that is isomorphic to a trivial Borel field of vector spaces.*

Admittedly, this does not have the ring of “vector bundle”, that should be trivial only “locally”. Still, if S is the finite union of Borel sets S_j , and for a Borel field of vector spaces $V \rightarrow S$ the restrictions $V|_{S_j}$ are isomorphic to trivial fields of the same rank, then V itself is isomorphic to a trivial Borel field of vector spaces.

Any topological vector bundle $\pi : V \rightarrow S$ over a compact base, with equidimensional fibers, determines a Borel vector field $(\pi : V \rightarrow S, \Xi)$: a section ξ of π is in Ξ if in any topological trivialization over some compact $K \subset S$, $V|_K \approx K \times A$, it corresponds to a bounded Borel function $K \rightarrow A$.—Further examples of Borel vector bundles can be obtained by pullback. Suppose $(\pi : V \rightarrow S, \Xi)$ is a Borel vector bundle, and $g : T \rightarrow S$ is a Borel map of topological spaces. Then $g^*V = \coprod_{t \in T} V_{g(t)}$, with the obvious projection $\rho : g^*V \rightarrow T$, and the $B(T)$ –submodule H generated by sections of form $\eta(t) = \xi(g(t))$, $\xi \in \Xi$, defines the pullback Borel vector bundle $(g^*V \rightarrow T, H)$, also denoted g^*V . (Here we identify fibers of the pullback of a bundle V with the fibers of V itself, something we will do elsewhere in the paper as well.) Equivalently, H consists of those sections η of ρ for which the map $T \ni t \mapsto \eta(t) \in V_{g(t)} \subset V$ is bounded and Borel, these properties measured using any trivialization $V \approx S \times A$.

A norm on a trivial Borel vector bundle $S \times A \rightarrow S$ is a Borel function $|\cdot| : S \times A \rightarrow [0, \infty)$ whose restrictions to the fibers $\{s\} \times A$ define norms $|\cdot|_s$ on A , and there is a $c \in \mathbb{R}$ such that $|\cdot|_s \leq c|\cdot|_t$ for all $s, t \in S$. Any isomorphism $F : S \times B \rightarrow S \times A$ pulls back the norm $|\cdot|$ to a norm $|\cdot| \circ F$ on $S \times B \rightarrow S$. Therefore we can define the notion of a norm on a general Borel vector bundle $V \rightarrow S$ as a function $V \rightarrow [0, \infty)$ that, under some (or an arbitrary) isomorphism $V \approx S \times A$, corresponds to a norm on $S \times A \rightarrow S$. Such a norm is measurable with respect to the Borel σ –algebra of V , defined as the pullback of the Borel σ –algebra of $S \times A$ by any isomorphism $V \approx S \times A$.

Given a Borel vector bundle $(V \rightarrow S, \Xi)$, it will be convenient to denote Ξ by¹ $B(V)$, and refer to it as the space of bounded Borel sections of V . A norm $|\cdot|$ on V turns $B(V)$ into a Banach space, with norm $\|\xi\| = \sup_S |\xi|$. The topology on $B(V)$ is independent of the choice of $|\cdot|$.

¹That we use similar notation $B(S)$, $B(V)$ —and later $C(S)$, $C(V)$ —for rather different objects should not confuse the reader. In the former S is a topological space and $B(S)$ the space bounded Borel functions on it; V is Borel vector bundle over S . Similarly, when S is a compact Hausdorff space and $V \rightarrow S$ a topological vector bundle, $C(S)$ is the space of continuous functions on S , and $C(V)$ is the space of continuous sections.

We finish this section by relating analysis on a Borel vector bundle $\pi : V \rightarrow S$ and on the Banach(able) space $B(V)$. Given $U \subset V$, a norm $|\cdot| = p$ on V , and $\varepsilon > 0$, set

$$U_{p,\varepsilon} = \{u \in U : p(u) < 1/\varepsilon \text{ and } U \text{ contains all } v \in V_{\pi u} \text{ with } p(u-v) < \varepsilon\}.$$

If q is another norm on V , then $U_{p,\varepsilon} \subset U_{q,c\varepsilon}$ with some $c > 0$ independent of ε . Define

$$(1.1) \quad \hat{U} = \text{int} \{ \xi \in B(V) : \xi(S) \subset U \} = \bigcup_{\varepsilon > 0} \{ \xi \in B(V) : \xi(S) \subset U_{p,\varepsilon} \}.$$

Consider another Borel vector bundle $W \rightarrow S$ and a fiberwise $H : U \rightarrow W$, i.e., $H(U_s) \subset W_s$, where $U_s = U \cap V_s$. Pick norms p, q on V, W .

Definition 1.2. *We say that H is of bounded type if q is bounded on $H(U_{p,\varepsilon})$ for all $\varepsilon > 0$.*

Clearly, bounded type are independent of the choice of p, q

Lemma 1.3. *Suppose the fibers $U_s \subset V_s$ are open. If $H : U \rightarrow W$ is Borel, of bounded type, and $H|_{U_s}$ are holomorphic for $s \in S$, then*

$$H^\circ : \hat{U} \ni \xi \mapsto H \circ \xi \in B(W)$$

is holomorphic.

Proof. That H° maps into $B(W)$ follows from the definition in a straightforward way. To prove holomorphy, we can assume that $V = S \times A$, $W = S \times B$ are trivial. We will write $d_A H$ for the differential of H along A . Cauchy's formula along one dimensional slices in $\{s\} \times A$ gives that partials of H along A are Borel and bounded on $U_{p,\varepsilon}$. Using this for first derivatives gives that H° is continuous; for second derivatives, that the directional derivatives

$$dH^\circ(\xi; \eta) = \lim_{\mathbb{R} \ni t \rightarrow 0} (H^\circ(\xi + t\eta) - H^\circ(\xi))/t,$$

exist for any $\xi \in \hat{U}$ and $\eta \in B(V)$, and $dH^\circ(\xi; \eta)(s)$ can be computed as $d_A H(\xi(s); \eta(s))$. Now $S \times TA \rightarrow S$ and $S \times TB \rightarrow S$ have a natural structure of a Borel vector bundle. Since the fiberwise tangent bundle $T_A U = \coprod_{s \in S} T(U \cap (\{s\} \times A))$ is a Borel subset of $S \times TA$, $dH_A : T_A U \rightarrow S \times TB$ is Borel, of bounded type, and holomorphic along the fibers, $dH^\circ(\xi; \eta)$ is a continuous function of ξ, η . It is also complex linear in η , which proves that H° is holomorphic.

2 The mapping spaces

Henceforward we work with a fixed finite dimensional, connected complex manifold X and a compact Hausdorff space S . Write $TX \rightarrow X$ for the real tangent bundle, on which the complex structure of X induces the structure of a holomorphic vector bundle. As in the Introduction, we let

$$B(S, X) = \{ \mathfrak{r} : S \rightarrow X \text{ is Borel} \mid \mathfrak{r}(S) \subset X \text{ has compact closure} \},$$

and endow it with the uniform topology. Basic neighborhoods of $\mathfrak{r} \in B(S, X)$ are determined by open neighborhoods $D \subset X \times X$ of the diagonal $\{(x, x) : x \in X\}$, the corresponding open neighborhood of \mathfrak{r} being

$$(2.1) \quad D(\mathfrak{r}) = \{\eta \in B(S, X) : (\mathfrak{r} \times \eta)(S) \text{ has compact closure in } D\}.$$

Here $(\mathfrak{r} \times \eta)(s) = (\mathfrak{r}(s), \eta(s))$.

This topology is metrizable. Take any metric d on X that induces its topology; then the metric

$$(2.2) \quad d_S(\mathfrak{r}, \eta) = \sup\{d(\mathfrak{r}(s), \eta(s)) : s \in S\}$$

induces the uniform topology of $B(S, X)$. (The sup in (2.2) is finite, because $\mathfrak{r}(S), \eta(S)$ are contained in some compact $K \subset X$ and d is continuous on $K \times K$.) The topology restricts to the compact–open topology on the subspace $C(S, X)$ of continuous maps.

The definition of a manifold structure on $B(S, X)$ involves the induced Borel vector bundles

$$\mathfrak{r}^*TX = \coprod_{s \in S} T_{\mathfrak{r}(s)}X \rightarrow S, \quad \mathfrak{r} \in B(S, X),$$

see section 1, and their spaces $B(\mathfrak{r}^*TX)$ of bounded Borel sections. The material in section 1 allows to treat topological vector bundles only over compact bases as Borel vector bundles, and then pull them back by Borel maps. Here in general we have to take any compact $K \subset X$ containing $\mathfrak{r}(S)$; then $TX|_K$ determines a Borel vector bundle and can be pulled back by \mathfrak{r} to produce a Borel vector bundle, that we denoted \mathfrak{r}^*TX .

If $D \subset X \times X$, F is a function defined on D , and $x \in X$, we let

$$(2.3) \quad D^x = \{y \in X : (x, y) \in D\} \quad \text{and} \quad F^x = F(x, \cdot).$$

Lemma 2.1. *There are a neighborhood $D \subset X \times X$ of the diagonal and a C^∞ diffeomorphism F between D and a neighborhood of the zero section in TX with the following properties:*

- (a) F^x maps D^x biholomorphically on a convex open subset of T_xX ;
- (b) $F^x(x) \in T_xX$ is the zero vector;
- (c) Identifying the tangent spaces of the vector space T_xX with T_xX itself, the differential $F_*^x|_x : T_xD^x = T_xX \rightarrow T_xX$ is the identity.

Moreover, if X is relatively compact and open in another complex manifold M , then F can be chosen real analytic.

The inverse of F is thus as good a substitute for a holomorphic exponential map $TX \rightarrow X \times X$ as it gets. Denoting by $\text{pr}_2 : X \times X \rightarrow X$ projection on the second factor, accordingly we define

$$(2.4) \quad \text{exp} = \text{pr}_2 \circ F^{-1} : TX \rightarrow X,$$

whose restriction to T_xX is a biholomorphism $F^x(D^x) \rightarrow D^x$, inverse to F^x .

Proof. The first part is [LS07, Lemma 1.1]. The second part is a special case of [L04, Lemma 2.1], that (when $k = \omega$) takes a compact real analytic manifold V , possibly with boundary, a real analytic map $\mathfrak{r} : V \rightarrow M$, and relates neighborhoods of the graph of \mathfrak{r} in $V \times M$ and of the zero section in the induced bundle \mathfrak{r}^*TM .—Note, though, the different use of X here and there.—To apply that result, following Grauert [G58] embed our M in some \mathbb{R}^N as a closed real analytic submanifold, choose $V \subset M$ a suitable sublevel set of the Euclidean norm function, and \mathfrak{r} the embedding of V in M .

In fact, a minor modification of the proof in [L04] gives that F can be chosen real analytic for *any* X , whether M as above exists or not.

We are now in the position to define holomorphic charts that turn $B(S, X)$ into a complex manifold. Given D, F as in the lemma, a coordinate neighborhood will be $D(\mathfrak{r})$ of (2.1). The map

$$(2.5) \quad \varphi_{\mathfrak{r}} : D(\mathfrak{r}) \ni \mathfrak{z} \mapsto F \circ (\mathfrak{r} \times \mathfrak{z}) \in B(\mathfrak{r}^*TX)$$

is a homeomorphism on a neighborhood of the zero section in the Banach space $B(\mathfrak{r}^*TX)$. Since the inverse of $\varphi_{\mathfrak{r}}$ is given by $\xi \mapsto \exp \circ \xi$, to show that the local coordinates (2.5) are holomorphically related we need to check that for $\mathfrak{r}, \eta \in B(S, X)$ the map

$$(2.6) \quad \varphi_{\eta} \circ \varphi_{\mathfrak{r}}^{-1} : \xi \mapsto F \circ (\eta \times (\exp \circ \xi))$$

is holomorphic wherever defined on $B(\mathfrak{r}^*TX)$. This is a special case of the following result, that we will need later on as well.

Lemma 2.2. *Let Y be another finite dimensional complex manifold, $u \subset S \times X$ and $U \subset S \times TX$ open,*

$$\psi : u \rightarrow Y \quad \text{and} \quad \Psi : U \rightarrow TY$$

*continuous. Assume that Ψ holomorphically maps the fibers $U \cap (\{s\} \times T_x X)$ to $T_{\psi(s,x)}Y$. Fix $\mathfrak{r} \in B(S, X)$ such that the closure of $(id_S \times \mathfrak{r})(S)$ is in u , and let $\mathfrak{U} \subset B(\mathfrak{r}^*TX)$ consist of those sections ξ for which the closure of $(id_S \times \xi)(S)$ is contained in U . Then $\eta = \psi \circ (id_S \times \mathfrak{r}) \in B(S, Y)$; for $\xi \in \mathfrak{U}$*

$$(2.7) \quad \eta = \Psi \circ (id_S \times \xi)$$

*is a bounded Borel section of η^*TY ; and the map*

$$\Psi^\circ : \mathfrak{U} \ni \xi \mapsto \Psi \circ (id_S \times \xi) \in B(\eta^*TY)$$

is holomorphic.

It is of course possible that \mathfrak{U} is empty, in which case the last statement is, well, vacuous.—The holomorphy of the map (2.6) follows from Lemma 2.2 if we let $Y = X$, $\psi : S \times X \rightarrow X$ the projection, and $\Psi(s, v) = F(\pi v, \exp(v))$, independently of s .

Proof of Lemma 2.2. The lemma can be reduced to Lemma 1.3, but it is easier to write out a direct proof, quite similar to the proof of Lemma 1.3. That η, η are Borel follows because the composition of Borel functions is Borel; they are also bounded, because ψ, Ψ are bounded on the relatively compact sets $(\text{id}_S \times \mathfrak{r})(S)$, respectively, $(\text{id}_S \times \xi)(S)$. Next, the map Ψ° is continuous at any $\xi_0 \in \mathfrak{U}$. This follows from the uniform continuity of Ψ over a compact neighborhood $K \subset U$ of the closure of $(\text{id}_S \times \xi_0)(S)$.

To prove Ψ° is complex differentiable at ξ_0 , we use the same K and another compact neighborhood $L \subset \text{int } K$. Cauchy estimates then give that partial derivatives of Ψ along the fibers of TX are bounded on L . By Taylor's formula this implies that, denoting by $d_2\Psi$ the differential of Ψ along the manifolds $\{s\} \times T_x X$, the derivative of Ψ° at ξ_0 in the direction $\xi \in B(\mathfrak{r}^*TX)$

$$d\Psi^\circ(\xi_0; \xi) = \lim_{\mathbb{C} \ni t \rightarrow 0} (\Psi^\circ(\xi_0 + t\xi) - \Psi^\circ(\xi_0))/t$$

exists, and can be computed as

$$d\Psi^\circ(\xi_0; \xi)(s) = d_2\Psi(s, \xi_0(s); \xi(s)),$$

cf. (2.7). One now argues, as with Ψ° itself, that $d\Psi^\circ(\xi_0; \cdot) : B(\mathfrak{r}^*TX) \rightarrow B(\eta^*TY)$ is continuous. Hence Ψ° is complex differentiable at ξ_0 , and therefore holomorphic on \mathfrak{U} .

As we saw, one consequence of Lemma 2.2 is that the local coordinates $\varphi_{\mathfrak{r}}$, $\mathfrak{r} \in B(S, X)$, are holomorphically related, and endow $B(S, X)$ with the structure of a complex manifold, modeled on Banach spaces of form $B(\mathfrak{r}^*TX)$.

For example, if $S = \{s_1, \dots, s_m\}$ is finite, then

$$B(S, X) \ni \mathfrak{r} \mapsto (\mathfrak{r}(s_1), \dots, \mathfrak{r}(s_m)) \in X^m$$

is a biholomorphism. In general, the correspondence $(S, X) \mapsto B(S, X)$ is natural: a continuous map $a : T \rightarrow S$ and a holomorphic map $X \rightarrow Y$ induce, by composition, a holomorphic map $B(S, X) \rightarrow B(T, Y)$. More generally instead of a holomorphic map we can use a family of holomorphic maps $\Phi : S \times X \rightarrow Y$. We formulate the result only when $a : T = S \rightarrow S$ is the identity.

Lemma 2.3. *Let $\Phi : S \times X \rightarrow Y$ be continuous. If for each $s \in S$ the map $\Phi(s, \cdot)$ is holomorphic, then*

$$\Phi^\circ : B(S, X) \ni \mathfrak{r} \mapsto \Phi \circ (\text{id}_S \times \mathfrak{r}) \in B(S, Y)$$

is holomorphic.

Proof. Let $D' \subset Y \times Y$ and $F' : D' \rightarrow TY$ be as in Lemma 2.1, but with Y rather than X . If $\varphi_{\mathfrak{r}}$ is a local coordinate on $B(S, X)$ as in (2.5), $\eta = \Phi^\circ(\mathfrak{r})$, and φ'_η is a local coordinate on $B(S, Y)$ at η , we need to check that the map

$$\varphi'_\eta \circ \Phi^\circ \circ \varphi_{\mathfrak{r}}^{-1} : \xi \mapsto F' \circ (\eta \times \Phi \circ (\text{id}_S \times \exp \circ \xi)) \in B(\eta^*TY)$$

is holomorphic where defined on $B(\mathfrak{r}^*TX)$. But this follows from Lemma 2.2 if we choose $\Psi(s, v) = F'(\pi v, \Phi(s, \exp v))$.

The complex structure on $B(S, X)$ was defined in terms of $D \subset X \times X$ and $F : D \rightarrow TX$. Since Lemma 2.3 applies with any choice of D, F (and D', F'), it follows by taking $Y = X$ and $\Phi(s, x) = x$ that, in the end, different choices of D, F give rise to the same complex structure on $B(S, X)$.

Continuous maps $S \rightarrow X$ form a complex submanifold $C(S, X) \subset B(S, X)$, modeled on Banach spaces $C(\mathfrak{r}^*TX)$ of continuous sections, $\mathfrak{r} \in C(S, X)$. It is closed as a subset of $B(S, X)$, but not a direct submanifold: while locally the pair $C(S, X) \subset B(S, X)$ is biholomorphic to the pair $C(\mathfrak{r}^*TX) \subset B(\mathfrak{r}^*TX)$, the space $C(\mathfrak{r}^*TX)$ will not in general have a closed complement in $B(\mathfrak{r}^*TX)$. It is this circumstance that makes even the local holomorphic extension problem from $C(S, X)$ to $B(S, X)$ difficult (but solved by the Aron–Berner extension theorem).

An $\mathfrak{r}_0 \in C(S, X)$ and a closed $A \subset S$ determine spaces of based maps

$$\mathfrak{B} = B(S, X, A, \mathfrak{r}_0) \subset B(S, X) \quad \text{and} \quad \mathfrak{C} = C(S, X, A, \mathfrak{r}_0) \subset C(S, X) \cap \mathfrak{B}.$$

$\mathfrak{B} = \{\mathfrak{r} \in B(S, X) : \mathfrak{r}|_A = \mathfrak{r}_0|_A\}$, and \mathfrak{C} denotes the connected component of \mathfrak{r}_0 in $\{\mathfrak{r} \in C(S, X) : \mathfrak{r}|_A = \mathfrak{r}_0|_A\}$. These are closed complex submanifolds of $B(S, X)$, resp. $C(S, X)$, modeled on subspaces

$$B_A(\mathfrak{r}^*TX) \subset B(\mathfrak{r}^*TX), \quad \text{respectively} \quad C_A(\mathfrak{r}^*TX) \subset C(\mathfrak{r}^*TX),$$

of sections that vanish on A . It is the manifolds \mathfrak{B} and $\mathfrak{C} \subset \mathfrak{B}$ that are the targets of our investigations.

3 Aron–Berner extension

Let $\mathfrak{X}, \mathfrak{Y}$ be Banach spaces, $x_0 \in \mathfrak{X}$, $r > 0$, and

$$B = \{x \in \mathfrak{X} : \|x - x_0\| < r\}, \quad B'' = \{\xi \in \mathfrak{X}'' : \|\xi - x_0\| < r\}.$$

In this section we review how to extend a bounded $f \in \mathcal{O}(B, \mathfrak{Y})$ to a bounded function $f^{\text{AB}} \in \mathcal{O}(B'', \mathfrak{Y}'')$, with the same bound.

The procedure starts with a(n always continuous) k -linear form $T : \mathfrak{X}^{\oplus k} \rightarrow \mathbb{C}$, $k = 0, 1, \dots$, and constructs its k -linear Aron–Berner extension $\bar{T} : \mathfrak{X}''^{\oplus k} \rightarrow \mathbb{C}$; the association $T \mapsto \bar{T}$ is linear. When $k = 0$, T is constant, and the extension is the same constant; when $k = 1$, $\bar{T} : \mathfrak{X}'' \rightarrow \mathbb{C}$ is the second transpose of T . In general, to define \bar{T} by recurrence, suppose \bar{T} exists for k -linear operators T ; then take a $(k + 1)$ -linear T . For any choice of $x_1, \dots, x_k \in \mathfrak{X}$ the second transpose of $T(x_1, \dots, x_k, \cdot) : \mathfrak{X} \rightarrow \mathbb{C}$ is a linear form $T''(x_1, \dots, x_k, \cdot) : \mathfrak{X}'' \rightarrow \mathbb{C}$. For each $\xi \in \mathfrak{X}''$

$$T_\xi = T''(\cdot, \dots, \cdot, \xi) : \mathfrak{X}^{\oplus k} \rightarrow \mathbb{C}$$

is k -linear, and depends linearly on ξ . The extensions $\bar{T}_\xi : \mathfrak{X}''^{\oplus k} \rightarrow \mathbb{C}$ then define

$$(3.1) \quad \bar{T}(\xi_1, \dots, \xi_{k+1}) = \bar{T}_{\xi_{k+1}}(\xi_1, \dots, \xi_k), \quad \xi_j \in \mathfrak{X}''.$$

More generally, if $T : \mathfrak{X}^{\oplus k} \rightarrow \mathfrak{Y}$ is a k -linear operator, for each $l \in \mathfrak{Y}'$ we consider the k -linear form $T_l = lT$. The extensions constructed above, for each $\xi_j \in \mathfrak{X}''$ produce

a linear form $l \mapsto \bar{T}_l(\xi_1, \dots, \xi_k)$ on \mathfrak{Y}' , i.e., an element $\bar{T}(\xi_1, \dots, \xi_k) \in \mathfrak{Y}''$. This defines the k -linear extension $\bar{T} : \mathfrak{X}''^{\oplus k} \rightarrow \mathfrak{Y}''$ of T . The standard cautionary remark here is that if the arguments of T are permuted to produce a k -linear operator T_1 , then \bar{T}_1 is not in general the same as \bar{T} with arguments correspondingly permuted.

The extension $\bar{T} : \mathfrak{X}'' \rightarrow \mathfrak{Y}''$ of a linear operator T is continuous when both $\mathfrak{X}'', \mathfrak{Y}''$ are endowed with the weak* topology. The extension of multilinear operators is continuous only in a weaker sense. Let us say that a function $F : \mathfrak{X}''^{\oplus k} \rightarrow \mathfrak{Y}''$ is slightly continuous to mean that for any $j = 1, \dots, k$, if $\xi_1, \dots, \xi_{j-1} \in \mathfrak{X}$ and $\xi_{j+1}, \dots, \xi_k \in \mathfrak{X}''$ are fixed, then $F(\xi_1, \dots, \xi_j, \dots, \xi_k) \in \mathfrak{Y}''$ depends continuously on $\xi_j \in \mathfrak{X}''$ in the weak* topology of $\mathfrak{X}'', \mathfrak{Y}''$. The following is not new:

Lemma 3.1. *The extension $\bar{T} : \mathfrak{X}''^{\oplus k} \rightarrow \mathfrak{Y}''$ of a k -linear $T : \mathfrak{X}^{\oplus k} \rightarrow \mathfrak{Y}$ is slightly continuous.*

Proof. It suffices to prove when $\mathfrak{Y} = \mathbb{C}$, that can be done by induction. When $k = 0$, \bar{T} is constant. When $k = 1$, \bar{T} is the second transpose of T , and the claimed continuity follows directly from the definitions. Suppose the lemma holds for a certain $k \geq 1$ and let T be $(k+1)$ -linear. To show that \bar{T} is slightly continuous, we have to check a certain continuity property for all $j = 1, \dots, k+1$. When $j \leq k$, this property follows from the induction hypothesis as $\bar{T}_{\xi_{k+1}}$ is slightly continuous, cf. (3.1). If $j = k+1$, and $\xi_i = x_i \in \mathfrak{X}$, $i \leq k$, then again $\bar{T}(x_1, \dots, x_k, \xi_{k+1}) = T_{\xi_{k+1}}(x_1, \dots, x_k) = T''(x_1, \dots, x_k, \xi_{k+1})$ depends continuously on $\xi_{k+1} \in \mathfrak{X}''$.

Since \mathfrak{X} is weak* dense in \mathfrak{X}'' (Goldstine's theorem, [DS88, p. 424]), it follows that in the weak* topologies of $\mathfrak{X}'', \mathfrak{Y}''$

$$(3.2) \quad \bar{T}(\xi_1, \dots, \xi_k) = \lim_{x_1 \rightarrow \xi_1} \lim_{x_2 \rightarrow \xi_2} \dots \lim_{x_k \rightarrow \xi_k} T(x_1, \dots, x_k), \quad \xi_1, \dots, \xi_k \in \mathfrak{X}'',$$

which is often taken as the definition of \bar{T} , e.g., in [AB78].

We return to the construction of the Aron–Bernstein extension of f . The construction is compatible with translations; for this reason we assume that the center of the balls B, B'' is $x_0 = 0$. Any homogeneous polynomial $P : \mathfrak{X} \rightarrow \mathfrak{Y}$ can be represented as $P(x) = T(x, \dots, x)$ with a unique symmetric multilinear $T : \mathfrak{X}^{\oplus k} \rightarrow \mathfrak{Y}$; we define $P^{\text{AB}}(\xi) = \bar{T}(\xi, \dots, \xi) \in \mathfrak{Y}''$, a homogeneous polynomial on \mathfrak{X}'' . The correspondence $P \mapsto P^{\text{AB}}$ can be extended by linearity to the space of all polynomials $P : \mathfrak{X} \rightarrow \mathfrak{Y}$. We need the fact that, using $\| \cdot \|$ for norm on \mathfrak{Y} and on its iterated dual spaces,

$$(3.3) \quad \sup_{B''} \|P^{\text{AB}}\| = \sup_B \|P\|.$$

This was proved by Davie and Gamelin [DG89] when $\mathfrak{Y} = \mathbb{C}$. Carando in [C99] observed that (3.3) holds for homogeneous polynomials; that it is true in general easily follows from [DG89]. Indeed, assume that $\sup_B \|P\| = 1$. If $l \in \mathfrak{Y}'$ has norm ≤ 1 , then $P_l = lP : \mathfrak{X} \rightarrow \mathbb{C}$ is a polynomial, $\sup_B |P_l| \leq 1$, and $P_l^{\text{AB}} = lP^{\text{AB}}$. In the latter formula l acts on \mathfrak{Y}'' by transposition. By [DG89], then

$$(3.4) \quad \sup_{B''} |lP^{\text{AB}}| \leq 1 \quad \text{for } l \in \mathfrak{Y}', \quad \|l\| \leq 1.$$

But by Goldstine's theorem [DS88, p. 424] the unit ball of \mathfrak{Y}' is weak* dense in the unit ball of \mathfrak{Y}''' , so that (3.4) holds for $l \in \mathfrak{Y}'''$. Hence $\sup_{B''} \|P^{AB}\| \leq 1$. The opposite inequality being obvious, (3.3) holds.

The following result, a slight improvement on [AB78, DG89], is known in the field:

Theorem 3.2. *Let $f \in \mathcal{O}(B, \mathfrak{Y})$ be bounded, $f = \sum_{k=0}^{\infty} P_k$ its expansion in homogeneous polynomials, $\deg P_k = k$. Then the series*

$$f^{AB} = \sum_{k=0}^{\infty} P_k^{AB}$$

converges uniformly on concentric balls $B''_{\rho} \subset B''$ of radius $\rho < r$; it represents the Aron–Bernstein extension $f^{AB} \in \mathcal{O}(B'', \mathfrak{Y}'')$ of f . In addition,

$$(3.5) \quad \sup_{B''} \|f^{AB}\| = \sup_B \|f\|.$$

Proof. P_k is given by

$$P_k(x) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{it}x) e^{-ikt} dt.$$

This and (3.3) imply

$$\sup_{B''} \|P_k^{AB}\| = \sup_B \|P_k\| \leq \sup_B \|f\|.$$

This in turn implies, in view of the homogeneity of P_k^{AB} , that $\sum_k P_k^{AB}$ converges uniformly on $B''_{\rho} \subset B''$, $\rho < r$, hence represents a holomorphic function $f^{AB} : B'' \rightarrow \mathfrak{Y}''$. The estimate (3.3), with P a partial sum, yields

$$\sup_{B''_{\rho}} \left\| \sum_{k=0}^m P_k^{AB} \right\| \leq \sup_B \left\| \sum_{k=0}^m P_k \right\|.$$

Letting first $m \rightarrow \infty$ then $\rho \rightarrow r$, (3.5) follows.

Of course, f^{AB} is not the unique holomorphic extension of f to B'' . But it is natural in a sense. To formulate a theorem, we need the notion of (Arens) regularity:

Definition 3.3. *A Banach space \mathfrak{Y} is regular if for any Banach space \mathfrak{Z} (or, equivalently, only for $\mathfrak{Z} = \mathbb{C}$), the extensions of any bilinear operator $T : \mathfrak{Y} \oplus \mathfrak{Y} \rightarrow \mathfrak{Z}$ and of $T_1(x, y) = T(y, x)$ satisfy $\bar{T}_1(\xi, \eta) = \bar{T}(\eta, \xi)$ for all $\xi, \eta \in \mathfrak{Y}$.*

Induction then shows that for any k -linear $T : \mathfrak{Y}^{\oplus k} \rightarrow \mathfrak{Z}$ and any permutation σ of $\{1, \dots, k\}$, the operator $T_1(x_1, \dots, x_k) = T(x_{\sigma(1)}, \dots, x_{\sigma(k)})$ satisfies $\bar{T}_1(\xi_1, \dots, \xi_k) = \bar{T}(\xi_{\sigma(1)}, \dots, \xi_{\sigma(k)})$. Since $\bar{T}(\cdot, \xi_2, \dots, \xi_k)$ is always continuous in the weak* topologies, in regular spaces \bar{T} is separately continuous in all its variables. Ülger shows in [U87] that any C* algebra is regular; later on we will need a minor generalization of this.

Theorem 3.4. *In addition to assumptions and notation above, let $y_0 \in \mathfrak{Y}$, $s > 0$, and $B_{\mathfrak{Y}} = \{y \in \mathfrak{Y} : \|y - y_0\| < s\}$. Suppose \mathfrak{Z} is a third Banach space, $f : B \rightarrow B_{\mathfrak{Y}}$ is holomorphic, $f(x_0) = y_0$, and $g : B_{\mathfrak{Y}} \rightarrow \mathfrak{Z}$ is bounded and holomorphic. Then $(g \circ f)^{AB} = g^{AB} \circ f^{AB}$, provided g is a first degree polynomial, or \mathfrak{Y} is regular.*

Proof. We will prove under the assumption that \mathfrak{Y} is regular. The other case follows by similar, but much simplified calculations. By translations we can arrange that $x_0 = 0$ and $f(x_0) = y_0 = 0$. First we assume that f and g are polynomials, g homogeneous of degree k . This means that there are symmetric multilinear operators $T : \mathfrak{Y}^{\oplus k} \rightarrow \mathfrak{Z}$ and $U_i : \mathfrak{X}^{\oplus i} \rightarrow \mathfrak{Y}$, $i = 0, 1, \dots, h$, such that

$$(3.6) \quad \begin{aligned} g(y) &= T(y, \dots, y), & f(x) &= \sum_{i=0}^h U_i(x, \dots, x), & \text{and therefore} \\ (g \circ f)(x) &= T(f(x), \dots, f(x)) = \sum T(U_{j_1}(x, \dots, x), U_{j_2}(x, \dots, x), \dots), \end{aligned}$$

summation over $j_1, \dots, j_k \leq h$. The function (3.6) is a polynomial in x . Its j 'th degree component is obtained from the symmetrization of the j -linear operator

$$S_j(x_1, \dots, x_j) = \sum_{j_1 + \dots + j_k = j} T(U_{j_1}(x_1, \dots, x_{j_1}), U_{j_2}(x_{j_1+1}, \dots, x_{j_1+j_2}), \dots)$$

by setting $x = x_1 = \dots = x_j$. The symmetrization $S_{j, \text{sym}}(x_1, \dots, x_j)$ is

$$(3.7) \quad \frac{1}{j!} \sum_{\sigma} \sum_{j_1 + \dots + j_k = j} T(U_{j_1}(x_{\sigma(1)}, \dots, x_{\sigma(j_1)}), U_{j_2}(x_{\sigma(j_1+1)}, \dots, x_{\sigma(j_1+j_2)}), \dots),$$

the first sum over all permutations σ of $\{1, \dots, j\}$. Now

$$(3.8) \quad \begin{aligned} g^{\text{AB}}(\eta) &= \bar{T}(\eta, \dots, \eta), & f^{\text{AB}}(\xi) &= \sum_i \bar{U}_i(\xi, \dots, \xi), & \text{and by (3.2)} \\ (g \circ f)^{\text{AB}}(\xi) &= \sum_j \lim_{x_1 \rightarrow \xi} \lim_{x_2 \rightarrow \xi} \dots \lim_{x_j \rightarrow \xi} S_{j, \text{sym}}(x_1, \dots, x_j). \end{aligned}$$

We compute this iterated limit for the generic term in (3.7). Since each U_i is symmetric, we may restrict attention to σ such that $\sigma(1) < \sigma(2) < \dots < \sigma(j_1)$, $\sigma(j_1 + 1) < \dots < \sigma(j_1 + j_2)$, and so on (shuffles). Then the arguments of T in this generic term tend to $\bar{U}_{j_1}(\xi, \dots, \xi)$, $\bar{U}_{j_2}(\xi, \dots, \xi)$, etc., and since T is separately continuous, the limit of the generic term is $\bar{T}(\bar{U}_{j_1}(\xi, \dots, \xi), \bar{U}_{j_2}(\xi, \dots, \xi), \dots)$. In light of (3.8) and (3.7) therefore the limit in (3.8) is indeed $\bar{T}(\sum_0^h \bar{U}_i(\xi, \dots, \xi), \dots, \sum_0^h \bar{U}_i(\xi, \dots, \xi))$.

Next we let f be arbitrary, but keep g k -homogeneous. We expand f in a homogeneous series. The partial sums f_j of the series, polynomials, tend to f as $j \rightarrow \infty$, uniformly on concentric balls $\subset B$ of radius $< r$. By (3.5) and by what we have proved,

$$(g \circ f)^{\text{AB}} = \lim_{j \rightarrow \infty} (g \circ f_j)^{\text{AB}} = \lim_{j \rightarrow \infty} g^{\text{AB}} \circ f_j^{\text{AB}} = g^{\text{AB}} \circ f^{\text{AB}}.$$

Finally, a general bounded holomorphic $g : B_{\mathfrak{Y}} \rightarrow \mathfrak{Z}$ can be expanded in a homogeneous series $g = \sum_k g_k$, converging uniformly on concentric balls $\subset B_{\mathfrak{Y}}$ of radius $< s$. Again using (3.5), on some neighborhood of $0 \in B''$

$$(g \circ f)^{\text{AB}} = \lim_{k \rightarrow \infty} \left(\sum_{j=0}^k g_j \circ f \right)^{\text{AB}} = \lim_{k \rightarrow \infty} \sum_{j=0}^k g_j^{\text{AB}} \circ f^{\text{AB}} = g^{\text{AB}} \circ f^{\text{AB}},$$

and $(g \circ f)^{\text{AB}} = g^{\text{AB}} \circ f^{\text{AB}}$ follows by analytic continuation.

4 Extensions from certain $C(S)$ –modules

In this section we restrict ourselves to certain Banach spaces \mathfrak{X} that are modules over $C(S)$. These modules arise from a topological complex vector bundle $V \rightarrow S$, of finite rank, over the compact space S ; they are the space $C(V)$ of continuous sections of V and certain subspaces. We embed the space $B(V)$ of bounded Borel sections into the bidual of $C(V)$, and study the restriction f^{ab} to $B(V)$ (and subspaces) of the Aron–Berner extension f^{AB} from $C(V)$ (and subspaces). The extension $f \mapsto f^{\text{ab}}$ has a certain naturality property. Since the mapping spaces $\mathfrak{B}, \mathfrak{C}$ are modeled on modules studied here, the results of this section directly apply to provide local Aron–Berner–type extensions in those mapping spaces.

By a norm on V we mean a continuous function $|\cdot| : V \rightarrow [0, \infty)$ that restricts to a vector space norm on each fiber $V_s, s \in S$. Given a closed $A \subset S$ and a norm on V , we consider the Banach spaces

$$B_A(V) = \{\text{bounded Borel sections } \xi \text{ of } V : \xi|_A = 0\}, \quad \|\xi\| = \sup_S |\xi|,$$

and $C_A(V) \subset B_A(V)$ the subspace of continuous sections. Multiplication by functions turns these spaces into modules over the Banach algebra $C(S)$. Thus $C_\emptyset(V) = C(V)$, $B_\emptyset(V) = B(V)$.

The main point will be the construction of a canonical embedding $B_A(V) \rightarrow C_A(V)''$. We start by describing the dual $C_A(V)'$ in terms of measures. Let $M(S \setminus A)$ denote the space of complex valued regular Borel measures on $S \setminus A$. This is also a $C(S)$ –module: the product fm of $f \in C(S)$ with a measure m is given by

$$(fm)(T) = \int_T f dm, \quad T \subset S \setminus A \text{ Borel};$$

thus $d(fm)/dm = f$. Let $C_0(S \setminus A)$ be the space of continuous complex functions on the locally compact space $S \setminus A$ that vanish at infinity, and $C_A(S) \subset C(S)$ the ideal of functions that vanish on A . The map

$$C_A(S) \ni f \mapsto f|(S \setminus A) \in C_0(S \setminus A)$$

is an isometric isomorphism. Let furthermore $V^* \rightarrow S$ be the bundle dual to V , endowed with the dual norm $|\cdot|^*$, and $\langle \cdot, \cdot \rangle$ the duality pairing between V^* and V . The tensor product

$$C(V^*) \otimes_{C(S)} M(S \setminus A)$$

is the $C(S)$ –module freely generated by $C(V^*) \times M(S \setminus A)$, modulo the the submodule generated by

$$(4.1) \quad (fg_1 + g_2, m) - f(g_1, m) - (g_2, m), \quad (g, fm_1 + m_2) - f(g, m_1) - (g, m_2),$$

$f \in C(S), g_1, g_2, g \in C(V^*), m_1, m_2, m \in M(S \setminus A)$. Write $g \otimes m$ for the class of (g, m) .

Lemma 4.1. *Associating with $t = \sum_1^k g_j \otimes m_j \in C(V^*) \otimes_{C(S)} M(S \setminus A)$ the linear form*

$$(4.2) \quad C_A(V) \ni \eta \mapsto \int_{S \setminus A} \sum_{j=1}^k \langle g_j, \eta \rangle dm_j \in \mathbb{C}$$

defines an isomorphism

$$(4.3) \quad C(V^*) \otimes_{C(S)} M(S \setminus A) \xrightarrow{\approx} C_A(V)'$$

of $C(S)$ -modules.

Proof. To see that different representations of t produce the same linear form (4.2) we have to check that (4.2) gives zero when the representations correspond to the vectors in (4.1); and this is obvious. Thus (4.2) indeed defines a homomorphism (4.3).

Next we verify that (4.3) is surjective. Let $l \in C_A(V)'$. Suppose $U \subset S$ is open and V is trivial over a neighborhood of \bar{U} . Fix $\xi_1, \dots, \xi_r \in C(V)$ and $g_1, \dots, g_r \in C(V^*)$ that form dual frames of V, V^* over U . If $\chi \in C(S)$ is supported in U , Riesz's theorem, generalized [R87], implies that the linear forms

$$C_0(S \setminus A) \approx C_A(S) \ni f \mapsto l(\chi f \xi_i) \in \mathbb{C}$$

can be represented as $l(\chi f \xi_i) = \int_{S \setminus A} f dm_i$ with $m_i \in M(S \setminus A)$. Hence

$$l(\chi \eta) = \sum_{i=1}^r l(\chi \langle g_i, \eta \rangle \xi_i) = \sum_{i=1}^r \int_{S \setminus A} \langle g_i, \eta \rangle dm_i, \quad \eta \in C_A(V).$$

If \mathfrak{U} is a finite cover of S consisting of U as above, $\chi = \chi_U$ form a subordinate partition of unity, and we choose $r = r^U$, $\xi_i = \xi_i^U$, $g_i = g_i^U$, $m_i = m_i^U$ accordingly, then

$$l(\eta) = \sum_{U \in \mathfrak{U}} l(\chi_U \eta) = \int_{S \setminus A} \sum_{U \in \mathfrak{U}} \sum_{i=1}^{r^U} \langle g_i^U, \eta \rangle dm_i^U, \quad \eta \in C_A(V).$$

Thus (4.3) sends $\sum_U \sum_i g_i^U \otimes m_i^U$ to l , and is therefore surjective.

Finally, suppose that $t = \sum h_j \otimes m_j$ is in the kernel of (4.3). Assume first that each h_j is supported in an open U , and there are $\xi_1, \dots, \xi_r \in C(V)$, $g_1, \dots, g_r \in C(V^*)$ that form dual frames over U . Then each h_j is in the submodule generated by the g_i , and t can be written as $\sum_i g_i \otimes n_i$, with $n_i \in M(S \setminus A)$ supported in U . Thus for any $f_1, \dots, f_r \in C_A(S)$ and $\eta = \sum_i f_i \xi_i \in C_A(V)$

$$\sum_i \int_{S \setminus A} f_i dn_i = \int_{S \setminus A} \sum_i \langle g_i, \eta \rangle dn_i = 0.$$

Hence each $n_i = 0$ and $t = 0$. Now with a general $t = \sum_j h_j \otimes m_j$ in the kernel of (4.3) and with a sufficiently fine partition of unity χ_U , $U \in \mathfrak{U}$, we have $t = \sum_U \chi_U t = 0$ by what we have already proved, since $\chi_U t$ is also in the kernel.

Next we define a bilinear pairing $P : B_A(V) \times C_A(V)' \rightarrow \mathbb{C}$. If $\eta \in B_A(V)$ and $l \in C_A(V)'$ corresponds, under (4.3), to $t = \sum_j g_j \otimes m_j \in C(V^*) \otimes_{C(S)} M(S \setminus A)$, set

$$(4.4) \quad P(\eta, l) = \int_{S \setminus A} \sum_j \langle g_j, \eta \rangle dm_j.$$

Again, different representations of t give the same integral in (4.4).

Lemma 4.2. $B_A(V) \ni \eta \mapsto P(\eta, \cdot) \in C_A(V)''$ is an isometric embedding, an extension of the canonical embedding $C_A(V) \rightarrow C_A(V)''$.

Proof. The statement about the extension is immediate from the definition of the canonical embedding; which embedding is isometric. Next, fix $\eta \in B_A(V)$ and $\sum_j g_j \otimes m_j \in C(V^*) \otimes_{C(S)} M(S \setminus A)$. Given $\varepsilon > 0$, it follows from Lusin's theorem that there is a $\xi \in C_A(V)$ such that

$$\sum_j \sup_S |g_j|^* \int_{S \setminus A} |\xi - \eta| d|m_j| < \varepsilon \quad \text{and} \quad \sup_S |\xi| \leq \sup |\eta|.$$

Hence $|P(\xi, l) - P(\eta, l)| = \left| \int_{S \setminus A} \sum_j \langle g_j, \xi - \eta \rangle dm_j \right| < \varepsilon$. Since $\|P(\xi, \cdot)\| = \sup_S |\xi|$, we obtain $\|P(\eta, \cdot)\| \leq \sup_S |\eta|$.

To prove the opposite inequality, let $s \in S \setminus A$ and δ_s the Dirac measure at s . Choose a linear $\lambda : V_s \rightarrow \mathbb{C}$ such that

$$|\lambda(v)| \leq |v| \quad \text{for all } v \in V_s, \quad \text{with equality if } v = \eta(s),$$

and define $l(\xi) = \lambda(\xi(s))$, $\xi \in C_A(V)$. Thus $l \in C(V)'$ corresponds under (4.3) to $g \otimes \delta_s$ with an arbitrary $g \in C_A(V^*)$ such that $g(s) = \lambda$. Hence $|P(\eta, l)| = |\eta(s)|$. Since $\|l\| \leq 1$, we have $\|P(\eta, \cdot)\| \geq |\eta(s)|$, and this being so for any $s \in S \setminus A$, in fact $\|P(\eta, \cdot)\| \geq \sup_S |\eta|$. This completes the proof.

Identifying $B_A(V)$ with its image in $C_A(V)''$ allows us to define a version of the Aron–Bernier extension from $C_A(V)$ to $B_A(V)$. If $\Gamma \subset B_A(V)$ is a ball with center in $C_A(V)$ and $f \in \mathcal{O}(\Gamma \cap C_A(V), \mathfrak{Y})$ is bounded, the extension f^{AB} discussed in section 3 can be restricted to Γ to produce a holomorphic

$$(4.5) \quad f^{\text{ab}} : \Gamma \rightarrow \mathfrak{Y}'',$$

with the same sup norm. Similarly, if $T : C_A(V)^{\oplus k} \rightarrow \mathfrak{Y}$ is k -linear, the restriction to $B_A(V)^{\oplus k}$ of its Aron–Bernier extension \bar{T} will be denoted T^{ab} .

Theorem 4.3. *In addition to $V \rightarrow S$, consider another topological complex vector bundle $W \rightarrow S$, $\xi_0 \in C_A(V)$, an open neighborhood $U \subset V$ of $\xi_0(S)$, and a ball $\Gamma \subset B_A(V)$ about ξ_0 such that $\xi \in \Gamma$ implies $\bar{\xi}(S) \subset U$. Suppose a continuous $H : U \rightarrow W$ maps fibers $U \cap V_s$ holomorphically into corresponding fibers W_s . Then the map*

$$H^\circ : \Gamma \ni \xi \mapsto H \circ \xi \in B(W)$$

is holomorphic.

If, furthermore, H° is bounded on $\Gamma \cap C_A(V)$, then

$$(4.6) \quad (H^\circ | \Gamma \cap C_A(V))^{\text{ab}} = H^\circ.$$

To prove Theorem 4.3 we need a lemma.

Lemma 4.4. *Suppose $J : V^{\oplus k} \rightarrow W$ is a continuous fiber map, fiberwise k -linear. Define a k -linear operator $J^\circ : B_A(V)^{\oplus k} \ni (\eta_1, \dots, \eta_k) \mapsto J \circ (\eta_1 \times \dots \times \eta_k) \in B_A(W)$. Then*

$$(J^\circ | C_A(V)^{\oplus k})^{\text{ab}} = J^\circ.$$

Proof. In this proof we will use the topologies on $B_A(V), B_A(W)$ inherited from the weak* topologies on $C_A(V)'', C_A(W)''$. By construction, these are induced by the pairings P in (4.4), respectively the corresponding $Q : B_A(W) \times C_A(W)' \rightarrow \mathbb{C}$. First we show that $J^\circ(\eta_1, \dots, \eta_k) \in B_A(W)$ depends continuously on each $\eta_j \in B_A(V)$. Let's prove for $j = 1$; fix η_2, \dots, η_k . We need to show that $Q(J^\circ(\eta_1, \dots, \eta_k), l)$, for any $l \in C_A(W)'$, depends continuously on η_1 . The map

$$K : V \ni v \mapsto J(v, \eta_2(s), \dots, \eta_k(s)) \in W, \quad v \in V_s, \quad s \in S,$$

is a bounded Borel homomorphism $V \rightarrow W$, with adjoint $K^* : W^* \rightarrow V^*$. If l corresponds to $\sum_i h_i \otimes m_i \in C(W^*) \otimes_{C(S)} M(S \setminus A)$, cf. (4.3), then

$$Q(J^\circ(\eta_1, \dots, \eta_k), l) = \int_{S \setminus A} \sum_i \langle h_i, K \circ \eta_1 \rangle dm_i = \int_{S \setminus A} \sum_i \langle K^* \circ h_i, \eta_1 \rangle dm_i.$$

The last expression is $l_1(\eta_1)$ with a certain $l_1 \in C_A(V)'$, and continuity follows.

In particular, J° is slightly continuous, as is $(J^\circ|C_A(V)^{\oplus k})^{\text{ab}}$ by Lemma 3.1. Since the two agree on $C_A(V)^{\oplus k}$, and $C_A(V)$ is dense in $B_A(V)$, they agree everywhere.

Proof of Theorem 4.3. By restricting to components of S , we can assume that V and W have equidimensional fibers. Then Lemma 1.3 implies that H° even has a holomorphic extension to a neighborhood of Γ in $B(V)$.

Next we turn to (4.6). By translations we can arrange that $\xi_0 = 0$ and $H^\circ(\xi_0) = 0$. In a neighborhood of the zero section of V expand H in a uniformly convergent series $H = \sum_{k=0}^{\infty} H_k$, with fiberwise k -homogeneous $H_k : V \rightarrow W$. Each H_k is induced by a symmetric $J : V^{\oplus k} \rightarrow W$ as in Lemma 4.4, $H_k(v) = J(v, \dots, v)$ for $v \in V$. By the lemma and by (3.5), in a neighborhood of $0 \in \Gamma$

$$(H^\circ| \Gamma \cap C_A(V))^{\text{ab}} = \sum_k (H_k^\circ| \Gamma \cap C_A(V))^{\text{ab}} = \sum_k H_k^\circ = H^\circ.$$

By analytic continuation therefore $(H^\circ| \Gamma \cap C_A(V))^{\text{ab}} = H^\circ$ on all of Γ .

We return to the notion of regularity, Definition 3.3.

Lemma 4.5. *The spaces $C_A(V)$ are regular.*

Proof. As a C^* algebra, $C_A(S)$ is regular by [U87, Theorem 2.8]. (That theorem is about regularity of an arbitrary bilinear operator $X \times X \rightarrow X$, when X is a C^* algebra. However, the justification given there applies to any bilinear operator on X with values in any Banach space.) It follows that $C_A(S)^{\oplus q}$ is also regular for any $q \in \mathbb{N}$, because if S_q is the disjoint union of q copies of S and $A_q \subset S_q$ is the disjoint union of q copies of A , then $C_A(S)^{\oplus q} \approx C_{A_q}(S_q)$. Now $C_A(V)$ is a quotient of $C_A(S)^{\oplus q}$ for some q . Indeed, if $\xi_1, \dots, \xi_q \in C(V)$ span each fiber V_s , then

$$C_A(S)^{\oplus q} \ni (f_1, \dots, f_q) \mapsto \sum_{i=1}^q f_i \xi_i \in C_A(V)$$

is surjective. Since regularity passes down to quotients according to Ülger, [U87, Corollary 2.4], $C_A(V)$ is indeed regular.

Putting this, Theorem 3.4, and Theorem 4.3 together we obtain:

Corollary 4.6. *Suppose that in Theorem 4.3 $H^\circ(\Gamma)$ is contained in a ball $\Delta \subset B_A(W)$, centered at $H^\circ(\xi_0)$. If $g \in \mathcal{O}(\Delta \cap C_A(W), \mathfrak{Y})$ is bounded, then*

$$(4.7) \quad (g \circ H^\circ)^{ab} = g^{ab} \circ H^\circ.$$

5 Extending germs in mapping spaces

The naturality property in Corollary 4.6 makes it possible to generalize the Aron–Berner extension from Banach spaces to Banach manifolds that are mapping spaces. Given $\mathfrak{r}_0 \in C(S, X)$ and a closed $A \subset X$, as in section 2, $\mathfrak{B} = \{\mathfrak{r} \in B(S, X) : \mathfrak{r}|_A = \mathfrak{r}_0|_A\}$, and \mathfrak{C} is the connected component of \mathfrak{r}_0 in $\{\mathfrak{r} \in C(S, X) : \mathfrak{r}|_A = \mathfrak{r}_0|_A\}$.

At this point we are extending germs of holomorphic functions only. Suppose $\mathfrak{r}_1 \in \mathfrak{C}$, \mathfrak{Y} is a Banach space, and $\mathbf{f} : (\mathfrak{C}, \mathfrak{r}_1) \rightarrow \mathfrak{Y}$ is a holomorphic germ. Let $D \subset X \times X$ and $F : D \rightarrow TX$ be as in Lemma 2.1. Suppose $\mathfrak{r} \in \mathfrak{C}$ is such that \mathfrak{r}_1 is contained in

$$(5.1) \quad D_A(\mathfrak{r}) = \{\mathfrak{z} \in \mathfrak{B} : (\mathfrak{r} \times \mathfrak{z})(S) \text{ has compact closure in } D\} \subset D(\mathfrak{r}),$$

cf. (2.1). Thus $D_A(\mathfrak{r})$ is a coordinate neighborhood in \mathfrak{B} , with local coordinate the restriction of $\varphi_{\mathfrak{r}}$ of (2.5),

$$(5.2) \quad \psi_{\mathfrak{r}} : D_A(\mathfrak{r}) \ni \mathfrak{z} \mapsto F \circ (\mathfrak{r} \times \mathfrak{z}) \subset B_A(\mathfrak{r}^*TX).$$

Say, \mathbf{f} is the germ of a bounded holomorphic function $f : V \rightarrow \mathfrak{Y}$. The pullback $f \circ \psi_{\mathfrak{r}}^{-1}$ is holomorphic in a neighborhood of $\psi_{\mathfrak{r}}(\mathfrak{r}_1) \in C_A(\mathfrak{r}^*TX)$; we can choose this neighborhood to be of form $\Gamma \cap C_A(\mathfrak{r}^*TX)$ with $\Gamma \subset B_A(\mathfrak{r}^*TX)$ a ball centered at $\psi_{\mathfrak{r}}(\mathfrak{r}_1)$. As we saw in section 4, $f \circ \psi_{\mathfrak{r}}^{-1}$ then has an Aron–Berner type extension to a holomorphic $(f \circ \psi_{\mathfrak{r}}^{-1})^{ab} : \Gamma \rightarrow \mathfrak{Y}''$. The germ of $(f \circ \psi_{\mathfrak{r}}^{-1})^{ab}$ at $\psi_{\mathfrak{r}}(\mathfrak{r}_1)$ is independent of which ball Γ is chosen.

Lemma 5.1. *The germ of $(f \circ \psi_{\mathfrak{r}}^{-1})^{ab} \circ \psi_{\mathfrak{r}}$ at \mathfrak{r}_1 is independent of the choice of \mathfrak{r} .*

Proof. Suppose $\mathfrak{r}_1 \in D_A(\eta)$ with some $\eta \in \mathfrak{C}$, and $g = f \circ \psi_{\eta}^{-1}$ is bounded and holomorphic on $\Delta \cap C_A(\eta^*TX)$, where $\Delta \subset B_A(\eta^*TX)$ is a ball about $\psi_{\eta}(\mathfrak{r}_1)$. The composition $h = \psi_{\eta} \circ \psi_{\mathfrak{r}}^{-1}$ is holomorphic in a neighborhood of $\psi_{\mathfrak{r}}(\mathfrak{r}_1) \in B_A(\mathfrak{r}^*TX)$; we can choose Γ above contained in this neighborhood, and so that $h(\Gamma) \subset \Delta$. (5.2) implies

$$h(\xi) = F \circ (\eta \times (\exp \circ \xi)), \quad \xi \in \Gamma,$$

see (2.4). Denote the bundle projection $\mathfrak{r}^*TX \rightarrow S$ by p , and for $v \in \mathfrak{r}^*TX$ in a neighborhood U of $\psi_{\mathfrak{r}}(\mathfrak{r}_1)(S)$ set

$$H(v) = F(\eta(pv), \exp(v)) \in \eta^*TX.$$

This H is continuous and fiberwise holomorphic, as in Theorem 4.3, and H° of that theorem agrees with h (on a sufficiently small Γ). By Corollary 4.6, on Γ

$$(f \circ \psi_{\eta}^{-1})^{ab} \circ \psi_{\eta} \circ \psi_{\mathfrak{r}}^{-1} = g^{ab} \circ H^\circ = (g \circ h)^{ab} = (f \circ \psi_{\mathfrak{r}}^{-1})^{ab},$$

and indeed $(f \circ \psi_{\eta}^{-1})^{ab} \circ \psi_{\eta} = (f \circ \psi_{\mathfrak{r}}^{-1})^{ab} \circ \psi_{\mathfrak{r}}$ in a neighborhood of \mathfrak{r}_1 .

Thus we can make the following definition:

Definition 5.2. *If $\mathbf{f} : (\mathfrak{C}, \mathfrak{r}_1) \rightarrow \mathfrak{Y}$ is a holomorphic germ, then $\mathbf{f}^{ab} : (\mathfrak{B}, \mathfrak{r}_1) \rightarrow \mathfrak{Y}''$ stands for its extension $(\mathbf{f} \circ \psi_{\mathfrak{r}}^{-1})^{ab} \circ \psi_{\mathfrak{r}}$. Here $\mathfrak{r} \in \mathfrak{C}$ is arbitrary as long as $\mathfrak{r}_1 \in D_A(\mathfrak{r})$, and $\psi_{\mathfrak{r}}$ is the local coordinate (5.2).*

6 Banach bundles and flat norms

In this section we introduce/review the notions in the title. The main result, Theorem 6.3, is that over a simply connected Banach manifold a holomorphic Banach bundle with a flat norm is trivial. Norms on vector bundles are also called Finsler metrics; we have already discussed them for finite rank bundles. We will not need the notion of general norms here, and will define only flat norms.

Definition 6.1. (a) *A holomorphic Banach bundle is a holomorphic map $\pi : E \rightarrow M$ of complex Banach manifolds, with each fiber $E_x = \pi^{-1}x$, $x \in M$, endowed with the structure of a complex vector space. It is required that for each $x \in M$ there be a neighborhood $U \subset M$, a complex Banach space $(W, \|\cdot\|)$, and a biholomorphism $g : \pi^{-1}U \rightarrow U \times W$ (local trivialization), that for all $y \in U$ maps E_y linearly to $\{y\} \times W$.*
 (b) *A flat norm on $E \rightarrow M$ is a function $p : E \rightarrow [0, \infty)$ such that the local trivializations g above can be chosen to satisfy*

$$p(e) = \|w\| \quad \text{if } e \in \pi^{-1}U, \quad g(e) = (y, w).$$

To describe the structure of flat normed holomorphic Banach bundles we need the following result. If W is a Banach space, $\text{End } W$ is the space of continuous linear operators $W \rightarrow W$, endowed with the operator norm, and $\text{GL}(W) \subset \text{End } W$ the set of invertibles.

Lemma 6.2. *Let U be a connected open subset of a complex Banach space Z , $(W, \|\cdot\|)$ another complex Banach space, and $F : U \rightarrow \text{GL}(W)$ holomorphic. If $\|F(\zeta)w\| = \|w\|$ for all $\zeta \in U$ and $w \in W$, then F is constant.*

Proof. It suffices to prove when $\dim Z = 1$, the general case then follows by restricting to one dimensional slices. So, let $Z = \mathbb{C}$. Denoting derivative by a dot, we will show that $\dot{F}(\zeta) = 0$ for arbitrary $\zeta \in U$. To simplify notation, we assume $\zeta = 0$, and denote $F(0) = P$, $\dot{F}(0) = 2Q$. Thus $F(\zeta) = P + 2Q\zeta + \dots$ for small $\zeta \in \mathbb{C}$. Let

$$K(\tau) = 1 + \cos 2\pi\tau = 1 + (e^{2\pi i\tau} + e^{-2\pi i\tau})/2 \geq 0, \quad \tau \in \mathbb{R},$$

be the first Fejér kernel. For $w \in W$ and small ζ

$$\|(P + \zeta Q)w\| = \left\| \int_0^1 K(\tau) F(e^{2\pi i\tau} \zeta) w \, d\tau \right\| \leq \int_0^1 K(\tau) \|F(e^{2\pi i\tau} \zeta) w\| \, d\tau = \|w\| = \|Pw\|.$$

Since the left hand side is a subharmonic function of ζ , the maximum principle implies $\|(P + \zeta Q)w\| = \|w\|$, when $|\zeta| < r$ with some $r > 0$.

Denote the norm on the dual space W' also by $\|\cdot\|$, and the transpose of P, Q by $P', Q' \in \text{End } W'$; thus P' is an isometry. Furthermore

$$(6.1) \quad \|(P' + \zeta Q')u\| = \|u\|, \quad |\zeta| < r, \quad u \in W',$$

as well. Indeed, with $\langle \cdot, \cdot \rangle$ the pairing between W' and W ,

$$\|(P' + \zeta Q')u\| = \sup_{\|w\|=1} |\langle (P' + \zeta Q')u, w \rangle| = \sup_{\|w\|=1} |\langle u, (P + \zeta Q)w \rangle| \leq \|u\| = \|P'u\|.$$

Again, the left hand side is a subharmonic function of ζ , and (6.1) follows.

Suppose u is an extreme point of the closed unit ball $\Delta \subset W'$. Then $P'u \in \Delta$ is also an extreme point. Comparing this with (6.1), we conclude $Q'u = 0$. Since in the weak* topology Δ is compact, and by the Krein–Milman theorem it is the closed convex hull of its extreme points, $Q'v = 0$ follows for all $v \in \Delta$. Hence $\dot{F}(0) = 2Q = 0$.

Theorem 6.3. *Suppose $\pi : E \rightarrow M$ is a holomorphic Banach bundle over a connected, simply connected base, and p is a flat norm on E . Then (E, p) is globally trivial: there is a Banach space $(W, \|\cdot\|)$ and a biholomorphism $g : E \rightarrow M \times W$ that maps each E_x linearly to $\{x\} \times W$, and satisfies*

$$p(e) = \|w\| \quad \text{if } e \in E, \quad g(e) = (x, w).$$

This is a standard consequence of Lemma 6.2. The proof depends on the following notion:

Definition 6.4. *Suppose $U \subset M$ is connected open and $E|U$ admits a trivialization $g : E|U \rightarrow U \times W$ as in Definition 6.1b. A section h of $E|U$ is called horizontal if $g \circ h : U \rightarrow U \times W$ is a constant section. (Lemma 6.2 implies that horizontality of h is independent of the choice of the local trivialization g .) A section h over a general open subset of M is called horizontal if it is locally such.*

Proof of Theorem 6.3. Define a Hausdorff topology on E by declaring that a neighborhood basis of $e \in E$ consists of sets of form $h(U)$, where $U \subset M$ is a neighborhood of $\pi(e)$ and h is a horizontal section of $E|U$. In this topology $\pi : E \rightarrow M$ is a covering, which must be trivial since M is connected and simply connected. This means that through every $e \in E$ there passes a unique global horizontal section h_e . Fix an $a \in M$ and let $(W, \|\cdot\|) = (E_a, p|E_a)$. This is a Banach space, and

$$g : E \ni e \mapsto (\pi(e), h_e(a)) \in M \times W$$

is the global trivialization sought.

If $E \rightarrow M$ is a holomorphic Banach bundle with a flat norm p over a connected base, the biduals of the fibers E_x constitute a holomorphic Banach bundle $E'' \rightarrow M$, on which the biduals of the norms $p|E_x$ define a flat norm p'' . Local trivializations g of (E, p) as in Definition 6.1b give rise to the corresponding local trivializations g'' of (E'', p'') : $g''(y)$ is just the second transpose of $g(y)$, $y \in U$.

These considerations lead to the Aron–Berner extension of germs of holomorphic sections of holomorphic Banach bundles. Suppose that \mathfrak{X} is a Banach space, $B \subset$

\mathfrak{X} is a ball centered at $x \in \mathfrak{X}$, $B'' \subset \mathfrak{X}''$ is the bidual ball, and $(E, p) \rightarrow B''$ is a holomorphic Banach bundle with a flat norm. Under the trivialization g of Theorem 6.3, a bounded holomorphic section f of $E|B$ induces a bounded holomorphic section $g \circ f$ of $B \times W \rightarrow B$, i.e., a bounded $\phi \in \mathcal{O}(B, W)$. The corresponding trivialization g'' of (E'', p'') and the Aron–Berner extension $\phi^{\text{AB}} \in \mathcal{O}(B'', W'')$, in turn, induce a bounded holomorphic section f^{AB} of (E'', p'') . Lemma 6.2 implies that any another norm respecting trivialization $E \rightarrow B'' \times W$ is of form $g_1 = (\text{id}_{B''} \times T) \circ g$, with $T \in \text{End } W$. This means that the corresponding trivialization of E'' is $g_1'' = (\text{id}_{B''} \times T'') \circ g''$. Since the corresponding $\phi_1 = T \circ \phi$, by Theorem 3.4 $\phi_1^{\text{AB}} = T'' \circ \phi^{\text{AB}}$; hence f^{AB} is independent of the choice of g .

More generally, given a flat holomorphic Banach bundle (E, p) over some open set in \mathfrak{X}'' , and a germ \mathbf{f} of a local holomorphic section of $E|\mathfrak{X}$ at some $x \in \mathfrak{X}$, the same prescription defines the extension \mathbf{f}^{AB} , a germ of a section of E'' . One can even consider bundles over more general manifolds. Combining the above construction with the discussion of mapping spaces $\mathfrak{C} \subset \mathfrak{B}$ in section 5, given a flat holomorphic Banach bundle $(E, p) \rightarrow \mathfrak{B}$, with any $E|\mathfrak{C}$ valued holomorphic germ \mathbf{f} at some $\mathfrak{r} \in \mathfrak{C}$ we can associate its Aron–Berner–type extension \mathbf{f}^{ab} , a holomorphic germ at \mathfrak{r} , valued in E'' .

7 Analytic continuation and extension

Consider a holomorphic Banach bundle $E \rightarrow M$ over a connected base, and let $\mathcal{E} \rightarrow M$ denote the sheaf of E valued holomorphic germs. Suppose $a < b$ are real numbers and $r : [a, b] \rightarrow M$ a path (that is, a continuous map).

Definition 7.1. *Given a germ $\mathbf{f} \in \mathcal{E}$ at $r(a)$, its analytic continuation along the path r is a continuous lift $[a, b] \ni t \mapsto \mathbf{f}_t \in \mathcal{E}_{r(t)}$ of r such that $\mathbf{f}_a = \mathbf{f}$.*

Recall that continuity at $t \in [a, b]$ means that for all $\tau \in [a, b]$ sufficiently close to t the \mathbf{f}_τ are germs of the same local section of E about $r(t)$. For this reason analytic continuation can be defined by a partition $a = t_0 < t_1 < \dots < t_k = b$, open neighborhoods $U_j \subset M$ of $r[t_{j-1}, t_j]$, $j = 1, \dots, k$, and holomorphic sections g_j of $E|U_j$ such that $g_j = g_{j+1}$ on a neighborhood of $r(t_j)$, and \mathbf{f} is the germ of g_1 at $r(a)$.

This shows that analytic continuation along r is unique; and that if the path is slightly perturbed, keeping its endpoints fixed, then the continuation of \mathbf{f} at $r(b)$ is not going to change. It follows that if a germ $\mathbf{f} \in \mathcal{E}_x$ admits analytic continuation along every path in M starting at x , then the analytic continuation at $y \in M$ depends only on the homotopy class of the path connecting x and y along which we continue. In particular, if M is also simply connected, then the analytic continuations define a holomorphic section f of E : $f(y) \in E_y$ will be the value at y of the analytic continuation from x to y .

Later on we will need the following.

Lemma 7.2. *Suppose $P : M \rightarrow N$ is a holomorphic and open map of complex Banach manifolds, $F \rightarrow N$ is a holomorphic Banach bundle, $[a, b] \ni t \mapsto x_t \in M$ is a path, and \mathbf{g}_t are F valued holomorphic germs at $P(x_t) \in N$. Let \mathbf{k}_t be the pullback of \mathbf{g}_t along P to x_t , holomorphic P^*F valued germs. If \mathbf{k}_t depend continuously on t , then so do \mathbf{g}_t .*

Proof. To prove continuity at some $t \in [a, b]$, let $V \subset N$ be a neighborhood of $P(x_t)$ and g a holomorphic section of $F|V$ whose germ at $P(x_t)$ is \mathbf{g}_t . Thus $g \circ P$ is a holomorphic section of $P^*F|P^{-1}V$ whose germ at x_t is \mathbf{k}_t . For $\tau \in [a, b]$ close to t , \mathbf{k}_τ is the germ of $g \circ P$ at x_τ . As P is open, this implies that the germ of g at $P(x_\tau)$ is \mathbf{g}_τ ; which shows that $t \mapsto g_t$ is indeed continuous.

In what follows, when we speak of analytic continuation of a germ $\mathbf{f} \in \mathcal{E}_x$ along an arbitrary path, we of course mean paths that start at x . Let p be a flat norm on E .

Definition 7.3. *We say that a germ $\mathbf{f} \in \mathcal{E}_x$ admits bounded analytic continuation along every path if first, it admits analytic continuations $t \mapsto \mathbf{f}_t \in \mathcal{E}_{r(t)}$ along every path r and second, there is a bound C such that for all such continuations $p(\mathbf{f}_t(r(t))) \leq C$.*

If M is simply connected, bounded analytic continuation along every path simply means that \mathbf{f} extends to a holomorphic section f of E and $p(f)$ is bounded.

Let us return to mapping spaces of finite dimensional connected complex manifolds X . As earlier, given a compact Hausdorff space S , a closed $A \subset S$, and a continuous $\mathfrak{r}_0 : S \rightarrow X$, we consider the spaces

$$(7.1) \quad \mathfrak{B} = B(S, X, A, \mathfrak{r}_0) \supset \mathfrak{C} = C(S, X, A, \mathfrak{r}_0);$$

$\mathfrak{B} = \{\mathfrak{r} \in B(S, X) : \mathfrak{r}|_A = \mathfrak{r}_0|_A\}$ and \mathfrak{C} is the connected component of \mathfrak{r}_0 in $\mathfrak{B} \cap C(S, X)$. Let furthermore $E \rightarrow \mathfrak{B}$ be a holomorphic Banach bundle. In section 6 we associated with an $E|\mathfrak{C}$ valued holomorphic germ \mathbf{f} at some $\eta \in \mathfrak{C}$ its Aron–Berner type extension \mathbf{f}^{ab} , an E'' valued germ at η .

Lemma 7.4. *If $t \mapsto \mathbf{f}_t$ is the analytic continuation of \mathbf{f} along a path in \mathfrak{C} , then $t \mapsto \mathbf{f}_t^{\text{ab}}$ is the continuation of \mathbf{f}^{ab} along the same path.*

Proof. It suffices to prove when the path is contained in a coordinate neighborhood. Then we might as well assume that \mathfrak{C} is a ball in a Banach space \mathfrak{X} , $\mathfrak{B} \subset \mathfrak{X}''$ is a ball in a subspace, $E \rightarrow \mathfrak{B}$ is trivial, and so E and E'' valued germs \mathbf{f} correspond to germs \mathbf{g} valued in fixed Banach spaces $\mathfrak{Y}, \mathfrak{Y}''$. Now a special case of a result of Zalduendo, that concerns the Aron–Berner extension from a Banach space to its bidual, [Z90, Section 2, Definition 3, Theorem 1], implies that for \mathfrak{Y} valued germs \mathbf{g} the analytic continuation of \mathbf{g}^{ab} is given by \mathbf{g}_t^{ab} , and this proves the lemma.

The following theorem generalizes Theorem 0.1:

Theorem 7.5. *Suppose that a holomorphic Banach bundle $E \rightarrow \mathfrak{B}$ is endowed with a flat norm p (Definition 6.1). Assume that an $E|\mathfrak{C}$ valued holomorphic germ \mathbf{f} at some $\mathfrak{r} \in \mathfrak{C}$ admits bounded analytic continuation along any path in \mathfrak{C} . Then the E'' valued germ \mathbf{f}^{ab} admits analytic continuation along any path in \mathfrak{B} , with the same bound as the continuations within \mathfrak{C} .*

The proof will be given in section 9, after preliminary work in section 8.

8 The key construction

The key to the proof of Theorem 7.5 is a construction of a family of holomorphic maps of balls into X , with the help of which we can pull back analytic continuation along certain paths in \mathfrak{B} to continuation in a Banach space. Endow X with a metric d , then, as in section 2, metrize the space $B(S, X)$ by the metric

$$d_S(\mathfrak{r}, \mathfrak{h}) = \sup\{d(\mathfrak{r}(s), \mathfrak{h}(s)) : s \in S\}.$$

Lemma 8.1. *Given $\mathfrak{r} \in C(S, X)$ and $\varepsilon > 0$, there is a $\delta > 0$ with the following property. Suppose $S = \coprod_0^J S_j$ is a Borel partition, $S_0 = A$ is closed, $a < b$ are real numbers, and $[a, b] \ni t \mapsto \mathfrak{h}_t \in B(S, X)$ is a continuous path. Suppose furthermore that $\mathfrak{h}_t = \mathfrak{r}$ on S_0 and \mathfrak{h}_t is constant on S_j , $j = 1, \dots, J$, $t \in T$. If $d_S(\mathfrak{r}, \mathfrak{h}_a) < \delta$, then there are a (finite rank) normed topological complex vector bundle $\pi : (V, | \cdot |) \rightarrow S$ (cf. section 4) with unit ball bundle $U = \{v \in V : |v| < 1\} \rightarrow S$, a $\mathfrak{v} \in B_A(V)$, $\sup_S |\mathfrak{v}| < 1$, and a continuous $\Psi : [a, b] \times U \rightarrow X \times \mathbb{C}$ such that for every $t \in [a, b]$ and $s \in S$*

- (a) Ψ maps the fibers $\{t\} \times U_s$ biholomorphically on open subsets of $X \times \mathbb{C}$;
- (b) the zero vector $0_s \in V_s$ satisfies $\Psi(t, 0_s) = (\mathfrak{r}(s), 0)$;
- (c) with $pr_X : X \times \mathbb{C} \rightarrow X$ the projection, $pr_X \circ \Psi(t, \mathfrak{v}) = \mathfrak{h}_t$;
- (d) $d(\mathfrak{r}(s), pr_X(\Psi(a, v))) < \varepsilon$ for $v \in U_s$.

The proof takes some preparation. Most of the construction will revolve around real analyticity. Let Y be a connected real analytic manifold. In what follows, we will drop ‘real’ from ‘real analytic’. By a theorem of Grauert, Y can be properly and analytically embedded in some Euclidean space [G58]; accordingly, we will assume $Y \subset \mathbb{R}^N$ is a closed analytic submanifold. For any $x \in \mathbb{R}^N$ in some neighborhood R of Y there is a unique nearest point $r(x)$ in Y ; the map $r : R \rightarrow Y$ is an analytic retraction.

Fix a topological space T and a compact interval $I \subset \mathbb{R}$. We will say that a map $g : T \times I \rightarrow Y$ is partly analytic, $g \in C^{0\omega}(T \times I, Y)$, if $T \times I \subset T \times \mathbb{C}$ has a neighborhood O to which g extends as a continuous function $\tilde{g} : O \rightarrow \mathbb{C}^N$ whose restrictions $\tilde{g}(t, \cdot)$ are holomorphic, $t \in T$. One can check that partial analyticity is independent of which embedding $Y \subset \mathbb{R}^N$ is chosen, and it does not even matter if Y is closed in \mathbb{R}^N , or only in some open subset of \mathbb{R}^N . It is clear that the composition of the partly analytic g with an analytic map $Y \rightarrow Y_1$ produces a partly analytic map.

Lemma 8.2. *Suppose M, P are finite dimensional complex manifolds, and $g \in C^{0\omega}((T \times M) \times I, P)$ is such that $g(t, \cdot, q) : M \rightarrow P$ is holomorphic for each $t \in T$, $q \in I$. Then $T \times M \times I \subset T \times M \times \mathbb{C}$ has a neighborhood O to which g extends as a continuous $f : O \rightarrow P$, whose restrictions $f(t, \cdot, \cdot)$ are holomorphic for each $t \in T$. If $O_{t,z} = \{q \in \mathbb{C} : (t, z, q) \in O\}$ are convex (that can always be achieved at the price of shrinking O), such extension f is unique.*

Proof. Assume first that M, P are open subsets of some $\mathbb{C}^m, \mathbb{C}^p$. With w_j, ζ_h the complex coordinates on \mathbb{C}^p , respectively \mathbb{C}^{2p} , consider the embedding α of \mathbb{C}^p (and so of P) into $\mathbb{R}^{2p} \subset \mathbb{C}^{2p}$, and its left inverse β ,

$$\alpha(w) = (\operatorname{Re} w_1, \operatorname{Im} w_1, \dots, \operatorname{Re} w_p, \operatorname{Im} w_p), \quad \beta(\zeta) = (\zeta_1 + i\zeta_2, \dots, \zeta_{2p-1} + i\zeta_{2p}).$$

By definition, $\alpha \circ g$ extends to a continuous $\tilde{g} : O \rightarrow \mathbb{C}^N$ whose restrictions $\tilde{g}(t, z, \cdot)$ are holomorphic. It follows that, possibly after shrinking O , $f = \beta \circ \tilde{g} : O \rightarrow P$ extends g and has holomorphic restrictions $f(t, z, \cdot)$; we can also arrange that the $O_{t,z}$ are convex. That even the restrictions $f(t, \cdot, \cdot)$ are holomorphic will follow by Morera's and Osgood's theorems once we show, denoting the coordinates on \mathbb{C}^m by z_1, \dots, z_m , that each $(t, z, q) \in O$ has a neighborhood $G \subset U$ such that $\int_{\Gamma} f(t, z, q) dz_k = 0$ whenever $\Gamma \subset M$ is a smooth loop contained in a complex line, $\{t\} \times \Gamma \times \{q\} \subset G$, and $k = 1, \dots, m$. But this is clear, as the integral is a holomorphic function of q , vanishing when $q \in I$.

The extension f is unique by the uniqueness theorem for holomorphic functions.

For a general P , cover $T \times M \times I$ by subsets $T_i \times M_i \times I_i$ that g maps into coordinate neighborhoods on P , T_i open, $M_i \subset M$ coordinate neighborhoods, $I_i \subset I$ compact intervals. We have just constructed extensions $f_i : O_i \rightarrow P$ to neighborhoods O_i of $T_i \times M_i \times I_i$. By uniqueness the f_i are compatible and produce the extension f claimed.

Lemma 8.3. *If T is compact and $L \subset I$ is finite, any $f \in C(T \times I, Y)$ can be approximated arbitrarily closely (in the uniform topology) by $g \in C^{0\omega}(T \times I, Y)$ such that $f = g$ on $T \times L$.*

Proof. We will avail ourselves of the following tools. There is a linear operator of interpolation, Λ , from the space $C(L, \mathbb{R}^N)$ of all functions $L \rightarrow \mathbb{R}^N$ to the space of polynomials $\mathbb{R} \rightarrow \mathbb{R}^N$ such that $(\Lambda\phi)|L = \phi$ for $\phi \in C(L, \mathbb{R}^N)$. Out of Λ we construct

$$\bar{\Lambda} : C(T \times L, \mathbb{R}^N) \rightarrow C^{0\omega}(T \times I, \mathbb{R}^N), \quad (\bar{\Lambda}\varphi)(t, \cdot) = \Lambda(\varphi(t, \cdot)),$$

continuous between the uniform topologies on $C(T \times L, \mathbb{R}^N)$ and $C^{0\omega}(T \times I, \mathbb{R}^N)$.

Furthermore, any $\varphi \in C(T \times I, \mathbb{R}^N)$ can be approximated uniformly by functions in $C^{0\omega}(T \times I, \mathbb{R}^N)$. (For this latter, extend φ to a compactly supported continuous $\tilde{\varphi} : T \times \mathbb{R} \rightarrow \mathbb{R}^N$. The approximating functions to φ can be obtained by convolutions, for example, with the Poisson kernel:

$$\varphi_\varepsilon(t, x) = \frac{\varepsilon}{\pi} \int_{\mathbb{R}} \frac{\tilde{\varphi}(t, y) dy}{|x - y|^2 + \varepsilon^2}, \quad (t, x) \in T \times I, \quad \varepsilon > 0.$$

Then $\varphi_\varepsilon \in C^{0\omega}(T \times I, \mathbb{R}^N)$ tend uniformly to φ as $\varepsilon \rightarrow 0$.)

To construct g of the lemma, first uniformly approximate f by $\phi \in C^{0\omega}(T \times I, \mathbb{R}^N)$. Next let $\psi = \phi + \bar{\Lambda}((f - \phi)|T \times L) \in C^{0\omega}(T \times I, \mathbb{R}^N)$, so that $\psi|T \times L = f|T \times L$. If the approximation by ϕ was good enough, then ϕ and ψ map $T \times I$ into the retract neighborhood R , and $g = r \circ \psi$ is the approximation of $f = r \circ f$ sought.

We return to the set up of Lemma 8.1. Let $I \subset \mathbb{R}$ be a compact interval and $q_0, \dots, q_J \in I$ distinct points.

Lemma 8.4. *Given $\mathfrak{x} \in C(S, X)$ and $\varepsilon > 0$, there is a $\delta > 0$ with the following property. Suppose $S = \coprod_0^J S_j$ is a Borel partition, S_0 is closed, $a < b$ are real numbers, and $[a, b] \ni t \mapsto \eta_t \in B(S, X)$ is a continuous path. Suppose furthermore that $\eta_t = \mathfrak{x}$ on S_0 , and each η_t is constant on S_j , $j = 1, \dots, J$. If $d_S(\mathfrak{x}, \eta_a) < \delta$, then there is a $\psi \in C^{0\omega}([a, b] \times S) \times I, X)$ such that*

- (a) $\psi(t, s, q_0) = \mathbf{r}(s)$ if $t \in [a, b]$, $s \in S$;
- (b) $\psi(t, s, q_j) = \boldsymbol{\eta}_t(s)$ if $t \in [a, b]$, $s \in S_j$, $j = 1, \dots, J$;
- (c) $d(\mathbf{r}(s), \psi(a, s, q)) < \varepsilon$ if $s \in S$, $q \in I$.

Proof. Embed X as a closed real analytic submanifold of some \mathbb{R}^N , with $r : R \rightarrow X$ an analytic retraction of a neighborhood $R \subset \mathbb{R}^N$. We can assume that the metric d of X is the restriction of the Euclidean metric; and also that $a = 0$ and $\varepsilon < 1$. Choose $\delta < \varepsilon/2$ so that the δ -neighborhood of

$$\{x \in X : |x| \leq 1 + \sup_S |\mathbf{r}|\} \subset \mathbb{R}^N$$

is contained in the retract neighborhood R ; and over this δ -neighborhood the retraction satisfies $|r - \text{id}| < \varepsilon/2$. Given $\boldsymbol{\eta}_t$, we will construct ψ in several steps. Consider

$$C = (S \times \{q_0\}) \cup \bigcup_{j=1}^J (\bar{S}_j \times \{q_j\}), \quad C_1 = [0, b] \times C, \quad C_2 = \{0\} \times S \times I;$$

C_1, C_2 are closed subsets of $[0, b] \times S \times I$. Define $\psi_1 : C_1 \rightarrow X$ by

$$\psi_1(t, s, q_j) = \begin{cases} \mathbf{r}(s) & \text{if } j = 0, \quad s \in S \\ \boldsymbol{\eta}_t(S_j) & \text{if } j \geq 1, \quad s \in \bar{S}_j. \end{cases}$$

In particular

$$(8.1) \quad d(\mathbf{r}(s), \psi_1(0, s, q)) \leq d_S(\mathbf{r}, \boldsymbol{\eta}_0) < \delta \quad \text{if } (0, s, q) \in C_1 \cap C_2.$$

Let $B_s \subset R$ denote the closed ball of radius δ about $\mathbf{r}(s)$. Extend $\psi_1|_{C_1 \cap C_2}$ to a continuous function $\psi_2 : C_2 \rightarrow \mathbb{R}^N$, then for $(s, q) \in S \times I$ define $\psi_3(0, s, q) \in B_s$ as the closest point to $\psi_2(0, s, q)$. Thus $\psi_3 : C_2 \rightarrow R$ is continuous and by (8.1) $\psi_1 = \psi_2 = \psi_3 = r \circ \psi_3$ on $C_1 \cap C_2$. Hence

$$\psi_4 = \begin{cases} \psi_1 & \text{on } C_1 \\ r \circ \psi_3 & \text{on } C_2 \end{cases}$$

defines a continuous function $C_1 \cup C_2 \rightarrow X$, which extends to a continuous $\psi_5 : G \rightarrow X$, with some neighborhood G of $C_1 \cup C_2$. Choose a neighborhood $G_1 \subset S \times I$ of C such that $[0, b] \times G_1 \subset G$, and let a continuous $\chi : S \times I \rightarrow [0, 1]$ be supported in G_1 , equal to 1 on $C_1 \cup C_2$. Then $\psi_6(t, s, q) = \psi_5(t\chi(s, q), s, q)$ defines a continuous function $[0, b] \times S \times I \rightarrow X$, that agrees with ψ_1 on C_1 and with $r \circ \psi_3$ on C_2 . Requirements (a,b) of the lemma hold for ψ_6 by the definition of ψ_1 , and (c) follows because $(0, s, q) \in C_2$ when $s \in S$, $q \in I$, and so

$$\begin{aligned} d(\mathbf{r}(s), \psi_6(0, s, q)) &= d(\mathbf{r}(s), r(\psi_3(0, s, q))) \\ &\leq d(\mathbf{r}(s), \psi_3(0, s, q)) + d(\psi_3(0, s, q), r(\psi_3(0, s, q))) < \delta + \varepsilon/2 < \varepsilon. \end{aligned}$$

A sufficiently good uniform approximation $\psi \in C^{0\omega}([a, b] \times S \times I, X)$ of ψ_6 such that $\psi(t, s, q_j) = \psi_6(t, s, q_j)$ for all t, s, j , as guaranteed by Lemma 8.3, will then do.

Proof of Lemma 8.1. To start, let $D \subset X \times X$ be a neighborhood of the diagonal and $F : D \rightarrow TX$ as in Lemma 2.1, and \exp as in (2.4). Possibly after shrinking X we can arrange that F and therefore \exp are analytic.

Consider the holomorphic vector bundle $\text{Hom} \rightarrow X \times X$ whose fibers are $\text{Hom}_{x,y} = \text{Hom}(T_x X, T_y X)$, and the open subset $\text{Iso} \subset \text{Hom}$ consisting of isomorphisms. The fiber bundle $p : \text{Iso} \rightarrow X \times X$ has a canonical analytic section, A , over D : this is so, because for fixed x the tangent spaces to $T_x X$ are canonically isomorphic, and the differential of $F(x, \cdot)$ sets up isomorphisms between $T_x X$, resp. $T_y X$ on the one hand, and the tangent spaces to $T_x X$ at $F(x, x), F(x, y)$ on the other. $A(x, y) \in \text{Hom}_{x,y}$ is then the composition of these isomorphisms or their inverses.

Let $I = [-1, 1]$ and $q_0 = 0, q_1, \dots, q_J \in I$ distinct points. We will show that the δ that Lemma 8.4. provides works for Lemma 8.1 as well. Given a, b, η_t as in Lemma 8.1, let $\psi \in C^{0\omega}([a, b] \times S \times I, X)$ be the map that Lemma 8.4 constructs. With $\psi_0 = \psi(a, \cdot, 0) : S \rightarrow X$ consider the sum of $\psi_0^* TX \rightarrow S$ and the trivial line bundle,

$$\pi : V = (\psi_0^* TX) \times \mathbb{C} \rightarrow S,$$

into which $(\psi_0^* TX) \times I$ is embedded.

Let furthermore $\tilde{\psi}(t, s, q) = \psi_0(s) = \psi(a, s, 0)$. The vector bundle $\tilde{\psi}^* TX$ is isomorphic to $\psi^* TX$, because $\psi, \tilde{\psi} : [a, b] \times S \times I \rightarrow X$ are homotopic. We use this isomorphism to construct a $C^{0\omega}$ map

$$(8.2) \quad \theta : ([a, b] \times S) \times I \rightarrow \text{Iso}, \quad p(\theta(t, s, q)) = (\psi_0(s), \psi(t, s, q)),$$

as follows. An isomorphism of the bundles translates to a continuous map θ_0 as in (8.2), that, by Lemma 8.3, can be approximated by $\theta_1 \in C^{0\omega}([a, b] \times S \times I, \text{Iso})$. Let $p \circ \theta_1 = \alpha \times \beta$, with $\alpha, \beta : [a, b] \times S \times I \rightarrow X$ close to $\tilde{\psi}$, resp. ψ . Then

$$\theta = (A \circ (\beta \times \psi)) \theta_1 (A \circ (\tilde{\psi} \times \alpha)) \in C^{0\omega}([a, b] \times S \times I, \text{Iso})$$

satisfies (8.2). (Above the values of θ are products of three linear transformations.)

With $\varpi : \psi_0^* TX \rightarrow S$ the bundle projection and $\Xi \subset \psi_0^* TX$ a suitable neighborhood of the zero section,

$$(8.3) \quad \phi : ([a, b] \times \Xi) \times I \ni (t, \xi, q) \mapsto (\exp(\theta(t, \varpi\xi, q)\xi), q) \in X \times \mathbb{C}$$

defines a $C^{0\omega}$ map that is holomorphic along the fibers of ϖ . We have

$$(8.4) \quad \phi(t, \xi_0, q) = (\psi(t, s, q), q) \quad \text{when } \xi_0 \in (\psi_0^* TX)_s \text{ is the zero vector,}$$

since then $\theta(t, \varpi\xi_0, q)\xi_0 \in T_{\psi(t,s,q)} X$ is the zero vector.

Locally in the base, the bundle $\psi_0^* TX$ is a direct product, and Ξ contains a neighborhood of the zero section that is a product of a piece of S with a complex manifold. Lemma 8.2 therefore gives, possibly after shrinking Ξ , an extension of ϕ to a neighborhood $O \subset [a, b] \times \Xi \times \mathbb{C} \subset [a, b] \times V$ of $[a, b] \times \Xi \times I$. The extension

$$\Phi = \Phi_1 \times \Phi_2 : O \rightarrow X \times \mathbb{C}$$

is continuous, and holomorphic along the fibers $O_{t,s} = O \cap (\{t\} \times V_s)$. By (8.3) $\Phi_2(t, \xi, q) = q$. We embed S into ψ_0^*TX as the zero section, and claim that—possibly after shrinking O to a smaller neighborhood of $[a, b] \times S \times I \subset [a, b] \times V$ —the restrictions $\Phi|_{O_{t,s}}$ map biholomorphically on open subsets of $X \times \mathbb{C}$.

Indeed, fix a zero vector $\xi_0 \in T_{\psi_0(s)}X$ and $q \in I$. According to the splitting $V = (\psi_0^*TX) \times \mathbb{C}$, the tangent space $T_{(t, \xi_0, q)}O_{t,s}$ also splits

$$(8.5) \quad T_{(t, \xi_0, q)}O_{t,s} \approx T_s\psi_0^*TX \oplus T_q\mathbb{C}.$$

In the splitting (8.5) the differential at (t, ξ_0, q) of $\Phi_1|_{O_{t,s}}$ is $\exp_*\theta(t, s, q) \oplus 0$, cf. (8.3), (8.4), and of $\Phi_2|_{O_{t,s}}$ it is $0 \oplus \text{id}_{T_q\mathbb{C}}$. Since \exp is biholomorphic on any T_xX , it follows that $(\Phi|_{O_{t,s}})_*$ is invertible at all points of $[a, b] \times S \times I$, hence also in a neighborhood in $[a, b] \times V$. As Φ is injective on $\{(t, s)\} \times I$, a simple indirect argument shows that if

$$O \subset [a, b] \times V = [a, b] \times (\psi_0^*TX) \times \mathbb{C}$$

is sufficiently shrunk, all $\Phi|_{O_{t,s}}$ will be biholomorphic.

We can take O of form $[a, b] \times O' \times \Omega$, with $O' \subset \psi_0^*TX$ a neighborhood of the zero section, and $\Omega \subset \mathbb{C}$ a convex neighborhood of I . Let $\Delta \subset \mathbb{C}$ be the unit disc and $f : \Delta \rightarrow \Omega$ biholomorphic, that maps 0 to 0. Choose a norm $|\cdot|_1$ on ψ_0^*TX whose unit ball bundle is contained in O' , and define a norm $|\cdot|$ on V by

$$|(\xi, \lambda)| = \max(|\xi|_1, |\lambda|), \quad \xi \in \psi_0^*TX, \quad \lambda \in \mathbb{C},$$

whose unit ball bundle U is contained in $O' \times \Delta$. The formula

$$(8.6) \quad \Psi(t, \xi, \lambda) = \Phi(t, \xi, f(\lambda)), \quad t \in [a, b], \quad (\xi, \lambda) \in U,$$

then defines a continuous $\Psi : [a, b] \times U \rightarrow X \times \mathbb{C}$, whose restrictions to $\{t\} \times U_s$ are biholomorphisms to open subsets of $X \times \mathbb{C}$, as (a) claims.

On $[a, b] \times S \times I$, by (8.4), $\text{pr}_X \circ \phi$ and $\text{pr}_X \circ \Phi$ agree with ψ . Therefore Lemma 8.4c implies that, if O above is taken a sufficiently small neighborhood of $[a, b] \times S \times I$, then $d(\mathfrak{r}(s), \text{pr}_X(\Phi(a, v))) < \varepsilon$ for v in the fiber $V_s \cap O$. Hence $d(\mathfrak{r}(s), \text{pr}_X(\Psi(a, v))) < \varepsilon$ when $v \in U_s$: (d) is also satisfied.

Finally, by (8.4) and by Lemma 8.4a,b (recall that $S \subset \psi_0^*TX$ and $S \times \{0\} \subset V$ are the zero sections)

$$\Phi(t, s, q_j) = \phi(t, s, q_j) = (\psi(t, s, q_j), q_j) = \begin{cases} (\mathfrak{r}(s), 0) & \text{if } s \in S, j = 0 \\ (\mathfrak{h}_t(s), q_j) & \text{if } s \in S_j, j = 1, \dots, J. \end{cases}$$

Therefore if we define $\mathfrak{v}(s) = f^{-1}(q_j)$ when $s \in S_j$, $j = 0, \dots, J$, Ψ will satisfy the remaining requirements (b,c) as well.

9 The extension theorem in mapping spaces

In this section we prove Theorem 7.5. Since the spaces $C(S, X, A, \mathfrak{r}_0) \subset B(S, X, A, \mathfrak{r}_0)$ do not change if \mathfrak{r}_0 is replaced by any other $\mathfrak{r} \in C(S, X, A, \mathfrak{r}_0)$, we will take $\mathfrak{r} = \mathfrak{r}_0$. Recall

the notion of basic neighborhoods $D_A(\eta) \subset \mathfrak{B}$, defined by a neighborhood $D \subset X \times X$ of the diagonal, see (5.1). If D is as in Lemma 2.1, then $D_A(\eta)$ is contractible.

Given the $E|\mathfrak{C}$ valued germ \mathbf{f} of Theorem 7.5, let $C \in \mathbb{R}$ be such that any analytic continuation \mathbf{h}_t of \mathbf{f} along any path \mathbf{x}_t in \mathfrak{C} satisfies $p(\mathbf{h}_t(\mathbf{x}_t)) \leq C$. Lemmas 9.1, 9.3 below continue \mathbf{f}^{ab} along special paths η_t in \mathfrak{B} . Fix D above so that \mathbf{f} is the germ of a holomorphic section f of $E|\mathfrak{C} \cap D_A(\mathbf{x})$ and f has an Aron–Berner–type extension f^{ab} to a section of $E''|D_A(\mathbf{x})$. Then \mathbf{f}^{ab} is the germ of f^{ab} .

Lemma 9.1. *Consider a path $[0, 1] \ni t \mapsto \eta_t \in \mathfrak{B}$ starting at \mathbf{x} . Assume $a \in (0, 1)$ is such that $\eta_t \in D_A(\mathbf{x})$ when $t \leq a$. Assume also that there are a normed topological vector bundle $\pi : (V, |\cdot|) \rightarrow S$ with unit ball bundle $U \rightarrow S$, a $\mathbf{v} \in B_A(V)$, $\sup_S |\mathbf{v}| < 1$, and a continuous $\Psi : [a, 1] \times U \rightarrow X$ such that for every $t \in [a, 1]$ and $s \in S$*

- (a) Ψ maps the fibers $\{t\} \times U_s$ biholomorphically on open subsets of X ;
- (b) the zero vector $0_s \in V_s$ satisfies $\Psi(t, 0_s) = \mathbf{x}(s)$;
- (c) $\Psi(t, \mathbf{v}) = \eta_t$;
- (d) $\Psi(a, v) \in D^{\mathbf{x}(s)}$ for $v \in U_s$, cf. (2.3).

Then \mathbf{f}^{ab} has an analytic continuation \mathbf{g}_t along η_t , such that $p(\mathbf{g}_t(\eta_t)) \leq C$.

Let $\Gamma \subset B_A(V)$ be the unit ball.

Lemma 9.2. *For $t \in [a, 1]$ the map*

$$(9.1) \quad \Psi_t^\circ : \Gamma \ni \mathbf{u} \mapsto \Psi(t, \mathbf{u}) \in D_A(\mathbf{x}), \quad \text{cf. (d) above,}$$

is biholomorphic on its image, an open subset of $D_A(\mathbf{x})$.

Proof. Fix $t \in [a, 1]$. First we claim that the map

$$G : U \ni v \mapsto (\pi v, \Psi(t, v)) \in S \times X$$

is injective and open. Injectivity follows from (a) of Lemma 9.1, since G maps U_s to $\{s\} \times X$ injectively. That G is open is a local property. Accordingly, we can assume that $V \rightarrow S$ is trivial, $V = S \times \mathbb{C}^r$. Suppose $\Psi(t, s_0, u_0) = z_0 \in X$ with some $(s_0, u_0) \in U$, and let $B \subset \mathbb{C}^r$ be a small compact neighborhood of u_0 . The topological degree $\deg(\Psi(t, s_0, \cdot), B, z_0) = 1$. Hence $\deg(\Psi(t, s, \cdot), B, z) = 1$ for s, z close to s_0, z_0 ; $\Psi(t, s, \cdot)$ attains the value z in B , and it follows that G is indeed open. This implies that Ψ_t° is open. It is also injective.

The inverse Φ of G is a homeomorphism between an open $O \subset S \times X$ and U , holomorphic on $O \cap (\{s\} \times X)$, and satisfies $\Psi(t, \Phi(s, z)) = z$, $s \in S$, $z \in D^{\mathbf{x}(s)}$. The map

$$(9.2) \quad \Phi^\circ : \Psi_t^\circ(\Gamma) \ni \mathfrak{z} \mapsto \Phi \circ (\text{id}_S \times \mathfrak{z}) \in \Gamma$$

is the inverse of Ψ_t° , and both Φ°, Ψ_t° are holomorphic. To check the holomorphy of Ψ_t , say, take a local coordinate $\psi_{\mathfrak{z}}$ on \mathfrak{B} as in (5.2) with $F : D \rightarrow TX$ of Lemma 2.1,

$$\psi_{\mathfrak{z}} : D_A(\mathfrak{z}) \ni \mathfrak{w} \mapsto F \circ (\mathfrak{z} \times \mathfrak{w}) \in B_A(\mathfrak{z}^*TX).$$

The formula $H(v) = F(\mathfrak{z}(\pi v), \Psi(t, v)) \in T_{\mathfrak{z}(\pi v)}X$ defines a map from a Borel subset of V to \mathfrak{z}^*TX that satisfies the assumptions of Lemma 1.3. Therefore $\psi_{\mathfrak{z}} \circ \Psi_t^\circ = F \circ (\mathfrak{z} \times \Psi_t^\circ)$,

the restriction of H° of that lemma, is holomorphic on the open subset of Γ where it is defined. This means Ψ_t° is indeed holomorphic. A similar argument proves for Φ° ; therefore both are biholomorphic.

Proof of Lemma 9.1. For values $t \leq a$ the germs of f^{ab} at η_t supply the continuation \mathbf{g}_t . To continue \mathbf{g}_a to values $t \geq a$, we will prove that with Ψ_t° of (9.1):

- (1) \mathbf{f}^{ab} is the germ of a holomorphic section g_t of $E''|_{\Psi_t^\circ(\Gamma)}$, $\sup p''(g_t) \leq C$;
- (2) the germ of g_a at η_a is \mathbf{g}_a ;
- (3) the germs \mathbf{g}_t of g_t at η_t , $t \in [a, 1]$, provide an analytic continuation of \mathbf{g}_a .

(1) Fix $t \in [a, 1]$. The pullback $\Psi_t^{\circ*}(E, p) = (\bar{E}, \bar{p}) \rightarrow \Gamma$ is a holomorphic Banach bundle with a flat norm \bar{p} . By Theorem 6.3 it is therefore trivial, and its (bounded) holomorphic sections can be viewed as (bounded) holomorphic functions with values in a fixed Banach space $(\mathfrak{Y}, \|\cdot\|)$. Since the germ $\mathbf{k} = \mathbf{f} \circ \Psi_t^\circ$ has bounded analytic continuation along any path in $\Gamma \cap \mathfrak{C}$ (supplied by the pullback along of Ψ_t° of the corresponding continuation of \mathbf{f}), it is the germ of a bounded holomorphic $k : \Gamma \cap \mathfrak{C} \rightarrow \mathfrak{Y}$. This k has an Aron–Berners–type extension k^{ab} to Γ with $\sup p''(k^{\text{ab}}) \leq C$, constructed in section 4. Corollary 4.6 implies that the germ of k^{ab} at 0 is $\mathbf{f}^{\text{ab}} \circ \Psi_t^\circ$, and this means that $g_t = k^{\text{ab}} \circ \Phi^\circ$ will do, cf. (9.2).

(2) Since $\Psi_a^\circ(\Gamma) \subset D_A(\mathfrak{x})$ is connected, and f^{ab}, g_a share the same germ at \mathfrak{x} , namely \mathbf{f}^{ab} , on $\Psi_a^\circ(\Gamma)$ we have $f^{\text{ab}} = g_a$. In particular, their germs at $\eta_a = \Psi_a^\circ(\mathfrak{v})$ agree.

(3) Given $t \in [a, 1]$, connect \mathfrak{x} with η_t by a path in $\Psi_t^\circ(\Gamma)$. By continuity of Ψ , for $\tau \in [a, 1]$ close to t , this path is contained in $\Psi_\tau^\circ(\Gamma)$. Since the germs of g_t, g_τ at \mathfrak{x} agree, g_t, g_τ agree in a neighborhood of the path, hence in a neighborhood of η_t . But this means that the germs \mathbf{g}_t depend continuously on t , and they continue \mathbf{g}_a analytically. This completes the proof.

Again we metrize X with a metric d , then d_S of (2.2) metrizes \mathfrak{B} .

Lemma 9.3. *There is a $\delta > 0$ with the following property. Let $S = \coprod_{j=0}^J S_j$ be a partition in Borel sets, $S_0 = A$. Suppose that $[0, 1] \ni t \mapsto \eta_t \in \mathfrak{B}$ is a path starting at \mathfrak{x} , and $a \in (0, 1)$ is such that*

- (a) $d_S(\mathfrak{x}, \eta_t) < \delta$ when $t \in [0, a]$;
- (b) $\eta_t|_{S_j}$ is constant when $t \in [a, 1]$, $j = 1, 2, \dots, J$.

Then \mathbf{f}^{ab} has an analytic continuation \mathbf{g}_t along η_t , such that $p(\mathbf{g}_t(\eta_t)) \leq C$.

Proof. With $X^+ = X \times \mathbb{C}$ and $\text{pr}_X : X^+ = X \times \mathbb{C} \rightarrow X$ the projection, let $D^+ = (\text{pr}_X \times \text{pr}_X)^{-1}D \subset X^+ \times X^+$, a neighborhood of the diagonal, $\mathfrak{x}^+ = \mathfrak{x} \times 0 \in C(S, X^+)$,

$$\mathfrak{B}^+ = \{\mathfrak{z} \in B(S, X^+) : \mathfrak{z}|_A = \mathfrak{x}^+|_A\},$$

and \mathfrak{C}^+ the component of \mathfrak{x}^+ in $\mathfrak{B}^+ \cap C(S, X^+)$. Choose $\varepsilon > 0$ so that $D(\mathfrak{x})$ contains the d_S metric ball of radius ε about \mathfrak{x} . Given \mathfrak{x} and ε , the $\delta > 0$ provided by Lemma 8.1 will do.

Indeed, if (a), (b) of our lemma hold, choose a normed vector bundle $(V, \|\cdot\|) \rightarrow S$ with unit ball bundle $U \subset V$, \mathfrak{v} in the unit ball $\Gamma \subset B_A(V)$, and a map $\Psi : [a, 1] \times U \rightarrow X^+$ as guaranteed by Lemma 8.1. Consider the projection $P : \mathfrak{B}^+ \rightarrow \mathfrak{B}$ given by $P(\mathfrak{z}) = \text{pr}_X \circ \mathfrak{z}$; it is holomorphic and maps \mathfrak{C}^+ to \mathfrak{C} . The pullback $f \circ P$ is a

holomorphic section of $P^*E|(\mathfrak{C}^+ \cap D^+(\mathfrak{r}^+))$, whose germ at \mathfrak{r}^+ we denote \mathbf{h} . Now \mathbf{h} analytically continues along any path r in \mathfrak{C}^+ , the continuations provided by the lifts of the continuations along $P \circ r$; these continuations are bounded, with the same bound C as for \mathbf{f} . Let us set $\mathfrak{z}_t = \Psi(t, \mathfrak{v}) \in \mathfrak{B}^+$, $t \in [0, 1]$. We are in the situation of Lemma 9.1, with X replaced by X^+ , \mathfrak{B} by \mathfrak{B}^+ , \mathfrak{v}_t by \mathfrak{z}_t . By the lemma \mathbf{h}^{ab} has an analytic continuation \mathbf{k}_t along \mathfrak{z}_t . It remains to show that the P^*E'' valued germs \mathbf{k}_t are pullbacks of germs \mathbf{g}_t that form the analytic continuation of \mathbf{f}^{ab} along \mathfrak{v}_t .

Let us call a holomorphic vector field ξ on \mathfrak{B}^+ vertical if it is tangential to the fibers $P^{-1}(\mathfrak{v})$, $\mathfrak{v} \in \mathfrak{B}$. Consider a Banach space \mathfrak{Y} and a \mathfrak{Y} valued holomorphic germ \mathbf{k} at some $\mathfrak{z} \in \mathfrak{B}^+$. It is of form $\mathbf{k} = \mathbf{g} \circ P$ with a \mathfrak{Y} valued holomorphic germ \mathbf{g} at $P(\mathfrak{z})$ if and only if the derivatives $\xi \mathbf{k} = 0$ for all vertical vector fields ξ . Similarly, if \mathbf{k} is a P^*E'' valued germ at $\mathfrak{z} \in \mathfrak{B}^+$, we can compute its vertical derivatives $\xi \mathbf{k}$, since the restriction of (a representative of) \mathbf{k} to a fiber $P^{-1}(\mathfrak{u})$ takes values in the fixed Banach space $E''_{P(\mathfrak{u})}$; and \mathbf{k} is a pullback if and only if $\xi \mathbf{k} = 0$ for all vertical ξ .

For the analytic continuation \mathbf{k}_t of $\mathbf{h}^{\text{ab}} = \mathbf{k}_0$ constructed above, $\xi \mathbf{k}_0 = 0$, because by Corollary 4.6 \mathbf{k}_0 is the pullback of \mathbf{f}^{ab} along P . Hence its analytic continuation $\xi \mathbf{k}_t$ also vanishes, and \mathbf{k}_t is the pullback of some germ \mathbf{g}_t at \mathfrak{v}_t . By Lemma 7.2 \mathbf{g}_t depend continuously on t , hence supply the analytic continuation of \mathbf{f}^{ab} .

We will reduce continuation along arbitrary paths to continuation along paths as in Lemma 9.3 in the following way. By a partition we will mean a decomposition $S = \coprod_{j=0}^J S_j$ into Borel sets such that $S_0 = A$, $S_j \neq \emptyset$, $j = 1, \dots, J$. Given such a partition, denoted Π , let

$$(9.3) \quad \mathfrak{B}_\Pi = \{\mathfrak{v} \in \mathfrak{B} : \mathfrak{v}|S_j = \text{const}, \text{ for } j = 1, \dots, J\}.$$

If Π' refines Π (notation: $\Pi' \succ \Pi$), then $\mathfrak{B}_{\Pi'} \supset \mathfrak{B}_\Pi$.

Lemma 9.4. (a) $\mathfrak{B}_\Pi \subset \mathfrak{B}$ is a closed submanifold, biholomorphic to X^J ;

(b) If Π_0 is a partition, then $\bigcup_{\Pi \succ \Pi_0} \mathfrak{B}_\Pi$ is dense in \mathfrak{B} ;

(c) Let $c \in \mathbb{R}$, $\mathfrak{U} \subset \mathfrak{B}$ open, $K \rightarrow \mathfrak{U}$ a holomorphic Banach bundle with a flat norm p , and Π_0 a partition. Suppose for each $\Pi \succ \Pi_0$ we are given a holomorphic section h_Π of $K|_{\mathfrak{B}_\Pi \cap \mathfrak{U}}$ such that $p(h_\Pi) \leq c$, and $\Pi' \succ \Pi$ implies $h_{\Pi'} = h_\Pi$ on $\mathfrak{B}_\Pi \cap \mathfrak{U}$. Then there is a unique holomorphic section h of $K|\mathfrak{U}$ such that $h|_{\mathfrak{B}_\Pi \cap \mathfrak{U}} = h_\Pi$.

(d) Let $K \rightarrow \mathfrak{B}$ be a holomorphic Banach bundle with a flat norm. If a K valued holomorphic germ \mathbf{h} at some $\mathfrak{z} \in \mathfrak{B}_{\Pi_0}$ has analytic continuation along any path in \mathfrak{B}_Π for $\Pi \succ \Pi_0$, and these continuations have the same bound c , then \mathbf{h} has an analytic continuation along any path in \mathfrak{B} , bounded by c .

Proof. (a) \mathfrak{B}_Π is clearly a closed subset. If $\mathfrak{v} \in \mathfrak{B}_\Pi$, local coordinates $\psi_{\mathfrak{v}}$ as in (5.2) send the neighborhood $D_A(\mathfrak{v})$ to an open $\mathfrak{V} \subset B_A(\mathfrak{v}^*TX)$. For all $j \geq 1$ the fibers of \mathfrak{v}^*TX over S_j are the same tangent spaces to X , hence one can talk about sections of \mathfrak{v}^*TX being constant over S_j . Those bounded Borel sections of \mathfrak{v}^*TX that vanish on S_0 and are constant over each S_j , $j \geq 1$, form a finite dimensional subspace

$$B_\Pi(\mathfrak{v}^*TX) \subset B_A(\mathfrak{v}^*TX),$$

and the local coordinate ψ_η sends $\mathfrak{B}_\Pi \cap D_A(\eta)$ to $B_\Pi(\eta^*TX) \cap \mathfrak{W}$. Therefore \mathfrak{B}_Π is indeed a submanifold of \mathfrak{B} . One can then check that the map that sends $(x_1, \dots, x_J) \in X^J$ to $\mathfrak{z} \in \mathfrak{B}_\Pi$ defined by $\mathfrak{z}(s) = x_j$ if $s \in S_j$, $j = 1, \dots, J$, is biholomorphic.

(b) Let $\eta \in \mathfrak{B}$ and $\varepsilon > 0$. To find a partition $\Pi \succ \Pi_0$ and $\mathfrak{z} \in \mathfrak{B}_\Pi$ in the ε neighborhood of η , cover the closure of $\eta(S) \subset X$ by finitely many balls of radius ε , choose $\Pi \succ \Pi_0$, $S = \coprod_{j=0}^J S_j$, so that each $\eta(S_j)$ is contained in one of those balls, centered at, say, $y_j \in X$, $j \geq 1$. Then $\mathfrak{z} \in \mathfrak{B}_\Pi$ given by

$$\mathfrak{z}(s) = \begin{cases} \mathfrak{r}(s) & \text{when } s \in S_0 = A \\ y_j & \text{when } s \in S_j, \quad i = 1, \dots, J \end{cases}$$

will do.

(c) We can assume that \mathfrak{U} is contained in a coordinate neighborhood about some $\eta \in \mathfrak{B}$ and that (K, p) is trivial over it. Then instead of sections one can talk about Banach space valued functions. The local coordinate ψ_η maps \mathfrak{U} to a neighborhood $\mathfrak{W} \subset B_A(\eta^*TX)$ of the zero section, and $\mathfrak{B}_\Pi \cap \mathfrak{U}$ into $B_\Pi(\eta^*TX)$ from the proof of part (a). A hermitian metric on X induces a norm on $B_A(\eta^*TX)$, and we can assume that \mathfrak{W} is the unit ball.

Accordingly, the claim will be proved once we show the following. Let \mathfrak{X} , $(\mathfrak{Y}, \|\cdot\|)$ be Banach spaces, $\mathfrak{W} \subset \mathfrak{X}$ the unit ball, and $c \in \mathbb{R}$. Suppose for each partition $\Pi \succ \Pi_0$ we are given a closed subspace $\mathfrak{X}_\Pi \subset \mathfrak{X}$ and $h_\Pi \in \mathcal{O}(\mathfrak{X}_\Pi \cap \mathfrak{W}, \mathfrak{Y})$ with the following properties. First, $\|h_\Pi\| \leq c$; second, $\mathfrak{X}_{\Pi'} \supset \mathfrak{X}_\Pi$ and $h_{\Pi'}|_{\mathfrak{X}_\Pi \cap \mathfrak{W}} = h_\Pi$ if $\Pi' \succ \Pi$; and finally, $\mathfrak{Z} = \bigcup_{\Pi \succ \Pi_0} \mathfrak{X}_\Pi$ is dense in \mathfrak{X} . Then there is a unique $h \in \mathcal{O}(\mathfrak{W}, \mathfrak{Y})$ such that $h|_{\mathfrak{X}_\Pi \cap \mathfrak{W}} = h_\Pi$ and $\|h\| \leq c$. (Of course, partitions here are irrelevant, Π could denote elements of an arbitrary directed set.)

The density of \mathfrak{Z} implies that h is unique. To construct h , define $g : \mathfrak{Z} \cap \mathfrak{W} \rightarrow \mathfrak{Y}$ by $g(x) = h_\Pi(x)$ if $x \in \mathfrak{X}_\Pi \cap \mathfrak{W}$. Cauchy estimates for the derivatives of h_Π imply that on a concentric ball $\mathfrak{W}_r \subset \mathfrak{W}$ of radius r the Lipschitz constant of h_Π is $\leq c/(1-r)$, hence the same holds for g . In particular, if $x \in \mathfrak{W}$ and a sequence $x_i \in \mathfrak{Z} \cap \mathfrak{W}$ converges to x , then $g(x_i)$ form a Cauchy sequence. Define $h(x) = \lim_i g(x_i)$; the limit is independent of the sequence $x_i \rightarrow x$. The function h constructed is continuous and its norm is bounded by c . To complete the proof we need to check that for any $x, y \in \mathfrak{X}$ and $\Lambda = \{\lambda \in \mathbb{C} : x + \lambda y \in \mathfrak{W}\}$, the function $f : \Lambda \ni \lambda \mapsto h(x + \lambda y)$ is holomorphic. Let $\Lambda_0 \subset \Lambda$ be open and relatively compact. Choose sequences $\Pi_i \succ \Pi_0$, and $x_i, y_i \in \mathfrak{X}_{\Pi_i}$ convergent to x, y . Then $f_i(\lambda) = h(x_i + \lambda y_i) = h_{\Pi_i}(x_i + \lambda y_i)$ is holomorphic on Λ_0 when i is sufficiently large. Since $\|f_i\| \leq c$ and $f_i \rightarrow f$ on Λ_0 , it follows that f is holomorphic, and therefore so is h .

(d) Let $r : [a, b] \rightarrow \mathfrak{B}$ be a path, \mathbf{h} a germ at $r(a) = \mathfrak{z}$. Let $\mathfrak{W}_t \subset D_A(r(t))$ be neighborhoods of $r(t)$ whose image under the local coordinate $\psi_{r(t)}$ is convex, $r(a) \in \mathfrak{W}_a$ for t in a neighborhood of a , and \mathbf{h} is the germ of a holomorphic section h^0 of $K|\mathfrak{W}_a$. Then $\mathfrak{B}_\Pi \cap \mathfrak{W}_t$ for each Π is contractible or empty. Choose $a = t_0 < t_1 < \dots < t_k = b$ so that $r[t_{j-1}, t_j] \subset \mathfrak{W}_{t_j}$ when $1 \leq j \leq k$. By recurrence we construct holomorphic sections h^j of $K|\mathfrak{W}_{t_j}$ that define the analytic continuation of \mathbf{h} along r . If $j \geq 1$ and h^i for $i < j$ has been constructed, choose a partition $\Pi_1 \succ \Pi_0$ and \mathfrak{w} in the connected component of $r(t_{j-1})$ in $\mathfrak{B}_{\Pi_1} \cap \mathfrak{W}_{t_{j-1}} \cap \mathfrak{W}_{t_j}$. The analytic continuations of the germ of h^{j-1} at \mathfrak{w} along paths in $\mathfrak{B}_\Pi \cap \mathfrak{W}_{t_j}$, $\Pi \succ \Pi_1$, define holomorphic sections h_Π of

$K|\mathfrak{B}_\Pi \cap \mathfrak{W}_{t_j}$. Since these sections satisfy the conditions in (c), they are restrictions of a holomorphic section h^j of $K|\mathfrak{W}_{t_j}$. By the uniqueness part of (c) the germs of h^{j-1} and h^j agree at \mathfrak{w} , therefore in the connected component of \mathfrak{w} in $\mathfrak{W}_{t_{j-1}} \cap \mathfrak{W}_{t_j}$. In particular, h^{j-1} and h^j agree near $r(t_{j-1})$. Thus the collection h^j indeed gives rise to an analytic continuation of \mathbf{h} along r .

Proof of Theorem 7.5. It suffices to construct continuations along paths $[0, 1] \ni t \mapsto \mathfrak{r}_t \in \mathfrak{B}$, $\mathfrak{r}_0 = \mathfrak{r}$. Let $\delta > 0$ as in Lemma 9.3 and $a \in (0, 1/2)$ be such that the δ -neighborhood $\mathfrak{U} \subset \mathfrak{B}$ of \mathfrak{r} is contained in $D_A(\mathfrak{r})$ and contains \mathfrak{r}_t when $t \in [0, 2a]$. Choose a partition $S = \coprod_{i=0}^l T_i$, denoted Π_0 , such that \mathfrak{B}_{Π_0} intersects \mathfrak{U} (Lemma 9.4b), say, $\eta \in \mathfrak{B}_{\Pi_0} \cap \mathfrak{U}$. Since the image of \mathfrak{U} under the local coordinate $\psi_{\mathfrak{r}}$ is convex, there is a path $[0, 2a] \ni t \mapsto \eta_t \in \mathfrak{U}$ from \mathfrak{r} to \mathfrak{r}_{2a} such that $\eta_a = \eta$; extend it to $2a \leq t \leq 1$ by $\eta_t = \mathfrak{r}_t$. The germs of f^{ab} provide continuation of \mathbf{f}^{ab} along both \mathfrak{r}_t, η_t , $t \in [0, 2a]$. These continuations give the same result at \mathfrak{r}_{2a} , that we denote \mathbf{g} . Let \mathbf{h} be the germ of f^{ab} at η_a . To produce analytic continuation of \mathbf{f}^{ab} along \mathfrak{r}_t , $t \in [0, 1]$, it suffices to show continuation of \mathbf{g} along \mathfrak{r}_t , $t \in [2a, 1]$, or of \mathbf{h} along η_t , $t \in [a, 1]$. But Lemma 9.3 guarantees that this latter continuation exists along paths η'_t for which $\eta'_t \in \mathfrak{B}_\Pi$ when $t \in [a, 1]$; here $\Pi \succ \Pi_0$ is arbitrary. Hence, by Lemma 9.4d, the continuation exists along η_t as well. This completes the proof.

10 The Monodromy theorem

In this section we prove the following generalization of Theorem 0.2. Consider mapping spaces $\mathfrak{B}, \mathfrak{C}$ determined by a compact Hausdorff space S , a complex manifold X , a continuous $\mathfrak{r} \in C(S, X)$ and a closed $A \subset S$, as before.

Theorem 10.1. *Suppose X is simply connected, $E \rightarrow \mathfrak{B}$ is a holomorphic Banach bundle with a flat norm p , and an $E|\mathfrak{C}$ valued holomorphic germ \mathbf{f} admits bounded analytic continuation along any path in \mathfrak{C} . Then \mathbf{f} is the germ of a holomorphic section f of $E|\mathfrak{C}$. If, furthermore, X is compact, then f is a horizontal section (Definition 6.4).*

Fix a partition $S = \coprod_{j=0}^J S_j$ as in section 9, denote it Π_0 , and let $\bar{\mathfrak{r}} \in \mathfrak{B}_{\Pi_0}$, see (9.3). Given $\nu = 0, 1, \dots$, inclusions induce a directed system of homotopy groups/sets and homomorphisms

$$(10.1) \quad \pi_\nu(\mathfrak{B}_\Pi, \bar{\mathfrak{r}}) \rightarrow \pi_\nu(\mathfrak{B}_P, \bar{\mathfrak{r}}), \quad P \succ \Pi \succ \Pi_0.$$

Lemma 10.2. *$\pi_\nu(\mathfrak{B}, \bar{\mathfrak{r}})$ is isomorphic to the direct limit of the system (10.1). Hence, if X is connected (our standing assumption), resp. simply connected, then so is \mathfrak{B} .*

Proof. We need to show that any continuous map $(S^\nu, *) \rightarrow (\mathfrak{B}, \bar{\mathfrak{r}})$, and any homotopy between two such maps, can be deformed to a map, respectively homotopy, into \mathfrak{B}_Π with some partition $\Pi \succ \Pi_0$; and that any homotopy in $(\mathfrak{B}, \bar{\mathfrak{r}})$ between continuous maps $f_1, f_2 : (S^\nu, *) \rightarrow (\mathfrak{B}_\Pi, \bar{\mathfrak{r}})$ can be deformed, fixing f_1, f_2 , to a homotopy in some $(\mathfrak{B}_P, \bar{\mathfrak{r}})$, $P \succ \Pi$. Once this done, we can define a homomorphism $\pi_\nu(\mathfrak{B}, \bar{\mathfrak{r}}) \rightarrow \varinjlim \pi_\nu(\mathfrak{B}_\Pi, \bar{\mathfrak{r}})$ by first sending the class in $\pi_\nu(\mathfrak{B}, \bar{\mathfrak{r}})$ of an $f : (S^\nu, *) \rightarrow (\mathfrak{B}, \bar{\mathfrak{r}})$ to an $f_1 : (S^\nu, *) \rightarrow (\mathfrak{B}_\Pi, \bar{\mathfrak{r}})$

homotopic to f , and then to the element of $\varinjlim \pi_\nu(\mathfrak{B}_P, \bar{\mathfrak{r}})$ that the class $[f_1] \in \pi_\nu(\mathfrak{B}_\Pi, \bar{\mathfrak{r}})$ induces. This homomorphism is automatically bijective.

The deformation statements are special cases of the following: Let T be a compact Hausdorff space, $C \subset T$ closed. For any continuous $\phi : (T, C) \rightarrow (\mathfrak{B}, \mathfrak{B}_\Pi)$ there is a continuous deformation $\phi_\tau : (T, C) \rightarrow (\mathfrak{B}, \mathfrak{B}_\Pi)$, $0 \leq \tau \leq 1$, such that $\phi_0 = \phi$, $\phi_\tau|_C = \phi|_C$ for all τ , and $\phi_1(T) \subset \mathfrak{B}_P$ with some $P \succ \Pi$. To verify this, we will use $D \subset X \times X$, D^x , $F : D \rightarrow TX$ of Lemma 2.1, and $\exp : TX \rightarrow X$ of (2.4). Since $\phi(t) \in \mathfrak{B}$ is a continuous function of $t \in T$, and each $\phi(t)(S) \subset X$ has compact closure, there is a compact $L \subset X$ that contains all $\phi(t)(S)$. Cover L by finitely many coordinate neighborhoods D^{x_i} , and choose a partition $P \succ \Pi$, given by $S = \coprod_{i=0}^I R_i$, such that $\phi(t)(R_i) \subset D^{x_i}$ for every $t \in T$ and $i = 1, \dots, I$. If $s_i \in R_i$, $i \geq 1$, then

$$\phi_\tau(t)(s) = \begin{cases} \phi(t)(s) = \mathfrak{r}(s) & \text{if } s \in R_0 = A \\ \exp \{ (1 - \tau)F(x_i, \phi(t)(s)) + \tau F(x_i, \phi(t)(s_i)) \} & \text{if } s \in R_i, \quad i = 1, \dots, I \end{cases}$$

is the deformation sought.

The second statement of the lemma follows because by Lemma 9.4a the spaces \mathfrak{B}_Π are homeomorphic to Cartesian powers of X , hence for (simply) connected X they themselves are (simply) connected.

Proof of Theorem 10.1. Since \mathfrak{B} is connected and simply connected by Lemma 10.2, (E, p) is trivial. Again, instead of sections of (E, p) we can talk about functions with values in some fixed Banach space $(\mathfrak{Y}, \|\cdot\|)$. Accordingly, we take \mathbf{f} to be a \mathfrak{Y} valued germ. By Theorem 7.5 \mathbf{f}^{ab} admits bounded analytic continuation along any path in \mathfrak{B} ; again because \mathfrak{B} is simply connected, this implies that \mathbf{f}^{ab} is the germ of a holomorphic $g : \mathfrak{B} \rightarrow \mathfrak{Y}''$. Hence \mathbf{f} is the germ of the holomorphic function $f = g|_{\mathfrak{C}}$. Since its germ \mathbf{f} takes values in $\mathfrak{Y} \subset \mathfrak{Y}''$, all the values that f takes are in \mathfrak{Y} .

If, in addition, X is compact, then the submanifolds $\mathfrak{B}_\Pi \approx X^J$ are also compact (and connected), hence the restrictions $g|_{\mathfrak{B}_\Pi}$ are constant. As $\bigcup_\Pi \mathfrak{B}_\Pi$ is dense in \mathfrak{B} , Lemma 9.4b, this implies that g itself is constant, and therefore f is constant.

11 Examples

In this section we have collected examples that show that some assumptions in Theorems 7.5, 10.1 cannot be dispensed with, or that the conclusions cannot be strengthened. We keep notation used in sections 7, 10.

Example 11.1. *There are mapping spaces $\mathfrak{C}, \mathfrak{B}$ and a bounded holomorphic function $f : \mathfrak{C} \rightarrow \mathbb{C}$ with germ \mathbf{f} at some \mathfrak{r}_0 such that \mathbf{f}^{ab} is not the germ of a single valued holomorphic function $g : \mathfrak{B} \rightarrow \mathbb{C}$.*

Let $r \in (1, \infty)$, $X = \{x \in \mathbb{C} : 1/r < |x| < r\}$ an annulus, $S = [0, 1]$, $A = \emptyset$, and $\mathfrak{r}_0 \equiv 1$. Thus $\mathfrak{B}, \mathfrak{C}$ are open sets in the Banach spaces $B(S)$, $C(S)$. If $\mathfrak{r} \in \mathfrak{C}$, define $f(\mathfrak{r}) = \sqrt{\mathfrak{r}(1)/\mathfrak{r}(0)}$, using the branch of the square root obtained by continuation from $\sqrt{1} = 1$ along the path $[0, 1] \ni t \mapsto \mathfrak{r}(t)/\mathfrak{r}(0)$. Let \mathbf{f} be the germ of f at \mathfrak{r}_0 . In a neighborhood of \mathfrak{r}_0 , we can write $f(\mathfrak{r}) = \sqrt{\phi(\mathfrak{r})/\psi(\mathfrak{r})}$ with holomorphic functions

$$(11.1) \quad \phi(\mathfrak{r}) = \mathfrak{r}(1), \quad \psi(\mathfrak{r}) = \mathfrak{r}(0)$$

on $C(S)$ and the principal branch of square root. Here ϕ, ψ are linear functions, and their extensions $\phi^{\text{ab}}, \psi^{\text{ab}}$ to $B(S)$ are given by the same formula (11.1). Since the Aron–Berner extension is compatible with multiplication, it follows that \mathbf{f}^{ab} is the germ of

$$g(\mathfrak{r}) = \sqrt{\mathfrak{r}(1)/\mathfrak{r}(0)}, \quad \mathfrak{r} \in \mathfrak{B} \text{ in a neighborhood of } \mathfrak{r}_0.$$

But \mathbf{f}^{ab} has no single valued continuation e.g. along the path $[0, 1] \ni t \rightarrow \mathfrak{r}_t \in \mathfrak{B}$,

$$\mathfrak{r}_t(s) = \begin{cases} e^{2\pi it} & \text{if } s = 1 \\ 1 & \text{if } s \neq 1. \end{cases}$$

Example 11.2. *There are mapping spaces $\mathfrak{B}, \mathfrak{C}$ and (unbounded) $f \in \mathcal{O}(\mathfrak{C})$ with germ \mathbf{f} at some $\mathfrak{r}_0 \in \mathfrak{C}$ such that \mathbf{f}^{ab} cannot be continued along some path in \mathfrak{B} .*

This is a minor reformulation of a result of Dineen. Let $X = \mathbb{C}$, $S = \{0, 1/n : n \in \mathbb{N}\} \subset \mathbb{R}$ with the topology inherited from \mathbb{R} , $A = \emptyset$, and $\mathfrak{r}_0 \equiv 0$. Thus \mathfrak{C} is biholomorphic to c , the space of convergent sequences of complex numbers, and $\mathfrak{B} \supset \mathfrak{C}$ is biholomorphic to $c'' = l^\infty \supset c$. In particular, \mathfrak{B} is simply connected, whence if a germ \mathbf{g} on it has analytic continuation along any path, then it is the germ of a holomorphic function on \mathfrak{B} . Since according to Dineen and Josefson [D71, J78] there are holomorphic functions f on c that do not extend to l^∞ , our claim follows. In fact, not only \mathbf{f}^{ab} cannot be continued analytically along some path, but no extension \mathbf{g} of \mathbf{f} to a germ on \mathfrak{B} has analytic continuation along all paths in \mathfrak{B} .

Example 11.3. (a) *Unless X is contractible, it has a mapping space \mathfrak{C} with a flat normed holomorphic Banach bundle $(E, p) \rightarrow \mathfrak{C}$, and there is an E valued holomorphic germ at some $\mathfrak{r}_0 \in \mathfrak{C}$ that has bounded analytic continuation along any path in \mathfrak{C} , yet does not extend to a holomorphic section of E . If X is simply connected, then E can be chosen a line bundle.*

(b) *If a Stein manifold X is not contractible, it has a corresponding mapping space \mathfrak{C} , and there is a \mathbb{C} valued holomorphic germ \mathbf{f} at some $\mathfrak{r}_0 \in \mathfrak{C}$ that continues analytically along any path in \mathfrak{C} , yet \mathbf{f} does not extend to a holomorphic function $\mathfrak{C} \rightarrow \mathbb{C}$.*

Both in (a) and (b) X can be simply connected. Thus (b) shows that in Theorem 10.1 the assumption that the analytic continuations are bounded cannot be simply dropped, and (a) shows that even boundedness is not sufficient unless it is known that (E, p) extends to a flat bundle over \mathfrak{B} .—If X is contractible, then its mapping spaces are also contractible, and the phenomena described in Example 11.3 will not occur.

If X is not contractible, according to Whitehead it has a nontrivial homotopy group $\pi_\nu(X, x_0)$, $\nu \geq 1$ [W49, Theorem 1]. Let $S = S^{\nu-1}$, $A \subset S$ a singleton, $\mathfrak{r}_0 \equiv x_0$, and $\mathfrak{C} = C(S, X, A, \mathfrak{r}_0)$. Since $\pi_1(\mathfrak{C}, \mathfrak{r}_0) \approx \pi_\nu(X, x_0) \neq 0$, there is a nontrivial covering $\pi : C \rightarrow \mathfrak{C}$ by a connected complex Banach manifold C . For $\mathfrak{r} \in \mathfrak{C}$ let $C_{\mathfrak{r}} = \pi^{-1}(\mathfrak{r})$, and let $E_{\mathfrak{r}}$ denote the Hilbert space of functions $e : C_{\mathfrak{r}} \rightarrow \mathbb{C}$ with l^2 norm $p(e) = (\sum_{v \in C_{\mathfrak{r}}} |e(v)|^2)^{1/2} < \infty$. On $E = \coprod_{\mathfrak{r} \in \mathfrak{C}} E_{\mathfrak{r}}$ the local trivializations of $C \rightarrow \mathfrak{C}$ induce the structure of a holomorphic Banach bundle over \mathfrak{C} , with flat norm p .

The pullback $\pi^*(E, p) \rightarrow C$ is trivial, and has a horizontal section g (Definition 6.4) whose value at $y \in C_{\mathfrak{r}}$ is the delta function $e \in E_{\mathfrak{r}}$ given by $e(y) = 1$, $e(v) = 0$ if $v \neq y$.

The germ \mathbf{g} of g at any point in $C_{\mathfrak{r}_0}$ pushes forward to a holomorphic E valued germ \mathbf{f} at \mathfrak{r}_0 ; the pushforwards of other germs of g provide bounded analytic continuations of \mathbf{f} along any path in \mathfrak{C} ; but \mathbf{f} is not the germ of a section f of E . If it were, g would be the pullback of f , and the singleton supports of the functions $f(\mathfrak{x}) : C_{\mathfrak{x}} \rightarrow \mathbb{C}$ would provide a global section of the covering $C \rightarrow \mathfrak{C}$, which is impossible.

If X is simply connected, then the above construction can be modified to produce a line bundle $E \rightarrow \mathfrak{C}$. Namely, $\nu \geq 2$ then, whence $G = \pi_1(\mathfrak{C}, \mathfrak{r}_0) \approx \pi_\nu(X, x_0)$ is Abelian. By Pontryagin duality it admits a nontrivial representation $\rho : G \rightarrow \mathbb{U}(1)$, [P66, Section 37]. This induces a flat line bundle $(E, p) \rightarrow \mathfrak{C}$, obtained from the trivial line bundle $\tilde{\mathfrak{C}} \times \mathbb{C} \rightarrow \tilde{\mathfrak{C}}$ over the universal cover of \mathfrak{C} , endowed with the trivial norm = Hermitian metric, by factoring out by the action

$$\tilde{\mathfrak{C}} \times \mathbb{C} \ni (\eta, \zeta) \mapsto (\gamma\eta, \rho(\gamma)\zeta) \in \tilde{\mathfrak{C}} \times \mathbb{C}, \quad \gamma \in G,$$

(G acts on $\tilde{\mathfrak{C}}$ by deck transformations). As ρ is nontrivial, (E, p) , as a flat line bundle, is nontrivial, either—although, as a holomorphic line bundle E may very well be trivial. The section $\tilde{\mathfrak{C}} \ni y \mapsto (y, 1) \in \tilde{\mathfrak{C}} \times \mathbb{C}$ induces a multivalued horizontal section of (E, p) , any of whose germs at \mathfrak{r}_0 is the \mathbf{f} sought.

A further modification is possible when X is Stein. In this case the universal cover $\tilde{\mathfrak{C}}$ can be embedded in a Banach space. To see this, we first embed X holomorphically and properly into some Euclidean space \mathbb{C}^N , and arrange that in the embedding x_0 is sent to 0. This exhibits \mathfrak{C} as a closed complex direct submanifold² of the Banach space $C_A(S)^{\oplus N}$. The latter space is isomorphic to a finite codimensional subspace of $C(S)^{\oplus N} \approx C[0, 1]^{\oplus N}$, hence has a Schauder basis, according to Miljutin and Schauder [M66, S27]. All this implies that $\tilde{\mathfrak{C}}$ can be embedded in a Banach space as a closed direct manifold. This is proved in [L10, Theorem 3.7], except the assumption there is that \mathfrak{C} should be embedded in a Banach space with unconditional basis. However, the unconditionality of the basis there was used only to be able to quote certain earlier results of cohomological nature by Patyi and by this author. All those results follow from cohomology vanishing in [LP07, Theorem 9.1], one of whose assumptions (“plurisubharmonic domination”) could at the time be proved only when unconditional bases existed. In the meantime, however, Patyi extended plurisubharmonic domination to spaces with a Schauder basis [P11, P15]. This means that embeddibility of the universal (or any) cover, [L10, Theorem 3.7], holds under the assumption of Schauder, rather than unconditional, basis. This takes care of the embedding of our $\tilde{\mathfrak{C}}$.

Once the embedding granted, for any pair of distinct points $a, b \in \tilde{\mathfrak{C}}$ there is a $g \in \mathcal{O}(\tilde{\mathfrak{C}})$ that takes different values at a, b . If we choose a, b both lie above \mathfrak{r}_0 , then the pushforward \mathbf{f} to \mathfrak{C} of the germ of g at a continues analytically along any path in \mathfrak{C} , but those continuations do not define a single valued function on \mathfrak{C} .

²A subset M of a complex Banach manifold N is a direct submanifold if for each $q \in N$ there are a neighborhood $U \subset N$ of q , a complemented subspace $A \subset B$ in a Banach space B , and an open $V \subset B$ such that the pairs $(U, U \cap M)$ and $(V, V \cap A)$ are biholomorphic.

References

- [AB78] Richard Aron, Paul Berner, A Hahn–Banach extension theorem for analytic mappings. *Bull. Soc. Math. France* 106 (1978) 3–24.
- [A⁺16] Richard Aron, Pablo Galindo, Damián Pinasco, Ignacio Zalduendo, Group-symmetric holomorphic functions on a Banach space. *Bull. Lond. Math. Soc.* 48 (2016) 779–796.
- [C99] Daniel Carando, Extendible polynomials on Banach spaces. *J. Math. Anal. Appl.* 233 (1999) 359–372.
- [C⁺08] Yun Sung Choi, Domingo Garcia, Sung Guen Kim, Manuel Maestre, Composition, numerical range and Aron–Berner extension. *Math. Scand.* 103 (2008) 97–110.
- [D⁺16] Garth Dales, Frederick Dashiell, Anthony Lau, Dona Strauss, Banach spaces of continuous functions as dual spaces. *CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC*. Springer, Cham, 2016.
- [DG89] Alexander Davie, Theodor Gamelin, A theorem on polynomial–star approximation. *Proc. Amer. Math. Soc.* 106 (1989) 351–356.
- [D71] Seán Dineen, Bounding subsets of a Banach space. *Math. Ann.* 192 (1971) 61–70.
- [D99] Seán Dineen, Complex analysis on infinite–dimensional spaces. *Springer Monographs in Mathematics*. Springer–Verlag London, Ltd., London, 1999.
- [DM98] Seán Dineen, Pauline Mellon, Holomorphic functions on symmetric Banach manifolds of compact type are constant. *Math. Z.* 229 (1998) 753–765.
- [DV04] Seán Dineen, Milena Venkova, Extending bounded type holomorphic mappings on a Banach space. Special issue dedicated to John Horváth. *J. Math. Anal. Appl.* 297 (2004) 645–658.
- [DS88] Nelson Dunford, Jacob Schwartz, Linear operators. Part I. General theory. With the assistance of William G. Bade and Robert G. Bartle. Reprint of the 1958 original. *Wiley Classics Library*. John Wiley & Sons, Inc., New York, 1988.
- [G66] Hugh Gordon, The maximal ideal space of a ring of measurable functions. *Amer. J. Math.* 88 (1966) 827–843.
- [G58] Hans Grauert, On Levi’s problem and imbedding of real analytic manifolds. *Ann. of Math(2)* 68 (1958) 460–472.
- [J78] Bengt Josefson, Bounding subsets of $l^\infty(A)$. *J. Math. Pures Appl. (9)* 57 (1978) 397–421.

- [K41] Shizuo Kakutani, Concrete representation of abstract (M) -spaces. (A characterization of the space of continuous functions.) *Ann. of Math. (2)* 42 (1941) 994–1024.
- [L98] László Lempert, The Dolbeault complex in infinite dimensions. I. *J. Amer. Math. Soc.* 11 (1998) 485–520.
- [L04] László Lempert, Holomorphic functions on (generalized) loop spaces. *Math. Proc. R. Ir. Acad.* 104A (2004) 35–46.
- [L10] László Lempert, Analytic continuation in mapping spaces. *Pure and Applied Math. Quart.* 6 (2010) 1051–1080.
- [LP07] László Lempert, Imre Patyi, Analytic sheaves in Banach spaces. *Ann. Sci. École Norm. Sup. (4)* 40 (2007) 453–486.
- [LS07] László Lempert, Endre Szabó, Rationally connected varieties and loop spaces. *Asian J. Math.* 11 (2007) 485–496.
- [M66] Alexei Miljutin, Isomorphism of the spaces of continuous functions over compact sets of the cardinality of the continuum. (Russian) *Teor. Funkcii Funkcional. Anal. i Priložen. Vyp. 2* (1966) 150–156.
- [M86] Jorge Mujica, Complex analysis in Banach spaces. Holomorphic functions and domains of holomorphy in finite and infinite dimensions. *North–Holland Mathematics Studies*, 120. *Notas de Matemática [Mathematical Notes]*, 107. North-Holland Publishing Co., Amsterdam, 1986.
- [P11] Imre Patyi, Plurisubharmonic domination in Banach spaces. *Adv. Math.* 227 (2011) 245–252.
- [P15] Imre Patyi, On holomorphic domination, II. *C. R. Math. Acad. Sci. Paris* 353 (2015) 501–503.
- [P66] Lev Pontryagin, Topological groups. Translated from the second Russian edition by Arlen Brown. Gordon and Breach Science Publishers, Inc., New York–London–Paris 1966
- [R87] Walter Rudin, Real and complex analysis. Third edition. McGraw-Hill Book Co., New York, 1987.
- [S27] Julius Schauder, Zur Theorie stetiger Abbildungen in Funktionalräumen. *Math. Z.* 26 (1927) 47–65.
- [U87] Ali Ülger, A. Weakly compact bilinear forms and Arens regularity. *Proc. Amer. Math. Soc.* 101 (1987) 697–704.
- [W49] J.C.H. Whitehead, Combinatorial homotopy I. *Bull. Amer. Math. Soc.* 55 (1949) 213–245.
- [Z90] Ignacio Zalduendo, A canonical extension for analytic functions on Banach spaces. *Trans. Amer. Math. Soc.* 320 (1990) 747–763.