

A dynamic risk score for early prediction of cardiogenic shock using machine learning

Yuxuan Hu¹, Albert Lui², Mark Goldstein³, Mukund Sudarshan³,
 Andrea Tinsay⁴, Cindy Tsui⁴, Samuel Maidman⁴, John Medamana⁴, Neil Jethani^{2,3},
 Aahlad Puli³, Vuthy Nguy⁵, Yindalon Aphinyanaphongs⁵, Nicholas Kiefer¹,
 Nathaniel Smilowitz¹, James Horowitz¹, Tania Ahuja⁶, Glenn I Fishman¹, Judith Hochman¹,
 Stuart Katz¹, Samuel Bernard¹, Rajesh Ranganath^{3,5,7}

¹ Leon. H. Charney Division of Cardiology, NYU Langone Health

² NYU Grossman School of Medicine

³ Courant Institute of Mathematics, New York University

⁴ Department of Medicine, NYU Langone Health

⁵ Department of Population Health, NYU Langone Health

⁶ Department of Pharmacy, NYU Langone Health

⁷ Center for Data Science, New York University

Abstract

Myocardial infarction and heart failure are major cardiovascular diseases that affect millions of people in the US. The morbidity and mortality are highest among patients who develop cardiogenic shock. Early recognition of cardiogenic shock is critical. Prompt implementation of treatment measures can prevent the deleterious spiral of ischemia, low blood pressure, and reduced cardiac output due to cardiogenic shock. However, early identification of cardiogenic shock has been challenging due to human providers' inability to process the enormous amount of data in the cardiac intensive care unit (ICU) and lack of an effective risk stratification tool. We developed a deep learning-based risk stratification tool, called CShock, for patients admitted into the cardiac ICU with acute decompensated heart failure and/or myocardial infarction to predict onset of cardiogenic shock. To develop and validate CShock, we annotated cardiac ICU datasets with physician adjudicated outcomes. CShock achieved an area under the receiver operator characteristic curve (AUROC) of 0.820, which substantially outperformed CardShock (AUROC 0.519), a well-established risk score for cardiogenic shock prognosis. CShock was externally validated in an independent patient cohort and achieved an AUROC of 0.800, demonstrating its generalizability in other cardiac ICUs.

Introduction

Cardiogenic shock is a life-threatening condition that is characterized by reduced cardiac output in the presence of adequate intravascular volume, resulting in tissue hypoxia. It complicates 5% to 12% of myocardial infarctions [1, 2, 3, 4] and 2% to 5% of acute decompensated heart failure [5, 6], the two most common etiologies for cardiogenic shock [7]. Mortality for cardiogenic shock remains high (30%-50%) [7, 8, 9, 10]. Early identification of cardiogenic shock can facilitate interventions that may mitigate the consequences of prolonged end-organ insult [11, 12], such as rapid employment of hemodynamic support with pharmacologic and nonpharmacologic agents, engagement of a dedicated shock team, and prompt transfer of patients at lower-acuity hospitals to a tertiary high-volume shock hub [3]. In the setting of acute myocardial infarction, early recognition of patients at high risk for cardiogenic shock allows providers to proceed with urgent revascularization of the infarct-related coronary artery [9]. Therefore, timely recognition of patients at high risk for cardiogenic shock is crucial for improving the care of patients and their outcomes.

systolic blood pressure less than 90mmHg for at least 30 minutes	AND	urine output less than 0.5cc/kg/hour for at least 6 hours or, increase in serum creatinine by 1.5 fold or 0.3mg/dl from baseline or, serum lactate above the upper limit of normal, which is > 2mmol/L
OR		
pharmacologic agents and/or mechanical circulatory support were initiated to maintain systolic blood pressure above 90mmHg		

Table 1: *Structured data used to determine time of shock onset*

Despite the potential benefits, early identification of cardiogenic shock has been challenging. Intensive care unit (ICU) providers are presented with tremendous amounts of data generated from multiple sources including laboratory measurements, vital signs, hemodynamics, and cardiac function studies. The limited ability of human providers to process, interpret, and act upon all the data stored in the electronic health record (EHR) in a timely fashion can lead to poor patient outcomes. Moreover, it is difficult to provide around-the-clock monitoring and risk assessment of cardiac ICU patients by human caregivers, especially in settings with low provider-to-patient ratios.

Furthermore, there lacks an effective risk stratification tool for development of cardiogenic shock. Current risk stratification strategies such as the Society for Cardiovascular Angiography and Intervention classification schema [13, 14] lack specific quantitative criteria, making it challenging to apply clinically. Well-established scores such as CardShock [15], IABP-shock II [16] and Acute Physiology and Chronic Health Evaluation (APACHE) II [17] were developed to predict outcomes such as in-hospital or 30-day mortality rather than the development of cardiogenic shock. Recent machine learning algorithms only classified patients who were initiated on inotropes and/or mechanical circulatory support as developing cardiogenic shock, but omitted those with low blood pressure and end-organ hypoperfusion [18, 19]. The best positive predictive value was 11% (with recall/sensitivity 27%) due to the low prevalence (1%) of cardiogenic shock patients in the study cohort [19], which would result in the algorithm having a high rate of false alarms, while also missing the majority of cardiogenic shock patients.

To improve risk stratification of cardiac critical care patients and provide early warning of cardiogenic shock for patients admitted into cardiac ICU with 1) acute decompensated heart failure and/or 2) myocardial infarction with or without ST-elevation, we develop a new dynamic risk score, called CShock, by training a machine learning model with a novel loss function designed specifically for risk scoring. We prepared a cardiac ICU dataset using MIMIC-III (Medical Information Mart for Intensive Care) database [20] by annotating with physician adjudicated outcomes and relevant features manually extracted from echocardiogram and catheterization reports. This dataset was then used to train CShock. We externally validated the risk model on the New York University (NYU) Langone Health cardiac ICU database that was also annotated with physician adjudicated outcomes.

Results

Preparation of a cardiac ICU dataset with physician adjudicated outcomes and cardiac features extracted from unstructured data

We used a publicly available critical care database MIMIC-III [20], which has 8188 admissions involving at least one cardiac ICU stay during the hospitalization (Fig. 1). 3220 admissions met the inclusion and exclusion criteria for chart review (Fig. 1): patients needed to have a possible diagnosis of acute decompensated heart failure and/or myocardial infarction with or without ST-elevation to be included; patients with age less than 18 years old or more than 89 years old, total hospital stay less than 24 hours, surgery ICU admission prior to cardiac ICU admission during the same hospitalization, shock on arrival to cardiac ICU were excluded (details in Methods, section Chart review). The discharge summary and echocardiogram reports

of the 3220 admissions were subsequently split among four physicians to review, which provided information regarding additional exclusion criteria, admission diagnoses, outcomes (no shock versus noncardiogenic shock only versus cardiogenic shock/mixed shock), left heart catheterization, and echocardiogram data (details in Methods, section Chart review). After applying the additional exclusion criteria (details in Methods, section Study cohort) obtained from chart review, the final study cohort comprised 1500 patients (Fig. 1). 1264 had no shock event before cardiac ICU discharge; 204 had cardiogenic/mixed shock; 32 had noncardiogenic shock only (Fig. 1). Patients’ outcomes were retrospectively adjudicated from chart review. Time of shock onset was determined as the earliest time after cardiac ICU admission when 1) systolic blood pressure was less than 90mmHg for at least 30 minutes and there was evidence of systemic hypoperfusion or 2) pharmacologic agents/mechanical circulatory support was initiated to maintain systolic blood pressure above 90mmHg (Table 1) [12]. If there was discrepancy between physician adjudicated outcomes and objective data, i.e. physician adjudicated outcome revealed no shock but there was shock based on structured data, a second physician reviewed both the discharge summary and the structured data to reconcile the discrepancy (details in Methods, section Chart review).




Figure 1: Study Cohort

Development of a dynamic risk score for prediction of cardiogenic shock

We developed CShock to determine the risk a patient has for developing cardiogenic shock after admission to the cardiac ICU. CShock is updated every hour after cardiac ICU admission. The features used to train the model for CShock included patient demographics, cardiac ICU admission diagnoses, routinely measured laboratory values and vital signs, and relevant features manually extracted from echocardiogram and left heart catheterization reports (Supplementary Information, List of features (Appendix A.3)).

The model that underlies CShock was based on a dilated causal convolutional neural network (CNN) architecture, often used for time series modeling (See Methods, Model development for more details on model architecture). There were 194 input variables, including 182 time-varying (physiological) and 12 static (e.g., demographic) features. The physiological time series of the 182 time-varying features from cardiac ICU admission to event (discharge if no shock; shock onset otherwise) were inputted into the CShock model. The

time-dependent variables also had missing indicators that took a value of 1 if the corresponding feature value was missing at each unique hour after cardiac ICU admission; 0 otherwise.

We designed a novel loss function specifically for risk scoring shown in Eq. (6) of the Supplementary Information. At a high level, optimizing the loss increases the soft-maximum risk across time for patients who developed cardiogenic shock during their stay while minimizing for those who did not develop cardiogenic shock. The study cohort (1500 patients) were split 50%, 25%, and 25% into training, validation, and testing in a 4-fold cross validation. To improve performance, the model was pretrained in an auxiliary task of predicting in-hospital mortality for ICU patients in MIMIC-III database.

The left panel of Fig. 2 demonstrates the physiological time series of five example features that were entered into the dilated CNN-based model, which then outputted the CShock score as shown by the blue lines in the right panel. The dashed blue line is the CShock score for a patient who develops no shock and gets discharged from the cardiac ICU at hour 37; the solid blue line shows the CShock score for a patient who goes into mixed cardiogenic/noncardiogenic shock at hour 32. Cardiogenic shock detection occurs if a patient’s CShock score exceeds the alarm threshold value. We followed MINIMAR (MINimum Information for Medical AI Reporting) [21] in reporting the results.




Figure 2: The physiological time series of five example features from cardiac ICU admission to event (discharge if no shock; shock onset otherwise) are shown for a patient developing mixed cardiogenic/noncardiogenic shock (left panel). Features displayed are Glasgow Coma Scale (a score for coma severity, see Appendix A.3 for the three components of the score), creatinine, lactate, heart rate and systolic blood pressure without support. The physiological time series were fed into the CNN-based model, which outputted CShock scores. CShock scores (blue lines in right panel) were calculated for each hour from cardiac ICU admission until event. Two patients’ CShock scores are shown here: one patient develops no shock and gets discharged from cardiac ICU at hour 37 (dashed blue line); the other patient goes into mixed shock at hour 32 (solid blue line). Cardiogenic shock detection occurs if a patient’s CShock score exceeds the alarm threshold value. The horizontal red line indicates the detection threshold corresponding to a sensitivity of 0.8.

Cardiogenic shock early warning system

We developed the CShock score for identifying patients at risk for cardiogenic shock before its overt onset. A patient was considered positive if the patient developed cardiogenic or mixed shock during their cardiac ICU stay; 204 patients were considered positive (Fig. 1). The 1296 patients who developed noncardiogenic shock only or no shock in the study cohort were treated as negative cases. The CShock score was able to achieve an area under the receiver operator characteristic curve (AUROC) of 0.820 (Fig. 3). The receiving operator characteristic (ROC) curve and its corresponding area under curve (AUC) were obtained by varying the alarm threshold that determined which patients were identified by the model as at risk for cardiogenic shock. At a positive predictive value of 0.33, CShock achieved a recall of 0.67 and was able to predict cardiogenic onset, on average, 36.5 hours before onset.

Current cardiogenic shock scores are mostly for outcome prediction once a diagnosis of cardiogenic shock has been made [22]. We decided to use the CardShock score [15] to compare its performance to CShock. Even though the CardShock score was not calibrated for our study objective, it is a widely accepted score for cardiogenic shock and was derived from a cohort of both acute coronary syndrome (ACS) and non-ACS etiology, similar to the population in our study. The AUROC for predicting development of cardiogenic shock using the CardShock score was 0.519 (Fig. 3). The basis for the poor performance of the CardShock score for this purpose, especially when compared to that for in-hospital mortality [15], could be multifold: 1) history of myocardial infarction (a variable in the score calculation) is not available for the study cohort and thus the CardShock score was underestimated for patients with a history of myocardial infarction 2) the score is meant for prognosis once the diagnosis of cardiogenic shock has been made and not for risk stratifying patients for development of cardiogenic shock, i.e. variables such as low glomerular filtration rate are associated with worse mortality outcome but are not necessarily the most predictive for development of cardiogenic shock.




Figure 3: ROC for predicting cardiogenic shock using the developed CShock score (blue) versus CardShock Score (orange)

Model performance in different subgroups

We performed subgroup analysis based on age and gender (Table 2). The model performed better for the younger group than the older group. This could be due to the fact that the elderly tend to have more comorbidities that were embedded only in the unstructured data of the EHR (e.g. clinical notes) and thus unaccounted for in the model. We also found that the model performed better for male patients (0.817)

		AUROC
Age (years)	below 25 percentile (less than 59)	0.855±0.028
	between 25-50 percentile (at least 59, less than 70)	0.801±0.060
	between 50-75 percentile (at least 70, less than 80)	0.775±0.035
	above 75 percentile (at least 80)	0.745±0.039
Gender	Female	0.756±0.038
	Male	0.817±0.026

Table 2: AUROC for each age group and gender

compared to female patients (0.756). This was possibly a result of the study cohort having more male patients (918) as compared to female patients (582) and thus the model was better trained for the male cohort.

External validation


The model was externally validated using a dataset that comprised all patients admitted to the cardiac ICU at NYU Langone Health who met the same inclusion and exclusion criteria as the study cohort (Fig. S1). Patients’ outcomes in the external validation cohort were retrospectively adjudicated by a physician. 58 of the 182 time-varying input features in the original model are routinely measured in the EHR at NYU Langone Health. Therefore, we developed the reduced model by training with 70 input features (58 were time-varying, 12 static) using MIMIC-III database (list of features is available in the Supplementary Information (Appendix A.3), denoted as included in the reduced model). The reduced model was able to achieve similar AUROC at 0.802 on the study cohort. It was then externally validated using the NYU Langone Health cardiac ICU dataset and obtained an AUROC of 0.800, demonstrating its generalizability and applicability in other cardiac ICUs.

Model interpretation

We used Shapley values [23] to provide explanations for how each feature influenced predictions. We employed FastSHAP [24], an efficient algorithm for calculating Shapley values [23]. The Shapley values were based on a FastSHAP’s surrogate model that was trained to produce CShock risk scores given different feature subsets by masked prediction [25, 26] rather than model the covariates [27] (See Appendix A.2 for more details on model interpretation). Shapley values were determined for the 70 features in the reduced model using the training dataset. Figure 4a presents the ten most important features, i.e. with the largest Shapley values; Fig. 4b shows how the AUROC changes as more features are added in descending order of importance into the FastSHAP surrogate model. AUROC improved as more features were included in the model and the best AUROCs were similar to that obtained using the reduced model. With only the 10 most important features (as displayed in Fig. 4a), FastSHAP surrogate model was able to achieve AUROC (yellow line in Fig. 4b) similar to that when more features were included.

Returning to Fig. 4a, each dot represents the feature values averaged over time of an individual patient from the training dataset with a higher value being more red and a lower value being more blue. Positive and negative Shapley values are associated with an increase or decrease in the CShock score, i.e. risk of cardiogenic shock development, respectively. For instance, heart rate is the most predictive feature of cardiogenic shock development and having an elevated heart rate is associated with an increased risk of cardiogenic shock development. This model interpretation analysis indicated that an admission diagnosis of myocardial infarction with ST-segment elevation is associated with lower risk of development of cardiogenic shock, whereas having an admission diagnosis of acute decompensated heart failure is associated with higher risk of development of cardiogenic shock. This likely reflects advancements in ACS management strategies over the years, which makes ACS patients less likely to develop cardiogenic shock. Among the 10 most important features, low Braden Scale (a risk score for identifying patients at risk for pressure ulcers [28], see Appendix A.3 for the six variables in Braden Scale), Glasgow Coma Scale, systolic blood pressure without

(a)



(b)




Figure 4: Fig. 4a shows the 10 most important features based on Shapley values in descending order of importance (1 Heart rate, 2 having an admission diagnosis of myocardial infarction with ST-segment elevation, 3 having an admission diagnosis of acute decompensated heart failure, 4 Braden Scale, 5 Glasgow Coma Scale, 6 Blood urea nitrogen, 7 Systolic blood pressure, 8 Serum chloride, 9 Serum sodium, 10 Arterial blood pH). Each dot represents the feature values averaged over time of an individual patient from the training dataset with a higher value being more red and a lower value being more blue. Positive and negative Shapley values are associated with an increase or decrease in risk of cardiogenic shock development, respectively. Fig. 4b AUROC obtained with increasing number of features included in the FastSHAP surrogate model using the evaluation dataset. Yellow line indicates the AUROC when only the 10 most important features (as displayed in Fig. 4a) were included in the FastSHAP surrogate model that was trained to predict on subsets of features.

support and serum sodium are predictive of cardiogenic shock development, as expected clinically [29, 30, 14]. Lactate, a commonly used lab measurement to assess for severity of cardiogenic shock, is not among the 10 most important features; conceivably other top features captured most of the predictive information that would be embedded in lactate.

Discussion

Myocardial infarction and heart failure affect 9 and 6 million people in the US respectively [31]. Cardiogenic shock is a common complication of myocardial infarction and heart failure and its occurrence is associated with substantial health and economic costs [1, 6]. Early recognition of cardiogenic shock is of paramount importance as it facilitates timely treatments that could potentially prevent the vicious spiral of cardiogenic shock and reduce the health and financial burden. However, early identification of cardiogenic shock has remained challenging [11]. We have demonstrated that the CShock score can predict cardiogenic shock with excellent AUROC. The early-warning system was tested in an independent patient cohort from a different hospital and showed comparable performance as in the development data.

Our CShock risk model is readily deployable. The reduced model with 70 features that are routinely captured in the EHR database at NYU Langone Health achieved an AUROC of 0.800. This model should be easily implemented at other facilities with modern EHR systems. It is interesting to note that the reduced model performed on par with the full model despite having fewer features. This could be because 1) it is difficult to manually extract features from unstructured data (e.g. echocardiogram and catheterization data) and the lack of data can jeopardize the performance of the machine learning model; it is possible if those

features were available as structured data, the performance of the full model would be even better. 2) The structured data captured most of the predictive information that would be embedded in unstructured data; incorporating unstructured data will not improve the performance of the machine learning model beyond what has been achieved with structured data only.

Previous machine learning studies relied on International Classification of Diseases (ICD) codes and natural language processing of clinical notes [32, 33] to determine outcomes such as sepsis and septic shock, which are often inaccurate [34]. In contrast, the outcome events for the cardiac ICU patients were carefully adjudicated by physicians in this study. This will allow the CShock score developed by this project to translate more easily into clinical implementation.

We constructed a novel loss function specifically for risk scoring. The dilated causal CNN architecture along with the novel loss function are of broad applicability and can be employed for risk stratification in multiple clinical settings such as sepsis and pulmonary embolism. Physiological time series models have not widely employed pretraining, a common technique used to improve model performance in deep learning [35, 36]. Pretraining is useful when there is access to a large dataset labeled for a task related to—but not identical to—the task of interest. In this case, we showed that pretraining with an auxiliary task of predicting mortality for ICU patients in MIMIC-III database improved the AUROC of the cardiogenic shock model from 0.752 to 0.820. We demonstrated that pretraining with a generic label such as mortality can be a useful strategy to improve physiological time series model performance when there is limited number of patients who meet all the inclusion and exclusion criteria of the study.

Shapley values are widely used to provide explanation for ‘blackbox’ models. However, they are computationally costly and are impractical to implement when large number of parameters are involved. We successfully implemented FastSHAP [24], a novel and efficient algorithm to estimate Shapley values to provide understanding about how each feature influenced cardiogenic shock prediction. Given the popularity of deep learning models in medicine and the ‘blackbox’ nature of these models, we demonstrated an efficient yet effective way to uncover important facets of the deep learning models.

There have been other recent efforts to predict cardiogenic shock with machine learning [18, 19]. However, they only considered patients who required inotropes/mechanical circulatory support as developing cardiogenic shock and used the time to initiate supportive measures as shock onset, which are not in alignment with the clinical criteria used in previous landmark trials [9, 37]; the early presentation of cardiogenic shock with low blood pressure and end-organ hypoperfusion would be missed by their algorithms. In addition, both studies lacked external validation to demonstrate generalizability in other populations. 1% of the study cohort developed cardiogenic shock in the study by Rahman et al. [19], which resulted in the best positive predictive value being only 11% (with recall/sensitivity 27%) and would lead to the algorithm sounding many false alarms secondary to extreme class imbalance [38] and missing the majority of cardiogenic shock patients. The study by Chang et al. [18] relied on ICD codes to determine outcomes, which would be inaccurate as discussed above; they excluded mixed cardiogenic/noncardiogenic shock patients from the study cohort, which could limit the algorithm’s applicability.

Given the performance of our CShock score, we hypothesize that this CNN-based early-warning model can help cardiac ICU teams clinically integrate complex data and more rapidly identify patients at risk for cardiogenic shock. A recent study using the Critical Care Cardiology Trials Network data showed that a multidisciplinary shock team approach improved outcomes in cardiogenic shock [11] and this machine learning based model could build upon this improvement, alerting shock teams to impending cardiogenic shock. Future prospective studies should be conducted to evaluate the generalizability of our findings across diverse health care systems and patient populations. In summary, we demonstrate that the CShock score we developed has the potential to provide automated detection and early warning for cardiogenic shock and improve the outcomes for the millions of patients who suffer from myocardial infarction and heart failure.

Acknowledgements

YH was supported by a NIH T32 grant (T32HL098129). RR was supported by NIH/NHLBI Award R01HL148248, NSF Award 1922658 NRT-HDR: FUTURE Foundations, Translation, and Responsibility for Data Science, NSF CAREER Award 2145542.

References

- [1] Robert J Goldberg, Frederick A Spencer, Joel M Gore, Darleen Lessard, and Jorge Yarzebski. Thirty-year trends (1975 to 2005) in the magnitude of, management of, and hospital death rates associated with cardiogenic shock in patients with acute myocardial infarction: a population-based perspective. *Circulation*, 119(9):1211–1219, 2009.
- [2] Dhaval Kolte, Sahil Khera, Wilbert S Aronow, Marjan Mujib, Chandrasekar Palaniswamy, Sachin Sule, Diwakar Jain, William Gotsis, Ali Ahmed, William H Frishman, et al. Trends in incidence, management, and outcomes of cardiogenic shock complicating st-elevation myocardial infarction in the u nited s tates. *Journal of the American Heart Association*, 3(1):e000590, 2014.
- [3] Sean Van Diepen, Jason N Katz, Nancy M Albert, Timothy D Henry, Alice K Jacobs, Navin K Kapur, Ahmet Kilic, Venu Menon, E Magnus Ohman, Nancy K Sweitzer, et al. Contemporary management of cardiogenic shock: a scientific statement from the american heart association. *Circulation*, 136(16):e232–e268, 2017.
- [4] Siddharth A Wayangankar, Sripal Bangalore, Lisa A McCoy, Hani Jneid, Faisal Latif, Wassef Karrowni, Konstantinos Charitakis, Dmitriy N Feldman, Habib A Dakik, Laura Mauri, et al. Temporal trends and outcomes of patients undergoing percutaneous coronary interventions for cardiogenic shock in the setting of acute myocardial infarction: a report from the cathpci registry. *Cardiovascular Interventions*, 9(4):341–351, 2016.
- [5] Ovidiu Chioncel, John Parissis, Alexandre Mebazaa, Holger Thiele, Steffen Desch, Johann Bauersachs, Veli-Pekka Harjola, Elena-Laura Antohi, Mattia Arrigo, Tuvia Ben Gal, et al. Epidemiology, pathophysiology and contemporary management of cardiogenic shock—a position statement from the heart failure association of the european society of cardiology. *European journal of heart failure*, 22(8):1315–1341, 2020.
- [6] Xavier Rossello, Héctor Bueno, Víctor Gil, Javier Jacob, Francisco Javier Martín-Sánchez, Pere Llorens, Pablo Herrero Puente, Aitor Alquézar-Arbé, Begoña Espinosa, Sergio Raposeiras-Roubín, et al. Synergistic impact of systolic blood pressure and perfusion status on mortality in acute heart failure. *Circulation: Heart Failure*, 14(3):e007347, 2021.
- [7] David D Berg, Erin A Bohula, Sean Van Diepen, Jason N Katz, Carlos L Alviar, Vivian M Baird-Zars, Christopher F Barnett, Gregory W Barsness, James A Burke, Paul C Cremer, et al. Epidemiology of shock in contemporary cardiac intensive care units: data from the critical care cardiology trials network registry. *Circulation: Cardiovascular Quality and Outcomes*, 12(3):e005618, 2019.
- [8] Judith S Hochman. Cardiogenic shock complicating acute myocardial infarction: expanding the paradigm. *Circulation*, 107(24):2998–3002, 2003.
- [9] Judith S Hochman, Lynn A Sleeper, John G Webb, Timothy A Sanborn, Harvey D White, J David Talley, Christopher E Buller, Alice K Jacobs, James N Slater, Jacques Col, et al. Early revascularization in acute myocardial infarction complicated by cardiogenic shock. *New England Journal of Medicine*, 341(9):625–634, 1999.
- [10] Patrick R Lawler, Erin Ann Bohula, Jeong-Gun Park, Jason Katz, Sean Van Diepen, David A Morrow, and Critical Care Cardiology Trials Network (CCCTN) Registry. Clinical staging in cardiogenic shock discriminates survival: data from the critical care cardiology trials network (ccctn) registry. *Journal of the American College of Cardiology*, 75(11_Supplement_1):1127–1127, 2020.

- [11] Alexander I Papolos, Benjamin B Kenigsberg, David D Berg, Carlos L Alviar, Erin Bohula, James A Burke, Anthony P Carnicelli, Sunit-Preet Chaudhry, Stavros Drakos, Daniel A Gerber, et al. Management and outcomes of cardiogenic shock in cardiac icus with versus without shock teams. *Journal of the American College of Cardiology*, 78(13):1309–1317, 2021.
- [12] Alex Reyentovich, Maya H Barghash, and Judith S Hochman. Management of refractory cardiogenic shock. *Nature Reviews Cardiology*, 13(8):481–492, 2016.
- [13] David A Baran, Cindy L Grines, Steven Bailey, Daniel Burkhoff, Shelley A Hall, Timothy D Henry, Steven M Hollenberg, Navin K Kapur, William O’Neill, Joseph P Ornato, et al. Scai clinical expert consensus statement on the classification of cardiogenic shock: this document was endorsed by the american college of cardiology (acc), the american heart association (aha), the society of critical care medicine (sccm), and the society of thoracic surgeons (sts) in april 2019. *Catheterization and Cardiovascular Interventions*, 94(1):29–37, 2019.
- [14] Srihari S Naidu, David A Baran, Jacob C Jentzer, Steven M Hollenberg, Sean van Diepen, Mir B Basir, Cindy L Grines, Deborah B Diercks, Shelley Hall, Navin K Kapur, et al. Scai shock stage classification expert consensus update: A review and incorporation of validation studies: This statement was endorsed by the american college of cardiology (acc), american college of emergency physicians (acep), american heart association (aha), european society of cardiology (esc) association for acute cardiovascular care (acvc), international society for heart and lung transplantation (ishlt), society of critical care medicine (sccm), and society of thoracic surgeons (sts) in december 2021. *Journal of the American College of Cardiology*, 79(9):933–946, 2022.
- [15] Veli-Pekka Harjola, Johan Lassus, Alessandro Sionis, Lars Køber, Tuukka Tarvasmäki, Jindrich Spinar, John Parissis, Marek Banaszewski, Jose Silva-Cardoso, Valentina Carubelli, et al. Clinical picture and risk prediction of short-term mortality in cardiogenic shock. *European journal of heart failure*, 17(5):501–509, 2015.
- [16] Janine Pöss, Jelena Köster, Georg Fuernau, Ingo Eitel, Suzanne de Waha, Taoufik Ouarrak, Johan Lassus, Veli-Pekka Harjola, Uwe Zeymer, Holger Thiele, et al. Risk stratification for patients in cardiogenic shock after acute myocardial infarction. *Journal of the American College of Cardiology*, 69(15):1913–1920, 2017.
- [17] William A Knaus, Elizabeth A Draper, Douglas P Wagner, and Jack E Zimmerman. Apache ii: a severity of disease classification system. *Critical care medicine*, 13(10):818–829, 1985.
- [18] Yale Chang, Corneliu Antonescu, Shreyas Ravindranath, Junzi Dong, Mingyu Lu, Francesco Vicario, Lisa Wondrely, Pam Thompson, Dennis Swearingen, and Deepak Acharya. Early prediction of cardiogenic shock using machine learning. *Frontiers in cardiovascular medicine*, page 1868, 2022.
- [19] Faisal Rahman, Noam Finkelstein, Anton Alyakin, Nisha A Gilotra, Jeff Trost, Steven P Schulman, and Suchi Saria. Using machine learning for early prediction of cardiogenic shock in patients with acute heart failure. *Journal of the Society for Cardiovascular Angiography & Interventions*, 1(3):100308, 2022.
- [20] Alistair EW Johnson, Tom J Pollard, Lu Shen, Li-wei H Lehman, Mengling Feng, Mohammad Ghassemi, Benjamin Moody, Peter Szolovits, Leo Anthony Celi, and Roger G Mark. MIMIC-III, a freely accessible critical care database. *Scientific data*, 3(1):1–9, 2016.
- [21] Tina Hernandez-Boussard, Selen Bozkurt, John PA Ioannidis, and Nigam H Shah. Minimar (minimum information for medical ai reporting): developing reporting standards for artificial intelligence in health care. *Journal of the American Medical Informatics Association*, 27(12):2011–2015, 2020.
- [22] Sanjog Kalra, Lauren S Ranard, Sehrish Memon, Prashant Rao, A Reshad Garan, Amirali Masoumi, WILLIAM O’NEILL, Navin K Kapur, Dimitri Karpaliotis, Justin A Fried, et al. Risk prediction in cardiogenic shock: current state of knowledge, challenges and opportunities. *Journal of cardiac failure*, 27(10):1099–1110, 2021.

- [23] Scott M Lundberg and Su-In Lee. A unified approach to interpreting model predictions. *Advances in neural information processing systems*, 30, 2017.
- [24] Neil Jethani, Mukund Sudarshan, Ian Connick Covert, Su-In Lee, and Rajesh Ranganath. Fastshap: Real-time shapley value estimation. In *International Conference on Learning Representations*, 2021.
- [25] Ian Covert, Scott M Lundberg, and Su-In Lee. Explaining by removing: A unified framework for model explanation. *J. Mach. Learn. Res.*, 22:209–1, 2021.
- [26] Neil Jethani, Mukund Sudarshan, Yindalon Aphinyanaphongs, and Rajesh Ranganath. Have we learned to explain?: How interpretability methods can learn to encode predictions in their interpretations. In *International Conference on Artificial Intelligence and Statistics*, pages 1459–1467. PMLR, 2021.
- [27] Mukund Sudarshan, Wesley Tansey, and Rajesh Ranganath. Deep direct likelihood knockoffs. *Advances in neural information processing systems*, 33:5036–5046, 2020.
- [28] Nancy Bergstrom et al. The braden scale for predicting pressure sore risk. *Nurs res*, 36(4):205–210, 1987.
- [29] Anu Kataja, Tuukka Tarvasmäki, Johan Lassus, Lars Køber, Alessandro Sionis, Jindrich Spinar, John Parissis, Valentina Carubelli, Jose Cardoso, Marek Banaszewski, et al. Altered mental status predicts mortality in cardiogenic shock—results from the cardshock study. *European Heart Journal: Acute Cardiovascular Care*, 7(1):38–44, 2018.
- [30] Carl V Leier, Livio Dei Cas, and Marco Metra. Clinical relevance and management of the major electrolyte abnormalities in congestive heart failure: hyponatremia, hypokalemia, and hypomagnesemia. *American heart journal*, 128(3):564–574, 1994.
- [31] Salim S Virani, Alvaro Alonso, Hugo J Aparicio, Emelia J Benjamin, Marcio S Bittencourt, Clifton W Callaway, April P Carson, Alanna M Chamberlain, Susan Cheng, Francesca N Delling, et al. Heart disease and stroke statistics—2021 update: a report from the american heart association. *Circulation*, 143(8):e254–e743, 2021.
- [32] Katharine E Henry, David N Hager, Peter J Pronovost, and Suchi Saria. A targeted real-time early warning score (trewscore) for septic shock. *Science translational medicine*, 7(299):299ra122–299ra122, 2015.
- [33] Ran Liu, Joseph L Greenstein, Stephen J Granite, James C Fackler, Melania M Bembea, Sridevi V Sarma, and Raimond L Winslow. Data-driven discovery of a novel sepsis pre-shock state predicts impending septic shock in the icu. *Scientific reports*, 9(1):1–9, 2019.
- [34] Benoît Misset, Didier Nakache, Aurélien Vesin, Mickael Darmon, Maïté Garrouste-Orgeas, Bruno Mourvillier, Christophe Adrie, Sébastien Pease, Marie-Aliette Costa de Beauregard, Dany Goldgran-Toledano, et al. Reliability of diagnostic coding in intensive care patients. *Critical care*, 12(4):1–7, 2008.
- [35] Jacob Devlin, Ming-Wei Chang, Kenton Lee, and Kristina Toutanova. Bert: Pre-training of deep bidirectional transformers for language understanding. *arXiv preprint arXiv:1810.04805*, 2018.
- [36] Matthew McDermott, Bret Nestor, Evan Kim, Wancong Zhang, Anna Goldenberg, Peter Szolovits, and Marzyeh Ghassemi. A comprehensive ehr timeseries pre-training benchmark. In *Proceedings of the Conference on Health, Inference, and Learning*, pages 257–278, 2021.
- [37] Holger Thiele, Uwe Zeymer, Franz-Josef Neumann, Mirosław Ferenc, Hans-Georg Olbrich, Jörg Hausleiter, Gert Richardt, Marcus Hennersdorf, Klaus Empen, Georg Fuernau, et al. Intraaortic balloon support for myocardial infarction with cardiogenic shock. *New England Journal of Medicine*, 367(14):1287–1296, 2012.
- [38] Yuzhe Yang and Zhi Xu. Rethinking the value of labels for improving class-imbalanced learning. *Advances in neural information processing systems*, 33:19290–19301, 2020.

Methods

Database and data extraction

We used a publicly available critical care database MIMIC-III, a single-center database that consists of patients admitted to critical care units at a large tertiary care hospital [20]. Data include vital signs, medications, laboratory measurements, notes by providers, and fluid balance. This database has been widely used to develop machine learning algorithms for risk stratifying critical care patients for sepsis and septic shock [32, 33]. EHR data was extracted from the MIMIC-III PostgreSQL database. The majority of data entries in MIMIC-III are comprised of timestamp-value pairs with a subject id, identifying which patient the data belong to, and an itemid identifying the meaning of the data.

Chart review

There are 8288 admissions in the MIMIC-III database that involve at least one cardiac ICU stay during the hospitalization (Fig. 1). To be eligible for chart review, the admissions needed to have possible diagnosis of 1) acute decompensated heart failure and/or 2) myocardial infarction with or without ST-elevation. We used the following criteria to identify those admissions: 1) positive troponin or 2) International Classification of Diseases, 9th Revision (ICD-9) or 3) received intravenous diuretics such as furosemide or bumetanide or 4) ICD-9 code for heart failure. The admissions that met the following criteria (based on structured data extracted from MIMIC-III database, Fig. 1) were excluded from chart review: age less than 18 years old or more than 89 years old, total hospital stay less than 24 hours, surgery ICU admission prior to cardiac ICU admission during the same hospitalization, patients who presented in shock on arrival to cardiac ICU. 3220 admissions met the above inclusion and exclusion criteria and the discharge summaries and echocardiogram reports were obtained and split between four physicians to review. Research electronic data capture (REDCap) database was set up for the chart review process.

Data obtained from chart review of discharge summaries provided the following information: whether patients met certain exclusion criteria as detailed below in section Study Cohort, cardiac ICU admission diagnoses 1) myocardial infarction with or without ST elevation and/or 2) acute decompensated heart failure, whether shock occurred during first cardiac ICU stay and if occurred, the etiology of the first shock (cardiogenic versus non cardiogenic versus mixed) and whether left heart catheterization was performed before the first cardiac ICU discharge and if performed, the locations of severe lesions, culprit lesion and any intervention undertaken.

Patients' outcomes (no shock versus noncardiogenic shock only versus cardiogenic shock/mixed shock) were adjudicated by a team of physicians from chart review of discharge summary. Time of shock onset was determined as the earliest time after cardiac ICU admission when the conditions in Table 1 were satisfied [12], as previously described in Results. If there was discrepancy between physician adjudicated outcomes and objective data, a second physician reviewed both the discharge summary and the structured data to reconcile the discrepancy. This could happen when there was a single low blood pressure reading and patient had acute kidney injury from recent contrast use in left heart catheterization. Based on structured data, the patient would be labeled as having a shock event but upon review of discharge summary, the patient had no shock event.

Data obtained from chart review of echocardiogram reports yielded information regarding left ventricular ejection fraction, right ventricular systolic function, right ventricular dilation, right ventricular systolic pressure, significant left sided valvular disease and presence of left ventricular aneurysm or pseudoaneurysm.

Study cohort

Based on information obtained from chart review of discharge summaries, the admissions that met the following criteria were excluded (Fig. 1): development of acute respiratory distress syndrome, massive pulmonary embolism, cardiac tamponade and mechanical complications of myocardial infarction, urgent coronary-artery bypass surgery, cardiac surgery in the 7 days prior to first cardiac ICU admission, cardiac arrest in the 7

days prior to first cardiac ICU admission and admission to cardiac ICU for diagnosis other than myocardial infarction with or without ST-elevation or acute decompensated heart failure. Patients who developed shock within 4 hours of cardiac ICU admission were also excluded due to the assumption that the patient was likely in a peri-shock state at the time of admission.

Our final study cohort comprised 1500 admissions (Fig. 1): 1264 patients did not develop shock; 204 patients developed cardiogenic or mixed shock and 32 patients developed noncardiogenic shock only.

Model development

Features. There were a total of 194 input features (182 were time-varying) that were either derived from routinely available measurements in the EHR or from chart review of discharge summaries and echocardiogram reports. The complete list of features is provided in the Supplementary Information (Appendix A.3). All features were standardized (zero mean, one standard deviation) before feeding into the model. The 182 time-varying variables had missing indicators, as described previously in Results. Missing values were imputed using population mean.

Model architecture. The model was based on dilated causal CNN, which has been shown to be easier to train compared to recurrent neural network (RNN) in time series modeling [39]. The model architecture consisted of 4 dilated causal convolution layers with each followed by a rectified linear unit (ReLU) activation [40], batch normalization, and dropout. The output of the dilated causal CNN captured the patient’s overall physiological state at each hour. This output was followed by a fully connected layer and sigmoid function to produce the time-varying CShock risk score for cardiogenic shock.

Training. We used the Adam [41] optimizer with a learning rate of 1×10^{-3} and batch size was set to 256. We evaluated loss (see Eq. (6) in the Supplementary Information) on the validation set for each epoch. Training was terminated after 50 epochs, which was shown to be sufficient for loss to plateau. The model was implemented using Pytorch. Hyperparameter optimization and model selection were conducted using 4-fold cross validation on the study cohort (1500 patients were split 50%, 25%, and 25% into training, validation, and testing). The training set was balanced for the minority class. To improve the performance of the model, we pretrained the model for an auxiliary task of predicting in-hospital mortality for ICU patients. The pretraining cohort included all patients admitted to cardiac and medical ICU in MIMIC-III database, excluding the study cohort used in developing CShock risk model. Other exclusion criteria for pretraining cohort included patients with age less than 18 years old or more than 89 years old, total hospital stay less than 24 hours, and surgery ICU admission during the same hospitalization. We used the pretrained weights to initialize the CShock model parameters. In addition, we tried several forms of data augmentation and contrastive learning, but we did not include them in the final model as their inclusion gave similar performance at the expense of longer run time.

Model validation. The best model was selected as the one achieving the best AUROC on validation set. Performance was then evaluated using the model on the testing set. The AUROC on test set across 4 folds was averaged and reported.

External validation

Compliance with ethical regulations

We followed NYU Langone Health human research protections policy, and completed the NYU self-certification form for research activities that may be classified as quality improvement. The work involving external validation met the NYU Grossman School of Medicine Institutional Review Boards (IRB) criteria for quality improvement work, not research involving human subjects, and thus this work did not require IRB review and no informed consent was required or obtained.

The model was externally validated using a dataset that comprised all patients who met the same inclusion and exclusion criteria as described in section Study cohort admitted to the cardiac ICU at NYU

Langone Health. The external validation dataset consisted of 131 patients with 25 patients experiencing cardiogenic/mixed shock (Fig. S1). The average age of the external validation cohort was 64.5 ± 14.9 . There were 85 male and 46 female patients. The average age of the study cohort in MIMIC-III was 68.1 ± 13.5 . There were 918 male and 582 female patients in MIMIC-III study cohort. Race, ethnicity and socioeconomic status were not provided for the study cohort and the external validation cohort. 58 of the 182 time-varying input features in the original model are routinely measured in the EHR at NYU Langone Health and therefore, the model was retrained with 70 input features including 58 time-varying features using MIMIC-III database (a list of features in the reduced model is available in the Supplementary Information, (Appendix A.3)). This model with reduced number of features was then externally validated using the dataset from the cardiac ICU at NYU Langone Health. The admission diagnoses and outcomes (cardiogenic/mixed shock versus noncardiogenic shock only versus no shock) for these patients were obtained in the same manner as those for the study cohort.

Data availability

The full MIMIC dataset can be downloaded from <http://physionet.org>. The labels for cardiogenic shock will be made available on request based on the MIMIC policies.

Code availability

The computer code used in this research will be available at www.github.com/yuxuanhu12/cshock under an open-source license.

Author contribution

Y.H., M.G., R.R. designed the experiments; Y.H., A.L. preprocessed and cleaned the MIMIC data; Y.H., A.T., C.T., S.M., J.M. reviewed the MIMIC charts; V.N., Y.A., N.K., J.Horowitz, T.A. selected and provided the NYU clinical data and context; Y.H. preprocessed, cleaned, and chart reviewed the NYU data; Y.H., M.G., R.R. developed the pipeline for CShock risk model; Y.H., M.G., implemented the CShock model; Y.H. implemented pretraining, analyzed subgroup performance and external validated the model with NYU data; Y.H., M.G., N.J., R.R. devised and implemented the model interpretation strategy; M.S., A.P., N.S., S.K. contributed to various analyses of the data; Y.H., A.L., M.G., R.R. conceived and directed the project; Y.H., M.G. made the figures; Y.H., M.G., G.F., J.Hochman, S.K., S.B., R.R. wrote the manuscript.

Methods References

- [39] Aaron van den Oord, Sander Dieleman, Heiga Zen, Karen Simonyan, Oriol Vinyals, Alex Graves, Nal Kalchbrenner, Andrew Senior, and Koray Kavukcuoglu. Wavenet: A generative model for raw audio. *arXiv preprint arXiv:1609.03499*, 2016.
- [40] Vinod Nair and Geoffrey E Hinton. Rectified linear units improve restricted boltzmann machines. In *Icml*, 2010.
- [41] Diederik P Kingma and Jimmy Ba. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.




Figure S1: NYU cardiac ICU Cohort for external validation

A Supplementary Information

A.1 Loss function

CShock’s loss function was constructed specifically for risk scoring. A risk score is calculated for each patient at each hour from time of admission until event of interest or discharge. For positive cases, the risk score is maximized for all times until the event of interest; for negative cases, the risk score is minimized until discharge.

Notation. We denote subject i ’s observed event type as $E_i \in \{\circ, \neq \circ\}$ where $E_i = \circ$ corresponds to cardiogenic/mixed shock and $E_i \neq \circ$ means the patient was not observed to develop cardiogenic shock, i.e. developed noncardiogenic shock only or no shock. Each patient has observed time $U_i \in \mathbb{R}_{\geq 0}$, which could be the time of shock onset or discharge. At each time $k \leq U_i$, features $X_{i,\leq k}$ from time 0 to time k are observed.

Model inputs. The CShock risk model r^θ with parameters θ takes in a time k and a variable-length feature list $X = [X_{i,0}, \dots, X_{i,k}]$ for $k \leq U_i$; it then produces CShock score for time k denoted $r^\theta(X_{\leq k}, k)$. Patient i ’s modeled CShock scores for all times from admission up to U_i are

$$r_i^\theta[\leq U_i] = \left\{ r_i^\theta(X_0, 0), \dots, \underbrace{r_i^\theta(X_{\leq k}, k)}_{\text{CShock score at time } k}, \dots, r_i^\theta(X_{\leq U_i}, U_i) \right\} \quad (1)$$

Loss for patients with cardiogenic/mixed shock. For a patient with observed cardiogenic/mixed shock at time U_i , it is sufficient for *any* CShock score until U_i to be large. We therefore maximize the maximum CShock score for patients with observed cardiogenic/mixed shock:

$$\max_{\theta} \sum_i \mathbb{1}[E_i = \circ] \left(\max_{k \in [0, \dots, U_i]} r_i^\theta[\leq U_i]_k \right) \quad (2)$$

To make this maximization amenable to gradient-based optimization, we maximize the CShock scores at all times, rather than just the maximum-risk-so-far. We weigh each risk by its proximity to the maximum risk and employ softmax for this purpose. For a patient with cardiogenic/mixed shock, the objective is:

$$\sum_k \text{Softmax}(r_i^\theta[\leq U_i])_k \cdot r_i^\theta[\leq U_i]_k = \text{Softmax}(r_i^\theta[\leq U_i])^\top r_i^\theta[\leq U_i] \quad (3)$$

Loss for patients with noncardiogenic shock only or no shock. The loss for non-cardiogenic shock only/no shock patients is similar: we minimize the maximum CShock score for times $[0, \dots, U_i]$ so that alarm does not sound for patients without cardiogenic shock. The optimization problem is:

$$\min_{\theta} \sum_i \mathbb{1}[E_i \neq \circ] \left(\max_{k \in [0, \dots, U_i]} r_i^\theta[\leq U_i]_k \right) \quad (4)$$

To relax this as a smooth optimization problem, we again use softmax weighting:

$$\sum_k \text{Softmax}(r_i^\theta[\leq U_i])_k \cdot r_i^\theta[\leq U_i]_k = \text{Softmax}(r_i^\theta[\leq U_i])^\top r_i^\theta[\leq U_i] \quad (5)$$

Combined loss function. Combining the loss for cardiogenic/mixed shock patients with that for noncardiogenic shock only/no shock patients, the loss, written as one minimization, is:

$$\min_{\theta} \sum_i \mathbb{1}[E_i \neq \circ] \text{Softmax}(\alpha r_i^\theta[\leq U_i])^\top r_i^\theta[\leq U_i] - \sum_i \mathbb{1}[E_i = \circ] \text{Softmax}(\alpha r_i^\theta[\leq U_i])^\top r_i^\theta[\leq U_i] \quad (6)$$

where α (set to 2) is a temperature scaling constant in the softmax, which makes the softmax behave more closely to a true maximum selection. Since cardiogenic shock detection occurs when CShock score exceeds the alarm threshold value, this loss function tightly maps to CShock’s alarm performance.

A.2 Model Interpretation: More Details

We used Shapley values, a popular explanation approach, to provide interpretations for model predictions. A Shapley value quantifies on average how much a *value function* changes when a feature is included versus when it is omitted in the presence of different subsets of the other features. Shapley values for the 70 features in the reduced model were estimated using FastSHAP [24].

Specifically, to construct a value function that scores different subsets of features, we train a surrogate model via masked prediction [26]. This surrogate model that takes as input a vector of masked feature pairs was trained by minimizing the same loss function as described in Eq. (6). Let \mathbf{s} be a subset of features e.g. $\mathbf{s} = \{1, 8, 11\}$. Let $\tilde{X}_t^{(\mathbf{s})}$ denote the features X_t at a given hour t , where only the features in \mathbf{s} are observed and the rest are masked. Masking features means zeroing their value and setting their mask indicator to indicate masking (-1 in this case). Features were masked at a frequency of 0.5 in the training of the surrogate model. Then, we can define $r_i^\theta[\leq U_i](\mathbf{s})$ to be equal to $r_i^\theta[\leq U_i]$ except with $\tilde{X}^{\mathbf{s}}$ instead of the original X . Finally, we can compute the original loss, but using $r_i^\theta[\leq U_i](\mathbf{s})$ instead of $r_i^\theta[\leq U_i]$:

$$\min_{\theta} \sum_i \mathbb{1}[E_i \neq \circ] \text{Softmax}(\alpha r_i^\theta[\leq U_i](\mathbf{s}))^\top r_i^\theta[\leq U_i](\mathbf{s}) - \sum_i \mathbb{1}[E_i = \circ] \text{Softmax}(\alpha r_i^\theta[\leq U_i](\mathbf{s}))^\top r_i^\theta[\leq U_i](\mathbf{s}) \quad (7)$$

The value function we use in FastSHAP is the contribution of each feature to the loss function without the label:

$$v(\mathbf{s}) = \text{Softmax}(\alpha r_i^\theta[\leq U_i](\mathbf{s}))^\top r_i^\theta[\leq U_i](\mathbf{s}) \quad (8)$$

This value function tracks how the soft-maximum risk score changes as a function of different subsets of features \mathbf{s} for the i -th patient. By using the trained surrogate model and the value function in Eq. (8), FastSHAP approximates Shapley values for the 70 features in the reduced model. The rankings for features based on Shapley values were computed by taking the mean absolute value across samples.

A.3 Features used in the models

See data file attached