

Generalized data thinning using sufficient statistics

Ameer Dharamshi¹, Anna Neufeld², Keshav Motwani¹, Lucy Gao³, Daniela Witten^{1,2}, and Jacob Bien⁴

¹Department of Biostatistics, University of Washington

²Department of Statistics, University of Washington

³Department of Statistics, University of British Columbia

⁴Department of Data Sciences and Operations, University of Southern California

March 24, 2023

Abstract

Our goal is to develop a general strategy to decompose a random variable X into multiple independent random variables, without sacrificing any information about unknown parameters. A recent paper showed that for some well-known natural exponential families, X can be *thinned* into independent random variables $X^{(1)}, \dots, X^{(K)}$, such that $X = \sum_{k=1}^K X^{(k)}$. In this paper, we generalize their procedure by relaxing this summation requirement and simply asking that some known function of the independent random variables exactly reconstruct X . This generalization of the procedure serves two purposes. First, it greatly expands the families of distributions for which thinning can be performed. Second, it unifies sample splitting and data thinning, which on the surface seem to be very different, as applications of the same principle. This shared principle is sufficiency. We use this insight to perform generalized thinning operations for a diverse set of families.

1 Introduction

Suppose that we want to *fit* and *validate* a model on the basis of a single dataset. Two example scenarios are as follows:

Scenario 1. We want to use the data both to generate and to test a hypothesis.

Scenario 2. We want to use the data both to fit a complicated model, and to obtain an accurate estimate of the expected prediction error.

In either case, it is clear that a naive approach that fits and validates a model on the same data is deeply problematic. In Scenario 1, testing a hypothesis on the same data used to generate it will lead to hypothesis tests that do not control the Type 1 error, and to confidence intervals that do not attain the nominal coverage (Fithian et al. 2014). And in Scenario 2, estimating the expected prediction error on the same data used to fit the model will lead to massive downward bias (see Tian 2020, Oliveira et al. 2021, for recent reviews).

In the case of Scenario 1, recent interest has focused on *selective inference*, a framework that enables a data analyst to generate and test a hypothesis on the same data (see, e.g., Taylor & Tibshirani 2015). The main idea is as follows: to test a hypothesis generated from the data, we should condition on the event that we selected this particular hypothesis. Despite promising applications of this framework to a number of problems, such as inference after regression (Lee et al. 2016), changepoint detection (Jewell et al. 2022, Hyun et al. 2021), clustering (Gao et al. 2022, Chen & Witten 2022, Yun & Barber 2023), and outlier detection (Chen & Bien 2020), it suffers from some drawbacks:

1. To perform selective inference, the procedure used to generate the null hypothesis must be fully-specified in advance. For instance, if a researcher wishes to cluster the data and then test for a difference in means between the clusters, as in Gao et al. (2022) and Chen & Witten (2022), then they must fully specify the clustering procedure (e.g., hierarchical clustering with squared Euclidean distance and complete linkage, cut to obtain K clusters) in advance.
2. Finite-sample selective inference typically requires an assumption of multivariate Gaussianity, though in some cases this can be relaxed to obtain asymptotic results (Taylor & Tibshirani 2018, Tian & Taylor 2017, Tibshirani et al. 2018, Tian & Taylor 2018).

Thus, it is clear that selective inference does not provide a flexible, “one-size-fits-all” approach to Scenario 1.

In the case of Scenario 2, proposals to de-bias the “in-sample” estimate of expected prediction error tend to be specialized to simple models, and thus do not provide an all-purpose tool that is broadly applicable to complex contemporary settings (Oliveira et al. 2021).

Sample splitting (Cox 1975) is an intuitive approach that is broadly applicable to a variety of settings, including Scenarios 1 and 2; see the left-hand panel of Figure 1. We split a dataset containing n observations into two sets, containing n_1 and n_2 observations, respectively (where $n_1 + n_2 = n$). Then we can generate a hypothesis based on the first set and test it on the second set (Scenario 1), or we can fit a model to the first set and estimate its error on the second set (Scenario 2). Sample splitting also forms the basis for cross-validation, an important tool for a practicing data scientist (Hastie et al. 2009).

While sample splitting often can adequately address both Scenarios 1 and 2, it also suffers from some drawbacks:

1. If the data contain outliers, then each outlier is assigned to a single subsample.

2. If the observations are not independent (for instance, if they correspond to a time series) then the subsamples that result from sample splitting are not independent, and so sample splitting does not provide a solution to either Scenario 1 or Scenario 2.
3. If one is interested in drawing conclusions at a per-observation level, then sample splitting is unsuitable. For example, if sample splitting is applied to a dataset consisting of the 50 states of the United States, then one can only conduct inference or perform validation on those states not used in fitting.
4. If the model of interest is fit using unsupervised learning, then sample splitting may not provide an adequate solution in either Scenario 1 or 2. The issue relates to #3 above. This is discussed in Gao et al. (2022), Chen & Witten (2022), and Neufeld et al. (2022) in the context of Scenario 1.

In recent work, Neufeld et al. (2023) proposed an approach for *convolution-closed data thinning* that addresses these drawbacks. They consider splitting, or *thinning*, a random variable X drawn from a convolution-closed family into K independent random variables $X^{(1)}, \dots, X^{(K)}$ such that $X = \sum_{k=1}^K X^{(k)}$, and $X^{(1)}, \dots, X^{(K)}$ come from the same family of distributions as X (see the right-hand panel of Figure 1). For instance, they show that $X \sim N(\mu, \sigma^2)$ can be thinned into two independent $N(\epsilon\mu, \epsilon\sigma^2)$ and $N((1 - \epsilon)\mu, (1 - \epsilon)\sigma^2)$ random variables that sum to X . Finally, and most critically, if X is drawn from a Gaussian, Poisson, negative binomial, binomial, multinomial, or gamma distribution, then they can thin it *even when parameters of its distribution are unknown*. Because the thinned random variables are independent, this provides a new approach to tackle Scenarios 1 and 2: one thins the dataset into two independent datasets. One then fits a model to one dataset, and validates it on the other.

On the surface, it is quite remarkable that one can break up a random variable X into two or more *independent* random variables that sum to X without knowing some (or sometimes any) of the parameters. In this paper, we seek to explain the underlying principles that make this possible. In doing so, we show that convolution-closed data thinning can be generalized so as to make it more flexible and much more widely applicable. The convolution-closed data thinning property $X = \sum_{k=1}^K X^{(k)}$ is desirable because it ensures that no information has been lost in the thinning process. However, clearly this would be equally true if we were to replace the summation by any other deterministic function. Likewise, the fact that $X^{(1)}, \dots, X^{(K)}$ are from the same family as X , while convenient, is nonessential.

Our generalization of convolution-closed data thinning is thus a procedure for splitting X into K random variables such that the following two properties hold:

- (i) $X = T(X^{(1)}, \dots, X^{(K)})$; and (ii) $X^{(1)}, \dots, X^{(K)}$ are mutually independent.

This generalization is broad enough to simultaneously encompass both convolution-closed data thinning and sample splitting. Furthermore, it greatly increases the scope of distributions that can be thinned. In the $K = 2$ case, this generalized goal has been stated before (see Leiner et al. 2022, “P1” property) as we will describe later. However, we are the first

to develop a widely applicable strategy for achieving this goal. Not only are we able to thin exponential families that were not previously possible (such as the beta family), but we can even thin outside of the exponential family. For example, generalized thinning enables us to thin $X \sim \text{Unif}(0, \theta)$ into $X^{(k)} \stackrel{\text{iid}}{\sim} \theta \cdot \text{Beta}\left(\frac{1}{K}, 1\right)$, for $k = 1, \dots, K$, in such a way that $X = \max\{X^{(1)}, \dots, X^{(K)}\}$.

The primary contributions of our paper are as follows:

1. We propose *generalized data thinning*, a general strategy for thinning a single random variable X into two or more independent random variables, $X^{(1)}, \dots, X^{(K)}$, without knowledge of the parameter value(s). Importantly, we show that *sufficiency* is the key property underlying the choice of the function $T(\cdot)$.
2. We show that it is possible to apply generalized data thinning to distributions far outside the scope of consideration of Neufeld et al. (2023): these include the beta, uniform, and shifted exponential distributions, among others. A summary of distributions covered by this work is provided in Table 1. In light of results by Darmois (1935), Koopman (1936), and Pitman (1936) (see the end of Section 2), we believe our examples are representative of the full range of cases to which this sufficiency-based approach can be applied.
3. We show that sample splitting — which, on its surface, bears little resemblance to convolution-closed data thinning — is in fact based on the same principle: both are special cases of generalized data thinning with different choices of the function $T(\cdot)$. In other words, our proposal is a direct *generalization* of sample splitting.

We are not the first to propose generalizations of sample splitting. Inspired by Tian & Taylor (2018)’s use of randomized responses, Rasines & Young (2022) define what they call the (U, V) -decomposition, which injects independent noise W to create two independent random variables $U = u(X, W)$ and $V = v(X, W)$ that together are jointly sufficient for the unknown parameters. However, they do not describe how to perform a (U, V) -decomposition other than in the special case of a Gaussian random vector with known covariance. Our generalized thinning framework achieves the goal set out in their paper, providing a concrete recipe for finding such decompositions in a broad set of examples. Another paper with a similar goal is Leiner et al. (2022). They define “data fission”, which seeks to find random variables $f(X)$ and $g(X)$ for which the distributions of $f(X)$ and $g(X) \mid f(X)$ are known and for which $X = h(f(X), g(X))$. When these two random variables are independent (which they describe as the “P1” property), their proposal aligns with generalized thinning. However, like Rasines & Young (2022), they do not provide a general strategy for performing P1-fission, and the only two examples they provide are the Gaussian vector with known covariance and the Poisson.

The rest of our paper is organized as follows. In Section 2, we define generalized data thinning, present our main theorem, and provide a simple recipe for thinning that is followed throughout the paper. In Section 3, we consider the case of thinning natural exponential families; this section also revisits the convolution-closed data thinning proposal of Neufeld




Figure 1: *Left*: Sample splitting assigns each observation into either a training set or a test set. *Right*: Generalized data thinning, the proposal of this paper, splits each observation into two parts that are independent and that can be used to recover the original observation $T(X^{(1)}, X^{(2)}) = X$. In some cases, they are drawn from the same distributional family as X .

et al. (2023), and clarifies the class of distributions that can be thinned using that approach. In Section 4, we show that we can apply data thinning to general exponential families. We consider distributions outside of the exponential family in Section 5. Section 6 contains examples of distributions that *cannot* be thinned using the approaches in this paper; these examples provide insight into the fact that sufficiency is the key property needed for (generalized) data thinning to “work”. We verify our results numerically in Section 7. Finally, we close with a discussion in Section 8; derivations and additional technical details are deferred to the appendix.

2 The generalized thinning proposal

Here and throughout, we write X to denote a random variable that can be scalar-, vector-, or matrix-valued (and likewise for $X^{(1)}, \dots, X^{(K)}$). We define *generalized data thinning* as follows.

Definition 1 (Generalized data thinning). *Consider a family of distributions $\mathcal{P} = \{P_\theta : \theta \in \Omega\}$. Suppose that there exists a distribution G_t , not depending on θ , and a deterministic function $T(\cdot)$ such that when we sample $(X^{(1)}, \dots, X^{(K)})|X$ from G_X , for $X \sim P_\theta$, the following properties hold:*

1. $X^{(1)}, \dots, X^{(K)}$ are mutually independent (and non-degenerate), and
2. $X = T(X^{(1)}, \dots, X^{(K)})$.

Then we say that \mathcal{P} is thinned by the function $T(\cdot)$.

When clear from context, sometimes we will say that “ P_θ is thinned” or “ X is thinned,” by which we mean that the corresponding family \mathcal{P} is thinned. Intuitively, we can think

Family	Distribution P_θ , where $X \sim P_\theta$.	Distribution $Q_\theta^{(k)}$ where $X^{(k)} \stackrel{ind.}{\sim} Q_\theta^{(k)}$.	Sufficient statistic T (sufficient for θ)	Reference / notes
Natural exponential family (in parameter θ)	$N(\theta, \sigma^2)$	$N(\epsilon_k \theta, \epsilon_k \sigma^2)$	$\sum_{k=1}^K X^{(k)}$	Neufeld et al. (2023)
	Poisson(θ)	Poisson($\epsilon_k \theta$)		
	NegBin(r, θ)	NegBin($\epsilon_k r, \theta$)		
	Binomial(r, θ)	Binomial($\epsilon_k r, \theta$)		
	Gamma(α, θ)	Gamma($\epsilon_k \alpha, \theta$)		
	$N_p(\boldsymbol{\theta}, \Sigma)$	$N_p(\epsilon_k \boldsymbol{\theta}, \epsilon_k \Sigma)$		
	Multinomial $_p(r, \boldsymbol{\theta})$	Multinomial $_p(\epsilon_k r, \boldsymbol{\theta})$		
	Gamma($K/2, \theta$)	$N(0, \frac{1}{2\theta})$	$\sum_{k=1}^K (X^{(k)})^2$	Example 3.3
	Gamma(K, θ)	Weibull($\theta^{-\frac{1}{\nu}}, \nu$)	$\sum_{k=1}^K (X^{(k)})^\nu$	Example 3.4
General exponential family (in parameter θ)	Beta(θ, β)	Beta($\frac{1}{K}\theta + \frac{k-1}{K}, \frac{1}{K}\beta$)	$(\prod_{k=1}^K X^{(k)})^{1/K}$	Example 4.1
	Beta(α, θ)	Beta($\frac{1}{K}\alpha, \frac{1}{K}\theta + \frac{k-1}{K}$)	$(\prod_{k=1}^K (1 - X^{(k)}))^{1/K}$	Text below Example 4.1
	Gamma(θ, β)	Gamma($\frac{1}{K}\theta + \frac{k-1}{K}, \frac{1}{K}\beta$)	$(\prod_{k=1}^K X^{(k)})^{1/K}$	Example 4.2
	Weibull(θ, ν)	Gamma($\frac{1}{K}, \theta^{-\nu}$)	$(\sum_{k=1}^K X^{(k)})^{1/\nu}$	Example 4.4
	Pareto(ν, θ)	Gamma($\frac{1}{K}, \theta$)	$\nu \times \text{Exp}(\sum_{k=1}^K X^{(k)})$	Example 4.4
		$N(\mu, \theta)$	Gamma($\frac{1}{2K}, \frac{1}{2\theta}$)	$X^2 = \sum_{k=1}^K X^{(k)}$
	$N_K(\theta_1 1_K, \theta_2 I_K)$	$N(\theta_1, \theta_2)$	sample mean and variance	Indirect only; Example 4.3
Truncated support family	Unif($0, \theta$)	$\theta \cdot \text{Beta}(\frac{1}{K}, 1)$	$\max(X^{(1)}, \dots, X^{(K)})$	Example 5.1
	$\theta \cdot \text{Beta}(\alpha, 1)$	$\theta \cdot \text{Beta}(\frac{\alpha}{K}, 1)$		Example B.1
	$\theta + \text{Exp}(\lambda)$	$\theta + \text{Exp}(\lambda/K)$		Example 5.2
Non-parametric	F^n	F^{n_k}	See Example 5.3	Example 5.3

Table 1: Examples of named families (indexed by an unknown parameter θ) that can be thinned into K components, where K is a positive integer, without knowledge of θ . In cases where they are used, $\epsilon_k > 0$ is a tuning parameter such that $\sum_{k=1}^K \epsilon_k = 1$, n_1, \dots, n_K are integers that sum to n , and $\nu > 0$.

of thinning as breaking X up into K independent pieces, but in a very particular way that ensures that none of the information about θ is lost. The fact that no information is lost is evident from the requirement that $X = T(X^{(1)}, \dots, X^{(K)})$.

It turns out that sample splitting (Cox 1975) can be viewed as a special case of generalized data thinning.

Remark 1 (Sample splitting). *Sample splitting, in which a sample of n independent and identically distributed random variables is partitioned into K subsamples, is a special case of generalized thinning. Here, $T(\cdot)$ is the function that takes in the subsamples as arguments, and concatenates and sorts their elements. For more details, see Section 5.2.*

Furthermore, Definition 1 is closely related to the proposal of Neufeld et al. (2023).

Remark 2 (Thinning convolution-closed families of distributions). *Neufeld et al. (2023) show that some well-known families of convolution-closed distributions, such as the binomial, negative binomial, gamma, Poisson, and Gaussian, can be thinned, in the sense of Definition 1, by addition: $T(x^{(1)}, \dots, x^{(K)}) = \sum_{k=1}^K x^{(k)}$.*

The two examples above do not resemble each other: the first involves a non-parametric family of distributions and applies quite generally, while the second depends on a specific property of the family of distributions. Furthermore, the functions $T(\cdot)$ are quite different from each other. It is natural to ask: How can we find families \mathcal{P} that can be thinned? Is there a unifying principle for the choice of $T(\cdot)$? How can we ensure that there exists a distribution G_t as in Definition 1 that does not depend on θ ? The following theorem answers these questions, and indicates that *sufficiency* is the key principle required to ensure that the distribution G_t does not depend on θ .

Theorem 1 (Main theorem). *Suppose \mathcal{P} is thinned by a function $T(\cdot)$ and, for $X \sim P_\theta$, let $Q_\theta^{(1)} \times \dots \times Q_\theta^{(K)}$ denote the distribution of the mutually independent random variables, $(X^{(1)}, \dots, X^{(K)})$, sampled as in Definition 1. Then, the following hold:*

(a) $T(X^{(1)}, \dots, X^{(K)})$ is a sufficient statistic for θ based on $(X^{(1)}, \dots, X^{(K)})$.

(b) The distribution G_t in Definition 1 is the conditional distribution

$$(X^{(1)}, \dots, X^{(K)}) | T(X^{(1)}, \dots, X^{(K)}) = t,$$

where $(X^{(1)}, \dots, X^{(K)}) \sim Q_\theta^{(1)} \times \dots \times Q_\theta^{(K)}$.

Theorem 1 is proven in Appendix A.1. This theorem further suggests a simple strategy for finding families of distributions \mathcal{P} and functions $T(\cdot)$ such that \mathcal{P} can be thinned by $T(\cdot)$.

Algorithm 1 (Finding distributions that can be thinned).

1. Choose K families of distributions, $\mathcal{Q}^{(k)} = \{Q_\theta^{(k)} : \theta \in \Omega\}$ for $k = 1, \dots, K$.

2. Let $(X^{(1)}, \dots, X^{(K)}) \sim Q_\theta^{(1)} \times \dots \times Q_\theta^{(K)}$, and let $T(X^{(1)}, \dots, X^{(K)})$ denote a sufficient statistic for θ .
3. Let P_θ denote the distribution of $T(X^{(1)}, \dots, X^{(K)})$.

By Theorem 1, the family $\mathcal{P} = \{P_\theta : \theta \in \Omega\}$ is thinned by $T(\cdot)$.

This recipe gives us a very succinct way to describe the distributions that can be thinned: *we can thin the distributions of sufficient statistics*. In particular, the recipe takes as input a joint distribution $Q_\theta^{(1)} \times \dots \times Q_\theta^{(K)}$, and requires us to choose a sufficient statistic for θ . Then, that statistic's distribution is the P_θ that can be thinned.

The next three sections of this paper focus on demonstrating the utility of Theorem 1 and Algorithm 1 in a series of examples. Outside of the exponential family, the sufficient statistic grows in dimension with the sample size, except for some distributions that have support controlled by a parameter (Darmois 1935, Koopman 1936, Pitman 1936). Motivated by this result, we focus on exponential families in Sections 3 and 4, and on examples outside of the exponential family in Section 5. The full set of examples considered in the coming sections is listed in Table 1.

3 Thinning natural exponential families

In Section 3.1, we show how to thin a natural exponential family into two or more natural exponential families. In Section 3.2, we show how the convolution-closed thinning proposal of Neufeld et al. (2023) can be understood in light of natural exponential family thinning. Finally, in Section 3.3, we show how natural exponential families can be thinned into more general (i.e., not necessarily natural) exponential families.

3.1 Thinning natural exponential families into natural exponential families

In this section, we apply Algorithm 1 to the important special case of natural exponential families (Lehmann & Romano 2005). A natural exponential family starts with a known probability distribution H , and then forms a family of distributions $\mathcal{P}^H = \{P_\theta^H : \theta \in \Omega\}$ based on H , as follows:

$$dP_\theta^H(x) = e^{x^\top \theta - \psi_H(\theta)} dH(x). \quad (1)$$

Here $e^{-\psi_H(\theta)}$ is the normalization constant needed to ensure that P_θ is a probability distribution, and we take Ω to be the set of θ for which this normalization is possible (i.e. for which $\psi_H(\theta) < \infty$).

The next theorem presents a property of H that is necessary and sufficient for the resulting natural exponential family \mathcal{P}^H to be thinned by addition into K natural exponential families. To streamline the statement of the theorem, we start with a definition.

Definition 2 (*K*-way convolution distribution). We say that a probability distribution H is the *K*-way convolution of distributions H_1, \dots, H_K if $\sum_{k=1}^K Y_k \sim H$ for $(Y_1, \dots, Y_K) \sim H_1 \times \dots \times H_K$.

Theorem 2 (Thinning natural exponential families by addition). Consider the natural exponential family \mathcal{P}^H defined in (1). This family can be thinned by the function $T(X^{(1)}, \dots, X^{(K)}) = \sum_{k=1}^K X^{(k)}$ into *K* natural exponential families $\mathcal{P}^{H_1}, \dots, \mathcal{P}^{H_K}$ if and only if H is the *K*-way convolution of H_1, \dots, H_K .

This theorem characterizes the set of natural exponential families that can be thinned by addition into *K* natural exponential families. These *K* natural exponential families can be different from each other, but they are all indexed by the same $\theta \in \Omega$ that was used in the original family \mathcal{P}^H .

The proof of Theorem 2 is in Appendix A.2. We provide a brief sketch here. We apply Algorithm 1, with each $\mathcal{Q}^{(k)}$ chosen to be a natural exponential family \mathcal{P}^{H_k} , where H_k is an arbitrary distribution. We then imagine observing $(X^{(1)}, \dots, X^{(K)}) \sim \mathcal{Q}_\theta^{(1)} \times \dots \times \mathcal{Q}_\theta^{(K)}$. The sum $X^{(1)} + \dots + X^{(K)}$ is sufficient for θ , so the distribution of this sum can be thinned by addition. Finally, we note that $X^{(1)} + \dots + X^{(K)} \sim P_\theta^H$, where H is the convolution of H_1, \dots, H_K .

Neufeld et al. (2023) show that it is possible to thin a Gaussian random variable by addition into *K* independent Gaussians. We now see that this result follows from Theorem 2.

Example 3.1 (Thinning $N_n(\theta, I_n)$). The family of $N_n(\theta, I_n)$ distributions is a natural exponential family indexed by $\theta \in \mathbb{R}^n$. It can be written in the notation of this section as \mathcal{P}^H , where H represents the $N_n(0_n, I_n)$ distribution. Furthermore, H is the *K*-way convolution of $H_k = N_n(0_n, \epsilon_k I_n)$ for $k = 1, 2, \dots, K$, where $\epsilon_1, \dots, \epsilon_K > 0$ and $\sum_{k=1}^K \epsilon_k = 1$. Thus, by Theorem 2, we can thin \mathcal{P}^H by addition into $\mathcal{P}^{H_1}, \dots, \mathcal{P}^{H_K}$, where $P_\theta^{H_k} = N_n(\epsilon_k \theta, \epsilon_k I_n)$.

The above example is particularly notable for its connection to a randomization strategy that has been frequently used in the literature.

Remark 3 (Connecting Example 3.1 to prior work). Other authors have considered obtaining two independent Gaussian random variables U and V from a single Gaussian random variable $X \sim N_n(\theta, I_n)$ by generating $W \sim N_n(0_n, \gamma I_n)$ for a tuning parameter $\gamma > 0$, and then setting $U = X + W$ and $V = X - \gamma^{-1}W$. Then, $U \sim N_n(\theta, (1 + \gamma)I_n)$ and $V \sim N_n(\theta, (1 + \gamma^{-1})I_n)$ are independent. Rasines & Young (2022) and Leiner et al. (2022) applied this decomposition to address Scenario 1 in Section 1. Additionally, Rasines & Young (2022) showed that this leads to asymptotically valid inference under certain regularity conditions, even when X is not normally distributed. Tian (2020) and Oliveira et al. (2021) applied this decomposition to address Scenario 2 in Section 1.

This decomposition is in fact identical to Example 3.1 up to scaling, with $X^{(1)} = \epsilon_1 U$, $X^{(2)} = \epsilon_2 V$, $\epsilon_1 = (1 + \gamma)^{-1}$, and $\epsilon_2 = 1 - \epsilon_1$. In particular, to thin $X \sim N_n(\theta, I_n)$ by addition into $(X^{(1)}, X^{(2)})$, we sample from G_X where G_t , defined in Theorem 1, can be shown to equal the singular multivariate normal distribution

$$N_{2n} \left(\begin{pmatrix} \epsilon_1 t \\ \epsilon_2 t \end{pmatrix}, \epsilon_1 \epsilon_2 \begin{pmatrix} I_n & -I_n \\ -I_n & I_n \end{pmatrix} \right).$$

Sampling from G_X is equivalent to sampling $W \sim N_n(0_n, \gamma I_n)$ (independent of X) and then generating $X^{(1)} = \epsilon_1(X + W)$ and $X^{(2)} = \epsilon_2(X - \gamma^{-1}W)$. These ideas can easily be generalized to thin $X \sim N_n(\theta, \Sigma)$ with a known positive definite covariance matrix Σ .

Furthermore, Theorem 2 implies that we can thin a natural exponential family by addition into natural exponential families *only if* H is the convolution of two or more distributions. Not all distributions satisfy this property. For example, the distribution $H = \text{Bernoulli}(0.5)$ cannot be written as the sum of two independent, non-constant random variables. Since $\mathcal{P}^{\text{Bernoulli}(0.5)}$ is the $\text{Bernoulli}([1 + e^{-\theta}]^{-1})$ natural exponential family, this tells us that Bernoulli random variables cannot be thinned by addition. In fact, in Section 6.1 we will prove a stronger statement: namely, that *no function* $T(\cdot)$ can thin the Bernoulli family.

3.2 Connections to Neufeld et al. (2023)

Neufeld et al. (2023) focus on convolution-closed families, i.e., those for which convolving two or more distributions (see Definition 2) that are in the family produces a distribution that is still in the family. They provide a recipe for decomposing a random variable X drawn from a distribution in such a family into independent random variables $X^{(1)}, \dots, X^{(K)}$ that sum to yield X . In this section, we will show that their decompositions of exponential dispersion families are encompassed by Theorem 2.

Exponential dispersion families (Jørgensen 1992, Jørgensen & Song 1998) are a subclass of convolution-closed families that include many families of interest. Given a distribution H with $\psi_H(\theta) < \infty$ for $\theta \in \Omega$ (as in (1)), we identify the set of distributions H_λ for which $\psi_{H_\lambda}(\cdot) = \lambda\psi_H(\cdot)$ (i.e., distributions whose cumulant generating function is a multiple of H 's cumulant generating function). We define Λ to be the set of λ for which such a distribution H_λ exists. Then, an (additive) *exponential dispersion family* is $\mathcal{P} = \bigcup_{\lambda \in \Lambda} \mathcal{P}^{H_\lambda}$, where \mathcal{P}^{H_λ} is the natural exponential family generated by H_λ (see (1)). The distributions in \mathcal{P} are indexed over $(\theta, \lambda) \in \Omega \times \Lambda$ and take the following form:

$$dP_\theta^{H_\lambda}(x) = e^{x^\top \theta - \lambda \psi_H(\theta)} dH_\lambda(x). \quad (2)$$

In words, an exponential dispersion family results from combining a collection of related natural exponential families. For example, starting with $H = \text{Bernoulli}(1/2)$, we can take $\Lambda = \mathbb{Z}^+$ since for any positive integer λ , $H_\lambda = \text{Binomial}(\lambda, 1/2)$ satisfies the necessary cumulant generating function relationship. Then, $\mathcal{P}^{\text{Binomial}(\lambda, 1/2)}$ corresponds to the binomial natural exponential family that results from fixing λ . Finally, allowing λ to vary gives the full binomial exponential dispersion family, which is the set of all binomial distributions (varying both of the parameters of the binomial distribution).

By construction, for any $\lambda_1, \dots, \lambda_K \in \Lambda$, convolving $P_\theta^{H_{\lambda_1}}, \dots, P_\theta^{H_{\lambda_K}}$ gives the distribution $P_\theta^{H_\lambda}$, where $\lambda = \sum_{k=1}^K \lambda_k$. The next corollary is an immediate application of Theorem 2 in the context of exponential dispersion families. Notably, the distributions $Q_\theta^{(k)}$ themselves still belong to the exponential dispersion family \mathcal{P} to which the distribution of X belongs.

Corollary 1 (Thinning while remaining inside an exponential dispersion family). *Consider an exponential dispersion family $\mathcal{P} = \bigcup_{\lambda \in \Lambda} \mathcal{P}^{H_\lambda}$ and suppose $\lambda_1, \dots, \lambda_K \in \Lambda$. Then for $\lambda = \sum_{k=1}^K \lambda_k$, we can thin the natural exponential family \mathcal{P}^{H_λ} by $T(x^{(1)}, \dots, x^{(K)}) = \sum_{k=1}^K x^{(k)}$ into the natural exponential families $\mathcal{P}^{H_{\lambda_1}}, \dots, \mathcal{P}^{H_{\lambda_K}}$.*

This result corresponds exactly to the data thinning proposal of Neufeld et al. (2023). We see from Corollary 1 that that proposal thins a natural exponential family, \mathcal{P}^{H_λ} , into a *different* set of natural exponential families, $\mathcal{P}^{H_{\lambda_1}}, \dots, \mathcal{P}^{H_{\lambda_K}}$. However, from the perspective of exponential dispersion families, it thins an exponential dispersion family into the same exponential dispersion family. Continuing the binomial example from above, the corollary tells us that we can thin the binomial family with λ as the number of trials into two or more binomial families with smaller numbers of trials, provided that $\lambda > 1$.

Neufeld et al. (2023) focus on convolution-closed families, not exponential dispersion families. However, Jørgensen & Song (1998) note that all convolution-closed families that have moment-generating functions can be written as exponential dispersion families. The Cauchy distribution is convolution-closed, but does not have a moment generating function and thus is not an exponential dispersion family. As we will see in Example 6.1, the Cauchy(θ_1, θ_2) distribution cannot be thinned by addition: decomposing it using the recipe of Neufeld et al. (2023) requires knowledge of both unknown parameters. Thus, not all convolution-closed distributions can be thinned by addition in the sense of Definition 1. However, Neufeld et al. (2023) claim that all convolution-closed distributions *can* be thinned. This apparent discrepancy is due to a slight difference in the definition of thinning between our paper and theirs: in Definition 1, we require that G_t not depend on θ ; however, Neufeld et al. (2023) have no such requirement. In practice, data thinning is useful only if G_t does not depend on θ , and so there is no meaningful difference between the two definitions.

3.3 Thinning natural exponential families into general exponential families

In this section, we apply Algorithm 1 in the case that $\mathcal{Q}^{(k)}$ are (possibly non-natural) exponential families, for which the sufficient statistic need not be the identity. In particular, for $k = 1, \dots, K$, we let $\mathcal{Q}^{(k)} = \{Q_\theta^{(k)} : \theta \in \Omega\}$ denote an exponential family based on a known distribution H_k and sufficient statistic $T^{(k)}(\cdot)$:

$$dQ_\theta^{(k)}(x) = \exp\{[T^{(k)}(x)]^\top \eta(\theta) - \psi_k(\theta)\} dH_k(x). \quad (3)$$

As in Section 3.1, $e^{-\psi_k(\theta)}$ is the normalization constant needed to ensure that $\int dQ_\theta^{(k)}(x) = 1$ and Ω is the set of θ for which $\psi_k(\theta) < \infty$. The function $\eta(\cdot)$ maps θ to the natural parameter. We note that $\sum_{k=1}^K T^{(k)}(X^{(k)})$ is a sufficient statistic for θ based on $(X^{(1)}, \dots, X^{(K)}) \sim Q_\theta^{(1)} \times \dots \times Q_\theta^{(K)}$. Then Algorithm 1 tells us that we can thin the distribution of this sufficient statistic. This leads to the next result.

Proposition 1 (Thinning natural exponential families with more general functions $T(\cdot)$). *Let $X^{(1)}, \dots, X^{(K)}$ be independent random variables with $X^{(k)} \sim Q_\theta^{(k)}$ for $k = 1, \dots, K$*

from any (i.e., possibly non-natural) exponential families $\mathcal{Q}^{(k)}$ as in (3). Let P_θ denote the distribution of $\sum_{k=1}^K T^{(k)}(X^{(k)})$. Then, $\mathcal{P} = \{P_\theta : \theta \in \Omega\}$ is a natural exponential family, and we can thin it into $X^{(1)}, \dots, X^{(K)}$ using the function $T(x^{(1)}, \dots, x^{(K)}) = \sum_{k=1}^K T^{(k)}(x^{(k)})$.

The fact that \mathcal{P} in this result is a natural exponential family follows from recalling that the sufficient statistic of an exponential family follows a natural exponential family (Lehmann & Romano 2005, Lemma 2.7.2(i)). Many named exponential families are not natural exponential families, involving non-identity functions $T^{(k)}(\cdot)$, such as the logarithm or polynomials. Therefore, to thin into those families, Proposition 1 will be useful. We illustrate the flexibility provided by Proposition 1 with a number of examples.

In particular, we demonstrate that a natural exponential family \mathcal{P} can be thinned by different functions $T(\cdot)$, leading to families of distributions $\mathcal{Q}^{(1)}, \dots, \mathcal{Q}^{(K)}$ different from \mathcal{P} . Specifically, the following examples demonstrate three possible K -fold thinning strategies for a gamma distribution when the shape, α , is known but the rate¹ θ , is unknown: the first uses the method in Neufeld et al. (2023), the second applies when α can be expressed as half of a natural number and decomposes the gamma observations into centred normal data, and the final applies when α is a natural number and decomposes the gamma observations into Weibull data with shape ν , a value to be chosen by the analyst.

Example 3.2 (Thinning $\text{Gamma}(\alpha, \theta)$ with α known, approach 1). *Following Algorithm 1, we start with $X^{(k)} \stackrel{iid}{\sim} \text{Gamma}(\frac{\alpha}{K}, \theta)$ for $k = 1, \dots, K$, and note that $T(X^{(1)}, \dots, X^{(K)}) = \sum_{k=1}^K X^{(k)}$ is sufficient for θ . Thus, we can thin the distribution of $\sum_{k=1}^K X^{(k)}$. A well-known property of the gamma distribution tells us that this is a $\text{Gamma}(\alpha, \theta)$ distribution. Sampling from G_t as in Theorem 1 corresponds exactly to the multi-fold gamma data thinning recipe of Neufeld et al. (2023) where $\epsilon_k = \frac{1}{K}$.*

The next two examples apply Proposition 1 to decompose the gamma family into different families.

Example 3.3 (Thinning $\text{Gamma}(\alpha, \theta)$ with $\alpha = K/2$ known, approach 2). *Starting with $X^{(k)} \stackrel{iid}{\sim} N(0, \frac{1}{2\theta})$, notice that $T^{(k)}(x^{(k)}) = (x^{(k)})^2$. We can thus apply Proposition 1 using the function*

$$T(x^{(1)}, \dots, x^{(K)}) = \sum_{k=1}^K (x^{(k)})^2$$

to thin the sufficient statistic, $\sum_{k=1}^K (X^{(k)})^2 \sim \frac{1}{2\theta} \chi_K^2 = \text{Gamma}(\frac{K}{2}, \theta)$, into $(X^{(1)}, \dots, X^{(K)})$. The function G_t from Theorem 1 is the conditional distribution

$$(X^{(1)}, \dots, X^{(K)}) \mid \sum_{k=1}^K (X^{(k)})^2 = t.$$

¹Although θ is often used in the gamma distribution to denote the scale parameter, here we use it to denote the rate parameter.

By the rotational symmetry of the $N_K(0, (2\theta)^{-1}I_K)$ distribution (which is the joint distribution of $(X^{(1)}, \dots, X^{(K)})$), G_t corresponds to the uniform distribution on the $(K-1)$ -sphere of radius $t^{1/2}$.

To sample from this conditional distribution, we generate $Z \sim N_K(0, I_K)$ and then take $(X^{(1)}, \dots, X^{(K)})|X$ to be $X^{1/2} \frac{Z}{\|Z\|_2}$.

The next example produces not just a single thinning function $T(\cdot)$ but a *family* of thinning functions, indexed by a parameter $\nu > 0$ that the analyst can choose.

Example 3.4 (Thinning $\text{Gamma}(\alpha, \theta)$ with $\alpha = K$ known, approach 3). *Recall that the Weibull distribution with known shape parameter ν (but varying θ) is a member of a general exponential family. Then, starting with $X^{(k)} \stackrel{iid}{\sim} \text{Weibull}(\lambda, \nu)$ for $k = 1, \dots, K$, we have that $T^{(k)}(x^{(k)}) = (x^{(k)})^\nu$. We can thus apply Proposition 1 using the function*

$$T(x^{(1)}, \dots, x^{(K)}) = \sum_{k=1}^K (x^{(k)})^\nu$$

to thin the distribution of $\sum_{k=1}^K (X^{(k)})^\nu$ into $(X^{(1)}, \dots, X^{(K)})$. In Appendix B.1, we show that $\sum_{k=1}^K (X^{(k)})^\nu \sim \text{Gamma}(K, \lambda^{-\nu})$, so then taking $\lambda = \theta^{-1/\nu}$ yields the desired result.

To generate $(X^{(1)}, \dots, X^{(K)})$, we can first apply the K -fold gamma thinning result discussed in Example 3.2 with $\epsilon_k = \frac{1}{K}$ to generate $Y^{(k)} \stackrel{iid}{\sim} \text{Exp}(\lambda^{-\nu})$, and then compute $X^{(k)} = (Y^{(k)})^{\frac{1}{\nu}}$.

Examples 3.2, 3.3, and 3.4 present three distinct strategies for thinning a gamma distribution with unknown rate parameter. The first approach applies most generally, allowing an analyst to thin any gamma distribution with known α , not just the subset with α parameters that are integers or halves of integers. Yet the second and third can have contextual advantages. For instance, if X is a sample variance that follows a scaled chi-squared distribution, then recovering an underlying sample of normal observations may be more useful than a sample of gamma data. Similarly, if X is the total waiting time for k units, then decomposing it into Weibull realizations may be preferable.

4 Indirect thinning of general exponential families

Sometimes rather than thinning X , we may choose to thin a function $S(X)$. When $S(X)$ is sufficient for θ based on X , the next proposition tells us that thinning $S(X)$ rather than X does not result in a loss of information about θ . We emphasize that we are using the concept of sufficiency in two ways here: (i) $S(X)$ is sufficient for θ based on $X \sim P_\theta$, and (ii) $T(X^{(1)}, \dots, X^{(K)})$ is sufficient for θ based on $(X^{(1)}, \dots, X^{(K)}) \sim Q_\theta^{(1)} \times \dots \times Q_\theta^{(K)}$.

Proposition 2 (Thinning a sufficient statistic preserves information). *Suppose $X \sim P_\theta \in \mathcal{P}$ has a sufficient statistic $S(X)$ for θ , and we thin $S(X)$ by $T(\cdot)$. That is, conditional on*

$S(X)$ (and without knowledge of θ) we are able to sample $X^{(1)}, \dots, X^{(K)}$ that are mutually independent and satisfy

$$S(X) = T(X^{(1)}, \dots, X^{(K)}). \quad (4)$$

Under regularity conditions on the distributions of X , $S(X)$, and $X^{(1)}, \dots, X^{(K)}$ needed for Fisher information to exist, we have that

$$I_X(\theta) = \sum_{k=1}^K I_{X^{(k)}}(\theta).$$

This proposition shows that thinning $S(X)$, rather than X , does not result in a loss of information about θ . Its proof (provided in Appendix A.3) follows easily from multiple applications of the fact that sufficient statistics preserve information.

We formalise this strategy in the following definition.

Definition 3 (Indirect thinning). *Consider $X \sim P_\theta \in \mathcal{P}$. Suppose we thin a sufficient statistic $S(X)$ for θ by a function $T(\cdot)$. We say that the family \mathcal{P} is indirectly thinned through $S(\cdot)$ by $T(\cdot)$.*

In light of Proposition 2, indirect thinning of a random variable X does not result in a loss of information.

When $S(\cdot)$ is invertible, then

$$X = S^{-1}(T(X^{(1)}, \dots, X^{(K)})),$$

which implies that we can thin X directly by $S^{-1}(T(\cdot))$. It turns out that, regardless of whether we thin X by $S^{-1}(T(\cdot))$ or indirectly thin X through $S(\cdot)$ by $T(\cdot)$, there is little difference between the resulting form of G_t in Theorem 1. In the former case, G_t is the conditional distribution of $(X^{(1)}, \dots, X^{(K)})$ given $S^{-1}(T(X^{(1)}, \dots, X^{(K)})) = t$. In the latter case, it is the conditional distribution of $(X^{(1)}, \dots, X^{(K)})$ given $T(X^{(1)}, \dots, X^{(K)}) = t$. Since $S(\cdot)$ is invertible, these two conditional distributions are identical following a reparameterization.

We now return to the setting of Proposition 2, where $S(\cdot)$ may or may not be invertible.

Remark 4 (Indirect thinning of general exponential families). *Let $\mathcal{P} = \{P_\theta : \theta \in \Omega\}$ be a general exponential family. That is,*

$$dP_\theta(x) = \exp\{[S(x)]^\top \eta(\theta) - \psi(\theta)\} dH(x),$$

where $e^{-\psi(\theta)}$ is the normalising constant. Since $S(X)$ is sufficient for θ , we can indirectly thin X through $S(\cdot)$ without a loss of Fisher information (Proposition 2). Furthermore, $S(X)$ belongs to a natural exponential family (Lehmann & Romano 2005, Lemma 2.7.2(i)). We can thus indirectly thin X through $S(\cdot)$ as follows:

1. Provided that the necessary and sufficient condition of Theorem 2 holds for $S(X)$, we can indirectly thin X through $S(\cdot)$ by addition into $X^{(1)}, \dots, X^{(K)}$ that follow natural exponential families, i.e. general exponential families where $T^{(k)}(\cdot)$ in (3) is the identity.

2. We now consider $X^{(1)}, \dots, X^{(K)}$ that belong to a general exponential family, where $T^{(k)}(\cdot)$ in (3) is not necessarily the identity. Suppose further that $S(X) \stackrel{D}{=} \sum_{k=1}^K T^{(k)}(X^{(k)})$. Then, by Proposition 1, we can indirectly thin X through $S(\cdot)$ into $X^{(1)}, \dots, X^{(K)}$, by $T(x^{(1)}, \dots, x^{(K)}) = \sum_{k=1}^K T^{(k)}(x^{(k)})$.

We see that 1) is a special case of 2).

We now demonstrate the usefulness of indirect thinning with some examples. First, we consider a $\text{Beta}(\theta, \beta)$ random variable, with β a known parameter. The beta family is not a natural exponential family, and so the results in Section 3 are not directly applicable. The beta distribution also differs from some of the other examples that we have seen in the following ways: (i) it is not convolution-closed; (ii) it has finite support; and (iii) the sufficient statistic for a sample of independent and identically distributed beta random variables has an unnamed distribution.

Example 4.1 (Thinning $\text{Beta}(\theta, \beta)$ with β known). We start with $X^{(k)} \stackrel{\text{ind}}{\sim} \text{Beta}\left(\frac{1}{K}\theta + \frac{k-1}{K}, \frac{1}{K}\beta\right)$, for $k = 1, \dots, K$; this is a general exponential family (3) with $T^{(k)}(x^{(k)}) = \frac{1}{K} \log(x^{(k)})$. Since $\sum_{k=1}^K T^{(k)}(X^{(k)})$ is sufficient for θ based on $X^{(1)}, \dots, X^{(K)}$, we can apply Proposition 1 to thin the distribution of $\sum_{k=1}^K T^{(k)}(X^{(k)})$ by the function

$$T(x^{(1)}, \dots, x^{(K)}) = \sum_{k=1}^K T^{(k)}(x^{(k)}) = \frac{1}{K} \sum_{k=1}^K \log(x^{(k)}) = \log \left[\left(\prod_{k=1}^K x^{(k)} \right)^{1/K} \right]. \quad (5)$$

Furthermore, we show in Appendix B.2 that $\exp(T(X^{(1)}, \dots, X^{(K)})) = \left(\prod_{k=1}^K X^{(k)} \right)^{1/K}$, the geometric mean of $X^{(1)}, \dots, X^{(K)}$, follows a $\text{Beta}(\theta, \beta)$ distribution. Therefore, we can indirectly thin a $\text{Beta}(\theta, \beta)$ random variable through $S(x) = \log(x)$ by $T(\cdot)$ defined in (5). This results in independent random variables $X^{(k)} \sim \text{Beta}\left(\frac{1}{K}\theta + \frac{k-1}{K}, \frac{1}{K}\beta\right)$, for $k = 1, \dots, K$.

It turns out that in this example, the thinning operation can be simplified using the fact that $S(x) = \log(x)$ is invertible: we can directly thin $X \sim \text{Beta}(\theta, \beta)$ by the function

$$T'(x^{(1)}, \dots, x^{(K)}) = S^{-1}(T(x^{(1)}, \dots, x^{(K)})) = \left(\prod_{k=1}^K x^{(k)} \right)^{1/K}. \quad (6)$$

To thin $\text{Beta}(\theta, \beta)$ using either of the strategies listed above, we need to sample from G_t defined in Theorem 1. This unnamed distribution can be easily sampled from using numerical methods, as detailed in Appendix B.2.

By symmetry of the beta distribution, we can also apply the thinning operations detailed in Example 4.1 to thin a $\text{Beta}(\alpha, \theta)$ random variable with α known.

Next, we consider thinning the gamma distribution with unknown shape parameter.

Example 4.2 (Thinning $\text{Gamma}(\theta, \beta)$ with β known). We start with $X^{(k)} \stackrel{\text{ind}}{\sim} \text{Gamma}\left(\frac{1}{K}\theta + \frac{k-1}{K}, \frac{1}{K}\beta\right)$, for $k = 1, \dots, K$; this is a general exponential family (3) with $T^{(k)}(X^{(k)}) = \frac{1}{K} \log(x^{(k)})$. Note that $T(X^{(1)}, \dots, X^{(K)}) = \sum_{k=1}^K T^{(k)}(X^{(k)})$ is sufficient for θ based on $X^{(1)}, \dots, X^{(K)}$. As $T^{(k)}(\cdot)$ is shared with Example 4.1, we can apply Proposition 1 to thin the distribution of $\sum_{k=1}^K T^{(k)}(X^{(k)})$ by the function defined in (5).

In Appendix B.3 we show that $\exp\left(T(X^{(1)}, \dots, X^{(K)})\right) = \left(\prod_{k=1}^K X^{(k)}\right)^{1/K}$ follows a $\text{Gamma}(\theta, \beta)$ distribution. Thus, we can indirectly thin a $\text{Gamma}(\theta, \beta)$ random variable through $S(x) = \log(x)$ by $T(\cdot)$ defined in (5). This produces independent random variables $X^{(k)} \sim \text{Gamma}\left(\frac{1}{K}\theta + \frac{k-1}{K}, \frac{1}{K}\beta\right)$ for $k = 1, \dots, K$. Once again, noting that $S(\cdot)$ is invertible, we can instead directly thin $X \sim \text{Gamma}(\theta, \beta)$ by the function defined in (6).

To apply either of these thinning strategies to a $\text{Gamma}(\theta, \beta)$ random variable, we must sample from G_t as defined in Theorem 1. This is an unnamed distribution over the set of K positive real numbers with geometric mean t . We can sample from this unnamed distribution using numerical methods as detailed in Appendix B.3.

We emphasize that Example 4.2 is different from the gamma thinning example from Neufeld et al. (2023): that involves thinning a $\text{Gamma}(\alpha, \theta)$ random variable with α known, whereas here we thin a $\text{Gamma}(\theta, \beta)$ random variable with β known.

Examples 4.1 and 4.2 enable us to thin a random variable into an arbitrary number of folds. However, unlike in the examples seen in Section 3, the resulting folds are not identically distributed: they are independent beta or gamma distributions with slightly different parameters.

In Examples 4.1 and 4.2, the function $S(\cdot)$ through which we indirectly thin X is invertible. We now consider a case in which $S(\cdot)$ is not invertible, and furthermore, is not scalar-valued. Specifically, we let $X = (X_1, \dots, X_n)$ represent a sample of n independent and identically distributed normal observations with unknown mean θ_1 and variance θ_2 . Note that the sample mean and sample variance are jointly sufficient for θ_1 and θ_2 . We will now apply ideas from Theorem 2 and Proposition 1 to thin X through the sample mean and sample variance.

Example 4.3 (Indirect thinning of $N_n(\theta_1 1_n, \theta_2 I_n)$ through the sample mean and sample variance). We assume that $X \sim N_n(\theta_1 1_n, \theta_2 I_n)$. Then $S(X)$ is sufficient for $\theta = (\theta_1, \theta_2)$, where

$$S(x) = \left(\frac{1}{n} \sum_{i=1}^n x_i, \frac{1}{n-1} \sum_{i=1}^n \left(x_i - \frac{1}{n} \sum_{i'=1}^n x_{i'} \right)^2 \right)$$

is the function that computes the sample mean and sample variance of the entries of a vector x . We will indirectly thin X through $S(\cdot)$ into $K = n$ univariate normals. To do so, we start with $X^{(k)} \stackrel{\text{iid}}{\sim} N(\theta_1, \theta_2)$ for $k = 1, \dots, K$. A sufficient statistic for θ based on $(X^{(1)}, \dots, X^{(K)})$ is $T(X^{(1)}, \dots, X^{(K)})$, where

$$T(x^{(1)}, \dots, x^{(K)}) = S((x^{(1)}, \dots, x^{(K)})^\top),$$

i.e., we concatenate the K entries into a vector and apply $S(\cdot)$. Furthermore, $T(X^{(1)}, \dots, X^{(K)})$ has the same distribution as $S(X)$, since $(X^{(1)}, \dots, X^{(K)})^\top$ and X have the same distribution (recall that we have taken $K = n$ in this example). This establishes that we can indirectly thin X through $S(\cdot)$ by $T(\cdot)$.

By Theorem 1, we define G_t to be the conditional distribution of $(X^{(1)}, \dots, X^{(K)})$ given $T(X^{(1)}, \dots, X^{(K)}) = t$, *i.e.* it is the distribution of a $N_K(\theta_1 1_K, \theta_2 I_K)$ random vector conditional on its sample mean and sample variance equalling $t = (t_1, t_2)$. We will show by a symmetry argument that this conditional distribution is uniform over the set of points in \mathbb{R}^K with sample mean t_1 and sample variance t_2 . To see this, note that G_t cannot depend on θ (by sufficiency), so we can take $\theta = (t_1, t_2)$ and equivalently describe G_t as the distribution of a $N_K(t_1 1_K, t_2 I_K)$ random vector conditional on its sample mean and sample variance equalling $t = (t_1, t_2)$. Such a distribution has constant density on a sphere centered at $t_1 1_K$. Thus, the conditional distribution is uniform over the set of points in \mathbb{R}^K with sample mean t_1 and sample variance t_2 .

Finally, we obtain $(X^{(1)}, \dots, X^{(K)})$ by sampling from G_X .

In effect, Example 4.3 shows that given a realization of $X \sim N_n(\theta_1 1_n, \theta_2 I_n)$, we can generate a new random vector $(X^{(1)}, \dots, X^{(n)})^\top$ with the identical distribution, and with the same sample mean and sample variance, without knowledge of the true mean or true variance. This might have applications in cases where the true values of the observations cannot be shared. Furthermore, the ideas in Example 4.3 can be applied in settings where only the sufficient statistics of a realization of $X \sim N_n(\theta_1 1_n, \theta_2 I_n)$ are available, and we wish to generate a “plausible” sample that could have led to those sufficient statistics.

Indirect thinning is a very flexible approach, and we briefly list a few more examples illustrating its use.

Example 4.4 (Additional examples of indirect thinning).

1. Suppose we observe $X \sim N(\mu, \theta)$ where μ is known; here μ denotes the mean and θ the variance. Then $S(X) = (X - \mu)^2 \sim \theta \chi_1^2 = \text{Gamma}(\frac{1}{2}, \frac{1}{2\theta})$. Thus, by applying the Gamma thinning strategy of Neufeld et al. (2023) discussed in Example 3.2 to $S(X)$, we can indirectly thin a normal distribution with unknown variance through $S(\cdot)$.
2. Suppose we observe $X \sim \text{Weibull}(\theta, \nu)$ where ν is known. Then, $S(X) = X^\nu \sim \text{Exp}(\theta^{-\nu})$. Thus, by applying the Gamma thinning strategy of Example 3.2 or 3.3 to $S(X)$, we can indirectly thin a Weibull distribution with unknown rate through $S(\cdot)$.
3. Suppose we observe $X \sim \text{Pareto}(\nu, \theta)$ where ν is known. Then $S(X) = \log(X/\nu) \sim \text{Exp}(\theta)$. Thus, by applying the Gamma thinning strategy of Example 3.2 or 3.3 to $S(X)$, we can indirectly thin a Pareto distribution with unknown shape through $S(\cdot)$.

In Definition 1, we specified that the thinning function $T(\cdot)$ is deterministic. We conclude with a remark noting that indirect thinning can be thought of as a means of relaxing this requirement.

Remark 5 (Indirect thinning through non-invertible functions). *Consider the setting where we indirectly thin a random variable X through a non-invertible function $S(\cdot)$. This could instead be viewed as directly thinning X by some stochastic function $T'(\cdot)$ where the randomness is over the set of values for X that map to $S(X)$. However, as noted in Proposition 2, as $S(X)$ is a sufficient statistic for θ , this randomness offers no information about θ . For simplicity, we thus conclude that our proposed indirect thinning method is preferable to such a stochastic direct thinning.*

5 Thinning outside of exponential families

In this section, we focus on thinning families of distributions that are not exponential families. Outside of the exponential family, only certain distributions with domains that vary with the parameter of interest have sufficient statistics that are bounded as the sample size increases (Darmois 1935, Koopman 1936, Pitman 1936). Thus, we first consider a setting where θ alters the support of the distribution (Section 5.1), and then one where the sufficient statistic's dimension grows as the sample size increases (Section 5.2). In the latter, we discuss sample splitting as a special case of the generalized data thinning framework.

5.1 Thinning distributions with varying support

We consider two examples in which the parameter of interest, θ , changes the support of a distribution. In Example 5.1, θ scales the support.

Example 5.1 (Thinning $\text{Unif}(0, \theta)$). *We start with $X^{(k)} \stackrel{iid}{\sim} \theta \cdot \text{Beta}(\frac{1}{K}, 1)$ for $k = 1, \dots, K$, and note that $T(X^{(1)}, \dots, X^{(K)}) = \max(X^{(1)}, \dots, X^{(K)})$ is sufficient for θ . Furthermore, $\max(X^{(1)}, \dots, X^{(K)}) \sim \text{Unif}(0, \theta)$. Thus, we define G_t to be the conditional distribution of $(X^{(1)}, \dots, X^{(K)})$ given $\max(X^{(1)}, \dots, X^{(K)}) = t$. Then, by Theorem 1, we can thin $X \sim \text{Unif}(0, \theta)$ by sampling from G_X .*

To sample from this conditional distribution, we first draw $C \sim \text{Categorical}_K(1/K, \dots, 1/K)$. Then, $X^{(k)} = C_k X + (1 - C_k) Z_k$ where $Z_k \stackrel{iid}{\sim} \text{Beta}(\frac{1}{K}, 1)$.

This is a special case of Example B.1, given in Appendix B.4.1, in which we show that one can thin the scale family $\theta \cdot \text{Beta}(\alpha, 1)$ where α is known. Setting $\alpha = 1$ yields Example 5.1.

We now consider an example in which θ shifts the support. In particular, we consider $X \sim \text{SExp}(\theta, \lambda)$, which is the location family generated by shifting an exponential random variable by an amount θ . It has density

$$p_{\theta, \lambda}(x) = \lambda e^{-\lambda(x-\theta)} 1\{x \geq \theta\}.$$

Example 5.2 (Thinning a $\text{SExp}(\theta, \lambda)$ random variable with known λ). *We begin with $X^{(k)} \stackrel{iid}{\sim} \text{SExp}(\theta, \lambda/K)$ for $k = 1, \dots, K$, and note that $T(X^{(1)}, \dots, X^{(K)}) = \min(X^{(1)}, \dots, X^{(K)})$ is sufficient for θ . Furthermore, $\min(X^{(1)}, \dots, X^{(K)}) \sim \text{SExp}(\theta, \lambda)$. We define G_t to be*

the conditional distribution of $(X^{(1)}, \dots, X^{(K)})$ given $\min(X^{(1)}, \dots, X^{(K)}) = t$. Then, by Theorem 1, we can thin $X \sim \text{SExp}(\theta, \lambda)$ by sampling from G_X .

To sample from G_X , we first draw $C \sim \text{Categorical}_K(1/K, \dots, 1/K)$. We then take $X^{(k)} = X + (1 - C_k)Z_k$, where $Z_k \stackrel{iid}{\sim} \text{Exp}(\lambda/K)$.

Derivations associated with Example 5.2 can be found in Appendix B.4.2.

5.2 Sample splitting as a special case of generalized data thinning

We now consider sample splitting, a well-known approach for splitting a sample of observations into two or more sets (Cox 1975). We show that sample splitting can be viewed as an instance of generalized thinning. In this setting, $X = (X_1, \dots, X_n)$ is a sample of independent and identically distributed random variables, $X_i \in \mathcal{X}$, each having distribution $F \in \mathcal{F}$, where \mathcal{F} is some (potentially non-parametric) family of distributions and \mathcal{X} is the set of values that the random variable X_i can take on (most commonly $\mathcal{X} = \mathbb{R}^p$). That is, $X \sim P_F \in \mathcal{P}$, where

$$\mathcal{P} = \{F^n : F \in \mathcal{F}\},$$

and $F^n = F \times \dots \times F$ denotes the joint distribution of n independent random variables each drawn from F . We will indirectly thin X through a sorting function $S(\cdot)$ by a particular function $T(\cdot)$.

Example 5.3 (Sample splitting is a special case of generalized data thinning). *We begin with $X^{(k)} := (X_1^{(k)}, \dots, X_{n_k}^{(k)}) \stackrel{iid}{\sim} F^{n_k}$, for $k = 1, \dots, K$. Here, $n_1, \dots, n_K > 0$, and $\sum_{k=1}^K n_k = n$. That is, for $k = 1, \dots, K$, $X^{(k)} \in \mathcal{X}^{n_k}$ denotes a set of n_k independent and identically distributed draws from F .*

We thin $S(X)$, where $S : \mathcal{X}^n \rightarrow \mathcal{X}^n$ sorts the entries of its input based on their values. Furthermore, we define $T : \mathcal{X}^{n_1} \times \dots \times \mathcal{X}^{n_K} \rightarrow \mathcal{X}^n$ as

$$T(x^{(1)}, \dots, x^{(K)}) = S((x^{(1)}, \dots, x^{(K)})),$$

the function that concatenates its arguments and then applies the sorting function $S(\cdot)$. Then $T(X^{(1)}, \dots, X^{(K)})$ is a sufficient statistic for F , and furthermore, $T(X^{(1)}, \dots, X^{(K)}) \stackrel{D}{=} S(X)$.

We define G_t to be the conditional distribution of $(X^{(1)}, \dots, X^{(K)})$ given $T(X^{(1)}, \dots, X^{(K)}) = t$. Suppose we observe $X \sim F^n$. Then, by Theorem 1, we can indirectly thin X through $S(\cdot)$ by $T(\cdot)$ by sampling from G_X .

This conditional distribution is uniform over all $\frac{n!}{n_1! \dots n_K!}$ assignments of n items to K groups of sizes n_1, \dots, n_K . Thus, to sample from G_X , we randomly partition the sample of size n into K groups of sizes n_1, \dots, n_K . This is precisely the same as sample splitting.

6 Counterexamples

In this section, we present two examples in which thinning strategies do not work. The first involves a natural exponential family that is based on a distribution that *cannot* be written

as the convolution of two distributions. In this case, Theorem 2 implies that we cannot thin it by addition. In fact, we will prove a stronger statement: namely, that there does not exist *any* function $T(\cdot)$ that can thin it. The second example involves a convolution-closed family outside of the natural exponential family in which addition is not sufficient. In this case, taking $T(\cdot)$ to be addition does not enable thinning, as Theorem 1 does not apply.

6.1 The Bernoulli family cannot be thinned

Let P_θ denote the Bernoulli(θ) distribution, where θ is the probability of success. While this distribution can be written as a natural exponential family (with natural parameter $\log(\frac{\theta}{1-\theta})$), it cannot be written as the sum of two independent, non-degenerate random variables. Thus, Theorem 2 does not apply, i.e. it cannot be thinned by addition into natural exponential families.

In this section, we prove a stronger result—namely, that there is *no function* $T(\cdot)$ that enables us to thin P_θ into distributions $Q_\theta^{(1)}$ and $Q_\theta^{(2)}$ with well-defined, positive Fisher information.

Based on Theorem 1 from Section 2, we can thin $X \sim P_\theta$ into independent $Q_\theta^{(1)}$ and $Q_\theta^{(2)}$ if and only if there exists a function $T(\cdot)$ such that $X = T(X^{(1)}, X^{(2)})$ and $T(X^{(1)}, X^{(2)})$ is sufficient for θ in the joint distribution of $X^{(1)}$ and $X^{(2)}$. The next result says that if such $Q_\theta^{(1)}$ and $Q_\theta^{(2)}$ have positive Fisher information, then no such function $T(\cdot)$ exists.

Theorem 3. *Suppose that*

- $T(X^{(1)}, X^{(2)}) \sim \text{Bernoulli}(\theta)$,
- $X^{(1)} \sim Q_\theta^{(1)}, X^{(2)} \sim Q_\theta^{(2)}$, and $X^{(1)}$ and $X^{(2)}$ are independent, where $Q_\theta^{(1)}$ and $Q_\theta^{(2)}$ are from families where Fisher information exists, and
- $I_{X^{(1)}}(\theta) > 0$ and $I_{X^{(2)}}(\theta) > 0$.

Then $T(X^{(1)}, X^{(2)})$ is not sufficient for θ based on $X^{(1)}$ and $X^{(2)}$.

The proof of Theorem 3 is in Appendix A.4. In summary, Bernoulli is a natural exponential family that cannot be thinned by any function into any distribution with well-defined Fisher information. A similar argument reveals that the categorical distribution (i.e., a multinomial with a single trial) also cannot be thinned into distributions with well-defined, positive Fisher information.

6.2 Not all convolution-closed families can be thinned by addition

Suppose now that our interest lies in a random variable $X = T(X^{(1)}, X^{(2)})$, where $T(X^{(1)}, X^{(2)})$ is *not* sufficient for the parameter θ based on $(X^{(1)}, X^{(2)})$. This means that the conditional distribution of $(X^{(1)}, X^{(2)})$ given $T(X^{(1)}, X^{(2)})$ depends on θ , and thus that we cannot thin X by $T(\cdot)$. We see this in the following example.

Example 6.1 (The trouble with thinning $\text{Cauchy}(\theta_1, \theta_2)$ by addition). *Recall that the Cauchy family, $\text{Cauchy}(\theta_1, \theta_2)$, indexed by $\theta = (\theta_1, \theta_2)$, is convolution-closed. In particular, if $X^{(1)}, X^{(2)} \stackrel{iid}{\sim} \text{Cauchy}(\frac{1}{2}\theta_1, \frac{1}{2}\theta_2)$, then $X^{(1)} + X^{(2)} \sim \text{Cauchy}(\theta_1, \theta_2)$. It is tempting therefore to try thinning this family by $T(x^{(1)}, x^{(2)}) = x^{(1)} + x^{(2)}$. However, the sum $X^{(1)} + X^{(2)}$ is not sufficient for θ , which means that Theorem 1 does not apply and in particular G_t , the conditional distribution of $(X^{(1)}, X^{(2)})$ given $X^{(1)} + X^{(2)} = t$, is a function of θ . Therefore, we cannot thin the Cauchy family with unknown parameters by addition.*

In fact, we can take this result a step further: for a sample of Cauchy random variables, the only sufficient statistic for θ is the vector of order statistics (Casella & Berger 2002, p. 275). Thus, there is no family \mathcal{P} and corresponding function $T(\cdot)$ that reduces the data for which thinning will produce independent Cauchy random variables. However, sample splitting (which as shown in Section 5.2 is based on a $T(\cdot)$ that does not reduce the data) where X is a sample of K independent Cauchy random variables would lead to thinning into K independent Cauchy random variables.

7 Numerical experiments

In this section, we illustrate some of the examples from Sections 3 and 4 through numerical simulations. Specifically, we thin a $\text{Gamma}(\alpha, \theta)$ distribution using the three different approaches described in Examples 3.2, 3.3, and 3.4, a $\text{Beta}(\theta, \beta)$ into two non-identical beta random variables as described in Example 4.1, and a $\text{Unif}(0, \theta)$ into scaled betas as described in Example 5.1. We take $K = 2$ throughout for ease of presentation.

In Figure 2, each row corresponds to one of the examples mentioned above. In the left-hand column, we display the empirical density of $B = 100,000$ realizations, x_b (for $b = 1, \dots, B$), of the $X \sim P_\theta$ that we wish to thin, overlaid with the true density of P_θ . In the center-left and center-right columns, we display B realizations of $(X^{(1)}, X^{(2)})$, where each realization $(x_b^{(1)}, x_b^{(2)})$ is obtained by sampling from G_{x_b} , the conditional distribution of $(X^{(1)}, X^{(2)})$ given $T(X^{(1)}, X^{(2)}) = x_b$. (This sampling is done without knowledge of θ .) We overlay the densities of the marginals $Q_\theta^{(1)}$ and $Q_\theta^{(2)}$. The right-hand column displays the empirical joint distribution of $(Q_\theta^{(1)}(X^{(1)}), Q_\theta^{(2)}(X^{(2)}))$.

In each case, our empirical findings corroborate our theoretical results: we see that the empirical distribution of $X \sim P_\theta$ agrees with its theoretical density (left-hand panel); that the empirical distributions of $X^{(1)}$ and $X^{(2)}$ sampled from G_X agree with $Q_\theta^{(1)}$ and $Q_\theta^{(2)}$ (even though the empirical distributions were obtained without knowledge of θ ; center-left and center-right columns); and that the joint distribution of $Q_\theta^{(1)}(X^{(1)})$ and $Q_\theta^{(2)}(X^{(2)})$ resembles a bivariate uniform (right-hand column).




Figure 2: Numerical examples of data thinning. The left-hand column displays a sample from P_{θ} , which we wish to thin. The center-left and center-right columns display the empirical distributions of $X^{(1)}$ and $X^{(2)}$ that result from thinning, overlaid with the theoretical distributions $Q_{\theta}^{(1)}$ and $Q_{\theta}^{(2)}$. The right-hand column displays the empirical joint distribution of $(Q_{\theta}^{(1)}(X^{(1)}), Q_{\theta}^{(2)}(X^{(2)}))$, providing visual evidence that they are independent.

8 Discussion

Our generalized data thinning proposal encompasses a diverse set of existing approaches for splitting a random variable into independent random variables, from convolution-closed data thinning (Neufeld et al. 2023) to sample splitting (Cox 1975). It provides a lens through which these existing approaches follow from the same simple principle — sufficiency — and can be derived through the same simple recipe (Algorithm 1).

The principle of sufficiency is key to generalized data thinning, as it enables a sampling mechanism that does not depend on unknown parameters. When no sufficient statistic that reduces the data is available, as in the non-parametric setting of Section 5.2 and the Cauchy example of Section 6.2, then sample splitting is still possible. Conversely, in a setting with $n = 1$ or where the elements of $X = (X_1, \dots, X_n)$ are not independent and identically distributed, sample splitting may not be possible, but other generalized thinning approaches may be available.

For example, consider a regression setting with a fixed design, in which each response Y_i has a potentially distinct distribution determined by its corresponding feature vector x_i , for $i = 1, \dots, n$. It is typical to recast this as random pairs $(x_1, Y_1), \dots, (x_n, Y_n)$ that are independent and identically distributed from some joint distribution, thereby justifying sample splitting. However, this amounts to viewing the model as arising from a random design, which may not match the reality of how the design matrix was generated, and may not be well-aligned with the goals of the data analysis. For instance, recall the example given in the introduction: given a dataset consisting of the $n = 50$ states of the United States, it is unrealistic to treat each state as an independent and identically distributed draw, and undesirable to perform inference only on the states that were “held out” of training. In this example, generalized data thinning could provide a more suitable alternative to sample splitting that stays true to the fixed design model underlying the data.

The starting place for any generalized thinning strategy—whether sampling splitting or otherwise—is the modeling assumption that the data are drawn from a distribution belonging to a family \mathcal{P} . The specification of the family \mathcal{P} allows us to identify the thinning function $T(\cdot)$ and sampling mechanism G_t . An important topic of future study is understanding the effect of model misspecification. In particular, if we assume that $X \sim P_\theta \in \mathcal{P}$, but in fact X ’s distribution belongs to a different family $\tilde{\mathcal{P}}$, what goes wrong? The random variables $X^{(1)}, \dots, X^{(K)}$ that we generate by thinning will still satisfy the property that $X = T(X^{(1)}, \dots, X^{(K)})$; however, $X^{(1)}, \dots, X^{(K)}$ may not be independent and may no longer have the intended marginals $Q_\theta^{(1)}, \dots, Q_\theta^{(K)}$. Can we quantify the effect of the model misspecification? I.e., if the true family is “close” to the assumed family, will $X^{(1)}, \dots, X^{(K)}$ be only weakly dependent, and will the marginals be close to $Q_\theta^{(1)}, \dots, Q_\theta^{(K)}$? Some initial answers to these questions can be found in Neufeld et al. (2023) and Rasines & Young (2022).

In the introduction, we noted that generalized thinning with $K = 2$ is a (U, V) -decomposition, as defined in Rasines & Young (2022). We elaborate on that connection here. The (U, V) -decomposition seeks independent random variables $U = u(X, W)$ and $V = v(X, W)$ such that U and V are jointly sufficient for the unknowns, where W is a random variable possibly depending on X . Suppose we can indirectly thin X through $S(\cdot)$ by $T(\cdot)$. This means we

have produced independent random variables $X^{(1)}$ and $X^{(2)}$ for which $S(X) = T(X^{(1)}, X^{(2)})$. Since $S(X)$ is sufficient for θ on the basis of X , this implies that $(X^{(1)}, X^{(2)})$ is jointly sufficient for θ . It follows that $(X^{(1)}, X^{(2)})$ is a (U, V) -decomposition of X . It is of interest to investigate whether there are (U, V) -decompositions that cannot be achieved through either direct or indirect generalized thinning.

Acknowledgments

We thank Nicholas Irons for helpful discussions that contributed to the results in Section 6.1. We acknowledge funding from the following sources: NIH R01 EB026908, NIH R01 DA047869, a Simons Investigator Award in Mathematical Modeling of Living Systems, and the Keck Foundation to DW; NIH R01 GM123993 to DW and JB; an Natural Sciences and Engineering Research Council of Canada Discovery Grant to LG.

References

- Abramowitz, M., Stegun, I. A. & Romer, R. H. (1972), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, American Association of Physics Teachers.
- Casella, G. & Berger, R. (2002), *Statistical Inference*, Thomson Learning.
- Chen, S. & Bien, J. (2020), ‘Valid inference corrected for outlier removal’, *Journal of Computational and Graphical Statistics* **29**(2), 323–334.
- Chen, Y. T. & Witten, D. M. (2022), ‘Selective inference for k-means clustering’, *arXiv preprint arXiv:2203.15267*.
- Cox, D. R. (1975), ‘A note on data-splitting for the evaluation of significance levels’, *Biometrika* **62**(2), 441–444.
- Darmois, G. (1935), ‘Sur les lois de probabilité à estimation exhaustive’, *Comptes Rendus de l’Académie des Sciences* **200**, 1265–1266.
- Feller, W. (1971), *An Introduction to Probability Theory and Its Applications. Vol. 2*, J. Willey and Sons, New York.
- Fithian, W., Sun, D. & Taylor, J. (2014), ‘Optimal inference after model selection’, *arXiv preprint arXiv:1410.2597*.
- Gao, L. L., Bien, J. & Witten, D. (2022), ‘Selective inference for hierarchical clustering’, *Journal of the American Statistical Association*.
- Hastie, T., Tibshirani, R., Friedman, J. H. & Friedman, J. H. (2009), *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*, Vol. 2, Springer.
- Hyun, S., Lin, K. Z., G’Sell, M. & Tibshirani, R. J. (2021), ‘Post-selection inference for changepoint detection algorithms with application to copy number variation data’, *Biometrics* **77**(3), 1037–1049.
- Jewell, S., Fearnhead, P. & Witten, D. (2022), ‘Testing for a change in mean after changepoint detection’, *Journal of the Royal Statistical Society Series B: Statistical Methodology* **84**(4), 1082–1104.
- Jørgensen, B. (1992), ‘Exponential dispersion models and extensions: A review’, *International Statistical Review/Revue Internationale de Statistique* pp. 5–20.
- Jørgensen, B. & Song, P. X.-K. (1998), ‘Stationary time series models with exponential dispersion model margins’, *Journal of Applied Probability* **35**(1), 78–92.
- Koopman, B. O. (1936), ‘On distributions admitting a sufficient statistic’, *Transactions of the American Mathematical Society* **39**(3), 399–409.

- Lee, J. D., Sun, D. L., Sun, Y. & Taylor, J. E. (2016), ‘Exact post-selection inference, with application to the lasso’, *The Annals of Statistics* **44**(3), 907 – 927.
- Lehmann, E. L. & Romano, J. P. (2005), *Testing Statistical Hypotheses: Third Edition*, Vol. 3, Springer.
- Leiner, J., Duan, B., Wasserman, L. & Ramdas, A. (2022), ‘Data fission: sample splitting a single sample’, *arXiv preprint arXiv:2112.11079* .
- Neufeld, A., Dharamshi, A., Gao, L. L. & Witten, D. (2023), ‘Data thinning for convolution-closed distributions’, *arXiv preprint arXiv:2301.07276* .
- Neufeld, A., Gao, L. L., Popp, J., Battle, A. & Witten, D. (2022), ‘Inference after latent variable estimation for single-cell RNA sequencing data’, *To appear in Biostatistics; arXiv preprint arXiv:2207.00554* .
- Oliveira, N. L., Lei, J. & Tibshirani, R. J. (2021), ‘Unbiased risk estimation in the normal means problem via coupled bootstrap techniques’, *arXiv preprint arXiv:2111.09447* .
- Pitman, E. J. G. (1936), ‘Sufficient statistics and intrinsic accuracy’, *Mathematical Proceedings of the Cambridge Philosophical Society* **32**(4), 567–579.
- Rasines, D. G. & Young, G. A. (2022), ‘Splitting strategies for post-selection inference’, *Biometrika* .
- Taylor, J. & Tibshirani, R. (2018), ‘Post-selection inference for-penalized likelihood models’, *Canadian Journal of Statistics* **46**(1), 41–61.
- Taylor, J. & Tibshirani, R. J. (2015), ‘Statistical learning and selective inference’, *Proceedings of the National Academy of Sciences* **112**(25), 7629–7634.
- Tian, X. (2020), ‘Prediction error after model search’, *The Annals of Statistics* **48**(2), 763 – 784.
- Tian, X. & Taylor, J. (2017), ‘Asymptotics of selective inference’, *Scandinavian Journal of Statistics* **44**(2), 480–499.
- Tian, X. & Taylor, J. (2018), ‘Selective inference with a randomized response’, *The Annals of Statistics* **46**(2), 679–710.
- Tibshirani, R. J., Rinaldo, A., Tibshirani, R. & Wasserman, L. (2018), ‘Uniform asymptotic inference and the bootstrap after model selection’, *The Annals of Statistics* **46**(3), 1255–1287.
- Yun, Y. & Barber, R. F. (2023), ‘Selective inference for clustering with unknown variance’, *arXiv preprint arXiv:2301.12999* .

Appendix

A Proofs

A.1 Proof of Theorem 1

Proof. By their construction in Definition 1, the random variables $(X^{(1)}, \dots, X^{(K)}) \sim Q_\theta^{(1)} \times \dots \times Q_\theta^{(K)}$ have conditional distribution

$$(X^{(1)}, \dots, X^{(K)}) | \{X = t\} \sim G_t.$$

Furthermore, Definition 1 tells us that $X = T(X^{(1)}, \dots, X^{(K)})$. This means that

$$(X^{(1)}, \dots, X^{(K)}) | \{T(X^{(1)}, \dots, X^{(K)}) = t\} \sim G_t,$$

which establishes part (b) of the theorem.

The distribution G_t in Definition 1 does not depend on θ (note that it is associated with the entire family \mathcal{P} , not a particular distribution P_θ). By the definition of sufficiency, the fact that the conditional distribution $(X^{(1)}, \dots, X^{(K)}) | T(X^{(1)}, \dots, X^{(K)})$ does not depend on θ implies that $T(X^{(1)}, \dots, X^{(K)})$ is sufficient for θ . This proves (a). \square

A.2 Proof of Theorem 2

Proof. We start by proving the \Leftarrow direction. Suppose H is the convolution of H_1, \dots, H_K . We follow the recipe given in Algorithm 1:

1. We choose $Q^{(k)} = \mathcal{P}^{H_k}$ for $k = 1, \dots, K$.
2. Let $(X^{(1)}, \dots, X^{(K)}) \sim P_\theta^{H_1} \times \dots \times P_\theta^{H_K}$. This joint distribution satisfies

$$\prod_{k=1}^K dP_\theta^{H_k}(x^{(k)}) = \exp \left\{ \left(\sum_{k=1}^K x^{(k)} \right)^\top \theta - \sum_{k=1}^K \psi_{H_k}(\theta) \right\} \prod_{k=1}^K dH_k(x^{(k)}).$$

By the factorization theorem, we find that $T(X^{(1)}, \dots, X^{(K)}) = \sum_{k=1}^K X^{(k)}$ is sufficient for θ .

3. It remains to determine the distribution of $U = T(X^{(1)}, \dots, X^{(K)})$. This random variable is the convolution of $P_\theta^{H_1} \times \dots \times P_\theta^{H_K}$, and its distribution μ is defined by the

following K -way integral:

$$\begin{aligned}
d\mu(u) &= \int \cdots \int 1 \left\{ \sum_{k=1}^K x^{(k)} = u \right\} \prod_{k=1}^K dP_{\theta}^{H_k}(x^{(k)}) \\
&= \exp \left\{ u^{\top} \theta - \sum_{k=1}^K \psi_{H_k}(\theta) \right\} \int \cdots \int 1 \left\{ \sum_{k=1}^K x^{(k)} = u \right\} \prod_{k=1}^K dH_k(x^{(k)}) \\
&= \exp \{ u^{\top} \theta - \psi_H(\theta) \} dH(u) \\
&= dP_{\theta}^H(u),
\end{aligned}$$

where in the second-to-last equality we use the assumption that H is the K -way convolution of H_1, \dots, H_K and the fact that the moment generating function of a convolution is the product of the individual moment generating functions (and recalling that ψ_H is the logarithm of the moment generating function of H). This establishes that $T(X^{(1)}, \dots, X^{(K)}) \sim P_{\theta}^H$. By Theorem 1, the family \mathcal{P}^H is thinned by this choice of $T(\cdot)$.

We now prove the \implies direction. Suppose that \mathcal{P}^H can be K -way thinned into $\mathcal{P}^{H_1}, \dots, \mathcal{P}^{H_K}$ using the summation function. Then applying Definition 1 with $\theta = 0$, we can take $X \sim P_0^H$ and produce $(X^{(1)}, \dots, X^{(K)}) \sim P_0^{H_1} \times \cdots \times P_0^{H_K}$ for which $X = X^{(1)} + \cdots + X^{(K)}$. Noting that $P_0^{H_k} = H_k$ for all k and $P_0^H = H$, this proves that H is a K -way convolution of H_1, \dots, H_K . \square

A.3 Proof of Proposition 2

Proof. The result follows from a chain of equalities:

$$\begin{aligned}
I_X(\theta) &= I_{S(X)}(\theta) \\
&= I_{T(X^{(1)}, \dots, X^{(K)})}(\theta) \\
&= I_{(X^{(1)}, \dots, X^{(K)})}(\theta) \\
&= \sum_{k=1}^K I_{X^{(k)}}(\theta).
\end{aligned}$$

The first equality is true because $S(X)$ is sufficient for θ based on X . The second equality follows from the definition of thinning $S(X)$ into $X^{(1)}, \dots, X^{(K)}$ using $T(\cdot)$. The third equality follows from Theorem 1, which tells us that $T(X^{(1)}, \dots, X^{(K)})$ is sufficient for θ based on $(X^{(1)}, \dots, X^{(K)})$. The final equality follows from independence. \square

A.4 Proof of Theorem 3

Proof. Suppose $0.5 < \theta < 1$. We prove the result assuming that $0.5 < \theta < 1$ since if $T(X^{(1)}, X^{(2)})$ cannot be sufficient when $\theta > 0.5$, it cannot be sufficient when $\theta \in (0, 1)$.

As $T(X^{(1)}, X^{(2)})$ is Bernoulli, we can write it as an indicator $T(X^{(1)}, X^{(2)}) = I((X^{(1)}, X^{(2)}) \in \mathcal{C})$ for some set \mathcal{C} . Let \mathcal{C}_1 denote the projection of \mathcal{C} onto the $X^{(1)}$ axis, meaning that $(X^{(1)}, X^{(2)}) \in \mathcal{C}$ implies $X^{(1)} \in \mathcal{C}_1$.

Next, let $U(X^{(1)}) = I(X^{(1)} \in \mathcal{C}_1)$. Note that this is also a Bernoulli random variable, and note that its success probability is $\theta + \epsilon$, where $\epsilon \geq 0$, because $(X^{(1)}, X^{(2)}) \in \mathcal{C}$ implies $X^{(1)} \in \mathcal{C}_1$.

First, suppose that $\theta + \epsilon = 1$. In this case, the function $T(X^{(1)}, X^{(2)})$ does not depend on $X^{(1)}$, and thus can be written as $\tilde{T}(X^{(2)})$. In this case, note that

$$\begin{aligned} I_{T(X^{(1)}, X^{(2)})}(\theta) &= I_{\tilde{T}(X^{(2)})}(\theta) \\ &\leq I_{X^{(2)}}(\theta) \\ &< I_{X^{(1)}}(\theta) + I_{X^{(2)}}(\theta) = I_{X^{(1)}, X^{(2)}}(\theta). \end{aligned}$$

The inequality in the second line follows because a statistic of a random variable can never contain more information than the variable itself. The strict inequality in the third line follows from our assumption that $I_{X^{(1)}}(\theta) > 0$, and the final equality follows from independence of $X^{(1)}$ and $X^{(2)}$. We conclude that $T(X^{(1)}, X^{(2)})$ is not sufficient for θ , because it contains strictly less information than $(X^{(1)}, X^{(2)})$.

Next, suppose that $\theta + \epsilon < 1$. Direct calculation yields that $I_{T(X^{(1)}, X^{(2)})}(\theta) = \frac{1}{\theta(1-\theta)}$, and that $I_{U(X^{(1)})}(\theta) = \frac{1}{(\theta+\epsilon)(1-(\theta+\epsilon))}$. Because we assumed that $0.5 \leq \theta \leq \theta + \epsilon < 1$, we see that $I_{T(X^{(1)}, X^{(2)})}(\theta) \leq I_{U(X^{(1)})}(\theta)$. This then implies that

$$\begin{aligned} I_{T(X^{(1)}, X^{(2)})}(\theta) &\leq I_{U(X^{(1)})}(\theta) \\ &\leq I_{X^{(1)}}(\theta) \\ &< I_{X^{(1)}}(\theta) + I_{X^{(2)}}(\theta) = I_{X^{(1)}, X^{(2)}}(\theta). \end{aligned}$$

The inequality in the second line follows from the fact that a statistic of a random variable can never contain more information than the variable itself. The strict inequality in the third line follows from our assumption that $I_{X^{(2)}}(\theta) > 0$. We conclude that the function $T(X^{(1)}, X^{(2)})$ cannot be sufficient, because it contains strictly less information about θ than $(X^{(1)}, X^{(2)})$. \square

B Derivations of decompositions

B.1 Weibull distribution

Proof of Example 3.4. We must prove that if $X^{(k)} \stackrel{iid}{\sim} \text{Weibull}(\lambda, \nu)$, for $k = 1, \dots, K$, then $\sum_{k=1}^K (X^{(k)})^\nu \sim \text{Gamma}(K, \lambda^{-\nu})$.

Recalling that the gamma distribution is convolution-closed in its shape parameter, it is sufficient to show for a single $X^{(k)} \sim \text{Weibull}(\lambda, \nu)$ random variable that $(X^{(k)})^\nu \sim \text{Gamma}(1, \lambda^{-\nu}) = \text{Exp}(\lambda^{-\nu})$, where $\nu > 0$. Denote $Z = (X^{(k)})^\nu$. Then,

$$\begin{aligned}
f_Z(z) &= f_{X^{(k)}}\left(z^{\frac{1}{\nu}}\right) \left| \frac{dx^{(k)}}{dz} \right| \\
&\propto \left(z^{\frac{1}{\nu}}\right)^{\nu-1} \exp\left(-\left(\frac{z^{\frac{1}{\nu}}}{\lambda}\right)^\nu\right) \left| \frac{1}{\nu} z^{-\frac{\nu-1}{\nu}} \right| \\
&\propto \exp(-\lambda^{-\nu} z).
\end{aligned}$$

The above implies that $(X^{(k)})^\nu \sim \text{Exp}(\lambda^{-\nu})$, and thus $\sum_{k=1}^K (X^{(k)})^\nu \sim \text{Gamma}(K, \lambda^{-\nu})$ as required. □

B.2 Beta distribution

Proof of Example 4.1. We must prove the following three claims:

1. If $X^{(k)} \sim \text{Beta}\left(\frac{1}{K}\theta + \frac{k-1}{K}, \frac{1}{K}\beta\right)$, for $k = 1, \dots, K$, and $X^{(k)}$ are mutually independent, then $\left(\prod_{k=1}^K X^{(k)}\right)^{\frac{1}{K}} \sim \text{Beta}(\theta, \beta)$.

Recall that the beta distribution is fully characterised by its moments due to its finite support (Feller 1971). Also recall that the expectation of the r th power of a $X \sim \text{Beta}(\theta, \beta)$ random variable is $E[X^r] = \frac{B(\theta+r, \beta)}{B(\theta, \beta)}$ where B is the beta function. Finally, note that the Gauss multiplication theorem (Abramowitz et al. 1972, page 256) says that

$$\prod_{k=1}^K \Gamma\left(z + \frac{k-1}{K}\right) = (2\pi)^{\frac{K-1}{2}} K^{\frac{1}{2}-Kz} \Gamma(Kz).$$

Then, the r th moment of $\left(\prod_{k=1}^K X^{(k)}\right)^{\frac{1}{K}}$ is

$$\begin{aligned}
E \left[\left(\left(\prod_{k=1}^K X^{(k)} \right)^{\frac{1}{K}} \right)^r \right] &= \prod_{k=1}^K E \left[(X^{(k)})^{\frac{r}{K}} \right] \\
&= \prod_{k=1}^K \frac{B(\frac{1}{K}\theta + \frac{k-1}{K} + \frac{r}{K}, \frac{1}{K}\beta)}{B(\frac{1}{K}\theta + \frac{k-1}{K}, \frac{1}{K}\beta)} \\
&= \prod_{k=1}^K \frac{\Gamma(\frac{1}{K}\theta + \frac{k-1}{K} + \frac{r}{K})\Gamma(\frac{1}{K}\beta)}{\Gamma(\frac{1}{K}\theta + \frac{k-1}{K} + \frac{r}{K} + \frac{1}{K}\beta)} \\
&= \prod_{k=1}^K \frac{\Gamma(\frac{1}{K}\theta + \frac{k-1}{K})\Gamma(\frac{1}{K}\beta)}{\Gamma(\frac{1}{K}\theta + \frac{k-1}{K} + \frac{1}{K}\beta)} \\
&= \prod_{k=0}^{K-1} \frac{\Gamma(\frac{1}{K}\theta + \frac{r}{K} + \frac{k}{K})\Gamma(\frac{1}{K}\theta + \frac{1}{K}\beta + \frac{k}{K})}{\Gamma(\frac{1}{K}\theta + \frac{1}{K}\beta + \frac{r}{K} + \frac{k}{K})\Gamma(\frac{1}{K}\theta + \frac{k}{K})} \\
&= \frac{\left[\prod_{k=0}^{K-1} \Gamma(\frac{1}{K}\theta + \frac{r}{K} + \frac{k}{K}) \right] \left[\prod_{k=0}^{K-1} \Gamma(\frac{1}{K}\theta + \frac{1}{K}\beta + \frac{k}{K}) \right]}{\left[\prod_{k=0}^{K-1} \Gamma(\frac{1}{K}\theta + \frac{1}{K}\beta + \frac{r}{K} + \frac{k}{K}) \right] \left[\prod_{k=0}^{K-1} \Gamma(\frac{1}{K}\theta + \frac{k}{K}) \right]} \\
&= \frac{\left[(2\pi)^{\frac{K-1}{2}} K^{\frac{1}{2}-(\theta+r)} \Gamma(\theta+r) \right] \left[(2\pi)^{\frac{K-1}{2}} K^{\frac{1}{2}-(\theta+\beta)} \Gamma(\theta+\beta) \right]}{\left[(2\pi)^{\frac{K-1}{2}} K^{\frac{1}{2}-(\theta+\beta+r)} \Gamma(\theta+\beta+r) \right] \left[(2\pi)^{\frac{K-1}{2}} K^{\frac{1}{2}-\theta} \Gamma(\theta) \right]} \\
&= \frac{\Gamma(\theta+r)\Gamma(\beta)\Gamma(\theta+\beta)}{\Gamma(\theta+\beta+r)\Gamma(\theta)\Gamma(\beta)} \\
&= \frac{B(\theta+r, \beta)}{B(\theta, \beta)}.
\end{aligned}$$

This matches the moments of a $\text{Beta}(\theta, \beta)$ distribution, implying that $\left(\prod_{k=1}^K X^{(k)} \right)^{\frac{1}{K}} \sim \text{Beta}(\theta, \beta)$ as required.

2. A sufficient statistic for θ in the joint distribution of $X^{(1)}, \dots, X^{(K)}$ is

$$T(X^{(1)}, \dots, X^{(K)}) = \left(\prod_{k=1}^K X^{(k)} \right)^{\frac{1}{K}}.$$

By the mutual independence of $X^{(k)}$, the joint density of $X^{(1)}, \dots, X^{(K)}$ can be written as

$$\begin{aligned}
f_{X^{(1)}, \dots, X^{(K)}}(x^{(1)}, \dots, x^{(K)}) &= \prod_{k=1}^K f_{X^{(k)}}(x^{(k)}) \\
&\propto \prod_{k=1}^K (x^{(k)})^{\frac{1}{K}\theta + \frac{k-1}{K} - 1} (1 - x^{(k)})^{\frac{1}{K}\beta - 1} \\
&= \left[\left(\prod_{k=1}^K x^{(k)} \right)^{\frac{1}{K}} \right]^{\theta} \left[\prod_{k=1}^K (x^{(k)})^{\frac{k-1}{K} - 1} \right] \left[\prod_{k=1}^K (1 - x^{(k)}) \right]^{\frac{1}{K}\beta - 1}.
\end{aligned}$$

By the factorization theorem, $T(X^{(1)}, \dots, X^{(K)}) = \left(\prod_{k=1}^K X^{(k)} \right)^{\frac{1}{K}}$ is a sufficient statistic for θ .

3. To sample from G_t , i.e., the distribution of $(X^{(1)}, \dots, X^{(K)}) | T(X^{(1)}, \dots, X^{(K)}) = t$, we first sample from $(X^{(1)}, \dots, X^{(K-1)}) | T(X^{(1)}, \dots, X^{(K)}) = t$ and then recover $X^{(K)}$. We will show that the conditional density $f_{X^{(1)}, \dots, X^{(K-1)} | T(X^{(1)}, \dots, X^{(K)}) = t}(x^{(1)}, \dots, x^{(K-1)})$ is, up to a normalization constant involving t ,

$$\left[\prod_{k=1}^{K-1} (x^{(k)})^{\frac{k-K}{K} - 1} \right] \left[\left(\prod_{k=1}^{K-1} (1 - x^{(k)}) \right) \left(1 - \frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}} \right) \right]^{\frac{1}{K}\beta - 1}.$$

We derive this as follows (where any factors not involving $x^{(1)}, \dots, x^{(K-1)}$ are omitted, and we write $\theta_k = \frac{\theta}{K} + \frac{k-1}{K}$):

$$\begin{aligned}
&f_{X^{(1)}, \dots, X^{(K-1)} | T(X^{(1)}, \dots, X^{(K)}) = t}(x^{(1)}, \dots, x^{(K-1)}) \\
&\propto f_{X^{(1)}, \dots, X^{(K-1)}, T(X^{(1)}, \dots, X^{(K)})}(x^{(1)}, \dots, x^{(K-1)}, t) \\
&= f_{X^{(1)}, \dots, X^{(K)}} \left(x^{(1)}, \dots, x^{(K-1)}, \frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}} \right) \left| \frac{\partial}{\partial t} \frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}} \right| \\
&= \left(\prod_{k=1}^{K-1} f_{X^{(k)}}(x^{(k)}) \right) f_{X^{(K)}} \left(\frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}} \right) \left| \frac{K t^{K-1}}{\prod_{k=1}^{K-1} x^{(k)}} \right| \\
&\propto \left(\prod_{k=1}^{K-1} (x^{(k)})^{\theta_k - 1} (1 - x^{(k)})^{\frac{1}{K}\beta - 1} \right) \left(\frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}} \right)^{\theta_{K-1}} \left(1 - \frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}} \right)^{\frac{1}{K}\beta - 1} \frac{1}{\prod_{k=1}^{K-1} x^{(k)}} \\
&\propto \left[\prod_{k=1}^{K-1} (x^{(k)})^{\theta_k - \theta_{K-1}} \right] \left[\left(\prod_{k=1}^{K-1} (1 - x^{(k)}) \right) \left(1 - \frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}} \right) \right]^{\frac{1}{K}\beta - 1}.
\end{aligned}$$

It remains to note that $\theta_k - \theta_{K-1} = \frac{k-K}{K}$.

After sampling from $(X^{(1)}, \dots, X^{(K-1)}) | T(X^{(1)}, \dots, X^{(K)}) = t$, we compute $X^{(K)} = \frac{t^K}{\prod_{k=1}^{K-1} X^{(k)}}$.

□

B.3 Gamma distribution

Proof of Example 4.2. We must prove the following three claims:

1. If for $k = 1, \dots, K$, $X^{(k)} \sim \text{Gamma}(\frac{1}{K}\theta + \frac{k-1}{K}, \frac{1}{K}\beta)$ and $X^{(k)}$ are mutually independent, then $(\prod_{k=1}^K X^{(k)})^{\frac{1}{K}} \sim \text{Gamma}(\theta, \beta)$.

Rather than work with gamma random variables directly, we will find it convenient to work with the logarithm of gamma random variables. We start by deriving the moment generating function of this distribution.

Lemma 1. Consider a random variable Y such that $e^Y \sim \text{Gamma}(\theta, \beta)$. Then the moment generating function of Y exists in a neighborhood around 0 and is given by

$$\Phi_Y(t) = \frac{\Gamma(\theta + t)}{\Gamma(\theta)\beta^t}.$$

Proof of Lemma 1. The density of Y is given by

$$\begin{aligned} f_Y(y) &= f_{\text{Gamma}(\theta, \beta)}(e^y)e^y \\ &= \frac{\beta^\theta}{\Gamma(\theta)} e^{y(\theta-1)} e^{-\beta e^y} e^y \\ &= \frac{\beta^\theta}{\Gamma(\theta)} e^{y\theta - \beta e^y}, \end{aligned}$$

where the extra factor of e^y in the first equality is the Jacobian of the transformation. For $t > -\theta$, the moment generating function for this random variable is given by

$$\begin{aligned} \Phi_Y(t) &= \mathbb{E}[e^{tY}] \\ &= \int e^{ty} \frac{\beta^\theta}{\Gamma(\theta)} e^{y\theta - \beta e^y} dy \\ &= \frac{\beta^\theta}{\Gamma(\theta)} \frac{\Gamma(\theta + t)}{\beta^{\theta+t}} \int \frac{\beta^{\theta+t}}{\Gamma(\theta + t)} e^{y(\theta+t) - \beta e^y} dy \\ &= \frac{\Gamma(\theta + t)}{\Gamma(\theta)\beta^t} \int \frac{\beta^{\theta+t}}{\Gamma(\theta + t)} e^{y(\theta+t) - \beta e^y} dy \\ &= \frac{\Gamma(\theta + t)}{\Gamma(\theta)\beta^t}. \end{aligned}$$

The assumption that $t > -\theta$ ensures that $\theta + t > 0$ so that the integrand in the second-to-last line is the density of the logarithm of a $\text{Gamma}(\theta + t, \beta)$ random variable. Since $\theta > 0$, we have established that the moment generating function exists in a neighborhood around 0. \square

Defining $Y^{(k)} = \log X^{(k)}$ and $\bar{Y}_K = \frac{1}{K} \sum_{k=1}^K Y^{(k)}$, observe that

$$\left(\prod_{k=1}^K X^{(k)} \right)^{\frac{1}{K}} = e^{\bar{Y}_K}.$$

Thus, our goal is to prove that $e^{\bar{Y}_K} \sim \text{Gamma}(\theta, \beta)$. Since the moment generating function completely characterizes a distribution, it is sufficient to show that $\Phi_{\bar{Y}_K}(t)$ matches the expression in Lemma 1. Applying Lemma 1 to $e^{Y^{(k)}} \sim \text{Gamma}(\theta_k, \beta_k)$, where $\theta_k = \theta/K + (k-1)/K$ and $\beta_k = \beta/K$, implies that

$$\Phi_{Y^{(k)}} = \frac{\Gamma(\theta_k + t)}{\Gamma(\theta_k)\beta_k^t}.$$

By independence of $Y^{(1)}, \dots, Y^{(K)}$ and standard properties of the moment generating function,

$$\begin{aligned} \Phi_{\bar{Y}_K}(t) &= \prod_{k=1}^K \Phi_{Y_{k/K}}(t) \\ &= \prod_{k=1}^K \Phi_{Y_k}(t/K) \\ &= \prod_{k=1}^K \frac{\Gamma(\theta_k + t/K)}{\Gamma(\theta_k)\beta_k^{t/K}}. \end{aligned}$$

Recalling the form of θ_k and β_k and applying the Gauss multiplication theorem (Abramowitz et al. 1972, page 256) to both the numerator and denominator gives

$$\Phi_{\bar{Y}_K}(t) = \frac{K^{-(\theta+t)}\Gamma(\theta+t)}{K^{-\theta}\Gamma(\theta)} \frac{1}{(\beta/K)^t} = \frac{\Gamma(\theta+t)}{\Gamma(\theta)\beta^t}.$$

This completes the proof.

2. A sufficient statistic for θ in the joint distribution of $X^{(1)}, \dots, X^{(K)}$ is $T(X^{(1)}, \dots, X^{(K)}) = \left(\prod_{k=1}^K X^{(k)} \right)^{\frac{1}{K}}$.

By the mutual independence of $X^{(k)}$, the joint density of $X^{(1)}, \dots, X^{(K)}$ can be written as,

$$\begin{aligned}
f_{X^{(1)}, \dots, X^{(K)}}(x^{(1)}, \dots, x^{(K)}) &= \prod_{k=1}^K f_{X^{(k)}}(x^{(k)}) \\
&\propto \prod_{k=1}^K (x^{(k)})^{\frac{1}{K}\theta + \frac{k-1}{K} - 1} \exp\left(-\frac{\beta}{K}x^{(k)}\right) \\
&= \left[\left(\prod_{k=1}^K x^{(k)} \right)^{\frac{1}{K}} \right]^\theta \left[\prod_{k=1}^K (x^{(k)})^{\frac{k-1}{K} - 1} \right] \exp\left(-\frac{\beta}{K} \sum_{k=1}^K x^{(k)}\right)
\end{aligned}$$

By the factorization theorem, $T(X^{(1)}, \dots, X^{(K)}) = \left(\prod_{k=1}^K X^{(k)} \right)^{\frac{1}{K}}$ is a sufficient statistic for θ as required.

3. To sample from G_t , the conditional distribution $(X^{(1)}, \dots, X^{(K)})|T(X^{(1)}, \dots, X^{(K)}) = t$, we first sample from $(X^{(1)}, \dots, X^{(K-1)})|T(X^{(1)}, \dots, X^{(K)}) = t$ and then recover $X^{(K)}$.

The conditional density $f_{X^{(1)}, \dots, X^{(K-1)}|T(X^{(1)}, \dots, X^{(K)})=t}(x^{(1)}, \dots, x^{(K-1)})$, up to a normalization constant depending on t , is

$$\left[\prod_{k=1}^{K-1} (x^{(k)})^{\frac{k-K}{K} - 1} \right] \exp\left(-\frac{\beta}{K} \left(\sum_{k=1}^{K-1} x^{(k)} + \frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}} \right)\right).$$

The derivation is as follows (where any factors not involving $x^{(1)}, \dots, x^{(K-1)}$ are omitted and we write $\theta_k = \frac{\theta}{K} + \frac{k-1}{K}$):

$$\begin{aligned}
&f_{X^{(1)}, \dots, X^{(K-1)}|T(X^{(1)}, \dots, X^{(K)})=t}(x^{(1)}, \dots, x^{(K-1)}) \\
&\propto f_{X^{(1)}, \dots, X^{(K-1)}, T(X^{(1)}, \dots, X^{(K)})}(x^{(1)}, \dots, x^{(K-1)}, t) \\
&= f_{X^{(1)}, \dots, X^{(K)}}\left(x^{(1)}, \dots, x^{(K-1)}, \frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}}\right) \left| \frac{\partial}{\partial t} \frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}} \right| \\
&= \left(\prod_{k=1}^{K-1} f_{X^{(k)}}(x^{(k)}) \right) f_{X^{(K)}}\left(\frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}}\right) \left| \frac{K t^{K-1}}{\prod_{k=1}^{K-1} x^{(k)}} \right| \\
&\propto \left(\prod_{k=1}^{K-1} (x^{(k)})^{\theta_k - 1} \exp\left(-\frac{\beta}{K}x^{(k)}\right) \right) \left(\frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}} \right)^{\theta_{K-1}} \exp\left(-\frac{\beta}{K} \frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}}\right) \frac{1}{\prod_{k=1}^{K-1} x^{(k)}} \\
&\propto \left(\prod_{k=1}^{K-1} (x^{(k)})^{\frac{k-K}{K} - 1} \right) \exp\left(-\frac{\beta}{K} \left(\sum_{k=1}^{K-1} x^{(k)} + \frac{t^K}{\prod_{k=1}^{K-1} x^{(k)}} \right)\right),
\end{aligned}$$

where in the last step we used that $\theta_k - \theta_K = \frac{k-K}{K}$.

After sampling from $(X^{(1)}, \dots, X^{(K-1)}) | T(X^{(1)}, \dots, X^{(K)}) = t$, compute $X^{(K)} = \frac{t^K}{\prod_{k=1}^{K-1} X^{(k)}}$.

□

B.4 Families with support controlled by an unknown parameter

B.4.1 Scaled beta distribution

We consider the family of distributions obtained by scaling a $\text{Beta}(\alpha, 1)$ distribution (with α fixed) by an unknown scale parameter $\theta > 0$. In the special case that $\alpha = 1$, this corresponds to the $\text{Unif}(0, \theta)$ family presented in Example 5.1.

Example B.1 (Thinning $\theta \cdot \text{Beta}(\alpha, 1)$ with α known). *We start with $X^{(k)} \stackrel{iid}{\sim} \theta \cdot \text{Beta}(\frac{\alpha}{K}, 1)$ for $k = 1, \dots, K$, and note that $T(X^{(1)}, \dots, X^{(K)}) = \max(X^{(1)}, \dots, X^{(K)})$ is sufficient for θ . Furthermore, $\max(X^{(1)}, \dots, X^{(K)}) \sim \theta \cdot \text{Beta}(\alpha, 1)$. Thus, we define G_t to be the conditional distribution of $(X^{(1)}, \dots, X^{(K)})$ given $\max(X^{(1)}, \dots, X^{(K)}) = t$. Then, by Theorem 1, we can thin $X \sim \theta \cdot \text{Beta}(\alpha, 1)$ by sampling from G_X .*

To sample from this conditional distribution, we first draw $C \sim \text{Categorical}_K(1/K, \dots, 1/K)$. Then, $X^{(k)} = C_k X + (1 - C_k) Z_k$ where $Z_k \stackrel{iid}{\sim} \text{Beta}(\frac{\alpha}{K}, 1)$.

Proof of Example B.1. We must prove the following three claims:

1. *If for $k = 1, \dots, K$, $X^{(k)} \stackrel{iid}{\sim} \theta \cdot \text{Beta}(\frac{\alpha}{K}, 1)$, then $\max(X^{(1)}, \dots, X^{(K)}) \sim \theta \cdot \text{Beta}(\alpha, 1)$.*

First, note that $\frac{1}{\theta} X^{(k)} \stackrel{iid}{\sim} \text{Beta}(\frac{\alpha}{K}, 1)$. The distribution of $\max(X^{(1)}, \dots, X^{(K)})$ can be derived using the CDF method as follows:

$$\begin{aligned}
 P(\max(X^{(1)}, \dots, X^{(K)}) \leq z) &= P(X^{(1)} \leq z, \dots, X^{(K)} \leq z) \\
 &= \prod_{k=1}^K P(X^{(k)} \leq z) \\
 &= \prod_{k=1}^K P\left(\frac{1}{\theta} X^{(k)} \leq \frac{z}{\theta}\right) \\
 &= \prod_{k=1}^K \left(\frac{z}{\theta}\right)^{\frac{\alpha}{K}} \\
 &= \left(\frac{z}{\theta}\right)^{\alpha},
 \end{aligned}$$

where we have used that $P(\text{Beta}(\alpha, 1) \leq x) = x^\alpha$ for $x \in (0, 1)$. The above implies that $\max(X^{(1)}, \dots, X^{(K)}) \sim \theta \cdot \text{Beta}(\alpha, 1)$ as required.

2. A sufficient statistic for θ based on $X^{(1)}, \dots, X^{(K)}$ is $\max(X^{(1)}, \dots, X^{(K)})$.

Using the independence of $X^{(k)}$, the joint distribution can be written as

$$\begin{aligned} & f_{X^{(1)}, \dots, X^{(K)}}(x^{(1)}, \dots, x^{(K)}) \\ &= \prod_{k=1}^K f_{X^{(k)}}(x^{(k)}) \\ &\propto \prod_{k=1}^K (x^{(k)})^{\frac{\alpha}{K}-1} I\{0 < x^{(k)} < \theta\} \\ &= \left(\prod_{k=1}^K x^{(k)} \right)^{\frac{\alpha}{K}-1} I\{\min(x^{(1)}, \dots, x^{(K)}) > 0\} I\{\max(x^{(1)}, \dots, x^{(K)}) < \theta\}. \end{aligned}$$

By the factorization theorem, we conclude that $\max(X^{(1)}, \dots, X^{(K)})$ is a sufficient statistic for θ as required.

3. We can sample from the conditional distribution $(X^{(1)}, \dots, X^{(K)}) | \max(X^{(1)}, \dots, X^{(K)}) = t$ by taking $X^{(k)} = C_k t + (1 - C_k) Z_k$ where $C \sim \text{Categorical}_K(1/K, \dots, 1/K)$ and $Z_k \sim \text{Beta}(\frac{\alpha}{K}, 1)$.

Without loss of generality, consider $X^{(k)}$. Given that $X^{(1)}, \dots, X^{(K)}$ are identically distributed, $P(X^{(k)} = t) = \frac{1}{K}$. Hence, in the first stage, we can draw one sample, $C \sim \text{Categorical}_K(1/K, \dots, 1/K)$ to determine if $X^{(k)}$ is the maximum. If $X^{(k)}$ is not the maximum then we know that $X^{(k)} \leq t$. We can compute the density of $Z_k \stackrel{D}{=} (X^{(k)} | X^{(k)} \leq t)$ as follows,

$$f_{X^{(k)} | X^{(k)} \leq t}(x^{(k)}) = \frac{f_{X^{(k)}}(x^{(k)})}{P(X^{(k)} \leq t)} = \frac{\frac{1}{\theta^{\frac{\alpha}{K}} B(\frac{\alpha}{K}, 1)} (x^{(k)})^{\frac{\alpha}{K}-1}}{\left(\frac{t}{\theta}\right)^{\frac{\alpha}{K}}} = \frac{1}{t^{\frac{\alpha}{K}} B(\frac{\alpha}{K}, 1)} (x^{(k)})^{\frac{\alpha}{K}-1}.$$

The above implies that $Z_k \sim t \cdot \text{Beta}(\frac{\alpha}{K}, 1)$ as required.

The result then follows from Theorem 1. □

B.4.2 Shifted exponential distribution

Proof of Example 5.2. We must prove the following three claims:

1. If for $k = 1, \dots, K$, $X^{(k)} \stackrel{iid}{\sim} \text{SExp}(\theta, \lambda/K)$, then $\min(X^{(1)}, \dots, X^{(K)}) \sim \text{SExp}(\theta, \lambda)$.

First, note that $X^{(k)} - \theta \stackrel{iid}{\sim} \text{Exp}(\lambda/K)$. The distribution of $\min(X^{(1)}, \dots, X^{(K)})$ can be derived using the CDF method as follows,

$$\begin{aligned}
P(\min(X^{(1)}, \dots, X^{(K)}) \geq z) &= P(X^{(1)} \geq z, \dots, X^{(K)} \geq z) \\
&= \prod_{k=1}^K P(X^{(k)} \geq z) \\
&= \prod_{k=1}^K P(X^{(k)} - \theta \geq z - \theta) \\
&= \prod_{k=1}^K \exp\left(-\frac{\lambda}{K}(z - \theta)\right) \\
&= \exp(-\lambda(z - \theta)).
\end{aligned}$$

The above implies that $\min(X^{(1)}, \dots, X^{(K)}) \sim \text{SExp}(\theta, \lambda)$ as required.

2. A sufficient statistic for θ based on $X^{(1)}, \dots, X^{(K)}$ is $\min(X^{(1)}, \dots, X^{(K)})$.

Using the independence of $X^{(k)}$, the joint distribution can be written as

$$\begin{aligned}
&f_{X^{(1)}, \dots, X^{(K)}}(x^{(1)}, \dots, x^{(K)}) \\
&= \prod_{k=1}^K f_{X^{(k)}}(x^{(k)}) \\
&\propto \prod_{k=1}^K \exp\left(-\frac{\lambda}{K}(x^{(k)} - \theta)\right) I\{x^{(k)} > \theta\} \\
&\propto \exp\left(-\frac{\lambda}{K} \sum_{k=1}^K x^{(k)}\right) I\{\min(x^{(1)}, \dots, x^{(K)}) > \theta\}.
\end{aligned}$$

Given that the joint distribution can be written such that θ only interacts with the data through the $I\{\min(x^{(1)}, \dots, x^{(K)}) > \theta\}$ term, we conclude that $\min(X^{(1)}, \dots, X^{(K)})$ is a sufficient statistic for θ as required.

3. We can sample from the conditional distribution $X^{(1)}, \dots, X^{(K)} | \min(X^{(1)}, \dots, X^{(K)}) = t$ by taking $X^{(k)} = t + (1 - C_k)Z_k$ where $C \sim \text{Categorical}_K(1/K, \dots, 1/K)$ and $Z_k \sim \text{Exp}(\lambda/K)$.

Without loss of generality, consider $X^{(k)}$. Given that $X^{(1)}, \dots, X^{(K)}$ are identically distributed, $P(X^{(k)} = X) = \frac{1}{K}$. Hence, in the first stage, we can draw one sample, $C \sim \text{Categorical}_K(1/K, \dots, 1/K)$ to determine if $X^{(k)}$ is the minimum. Otherwise, we require that $X^{(k)} \geq t$. We know that the density of $Z_k \stackrel{D}{=} (X^{(k)} | X^{(k)} \geq t) \stackrel{D}{=} X^{(k)}$ by the memoryless property of the exponential distribution. Thus, $Z_k \sim \text{Exp}(\lambda/K)$ as required.

The result then follows from Theorem 1.

□