

GORENSTEIN  $\mathcal{FP}_n$ -FLAT MODULES AND WEAK GLOBAL DIMENSIONS

VÍCTOR BECERRIL

ABSTRACT. In this paper we characterize the relative Gorenstein weak global dimension of the generalized Gorenstein  $\mathcal{Y}$ -flat modules recently studied by S. Estrada, A. Iacob, and M. A. Pérez. As application we prove that the weak global dimension that comes from the Gorenstein  $\mathcal{FP}_n$ -flat modules is finite over a Gorenstein  $n$ -coherent ring and coincide with the flat dimension of the right  $\mathcal{FP}_n$ -injective modules. This result extends the known for Gorenstein flat modules over Iwanaga-Gorenstein and Ding-Chen rings.

## 1. INTRODUCTION

Enochs, Jenda and Torrecillas [10, 11] introduced Gorenstein projective, injective and flat modules for any ring and since then a Gorenstein homological algebra has been developing intensively. Most recently, there has been an increasing interest for generalizations of this families of modules. Among them are the Ding projective and Ding injective modules presented by Mao, Ding and Gillespie [13, 15]. Also the named Gorenstein AC-projective and Gorenstein AC-injective modules are presented by Bravo, Gillespie and Hovey in [4]. Later are introduced the Gorenstein AC-flat modules by Bravo, Estrada and Iacob in [3]. More recently, this notions are encompassed by Iacob in [14], with the name of Gorenstein  $\mathcal{FP}_n$ -projective and Gorenstein  $\mathcal{FP}_n$ -injective. Under this impetus of generality in this paper we will characterize the Gorenstein weak global dimension relative to a pair  $(\mathcal{F}(R), \mathcal{Y}) \subseteq \text{Mod}(R) \times \text{Mod}(R^{\text{op}})$  (where  $\mathcal{F}(R)$  denotes the class of flat left  $R$ -modules), with application to the corresponding class of Gorenstein  $\mathcal{FP}_n$ -flat modules studied by [12, Example 2.21 (3)], but not limited to such family [12, Definition 2.1 (1)]. We know from Wang and Zhang [18, Lemma 1.2] that a ring  $R$  is Gorenstein (resp. Ding-Chen) if and only if  $R$  is Noetherian on both sides (resp. coherent) with finite Gorenstein weak global dimension. Also, without condition over the ring  $R$ , the Gorenstein weak global dimension has been characterized by Emmanouil [9] and is proved by Christensen, Estrada and Thompson that is symmetric [7]. In what follows we prove that over  $R$  a Gorenstein  $n$ -coherent ring [17, §4] the left Gorenstein weak global dimension relative to the pair  $(\mathcal{F}(R), \mathcal{FP}_k\text{-Inj}(R^{\text{op}}))$  is always finite and coincide with the flat dimension of  $\mathcal{FP}_k\text{-Inj}(R^{\text{op}})$  for all  $k \in \{0, 1, \dots, n\}$ . The above is a consequence of a general result that characterizes the relative Gorenstein weak global dimension. We will also see that for the Gorenstein  $\mathcal{FP}_n$ -flat modules the symmetry of the relative Gorenstein weak global dimension makes sense.

## 2. PRELIMINARIES

In what follows, we shall work with categories of modules over an associative ring  $R$  with identity. By  $\text{Mod}(R)$  and  $\text{Mod}(R^{\text{op}})$  we denote the categories of left and right  $R$ -modules. If  $\mathcal{X}$  is a class of left (or right)  $R$ -modules, then  $\text{Ch}(\mathcal{X})$  denotes the class of chain complexes of modules in  $\mathcal{X}$ , that is,  $X_\bullet \in \text{Ch}(\mathcal{X})$  if  $X_m \in \mathcal{X}$  for every  $m \in \mathbb{Z}$ .

---

2010 *Mathematics Subject Classification*. Primary 18G10, 18G20, 18G25. Secondary 16E10.  
Key Words: Gorenstein  $n$ -coherent,  $\mathcal{FP}_n$ -injective, Gorenstein weak global dimension.

Projective, injective and flat modules will be important to present some definitions, remarks and examples. The classes of projective left and right  $R$ -modules will be denoted by  $\mathcal{P}(R)$  and  $\mathcal{P}(R^{\text{op}})$ , respectively. Similarly, we shall use the notations  $\mathcal{I}(R)$ ,  $\mathcal{I}(R^{\text{op}})$ ,  $\mathcal{F}(R)$  and  $\mathcal{F}(R^{\text{op}})$  for the classes of injective and flat modules in  $\text{Mod}(R)$  and  $\text{Mod}(R^{\text{op}})$ , respectively.

Concerning functors defined on modules,  $\text{Ext}_R^i(-, -)$  denotes the right  $i$ -th derived functor of  $\text{Hom}_R(-, -)$ . If  $M \in \text{Mod}(R^{\text{op}})$  and  $N \in \text{Mod}(R)$ ,  $M \otimes_R N$  denotes the tensor product of  $M$  and  $N$ . Recall the construction of this tensor products defines a bifunctor  $-\otimes_R - : \text{Mod}(R^{\text{op}}) \times \text{Mod}(R) \rightarrow \text{Mod}(\mathbb{Z})$ , where  $\text{Mod}(\mathbb{Z})$  is the category of abelian groups.

Recall that a left  $R$ -module  $G$  is said to be *Gorenstein flat*, if it is isomorphic to  $\text{Ker}(F^0 \rightarrow F^1)$  of an exact complex of flat left  $R$ -modules

$$\cdots \rightarrow F_1 \rightarrow F_0 \rightarrow F^0 \rightarrow F^1 \rightarrow \cdots$$

which remains acyclic whenever that the functor  $I \otimes_R -$  is applied to it for any  $I \in \mathcal{I}(R^{\text{op}})$ . The class of all Gorenstein flat modules is denoted by  $\mathcal{GF}(R)$ . Given a class  $\mathcal{Y} \subseteq \text{Mod}(R^{\text{op}})$ , we say that  $G$  is Gorenstein  $\mathcal{Y}$ -flat, if there exists an exact complex  $F_\bullet \in \text{Ch}(\mathcal{F}(R))$  which is  $(Y \otimes_R -)$ -acyclic for all  $Y \in \mathcal{Y}$  and  $G \cong Z_0(F_\bullet)$  [12, Definition 2.1], we denote to the class of all Gorenstein  $\mathcal{Y}$ -flat modules by  $\mathcal{GF}_{(\mathcal{F}(R), \mathcal{Y})}$ , as in [1]. For any  $M \in \text{Mod}(R)$  we define the dimension Gorenstein  $\mathcal{Y}$ -flat dimension of  $M$ , denoted  $\text{Gfd}_{(\mathcal{F}(R), \mathcal{Y})}(M)$  as the minimum length over all the exact sequences  $0 \rightarrow G_m \rightarrow \cdots \rightarrow G_0 \rightarrow M \rightarrow 0$ , with  $G_i \in \mathcal{GF}_{(\mathcal{F}(R), \mathcal{Y})}$ .

### 3. GORENSTEIN WEAK GLOBAL DIMENSIONS

The fundamental idea in the following result is the one in the proof of [9, Theorem 5.3], we believe it is illustrative to make the proof of this fact. For a class  $\mathcal{X} \subseteq \text{Mod}(R)$ , the notation  $\text{fd}(\mathcal{X})$ , means *the flat dimension of the class  $\mathcal{X}$* , this is  $\sup\{\text{fd}(X) : X \in \mathcal{X}\}$ .

**Lemma 3.1.** *Consider  $\mathcal{Y} \subseteq \text{Mod}(R^{\text{op}})$  such that  $\text{fd}(\mathcal{Y}) < \infty$ . Then, all exact complex of flat  $R$ -modules is  $(\mathcal{Y} \otimes_R -)$ -exact.*

*Proof.* Assume that  $m := \text{fd}(\mathcal{Y}) < \infty$  and take  $F^\bullet : \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow F^0 \rightarrow F^1 \rightarrow \cdots$  an exact complex of flat  $R$ -modules. Since  $\text{fd}(\mathcal{Y}) = m$ , for  $A \in \mathcal{Y}$  there is an exact sequence  $0 \rightarrow P_m \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow A \rightarrow 0$  with  $P_i \in \mathcal{F}(R)$ . For each  $j \in \{0, 1, \dots, m-1\}$  consider the short exact sequences  $0 \rightarrow K_j \rightarrow P_j \rightarrow K_{j-1} \rightarrow 0$  where  $K_{m-1} := P_m$  and  $K_{-1} := A$ . For each  $j$  we have an exact sequence of complexes  $0 \rightarrow K_j \otimes_R F^\bullet \rightarrow P_j \otimes_R F^\bullet \rightarrow K_{j-1} \otimes_R F^\bullet \rightarrow 0$ . Since  $K_{m-1} = P_m$  we get that  $K_{m-2} \otimes_R F^\bullet$  is exact, so continuing iteratively we obtain that  $A \otimes_R F^\bullet$  is exact.  $\square$

We have the following result whose fundamental idea comes from [9, Lemma 5.2 and Theorem 5.3], but here we explain it in a different way. Recall the notion of *left Gorenstein weak global dimension* [1, §4] relative to the pair  $(\mathcal{F}(R), \mathcal{Y})$

$$\text{l.w.Ggl}_{(\mathcal{F}(R), \mathcal{Y})}(R) := \sup\{\text{Gfd}_{(\mathcal{F}(R), \mathcal{Y})}(M) \mid M \in \text{Mod}(R)\}.$$

**Proposition 3.2.** *Consider  $\mathcal{Y} \subseteq \text{Mod}(R^{\text{op}})$  such that  $\mathcal{GF}_{(\mathcal{F}(R), \mathcal{Y})}$  is closed by extensions and  $\text{fd}(\mathcal{Y}) < \infty$ . Then  $\text{fd}(\mathcal{I}(R)) \leq n$  implies that  $\text{l.w.Ggl}_{(\mathcal{F}(R), \mathcal{Y})}(R) \leq n$ .*

*Proof.* Take  $M \in \text{Mod}(R)$ , we can consider an exact sequence  $0 \rightarrow M \rightarrow I^0 \rightarrow I^1 \rightarrow \cdots$ , with  $I^j \in \mathcal{I}(R)$  for all  $j \geq 0$ . From the inequality  $\text{fd}(\mathcal{I}(R)) \leq n$  and [6, Ch. XVII, §1 Proposition 1.3]

we can construct the following commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & Q & \xrightarrow{C} & P_n^0 & \longrightarrow & P_n^1 & \longrightarrow & \cdots \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & Q_{n-1} & \longrightarrow & P_{n-1}^0 & \longrightarrow & P_{n-1}^1 & \longrightarrow & \cdots \\
& & \downarrow & & \downarrow & & \downarrow & & \\
& & \vdots & & \vdots & & \vdots & & \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & Q_0 & \longrightarrow & P_0^0 & \longrightarrow & P_0^1 & \longrightarrow & \cdots \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & M & \xrightarrow{C} & I^0 & \longrightarrow & I^1 & \longrightarrow & \cdots
\end{array}$$

such that  $P_i^j \in \mathcal{P}(R)$  for all  $i \in \{0, 1, \dots, n-1\}$  and all  $j \geq 0$ , and where  $P_n^j \in \mathcal{F}(R)$  for all  $j > 0$ . With  $Q_i := \text{Ker}(P_i^0 \rightarrow P_i^1) \in \mathcal{P}(R)$  for all  $i \in \{0, \dots, n-1\}$  and

$$\begin{aligned}
\lambda &: 0 \rightarrow Q \rightarrow P_n^0 \rightarrow P_n^1 \rightarrow \cdots \\
\theta &: 0 \rightarrow Q \rightarrow Q_{n-1} \rightarrow \cdots \rightarrow Q_0 \rightarrow M \rightarrow 0
\end{aligned}$$

exact complexes. We can complete  $\lambda$  to an exact complex of flat  $R$ -modules by consider an projective resolution of  $Q$ ,

$$\cdots \rightarrow P_1 \rightarrow P_0 \rightarrow Q \rightarrow 0.$$

Since  $\text{fd}(\mathcal{Y}) < \infty$  we know from Lemma 3.1 that the exact complex of flat  $R$ -modules

$$\cdots \rightarrow P_1 \rightarrow P_0 \rightarrow P_n^0 \rightarrow P_n^1 \rightarrow \cdots,$$

is  $(\mathcal{Y} \otimes_R -)$ -exact. Therefore  $Q \in \mathcal{GF}_{(\mathcal{F}, \mathcal{Y})}$ . Thus from the exact sequence  $\theta$  we conclude that  $\text{Gfd}_{(\mathcal{F}(R), \mathcal{Y})}(M) \leq n$ .  $\square$

From the results in [1], we have additionally the following characterization of the left Gorenstein weak global dimension relative to the pair  $(\mathcal{F}(R), \mathcal{Y})$ .

**Proposition 3.3.** *Consider  $\mathcal{Y} \subseteq \text{Mod}(R^{\text{op}})$  such that  $\mathcal{GF}_{(\mathcal{F}, \mathcal{Y})}$  is closed by extensions and  $\mathcal{I}(R^{\text{op}}) \subseteq \mathcal{Y}$ . Then  $\text{l.w.Ggl}_{(\mathcal{F}(R), \mathcal{Y})}(R) < \infty$  if and only if  $\text{fd}(\mathcal{Y}), \text{fd}(\mathcal{I}(R)) < \infty$ .*

*Proof.* From the previous result we only need to prove  $\Rightarrow$ . For do this, from [1, Proposition 4.1] we have that  $\text{fd}(\mathcal{Y}) < \infty$ . Since  $\mathcal{I}(R^{\text{op}}) \subseteq \mathcal{Y}$ , then the containment  $\mathcal{GF}_{(\mathcal{F}(R), \mathcal{Y})} \subseteq \mathcal{GF}_{(\mathcal{F}(R), \mathcal{I})}$  is given, thus for all  $M \in \text{Mod}(R)$  their dimension are such that  $\text{Gfd}_{(\mathcal{F}(R), \mathcal{I})}(M) \leq \text{Gfd}_{(\mathcal{F}(R), \mathcal{Y})}(M)$ , thus the finiteness of  $\text{l.w.Ggl}_{(\mathcal{F}(R), \mathcal{Y})}(R)$  implies the finiteness of the classical weak global Gorenstein flat dimension  $\text{l.w.Ggl}(R)$ . From [7, Theorem 2.4] we obtain in particular that  $\text{fd}(\mathcal{I}(R)) < \infty$ .  $\square$

Recall the notion of *global Gorenstein projective dimension* [2]

$$\text{gl.GPD}(R) := \sup\{\text{Gpd}(M) \mid M \in \text{Mod}(R)\}.$$

From the previous Proposition 3.3 we see that when the classical global Gorenstein projective dimension  $\text{gl.GPD}(R)$  and  $\text{fd}(\mathcal{Y})$  are finite, then the dimension  $\text{l.w.Ggl}_{(\mathcal{F}(R), \mathcal{Y})}(R)$  is also finite. From [2, Corollary 2.7] we obtain that  $\text{fd}(\mathcal{I}(R)) < \infty$ . Finally from Proposition 3.3, it follows

that  $\text{l.w.Ggl}_{(\mathcal{F}(R), \mathcal{Y})}(R)$  is finite.

### 3.1. Gorenstein $\mathcal{FP}_n$ -flat modules.

**Example 3.4.** *We will consider some interesting classes of modules.*

- (1) An  $R$ -module  $F$  is of **type  $\mathbf{FP}_\infty$**  [4] if it has a projective resolution  $\cdots \rightarrow P_1 \rightarrow P_0 \rightarrow F \rightarrow 0$ , where each  $P_i$  is finitely generated. An  $R$ -module  $A$  is **absolutely clean** if  $\text{Ext}_R^1(F, A) = 0$  for all  $R$ -modules  $F$  of type  $\mathbf{FP}_\infty$ . Also an  $R$ -module  $L$  is **level** if  $\text{Tor}_1^R(F, L) = 0$  for all (right)  $R$ -module  $F$  of type  $\mathbf{FP}_\infty$ . We denote by  $\text{AC}(R)$  (also denoted  $\mathcal{FP}_\infty\text{-Inj}(R)$  [5, Definition 3.1]) the class of all absolutely clean  $R$ -modules, and by  $\text{Lev}(R)$  (also denoted  $\mathcal{FP}_\infty\text{-Flat}(R)$  [5, Definition 3.2]) the class of all level  $R$ -modules. The class  $\mathcal{GF}_{(\mathcal{F}(R), \text{AC}(R^{\text{op}}))}$  is called **Gorenstein AC-flat** [3].
- (2)  $M \in \text{Mod}(R)$  is called **finitely  $n$ -presented** if  $M$  possesses a projective resolution  $0 \rightarrow P_n \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$  for which each  $P_i$  is finitely generated free. We denote by  $\mathcal{FP}_n(R)$  the class of all finite  $n$ -presented  $R$ -modules. An  $R$ -module  $E$  is called  **$\mathcal{FP}_n$ -injective** if for all  $M \in \mathcal{FP}_n(R)$ ,  $\text{Ext}_R^1(M, E) = 0$ . Also an  $R^{\text{op}}$ -module  $H$  is called  **$\mathcal{FP}_n$ -flat** if for all  $M \in \mathcal{FP}_n(R)$ ,  $\text{Tor}_1^R(H, M) = 0$ . These classes are denoted by  $\mathcal{FP}_n\text{-Inj}(R)$  and  $\mathcal{FP}_n\text{-Flat}(R^{\text{op}})$ , respectively.

The family of generalized Gorenstein modules that comes from the classes  $\mathcal{FP}_n\text{-Flat}(R)$  and  $\mathcal{FP}_n\text{-Inj}(R)$  was studied for the notions of Gorenstein projective and Gorenstein injective in [14]. From this reason we call Gorenstein  $\mathcal{FP}_n\text{-Inj}(R^{\text{op}})$ -flat modules to the modules in the class  $\mathcal{GF}_{(\mathcal{F}(R), \mathcal{FP}_n\text{-Inj}(R^{\text{op}}))}$ .

We can conclude from [12, Corollary 2.20], that the class  $\mathcal{GF}_{(\mathcal{F}, \mathcal{FP}_n\text{-Inj}(R^{\text{op}}))}$  is closed by extensions.

Can be easily seen that always there is a contention  $\mathcal{FP}_n(R) \supseteq \mathcal{FP}_{n-1}(R)$ , this is, there is an descending chain

$$\mathcal{FP}_0(R) \supseteq \mathcal{FP}_1(R) \supseteq \cdots \supseteq \mathcal{FP}_{n+1}(R) \supseteq \cdots \supseteq \mathcal{FP}_\infty(R),$$

can be seen also that  $R$  is left coherent, if and only if, the containment  $\mathcal{FP}_1(R) \subseteq \mathcal{FP}_2(R)$  is given [5, Proposition 2.1]. Also know that  $R$  is left Noetherian if and only if every module finitely generated is finitely presented, this is,  $\mathcal{FP}_0(R) \subseteq \mathcal{FP}_1(R)$ . From this we have the following notion.

**Definition 3.5.** [5, Definition 2.2] *A ring  $R$  is left  $n$ -coherent if every finitely  $n$ -presented left  $R$ -module is finitely  $(n+1)$ -presented, that is,  $\mathcal{FP}_n(R) \subseteq \mathcal{FP}_{n+1}(R)$ .*

For  $n \in \mathbb{Z}_{n \geq 0}$ , as in [17], a ring  $R$  is called *Goresntein  $n$ -coherent* if  $R$  is  $n$ -coherent and there exists  $m \in \mathbb{Z}_{\geq 0}$  such that  $\text{fd}(M) \leq m$  for all  $M \in \mathcal{FP}_n\text{-Inj}(R)$  (see also [16, §4]). This last condition implies also that  $\text{fd}(N) \leq m$  for all  $N \in \mathcal{FP}_n\text{-Inj}(R^{\text{op}})$  [16, Proposition 4.5]. In this situation and by [16, Proposition 4.5], since  $\mathcal{I}(R) \subseteq \mathcal{FP}_n\text{-Inj}(R)$ , then  $\text{fd}(\mathcal{I}(R)) < \infty$ . Therefore, from Proposition 3.3 we can conclude that the dimension  $\text{l.w.Ggl}_{(\mathcal{F}(R), \mathcal{FP}_n\text{-Inj}(R^{\text{op}}))}(R)$  is finite. We declare this as follows.

**Proposition 3.6.** *Let  $R$  be a Gorenstein  $n$ -coherent ring and  $\mathcal{FP}_n\text{-Inj}(R^{\text{op}})$  the class of right  $\mathcal{FP}_n$ -injective  $R$ -modules. Then the left Gorenstein weak global dimension relative to the pair  $(\mathcal{F}(R), \mathcal{FP}_n\text{-Inj}(R^{\text{op}}))$*

$$\text{l.w.Ggl}_{(\mathcal{F}(R), \mathcal{FP}_n\text{-Inj}(R^{\text{op}}))}(R)$$

*is finite and coincides with  $\text{fd}(\mathcal{FP}_n\text{-Inj}(R^{\text{op}}))$ .*

Note that in the sense of Definition 3.5 a Gorenstein ring is precisely a Gorenstein 0-coherent ring, and a Gorenstein 1-coherent ring is precisely a Ding-Chen ring [8]. Thus, the previous result recovers and extends the known in [18, Lemma 1.2].

**Example 3.7.** *There are rings  $R$  where  $\text{fd}(\mathcal{FP}_\infty\text{-Inj}(R^{\text{op}})) = \infty$  (see [16, Remark. 4.4]), such is the case when is considered  $\text{AC}(R^{\text{op}})$  over the ring defined by  $R := K \rtimes E$  the trivial extension of  $K$  by  $E$  where  $K$  is a field and  $E$  a  $K$ -vector space with infinite rank. On this ring the right regular module  $R_R$  admits infinite global weak dimension, but  $R_R \in \text{AC}(R^{\text{op}})$ , thus  $\text{fd}(\text{AC}(R^{\text{op}})) = \infty$ . This implies that the Gorenstein weak global dimension  $\text{l.w.Ggl}_{(\mathcal{F}(R), \text{AC}(R^{\text{op}}))}$  is also unbounded.*

We know that there exists an ascending chain of inclusions

$$\mathcal{FP}_0\text{-Inj}(R) \subseteq \mathcal{FP}_1\text{-Inj}(R) \subseteq \cdots \subseteq \mathcal{FP}_n\text{-Inj}(R) \subseteq \cdots \subseteq \mathcal{FP}_\infty\text{-Inj}(R),$$

of classes of modules [5, §3]. If  $R$  is a fix Gorenstein  $n$ -coherent ring (in particular if the dimension  $\text{fd}(\mathcal{FP}_n\text{-Inj}(R))$  is finite), then for all  $k \leq n$  we have that the dimension  $\text{fd}(\mathcal{FP}_k\text{-Inj}(R))$  is finite, since  $\text{fd}(\mathcal{FP}_k\text{-Inj}(R)) \leq \text{fd}(\mathcal{FP}_n\text{-Inj}(R))$  for all  $k \in \{0, 1, \dots, n\}$ . With this we have the following.

**Corollary 3.8.** *Let  $R$  be a Gorenstein  $n$ -coherent ring. Then for all  $k \in \{0, 1, \dots, n\}$  the left Gorenstein weak global dimension that comes from the class  $\mathcal{GF}_{(\mathcal{F}(R), \mathcal{FP}_k\text{-Inj}(R^{\text{op}}))}$  is finite and coincides with  $\text{fd}(\mathcal{FP}_k\text{-Inj}(R^{\text{op}}))$ .*

**3.2. Symmetry situations.** Recently has been proved by Christensen, Estrada and Thompson [7] that without conditions over the ring  $R$ , the classical Gorenstein weak global dimension is symmetric, this is, the quantities  $\text{l.w.Ggl}_{(\mathcal{F}(R), \mathcal{I}(R^{\text{op}}))}$  and  $\text{r.w.Ggl}_{(\mathcal{F}(R^{\text{op}}), \mathcal{I}(R))}$  always agree. Actually we have two classes of  $\mathcal{FP}_n$ -injective modules; one in  $\text{Mod}(R)$  and other in  $\text{Mod}(R^{\text{op}})$ , denoted  $\mathcal{FP}_n\text{-Inj}(R)$  and  $\mathcal{FP}_n\text{-Inj}(R^{\text{op}})$ , resp. From [16, Example 4.3 (4)] and applying [16, Proposition 4.5 (6)], we get that  $\text{fd}(\mathcal{FP}_n\text{-Inj}(R))$  and  $\text{fd}(\mathcal{FP}_n\text{-Inj}(R^{\text{op}}))$  are both finite when  $R$  is a Gorenstein  $n$ -coherent ring. The author is not sure if these quantities match. For now we only we can only assume it.

**Remark 3.9.** *Let  $R$  be a Gorenstein  $n$ -coherent ring, if the dimensions  $\text{fd}(\mathcal{FP}_n\text{-Inj}(R))$  and  $\text{fd}(\mathcal{FP}_n\text{-Inj}(R^{\text{op}}))$  match, then Global weak global dimension relative to  $(\mathcal{F}(R), \mathcal{FP}_n\text{-Inj}(R^{\text{op}}))$  is symmetric, this is, the following equality is given.*

$$\text{l.w.Ggl}_{(\mathcal{F}(R), \mathcal{FP}_n\text{-Inj}(R^{\text{op}}))}(R) = \text{r.w.Ggl}_{(\mathcal{F}(R^{\text{op}}), \mathcal{FP}_n\text{-Inj}(R))}(R).$$

Note that this ideas can be thought of more generally over  $(\mathcal{L}, \mathcal{A})$ -Gorenstein rings [16, §4], obtaining similar results to the ones declared here, but for the corresponding class of Gorenstein  $\mathcal{A}$ -flat modules  $\mathcal{GF}_{(\mathcal{F}(R), \mathcal{A})}$ .

**Acknowledgements** The author want to thank to professors Marco A. Pérez and Raymundo Bautista, for several helpful discussions on the results of this article.

**Funding** The author was fully supported by a CONACyT posdoctoral fellowship CVU 443002, managed by the Universidad Michoacana de San Nicolas de Hidalgo at the Centro de Ciencias Matemáticas, UNAM.

## REFERENCES

- [1] V. Becerril, *( $\mathcal{F}, \mathcal{A}$ )-Gorenstein flat homological dimensions*, Journal of the Korean Mathematical Society, 2022, Vol 59, no. 6, p. 1203-1227.
- [2] D. Bennis, N. Mahdou, *Global Gorenstein dimensions*, Proceedings of the American Mathematical Society, 2010, vol. 138, no 2, p. 461-465.
- [3] D. Bravo, S. Estrada, A. Iacob, *FPn-injective, FPn-flat covers and preenvelopes, and Gorenstein AC-flat covers*, Algebra Colloq. 25 (2) (2018) 319-334.
- [4] D. Bravo, J. Gillespie and M. Hovey, *The stable module category of a general ring*, Preprint, 2014, arXiv:1405.5768
- [5] D. Bravo, M. A. Pérez, *Finiteness conditions and cotorsion pairs*, Journal of Pure and Applied Algebra, 2017, vol. 221, no 6, p. 1249-1267.
- [6] H. Cartan, S. Eilenberg, *Homological Algebra*, Princeton Univ. Press, Princeton, N.J., 1956.
- [7] L. W. Christensen, S. Estrada, P. Thompson, *Gorenstein weak global dimension is symmetric*, Mathematische Nachrichten, 2021, vol. 294, no 11, p. 2121-2128.
- [8] N. Q. Ding, J. L. Chen, *Coherent rings with finite self-FP-injective dimension*, Communications in algebra, 1996, vol. 24, no 9, p. 2963-2980.
- [9] I. Emmanouil, *On the finiteness of Gorenstein homological dimensions*, Journal of Algebra, 2012, vol. 372, p. 376-396.
- [10] E. E. Enochs and O. M. G. Jenda, *Gorenstein injective and projective modules*, Math. Z. 220 (1995) 611-633
- [11] E. E. Enochs, O. M. G. Jenda and B. Torrecillas, *Gorenstein flat modules*, Nanjing Daxue Xuebao Shuxue Bannian Kan 10 (1993) 1-9.
- [12] S. Estrada, A. Iacob, and M. A. Pérez, *Model structures and relative Gorenstein flat modules and chain complexes*, in Categorical, homological and combinatorial methods in algebra, 135-175, Contemp. Math., 751, Amer. Math. Soc., RI, 2020.
- [13] J. Gillespie, *Model structures on modules over Ding-Chen rings*, Homol. Homotopy Appl. 12 (2010) 61-73.
- [14] A. Iacob, *Generalized Gorenstein modules*, Algebra Colloquium. World Scientific Publishing Company, 2022. p. 651-662.
- [15] L. X. Mao and N. Q. Ding, *Gorenstein FP-injective and Gorenstein flat modules*, J. Algebra Appl. 4 (2008) 497-506.
- [16] J. Wang, Z. Di, *Relative Gorenstein rings and duality pairs*, Journal of Algebra and Its Applications, 2020, vol. 19, no 08, pp. 2050147-1-21.
- [17] J. P. Wang, Z.K. Liu, X.Y. Yang, *A negative answer to a question of Gillespie*, Sci. Sinica Math. 48 (9) (2018) 1121-1130.
- [18] J. Wang, X. Zhang, *On rings with finite Gorenstein weak global dimension*, Journal of Algebra and Its Applications, 2022, p. 2350111.

(V. Becerril) CENTRO DE CIENCIAS MATEMÁTICAS. UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO. CP58089. MORELIA, MICHOACÁN, MÉXICO

Email address: victorbecerril@matmor.unam.mx