

QUASI-HOMOMORPHISMS OF QUANTUM CLUSTER ALGEBRAS

WEN CHANG, MIN HUANG, JIAN-RONG LI

ABSTRACT. In this paper, we study quasi-homomorphisms of quantum cluster algebras, which are quantum analogy of quasi-homomorphisms of cluster algebras introduced by Fraser. Quasi-homomorphisms of quantum cluster algebras appeared in a work by Kimura, Qin, and Wei in 2022 which they called “variation maps”.

For a quantum Grassmannian cluster algebra $\mathbb{C}_q[\text{Gr}(k, n)]$, we show that there is an associated braid group and each generator σ_i of the braid group preserves the quasi-commutative relations of quantum Plücker coordinates and exchange relations of the quantum Grassmannian cluster algebra. We conjecture that σ_i also preserves r -term ($r \geq 4$) quantum Plücker relations of $\mathbb{C}_q[\text{Gr}(k, n)]$. Up to this conjecture, we show that σ_i is a quasi-automorphism of $\mathbb{C}_q[\text{Gr}(k, n)]$ and the braid group acts on $\mathbb{C}_q[\text{Gr}(k, n)]$.

CONTENTS

1. Introduction	1
2. Quasi-homomorphisms of quantum cluster algebras	3
3. Braid group action on quantum Grassmannian cluster algebras	7
4. Proof of Theorem 3.15	20
References	20

1. INTRODUCTION

Cluster algebras were introduced by Fomin and Zelevinsky in [FZ02]. To study the symmetries of the cluster algebras, Assem, Schiffler, and Shramchenko introduced cluster automorphisms in [ASS12]. The concept was generalized to quasi-automorphisms of cluster algebras by Fraser in [F16]. More relations between cluster automorphisms and quasi-automorphisms were discovered in [CS19]. Both the automorphisms and quasi-automorphisms of cluster algebras are essential tools in the study of symmetries of cluster algebras.

It is natural to consider quantum version of “quasi-automorphisms of cluster algebras”. Quasi-homomorphisms of quantum cluster algebras appeared in [KQW22] by Kimura, Qin, and Wei in 2022 which they called “variation maps”. They also referred to “quantum quasi-automorphisms of a quantum cluster algebra” as “(quantum) twist automorphisms”.

The goal of this paper is to study quasi-homomorphisms of quantum cluster algebras and to study braid group action on the quantum Grassmannian cluster algebra $\mathbb{C}_q[\text{Gr}(k, n)]$.

For $k \leq n$, the Grassmannian $\text{Gr}(k, n)$ is the set of k -dimensional subspaces in an n -dimensional vector space. Denote by $\mathbb{C}[\text{Gr}(k, n)]$ its coordinate ring. Scott proved that there is a cluster algebra structure on $\mathbb{C}[\text{Gr}(k, n)]$ [Sco06]. The coordinate ring $\mathbb{C}[\text{Gr}(k, n)]$ is called a *Grassmannian cluster algebra*. Fraser [F17] has studied braid group action on $\mathbb{C}[\text{Gr}(k, n)]$. He introduced regular automorphisms $\sigma_i, 1 \leq i \leq d - 1$, on $\text{Gr}(k, n)$, which induce quasi-automorphisms σ_i^* on the coordinate ring $\mathbb{C}[\text{Gr}(k, n)]$, where $d = \gcd(k, n)$, and he proved that these automorphisms give a braid group action on $\mathbb{C}[\text{Gr}(k, n)]$.

In this paper we take $\mathbb{C}_q = \mathbb{C}(q^{\pm 1/2})$. The quantized coordinate ring $\mathbb{C}_q[\text{Gr}(k, n)]$ is the subalgebra of the quantum matrix algebra $\mathbb{C}_q[M(k, n)]$ generated by the quantum Plücker coordinates, see [TT91, GL14, Lau06, Lau10] and Section 3.1. Grabowski and Launois

[GL11, GL14] proved that the quantum deformation $\mathbb{C}_q[\text{Gr}(k, n)]$ of $\mathbb{C}[\text{Gr}(k, n)]$ has a quantum algebra structure.

Jensen, King and Su showed (see the paragraphs before Theorem 1.3 in [JKS22]) that there is a quantum cluster algebra containing cluster variables X_J for all k -subsets J of $[n]$, whose quasi-commutation rules are the same as the quasi-commutation rules of the quantum minors Δ_q^J of $\mathbb{C}_q[\text{Gr}(k, n)]$. They showed that the quantum cluster algebra is isomorphic to the quantum coordinate ring $\mathbb{C}_q[\text{Gr}(k, n)]$ by identifying X_J with Δ_q^J for all J , see Theorem 1.3 in [JKS22] (note that we take $\mathbb{C}_q = \mathbb{C}(q^{\pm 1/2})$).

There is a bijection φ between cluster variables in $\mathbb{C}[\text{Gr}(k, n)]$ and cluster variables in $\mathbb{C}_q[\text{Gr}(k, n)]$, see Section 3.3. It is shown in [F17] that in $\mathbb{C}[\text{Gr}(k, n)]$, for every cluster variable x , $\sigma_i(x) = y_1 y_2 \cdots y_r \tilde{x}$, where r is a positive integer, y_i is a frozen variable or the inverse of a frozen variable, and \tilde{x} is a cluster variable. In $\mathbb{C}_q[\text{Gr}(k, n)]$, we define $\sigma_i(\varphi(x))$ to be $\varphi(y_1)\varphi(y_2)\cdots\varphi(y_r)\varphi(\tilde{x})$ as a commutative quantum cluster monomial (see Sections 3.2 and 3.3).

We prove that every map σ_i extends to a map (also denoted by σ_i) that preserves the quasi-commutative relations of quantum Plücker coordinates and exchange relations of the quantum Grassmannian cluster algebra $\mathbb{C}_q[\text{Gr}(k, n)]$. We conjecture that σ_i also preserves r -term ($r \geq 4$) quantum Plücker relations of $\mathbb{C}_q[\text{Gr}(k, n)]$. Up to this conjecture, we show that σ_i is a quasi-automorphism of $\mathbb{C}_q[\text{Gr}(k, n)]$ and the braid group acts on $\mathbb{C}_q[\text{Gr}(k, n)]$, see Theorem 3.15. This braid group action gives a way to discover many quantum cluster variables and quantum clusters of the quantum Grassmannian cluster algebra.

Symmetries of quantum Grassmannians have been studied in [LL11], [AG12]. Launois and Lenagan proved that a certain cycling map ρ defines a \mathbb{C}_q -algebra isomorphism from $\mathbb{C}_q[\text{Gr}(k, n)]$ to its twisted algebra $T(\mathbb{C}_q[\text{Gr}(k, n)])$ and they showed that the cycling map ρ is not a \mathbb{C}_q -algebra automorphism of $\mathbb{C}_q[\text{Gr}(k, n)]$. For example, in $\mathbb{C}_q[\text{Gr}(2, 4)]$, the map σ_1 is similar to but different from the cycling map ρ . Unlike ρ , the map σ_1 is a \mathbb{C}_q -algebra automorphism on $\mathbb{C}_q[\text{Gr}(2, 4)]$ (see Section 3.4). The braid group action gives a large family of \mathbb{C}_q -algebra automorphisms on $\mathbb{C}_q[\text{Gr}(k, n)]$ (quasi-automorphisms on the quantum cluster algebra $\mathbb{C}_q[\text{Gr}(k, n)]$ are \mathbb{C}_q -algebra automorphisms, see Lemma 3.14).

Marsh and Scott [MS16] defined a twist automorphism on $\mathbb{C}[\text{Gr}(k, n)]$ which sends cluster variables to cluster variables up to multiplying frozen variables. This twist automorphism is closely related to Berenstein–Fomin–Zelevinsky twist automorphisms on unipotent cells [BFZ96]. We expect that there is a quantum version of Marsh–Scott’s twist map on $\mathbb{C}_q[\text{Gr}(k, n)]$ which should be closely related to the quantum twist map studied in [KO19].

The paper is organized as follows. In Section 2, we introduce the concept of quasi-homomorphisms of quantum cluster algebras. In Section 3, we study braid group actions on quantum Grassmannian cluster algebras. In Section 4, we prove our main result: Theorem 3.15.

Acknowledgements. We would like to thank Chris Fraser, Jan E. Grabowski, and Bach Nguyen for very helpful discussions. We would like to thank Fan Qin for telling us that “quasi-homomorphisms of quantum cluster algebras” appeared in [KQW22] by Kimura, Qin, and Wei in 2022 which they called “variation maps” after our paper was uploaded to Arxiv.

The first author is supported by Shaanxi Normal University and NSF of China: 12271321, and the project of Youth Innovation Team of Universities of Shandong Province (No. 2022KJ314). The second author is supported by NSF of China: 12101617 and Guangdong Basic and Applied Basic Research Foundation 2021A1515012035. The third author is supported by the Austrian Science Fund (FWF): P-34602.

Notation. Throughout the paper, we use \mathbb{Z} as the set of integers and use the notation $[b]_+ = \max(b, 0)$ for $b \in \mathbb{Z}$. We use $\text{Mat}_{m \times n}(\mathbb{Z})$ to denote the set of $m \times n$ matrices over \mathbb{Z} .

For a matrix $M = (b_{ij}) \in \text{Mat}_{m \times n}(\mathbb{Z})$, we also use the notation $[M]_+ = ([b_{ij}]_+)$. For any positive integer m , we denote by $[1, m]$ the (ordered) set of positive integers smaller than $m + 1$. We will use $|I|$ to denote the cardinality of a set I .

2. QUASI-HOMOMORPHISMS OF QUANTUM CLUSTER ALGEBRAS

We will introduce the notion of quasi-homomorphisms of quantum cluster algebras in this section. We start with briefly recalling some background which will be used in the following, see details in [BZ05].

2.1. Background on quantum cluster algebras.

2.1.1. *Quantum seeds.* Let m and n be two positive integers. A quantum seed is a triple $\Sigma = (\mathbf{x}, \tilde{B}, \Lambda)$, where

- $\mathbf{x} = (x_1, \dots, x_m)$ is an ordered set with m elements;
- $\tilde{B} = \begin{pmatrix} B \\ C \end{pmatrix} \in \text{Mat}_{m \times n}(\mathbb{Z})$ is an extended skew-symmetrizable matrix, that is, there exists a diagonal matrix D with positive integer entries such that DB is skew-symmetric, where B is the $n \times n$ submatrix of \tilde{B} consisting of the first n rows;
- $\Lambda \in \text{Mat}_{m \times m}(\mathbb{Z})$ is a skew-symmetric matrix such that (\tilde{B}, Λ) is a compatible pair, that is $\tilde{B}^t \Lambda = (D, 0_{n \times (m-n)})$.

The set \mathbf{x} is the *cluster* of the quantum seed Σ . The elements in $\mathbf{p} = \{x_{n+1}, \dots, x_m\} \subset \mathbf{x}$ are called the *coefficient variables* and the elements in $\mathbf{x} \setminus \mathbf{p}$ are called the *cluster variables* of Σ . The $n \times n$ matrix B is called the *exchange matrix* of Σ ; its rows are called *exchangeable rows*, and the remaining rows of \tilde{B} are called *frozen rows*. We always assume that both \tilde{B} and B are indecomposable matrices, and we also assume that $n > 1$ for convenience.

2.1.2. *Quantum torus.* Denote $\mathbb{C}_q = \mathbb{C}(q^{\pm \frac{1}{2}})$, where q is an indeterminate. A quantum torus \mathbb{T} of rank $m \in \mathbb{Z}_{>0}$ is a \mathbb{C}_q -algebra which has basis $\{X^a \mid a \in \mathbb{Z}^m\}$ with multiplication:

$$(2.1) \quad X^a X^b = q^{\frac{1}{2}\Lambda(a,b)} X^{a+b},$$

where $\Lambda(-, -)$ is the bilinear form given by a skew-symmetric matrix Λ .

Let $\{e_1, \dots, e_m\}$ be the standard basis of \mathbb{Z}^m , then we have

$$(2.2) \quad X^{e_i} X^{e_j} = q^{\frac{1}{2}\Lambda(e_i, e_j)} X^{e_i + e_j}, \quad X^{e_i} X^{e_j} = q^{\Lambda(e_i, e_j)} X^{e_j} X^{e_i}.$$

We associate with \mathbb{T} the \mathbb{C} -linear *bar-involution* by setting

$$\overline{q^{\frac{1}{2}} X^a} = q^{-\frac{1}{2}} X^a$$

on each basis element. An element X in \mathbb{T} is a *bar-invariant* if it is maintained by the bar-involution, that is, $\overline{X} = X$.

Given a quantum seed $\Sigma = (\mathbf{x}, \tilde{B}, \Lambda)$ with notations as above, we associate to it a quantum torus $\mathbb{T}(\Sigma)$, where $X^{e_i} = x_i$ and $\Lambda(-, -)$ is the bilinear form given by Λ . We denote by \mathbb{P} the monoid (the multiplication is inherited from $\mathbb{T}(\Sigma)$) consisting of elements $X^{(0,a)}$ with $0 \in \mathbb{Z}^n$ and $a \in \mathbb{Z}^{m-n}$.

2.1.3. *Quantum mutations.* Denote by b_{ij} the entry in \tilde{B} located at position (i, j) for $1 \leq i \leq m$, $1 \leq j \leq n$ and by b_k the k -th column of \tilde{B} for $1 \leq k \leq n$. Similarly we use notations λ_{ij} and λ_k for Λ with $1 \leq i, j, k \leq m$. The quantum seed $\mu_k(\Sigma) = (\mu_k(\mathbf{x}), \mu_k(\tilde{B}), \mu_k(\Lambda))$ obtained by the *mutation* of Σ in direction k is given by:

- $\mu_k(\mathbf{x}) = (\mathbf{x} \setminus \{x_k\}) \sqcup \{\mu_k(x_k)\}$ where $\mu_k(x_k)$ is an element in $\mathbb{T}(\Sigma)$ defined by

$$(2.3) \quad \mu_k(x_k) = X^{-e_k + [b_k]_+} + X^{-e_k + [-b_k]_+}.$$

• $\mu_k(\tilde{B}) = (b'_{ji})_{m \times n} \in M_{m \times n}(\mathbb{Z})$ is given by

$$(2.4) \quad b'_{ji} = \begin{cases} -b_{ji} & \text{if } i = k \text{ or } j = k ; \\ b_{ji} + [-b_{jk}]_+ b_{ki} + b_{jk} [b_{ki}]_+ & \text{otherwise.} \end{cases}$$

• $\mu_k(\Lambda) = (\lambda'_{ji})_{m \times m} \in M_{m \times m}(\mathbb{Z})$ is given by

$$(2.5) \quad \lambda'_{ji} = \begin{cases} \lambda_{jk} + \lambda_j^t [b_k]_+ & \text{if } i = k ; \\ \lambda_{ki} - \lambda_i^t [b_k]_+ & \text{if } j = k ; \\ \lambda_{ji} & \text{otherwise.} \end{cases}$$

It can be check that $(\mu_k(\tilde{B}), \mu_k(\Lambda))$ is still a compatible pair and $\mu(\Sigma)$ is a quantum seed.

2.1.4. *Quantum cluster algebra.* Let \mathcal{X} be the set of cluster variables in the quantum seeds obtained by iterated mutations from a quantum seed Σ . Then the algebra

$$\mathcal{A}_q = \mathcal{A}_q(\Sigma) := \mathbb{C}_q \mathbb{P}[\mathcal{X}]$$

algebraically generated by elements in \mathcal{X} over $\mathbb{C}_q \mathbb{P}$ is called the *quantum cluster algebra* of rank n defined by Σ , which is a $\mathbb{C}_q \mathbb{P}$ -subalgebra of the quantum torus $\mathbb{T}(\Sigma)$. The elements in \mathcal{X} are called *cluster variables*, and the elements in $\mathbb{C}_q \mathbb{P}$ are called *coefficients* of \mathcal{A}_q .

2.2. **Quasi-homomorphisms of quantum cluster algebras.** Let \mathcal{A}_q and \mathcal{A}'_q be two quantum cluster algebras of rank n . Assume that $\Sigma = \{\mathbf{x} = \{x_1, \dots, x_m\}, \tilde{B} = \begin{pmatrix} B \\ C \end{pmatrix}, \Lambda\}$ and $\Sigma' = \{\mathbf{x}' = \{x'_1, \dots, x'_{m'}\}, \tilde{B}' = \begin{pmatrix} B' \\ C' \end{pmatrix}, \Lambda'\}$ be the initial seeds of them respectively. Correspondingly, we have quantum torus \mathbb{T} and \mathbb{T}' .

Definition 2.1. We call a \mathbb{C}_q -algebra homomorphism $f : \mathcal{A}_q \rightarrow \mathcal{A}'_q$ a quantum quasi-homomorphism (with respect to Σ and Σ') if the following conditions are satisfied:

- (1) $B = B'$;
- (2) $f(x_i) = X^{e_i + \tilde{h}_i}$ for all $1 \leq i \leq n$, and $f(x_j) = X^{\tilde{l}_j}$ for all $n+1 \leq i \leq m$, where $\tilde{h}_i = \begin{pmatrix} 0 \\ h_i \end{pmatrix}$ and $\tilde{l}_j = \begin{pmatrix} 0 \\ l_j \end{pmatrix}$ with $0 \in \mathbb{Z}^n$, $h_i \in \mathbb{Z}^{m'-n}$ and $l_j \in \mathbb{Z}^{m'-n}$ such that

$$f(\hat{y}_i) = \hat{y}'_i$$

for all $1 \leq i \leq n$, where $\hat{y}_i = X^{b_i}$ and $\hat{y}'_i = X^{b'_i}$.

Remark 2.2. In Definition 3.1.3 of [KQW22], the variation map between Laurent polynomial rings preserving the Poisson structure was defined. It commutes with mutation sequences by Proposition 3.3.2 in [KQW22]. It restricts to a map between upper cluster algebras by Lemma 3.3.4 in [KQW22]. Since the Poisson structure is preserved, the map has a natural quantization (see the first paragraph in Section 8.2 in [KQW22]). The quantization coincides with the quantum quasi-homomorphism in Definition 2.1. In [KQW22], “quantum quasi-automorphisms of a quantum cluster algebra” are called “(quantum) twist automorphisms”. We would like to thank Fan Qin for telling us these after our paper was uploaded to Arxiv.

Let

$$L = (l_1, \dots, l_{m-n}), \quad H = (h_1, \dots, h_n), \quad R = \begin{pmatrix} I_{n \times n} & 0 \\ H & L \end{pmatrix} \in \text{Mat}_{m' \times m}(\mathbb{Z}).$$

Denote by r_i the i -th column of R . Then for any $1 \leq i \leq m$,

$$(2.6) \quad f(X^{e_i}) = X^{r_i}.$$

Lemma 2.3. For a quantum quasi-homomorphism $f : \mathcal{A}_q \rightarrow \mathcal{A}'_q$, we have that

$$R^t \Lambda' R = \Lambda.$$

Proof. For any $i, j \in [m]$, we have $X^{e_i}X^{e_j} = q^{\Lambda(e_i, e_j)}X^{e_j}X^{e_i}$. As f is a \mathbb{C}_q -algebra homomorphism, we obtain

$$f(X^{e_i})f(X^{e_j}) = q^{\Lambda(e_i, e_j)}f(X^{e_j})f(X^{e_i}).$$

Then by equalities (2.2) and (2.6), we have $\frac{1}{2}\Lambda'(r_i, r_j) = \Lambda(e_i, e_j) + \frac{1}{2}\Lambda'(r_j, r_i)$. Consequently, $r_i^t \Lambda' r_j = \Lambda'(r_i, r_j) = \Lambda(e_i, e_j) = \lambda_{ij}$ for $i, j \in [1, m]$. This implies the required equality. \square

For any $i \in \{1, \dots, n\}$, denote by $R' \in \text{Mat}_{m' \times m}(\mathbb{Z})$ the matrix with the k -th column r'_k , $k = 1, \dots, m$, where

$$r'_k = \begin{cases} e'_i + R[b_i]_+ - [b'_i]_+ & \text{if } k = i; \\ r_k & \text{if } k \neq i. \end{cases}$$

Note that if we set q to be one, then the definition degenerates back to the non-quantum case introduced by [F16]. In particular, the following lemma follows from the commutative case.

Lemma 2.4. *For a quantum quasi-homomorphism $f : \mathcal{A}_q \rightarrow \mathcal{A}'_q$, we have that*

- (1) $R\widetilde{B} = \widetilde{B}'$,
- (2) $R'\mu_i\widetilde{B} = \mu_i\widetilde{B}'$.

For any quasi-homomorphism $f : \mathcal{A}_q(\Sigma) \rightarrow \mathcal{A}_q(\Sigma')$, it is clear that f induces a ring homomorphism $f : \mathbb{T}(\Sigma) \rightarrow \mathbb{T}(\Sigma')$.

Lemma 2.5. *Let $f : \mathcal{A}_q \rightarrow \mathcal{A}'_q$ be a quasi-homomorphism. Then for any $a = (a_1, \dots, a_m)^t \in \mathbb{Z}^m$, we have that*

$$f(X^a) = X'^{Ra}.$$

In particular, $f(X^a)$ is bar-invariant.

Proof. By direct calculation, we have

$$\begin{aligned} f(X^a) &= f(q^{-\frac{1}{2}\sum_{i<j} a_i a_j \lambda_{ij}} x_1^{a_1} \dots x_m^{a_m}) = q^{-\frac{1}{2}\sum_{i<j} a_i a_j \lambda_{ij}} X'^{a_1 r_1} \dots X'^{a_m r_m} \\ &= q^{-\frac{1}{2}\sum_{i<j} a_i a_j \lambda_{ij}} q^{\frac{1}{2}\sum_{i<j} a_i a_j \Lambda'(r_i, r_j)} X'^{a_1 r_1 + \dots + a_m r_m} = X'^{a_1 r_1 + \dots + a_m r_m} \\ &= X'^{Ra}, \end{aligned}$$

where the last but one equality follows by Lemma 2.3. \square

Proposition 2.6. *Let $f : \mathcal{A}_q \rightarrow \mathcal{A}'_q$ be a quasi-homomorphism. Then f maps a bar-invariant element in \mathcal{A}_q to a bar-invariant element in \mathcal{A}'_q , that is, $\overline{f(X)} = f(\overline{X})$ if $\overline{X} = X \in \mathcal{A}_q$.*

Proof. It follows from the second condition in Definition 2.1 that $f(x_i), i \in \{1, \dots, m\}$ are bar-invariant. For any $X \in \mathcal{A}_q$ such that $\overline{X} = X$, by the quantum Laurent phenomenon, we have $X = \sum_a \lambda(a)X^a$ for some $\lambda(a) \in \mathbb{C}_q$. Then we have $f(X) = \sum_a \lambda(a)f(X^a)$ and $\overline{f(X)} = \sum_a \overline{\lambda(a)}\overline{f(X^a)}$. As $\overline{X} = X$, we have $\overline{\lambda(a)} = \lambda(a)$ for each a . Then by Lemma 2.5, we see that $\overline{f(X)} = f(\overline{X}) = f(X)$, that is, $f(X)$ is bar-invariant. \square

The following proposition shows that the definition of quasi-homomorphism of quantum cluster algebras is compatible with the mutations.

Proposition 2.7. *Let $f : \mathcal{A}_q \rightarrow \mathcal{A}'_q$ be a quantum quasi-homomorphism with respect to seeds Σ and Σ' . Then for any sequence (i_1, i_2, \dots, i_s) , $i_j \in \{1, \dots, n\}$, we have that $f : \mathcal{A}_q \rightarrow \mathcal{A}'_q$ is a quantum quasi-homomorphism with respect to the seeds $\mu_{i_s} \dots \mu_{i_1} \Sigma$ and $\mu_{i_s} \dots \mu_{i_1} \Sigma'$.*

Proof. By induction, it suffices to prove the case that $s = 1$. Denote $i = i_1$. Then by Lemma 2.5 we have

$$\begin{aligned} f(\mu_i x_i) &= f(X^{-e_i + [b_i]_+} + X^{-e_i + [-b_i]_+}) = X'^{R(-e_i + [b_i]_+)} + X'^{R(-e_i + [-b_i]_+)} \\ &= X'^{-e'_i + R[b_i]_+} + X'^{-e'_i + R[-b_i]_+}. \end{aligned}$$

By Lemma 2.4 (1), we have $[b'_i]_+ - [-b'_i]_+ = b'_i = Rb_i = R[b_i]_+ - R[-b_i]_+$. As $\mu_i(x'_i) = X'^{-e'_i + [b'_i]_+} + X'^{-e'_i + [-b'_i]_+}$, it follows that

$$f(\mu_i x_i) = (\mu_i(X'))^{e'_i + R[b_i]_+ - [b'_i]_+}.$$

As $R = \begin{pmatrix} I_{n \times n} & 0 \\ H & L \end{pmatrix}$ and $B = B'$, we see that the first n coordinates of $R[b_i]_+ - [b'_i]_+$ are 0.

For any $i \neq j \in \{1, \dots, n\}$, as the first n coordinates of \tilde{h}_j are 0, we have

$$f(x_j) = X'^{e'_j + \tilde{h}_j} = (\mu_i(X'))^{e'_j + \tilde{h}_j}.$$

For any $j \in \{n+1, \dots, m\}$, as the first n coordinates of \tilde{l}_j are 0, we have

$$f(x_j) = X'^{\tilde{l}_j} = (\mu_i(X'))^{\tilde{l}_j}.$$

Denote by d_j and d'_j the j -th column of the matrix $\mu_i \tilde{B}$ and $\mu_i \tilde{B}'$, respectively. Denote $\mu_i(\hat{y}_j) = (\mu_i(X))^{d_j}$ and $\mu_i(\hat{y}'_j) = (\mu_i(X'))^{d'_j}$ for any $j \in \{1, \dots, n\}$.

Next, we show that $f(\mu_i(\hat{y}_j)) = \mu_i(\hat{y}'_j)$. If $j = i$, we have $\mu_i(\hat{y}_i) = \hat{y}_i^{-1}$ and $\mu_i(\hat{y}'_i) = \hat{y}'_i^{-1}$, and thus

$$f(\mu_i(\hat{y}_i)) = f(\hat{y}_i^{-1}) = \hat{y}'_i^{-1} = \mu_i(\hat{y}'_i).$$

If $j \neq i$, we have $\mu_i(\hat{y}_j) = (\mu_i(X))^{\sum_k d_{kj}}$. Then by Proposition 2.6 and Lemma 2.4 (2), we have

$$\begin{aligned} f(\mu_i(\hat{y}_j)) &= (\mu_i(X'))^{d_{ij}(e'_i + R[b_i]_+ - [b'_i]_+) + \sum_{1 \leq k(\neq i) \leq n} d_{kj}(e'_k + \tilde{h}_k) + \sum_{n+1 \leq k \leq m} d_{kj} \tilde{l}_j} \\ &= (\mu_i(X'))^{d'_j} = \mu_i(\hat{y}'_j). \end{aligned}$$

The proof is complete. \square

So roughly speaking, a quasi-homomorphism of quantum cluster algebras is an algebra homomorphism between quantum cluster algebras of the same type which sends a cluster variable to a cluster variable possibly multiplying a Laurent monomial in frozen variables, sends a frozen variable to a Laurent monomial in frozen variables, in a way that keeps the hatted y -variables and commutes with the mutations. On the other hand, from a view point of linear algebra, the following proposition says that a quasi-homomorphism between quantum cluster algebras is equivalent to two matrix equations.

Proposition 2.8. *Let \mathbf{x} and \mathbf{x}' respectively be clusters of \mathcal{A}_q and \mathcal{A}'_q with the same exchange matrix. A map $f : \mathbf{x} \mapsto \mathbf{x}'$ induces a quasi-homomorphism from \mathcal{A}_q to \mathcal{A}'_q if and only if the following conditions are satisfied,*

- (1) $f(x_i) = X'^{e'_i + \tilde{h}_i}$, $1 \leq i \leq n$, where $\tilde{h}_i = \begin{pmatrix} 0 \\ h_i \end{pmatrix}$ with $h_i \in \mathbb{Z}^{m'-n}$;
- (2) $f(x_{n+j}) = X'^{\tilde{l}_j}$, $1 \leq j \leq m-n$, where $\tilde{l}_j = \begin{pmatrix} 0 \\ l_j \end{pmatrix}$ with $l_j \in \mathbb{Z}^{m'-n}$;
- (3) $R\tilde{B} = \tilde{B}'$, where $L = (l_1, \dots, l_{m-n})$, $H = (h_1, \dots, h_n)$ and $R = \begin{pmatrix} I_{n \times n} & 0 \\ H & L \end{pmatrix}$;
- (4) $R^t \Lambda' R = \Lambda$.

Proof. “Only If Part”: (1) and (2) follow by condition (2) in Definition 2.1. (3) follows by Lemma 2.4 (1), and (4) follows by Proposition 2.3.

“If Part”: from the proof of Proposition 2.3, we know that (4) is equivalent to $f(x_i)f(x_j) = q^{\Lambda(e_i, e_j)} f(x_j)f(x_i)$ for any $i, j \in [1, m]$. Therefore, f can be extended to a well-defined algebra homomorphism from \mathbb{T} to \mathbb{T}' .

$$\begin{aligned} f(\hat{y}_k) &= f(q^{-\frac{1}{2} \sum_{i < j} b_{ik} b_{jk} \lambda_{ij}} x_1^{b_{1k}} \dots x_m^{b_{mk}}) = q^{-\frac{1}{2} \sum_{i < j} b_{ik} b_{jk} \lambda_{ij}} X'^{b_{1k} r_1} \dots X'^{b_{mk} r_m} \\ &= q^{-\frac{1}{2} \sum_{i < j} b_{ik} b_{jk} \lambda_{ij}} q^{\frac{1}{2} \sum_{i < j} b_{ik} b_{jk} \Lambda'(r_i, r_j)} X'^{b_{1k} r_1 + \dots + b_{mk} r_m} = X'^{b_{1k} r_1 + \dots + b_{mk} r_m} \\ &= X'^{Rb_k} = X'^{b'_k} = \hat{y}'_k, \end{aligned}$$

where the last but one equality follows from (3).

Furthermore, by a similar argument used in Proposition 2.7, f maps any cluster variable of \mathcal{A}_q to a cluster variable possibly multiplying a coefficient monomial of \mathcal{A}'_q . In particular, f induces a quasi-homomorphism from \mathcal{A}_q to \mathcal{A}'_q . \square

To introduce a concept of quasi-isomorphism of quantum cluster algebras, we need the following proportional relation, which is a quantum analogy of the one introduced by Fraser [F16]. For elements x, y in a quantum cluster algebra \mathcal{A}_q , we say that x is *proportional to* y , writing $x \propto y$, if $x = q^{\frac{\ell}{2}}py$ for some $\ell \in \mathbb{Z}$ and some coefficient monomial $p \in \mathbb{P}$. Likewise, let \mathcal{A}_q and \mathcal{A}'_q be a pair of quantum cluster algebras, and let f_1 and f_2 be two quasi-homomorphisms from \mathcal{A}_q to \mathcal{A}'_q , we say that f_1 is *proportional to* f_2 , and write $f_1 \propto f_2$, if $f_1(x) \propto f_2(x)$ for every cluster variable $x \in \mathcal{A}_q$.

A quasi-homomorphism $f: \mathcal{A}_q \rightarrow \mathcal{A}'_q$ is called a *quasi-isomorphism* if there is a quasi-homomorphism $g: \mathcal{A}'_q \rightarrow \mathcal{A}_q$ such that the composite $g \circ f$ is proportional to the identity map on \mathcal{A}_q and the composite $f \circ g$ is proportional to the identity map on \mathcal{A}'_q . A *quasi-automorphism* is a quasi-isomorphism from a quantum cluster algebra to itself. Then we are able to consider the *quasi-automorphism group* $\text{QAut}(\mathcal{A}_q)$ of a quantum cluster algebra \mathcal{A}_q , which consists of the equivalence classes of quasi-automorphisms on \mathcal{A}_q up to proportionality.

Note that for a classical (non-quantum) cluster algebra, Assem, Schiffler, and Shramchenko introduced cluster automorphisms in [ASS12]. Similarly, one may consider the quantum version, a *quantum cluster automorphism*, that is, a \mathbb{C}_q -algebra automorphism of a quantum cluster algebra which sends a cluster to a cluster, sends a cluster variable to a cluster variable (without multiplying frozen variables), and sends a frozen variable to a frozen variable. Then it is not hard to see that a quantum cluster automorphism is a special quantum quasi-automorphism, and a quantum quasi-homomorphism f is a quantum cluster automorphism if and only if $(\widetilde{B}, \Lambda) = (\widetilde{B}', \Lambda')$.

It would be interesting to compute the *automorphism group* $\text{Aut}(\mathcal{A}_q)$ of a quantum cluster algebra, which consists of all the quantum cluster automorphisms, and compare it with the group $\text{QAut}(\mathcal{A}_q)$. This is in some sense trivial for the quantum cluster algebras arising from the surfaces. More precisely, let \mathcal{A} be the (classical non-quantum) cluster algebra arising from a unpunctured surface (S, M) with coefficients arising from the boundary segments, and let $\mathcal{A}_{\text{triv}}$ be the associated cluster algebra with trivial coefficients. Since $\text{Aut}(\mathcal{A}) \cong \text{Aut}(\mathcal{A}_{\text{triv}}) \cong \text{MCG}(S, M)$ [BS15, CZ16], where $\text{MCG}(S, M)$ is the mapping class group of the surface, triangulations T and T' are homotopic if and only if the associated matrices \widetilde{B} and \widetilde{B}' coincide. Moreover, the Lambda matrix of the associated quantum surface cluster algebra \mathcal{A}_q comes from the triangulation in the sense of Muller [M16], thus the matrices Λ and Λ' arising from T and T' coincide: $\Lambda = \Lambda'$. Therefore we have an isomorphism $\text{Aut}(\mathcal{A}_q) \cong \text{Aut}(\mathcal{A})$. On the other hand, it is proved in [CS19] that $\text{QAut}(\mathcal{A}) \cong \text{Aut}(\mathcal{A})$. By using the fact that the Lambda matrix is arising from the triangulation, the method used in [CS19] can be extended to the quantum case, and we have an isomorphism of groups: $\text{QAut}(\mathcal{A}_q) \cong \text{Aut}(\mathcal{A}_q)$. To sum up, for (almost all) the surface cluster algebras, both for the quantum case and the non-quantum case, the groups we mentioned above are all isomorphic.

3. BRAID GROUP ACTION ON QUANTUM GRASSMANNIAN CLUSTER ALGEBRAS

In this section, we introduce a braid group action on the quantum Grassmannian cluster algebra $\mathbb{C}_q[\text{Gr}(k, n)]$. We collect some background in the following, [TT91, LZ98, BG02, S05, Lau06, Lau10, GL14]. For $\mathbb{C}_q[\text{Gr}(k, n)]$, we assume that q is not a root of unity.

3.1. Background on the quantum Grassmannian coordinate ring $\mathbb{C}_q[\text{Gr}(k, n)]$. The quantum matrix algebra $\mathbb{C}_q[M(k, n)]$ is a \mathbb{C}_q -algebra generated by $\{x_{ij} \mid 1 \leq i \leq k, 1 \leq j \leq n\}$.

$n\}$ subject to the following relations.

$$\begin{aligned} x_{ij}x_{il} &= qx_{il}x_{ij}, & j < l, \\ x_{ij}x_{rj} &= qx_{rj}x_{ij}, & i < r, \\ x_{ij}x_{rl} &= x_{rl}x_{ij}, & i < r, j > l, \\ x_{ij}x_{rl} &= x_{rl}x_{ij} + (q - q^{-1})x_{il}x_{rj}, & i < r, j < l. \end{aligned}$$

Let $m, n \in \mathbb{Z}_{>0}$. For any $I \subset [1, m]$, $J \subset [1, n]$ with $|I| = |J| = l > 0$, a quantum minor $\Delta_{I,J}$ is defined by

$$\Delta_q^{I,J} = \sum_{\sigma \in S_l} (-q)^{\ell(\sigma)} x_{i_1, j_{\sigma(1)}} \cdots x_{i_l, j_{\sigma(l)}},$$

where $\{i_1 < \cdots < i_l\} = I$, $\{j_1 < \cdots < j_l\} = J$, and $\ell(\sigma)$ is the length of the permutation σ .

For the Grassmannian $\text{Gr}(k, n)$, its quantized coordinate ring $\mathbb{C}_q[\text{Gr}(k, n)]$ is the subalgebra of the quantum matrix algebra $\mathbb{C}_q[M(k, n)]$ generated by the quantum Plücker coordinates

$$\Delta_q^J := \Delta_q^{I,J},$$

where $I = \{1, \dots, k\}$ and $J \subset [1, n]$ with $|J| = k$, see [TT91, GL14]. We assume that the frozen quantum Plücker coordinates Δ_q^J are invertible, where J consists of k consecutive integers (up to cyclic shift).

It is shown in [LZ98] that two quantum Plücker coordinates Δ_q^I, Δ_q^J quasi-commute if and only if I, J are weakly separated. We recall the definition of “weakly separated” in the following.

Definition 3.1. Given two k -subsets I and J of $[1, n]$, denote by $\min(J)$ the minimal element in J and by $\max(I)$ the maximal element in I , we write $I \prec J$ if $\max(I) < \min(J)$. The sets I and J are called weakly separated if at least one of the following two conditions holds:

- (1) $J - I$ can be partitioned into a disjoint union $J - I = J' \sqcup J''$ so that $J' \prec I - J \prec J''$;
- (2) $I - J$ can be partitioned into a disjoint union $I - J = I' \sqcup I''$ so that $I' \prec J - I \prec I''$.


In [S05, Theorem 2], Scott computed quasi-commutation relations on weakly separated subsets.

Theorem 3.2 ([S05, Theorem 2]). *Suppose I and J are weakly separated k -subsets of $[1, n]$. Then we have $\Delta_q^I \Delta_q^J = q^{\Lambda(\Delta_q^I, \Delta_q^J)} \Delta_q^J \Delta_q^I$, where*

$$\Lambda(\Delta_q^I, \Delta_q^J) = \begin{cases} |J''| - |J'|, & \text{if (1) in Definition 3.1 is satisfied,} \\ |I'| - |I''|, & \text{if (2) in Definition 3.1 is satisfied.} \end{cases}$$

Due to Postnikov, every maximal weakly separated collection determines a seed in the cluster algebra $\mathbb{C}[\text{Gr}(k, n)]$, see [S05], where the cluster in the seed is the set of Plücker coordinates parameterized by the subsets in the collection. Therefore each maximal weakly separated collection also determines a seed in the quantum cluster algebra $\mathbb{C}_q[\text{Gr}(k, n)]$, where the Lambda matrix is given by the quasi-commutation relations of the associated Plücker coordinates in Theorem 3.2.

3.2. Commutative quantum cluster monomials and quantum cluster mutations on $\mathbb{C}_q[\text{Gr}(k, n)]$. For two cluster variables x_1, x_2 in the same cluster of a cluster algebra, we have that $x_1 * x_2 = q^{\Lambda(x_1, x_2)} x_2 * x_1$ for some $\Lambda(x_1, x_2)$. We define $x_1 x_2 = x_2 x_1$ by $x_1 * x_2 = q^{\frac{1}{2}\Lambda(x_1, x_2)} x_1 x_2 = q^{\frac{1}{2}\Lambda(x_1, x_2)} x_2 x_1$. Then $x_2 * x_1 = q^{-\frac{1}{2}\Lambda(x_1, x_2)} x_1 x_2 = q^{-\frac{1}{2}\Lambda(x_1, x_2)} x_2 x_1$. Let x_1, \dots, x_m be cluster variables in the same cluster, where we abuse notation and allow


 FIGURE 1. An initial cluster for $\mathbb{C}_q[\text{Gr}(3, 6)]$.

x_i to be a frozen variable. Following Section 2.1.2, we define $x_1^{a_1} x_2^{a_2} \cdots x_n^{a_m}$ (the order of $x_1^{a_1} x_2^{a_2} \cdots x_n^{a_m}$ can change arbitrarily) by

$$x_1^{*a_1} * x_2^{*a_2} * \cdots * x_r^{*a_m} = q^{\frac{1}{2} \sum_{1 \leq i < j \leq m} a_i a_j \Lambda(x_i, x_j)} x_1^{a_1} x_2^{a_2} \cdots x_r^{a_m}$$

and we call $x_1^{a_1} x_2^{a_2} \cdots x_n^{a_m}$ a *commutative quantum cluster monomial*.

While noticing that the commutative quantum cluster monomial is just the element X^a , for an integer vector $a = (a_1, \cdots, a_m)$, in the quantum torus associated to a quantum seed with cluster $\mathbf{x} = \{x_1, \cdots, x_m\}$, this notion has advantages when we extend a map on cluster variables to a map on quantum cluster monomial, see for example the definition of φ given by equation (3.1).

Quantum Grassmannian cluster algebras was studied in [GL11, GL14]. We recall their results in the case of $\mathbb{C}_q[\text{Gr}(3, 6)]$. We compute quantum cluster variables of $\mathbb{C}_q[\text{Gr}(3, 6)]$, starting from the cluster in Figure 1.

We have relations

$$\begin{aligned} \Delta_q^{124} * \Delta_q^{123} &= \frac{1}{q} \Delta_q^{123} * \Delta_q^{124}, & \Delta_q^{124} * \Delta_q^{145} &= q \Delta_q^{145} * \Delta_q^{124}, & \Delta_q^{124} * \Delta_q^{125} &= q \Delta_q^{125} * \Delta_q^{124}, \\ \Delta_q^{124} * \Delta_q^{134} &= q \Delta_q^{134} * \Delta_q^{124}, & \Delta_q^{125} * \Delta_q^{134} &= \Delta_q^{134} * \Delta_q^{125}, & \Delta_q^{123} * \Delta_q^{145} &= q^2 \Delta_q^{145} * \Delta_q^{123}. \end{aligned}$$

Mutating at Δ_q^{124} , we have that

$$(\Delta_q^{124})' = (\Delta_q^{124})^{-1} \Delta_q^{125} \Delta_q^{134} + \Delta_q^{123} (\Delta_q^{124})^{-1} \Delta_q^{145}.$$

On the other hand, we have the relation

$$\Delta_q^{124} * \Delta_q^{135} = q \Delta_q^{125} * \Delta_q^{134} + \frac{1}{q} \Delta_q^{123} * \Delta_q^{145} = q \Delta_q^{125} \Delta_q^{134} + \Delta_q^{123} \Delta_q^{145}.$$

Therefore

$$\Delta_q^{135} = (\Delta_q^{124})^{-1} * (q \Delta_q^{125} \Delta_q^{134} + \Delta_q^{123} \Delta_q^{145}) = (\Delta_q^{124})^{-1} \Delta_q^{125} \Delta_q^{134} + \Delta_q^{123} (\Delta_q^{124})^{-1} \Delta_q^{145}.$$

It follows that $(\Delta_q^{124})' = \Delta_q^{135}$. Using quantum mutations, we see that the quantum cluster variables are:

$$\begin{aligned} &\Delta_q^{124}, \Delta_q^{125}, \Delta_q^{134}, \Delta_q^{135}, \Delta_q^{136}, \Delta_q^{145}, \Delta_q^{146}, \Delta_q^{235}, \Delta_q^{236}, \Delta_q^{245}, \Delta_q^{246}, \Delta_q^{256}, \Delta_q^{346}, \Delta_q^{356}, \\ &q^{-\frac{3}{2}} (\Delta_q^{124} * \Delta_q^{356} - \frac{1}{q} \Delta_q^{123} * \Delta_q^{456}), q^{-\frac{1}{2}} (\Delta_q^{145} * \Delta_q^{236} - \frac{1}{q^2} \Delta_q^{123} * \Delta_q^{456}). \end{aligned}$$

3.3. The maps σ_i and the main results. We define a map φ from the set of cluster variables of $\mathbb{C}[\text{Gr}(k, n)]$ to the set of cluster variables in $\mathbb{C}_q[\text{Gr}(k, n)]$ as follows. First for Δ^J (including frozen variable) in the initial cluster, we define $\varphi(\Delta^J) = \Delta_q^J$. We use induction to define $\varphi(x)$ for other cluster variables in $\mathbb{C}[\text{Gr}(k, n)]$. We apply the same mutation sequence to $\mathbb{C}[\text{Gr}(k, n)]$ and $\mathbb{C}_q[\text{Gr}(k, n)]$. At each step of mutation at some vertex with cluster variable x , we define $\varphi(x') = (\varphi(x))'$, where x' is the cluster variable of $\mathbb{C}[\text{Gr}(k, n)]$ obtained by mutation at x and $(\varphi(x))'$ is the cluster variable obtained by mutation at $\varphi(x)$ in $\mathbb{C}_q[\text{Gr}(k, n)]$. For a cluster variable or frozen variable x , we define $\varphi(x^{-1}) = (\varphi(x))^{-1}$. Starting from the same initial quiver, the mutation rules of quivers for $\mathbb{C}[\text{Gr}(k, n)]$ and $\mathbb{C}_q[\text{Gr}(k, n)]$ are the same. The map φ is clearly a bijection between the set of cluster variables in $\mathbb{C}[\text{Gr}(k, n)]$ and the set of cluster variables in $\mathbb{C}_q[\text{Gr}(k, n)]$. Therefore every cluster variable or frozen variable in $\mathbb{C}_q[\text{Gr}(k, n)]$ is of the form $\varphi(x)$ for some cluster variable x in $\mathbb{C}[\text{Gr}(k, n)]$.

For a cluster monomial $x_1 \cdots x_r$ in $\mathbb{C}[\text{Gr}(k, n)]$ and for $a_1, \dots, a_r \in \mathbb{Z}$, we define

$$(3.1) \quad \varphi(x_1^{a_1} \cdots x_r^{a_r}) = \varphi(x_1)^{a_1} \cdots \varphi(x_r)^{a_r}$$

as a commutative quantum cluster monomial. Here we slightly abuse of terminology. We call $\varphi(x_1)^{a_1} \cdots \varphi(x_r)^{a_r}$ a commutative quantum cluster monomial only requiring that the quantum cluster variables $\varphi(x_1), \dots, \varphi(x_r)$ appear in the same cluster and a_i can be negative.

In [GL11, GL14], it is shown that $\mathbb{C}_q[\text{Gr}(k, n)]$ has a quantum cluster algebra structure. The exchange relations in $\mathbb{C}_q[\text{Gr}(k, n)]$ are of the form $x'_i = x_i^{-1} \prod_{j \rightarrow i} x_j + x_i^{-1} \prod_{i \rightarrow j} x_j$, where the summands on the right hand side are commutative quantum cluster monomials. Moreover, the mutation of quivers in $\mathbb{C}_q[\text{Gr}(k, n)]$ and $\mathbb{C}[\text{Gr}(k, n)]$ are the same. This implies that for any exchange relation $\tilde{x}'_i = \tilde{x}_i^{-1} \prod_{j \rightarrow i} \tilde{x}_j + \tilde{x}_i^{-1} \prod_{i \rightarrow j} \tilde{x}_j$ in $\mathbb{C}[\text{Gr}(k, n)]$, if we replace \tilde{x}_j 's by $\varphi(\tilde{x}_j)$, we obtain an exchange relation $\varphi(\tilde{x}'_i) = \varphi(\tilde{x}_i)^{-1} \prod_{j \rightarrow i} \varphi(\tilde{x}_j) + \varphi(\tilde{x}_i)^{-1} \prod_{i \rightarrow j} \varphi(\tilde{x}_j)$ in $\mathbb{C}_q[\text{Gr}(k, n)]$, where the summands on the right hand side are written as commutative quantum cluster monomials.

Remark 3.3. The algebras $\mathbb{C}[\text{Gr}(k, n)]$, $\mathbb{C}_q[\text{Gr}(k, n)]$ are not isomorphic. Therefore φ cannot be extended to an algebra isomorphism between $\mathbb{C}[\text{Gr}(k, n)]$ and $\mathbb{C}_q[\text{Gr}(k, n)]$.

It is shown in [F17] that in $\mathbb{C}[\text{Gr}(k, n)]$, for every cluster variable or a frozen variable x , $\sigma_i(x) = y_1 y_2 \cdots y_r \tilde{x}$, where y_i is a frozen variable or the inverse of a frozen variable, and \tilde{x} is a cluster variable. In $\mathbb{C}_q[\text{Gr}(k, n)]$, we define $\sigma_i(\varphi(x))$ to be $\varphi(y_1) \varphi(y_2) \cdots \varphi(y_r) \varphi(\tilde{x})$ as a commutative quantum cluster monomial. We define $\sigma_i(\varphi(x^{-1})) = (\sigma_i(\varphi(x)))^{-1}$. We extend σ_i to $\mathbb{C}_q[\text{Gr}(k, n)]$ by defining $\sigma_i(x * y) = \sigma_i(x) * \sigma_i(y)$ and $\sigma_i(ax + by) = a\sigma_i(x) + b\sigma_i(y)$ for any cluster monomials x, y and $a, b \in \mathbb{C}_q$.


For a cluster monomial x in $\mathbb{C}_q[\text{Gr}(k, n)]$, we define $\sigma_i^{-1}(x)$ in the same way as above by replacing σ_i by σ_i^{-1} .

In the following, we describe the map σ_i explicitly on some particular initial cluster. We recall a nice initial cluster \mathbf{x} in $\mathbb{C}_q[\text{Gr}(k, n)]$, which is used frequently when consider Grassmannian cluster algebras, see for example [F17] and the quantum version in [GL14]. The cluster is defined as

$$(3.2) \quad \mathbf{x} = \{\Delta_q^{\dot{I}} : \dot{I} = [1, a] \cup [a + b + 1, b + k], 0 \leq a \leq k, 0 \leq b \leq n - k\}.$$

The quiver of the associated matrix of \mathbf{x} is given in Figure 2, where the coordinate at position (a, b) is

$$(3.3) \quad \Delta_q^{\dot{I}(a,b)}, \dot{I}(a,b) = [1, a] \cup [a + b + 1, b + k].$$


 FIGURE 2. The quiver of the cluster \mathbf{x} in (3.2).

The frozen variables of $\mathbb{C}_q[\mathrm{Gr}(k, n)]$ correspond to k -subsets of the form $[j, j + k - 1]$ for some $j \in [1, n - k + 1]$ or $[1, j] \cup [n - k + j + 1, n]$ for some $j \in [1, k - 1]$.

Let $d = \gcd(k, n)$. Note that $d = 1$ is trivial for our purpose, so we always assume that $d \geq 2$. For \dot{I} in $\Delta_q^i \in \mathbf{x}$, we set

$$(3.4) \quad I = \begin{cases} \dot{I}, & \text{if } \dot{I} \text{ is frozen;} \\ \dot{I}, & \text{if } a + b \not\equiv i \pmod{d}; \\ [1, a - 1] \cup \{a + b\} \cup [a + b + 2, b + k + 1], & \text{if } a + b \equiv i \pmod{d}. \end{cases}$$

Then this modified subset is still a weakly separated collection and the associated set $\mathbf{x}(i) = \{\Delta_q^I\}$ is a cluster, see [F17].

Let \mathfrak{S}_n denote the symmetric group on n symbols. Define $\bar{\sigma}_i \in \mathfrak{S}_n$ as the product of commuting transpositions $\bar{\sigma}_i = \prod_{j=0}^{\frac{n}{d}-1} (jd + i, jd + i + 1)$. This permutation acts by switching adjacent numbers that are equivalent to i and $i + 1 \pmod{d}$.

For $\mathrm{Gr}(k, n)$, in the case that $n < a + k - 1 \leq n + k - 1$, the interval $[a, a + k - 1]$ means $[1, 2, \dots, a + k - n - 1, a, a + 1, \dots, n]$. For example, in the case of $\mathrm{Gr}(3, 6)$, $[5, 6, 7]$ means $[1, 5, 6]$. The following lemma follows from [F17, Section 5] and the definition of σ_i on $\mathbb{C}_q[\mathrm{Gr}(k, n)]$ in the beginning of this subsection.

Lemma 3.4. *The map σ_i on $\mathbf{x}(i)$ is given by:*

$$(3.5) \quad \sigma_i(\Delta_q^I) = \begin{cases} \Delta_q^{[j-1, j+k-2]} (\Delta_q^{[j, j+k-1]})^{-1} \Delta_q^{[j+1, j+k]}, & \text{if } I = [j, j + k - 1] \text{ with } j \equiv i + 1 \pmod{d}; \\ \Delta_q^I, & \text{if } I = [j, j + k - 1] \text{ with } j \not\equiv i + 1 \pmod{d}; \\ \Delta_q^{\bar{\sigma}_i(I)}, & \text{otherwise,} \end{cases}$$

where terms on the right hand side are commutative quantum cluster monomials.

Example 3.5. In $\mathbb{C}_q[\mathrm{Gr}(3, 6)]$, $\sigma_1(\Delta_q^{124}) = \Delta_q^{125}$, and

$$\begin{aligned} \sigma_1(\Delta_q^{234}) &= \Delta_q^{123} (\Delta_q^{234})^{-1} \Delta_q^{345} = \Delta_q^{123} * (\Delta_q^{234})^{-1} * \Delta_q^{345} = \Delta_q^{345} * (\Delta_q^{234})^{-1} * \Delta_q^{123} \\ &= \frac{1}{q} (\Delta_q^{234})^{-1} * \Delta_q^{123} * \Delta_q^{345} = \frac{1}{q} \Delta_q^{123} * \Delta_q^{345} * (\Delta_q^{234})^{-1} \\ &= q (\Delta_q^{234})^{-1} * \Delta_q^{345} * \Delta_q^{123} = q \Delta_q^{345} * \Delta_q^{123} * (\Delta_q^{234})^{-1}. \end{aligned}$$

Lemma 3.6. *Let $i \in [d - 1]$ and let x_1, x_2 be any pair of cluster variables (including frozen variables) in a cluster. Then for $\tilde{x}_1 \in \{x_1, x_1^{-1}\}$, $\tilde{x}_2 \in \{x_2, x_2^{-1}\}$, we have that $\tilde{x}_1 * \tilde{x}_2 =$*

$q^c \tilde{x}_2 * \tilde{x}_1$ for some c implies that $\sigma_i(\tilde{x}_1) * \sigma_i(\tilde{x}_2) = q^c \sigma_i(\tilde{x}_2) * \sigma_i(\tilde{x}_1)$, and $\sigma_i^{-1}(\tilde{x}_1) * \sigma_i^{-1}(\tilde{x}_2) = q^c \sigma_i^{-1}(\tilde{x}_2) * \sigma_i^{-1}(\tilde{x}_1)$.

Proof. We will prove the statement for σ_i . The proof of the result for σ_i^{-1} is the similar.

Fraser [F17] has proved that in non-quantum case, σ_i sends a cluster to a cluster up to frozen variables. Therefore by the definition of σ_i in quantum case in the beginning of this subsection, we have that for two cluster variables x_1, x_2 in the same cluster, $\sigma_i(x_1), \sigma_i(x_2)$ are cluster variables (possibly multiply by Laurent monomials of frozen variables).

It suffices to prove the result for x_1, x_2 in one particular cluster. Indeed, suppose that the result is true for the cluster $\mathbf{x} = (x_1, \dots, x_m)$. Let x'_r be the quantum cluster variable obtained by mutating x_r and let $\mathbf{x}' = (x_1, \dots, x'_r, \dots, x_m)$. Then $x'_r = x_r^{-1} x_{j_1} \cdots x_{j_t} + x_r^{-1} x_{e_1} \cdots x_{e_s}$, where the right hand side are commutative quantum cluster monomials, s, t, j_l, e_l are some integers. Therefore $x'_r = q^a x_r^{-1} * x_{j_1} * \cdots * x_{j_t} + q^b x_r^{-1} * x_{e_1} * \cdots * x_{e_s}$ for some a, b . Suppose that $x_f * x'_r = q^c x'_r * x_f$ for some c and some $x_f \in \mathbf{x}$. We want to show that $\sigma_i(x_f) * \sigma_i(x'_r) = q^c \sigma_i(x'_r) * \sigma_i(x_f)$. We have

$$x_f * x'_r = x_f * (q^a x_r^{-1} * x_{j_1} * \cdots * x_{j_t} + q^b x_r^{-1} * x_{e_1} * \cdots * x_{e_s}).$$

Since $x_f, x_r, x_{j_l}, x_{e_l}$ are in the same cluster, we have

$$x_f * x'_r = (q^{a+d} x_r^{-1} * x_{j_1} * \cdots * x_{j_t} + q^{b+d'} x_r^{-1} * x_{e_1} * \cdots * x_{e_s}) * x_f,$$

for some d, d' . Since we have assumed that $x_f * x'_r = q^c x'_r * x_f$, we have $d = d' = c$. Since we have assumed that the result is true for the cluster $\mathbf{x} = (x_1, \dots, x_m)$, we have that

$$\begin{aligned} \sigma_i(x_f) * \sigma_i(x'_r) &= \sigma_i(x_f) * \sigma_i(q^a x_r^{-1} * x_{j_1} * \cdots * x_{j_t} + q^b x_r^{-1} * x_{e_1} * \cdots * x_{e_s}) \\ &= \sigma_i(x_f) * (q^a \sigma_i(x_r^{-1}) * \sigma_i(x_{j_1}) * \cdots * \sigma_i(x_{j_t}) + q^b \sigma_i(x_r^{-1}) * \sigma_i(x_{e_1}) * \cdots * \sigma_i(x_{e_s})) \\ &= (q^{a+c} \sigma_i(x_r^{-1}) * \sigma_i(x_{j_1}) * \cdots * \sigma_i(x_{j_t}) + q^{b+c} \sigma_i(x_r^{-1}) * \sigma_i(x_{e_1}) * \cdots * \sigma_i(x_{e_s})) * \sigma_i(x_f). \end{aligned}$$

Therefore $\sigma_i(x_f) * \sigma_i(x'_r) = q^c \sigma_i(x'_r) * \sigma_i(x_f)$. It follows that $x_f^{\pm 1} * x'_r^{\pm 1} = q^c x'_r^{\pm 1} * x_f^{\pm 1}$ for some c implies that $\sigma_i(x_f^{\pm 1}) * \sigma_i(x'_r^{\pm 1}) = q^c \sigma_i(x'_r^{\pm 1}) * \sigma_i(x_f^{\pm 1})$.

We now prove the following: for cluster variables (including frozen variables) Δ_q^J, Δ_q^K in $\mathbf{x}(i)$ (see Equation (3.4)), we have that $\Delta_q^J * \Delta_q^K = q^{\lambda_{J,K}} \Delta_q^K * \Delta_q^J$ implies that $\sigma_i(\Delta_q^J) * \sigma_i(\Delta_q^K) = q^{\lambda_{J,K}} \sigma_i(\Delta_q^K) * \sigma_i(\Delta_q^J)$. From this, it follows that $(\Delta_q^J)^{\pm 1} * (\Delta_q^K)^{\pm 1} = q^{\lambda_{J,K}} (\Delta_q^K)^{\pm 1} * (\Delta_q^J)^{\pm 1}$ implies that $\sigma_i((\Delta_q^J)^{\pm 1}) * \sigma_i((\Delta_q^K)^{\pm 1}) = q^{\lambda_{J,K}} \sigma_i((\Delta_q^K)^{\pm 1}) * \sigma_i((\Delta_q^J)^{\pm 1})$.

The proof is straightforward by checking all cases in (3.5). Note that the cluster $\mathbf{x}(i)$ is a very special cluster.

Case 1. Both of Δ_q^J and Δ_q^K are frozen variables. According to the formula (3.5), σ_i sends a frozen Δ_q^J either to itself or to $\Delta_q^{J'} (\Delta_q^J)^{-1} \Delta_q^{J''}$, where J' is the shift of J to the left by 1 and J'' is the shift of J to the right by 1.

If both of Δ_q^J and Δ_q^K are not change under σ_i , it is clear that the quasi-commutative relation is preserved.

Suppose that under the map σ_i , Δ_q^J is not changed, and Δ_q^K is changed to $\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}$. We have that $K = [a, b]$, $K' = [a-1, b-1]$, $K'' = [a+1, b+1]$ for some a, b (here we only write the case that $a-1 \geq 1$, $b+1 \leq n$; the cases that $a-1 = 0 \equiv n \pmod{n}$ or $b+1 = n+1 \equiv 1 \pmod{n}$ are similar). According to Theorem 3.2 and the expression of K, K', K'' , we have that

$$\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''} = \Delta_q^{K'} * (\Delta_q^K)^{-1} * \Delta_q^{K''},$$

and

$$\begin{aligned} \Delta_q^J * (\Delta_q^K)^{-1} &= q^{-\lambda_{J,K}} (\Delta_q^K)^{-1} * \Delta_q^J, & \Delta_q^J * \Delta_q^{K'} &= q^{\lambda_{J,K-1}} \Delta_q^{K'} * \Delta_q^J, \\ \Delta_q^J * \Delta_q^{K''} &= q^{\lambda_{J,K+1}} \Delta_q^{K''} * \Delta_q^J. \end{aligned}$$

Therefore

$$\Delta_q^J * (\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}) = q^{\lambda_{J,K}} (\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}) * \Delta_q^J.$$

For example, in $\mathbb{C}[\text{Gr}(3, 6)]$ we have that $\Delta_q^{123} * \Delta_q^{345} = q^2 \Delta_q^{345} * \Delta_q^{123}$. Apply σ_2 , we obtain

$$\sigma_2(\Delta_q^{123}) = \Delta_q^{123}, \quad \sigma_2(\Delta_q^{345}) = \Delta_q^{234} (\Delta_q^{345})^{-1} \Delta_q^{456}.$$

Since

$$\Delta_q^{123} * \Delta_q^{234} = q \Delta_q^{234} * \Delta_q^{123}, \quad \Delta_q^{123} * (\Delta_q^{345})^{-1} = q^{-2} (\Delta_q^{345})^{-1} * \Delta_q^{123}, \quad \Delta_q^{123} * \Delta_q^{456} = q^3 \Delta_q^{456} * \Delta_q^{123},$$

we have that $\sigma_2(\Delta_q^{123}) * \sigma_2(\Delta_q^{345}) = q^2 \sigma_2(\Delta_q^{345}) * \sigma_2(\Delta_q^{123})$.

Suppose that under the map σ_i , Δ_q^J is changed to $\Delta_q^{J'} (\Delta_q^J)^{-1} \Delta_q^{J''}$, and Δ_q^K is changed to $\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}$. We have that $J = [a, b]$, $J' = [a-1, b-1]$, $J'' = [a+1, b+1]$ for some a, b and $K = [c, d]$, $K' = [c-1, d-1]$, $K'' = [c+1, d+1]$ for some c, d (here we only write the cases of $a-1 > 0$, $c-1 > 0$, $b+1 \leq n$, $d+1 \leq n$; the cases of $a=1$ or $c=1$ or $b=n$ or $d=n$ are similar).

Using the result in the case above, since $(\Delta_q^J)^{-1} * \Delta_q^K = q^{-\lambda_{J,K}} \Delta_q^K * (\Delta_q^J)^{-1}$, we have that

$$(\Delta_q^J)^{-1} * (\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}) = q^{-\lambda_{J,K}} (\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}) * (\Delta_q^J)^{-1}.$$

We have $\Delta_q^{J'} * \Delta_q^K = q^{\lambda_{J,K}+1} \Delta_q^K * \Delta_q^{J'}$. Therefore

$$\Delta_q^{J'} * (\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}) = q^{\lambda_{J,K}+1} (\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}) * \Delta_q^{J'}.$$

We have $\Delta_q^{J''} * \Delta_q^K = q^{\lambda_{J,K}-1} \Delta_q^K * \Delta_q^{J''}$. Therefore

$$\Delta_q^{J''} * (\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}) = q^{\lambda_{J,K}-1} (\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}) * \Delta_q^{J''}.$$

It follows that

$$(\Delta_q^{J'} (\Delta_q^J)^{-1} \Delta_q^{J''}) * (\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}) = q^{\lambda_{J,K}} (\Delta_q^{K'} (\Delta_q^K)^{-1} \Delta_q^{K''}) * (\Delta_q^{J'} (\Delta_q^J)^{-1} \Delta_q^{J''}).$$

Case 2. Δ_q^J is a frozen variable and Δ_q^K is a cluster variable. Here we only write the case that σ_i does not change Δ_q^J , J is an interval, and σ_i changes Δ_q^K to $\Delta_q^{\bar{\sigma}_i(K)}$. The other cases are similar.

By Theorem 3.2, we either have $\lambda_{J,K} = |K_1| - |K_2|$, where $K - J = K_1 \sqcup K_2$, $K_1 \prec J - K \prec K_2$, or $\lambda_{J,K} = |J_2| - |J_1|$, where $J - K = J_1 \sqcup J_2$, $J_1 \prec K - J \prec J_2$.

Recall that $\bar{\sigma}_i$ is the product $(i, i+1)(d+i, d+i+1) \cdots (n-d+i, n-d+i+1)$ of simple reflections. After applying σ_i , we have either $\bar{\sigma}_i(K) - J = \bar{\sigma}_i(K_1) \sqcup \bar{\sigma}_i(K_2)$, $\bar{\sigma}_i(K_1) \prec J - \bar{\sigma}_i(K) \prec \bar{\sigma}_i(K_2)$ or $J - \bar{\sigma}_i(K) = J_1 \sqcup J_2$, $J_1 \prec \bar{\sigma}_i(K) - J \prec J_2$. Since $|\bar{\sigma}_i(K_j)| = |K_j|$, $j = 1, 2$, we have $|\bar{\sigma}_i(K_1)| - |\bar{\sigma}_i(K_2)| = |K_1| - |K_2|$ and hence the quasi-commutative relation does not change.

For example, in $\mathbb{C}[\text{Gr}(3, 6)]$ we have that $\Delta_q^{123} * \Delta_q^{146} = q^2 \Delta_q^{146} * \Delta_q^{123}$. We have that

$$\sigma_1(\Delta_q^{123}) = \Delta_q^{123}, \quad \sigma_1(\Delta_q^{146}) = \Delta_q^{256},$$

and $\Delta_q^{123} * \Delta_q^{256} = q^2 \Delta_q^{256} * \Delta_q^{123}$. Therefore $\sigma_1(\Delta_q^{123}) * \sigma_1(\Delta_q^{256}) = q^2 \sigma_1(\Delta_q^{256}) * \sigma_1(\Delta_q^{123})$.

Case 3. Both of Δ_q^J and Δ_q^K are cluster variables. In this case, Δ_q^J is sent to $\Delta_q^{\bar{\sigma}_i(J)}$ and Δ_q^K is sent to $\Delta_q^{\bar{\sigma}_i(K)}$. This case is very similar to Case 2.

By Theorem 3.2, we either have $\lambda_{J,K} = |K_1| - |K_2|$, where $K - J = K_1 \sqcup K_2$, $K_1 \prec J - K \prec K_2$, or $\lambda_{J,K} = |J_2| - |J_1|$, where $J - K = J_1 \sqcup J_2$, $J_1 \prec K - J \prec J_2$.

After applying σ_i , we have either $\bar{\sigma}_i(K) - \bar{\sigma}_i(J) = \bar{\sigma}_i(K_1) \sqcup \bar{\sigma}_i(K_2)$, $\bar{\sigma}_i(K_1) \prec \bar{\sigma}_i(J) - \bar{\sigma}_i(K) \prec \bar{\sigma}_i(K_2)$ or $\bar{\sigma}_i(J) - \bar{\sigma}_i(K) = \bar{\sigma}_i(J_1) \sqcup \bar{\sigma}_i(J_2)$, $\bar{\sigma}_i(J_1) \prec \bar{\sigma}_i(K) - \bar{\sigma}_i(J) \prec \bar{\sigma}_i(J_2)$. Since $|\bar{\sigma}_i(K_j)| = |K_j|$ and $|\bar{\sigma}_i(J_j)| = |J_j|$, $j = 1, 2$, we have $|\bar{\sigma}_i(K_1)| - |\bar{\sigma}_i(K_2)| = |K_1| - |K_2|$ and $|\bar{\sigma}_i(J_2)| - |\bar{\sigma}_i(J_1)| = |J_2| - |J_1|$, and hence the quasi-commutative relation does not change.

For example, in $\mathbb{C}[\text{Gr}(3, 6)]$ we have that $\Delta_q^{124} * \Delta_q^{134} = q\Delta_q^{134} * \Delta_q^{124}$. We have that

$$\sigma_1(\Delta_q^{124}) = \Delta_q^{125}, \quad \sigma_1(\Delta_q^{134}) = \Delta_q^{235},$$

and $\Delta_q^{125} * \Delta_q^{235} = q\Delta_q^{235} * \Delta_q^{125}$. Therefore $\sigma_1(\Delta_q^{125}) * \sigma_1(\Delta_q^{235}) = q\sigma_1(\Delta_q^{235}) * \sigma_1(\Delta_q^{125})$. \square

Lemma 3.7. *In $\mathbb{C}_q[\text{Gr}(k, n)]$, for a commutative quantum cluster monomial $x_1^{a_1} x_2^{a_2} \cdots x_r^{a_r}$, where x_1, \dots, x_r are cluster variables or frozen variables which are in the same cluster, $a_i \in \mathbb{Z}$, we have that $\sigma_i(x_1^{a_1} x_2^{a_2} \cdots x_r^{a_r}) = \sigma_i(x_1)^{a_1} \cdots \sigma_i(x_r)^{a_r}$ as a commutative quantum cluster monomial.*

Proof. By [F17] and the definition of σ_i in the quantum setting in the beginning of this subsection, we have that $\sigma_i(x_1), \dots, \sigma_i(x_r)$ are cluster variables in the same cluster (possibly multiply by Laurent monomials of frozen variables).

By definition of commutative quantum cluster monomial in Section 3.2, we have that

$$x_1^{a_1} x_2^{a_2} \cdots x_r^{a_r} = q^{-\frac{1}{2} \sum_{1 \leq s < t \leq r} a_s a_t \Lambda(x_s, x_t)} x_1^{*a_1} * \cdots * x_r^{*a_r},$$

and

$$\sigma_i(x_1)^{a_1} \sigma_i(x_2)^{a_2} \cdots \sigma_i(x_r)^{a_r} = q^{-\frac{1}{2} \sum_{1 \leq s < t \leq r} a_s a_t \Lambda(\sigma_i(x_s), \sigma_i(x_t))} \sigma_i(x_1)^{*a_1} * \cdots * \sigma_i(x_r)^{*a_r}.$$

By the definition of σ_i in the beginning of this subsection, we have that $\sigma_i(x_1^{*a_1} * \cdots * x_r^{*a_r}) = \sigma_i(x_1)^{*a_1} * \cdots * \sigma_i(x_r)^{*a_r}$. By Lemma 3.6, for any $1 \leq s < t \leq r$, we have that $\Lambda(x_s, x_t) = \Lambda(\sigma_i(x_s), \sigma_i(x_t))$. Therefore $\sigma_i(x_1^{a_1} x_2^{a_2} \cdots x_r^{a_r}) = \sigma_i(x_1)^{a_1} \cdots \sigma_i(x_r)^{a_r}$. \square

Lemma 3.8. *For any cluster monomial x (x can contain frozen variables or inverses of frozen variables) in $\mathbb{C}[\text{Gr}(k, n)]$ and any $i \in [d - 1]$, $d = \text{gcd}(k, n)$, we have $\varphi(\sigma_i(x)) = \sigma_i(\varphi(x))$ and $\varphi(\sigma_i^{-1}(x)) = \sigma_i^{-1}(\varphi(x))$.*

Proof. For a cluster variable x in $\mathbb{C}[\text{Gr}(k, n)]$, by the definition in the beginning of this subsection, we have that

$$(3.6) \quad \sigma_i(\varphi(x)) = \varphi(\sigma_i(x)).$$

Suppose that $x = x_1 \cdots x_r$, where each x_i is a cluster variable or a frozen variable or the inverse of a frozen variable. By Lemma 3.7, we have that $\sigma_i(x) = \sigma_i(x_1) \cdots \sigma_i(x_r)$. By definition (3.1),

$$\varphi(\sigma_i(x)) = \varphi(\sigma_i(x_1) \cdots \sigma_i(x_r)) = \varphi(\sigma_i(x_1)) \cdots \varphi(\sigma_i(x_r)) = \sigma_i(\varphi(x_1)) \cdots \sigma_i(\varphi(x_r)),$$

where in the last step we used (3.6). Since $\varphi(x_1) \cdots \varphi(x_r)$ is a commutative quantum cluster monomial, by Lemma 3.7 and the definition of φ , we have that

$$\sigma_i(\varphi(x_1)) \cdots \sigma_i(\varphi(x_r)) = \sigma_i(\varphi(x_1) \cdots \varphi(x_r)) = \sigma_i(\varphi(x_1 \cdots x_r)) = \sigma_i(\varphi(x)).$$

The prove of $\varphi(\sigma_i^{-1}(x)) = \sigma_i^{-1}(\varphi(x))$ is the similar. \square

Remark 3.9. Consider a quantum cluster variable

$$(3.7) \quad x'_r = x_r^{-1} \prod_{j \rightarrow l} x_l + x_r^{-1} \prod_{l \rightarrow j} x_l,$$

obtained by mutating x_r in a cluster $\mathbf{x} = (x_1, \dots, x_m)$. On the one hand, $\sigma_i(x'_r)$ is defined to be $\varphi(\sigma_i(\tilde{x}'_r))$, where \tilde{x}'_r is the cluster variable in $\mathbb{C}[\text{Gr}(k, n)]$ corresponding to x'_r , where the map φ is defined in the beginning of this subsection. On the other hand, $\sigma_i(x'_r)$ should also be $\sigma_i(x_r^{-1} \prod_{j \rightarrow l} x_l + x_r^{-1} \prod_{l \rightarrow j} x_l)$. By Lemma 3.7, $\sigma_i(x_r^{-1} \prod_{j \rightarrow l} x_l + x_r^{-1} \prod_{l \rightarrow j} x_l) = \sigma_i(x_r)^{-1} \prod_{j \rightarrow l} \sigma_i(x_l) + \sigma_i(x_r)^{-1} \prod_{l \rightarrow j} \sigma_i(x_l)$.

We now verify that $\sigma_i(x_r)^{-1} \prod_{j \rightarrow l} \sigma_i(x_l) + \sigma_i(x_r)^{-1} \prod_{l \rightarrow j} \sigma_i(x_l) = \varphi(\sigma_i(\tilde{x}'_r))$. In the beginning of this subsection, we see that the map φ sends a cluster in $\mathbb{C}[\text{Gr}(k, n)]$ to a cluster in $\mathbb{C}_q[\text{Gr}(k, n)]$. Let \tilde{x}_j be cluster variables in $\mathbb{C}[\text{Gr}(k, n)]$ such that $\varphi(\tilde{x}_j) = x_j$. The exchange

relation in $\mathbb{C}[\text{Gr}(k, n)]$ corresponding to (3.7) is $\tilde{x}'_r = \tilde{x}_r^{-1} \prod_{j \rightarrow l} \tilde{x}_l + \tilde{x}_r^{-1} \prod_{l \rightarrow j} \tilde{x}_l$. We have that

$$\sigma_i(\tilde{x}'_r) = \sigma_i(\tilde{x}_r)^{-1} \prod_{j \rightarrow l} \sigma_i(\tilde{x}_l) + \sigma_i(\tilde{x}_r)^{-1} \prod_{l \rightarrow j} \sigma_i(\tilde{x}_l).$$

This is an exchange relation in $\mathbb{C}[\text{Gr}(k, n)]$ up to multiplying frozen variables. As discussed in the beginning of this subsection, if we replace $\sigma_i(\tilde{x}_r)$ by $\varphi(\sigma_i(\tilde{x}_r))$, then we obtain an exchange relation in $\mathbb{C}_q[\text{Gr}(k, n)]$ up to multiplying frozen variables. That is, $\varphi(\sigma_i(\tilde{x}'_r))$ is equal to

$$\varphi(\sigma_i(\tilde{x}_r))^{-1} \prod_{j \rightarrow l} \varphi(\sigma_i(\tilde{x}_l)) + \varphi(\sigma_i(\tilde{x}_r))^{-1} \prod_{l \rightarrow j} \varphi(\sigma_i(\tilde{x}_l)).$$

By Lemma 3.8, this is equal to

$$\sigma_i(\varphi(\tilde{x}_r))^{-1} \prod_{j \rightarrow l} \sigma_i(\varphi(\tilde{x}_l)) + \sigma_i(\varphi(\tilde{x}_r))^{-1} \prod_{l \rightarrow j} \sigma_i(\varphi(\tilde{x}_l)),$$

which is equal to $\sigma_i(x_r)^{-1} \prod_{j \rightarrow l} \sigma_i(x_l) + \sigma_i(x_r)^{-1} \prod_{l \rightarrow j} \sigma_i(x_l)$.

Lemma 3.10. *For $i \in [d-1]$, σ_i and σ_i^{-1} preserve exchange relations in $\mathbb{C}_q[\text{Gr}(k, n)]$.*

Proof. We will prove the result for σ_i . The proof of the result for σ_i^{-1} is similar.

It is shown in [GL11, GL14] that $\mathbb{C}_q[\text{Gr}(k, n)]$ has a quantum cluster algebra structure. Let x be a quantum cluster variable in a cluster. Mutate at x we obtain that

$$(3.8) \quad x' = x^{-1} \prod_{x_j \rightarrow x} x_j + x^{-1} \prod_{x \rightarrow x_j} x_j,$$

where $\prod_{x_j \rightarrow x} x_j$ and $\prod_{x \rightarrow x_j} x_j$ are commutative quantum cluster monomials. We rewrite the identity using non-commutative product:

$$(3.9) \quad x' = q^a x^{-1} * c_1 * \cdots * c_r + q^b x^{-1} * d_1 * \cdots * d_s,$$

where $c_1, \dots, c_r, d_1, \dots, d_s$ are x_j 's in (3.8) in some order, and $a, b \in \frac{1}{2}\mathbb{Z}$. Therefore

$$x * x' = q^a c_1 * \cdots * c_r + q^b d_1 * \cdots * d_s.$$

This is an exchange relation in $\mathbb{C}_q[\text{Gr}(k, n)]$. We now verify that

$$(3.10) \quad \sigma_i(x) * \sigma_i(x') = q^a \sigma_i(c_1) * \cdots * \sigma_i(c_r) + q^b \sigma_i(d_1) * \cdots * \sigma_i(d_s).$$

By Lemma 3.7 and Remark 3.9, apply σ_i to (3.8), we have that

$$(3.11) \quad \sigma_i(x') = \sigma_i(x)^{-1} \prod_{x_j \rightarrow x} \sigma_i(x_j) + \sigma_i(x)^{-1} \prod_{x \rightarrow x_j} \sigma_i(x_j)$$

where the right hand side are commutative quantum cluster monomials.

Note that $c_1, \dots, c_r, d_1, \dots, d_s$ are x_j 's in (3.8) in some order. There are some a', b' such that the right hand side of (3.11) is

$$q^{a'} \sigma_i(x)^{-1} * \sigma_i(c_1) * \cdots * \sigma_i(c_r) + q^{b'} \sigma_i(x)^{-1} * \sigma_i(d_1) * \cdots * \sigma_i(d_s).$$

By Lemma 3.6, $c_j * c_{j'} = q^{\lambda_{j,j'}} c_j * c_{j'}$ implies that $\sigma_i(c_j) * \sigma_i(c_{j'}) = q^{\lambda_{j,j'}} \sigma_i(c_j) * \sigma_i(c_{j'})$. Therefore (3.8), (3.9), and (3.11) implies that $a = a', b = b'$. Hence

$$\sigma_i(x') = q^a \sigma_i(x)^{-1} * \sigma_i(c_1) * \cdots * \sigma_i(c_r) + q^b \sigma_i(x)^{-1} * \sigma_i(d_1) * \cdots * \sigma_i(d_s).$$

Multiply $\sigma_i(x)$ from the left hand side, we obtain (3.10). □

The 3-term quantum Plücker relations are special exchange relations. We conjecture that σ_i and σ_i^{-1} also preserve r -term ($r \geq 4$) quantum Plücker relations.

Conjecture 3.11. *The maps σ_i and σ_i^{-1} also preserve r -term quantum Plücker relations for $r \geq 4$ in $\mathbb{C}_q[\text{Gr}(k, n)]$.*

Remark 3.12. Let $\text{Gr}^\circ(k, n) \subset \text{Gr}(k, n)$ be the quasi-affine variety defined by non-vanishing of the cyclically consecutive Plücker coordinates $P_{i+1, \dots, i+k}$, $i \in [n]$, and the indices are treated modulo n . The coordinate ring $\mathbb{C}[\text{Gr}^\circ(k, n)]$ is the localization of $\mathbb{C}[\text{Gr}(k, n)]$ at the cyclically consecutive Plücker coordinates. The distinction between $\mathbb{C}[\text{Gr}^\circ(k, n)]$ and $\mathbb{C}[\text{Gr}(k, n)]$ is unimportant from a combinatorial perspective, and in [F17] Fraser freely translate between them, see Section 3 in [F17]. In [F17], Fraser proved that σ_i 's are regular automorphisms of $\text{Gr}^\circ(k, n)$. These automorphisms induce algebra automorphisms on $\mathbb{C}[\text{Gr}^\circ(k, n)]$.

In [FWZ], Section 6.8, Example 6.8.6, it is observed that in general, the ideal of relations among the cluster variables of $\mathbb{C}[\text{Gr}(k, n)]$ is not generated by the exchange relations. Therefore in quantum case, in general, the ideal of relations among the cluster variables of $\mathbb{C}_q[\text{Gr}(k, n)]$ is not generated by quasi-commutative relations and the exchange relations.

In this paper, we only proved that σ_i 's preserve quasi-commutative relations and exchange relations, and σ_i 's are automorphisms on $\mathbb{C}_q[\text{Gr}(k, n)]$.

We give an example that σ_i preserves a 4-term quantum Plücker relation to show evidence of Conjecture 3.11.

Example 3.13. In $\mathbb{C}_q[\text{Gr}(3, 6)]$, we have that

$$\Delta_q^{124} * \Delta_q^{356} = \frac{1}{q} \Delta_q^{123} * \Delta_q^{456} + q \Delta_q^{125} * \Delta_q^{346} - q^2 \Delta_q^{126} * \Delta_q^{345}.$$

We now verify that σ_1 preserves the above identity. That is, by Table 1, we need to verify that

$$\Delta_q^{125} * (\Delta_q^{136} (\Delta_q^{156})^{-1} \Delta_q^{456}) = \frac{1}{q} \Delta_q^{123} * \Delta_q^{456} + q (\Delta_q^{126} (\Delta_q^{156})^{-1} \Delta_q^{145}) * \Delta_q^{356} - q^2 \Delta_q^{126} * \Delta_q^{345}.$$

We have that

$$\begin{aligned} \Delta_q^{136} (\Delta_q^{156})^{-1} \Delta_q^{456} &= q^{-1} \Delta_q^{136} * \Delta_q^{456} * (\Delta_q^{156})^{-1}, \\ \Delta_q^{126} (\Delta_q^{156})^{-1} \Delta_q^{145} &= q \Delta_q^{126} * \Delta_q^{145} * (\Delta_q^{156})^{-1}. \end{aligned}$$

Therefore we need to verify

$$q^{-1} \Delta_q^{125} * \Delta_q^{136} * \Delta_q^{456} * (\Delta_q^{156})^{-1} = \frac{1}{q} \Delta_q^{123} * \Delta_q^{456} + q^2 \Delta_q^{126} * \Delta_q^{145} * (\Delta_q^{156})^{-1} * \Delta_q^{356} - q^2 \Delta_q^{126} * \Delta_q^{345}.$$

We have that $(\Delta_q^{156})^{-1} * \Delta_q^{356} = q^{-1} \Delta_q^{356} * (\Delta_q^{156})^{-1}$. Therefore we need to verify that

$$q^{-1} \Delta_q^{125} * \Delta_q^{136} * \Delta_q^{456} * (\Delta_q^{156})^{-1} = \frac{1}{q} \Delta_q^{123} * \Delta_q^{456} + q \Delta_q^{126} * \Delta_q^{145} * \Delta_q^{356} * (\Delta_q^{156})^{-1} - q^2 \Delta_q^{126} * \Delta_q^{345}.$$

That is,

$$\Delta_q^{125} * \Delta_q^{136} * \Delta_q^{456} = \Delta_q^{123} * \Delta_q^{456} * \Delta_q^{156} + q^2 \Delta_q^{126} * \Delta_q^{145} * \Delta_q^{356} - q^3 \Delta_q^{126} * \Delta_q^{345} * \Delta_q^{156}.$$

This is checked directly on computer.

Lemma 3.14. *Up to Conjecture 3.11, the maps σ_i and σ_i^{-1} are \mathbb{C}_q -algebra automorphisms of the quantum cluster algebra $\mathbb{C}_q[\text{Gr}(k, n)]$.*

Proof. Assume that Conjecture 3.11 is true. Then lemmas 3.6 and 3.10 imply that σ_i and σ_i^{-1} are algebra homomorphisms of $\mathbb{C}_q[\text{Gr}(k, n)]$.

We now check that σ_i and σ_i^{-1} are inverses of each other on $\mathbb{C}_q[\text{Gr}(k, n)]$. In the beginning of this subsection, we see that every quantum Plücker coordinate of $\mathbb{C}_q[\text{Gr}(k, n)]$ is of the form $\varphi(x)$, where x is a Plücker coordinate of $\mathbb{C}[\text{Gr}(k, n)]$. By Lemma 3.8, we have that $\sigma_i^{-1}(\sigma_i(\varphi(x))) = \sigma_i^{-1}(\varphi(\sigma_i(x))) = \varphi(\sigma_i^{-1}(\sigma_i(x)))$. It is shown in [F17] that σ_i, σ_i^{-1} are inverses

of each other on $\mathbb{C}[\text{Gr}(k, n)]$. Therefore $\varphi(\sigma_i^{-1}(\sigma_i(x))) = \varphi(x)$ and hence $\sigma_i^{-1}(\sigma_i(\varphi(x))) = \varphi(x)$. Similarly, $\sigma_i(\sigma_i^{-1}(\varphi(x))) = \varphi(x)$.

Every element in $\mathbb{C}_q[\text{Gr}(k, n)]$ is a \mathbb{C}_q -linear combination of products of quantum Plücker coordinates of $\mathbb{C}_q[\text{Gr}(k, n)]$. To show that σ_i and σ_i^{-1} are inverses of each other on $\mathbb{C}_q[\text{Gr}(k, n)]$, it suffices to show that $\sigma_i^{-1}(\sigma_i(x_1 * \cdots * x_r)) = x_1 * \cdots * x_r$ and $\sigma_i(\sigma_i^{-1}(x_1 * \cdots * x_r)) = x_1 * \cdots * x_r$, where x_i 's are quantum Plücker coordinates. This follows from the fact that σ_i, σ_i^{-1} are algebra homomorphisms on $\mathbb{C}_q[\text{Gr}(k, n)]$ and $\sigma_i^{-1}(\sigma_i(x_i)) = x_i, \sigma_i(\sigma_i^{-1}(x_i)) = x_i$ for quantum Plücker coordinates x_i . \square

Our main result of this section is the following.

Theorem 3.15. *Let $k \leq n$ and $d = \gcd(k, n)$.*

- (i) *Assume that Conjecture 3.11 is true. Then for any $1 \leq i \leq d - 1$, the σ_i is a quasi-automorphism of $\mathbb{C}_q[\text{Gr}(k, n)]$.*
- (ii) *The maps σ_i 's give rise to a braid group action on $\mathbb{C}_q[\text{Gr}(k, n)]$, that is, they satisfy relations*

$$\begin{aligned} \sigma_i \sigma_j &= \sigma_j \sigma_i, & |i - j| > 1, \\ \sigma_i \sigma_j \sigma_i &= \sigma_j \sigma_i \sigma_j, & |i - j| = 1. \end{aligned}$$

Remark 3.16. In the case that $k = 2$, the Plücker relations in $\mathbb{C}[\text{Gr}(2, n)]$ are all 3-term relations. Therefore we have that Theorem 3.15 is true without the assumption that Conjecture 3.11 is true.

We first give some examples in the next two subsections and we prove Theorem 3.15 in Section 4.

3.4. Braid group action on $\mathbb{C}_q[\text{Gr}(2, 4)]$. By definition, σ_1 sends

$$\Delta_q^{12} \mapsto \Delta_q^{12}, \Delta_q^{13} \mapsto \Delta_q^{24}, \Delta_q^{14} \mapsto \Delta_q^{12} (\Delta_q^{14})^{-1} \Delta_q^{34}, \Delta_q^{23} \mapsto \Delta_q^{12} (\Delta_q^{23})^{-1} \Delta_q^{34}, \Delta_q^{34} \mapsto \Delta_q^{34}.$$

We have that

$$\begin{aligned} \Delta_q^{24} &= (\Delta_q^{13})' = (\Delta_q^{13})^{-1} \Delta_q^{12} \Delta_q^{34} + (\Delta_q^{13})^{-1} \Delta_q^{14} \Delta_q^{23} \\ &= \frac{1}{q} (\Delta_q^{13})^{-1} * \Delta_q^{12} * \Delta_q^{34} + q (\Delta_q^{13})^{-1} * \Delta_q^{14} * \Delta_q^{23}. \end{aligned}$$

By Lemma 3.7,

$$\begin{aligned} \sigma_1(\Delta_q^{24}) &= \sigma_1((\Delta_q^{13})^{-1}) \sigma_1(\Delta_q^{12}) \sigma_1(\Delta_q^{34}) + \sigma_1((\Delta_q^{13})^{-1}) \sigma_1(\Delta_q^{14}) \sigma_1(\Delta_q^{23}) \\ &= (\Delta_q^{24})^{-1} \Delta_q^{12} \Delta_q^{34} + (\Delta_q^{24})^{-1} \Delta_q^{12} (\Delta_q^{14})^{-1} \Delta_q^{34} \Delta_q^{12} (\Delta_q^{23})^{-1} \Delta_q^{34} \\ &= q^{-1} (\Delta_q^{24})^{-1} * \Delta_q^{12} * \Delta_q^{34} + q (\Delta_q^{24})^{-1} * \Delta_q^{12} * (\Delta_q^{14})^{-1} * \Delta_q^{34} * \Delta_q^{12} * (\Delta_q^{23})^{-1} * \Delta_q^{34}. \end{aligned}$$

We now verify that σ_1 preserve the Plücker relation:

$$\Delta_q^{13} * \Delta_q^{24} = q^{-1} \Delta_q^{12} * \Delta_q^{34} + q \Delta_q^{14} * \Delta_q^{23}.$$

On the left hand side,

$$\begin{aligned} \sigma_1(\Delta_q^{13}) * \sigma_1(\Delta_q^{24}) &= \Delta_q^{24} * (q^{-1} (\Delta_q^{24})^{-1} * \Delta_q^{12} * \Delta_q^{34} \\ &\quad + q (\Delta_q^{24})^{-1} * \Delta_q^{12} * (\Delta_q^{14})^{-1} * \Delta_q^{34} * \Delta_q^{12} * (\Delta_q^{23})^{-1} * \Delta_q^{34}) \\ &= q^{-1} \Delta_q^{12} * \Delta_q^{34} + q \Delta_q^{12} * (\Delta_q^{14})^{-1} * \Delta_q^{34} * \Delta_q^{12} * (\Delta_q^{23})^{-1} * \Delta_q^{34}. \end{aligned}$$

On the right hand side,

$$\begin{aligned} &q^{-1} \sigma_1(\Delta_q^{12}) * \sigma_1(\Delta_q^{34}) + q \sigma_1(\Delta_q^{14}) * \sigma_1(\Delta_q^{23}) \\ &= q^{-1} \Delta_q^{12} * \Delta_q^{34} + q \Delta_q^{12} (\Delta_q^{14})^{-1} \Delta_q^{34} * \Delta_q^{12} (\Delta_q^{23})^{-1} \Delta_q^{34} \\ &= q^{-1} \Delta_q^{12} * \Delta_q^{34} + q \Delta_q^{12} * (\Delta_q^{14})^{-1} * \Delta_q^{34} * \Delta_q^{12} * (\Delta_q^{23})^{-1} * \Delta_q^{34}. \end{aligned}$$

Therefore

$$\sigma_1(\Delta_q^{13}) * \sigma_1(\Delta_q^{24}) = q^{-1}\sigma_1(\Delta_q^{12}) * \sigma_1(\Delta_q^{34}) + q\sigma_1(\Delta_q^{14}) * \sigma_1(\Delta_q^{23}).$$

We now check that the relation $\Delta_q^{14} * \Delta_q^{23} = \Delta_q^{23} * \Delta_q^{14}$ is preserved by σ_1 . On the left hand side,

$$\begin{aligned} \sigma_1(\Delta_q^{14}) * \sigma_1(\Delta_q^{23}) &= (\Delta_q^{12}(\Delta_q^{14})^{-1}\Delta_q^{34}) * (\Delta_q^{12}(\Delta_q^{23})^{-1}\Delta_q^{34}) \\ &= \Delta_q^{12} * (\Delta_q^{14})^{-1} * \Delta_q^{34} * \Delta_q^{12} * (\Delta_q^{23})^{-1} * \Delta_q^{34} \\ &= \Delta_q^{12}(\Delta_q^{14})^{-1}\Delta_q^{34}\Delta_q^{12}(\Delta_q^{23})^{-1}\Delta_q^{34}, \end{aligned}$$

where the last expression is a commutative quantum cluster monomial. On the right hand side,

$$\begin{aligned} \sigma_1(\Delta_q^{23}) * \sigma_1(\Delta_q^{14}) &= (\Delta_q^{12}(\Delta_q^{23})^{-1}\Delta_q^{34}) * (\Delta_q^{12}(\Delta_q^{14})^{-1}\Delta_q^{34}) \\ &= \Delta_q^{12} * (\Delta_q^{23})^{-1} * \Delta_q^{34} * \Delta_q^{12} * (\Delta_q^{14})^{-1} * \Delta_q^{34} \\ &= \Delta_q^{12}(\Delta_q^{23})^{-1}\Delta_q^{34}\Delta_q^{12}(\Delta_q^{14})^{-1}\Delta_q^{34}, \end{aligned}$$

where the last expression is a commutative quantum cluster monomial. Therefore $\sigma_1(\Delta_q^{14}) * \sigma_1(\Delta_q^{23}) = \sigma_1(\Delta_q^{23}) * \sigma_1(\Delta_q^{14})$.

Remark 3.17. It is shown in [LL11, Section 3] that the cycling map is not an automorphism of the quantum Grassmannian $\mathbb{C}_q[\text{Gr}(k, n)]$.

In $\mathbb{C}_q[\text{Gr}(2, 4)]$, the map σ_1 is similar but different from the cycling map ρ but different. In $\mathbb{C}_q[\text{Gr}(2, 4)]$, the cycling map is given by $\Delta_q^{12} \mapsto \Delta_q^{23}$, $\Delta_q^{13} \mapsto \Delta_q^{24}$, $\Delta_q^{23} \mapsto \Delta_q^{34}$, $\Delta_q^{24} \mapsto \Delta_q^{13}$, $\Delta_q^{34} \mapsto \Delta_q^{14}$, $\Delta_q^{14} \mapsto \Delta_q^{12}$. We have that $\Delta_q^{23} * \Delta_q^{14} = \Delta_q^{14} * \Delta_q^{23}$. Applying the cycling map to $\Delta_q^{23}, \Delta_q^{14}$, we obtain $\Delta_q^{34}, \Delta_q^{12}$ respectively. Although $\Delta_q^{23}, \Delta_q^{14}$ commute, $\Delta_q^{34}, \Delta_q^{12}$ do not commute and we have $\Delta_q^{12} * \Delta_q^{34} = q^2\Delta_q^{34} * \Delta_q^{12}$. No matter how we choose $a_{ijj'j'}$ in the deformed cycling map $\Delta_q^{ij} \mapsto q^{a_{ijj'j'}}\Delta_q^{i'j'}$, the deformed cycling map always sends the relation $\Delta_q^{23} * \Delta_q^{14} = \Delta_q^{14} * \Delta_q^{23}$ to the relation $\Delta_q^{12} * \Delta_q^{34} = \Delta_q^{34} * \Delta_q^{12}$ which does not hold in $\mathbb{C}_q[\text{Gr}(2, 4)]$.

3.5. Braid group action on $\mathbb{C}_q[\text{Gr}(3, 6)]$.

We list the images of quantum cluster variables of $\mathbb{C}_q[\text{Gr}(3, 6)]$ under the maps σ_i , $i = 1, 2$, in Table 1.

We now check that quantum Plücker relations are preserved by σ_1, σ_2 . We check one example and others are similar. Consider the quantum Plücker relation

$$(3.12) \quad \Delta_q^{124} * \Delta_q^{136} = \frac{1}{q}\Delta_q^{123} * \Delta_q^{146} + q\Delta_q^{126} * \Delta_q^{134}.$$

We have that

$$\Delta_q^{136} = (\Delta_q^{124})^{-1}\Delta_q^{123}\Delta_q^{146} + (\Delta_q^{124})^{-1}\Delta_q^{126}\Delta_q^{134},$$

and

$$\begin{aligned} \sigma_1(\Delta_q^{136}) &= \sigma_1((\Delta_q^{124})^{-1})\sigma_1(\Delta_q^{123})\sigma_1(\Delta_q^{146}) + \sigma_1((\Delta_q^{124})^{-1})\sigma_1(\Delta_q^{126})\sigma_1(\Delta_q^{134}) \\ &= (\Delta_q^{125})^{-1}\Delta_q^{123}\Delta_q^{256} + (\Delta_q^{125})^{-1}\Delta_q^{126}\Delta_q^{235} \\ &= \frac{1}{q}(\Delta_q^{125})^{-1} * \Delta_q^{123} * \Delta_q^{256} + q(\Delta_q^{125})^{-1} * \Delta_q^{126} * \Delta_q^{235}. \end{aligned}$$

variable x	$\sigma_1(x)$	$\sigma_2(x)$
Δ_q^{123}	Δ_q^{123}	Δ_q^{123}
Δ_q^{234}	$\Delta_q^{123} (\Delta_q^{234})^{-1} \Delta_q^{345}$	Δ_q^{234}
Δ_q^{345}	Δ_q^{345}	$\Delta_q^{234} (\Delta_q^{345})^{-1} \Delta_q^{456}$
Δ_q^{456}	Δ_q^{456}	Δ_q^{456}
Δ_q^{126}	Δ_q^{126}	$\Delta_q^{123} (\Delta_q^{126})^{-1} \Delta_q^{156}$
Δ_q^{156}	$\Delta_q^{126} (\Delta_q^{156})^{-1} \Delta_q^{456}$	Δ_q^{156}
Δ_q^{124}	Δ_q^{125}	Δ_q^{134}
Δ_q^{125}	$\Delta_q^{126} (\Delta_q^{156})^{-1} \Delta_q^{145}$	Δ_q^{136}
Δ_q^{134}	Δ_q^{235}	$\Delta_q^{145} (\Delta_q^{345})^{-1} \Delta_q^{234}$
Δ_q^{135}	$(\Delta_q^{156})^{-1} z$	$(\Delta_q^{345})^{-1} z$
Δ_q^{136}	Δ_q^{236}	$\Delta_q^{123} \Delta_q^{156} \Delta_q^{245} (\Delta_q^{126})^{-1} (\Delta_q^{345})^{-1}$
Δ_q^{145}	Δ_q^{245}	Δ_q^{146}
Δ_q^{146}	Δ_q^{256}	$\Delta_q^{124} (\Delta_q^{126})^{-1} \Delta_q^{156}$
Δ_q^{235}	$\Delta_q^{123} \Delta_q^{146} (\Delta_q^{156})^{-1} (\Delta_q^{234})^{-1} \Delta_q^{345}$	Δ_q^{236}
Δ_q^{236}	$\Delta_q^{123} (\Delta_q^{234})^{-1} \Delta_q^{346}$	$\Delta_q^{123} (\Delta_q^{126})^{-1} \Delta_q^{256}$
Δ_q^{245}	$\Delta_q^{124} (\Delta_q^{234})^{-1} \Delta_q^{345}$	Δ_q^{346}
Δ_q^{246}	$(\Delta_q^{234})^{-1} y$	$(\Delta_q^{126})^{-1} y$
Δ_q^{256}	$\Delta_q^{126} \Delta_q^{134} \Delta_q^{456} (\Delta_q^{156})^{-1} (\Delta_q^{234})^{-1}$	Δ_q^{356}
Δ_q^{346}	Δ_q^{356}	$\Delta_q^{125} (\Delta_q^{126})^{-1} \Delta_q^{234} (\Delta_q^{345})^{-1} \Delta_q^{456}$
Δ_q^{356}	$\Delta_q^{136} (\Delta_q^{156})^{-1} \Delta_q^{456}$	$\Delta_q^{235} (\Delta_q^{345})^{-1} \Delta_q^{456}$
y	$\Delta_q^{126} \Delta_q^{135} (\Delta_q^{156})^{-1} \Delta_q^{456}$	$\Delta_q^{135} \Delta_q^{234} (\Delta_q^{345})^{-1} \Delta_q^{456}$
z	$\Delta_q^{123} (\Delta_q^{234})^{-1} \Delta_q^{246} \Delta_q^{345}$	$\Delta_q^{123} (\Delta_q^{126})^{-1} \Delta_q^{156} \Delta_q^{246}$

TABLE 1. Braid group action on quantum cluster variables in $\mathbb{C}_q[\text{Gr}(3, 6)]$. The expressions are commutative quantum cluster monomials, where $y = q^{-\frac{3}{2}}(\Delta_q^{124} * \Delta_q^{356} - \frac{1}{q} \Delta_q^{123} * \Delta_q^{456})$ and $z = q^{-\frac{1}{2}}(\Delta_q^{145} * \Delta_q^{236} - \frac{1}{q^2} \Delta_q^{123} * \Delta_q^{456})$.

On the left hand side,

$$\begin{aligned} \sigma_1(\Delta_q^{124}) * \sigma_1(\Delta_q^{136}) &= \Delta_q^{125} * \left(\frac{1}{q} (\Delta_q^{125})^{-1} * \Delta_q^{123} * \Delta_q^{256} + q (\Delta_q^{125})^{-1} * \Delta_q^{126} * \Delta_q^{235} \right) \\ &= \frac{1}{q} \Delta_q^{123} * \Delta_q^{256} + q \Delta_q^{126} * \Delta_q^{235}. \end{aligned}$$

On the right hand side,

$$\frac{1}{q} \sigma_1(\Delta_q^{123}) * \sigma_1(\Delta_q^{146}) + q \sigma_1(\Delta_q^{126}) * \sigma_1(\Delta_q^{134}) = \frac{1}{q} \Delta_q^{123} * \Delta_q^{256} + q \Delta_q^{126} * \Delta_q^{235}.$$

Therefore σ_1 preserves the Plücker relation (3.12).

We now verify that σ_1 and σ_2 satisfy the braid relation $\sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2$. Consider $\sigma_1 \sigma_2 \sigma_1(\Delta_q^{145})$ and $\sigma_2 \sigma_1 \sigma_2(\Delta_q^{145})$. We have that

$$\begin{aligned} \sigma_1 \sigma_2 \sigma_1(\Delta_q^{145}) &= \sigma_1 \sigma_2(\Delta_q^{245}) = \sigma_1(\Delta_q^{346}) = \Delta_q^{356}, \\ \sigma_2 \sigma_1 \sigma_2(\Delta_q^{145}) &= \sigma_2 \sigma_1(\Delta_q^{146}) = \sigma_2(\Delta_q^{256}) = \Delta_q^{356}. \end{aligned}$$

Similarly, we have $\sigma_1 \sigma_2 \sigma_1(x_i) = \sigma_2 \sigma_1 \sigma_2(x_i)$ for other quantum cluster variables x_i in the initial cluster.

4. PROOF OF THEOREM 3.15

In this section, we prove our main result Theorem 3.15.

4.1. Proof of (i) of Theorem 3.15. By Lemma 3.14, the maps σ_i and σ_i^{-1} are algebra automorphisms of the quantum cluster algebra $\mathbb{C}_q[\text{Gr}(k, n)]$.

By the definition of σ_i , the matrix associated to σ_i is $R = \begin{pmatrix} I_{n \times n} & 0 \\ 0 & L \end{pmatrix}$, where $L = (l_{jk})$ is an $n \times n$ matrix with entries

$$(4.1) \quad l_{jk} = \begin{cases} 1, & \text{if } j = l \not\equiv i + 1, j + 1 = l \equiv i + 1, \text{ or } j - 1 = l \equiv i + 1, \\ -1, & \text{if } j = l \equiv i + 1, \\ 0, & \text{otherwise.} \end{cases}$$

It is not hard to see that $L^{-1} = L$ and $R^{-1} = R$. Here we place the frozen variables, i.e., consecutive quantum Plücker coordinates, in the order of $[1, k], [2, k + 1], \dots, [n - k + 1, n], [n - k + 2, n + 1], \dots, [n, n + k - 1]$, where the addition is taken in \mathbb{Z}_n .

It is clear that by the definition, the maps σ_i and σ_i^{-1} satisfies the condition (2) in Proposition 2.8. On the other hand, the conditions (1) and (3) in Proposition 2.8 follow from [F17, Lemma 5.11]. So to prove Theorem 3.15 (i) we only need to check the condition (4) in Proposition 2.8, that is, $R^t \bar{\Lambda} R = \Lambda$. This follows from Lemma 3.6.

4.2. Proof of (ii) of Theorem 3.15. Recall that in Section 3.3, we defined a map φ sending a cluster monomial in $\mathbb{C}[\text{Gr}(k, n)]$ to a quantum cluster monomial in $\mathbb{C}_q[\text{Gr}(k, n)]$. Every quantum cluster variable in $\mathbb{C}_q[\text{Gr}(k, n)]$ is of the form $\varphi(x)$, where x is a cluster variable in $\mathbb{C}[\text{Gr}(k, n)]$. By Lemma 3.8, we have that for every i and every cluster monomial x (x can contain frozen variables or inverses of frozen variables) in $\mathbb{C}[\text{Gr}(k, n)]$, $\sigma_i(\varphi(x)) = \varphi(\sigma_i(x))$. In particular, for any Plücker coordinate P , we have that $\sigma_i(\varphi(P)) = \varphi(\sigma_i(P))$. It is shown in [F17] that for $|i - j| = 1$, $\sigma_i \sigma_j \sigma_i(P) = \sigma_j \sigma_i \sigma_j(P)$. Therefore

$$\sigma_i \sigma_j \sigma_i(\varphi(P)) = \sigma_i \sigma_j(\varphi(\sigma_i(P))) = \varphi(\sigma_i \sigma_j \sigma_i(P)) = \varphi(\sigma_j \sigma_i \sigma_j(P)) = \sigma_j \sigma_i \sigma_j(\varphi(P)).$$

To show that $\sigma_i \sigma_j \sigma_i = \sigma_j \sigma_i \sigma_j$ on $\mathbb{C}_q[\text{Gr}(k, n)]$, it suffices to shown that $\sigma_i \sigma_j \sigma_i(x_1 * \dots * x_r) = \sigma_j \sigma_i \sigma_j(x_1 * \dots * x_r)$, where x_i 's are quantum Plücker coordinates. This follows from the fact that σ_i, σ_j are algebra automorphisms on $\mathbb{C}_q[\text{Gr}(k, n)]$ and $\sigma_i \sigma_j \sigma_i(x_i) = \sigma_j \sigma_i \sigma_j(x_i)$ for quantum Plücker coordinates x_i .

The proof of $\sigma_i(\sigma_j(x)) = \sigma_j(\sigma_i(x))$ for every $x \in \mathbb{C}_q[\text{Gr}(k, n)]$, $|i - j| > 1$, is similar.

REFERENCES

- [ASS12] I. Assem, R. Schiffler, and V. Shramchenko, *Cluster automorphisms*, Proc. Lond. Math. Soc. (3) **104** (2012), no. 6, 1271–1302.
- [AG12] J. Allman and J. E. Grabowski, *A quantum analogue of the dihedral action on Grassmannians*, J. Algebra **359** (2012), 49–68.
- [BFZ96] A. Berenstein, S. Fomin, and A. Zelevinsky, *Parametrizations of canonical bases and totally positive matrices*, Adv. Math. **122** (1996), no. 1, 49–149.
- [BG02] K. A. Brown and K. R. Goodearl, *Lectures on algebraic quantum groups*, Advanced Courses in Mathematics, CRM Barcelona, Birkhäuser Verlag, Basel, 2002.
- [BS15] T. Bridgeland and I. Smith, *Quadratic differentials as stability conditions*, Publ. Math. Inst. Hautes Études Sci. **121** (2015), 155–278.
- [BZ05] A. Berenstein and A. Zelevinsky, *Quantum cluster algebras*, Adv. Math. **195** (2005), no. 2, 405–455.
- [CS19] W. Chang and R. Shiffler, *Cluster automorphisms and quasi-automorphisms*, Advances in Applied Mathematics, **110** (2019), 342–374.
- [CZ16] W. Chang and B. Zhu, *Cluster automorphism groups of cluster algebras with coefficients*, Sci. China Math. **59** (2016), no. 10, 1919–1936.
- [F16] C. Fraser, *Quasi-homomorphisms of cluster algebras*, Advances in Applied Mathematics, **81** (2016), 40–77.

- [F17] C. Fraser, *Braid group symmetries of Grassmannian cluster algebras*, Sel. Math. New Ser. **26** (2020), article number: 17.
- [FWZ] S. Fomin, L. Williams, and A. Zelevinsky, *Introduction to Cluster Algebras, Chapter 6*, preprint (2020), arXiv:2008.09189 [math.CO].
- [FZ02] S. Fomin, A. Zelevinsky, *Cluster algebras. I. Foundations*. Journal of the American Mathematical Society, 2002, 15(2), 497–529.
- [GL11] J. E. Grabowski and S. Launois, *Quantum cluster algebra structures on quantum Grassmannians and their quantum Schubert cells: the finite-type cases*, International Mathematics Research Notices, Volume **2011**, issue 10, 2011, 2230–2262.
- [GL14] J. E. Grabowski and S. Launois, *Graded quantum cluster algebras and an application to quantum Grassmannians*, Proc. Lond. Math. Soc., **109** (2014), no. 3, 697–732.
- [JKS22] B. T. Jensen, A. King, X. Su, *Categorification and the quantum Grassmannian*, Advances in Mathematics, Volume **406**, 2022, 108577.
- [KO19] Y. Kimura and H. Oya, *Twist automorphisms on quantum unipotent cells and dual canonical bases*, International Mathematics Research Notices, volume **2021**, issue 9, 2021, 6772–6847.
- [KQW22] Y. Kimura, F. Qin, and Q.l. Wei, *Twist automorphisms and Poisson structures*, arXiv:2201.10284.
- [Lau06] A. Lauve, *Quantum- and quasi-Plücker coordinates*, Journal of Algebra, **296** (2006), 440–461.
- [Lau10] A. Lauve, *Quasi-determinants and q -commuting minors*, Glasgow Math. J. **52** (2010), 663–675.
- [LL11] S. Launois and T. H. Lenagan, *Twisting the quantum Grassmannian*, Proc. Amer. Math. Soc. **139** (2011), no. 1, 99–110.
- [LZ98] B. Leclerc and A. Zelevinsky, *Quasicommuting families of quantum Plücker coordinates*, Kirillov’s seminar on representation theory, 85–108, Amer. Math. Soc. Transl. Ser. 2, **181**, Adv. Math. Sci., **35**, Amer. Math. Soc., Providence, RI, 1998.
- [M16] G. Muller, *Skein and cluster algebras of marked surfaces*, Quantum Topol. **7**, (2016), no. 3, 435–503.
- [MS16] B.R. Marsh and J.S. Scott, *Twists of Plücker coordinates as dimer partition functions*, Commun. Math. Phys. **341**, 821–884 (2016).
- [S05] J. Scott, *Quasi-commuting families of quantum minors*, Journal of Algebra, **290** (2005), 204–220.
- [Sco06] J. Scott, *Grassmannians and cluster algebras*, Proc. London Math. Soc. (3) **92** (2006), no. 2, 345–380.
- [TT91] E. Taft and J. Towber, *Quantum deformation of flag schemes and Grassmann schemes, I. A q -deformation of the shape-algebra for $GL(n)$* , J. Algebra **142** (1991), no. 1, 1–36.

WEN CHANG, SCHOOL OF MATHEMATICS AND STATISTICS, SHAANXI NORMAL UNIVERSITY, XI’AN, CHINA

Email address: changwen161@163.com

MIN HUANG, SCHOOL OF MATHEMATICS (ZHUHAI), SUN YAT-SEN UNIVERSITY, ZHUHAI, CHINA

Email address: huangm97@mail.sysu.edu.cn

JIAN-RONG LI, FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, OSKAR-MORGENSTERN-PLATZ 1, 1090 VIENNA, AUSTRIA

Email address: lijr07@gmail.com