

TORIC RINGS OF $(0, 1)$ -POLYTOPES WITH SMALL RANK

KOJI MATSUSHITA

ABSTRACT. The rank of a d -dimensional polytope P is defined by $F - (d + 1)$, where F denotes the number of facets of P . In this paper, We focus on the toric rings of $(0, 1)$ -polytopes with small rank. We study their normality, the torsionfreeness of their divisor class groups and the classification of their isomorphism classes.

1. INTRODUCTION

Throughout the present paper, let \mathbb{k} be an algebraically closed field of characteristic 0, for simplicity.

1.1. **Backgrounds.** For an integral polytope $P \subset \mathbb{R}^d$, let $\mathbb{k}[P]$ denote the toric ring of P , i.e.,

$$\mathbb{k}[P] = \mathbb{k}[t_1^{v_1} \cdots t_d^{v_d} t_0 : (v_1, \dots, v_d) \in P \cap \mathbb{Z}^d].$$

Toric rings of integral polytopes are homogenous affine semigroup rings and their algebraic properties have been well investigated. In particular, normality provides various important properties to toric rings and some algebraic objects can be described in the terms associated with polytopes. In fact, it is well known that normal toric rings are Cohen-Macaulay for any field \mathbb{k} ([18]) and that the canonical module of the normal toric ring is described as the ideal generated by the monomials corresponding to the elements in the interior of $\mathbb{Z}_{\geq 0}\mathcal{A}(P)$ ([28]), where $\mathcal{A}(P) = \{(\mathbf{v}, 1) \in \mathbb{Z}^{d+1} : \mathbf{v} \in P \cap \mathbb{Z}^d\}$ and $\mathbb{Z}_{\geq 0}\mathcal{A}(P) = \{a_1\mathbf{x}_1 + \cdots + a_n\mathbf{x}_n : \mathbf{x}_1, \dots, \mathbf{x}_n \in \mathcal{A}(P), a_1, \dots, a_n \in \mathbb{Z}_{\geq 0}\}$. Moreover, the toric ring $\mathbb{k}[P]$ is normal if and only if it is a Krull ring (cf. [33, Theorem 9.8.13]). In this case, the divisor class group of $\mathbb{k}[P]$ can be easily computed by using the generators and the support forms of $\mathbb{Z}_{\geq 0}\mathcal{A}(P)$ (see Section 2.1).

The divisor class group $\text{Cl}(\mathbb{k}[P])$ of a normal toric ring $\mathbb{k}[P]$, which is a finitely generated abelian group, is one of the most interesting invariants. Its elements are identified with the isomorphism classes of the divisorial ideals of $\mathbb{k}[P]$. Recently, conic divisorial ideals, which are a certain class of divisorial and a special kind of maximal Cohen-Macaulay modules of rank one, are well studied (see, e.g., [1, 2]). Indeed, they play beneficial roles in the theory of non-commutative algebraic geometry as well as commutative rings. For example, we can construct non-commutative crepant resolutions (NCCRs, for short), which were introduced by Van den Bergh ([32]), by considering the endomorphism ring of the direct sum of some conic divisorial ideals. In fact, they are constructed for many classes of toric rings (see, e.g., [15, 17, 21, 22, 27, 32], and so on). However, in general, the existence of NCCRs for Gorenstein toric rings is still open. In solving this problem, it is natural to consider the case of the toric rings whose divisor class groups have small rank. On the other hand, it was shown in [7] that if a Cohen-Macaulay normal domain R has an NCCR,

2020 *Mathematics Subject Classification.* Primary 13F65; Secondary 13C20, 52B20.

Key words and phrases. Toric rings, $(0, 1)$ -polytopes, Divisor class groups.

then R is \mathbb{Q} -Gorenstein. In particular, if the divisor class group of R is torsionfree and R is not Gorenstein, then R does not have an NCCR. Therefore, it is important to decide whether divisor class groups are torsionfree.

The convex hull of a finite set of $(0, 1)$ -vectors is called a $(0, 1)$ -polytope. It arises from various combinatorial objects such as partially ordered sets (posets, for short), graphs and matroids. For instance, the following are families of $(0, 1)$ -polytopes and their toric rings are well studied:

- Order polytopes (see [30] and Section 2.2);
- Stable set polytopes ([6]) \supset Chain polytopes ([30]);
- Edge polytopes (see e.g., [24, 26]);

Of course, these are not the only ones, there are much more; cut polytopes ([8]), perfect matching polytopes ([9]), matroid polytopes ([23]), and so on. In addition, compressed polytopes introduced by Stanley [29] are a natural class of integral polytopes and include some of the $(0, 1)$ -polytopes mentioned above. It is known that the toric ring of a compressed polytope is isomorphic to that of a $(0, 1)$ -polytope ([31, Theorem 2.4]).

By studying the commutative ring-theoretic properties of their toric rings, we can solve problems related to underlying combinatorial objects.

In the present paper, we focus on the *rank* of an integral polytope P , which is defined by

$$\text{rank } P = F - (\dim P + 1),$$

where F denotes the number of facets of P . Note that $\text{rank } P$ is a nonnegative integer. Actually, the rank of P coincide with that of $\text{Cl}(\mathbb{k}[P])$ if P is normal (see Section 2.1). Namely, we deal with toric rings whose divisor class groups have small rank.

The goal of the present paper is to analyze the normality and torsionfreeness of $(0, 1)$ -polytopes, and to classify the isomorphism classes of their toric rings in the case where the underlying polytopes have small rank.

1.2. Normality. We say that an integral polytope P is normal if so is $\mathbb{k}[P]$. The normality of an integral polytope P can be characterized in terms of the lattice points generated by $\mathcal{A}(P)$ (see Section 2.1). In general, it is not easy to determine if P is normal or not, but there is a useful sufficient condition for P to be normal; P is normal if P possesses a unimodular triangulation (cf. [13, Corollary 4.12]). This implies that order polytopes and stable set polytopes of perfect graphs including chain polytopes are normal since these are compressed polytopes (see [25]), which possess a unimodular triangulation. Meanwhile, it is known that the edge polytope of a graph G is normal if and only if G satisfies the odd cycle condition ([24, 26]).

As this shows, $(0, 1)$ -polytopes are not necessarily normal. However, non-normal $(0, 1)$ -polytopes seem to have large rank. In fact, we can observe that the rank of a non-normal edge polytope is at least 4 by [16, Proposition 4.4] and some arguments. Therefore, we are interested in the normality of integral polytopes with small rank and consider the following problem:

Problem 1. *Examine the normality of $(0, 1)$ -polytopes with small rank.*

1.3. Torsionfreeness. We say that a normal integral polytope P is torsionfree if so is $\text{Cl}(\mathbb{k}[P])$. The torsionfreeness of a normal integral polytope P can be determined by computing the Smith normal form of the matrix constructed from the lattice points in P and the facet defining inequalities of P (see Section 2.1). In [20], a sufficient condition for P to be torsionfree is given. As a corollary, it follows that compressed polytopes are

torsionfree ([20, Corollary 3.4]), and so are order polytopes and stable set polytopes of perfect graphs. Moreover, normal edge polytopes are also torsionfree ([16, Theorem 3.6]). Furthermore, recently, the divisor class groups of toric face rings, which can be regarded as the toric rings of $(0, 1)$ -polytopes arising from simplicial complexes, are investigated in [12]. It is shown that they are torsionfree if their underlying toric face rings are normal ([12, Corollary 1.8]).

These facts pose us the following problem:

Problem 2. *Are normal $(0, 1)$ -polytopes always torsionfree?*

1.4. Classification. The classification of the isomorphism classes and relationships among certain classes of the toric rings whose divisor class groups have small rank have been studied in [16] as follows. Let \mathbf{Order}_n (resp. \mathbf{Stab}_n , \mathbf{Edge}_n) denote the sets of isomorphic classes of the toric rings of order polytopes (resp. stable set polytopes of perfect graphs, normal edge polytopes) with rank n . Then, the following relationships hold:

- $\mathbf{Order}_n = \mathbf{Stab}_n = \mathbf{Edge}_n$ if $n \leq 1$.
- $\mathbf{Stab}_2 \cup \mathbf{Edge}_2 = \mathbf{Order}_2$ and no inclusion between \mathbf{Stab}_2 and \mathbf{Edge}_2 ;
- there is no inclusion among \mathbf{Order}_n , \mathbf{Stab}_n and \mathbf{Edge}_n if $n \geq 3$.

In particular, the relationship $\mathbf{Stab}_n \cup \mathbf{Edge}_n \subset \mathbf{Order}_n$ holds if $n \leq 2$.

Now, we consider a generalization of this result. Let $(\mathbf{0}, \mathbf{1})_n$ denote the sets of isomorphic classes of the toric rings of $(0, 1)$ -polytopes with rank n . Clearly, the relationship $\mathbf{Order}_n \cup \mathbf{Stab}_n \cup \mathbf{Edge}_n \subset (\mathbf{0}, \mathbf{1})_n$ holds. It follows from the third relationship of the above three families that one has $\mathbf{Order}_n \subsetneq (\mathbf{0}, \mathbf{1})_n$ if $n \geq 3$. On the other hand, in the case $n \leq 2$, the relationship between $(\mathbf{0}, \mathbf{1})_n$ and \mathbf{Order}_n is very close.

Problem 3. *Determine the set $(\mathbf{0}, \mathbf{1})_n$. Also, does the relationship $(\mathbf{0}, \mathbf{1})_n = \mathbf{Order}_n$ hold if $n \leq 2$?*

In classifying isomorphism classes of the toric rings of $(0, 1)$ -polytopes, to consider Gale-diagrams, which is an effective way to investigate the combinatorial types of polytopes with “few vertices”, is compatible with our study. In this paper, we treat polytopes with small rank, i.e., “few facets”, so we can apply the method of Gale-diagrams by considering the dual polytopes. We present the following problem:

Problem 4. *Does the combinatorial equivalence of two $(0, 1)$ -polytopes imply the isomorphism of their toric rings?*

1.5. Results. Let P be a $(0, 1)$ -polytope. We give complete or partial answers to Problems 1, 2, 3 and 4 in each case;

$$(r1) \quad \text{rank } P = 0 \text{ or } 1, \quad (r2) \quad \text{rank } P = 2, \quad (r3) \quad \text{rank } P \geq 3.$$

First, we provide the complete answers to the above problems in case (r1). Actually, most of this result has been already given in [20]:

Theorem 1.1 (Proposition 4.1 and Theorem 4.3). *Let P be a $(0, 1)$ -polytope with $\text{rank } P \leq 1$. Then, P is normal and torsionfree. Moreover, the following relationships hold:*

$$(\mathbf{0}, \mathbf{1})_0 = \mathbf{Order}_0 = \{\mathbb{k}[x_1, \dots, x_n] : n \in \mathbb{Z}_{>0}\} \text{ and}$$

$(\mathbf{0}, \mathbf{1})_1 = \mathbf{Order}_1 = \{(\mathbb{k}[x_1, \dots, x_{n+1}] \# \mathbb{k}[y_1, \dots, y_{m+1}]) \otimes_{\mathbb{k}} \mathbb{k}[z_1, \dots, z_{l-1}] : n, m, l \in \mathbb{Z}_{>0}\}$.
Furthermore, for two $(0, 1)$ -polytopes P_1 and P_2 with $\text{rank } P_i \leq 1$, these polytopes have the same combinatorial type if and only if $\mathbb{k}[P_1] \cong \mathbb{k}[P_2]$.

Next, we discuss case (r2). Before that, we need to introduce a new family of $(0, 1)$ -polytopes P_{n_1, \dots, n_k} .

Let \mathcal{B}_n denote the standard basis of \mathbb{R}^n . For $n_1, \dots, n_k \in \mathbb{Z}_{>0}$, the $(0, 1)$ -polytope P_{n_1, \dots, n_k} is defined as the convex hull of the origin, \mathcal{B}_d , where $d := n_1 + \dots + n_k$, and the product $\mathcal{B}_{n_1} \times \dots \times \mathcal{B}_{n_k}$.

In Section 3, we study several properties of P_{n_1, \dots, n_k} . Specifically, we describe the facet defining inequalities of P_{n_1, \dots, n_k} (Proposition 3.2). Moreover, we show that $\mathbb{k}[P_{n_1, \dots, n_k}]$ possesses a squarefree initial ideal (Proposition 3.4). Furthermore, the following facts hold (Theorem 3.6):

- P_{n_1, \dots, n_k} has the integer decomposition property, and hence it is normal.
- $\mathbb{k}[P_{n_1, \dots, n_k}]$ is Gorenstein if and only if $n_1 = n_2 = \dots = n_k$.
- $\text{Cl}(\mathbb{k}[P_{n_1, \dots, n_k}]) \cong \mathbb{Z}^{k-1}$.
- If $k \geq 3$, then $\mathbb{k}[P_{n_1, \dots, n_k}] \notin \mathbf{Order}_{k-1}$ for any $n_1, \dots, n_k \in \mathbb{Z}_{>0}$.

We provide various observations on Problems 3 and 4 in case (r2):

- The above facts imply that $\text{rank } P_{n_1, n_2, n_3} = 2$ and $\mathbb{k}[P_{n_1, n_2, n_3}] \notin \mathbf{Order}_2$ for any $n_1, n_2, n_3 \in \mathbb{Z}_{>0}$, this is a counterexample to the equality $(\mathbf{0}, \mathbf{1})_2 = \mathbf{Order}_2$ of Problem 3.
- We study the combinatorial types of the dual polytopes of $\mathcal{O}_{\Pi_1}, \dots, \mathcal{O}_{\Pi_4}$ (these are the order polytopes with rank 2) and P_{n_1, n_2, n_3} by drawing their standard Gale-diagrams (see Figures 5, 6, 7, 8 and 9). In particular, Problem 4 has positive answers for $(0, 1)$ -polytopes that the standard Gale-diagrams of their dual polytopes coincide with Gale_1 or Gale_2 . (Theorems 4.4 and 4.5)

Finally, we give answers to the above problems in case (r3), although they are negative:

Theorem 1.2 (Propositions 4.6, 4.7 and 4.8). *For any positive integer $r \geq 3$, there exist non-normal $(0, 1)$ -polytope P and non-torsionfree normal $(0, 1)$ -polytope P' with $\text{rank } P = \text{rank } P' = r$, respectively. Moreover, there exist two $(0, 1)$ -polytopes Q and Q' with the same combinatorial type and $\text{rank } Q = \text{rank } Q' = r$ such that their toric rings are not isomorphic to each other.*

1.6. Organization. In Section 2, we recall the definitions and notation of toric rings of integral polytopes and give an algorithm to compute divisor class groups of toric rings of normal integral polytopes. We also recall the toric rings of certain $(0, 1)$ -polytopes, especially the order polytopes, and recall its several properties. Moreover, we recall the notion of Gale-diagrams and provide a method to obtain the Gale-diagram of the dual polytope of a given polytope. In Section 3, we introduce a new family of $(0, 1)$ -polytopes and study its properties. In Section 4, we give complete or partial answers to Problems 1, 2, 3 and 4 in each case; $\text{rank } P = 0$ or 1, $\text{rank } P = 2$, and $\text{rank } P \geq 3$.

Acknowledgement. The author would like to thank Akihiro Higashitani for a lot of his helpful comments and instructive discussions. The author is partially supported by Grant-in-Aid for JSPS Fellows Grant JP22J20033.

2. PRELIMINARIES

2.1. Toric rings of integral polytopes and their divisor class groups. In this subsection, we recall the definitions and notation of polytopes and toric rings. We refer the readers to e.g., [3] or [33], for the introduction.

First, we introduce toric rings of integral polytopes. An *integral polytope* $P \subset \mathbb{R}^d$ is a polytope whose vertices sit in \mathbb{Z}^d . For an integral polytope $P \subset \mathbb{R}^d$, we define ϕ_P as the morphism of \mathbb{k} -algebras:

$$\phi_P : \mathbb{k}[x_{\mathbf{v}} : \mathbf{v} \in P \cap \mathbb{Z}^d] \rightarrow \mathbb{k}[t_0, t_1^{\pm 1}, \dots, t_d^{\pm 1}], \text{ induced by } \phi_P(x_{\mathbf{v}}) = \mathbf{t}^{\mathbf{v}} t_0,$$

where $\mathbf{t}^{\mathbf{v}} = t_1^{v_1} \cdots t_d^{v_d}$ for $\mathbf{v} = (v_1, \dots, v_d) \in \mathbb{Z}^d$. Then, the kernel of ϕ_P , denoted by I_P , is called the *toric ideal* of P . Moreover, the image of ϕ_P , denoted by $\mathbb{k}[P]$, is called the *toric ring* of P . Note that $\mathbb{k}[P] \cong \mathbb{k}[x_{\mathbf{v}} : \mathbf{v} \in P \cap \mathbb{Z}^d]/I_P$.

It is well known that the toric ideal I_P is generated by homogeneous binomials. The toric ring $\mathbb{k}[P]$ is a standard graded \mathbb{k} -subalgebra of $\mathbb{k}[t_0, t_1^{\pm 1}, \dots, t_d^{\pm 1}]$ by setting $\deg(\mathbf{t}^{\mathbf{v}} t_0) = 1$ for each $\mathbf{v} \in P \cap \mathbb{Z}^d$. The Krull dimension of $\mathbb{k}[P]$ is equal to the dimension of P plus 1.

For an integral polytope $P \in \mathbb{R}^d$, let $\mathcal{A}(P) = \{(\mathbf{v}, 1) \in \mathbb{Z}^{d+1} : \mathbf{v} \in P \cap \mathbb{Z}^d\}$ and let $\mathbb{Z}_{\geq 0}\mathcal{A}(P) = \{a_1 \mathbf{x}_1 + \cdots + a_n \mathbf{x}_n : \mathbf{x}_1, \dots, \mathbf{x}_n \in \mathcal{A}(P), a_1, \dots, a_n \in \mathbb{Z}_{\geq 0}\}$ (we define $\mathbb{Z}\mathcal{A}(P)$ and $\mathbb{R}_{\geq 0}\mathcal{A}(P)$ analogously).

Then, the following facts hold:

- P is normal (i.e., $\mathbb{k}[P]$ is normal) if and only if $\mathbb{Z}_{\geq 0}\mathcal{A}(P) = \mathbb{Z}\mathcal{A}(P) \cap \mathbb{R}_{\geq 0}\mathcal{A}(P)$ (cf. [33, Theorem 9.8.13]).
- Suppose that $\mathbb{Z}\mathcal{A}(P) = \mathbb{Z}^{d+1}$. Then, P is normal if and only if P has the *integer decomposition property* (IDP, for short), that is, for any $n \in \mathbb{Z}_{>0}$ and any $\mathbf{a} \in nP \cap \mathbb{Z}^d$, there exist $\mathbf{a}_1, \dots, \mathbf{a}_n \in P \cap \mathbb{Z}^d$ such that $\mathbf{a} = \mathbf{a}_1 + \cdots + \mathbf{a}_n$ (cf. [13, Theorem 4.7]).

We also define the *product* of two polytopes $P \subset \mathbb{R}^d$ and $Q \subset \mathbb{R}^e$ as

$$P \times Q = \{(\mathbf{x}, \mathbf{y}) : \mathbf{x} \in P, \mathbf{y} \in Q\} \subset \mathbb{R}^{d+e}.$$

We can see that $P \times Q$ is a polytope of dimension $\dim(P) + \dim(Q)$, whose nonempty faces are the products of nonempty faces (including itself) of P and Q . In particular, the number of facets of $P \times Q$ is equal to $|\Psi(P)| + |\Psi(Q)|$, where $\Psi(P)$ denotes the set of facets of P . Therefore, we have $\text{rank } P \times Q = \text{rank } P + \text{rank } Q + 1$.

The toric ring of the product of two integral polytopes corresponds to the ‘‘Segre product’’ of these toric rings; let P_1 and P_2 be two integral polytopes, then $\mathbb{k}[P_1 \times P_2]$ is isomorphic to the Segre product of $\mathbb{k}[P_1]$ and $\mathbb{k}[P_2]$. Here, for two standard \mathbb{k} -algebras $R = \bigoplus_{n \geq 0} R_n$ and $S = \bigoplus_{n \geq 0} S_n$, we define their *Segre product* $R \# S$ as the graded \mathbb{k} -algebra:

$$R \# S = (R_0 \otimes_{\mathbb{k}} S_0) \oplus (R_1 \otimes_{\mathbb{k}} S_1) \oplus \cdots \subset R \otimes_{\mathbb{k}} S.$$

Next, we give an algorithm to compute divisor class groups of toric rings of normal integral polytopes. We use theories in [33, Section 9.8].

Let $\langle \cdot, \cdot \rangle$ denote the natural inner product of \mathbb{R}^d . For $\mathbf{a} \in \mathbb{R}^d$ and $b \in \mathbb{R}$, we denote by $H^+(\mathbf{a}; b)$ (resp. $H(\mathbf{a}; b)$) a closed half-space $\{\mathbf{u} \in \mathbb{R}^d : \langle \mathbf{u}, \mathbf{a} \rangle + b \geq 0\}$ (resp. an affine hyperplane $\{\mathbf{u} \in \mathbb{R}^d : \langle \mathbf{u}, \mathbf{a} \rangle + b = 0\}$). In particular, we denote the linear hyperplane $H(\mathbf{a}; 0)$ by $H_{\mathbf{a}}$.

For each $F \in \Psi(P)$, there exist a vector $\mathbf{a}_F \in \mathbb{Q}^d$ and a rational number b_F with the following conditions:

- (d1) $H(\mathbf{a}_F; b_F)$ is a support hyperplane associated with F and $P \subset H^+(\mathbf{a}_F; b_F)$;
- (d2) $d_F(\mathbf{v}) \in \mathbb{Z}$ for any $\mathbf{v} \in P \cap \mathbb{Z}^d$;
- (d3) $\sum_{\mathbf{v} \in P \cap \mathbb{Z}^d} d_F(\mathbf{v}) \mathbb{Z} = \mathbb{Z}$,

where $d_F(\mathbf{v}) = \langle \mathbf{v}, \mathbf{a}_F \rangle + b_F$ for $\mathbf{v} \in \mathbb{Z}^d$. We can see that $d_F(\mathbf{v})$ for $\mathbf{v} \in P \cap \mathbb{Z}^d$ is independent of the choice of \mathbf{a}_F and b_F .

In this situation, P has the irreducible representation:

$$\mathbb{R}_{\geq 0}\mathcal{A}(P) = \text{aff}(\mathcal{A}(P)) \cap \left(\bigcap_{F \in \Psi(P)} H_{\mathbf{c}_F}^+ \right), \quad (2.1)$$

where $\mathbf{c}_F := (\mathbf{a}_F, b_F) \in \mathbb{Q}^{d+1}$. Given $v \in P \cap \mathbb{Z}^d$, we define \mathbf{w}_v belonging to a free abelian group $\mathcal{F} = \bigoplus_{F \in \Psi(P)} \mathbb{Z}\epsilon_F$ with its basis $\{\epsilon_F\}_{F \in \Psi(P)}$ as follows:

$$\mathbf{w}_v = \sum_{F \in \Psi(P)} \langle (v, 1), \mathbf{c}_F \rangle \epsilon_F = \sum_{F \in \Psi(P)} d_F(v) \epsilon_F.$$

Let

$$\mathcal{S} = \sum_{\mathbf{v} \in P \cap \mathbb{Z}^d} \mathbb{Z}\mathbf{w}_v = \left\{ \sum_{F \in \Psi(P)} \langle \mathbf{v}', \mathbf{c}_F \rangle \epsilon_F : \mathbf{v}' \in \mathbb{Z}\mathcal{A}(P) \right\} \quad (2.2)$$

and let \mathcal{M}_P be the matrix whose column vectors consist of \mathbf{w}_v for $\mathbf{v} \in P \cap \mathbb{Z}^d$, that is, $\mathcal{M}_P = (d_F(\mathbf{v}))_{F \in \Psi(P), \mathbf{v} \in P \cap \mathbb{Z}^d}$. Then we can compute the divisor class group of $\mathbb{k}[P]$ as follows:

Theorem 2.1 (cf. [33, Theorem 9.8.19]). *Work with the same notation as above and suppose that P is normal. Then, we have*

$$\text{Cl}(\mathbb{k}[P]) \cong \mathcal{F}/\mathcal{S}.$$

In particular, we have

$$\text{Cl}(\mathbb{k}[P]) \cong \mathbb{Z}^r \oplus \mathbb{Z}/s_1\mathbb{Z} \oplus \cdots \oplus \mathbb{Z}/s_m\mathbb{Z},$$

where $m = \dim P + 1$, $r = |\Psi(P)| - m$ and s_1, \dots, s_m are positive integers appearing in the diagonal of the Smith normal form of \mathcal{M}_P .

The integers s_1, \dots, s_m are called the *invariant factors* of \mathcal{M}_P .

Let $P \subset \mathbb{R}^d$ be an integral polytope and let k_P be a maximal nonnegative integer satisfying the following statement:

- (*) There exist distinct integral points $\mathbf{v}_1, \dots, \mathbf{v}_{k_P} \in P \cap \mathbb{Z}^d$ and distinct facets F_1, \dots, F_{k_P} of P such that $v_i \in \bigcap_{l=1}^{i-1} F_l$ for each $1 < i \leq k_P$ and $d_{F_i}(\mathbf{v}_i) = 1$ for each $1 \leq i \leq k_P$.

Theorem 2.2 ([20, Theorem 3.2]). *Let $P \subset \mathbb{R}^d$ be a normal integral polytope and let s_1, \dots, s_m be the invariant factors of \mathcal{M}_P . Then, $s_1 = \cdots = s_{k_P} = 1$. In particular, $\text{Cl}(\mathbb{k}[P])$ is torsionfree if $k_P = \dim P + 1$.*

Finally, we explain that normal toric rings whose divisor class groups are torsionfree can be described as the ring of invariants under an action of torus on polynomial rings. Suppose that $P \subset \mathbb{R}^d$ is a d -dimension normal integral polytope and satisfies $\mathbb{Z}\mathcal{A}(P) = \mathbb{Z}^{d+1}$. Moreover, we assume that $\text{Cl}(\mathbb{k}[P]) \cong \mathbb{Z}^r$. We fix an isomorphism $\iota : \mathcal{F}/\mathcal{S} \rightarrow \mathbb{Z}^r$ and let $\beta_F := \iota(\epsilon_F)$ for each $F \in \Psi(P)$. In this paper, we call β_F 's *weights* of $\mathbb{k}[P]$. We consider the action of the algebraic torus $G = (\mathbb{k}^\times)^r$ on the polynomial ring $S = \mathbb{k}[x_F : F \in \Psi(P)]$, which is the action induced by $g \cdot x_F = \beta_F(g)x_F$ for $g \in G$. Then, we have $R = S^G$ (see e.g., [2, Theorem 2.1]).

2.2. Toric rings of certain $(0, 1)$ -polytopes. In this subsection, we introduce some properties associated with toric rings of certain $(0, 1)$ -polytopes.

First, we introduce certain polytopes and propositions associated with their toric rings.

- A polytopes P is called a $(0, 1)$ -polytope if its all vertices are $(0, 1)$ -vectors.
- A pyramid $P \subset \mathbb{R}^d$ is the convex hull of the union of a polytope $Q \subset \mathbb{R}^d$ (basis of P) and a point $v_0 \in \mathbb{R}^d$ (apex of P), where v_0 does not belong to $\text{aff}(Q)$. Note that the basis of a pyramid P is a facet of P .
- A polytope $P \subset \mathbb{R}^d$ is *simple* if each vertex of P is contained in precisely $\dim P$ facets.

Proposition 2.3 ([19, Theorem 1]). *A $(0, 1)$ -polytope $P \subset \mathbb{R}^d$ is simple if and only if it is equal to a product of $(0, 1)$ -simplices.*

Proposition 2.4. *Let $P \subset \mathbb{R}^d$ be a $(0, 1)$ -pyramid with basis Q . Then, $\mathbb{k}[P]$ is the polynomial extension of $\mathbb{k}[Q]$. In particular, if P is normal, then so is Q and we have*

$$\text{Cl}(\mathbb{k}[P]) \cong \text{Cl}(\mathbb{k}[Q]).$$

Next, we introduce the certain class of $(0, 1)$ -polytopes, which are arising from posets. These are called order polytopes ([30]) and play an important role in this paper.

Let Π be a finite poset equipped with a partial order \preceq . For a subset $I \subset \Pi$, we say that I is a *poset ideal* of Π if $p \in I$ and $q \preceq p$ then $q \in I$. Note that \emptyset is regarded as a poset ideal. Let $\hat{\Pi} = \Pi \sqcup \{\hat{0}, \hat{1}\}$, where $\hat{0}$ (resp. $\hat{1}$) is a new minimal (resp. maximal) element not belonging to Π .

For a poset $\Pi = \{p_1, \dots, p_d\}$, let

$$\mathcal{O}_\Pi = \{(x_1, \dots, x_d) \in \mathbb{R}^d : x_i \geq x_j \text{ if } p_i \preceq p_j \text{ in } \Pi, 0 \leq x_i \leq 1 \text{ for } i = 1, \dots, d\}.$$

A convex polytope \mathcal{O}_Π is called the *order polytope* of Π . It is known that \mathcal{O}_Π is a $(0, 1)$ -polytope and has IDP. According to [30], the vertices of \mathcal{O}_Π one-to-one correspond to the poset ideals of Π . In fact, a $(0, 1)$ -vector (a_1, \dots, a_d) is a vertex of \mathcal{O}_Π if and only if $\{p_i \in \Pi : a_i = 1\}$ is a poset ideal. Moreover, the facets of \mathcal{O}_Π one-to-one correspond to the edges of the Hasse diagram of $\hat{\Pi}$. In fact, for $p_i, p_j \in \Pi$ with $p_i \prec p_j$, $\{p_i, p_j\}$ is an edge of the Hasse diagram of $\hat{\Pi}$ if and only if the inequality $x_i \geq x_j$ defines a facet of \mathcal{O}_Π , where we let $x_i = 1$ (resp. $x_j = 0$) if $p_i = \hat{0}$ (resp. $p_j = \hat{1}$).

The toric ring $\mathbb{k}[\mathcal{O}_\Pi]$ is called the *Hibi ring* of Π , which was originally introduced in [14]. We denote the Hibi ring of Π by $\mathbb{k}[\Pi]$ (instead of $\mathbb{k}[\mathcal{O}_\Pi]$ for short).

Let P and Q be two posets with $P \cap Q = \emptyset$. The *disjoint union* of P and Q is the poset $P + Q$ on $P \cup Q$ such that $x \preceq y$ in $P + Q$ if (a) $x, y \in P$ and $x \preceq y$ in P , or (b) $x, y \in Q$ and $x \preceq y$ in Q . The *ordinal sum* of P and Q is the poset $P \oplus Q$ on $P \cup Q \cup \{z\}$ such that $x \preceq y$ in $P \oplus Q$ if (a) $x, y \in P$ and $x \preceq y$ in P , (b) $x, y \in Q$ and $x \preceq y$ in Q , (c) $x \in P$ and $y = z$, or (d) $x = z$ and $y \in Q$, where z is a new element which is not contained in $P \cup Q$. By observing poset ideals of $P + Q$ and $P \oplus Q$, the following proposition holds:

Proposition 2.5. *Let P and Q be two posets with $P \cap Q = \emptyset$.*

- One has $\mathbb{k}[P + Q] \cong \mathbb{k}[P] \# \mathbb{k}[Q]$.*
- One has $\mathbb{k}[P \oplus Q] \cong \mathbb{k}[Q \oplus P] \cong \mathbb{k}[P] \otimes_{\mathbb{k}} \mathbb{k}[Q]$.*

We provide the description of the divisor class groups of Hibi rings, which is given in [11]. Let n be the number of edges of the Hasse diagram of $\widehat{\Pi}$. Then, it is proved in [11] that

$$\text{Cl}(\mathbb{k}[\Pi]) \cong \mathbb{Z}^{n-d-1}. \quad (2.3)$$

In particular, the rank of \mathcal{O}_{Π} is equal to $n - (d + 1)$.

The characterization of posets Π with $\text{rank } \mathcal{O}_{\Pi} \leq 2$ has been investigated in [22, 16]:

Theorem 2.6 ([22, Lemma 3.2] and [16, Proposition 4.1]). *Let Π be a poset. Then,*

(i) *rank $\mathcal{O}_{\Pi} = 0$ if and only if (the Hasse diagram of) Π is a chain.*

Moreover, assume that $\mathbb{k}[\Pi]$ is not a polynomial extension of a Hibi ring. Then,

(ii) *rank $\mathcal{O}_{\Pi} = 1$ if and only if Π is the disjoint union of two chains.*

(iii) *rank $\mathcal{O}_{\Pi} = 2$ if and only if Π is one of the posets Π_1, Π_2, Π_3 or Π_4 (see Figures 1, 2, 3 and 4).*




FIGURE 1. The poset $\widehat{\Pi}_1$




FIGURE 2. The poset $\widehat{\Pi}_2$




FIGURE 3. The poset $\widehat{\Pi}_3$




FIGURE 4. The poset $\widehat{\Pi}_4$

Here, n_1, n_2, n_3, m_1 and m_2 are the number of edges. Note that the Hibi ring of \mathcal{O}_{Π} with $\text{rank } \mathcal{O}_{\Pi} = 0$ is isomorphic to a polynomial ring. Moreover, it follows from Proposition 2.5 (i) that the Hibi ring of \mathcal{O}_{Π} with $\text{rank } \mathcal{O}_{\Pi} = 1$ is isomorphic to the Segre product of two polynomial rings.

Weights of the Hibi rings whose divisor class groups have rank 2 are computed in [22, Sections 3.2 and 3.3]. The following table summarizes the weights of the Hibi rings associated with the posets in Figures 1, 2, 3 and 4.

Poset	Π_1	Π_2	Π_3	Π_4
Weights	$(1, 0) \times n_1$	$(1, 0) \times n_1$	$(1, 0) \times n_1$	$(1, 0) \times n_1$
	$(0, 1) \times n_2$	$(-1, 0) \times n_2$	$(-1, -1) \times n_2$	$(-1, 0) \times n_2$
	$(-1, -1) \times n_3$	$(0, 1) \times m_1$	$(0, 1) \times n_3$	$(-1, 1) \times m_1$
		$(0, -1) \times m_2$	$(0, -1) \times m_1$	$(0, 1) \times l_1$
				$(0, -1) \times l_2$

Here $\times n_i$, $\times m_i$ and $\times l_i$ stand for the multiplicities.

2.3. Gale-diagrams. In this subsection, we recall the notation of Gale-diagrams of a polytope, which helps us to classify the isomorphic classes of toric rings.

Throughout this subsection, let $P \subset \mathbb{R}^d$ be a d -dimensional polytope with the vertex set $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ and suppose that P has the irreducible representation (2.1).

We consider

$$D(P) = \left\{ (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n : \sum_{i \in [n]} \alpha_i \mathbf{v}_i = \mathbf{0} \text{ and } \sum_{i \in [n]} \alpha_i = 0 \right\}.$$

We can see that $D(P)$ is a $(n - d - 1)$ -dimensional subspace of \mathbb{R}^d . Let $\mathbf{b}_1, \dots, \mathbf{b}_{n-d-1}$ be a basis of $D(P)$. Moreover, for $i \in [n]$, let $\bar{\mathbf{b}}_i$ denote the i -th column vector of the

$(n - d - 1) \times n$ matrix $\begin{pmatrix} \mathbf{b}_1 \\ \vdots \\ \mathbf{b}_{n-d-1} \end{pmatrix}$, that is, $\bar{\mathbf{b}}_i = (\mathbf{b}_1^{(i)}, \dots, \mathbf{b}_{n-d-1}^{(i)})$, where for a vector

$\mathbf{v} \in \mathbb{R}^n$, $\mathbf{v}^{(i)}$ denotes the i -th coordinate of \mathbf{v} .

Then, the n -tuple $(\bar{\mathbf{b}}_1, \dots, \bar{\mathbf{b}}_n)$ is called the *Gale-transform* of $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ (or of P). Furthermore, let

$$\widehat{\mathbf{b}}_i = \begin{cases} \bar{\mathbf{b}}_i / \|\bar{\mathbf{b}}_i\| & \text{if } \bar{\mathbf{b}}_i \neq \mathbf{0}, \\ \mathbf{0} & \text{if } \bar{\mathbf{b}}_i = \mathbf{0} \end{cases}$$

for each $i \in [n]$, where $\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$ for a vector $\mathbf{v} \in \mathbb{R}^n$. Then, the n -tuple $(\widehat{\mathbf{b}}_1, \dots, \widehat{\mathbf{b}}_n)$ is called the *Gale-diagram* of $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ (or of P). Gale-transforms and Gale-diagrams depend on the choice of the basis of $D(P)$.

Especially, we are interested in the case $n - d - 1 = 2$, that is, P just has $d + 3$ vertices. In this case, we can draw a *standard* (or *reduced*) Gale-diagram from the Gale-diagram $(\widehat{\mathbf{b}}_1, \dots, \widehat{\mathbf{b}}_n)$, which consists of the unit circle centered at the origin of \mathbb{R}^2 and diameters having at least one endpoint with multiplicity. See [10, Section 6.3] for the precise way to draw it.

We say that two polytopes Q and Q' are *combinatorially equivalent* or of the same *combinatorial type* (resp. *dual* to each other) if there exists a one-to-one mapping Φ between the set of all faces of Q and the set of all faces of Q' such that Φ is inclusion-preserving (resp. inclusion-reversing). Note that the classes of simplicial polytopes, which are polytopes whose facets are simplices, and simple polytopes are dual to each other.

According to [10, Sections 5.4 and 6.3], the following facts are known:

- Two d -dimensional polytopes with $d + 3$ vertices are combinatorially equivalent if and only if their standard Gale-diagrams are orthogonally equivalent (i.e., isomorphic under an orthogonal linear transformation of \mathbb{R}^2 onto itself).
- P is simplicial if and only if no diameter of the standard Gale-diagram has both endpoints.

Clearly, all polytopes which are dual to P have the same combinatorial type. Whenever we consider dual polytopes, we focus on their combinatorial type, so we call them *the* dual polytope of P .

For the remainder of this subsection, we will explain how to obtain a Gale-diagram of the dual polytope P . The *polar* P° of P is defined as

$$P^\circ = \{\mathbf{x} \in \mathbb{R}^d : \langle \mathbf{x}, \mathbf{y} \rangle \geq -1 \text{ for any } \mathbf{y} \in P\}.$$

It is known that if the origin is an interior point of P , then P° is a polytope and is dual to P . Moreover, $\{\mathbf{a}_F/b_F : F \in \Psi(P)\}$ is the vertex set of P° .

The following theorem means that we can get a Gale-diagram of the dual polytope of P from weights of its toric ring.

Theorem 2.7. *Suppose that P is normal, $\mathbb{Z}\mathcal{A}(P) = \mathbb{Z}^{d+1}$ and $\text{Cl}(\mathbb{k}[P]) \cong \mathbb{Z}^r$. Let $\{\beta_F\}_{F \in \Psi(P)}$ be weights of $\mathbb{k}[P]$. Then, $(\beta_F/\|\beta_F\|)_{F \in \Psi(P)}$ is a Gale-diagram of the dual polytope of P .*

Before we give the proof, we prepare the following lemma:

Lemma 2.8. *Work with the same assumption and notation as Theorem 2.7. Then, we have $\sum_{F \in \Psi(P)} \beta_F^{(j)} \mathbf{a}_F = \mathbf{0}$ and $\sum_{F \in \Psi(P)} \beta_F^{(j)} b_F = 0$ for each $j \in [r]$.*

Proof. Let \mathbf{e}_i be the i -th unit vector of \mathbb{Z}^{d+1} . Since $\mathbf{e}_i \in \mathbb{Z}\mathcal{A}(P)$ for each $i \in [d + 1]$, $\sum_{F \in \Psi(P)} \langle \mathbf{e}_i, \mathbf{c}_F \rangle \epsilon_F$ belongs to \mathcal{S} from (2.2). Thus, by considering its image in \mathbb{Z}^r , we can obtain that $\sum_{F \in \Psi(P)} \mathbf{c}_F^{(i)} \beta_F = \mathbf{0}$ for each $i \in [d + 1]$. This fact is equivalent to $\sum_{F \in \Psi(P)} \beta_F^{(j)} \mathbf{c}_F = \mathbf{0}$ for each $j \in [r]$. Therefore, we obtain the desired equations. \square

Proof of Theorem 2.7. Let $\mathbf{w} \in \mathbb{R}^d$ be a vector such that the interior of $P + \mathbf{u}$ has $\mathbf{0}$. Then, $Q = (P + \mathbf{u})^\circ$ is the dual polytope of P . Moreover, $P + \mathbf{u}$ has the irreducible representation

$$P + \mathbf{u} = \bigcap_{F \in \Psi(P)} H^+(\mathbf{a}_F; b'_F),$$

where $b'_F = b_F - \langle \mathbf{a}_F, \mathbf{u} \rangle$. Note that $b'_F \neq 0$ for any $F \in \Psi(P)$ since $\mathbf{0}$ belongs to the interior of $P + \mathbf{u}$.

We show that $(b'_F \beta_F)_{F \in \Psi(P)}$ is a Gale-transform of Q . For each $j \in [r]$, we define the vector \mathbf{b}_j of $\mathbb{R}^{|\Psi(P)|}$ as $\mathbf{b}_j := (b'_F \beta_F^{(j)})_{F \in \Psi(P)}$. Then, we can see that

$$\overline{\mathbf{b}}_j = (b'_F \beta_F^{(1)}, \dots, b'_F \beta_F^{(r)}) = b'_F \beta_F \quad \text{and} \quad \widehat{\mathbf{b}}_j = \beta_F / \|\beta_F\|.$$

Hence, it is enough to show that \mathbf{b}_j 's satisfy the following:

- (i) for each $j \in [r]$, \mathbf{b}_j is in $D(Q)$ and (ii) $\mathbf{b}_1, \dots, \mathbf{b}_r$ form a basis of $D(Q)$.

(i) Note that the vertex set of Q is $\{\mathbf{a}_F/b'_F : F \in \Psi(P)\}$. From Lemma 2.8, for each $j \in [r]$,

$$\sum_{F \in \Psi(P)} b'_F \beta_F^{(j)} \cdot \mathbf{a}_F/b'_F = \sum_{F \in \Psi(P)} \beta_F^{(j)} \mathbf{a}_F = \mathbf{0}.$$

Moreover,

$$\begin{aligned} \sum_{F \in \Psi(P)} b'_F \beta_F^{(j)} &= \sum_{F \in \Psi(P)} b_F \beta_F^{(j)} - \sum_{F \in \Psi(P)} \langle \mathbf{a}_F, \mathbf{u} \rangle \beta_F^{(j)} \\ &= 0 - \langle \sum_{F \in \Psi(P)} \beta_F^{(j)} \mathbf{a}_F, \mathbf{u} \rangle = -\langle \mathbf{0}, \mathbf{u} \rangle = 0. \end{aligned}$$

Therefore, we have $\mathbf{b}_j \in D(Q)$ for all $j \in [r]$.

(ii) Note that $D(Q)$ is a r -dimensional subspace of $\mathbb{R}^{|\Psi(P)|}$. Thus, it is enough to show that $\mathbf{b}_1, \dots, \mathbf{b}_r$ are linearly independent. Suppose that there exists $\mathbf{a} = (a_1, \dots, a_r) \in \mathbb{R}^r$ such that

$$a_1 \mathbf{b}_1 + \dots + a_r \mathbf{b}_r = \mathbf{0}.$$

Then, for each $F \in \Psi(P)$, one has

$$a_1 (b'_F \beta_F^{(1)}) + \dots + a_r (b'_F \beta_F^{(r)}) = b'_F \langle \mathbf{a}, \beta_F \rangle = 0.$$

Since $b'_F \neq 0$, we have $\langle \mathbf{a}, \beta_F \rangle = 0$ for any $F \in \Psi(P)$, equivalently, β_F 's lie on the hyperplane $H_{\mathbf{a}}$. This is a contradiction to the fact that β_F 's span $\mathcal{F}/\mathcal{S} \cong \mathbb{Z}^r$ as a semigroup ([5, Theorem 2]). \square

3. A NEW FAMILY OF $(0, 1)$ -POLYTOPES

In this section, we construct a new class of $(0, 1)$ -polytopes, denoted by P_{n_1, \dots, n_k} , and study its properties. We show that P_{n_1, \dots, n_k} is normal and torsionfree. Moreover, we see that if $k \geq 3$, then $\mathbb{k}[P_{n_1, \dots, n_k}]$ is not isomorphic to any Hibi ring.

Let n_1, \dots, n_k be positive integers and let $d = n_1 + \dots + n_k$. Moreover, let \mathcal{B}_n be the standard basis of \mathbb{R}^n and let $\mathcal{B}_{n_1, \dots, n_k} = \mathcal{B}_{n_1} \times \dots \times \mathcal{B}_{n_k}$. Then, we define the subset V_{n_1, \dots, n_k} of \mathbb{Z}^d as

$$V_{n_1, \dots, n_k} = \{\mathbf{0}\} \cup \mathcal{B}_d \cup \mathcal{B}_{n_1, \dots, n_k}$$

and define the $(0, 1)$ -polytope $P_{n_1, \dots, n_k} = \text{conv}(V_{n_1, \dots, n_k})$.

Example 3.1. We can see that

$$P_{1,1,1} = \text{conv}(\{(0, 0, 0), (1, 0, 0), (0, 1, 0), (0, 0, 1), (1, 1, 1)\})$$

and

$$\mathbb{k}[P_{1,1,1}] \cong \mathbb{k}[x_1, x_2, x_3, x_4, x_5]/(x_2 x_3 x_4 - x_0^2 x_5).$$

In addition, we can see that

$$\begin{aligned} P_{1,1,1,2} &= \text{conv}(\{(0, 0, 0, 0, 0), (1, 0, 0, 0, 0), (0, 1, 0, 0, 0), (0, 0, 1, 0, 0), \\ &\quad (0, 0, 0, 1, 0), (0, 0, 0, 0, 1), (1, 1, 1, 1, 0), (1, 1, 1, 0, 1)\}) \end{aligned}$$

and

$$\mathbb{k}[P_{1,1,1,2}] \cong \mathbb{k}[x_1, \dots, x_8]/(x_6 x_7 - x_5 x_8, x_2 x_3 x_4 x_6 - x_1^3 x_8, x_2 x_3 x_4 x_5 - x_1^3 x_7).$$

First, we describe the facet defining inequalities of P_{n_1, \dots, n_k} . By considering the embedding

$$\mathcal{B}_{n_p} \hookrightarrow \mathbb{R}^d \quad \mathbf{i}_p \mapsto (\underbrace{0, \dots, 0}_{n_1 + \dots + n_{p-1}}, \mathbf{i}_p, \underbrace{0, \dots, 0}_{n_{p+1} + \dots + n_k}),$$

we regard $\mathbf{i}_p \in \mathcal{B}_{n_p}$ as the element of \mathbb{R}^d .

For $p \in [k]$, let $\mathbf{f}_p = (k-2)\mathbf{1}_p - \sum_{q \in [k] \setminus \{p\}} \mathbf{1}_q$, where $\mathbf{1}_p = \sum_{\mathbf{i}_p \in \mathcal{B}_{n_p}} \mathbf{i}_p$.

Proposition 3.2. *The $(0, 1)$ -polytope P_{n_1, \dots, n_k} has the irreducible representation:*

$$P_{n_1, \dots, n_k} = \left(\bigcap_{\mathbf{e} \in \mathcal{B}_d} H_{\mathbf{e}}^+ \right) \cap \left(\bigcap_{p \in [k]} H^+(\mathbf{f}_p; 1) \right).$$

Proof. If $k = 1$, then $P_{n_1} = \text{conv}(\{\mathbf{0}\} \cup \mathcal{B}_{n_1})$ and we can see easily that P_{n_1} has the above irreducible representation. In what follows, suppose that $k \geq 2$.

We can find d affinely independent vectors in V_{n_1, \dots, n_k} on each hyperplane. Thus, these define facets of P_{n_1, \dots, n_k} .

Let $P' = \left(\bigcap_{\mathbf{e} \in \mathcal{B}_d} H_{\mathbf{e}}^+ \right) \cap \left(\bigcap_{p \in [k]} H^+(\mathbf{f}_p; 1) \right)$ and we show $P = P'$. It is clear that $P \subset P'$ since $V_{n_1, \dots, n_k} \subset P'$. Also, each vertex of P_{n_1, \dots, n_k} is on d affinely independent hyperplanes $H_{\mathbf{e}}$ or $H(\mathbf{f}_p; 1)$, that is, the set of vertices of P' , denoted by V , contains V_{n_1, \dots, n_k} . Therefore, it is enough to show that $V \subset V_{n_1, \dots, n_k}$.

Note that if $\mathbf{x} \in P'$, then $\mathbf{x} \in H^+(\mathbf{f}_p; 1)$, i.e., $\langle \mathbf{x}, \mathbf{f}_p \rangle + 1 \geq 0$ for all $p \in [k]$. Thus, for any $p \in [k]$, we have

$$\sum_{q \in [k] \setminus \{p\}} (\langle \mathbf{x}, \mathbf{f}_q \rangle + 1) = -(k-1)\langle \mathbf{x}, \mathbf{1}_p \rangle + (k-1) \geq 0 \Rightarrow \langle \mathbf{x}, \mathbf{1}_p \rangle \leq 1. \quad (3.1)$$

Moreover, since $\mathbf{x} \in H_{\mathbf{e}}^+$ for all $\mathbf{e} \in \mathcal{B}_d$, we see that $V \subset P' \subset [0, 1]^d$.

Assume that there exists an element \mathbf{v} in $V \setminus V_{n_1, \dots, n_k}$. First, suppose that $\mathbf{v} \in \{0, 1\}^d$. In this case, we see that $\langle \mathbf{v}, \mathbf{1}_p \rangle = 0$ or 1 holds for all $p \in [k]$ from (3.1). One of the following three cases happens;

- (i) $\langle \mathbf{v}, \mathbf{1}_p \rangle = 0$ for all $p \in [k]$ or $\langle \mathbf{v}, \mathbf{1}_p \rangle = 1$ for all $p \in [k]$.
- (ii) $\langle \mathbf{v}, \mathbf{1}_p \rangle = 1$ for some $p \in [k]$ and $\langle \mathbf{v}, \mathbf{1}_q \rangle = 0$ for all $q \in [k] \setminus \{p\}$.
- (iii) There exist $p_1, p_2, p_3 \in [k]$ such that $\langle \mathbf{v}, \mathbf{1}_{p_1} \rangle = \langle \mathbf{v}, \mathbf{1}_{p_2} \rangle = 1$ and $\langle \mathbf{v}, \mathbf{1}_{p_3} \rangle = 0$.

In the cases (i) and (ii), \mathbf{v} must lie in V_{n_1, \dots, n_k} . In case (iii), we have $\langle \mathbf{v}, \mathbf{f}_{p_3} \rangle + 1 < 0$, that is, $\mathbf{v} \notin H^+(\mathbf{f}_{p_3}; 1)$, a contradiction.

Now, suppose that $\mathbf{v} \notin \{0, 1\}^d$. If there are two elements $\mathbf{i}_p, \mathbf{j}_p \in \mathcal{B}_{n_p}$ with $\langle \mathbf{v}, \mathbf{i}_p \rangle, \langle \mathbf{v}, \mathbf{j}_p \rangle > 0$ for some $p \in [k]$, then \mathbf{v} is not a vertex of P' . Indeed, let

$$\mathbf{v}' = \mathbf{v} + \epsilon \mathbf{i}_p - \epsilon \mathbf{j}_p, \quad \mathbf{v}'' = \mathbf{v} - \epsilon \mathbf{i}_p + \epsilon \mathbf{j}_p,$$

where $\epsilon > 0$ is sufficiently small. Then, we can check that $\mathbf{v}', \mathbf{v}'' \in P'$ and $\mathbf{v} = \frac{1}{2}(\mathbf{v}' + \mathbf{v}'')$. Therefore, we assume that for each $p \in [k]$, $\langle \mathbf{v}, \mathbf{i}_p \rangle \geq 0$ for some $\mathbf{i}_p \in \mathcal{B}_{n_p}$ and $\langle \mathbf{v}, \mathbf{j}_p \rangle = 0$ for all $\mathbf{j}_p \in \mathcal{B}_{n_p} \setminus \{\mathbf{i}_p\}$. Let $t = \sum_{p \in [k]} \langle \mathbf{v}, \mathbf{i}_p \rangle$. In the case $t \leq 1$, let $t_p = \langle \mathbf{v}, \mathbf{i}_p \rangle$ for $p \in [k]$ and $t_{k+1} = 1 - t$. Then, we have $\mathbf{v} = \sum_{p \in [k]} t_p \mathbf{i}_p + t_{k+1} \mathbf{0}$ and $\sum_{p \in [k+1]} t_p = 1$, and hence $\mathbf{v} \notin V$. Finally, we consider the case of $t > 1$. If there exists $p \in [k]$ such that $\langle \mathbf{v}, \mathbf{i}_p \rangle = 0$ then we have $\langle \mathbf{v}, \mathbf{f}_p \rangle + 1 = -\sum_{q \in [k] \setminus \{p\}} \langle \mathbf{v}, \mathbf{i}_q \rangle + 1 = -t + 1 < 0$, a contradiction. Thus, we can see that $\langle \mathbf{v}, \mathbf{i}_p \rangle > 0$ for all $p \in [k]$. Moreover, since \mathbf{v} must lie on d affinely independent hyperplanes $H_{\mathbf{e}}$ or $H(\mathbf{f}_p; 1)$, we have $\mathbf{v} \in H(\mathbf{f}_p; 1)$ for all $p \in [k]$, that is, $\langle \mathbf{v}, \mathbf{f}_p \rangle + 1 = 0$

holds for any $p \in [k]$. We may assume that $\langle \mathbf{v}, \mathbf{i}_1 \rangle = \min\{\langle \mathbf{v}, \mathbf{i}_p \rangle : p \in [k]\}$, and let $t_p = \begin{cases} \langle \mathbf{v}, \mathbf{i}_p \rangle - \langle \mathbf{v}, \mathbf{i}_1 \rangle & \text{if } p \neq 1, \\ \langle \mathbf{v}, \mathbf{i}_1 \rangle & \text{if } p = 1 \end{cases}$ for $p \in [k]$. Then, we obtain

$$\mathbf{v} = \sum_{p \in [k] \setminus \{1\}} t_p \mathbf{i}_p + t_1 \sum_{p \in [k]} \mathbf{i}_p$$

and

$$\sum_{p \in [k]} t_p = -(k-2)\langle \mathbf{v}, \mathbf{i}_1 \rangle + \sum_{p \in [k] \setminus \{1\}} \langle \mathbf{v}, \mathbf{i}_p \rangle = -(k-2)\langle \mathbf{v}, \mathbf{1}_1 \rangle + \sum_{p \in [k] \setminus \{1\}} \langle \mathbf{v}, \mathbf{1}_p \rangle = -\langle \mathbf{v}, \mathbf{f}_1 \rangle = 1.$$

Therefore, v is not a vertex of P' , and hence V coincides with V_{n_1, \dots, n_k} . \square

Example 3.3. We consider $P_{1,1,1}$. Then,

$$x_i \geq 0 \quad (i = 1, 2, 3),$$

$$\langle \mathbf{f}_p, \mathbf{x} \rangle + 1 = x_p - (x_{q_1} + x_{q_2}) + 1 \geq 0 \quad (p = 1, 2, 3 \text{ and } \{p, q_1, q_2\} = \{1, 2, 3\})$$

are the facet defining inequalities of $P_{1,1,1}$.

Moreover, we consider $P_{1,1,1,2}$. Then, the following are the facet defining inequalities of $P_{1,1,1,2}$:

$$x_i \geq 0 \quad (i = 1, \dots, 5),$$

$$2x_p - (x_{q_1} + \dots + x_{q_4}) + 1 \geq 0 \quad (p = 1, 2, 3 \text{ and } \{p, q_1, \dots, q_4\} = \{1, \dots, 5\}) \text{ and}$$

$$2(x_4 + x_5) - (x_1 + x_2 + x_3) + 1 \geq 0.$$

Next, we investigate the initial ideal of the toric ideal $I_{P_{n_1, \dots, n_k}}$ with respect to a monomial order and provide its Gröbner basis, which allows us to study the normality. For the fundamental materials on initial ideals and Gröbner bases, consult, e.g., [13].

For $x_{\mathbf{v}} \in T = \mathbb{k}[x_{\mathbf{v}} : \mathbf{v} \in P_{n_1, \dots, n_k} \cap \mathbb{Z}^d]$, we denote by $x_{\mathbf{i}_1, \dots, \mathbf{i}_k}$ instead of $x_{(\mathbf{i}_1, \dots, \mathbf{i}_k)}$ for $(\mathbf{i}_1, \dots, \mathbf{i}_k) \in \mathcal{B}_{n_1, \dots, n_k}$. Let $<$ denote the reverse lexicographic order on T induced by the ordering of the variables as follows:

- $x_{\mathbf{e}} < x_{\mathbf{i}_1, \dots, \mathbf{i}_k}$ for any $\mathbf{e} \in \{\mathbf{0}\} \cup \mathcal{B}_d$ and any $(\mathbf{i}_1, \dots, \mathbf{i}_k) \in \mathcal{B}_{n_1, \dots, n_k}$;
- For $\mathbf{e}, \mathbf{e}' \in \{\mathbf{0}\} \cup \mathcal{B}_d$ (resp. for $(\mathbf{i}_1, \dots, \mathbf{i}_k), (\mathbf{j}_1, \dots, \mathbf{j}_k) \in \mathcal{B}_{n_1, \dots, n_k}$), $x_{\mathbf{e}} < x_{\mathbf{e}'}$ (resp. $x_{\mathbf{i}_1, \dots, \mathbf{i}_k} < x_{\mathbf{j}_1, \dots, \mathbf{j}_k}$) if and only if $\mathbf{e} < \mathbf{e}'$ (resp. $(\mathbf{i}_1, \dots, \mathbf{i}_k) < (\mathbf{j}_1, \dots, \mathbf{j}_k)$), which means that the leftmost nonzero component of the vector $\mathbf{e}' - \mathbf{e}$ (resp. $(\mathbf{j}_1 - \mathbf{i}_1, \dots, \mathbf{j}_k - \mathbf{i}_k)$) is positive.

Moreover, let $\mathcal{G}_{n_1, \dots, n_k}$ be the sets of the following binomials in T :

$$(b1) \quad x_{\mathbf{i}_1} x_{\mathbf{i}_2} \cdots x_{\mathbf{i}_k} - x_{\mathbf{0}}^{k-1} x_{\mathbf{i}_1, \dots, \mathbf{i}_k} \text{ for } (\mathbf{i}_1, \dots, \mathbf{i}_k) \in \mathcal{B}_{n_1, \dots, n_k};$$

$$(b2) \quad x_{\mathbf{j}_p} x_{\mathbf{i}_1, \dots, \mathbf{i}_p, \dots, \mathbf{i}_k} - x_{\mathbf{i}_p} x_{\mathbf{i}_1, \dots, \mathbf{j}_p, \dots, \mathbf{i}_k} \text{ for } (\mathbf{i}_1, \dots, \mathbf{i}_k) \in \mathcal{B}_{n_1, \dots, n_k}, \mathbf{j}_p \in \mathcal{B}_{n_p} \text{ with } \mathbf{i}_p < \mathbf{j}_p;$$

$$(b3) \quad x_{\mathbf{i}_1, \dots, \mathbf{i}_k} x_{\mathbf{j}_1, \dots, \mathbf{j}_k} - x_{\mathbf{i}'_1, \dots, \mathbf{i}'_k} x_{\mathbf{j}'_1, \dots, \mathbf{j}'_k} \text{ for } (\mathbf{i}_1, \dots, \mathbf{i}_k), (\mathbf{j}_1, \dots, \mathbf{j}_k) \in \mathcal{B}_{n_1, \dots, n_k},$$

where we define $\mathbf{i}'_p = \begin{cases} \mathbf{i}_p & \text{if } x_{\mathbf{i}_p} < x_{\mathbf{j}_p}, \\ \mathbf{j}_p & \text{else} \end{cases}$ and define \mathbf{j}'_p as satisfying $\{\mathbf{i}'_p, \mathbf{j}'_p\} = \{\mathbf{i}_p, \mathbf{j}_p\}$.

Note that each leading term of these binomials is the initial monomial with respect to $<$. Furthermore, these binomials belong to the toric ideal of P_{n_1, \dots, n_k} , that is, $\mathcal{G}_{n_1, \dots, n_k} \subset I_{P_{n_1, \dots, n_k}}$.

Proposition 3.4. *Let the notation be the same as above. Then, $\mathcal{G}_{n_1, \dots, n_k}$ is a Gröbner basis of $I_{\mathcal{P}_{n_1, \dots, n_k}}$ with respect to $<$.*

To show this proposition, we use the following lemma:

Lemma 3.5 ([13, Theorem 3.11]). *Let $I \subset S = \mathbb{k}[x_1, \dots, x_n]$ be the toric ideal of an integral polytope and $\mathcal{G} = \{g_1, \dots, g_s\}$ the set of binomials in I . Fix a monomial order $<$ on S and let $\text{in}_<(\mathcal{G})$ be the ideal of S generated by the initial monomials $\text{in}_<(g_1), \dots, \text{in}_<(g_s)$, that is, $\text{in}_<(\mathcal{G}) = (\{\text{in}_<(g) : g \in \mathcal{G}\})$. Then, the following conditions are equivalent:*

- (i) \mathcal{G} is a Gröbner basis of I with respect to $<$.
- (ii) For monomials $u, v \in S$, if $u \notin \text{in}_<(\mathcal{G})$, $v \notin \text{in}_<(\mathcal{G})$ and $u \neq v$ then $u - v \notin I$.

Proof of Proposition 3.4. We show that $\mathcal{G}_{n_1, \dots, n_k}$ satisfies the condition (ii) in Lemma 3.5. Let u be a monomial in T . Then, u can be written by

$$x_0^a x_{\mathbf{e}_1} x_{\mathbf{e}_2} \cdots x_{\mathbf{e}_s} x_{\mathbf{i}_{11}, \dots, \mathbf{i}_{1k}} \cdots x_{\mathbf{i}_{t1}, \dots, \mathbf{i}_{tk}},$$

where $a \in \mathbb{Z}_{\geq 0}$, $\mathbf{e}_1, \dots, \mathbf{e}_s \in \mathcal{B}_d$ and $\mathbf{i}_{1p}, \dots, \mathbf{i}_{tp} \in \mathcal{B}_{n_p}$ for each $p \in [k]$. Let $M_p = \{\mathbf{e}_1, \dots, \mathbf{e}_s\} \cap \mathcal{B}_{n_p}$ and $N_p = |M_p|$ for $p \in [k]$. Then, if $u \notin \text{in}_<(\mathcal{G}_{n_1, \dots, n_k})$, we can see that:

- (a) there exists $p \in [k]$ such that $M_p = \emptyset$ from (b1);
- (b) for each $p \in [k]$, we have $\mathbf{e}_i \leq \mathbf{i}_{lp}$ for any $\mathbf{e} \in M_p$ and for any $l \in [t]$ from (b2);
- (c) $\mathbf{i}_{1p} \leq \mathbf{i}_{2p} \leq \cdots \leq \mathbf{i}_{tp}$ for all $p \in [k]$ by permuting $x_{\mathbf{i}_{11}, \dots, \mathbf{i}_{1k}}, \dots, x_{\mathbf{i}_{t1}, \dots, \mathbf{i}_{tk}}$ from (b3).

Let $\phi_{P_{n_1, \dots, n_k}}(u) = t_1^{r_1} t_2^{r_2} \cdots t_d^{r_d} t_0^{r_0}$. Now, we may assume that $M_1 = \emptyset$. Then, we can see that $t = r_1 + r_2 + \cdots + r_{n_1}$. Similarly, N_p and a can also be represented by r_i 's. Moreover, it follows from (b), (c) that \mathbf{e}_i and $\mathbf{i}_{1p}, \dots, \mathbf{i}_{tp}$ can be determined uniquely from r_0, \dots, r_d . Therefore, we can recover u from $t_1^{r_1} t_2^{r_2} \cdots t_d^{r_d} t_0^{r_0}$. This is equivalent to (ii) in Lemma 3.5. \square

Now, we give the main theorem in this section.

Theorem 3.6. *The $(0, 1)$ -polytope P_{n_1, \dots, n_k} has the following properties:*

- (i) P_{n_1, \dots, n_k} has IDP.
- (ii) $\mathbb{k}[P_{n_1, \dots, n_k}]$ is Gorenstein if and only if $n_1 = n_2 = \cdots = n_k$.
- (iii) $\text{Cl}(\mathbb{k}[P_{n_1, \dots, n_k}]) \cong \mathbb{Z}^{k-1}$.
- (iv) If $k \geq 3$, then $\mathbb{k}[P_{n_1, \dots, n_k}] \notin \mathbf{Order}_{k-1}$ for any $n_1, \dots, n_k \in \mathbb{Z}_{>0}$.

Proof. (i) It follows from Proposition 3.4 that the initial ideal of I_{n_1, \dots, n_k} is squarefree, and hence P_{n_1, \dots, n_k} possesses a regular unimodular triangulation (cf. [13, Theorem 4.17]). This implies that P_{n_1, \dots, n_k} is normal. Moreover, $\mathbb{Z}\mathcal{A}(P_{n_1, \dots, n_k})$ coincides with \mathbb{Z}^{d+1} since $\{\mathbf{0}\} \cup \mathcal{B}_d \subset P_{n_1, \dots, n_k}$, and hence P_{n_1, \dots, n_k} has IDP.

(ii) Since $\mathbb{k}[P_{n_1, \dots, n_k}]$ is a normal affine semigroup ring, the canonical module $\omega_{\mathbb{k}[P_{n_1, \dots, n_k}]}$ is isomorphic to the module generated by all monomials whose exponent vector is a lattice point in $\left(\bigcap_{\mathbf{e} \in \mathcal{B}_d} H^+(\mathbf{e}, 0); 1\right) \cap \left(\bigcap_{p \in [k]} H^+(\mathbf{f}_p, 1); 1\right)$. By the parallel translation by $(1, \dots, 1, \alpha)$ for some integer α , we can see that it is also isomorphic to the module generated by all monomials whose exponent vector is a lattice point in $\left(\bigcap_{\mathbf{e} \in \mathcal{B}_d} H^+(\mathbf{e}, 0); 0\right) \cap \left(\bigcap_{p \in [k]} H^+(\mathbf{f}_p, 1); m_p\right)$, where $m_p = 1 - \alpha - (k - 2)n_p + \sum_{q \in [k] \setminus \{p\}} n_q$.

Thus, if $\mathbb{k}[P_{n_1, \dots, n_k}]$ is Gorenstein, then the equality $m_1 = \cdots = m_k = 0$ must hold. This implies $n_1 = \cdots = n_k$. Conversely, if n_1, \dots, n_k are equal, then we can see that $\omega_{\mathbb{k}[P_{n_1, \dots, n_k}]}$ is isomorphic to $\mathbb{k}[P_{n_1, \dots, n_k}]$ by setting $\alpha = n_1 + 1$, that is, $\mathbb{k}[P_{n_1, \dots, n_k}]$ is Gorenstein.

(iii) Let $\mathbf{v}_1, \dots, \mathbf{v}_d$ be vectors in \mathbb{Z}^d satisfying $\{\mathbf{v}_1, \dots, \mathbf{v}_d\} = \mathcal{B}_d$ and let $\mathbf{v}_{d+1} = \mathbf{0}$. Moreover, for $i \in [d]$, let F_i be the facet of P_{n_1, \dots, n_k} defined by $H(\mathbf{v}_i; 0)$ and let F_{d+1} be the facet of P_{n_1, \dots, n_k} defined by $H(\mathbf{f}_1; 1)$. Then, we can see that these sequences satisfy the statement (*) and we have $k_{P_{n_1, \dots, n_k}} = d + 1$. Therefore, we have $\text{Cl}(\mathbb{k}[P_{n_1, \dots, n_k}]) \cong \mathbb{Z}^{k-1}$ from Theorem 2.2.

(iv) Since $\mathcal{G}_{n_1, \dots, n_k}$ is a Gröbner basis, the binomials (b1), (b2) and (b3) generate $I_{P_{n_1, \dots, n_k}}$. Consider a minimal generators \mathcal{G}' of $I_{P_{n_1, \dots, n_k}}$ contained these binomials. If \mathcal{G}' does not contain any binomial $x_{i_1} x_{i_2} \cdots x_{i_k} - x_{\mathbf{0}}^{k-1} x_{i_1, \dots, i_k}$ in (b1), then there exists an expression:

$$x_{i_1} x_{i_2} \cdots x_{i_k} - x_{\mathbf{0}}^{k-1} x_{i_1, \dots, i_k} = \sum_{i=1}^s \alpha_i \mathbf{x}^{\mathbf{w}_i} f_i,$$

where $\alpha_i \in \mathbb{Z}$, $\mathbf{x}^{\mathbf{w}_i}$ is a monomial of the polynomial ring $\mathbb{k}[x_{\mathbf{v}} : \mathbf{v} \in P \cap \mathbb{Z}^d]$ and f_i 's are binomials in (b2) or (b3) (cf. [13, Lemma 3.7]). However, it is impossible because the variable x_{i_1, \dots, i_k} for some $(\mathbf{i}_1, \dots, \mathbf{i}_k) \in V_{n_1, \dots, n_k}$ must appear in the terms of f_i . Therefore, \mathcal{G}' contains $x_{i_1} x_{i_2} \cdots x_{i_k} - x_{\mathbf{0}}^{k-1} x_{i_1, \dots, i_k}$ for some $\mathbf{i}_1, \dots, \mathbf{i}_k$. This implies that I_{n_1, \dots, n_k} cannot be generated by quadratic binomials.

On the other hand, the toric ideals of Hibi rings are generated by quadratic binomials ([14]). Thus, $\mathbb{k}[P_{n_1, \dots, n_k}]$ is not isomorphic to any Hibi ring. \square

Finally, we compute weights of $\mathbb{k}[P_{n_1, \dots, n_k}]$. Let $F_{\mathbf{e}}$ denote the facet $P_{n_1, \dots, n_k} \cap H(\mathbf{e}; 0)$ for $\mathbf{e} \in \mathcal{B}_d$ and let F_i denote the facet $P_{n_1, \dots, n_k} \cap H(\mathbf{f}_p; 1)$ for $p \in [k]$. From (2.2) and Theorem 3.6 (i), the following elements in \mathcal{F} belong to \mathcal{S} :

$$\sum_{F \in \Psi(P_{n_1, \dots, n_k})} \langle (\mathbf{i}_p, 0), \mathbf{c}_F \rangle \epsilon_F = \epsilon_{F_{i_p}} + (k-2)\epsilon_{F_p} - \sum_{q \in [k] \setminus \{p\}} \epsilon_{F_q} \quad (3.2)$$

for each $p \in [k]$ and $\mathbf{i}_p \in \mathcal{B}_{n_p}$ and

$$\sum_{F \in \Psi(P_{n_1, \dots, n_k})} \langle (\mathbf{0}, 1), \mathbf{c}_F \rangle \epsilon_F = \sum_{p \in [k]} \epsilon_{F_p}. \quad (3.3)$$

We consider the map $\iota : \mathcal{F}/\mathcal{S} \rightarrow \mathbb{Z}^{k-1}$; let $\iota(\epsilon_{F_i}) = \mathbf{e}_i$ for $i \in [k-1]$, where \mathbf{e}_i denotes the i -th unit vector of \mathbb{Z}^{k-1} . This induces an isomorphism $\iota : \mathcal{F}/\mathcal{S} \rightarrow \mathbb{Z}^{k-1}$ and we can calculate the remaining weight from (3.2) and (3.3):

$$\begin{aligned} \iota(\epsilon_{F_k}) &= -(\mathbf{e}_1 + \cdots + \mathbf{e}_{k-1}); \\ \iota(\epsilon_{F_{i_p}}) &= -(k-1)\mathbf{e}_p \quad \text{for each } p \in [k-1] \text{ and any } \mathbf{i}_p \in \mathcal{B}_{n_p}; \\ \iota(\epsilon_{F_{i_k}}) &= (k-1)(\mathbf{e}_1 + \cdots + \mathbf{e}_{k-1}) \quad \text{for any } \mathbf{i}_k \in \mathcal{B}_{n_k}. \end{aligned}$$

In particular, we can get the weights of the case $k = 3$ as follows:

$$(1, 0), \quad (0, 1), \quad (-1, -1), \quad (-2, 0) \times n_1, \quad (0, -2) \times n_2, \quad (2, 2) \times n_3. \quad (3.4)$$

4. ANSWERS TO THE PROBLEMS

Throughout this section, let $P \subset \mathbb{R}^d$ be a $(0, 1)$ -polytope. We give complete or partial answers to Problems 1, 2, 3 and 4 in each case;

$$(r1) \quad \text{rank } P = 0 \text{ or } 1, \quad (r2) \quad \text{rank } P = 2, \quad (r3) \quad \text{rank } P \geq 3.$$

4.1. Case (r1). First, we discuss the case $\text{rank } P = 0$, this case is trivial. Notice that P has rank 0 if and only if P is a simplex. Moreover, it is known that the toric ring of d -dimensional $(0, 1)$ -simplex is isomorphic to the polynomial ring with $d + 1$ variables over \mathbb{k} (cf. [33, Lemma 3.1.5]).

Clearly, polynomial rings are normal and their divisor class groups are torsionfree. Moreover, for two $(0, 1)$ -simplices P and P' , one has $\mathbb{k}[P] \cong \mathbb{k}[P']$ if and only if P and P' are combinatorially equivalent (equivalently, they have the same number of vertices). Therefore, we get the following theorem:

Proposition 4.1. *All $(0, 1)$ -polytopes with rank 0 are normal and torsionfree. Moreover, the following relationship holds:*

$$(\mathbf{0}, \mathbf{1})_0 = \mathbf{Order}_0 = \{\mathbb{k}[x_1, \dots, x_k] : k \in \mathbb{Z}_{>0}\}.$$

Furthermore, for two $(0, 1)$ -polytopes P_1 and P_2 with $\text{rank } P_i = 0$, they have the same combinatorial type if and only if $\mathbb{k}[P_1] \cong \mathbb{k}[P_2]$.

Next, we discuss the case $\text{rank } P = 1$. This case has been investigated in [20] as follows:

Theorem 4.2 ([20, Theorem 3.7]). *The following conditions are equivalent:*

- (i) P has rank 1, that is, P has just $\dim P + 2$ facets;
- (ii) $\mathbb{k}[P]$ is isomorphic to the Segre product of two polynomial rings $\mathbb{k}[x_1, \dots, x_{n+1}]$ and $\mathbb{k}[y_1, \dots, y_{m+1}]$ for some $n, m \in \mathbb{Z}_{>0}$ or its polynomial extension;
- (iii) P is normal and $\text{Cl}(\mathbb{k}[P]) \cong \mathbb{Z}$.

Therefore, $(0, 1)$ -polytopes which have rank 1 are normal and torsionfree. Moreover, the equivalence (i) and (ii) imply that for two $(0, 1)$ -polytopes P and P' with $\text{rank } P = \text{rank } P' = 1$, one has $\mathbb{k}[P] \cong \mathbb{k}[P']$ if and only if P and P' are combinatorially equivalent. Hence, the following theorem holds:

Theorem 4.3. *All $(0, 1)$ -polytopes with rank 1 are normal and torsionfree. Moreover, the following relationship holds:*

$$(\mathbf{0}, \mathbf{1})_1 = \mathbf{Order}_1 = \{(\mathbb{k}[x_1, \dots, x_{n+1}] \# \mathbb{k}[y_1, \dots, y_{m+1}]) \otimes_{\mathbb{k}} \mathbb{k}[z_1, \dots, z_{l-1}] : n, m, l \in \mathbb{Z}_{>0}\}.$$

Furthermore, for two $(0, 1)$ -polytopes P_1 and P_2 with $\text{rank } P_i = 1$, they have the same combinatorial type if and only if $\mathbb{k}[P_1] \cong \mathbb{k}[P_2]$.

4.2. Case (r2). By Proposition 2.4, in what follows, we may assume that P is not pyramidal.

By Theorem 2.7, we can obtain the standard Gale-diagrams of the dual polytopes of the order polytopes which have rank 2 and P_{n_1, n_2, n_3} . We draw the standard Gale-diagrams as follows; the dual polytopes of $\mathcal{O}_{\Pi_1}, \mathcal{O}_{\Pi_2}, \mathcal{O}_{\Pi_3}, \mathcal{O}_{\Pi_4}$ and P_{n_1, n_2, n_3} correspond to the standard Gale-diagrams $\text{Gale}_1, \text{Gale}_2, \text{Gale}_3, \text{Gale}_4$ and Gale_5 , respectively.

Theorem 4.4. *The following are equivalent:*

- (i) The standard Gale-diagram of the dual polytope of P is orthogonally equivalent to Gale_1 .
- (ii) The toric ring of P is isomorphic to the Segre product of three polynomial rings over \mathbb{k} .

In particular, let P_1 and P_2 be two $(0, 1)$ -polytopes which have a standard Gale-diagram Gale_1 , then P_1 and P_2 are combinatorially equivalent if and only if $\mathbb{k}[P_1] \cong \mathbb{k}[P_2]$.




FIGURE 5.
The Gale-diagram Gale_1




FIGURE 6.
The Gale-diagram Gale_2




FIGURE 7.
The Gale-diagram Gale_3




FIGURE 8.
The Gale-diagram Gale_4




FIGURE 9.
The Gale-diagram Gale_5

Proof. (i) \Rightarrow (ii) : Since no diameter of the standard Gale-diagram has both endpoints, the dual polytope of P is simplicial, that is, P is simple. Therefore, P is the product of just three $(0, 1)$ -simplices from Proposition 2.3, and hence $\mathbb{k}[P]$ is isomorphic to the Segre product of three polynomial rings.

(ii) \Rightarrow (i) : It follows from Proposition 2.5 that the Segre product of three polynomial rings can be realized as the Hibi ring of a poset Π_1 . Its weights have been already given in Section 2.2. By Theorem 2.7, we can obtain the standard Gale-diagram of the dual polytope of P which is orthogonally equivalent to Gale_1 .

The last statement follows from the equivalence (i) and (ii). □

Theorem 4.5. *The following are equivalent:*

- (i) *The standard Gale-diagram of the dual polytope of P is orthogonally equivalent to Gale_2 .*

(ii) *The toric ring of P is isomorphic to $R_1 \otimes_{\mathbb{k}} R_2$, where R_i is the Segre product of two polynomial rings over \mathbb{k} .*

In particular, let P_1 and P_2 be two $(0,1)$ -polytopes which have a standard Gale-diagram Gale_2 , then P_1 and P_2 are combinatorially equivalent if and only if $\mathbb{k}[P_1] \cong \mathbb{k}[P_2]$.

Proof. (i) \Rightarrow (ii) : Since the standard Gale-diagram of the dual polytope of P is orthogonally equivalent to Gale_2 , P is combinatorially equivalent to \mathcal{O}_{Π_2} . Therefore, there exists a one-to-one mapping Φ between the set of all faces of \mathcal{O}_{Π_2} and the set of all faces of P such that Φ is inclusion preserving.

Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ denote the vertices of \mathcal{O}_{Π_2} corresponding to the poset ideals of Π_2 including the element which is comparable with any other element of Π_2 , and let $\mathbf{v}_1, \dots, \mathbf{v}_m$ denote the remaining vertices of \mathcal{O}_{Π_2} . Notice that any facet F of \mathcal{O}_{Π_2} contains $\{\mathbf{u}_1, \dots, \mathbf{u}_n\}$ or $\{\mathbf{v}_1, \dots, \mathbf{v}_m\}$, that is, $d_F(\mathbf{u}_i) = 0$ for all $i \in [n]$, or $d_F(\mathbf{v}_j) = 0$ for all $j \in [m]$. Moreover, we can see that $O_1 = \text{conv}(\{\mathbf{u}_1, \dots, \mathbf{u}_n\})$ and $O_2 = \text{conv}(\{\mathbf{v}_1, \dots, \mathbf{v}_m\})$ are $(0,1)$ -polytopes with $\text{rank } O_1 = \text{rank } O_2 = 1$.

Let $\mathbf{w}_i = \Phi(\mathbf{u}_i)$, $\mathbf{z}_j = \Phi(\mathbf{v}_j)$, $Q_1 = \Phi(O_1) = \text{conv}(\{\mathbf{w}_1, \dots, \mathbf{w}_n\})$ and $Q_2 = \Phi(O_2) = \text{conv}(\{\mathbf{z}_1, \dots, \mathbf{z}_m\})$. We show that $\mathbb{k}[P] \cong \mathbb{k}[Q_1] \otimes_{\mathbb{k}} \mathbb{k}[Q_2]$, equivalently, $I_P = RI_{Q_1} + RI_{Q_2}$, where $R = \mathbb{k}[x_{\mathbf{v}} : \mathbf{v} \in P \cap \mathbb{Z}^d]$. Since $I_P \supset RI_{Q_1}$ and $I_P \supset RI_{Q_2}$, we have $I_P \supset RI_{Q_1} + RI_{Q_2}$. To prove the reverse inclusion, it is enough to show that a binomial $b = x_{\mathbf{w}_{i_1}} \cdots x_{\mathbf{w}_{i_p}} x_{\mathbf{z}_{j_1}} \cdots x_{\mathbf{z}_{j_q}} - x_{\mathbf{w}_{h_1}} \cdots x_{\mathbf{w}_{h_s}} x_{\mathbf{z}_{g_1}} \cdots x_{\mathbf{z}_{g_t}} \in I_P$ belongs to $RI_{Q_1} + RI_{Q_2}$.

In this situation, we have

$$\mathbf{w}'_{i_1} + \cdots + \mathbf{w}'_{i_p} + \mathbf{z}'_{j_1} + \cdots + \mathbf{z}'_{j_q} = \mathbf{w}'_{h_1} + \cdots + \mathbf{w}'_{h_s} + \mathbf{z}'_{g_1} + \cdots + \mathbf{z}'_{g_t},$$

where for $\mathbf{v} \in \mathbb{Z}^d$, we define $\mathbf{v}' = (\mathbf{v}, 1)$.

Since for any $F' \in \Psi(P)$, $d_{F'}(\mathbf{w}_i) = 0$ for all $i \in [n]$, or $d_{F'}(\mathbf{z}_j) = 0$ for all $j \in [m]$, we can see that $\langle \sum_{k \in [p]} \mathbf{w}'_{i_k} - \sum_{l \in [s]} \mathbf{w}'_{h_l}, \mathbf{c}_{F'} \rangle = 0$ for all $F' \in \Psi(P)$. Indeed, if F' contains \mathbf{w}_i for all $i \in [n]$, then

$$\begin{aligned} \langle \sum_{k \in [p]} \mathbf{w}'_{i_k} - \sum_{l \in [s]} \mathbf{w}'_{h_l}, \mathbf{c}_{F'} \rangle &= \sum_{k \in [p]} \langle \mathbf{w}'_{i_k}, \mathbf{c}_{F'} \rangle - \sum_{l \in [s]} \langle \mathbf{w}'_{h_l}, \mathbf{c}_{F'} \rangle \\ &= \sum_{k \in [p]} d_{F'}(\mathbf{w}_{i_k}) - \sum_{l \in [s]} d_{F'}(\mathbf{w}_{h_l}) = 0. \end{aligned}$$

Moreover, if F' contains \mathbf{z}_j for all $j \in [m]$, then

$$\begin{aligned} \langle \sum_{k \in [p]} \mathbf{w}'_{i_k} - \sum_{l \in [s]} \mathbf{w}'_{h_l}, \mathbf{c}_{F'} \rangle &= \langle \sum_{k \in [p]} \mathbf{w}'_{i_k} + \sum_{k \in [q]} \mathbf{z}'_{j_k} - \sum_{l \in [s]} \mathbf{w}'_{h_l} - \sum_{l \in [t]} \mathbf{z}'_{g_l}, \mathbf{c}_{F'} \rangle \\ &= \langle \mathbf{0}, \mathbf{c}_{F'} \rangle = 0. \end{aligned}$$

The homomorphism from $\mathbb{Z}\mathcal{A}(P)$ onto \mathcal{S} given by $\mathbf{v}' \mapsto \sum_{F' \in \Psi(P)} \langle \mathbf{v}', \mathbf{c}_{F'} \rangle \epsilon_{F'}$ is an isomorphism (cf. [33, Proposition 9.8.17]). Therefore, we have $\sum_{k \in [p]} \mathbf{w}'_{i_k} - \sum_{l \in [s]} \mathbf{w}'_{h_l} = \mathbf{0}$. Similarly, we have $\sum_{k \in [q]} \mathbf{z}'_{j_k} - \sum_{l \in [t]} \mathbf{z}'_{g_l} = \mathbf{0}$, and hence

$$\begin{aligned} b &= x_{\mathbf{w}_{i_1}} \cdots x_{\mathbf{w}_{i_p}} (x_{\mathbf{z}_{j_1}} \cdots x_{\mathbf{z}_{j_q}} - x_{\mathbf{z}_{g_1}} \cdots x_{\mathbf{z}_{g_t}}) + x_{\mathbf{z}_{g_1}} \cdots x_{\mathbf{z}_{g_t}} (x_{\mathbf{w}_{i_1}} \cdots x_{\mathbf{w}_{i_p}} - x_{\mathbf{w}_{h_1}} \cdots x_{\mathbf{w}_{h_s}}) \\ &\in RI_{Q_1} + RI_{Q_2}. \end{aligned}$$

Since Q_1 and Q_2 have rank 1, it follows from Theorem 4.2 that $\mathbb{k}[Q_1]$ and $\mathbb{k}[Q_2]$ are the Segre products of two polynomial rings, we get desired.

(ii) \Rightarrow (i) : By Proposition 2.5 (i) and (ii), $\mathbb{k}[P]$ can be realized as the Hibi ring of a poset Π_2 . From its weights given in Section 2.2 and Theorem 2.7, we can see that the standard Gale-diagram of the dual polytope of P is orthogonally equivalent to Gale_2 .

The last statement follows from the equivalence (i) and (ii). \square

These theorems do not give a complete answer to our problems. We are left with the following questions:

Question 1. For any $(0, 1)$ -polytope P with $\text{rank } P = 2$, is $\mathbb{k}[P]$ isomorphic to a Hibi ring or $\mathbb{k}[P_{n_1, n_2, n_3}]$ for some $n_1, n_2, n_3 \in \mathbb{Z}_{>0}$? In other words, does the relationship $(\mathbf{0}, \mathbf{1})_2 = \text{Order}_2 \sqcup \{\mathbb{k}[P_{n_1, n_2, n_3}] : n_1, n_2, n_3 \in \mathbb{Z}_{>0}\}$ hold?

Question 2. For any $(0, 1)$ -polytope P with $\text{rank } P = 2$, is the standard Gale-diagram of the dual polytope of P orthogonally equivalent to one of the Gale_i 's? Also, let P and P' be two $(0, 1)$ -polytopes whose dual polytopes have the standard Gale-diagrams $\text{Gale}_3, \text{Gale}_4$ or Gale_5 , then does the combinatorial equivalence of P and P' imply the isomorphism of their toric rings?

If Question 1 has a positive answer, then all $(0, 1)$ -polytopes with rank 2 are normal and torsionfree.

4.3. Case (r3). Finally, we discuss the normality, torsionfreeness and classification in the case $\text{rank } P \geq 3$. In fact, unlike the previous cases, these properties are not guaranteed.

Proposition 4.6. For any positive integer $r \geq 3$, there exists a non-normal $(0, 1)$ -polytope P with $\text{rank } P = r$.

Proof. Let

$$Q_1 = \text{conv}(\{(0, 0, 0, 0), (1, 1, 0, 0), (1, 0, 1, 0), (0, 1, 1, 0), (0, 0, 0, 1), (1, 1, 1, 1)\}).$$

Then, we can see that Q_1 has rank 3 and is not normal. Indeed, we can check that $\dim Q_1 = 4$ and Q_1 has 8 facets by using Magma ([4]), thus $\text{rank } Q_1 = 3$. Moreover, one has $(1, 1, 1, 0, 2) \in \mathbb{Z}\mathcal{A}(Q_1) \cap \mathbb{R}_{\geq 0}\mathcal{A}(Q_1)$ while $(1, 1, 1, 0, 2) \notin \mathbb{Z}_{\geq 0}\mathcal{A}(Q_1)$.

Furthermore, $Q_1 \times [0, 1]^{r-3}$, where $[0, 1]^d$ denotes the d dimensional unit cube, is also a non-normal $(0, 1)$ -polytope and its rank is equal to r . \square

Proposition 4.7. For any positive integer $r \geq 3$, there exists a non-torsionfree normal $(0, 1)$ -polytope P with $\text{rank } P = r$.

Proof. Let

$$Q_2 = \text{conv}(\{(0, 0, 0, 0), (1, 0, 0, 0), (0, 1, 0, 1), (0, 0, 1, 1), (1, 1, 1, 0), (1, 1, 1, 1)\}).$$

Then, we can see that Q_2 has rank 3 and is normal but not torsionfree. Indeed, we can see that $\mathbb{Z}\mathcal{A}(Q_2) = \mathbb{Z}^5$ and Q_2 has IDP by using Magma, so Q_2 is normal. Moreover, we can compute \mathcal{M}_{Q_2} and its Smith normal form as follows:

$$\mathcal{M}_{Q_2} = \begin{pmatrix} 1 & 0 & 2 & 0 & 0 & 0 \\ 1 & 0 & 0 & 2 & 0 & 0 \\ 1 & 0 & 0 & 0 & 2 & 0 \\ 1 & 0 & 0 & 0 & 0 & 2 \\ 0 & 1 & 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 2 & 0 & 0 \\ 0 & 1 & 0 & 0 & 2 & 0 \\ 0 & 1 & 0 & 0 & 0 & 2 \end{pmatrix} \longrightarrow \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

- [9] J. Edmonds, Maximum matching and a polyhedron with 0, 1-vertices, *J. Res. Nat. Bur. Standards Sect B*, **69B**, (1965), 125–130.
- [10] B. Grünbaum, “Convex Polytopes” Second edition, Graduate Texts in Mathematics, **221**, Springer, New York, (2003).
- [11] M. Hashimoto, T. Hibi and A. Noma, Divisor class groups of affine semigroup rings associated with distributive lattices, *J. Algebra* **149** (2), (1992), 352–357.
- [12] J. Herzog, T. Hibi, S. Moradi, and A. A. Qureshi, The toric face ring of a simplicial complex, arXiv:2302.03653.
- [13] J. Herzog, T. Hibi and H. Ohsugi, Binomial ideals, Graduate Texts in Mathematics, **279**, Springer, Cham, (2018).
- [14] T. Hibi, Distributive lattices, affine semigroup rings and algebras with straightening laws. In: Nagata, M., Matsumura, H. (eds.) Commutative Algebra and Combinatorics. Advanced Studies in Pure Mathematics, vol. 11, pp. 93–109. North-Holland, Amsterdam (1987).
- [15] A. Higashitani and K. Matsushita, Conic divisorial ideals and non-commutative crepant resolutions of edge rings of complete multipartite graphs, *J. Algebra* **594** (2022), 685–711.
- [16] A. Higashitani and K. Matsushita, Three families of toric rings arising from posets or graphs with small class groups, *J. Pure Appl. Algebra*, **226** no. 10, (2022), 107079.
- [17] A. Higashitani and Y. Nakajima, Conic divisorial ideals of Hibi rings and their applications to non-commutative crepant resolutions, *Selecta Math.* **25** (2019), 25pp.
- [18] M. Hochster, Rings of invariants of tori, Cohen–Macaulay rings generated by monomials, and polytopes. *Ann. Math.* **96** (1972), 318–337.
- [19] V. KAIBEL and M. WOLFF, Simple 0/1-polytopes, *European J. Combinatorics* **21** (2000), 139–144.
- [20] K. Matsushita, Torsionfreeness for divisor class groups of toric rings of integral polytopes, arXiv:2202.10686.
- [21] K. Matsushita, Conic divisorial ideals of toric rings and applications to Hibi rings and stable set rings, arXiv:2210.02031.
- [22] Y. Nakajima, Non-commutative crepant resolutions of Hibi rings with small class groups, *J. Pure Appl. Algebra* **223** (2019), 3461–3484.
- [23] J. Oxley, Matroid theory, Oxford Graduate Texts in Mathematics, Vol. 21, Oxford University Press, Oxford, second edition, 2011.
- [24] H. Ohsugi and T. Hibi, Normal polytopes arising from finite graphs, *J. Algebra* **207** (1998), 409–426.
- [25] H. Ohsugi and T. Hibi, Convex polytopes all of whose reverse lexicographic initial ideals are squarefree, *Proc. Amer. Math. Soc.*, **129** (2001), No.9, 2541–2546.
- [26] A. Simis, W. V. Vasconcelos and R. H. Villarreal, The integral closure of subrings associated to graphs, *J. Algebra* **199** (1998), 281–289.
- [27] Š. Špenko and M. Van den Bergh, Non-commutative resolutions of quotient singularities for reductive groups, *Invent. Math.* **210** (2017), no. 1, 3–67.
- [28] R.P. Stanley. Hilbert functions of graded algebras. *Adv. Math.* **28** (1978), 57–83.
- [29] R. P. Stanley, Decompositions of rational convex polytopes, *Ann. Discrete Math.* **6** (1980), 333–342. MR **82a**:52007.
- [30] R. P. Stanley, Two Poset Polytopes, *Discrete Comput. Geom.* **1** (1986), 9–23.
- [31] S. Sullivant, Compressed polytopes and statistical disclosure limitation, *Tohoku Math. J.* **58** (2006), 433–445.
- [32] M. Van den Bergh, Non-Commutative Crepant Resolutions, The Legacy of Niels Henrik Abel, pp. 749–770. Springer, Berlin (2004).
- [33] R. H. Villarreal, “Monomial algebras”, Monographs and Research Notes in Mathematics. CRC Press, Boca Raton, FL, 2015.

DEPARTMENT OF PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, OSAKA UNIVERSITY, SUITA, OSAKA 565-0871, JAPAN
 Email address: k-matsushita@ist.osaka-u.ac.jp