

Some structural similarities between uncountable sets, powersets and the universe

Athanassios Tzouvaras

Department of Mathematics
Aristotle University of Thessaloniki
541 24 Thessaloniki, Greece
e-mail: tzouvara@math.auth.gr

Abstract

We establish some similarities/analogies between uncountable cardinals or powersets and the class V of all sets. They concern mainly the Boolean algebras $\mathcal{P}(\kappa)$, for a regular cardinal κ , and $\mathcal{C}(V)$ (the class of subclasses of the universe V), endowed with some ideals, especially the ideal $[\kappa]^{<\kappa}$ for $\mathcal{P}(\kappa)$, and the ideal of sets V for $\mathcal{C}(V)$.

Mathematics Subject Classification (2020): 03E05, 03E20.

Keywords: Uncountable cardinal, access ideal for a set/class, monadic second-order language of ordering.

1 Introduction

This article belongs roughly to the same line of thought as [6] and [8], which on the one hand try to shed some light to the structure of powersets of infinite sets, and on the other, at philosophical level, express a skeptical attitude towards the powerset axiom (*Pow*), that is the claim that the collection of subsets of an infinite set is a set (rather than a proper class). To the same vein belongs also [9], where it is proposed that the powerset and replacement axioms (especially the former one) would be better to be replaced by local (and thus relativized) versions of them that lead to a local version of the entire ZFC. The approach of the present article differs, however, in that we focus on the relationship and similarities between *uncountable* sets in

general, rather than powersets alone, and the universe. Since the connection between the existence of powersets and the existence of uncountable sets seems somewhat confused, let me elaborate briefly on this point before coming to the main body of the paper.

It is well-known that powersets and uncountable sets are closely related through the famous Cantor's theorem that every infinite powerset is uncountable. However this connection seems to be only one-way: as is well-known, the existence of uncountable sets is (relatively) consistent with the failure of *Pow*. For example if $M \models \text{ZFC} + 2^{\aleph_0} = \aleph_2$, then

$$H_{\omega_2}^M \models \text{ZFC}^- + \text{"}\omega_1 \text{ exists"} + \text{"}\mathcal{P}(\omega) \text{ does not exist"} ,$$

where $H_{\omega_2} = \{x : |TC(x)| < \aleph_2\}$ and $\text{ZFC}^- = \text{ZFC} - \{\text{Pow}\}$. So, if *Unc* denotes the existence of uncountable sets, $\text{ZFC}^- + \text{Unc} \not\vdash \text{Pow}$, i.e., *Pow* is *formally* independent of *Unc* over ZFC^- .

The problem is that a resident of $H_{\omega_2}^M$, in contrast to a resident of M , has no clue as to *how* the uncountable ω_1 has emerged. The resident of M has a natural way to construct ω_1 , as well as the other uncountable cardinals: they consider the collection $WO(\omega)$ of all well-orderings of ω , which is a subcollection of $\mathcal{P}(\omega \times \omega)$. Since the latter is a *set* by *Pow*, so is $WO(\omega)$ by Separation. Then taking the set $WO(\omega)/\cong$ of isomorphism types of $WO(\omega)$, they arrive at ω_1 . But for the resident of $H_{\omega_2}^M$, $\mathcal{P}(\omega \times \omega)$ is only a proper class. It is true that they can define $WO(\omega)$ in $H_{\omega_2}^M$, as a subclass of $\mathcal{P}(\omega \times \omega)$. But they cannot even define $WO(\omega)/\cong$, since each isomorphism type of a well-ordering of ω has continuum many members, so it is a proper class too. Thus ω_1 is a completely *accidental* object of $H_{\omega_2}^M$, which seems to be there by mere chance.

If, alternatively, we start with a model $N \models \text{ZFC} + 2^{\aleph_0} = \aleph_1$, then

$$H_{\omega_2}^N \models \text{ZFC}^- + \text{"}\mathcal{P}(\omega) \text{ exists"} + \text{"}\mathcal{P}(\mathcal{P}(\omega)) \text{ does not exist"} ,$$

and the problem now for a resident of $H_{\omega_2}^N$ is to explain how the uncountable set $\mathcal{P}(\omega)$ has occurred, since the operation $\mathcal{P}(x)$ is not defined for all sets.

The role of *Pow* in ZF is quite analogous to the role of exponentiation axiom (*Exp*) in (subsystems) of Peano arithmetic (PA). *Exp* is the sentence (in the language of PA) $\forall x, y \exists z (x^y = z)$, or more simply $\forall x \exists y (2^x = y)$. If $\text{I}\Sigma_n$ denotes the fragment of PA that consists of the elementary axioms of PA and the induction scheme restricted to Σ_n formulas alone, then it is well-known that *Exp* is provable in $\text{I}\Sigma_1$ but unprovable in $\text{I}\Sigma_0$. More precisely, $\text{I}\Sigma_0 < \text{I}\Sigma_0 + \text{Exp} < \text{I}\Sigma_1$ (where for theories T_1, T_2 , $T_1 < T_2$ means that T_2 is strictly stronger than T_1). Roughly, *Exp* is with respect to $\text{I}\Sigma_0$, what *Pow*

is with respect to ZF^- , and uncountable sets are analogous to “very big” numbers of arithmetic. The only difference is that uncountable sets have a precise definition (sets of size strictly larger than \aleph_0), while very big numbers do not. Intuitively a is “very big” if $a \geq 2^b$ for some “big” b . As is well-known the models of $I\Sigma_0$ are closed under $+$ and \cdot but not, in general, under exponentiation. In particular, the operation x^y is only partially defined in them. Nevertheless, there are (nonstandard) models of $I\Sigma_0 + \neg Exp$ (see e.g. [5], Theorems 4.3, 4.4) which, despite the failure of Exp , may contain 2^a for some arbitrary nonstandard number a . In the absence of the full exponentiation operation, the existence of 2^a can be seen by a resident of the model only as *accidental*, exactly as was seen the cardinal ω_1 in the model $H_{\omega_2}^M$ of ZFC^- mentioned above.

The conclusion of the preceding discussion is that the connection between powersets and the uncountable is not just one-way. It is in fact a two-way dependence, if behind every encounter with an uncountable set we seek the reason and source of its occurrence.

Having argued that Pow is, directly or indirectly, *responsible* for the existence of uncountable sets, a reasonable way to reinforce the skeptical attitude towards this axiom is to present concrete *mathematical facts* that establish specific similarities/analogies between uncountable sets and the universe of all sets V . This is what we are doing in this article. Namely, we compare the Boolean algebra $\mathcal{P}(\kappa)$ of an infinite cardinal κ , with the Boolean algebra $\mathcal{C}(V)$ of subclasses of V , and point out certain similarities. The similarities are of three kinds: (a) set-theoretic, (b) algebraic, and (c) order-theoretic. In the comparison of set-theoretic and algebraic facts some ideals play a key role, namely the ideals $[\kappa]^{<\lambda}$, for $\lambda \leq \kappa$, and in particular $[\kappa]^{<\kappa}$, for the case of cardinals, and the ideal of sets V for $\mathcal{C}(V)$. We deal with each of the three kinds of similarities in separate sections.

Summary of contents. In section 2 we deal with a specific set-theoretic similarity. The main results are the following:

(a) The ideal V of $\mathcal{C}(V)$ is *minimal* in a natural and precisely defined way. Moreover, it is the *unique* minimal ideal of this kind (Theorem 2.5).

(b) For every regular cardinal κ , the ideal $[\kappa]^{<\kappa}$ of $\mathcal{P}(\kappa)$ is *minimal* in the above sense, and also the *unique* minimal ideal of this kind (Theorem 2.6).

(c) (In sharp contrast to (b)), for every singular cardinal κ , there is *no* minimal ideal of $\mathcal{P}(\kappa)$ in the above sense (Theorem 2.11).

In section 3 we deal with the algebraic similarities. Specifically we compare the Boolean algebra $\mathcal{P}(\kappa)$ equipped with an ideal $[\kappa]^{<\lambda}$, for any

$\aleph_0 \leq \lambda \leq \kappa$, with the Boolean algebra $\mathcal{C}(V)$ equipped with the ideal V . Using Ehrenfeucht-Fraïssé games, it is shown that these structures are elementarily equivalent, that is, $\langle \mathcal{P}(\kappa), [\kappa]^{<\lambda} \rangle \equiv_{L_b(S)} \langle \mathcal{C}(V), V \rangle$, where $L_b(S)$ is the language of Boolean algebras augmented with a unary predicate $S(\cdot)$.

In section 4 we deal with the order-theoretic similarities. For convenience we identify V with the class of ordinals On by the help of von Neumann's global choice vN , and compare the first-order structures $\langle \kappa, < \rangle$ and $\langle On, < \rangle$, where $<$ is the natural well-orderings of On and κ (i.e., \in). Extending results of [2] and [7] we show that $\langle \kappa, < \rangle \equiv_{L_{ord}} \langle On, < \rangle$, where L_{ord} is the first-order language of ordering. This equivalence is further strengthened to $\langle \kappa, <, \text{Cof}(\kappa) \rangle \equiv_{\forall_{pos}^1} \langle On, <, \text{Cof}(On) \rangle$, where $\text{Cof}(\kappa)$ (resp. $\text{Cof}(On)$) denotes the class of cofinal subsets of $\langle \kappa, < \rangle$ (resp. cofinal subclasses of $\langle On, < \rangle$) and $\equiv_{\forall_{pos}^1}$ denotes elementary equivalence with respect to all positive \forall_1^1 formulas of the monadic (second-order) language L_{mon} , which is L_{ord} augmented with set variables and \in .

Some notational conventions. In order to talk about subclasses of V we need to work in a class theory (rather than ZFC), like Gödel-Bernays theory with choice GBC . So throughout our discussion will be held having as ground theory either GBC or ZFC . Concerning the collection $\mathcal{C}(V)$ of all subclasses of V when working in GBC , this is actually a “virtual”, i.e. syntactic, entity, since in order to accommodate it literally as an extensional object, we would need a third-order set theory. However this is not a serious problem. We can always treat $\mathcal{C}(V)$ in model-theoretic terms, that is, think that we work in an arbitrary model $\mathfrak{M} = \langle M, \mathcal{M} \rangle$ of GBC , where M contains the sets and \mathcal{M} contains the classes of \mathfrak{M} . Then $V^{\mathfrak{M}} = M$ while $\mathcal{C}(V)^{\mathfrak{M}} = \mathcal{M}$. Thus $\mathcal{C}(V)$ is simply the range of class variables and, like $\mathcal{P}(A)$, an atomic Boolean algebra. On the other hand, when we talk about *ideals* of $\mathcal{C}(V)$, which are collections $\mathcal{I} \subseteq \mathcal{C}(V)$, these will always be *definable* by a formula $\phi(X)$ with one free class variable.

We have already used above the notation $[\kappa]^{<\lambda}$. In general, for any set A and cardinal $\kappa \leq |A|$ we let

$$[A]^{<\kappa} = \{x \subseteq A : |x| < \kappa\}^1, \quad [A]^{\leq \lambda} = \{x \subseteq A : |x| \leq \lambda\}.$$

Also, in view of AC , we can identify every powerset $\mathcal{P}(A)$ with a cardinal 2^κ , and write $\mathcal{P}(2^\kappa)$ instead of $\mathcal{P}(\mathcal{P}(A))$, $[2^\kappa]^{<\lambda}$ instead of $[\mathcal{P}(A)]^{<\lambda}$, and so on.

¹This set is usually denoted also $\mathcal{P}_\kappa(A)$.

2 A set-theoretic similarity between V and regular cardinals: minimality and uniqueness of access ideals

Throughout upper case letters X, Y are used as class variables for the language of GBC, while lower case letters x, y denote sets (elements of classes). For two classes X, Y , let $|X| = |Y|$ denote the fact that X and Y are equipollent, i.e. there is a bijection between them. The axiom vN says that all proper classes are equipollent, or equivalently $|V| = |On|$. Following the conventions agreed in the Introduction we shall write 2^{\aleph_0} instead of $\mathcal{P}(\omega)$ and in general 2^κ instead of $\mathcal{P}(\kappa)$.

Definable classes are produced as usual by formulas $\phi(x)$ with a set variable, i.e., $X = \{x : \phi(x)\}$. But also classes of classes are permitted, provided they are defined by formulas $\phi(X)$ with a class free variable. We denote such classes by calligraphic letters \mathcal{I}, \mathcal{J} and treat them as syntactic objects, namely \mathcal{I}, \mathcal{J} can be thought of as formulas, and writing $\mathcal{I} \subseteq \mathcal{J}$ we just mean that $(\forall X)(\mathcal{I}(X) \rightarrow \mathcal{J}(X))$. For any such \mathcal{I} we have $\mathcal{I} \subseteq \mathcal{C}(V)$, where $\mathcal{C}(V)$ is the class of all subclasses of V . Given a class X , an *ideal* of $\mathcal{C}(X)$ (or an ideal on X) is a class $\mathcal{I} \subseteq \mathcal{C}(X)$ such that:

- (i) If $Y_1, Y_2 \in \mathcal{I}$, then $Y_1 \cup Y_2 \in \mathcal{I}$,
- (ii) If $Y \in \mathcal{I}$ and $Z \subseteq Y$, then $Z \in \mathcal{I}$,
- (iii) \mathcal{I} is proper, i.e., $X \notin \mathcal{I}$.

Trivially V is an ideal of $\mathcal{C}(V)$.

2.1 Accessibility of classes and sets by ideals

A (total) *preordering* on a class X is a binary relation \preceq which is reflexive, transitive and total, where the latter means that for all $x, y \in X$ $x \preceq y$ or $y \preceq x$. $x \prec y$ means $x \preceq y$ and $x \neq y$. If we add the antisymmetric property, the preordering becomes a linear ordering. To avoid trivialities we shall assume throughout that no preordered class $\langle X, \preceq \rangle$ has a last element. Given a preordering $\langle X, \preceq \rangle$ and $x \in X$, \preceq_x denotes the initial segment of X determined by x , i.e., $\preceq_x = \{y \in X : y \preceq x\}$. A $Y \subseteq X$ is said to be *bounded* w.r.t. \preceq if there is $x \in X$ such that $Y \subseteq \preceq_x$. An unbounded $Y \subseteq X$ is called also *cofinal*. For every preordering \preceq of X , the bounded subsets of X form a natural ideal of $\mathcal{C}(X)$ denoted $Seg(\preceq)$, that is

$$Seg(\preceq) = \{Y \subset X : \exists x \in X (Y \subseteq \preceq_x)\}.$$

The fact that $\langle X, \preceq \rangle$ has no last element guarantees that the ideal $Seg(\preceq)$ is proper.

Definition 2.1 (GBC or ZFC) Let \mathcal{I} be an ideal on a class or set X .

(1) \mathcal{I} is said to be an *access ideal for X* , or *X is \mathcal{I} -accessible*, if there is a preordering \preceq of X such that $\text{Seg}(\preceq) \subseteq \mathcal{I}$. In that case \preceq is said to be an *\mathcal{I} -preordering*.

(2) The ideal \mathcal{I} is said to be a *minimal access ideal*, or just *minimal for X* , if it is an access ideal for X and for any ideal $\mathcal{J} \subsetneq \mathcal{I}$, X is not \mathcal{J} -accessible.

Trivially every preordering \preceq of X is a $\text{Seg}(\preceq)$ -preordering. Notice that, although the preceding definitions are expressible in GBC or ZFC (when X and \mathcal{I} are definable objects), it is not to be meant that it is also *decidable* in these theories whether, for given X and \mathcal{I} , (a) X is \mathcal{I} -accessible or not, and (b) \mathcal{I} is minimal for X or not. These are relative facts, in general, depending on the model of GBC/ZFC we work in. For example, as we shall see below, the cardinal 2^{\aleph_0} is accessible by the ideal $[2^{\aleph_0}]^{\leq \aleph_0}$ of its countable subsets if and only if $2^{\aleph_0} = \aleph_1$. Moreover, in the latter case $[2^{\aleph_0}]^{\leq \aleph_0}$ is the unique minimal access ideal for 2^{\aleph_0} . More generally, if $2^{\aleph_0} = \aleph_\alpha$ holds in a model \mathfrak{M} and \aleph_α is regular, then $[2^{\aleph_0}]^{< \aleph_\alpha}$ is the unique minimal access ideal for 2^{\aleph_0} in \mathfrak{M} . In contrast, if \aleph_α is singular, then 2^{\aleph_0} is $[2^{\aleph_0}]^{< \aleph_\alpha}$ -accessible but there is no minimal access ideal for 2^{\aleph_0} .

On the other hand, the class V is V -accessible *independently* of the underlying model, in view of the axiom of regularity and the induced V -preordering $x \preceq_r y \Leftrightarrow \text{rank}(x) \leq \text{rank}(y)$ of the universe.² Similarly, for every *regular* cardinal κ , κ is provably $[\kappa]^{< \kappa}$ -accessible. More important are the facts shown below, that the ideals V and $[\kappa]^{< \kappa}$ are both *provably* (a) *minimal* access ideals, and (b) the *unique* minimal access ideals for V and any regular κ , respectively.

Needles to say that for V , as a well as for any cardinal κ , there is an abundance of access ideals that arise in the form of $\text{Seg}(\preceq)$ for the various preorderings \preceq of V or κ , so the proof of existence of *minimal* such ideals and, further, of *unique minimal* ones is by no means trivial. For example partition κ into finitely or infinitely many pieces and then rearrange the pieces into an arbitrary total ordering in which each piece keeps its standard ordering. Then a total ordering of κ arises and hence an access ideal for it.

According to Definition 2.1, if a class X is \mathcal{I} -accessible for an ideal \mathcal{I} within a model \mathfrak{M} , then there is a preordering \preceq of X in \mathfrak{M} such that every

²Actually this fact is the reason that in the definition of accessibility 2.1 given above we use preorderings rather than total orderings or well-orderings.

\preceq -bounded set of X belongs to \mathcal{I} , i.e.,

$$\mathfrak{M} \models (\forall Y \subseteq X)(Y \text{ is } \preceq\text{-bounded} \rightarrow Y \in \mathcal{I}). \quad (1)$$

This definition obviously allows the ideal \mathcal{I} to contain also unbounded subsets of X . If, however, \mathcal{I} is minimal, then the converse of (1) holds too. Namely the following holds.

Lemma 2.2 *Let $\mathfrak{M} \models \text{GBC}$ and let \mathcal{I} be a minimal access ideal for a definable class X in \mathfrak{M} . Then for every \mathcal{I} -preordering $\preceq \in \mathfrak{M}$ of X :*

$$\mathfrak{M} \models (\forall Y \subseteq X)(Y \text{ is } \preceq\text{-bounded} \leftrightarrow Y \in \mathcal{I}).$$

Proof. One direction follows from Definition 2.1. For the converse, assume that there is an \mathcal{I} -preordering $\preceq \in \mathfrak{M}$ of X such that \mathcal{I} contains a set $Y_0 \subseteq X$ unbounded in $\langle X, \preceq \rangle$. Then \mathcal{I} is not a minimal access ideal for X . Because X is $\text{Seg}(\preceq)$ -accessible, and $\text{Seg}(\preceq) \subsetneq \mathcal{I}$, since $Y_0 \in \mathcal{I} \setminus \text{Seg}(\preceq)$. \dashv

Minimality and uniqueness of V as an access ideal for V

Theorem 2.3 (GBC) *The class V is a minimal access ideal for V .*

Proof. By the regularity axiom of GBC, the relation $x \preceq y$ iff $\text{rank}(x) \leq \text{rank}(y)$ is a V -preordering, thus V is V -accessible in GBC. To prove the minimality of V , let \mathcal{J} be any ideal on V such that $\mathcal{J} \subsetneq V$. It suffices to see that V is not \mathcal{J} -accessible. Assume the contrary, and let \preceq_1 be a preordering of V such that for every $x \in V$, $\preceq_{1,x} \in \mathcal{J}$. Pick a $z \in V \setminus \mathcal{J}$. If z were cofinal in $\langle V, \preceq_1 \rangle$, then $V = \bigcup_{x \in z} \preceq_{1,x}$, and since for every x , $\preceq_{1,x} \in \mathcal{J} \subseteq V$, by Replacement and Union $\bigcup_{x \in z} \preceq_{1,x} = V$ would be a set, a contradiction. If, on the other hand, z were not cofinal in $\langle V, \preceq_1 \rangle$, we should have $z \subseteq \preceq_{1,x}$ for some x , so $z \in \mathcal{J}$ since $\preceq_{1,x} \in \mathcal{J}$, a contradiction again. \dashv

In fact V is not just a minimal access ideal for V , but it is the *unique* minimal access ideal. We shall need, first, the following simple consequence of the axiom of choice AC . This is rather folklore but we include its proof here for the reader's convenience.

Lemma 2.4 (GBC) *Let X be a proper class. Then for every cardinal κ there is set $b \subset X$ such that $|b| = \kappa$.*

Proof. Given the proper class X , let $R = \{\text{rank}(x) : x \in X\}$. R is a proper class of ordinals and for every such class there is in GB a strictly increasing enumeration $R = \{\alpha_\xi : \xi \in \text{On}\}$, i.e., a bijection $F : \text{On} \rightarrow R$ such that $\xi < \zeta \Leftrightarrow F(\xi) < F(\zeta)$. (This can be shown as follows: let S be the class of set functions f such that $\text{dom}(f) = \xi \in \text{On}$, $\text{rng}(f) \subset R$ and for every $\delta \in \text{dom}(f)$, $f(\delta) = \min(R \setminus \{f(\gamma) : \gamma < \delta\})$. S is defined by a normal formula so it is a class of GB. Then $F : \text{On} \rightarrow R$ is defined as follows: $F(\xi) = \alpha \Leftrightarrow (\exists f \in S)(\xi \in \text{dom}(f) \wedge f(\xi) = \alpha)$.) If $d_\xi = X \cap (V_{\alpha_{\xi+1}} \setminus V_{\alpha_\xi})$, then $\{d_\xi : \xi \in \text{On}\}$, is a proper class of nonempty disjoint sets. Let $s = \{d_\xi : \xi < \kappa\}$. s is a set of nonempty sets, so by AC it has a choice function f . If $b = f[s]$, then b is as required. \dashv

Theorem 2.5 (GBC) V is the unique minimal access ideal for V .

Proof. Towards reaching a contradiction assume that $\mathcal{I} \neq V$ is another minimal access ideal for V . Then we cannot have $V \subseteq \mathcal{I}$, for since $\mathcal{I} \neq V$, that would mean that $V \subsetneq \mathcal{I}$, which contradicts the minimality of \mathcal{I} . Thus $V \not\subseteq \mathcal{I}$. Pick a set $a \in V \setminus \mathcal{I}$ and fix an \mathcal{I} -preordering \preceq of V . (Remember that all preorderings used throughout have no last element.) It suffices to construct a preordering \preceq_1 of V such that $\text{Seg}(\preceq_1) \subsetneq \mathcal{I}$, contradicting thus the minimality of \mathcal{I} .

Notice first that the set $a \in V \setminus \mathcal{I}$ must be cofinal in $\langle V, \preceq \rangle$, otherwise $a \subseteq \preceq_x$ for some x , and hence $a \in \mathcal{I}$ since $\preceq_x \in \mathcal{I}$. Also for some $x \in a$, \preceq_x must be a proper class, otherwise $V = \bigcup_{x \in a} \preceq_x$ would be a set. Fix an element $x_0 \in a$ such that \preceq_{x_0} is a proper class and let $X = \preceq_{x_0}$. Let also $Y = V \setminus X$ be the final segment of (V, \preceq) determined by x_0 . Then we can write

$$V = X \uplus Y \tag{2}$$

for the ordered disjoint union of the initial and final segments to which x_0 splits $\langle V, \preceq \rangle$. We shall construct a preordering \preceq_1 such that if $J = \text{Seg}(\preceq_1)$, then $J \subsetneq \mathcal{I}$. Let us set $a^* = a \cap Y$. Since a is cofinal in \preceq and infinite (because \preceq does not have a last element), so is a^* . Let $|a^*| = \kappa$ and let $a^* = \{y_\xi : \xi < \kappa\}$. Using Lemma 2.4, pick a subset b of the proper class X such that $|b| = |a^*| = \kappa$ and let $b = \{z_\xi : \xi < \kappa\}$. The idea is to cut b off X and make it cofinal in the new preordering \preceq_1 , retaining \preceq to the rest parts of V . This can be obtained if we define \preceq_1 so that \preceq_1 agrees with \preceq on $V \setminus b$, while b and a^* both run cofinally to V (and to each other) with respect to \preceq_1 . So let

$$R = [\preceq \cap (V \setminus b)^2] \cup \{\langle y_\xi, z_\xi \rangle, \langle z_\xi, y_\xi \rangle : \xi < \kappa\},$$

and let \preceq_1 be the transitive closure of the binary relation R , i.e., $\langle x, z \rangle \in \preceq_1$ whenever there is y such that $\langle x, y \rangle$ and $\langle y, z \rangle$ belong to \preceq_1 . By this definition $y_\xi \preceq_1 z_\xi$ and $z_\xi \preceq_1 y_\xi$ for all $\xi < \kappa$, and \preceq_1 is a preordering of V which, in analogy to (2), splits V into the ordered union

$$V = (X \setminus b) \uplus (Y \cup b), \quad (3)$$

with $X \setminus b$, Y and b being pairwise disjoint. Moreover $a^* \subseteq Y$, hence $a^* \cap b = \emptyset$ and a^* is cofinal in $\langle V, \preceq_1 \rangle$. Let

$$\mathcal{J} = \text{Seg}(\preceq_1) = \{Z : (\exists w)(Z \subseteq \preceq_{1,w})\}.$$

We show first that $\mathcal{J} \subseteq \mathcal{I}$. Let $Z \in \mathcal{J}$ and let $Z \subseteq \preceq_{1,w}$ for some w . We shall show that $Z \in \mathcal{I}$. By the picture of V in (3) as an ordered union, Z is written as a disjoint union $Z = X_1 \cup X_2 \cup x_3$, where $X_1 \subseteq X \setminus b$, $X_2 \subseteq Y$ and $x_3 \subseteq b$. Since $X \setminus b$ and b already belong to \mathcal{I} , it follows that $X_1, x_3 \in \mathcal{I}$. It remains to show that $X_2 \in \mathcal{I}$. We have $X_2 \subseteq Y$ and $X_2 \subseteq \preceq_{1,w}$. Clearly either $w \in Y$ or $w \in b$. If $w \in Y$, then since \preceq and \preceq_1 agree on Y , we have $X_2 \subseteq \preceq_w$, so $X_2 \in \mathcal{I}$ because \preceq is an \mathcal{I} -preordering. Assume $w \in b$, so $w = z_\xi$ for some $\xi < \kappa$. By definition $z_\xi \preceq_1 y_\xi$, where $y_\xi \in a^* \subseteq Y$, i.e., $y_\xi \in Y$. Then $X_2 \subseteq \preceq_{1,z_\xi} \subseteq \preceq_{1,y_\xi}$. Since $y_\xi \in Y$ as before $X_2 \subseteq \preceq_{y_\xi}$, whence again $X_2 \in \mathcal{I}$. Thus all three pieces X_1, X_2, x_3 of Z belong to \mathcal{I} , and hence $Z \in \mathcal{I}$. This proves $\mathcal{J} \subseteq \mathcal{I}$. On the other hand $b \notin \mathcal{J}$ because it is cofinal in \preceq_1 , while $b \in \mathcal{I}$ as a subset of X which belongs to \mathcal{I} . Therefore $\mathcal{J} \subsetneq \mathcal{I}$. This contradicts the minimality of \mathcal{I} and completes the proof. \dashv

Minimality and uniqueness of access ideals on cardinals

Before considering access ideals on powersets, i.e., on cardinals 2^κ , we must consider access ideals on general uncountable cardinals. We shall see that there is a sharp difference between regular and singular cardinals with respect to this matter. We deal first with regular cardinals.

Theorem 2.6 (ZFC) *Let κ be an uncountable cardinal and $<$ be its natural well-ordering. Then:*

- (a) $\text{Seg}(<) = [\kappa]^{<\kappa}$ iff κ is regular.
- (b) If κ is regular, then $[\kappa]^{<\kappa}$ is a minimal access ideal for κ .
- (c) Moreover if κ is regular, $[\kappa]^{<\kappa}$ is the unique minimal access ideal for κ .

Proof. (a) Firstly, for every κ , $\text{Seg}(<) \subseteq [\kappa]^{<\kappa}$. Suppose κ is regular and $x \in [\kappa]^{<\kappa}$. Then x is bounded in $\langle \kappa, < \rangle$, thus $x \in \text{Seg}(<)$, so $[\kappa]^{<\kappa} \subseteq \text{Seg}(<)$.

$Seg(<)$ and therefore $Seg(<) = [\kappa]^{<\kappa}$. Conversely, if κ is not regular there is $x \subset \kappa$ cofinal in $\langle \kappa, < \rangle$ with $|x| < \kappa$. Thus $x \in [\kappa]^{<\kappa} \setminus Seg(<)$ and hence $Seg(<) \subsetneq [\kappa]^{<\kappa}$.

(b) Let κ be regular and assume $[\kappa]^{<\kappa}$ is not minimal for κ , i.e., there is an ideal $\mathcal{I} \subsetneq [\kappa]^{<\kappa}$ and an \mathcal{I} -preordering \preceq for κ . Pick $x \in [\kappa]^{<\kappa} \setminus \mathcal{I}$. Then x is cofinal in $\langle \kappa, \preceq \rangle$, because otherwise $x \subseteq \preceq_{\xi} \in \mathcal{I}$, for some $\xi \in \kappa$, and hence $x \in \mathcal{I}$, contradiction. This implies that $\kappa = \bigcup_{\xi \in x} \preceq_{\xi}$. But for each ξ , $|\preceq_{\xi}| < \kappa$, since $\preceq_{\xi} \in \mathcal{I} \subseteq [\kappa]^{<\kappa}$, and also $|x| < \kappa$. This clearly contradicts the regularity for κ .

(c) Let κ be regular. The proof that $[\kappa]^{<\kappa}$ is the unique minimal ideal for κ is quite similar to the proof of Theorem 2.5 for the corresponding fact about V , so we simply sketch it. Let $\mathcal{I} \neq [\kappa]^{<\kappa}$ be another minimal access ideal for κ and fix an \mathcal{I} -preordering \preceq of κ . If $[\kappa]^{<\kappa} \subseteq \mathcal{I}$, then $[\kappa]^{<\kappa} \subsetneq \mathcal{I}$, so \mathcal{I} would not be minimal. Therefore $[\kappa]^{<\kappa} \not\subseteq \mathcal{I}$. Pick $a \in [\kappa]^{<\kappa} \setminus \mathcal{I}$. Then, as in the proof of 2.5, a is cofinal in $\langle \kappa, \preceq \rangle$, and for some $x \in a$, $|\preceq_x| = \kappa$. Because $\kappa = \bigcup_{x \in a} \preceq_x$, and $|a| < \kappa$, so if $|\preceq_x| < \kappa$ for all $x \in \kappa$, κ would be singular. Fix such a $x_0 \in \kappa$ and let $X = \preceq_{x_0}$, so $|X| = \kappa$, $Y = \kappa \setminus X$, and $a^* = a \cap Y$. Let $|a^*| = \lambda < \kappa$, and pick $b \subset X$, with $|b| = |a^*| = \lambda$. Let also $a^* = \{y_{\xi} : \xi < \lambda\}$ and $b = \{z_{\xi} : \xi < \lambda\}$. We define a preordering \preceq_1 of κ in complete analogy to that of 2.5, making b cofinal in $\langle \kappa, \preceq_1 \rangle$, and so that κ is written

$$\kappa = (X \setminus b) \uplus (Y \cup b),$$

with $X \setminus b$, Y and b being pairwise disjoint. If $\mathcal{J} = Seg(\preceq_1)$, it is shown as before that $\mathcal{J} \subsetneq \mathcal{I}$, which contradicts the minimality of \mathcal{I} . \dashv

Now when coming to powersets, Theorem 2.6 (c) gives its place to the following variant that claims consistency rather than provability.

Theorem 2.7 (ZFC) *For any $\kappa \geq \aleph_0$ and any regular $\lambda > \kappa$, it is (relatively) consistent to believe that $[2^{\kappa}]^{<\lambda}$ is the unique minimal access ideal on 2^{κ} .*

Proof. It is well-known that the following is consistent with ZFC: $\lambda > \kappa$ is regular and $2^{\kappa} = \lambda$. This is obtained by a standard Cohen forcing extension \mathfrak{M} in which $\lambda = 2^{\kappa}$ is still regular, so it follows from 2.6 (c) that in \mathfrak{M} , $[2^{\kappa}]^{<\lambda} = [\lambda]^{<\lambda}$ is the unique minimal access ideal on 2^{κ} . \dashv

We come next to the \mathcal{I} -accessibility of singular cardinals. First recall that for every preordered set $\langle A, \preceq \rangle$, the *cofinality* of $\langle A, \preceq \rangle$, denoted $\text{cof}(A, \preceq)$, is the least cardinal λ such that: There is a set $X \subseteq A$ such that $|X| = \lambda$,

and $\langle X, \preceq \rangle$ is well-ordered and cofinal to $\langle A, \preceq \rangle$. Clearly, for every infinite set preordering $\langle A, \preceq \rangle$ having no last element, $\text{cof}(A, \preceq)$ is an infinite regular cardinal. Therefore, as an immediate consequence we have the following.

Fact 2.8 *If κ is a singular cardinal, then for every preordering \preceq of κ , there is a regular $\lambda < \kappa$ such that $\text{cof}(\kappa, \preceq) = \lambda$.*

Note that for any κ , either regular or singular, $\text{cof}(\kappa, \preceq)$ need not be equal to $\text{cf}(\kappa)$. $\text{cof}(\kappa, \preceq)$ depends heavily on \preceq and may be less or greater than $\text{cf}(\kappa)$. For example for every κ there is \preceq such that $\text{cof}(\kappa, \preceq) = \aleph_0$. On the other hand $\text{cf}(\aleph_{\omega_1}) = \aleph_1$, but we can easily find orderings of \aleph_{ω_1} with cofinalities $\aleph_n > \aleph_1$. More generally, given any regular λ and $\kappa > \lambda$, we can find a preordering \preceq_λ of κ such that $\text{cof}(\kappa, \preceq_\lambda) = \lambda$. (This can be done by splitting κ into λ (disjoint) pieces x_i , $i < \lambda$, e.g. by the help of a surjection $f : \kappa \rightarrow \lambda$, and then define \preceq_λ by totally ordering the pieces x_i so that $x_i \prec_\lambda x_j$ iff $i < j$.)

Next we show that, even when κ is singular, for every $\lambda < \kappa$, κ is not $[\kappa]^{<\lambda}$ -accessible, i.e., there is no $[\kappa]^{<\lambda}$ -preordering of κ . This is due to the following fact.

Lemma 2.9 *Let κ be an uncountable cardinal and let $\aleph_0 \leq \lambda < \kappa$. For every preordering \preceq of κ , there is an $x \in \kappa$ such that $|\preceq_x| \geq \lambda$.*

Proof. Assume for contradiction that for a given $\lambda < \kappa$, there is a preordering \preceq of κ such that $|\preceq_x| < \lambda$ for every $x \in \kappa$. For each $x \in \kappa$, let $[x] = \{y \in \kappa : x \preceq y \ \& \ y \preceq x\}$ and let $\kappa/\sim = \{[x] : x \in \kappa\}$. Then \preceq is (essentially) a linear ordering of κ/\sim . Since $|\preceq_x| < \lambda$ for all x , we have that $|[x]| < \lambda$ for all $x \in \kappa$, and therefore $|\kappa/\sim| = \kappa$. So it suffices to prove the claim for the linearly ordered set $\langle \kappa/\sim, \preceq \rangle$. For simplicity we pick an element s_x from each class $[x]$, and we set $S = \{s_x : x \in \kappa\}$. Then $|S| = \kappa$, and we prove the claim for the isomorphic set $\langle S, \preceq \rangle$. Now either S contains a cofinal subset Y of cardinality λ , or not. In the first case $S = \bigcup_{s \in Y} \preceq_s$, where $|Y| = \lambda$ and $|\preceq_s| < \lambda$, for every $s \in Y$. But then $|S| = \lambda$, a contradiction. In the other case, S does not contain any cofinal set of cardinality λ . So if we pick a $Y \subseteq S$ with $|Y| = \lambda$, then there is an $s \in S$ such that $Y \subseteq \preceq_s$. But then $|\preceq_s| \geq \lambda$, a contradiction again. \dashv

Corollary 2.10 *Let $\aleph_0 \leq \lambda < \kappa$. Then for every $\mathcal{I} \subseteq [\kappa]^{<\lambda}$, κ is not \mathcal{I} -accessible, i.e., there is no \mathcal{I} -preordering of κ .*

Proof. If \preceq were a $[\kappa]^{<\lambda}$ -preordering of κ , then $|\preceq_x| < \lambda$ for every $x \in \kappa$, which contradicts 2.9. \dashv

Next theorem is in sharp contrast to Theorem 2.6 (c).

Theorem 2.11 (ZFC) *If κ is a singular cardinal, then there is no minimal access ideal on κ . In particular, $Seg(<)$ is not minimal for κ .*

Proof. Let κ be singular. We have to show that for every preordering \preceq of κ , $Seg(\preceq)$ is not minimal. That is, there is a preordering \preceq_1 of κ such that $Seg(\preceq_1) \subsetneq Seg(\preceq)$. It is remarkable that the construction of \preceq_1 follows exactly the same steps as the corresponding constructions in the proofs of theorems 2.5 and 2.6 (c), although the latter theorems make claims to the opposite direction from that of the present one.

Fix a preordering \preceq of κ and let $\lambda = \text{cof}(\kappa, \preceq)$. By Fact 2.8, $\lambda < \kappa$. Also by Lemma 2.9 there is $x_0 \in \kappa$ with $|\preceq_{x_0}| \geq \lambda$. We construct \preceq_1 imitating the proofs of 2.5 and 2.6 (c). Namely, let $X = \preceq_{x_0}$, let $b \subseteq X$ with $|b| = \lambda$, and let $Y = \kappa \setminus X$. Pick also a set a with $|a| = \lambda$, which is well-ordered by \preceq with order-type λ , and $\langle a, \preceq \rangle$ is cofinal to $\langle \kappa, \preceq \rangle$. So we can write $a = \{\alpha_\xi : \xi < \lambda\}$, where $\alpha_\xi < \alpha_\zeta \Leftrightarrow \xi < \zeta$. Without loss of generality we may assume that $a \subseteq Y$. Let $b = \{\beta_\xi : \xi < \lambda\}$. As in the previous proofs we cut off b from X and make it cofinal to a . Namely, we let

$$R = [\preceq \cap (\kappa \setminus b)^2] \cup \{\langle \alpha_\xi, \beta_\xi \rangle, \langle \beta_\xi, \alpha_\xi \rangle : \xi < \lambda\}.$$

Also we set $\preceq_1 =$ the transitive closure of the binary relation R . Then \preceq_1 is as required. For exactly as in the proof of 2.5 we see that $Seg(\preceq_1) \subseteq Seg(\preceq)$, while on the other hand $b \in Seg(\preceq) \setminus Seg(\preceq_1)$. Thus $Seg(\preceq_1) \subsetneq Seg(\preceq)$ and this completes the proof. \dashv

As a complement to Theorem 2.7 we have the following.

Theorem 2.12 (ZFC) *For any cardinal $\kappa \geq \aleph_0$, it is consistent to believe that there is no minimal access ideal for 2^κ .*

Proof. Given $\kappa \geq \aleph_0$, it is well-known that the following is consistent with ZFC: $\kappa < \text{cf}(\lambda) < \lambda$ and $2^\kappa = \lambda$. Fixing any singular cardinal λ with $\kappa < \text{cf}(\lambda) < \lambda$, if \mathfrak{M} is a model of ZFC obtained by a κ -cc forcing which makes $2^\kappa = \lambda$, then λ remains singular in \mathfrak{M} . So it follows from Theorem 2.11 that in \mathfrak{M} no minimal access ideal exists for 2^κ . \dashv

3 Algebraic similarity of $\mathcal{P}(\kappa)$ and $\mathcal{C}(V)$ equipped with ideals

Given infinite cardinals $\lambda \leq \kappa$, let $\langle \mathcal{P}(\kappa), [\kappa]^{<\lambda} \rangle$ and $\langle \mathcal{C}(V), V \rangle$ be the Boolean algebras of all subcollections of κ and V , equipped with the ideals $[\kappa]^{<\lambda}$ and V , respectively, representing some “small sets/classes” of the algebras. The language of the structures is $L_b(S) = L_b \cup \{S\}$, where L_b is the language of Boolean algebras, and $S(\cdot)$ is a unary predicate symbol for the elements of the ideals. The purpose of this section is to prove the following.

Theorem 3.1 (GBC) *For all infinite cardinals $\lambda \leq \kappa$, $\langle \mathcal{P}(\kappa), [\kappa]^{<\lambda} \rangle \equiv_{L_b(S)} \langle \mathcal{C}(V), V \rangle$. In particular for $\kappa = 2^\lambda$ we get $\langle \mathcal{P}(2^\lambda), [2^\lambda]^{\leq \lambda} \rangle \equiv_{L_b(S)} \langle \mathcal{C}(V), V \rangle$.*

It is already known that any two powerset Boolean algebras are elementarily equivalent. This is an easy corollary of a general result concerning algebras with the same “invariants” (see [3, p. 296], §18). However the algebras here are endowed with ideals so a stronger argument is needed. The method of proof is an application of the Ehrenfeucht-Fraïssé games (see e.g. [4], pp. 52f). We recall briefly some definitions.

Let L be a first-order language and let $\mathcal{M} = \langle M, \dots \rangle$, $\mathcal{N} = \langle N, \dots \rangle$ be two L -structures. We define first what an infinite game is between \mathcal{M} , \mathcal{N} . This is a two-player game denoted $G_\omega(\mathcal{M}, \mathcal{N})$. The players, called I and II, play at stages $1, 2, \dots$, by choosing at each stage i , alternately elements $a_i \in M$ and $b_i \in N$. A play consists of ω stages. Player I may choose an element either from M or from N , and then II must respond by choosing an element from the other structure. Thus if I chooses $a_i \in M$ (resp. $b_i \in N$), then II must choose $b_i \in N$ (resp. $a_i \in M$). Player II wins the play if $f = \{\langle a_i, b_i \rangle : i = 1, 2, \dots\}$ is the graph of a partial embedding from \mathcal{M} into \mathcal{N} . Otherwise I wins. A strategy for player II in $G_\omega(\mathcal{M}, \mathcal{N})$ is a function τ such that for each i , if u_1, \dots, u_i are the first i moves of player I, then the i -th move of player II is $\tau(u_1, \dots, u_i)$. The strategy τ is a *winning* strategy for II, if for any sequence u_1, u_2, \dots of moves of player I, II wins the play if he/she follows the strategy τ .

Now an Ehrenfeucht-Fraïssé (E-F) game between \mathcal{M} , \mathcal{N} is a game of the above general form, except that it consists only of *finitely* many stages instead of infinite ones. It is denoted $G_n(\mathcal{M}, \mathcal{N})$ if it stops after the n -th stage. The winning strategy for player II or I is defined as before. E-F games can be used to characterize elementary equivalence of two structures. Specifically the following holds.

Theorem 3.2 ([4, Thm 2.4.6]) *Let L be a finite language without function symbols and let \mathcal{M}, \mathcal{N} be L -structures. Then $\mathcal{M} \equiv \mathcal{N}$ if and only if player II has a winning strategy in the game $G_n(\mathcal{M}, \mathcal{N})$, for all $n > 0$.*

We shall use 3.2 to prove Theorem 3.1. In the proof we shall need the following simple fact which is provable in GB.

Lemma 3.3 (GB) *Every proper class can be partitioned into two proper disjoint subclasses.*

Proof. Let X be a proper class and, as in Lemma 2.4, consider the class $R = \{\text{rank}(x) : x \in X\}$ which is proper and can be written $R = \{\alpha_\xi : \xi \in On\}$ in its natural well-ordering. Let $R_0 = \{\alpha_\xi : \xi \text{ is a limit ordinal}\}$, and $R_1 = \{\alpha_\xi : \xi \text{ is a successor ordinal}\}$. If for $i = 0, 1$ we set $X_i = \{x \in X : \text{rank}(x) \in R_i\}$, X_0, X_1 is the required partition. \dashv

Proof of Theorem 3.1. Fix infinite cardinals $\lambda \leq \kappa$ and let $\mathcal{M} = \langle \mathcal{P}(\kappa), [\kappa]^{<\lambda} \rangle$ and $\mathcal{N} = \langle \mathcal{C}(V), V \rangle$. (Remember that we can always think we are working in a model $\mathfrak{K} = \langle K, \mathcal{K} \rangle$ of GBC, with $V^{\mathfrak{K}} = K$ and $\mathcal{C}(V)^{\mathfrak{K}} = \mathcal{K}$, so in this case $\mathcal{N} = \langle \mathcal{K}, K \rangle$.) In order to comply with the condition of Theorem 3.2 that the language of the structures must be relational, we shall take the language L_b of Boolean algebras to be $L_b = \{\sqsubseteq\}$ (instead of the more often used algebraic language $\{+, \cdot, -, 0, 1\}$). According to Theorem 3.2, we have to show that for each n , Player II has a winning strategy in $G_n(\mathcal{M}, \mathcal{N})$. In fact we can show that player II has a winning strategy not only in the finite games $G_n(\mathcal{M}, \mathcal{N})$, but also in the infinite game $G_\omega(\mathcal{M}, \mathcal{N})$. To see this it suffices to define, for every n , what the move of player II will be after the completion of the first n stages. Now since our ground structures \mathcal{M}, \mathcal{N} are Boolean algebras, every finite subset A of each of the algebras generates a finite Boolean subalgebra *fully determined* by the set of its atoms which are the intersections of the elements of A . And whenever a player chooses a new element from the ground algebra, this element expands the previous subalgebra to a larger finite one by refining the set of its atoms. Thus in our case each stage of the game is not obtained from the previous one by the attachment of a single new pair of elements from each algebra, but of several new pairs that correspond to the generated new atoms. So we can consider the rounds of the game as partial embeddings $f_i : C_i \rightarrow D_i$, $i \geq 1$, where C_i, D_i are finite Boolean subalgebras of $\mathcal{P}(\kappa)$ and $\mathcal{C}(V)$ respectively, and for every i , $f_i \subsetneq f_{i+1}$. Moreover, f_i respect the predicate $S(\cdot)$ by preserving the ideals, i.e., satisfying the condition

(*) For every $a \in C_i$, $a \in [\kappa]^{<\lambda} \Leftrightarrow f_i(a) \in V$.

Assume that the first n rounds of the game have been completed, i.e., embeddings $f_i : C_i \rightarrow D_i$ which satisfy (*) have been defined for all $i \leq n$, and suppose player I starts the $(n + 1)$ -th round by choosing either some $a \in \mathcal{P}(\kappa)$ or some $b \in \mathcal{C}(V)$. We have to define the response of player II, i.e. a $b \in \mathcal{C}(V)$, if player I chooses $a \in \mathcal{P}(\kappa)$, or an $a \in \mathcal{P}(\kappa)$, if player I chooses $b \in \mathcal{C}(V)$, and the extension $f_{n+1} : C_{n+1} \rightarrow D_{n+1}$ of f_n so that $\langle a, b \rangle \in f_{n+1}$ and f_{n+1} satisfies (*). Without loss of generality we assume that $a \in \mathcal{P}(\kappa)$. We shall describe the definition of b and f_{n+1} , and this will be the description of the winning strategy of player II. A quite similar strategy would be described if player I had started with a $b \in \mathcal{C}(V)$ and player II should choose $a \in \mathcal{P}(\kappa)$.

The strategy of player II: Let c_1, \dots, c_k be the atoms of the algebra C_n which form a partition of κ . Then $f_n(c_1), \dots, f_n(c_k)$ are the atoms of D_n which form a partition of V . For each $i = 1, \dots, k$, let $x_i = c_i \cap a$ and $x'_i = c_i \cap (\kappa \setminus a)$ and let

$$X = \{x_i, x'_i \neq \emptyset : i = 1, \dots, k\}.$$

X is a new partition of κ which refines the partition of c_i 's and determines a finite subalgebra C_{n+1} of $\mathcal{P}(\kappa)$ such that $a \in C_{n+1} \supseteq C_n$. In order to define f_{n+1} and $f_{n+1}(a) = b$, clearly it suffices to define $f_{n+1}(x_i) = y_i$ and $f_{n+1}(x'_i) = y'_i$ for all $x_i, x'_i \in X$, so that y_i, y'_i form a suitable partition of $f_n(c_i)$. In particular we must have

$$|x_i| < \lambda \text{ (} |x'_i| < \lambda \text{)} \Leftrightarrow y_i \in V \text{ (} y'_i \in V \text{)}. \quad (4)$$

If for some i , $c_i \subseteq a$, or $c_i \subseteq \kappa \setminus a$, then $x_i = c_i$ or $x'_i = c_i$, respectively, for which f_n is already defined. So if $x_i = c_i$ or $x'_i = c_i$, we just let $f_{n+1}(x_i) = f_n(c_i)$ and $f_{n+1}(x'_i) = f_n(c_i)$. In all other cases x_i, x'_i form a proper partition of c_i and we consider the following cases with respect to the cardinalities of x_i and x'_i .

1) $|x_i|, |x'_i| \geq \lambda$. Then $|c_i| \geq \lambda$ so, by (*) for f_n , $f_n(c_i)$ is a proper class. In this case it suffices to take, using Lemma 3.3, y_i, y'_i to be complementary proper subclasses of $f_n(c_i)$.

2) $|x_i| < \lambda$ and $|x'_i| \geq \lambda$ (or vice versa). Then $|c_i| \geq \lambda$, hence $f_n(c_i)$ is a proper class. Choose $\emptyset \neq y_i \subseteq f_n(c_i)$ to be a set and let $y'_i = f_n(c_i) \setminus y_i$. In particular, if x_i is finite, we pick y_i so that $|y_i| = |x_i|$.

3) $|x_i|, |x'_i| < \lambda$. Then $|c_i| < \lambda$. Therefore $f_n(c_i)$ is a set. If x_i, x'_i are both infinite, we pick any infinite y_i, y'_i that form a partition of $f_n(c_i)$. If

either x_i or/and x'_i is finite we take the partition y_i, y'_i so that $|y_i| = |x_i|$ or/and $|y'_i| = |x'_i|$, respectively. (This is always possible, even when c_i is finite. Because the same strategy has been followed by player II at the previous stages, so if c_i is finite the strategy has yielded $f_n(c_i)$ such that $|f_n(c_i)| = |c_i|$.) Let

$$Y = \{y_i, y'_i : i = 1, \dots, k\}.$$

Y is the set of atoms of a subalgebra $D_{n+1} \supseteq D_n$ of $\mathcal{C}(V)$ and the function $f_{n+1} : C_{n+1} \rightarrow D_{n+1}$ for which $f_{n+1}(x_i) = y_i$ and $f_{n+1}(x'_i) = y'_i$ induces a partial isomorphism that extends f_n . Thus clearly f_{n+1} satisfies (4). It satisfies also condition (*) above. Because by (4), f_{n+1} already satisfies (*) for the atoms of C_{n+1} . If now x is any element of C_{n+1} and z_1, \dots, z_m are the atoms of C_{n+1} below x , then $x = z_1 \cup \dots \cup z_m$, hence $f_{n+1}(x) = f_{n+1}(z_1) \cup \dots \cup f_{n+1}(z_m)$. So clearly $|x| < \lambda$ iff $|z_i| < \lambda$ for all $i = 1, \dots, m$, iff $f_{n+1}(z_i)$ is a set for all $i = 1, \dots, m$, iff $f_{n+1}(x)$ is a set. Thus f_{n+1} satisfies (*).

This completes the description of the strategy of player II. It remains to prove that the strategy is winning, that is, all these f_i are partial embeddings from \mathcal{M} into \mathcal{N} . Namely it suffices to show the following:

Claim. Let F be the set of all finite functions from \mathcal{M} into \mathcal{N} which are defined through E-F games, in which player II follows the above described strategy. Then every $f \in F$ is a partial embedding, i.e., for every $\phi(x_1, \dots, x_n)$ of $L_b(S)$ and every $a_1, \dots, a_n \in \mathcal{P}(\kappa)$ such that $a_1, \dots, a_n \in \text{dom}(f)$,

$$\mathcal{M} \models \phi(a_1, \dots, a_n) \Leftrightarrow \mathcal{N} \models \phi(f(a_1), \dots, f(a_n)). \quad (5)$$

Proof of Claim. We prove that (5) holds for every $f \in F$ and for every ϕ by induction on the length of ϕ . The atomic formulas of $L_b(S)$ are $x \sqsubseteq y$ and $S(x)$ and clearly the ordering \sqsubseteq is preserved by all $f \in F$. Also, we saw above that every $f \in F$ satisfies condition (*), so for every $a \in \text{dom}(f)$

$$\mathcal{M} \models S(a) \Leftrightarrow a \in [\kappa]^{<\lambda} \Leftrightarrow f(a) \in V \Leftrightarrow \mathcal{N} \models S(f(a)).$$

The induction steps for the connectives are straightforward for all $f \in F$, and the only nontrivial step of the induction is the quantifier step. So given a $\phi(x, x_1, \dots, x_n)$ we assume that all $f \in F$ satisfy (5) for ϕ , and prove this for $\exists x \phi(x, x_1, \dots, x_n)$, i.e., for all a_1, \dots, a_n in $\text{dom}(f)$

$$\mathcal{M} \models \exists x \phi(x, a_1, \dots, a_n) \Leftrightarrow \mathcal{N} \models \exists x \phi(x, f(a_1), \dots, f(a_n)). \quad (6)$$

Suppose $\mathcal{M} \models \exists x \phi(x, a_1, \dots, a_n)$ for given a_1, \dots, a_n in $\text{dom}(f)$. Then $\mathcal{M} \models \phi(a, a_1, \dots, a_n)$ for some $a \in \mathcal{P}(\kappa)$. Now, by definition, f is the n -th

stage of some E-F game between \mathcal{M} and \mathcal{N} , so a can be the move of player I at the $(n+1)$ -th stage of the game. Then player II responds following the above strategy by picking a $b \in \mathcal{C}(V)$ and extending f to $g \supseteq f \cup \{\langle a, b \rangle\}$ so that $g \in F$. By our induction assumption g satisfies (5) for ϕ , hence $\mathcal{N} \models \phi(g(a), g(a_1), \dots, g(a_n))$, or $\mathcal{N} \models \phi(b, f(a_1), \dots, f(a_n))$, therefore $\mathcal{N} \models \exists x \phi(x, f(a_1), \dots, f(a_n))$. This proves “ \Rightarrow ” of (6). For the converse assume $\mathcal{N} \models \exists x \phi(x, f(a_1), \dots, f(a_n))$ and $\mathcal{N} \models \phi(b, f(a_1), \dots, f(a_n))$ for a $b \in \mathcal{C}(V)$. As before b can also be the move of player I at the $(n+1)$ -th stage, and then player II responds by choosing an $a \in \mathcal{P}(\kappa)$ and $g \supseteq f \cup \{\langle a, b \rangle\}$ so that g belongs to F . Then $\mathcal{N} \models \phi(g(a), g(a_1), \dots, g(a_n))$ and by the induction assumption $\mathcal{M} \models \phi(a, a_1, \dots, a_n)$, whence $\mathcal{M} \models \exists x \phi(a_1, \dots, a_n)$. This proves the Claim and completes the proof of the Theorem. \dashv

The similarity of structures $\langle \mathcal{P}(\kappa), [\kappa]^{<\lambda} \rangle$ and $\langle \mathcal{C}(V), V \rangle$ expressed by Theorem 3.1 is of a rather algebraic character, since it is about Boolean algebras equipped with ideals. However it can be transcribed into one with more set-theoretic flavor, i.e., one using the language of set theory. Specifically, the elements of the ground set κ and class V can be treated as *atoms* (or *urelements*), while $\mathcal{P}(\kappa)$ and $\mathcal{C}(V)$ as collections of sets/classes of atoms. So let $L_1(S)$ be the language consisting of \in , $S(\cdot)$, variables p, q, \dots for urelements, and variables x, y, \dots for sets. The atomic formulas of $L_1(S)$ are: $p = q$, $p \in x$, and $S(x)$. The intended $L_1(S)$ -structures are $\langle \kappa, \mathcal{P}(\kappa), [\kappa]^{<\lambda} \rangle$, for sets, and $\langle V, \mathcal{C}(V), V \rangle$ for the universe. However a technical problem here is the fact that the two kinds of entities, atoms and sets, should be disjoint, while $\kappa \cap \mathcal{P}(\kappa) \neq \emptyset$ (and similarly for V). We can bypass this obstacle by working with suitable copies of κ and V . Namely we set

$$\kappa^* = \{\langle x, 0 \rangle : x \in \kappa\} \text{ and } V^* = \{\langle x, 0 \rangle : x \in V\}.$$

Then $\kappa^* \cap \mathcal{P}(\kappa^*) = \emptyset$ and $V^* \cap \mathcal{C}(V^*) = \emptyset$. Under this coding the $L_b(S)$ structures $\langle \mathcal{P}(\kappa), [\kappa]^{<\lambda} \rangle$ and $\langle \mathcal{C}(V), V \rangle$ become isomorphic to $\langle \mathcal{P}(\kappa^*), [\kappa^*]^{<\lambda} \rangle$ and $\langle \mathcal{C}(V^*), \mathcal{P}(V^*) \rangle$, respectively, where $\mathcal{P}(V^*)$ is the class of subsets of V^* . Then in view of Theorem 3.1 we have: For all infinite cardinals $\lambda \leq \kappa$

$$\langle \mathcal{P}(\kappa^*), [\kappa^*]^{<\lambda} \rangle \equiv_{L_b(S)} \langle \mathcal{C}(V^*), \mathcal{P}(V^*) \rangle. \quad (7)$$

Based on (7) we can easily deduce the following.

Theorem 3.4 *For all infinite $\lambda \leq \kappa$,*

$$\langle \kappa^*, \mathcal{P}(\kappa^*), [\kappa^*]^{<\lambda} \rangle \equiv_{L_1(S)} \langle V^*, \mathcal{C}(V^*), \mathcal{P}(V^*) \rangle.$$

Proof. Define an interpretation $\phi \mapsto \phi^+$ from the atomic formulas of $L_1(S)$ to formulas of $L_b(S)$ as follows: Fix enumerations $p_n, x_n, n \in \mathbb{N}$, of the variables of $L_1(S)$ for atoms and sets, respectively, and an enumeration $y_n, n \in \mathbb{N}$, of the variables of $L_b(S)$. The atomic formulas of $L_1(S)$ are $p_i = p_j, p_i \in x_j$ and $S(x_j)$. Define

- (a) $(p_i = p_j)^+ := (y_{2i} = y_{2j})$,
- (b) $(p_i \in x_j)^+ := \text{Atom}(y_{2i}) \wedge y_{2i} \sqsubseteq y_{2j+1}$,
- (c) $S(x_j)^+ = S(y_{2j+1})$,

where $\text{Atom}(y)$ is the formula of $L_b(S)$ saying that “ y is an atom”. This interpretation is also extended to all formulas of $L_1(S)$ as expected. (As follows from clauses (a)-(c) above, the variable p_i translates to y_{2i} while x_j translates to y_{2j+1} . This translation determines also of course the interpretation of quantifiers.) Then it is easy to see that for all $\phi \in L_1(S)$,

$$\langle \kappa^*, \mathcal{P}(\kappa^*), [\kappa^*]^{<\lambda} \rangle \models \phi \Leftrightarrow \langle \mathcal{P}(\kappa^*), [\kappa^*]^{<\lambda} \rangle \models \phi^+,$$

and

$$\langle V^*, \mathcal{C}(V^*), \mathcal{P}(V^*) \rangle \models \phi \Leftrightarrow \langle \mathcal{C}(V^*), \mathcal{P}(V^*) \rangle \models \phi^+.$$

By (7),

$$\langle \mathcal{P}(\kappa^*), [\kappa^*]^{<\lambda} \rangle \models \phi^+ \Leftrightarrow \langle \mathcal{C}(V^*), \mathcal{P}(V^*) \rangle \models \phi^+,$$

therefore

$$\langle \kappa^*, \mathcal{P}(\kappa^*), [\kappa^*]^{<\lambda} \rangle \models \phi \Leftrightarrow \langle V^*, \mathcal{C}(V^*), \mathcal{P}(V^*) \rangle \models \phi.$$

–

In fact Theorems 3.1 and 3.4 are essentially equivalent, since a proof of 3.1 from 3.4 can also be obtained by defining a converse interpretation $\phi \mapsto \phi'$ from atomic formulas of $L_b(S)$ to formulas of $L_1(S)$ as follows:

- $(x \sqsubseteq y)' := \forall p(p \in x \rightarrow p \in y)$,
- $(x = y)' := (x = y)$, and
- $S(x)' := S(x)$.

This interpretation, extended to all formulas, yields: For every ϕ of $L_b(S)$,

$$\langle \mathcal{P}(\kappa^*), [\kappa^*]^{<\lambda} \rangle \models \phi \Leftrightarrow \langle \kappa^*, \mathcal{P}(\kappa^*), [\kappa^*]^{<\lambda} \rangle \models \phi'$$

and

$$\langle \mathcal{C}(V^*), \mathcal{P}(V^*) \rangle \models \phi \Leftrightarrow \langle V^*, \mathcal{C}(V^*), \mathcal{P}(V^*) \rangle \models \phi'.$$

Thus 3.4 implies 3.1.

4 Order-theoretic similarity of uncountable cardinals with the class of ordinals

By the axiom of choice every uncountable set A can be identified with an uncountable cardinal κ , and by the axiom of global choice vN the universe V can be identified with the class of ordinals On . However these identifications *add structure* to A and V , namely the well-orderings of κ and On , which we denote by the same symbol $<$. Admitting the naturalness of choice principles, it is then reasonable to compare the first-order structures $\langle \kappa, < \rangle$, $\langle \lambda, < \rangle$ for uncountable κ, λ , as well as $\langle \kappa, < \rangle$ and $\langle On, < \rangle$. In this section we shall show how the main result of [7] (Theorem 2.1), which says that the monadic \forall_1^1 positive theory of linear ordering is the same for all uncountable cardinals, can be extended also over $\langle On, < \rangle$.

The starting point of [7] was the following result, which is a special case of [2, Cor. 44]:

Theorem 4.1 ([7, Thm 1.1]) *For all uncountable cardinals κ, λ , $\langle \kappa, < \rangle \equiv_{L_{ord}} \langle \lambda, < \rangle$, where $L_{ord} = \{<\}$ is the first-order language of ordering.*

Remember that the ordering $<$ of ordinals is a well-ordering, that is a second-order property, while the equivalence $\equiv_{L_{ord}}$ refers only to the first-order properties of $<$. Theorem 4.1 is in fact a consequence of Corollary 44 of [2] and, essentially, the same argument that we used in order to justify it in [7] can be applied also to justify the following extension.

Theorem 4.2 *For every uncountable cardinal κ , $\langle \kappa, < \rangle \equiv_{L_{ord}} \langle On, < \rangle$.*

Proof. (Sketch) By Corollary 44 of [2], for any ordinals α, β , $\langle \alpha, < \rangle \equiv \langle \beta, < \rangle$ if and only if α, β are *congruent modulo ω^ω* , i.e., there are ξ, η, δ such that $\delta < \omega^\omega$, $\alpha = \omega^\omega \cdot \xi + \delta$, $\beta = \omega^\omega \cdot \eta + \delta$ and either $\xi = \eta = 0$ or $\xi \neq 0$ and $\eta \neq 0$.³ Since for all cardinals $\kappa, \lambda > \omega$, $\kappa = \omega^\omega \cdot \kappa$ and $\lambda = \omega^\omega \cdot \lambda$, it follows that κ, λ are congruent modulo ω^ω . This already proves Theorem 4.1.

Concerning the justification of the present claim, without any risk of fallacy we can treat On in this particular situation as the “greatest (limit) ordinal”, and extend the usual (right) multiplication of ordinals over $On \cup \{On\}$ in the obvious way. Specifically, just as for any α and any limit β we

³Important notice: when I was preparing [7], Professor Doner had kindly informed me in a private communication, that the definition of *congruence modulo ω^ω* as given in [2], p. 51, is mistaken. The correct definition is that given above and can be found in [1, p. 6].

have $\alpha \cdot \beta = \sup\{\alpha \cdot \gamma : \gamma < \beta\}$, simply define $\alpha \cdot On = \sup\{\alpha \cdot \gamma : \gamma < On\}$. Then trivially $\omega^\omega \cdot On = On$, so for every uncountable cardinal κ , κ and On are congruent modulo ω^ω . Applying again Corollary 44 of [2] to $\langle \kappa, < \rangle$ and $\langle On, < \rangle$, we are done. (As for the details why the relation of congruence modulo ω^ω entails the elementary equivalence of well-ordered sets or classes, one should consult [2], especially the theory \mathcal{W} (Def. 2) of “weak well-orderings”, the notion of “canonical order type” (Def. 39), and Theorem 41 which relates these notions with elementarily equivalent ordinals.) \dashv

We proceed next to extend the main theorem of [7], which establishes a similarity significantly stronger than that of Theorem 4.1, so as to hold between every uncountable cardinal and On . We need first some terminology and notation.

We first extend the language of ordering $L_{ord} = \{<\}$ used previously for the structures $\langle \kappa, < \rangle$ (with lower-case individual variables x, y, z, \dots) to the *monadic* (second-order) language L_{mon} which is L_{ord} augmented with \in and upper-case set variables X, Y, Z, \dots . A formula of L_{mod} without quantified set variables is called *normal*. A formula of the form $\forall X\phi$ (resp. $\exists X\phi$), where ϕ is normal is called a \forall_1^1 (resp. \exists_1^1) formula. The standard interpretations of \mathcal{L}_{mon} are the structures $\langle A, <, \mathcal{P}(A) \rangle$ where $\langle A, < \rangle$ is an ordered set that interprets the formulas of L_{ord} , and $\mathcal{P}(A)$ is the range of set variables. However there are more general interpretations. They are of the form $\langle A, <, \mathcal{A} \rangle$, where $\mathcal{A} \subset \mathcal{P}(A)$. \mathcal{A} may be some class of “small” sets, i.e., an ideal of $\mathcal{P}(A)$, or, in the opposite direction, a class of “large” sets. Most often these large sets are the sets cofinal in $\langle A, < \rangle$, so let us set

$$\text{Cof}(A) = \{X \subseteq A : X \text{ is cofinal in } \langle A, < \rangle\}.$$

A can be also a proper class and \mathcal{A} be a class of subclasses of A . When we deal with an ordered collection which is a proper class, like On , then, in order to interpret the set variables X, Y , we must work again in a class theory. Then as usual we write $\mathcal{C}(On)$ for the collection of all subclasses of On . In particular below we shall deal with the analogue of $\text{Cof}(\kappa)$ above, which is

$$\text{Cof}(On) = \{X \subseteq On : X \text{ is cofinal in } \langle On, < \rangle\}$$

and consists of the proper subclasses of On .

Definition 4.3 Let $\phi(X)$ be a normal formula of \mathcal{L}_{mon} with at most one (free) set variable X . $\phi(X)$ is said to be *positive in X* (or just *positive*) if it belongs to the smallest class of formulas which (a) contains all formulas

of L_{ord} , (b) contains the atomic formulas $x \in X$ and (c) is closed under the logical operations \wedge , \vee , and the first-order quantifiers \exists and \forall .

The main result of [7] is that the structures $\langle \kappa, <, \text{Cof}(\kappa) \rangle$, for all uncountable cardinals κ , are elementarily equivalent with respect to \forall_1^1 positive formulas of L_{mod} .

Theorem 4.4 ([7, Thm. 2.1]) *In ZFC the following holds: For all uncountable cardinals κ, λ ,*

$$\langle \kappa, <, \text{Cof}(\kappa) \rangle \equiv_{pos}^{\forall_1^1} \langle \lambda, <, \text{Cof}(\lambda) \rangle,$$

i.e., for every positive $\phi(X)$,

$$\langle \kappa, <, \text{Cof}(\kappa) \rangle \models (\forall X)\phi(X) \Leftrightarrow \langle \lambda, <, \text{Cof}(\lambda) \rangle \models (\forall X)\phi(X).$$

Here we strengthen 4.4 to the following.

Theorem 4.5 (GBC) *For every uncountable cardinal κ ,*

$$\langle \kappa, <, \text{Cof}(\kappa) \rangle \equiv_{pos}^{\forall_1^1} \langle On, <, \text{Cof}(On) \rangle.$$

Proof. (Sketch) The proof goes exactly along the lines of the proof of 4.4 and is based on the *syntactic form* of positive (normal) formulas of L_{mod} which has been discovered by Y. Moschovakis. Namely, every formula $\phi(X)$ positive in X is equivalent over any structure $\langle A, <, \mathcal{P}(A) \rangle$, for all $X \neq A$, to a formula having prenex form $(\overline{Q}\overline{w})(\forall u)(\theta(\overline{w}, u) \rightarrow u \in X)$, where $\overline{w} = \langle w_1, \dots, w_n \rangle$ is a string of variables, $\overline{Q} = \langle Q_1, \dots, Q_n \rangle$ a string of quantifiers and $\theta(\overline{w}, u)$ is quantifier-free. (Note that, since $\phi(X)$ does not contain set quantifiers, the above prenex form for $\phi(X)$ is valid also over every structure $\langle A, <, \mathcal{A} \rangle$, with $\mathcal{A} \subseteq \mathcal{P}(A)$.) In view of the preceding syntactic characterization of \forall_1^1 positive formulas, the proof splits into several cases that correspond to the various forms of the string $\overline{Q}\overline{w}$, as well as to the forms of θ as a Boolean combination of atoms and negated atoms. Then, through an exhaustive inspection of all these cases, it is shown that we can essentially reduce the truth of $(\forall X)\phi(X)$ to the truth of a *first-order* formula. But this means, in view of Theorem 4.2, that $(\forall X)\phi(X)$ is absolute between $\langle \kappa, <, \text{Cof}(\kappa) \rangle$ and $\langle On, <, \text{Cof}(On) \rangle$. \dashv

If we work in $\text{GB} + vN$, where a bijection between On and V exists, the well-ordering of On is transferred to a well-ordering of V so the following holds.

Corollary 4.6 *In GB + vN it is provable that there is a well-ordering \prec of V such that for every uncountable cardinal κ ,*

$$\langle \kappa, <, \text{Cof}(\kappa) \rangle \equiv_{\text{pos}}^{\forall_1^1} \langle V, \prec, \text{Cof}(V) \rangle.$$

Concerning the optimality of Theorem 4.5 with respect to the indistinguishability of structures $\langle \kappa, <, \text{Cof}(\kappa) \rangle$ and $\langle On, <, \text{Cof}(On) \rangle$, and the analogous question about structures $\langle \kappa, <, \text{Cof}(\kappa) \rangle$ and $\langle \lambda, <, \text{Cof}(\lambda) \rangle$ for uncountable κ, λ , the reader is referred to the corresponding discussion in [7]: It is not known whether $\langle \kappa, <, \text{Cof}(\kappa) \rangle$ and $\langle \lambda, <, \text{Cof}(\lambda) \rangle$ are indistinguishable with respect to all \forall_1^1 and \exists_1^1 formulas. What we know is that the structures $\langle \omega_1, <, \text{Cof}(\omega_1) \rangle$ and $\langle \omega_2, <, \text{Cof}(\omega_2) \rangle$ can be distinguished by a sentence ϕ_{ω_1} of the form $(\forall x)(\exists X)\psi$, where ψ is normal but not positive (see section 5 of [7]). This is easily generalized to all ω_m, ω_n . A fortiori ϕ_{ω_1} distinguishes also the structures $\langle \omega_1, <, \text{Cof}(\omega_1) \rangle$ and $\langle On, <, \text{Cof}(On) \rangle$.

Acknowledgement My thanks go to an anonymous referee whose remarks helped me fix some unclarities and improve considerably the presentation of the article.

References

- [1] J.E. Doner, Definability in extended arithmetic of ordinal numbers, *Dissertationes Math. (Rozprawy Mat.)* **96** (1972), 50 pp.
- [2] J.E. Doner, A. Mostowski and A. Tarski, The elementary theory of well-ordering – a metamathematical study, in: *Logic Colloquium '77*, A. Macintyre, L. Pacholski and J. Paris (eds.), North Holland, 1978, pp. 1-54.
- [3] S. Koppelberg, Metamathematics (of Boolean algebras), in: *Handbook of Boolean Algebras*, J.D. Monk and R. Bonnet (eds.), vol. 1, Elsevier Science Publishers B.V. 1989, pp. 287-307.
- [4] D. Marker, *Model Theory: An Introduction*, Springer, 2002.
- [5] R. Parikh, Existence and feasibility in arithmetic, *J. Symb. Logic* **36** (1971), no. 3, 494-508.
- [6] A. Tzouvaras, Positive set-operators of low complexity, *Math. Log. Quar.* **49** (2003), no. 3, 284-292.

- [7] A. Tzouvaras, Uncountable cardinals have the same monadic \forall_1^1 -positive theory over large sets, *Fund. Math.* **181** (2004), no. 2, 125-142.
- [8] A. Tzouvaras, What is so special with the power set operation?, *Arch. Math. Logic* **43** (2004), no. 6, 723-737.
- [9] A. Tzouvaras, Localizing the axioms, *Arch. Math. Logic* **49** (2010), no. 5, 571-601.