

# Integer group determinants for $C_4 \rtimes C_4$

Yuka Yamaguchi and Naoya Yamaguchi

March 31, 2023

## Abstract

Let  $C_4$  be the cyclic group of order 4. We determine all possible values of the integer group determinant of  $C_4 \rtimes C_4$ .

## 1 Introduction

For a finite group  $G$ , let  $\Theta_G(x_g)$  be the group determinant of  $G$  and let

$$S(G) := \{\Theta_G(x_g) \mid x_g \in \mathbb{Z}\}.$$

In this paper, we determine  $S(C_4 \rtimes C_4)$ , where

$$C_4 \rtimes C_4 := \langle g_1, g_2 \mid g_1^4 = g_2^4 = e, g_2g_1 = g_1^3g_2 \rangle = \{g_1^s g_2^t \mid 0 \leq s, t \leq 3\}.$$

Here,  $e$  denotes the unit element.

**Theorem 1.1.** *Let*

$$P := \{p \mid p \equiv -3 \pmod{8} \text{ is a prime number}\},$$

$$P' := \{p \mid p \equiv -1 \pmod{4} \text{ is a prime number}\},$$

$$A := \{(8k-3)(8l-3)(4m-1)^2 \mid k, m \in \mathbb{Z}, m \geq 1, 8l-3 \in P, k+l \equiv m \pmod{2}\} \\ \subsetneq \{16m-7 \mid m \in \mathbb{Z}\}.$$

*Then we have*

$$S(C_4 \rtimes C_4) = \{16m+1, 2^{14}p(2m+1), 2^{14}q^2(2m+1), 2^{15}m \mid m \in \mathbb{Z}, p \in P, q \in P'\} \cup A.$$

We recall the definition of the group determinant. For a finite group  $G$ , let  $x_g$  be a variable for each  $g \in G$  and let  $\mathbb{Z}[x_g]$  be the multivariate polynomial ring in  $x_g$  over  $\mathbb{Z}$ . The group determinant  $\Theta_G(x_g)$  of  $G$  was defined by Dedekind as follows:

$$\Theta_G(x_g) := \det (x_{gh^{-1}})_{g,h \in G} \in \mathbb{Z}[x_g].$$

The irreducible factorization of  $\Theta_G(x_g)$  over  $\mathbb{C}$  was given by Frobenius [2]: *Let  $G$  be a finite group and let  $\widehat{G}$  be a complete set of representatives of the equivalence classes of irreducible representations of  $G$  over  $\mathbb{C}$ . Then*

$$\Theta_G(x_g) = \prod_{\varphi \in \widehat{G}} \det \left( \sum_{g \in G} \varphi(g)x_g \right)^{\deg \varphi},$$

where  $\deg \varphi$  is the degree of  $\varphi$ . This is called Frobenius' theorem.

A group determinant called an integer group determinant when its variables are integers. At the meeting of the American Mathematical Society in Hayward, California, in April 1977, Olga Taussky-Todd [15] asked whether one could characterize the values of the integer group determinant [12, p.1]. For some groups,  $S(G)$  was determined in [1, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 19, 20, 18, 21, 22]. As a result, for every group  $G$  of order at most 15, a complete description was obtained for  $S(G)$ . Also, for all five abelian groups of order 16, and the non-abelian groups  $D_{16}, D_8 \times C_2, Q_8 \times C_2, Q_{16}, C_2^2 \rtimes C_4$ ,  $S(G)$  have been determined. There are fourteen groups of order 16 up to isomorphism [3, 23]. Theorem 1.1 leaves three unresolved groups of order 16.

## 2 Preliminaries

For any  $\bar{r} \in C_4$  with  $r \in \{0, 1, 2, 3\}$ , we denote the variable  $x_{\bar{r}}$  by  $x_r$ , and let

$$D_4(x_0, x_1, x_2, x_3) := \det (x_{gh^{-1}})_{g,h \in C_4}.$$

For any  $(\bar{r}, \bar{s}) \in C_4 \times C_2$  with  $r \in \{0, 1, 2, 3\}$  and  $s \in \{0, 1\}$ , we denote the variable  $y_{(\bar{r}, \bar{s})}$  by  $y_j$ , where  $j := r + 4s$ , and let

$$D_{4 \times 2}(y_0, y_1, \dots, y_7) := \det (y_{gh^{-1}})_{g,h \in C_4 \times C_2}.$$

From the  $H = C_4$  and  $K = C_2$  case of [17, Theorem 1.1], we have the following corollary.

**Corollary 2.1.** *We have*

$$D_{4 \times 2}(y_0, y_1, \dots, y_7) = D_4(y_0 + y_4, y_1 + y_5, y_2 + y_6, y_3 + y_7) D_4(y_0 - y_4, y_1 - y_5, y_2 - y_6, y_3 - y_7).$$

Let  $G := C_4 \rtimes C_4$ . For any  $g = g_1^s g_2^t \in G$  ( $0 \leq s, t \leq 3$ ), we denote the variable  $z_g$  by  $z_j$ , where  $j := t + 4s$ , and let

$$D_G(z_0, z_1, \dots, z_{15}) := \det (z_{gh^{-1}})_{g,h \in G}.$$

Let  $[x] := \max \{n \in \mathbb{Z} \mid n \leq x\}$  and  $\widehat{G} := \{\varphi_k \mid 0 \leq k \leq 9\}$ , where

$$\varphi_k(g_1) := (-1)^{\lfloor \frac{k}{4} \rfloor}, \quad \varphi_k(g_2) := \sqrt{-1}^k$$

for any  $0 \leq k \leq 7$  and

$$\varphi_k(g_1) := (-1)^k \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \varphi_k(g_2) := \sqrt{-1}^k \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

for any  $8 \leq k \leq 9$ . Then,  $\widehat{G}$  is a complete set of representatives of the equivalence classes of irreducible representations of  $G$  over  $\mathbb{C}$ . Let  $F$  be the eight variable polynomial defined by

$$F(w_0, w_1, \dots, w_7) := f_0(w_0 - w_2, w_4 - w_6, w_1 - w_3, w_5 - w_7) f_1(w_0 + w_2, w_4 + w_6, w_1 + w_3, w_5 + w_7),$$

where  $f_k(x, y, z, w) := x^2 + y^2 + (-1)^k z^2 + (-1)^k w^2$  ( $0 \leq k \leq 1$ ). Then we have the following.

**Lemma 2.2.** *We have*

$$D_G(z_0, z_1, \dots, z_{15}) = D_{4 \times 2}(z_0 + z_8, z_1 + z_9, \dots, z_7 + z_{15}) F(z_0 - z_8, z_1 - z_9, \dots, z_7 - z_{15})^2.$$

*Proof.* From Frobenius' theorem, we have

$$D_G(z_0, z_1, \dots, z_{15}) = \prod_{\varphi \in \widehat{G}} \det \left( \sum_{g \in G} \varphi(g) z_g \right)^{\deg \varphi} = \prod_{k=0}^9 \det M_k^{\deg \varphi_k},$$

where  $M_k := \sum_{s=0}^3 \sum_{t=0}^3 \varphi_k(g_1^s g_2^t) z_{t+4s}$ . Note that  $\{\chi_k \mid 0 \leq k \leq 7\}$  is the dual group of  $C_4 \times C_2$ , where  $\chi_k((\bar{1}, \bar{0})) := \sqrt{-1}^k$  and  $\chi_k((\bar{0}, \bar{1})) := (-1)^{\lfloor \frac{k}{4} \rfloor}$  for  $0 \leq k \leq 7$ . Then we have

$$\begin{aligned} \prod_{k=0}^7 \det M_k^{\deg \varphi_k} &= \prod_{k=0}^7 M_k \\ &= \prod_{k=0}^7 \sum_{s=0}^3 \sum_{t=0}^3 (-1)^{\lfloor \frac{k}{4} \rfloor s} \sqrt{-1}^{kt} z_{t+4s} \\ &= \prod_{k=0}^7 \sum_{t=0}^3 \sum_{s=0}^1 \sqrt{-1}^{kt} (-1)^{\lfloor \frac{k}{4} \rfloor s} (z_{t+4s} + z_{t+4(s+2)}) \\ &= \prod_{k=0}^7 \sum_{t=0}^3 \sum_{s=0}^1 \chi_k((\bar{t}, \bar{s})) (z_{t+4s} + z_{t+4s+8}) \\ &= D_{4 \times 2}(z_0 + z_8, z_1 + z_9, \dots, z_7 + z_{15}). \end{aligned}$$

Also, for any  $8 \leq k \leq 9$ , we have

$$\begin{aligned}
M_k &= \sum_{s=0}^3 \sum_{t=0}^3 (-1)^{ks} \sqrt{-1}^{kt} z_{t+4s} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}^s \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^t \\
&= \sum_{t=0}^3 \sum_{s=0}^1 \sqrt{-1}^{kt} (-1)^{ks} (z_{t+4s} - z_{t+4s+8}) \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}^s \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^t \\
&= \sum_{t=0}^3 \left\{ \sqrt{-1}^{kt} (z_t - z_{t+8}) \begin{pmatrix} 1 & 0 \\ 0 & (-1)^t \end{pmatrix} + \sqrt{-1}^{kt} (-1)^k (z_{t+4} - z_{t+12}) \begin{pmatrix} 0 & (-1)^{t+1} \\ 1 & 0 \end{pmatrix} \right\} \\
&= \begin{pmatrix} \sum_{t=0}^3 \sqrt{-1}^{kt} \tilde{z}_t & \sum_{t=0}^3 (-1)^{k+t+1} \sqrt{-1}^{kt} \tilde{z}_{t+4} \\ \sum_{t=0}^3 (-1)^k \sqrt{-1}^{kt} \tilde{z}_{t+4} & \sum_{t=0}^3 (-1)^t \sqrt{-1}^{kt} \tilde{z}_t \end{pmatrix},
\end{aligned}$$

where  $\tilde{z}_t := z_t - z_{t+8}$  for  $0 \leq t \leq 7$ . Therefore,

$$\begin{aligned}
\det M_k &= \sum_{s=0}^3 \sqrt{-1}^{ks} \tilde{z}_s \sum_{t=0}^3 (-1)^t \sqrt{-1}^{kt} \tilde{z}_t + \sum_{s=0}^3 \sqrt{-1}^{ks} \tilde{z}_{s+4} \sum_{t=0}^3 (-1)^t \sqrt{-1}^{kt} \tilde{z}_{t+4} \\
&= \{\tilde{z}_0 + (-1)^k \tilde{z}_2\}^2 - (-1)^k \{\tilde{z}_1 + (-1)^k \tilde{z}_3\}^2 \\
&\quad + \{\tilde{z}_4 + (-1)^k \tilde{z}_6\}^2 - (-1)^k \{\tilde{z}_5 + (-1)^k \tilde{z}_7\}^2.
\end{aligned}$$

That is,

$$\det M_8 = f_1(\tilde{z}_0 + \tilde{z}_2, \tilde{z}_4 + \tilde{z}_6, \tilde{z}_1 + \tilde{z}_3, \tilde{z}_5 + \tilde{z}_7), \quad \det M_9 = f_0(\tilde{z}_0 - \tilde{z}_2, \tilde{z}_4 - \tilde{z}_6, \tilde{z}_1 - \tilde{z}_3, \tilde{z}_5 - \tilde{z}_7).$$

This completes the proof.  $\square$

Throughout this paper, we assume that  $a_0, a_1, \dots, a_{15} \in \mathbb{Z}$  and let

$$\begin{aligned}
b_i &:= (a_i + a_{i+8}) + (a_{i+4} + a_{i+12}) & (0 \leq i \leq 3), \\
c_i &:= (a_i + a_{i+8}) - (a_{i+4} + a_{i+12}) & (0 \leq i \leq 3), \\
d_i &:= a_i - a_{i+8} & (0 \leq i \leq 7).
\end{aligned}$$

Also, let

$$\mathbf{a} := (a_0, a_1, \dots, a_{15}), \quad \mathbf{b} := (b_0, b_1, b_2, b_3), \quad \mathbf{c} := (c_0, c_1, c_2, c_3), \quad \mathbf{d} := (d_0, d_1, \dots, d_7).$$

Then, from Corollary 2.1 and Lemma 2.2, we have

$$\begin{aligned}
D_G(\mathbf{a}) &= D_{4 \times 2}(a_0 + a_8, a_1 + a_9, \dots, a_7 + a_{15}) F(a_0 - a_8, a_1 - a_9, \dots, a_7 - a_{15})^2 \\
&= D_4(\mathbf{b}) D_4(\mathbf{c}) F(\mathbf{d})^2.
\end{aligned}$$

**Remark 2.3.** For any  $0 \leq i \leq 3$ , we have  $b_i \equiv c_i \equiv d_i + d_{i+4} \pmod{2}$ .

**Lemma 2.4.** The following hold:

$$(1) D_4(\mathbf{b}) \equiv b_0 + b_1 + b_2 + b_3 \pmod{2};$$

$$(2) D_4(\mathbf{c}) \equiv c_0 + c_1 + c_2 + c_3 \pmod{2};$$

$$(3) F(\mathbf{d}) \equiv d_0 + d_1 + \cdots + d_7 \pmod{2}.$$

*Proof.* We obtain (1) and (2) from the following: for any  $x_0, x_1, x_2, x_3 \in \mathbb{Z}$ ,

$$\begin{aligned} D_4(x_0, x_1, x_2, x_3) &= \{(x_0 + x_2)^2 - (x_1 + x_3)^2\} \{(x_0 - x_2)^2 + (x_1 - x_3)^2\} \\ &\equiv (x_0^2 + x_2^2 + x_1^2 + x_3^2)^2 \\ &\equiv x_0 + x_1 + x_2 + x_3 \pmod{2}. \end{aligned}$$

We prove (3). Since  $f_k(x, y, z, w) = x^2 + y^2 + (-1)^k z^2 + (-1)^k w^2 \equiv x + y + z + w \pmod{2}$  for any  $0 \leq k \leq 1$  and  $x, y, z, w \in \mathbb{Z}$ , we have

$$\begin{aligned} F(\mathbf{d}) &= f_0(d_0 - d_2, d_4 - d_6, d_1 - d_3, d_5 - d_7) f_1(d_0 + d_2, d_4 + d_6, d_1 + d_3, d_5 + d_7) \\ &\equiv (d_0 + d_1 + \cdots + d_7)^2 \\ &\equiv d_0 + d_1 + \cdots + d_7 \pmod{2}. \end{aligned}$$

□

From Remark 2.3 and Lemma 2.4, we have the following lemma.

**Lemma 2.5.** *We have  $D_G(\mathbf{a}) \equiv D_4(\mathbf{b}) \equiv D_4(\mathbf{c}) \equiv F(\mathbf{d}) \pmod{2}$ .*

### 3 Impossible odd numbers

In this section, we consider impossible odd numbers. Let  $\mathbb{Z}_{\text{odd}}$  be the set of all odd numbers. Let  $P := \{p \mid p \equiv -3 \pmod{8} \text{ is a prime number}\}$  and let

$$A := \{(8k - 3)(8l - 3)(4m - 1)^2 \mid k, m \in \mathbb{Z}, m \geq 1, 8l - 3 \in P, k + l \equiv m \pmod{2}\}.$$

**Lemma 3.1.** *We have  $S(C_4 \rtimes C_4) \cap \mathbb{Z}_{\text{odd}} \subset \{16m + 1 \mid m \in \mathbb{Z}\} \cup A$ .*

Let

$$\begin{aligned} n_0 &:= f_0(d_0 - d_2, d_4 - d_6, d_1 - d_3, d_5 - d_7), & n_1 &:= f_1(d_0 + d_2, d_4 + d_6, d_1 + d_3, d_5 + d_7), \\ d &:= (d_0 + d_2)(d_4 + d_6) + (d_1 + d_3)(d_5 + d_7), & d^* &:= d_0 d_2 + d_4 d_6 + d_1 d_3 + d_5 d_7. \end{aligned}$$

To prove Lemma 3.1, we use the following two remarks and four lemmas.

**Remark 3.2.** *From [21, Lemma 2.10 (1)], we have  $b_0 b_2 + b_1 b_3 + c_0 c_2 + c_1 c_3 \equiv 2d^* \pmod{4}$ .*

**Remark 3.3.** Let  $\beta := (d_0 + d_2 + \sqrt{-1}d_4 + \sqrt{-1}d_6)^2 - (d_1 + d_3 + \sqrt{-1}d_5 + \sqrt{-1}d_7)^2$ . Suppose that  $b_0 + b_2 \not\equiv b_1 + b_3 \pmod{2}$ . Then,  $d_0 + d_2 + d_4 + d_6 \not\equiv d_1 + d_3 + d_5 + d_7 \pmod{2}$ . Thus, from [21, Lemma 2.8 (1)], we have

$$\begin{aligned} \beta\bar{\beta} \equiv -3 \pmod{8} &\iff \text{Exactly one of } d_0 + d_2, d_4 + d_6, d_1 + d_3, d_5 + d_7 \text{ is even} \\ &\iff d = (d_0 + d_2)(d_4 + d_6) + (d_1 + d_3)(d_5 + d_7) \equiv 1 \pmod{2}. \end{aligned}$$

Therefore, the assumption  $\beta\bar{\beta} \equiv -3 \pmod{8}$  in [21, Lemma 4.4] can be replaced by  $d \equiv 1 \pmod{2}$ .

**Lemma 3.4.** If  $b_0 + b_2 \not\equiv b_1 + b_3 \pmod{2}$ , then  $D_4(\mathbf{b})D_4(\mathbf{c}) \equiv 1 - 8d^* \pmod{16}$ .

Lemma 3.4 is immediately obtained from [21, Lemma 4.3] and Remark 3.2.

**Lemma 3.5.** Suppose that  $b_0 + b_2 \not\equiv b_1 + b_3$ ,  $d \equiv 1 \pmod{2}$ . Then the following hold:

(1) If  $d^* \equiv 0 \pmod{2}$ , then

$$D_4(\mathbf{b})D_4(\mathbf{c}) \in \{(8k - 3)(8l - 3) \mid k \in \mathbb{Z}, 8l - 3 \in P, k \not\equiv l \pmod{2}\};$$

(2) If  $d^* \equiv 1 \pmod{2}$ , then

$$D_4(\mathbf{b})D_4(\mathbf{c}) \in \{(8k - 3)(8l - 3) \mid k \in \mathbb{Z}, 8l - 3 \in P, k \equiv l \pmod{2}\}.$$

Lemma 3.5 is immediately obtained from [21, Lemma 4.4] and Remarks 3.2 and 3.3.

**Lemma 3.6.** If  $b_0 + b_2 \not\equiv b_1 + b_3 \pmod{2}$ , then  $F(\mathbf{d})^2 \equiv 1 - 8d^* - 8d \pmod{16}$ .

*Proof.* Let  $b_0 + b_2 \not\equiv b_1 + b_3 \pmod{2}$ . Then

$$d_0 + d_2 + d_4 + d_6 \not\equiv d_1 + d_3 + d_5 + d_7 \pmod{2}.$$

Therefore, from  $n_0 \equiv n_1 \equiv 1 \pmod{2}$ , we have  $n_0^2 \equiv n_1^2 \equiv 1 \pmod{8}$ . Also, from

$$\begin{aligned} n_0 + n_1 &= 2(d_0 + d_2 + d_4 + d_6)^2 - 4(d_0 + d_2)(d_4 + d_6) - 4d^*, \\ n_0 - n_1 &= 2(d_1 + d_3 + d_5 + d_7)^2 - 4(d_1 + d_3)(d_5 + d_7) - 4d^*, \end{aligned}$$

we have

$$\begin{aligned} n_0^2 - n_1^2 &= (n_0 + n_1)(n_0 - n_1) \\ &\equiv 8(d_0 + d_2 + d_4 + d_6)^2(d_1 + d_3)(d_5 + d_7) \\ &\quad + 8(d_1 + d_3 + d_5 + d_7)^2(d_0 + d_2)(d_4 + d_6) \\ &\quad + 8\{(d_0 + d_2 + d_4 + d_6)^2 + (d_1 + d_3 + d_5 + d_7)^2\}d^* \\ &\equiv 8(d_0 + d_2 + d_4 + d_6)(d_1 + d_3)(d_5 + d_7) \\ &\quad + 8(d_1 + d_3 + d_5 + d_7)(d_0 + d_2)(d_4 + d_6) + 8d^* \\ &\equiv 8\{(d_0 + d_2 + d_4 + d_6) + (d_1 + d_3 + d_5 + d_7)\}d \\ &\quad - 8(d_0 + d_2 + d_4 + d_6)(d_0 + d_2)(d_4 + d_6) \\ &\quad - 8(d_1 + d_3 + d_5 + d_7)(d_1 + d_3)(d_5 + d_7) + 8d^* \\ &\equiv 8d + 8d^* \pmod{16}. \end{aligned}$$

From the above,  $F(\mathbf{d})^2 = n_0^2 n_1^2 \equiv n_0^2(n_0^2 - 8d - 8d^*) \equiv 1 - 8d - 8d^* \pmod{16}$ .  $\square$

**Lemma 3.7.** *Suppose that  $b_0 + b_2 \not\equiv b_1 + b_3$ ,  $d \equiv 1 \pmod{2}$ . Then we have the following:*

(1) *If  $d^* \equiv 0 \pmod{2}$ , then*

$$F(\mathbf{d})^2 \in \{(4m-1)^2(8n+1) \mid m, n \in \mathbb{Z}, m \geq 1, m \not\equiv n \pmod{2}\};$$

(2) *If  $d^* \equiv 1 \pmod{2}$ , then*

$$F(\mathbf{d})^2 \in \{(4m-1)^2(8n+1) \mid m, n \in \mathbb{Z}, m \geq 1, m \equiv n \pmod{2}\}.$$

*Proof.* From  $b_0 + b_2 \not\equiv b_1 + b_3$ ,  $d \equiv 1 \pmod{2}$ , exactly one of  $d_0 + d_2$ ,  $d_4 + d_6$ ,  $d_1 + d_3$ ,  $d_5 + d_7$  is even. Therefore, we have  $n_0 \equiv -1 \pmod{4}$  and  $n_1 \equiv 1 \pmod{2}$ . From  $n_0 \geq 0$ , there exists  $m \geq 1$  satisfying  $n_0 = 4m - 1$ . Also, from  $n_1^2 \equiv 1 \pmod{8}$ , there exists  $n \in \mathbb{Z}$  satisfying  $n_1^2 = 8n + 1$ . Therefore,  $F(\mathbf{d})^2 = n_0^2 n_1^2 = (4m-1)^2(8n+1) \equiv 8(m+n) + 1 \pmod{16}$ . On the other hand, from Lemma 3.6, we have  $F(\mathbf{d})^2 \equiv 1 - 8d^* - 8d \equiv 9 - 8d^* \pmod{16}$ . From the above, the lemma is proved.  $\square$

*Proof of Lemma 3.1.* Suppose that  $D_G(\mathbf{a}) = D_4(\mathbf{b})D_4(\mathbf{c})F(\mathbf{d})^2 \in \mathbb{Z}_{\text{odd}}$ . Then  $D_4(\mathbf{b})$  is odd. Thus, from Lemma 2.4 (1),  $b_0 + b_2 \not\equiv b_1 + b_3 \pmod{2}$ . Therefore, from Lemmas 3.4 and 3.6,

$$D_4(\mathbf{b})D_4(\mathbf{c})F(\mathbf{d})^2 \equiv (1 - 8d^*)(1 - 8d^* - 8d) \equiv 1 - 8d \pmod{16}.$$

From this, if  $d \equiv 0 \pmod{2}$ , then  $D_G(\mathbf{a}) \in \{16m + 1 \mid m \in \mathbb{Z}\}$ . On the other hand, if  $d \equiv 1 \pmod{2}$ , then  $D_G(\mathbf{a}) \in A'$  from Lemmas 3.5 and 3.7, where

$$A' := \{(8k-3)(8l-3)(4m-1)^2(8n+1) \mid k, m, n \in \mathbb{Z}, m \geq 1, \\ 8l-3 \in P, k+l \equiv m+n \pmod{2}\}$$

Since  $A = A'$  holds, the lemma is proved.  $\square$

## 4 Impossible even numbers

In this section, we consider impossible even numbers. Let

$$P' := \{p \mid p \equiv -1 \pmod{4} \text{ is a prime number}\}.$$

**Lemma 4.1.** *The following hold:*

- (1)  $S(\mathbb{C}_4 \rtimes \mathbb{C}_4) \cap 2\mathbb{Z} \subset 2^{14}\mathbb{Z}$ ;
- (2)  $S(\mathbb{C}_4 \rtimes \mathbb{C}_4) \cap 2^{14}\mathbb{Z}_{\text{odd}} \subset \{2^{14}p(2m+1), 2^{14}q^2(2m+1) \mid m \in \mathbb{Z}, p \in P, q \in P'\}$ .

To prove Lemma 4.1, we use the following three lemmas.

**Lemma 4.2.** *The following hold:*

(1) If  $b_0 + b_2 \equiv b_1 + b_3 \equiv 0 \pmod{2}$ , then

$$D_4(\mathbf{b})D_4(\mathbf{c}) \in \begin{cases} 2^{11}\mathbb{Z}, & b_0 + b_1 + b_2 + b_3 \not\equiv c_0 + c_1 + c_2 + c_3 \pmod{4}, \\ 2^8\mathbb{Z}, & b_0 + b_1 + b_2 + b_3 \equiv c_0 + c_1 + c_2 + c_3 \pmod{4}; \end{cases}$$

(2) If  $b_0 + b_2 \equiv b_1 + b_3 \equiv 1 \pmod{2}$ , then

$$D_4(\mathbf{b})D_4(\mathbf{c}) \in \begin{cases} 2^8\mathbb{Z}_{\text{odd}}, & (b_0 + b_2)(b_1 + b_3) \equiv \pm 3, (c_0 + c_2)(c_1 + c_3) \equiv \pm 3 \pmod{8}, \\ 2^9\mathbb{Z}, & \text{otherwise.} \end{cases}$$

Lemma 4.2 is immediately obtained from [21, Lemma 5.2].

**Lemma 4.3.** *The following hold:*

(1) If  $b_0 + b_2 \equiv b_1 + b_3 \equiv 0 \pmod{2}$ , then

$$F(\mathbf{d}) \in \begin{cases} 2^2\mathbb{Z}_{\text{odd}}, & b_0 + b_1 + b_2 + b_3 \not\equiv c_0 + c_1 + c_2 + c_3 \pmod{4}, \\ 2^4\mathbb{Z}, & b_0 + b_1 + b_2 + b_3 \equiv c_0 + c_1 + c_2 + c_3 \pmod{4}; \end{cases}$$

(2) If  $b_0 + b_2 \equiv b_1 + b_3 \equiv 1 \pmod{2}$ , then

$$F(\mathbf{d}) \in \begin{cases} 2^3\mathbb{Z}_{\text{odd}}, & d \equiv 2 \pmod{4}, \\ 2^4\mathbb{Z}, & d \equiv 0 \pmod{4}, \end{cases}$$

where  $d := (d_0 + d_2)(d_4 + d_6) + (d_1 + d_3)(d_5 + d_7)$ .

*Proof.* We prove (1). Let  $b_0 + b_2 \equiv b_1 + b_3 \equiv 0 \pmod{2}$ . Then,

$$d_0 + d_2 + d_4 + d_6 \equiv d_1 + d_3 + d_5 + d_7 \equiv 0 \pmod{2}.$$

Also, we have

$$\begin{aligned} b_0 + b_1 + b_2 + b_3 &\equiv c_0 + c_1 + c_2 + c_3 \pmod{4} \\ \iff b_0 + b_2 + c_0 + c_2 &\equiv b_1 + b_3 + c_1 + c_3 \pmod{4} \\ \iff d_0 + d_2 &\equiv d_1 + d_3 \pmod{2}. \end{aligned}$$

Therefore, if  $b_0 + b_1 + b_2 + b_3 \not\equiv c_0 + c_1 + c_2 + c_3 \pmod{4}$ , then  $d_0 + d_2 \equiv d_4 + d_6 \not\equiv d_1 + d_3 \equiv d_5 + d_7 \pmod{2}$ . From this,  $n_0 \equiv n_1 \equiv 2 \pmod{4}$  holds. Thus we have  $F(\mathbf{d}) = n_0 n_1 \in 2^2\mathbb{Z}_{\text{odd}}$ . On the other hand, if  $b_0 + b_1 + b_2 + b_3 \equiv c_0 + c_1 + c_2 + c_3 \pmod{4}$ , then  $d_0 + d_2 \equiv d_4 + d_6 \equiv d_1 + d_3 \equiv d_5 + d_7 \pmod{2}$ . From this,  $n_0 \equiv n_1 \equiv 0 \pmod{4}$  holds. Thus we have  $F(\mathbf{d}) = n_0 n_1 \in 2^4\mathbb{Z}$ . We prove (2). Let  $b_0 + b_2 \equiv b_1 + b_3 \equiv 1 \pmod{2}$ . Then,

$$d_0 + d_2 + d_4 + d_6 \equiv d_1 + d_3 + d_5 + d_7 \equiv 1 \pmod{2}.$$

Thus we have  $n_0 \equiv 2 \pmod{4}$ . Also, we have

$$\begin{aligned} n_1 &= (d_0 + d_2 + d_4 + d_6)^2 - (d_1 + d_3 + d_5 + d_7)^2 - 2(d_0 + d_2)(d_4 + d_6) + 2(d_1 + d_3)(d_5 + d_7) \\ &\equiv 2d \pmod{8}. \end{aligned}$$

Therefore, if  $d \equiv 2 \pmod{4}$ , then  $F(\mathbf{d}) = n_0 n_1 \in 2^3 \mathbb{Z}_{\text{odd}}$ , and if  $d \equiv 0 \pmod{4}$ , then  $F(\mathbf{d}) = n_0 n_1 \in 2^4 \mathbb{Z}$ .  $\square$

**Lemma 4.4.** *If  $D_G(\mathbf{a}) \in 2^{14} \mathbb{Z}_{\text{odd}}$ , then*

$$D_G(\mathbf{a}) \in \{2^{14} p(2m+1), 2^{14} q^2(2m+1) \mid m \in \mathbb{Z}, p \in P, q \in P'\}.$$

*Proof.* From Lemmas 4.2 and 4.3, we have

$$b_0 + b_2 \equiv b_1 + b_3 \equiv 1 \pmod{2}, \quad D_4(\mathbf{b})D_4(\mathbf{c}) \in 2^8 \mathbb{Z}_{\text{odd}}, \quad F(\mathbf{d}) \in 2^3 \mathbb{Z}_{\text{odd}}.$$

That is,

$$(b_0 + b_2)(b_1 + b_3) \equiv 1 \pmod{2}, \quad (c_0 + c_2)(c_1 + c_3) \equiv \pm 3 \pmod{8}, \quad d \equiv 2 \pmod{4},$$

where  $d := (d_0 + d_2)(d_4 + d_6) + (d_1 + d_3)(d_5 + d_7)$ . We divide the proof into the following cases:

- (i)  $(b_0 b_3 + b_2 b_1, c_0 c_3 + c_2 c_1) \equiv (0, 0), (0, 2) \text{ or } (2, 0) \pmod{4}$ ;
- (ii)  $(b_0 b_3 + b_2 b_1, c_0 c_3 + c_2 c_1) \equiv (2, 2) \pmod{4}$ ;
- (iii)  $(b_0 b_1 + b_2 b_3, c_0 c_1 + c_2 c_3) \equiv (0, 0), (0, 2) \text{ or } (2, 0) \pmod{4}$ ;
- (iv)  $(b_0 b_1 + b_2 b_3, c_0 c_1 + c_2 c_3) \equiv (2, 2) \pmod{4}$ .

First, we consider the case (i). Suppose that  $b_0 b_3 + b_2 b_1 \equiv 0 \pmod{4}$ . Then

$$(b_0 - b_2)(b_1 - b_3) = (b_0 + b_2)(b_1 + b_3) - 2(b_0 b_3 + b_2 b_1) \equiv \pm 3 \pmod{8}.$$

Thus,  $(b_0 - b_2)^2 + (b_1 - b_3)^2 \equiv 10 \pmod{16}$ . It implies that  $(b_0 - b_2)^2 + (b_1 - b_3)^2$  has at least one prime factor of the form  $8k - 3$ . That is,  $D_4(\mathbf{b})$  has at least one prime factor of the form  $8k - 3$ . In the same way, we can prove that  $D_4(\mathbf{c})$  has at least one prime factor of the form  $8k - 3$  when  $c_0 c_3 + c_2 c_1 \equiv 0 \pmod{4}$ . From the above, in the case (i), it holds that  $D_4(\mathbf{b})D_4(\mathbf{c})$  has at least one prime factor of the form  $8k - 3$ . We can obtain the same conclusion for the case (iii). Next, we consider the case (ii). We prove that  $F(\mathbf{d})$  has at least one prime factor of the form  $4k - 1$ . From  $b_0 + b_2 \equiv b_1 + b_3 \equiv 1 \pmod{2}$ , we have  $d_0 + d_2 + d_4 + d_6 \equiv d_1 + d_3 + d_5 + d_7 \equiv 1 \pmod{2}$ . Also, since  $F(\mathbf{d})$  is invariant under the each replacing  $(d_0, d_2) \leftrightarrow (d_4, d_6)$  and  $(d_1, d_3) \leftrightarrow (d_5, d_7)$ , we may assume without loss of generality that  $d_0 + d_2 \equiv d_1 + d_3 \equiv 0, d_4 + d_6 \equiv d_5 + d_7 \equiv 1 \pmod{2}$ . Then, we have

$$\begin{aligned} (d_0 - d_2) + (d_1 - d_3) &\equiv (d_0 - d_2)(d_5 - d_7) + (d_1 - d_3)(d_4 - d_6) \\ &\equiv (d_0 + d_2)(d_5 + d_7) + (d_1 + d_3)(d_4 + d_6) \\ &\quad + 2(d_0 d_7 + d_2 d_5 + d_4 d_3 + d_6 d_1) \\ &\equiv d + (b_0 b_3 + b_2 b_1) - (c_0 c_3 + c_2 c_1) \\ &\equiv 2 \pmod{4} \end{aligned}$$

from [21, Lemma 2.10 (2)]. Therefore,  $n_0 \equiv 6 \pmod{8}$  holds. From this and  $n_0 \neq -2$ , it holds that  $n_0$  has at least one prime factor of the form  $4k - 1$ . We can obtain the same conclusion for the case (iv).  $\square$

*Proof of Lemma 4.1.* Suppose that  $D_G(\mathbf{a}) = D_4(\mathbf{b})D_4(\mathbf{c})F(\mathbf{d})^2 \in 2\mathbb{Z}$ . Then, from Lemmas 2.4 (1) and 2.5, we have  $b_0 + b_2 \equiv b_1 + b_3 \pmod{2}$ . Therefore, from Lemmas 4.2 and 4.3, we obtain (1). Also, from Lemma 4.4, we obtain (2).  $\square$

## 5 Possible integers

In this section, we determine all possible integers. Lemmas 3.1 and 4.1 imply that  $S(C_4 \rtimes C_4)$  does not include every integer that is not mentioned in the following Lemmas 5.1–5.4.

**Lemma 5.1.** *For any  $m \in \mathbb{Z}$ , the following are elements of  $S(C_4 \rtimes C_4)$ :*

- (1)  $16m + 1$ ;
- (2)  $2^{15}(2m + 1)$ ;
- (3)  $2^{16}m$ .

**Lemma 5.2.** *For any  $k, m \in \mathbb{Z}$  with  $m \geq 1$  and  $16l - 3, 16l + 5 \in P$ , the following are elements of  $S(C_4 \rtimes C_4)$ :*

- (1)  $(16k - 3)(16l - 3)(8m - 1)^2$ ;
- (2)  $(16k + 5)(16l + 5)(8m - 1)^2$ ;
- (3)  $(16k + 5)(16l - 3)(8m + 3)^2$ ;
- (4)  $(16k - 3)(16l + 5)(8m + 3)^2$ .

**Lemma 5.3.** *For any  $m \in \mathbb{Z}$  and  $p \in P$ , we have  $2^{14}p(2m + 1) \in S(C_4 \rtimes C_4)$ .*

**Lemma 5.4.** *For any  $m \in \mathbb{Z}$  and  $q \in P'$ , we have  $2^{14}q^2(2m + 1) \in S(C_4 \rtimes C_4)$ .*

*Proof of Lemma 5.1.* We obtain (1) from

$$\begin{aligned}
 D_G(m + 1, m, m, \dots, m) &= D_{4 \times 2}(2m + 1, 2m, 2m, \dots, 2m)F(1, 0, 0, \dots, 0)^2 \\
 &= D_4(4m + 1, 4m, 4m, 4m)D_4(1, 0, 0, 0)f_0(1, 0, 0, 0)^2f_1(1, 0, 0, 0)^2 \\
 &= (8m + 1)^2 - (8m)^2 \\
 &= 16m + 1.
 \end{aligned}$$

We obtain (2) from

$$\begin{aligned}
& D_G(m+1, m+1, m+1, m+1, m, m+1, m+1, m, m+1, m, m+1, m, m, m, m, m) \\
&= D_{4 \times 2}(2m+2, 2m+1, 2m+2, 2m+1, 2m, 2m+1, 2m+1, 2m) F(0, 1, 0, 1, 0, 1, 1, 0)^2 \\
&= D_4(4m+2, 4m+2, 4m+3, 4m+1) D_4(2, 0, 1, 1) f_0(0, -1, 0, 1)^2 f_1(0, 1, 2, 1)^2 \\
&= 2 \{ (8m+5)^2 - (8m+3)^2 \} \cdot 16 \cdot 2^2 \cdot (-4)^2 \\
&= 2^{15}(2m+1).
\end{aligned}$$

We obtain (3) from

$$\begin{aligned}
& D_G(m+1, m, m+1, m, m, m, m, m, m, m, m, m-1, m, m, m-1) \\
&= D_{4 \times 2}(2m+1, 2m, 2m+1, 2m, 2m-1, 2m, 2m, 2m-1) F(1, 0, 1, 0, 1, 0, 0, 1)^2 \\
&= D_4(4m, 4m, 4m+1, 4m-1) D_4(2, 0, 1, 1) f_0(0, 1, 0, -1)^2 f_1(2, 1, 0, 1)^2 \\
&= 2 \{ (8m+1)^2 - (8m-1)^2 \} \cdot 16 \cdot 2^2 \cdot 4^2 \\
&= 2^{16}m.
\end{aligned}$$

□

*Proof of Lemma 5.2.* Note that the following hold:

- (i) For any  $16l-3 \in P$ , there exist  $r, s \in \mathbb{Z}$  satisfying  $16l-3 = (8r+3)^2 + (8s+2)^2$ ;
- (ii) For any  $16l+5 \in P$ , there exist  $r, s \in \mathbb{Z}$  satisfying  $16l+5 = (8r+1)^2 + (8s+2)^2$ ;
- (iii) For any  $m \geq 1$ , there exist  $t, u, v, w \in \mathbb{Z}$  satisfying

$$8m-1 = (4t+1)^2 + (4u+1)^2 + (4v+1)^2 + (4w+2)^2;$$

- (iv) For any  $m \geq 1$ , there exist  $t, u, v, w \in \mathbb{Z}$  satisfying

$$8m+3 = (4t+1)^2 + (4u+1)^2 + (4v+1)^2 + (4w)^2.$$

Let

$$\begin{aligned}
a_0 &= k+r+w+1, & a_1 &= k+s+u, & a_2 &= k-r-w-1, & a_3 &= k-s-u-1, \\
a_4 &= k+r+v, & a_5 &= k+s+t, & a_6 &= k-r-v-1, & a_7 &= k-s-t-1, \\
a_8 &= k+r-w, & a_9 &= k+s-u, & a_{10} &= k-r+w, & a_{11} &= k-s+u, \\
a_{12} &= k+r-v, & a_{13} &= k+s-t, & a_{14} &= k-r+v, & a_{15} &= k-s+t.
\end{aligned}$$

Then we obtain (1) from (i), (iii) and

$$D_G(\mathbf{a}) = (16k-3) \{ (8r+3)^2 + (8s+2)^2 \} \{ (4w+2)^2 + (4v+1)^2 + (4u+1)^2 + (4t+1)^2 \}^2.$$

Let

$$\begin{aligned}
a_0 &= k - r + w + 1, & a_1 &= k - s + u, & a_2 &= k + r - w, & a_3 &= k + s - u, \\
a_4 &= k - r + v, & a_5 &= k - s + t, & a_6 &= k + r - v, & a_7 &= k + s - t, \\
a_8 &= k - r - w, & a_9 &= k - s - u, & a_{10} &= k + r + w + 1, & a_{11} &= k + s + u + 1, \\
a_{12} &= k - r - v, & a_{13} &= k - s - t, & a_{14} &= k + r + v + 1, & a_{15} &= k + s + t + 1.
\end{aligned}$$

Then we obtain (2) from (ii), (iii) and

$$D_G(\mathbf{a}) = (16k + 5) \{(8r + 1)^2 + (8s + 2)^2\} \{(4w + 2)^2 + (4v + 1)^2 + (4u + 1)^2 + (4t + 1)^2\}^2.$$

Let

$$\begin{aligned}
a_0 &= k + r + w + 1, & a_1 &= k + s + u + 1, & a_2 &= k - r - w, & a_3 &= k - s - u, \\
a_4 &= k + r + v + 1, & a_5 &= k + s + t + 1, & a_6 &= k - r - v, & a_7 &= k - s - t, \\
a_8 &= k + r - w + 1, & a_9 &= k + s - u, & a_{10} &= k - r + w, & a_{11} &= k - s + u, \\
a_{12} &= k + r - v, & a_{13} &= k + s - t, & a_{14} &= k - r + v, & a_{15} &= k - s + t.
\end{aligned}$$

Then we obtain (3) from (i), (iv) and

$$D_G(\mathbf{a}) = (16k + 5) \{(8r + 3)^2 + (8s + 2)^2\} \{(4w)^2 + (4v + 1)^2 + (4u + 1)^2 + (4t + 1)^2\}^2.$$

Let

$$\begin{aligned}
a_0 &= k - r + w, & a_1 &= k - s + u, & a_2 &= k + r - w, & a_3 &= k + s - u, \\
a_4 &= k - r + v, & a_5 &= k - s + t, & a_6 &= k + r - v, & a_7 &= k + s - t, \\
a_8 &= k - r - w, & a_9 &= k - s - u - 1, & a_{10} &= k + r + w, & a_{11} &= k + s + u, \\
a_{12} &= k - r - v - 1, & a_{13} &= k - s - t - 1, & a_{14} &= k + r + v, & a_{15} &= k + s + t.
\end{aligned}$$

Then we obtain (4) from (ii), (iv) and

$$D_G(\mathbf{a}) = (16k - 3) \{(8r + 1)^2 + (8s + 2)^2\} \{(4w)^2 + (4v + 1)^2 + (4u + 1)^2 + (4t + 1)^2\}^2.$$

□

*Proof of Lemma 5.3.* For any  $p \in P$ , there exist  $k, l \in \mathbb{Z}$  satisfying  $2p = (8k + 3)^2 + (8l + 1)^2$ .

Let

$$\begin{aligned}
a_0 &= m + k + 1, & a_1 &= m + l, & a_2 &= m - k, & a_3 &= m - l, \\
a_4 &= m - k + 1, & a_5 &= m - l, & a_6 &= m + k + 1, & a_7 &= m + l + 1, \\
a_8 &= m + k, & a_9 &= m + l, & a_{10} &= m - k - 1, & a_{11} &= m - l, \\
a_{12} &= m - k, & a_{13} &= m - l, & a_{14} &= m + k + 1, & a_{15} &= m + l.
\end{aligned}$$

Then we have

$$D_G(\mathbf{a}) = 2 \{(8m + 3)^2 - (8m + 1)^2\} \cdot 8 \{(8k + 3)^2 + (8l + 1)^2\} \cdot 2^2 \cdot 4^2 = 2^{14}p(4m + 1).$$

Let

$$\begin{aligned} a_0 &= m + k, & a_1 &= m + l, & a_2 &= m - k - 1, & a_3 &= m - l, \\ a_4 &= m - k, & a_5 &= m - l, & a_6 &= m + k + 1, & a_7 &= m + l, \\ a_8 &= m + k, & a_9 &= m + l - 1, & a_{10} &= m - k - 1, & a_{11} &= m - l - 1, \\ a_{12} &= m - k, & a_{13} &= m - l - 1, & a_{14} &= m + k, & a_{15} &= m + l. \end{aligned}$$

Then we have

$$D_G(\mathbf{a}) = 2 \{(8m - 1)^2 - (8m - 3)^2\} \cdot 8 \{(8k + 3)^2 + (8l + 1)^2\} \cdot 2^2 \cdot (-4)^2 = 2^{14}p(4m - 1).$$

□

*Proof of Lemma 5.4.* For any  $q \in P'$ , there exist  $r, s, t, u \in \mathbb{Z}$  satisfying  $2q = (4r + 1)^2 + (4s)^2 + (4t + 1)^2 + (4u + 2)^2$ . Let

$$\begin{aligned} a_0 &= m + r + 1, & a_1 &= m + t, & a_2 &= m - r, & a_3 &= m - t, \\ a_4 &= m + s, & a_5 &= m + u + 1, & a_6 &= m - s, & a_7 &= m - u, \\ a_8 &= m - r + 1, & a_9 &= m - t, & a_{10} &= m + r + 1, & a_{11} &= m + t + 1, \\ a_{12} &= m - s, & a_{13} &= m - u - 1, & a_{14} &= m + s, & a_{15} &= m + u. \end{aligned}$$

Then we have

$$D_G(\mathbf{a}) = 2 \{(8m + 3)^2 - (8m + 1)^2\} \cdot 16 \cdot (2q)^2 \cdot (-4)^2 = 2^{14}q^2(4m + 1).$$

Let

$$\begin{aligned} a_0 &= m + r + 1, & a_1 &= m + t, & a_2 &= m - r, & a_3 &= m - t, \\ a_4 &= m + s, & a_5 &= m + u, & a_6 &= m - s, & a_7 &= m - u - 1, \\ a_8 &= m - r, & a_9 &= m - t - 1, & a_{10} &= m + r, & a_{11} &= m + t, \\ a_{12} &= m - s - 1, & a_{13} &= m - u - 1, & a_{14} &= m + s - 1, & a_{15} &= m + u. \end{aligned}$$

Then we have

$$D_G(\mathbf{a}) = 2 \{(8m - 1)^2 - (8m - 3)^2\} \cdot 16 \cdot (2q)^2 \cdot 4^2 = 2^{14}q^2(4m - 1).$$

□

From Lemmas 3.1, 4.1 and 5.1–5.4, Theorem 1.1 is proved.

## References

- [1] Ton Boerkoel and Christopher Pinner. Minimal group determinants and the Lind-Lehmer problem for dihedral groups. *Acta Arith.*, 186(4):377–395, 2018.
- [2] Ferdinand Georg Frobenius. Über die Primfactoren der Gruppendedeterminante. *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin*, pages 1343–1382, 1896. Reprinted in *Gesammelte Abhandlungen, Band III*. Springer-Verlag Berlin Heidelberg, New York, 1968, pages 38–77.
- [3] Otto Hölder. Die Gruppen der Ordnungen  $p^3$ ,  $pq^2$ ,  $pqr$ ,  $p^4$ . *Math. Ann.*, 43(2-3):301–412, 1893.
- [4] Norbert Kaiblinger. Progress on Olga Taussky-Todd’s circulant problem. *Ramanujan J.*, 28(1):45–60, 2012.
- [5] H. Turner Laquer. Values of circulants with integer entries. In *A collection of manuscripts related to the Fibonacci sequence*, pages 212–217. Fibonacci Assoc., Santa Clara, Calif., 1980.
- [6] Michael J. Mossinghoff and Christopher Pinner. Prime power order circulant determinants, 2022. arXiv:2205.12439v2 [math.NT].
- [7] Morris Newman. Determinants of circulants of prime power order. *Linear and Multilinear Algebra*, 9(3):187–191, 1980.
- [8] Morris Newman. On a problem suggested by Olga Taussky-Todd. *Illinois J. Math.*, 24(1):156–158, 1980.
- [9] Bishnu Paudel and Chris Pinner. Integer circulant determinants of order 15. *Integers*, 22:Paper No. A4, 21, 2022.
- [10] Bishnu Paudel and Christopher Pinner. Minimal group determinants for dicyclic groups. *Mosc. J. Comb. Number Theory*, 10(3):235–248, 2021.
- [11] Bishnu Paudel and Christopher Pinner. The group determinants for  $\mathbb{Z}_n \times H$ , 2022. arXiv:2211.09930v3 [math.NT].
- [12] Bishnu Paudel and Christopher Pinner. The integer group determinants for  $Q_{16}$ , 2023. arXiv:2302.11688v1 [math.NT].
- [13] Christopher Pinner. The integer group determinants for the symmetric group of degree four. *Rocky Mountain J. Math.*, 49(4):1293–1305, 2019.
- [14] Christopher Pinner and Christopher Smyth. Integer group determinants for small groups. *Ramanujan J.*, 51(2):421–453, 2020.

- [15] Olga Taussky Todd. Integral group matrices. In *Notices of the American Mathematical Society*, volume 24, United States of America, April 1977. American Mathematical Society. Abstract no. 746-A15, 746th Meeting, Hayward, California, Apr. 22–23, 1977.
- [16] Naoya Yamaguchi and Yuka Yamaguchi. Generalized Dedekind’s theorem and its application to integer group determinants, 2022. arXiv:2203.14420v2 [math.RT].
- [17] Naoya Yamaguchi and Yuka Yamaguchi. Remark on Laquer’s theorem for circulant determinants. *International Journal of Group Theory*, 12(4):265–269, 2023.
- [18] Yuka Yamaguchi and Naoya Yamaguchi. Integer group determinants for  $C_2^4$  (in press). *Ramanujan J.* Available also at arXiv:2209.12446.
- [19] Yuka Yamaguchi and Naoya Yamaguchi. Integer circulant determinants of order 16. *Ramanujan J.*, 2022. Advance online publication, <https://doi.org/10.1007/s11139-022-00599-9>.
- [20] Yuka Yamaguchi and Naoya Yamaguchi. Integer group determinants for abelian groups of order 16, 2022. arXiv:2211.14761v1 [math.NT].
- [21] Yuka Yamaguchi and Naoya Yamaguchi. Integer group determinants for  $C_4^2$ , 2022. arXiv:2211.01597v2 [math.NT].
- [22] Yuka Yamaguchi and Naoya Yamaguchi. Integer group determinants for  $C_2^2 \rtimes C_4$ , 2023. arXiv:2303.08489v2 [math.NT].
- [23] J. W. A. Young. On the Determination of Groups Whose Order is a Power of a Prime. *Amer. J. Math.*, 15(2):124–178, 1893.

Faculty of Education, University of Miyazaki, 1-1 Gakuen Kibanadai-nishi, Miyazaki 889-2192, Japan

*Email address*, Yuka Yamaguchi: y-yamaguchi@cc.miyazaki-u.ac.jp

*Email address*, Naoya Yamaguchi: n-yamaguchi@cc.miyazaki-u.ac.jp