

The lifespan estimates of radially symmetric solutions to systems of nonlinear wave equations in even space dimensions

Yuki Kurokawa *

MSC 2020 : 35L71, 35L05, 35E15

Key words : system, semilinear wave equation, radially symmetric solution, lifespan

Running head : Lifespan estimates of radially symmetric solutions

Abstract

The optimal lifespan estimates of a solution to weakly coupled systems of wave equations have been investigated by many works, except for the lower bound in even space dimensions. Our aim is to prove the open part under the assumption of radial symmetry on the solution. The odd dimensional case was already obtained in our previous paper [18] by long time existence of the solution in weighted L^∞ space. In this paper, we employ similar methods. The main difficulty is found in estimating the integral kernel which is completely different from odd dimensional case.

1 Introduction

We consider the following Cauchy problem of the systems of semilinear wave equations :

$$\begin{cases} u_{tt} - \Delta u = |v|^p, \\ v_{tt} - \Delta v = |u|^q, \end{cases} \quad \text{in } \mathbf{R}^n \times (0, \infty), \quad (1.1)$$

for $p, q > 1$ with the initial data

$$\begin{cases} u(x, 0) = \varepsilon\varphi_1(x), \quad u_t(x, 0) = \varepsilon\psi_1(x), \\ v(x, 0) = \varepsilon\varphi_2(x), \quad v_t(x, 0) = \varepsilon\psi_2(x), \end{cases} \quad \text{for } x \in \mathbf{R}^n, \quad (1.2)$$

where $\varepsilon > 0$ is a sufficiently small parameter and φ_i, ψ_i ($i = 1, 2$), which present the shape of the data, are sufficiently smooth and have compact support.

For this problem, Del Santo, Georgiev and Mitidieri [4] proved that

- $F(p, q; n) < 0, 2 \leq p, q \leq 3 \implies$ Existence of a global-in-time solution for $\varepsilon \ll 1$,
- $F(p, q; n) > 0$, some positivity on data \implies Nonexistence of a global-in-time solution,

*Hokkaido University of Education, Kushiro 085-8580, Japan. e-mail : kurokawa.yuki@k.hokkyodai.ac.jp.

for $n \geq 2$, where

$$F(p, q; n) := \max \left\{ \frac{p+2+q^{-1}}{pq-1}, \frac{q+2+p^{-1}}{pq-1} \right\} - \frac{n-1}{2}. \quad (1.3)$$

Their restrictions “ $p, q \leq 3$ ” and “some positivity on data” were relaxed by Del Santo [3] in three space dimensions. Thus, we see that there exists a critical curve $F(p, q; n) = 0$ on the p - q plane, which is symmetric across the line $p = q$. Independently, Deng [6], [7] proved the blow-up of a weak solution for the case of $F(p, q; n) > 0$ with $p \leq q$ and $n \geq 2$ for some positive data. The blow-up of a solution for the critical case $F(p, q; n) = 0$ was proved by Del Santo and Mitidieri [5] in three space dimensions under some positivity on data.

As for the blow-up case *i.e.* $F(p, q) \geq 0$, there are several results on “lifespan”, the maximal existence time of a solution. Kubo and Ohta [17] obtained the upper bound of the lifespan of the classical solution in two or three space dimensions for some positive data. The optimal lifespan of a classical solution for $n = 2$ or 3 has been proved for all cases by Agemi, Kurokawa and Takamura [1] and Kurokawa and Takamura [19]. Thus, we have

$$\begin{aligned} T_0(\varepsilon) &= \infty && \text{if } F(p, q; n) < 0, \\ \exp(c\varepsilon^{-\min\{p(pq-1), q(pq-1)\}}) &\leq T_0(\varepsilon) \leq \exp(c'\varepsilon^{-\min\{p(pq-1), q(pq-1)\}}) && \text{if } F(p, q; n) = 0, p \neq q, \\ \exp(c\varepsilon^{-p(p-1)}) &\leq T_0(\varepsilon) \leq \exp(c'\varepsilon^{-p(p-1)}) && \text{if } F(p, q; n) = 0, p = q, \\ c\varepsilon^{-F(p, q; n)^{-1}} &\leq T_0(\varepsilon) \leq c'\varepsilon^{-F(p, q; n)^{-1}} && \text{if } F(p, q; n) > 0, \end{aligned}$$

for $n = 2, 3$, where c and c' are positive constants independent of ε and $T_0(\varepsilon)$ means the lifespan of the classical solution to (1.1) with (1.2).

The higher dimensional case was first investigated by Georgiev, Takamura and Zhou in [9] for the sub-critical case. They showed that for some data of Sobolev functions the lifespan T_ε of a certain weak solution in Lebesgue space can be pinched from both side as follows.

$$C_1\varepsilon^{-(F(p, q; n) + \sigma)^{-1}} \leq T_\varepsilon \leq C_2\varepsilon^{-F(p, q; n)^{-1}}, \quad (1.4)$$

where C_1, C_2 are positive constants independent of ε and $\sigma > 0$ is a small error. Moreover, Remark 1.2 in [9], they said that they can get the similar lower bound of the lifespan for the case of $F(p, q; n) = 0$:

$$C_1\varepsilon^{-1/\sigma} \leq T_\varepsilon. \quad (1.5)$$

Our previous paper, Kurokawa [18], succeeded to remove the error σ in the lower bound of the lifespan for $F(p, q; n) \geq 0$ in odd space dimensions under the assumption of radial symmetry of a solution. The present paper is devoted to remove the error for a radially symmetric solution in even space dimensions as well.

The upper bound of the lifespan for the critical case $F(p, q; n) = 0$ was obtained by Kurokawa, Takamura and Wakasa [20] for $n \geq 4$. Their estimate is same as the one in low space dimensions. Recently there are some alternative proof for the upper bound of the lifespan. Ikeda, Sobajima and Wakasa [12] introduced new test functions in the weak form to obtain the same results for $n \geq 2$ and $F \geq 0$. Palmieri and Takamura [25] studied the systems of semilinear damped wave equations :

$$\begin{cases} u_{tt} - \Delta u + b_1(t)u_t = |v|^p, & x \in \mathbf{R}^n, t > 0, \\ v_{tt} - \Delta v + b_2(t)v_t = |u|^q, & x \in \mathbf{R}^n, t > 0, \\ (u, u_t, v, v_t)(0, x) = (\varepsilon u_0, \varepsilon u_1, \varepsilon v_0, \varepsilon v_1)(x) & x \in \mathbf{R}^n \end{cases} ,$$

for $b_1, b_2 \in C([0, \infty)) \cap L^1([0, \infty))$ and they get the upper bound of the lifespan of an energy solution for $n \geq 2$ and $F \geq 0$. This work includes the case of $b_1 \equiv 0$ and $b_2 \equiv 0$ *i.e.* our systems (1.1).

From now on, we consider the following radially symmetric version of the systems of wave equations for $r = |x|$ and $T > 0$:

$$\begin{cases} u_{tt} - u_{rr} - \frac{n-1}{r}u_r = |v|^p, \\ v_{tt} - v_{rr} - \frac{n-1}{r}v_r = |u|^q, \end{cases} \quad \text{in } (0, \infty) \times (0, T) \quad (1.6)$$

for $p, q > 1$ with the initial data

$$\begin{cases} u(r, 0) = \varepsilon f_1(r), & u_t(r, 0) = \varepsilon g_1(r), \\ v(r, 0) = \varepsilon f_2(r), & v_t(r, 0) = \varepsilon g_2(r), \end{cases} \quad (1.7)$$

where, $\varepsilon > 0$ and f_i, g_i ($i = 1, 2$) are sufficiently smooth functions of compact support. More precisely, we assume that

$$\text{supp } \{f_1, f_2, g_1, g_2\} \subset \{r \leq t + k\} \quad \text{with } k > 1. \quad (1.8)$$

In this paper, we study the following system of the integral equations which is equivalent to (1.6) with (1.7).

$$\begin{cases} u(r, t) = \varepsilon u^0(r, t) + L(|v|^p)(r, t), \\ v(r, t) = \varepsilon v^0(r, t) + L(|u|^q)(r, t), \end{cases} \quad \text{for } (r, t) \in (0, \infty) \times (0, T), \quad (1.9)$$

where u^0, v^0 are solutions to free wave equation with the same data as the ones of u, v in (1.7) without ε . L is an integral operator associated to Duhamel's principle which will be defined in the next section. In odd space dimensions, (1.6) with (1.7) is equivalent to (1.9), since we can construct a C^2 solution of (1.9) (See [15], [18]). On the other hand, we do not expect to obtain C^2 solution due to the lack of the differentiability of the kernel of the integral expressions. Therefore, our solution is defined as a C^1 solution of (1.9). Namely, the lifespan $T(\varepsilon)$ in the present paper is defined by

$$T(\varepsilon) := \sup\{T \in (0, \infty] : \text{There exists a radially symmetric solution } (u, v) \in (C^1((0, \infty) \times [0, T]))^2 \text{ to the system (1.9) with any (1.7)}.\}. \quad (1.10)$$

Remark 1.1 *The solution defined above can be a weak solution of (1.1) in the sense of $(u, v) \in (C^1((0, \infty) \times [0, T]))^2 \cap (C^2([0, T]; \mathcal{D}'(\mathbf{R}^n)))^2$. See the next section. More precisely, see Theorem 2.1 and Theorem 2.2 in [16].*

Now our main result is the following.

Theorem 1 *Let $n = 2m + 2$ ($m = 2, 3, 4, \dots$), $F(p, q; n) \geq 0$ and*

$$p, q \in \left(\frac{n+1}{n-1}, \frac{n+3}{n-1} \right).$$

Assume that $f_i \in C^2([0, \infty))$ and $g_i \in C^1([0, \infty))$ ($i = 1, 2$) satisfy the support property (1.8). Then, there exists a positive constant $\varepsilon_0 = \varepsilon_0(p, q, f_1, f_2, g_1, g_2, n, k)$ such that the lifespan $T(\varepsilon)$, defined in (1.10), satisfies

$$T(\varepsilon) \geq \begin{cases} \exp(C\varepsilon^{-\min\{p(pq-1), q(pq-1)\}}) & \text{if } F(p, q; n) = 0, p \neq q, \\ \exp(C\varepsilon^{-p(p-1)}) & \text{if } F(p, q; n) = 0, p = q, \\ C\varepsilon^{-F(p, q; n)^{-1}} & \text{if } F(p, q; n) > 0 \end{cases}$$

for any $\varepsilon \in (0, \varepsilon_0]$, where C is a positive constant independent of ε and $F(p, q; n)$ is the one in (1.3).

Remark 1.2 By the symmetricity of the systems (1.9), it is enough to prove the case of $p \leq q$ only.

Remark 1.3 In this paper, $m = 1$ i.e. four space dimensional case is excluded. Because we employ the methods in Kubo and Kubota [16] which study the asymptotic behavior of radially symmetric solutions to the corresponding single wave equations in even space dimensions with $n \geq 6$. Thus, the similar results for $n = 4$ has not been obtained yet, while Li and Zhou [22] and Lindblad and Sogge [24] already get the sharp lower bound for the single equation in four space dimensions.

Remark 1.4 The corresponding single equation :

$$\begin{cases} u_{tt} - \Delta u = |u|^p & \text{in } \mathbf{R}^n \times [0, \infty), \\ u(x, 0) = \varepsilon f(x), \quad u_t(x, 0) = \varepsilon g(x) & \text{for } x \in \mathbf{R}^n \end{cases} \quad (1.11)$$

has been investigated by many works and we now know that the critical power $p_0(n)$ is the positive root of

$$\gamma(p, n) \equiv 2 + (n + 1)p - (n - 1)p^2 = 0.$$

Namely, if $p > p_0(n)$, there exists a global-in time solution to (1.11) for sufficiently small ε . On the other hand, a blow-up occurs, for some initial data, if $p \leq p_0(n)$ (See [14], [29], [11], [10], [27], [28], [26], [8], [32], [36]).

The lifespan estimates are also well-studied by many works. See [33], [23], [34], [35], [30], [13] for low space dimensions and [22], [24], [31], [37], [21] for high space dimensions. For the details, see [13] and its introduction. Lindblad and Sogge [24] has an important relation to our problems. They studied (1.11) with radially symmetric data and showed that for $n \geq 3$ and $(n + 1)/(n - 1) \leq p \leq p_0(n)$ (1.11) has a unique solution u verifying

$$t^{(n-1)/2-(n+1)/2p} r^{(n+1)/2p} u \in L_t^\infty L_r^p([0, \tilde{T}_\varepsilon] \times \mathbf{R}_+),$$

where

$$\tilde{T}_\varepsilon = \begin{cases} c_0 \varepsilon^{-2p(p-1)/\gamma(p, n)} & \text{if } \frac{n+1}{n-1} \leq p < p_0(n), \\ \exp(c_0 \varepsilon^{-p(p-1)}) & \text{if } p = p_0(n), \end{cases},$$

provided c_0 is a sufficiently small positive constant. Namely, \tilde{T}_ε implies the lower bound of the lifespan for (1.11). Note that our estimates for $p = q$ in Theorem 1 have the same form as their \tilde{T}_ε , since we have

$$F(p, p; n) = \frac{\gamma(p, n)}{2p(p-1)}.$$

This article is organized as follows. In section 2, we recall the representation formulas introduced by Kubo and Kubota in [16]. We will consider the decay estimates of a solution to free wave equations in section 3. In section 4, we will state a priori estimates. Using them, we will construct a local solution and estimate the lifespan in section 5. Finally section 6 is devoted to the proof of a priori estimates.

2 Representation formula

Theorem 1 will be proved by the construction of a local-in-time solution to (1.9) (1.7) by the iteration argument in a weighted L^∞ space with the sharp pointwise estimates of a solution. This is based on the way by Kubo and Kubota [16]. In [16], they get two useful representation formulas starting from the one by Courant and Hilbert [2]. We shall use their formulas and apply their methods into our problem for the compactly supported data.

Let $n = 2m + 2$ ($m \geq 2$). We define in (1.9) that

$$u^0(r, t) := \frac{1}{c_n} \{ \Theta(g_1)(r, t) + D_t \Theta(f_1)(r, t) \}, \quad (2.1)$$

$$v^0(r, t) := \frac{1}{c_n} \{ \Theta(g_2)(r, t) + D_t \Theta(f_2)(r, t) \}, \quad (2.2)$$

$$L(G)(r, t) := \frac{1}{c_n} \int_0^t \Theta(G(\cdot, \tau))(r, t - \tau) d\tau, \quad (2.3)$$

where $c_n := \sqrt{\pi} \Gamma((n-1)/2)$ and

$$\Theta(g)(r, t) := \frac{1}{r^{2m}} (J_1(g)(r, t) + J_2(g)(r, t)). \quad (2.4)$$

Here J_1 and J_2 are defined by

$$J_1(r, t) := \int_{|t-r|}^{t+r} \lambda^{2m+1} g(\lambda) K_m(\lambda, r, t) d\lambda, \quad (2.5)$$

$$J_2(r, t) := \int_0^{(t-r)_+} \lambda^{2m+1} g(\lambda) \tilde{K}_m(\lambda, r, t) d\lambda, \quad (2.6)$$

$$(2.7)$$

where

$$K_j(\lambda, r, t) := \int_\lambda^{t+r} \frac{H_j(r, t)(\rho, r, t)}{\sqrt{\rho^2 - \lambda^2}} d\rho \quad \text{for } 0 \leq j \leq m, \quad (2.8)$$

$$\tilde{K}_j(\lambda, r, t) := \int_{t-r}^{t+r} \frac{H_j(r, t)(\rho, r, t)}{\sqrt{\rho^2 - \lambda^2}} d\rho \quad \text{for } 0 \leq j \leq m, \quad (2.9)$$

with

$$H_j(\rho, r, t) := \left\{ D_\rho \left(-\frac{1}{2\rho} \right) \right\}^j (r^2 - (\rho - t)^2)^{m-\frac{1}{2}} \quad \text{for } |\rho - t| < r, \quad 0 \leq j \leq m.$$

This expression was introduced by [16], while they denote J_i by W_i in [16]. First we mention the following property.

Proposition 2.1 *Let the assumptions of Theorem 1 be fulfilled. Let $(u(r, t), v(r, t))$ be C^1 solution of (1.9) with (1.7). If we set $\varphi_i(x) = f_i(|x|)$, $\psi_i(x) = g_i(|x|)$ ($i = 1, 2$), $\tilde{u}(x, t) = u(|x|, t)$, $\tilde{v}(x, t) = v(|x|, t)$ for $(x, t) \in (\mathbf{R}^n \setminus \{0\}) \times [0, T]$, then, (\tilde{u}, \tilde{v}) is a weak solution of (1.1) with (1.2) in a distribution sense and it belongs to $((C^2([0, T]; \mathcal{D}'(\mathbf{R}^n)))^2$.*

(Proof)

This can be proved by the same way as Theorem 2.1 and Theorem 2.2 in [16]. So we shall omit the proof. \square

In what follows, we shall summarize the properties of K_j in (2.8) and \tilde{K}_j in (2.9) mentioned by Kubo and Kubota in [16] which will be used in our problem.

Lemma 2.1 (Kubo and Kubota [16] §4.1 p.215)

For $0 \leq j \leq m - 1$, we have

$$K_j(\lambda, r, t) = \tilde{K}(\lambda, r, t) \quad \text{for } \lambda = t - r > 0, \quad (2.10)$$

$$K_j(\lambda, r, t) = 0 \quad \text{for } \lambda = t + r. \quad (2.11)$$

Lemma 2.2 (Kubo and Kubota [16] §4.1 Lemma 4.1)

Let $1 \leq j \leq m$ and $(r, t) \in \Omega$. Then we have

$$K_j(\lambda, r, t) = -D_{\lambda^2} K_{j-1}(\lambda, r, t) \quad \text{for } |t - r| < \lambda < t + r, \quad (2.12)$$

$$\tilde{K}_j(\lambda, r, t) = -D_{\lambda^2} \tilde{K}_{j-1}(\lambda, r, t) \quad \text{for } 0 < \lambda < t - r, \quad (2.13)$$

Lemma 2.3 (Kubo and Kubota [16] §4.1 Lemma 4.2)

Let $\eta = 0$ or $\frac{1}{2}$ and let $(r, t) \in \Omega$. Denote (D_r, D_t, D_λ) by D . If $j + |\alpha| \leq m$, we have

$$|D^\alpha K_j(\lambda, r, t)| \leq Cr^{2m-j-|\alpha|+\eta-\frac{1}{2}} \lambda^{-j-\eta} (\lambda - t + r)^{-\frac{1}{2}} \quad \text{for } |t - r| < \lambda < t + r, \quad t \geq 0, \quad (2.14)$$

$$|D^\alpha \tilde{K}_j(\lambda, r, t)| \leq Cr^{2m-j-|\alpha|+\eta-\frac{1}{2}} (t - r)^{-j-\eta} (t - r - \lambda)^{-\frac{1}{2}} \quad \text{for } 0 < \lambda < t - r. \quad (2.15)$$

Lemma 2.4 (Kubo and Kubota [16] §4.1 Corollary 4.3)

Let $\eta = 0$ or $\frac{1}{2}$ and let $(r, t) \in \Omega$. If $j + |\alpha| \leq m$, we have

$$|D_\lambda D_{r,t}^\alpha \tilde{K}_j(\lambda, r, t)| \leq Cr^{2m-j-|\alpha|+\eta-\frac{1}{2}} (t - r)^{-j-\eta} (t - r - \lambda)^{-\frac{3}{2}} \quad \text{for } 0 < \lambda < t - r. \quad (2.16)$$

Lemma 2.5 (Kubo and Kubota [16] §4.1 Lemma 4.4)

Let $(r, t) \in \Omega$ and $0 < \lambda < t - r$. Let $\eta = 0$ or $\frac{1}{2}$. If $0 \leq j \leq m$, $|\alpha| \leq m$ and $j + |\alpha| \geq m$, we have

$$|D_{r,t}^\alpha \tilde{K}_j(\lambda, r, t)| \leq Cr^{m-\eta} (t - r)^{-j-\frac{1}{2}+\eta} (t - r - \lambda)^{m-j-|\alpha|-\frac{1}{2}}. \quad (2.17)$$

Moreover, if $0 \leq j \leq m$, $|\alpha| \leq m - 1$ and $j + |\alpha| \geq m - 1$, we have

$$|D_{r,t}^\alpha \tilde{K}_j(\lambda, r, t)| \leq Cr^{m+1-\eta} (t - r)^{-j-\frac{1}{2}+\eta} (t - r - \lambda)^{m-j-|\alpha|-\frac{3}{2}}. \quad (2.18)$$

Lemma 2.6 (Kubo and Kubota [16] §4.1 Lemma 4.5)

Let $\eta = 0$ or $\frac{1}{2}$ and let $(r, t) \in \Omega$. If $0 < r - t \leq \lambda < r + t$, we have

$$|D_{r,t,\lambda}^\alpha K_m(\lambda, r, t)| \leq Cr^{m-\frac{1}{2}+\eta} \lambda^{-m-|\alpha|-\frac{1}{2}-\eta}. \quad (2.19)$$

Lemma 2.7 (*Kubo and Kubota [16] §4.1 p.213 REMARK*)

When $0 < t < r$, we have for $\lambda = r + t$

$$|K_m(\lambda, r, t)| \leq Cr^{m-\frac{1}{2}+\mu}\lambda^{-m-\frac{1}{2}-\mu} \quad (2.20)$$

$$|D_{r,t,\lambda}K_m(\lambda, r, t)| \leq Cr^{m-\frac{1}{2}}\lambda^{-m-\frac{3}{2}}. \quad (2.21)$$

Using these properties, Kubo and Kubota obtained another representation formula for Θ as follows by Lemma 4.6 and Lemma 4.7 in [16].

First, assume that

$$g \in C^1((0, \infty)) \quad \text{and} \quad g^{(j)}(\lambda) = O(\lambda^{-2m+\delta-j}) \quad \text{as} \quad \lambda \downarrow 0 \quad (2.22)$$

hold for $j = 0, 1$ and some positive constant δ . Then we have

$$2r^{2m}\Theta(g)(r, t) = J_3(r, t) + J_4(r, t), \quad (2.23)$$

where

$$J_3(r, t) := \int_{|t-r|}^{t+r} (\lambda^{2m}g(\lambda))'K_{m-1}(\lambda, r, t), \quad (2.24)$$

$$J_4(r, t) := \begin{cases} \int_0^{t-r} (\lambda^{2m}g(\lambda))'\tilde{K}_{m-1}(\lambda, r, t)d\lambda, & \text{if } t-r > 0, \\ (r-t)^{2m}g(r-t)K_{m-1}(r-t, r, t) & \text{if } t-r < 0. \end{cases} \quad (2.25)$$

Moreover, we have

$$D_\alpha(2r^{2m}\Theta(g)(r, t)) = J_5(r, t) + J_6(r, t), \quad (2.26)$$

where $D_\alpha = D_r$ or D_t and

$$J_5(r, t) := \int_{|t-r|}^{t+r} (\lambda^{2m}g(\lambda))'D_\alpha K_{m-1}(\lambda, r, t), \quad (2.27)$$

$$J_6(r, t) := \begin{cases} \int_0^{t-r} (\lambda^{2m}g(\lambda))'D_\alpha \tilde{K}_{m-1}(\lambda, r, t)d\lambda, & \text{if } t-r > 0, \\ (r-t)^{2m}g(r-t)D_\alpha K_{m-1}(r-t, r, t) & \text{if } t-r < 0. \end{cases} \quad (2.28)$$

Next, assume that

$$f \in C^2((0, \infty)) \quad \text{and} \quad f^{(j)}(\lambda) = O(\lambda^{-2m+2+\delta-j}) \quad \text{as} \quad \lambda \downarrow 0 \quad (2.29)$$

for $j = 0, 1, 2$ and some positive constant δ . When $t - r > 0$, we have

$$D^\beta D_t(2r^{2m}\Theta(f)(r, t)) = J_7(r, t) + J_8(r, t), \quad (2.30)$$

where $D = (D_r, D_t)$ and

$$J_7(r, t) := \int_{t-r}^{t+r} D_\lambda D_{\lambda^2}(\lambda^{2m}f(\lambda)) \cdot D^\beta D_t K_{m-2}(\lambda, r, t), \quad (2.31)$$

$$J_8(r, t) := \int_0^{t-r} D_\lambda D_{\lambda^2}(\lambda^{2m}f(\lambda)) \cdot D^\beta D_t \tilde{K}_{m-2}(\lambda, r, t). \quad (2.32)$$

$$(2.33)$$

On the other hand, when $t - r < 0$, we have

$$D_t(2r^{2m}\Theta(f)(r, t)) = J_9(r, t) + J_{10}(r, t), \quad (2.34)$$

where

$$J_9(r, t) := \int_{r-t}^{r+t} \lambda^{2m+1} f(\lambda) \cdot D_t K_m(\lambda, r, t), \quad (2.35)$$

$$J_{10}(r, t) := (\lambda^{2m+1} f(\lambda) K_m(\lambda, r, t))|_{\lambda=r+t} + (\lambda^{2m+1} f(\lambda) K_m(\lambda, r, t))|_{\lambda=r-t}. \quad (2.36)$$

Our assumptions on g_i and f_i ($i = 1, 2$) in Theorem 1 can play the same roles as (2.22) and (2.29) in the proof of the above representation formulas in Kubo and Kubota [16]. Therefore, we will use these representation formulas (2.4), (2.23), (2.26), (2.30), (2.34) suitably in the forthcoming sections.

3 Decay estimates of free solution

In this section, we show the decay estimates of u^0 and v^0 . We have the following.

Proposition 3.1 *Let the assumptions on f_i, g_i ($i = 1, 2$) in Theorem 1 be fulfilled. Let u^0 and v_0 are the one in (2.1) and (2.2), respectively. Then we have $\text{supp} \{u^0, v^0\} \subset \{r \leq t + k\}$ and*

$$|D^\beta u^0(r, t)| \leq \tilde{C} \left(\frac{r}{k}\right)^{-m+1-|\beta|} \left(\frac{r+2k}{k}\right)^{-1+|\beta|} \tau_+(r, t)^{-\frac{1}{2}} \phi_i(r, t),$$

$$|D^\beta v^0(r, t)| \leq \tilde{C} \left(\frac{r}{k}\right)^{-m+1-|\beta|} \left(\frac{r+2k}{k}\right)^{-1+|\beta|} \tau_+(r, t)^{-\frac{1}{2}} \phi_i(r, t),$$

for $(r, t) \in \Omega := \{(r, t) \in (0, \infty) \times [0, \infty) : r \leq t + k\}$, where $D = (D_r, D_t)$, $|\beta| \leq 1$, $\tau_\pm(r, t) = (t \pm r + 2k)/k$, $\phi_i(r, t) = \tau_-^{-m-\frac{1}{2}-|\beta|}$ and \tilde{C} is a positive constant independent of ε .

Proof. We shall consider u^0 only, because v_0 has the same form. The support property of u^0 can be readily obtained by the representation formula (2.1), (2.4), (2.5) and (2.6). In what follows, we prove the decay estimates of $D^\beta u^0$ by the dividng into three cases. This division is due Kubo and Kubota in ([16]). Here and hereafter, we denote any positive constant independent of ε by C . Namely, C can mean the different values in each line. Besides, we denote $\tau_\pm(r, t)$ by τ_\pm simply, since we fix (r, t) in each cases. Similarly, we shall omit the variable (r, t) of the other functions as well when not misleading.

3.1 Case 1, $t \geq 2r > 0$ or $0 < r < \min\{t, k\}$

In this case, it holds that

$$t \geq 2r > 0 \implies 3\tau_- \geq \tau_+ \geq \frac{r+2k}{k},$$

$$0 < r < \min\{t, r\} \implies 2 < \frac{r+2k}{k}, \tau_+, \tau_- \leq 5,$$

for $(r, t) \in \Omega$. Hence, it is enough to show

$$|D^\beta D_t(2r^{2m}\Theta(f))(r, t)| \leq Cr^{m+1-|\beta|}\tau_-^{-m-2}, \quad (3.1)$$

$$|D^\beta(2r^{2m}\Theta(g))(r, t)| \leq Cr^{m+1-|\beta|}\tau_-^{-m-2}, \quad (3.2)$$

for $|\beta| = 0, 1$ in Ω .

For (3.1), we use the representation formula (2.30) both for $|\beta| = 0$ and 1. By (2.14) with $\eta = 0$, we have

$$|J_7(r, t)| \leq Cr^{m+\frac{1}{2}-|\beta|} \int_{t-r}^{t+r} \frac{\lambda^{m-1}|f(\lambda)| + \lambda^m|f'(\lambda)| + \lambda^{m+1}|f''(\lambda)|}{\sqrt{\lambda-t+r}} d\lambda.$$

When $t-r \geq k$, $J_7 = 0$ because of the assumption of the support property of data, *i.e.*

$$f(r) = g(r) = 0 \quad \text{if } r \geq k. \quad (3.3)$$

When $0 < t-r \leq k$, it follows from (3.3) and $2 \leq \tau_- \leq 3$ that

$$|J_7(r, t)| \leq Cr^{m+\frac{1}{2}-|\beta|} \int_{t-r}^{\min\{k, t+r\}} \frac{1}{\sqrt{\lambda-t+r}} d\lambda \leq Cr^{m+1-|\beta|}\tau_-^{-m-2}.$$

As to J_8 for $0 < t-r \leq 2k$, we use (2.15) $_{m-2}$ with $\mu = \frac{1}{2}$. Then, it follows from (3.3) and $2 \leq \tau_- \leq 4$ that

$$\begin{aligned} |J_8(r, t)| &\leq Cr^{m+1-|\beta|}(t-r)^{-m+\frac{3}{2}} \int_0^{t-r} \frac{\lambda^{2m-3}|f(\lambda)| + \lambda^{2m-2}|f'(\lambda)| + \lambda^{2m-1}|f''(\lambda)|}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m+1-|\beta|} \int_0^{t-r} \frac{\lambda^{m-\frac{3}{2}}|f(\lambda)| + \lambda^{m-\frac{1}{2}}|f'(\lambda)| + \lambda^{m+\frac{1}{2}}|f''(\lambda)|}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m+1-|\beta|} \int_0^{t-r} \frac{1}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m+1-|\beta|}\tau_-^{-m-2}. \end{aligned}$$

When $t-r \geq 2k$, (3.3) gives us

$$J_8(r, t) = \int_0^{\frac{t-r}{2}} D_\lambda D_{\lambda^2}(\lambda^{2m}f(\lambda)) \cdot D^\beta D_t \tilde{K}_{m-2}(\lambda, r, t) d\lambda.$$

By employing the integration by parts and (2.13) twice together with (3.3), we have

$$\begin{aligned} J_8(r, t) &= \left[D_{\lambda^2}(\lambda^{2m}f(\lambda)) D^\beta D_t \tilde{K}_{m-2}(\lambda, r, t) \right]_{\lambda=0}^{\frac{t-r}{2}} \\ &\quad + \left[\lambda^{2m}f(\lambda) D^\beta D_t \tilde{K}_{m-1}(\lambda, r, t) \right]_{\lambda=0}^{\frac{t-r}{2}} \\ &\quad + 2 \int_0^{\frac{t-r}{2}} \lambda^{2m+1}f(\lambda) D^\beta D_t \tilde{K}_m(\lambda, r, t) d\lambda. \end{aligned} \quad (3.4)$$

By (3.3), the first and the second line of (3.4) vanishes. Using (2.18)_m with $\mu = 0$ for $|\beta| = 0$ and (2.17)_m with $\mu = 0$ for $|\beta| = 1$, we have

$$|J_8(r, t)| \leq Cr^{m+1-|\beta|}(t-r)^{-m-3} \int_0^{\frac{t-r}{2}} \lambda^{2m+1}|f(\lambda)|d\lambda \leq Cr^{m+1-|\beta|}\tau_-^{-m-2}.$$

For (3.2) with $|\beta| = 0$, we use (2.23). Using (2.14)_{m-1} with $\mu = 0$, we have

$$|J_3(r, t)| \leq Cr^{m+\frac{1}{2}} \int_{t-r}^{t+r} \frac{\lambda^m|g(\lambda)| + \lambda^{m+1}|g'(\lambda)|}{\sqrt{\lambda-t+r}}d\lambda.$$

Then, we can treat it by the same way as J_7 for (3.1) and get the desired estimate. As to J_4 for $t-r \leq 2k$, we use (2.15)_{m-1} with $\mu = \frac{1}{2}$. Then, it follows from $2 \leq \tau_- \leq 4$ that

$$\begin{aligned} |J_4(r, t)| &\leq Cr^{m+1}(t-r)^{-m+\frac{1}{2}} \int_0^{t-r} \frac{\lambda^{2m+1}|g(\lambda)| + \lambda^{2m}|g'(\lambda)|}{\sqrt{t-r-\lambda}}d\lambda \\ &\leq Cr^{m+1}(t-r)^{\frac{1}{2}} \int_0^{t-r} \frac{\lambda^{m+1}|g(\lambda)| + \lambda^m|g'(\lambda)|}{\sqrt{t-r-\lambda}}d\lambda \\ &\leq Cr^{m+1}(t-r)^{\frac{1}{2}} \int_0^{t-r} \frac{1}{\sqrt{t-r-\lambda}}d\lambda \\ &\leq Cr^{m+1}\tau_-^{-m-2}. \end{aligned}$$

When $t-r \geq 2k$, using (3.3), the integration by parts and (2.13), we have

$$\begin{aligned} J_4(r, t) &= \int_0^{\frac{t-r}{2}} (\lambda^{2m}g(\lambda))' \tilde{K}_{m-1}(\lambda, r, t)d\lambda \\ &= - \int_0^{\frac{t-r}{2}} \lambda^{2m}g(\lambda)D_\lambda \tilde{K}_{m-1}(\lambda, r, t)d\lambda \\ &= -2 \int_0^{\frac{t-r}{2}} \lambda^{2m+1}g(\lambda)D_{\lambda^2} \tilde{K}_{m-1}(\lambda, r, t)d\lambda \\ &= 2 \int_0^{\frac{t-r}{2}} \lambda^{2m+1}g(\lambda) \tilde{K}_m(\lambda, r, t)d\lambda. \end{aligned}$$

By(2.18)_m with $\mu = 0$, we get

$$\begin{aligned} |J_4(r, t)| &\leq Cr^{m+1}(t-r)^{-m-\frac{1}{2}} \int_0^{\frac{t-r}{2}} \frac{\lambda^{2m+1}|g(\lambda)|}{(t-r-\lambda)^{\frac{3}{2}}}d\lambda \\ &\leq Cr^{m+1}(t-r)^{-m-2} \int_0^k \lambda^{2m+1}|g(\lambda)|d\lambda \\ &\leq Cr^{m+1}\tau_-^{-m-2}. \end{aligned}$$

For (3.2) with $|\beta| = 1$, we use (2.26). If we use

$$\begin{aligned} &(2.14)_{m-1} \text{ with } \mu = 0 \quad \text{for } J_5, \\ &(2.15)_{m-1} \text{ with } \mu = \frac{1}{2} \quad \text{for } J_6 \text{ with } t-r \leq 2k, \\ &(2.13) \text{ and } (2.17)_m \text{ with } \mu = 0 \quad \text{for } J_6 \text{ with } t-r \geq 2k, \end{aligned}$$

we can treat J_5 and J_6 by the same way as J_7 and J_8 for (3.1), respectively.

3.2 Case 2, $k \leq r \leq t \leq 2r$

In this case, it holds that

$$\tau_- \leq \tau_+ \leq 3\frac{r+2k}{k} \leq \frac{9r}{k}$$

for $(r, t) \in \Omega$. Hence, it is enough to show

$$|D^\beta D_t(2r^{2m}\Theta(f)(r, t))| \leq Cr^{m-\frac{1}{2}}\tau_-^{-m-\frac{1}{2}-|\beta|}, \quad (3.5)$$

$$|D^\beta(2r^{2m}\Theta(g)(r, t))| \leq Cr^{m-\frac{1}{2}}\tau_-^{-m-\frac{1}{2}-|\beta|}, \quad (3.6)$$

for $|\beta| = 0, 1$ in Ω .

For (3.5) with $|\beta| = 0$, we use (2.26). Using $(2.14)_{m-1}$ with $\mu = 0$,

$$J_5(r, t) = Cr^{m-\frac{1}{2}} \int_{t-r}^{t+r} \frac{\lambda^m |f(\lambda)| + \lambda^{m+1} |f'(\lambda)|}{\sqrt{\lambda - t + r}} d\lambda.$$

Since $J_5(r, t) = 0$ for $t - r \geq k$ by (3.3), we consider it when $t - r \leq k$. Then, we have by (3.3)

$$\begin{aligned} J_5(r, t) &\leq Cr^{m-\frac{1}{2}} \\ &\leq Cr^{m-\frac{1}{2}}\tau_-^{-m-\frac{1}{2}}. \end{aligned}$$

Next we consider J_6 . When $t - r \leq 2k$, using $(2.17)_{m-1}$ with $\mu = \frac{1}{2}$ and (3.3), we have

$$\begin{aligned} J_6(r, t) &\leq Cr^{m-\frac{1}{2}}(t-r)^{-m+1} \int_0^{t-r} \frac{\lambda^{2m-1}|f(\lambda)| + \lambda^{2m}|f'(\lambda)|}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}}(t-r) \int_0^{t-r} \frac{\lambda^{m-1}|f(\lambda)| + \lambda^m|f'(\lambda)|}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \\ &\leq Cr^{m-\frac{1}{2}}\tau_-^{-m-\frac{1}{2}}. \end{aligned}$$

When $t - r \geq 2k$, by the integration by parts, (3.3) and (2.13), we have

$$\begin{aligned} J_6(r, t) &= \int_0^{\frac{t-r}{2}} (\lambda^{2m} f(\lambda))' D_t \tilde{K}_{m-1}(\lambda, r, t) d\lambda \\ &= - \int_0^{\frac{t-r}{2}} \lambda^{2m} f(\lambda) D_\lambda D_t \tilde{K}_{m-1}(\lambda, r, t) d\lambda. \\ &= 2 \int_0^{\frac{t-r}{2}} \lambda^{2m+1} f(\lambda) D_t \tilde{K}_m(\lambda, r, t) d\lambda. \end{aligned}$$

Hence, by (2.17)_{m-1} with $\eta = \frac{1}{2}$, we have

$$\begin{aligned} |J_6(r, t)| &\leq Cr^{m-\frac{1}{2}}(t-r)^{-m} \int_0^{\frac{t-r}{2}} \frac{\lambda^{2m+1} f(\lambda)}{(t-r-\lambda)^{\frac{3}{2}}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}}(t-r)^{-m-\frac{3}{2}} \int_0^{\frac{t-r}{2}} \lambda^{2m+1} f(\lambda) d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \tau_-^{-m-\frac{1}{2}}. \end{aligned}$$

For (3.5) with $|\beta| = 1$, we use (2.30). Using (2.14)_{m-2} with $\mu = 0$, we have

$$|J_7(r, t)| \leq Cr^{m-\frac{1}{2}} \int_{t-r}^{t+r} \frac{\lambda^{m-1} |f(\lambda)| + \lambda^m |f'(\lambda)| + \lambda^{m+1} |f''(\lambda)|}{\sqrt{\lambda - t + r}} d\lambda.$$

When $t - r \geq k$, $J_7 = 0$. When $t - r \leq k$, we can treat it as before.

As to J_8 , dividing it as

$$\int_0^{t-r} d\lambda = \int_0^{\frac{t-r}{2}} d\lambda + \int_{\frac{t-r}{2}}^{t-r} d\lambda.$$

and calculating the first term by the same way as in (3.4), we have

$$\begin{aligned} J_8(r, t) &= \left[D_{\lambda^2}(\lambda^{2m} f(\lambda)) D^\beta D_t \tilde{K}_{m-2}(\lambda, r, t) \right]_{\lambda=0}^{\frac{t-r}{2}} \\ &\quad + \left[\lambda^{2m} f(\lambda) D^\beta D_t \tilde{K}_{m-1}(\lambda, r, t) \right]_{\lambda=0}^{\frac{t-r}{2}} \\ &\quad + 2 \int_0^{\frac{t-r}{2}} \lambda^{2m+1} f(\lambda) D^\beta D_t \tilde{K}_m(\lambda, r, t) d\lambda \\ &\quad + \int_{\frac{t-r}{2}}^{t-r} D_\lambda D_{\lambda^2}(\lambda^{2m} f(\lambda)) \cdot D^\beta D_t \tilde{K}_{m-2}(\lambda, r, t) d\lambda \\ &\equiv A_1 + A_2 + A_3 + A_4. \end{aligned}$$

For A_1 , using (2.15)_{m-2} with $\eta = 0$ to the term of $\lambda = \frac{t-r}{2}$, we have

$$|A_1| \leq Cr^{m-\frac{1}{2}} \left((t-r)^{m-\frac{1}{2}} \left| f\left(\frac{t-r}{2}\right) \right| + (t-r)^{m+\frac{1}{2}} \left| f'\left(\frac{t-r}{2}\right) \right| \right).$$

When $t - r \geq 2k$, $A_1 = 0$. When $t - r \leq 2k$, we have $|A_1| \leq Cr^{m-\frac{1}{2}}$.

For A_2 , using (2.17) with $\eta = \frac{1}{2}$ to the term of $\lambda = \frac{t-r}{2}$, we have

$$|A_2| \leq Cr^{m-\frac{1}{2}}(t-r)^{m-\frac{1}{2}} \left| f\left(\frac{t-r}{2}\right) \right|.$$

When $t - r \geq 2k$, $A_2 = 0$. When $t - r \leq 2k$, we have $|A_2| \leq Cr^{m-\frac{1}{2}}$.

For A_3 , using (2.17) with $\eta = \frac{1}{2}$, we have

$$\begin{aligned} |A_3| &\leq Cr^{m-\frac{1}{2}}(t-r)^{-m} \int_0^{\frac{t-r}{2}} \frac{\lambda^{2m+1} |f(\lambda)|}{(t-r-\lambda)^{\frac{5}{2}}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}}(t-r)^{-m-\frac{5}{2}} \int_0^{\frac{t-r}{2}} \lambda^{2m+1} |f(\lambda)| d\lambda. \end{aligned}$$

When $t - r \geq 2k$, we have

$$\begin{aligned} |A_3| &\leq Cr^{m-\frac{1}{2}}(t-r)^{-m-\frac{5}{2}} \int_0^k \lambda^{2m+1}|f(\lambda)|d\lambda \\ &\leq Cr^{m-\frac{1}{2}}\tau_-^{-m-\frac{5}{2}}. \end{aligned}$$

When $t - r \leq 2k$, we have

$$\begin{aligned} |A_3| &\leq Cr^{m-\frac{1}{2}}(t-r)^{m-\frac{3}{2}} \int_0^{\frac{t-r}{2}} |f(\lambda)|d\lambda \\ &\leq Cr^{m-\frac{1}{2}}. \end{aligned}$$

For A_4 , using (2.15) $_{m-2}$ with $\eta = 0$, we have

$$A_4 \leq Cr^{m-\frac{1}{2}}(t-r)^{-m+2} \int_{\frac{t-r}{2}}^{t-r} \frac{\lambda^{2m-3}|f(\lambda)| + \lambda^{2m-2}|f'(\lambda)| + \lambda^{2m-1}|f''(\lambda)|}{\sqrt{t-r-\lambda}} d\lambda.$$

When $t - r \geq 2k$, $B = 0$. When $t - r \leq 2k$, we have

$$\begin{aligned} A_4 &\leq Cr^{m-\frac{1}{2}} \int_{\frac{t-r}{2}}^{t-r} \frac{\lambda^{m-1}|f(\lambda)| + \lambda^m|f'(\lambda)| + \lambda^{m+1}|f''(\lambda)|}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \int_{\frac{t-r}{2}}^{t-r} \frac{1}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \end{aligned}$$

Therefore we get

$$\begin{aligned} |J_8(r, t)| &\leq \begin{cases} Cr^{m-\frac{1}{2}}\tau_-^{-m-\frac{5}{2}} & \text{if } t-r \geq 2k, \\ Cr^{m-\frac{1}{2}} & \text{if } t-r \leq 2k, \end{cases} \\ &\leq Cr^{m-\frac{1}{2}}\tau_-^{-m-\frac{3}{2}}. \end{aligned}$$

For (3.6) with $|\beta| = 0$, we use (2.4). Usin (2.14) $_m$ with $\eta = 0$, we have

$$|J_1(r, t)| \leq Cr^{m-\frac{1}{2}} \int_{t-r}^{t+r} \frac{\lambda^{m+1}|g(\lambda)|}{\sqrt{\lambda-t+r}} d\lambda.$$

When $t - r \geq k$, $J_1 = 0$. When $t - r \leq k$, we have

$$\begin{aligned} |J_1(r, t)| &\leq Cr^{m-\frac{1}{2}} \int_{t-r}^k \frac{1}{\sqrt{\lambda-t+r}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \\ &\leq Cr^{m-\frac{1}{2}}\tau_-^{-m-\frac{1}{2}}. \end{aligned}$$

Using (2.15)_m with $\eta = 0$, we have

$$|J_2(r, t)| \leq Cr^{m-\frac{1}{2}}(t-r)^{-m} \int_0^{t-r} \frac{\lambda^{2m+1}|g(\lambda)|}{\sqrt{t-r-\lambda}} d\lambda.$$

When $t-r \leq 2k$,

$$\begin{aligned} |J_2(r, t)| &\leq Cr^{m-\frac{1}{2}} \int_0^{t-r} \frac{\lambda^{m+1}|g(\lambda)|}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \int_0^{t-r} \frac{1}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \\ &\leq Cr^{m-\frac{1}{2}} \tau_-^{-m-\frac{1}{2}}. \end{aligned}$$

When $t-r \geq 2k$, we have by (3.3)

$$\begin{aligned} |J_2(r, t)| &\leq Cr^{m-\frac{1}{2}}(t-r)^{-m} \int_0^{\frac{t-r}{2}} \frac{\lambda^{m+1}|g(\lambda)|}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}}(t-r)^{-m-\frac{1}{2}} \int_0^k \lambda^{m+1}|g(\lambda)| d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \tau_-^{-m-\frac{1}{2}}. \end{aligned}$$

For (3.6) with $|\beta| = 1$, we use (2.26). Using (2.14)_{m-1} with $\eta = 0$, we have

$$|J_5(r, t)| \leq Cr^{m-\frac{1}{2}} \int_{t-r}^{t+r} \frac{\lambda^m|g(\lambda)| + \lambda^{m+1}|g'(\lambda)|}{\sqrt{\lambda-t+r}} d\lambda.$$

When $t-r \geq k$, $J_5 = 0$. When $t-r \leq k$,

$$\begin{aligned} |J_5(r, t)| &\leq Cr^{m-\frac{1}{2}} \int_{t-r}^k \frac{1}{\sqrt{\lambda-t+r}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \\ &\leq Cr^{m-\frac{1}{2}} \tau_-^{-m-\frac{3}{2}}. \end{aligned}$$

As to J_6 for $t-r \leq 2k$, (2.15)_{m-1} with $\eta = 0$ gives us

$$\begin{aligned} |J_6(r, t)| &\leq Cr^{m-\frac{1}{2}}(t-r)^{-m+1} \int_0^{t-r} \frac{\lambda^{2m-1}|g(\lambda)| + \lambda^{2m}|g'(\lambda)|}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \int_0^{t-r} \frac{\lambda^m|g(\lambda)| + \lambda^{m+1}|g'(\lambda)|}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \int_0^{t-r} \frac{1}{\sqrt{t-r-\lambda}} d\lambda \\ &\leq Cr^{m-\frac{1}{2}} \\ &\leq Cr^{m-\frac{1}{2}} \tau_-^{-m-\frac{3}{2}}. \end{aligned}$$

When $t - r \geq 2k$, using (3.3), (2.13) and the integration by parts, we have

$$\begin{aligned}
J_6(r, t) &= \int_0^{\frac{t-r}{2}} (\lambda^{2m} g(\lambda))' D_\alpha \tilde{K}_{m-1}(\lambda, r, t) d\lambda \\
&= - \int_0^{\frac{t-r}{2}} \lambda^{2m} g(\lambda) D_\lambda D_\alpha \tilde{K}_{m-1}(\lambda, r, t) d\lambda \\
&= 2 \int_0^{\frac{t-r}{2}} \lambda^{2m+1} g(\lambda) D_\alpha \tilde{K}_m(\lambda, r, t) d\lambda.
\end{aligned}$$

Then, by (2.17) with $\eta = \frac{1}{2}$, we have

$$\begin{aligned}
|J_6(r, t)| &\leq Cr^{m-\frac{1}{2}}(t-r)^{-m} \int_0^{\frac{t-r}{2}} \frac{\lambda^{2m+1}|g(\lambda)|}{(t-r-\lambda)^{\frac{3}{2}}} d\lambda \\
&\leq Cr^{m-\frac{1}{2}}(t-r)^{-m-\frac{3}{2}} \int_0^k \lambda^{2m+1}|g(\lambda)| d\lambda \\
&\leq Cr^{m-\frac{1}{2}}\tau_-^{-m-\frac{3}{2}}.
\end{aligned}$$

3.3 Case 3, $r > t$

In this case, it holds that $1 < \tau_- \leq 2$ and

$$\begin{aligned}
1 \leq \frac{r+2k}{k}, \quad \tau_-, \tau_+ \leq 4 &\quad \text{if } 0 < r \leq k, \\
\tau_+ \leq 2\frac{r+2k}{k} \leq \frac{6r}{k} &\quad \text{if } r \geq k,
\end{aligned}$$

for $(r, t) \in \Omega$. Hence, it is enough to show

$$|D^\beta D_t(2r^{2m}\Theta(f)(r, t))| \leq \begin{cases} Cr^{m+1-|\beta|} & \text{if } 0 < r \leq k, \\ Cr^{m-\frac{1}{2}} & \text{if } r \geq k, \end{cases} \quad (3.7)$$

$$|D^\beta(2r^{2m}\Theta(g)(r, t))| \leq \begin{cases} Cr^{m+1-|\beta|} & \text{if } 0 < r \leq k, \\ Cr^{m-\frac{1}{2}} & \text{if } r \geq k, \end{cases} \quad (3.8)$$

for $|\beta| = 0, 1$ in Ω .

For (3.7), we use (2.34) for $|\beta| = 0$ and the derivative of (2.34) for $|\beta| = 1$. Using (2.19) with $\eta = 0$, we have

$$|J_9(r, t)| \leq Cr^{m-\frac{1}{2}} \int_{r-t}^{r+t} \lambda^{m-\frac{1}{2}} |f(\lambda)| d\lambda.$$

When $0 < r \leq k$, we have

$$|J_9(r, t)| \leq Cr^{m-\frac{1}{2}}(r+t)^{\frac{3}{2}} \int_{r-t}^{r+t} \lambda^{m-2} |f(\lambda)| d\lambda \leq Cr^{m+1}.$$

When $r \geq k$, we have

$$|J_9(r, t)| \leq Cr^{m-\frac{1}{2}} \int_0^k \lambda^{m-\frac{1}{2}} |f(\lambda)| d\lambda \leq Cr^{m-\frac{1}{2}}.$$

Next, we consider $D_\alpha J_9(r, t)$. Differentiating J_9 and using (2.19) with $\mu = 0$, we have

$$\begin{aligned} |DJ_9(r, t)| &\leq \int_{r-t}^{r+t} \lambda^{2m+1} |f(\lambda)| |(D_\alpha \partial_t K_m)(\lambda, r, t)| d\lambda \\ &\quad + (r+t)^{2m+1} |f(r+t)| |(\partial_t K_m)(r+t, r, t)| \\ &\quad + (r-t)^{2m+1} |f(r-t)| |(\partial_t K_m)(r-t, r, t)| \\ &\leq Cr^{m-\frac{1}{2}} \left\{ \int_{r-t}^{r+t} \lambda^{m-\frac{3}{2}} |f(\lambda)| d\lambda + (r+t)^{m-\frac{1}{2}} |f(r+t)| + (r-t)^{m-\frac{1}{2}} |f(r-t)| \right\}. \end{aligned}$$

It follows from (3.3) that

$$|DJ_9(r, t)| \leq Cr^m \left\{ \int_{r-t}^{r+t} \lambda^{m-2} |f(\lambda)| d\lambda + (r+t)^{m-1} |f(r+t)| + (r-t)^{m-1} |f(r-t)| \right\} \leq Cr^m$$

for $0 < r \leq k$ and

$$|DJ_9(r, t)| \leq Cr^{m-\frac{1}{2}}.$$

for $r \geq k$.

Third, we consider J_{10} . Using (2.20) for $K_m(r+t, r, t)$ and (2.19) with $\eta = 0$ for $K_m(r-t, r, t)$, we have

$$|J_{10}(r, t)| \leq Cr^{m-\frac{1}{2}} \{ (r+t)^{m+\frac{1}{2}} |f(r+t)| + (r-t)^{m+\frac{1}{2}} |f(r-t)| \}.$$

It follows from (3.3) that

$$|J_{10}(r, t)| \leq Cr^{m+1} \{ (r+t)^{m-1} |f(r+t)| + (r-t)^{m-1} |f(r-t)| \} \leq Cr^{m+1}$$

for $0 < r \leq k$ and

$$|J_{10}(r, t)| \leq Cr^{m-\frac{1}{2}}$$

for $r \geq k$. Finally we consider $D_\alpha J_{10}$. Differentiating J_{10} , we have

$$\begin{aligned} |DJ_{10}(r, t)| &\leq \{ 2m(r+t)^{2m} |f(r+t)| + (r+t)^{2m+1} |f'(r+t)| \} |K_m(r+t, r, t)| \\ &\quad + (r+t)^{2m+1} |f(r+t)| \{ |(D_\lambda K_m)(r+t, r, t)| + |DK_m(r+t, r, t)| \} \\ &\quad + \{ 2m(r-t)^{2m} |f(r-t)| + (r-t)^{2m+1} |f'(r-t)| \} |K_m(r-t, r, t)| \\ &\quad + (r-t)^{2m+1} |f(r-t)| \{ |(D_\lambda K_m)(r-t, r, t)| + |DK_m(r-t, r, t)| \}. \end{aligned}$$

Using (2.20) for $D_\lambda K_m(r+t, r, t)$, $DK_m(r+t, r, t)$ and (2.19) with $\eta = 0$ for $D_\lambda K_m(r-t, r, t)$, $DK_m(r-t, r, t)$, we have

$$\begin{aligned} |D_\alpha J_{10}(r, t)| &\leq Cr^{m-\frac{1}{2}} \{ (r+t)^{m-\frac{1}{2}} |f(r+t)| + (r+t)^{m+\frac{1}{2}} |f'(r+t)| \\ &\quad + (r-t)^{m-\frac{1}{2}} |f(r-t)| + (r-t)^{m+\frac{1}{2}} |f'(r-t)| \} \end{aligned}$$

Treating this by the same way as J_{10} , we have the desired estimates.

For (3.8), we use (2.4) for $|\beta| = 0$ and the derivative of (2.4) for $|\beta| = 1$. By $r > t$, $J_2 \equiv 0$. So, we shall estimate J_1 and DJ_1 . Using (2.14) $_m$ with $\mu = 0$, we have

$$|J_1(r, t)| \leq Cr^{m-\frac{1}{2}} \int_{r-t}^{r+t} \lambda^{m+1} |g(\lambda)| d\lambda.$$

It follows from (3.3) that

$$|J_1(r, t)| \leq Cr^{m+1} \int_{r-t}^{r+t} \lambda^{m-\frac{1}{2}} |g(\lambda)| d\lambda \leq Cr^{m+1}$$

for $r \leq k$, while we simply have for $r \geq k$

$$|J_1(r, t)| \leq Cr^{m-\frac{1}{2}}.$$

Differentiating J_1 , we have

$$\begin{aligned} |D_\alpha J_1(r, t)| &\leq \int_{r-t}^{r+t} \lambda^{2m+1} |g(\lambda)| |(D_\alpha K_m)(\lambda, r, t)| d\lambda \\ &\quad + (r+t)^{2m+1} |g(r+t)| |K_m(r+t, r, t)| + (r-t)^{2m+1} |g(r-t)| |K_m(r-t, r, t)|. \end{aligned}$$

Hence using (2.20) for the second line and (2.19) with $\mu = 0$ for the others, we have

$$|D_\alpha J_1(r, t)| \leq Cr^{m-\frac{1}{2}} \left\{ \int_{r-t}^{r+t} \lambda^{m-\frac{1}{2}} |g(\lambda)| d\lambda + (r+t)^{m-\frac{1}{2}} |g(r+t)| + (r-t)^{m-\frac{1}{2}} |g(r-t)| \right\}.$$

Then, we can treat this by the same way as J_1 and get the desired estimate. This completes the proof of Proposition 3.1. \square

4 A priori estimates

In this section, we consider the nonlinear problems (1.9). Here and hereafter, we assume $p \leq q$ without loss of generality by the symmetricity of the systems. We will construct a solution to (1.9) on $\Omega_T := (0, \infty) \times [0, T]$ in a function space X defined by

$$\begin{aligned} X := \{ (u, v) \in (C((0, \infty) \times [0, T]))^2 : (\partial_r u, \partial_r v) \in (C((0, \infty) \times [0, T]))^2 \\ \text{supp } u \cup \text{supp } v \subset \{r \leq t + k\}, \quad \|u\|_1 + \|v\|_2 < \infty \}, \end{aligned}$$

where

$$\|u\|_1 := \sum_{i=0}^1 \sup_{(r,t) \in \Omega_T} w_i(r, t) |D_r^i u(r, t)|, \quad (4.1)$$

$$\|v\|_2 := \sum_{i=0}^1 \sup_{(r,t) \in \Omega_T} z_i(r, t) |D_r^i v(r, t)|. \quad (4.2)$$

The weight functions w_i, z_i are defined by

$$w_i(r, t) := \left(\frac{r}{k}\right)^{m-1+i} \left(\frac{r+2k}{k}\right)^{1-i} \tau_+^{\frac{1}{2}} W(r, t), \quad (4.3)$$

$$z_i(r, t) := \left(\frac{r}{k}\right)^{m-1+i} \left(\frac{r+2k}{k}\right)^{1-i} \tau_+^{\frac{1}{2}} Z(r, t) \quad (4.4)$$

$$(4.5)$$

for $i = 0, 1$, where

$$W(r, t) := \tau_-^{(m+\frac{1}{2})p-(m+\frac{3}{2})}, \quad (4.6)$$

$$Z(r, t) := \begin{cases} \tau_-^{\frac{1}{p}} (\log 3\tau_-)^\nu & \text{if } F = 0, p \neq q \\ \tau_-^\mu & \text{otherwise,} \end{cases} \quad (4.7)$$

with

$$\mu := \frac{1}{p} - (q-1)F(p, q; n), \quad (4.8)$$

$$\nu := \frac{q(p-1)}{p(pq-1)}. \quad (4.9)$$

Let us consider the sequences $\{U_j, V_j\}$ defined by

$$\begin{aligned} U_1 &= 0, & U_{j+1} &= L(|v_0 + V_j|^p) \quad (j \geq 1), \\ V_1 &= 0, & V_{j+1} &= L(|u_0 + U_j|^q) \quad (j \geq 1), \end{aligned} \quad (4.10)$$

where u_0, v_0 and L are defined in (2.1), (2.2) and (2.3), respectively. We will show the convergence of these sequences in X . If we prove the existence of the limit functions U and V , we can show the existence of solution u and v to (1.9), since $u = \varepsilon u^0 + U, v = \varepsilon v^0 + V$.

First, recall the representation formula of $L(G)$ in (2.3) and the expressions of Θ (2.4), (2.23), (2.26). Then, we have

$$r^{2m} L(G)(r, t) = \int_0^t \left\{ \tilde{J}_1(G(\cdot, \tau))(r, t - \tau) + \tilde{J}_2(G(\cdot, \tau))(r, t - \tau) \right\} d\tau, \quad (4.11)$$

$$2r^{2m} L(G)(r, t) = \int_0^t \left\{ \tilde{J}_3(G(\cdot, \tau))(r, t - \tau) + \tilde{J}_4(G(\cdot, \tau))(r, t - \tau) \right\} d\tau, \quad (4.12)$$

$$\partial_r(2r^{2m} L(G)(r, t)) = \int_0^t \left\{ \tilde{J}_5(G(\cdot, \tau))(r, t - \tau) + \tilde{J}_6(G(\cdot, \tau))(r, t - \tau) \right\} d\tau, \quad (4.13)$$

where \tilde{J}_i ($i = 1, 2, \dots, 6$) is defined by replacing g by $G(\cdot, \tau)$ in (2.5), (2.6), (2.24), (2.25), (2.27), (2.28). According to the basic attitude and the choice of the estimates of K, \tilde{K} in the proof of Lemma 6.3, Lemma 6.4, Lemma 6.5 of [16], we have

$$\begin{aligned} |L(G)(r, t)| &\leq \frac{C}{r^{m+1/2-\eta}} \int_0^t d\tau \int_{|\lambda_-|}^{\lambda_+} \frac{\lambda^{m+1-\eta} |G(\lambda, \tau)|}{\sqrt{\tau + \lambda - t + r}} d\lambda \\ &\quad + \frac{C}{r^{m+1/2-\eta}} \int_0^{(t-r)_+} (t-r-\tau)^{-m-\eta} d\tau \int_0^{\lambda_-} \frac{\lambda^{2m+1} |G(\lambda, \tau)|}{\sqrt{t-r-\tau-\lambda}} d\lambda, \end{aligned} \quad (4.14)$$

by (4.11) and

$$\begin{aligned}
|\partial_r^i L(G)(r, t)| &\leq \frac{C}{r^{m-1/2-\eta+i}} \int_0^t d\tau \int_{|\lambda_-|}^{\lambda_+} \frac{|\partial_\lambda(\lambda^{2m} G(\lambda, \tau))| \lambda^{-m+1-\eta}}{\sqrt{\tau + \lambda - t + r}} d\lambda \\
&+ \frac{C}{r^{m-1/2-\eta+i}} \int_0^{(t-r)_+} (t-r-\tau)^{m+1/2-\eta} |G(\lambda_-/2, \tau)| d\tau \\
&+ \frac{C}{r^{m-1/2-\eta+i}} \int_0^{(t-r)_+} (t-r-\tau)^{-m+1-\eta} d\tau \int_0^{\lambda_-/2} \frac{\lambda^{2m} |G(\lambda, \tau)|}{(t-r-\tau-\lambda)^{3/2}} d\lambda \\
&+ \frac{C}{r^{m-1/2-\eta+i}} \int_0^{(t-r)_+} (t-r-\tau)^{-m+1-\eta} d\tau \int_{\lambda_-/2}^{\lambda_-} \frac{|\partial_\lambda(\lambda^{2m} G(\lambda, \tau))|}{\sqrt{\lambda_- - \lambda}} d\lambda \\
&+ \frac{C}{r^{m-1/2-\eta+i}} \int_{t-r}^t (\tau+r-t)^{m+1/2-\eta} |G(\tau+r-t, \tau)| d\tau \tag{4.15}
\end{aligned}$$

by (4.12) and (4.13), where $\eta = 0$ or $\frac{1}{2}$, $\lambda_\pm = t - \tau \pm r$ and $(a)_+ = \max\{a, 0\}$.

To show that the sequence $(\{U_j, V_j\})$ in (4.10) converges in X , it is necessary to consider (4.14) and (4.15) with $G = G_1, G_2, G_3, G_4$ below.

$$\begin{aligned}
G_1 &:= |v_0 + V_j|^p, \\
G_2 &:= |u_0 + U_j|^q, \\
G_3 &:= |v_0 + V_j|^p - |v_0 + V_{j-1}|^p, \\
G_4 &:= |u_0 + U_j|^q - |u_0 + U_{j-1}|^q.
\end{aligned}$$

First, it holds that

$$\begin{aligned}
|G_1| &\leq C(|v_0|^p + |V_j|^p), \\
|G_2| &\leq C(|u_0|^q + |U_j|^q), \\
|G_3| &\leq C|V_j - V_{j-1}|(|v_0|^{p-1} + \tilde{V}_j^{p-1}), \\
|G_4| &\leq C|U_j - U_{j-1}|(|u_0|^{q-1} + \tilde{U}_j^{q-1}),
\end{aligned}$$

and

$$\begin{aligned}
|\partial_\lambda G_1| &\leq C(|v_0|^{p-1} + |V_j|^{p-1})(|\partial_\lambda v_0| + |\partial_\lambda V_j|), \\
|\partial_\lambda G_2| &\leq C(|u_0|^{q-1} + |U_j|^{q-1})(|\partial_\lambda u_0| + |\partial_\lambda U_j|), \\
|\partial_\lambda G_3| &\leq C|\partial_\lambda V_j - \partial_\lambda V_{j-1}|(|v_0|^{p-1} + |V_j|^{p-1}) + C(|\partial_\lambda v_0| + |\partial_\lambda V_{j-1}|)|V_j - V_{j-1}|^{p-1}, \\
|\partial_\lambda G_4| &\leq C|\partial_\lambda U_j - \partial_\lambda U_{j-1}|(|u_0|^{q-1} + |U_j|^{q-1}) + C(|\partial_\lambda u_0| + |\partial_\lambda U_{j-1}|)|U_j - U_{j-1}|^{q-1},
\end{aligned}$$

where $\tilde{V}_j = \max\{|V_j|, |V_{j-1}|\}$, $\tilde{U}_j = \max\{|U_j|, |U_{j-1}|\}$. Here the second terms of $\partial_\lambda G_3$ and $\partial_\lambda G_4$ come from Hölder continuity, since $1 < p, q < 2$.

To close the iteration, we introduce the auxiliary norm $\|\cdot\|$ as

$$\|u\|_1 := \sup_{(r,t) \in \Omega_T} w_1(r, t) |u(r, t)|, \tag{4.16}$$

$$\|v\|_2 := \sup_{(r,t) \in \Omega_T} z_1(r, t) |v(r, t)|. \tag{4.17}$$

according to [16]. Hence, Using Proposition 3.1, (4.1), (4.2), (4.16) and (4.17), we get the followings by (4.14).

$$|U_{j+1}| \leq \frac{C}{r^{m+\frac{1}{2}-\eta}} \left[\varepsilon^p \sum_{k=1}^2 I_k(\phi_0^{-p}, p) + \|V_j\|_2^p \sum_{k=1}^2 I_k(Z^{-p}, p) \right] \quad (4.18)$$

$$|V_{j+1}| \leq \frac{C}{r^{m+\frac{1}{2}-\eta}} \left[\varepsilon^q \sum_{k=1}^2 I_k(\phi_0^{-q}, q) + \|U_j\|_1^q \sum_{k=1}^2 I_k(W^{-q}, q) \right] \quad (4.19)$$

$$|U_{j+1} - U_j| \leq \frac{C}{r^{m+\frac{1}{2}-\eta}} \|V_j - V_{j-1}\|_2 \left[\varepsilon^{p-1} \sum_{k=1}^2 I_k(\phi_0^{1-p} Z^{-1}, p) + \|\tilde{V}_j\|_2^{p-1} \sum_{k=1}^2 I_k(Z^{-p}, p) \right] \quad (4.20)$$

$$|V_{j+1} - V_j| \leq \frac{C}{r^{m+\frac{1}{2}-\eta}} \|U_j - U_{j-1}\|_1 \left[\varepsilon^{q-1} \sum_{k=1}^2 I_k(\phi_0^{1-q} W^{-1}, q) + \|\tilde{U}_j\|_1^{q-1} \sum_{k=1}^2 I_k(W^{-q}, q) \right] \quad (4.21)$$

$$|U_{j+1} - U_j| \leq \frac{C}{r^{m+\frac{1}{2}-\eta}} \|V_j - V_{j-1}\|_2 \left[\varepsilon^{p-1} \sum_{k=1}^2 I_k(\phi_0^{1-p} Z^{-1}, p) + \|\tilde{V}_j\|_2^{p-1} \sum_{k=1}^2 I_k(Z^{-p}, p) \right] \quad (4.22)$$

$$|V_{j+1} - V_j| \leq \frac{C}{r^{m+\frac{1}{2}-\eta}} \|U_j - U_{j-1}\|_1 \left[\varepsilon^{q-1} \sum_{k=1}^2 I_k(\phi_0^{1-q} W^{-1}, q) + \|\tilde{U}_j\|_1^{q-1} \sum_{k=1}^2 I_k(W^{-q}, q) \right], \quad (4.23)$$

where

$$I_1(\Phi, p)(r, t) := \int_{-k}^{t-r} d\beta \int_{|t-r|}^{t+r} \frac{\left(\frac{\alpha-\beta}{k}\right)^{m+1-mp-\eta} \left(\frac{\alpha+2k}{k}\right)^{-p/2} \Phi\left(\frac{\alpha-\beta}{2}, \frac{\alpha+\beta}{2}\right)}{\sqrt{\alpha-t+r}} d\alpha \quad (4.24)$$

$$I_2(\Phi, p)(r, t) := \int_0^{t-r} \frac{\left(\frac{\alpha+2k}{k}\right)^{-p/2}}{\sqrt{t-r-\alpha}} d\alpha \quad (4.25)$$

$$\times \int_{-k}^{\alpha} \left(\frac{\alpha-\beta}{k}\right)^{2m-mp} \left(t-r-\frac{\alpha+\beta}{2}\right)^{-m+1-\eta} \Phi\left(\frac{\alpha-\beta}{2}, \frac{\alpha+\beta}{2}\right) d\beta, \quad (4.26)$$

provided $\text{supp } U_j \cup \text{supp } V_j \subset \{r \leq t+k\}$ ($j \geq 0$). Similarly, it follows from (4.15) that

$$|D_{r,t}^i U_{j+1}| \leq \frac{C}{r^{m-\frac{1}{2}-\eta+i}} \quad (4.27)$$

$$\begin{aligned} & \times \left[\varepsilon^p \sum_{k=1}^4 I_k(\phi_0^{-p}, p) + \varepsilon^p \sum_{k=1}^2 I_k(\phi_0^{1-p} \phi_1^{-1}, p) + \varepsilon^{p-1} \|V_j\|_2 \sum_{k=1}^2 I_k(\phi_0^{1-p} Z^{-1}, p) \right. \\ & \left. + \varepsilon \|V_j\|_2^{p-1} \sum_{k=1}^2 I_k(\phi_0^{-1} Z^{1-p}, p) + \|V_j\|_2^p \sum_{k=1}^4 I_k(Z^{-p}, p) \right], \quad (4.28) \end{aligned}$$

$$|D_{r,t}^i V_{j+1}| \leq \frac{C}{r^{m-\frac{1}{2}-\eta+i}} \quad (4.29)$$

$$\begin{aligned} & \times \left[\varepsilon^q \sum_{k=1}^4 I_k(\phi_0^{-q}, q) + \varepsilon^q \sum_{k=1}^2 I_k(\phi_0^{1-q} \phi_1^{-1}, q) + \varepsilon^{q-1} \|U_j\|_1 \sum_{k=1}^2 I_k(\phi_0^{1-q} W^{-1}, q) \right. \\ & \left. + \varepsilon \|U_j\|_1^{q-1} \sum_{k=1}^2 I_k(\phi_0^{-1} W^{1-q}, q) + \|U_j\|_1^q \sum_{k=1}^4 I_k(W^{-q}, q) \right]. \quad (4.30) \end{aligned}$$

$$\begin{aligned} & |D_{r,t}^i U_{j+1} - D_{r,t}^i U_j| \\ & \leq \frac{C}{r^{m-\frac{1}{2}-\eta+i}} \|V_j - V_{j-1}\|_2 \left[\varepsilon^{p-1} \sum_{k=1}^4 I_k(\phi_0^{1-p} Z^{-1}, p) + \|V_j\|_2^{p-1} \sum_{k=1}^4 I_k(Z^{-p}, p) \right] \\ & \quad + \frac{C}{r^{m-\frac{1}{2}-\eta+i}} \|V_j - V_{j-1}\|_2^{p-1} \left[\varepsilon \sum_{k=1}^2 I_k(\phi_1^{-1} Z^{1-p}, p) + \|V_{j-1}\|_2 \sum_{k=1}^2 I_k(Z^{-p}, p) \right], \quad (4.31) \end{aligned}$$

$$\begin{aligned} & |D_{r,t}^i V_{j+1} - D_{r,t}^i V_j| \\ & \leq \frac{C}{r^{m-\frac{1}{2}-\eta+i}} \|U_j - U_{j-1}\|_1 \left[\varepsilon^{q-1} \sum_{k=1}^4 I_k(\phi_0^{1-q} W^{-1}, q) + \|U_j\|_1^{q-1} \sum_{k=1}^4 I_k(W^{-q}, q) \right] \\ & \quad + \frac{C}{r^{m-\frac{1}{2}-\eta+i}} \|U_j - U_{j-1}\|_1^{q-1} \left[\varepsilon \sum_{k=1}^2 I_k(\phi_1^{-1} W^{1-q}, q) + \|U_{j-1}\|_1 \sum_{k=1}^2 I_k(W^{-q}, q) \right], \quad (4.32) \end{aligned}$$

where

$$I_3(\Phi, p)(r, t) := \int_{\frac{t-r-2k}{3}}^{t-r} \left(\frac{\lambda_-}{k} \right)^{m+\frac{1}{2}-(m-1)p-\eta} \left(\frac{\lambda_- + 2k}{k} \right)^{-p} \left(\frac{\tau + \frac{\lambda_-}{2} + 2k}{k} \right)^{-p/2} \Phi \left(\frac{\lambda_-}{2}, \tau \right) d\tau \quad (4.33)$$

$$I_4(\Phi, p)(r, t) := \int_{t-r}^t \left(\frac{-\lambda_-}{k} \right)^{m+\frac{1}{2}-(m-1)p-\eta} \left(\frac{-\lambda_- + 2k}{k} \right)^{-p} \left(\frac{\tau - \lambda_- + 2k}{k} \right)^{-p/2} \Phi(-\lambda_-, \tau) d\tau, \quad (4.34)$$

provided $\text{supp } U_j \cup \text{supp } V_j \subset \{r \leq t + k\}$ ($j \geq 0$). We have the following estimates for I_k ($k = 1, 2, 3, 4$).

Proposition 4.1 *Let $m \geq 2$, $\frac{2m+3}{2m+1} < p, q < \frac{2m+5}{2m+1}$ and $F(p, q; n) \geq 0$. Then, we have*

$$\begin{aligned} |I_k(\phi_0^{-p}, p)| &\leq C\tau_-^{-\eta} W(r, t)^{-1}, & |I_k(\phi_0^{-q}, q)| &\leq C\tau_-^{-\eta} Z(r, t)^{-1}, \\ |I_k(\phi_0^{1-p} \phi_1^{-1}, p)| &\leq C\tau_-^{-\eta} W(r, t)^{-1}, & |I_k(\phi_0^{1-q} \phi_1^{-1}, q)| &\leq C\tau_-^{-\eta} Z(r, t)^{-1}, \\ |I_k(\phi_0^{1-p} Z^{-1}, p)| &\leq C\tau_-^{-\eta} W(r, t)^{-1}, & |I_k(\phi_0^{1-q} W^{-1}, q)| &\leq C\tau_-^{-\eta} Z(r, t)^{-1}, \\ |I_k(\phi_0^{-1} Z^{1-p}, p)| &\leq C\tau_-^{-\eta} W(r, t)^{-1}, & |I_k(\phi_0^{-1} W^{1-q}, q)| &\leq C\tau_-^{-\eta} Z(r, t)^{-1}, \\ |I_k(\phi_1^{-1} Z^{1-p}, p)| &\leq C\tau_-^{-\eta} W(r, t)^{-1}, & |I_k(\phi_1^{-1} W^{1-q}, q)| &\leq C\tau_-^{-\eta} Z(r, t)^{-1}, \\ |I_k(Z^{-p}, p)| &\leq C\tau_-^{-\eta} W(r, t)^{-1} E_1(T), & |I_k(W^{-q}, q)| &\leq C\tau_-^{-\eta} Z(r, t)^{-1} E_2(T), \end{aligned}$$

in Ω_T for $\eta = 0, \frac{1}{2}$ and $k = 1, 2$, where

$$E_1(T) := \begin{cases} \left(\log \frac{T+2k}{k}\right)^{1-p\nu} & \text{if } F(p, q; n) = 0, p \neq q, \\ \log \frac{T+2k}{k} & \text{if } F(p, q; n) = 0, p = q, \\ \left(\frac{T+2k}{k}\right)^{p(q-1)F(p, q; n)} & \text{if } F(p, q; n) > 0, \end{cases} \quad (4.35)$$

$$E_2(T) := \begin{cases} \left(\log \frac{T+2k}{k}\right)^\nu & \text{if } F(p, q; n) = 0, p \neq q, \\ \log \frac{T+2k}{k} & \text{if } F(p, q; n) = 0, p = q, \\ \left(\frac{T+2k}{k}\right)^{q(p-1)F(p, q; n)} & \text{if } F(p, q; n) > 0, \end{cases} \quad (4.36)$$

Proposition 4.2 *Let $m \geq 2$, $\frac{2m+3}{2m+1} < p, q < \frac{2m+5}{2m+1}$ and $F(p, q; n) \geq 0$. Then, we have*

$$\begin{aligned} |I_k(\phi_0^{-p}, p)| &\leq C\tau_-^{-\eta}W(r, t)^{-1}, & |I_k(\phi_0^{-q}, q)| &\leq C\tau_-^{-\eta}Z(r, t)^{-1}, \\ |I_k(\phi_0^{1-p}\phi_1^{-1}, p)| &\leq C\tau_-^{-\eta}W(r, t)^{-1}, & |I_k(\phi_0^{1-q}\phi_1^{-1}, q)| &\leq C\tau_-^{-\eta}Z(r, t)^{-1}, \\ |I_k(Z^{-p}, p)| &\leq C\tau_-^{-\eta}W(r, t)^{-1}E_1(T), & |I_k(S(W^{-q}, q))| &\leq C\tau_-^{-\eta}Z(r, t)^{-1}E_2(T), \end{aligned}$$

in Ω_T for $\eta = 0, \frac{1}{2}$ and $k = 3, 4$, where $E_1(T)$ and $E_2(T)$ are defined in (4.35) and (4.36), respectively.

Proposition 4.1 and Proposition 4.2 will be proved in §6. Here, we shall use two Propositions and proceed to a priori estimates.

In (4.18)-(4.23) together with Proposition 4.1, we can find $r^{-m-\frac{1}{2}+\eta}\tau_-^{-\eta}$ in all the inequalities. Similarly, (4.28)-(4.32) with Proposition 4.1, Proposition 4.2 has $r^{-m+\frac{1}{2}+\eta-i}$ in common. For them, we can treat as follows.

Lemma 4.1 *Let $(r, t) \in \Omega_T$ with $t - r \geq -k$. Assume that I, J_i ($i = 0, 1$) satisfy*

$$I \leq r^{-m-\frac{1}{2}+\eta}\tau_-^{-\eta} \quad \text{for } \eta = 0, \frac{1}{2} \quad (4.37)$$

$$J_i \leq r^{-m+\frac{1}{2}+\eta-i}\tau_-^{-\eta} \quad \text{for } \eta = 0, \frac{1}{2}, \quad i = 0, 1 \quad (4.38)$$

for $\eta = 0, \frac{1}{2}$. Then, we have

$$I \leq C \left(\frac{r}{k}\right)^{-m+1} \left(\frac{r+2k}{k}\right)^{-1} \tau_+^{-\frac{1}{2}} \quad \text{for } r \geq k, \quad (4.39)$$

$$I \leq C \left(\frac{r}{k}\right)^{-m} \tau_+^{-\frac{1}{2}} \quad \text{for } r > 0, \quad (4.40)$$

$$J_i \leq C \left(\frac{r}{k}\right)^{-m+1-i} \left(\frac{r+2k}{k}\right)^{-1+i} \tau_+^{-\frac{1}{2}} \quad \text{for "i = 0 and } r \leq k" \text{ or } i = 1. \quad (4.41)$$

Proof. Taking η suitably as in the table below, one can easily prove them.

Area of (r, t)	$r \geq k, t \leq 2r$	$r \geq k, t \geq 2r$	$r \leq k, t \leq 2r$	$r \leq k, t \geq 2r$
features	$\frac{9r}{k} \geq \frac{3(r+2k)}{k} \geq \tau_+$	$\frac{3r}{k} \geq \frac{r+2k}{k}$ $3\tau_- \geq \tau_+$	$1 \leq \tau_- \leq \tau_+$ $\leq \frac{3(r+2k)}{k} \leq 9$	$2 \leq \frac{r+2k}{k} \leq 3$ $3\tau_- \geq \tau_+$
For (4.39)	(4.37), $\eta = 0$	(4.37), $\eta = \frac{1}{2}$	-	-
For (4.40)			(4.37), $\eta = \frac{1}{2}$	(4.37), $\eta = \frac{1}{2}$
For (4.41), $j = 0$	-	-	(4.38), $\eta = \frac{1}{2}$	(4.38), $\eta = \frac{1}{2}$
For (4.41), $j = 1$	(4.38), $\eta = 0$	(4.38), $\eta = \frac{1}{2}$		

Now we are in a position to state a priori estimates which will give us the local existence of solution and the lower bound of the lifespan. For the norms (4.1) and (4.2), we shall apply When $r \geq k$, we use (4.18), (4.19), (4.20) and (4.21) and apply Proposition 4.1, (4.39) and the definition of weight functions (4.3), (4.4) to them. For “ $i = 0$ and $r \leq k$ ” or $i = 1$ in (4.28)(4.30)(4.31)(4.32), we shall apply Proposition 4.1, Proposition 4.2 and (4.41) to them. As to the auxiliary norms (4.16), (4.17), we use (4.22)(4.23) with Proposition 4.1, (4.40). Then, we have the following.

Proposition 4.3 *Let $(U_j, V_j) \in X$ be the one in (4.10) for $j \geq 1$. Assume that $\frac{2m+3}{2m+1} < p \leq q < \frac{2m+5}{2m+1}$ and $F(p, q; n) \geq 0$. Then, the following inequalities hold.*

$$\|U_{j+1}\|_1 \leq A\varepsilon^p + C\varepsilon^{p-1}\|V_j\|_2 + C\varepsilon\|V_j\|_2^{p-1} + C\|V_j\|_2^p E_1(T), \quad (4.42)$$

$$\|V_{j+1}\|_2 \leq A\varepsilon^q + C\varepsilon^{q-1}\|U_j\|_1 + C\varepsilon\|U_j\|_1^{q-1} + C\|U_j\|_1^q E_2(T), \quad (4.43)$$

$$\begin{aligned} \|U_{j+1} - U_j\|_1 &\leq C\|V_j - V_{j-1}\|_2(\varepsilon^{p-1} + \|\tilde{V}_j\|_2^{p-1} E_1(T)) \\ &\quad + C\|V_j - V_{j-1}\|_2^{p-1}(\varepsilon + \|V_{j-1}\|_2 E_1(T)), \end{aligned} \quad (4.44)$$

$$\begin{aligned} \|V_{j+1} - V_j\|_2 &\leq C\|U_j - U_{j-1}\|_1(\varepsilon^{q-1} + \|\tilde{U}_j\|_1^{q-1} E_2(T)) \\ &\quad + C\|U_j - U_{j-1}\|_1^{q-1}(\varepsilon + \|U_{j-1}\|_1 E_2(T)), \end{aligned} \quad (4.45)$$

$$\|U_{j+1} - U_j\|_1 \leq C\|V_j - V_{j-1}\|_2(\varepsilon^{p-1} + \|\tilde{V}_j\|_2^{p-1} E_1(T)), \quad (4.46)$$

$$\|V_{j+1} - V_j\|_2 \leq C\|U_j - U_{j-1}\|_1(\varepsilon^{q-1} + \|\tilde{U}_j\|_1^{q-1} E_2(T)), \quad (4.47)$$

where $A \geq 1$ and $C \geq 1$ are constants independent of ε .

5 Construction of a solution and Lifespan

Theorem 1 can be proved by the iteration methods using Proposition 4.3. For this, we shall define a closed subspace Y of X as follows.

$$Y := \{(U, V) \in X : \|U\|_1 \leq 2A\varepsilon^p, \|V\|_2 \leq 2A\varepsilon^q\},$$

where A is the one in Proposition 4.3. We will find the conditions which guarantee that $\{U_j, V_j\}$ is a convergent sequence in Y .

First, we note that

$$(U_1, V_1) = (0, 0) \in Y.$$

Assume that $\{(U_j, V_j)\} \subset Y$. Then, we have

$$(U_{j+1}, V_{j+1}), (\partial_r U_{j+1}, \partial_r V_{j+1}) \in (C((0, \infty) \times [0, T]))^2, \quad (5.1)$$

$$\text{supp } U_{j+1} \cup \text{supp } V_{j+1} \subset \{r \leq t + k\}, \quad (5.2)$$

by the definition of the sequences (4.10), Proposition 3.1 and the representation formulas (4.12), (4.13) with (2.24), (2.25), (2.27), (2.28). Moreover, we have by (4.42), (4.43),

$$\begin{aligned} \|U_{j+1}\|_1 &\leq A\varepsilon^p + C\varepsilon^{p-1}2A\varepsilon^q + C\varepsilon(2A\varepsilon^q)^{p-1} + C(2A\varepsilon^q)^p E_1(T), \\ \|V_{j+1}\|_2 &\leq A\varepsilon^q + C\varepsilon^{q-1}2A\varepsilon^p + C\varepsilon(2A\varepsilon^p)^{q-1} + C(2A\varepsilon^p)^q E_2(T). \end{aligned}$$

Thus, if we assume that

$$C\varepsilon^{p-1}2A\varepsilon^q + C\varepsilon(2A\varepsilon^q)^{p-1} + C(2A\varepsilon^q)^p E_1(T) \leq A\varepsilon^p, \quad (5.3)$$

$$C\varepsilon^{q-1}2A\varepsilon^p + C\varepsilon(2A\varepsilon^p)^{q-1} + C(2A\varepsilon^p)^q E_2(T) \leq A\varepsilon^q, \quad (5.4)$$

then, $(U_{j+1}, V_{j+1}) \in Y$. By the induction,

$$(U_j, V_j) \in Y \quad \text{for any } j \geq 1. \quad (5.5)$$

Hereafter, we assume that (5.3) and (5.4) are fulfilled.

Next, we consider the differences of $\{(U_m, V_m)\}$. As to the norm $\|\cdot\|$, (4.46), (4.47) and (5.5) give us

$$\begin{aligned} \|U_{j+1} - U_j\|_1 &\leq C\|V_j - V_{j-1}\|_2(\varepsilon^{p-1} + (2A\varepsilon^q)^{p-1}E_1(T)), \\ \|V_{j+1} - V_j\|_2 &\leq C\|U_j - U_{j-1}\|_1(\varepsilon^{q-1} + (2A\varepsilon^p)^{q-1}E_2(T)). \end{aligned}$$

Thus, if we assume that

$$C^2\{\varepsilon^{p-1} + C(2A\varepsilon^q)^{p-1}E_1(T)\}\{\varepsilon^{q-1} + C(2A\varepsilon^p)^{q-1}E_2(T)\} \leq \frac{1}{4}, \quad (5.6)$$

then, we have

$$\|U_{j+1} - U_j\|_1 \leq \frac{1}{4}\|U_{j-1} - U_{j-2}\|_1, \quad \|V_{j+1} - V_j\|_2 \leq \frac{1}{4}\|V_{j-1} - V_{j-2}\|_2$$

which means that

$$\|U_{j+1} - U_j\|_1 \leq \frac{B}{2^j}, \quad \|V_{j+1} - V_j\|_2 \leq \frac{B}{2^j}, \quad (5.7)$$

where we set $B = \max\{1, 4\|U_3 - U_2\|_1, 2\|U_2 - U_1\|_1, 4\|V_3 - V_2\|_2, 2\|V_2 - V_1\|_2\} \geq 1$. Hereafter, we assume (5.7).

As to the norm $\|\cdot\|$, by (4.44), (4.46), (5.5) and (5.7), we have

$$\begin{aligned} \|U_{j+1} - U_j\|_1 &\leq C\|V_j - V_{j-1}\|_2(\varepsilon^{p-1} + (2A\varepsilon^q)^{p-1}E_1(T)) + C\left(\frac{B}{2^{j-1}}\right)^{p-1}(\varepsilon + 2A\varepsilon^q E_1(T)), \\ \|V_{j+1} - V_j\|_2 &\leq C\|U_j - U_{j-1}\|_1(\varepsilon^{q-1} + (2A\varepsilon^p)^{q-1}E_2(T)) + C\left(\frac{B}{2^{j-1}}\right)^{q-1}(\varepsilon + 2A\varepsilon^p E_2(T)). \end{aligned}$$

Here we set $s = 2^{1-p} < 1$. If we assume that

$$C^2(\varepsilon^{p-1} + (2A\varepsilon^q)^{p-1}E_1(T))(\varepsilon^{q-1} + (2A\varepsilon^p)^{q-1}E_2(T)) \leq s^2 \quad (5.8)$$

$$C^2B^{q-1}(\varepsilon + 2A\varepsilon^pE_2(T))(\varepsilon^{p-1} + (2A\varepsilon^q)^{p-1}E_1(T)) \leq \frac{s^2}{2}, \quad (5.9)$$

$$CB^{p-1}(\varepsilon + 2A\varepsilon^qE_1(T)) \leq \frac{s}{2} \quad (5.10)$$

$$C^2B^{p-1}(\varepsilon + 2A\varepsilon^qE_1(T))(\varepsilon^{q-1} + (2A\varepsilon^p)^{q-1}E_2(T)) \leq \frac{s^2}{2}, \quad (5.11)$$

$$CB^{q-1}(\varepsilon + 2A\varepsilon^pE_2(T)) \leq \frac{s}{2}, \quad (5.12)$$

then, we have

$$\|U_{j+1} - U_j\|_1 \leq s^2\|U_{j-1} - U_{j-2}\|_1 + s^j, \quad \|V_{j+1} - V_j\|_2 \leq s^2\|V_{j-1} - V_{j-2}\|_2 + s^j,$$

which mean that

$$\|U_{j+1} - U_j\|_1 \leq B's^j + js^j, \quad \|V_{j+1} - V_j\|_2 \leq B's^j + js^j,$$

where we set $B' = \max\{s^{-2}\|U_3 - U_2\|_1, s^{-1}\|U_2 - U_1\|_1, s^{-2}\|V_3 - V_2\|_2, s^{-1}\|V_2 - V_1\|_2\}$. Since the right-hand sides in the above inequalities converge, we find that $\{U_j\}$ and $\{V_j\}$ converge with respect to the norm $\|\cdot\|$. Hence, we now know that $\{(U_j, V_j)\}$ is a convergent sequence in Y , provided (5.3), (5.4), (5.6), (5.8), (5.9), (5.10), (5.11) and (5.12) are satisfied.

Let us define

$$\varepsilon_1 := \min \left\{ (6C)^{-\frac{1}{p-1}}, (2^{q-1}3CA^{q-2})^{-\frac{1}{(p-1)(q-1)}}, (8C^2B^{p-1}s^{-2})^{-\frac{1}{q}}, (4CB^{q-1}s^{-1})^{-1} \right\}$$

and

$$E := \min \left\{ (2^q3A^{q-1}C)^{-1}, (2^{q+1}A^{q-1}C^2s^{-2})^{-1}, (2^{p+2}A^{p-1}C^2B^{q-1}s^{-2})^{-1}, (2^{q+2}A^{q-1}C^2B^{p-1}s^{-2})^{-1}, (8ACB^{q-1}s^{-1})^{-1} \right\} < 1. \quad (5.13)$$

Then, we see that for any $\varepsilon \in (0, \varepsilon_1)$, all of the conditions above are satisfied, provided

$$\varepsilon^{p(q-1)}E_1(T) \leq E, \quad (5.14)$$

$$\varepsilon^{q(p-1)}E_2(T) \leq E. \quad (5.15)$$

Note that we used the fact that $\varepsilon < \varepsilon^{p-1}$ for $1 < p < 2$ and $0 < \varepsilon < 1$ in the calculations above.

Now we can get the lower bound of the lifespan by (5.14) and (5.15). For example, let us see the case where $F = 0$, $p \neq q$. (5.14) and (5.15) can be rewritten by (4.35) and (4.36) as

$$\varepsilon^{p(q-1)} \left(\log \frac{T+2k}{k} \right)^{1-p\nu} \leq E, \quad \varepsilon^{q(p-1)} \left(\log \frac{T+2k}{k} \right)^\nu \leq E. \quad (5.16)$$

By the definition of ν (4.9), one can see that these are guaranteed by

$$T \leq k \exp \left(E^{\frac{p(pq-1)}{q(p-1)}} \varepsilon^{-p(pq-1)} \right) - 2k, \quad (5.17)$$

since $0 < E < 1$ and $p \leq q$. Besides, defining ε_2 so that

$$k \exp\left(E^{\frac{p(pq-1)}{q(p-1)}} \varepsilon_2^{-p(pq-1)}\right) - 2k = \exp\left(E^{\frac{p(pq-1)}{q(p-1)}} \varepsilon_2^{-p(pq-1)}\right),$$

we see that

$$T \leq \exp\left(E^{\frac{p(pq-1)}{q(p-1)}} \varepsilon^{-p(pq-1)}\right) \quad \text{for any } \varepsilon \in (0, \min\{\varepsilon_1, \varepsilon_2\}],$$

is a sufficient condition for (5.17).

Similarly, we can consider the other cases. Therefore, we find that there exists a positive constant $\varepsilon_0 = \varepsilon_0(p, q, f_1, f_2, g_1, g_2, n, k)$ such that, for any ε satisfying $0 < \varepsilon \leq \varepsilon_0$, a local-in-time solution to (1.9) in Y exists, provided

$$T \leq \begin{cases} \exp(\tilde{C}\varepsilon^{-p(pq-1)}) & \text{if } F = 0, p \neq q, \\ \exp(\tilde{C}\varepsilon^{-p(p-1)}) & \text{if } F = 0, p = q, \\ \tilde{C}\varepsilon^{-F(p,q;n)^{-1}} & \text{if } F > 0. \end{cases}$$

This implies the lower bound of the lifespan.

6 Proof of A priori estimates

In this section, we shall prove Proposition 4.1 and Proposition 4.2. As to Proposition 4.1, it is enough to consider I_k ($k = 1, 2, 3, 4$) only for

$$\begin{aligned} \phi_0^{-p}, \quad \phi_0^{1-p}Z^{-1}, \quad \phi_0^{-1}Z^{1-p}, \quad Z^{-p}, \\ \phi_0^{-q}, \quad \phi_0^{1-q}W^{-1}, \quad \phi_0^{-1}W^{1-q}, \quad W^{-q}, \end{aligned} \quad (6.1)$$

since $\phi_1^{-1} = \tau_-^{-(m+\frac{3}{2})} \leq \tau_-^{-(m+\frac{1}{2})} = \phi_0^{-1}$. The calculation in this section is essentially based on the way of Proposition 6.6 in [16].

Before the proofs, we remark the properties of μ in (4.8) and ν in (4.9).

Remark 6.1 *Let $F(p, q; n)$ with $\frac{2m+3}{2m+1} < p \leq q < \frac{2m+5}{2m+1}$. Then, we have*

$$0 < \mu < \frac{1}{p}, \quad 0 < \nu < \frac{1}{p}, \quad (6.2)$$

$$1 - p\mu = p(q-1)F(p, q; n), \quad 1 - p\nu = \frac{q-1}{pq-1}, \quad (6.3)$$

$$m + \frac{3}{2} - \left(m + \frac{1}{2}\right)q + \mu = \frac{(p-q)(pq+1)}{p(pq-1)} \leq 0, \quad (6.4)$$

$$- \left\{ \left(m + \frac{1}{2}\right)p - \left(m + \frac{3}{2}\right) \right\} q + m + \frac{5}{2} - \left(m + \frac{1}{2}\right)q = -\mu + q(p-1)F(p, q; n). \quad (6.5)$$

One can readily prove them by the definitions of μ and ν .

Hereafter, we assume that $m \geq 2$, $\frac{2m+3}{2m+1} < p \leq q < \frac{2m+5}{2m+1}$ and $F(p, q; n) \geq 0$. Proposition 4.1 will be proved in **6.1**, **6.2** and Proposition 4.2 will be proved in **6.3**, **6.4**.

6.1 Estimates for I_1

The first line of (6.1)

When $F = 0$ with $p \neq q$, we have by (4.24), (4.7) and $\phi_0 = \tau_-^{-m-\frac{1}{2}}$,

$$|I_1(\phi_0^{-p}, p)| \leq \int_{-k}^{t-r} \left(\frac{\beta + 2k}{k} \right)^{-(m+\frac{1}{2})p} d\beta \int_{|t-r|}^{t+r} \frac{\left(\frac{\alpha-\beta}{k} \right)^{m+1-mp-\eta} \left(\frac{\alpha+2k}{k} \right)^{-\frac{p}{2}}}{\sqrt{\alpha - (t-r)}} d\alpha \quad (6.6)$$

$$\begin{aligned} |I_1(\phi_0^{1-p} Z^{-1}, p)| &\leq \int_{-k}^{t-r} \left(\frac{\beta + 2k}{k} \right)^{-(m+\frac{1}{2})(p-1)-\frac{1}{p}} \left(\log 3 \frac{\beta + 2k}{k} \right)^{-\nu} d\beta \\ &\quad \times \int_{|t-r|}^{t+r} \frac{\left(\frac{\alpha-\beta}{k} \right)^{m+1-mp-\eta} \left(\frac{\alpha+2k}{k} \right)^{-\frac{p}{2}}}{\sqrt{\alpha - (t-r)}} d\alpha \end{aligned} \quad (6.7)$$

$$\begin{aligned} |I_1(\phi_0^{-1} Z^{1-p}, p)| &\leq \int_{-k}^{t-r} \left(\frac{\beta + 2k}{k} \right)^{-(m+\frac{1}{2})-\frac{p-1}{p}} \left(\log 3 \frac{\beta + 2k}{k} \right)^{-(p-1)\nu} d\beta \\ &\quad \times \int_{|t-r|}^{t+r} \frac{\left(\frac{\alpha-\beta}{k} \right)^{m+1-mp-\eta} \left(\frac{\alpha+2k}{k} \right)^{-\frac{p}{2}}}{\sqrt{\alpha - (t-r)}} d\alpha \end{aligned} \quad (6.8)$$

$$\begin{aligned} |I_1(Z^{-p}, p)| &\leq \int_{-k}^{t-r} \left(\frac{\beta + 2k}{k} \right)^{-1} \left(\log 3 \frac{\beta + 2k}{k} \right)^{-p\nu} d\beta \\ &\quad \times \int_{|t-r|}^{t+r} \frac{\left(\frac{\alpha-\beta}{k} \right)^{m+1-mp-\eta} \left(\frac{\alpha+2k}{k} \right)^{-\frac{p}{2}}}{\sqrt{\alpha - (t-r)}} d\alpha. \end{aligned} \quad (6.9)$$

We see that the α -integrals in four inequqlities are common. By

$$\left(\frac{\alpha + 2k}{k} \right)^{-\frac{p}{2}} \leq \tau_-^{-\eta} \left(\frac{\alpha - \beta}{k} \right)^{-\frac{p}{2}+\eta}$$

and the integration by parts, we have

$$\begin{aligned} (\alpha\text{-integral}) &\leq C\tau_-^{-\eta} \left\{ \sqrt{r} \left(\frac{t+r-\beta}{k} \right)^{m+1-(m+\frac{1}{2})p} + \int_{|t-r|}^{t+r} \left(\frac{\alpha-\beta}{k} \right)^{m+\frac{1}{2}-(m+\frac{1}{2})p} d\alpha \right\} \\ &\leq C\tau_-^{-\eta} \left(\frac{t-r-\beta}{k} \right)^{m+\frac{3}{2}-(m+\frac{1}{2})p}, \end{aligned}$$

since $-1 < m + \frac{3}{2} - (m + \frac{1}{2})p < 0$. Thus, for example, (6.6) becomes

$$|I_1(\phi_0^{-p}, p)| \leq C\tau_-^{-\eta} \int_{-k}^{t-r} \left(\frac{\beta + 2k}{k} \right)^{-(m+\frac{1}{2})p} \left(\frac{t-r-\beta}{k} \right)^{m+\frac{3}{2}-(m+\frac{1}{2})p} d\beta. \quad (6.10)$$

Here, the following Lemma is useful.

Lemma 6.1 *Let $k > 0$, $-k < a$, $l > 0$ and $0 < h < 1$. Then we have*

$$\int_{-k}^a \left(\frac{\beta + 2k}{k} \right)^{-l} \left(\frac{a-\beta}{k} \right)^{-h} d\beta \leq Ck \left(\frac{a+2k}{k} \right)^{-h} \tilde{E}(a),$$

where

$$\tilde{E}(a) := \begin{cases} 1 & \text{if } l > 1, \\ \log \frac{a+2k}{k} & \text{if } l = 1, \\ \left(\frac{a+2k}{k}\right)^{1-l} & \text{if } l < 1. \end{cases}$$

Proof. Lemma 4.4 in [18] is similar to this lemma and we can find its proof there. So, we omit the proof here. \square

Using Lemma 6.1 with $a = t - r$, $h = (m + \frac{1}{2})p - (m + \frac{3}{2}) \in (0, 1)$, $l = (m + \frac{1}{2})p > 1$ for (6.10), we have

$$|I_1(\phi_0^{-p}, p)| \leq C\tau_-^{-\eta+m+\frac{3}{2}-(m+\frac{1}{2})p} = C\tau_-^{-\eta}W(r, t)^{-1}.$$

As to (6.7) and (6.8), removing $\log \frac{\beta+2k}{k}$ and taking $l = (m+\frac{1}{2})(p-1)+\frac{1}{p} > 1$, $l = m+\frac{1}{2}+\frac{p-1}{p} > 1$ respectively in applying Lemma 6.1, we have the same conclusion as (6.6). (6.9) became needs the following Lemma.

Lemma 6.2 *Let $k > 0$, $-k < a$, $0 < l < 1$ and $0 < h < 1$. Then we have*

$$\int_{-k}^a \left(\frac{\beta+2k}{k}\right)^{-1} \left(\log 3\frac{\beta+2k}{k}\right)^{-l} \left(\frac{a-\beta}{k}\right)^{-h} d\beta \leq Ck \left(\frac{a+2k}{k}\right)^{-h} \left(\log 3\frac{a+2k}{k}\right)^{1-l}.$$

Proof. This lemma can be proved by the same division as the one in the proof of Lemma 6.1. So, we omit the proof here. \square

Applying Lemma 6.2 with $a = t - r$, $h = (m + \frac{1}{2})p - (m + \frac{3}{2}) \in (0, 1)$, $l = p\nu \in (0, 1)$, we get

$$|I_1(Z^{-p}, p)| \leq C\tau_-^{-\eta+m+\frac{3}{2}-(m+\frac{1}{2})p+1-p\nu} \leq C\tau_-^{-\eta}W(r, t)^{-1}E_1(T).$$

When $F = 0$ with $p = q$ or $F > 0$, we have

$$|I_1(\phi_0^{-p}, p)|, |I_1(\phi_0^{1-p}Z^{-1}, p)|, |I_1(\phi_0^{-1}Z^{1-p}, p)| \leq C\tau_-^{-\eta}W(r, t)^{-1},$$

by the similar calculation. $I_1(Z^{-p}, p)$ has the following form.

$$|I_1(Z^{-p}, p)| \leq \int_{-k}^{t-r} \left(\frac{\beta+2k}{k}\right)^{-p\mu} d\beta \int_{|t-r|}^{t+r} \frac{\left(\frac{\alpha-\beta}{k}\right)^{m+1-mp-\eta} \left(\frac{\alpha+2k}{k}\right)^{-\frac{p}{2}}}{\sqrt{\alpha-(t-r)}} d\alpha. \quad (6.11)$$

After the same treatment for the α -integral, we shall apply Lemma 6.1 with $a = t - r$, $h = (m + \frac{1}{2})p - (m + \frac{3}{2}) \in (0, 1)$ and $l = p\mu$ to the β -integral. Noting that $l = 1$ if $F(p, q; n) = 0$ and $l < 1$ if $F(p, q; n) > 0$ because of (6.3), we have

$$\begin{aligned} |I_1(Z^{-p}, p)| &\leq C\tau_-^{-\eta+m+\frac{3}{2}-(m+\frac{1}{2})p} \times \begin{cases} \log \tau_- & \text{if } F(p, q; n) = 0, p = q \\ \tau_-^{1-p\mu} & \text{if } F(p, q; n) > 0 \end{cases} \\ &\leq C\tau_-^{-\eta}W(r, t)^{-1}E_1(T). \end{aligned}$$

The second line of (6.1)

By the similar way to the above case, we have

$$|I_1(\phi_0^{-q}, q)|, |I_1(\phi_0^{1-q}W^{-1}, q)|, |I_1(\phi_0^{-1}W^{1-q}, q)| \leq C\tau_-^{-\eta+m+\frac{3}{2}-(m+\frac{1}{2})q}.$$

Hence, we get the desired estimates by (6.4). As to $I_1(W^{-q}, q)$, in applying Lemma 6.1, we take $l = \{(m + \frac{1}{2})p - (m + \frac{3}{2})\}q$. Using (6.5) and (6.4) in turn, we see that

$$\begin{aligned} l &= m + \frac{5}{2} - (m + \frac{1}{2})q + \mu - q(p-1)F(p, q; n) \\ &= 1 + \frac{(p-q)(pq+1)}{p(pq-1)} - q(p-1)F(p, q; n) \end{aligned} \quad (6.12)$$

which yields that $l = 1$ if $F(p, q; n) = 0$, $p = q$ and $l < 1$ if $F(p, q; n) > 0$. Thus we get

$$|I_1(W^{-q}, q)| \leq C\tau_-^{-\eta+m+\frac{3}{2}-(m+\frac{1}{2})q} \times \begin{cases} \log \tau_- & \text{if } F(p, q; n) = 0, p = q \\ \tau_-^{1-\{(m+\frac{1}{2})p-(m+\frac{3}{2})\}q} & \text{Otherwise.} \end{cases}$$

Using (6.4) if $F(p, q; n) = 0$, $p = q$ and (6.5) if $F(p, q; n) > 0$, this is dominated by $C\tau_-^{-\eta}Z(r, t)^{-1}E_2(T)$. When $F = 0$, $p \neq q$, noting that $\mu = \frac{1}{p}$ by (6.3) and using (6.5), we have

$$\begin{aligned} |I_1(W^{-q}, q)| &\leq C\tau_-^{-\eta-\frac{1}{p}} \\ &= C\tau_-^{-\eta-\frac{1}{p}}(\log 3\tau_-)^{-\nu}(\log 3\tau_-)^{\nu} \\ &\leq C\tau_-^{-\eta}Z(r, t)^{-1}E_2(T). \end{aligned}$$

Therefore Proposition 4.1 for I_1 has been proved.

6.2 Estimates for I_2

Here, we assume that $t - r > 0$. We shall divide the α -integral in (4.26) as follows.

$$\int_0^{t-r} d\alpha = \int_0^{(\frac{t-r}{2}-k)_+} + \int_{\frac{t-r}{2}-k}^{t-r} d\alpha. \quad (6.13)$$

Since it holds that

$$\begin{aligned} t-r - \frac{\alpha + \beta}{2} &\geq t-r - \alpha, \\ t-r - \frac{\alpha + \beta}{2} &\geq \frac{\alpha - \beta}{2}, \\ t-r - \frac{\alpha + \beta}{2} &\leq C\tau_- \end{aligned}$$

in the domain of the integration, we have two estimates:

$$\begin{aligned} &\left(\frac{\alpha - \beta}{k}\right)^{2m-mp} \left(t-r - \frac{\alpha + \beta}{2}\right)^{-m+1-\eta} \\ &\leq \begin{cases} C\tau_-^{\frac{1}{2}-\eta}(t-r - \alpha)^{m+\frac{3}{2}-(m+\frac{1}{2})p-\delta} \left(\frac{\alpha - \beta}{k}\right)^{\frac{p}{2}-1+\delta}, \\ C\tau_-^{m+\frac{5}{2}-(m+\frac{1}{2})p}(t-r - \alpha)^{-\frac{1}{2}+\delta} \left(\frac{\alpha - \beta}{k}\right)^{\frac{p}{2}-1-\eta-\delta}, \end{cases} \end{aligned} \quad (6.14)$$

for $0 < \delta < 1/2$ and $\eta = 0$ or $1/2$. Using the first line of (6.14) for the first term in (6.13) and the second line of (6.14) for the second term in (6.13), we have

$$\begin{aligned}
|I_2(\Phi, p)| &= C\tau_-^{\frac{1}{2}-\eta} \int_0^{(\frac{t-r}{2}-k)_+} \left(\frac{\alpha+2k}{k}\right)^{-p/2} (t-r-\alpha)^{m+1-(m+\frac{1}{2})p-\delta} d\alpha \\
&\quad \times \int_{-k}^\alpha \left(\frac{\alpha-\beta}{k}\right)^{\frac{p}{2}-1+\delta} \Phi\left(\frac{\alpha-\beta}{2}, \frac{\alpha+\beta}{2}\right) d\beta \\
&+ C\tau_-^{m+\frac{5}{2}-(m+\frac{1}{2})p} \int_{\frac{t-r}{2}-k}^{t-r} \left(\frac{\alpha+2k}{k}\right)^{-p/2} (t-r-\alpha)^{-1+\delta} d\alpha \\
&\quad \times \int_{-k}^\alpha \left(\frac{\alpha-\beta}{k}\right)^{\frac{p}{2}-1-\eta-\delta} \Phi\left(\frac{\alpha-\beta}{2}, \frac{\alpha+\beta}{2}\right) d\beta. \tag{6.15}
\end{aligned}$$

The first line of (6.1)

When $F = 0$, $p \neq q$, we have by (6.15), (4.7) and $\phi_0 = \tau_-^{-m-\frac{1}{2}}$,

$$\begin{aligned}
|I_2(\phi_0^{-p}, p)| &= C\tau_-^{\frac{1}{2}-\eta} \int_0^{(\frac{t-r}{2}-k)_+} \left(\frac{\alpha+2k}{k}\right)^{-p/2} (t-r-\alpha)^{m+1-(m+\frac{1}{2})p-\delta} d\alpha \\
&\quad \times \int_{-k}^\alpha \left(\frac{\alpha-\beta}{k}\right)^{\frac{p}{2}-1+\delta} \left(\frac{\beta+2k}{k}\right)^{-(m+\frac{1}{2})p} d\beta \\
&+ C\tau_-^{m+\frac{5}{2}-(m+\frac{1}{2})p} \int_{\frac{t-r}{2}-k}^{t-r} \left(\frac{\alpha+2k}{k}\right)^{-p/2} (t-r-\alpha)^{-1+\delta} d\alpha \\
&\quad \times \int_{-k}^\alpha \left(\frac{\alpha-\beta}{k}\right)^{\frac{p}{2}-1-\eta-\delta} \left(\frac{\beta+2k}{k}\right)^{-(m+\frac{1}{2})p} d\beta, \tag{6.16}
\end{aligned}$$

$$\begin{aligned}
|I_2(\phi_0^{1-p}Z^{-1}, p)| &= C\tau_-^{\frac{1}{2}-\eta} \int_0^{(\frac{t-r}{2}-k)_+} \left(\frac{\alpha+2k}{k}\right)^{-p/2} (t-r-\alpha)^{m+1-(m+\frac{1}{2})p-\delta} d\alpha \\
&\quad \times \int_{-k}^\alpha \left(\frac{\alpha-\beta}{k}\right)^{\frac{p}{2}-1+\delta} \left(\frac{\beta+2k}{k}\right)^{-(m+\frac{1}{2})(p-1)-\frac{1}{p}} \left(\log 3\frac{\beta+2k}{k}\right)^{-\nu} d\beta \\
&+ C\tau_-^{m+\frac{5}{2}-(m+\frac{1}{2})p} \int_{\frac{t-r}{2}-k}^{t-r} \left(\frac{\alpha+2k}{k}\right)^{-p/2} (t-r-\alpha)^{-1+\delta} d\alpha \\
&\quad \times \int_{-k}^\alpha \left(\frac{\alpha-\beta}{k}\right)^{\frac{p}{2}-1-\eta-\delta} \left(\frac{\beta+2k}{k}\right)^{-(m+\frac{1}{2})(p-1)-\frac{1}{p}} \left(\log 3\frac{\beta+2k}{k}\right)^{-\nu} d\beta, \tag{6.17}
\end{aligned}$$

$$\begin{aligned}
|I_2(\phi_0^{-1}Z^{1-p}, p)| &= C\tau_-^{\frac{1}{2}-\eta} \int_0^{(\frac{t-r}{2}-k)^+} \left(\frac{\alpha+2k}{k}\right)^{-p/2} (t-r-\alpha)^{m+1-(m+\frac{1}{2})p-\delta} d\alpha \\
&\quad \times \int_{-k}^\alpha \left(\frac{\alpha-\beta}{k}\right)^{\frac{p}{2}-1+\delta} \left(\frac{\beta+2k}{k}\right)^{-(m+\frac{1}{2})-\frac{p-1}{p}} \left(\log 3\frac{\beta+2k}{k}\right)^{-(p-1)\nu} d\beta \\
&+ C\tau_-^{m+\frac{5}{2}-(m+\frac{1}{2})p} \int_{\frac{t-r}{2}-k}^{t-r} \left(\frac{\alpha+2k}{k}\right)^{-p/2} (t-r-\alpha)^{-1+\delta} d\alpha \\
&\quad \times \int_{-k}^\alpha \left(\frac{\alpha-\beta}{k}\right)^{\frac{p}{2}-1-\eta-\delta} \left(\frac{\beta+2k}{k}\right)^{-(m+\frac{1}{2})-\frac{p-1}{p}} \left(\log 3\frac{\beta+2k}{k}\right)^{-(p-1)\nu} d\beta,
\end{aligned} \tag{6.18}$$

$$\begin{aligned}
|I_2(Z^{-p}, p)| &= C\tau_-^{\frac{1}{2}-\eta} \int_0^{(\frac{t-r}{2}-k)^+} \left(\frac{\alpha+2k}{k}\right)^{-p/2} (t-r-\alpha)^{m+1-(m+\frac{1}{2})p-\delta} d\alpha \\
&\quad \times \int_{-k}^\alpha \left(\frac{\alpha-\beta}{k}\right)^{\frac{p}{2}-1+\delta} \left(\frac{\beta+2k}{k}\right)^{-1} \left(\log 3\frac{\beta+2k}{k}\right)^{-p\nu} d\beta \\
&+ C\tau_-^{m+\frac{5}{2}-(m+\frac{1}{2})p} \int_{\frac{t-r}{2}-k}^{t-r} \left(\frac{\alpha+2k}{k}\right)^{-p/2} (t-r-\alpha)^{-1+\delta} d\alpha \\
&\quad \times \int_{-k}^\alpha \left(\frac{\alpha-\beta}{k}\right)^{\frac{p}{2}-1-\eta-\delta} \left(\frac{\beta+2k}{k}\right)^{-1} \left(\log 3\frac{\beta+2k}{k}\right)^{-p\nu} d\beta.
\end{aligned} \tag{6.19}$$

for $0 < \delta < \frac{1}{2}$.

Let us Remove $\log \frac{\beta+2k}{k}$ in (6.17), (6.18). Then, (6.16), (6.17) and (6.18) can be treated by the same way. As to the first term, setting $a = \alpha$, $h = -\frac{p}{2} + 1 - \delta$, we have $l > 1$, $h \in (0, 1)$ provided $0 < \delta < \frac{1}{10}$. Thus, applying Lemma 6.1 with a, h in common, and $l = (m + \frac{1}{2})p > 1$, $l = (m + \frac{1}{2})(p - 1) + \frac{1}{p} > 1$, $l = m + \frac{1}{2} + \frac{p-1}{p} > 1$ in turn, we can calculate them as follows.

$$\begin{aligned}
&(\text{the first tems of 6.16), (6.17), (6.18)}) \\
&\leq C\tau_-^{\frac{1}{2}-\eta} \int_0^{(\frac{t-r}{2}-k)^+} \left(\frac{\alpha+2k}{k}\right)^{-1+\delta} (t-r-\alpha)^{m+1-(m+\frac{1}{2})p-\delta} d\alpha \\
&\leq C\tau_-^{-\eta+m+\frac{3}{2}-(m+\frac{1}{2})p-\delta} \int_0^{(\frac{t-r}{2}-k)^+} \left(\frac{\alpha+2k}{k}\right)^{-1+\delta} d\alpha \\
&\leq C\tau_-^{-\eta+m+\frac{3}{2}-(m+\frac{1}{2})p} \\
&\leq C\tau_-^{-\eta} W(r, t)^{-1}.
\end{aligned}$$

As to the second term, setting $a = \alpha$ and $h = -\frac{p}{2} + 1 + \eta + \delta$, we have $h \in (0, 1)$ provided

$0 < \delta < \frac{1}{2m+1}$. Taking same l as the first terms and applying Lemma 6.1, we have

$$\begin{aligned}
& \text{(the second term of 6.16), (6.17), (6.18))} \\
& \leq C\tau_-^{m+\frac{5}{2}-(m+\frac{1}{2})p} \int_{\frac{t-r}{2}-k}^{t-r} \left(\frac{\alpha+2k}{k}\right)^{-1-\eta-\delta+\delta'} (t-r-\alpha)^{-1+\delta} d\alpha \\
& \leq C\tau_-^{-\mu+m+\frac{3}{2}-(m+\frac{1}{2})p-\delta} \int_{\frac{t-r}{2}-k}^{t-r} (t-r-\alpha)^{-1+\delta} d\alpha \\
& \leq C\tau_-^{-\mu+m+\frac{3}{2}-(m+\frac{1}{2})p} \\
& = C\tau_-^{-\eta}W(r,t)^{-1}.
\end{aligned}$$

Therefore we get

$$|I_2(\phi_0^{-p}, p)|, |I_2(\phi_0^{1-p}Z^{-1}, p)|, |I_2(\phi_0^{-1}Z^{1-p}, p)| \leq C\tau_-^{-\eta}W(r,t)^{-1}.$$

As to (6.19), we do not remove $\log \frac{\beta+2k}{k}$ and apply Lemma 6.2. Since $-\frac{p}{2} + 1 - \delta$ and $-\frac{p}{2} + 1 + \eta + \delta$ belong to $(0, 1)$ provided $0 < \delta < \frac{1}{2m+1}$, we set them as h for the first and the second term, respectively. Besides taking $a = \alpha$ and $l = p\nu < 1$ which comes from (6.2), we can apply Lemma 6.2 to the β -integrals and calculate (6.19) as follows.

$$\begin{aligned}
& |I_2(Z^{-p}, p)| \\
& = C\tau_-^{\frac{1}{2}-\eta} \int_0^{(\frac{t-r}{2}-k)_+} \left(\frac{\alpha+2k}{k}\right)^{-1+\delta} (t-r-\alpha)^{m+1-(m+\frac{1}{2})p-\delta} \left(\log 3\frac{\alpha+2k}{k}\right)^{1-p\nu} d\alpha \\
& \quad + C\tau_-^{m+\frac{5}{2}-(m+\frac{1}{2})p} \int_{\frac{t-r}{2}-k}^{t-r} \left(\frac{\alpha+2k}{k}\right)^{-1-\eta-\delta} (t-r-\alpha)^{-1+\delta} \left(\log 3\frac{\alpha+2k}{k}\right)^{1-p\nu} d\alpha \\
& \leq C\tau_-^{-\eta+m+\frac{3}{2}-(m+\frac{1}{2})p-\delta} (\log 3\tau_-)^{1-p\nu} \int_0^{(\frac{t-r}{2}-k)_+} \left(\frac{\alpha+2k}{k}\right)^{-1+\delta} d\alpha \\
& \quad + C\tau_-^{-\eta+m+\frac{3}{2}-(m+\frac{1}{2})p-\delta} (\log 3\tau_-)^{1-p\nu} \int_{\frac{t-r}{2}-k}^{t-r} (t-r-\alpha)^{-1+\delta} d\alpha \\
& \leq C\tau_-^{-\eta}W(r,t)^{-1}E_1(T).
\end{aligned}$$

When $F(p, q; n) = 0$ with $p = q$ or $F(p, q; n) > 0$, we can treat them similarly, since ϕ_0 and $Z(r, t)$ consist of τ_- only. Using Lemma 6.1 for all of four, we can get the desired estimates. The second line of (6.1)

Since ϕ_0 and W consist of τ_- only, we can calculate them by the same way as above and use (6.12) for $I_2(W^{-q}, q)$. Then, we have

$$|I_2(\phi_0^{-q}, q)|, |I_2(\phi_0^{1-q}W^{-1}, q)|, |I_2(\phi_0^{-1}W^{1-q}, q)| \leq C\tau_-^{-\eta+m+\frac{3}{2}-(m+\frac{1}{2})q}, \quad (6.20)$$

$$|I_2(W^{-q}, q)| \leq C\tau_-^{-\eta+m+\frac{3}{2}-(m+\frac{1}{2})q} \times \begin{cases} \log \tau_- & \text{if } F = 0, p = q \\ \tau_-^{1-\{(m+\frac{1}{2})p-(m+\frac{3}{2})\}q} & \text{Otherwise.} \end{cases} \quad (6.21)$$

We shall use (6.4) for the first three estimates. As to the fourth, we can treat it by the same way for the corresponding case for I_1 . Then we get the desired estimates and prove Proposition 4.1 for I_2 . This completes the proof of Proposition 4.1.

6.3 Estimates for I_3

In this subsection, we assume $t - r > 0$. We can calculate all the inequalities by the same way, we consider $I_3(\phi_0^{-p}, p)$ for the case of $F = 0$ with $p \neq q$ as an example. By (4.33) and $\phi_0 = \tau^{-m-\frac{1}{2}}$, we have

$$|I_3(\phi_0^{-p}, p)| \leq \int_{\frac{t-r-2k}{3}}^{t-r} \left(\frac{t-r-\tau}{k} \right)^{m+\frac{1}{2}-(m-1)p-\eta} \left(\frac{t-r-\tau+2k}{k} \right)^{-p} \\ \times \left(\frac{\tau + \frac{\lambda_-}{2} + 2k}{k} \right)^{-\frac{p}{2}} \left(\frac{\tau - \frac{\lambda_-}{2} + 2k}{k} \right)^{-(m+\frac{1}{2})p} d\tau. \quad (6.22)$$

By $t - r > 0$, we shall divide it as follows.

$$\int_{\frac{t-r-2k}{3}}^{t-r} d\tau = \int_{\frac{t-r-2k}{3}}^{t-r-\frac{2}{3}k} d\tau + \int_{t-r-\frac{2}{3}k}^{t-r} d\tau =: A + B \quad (6.23)$$

In the domain of integration of A , $t - r - \tau$ is equivalent to $t - r - \tau + 2k$. By

$$\left(\frac{\tau + \frac{\lambda_-}{2} + 2k}{k} \right)^{-\frac{p}{2}} \leq C\tau_-^{-\eta} \left(\frac{t-r-\tau+2k}{k} \right)^{-\frac{p}{2}+\eta},$$

we have

$$A \leq C\tau_-^{-\eta} \int_{\frac{t-r-2k}{3}}^{t-r-\frac{2}{3}k} \left(\frac{t-r-\tau+2k}{k} \right)^{m+\frac{1}{2}-(m+\frac{1}{2})p} \left(\frac{3\tau - (t-r) + 4k}{k} \right)^{-(m+\frac{1}{2})p} d\tau.$$

Moreover, dividing it as

$$\int_{\frac{t-r-2k}{3}}^{t-r-\frac{2}{3}k} d\tau = \int_{\frac{t-r-2k}{3}}^{\frac{t-r}{2}-\frac{2}{3}k} d\tau + \int_{\frac{t-r}{2}-\frac{2}{3}k}^{t-r-\frac{2}{3}k} d\tau =: A_1 + A_2,$$

each integral can be calculated as follows.

$$A_1 \leq C\tau_-^{-\eta+m+\frac{1}{2}-(m+\frac{1}{2})p} \int_{\frac{t-r-2k}{3}}^{\frac{t-r}{2}-\frac{2}{3}k} \left(\frac{3\tau - (t-r) + 4k}{2k} \right)^{-(m+\frac{1}{2})p} d\tau \\ \leq C\tau_-^{-\eta+m+\frac{1}{2}-(m+\frac{1}{2})p} \\ = C\tau_-^{-\eta} W(r, t)^{-1}.$$

$$A_2 \leq C\tau_-^{-\eta-(m+\frac{1}{2})p} \int_{\frac{t-r}{2}-\frac{2}{3}k}^{t-r-\frac{2}{3}k} \left(\frac{t-r-\tau+2k}{k} \right)^{m+\frac{1}{2}-(m+\frac{1}{2})p} d\tau \\ \leq C\tau_-^{-\eta-(m+\frac{1}{2})p} \\ = C\tau_-^{-\eta} W(r, t)^{-1}.$$

As to B in (6.23), we have

$$\begin{aligned}
B &= \int_{t-r-\frac{2}{3}k}^{t-r} \left(\frac{t-r-\tau}{k} \right)^{m+\frac{1}{2}-(m-1)p-\eta} \left(\frac{t-r-\tau+2k}{k} \right)^{-p} \\
&\quad \times \left(\frac{\tau+\frac{\lambda_-}{2}+2k}{k} \right)^{-\frac{p}{2}} \left(\frac{\tau-\frac{\lambda_-}{2}+2k}{k} \right)^{-(m+\frac{1}{2})p} d\tau \\
&\leq C\tau_-^{-\frac{p}{2}-(m+\frac{1}{2})p} \int_{t-r-\frac{2}{3}k}^{t-r} \left(\frac{t-r-\tau}{k} \right)^{m+\frac{1}{2}-(m-1)p-\eta} d\tau.
\end{aligned}$$

Since

$$m + \frac{3}{2} - (m-1)p - \eta > \frac{1}{2} + \frac{6}{2m+1} - \eta > 0 \quad \text{for } p < \frac{2m+5}{2m+1}, \quad \eta = 0 \text{ or } \frac{1}{2}, \quad (6.24)$$

we can integrate it as follows.

$$B \leq C\tau_-^{-\frac{p}{2}-(m+\frac{1}{2})p} = C\tau_-^{-\eta}W(r, t)^{-1}.$$

Therefore, we get

$$|I_3(\phi_0^{-p}, p)| \leq C\tau_-^{-\eta}W(r, t)^{-1}.$$

By the similar calculation, we have

$$\begin{aligned}
|I_3(\phi_0^{1-p}Z^{-1}, p)| &\leq C\tau_-^{-\eta+m+\frac{1}{2}-(m+\frac{1}{2})p} + C\tau_-^{-\eta-(m+\frac{1}{2})(p-1)-\frac{1}{p}} + C\tau_-^{-\frac{p}{2}-(m+\frac{1}{2})(p-1)-\frac{1}{p}}, \\
|I_3(Z^{-p}, p)| &\leq C\tau_-^{-\eta+m+\frac{1}{2}-(m+\frac{1}{2})p} \log 3 \frac{t-r+4k}{4k} + C\tau_-^{-\eta-1} + C\tau_-^{-\frac{p}{2}-1}. \quad (6.25)
\end{aligned}$$

for the case of $F(p, q; n) = 0$ with $p \neq q$,

$$\begin{aligned}
|I_3(\phi_0^{-p}, p)| &\leq C\tau_-^{-\eta+m+\frac{1}{2}-(m+\frac{1}{2})p} + C\tau_-^{-\eta-(m+\frac{1}{2})p} + C\tau_-^{-\frac{p}{2}-(m+\frac{1}{2})p}, \\
|I_3(\phi_0^{1-p}Z^{-1}, p)| &\leq C\tau_-^{-\eta+m+\frac{1}{2}-(m+\frac{1}{2})p} + C\tau_-^{-\eta-(m+\frac{1}{2})(p-1)-\mu} + C\tau_-^{-\frac{p}{2}-(m+\frac{1}{2})(p-1)-\mu}, \\
|I_3(Z^{-p}, p)| &\leq C\tau_-^{-\eta+m+\frac{1}{2}-(m+\frac{1}{2})p} \times \begin{cases} \log \tau_- & \text{if } F = 0, p = q \\ \tau_-^{1-p\mu} & \text{otherwise} \end{cases} \\
&\quad + C\tau_-^{-\eta-p\mu} + C\tau_-^{-\frac{p}{2}-p\mu}.
\end{aligned}$$

for the case of $F(p, q; n) = 0$ with $p = q$ or $F(p, q; n) > 0$. Since it holds that

$$\log X \leq \frac{1}{\delta} X^\delta \quad (X \geq 1),$$

for any $\delta > 0$, we shall apply it to (6.25) with sufficiently small δ . Thus, we have proved all the inequalities in the left column in Proposition 4.2 for all cases.

As for the right column in Proposition 4.2, calculating them by the same way as above, we have

$$\begin{aligned}
|I_3(\phi_0^{-q}, q)| &\leq C\tau_-^{-\eta+m+\frac{1}{2}-(m+\frac{1}{2})q} + C\tau_-^{-\eta-(m+\frac{1}{2})q} + C\tau_-^{-\frac{q}{2}-(m+\frac{1}{2})q}, \\
|I_3(\phi_0^{1-q}W^{-1}, q)| &\leq C\tau_-^{-\eta+m+\frac{1}{2}-(m+\frac{1}{2})q} + C\tau_-^{-\eta-(m+\frac{1}{2})(q-1)-\{(m+\frac{1}{2})p-(m+\frac{3}{2})\}} \\
&\quad + C\tau_-^{-\frac{p}{2}-(m+\frac{1}{2})(q-1)-\{(m+\frac{1}{2})p-(m+\frac{3}{2})\}}, \\
|I_3(W^{-q}, q)| &\leq C\tau_-^{-\eta+m+\frac{1}{2}-(m+\frac{1}{2})q} \times \begin{cases} \log \tau_- & \text{if } F = 0, p = q \\ \tau_-^{1-\{(m+\frac{1}{2})p-(m+\frac{3}{2})\}q} & \text{otherwise} \end{cases} \\
&\quad + C\tau_-^{-\eta-\{(m+\frac{1}{2})p-(m+\frac{3}{2})\}q} + C\tau_-^{-\frac{p}{2}-\{(m+\frac{1}{2})p-(m+\frac{3}{2})\}q}.
\end{aligned}$$

Here we used (6.12) for the third. By $m + \frac{5}{2} - (m + \frac{1}{2})p > 0$ and (6.5), the right column in Proposition 4.2 has been proved for all the cases.

6.4 Estimates for I_4

If $\Phi = \Phi(\tau_-)$, we have $|I_4(\Phi, p)| \leq C\Phi(\tau_-)\tilde{I}_4$ by (4.34), where

$$\tilde{I}_4 := \int_{t-r}^t \left(\frac{\tau + r - t}{k} \right)^{m+\frac{1}{2}-(m-1)p-\eta} \left(\frac{\tau + r - t + 2k}{k} \right)^{-p} \left(\frac{2\tau + r - t + 2k}{k} \right)^{-\frac{p}{2}} d\tau$$

Since W, Z and ϕ_0 consists of τ_- only, we shall consider \tilde{I}_4 .

When $t - r + 2k \geq t$, extending the upper limit, we have by (6.24)

$$\tilde{I}_4 \leq C\tau_-^{-\frac{p}{2}} \int_{t-r}^{t-r+2k} \left(\frac{\tau + r - t}{k} \right)^{m+\frac{1}{2}-(m-1)p-\eta} d\tau \leq C\tau_-^{-\frac{p}{2}}.$$

When $t - r + 2k \leq t$, we divide the integral as

$$\int_{t-r}^t d\tau = \int_{t-r}^{t-r+2k} d\tau + \int_{t-r+2k}^t d\tau.$$

The first term is same as the one for $t - r + 2k \leq t$. By

$$\left(\frac{2\tau + r - t + 2k}{k} \right)^{-\frac{p}{2}} \leq \tau_-^{-\eta} \left(\frac{\tau + r - t}{k} \right)^{-\frac{p}{2}+\eta}$$

the second term is calculated as below.

$$(\text{the second term}) \leq \tau_-^{-\eta} \int_{t-r+2k}^t \left(\frac{\tau + r - t}{k} \right)^{m+\frac{1}{2}-(m+\frac{1}{2})p} d\tau \leq C\tau_-^{-\eta}.$$

Therefore, we get

$$\begin{aligned}
|I_4(\phi_0^{-p}, p)| &\leq C\tau_-^{-(m+\frac{1}{2})p-\frac{p}{2}} + C\tau_-^{-(m+\frac{1}{2})p-\eta} \\
|I_4(\phi_0^{1-p}Z^{-1}, p)| &\leq C\tau_-^{-(m+\frac{1}{2})(p-1)-\frac{1}{p}-\frac{p}{2}}(\log 3\tau_-)^{-\nu} + C\tau_-^{-(m+\frac{1}{2})(p-1)-\frac{1}{p}-\eta}(\log 3\tau_-)^{-\nu} \\
|I_4(\phi_0^{-1}Z^{1-p}, p)| &\leq C\tau_-^{-1-\frac{p}{2}}(\log 3\tau_-)^{-p\nu} + C\tau_-^{-1-\eta}(\log 3\tau_-)^{-p\nu},
\end{aligned}$$

if $F(p, q; n)$ with $p \neq q$,

$$\begin{aligned} |I_4(\phi_0^{-p}, p)| &\leq C\tau_-^{-(m+\frac{1}{2})p-\frac{p}{2}} + C\tau_-^{-(m+\frac{1}{2})p-\eta} \\ |I_4(\phi_0^{1-p}Z^{-1}, p)| &\leq C\tau_-^{-(m+\frac{1}{2})(p-1)-\mu-\frac{p}{2}} + C\tau_-^{-(m+\frac{1}{2})(p-1)-\mu-\eta} \\ |I_4(\phi_0^{-1}Z^{1-p}, p)| &\leq C\tau_-^{-p\mu-\frac{p}{2}} + C\tau_-^{-p\mu-\eta}, \end{aligned}$$

if $F(p, q; n) = 0$ with $p = 1$ or $F(p, q; n) > 0$. Since $m + \frac{5}{2} - (m + \frac{1}{2})p > 0$, all of them can be dominated by $C\tau_-^{-\eta}W(r, t)^{-1}$. Similary, we have

$$\begin{aligned} |I_4(\phi_0^{-q}, q)| &\leq C\tau_-^{-(m+\frac{1}{2})q-\frac{q}{2}} + C\tau_-^{-(m+\frac{1}{2})q-\eta} \\ |I_4(\phi_0^{1-q}W^{-1}, q)| &\leq C\tau_-^{-(m+\frac{1}{2})(q-1)-\{(m+\frac{1}{2})p-(m+\frac{3}{2})\}-\frac{q}{2}} + C\tau_-^{-(m+\frac{1}{2})(q-1)-\{(m+\frac{1}{2})p-(m+\frac{3}{2})\}-\eta} \\ |I_4(\phi_0^{-1}W^{1-q}, q)| &\leq C\tau_-^{-\{(m+\frac{1}{2})p-(m+\frac{3}{2})\}q-\frac{q}{2}} + C\tau_-^{-\{(m+\frac{1}{2})p-(m+\frac{3}{2})\}q-\eta}. \end{aligned}$$

By $m + \frac{5}{2} - (m + \frac{1}{2})q > 0$ and (6.4), the first two are dominated by $C\tau_-^{-\eta}W(r, t)^{-1}$. Moreover, using (6.5), we have

$$\begin{aligned} |I_4(\phi_0^{-1}W^{1-q}, q)| &\leq C\tau_-^{-\eta-\mu+q(p-1)F} \\ &\leq C\tau_-^{-\eta} \times \begin{cases} \tau_-^{\frac{1}{p}}(\log 3\tau_-)^{-\nu}(\log 3\tau_-)^{\nu} & \text{if } F = 0, p \neq q, \\ \tau_-^{-\mu} & \text{if } F = 0, p = q, \\ \tau_-^{-\mu+q(p-1)F} & \text{if } F > 0 \end{cases} \\ &\leq C\tau_-^{-\eta}Z(r, t)^{-1}E_2(T) \end{aligned}$$

This completes the proof of Proposition 4.2.

Acknowledgements

The author thanks Professor Hiroyuki Takamura (Tohoku Univ., Japan) for proposing this problem, stimulating discussions on the blow-up and local-existence of L^p solutions and his valuable guidance.

References

- [1] R. Agemi, Y. Kurokawa and H. Takamura, *Critical curve for p-q systems of nonlinear wave equations in three space dimensions*, J. Differential Equations **167** (2000), 87-133.
- [2] R. Courant and D. Hilbert, *Methods of Mathematical physics*, II, Interscience, New York, (1962).
- [3] D. Del Santo, *Global existence and blow-up for a hyperbolic system in three space dimensions*, Rend. Istit. Mat. Univ. Trieste 29 (1997), no. 1-2, 115-140.
- [4] D. Del Santo, V. Georgiev and E. Mitidieri, *Global existence of the solutions and formation of singularities for a class of hyperbolic systems*, Geometrical optics and related topics (Cortona, 1996), 117-140, Progr. Nonlinear Differential Equations Appl., 32, Birkhäuser Boston, Boston, MA, 1997.

- [5] D. Del Santo and E. Mitidieri, *Blow-up of solutions of a hyperbolic system : the critical case*, Differential Equations, **34**, (1998), 1157-1163.
- [6] K. Deng, *Nonexistence of global solutions of a nonlinear hyperbolic system*, Trans. Amer. Math. Soc. 349 (1997), no. 4, 1685–1696.
- [7] K. Deng, *Blow-up of solutions of some nonlinear hyperbolic systems*, Rocky Mountain J. Math. 29 (1999), no. 3, 807–820.
- [8] V. Georgiev, H. Lindblad and C.D. Sogge, *Weighted Strichartz estimates and global existence for semilinear wave equations*, Amer. J. Math. 119 (1997), no. 6, 1291–1319.
- [9] V. Georgiev, H. Takamura and Y. Zhou, *The lifespan of solutions to nonlinear systems of a high-dimensional wave equation*, Nonlinear Anal. 64 (2006), no. 10, 2215–2250.
- [10] R. T. Glassey, *Finite-time blow-up for solutions of nonlinear wave equations*, Math. Z. 177 (1981), no. 3, 323–340.
- [11] R. T. Glassey, *Existence in the large for $\square u = F(u)$ in two space dimensions*, Math. Z. 178 (1981), no. 2, 233–261.
- [12] M. Ikeda, M. Sobajima and K. Wakasa, *Blow-up phenomena of semilinear wave equations and their weakly coupled systems*, J. Differential Equations 267 (2019), no. 9, 5165–5201.
- [13] T. Imai, M. Kato, H. Takamura and K. Wakasa, *The sharp lower bound of the lifespan of solutions to semilinear wave equations with low powers in two space dimensions*, . Asymptotic analysis for nonlinear dispersive and wave equations, 31–53, Adv. Stud. Pure Math., 81, Math. Soc. Japan, Tokyo.
- [14] F. John, *Blow-up of solutions of nonlinear wave equations in three space dimensions*, Proc. Nat. Acad. Sci. U.S.A. 76 (1979), no. 4, 1559–1560.
- [15] H. Kubo and K. Kubota, *Asymptotic behaviors of radially symmetric solutions of $\square u = |u|^p$ for super critical values p in odd space dimensions*, Hokkaido Math. J. 24 (1995), no. 2, 287–336.
- [16] H. Kubo and K. Kubota, *Asymptotic behavior of radially symmetric solutions of $\square u = |u|^p$ for super critical values p in even space dimensions*, Japan J.Math. **24** no.2 (1998), 191-256.
- [17] H. Kubo and M. Ohta, *Critical blow-up for systems of semilinear wave equations in low space dimensions*, J. Math. Anal. Appl., **240**, (1999), 340-360.
- [18] Y. Kurokawa, “The lifespan of radially symmetric solutions to nonlinear systems of odd dimensional wave equations.” Nonlinear Anal. 60 (2005), no. 7, 1239–1275.
- [19] Y. Kurokawa and H. Takamura, *A weighted pointwise estimate for two dimensional wave equations and its application to nonlinear systems*, Tsukuba J. Math. **27** no.2, (2003), 417-448.

- [20] Y. Kurokawa, H. Takamura and K. Wakasa, *The blow-up and lifespan of solutions to systems of semilinear wave equation with critical exponents in high dimensions*, Differential Integral Equations 25 (2012), no. 3-4, 363–382.
- [21] N.-A. Lai and Y. Zhou, *An elementary proof of Strauss conjecture*, J. Funct. Anal. 267 (2014), no. 5, 1364–1381.
- [22] T. T. Li and Y. Zhou, *A note on the life-span of classical solutions to nonlinear wave equations in four space dimensions*, Indiana Univ. Math. J. 44 (1995), no. 4, 1207–1248.
- [23] H. Lindblad, *Blow-up for solutions of $\square u = |u|^p$ with small initial data*, Comm. Partial Differential Equations 15 (1990), no. 6, 757–821.
- [24] H. Lindblad and C. D. Sogge, *Long-time existence for small amplitude semilinear wave equations*, Amer. J. Math., **118**, (1996), no.5, 1047-1135.
- [25] A. Palmieri and H. Takamura, *Blow-up for a weakly coupled system of semilinear damped wave equations in the scattering case with power nonlinearities*, Nonlinear Anal. 187 (2019), 467–492.
- [26] M. A. Rammaha, *Nonlinear wave equations in high dimensions*, Proceedings of the International Conference on Theory and Applications of Differential Equations, Ohio University Press, **II** (1988), 322-326.
- [27] J. Schaeffer, *The equation $u_{tt} - \Delta u = |u|^p$ for the critical value of p* , Proc. Roy. Soc. Edinburgh Sect. A 101 (1985), no. 1-2, 31–44.
- [28] T. C. Sideris, *Nonexistence of global solutions to semilinear wave equations in high dimensions*, J. Partial Differential Equations, **52**, (1984), 378-406.
- [29] W. A. Strauss, *Nonlinear scattering theory at low energy*, J. Funct. Anal. 41, 110-133 (1981).
- [30] H. Takamura, *Improved Kato’s lemma on ordinary differential inequality and its application to semilinear wave equations*, Nonlinear Anal. 125 (2015), 227–240.
- [31] H. Takamura and K. Wakasa, *The sharp upper bound of the lifespan of solutions to critical semilinear wave equations in high dimensions*, J. Differential Equations 251 (2011), no. 4-5, 1157–1171.
- [32] B. Yordanov and Q. S. Zhang, *Finite time blow up for critical wave equations in high dimensions*, J. Funct. Anal. 231 (2006), no. 2, 361–374.
- [33] Y. Zhou, *Life span of classical solutions to $u_{tt} - u_{xx} = |u|^{1+\alpha}$* , Chinese Ann. Math. Ser. B 13 (1992), no. 2, 230–243.
- [34] Y. Zhou, *Blow up of classical solutions to $\square u = |u|^{1+\alpha}$ in three space dimensions*, J. Partial Differential Equations 5 (1992), no. 3, 21–32.

- [35] Y. Zhou, *Life span of classical solutions to $\square u = |u|^p$ in two space dimensions*, Chinese Ann. Math. Ser. B 14 (1993), no. 2, 225–236.
- [36] Y. Zhou, *Blow up of solutions to semilinear wave equations with critical exponent in high dimensions*, Chinese Ann. Math. Ser. B 28 (2007), no. 2, 205–212.
- [37] Y. Zhou and W. Han, *Life-span of solutions to critical semilinear wave equations*, Comm. Partial Differential Equations 39 (2014), no. 3, 439–451.