

# Quartic diophantine equation $X^4 - Y^4 = R^2 - S^2$

S.Muthuvel, R.Venkatraman\*

*Department of Mathematics, College of Engineering and Technology, SRM Institute of Science and Technology, Vadapalani Campus, No.1 Jawaharlal Nehru Salai, Vadapalani, Chennai-600026, Tamilnadu, India.*

---

## Abstract

In this paper, we deal with the quartic diophantine equation  $X^4 - Y^4 = R^2 - S^2$  to present its infinitely many integer solutions.

*Keywords:* Quartic diophantine equation, Elementary method

*2020 MSC:* 11D25, 11D45

---

## 1. Introduction

In the general form of Diophantine equation

$$x^n + y^n = u^n + v^n, \quad n \in \mathbb{N}$$

The case  $n = 2$  has been expressed in [6, 12, 13]. For  $n = 4$ , the parametric solutions to the aforementioned equation are described in [4, 8, 9]. More general diophantine equations with more variable or with integer coefficients that are not all equal to one were taken into consideration by several researchers [5, 7, 10, 11]. The authors provided an infinite number of positive integral solutions for various powers as illustrated in [3]. The objective of this work is to obtain infinitely many integral solutions of

$$X^4 - Y^4 = R^2 - S^2 \tag{1}$$

---

\*Corresponding author

*Email addresses:* muthushan15@gmail.com, ms3081@srmist.edu.in (S.Muthuvel), venkatrr1@srmist.edu.in (R.Venkatraman )

for each one of the parametric method. In [2], the equations

$$a \left( X_1'^5 + X_2'^5 \right) + \sum_{i=0}^m a_i X_i^5 = b \left( Y_1'^3 + Y_2'^3 \right) + \sum_{i=0}^n b_i Y_i^3 \quad (2)$$

where  $m, n \in \mathbb{N} \cup \{0\}$  and  $a, b \neq 0$ ,  $a_i, b_i$  are fixed arbitrary rational numbers are examined. The solution to (2), which is converted into a cubic or a quartic elliptic curve with a positive rank, is found using the theory of elliptic curves. Authors demonstrate that in [[1], Main Theorem 2]

$$\sum_{i=1}^n p_i x_i^{a_i} = \sum_{j=1}^m q_j y_j^{b_j}$$

$m, n, a_i, b_j \in \mathbb{N}$ ,  $p_i, q_j \in \mathbb{Z}$ ,  $i = 1, 2, \dots, n$ ,  $j = 1, 2, \dots, m$  has a parametric solution and infinitely many solutions in nonzero integers if there exists an  $i$  such that  $p_i = 1$  and  $(a_i, a_1 a_2 \dots a_{i-1} a_{i+1} \dots a_n b_1 b_2 \dots b_m) = 1$  or there exists a  $j$  such that  $q_j = 1$  and  $(b_j, a_1 \dots a_n b_1 \dots b_{j-1} b_{j+1} \dots b_m) = 1$ . Although linear transformations are also employed in this article, we propose a different strategy and some different conditions for the integer coefficients in order to solve (1).

## 2. Solving the Diophantine equation $X^4 - Y^4 = R^2 - S^2$

The trivial solution of the equation (1) is  $(X, Y, R, S) = (m, n, m^2, n^2)$  for  $m, n \in \mathbb{Z}$ . Four different linear transformations are considered and for each one of them, we give a different class of infinitely many integer solutions of equation (1).

### 2.1. Method-1

Consider the linear transformations,

$$X = px + u, \quad Y = qx - u, \quad R = x + v, \quad S = px - v \quad (3)$$

$p, x, u, v \in \mathbb{Z}$ . Introducing (3) in (1), we get

$$\alpha x^4 + \beta x^3 + \gamma x^2 + \delta x = 0 \quad (4)$$

where

$$\begin{aligned}\alpha &= p^4 - q^4, & \beta &= 4p^3u + 4q^3u, \\ \gamma &= 6p^2u^2 - 6q^2u^2 + p^2 - 1, & \delta &= 4pu^3 + 4qu^3 - 2v + 2pv\end{aligned}\quad (5)$$

For  $\delta = 0$  in (5), we obtain

$$(2p + 2q)u^3 = v(1 - p)$$

Further, we put  $u = t, v = t^3$  and get  $p = \frac{1-2q}{3}$ .

In (5), equating the like terms  $\gamma = 0$

$$(q + 1)[(-15t^2 + 2)q + (3t^2 - 4)] = 0$$

Simplifying the above expression, we have  $q = \frac{4-3t^2}{2-15t^2}$  and therefore (4) becomes

$$\begin{aligned}\alpha x^4 + \beta x^3 &= 0 \\ x &= \frac{-t(405t^8 + 459t^6 - 1404t^4 + 600t^2 - 56)}{3(27t^6 - 27t^4 + 36t^2 - 10)}\end{aligned}$$

Plugging the values  $p, q, u, v$  in (3), we acquire

$$\begin{aligned}X &= \frac{t(1215t^{10} - 1782t^8 + 4671t^6 - 774t^4 - 366t^2 + 52)}{3(2 - 15t^2)(27t^6 - 27t^4 + 36t^2 - 10)} \\ Y &= \frac{t(1215t^{10} - 1782t^8 + 4671t^6 - 5634t^4 + 1902t^2 - 164)}{3(2 - 15t^2)(27t^6 - 27t^4 + 36t^2 - 10)} \\ R &= \frac{-t(324t^8 - 378t^6 + 1296t^4 - 570t^2 + 56)}{3(27t^6 - 27t^4 + 36t^2 - 10)} \\ S &= \frac{t(810t^8 + 1512t^6 + 1674t^4 - 1092t^2 + 112)}{3(2 - 15t^2)(27t^6 - 27t^4 + 36t^2 - 10)}\end{aligned}\quad (6)$$

Eliminating the denominators from the above equations,

$$\begin{aligned}X &= 1215t^{11} - 1782t^9 + 4671t^7 - 774t^5 - 366t^3 + 52t \\ Y &= 1215t^{11} - 1782t^9 + 4671t^7 - 5634t^5 + 1902t^3 - 164t \\ R &= 5904900t^{19} - 14368590t^{17} + 41898546t^{15} - 55842858t^{13} + 58236894t^{11} \\ &\quad - 36547200t^9 + 12314916t^7 - 2186784t^5 + 193392t^3 - 6720t \\ S &= 984150t^{17} + 721710t^{15} + 1395306t^{13} - 1476954t^{11} + 3664440t^9 - 3124332t^7 \\ &\quad + 1027296t^5 - 140112t^3 + 6720t\end{aligned}$$

We get a integer solution  $(X, Y, R, S)$  of equation (1) for every  $t \in \mathbb{Z}$ . So, the presented method generates infinitely many integer solutions of the initial equation (1).

## 2.2. Method-2

In this method, we deal with different transformation in (1). Let

$$X = px + u, \quad Y = qx - u, \quad R = x + v, \quad S = px - v \quad (7)$$

$p, x, u, v \in \mathbb{Z}$ . In previous subsection, by introducing these linear transformations in (1), leads us to the equation of the form

$$Ax^4 + Bx^3 + Cx^2 + Dx = 0 \quad (8)$$

where

$$\begin{aligned} A &= p^4 - q^4, & B &= 4p^3u + 4q^3u, \\ C &= 6p^2u^2 - 6q^2u^2 + p^2 - 1, & D &= 4pu^3 + 4qu^3 - 2v - 2pv \end{aligned} \quad (9)$$

For  $D = 0$  in (9), we get

$$(2p + 2q)u^3 = v(1 + p)$$

Further, we set  $u = t, v = t^3$  and attain  $p = 1 - 2q$ . In (9), equating the like terms  $C = 0$

$$(q - 1) [(9t^2 + 2)q - 3t^2] = 0$$

Using the above equation, we obtain  $q = \frac{3t^2}{9t^2 + 2}$  and therefore (8) becomes

$$\begin{aligned} Ax^4 + Bx^3 &= 0 \\ x &= \frac{-t(243t^8 + 297t^6 + 216t^4 + 72t^2 + 8)}{(27t^6 + 27t^4 + 12t^2 + 2)} \end{aligned}$$

Applying the values  $p, q, u, v$  in (7), we acquire

$$\begin{aligned} X &= \frac{-3t(27t^8 + 36t^6 + 27t^4 + 12t^2 + 2)}{(27t^6 + 27t^4 + 12t^2 + 2)} \\ Y &= \frac{-t(81t^8 + 108t^6 + 81t^4 + 24t^2 + 2)}{(27t^6 + 27t^4 + 12t^2 + 2)} \\ R &= \frac{-2t(108t^8 + 135t^6 + 102t^4 + 35t^2 + 4)}{(27t^6 + 27t^4 + 12t^2 + 2)} \\ S &= \frac{-2t(54t^8 + 81t^6 + 60t^4 + 25t^2 + 4)}{(27t^6 + 27t^4 + 12t^2 + 2)} \end{aligned} \quad (10)$$

By cancelling the denominators in (10),

$$X = 2187t^{15} + 5103t^{13} + 6075t^{11} + 4617t^9 + 2322t^7 + 756t^5 + 144t^3 + 12t$$

$$Y = 2187t^{15} + 5103t^{13} + 6075t^{11} + 4293t^9 + 1890t^7 + 504t^5 + 72t^3 + 4t$$

$$R = 4251528t^{27} + 18068994t^{25} + 38381850t^{23} + 52986636t^{21} + 52435512t^{19} \\ + 38959218t^{17} + 22221378t^{15} + 9803592t^{13} + 3331692t^{11} + 858168t^9 \\ + 162216t^7 + 21216t^5 + 1712t^3 + 64t$$

$$S = 2125764t^{27} + 9565938t^{25} + 21139542t^{23} + 30154356t^{21} + 30784212t^{19} \\ + 23632722t^{17} + 13977846t^{15} + 6426864t^{13} + 2290356t^{11} + 623160t^9 \\ + 125496t^7 + 17664t^5 + 1552t^3 + 64t$$

For any  $t \in \mathbb{Z}$ , we obtain an integer solution  $(X, Y, R, S)$  to equation (1). Consequently, the proposed method yields an infinite number of integer solutions to the starting equation (1).

### 2.3. Method-3

In this method, we deal with different transformation in (1). Let

$$X = v, \quad Y = px + v, \quad R = qx + u, \quad S = x + u \quad (11)$$

$p, x, u, v \in \mathbb{Z}$ . In subsection-1, by introducing these linear transformations in (1), leads us to the equation of the form

$$ax^4 + bx^3 + cx^2 + dx = 0 \quad (12)$$

where

$$a = p^4, \quad b = 4p^3v, \quad (13) \\ c = 6p^2v^2 + q^2 - 1, \quad d = 4pv^3 - 2u - 2qu$$

For  $d = 0$  in (13), we obtain

$$(2p)v^3 = u(1 - q)$$

Additionally, we put  $u = t^3, v = t$  and get  $p = \frac{1-q}{2}$ . In (13), equating the like terms  $c = 0$

$$(q - 1) [(3t^2 + 2)q - (3t^2 - 2)] = 0$$

Thus, we get  $q = \frac{3t^2-2}{3t^2+2}$  and therefore (12) becomes

$$\begin{aligned} ax^4 + bx^3 &= 0 \\ x &= \frac{-2t(81t^8 + 216t^6 + 216t^4 + 96t^2 + 16)}{(27t^6 + 54t^4 + 36t^2 + 8)} \end{aligned}$$

Taking the values  $p, q, u, v$  and applying it in (11), we get

$$\begin{aligned} X &= t \\ Y &= \frac{-3t(81t^8 + 216t^6 + 216t^4 + 96t^2 + 16)}{(3t^2 + 2)(27t^6 + 54t^4 + 36t^2 + 8)} \\ R &= \frac{-t(405t^{10} + 756t^8 + 216t^6 - 384t^4 - 304t^2 - 64)}{(3t^2 + 2)(27t^6 + 54t^4 + 36t^2 + 8)} \\ S &= \frac{-t(135t^8 + 358t^6 + 396t^4 + 184t^2 + 32)}{(27t^6 + 54t^4 + 36t^2 + 8)} \end{aligned} \tag{14}$$

Neglecting the denominators from the above equation,

$$\begin{aligned} X &= 81t^{10} + 216t^8 + 216t^6 + 96t^4 + 16t^2 \\ Y &= 243t^{10} + 648t^8 + 648t^6 + 288t^4 + 48t^2 \\ R &= 32805t^{21} + 148716t^{19} + 268272t^{17} + 217728t^{15} + 18144t^{13} - 120960t^{11} \\ &\quad - 112896t^9 - 49152t^7 - 11008t^5 - 1024t^3 \\ S &= 32805t^{21} + 196344t^{19} + 532008t^{17} + 849312t^{15} + 874656t^{13} + 600000t^{11} \\ &\quad + 273536t^9 + 79872t^7 + 13568t^5 + 1024t^3 \end{aligned}$$

For each  $t \in \mathbb{Z}$ , equation (1) has an integer solution  $(X, Y, R, S)$ . The resulting method produces an infinite number of integer solutions to the initial equation (1).

#### 2.4. Method-4

In this method, we deal with different transformation in (1). Let

$$X = -v, \quad Y = px - v, \quad R = qx - u, \quad S = x + u \tag{15}$$

$p, x, u, v \in \mathbb{Z}$ . In subsection-1, by introducing these linear transformations in (1), leads us to the equation of the form

$$Mx^4 + Nx^3 + Px^2 + Qx = 0 \quad (16)$$

where

$$\begin{aligned} M &= p^4, & N &= -4p^3v, \\ P &= 6p^2v^2 + q^2 - 1, & Q &= -4pv^3 - 2u - 2qu \end{aligned} \quad (17)$$

For  $Q = 0$  in (17), we obtain

$$(2p)v^3 = u(-1 - q)$$

Further, we put  $u = t^3, v = t$  and get  $q = -1 - 2p$ . In (17) equating the like terms  $Q = 0$

$$p = \frac{-2}{3t^2 + 2}$$

Therefore (16) becomes

$$\begin{aligned} Mx^4 + Nx^3 &= 0 \\ x &= -2t(3t^2 + 2) \end{aligned}$$

Substituting the values  $p, q, u, v$  in (15), we obtain

$$\begin{aligned} X &= -t \\ Y &= 3t \\ R &= 5t^3 - 4t \\ S &= -5t^3 - 4t \end{aligned}$$

We get a integer solution  $(X, Y, R, S)$  of equation (1) for every  $t \in \mathbb{Z}$ . So, the presented method generates infinitely many integer solutions of the initial equation (1).

## References

- [1] M. Baghlaghdam, F. Izadi, A note on the high power diophantine equations, Proc. Math. Sci., 129(14), (2019).
- [2] M. Baghlaghdam, F. Izadi, On the diophantine equation in the form that a sum of cubes equals a sum of quantics, Math. J. Okaama Univ., 61, (2019) 75-84.
- [3] S. Bujačić Babić, K. Nabardi, On some Diophantine equations, Miskolc Mathematical Notes, 22(1), (2021), 65-75.
- [4] A. Choudhry, The diophantine equation  $A^4 + B^4 = C^4 + D^4$ , Indian J. Pure Appl. Math., 22(1), (1991) 9-11.
- [5] A. Choudhry, On the Diophantine equation  $A^4 + hB^4 = C^4 + hD^4$ , Indian J. Pure Appl. Math., 26(11), (1995) 1057-1061.
- [6] H.B. Davies, On generating solutions of the Diophantine equation  $x^2 + y^2 = u^2 + v^2$ , Int. J. Math. Educ. Sci. Technol., 15(1), (1984) 43-46.
- [7] N. Elkies, On  $A^4 + B^4 + C^4 = D^4$ , Math. Comput., 51(184), (1988) 825-835.
- [8] L. Euler, Novi Comm. Acad. Petrop. v(17), (1772).
- [9] G. H. Hardy, E. M. Wright, An Introduction to the Theory of Numbers, Oxford University Press, London (1960).
- [10] F. Izadi, K. Nabardi, Diophantine equation  $X^4 + Y^4 = 2(U^4 + V^4)$ , Math. Slovaca., 66(3), (2016) 557-560.
- [11] A. S. Janfada, N. Nabardi, On Diophantine equation  $x^4 + y^4 = n(u^4 + v^4)$ , Math. Slovaca., 69(6), (2019) 1245-1248.
- [12] J. Kersey, The Elements of Algebra, London (1674).
- [13] P. Pasternak, Zeitschr, Math. Naturw. Unterricht., 37, (1906) 33-35.