

Last updated March 24, 2023.

Golden Ratio Base Expansions of the Logarithm and Inverse Tangent of Fibonacci and Lucas Numbers

Kunle Adegoke

Department of Physics and Engineering Physics,
Obafemi Awolowo University, Ile-Ife
Nigeria

adegoke00@gmail.com

Jaume Oliver Lafont

Conselleria d'Educacio i Cultura, Govern de les Illes Balears,
Palma, Spain

joliverlafont@gmail.com

Abstract

Let $\alpha = (1 + \sqrt{5})/2$, the golden ratio, and $\beta = -1/\alpha = (1 - \sqrt{5})/2$. Let F_n and L_n be the Fibonacci and Lucas numbers, defined by $F_n = (\alpha^n - \beta^n)/\sqrt{5}$ and $L_n = \alpha^n + \beta^n$, for all non-negative integers. We derive base α expansions of $\log F_n$, $\log L_n$, $\arctan \frac{1}{F_n}$ and $\arctan \frac{1}{L_n}$ for all positive integers n .

Keywords: Fibonacci number, Lucas number, logarithm, arctangent, inverse tangent, golden ratio, non-integer base expansion, BBP-type formula.

1 Introduction

Let α denote the golden ratio; that is $\alpha = (1 + \sqrt{5})/2$. Let $\beta = -1/\alpha = (1 - \sqrt{5})/2$. Thus $\alpha\beta = -1$ and $\alpha + \beta = 1$. Let F_n and L_n be the Fibonacci and Lucas numbers, defined by $F_n = (\alpha^n - \beta^n)/\sqrt{5}$ and $L_n = \alpha^n + \beta^n$, for all non-negative integers n .

Let b be any non-zero number whose magnitude is greater than unity. Let n and s be positive integers. A convergent series of the form

$$C = \sum_{k=0}^{\infty} \frac{1}{b^k} \left(\frac{a_1}{(kn+1)^s} + \frac{a_2}{(kn+2)^s} + \cdots + \frac{a_n}{(kn+n)^s} \right), \quad (1.1)$$

where a_1, a_2, \dots, a_n are certain numbers, defines a base b , length n and degree s expansion of the mathematical constant C .

If b is an integer and a_k are rational numbers, then (1.1) is referred to as a BBP-type formula, after the initials of the authors of the paper [4] in which such an expansion was first presented for π and some other mathematical constants. Any mathematical constant that possesses a base b BBP-type formula has the property that its n -th digit in base b could be calculated directly, without needing to compute any of the previous $n-1$ digits. Although the infinite series presented in this paper have the structure of BBP-type formulas, it must be clearly stated that the series do not yield digit extraction since here the base $b = \alpha^n$ is not an integer and the a_k are not rational numbers; rather the series correspond to base α expansions of the mathematical constants concerned.

Our goal in this paper is to derive base α expansion formulas for the logarithm and the inverse tangent of all Fibonacci and Lucas numbers. We will often give the expansion using the compact P -notation for BBP-type formulas, introduced by Bailey and Crandall [5], namely,

$$C = P(s, b, n, A) = \sum_{k=0}^{\infty} \frac{1}{b^k} \sum_{j=1}^n \frac{a_j}{(kn+j)^s},$$

where s and n are integers and, in this present paper, b is an integer power of α and $A = (a_1, a_2, \dots, a_n)$ is a vector of rational multiples of powers of β . For example, we will show (see (2.20)) that

$$\log F_3 = \log 2 = \sum_{k=0}^{\infty} \frac{1}{\alpha^{12k}} \left(\frac{\beta^2}{6k+1} + \frac{3\beta^4}{6k+2} + \frac{4\beta^6}{6k+3} + \frac{3\beta^8}{6k+4} + \frac{\beta^{10}}{6k+5} \right),$$

which, in the P -notation, can be written as

$$\log F_3 = \log 2 = P(1, \alpha^{12}, 6, (\beta^2, 3\beta^4, 4\beta^6, 3\beta^8, \beta^{10}, 0)).$$

Base α expansions have also been studied or reported by Bailey and Crandall [5], Chan [7, 8], Zhang [12], Borwein and Chamberland [6], Cloitre [9], Adegoke [2], Wei [11], and more recently Kristensen and Mathiasen [10].

2 Base α expansions of logarithms

The base α expansions of the logarithms of Fibonacci and Lucas numbers are presented in Theorems 2.3 and 2.4 but first we state a couple of Lemmata upon which the results are based.

Let

$$\text{Li}_1(x) = -\log(1-x) = \sum_{k=1}^{\infty} \frac{x^k}{k} = \sum_{k=0}^{\infty} x^k \frac{x}{k+1}, \quad -1 \leq x < 1.$$

Lemma 2.1. *If $|b| > 1$, $t > 0$ and m and n are arbitrary positive integers, then,*

$$\text{Li}_1\left(\frac{1}{b^t}\right) = \sum_{k=0}^{\infty} \frac{1}{b^{tmk}} \sum_{j=1}^m \frac{1/b^{tj}}{(mk+j)}, \quad (2.1)$$

$$\text{Li}_1\left(-\frac{1}{b^t}\right) = \sum_{k=0}^{\infty} \frac{1}{b^{2tnk}} \sum_{j=1}^{2n} \frac{(-1)^j/b^{tj}}{(2nk+j)}. \quad (2.2)$$

Proof. We have

$$\text{Li}_1\left(\frac{1}{b^t}\right) = \sum_{k=0}^{\infty} \frac{1}{b^{tk}} \frac{1/b^t}{k+1},$$

from which (2.1) follows upon using the identity

$$\sum_{k=0}^{\infty} f_k = \sum_{k=0}^{\infty} \sum_{j=1}^m f_{mk+j-1}, \quad (2.3)$$

with

$$f_k = \frac{1}{b^{tk+tt}} \frac{1}{k+1}.$$

The proof of (2.2) is similar, with $m = 2n$ in (2.3). \square

Lemma 2.2. *If r is an integer, then,*

$$\log L_r = r \text{Li}_1\left(\frac{1}{\alpha^2}\right) - \text{Li}_1\left(\frac{(-1)^{r+1}}{\alpha^{2r}}\right), \quad (2.4)$$

$$\log F_r = (r-2) \text{Li}_1\left(\frac{1}{\alpha^2}\right) + \text{Li}_1\left(\frac{1}{\alpha^4}\right) - \text{Li}_1\left(\frac{(-1)^r}{\alpha^{2r}}\right), \quad r \neq 0. \quad (2.5)$$

Proof. We have

$$\text{Li}_1\left(-\frac{\beta^r}{\alpha^r}\right) = -\log\left(\frac{\alpha^r + \beta^r}{\alpha^r}\right) = -\log\left(\frac{L_r}{\alpha^r}\right) = -\log L_r + r \log \alpha, \quad (2.6)$$

in which setting $r = 1$ gives

$$\log \alpha = \text{Li}_1\left(\frac{1}{\alpha^2}\right). \quad (2.7)$$

Using (2.7) in (2.6) gives (2.4).

Also,

$$\text{Li}_1\left(\frac{\beta^r}{\alpha^r}\right) = -\log\left(\frac{\alpha^r - \beta^r}{\alpha^r}\right) = -\log\left(\frac{F_r\sqrt{5}}{\alpha^r}\right) = -\log F_r + r \log \alpha - \log \sqrt{5}, \quad (2.8)$$

in which setting $r = 2$ gives

$$\log \sqrt{5} = 2 \log \alpha - \text{Li}_1\left(\frac{1}{\alpha^4}\right) = 2 \text{Li}_1\left(\frac{1}{\alpha^2}\right) - \text{Li}_1\left(\frac{1}{\alpha^4}\right), \quad (2.9)$$

where we used (2.7). Identity (2.5) follows from (2.8) and (2.9). \square

Theorem 2.3. *If r is an integer, then,*

$$\log F_r = \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^r \frac{\beta^{4j-2}(r-2+r\delta_{j,(r+1)/2})}{2rk+2j-1} + \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^{r-1} \frac{\beta^{4j}r}{2rk+2j}, \quad r \text{ odd}, \quad (2.10)$$

$$\log F_r = \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^r \frac{(r-2)\beta^{4j-2}}{2rk+2j-1} + \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^{r-1} \frac{\beta^{4j}r(1-\delta_{j,r/2})}{2rk+2j}, \quad r \text{ even}. \quad (2.11)$$

Here and throughout this paper, δ_{mn} denotes the Kronecker delta symbol whose value is unity when m equals n and zero otherwise.

Proof. We prove (2.11). When r is even, (2.5) reads

$$\log F_r = (r-2) \text{Li}_1\left(\frac{1}{\alpha^2}\right) + \text{Li}_1\left(\frac{1}{\alpha^4}\right) - \text{Li}_1\left(\frac{1}{\alpha^{2r}}\right), \quad r \neq 0. \quad (2.12)$$

We proceed to write the three Li_1 terms in a common base α^{4r} , using (2.1) with appropriate t and m choices. Thus,

$$\text{Li}_1\left(\frac{1}{\alpha^2}\right) = \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^{2r} \frac{1/\alpha^{2j}}{2rk+j}, \quad (2.13)$$

$$\text{Li}_1\left(\frac{1}{\alpha^4}\right) = \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^r \frac{1/\alpha^{4j}}{rk+j}, \quad (2.14)$$

$$\text{Li}_1\left(\frac{1}{\alpha^{2r}}\right) = \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \left(\frac{1/\alpha^{2r}}{2k+1} + \frac{1/\alpha^{4r}}{2k+2}\right). \quad (2.15)$$

Using (2.13), (2.14) and (2.15) in (2.12) gives

$$\begin{aligned} \log F_r &= \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^{2r} \frac{\beta^{2j}(r-2)}{2rk+j} + \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^r \frac{2\beta^{4j}}{2rk+2j} \\ &\quad - \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \left(\frac{r\beta^{2r}}{2rk+r} + \frac{r\beta^{4r}}{2rk+2r}\right). \end{aligned} \quad (2.16)$$

Using the summation identity

$$\sum_{j=1}^{2r} f_j = \sum_{j=1}^r f_{2j} + \sum_{j=1}^r f_{2j-1} \quad (2.17)$$

to write its inner sum, the first term on the right hand side of (2.16) can be written as

$$\sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^{2r} \frac{\beta^{2j}(r-2)}{2rk+j} = \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^r \frac{\beta^{4j}(r-2)}{2rk+2j} + \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^r \frac{\beta^{4j-2}(r-2)}{2rk+2j-1}. \quad (2.18)$$

Using (2.18) in (2.16) yields (2.11). □

Identities (2.10) and (2.11) written in the P -notation are

$$\log F_r = P(1, \alpha^{4r}, 2r, (a_1, a_2, \dots, a_{2r})) \quad (2.19)$$

where for $1 \leq j \leq r$,

$$a_{2j-1} = \beta^{4j-2}(r-2 + r\delta_{j,(r+1)/2}), \quad a_{2j} = \beta^{4j}r(1 - \delta_{rj}), \quad r \text{ odd};$$

and

$$a_{2j-1} = (r-2)\beta^{4j-2}, \quad a_{2j} = \beta^{4j}r(1 - \delta_{j,r/2} - \delta_{j,r}), \quad r \text{ even}.$$

Examples.

$$\log F_3 = \log 2 = P(1, \alpha^{12}, 6, (\beta^2, 3\beta^4, 4\beta^6, 3\beta^8, \beta^{10}, 0)), \quad (2.20)$$

$$\log F_5 = \log 5 = P(1, \alpha^{20}, 10, (3\beta^2, 5\beta^4, 3\beta^6, 5\beta^8, 8\beta^{10}, 5\beta^{12}, 3\beta^{14}, 5\beta^{16}, 3\beta^{18}, 0)), \quad (2.21)$$

$$\log F_4 = \log 3 = P(1, \alpha^{16}, 8, (2\beta^2, 4\beta^4, 2\beta^6, 0, 2\beta^{10}, 4\beta^{12}, 2\beta^{14}, 0)), \quad (2.22)$$

$$\log F_8 = \log 21 = P(1, \alpha^{32}, 16, (6\beta^2, 8\beta^4, 6\beta^6, 8\beta^8, 6\beta^{10}, 8\beta^{12}, 6\beta^{14}, 0, 6\beta^{18}, 8\beta^{20}, 6\beta^{22}, 8\beta^{24}, 6\beta^{26}, 8\beta^{28}, 6\beta^{30}, 0)), \quad (2.23)$$

$$\log F_{12} = \log 144 = P(1, \alpha^{48}, 24, (10\beta^2, 12\beta^4, 10\beta^6, 12\beta^8, 10\beta^{10}, 12\beta^{12}, 10\beta^{14}, 12\beta^{16}, 10\beta^{18}, 12\beta^{20}, 10\beta^{22}, 0, 10\beta^{26}, 12\beta^{28}, 10\beta^{30}, 12\beta^{32}, 10\beta^{34}, 12\beta^{36}, 10\beta^{38}, 12\beta^{40}, 10\beta^{42}, 12\beta^{44}, 10\beta^{46}, 0)). \quad (2.24)$$

Theorem 2.4. *If r is an integer, then,*

$$\log L_r = \sum_{k=0}^{\infty} \frac{1}{\alpha^{2rk}} \sum_{j=1}^{r-1} \frac{\beta^{2j} r}{rk+j}, \quad r \text{ odd}, \quad (2.25)$$

$$\log L_r = \sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \sum_{j=1}^{2r-1} \frac{\beta^{2j} r (1 + \delta_{rj})}{2rk+j}, \quad r \text{ even}. \quad (2.26)$$

Proof. We prove (2.25). If r is an odd integer, (2.4) gives

$$\log L_r = r \operatorname{Li}_1 \left(\frac{1}{\alpha^2} \right) - \operatorname{Li}_1 \left(\frac{1}{\alpha^{2r}} \right). \quad (2.27)$$

With

$$\operatorname{Li}_1 \left(\frac{1}{\alpha^2} \right) = \sum_{k=0}^{\infty} \frac{1}{\alpha^{2rk}} \sum_{j=1}^r \frac{1/\alpha^{2j}}{rk+j}$$

and

$$\operatorname{Li}_1 \left(\frac{1}{\alpha^{2r}} \right) = \sum_{k=0}^{\infty} \frac{1}{\alpha^{2rk}} \frac{r/\alpha^{2r}}{rk+r}$$

in (2.27); identity (2.25) follows. \square

Identities (2.25) and (2.26) in the P -notation are

$$\log L_r = P(1, \alpha^{2r}, r, (a_1, a_2, \dots, a_r)), \quad r \text{ odd}, \quad (2.28)$$

with

$$a_j = r\beta^{2j}(1 - \delta_{rj}), \quad 1 \leq j \leq r,$$

and

$$\log L_r = P(1, \alpha^{4r}, 2r, (a_1, a_2, \dots, a_{2r})), \quad r \text{ even}, \quad (2.29)$$

with

$$a_j = r\beta^{2j}(1 + \delta_{jr} - \delta_{j,2r}), \quad 1 \leq j \leq 2r.$$

Examples.

$$\log L_2 = \log 3 = \sum_{k=0}^{\infty} \frac{1}{\alpha^{8k}} \left(\frac{2\beta^2}{4k+1} + \frac{4\beta^4}{4k+2} + \frac{2\beta^6}{4k+3} \right); \quad (2.30)$$

that is,

$$\log 3 = P(1, \alpha^8, 4, (2\beta^2, 4\beta^4, 2\beta^6, 0)). \quad (2.31)$$

$$\log L_3 = \log 4 = P(1, \alpha^6, 3, (3\beta^2, 3\beta^4, 0)), \quad (2.32)$$

$$\log L_4 = \log 7 = 4\beta^2 P(1, \alpha^{16}, 8, (1, \beta^2, \beta^4, 2\beta^6, \beta^8, \beta^{10}, \beta^{12}, 0)), \quad (2.33)$$

$$\log L_6 = \log 18 = P(1, \alpha^{24}, 12, (6\beta^2, 6\beta^4, 6\beta^6, 6\beta^8, 6\beta^{10}, 12\beta^{12}, 6\beta^{14}, 6\beta^{16}, 6\beta^{18}, 6\beta^{20}, 6\beta^{22}, 0)). \quad (2.34)$$

3 Base α expansions of inverse tangents

The base α expansions of the inverse tangent of Fibonacci and Lucas numbers are stated in Theorems 3.4–3.8 but first we collect some required identities in Lemmata 3.1–3.3.

Lemma 3.1. *If r is an integer, then,*

$$\alpha^r - \alpha^{-r} = \begin{cases} F_r \sqrt{5}, & r \text{ even}; \\ L_r, & r \text{ odd}; \end{cases} \quad (3.1)$$

$$\alpha^r + \alpha^{-r} = \begin{cases} L_r & r \text{ even}; \\ F_r \sqrt{5}, & r \text{ odd}. \end{cases} \quad (3.2)$$

Lemma 3.2. *If r and m are integers, then,*

$$\arctan \frac{1}{\alpha^{r-m}} - \arctan \frac{1}{\alpha^{r+m}} = \begin{cases} \arctan(L_m/(F_r \sqrt{5})), & m \text{ odd}, r \text{ odd}, \\ \arctan(L_m/L_r), & m \text{ odd}, r \text{ even}, \\ \arctan(F_m/F_r), & m \text{ even}, r \text{ odd}, \\ \arctan(F_m \sqrt{5}/L_r), & m \text{ even}, r \text{ even}; \end{cases} \quad (3.3)$$

$$\arctan \frac{1}{\alpha^{r-m}} + \arctan \frac{1}{\alpha^{r+m}} = \begin{cases} \arctan(F_m \sqrt{5}/L_r), & m \text{ odd}, r \text{ odd}, \\ \arctan(F_m/F_r), & m \text{ odd}, r \text{ even}, \\ \arctan(L_m/L_r), & m \text{ even}, r \text{ odd}, \\ \arctan(L_m/(F_r \sqrt{5})), & m \text{ even}, r \text{ even}. \end{cases} \quad (3.4)$$

Proof. The arctangent subtraction and addition formulas give

$$\arctan \frac{1}{\alpha^{r-m}} - \arctan \frac{1}{\alpha^{r+m}} = \arctan \left(\frac{\alpha^r (\alpha^m - \alpha^{-m})}{\alpha^{2r} + 1} \right),$$

$$\arctan \frac{1}{\alpha^{r-m}} + \arctan \frac{1}{\alpha^{r+m}} = \arctan \left(\frac{\alpha^r (\alpha^m + \alpha^{-m})}{\alpha^{2r} - 1} \right);$$

and hence the stated identities upon the use of Lemma 3.1. \square

Lemma 3.3. *If r is an integer, then,*

$$\alpha^{2r} - 1 = \alpha^r L_r, \quad \beta^{2r} - 1 = \beta^r L_r, \quad r \text{ odd}, \quad (3.5)$$

$$\alpha^{2r} - 1 = \alpha^r F_r \sqrt{5}, \quad \beta^{2r} - 1 = -\beta^r F_r \sqrt{5}, \quad r \text{ even}, \quad (3.6)$$

$$\alpha^{2r} + 1 = \alpha^r F_r \sqrt{5}, \quad \beta^{2r} + 1 = -\beta^r F_r \sqrt{5}, \quad r \text{ odd}, \quad (3.7)$$

$$\alpha^{2r} + 1 = \alpha^r L_r, \quad \beta^{2r} + 1 = \beta^r L_r, \quad r \text{ even}. \quad (3.8)$$

Theorem 3.4. *If r is an odd integer greater than unity, then,*

$$\arctan \frac{1}{F_r} = P(1, \alpha^{4(r^2-4)}, 4(r^2-4), (a_1, a_2, \dots, a_{4(r^2-4)})), \quad (3.9)$$

where the only non-zero constants a_j are given by

$$\begin{aligned} a_{(r-2)(4j-3)} &= -\beta^{(r-2)(4j-3)}(r-2), \quad j = 1, 2, \dots, r+2, \\ a_{(r-2)(4j-1)} &= \beta^{(r-2)(4j-1)}(r-2), \quad j = 1, 2, \dots, r+2, \\ a_{(r+2)(4j-3)} &= \beta^{(r+2)(4j-3)}(r+2), \quad j = 1, 2, \dots, r-2, \\ a_{(r+2)(4j-1)} &= -\beta^{(r+2)(4j-1)}(r+2), \quad j = 1, 2, \dots, r-2, \\ a_{(r-2)(r+2)} &= (-1)^{(r+1)/2} 4\beta^{r^2-4} \end{aligned}$$

and

$$a_{3(r-2)(r+2)} = (-1)^{(r-1)/2} 4\beta^{3(r^2-4)}.$$

Proof. With r an odd number, $m = 2$ in (3.3) gives

$$\arctan \frac{1}{F_r} = \arctan \frac{1}{\alpha^{r-2}} - \arctan \frac{1}{\alpha^{r+2}}.$$

The following identity, proved in [1, Identity (10)]:

$$n\sqrt{n} \arctan \left(\frac{1}{\sqrt{n}} \right) = \sum_{k=0}^{\infty} \frac{1}{n^{2k}} \left(\frac{n}{4k+1} - \frac{1}{4k+3} \right) \quad (3.10)$$

gives

$$\arctan \frac{1}{\alpha^{r-2}} = \sum_{k=0}^{\infty} \frac{1}{\alpha^{(r-2)(4k+3)}} \left(\frac{\alpha^{2r-4}}{4k+1} - \frac{1}{4k+3} \right)$$

and

$$\arctan \frac{1}{\alpha^{r+2}} = \sum_{k=0}^{\infty} \frac{1}{\alpha^{(r+2)(4k+3)}} \left(\frac{\alpha^{2r+4}}{4k+1} - \frac{1}{4k+3} \right),$$

or, by (2.3),

$$\arctan \frac{1}{\alpha^{r-2}} = \sum_{k=0}^{\infty} \frac{1}{\alpha^{(4r^2-16)k}} \sum_{j=1}^{r+2} \left(\frac{\alpha^{-(r-2)(4j-3)}}{4(r+2)k+4j-3} - \frac{\alpha^{-(r-2)(4j-1)}}{4(r+2)k+4j-1} \right)$$

and

$$\arctan \frac{1}{\alpha^{r+2}} = \sum_{k=0}^{\infty} \frac{1}{\alpha^{(4r^2-16)k}} \sum_{j=1}^{r-2} \left(\frac{\alpha^{-(r+2)(4j-3)}}{4(r-2)k+4j-3} - \frac{\alpha^{-(r+2)(4j-1)}}{4(r-2)k+4j-1} \right).$$

Thus,

$$\arctan \frac{1}{F_r} = \sum_{k=0}^{\infty} \frac{1}{\alpha^{(4r^2-16)k}} \times \left(\sum_{j=1}^{r+2} \left(\frac{-\beta^{(r-2)(4j-3)}(r-2)}{4(r^2-4)k + (r-2)(4j-3)} + \frac{\beta^{(r-2)(4j-1)}(r-2)}{4(r^2-4)k + (r-2)(4j-1)} \right) + \sum_{j=1}^{r-2} \left(\frac{\beta^{(r+2)(4j-3)}(r+2)}{4(r^2-4)k + (r+2)(4j-3)} - \frac{\beta^{(r+2)(4j-1)}(r+2)}{4(r^2-4)k + (r+2)(4j-1)} \right) \right). \quad (3.11)$$

Identity (3.9) is (3.11) expressed in the P -notation. \square

Examples.

$$\arctan \frac{1}{F_3} = \arctan \frac{1}{2} = P(1, \alpha^{20}, 20, (-\beta, 0, \beta^3, 0, 4\beta^5, 0, \beta^7, 0, -\beta^9, 0, \beta^{11}, 0, -\beta^{13}, 0, -4\beta^{15}, 0, -\beta^{17}, 0, \beta^{19}, 0)), \quad (3.12)$$

$$\arctan \frac{1}{F_5} = \arctan \frac{1}{5} = P(1, \alpha^{84}, 84, (0, 0, -3\beta^3, 0, 0, 0, 7\beta^7, 0, 3\beta^9, 0, 0, 0, 0, 0, -3\beta^{15}, 0, 0, 0, 0, 0, -4\beta^{21}, 0, 0, 0, 0, 0, -3\beta^{27}, 0, 0, 0, 0, 0, 3\beta^{33}, 0, 7\beta^{35}, 0, 0, 0, -3\beta^{39}, 0, 0, 0, 0, 0, 3\beta^{45}, 0, 0, 0, -7\beta^{49}, 0, -3\beta^{51}, 0, 0, 0, 0, 0, 3\beta^{57}, 0, 0, 0, 0, 0, 4\beta^{63}, 0, 0, 0, 0, 0, 3\beta^{69}, 0, 0, 0, 0, 0, -3\beta^{75}, 0, -7\beta^{77}, 0, 0, 0, 3\beta^{81}, 0, 0, 0)), \quad (3.13)$$

$$\arctan \frac{1}{F_7} = \arctan \frac{1}{13} = P(1, \alpha^{180}, 180, (0, 0, 0, 0, -5\beta^5, 0, 0, 0, 9\beta^9, 0, 0, 0, 0, 0, 5\beta^{15}, 0, 0, 0, 0, 0, 0, 0, 0, 0, -5\beta^{25}, 0, -9\beta^{27}, 0, 0, 0, 0, 0, 0, 0, 5\beta^{35}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 4\beta^{45}, 0, 0, 0, 0, 0, 0, 0, 0, 5\beta^{55}, 0, 0, 0, 0, 0, 0, -9\beta^{63}, 0, -5\beta^{65}, 0, 0, 0, 0, 0, 0, 0, 0, 5\beta^{75}, 0, 0, 0, 0, 0, 9\beta^{81}, 0, 0, 0, -5\beta^{85}, 0, 0, 0, 0, 0, 0, 0, 0, 5\beta^{95}, 0, 0, 0, -9\beta^{99}, 0, 0, 0, 0, 0, -5\beta^{105}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 5\beta^{115}, 0, 9\beta^{117}, 0, 0, 0, 0, 0, 0, 0, -5\beta^{125}, 0, 0, 0, 0, 0, 0, 0, 0, 0, -4\beta^{135}, 0, 0, 0, 0, 0, 0, 0, 0, -5\beta^{145}, 0, 0, 0, 0, 0, 0, 0, 9\beta^{153}, 0, 5\beta^{155}, 0, 0, 0, 0, 0, 0, 0, 0, -5\beta^{165}, 0, 0, 0, 0, 0, -9\beta^{171}, 0, 0, 0, 5\beta^{175}, 0, 0, 0, 0, 0)). \quad (3.14)$$

Theorem 3.5. *If r is a positive even integer, then,*

$$\arctan \frac{1}{F_r} = P(1, \alpha^{4(r^2-1)}, 4(r^2-1), (a_1, a_2, \dots, a_{4(r^2-1)})), \quad (3.15)$$

where the only non-zero constants a_j are given by

$$\begin{aligned} a_{(r-1)(4j-3)} &= -\beta^{(r-1)(4j-3)}(r-1), & j = 1, 2, \dots, r+1, \\ a_{(r-1)(4j-1)} &= \beta^{(r-1)(4j-1)}(r-1), & j = 1, 2, \dots, r+1, \\ a_{(r+1)(4j-3)} &= -\beta^{(r+1)(4j-3)}(r+1), & j = 1, 2, \dots, r-1, \\ a_{(r+1)(4j-1)} &= \beta^{(r+1)(4j-1)}(r+1), & j = 1, 2, \dots, r-1, \\ a_{(r-1)(r+1)} &= (-1)^{r/2} 2\beta^{r^2-1} \end{aligned}$$

and

$$a_{3(r-1)(r+1)} = (-1)^{(r+2)/2} 2\beta^{3(r^2-1)}.$$

Proof. Setting $m = 1$ in (3.4) gives

$$\arctan \frac{1}{F_r} = \arctan \frac{1}{\alpha^{r-1}} + \arctan \frac{1}{\alpha^{r+1}}, \quad r \text{ even.}$$

The proof now proceeds as in that of Theorem 3.4. □

Examples.

$$\arctan \frac{1}{F_2} = \frac{\pi}{4} = P(1, \alpha^{12}, 12, (-\beta, 0, -2\beta^3, 0, -\beta^5, 0, \beta^7, 0, 2\beta^9, 0, \beta^{11}, 0)), \quad (3.16)$$

$$\begin{aligned} \arctan \frac{1}{F_4} = \arctan \frac{1}{3} = P(1, \alpha^{60}, 60, (0, 0, -3\beta^3, 0, -5\beta^5, 0, 0, 0, 3\beta^9, 0, 0, 0, 0, 0, 2\beta^{15}, \\ 0, 0, 0, 0, 0, 3\beta^{21}, 0, 0, 0, -5\beta^{25}, 0, -3\beta^{27}, 0, 0, 0, 0, 0, 3\beta^{33}, \\ 0, 5\beta^{35}, 0, 0, 0, -3\beta^{39}, 0, 0, 0, 0, 0, -2\beta^{45}, 0, 0, 0, 0, 0, -3\beta^{51}, \\ 0, 0, 0, 5\beta^{55}, 0, 3\beta^{57}, 0, 0, 0)). \end{aligned} \quad (3.17)$$

Theorem 3.6. *If r is a positive even integer, then,*

$$\arctan \frac{1}{L_r} = P(1, \alpha^{4(r^2-1)}, 4(r^2-1), (a_1, a_2, \dots, a_{4(r^2-1)})), \quad (3.18)$$

where the only non-zero constants a_j are given by

$$a_{(r-1)(4j-3)} = -\beta^{(r-1)(4j-3)}(r-1), \quad j = 1, 2, \dots, r+1,$$

$$\begin{aligned}
a_{(r-1)(4j-1)} &= \beta^{(r-1)(4j-1)}(r-1), \quad j = 1, 2, \dots, r+1, \\
a_{(r+1)(4j-3)} &= \beta^{(r+1)(4j-3)}(r+1), \quad j = 1, 2, \dots, r-1, \\
a_{(r+1)(4j-1)} &= -\beta^{(r+1)(4j-1)}(r+1), \quad j = 1, 2, \dots, r-1, \\
a_{(r-1)(r+1)} &= (-1)^{(r+2)/2} 2r \beta^{r^2-1}
\end{aligned}$$

and

$$a_{3(r-1)(r+1)} = (-1)^{r/2} 2r \beta^{3(r^2-1)}.$$

Proof. Setting $m = 1$ in (3.3) gives

$$\arctan \frac{1}{L_r} = \arctan \frac{1}{\alpha^{r-1}} - \arctan \frac{1}{\alpha^{r+1}}, \quad r \text{ even.}$$

The proof now proceeds as in that of Theorem 3.4. \square

Examples.

$$\arctan \frac{1}{L_2} = \arctan \frac{1}{3} = (1, \alpha^{12}, 12, (-\beta, 0, 4\beta^3, 0, -\beta^5, 0, \beta^7, 0, -4\beta^9, 0, \beta^{11}, 0)), \quad (3.19)$$

$$\begin{aligned}
\arctan \frac{1}{L_4} = \arctan \frac{1}{7} &= (1, \alpha^{60}, 60, (0, 0, -3\beta^3, 0, 5\beta^5, 0, 0, 0, 3\beta^9, 0, 0, 0, 0, 0, -8\beta^{15}, \\
&\quad 0, 0, 0, 0, 3\beta^{21}, 0, 0, 0, 5\beta^{25}, 0, -3\beta^{27}, 0, 0, 0, 0, 0, 3\beta^{33}, \\
&\quad 0, -5\beta^{35}, 0, 0, 0, -3\beta^{39}, 0, 0, 0, 0, 0, 8\beta^{45}, 0, 0, 0, 0, 0, \\
&\quad -3\beta^{51}, 0, 0, 0, -5\beta^{55}, 0, 3\beta^{57}, 0, 0, 0)). \quad (3.20)
\end{aligned}$$

Theorem 3.7. *If r is an integer, then,*

$$\begin{aligned}
&\sum_{k=0}^{\infty} \frac{1}{\alpha^{12rk}} \left(\frac{\beta^r}{12k+1} + \frac{2\beta^{3r}}{12k+3} + \frac{\beta^{5r}}{12k+5} - \frac{\beta^{7r}}{12k+7} - \frac{2\beta^{9r}}{12k+9} - \frac{\beta^{11r}}{12k+11} \right) \\
&= \begin{cases} -\arctan \left(\frac{1}{L_r} \right), & r \text{ odd,} \\ \arctan \left(\frac{1}{F_r \sqrt{5}} \right), & r \text{ even;} \end{cases} \quad (3.21)
\end{aligned}$$

that is,

$$\begin{aligned}
&P(1, \alpha^{12r}, 12, (\beta^r, 0, 2\beta^{3r}, 0, \beta^{5r}, 0, -\beta^{7r}, 0, -2\beta^{9r}, 0, -\beta^{11r}, 0)) \\
&= \begin{cases} -\arctan \left(\frac{1}{L_r} \right), & r \text{ odd,} \\ \arctan \left(\frac{1}{F_r \sqrt{5}} \right), & r \text{ even.} \end{cases} \quad (3.22)
\end{aligned}$$

Proof. In [1, Identity (27)], it was shown that

$$n^2 \sqrt{n} \arctan \left(\frac{\sqrt{n}}{n-1} \right) = \sum_{k=0}^{\infty} \frac{1}{(-n^3)^k} \left(\frac{n^2}{6k+1} + \frac{2n}{6k+3} + \frac{1}{6k+5} \right).$$

In base n^6 , length 12, this is

$$\begin{aligned} n^2 \sqrt{n} \arctan \left(\frac{\sqrt{n}}{n-1} \right) \\ = \sum_{k=0}^{\infty} \frac{1}{n^{6k}} \left(\frac{n^2}{12k+1} + \frac{2n}{12k+3} + \frac{1}{12k+5} - \frac{1/n}{12k+7} - \frac{2/n^2}{12k+9} - \frac{1/n^3}{12k+11} \right). \end{aligned} \quad (3.23)$$

Identity (3.21) follows upon setting $n = \alpha^{2r}$ in (3.23) and making use of (3.5) and (3.6). \square

Examples.

$$\frac{\pi}{4} = P(1, \alpha^{12}, 12, (-\beta, 0, -2\beta^3, 0, -\beta^5, 0, \beta^7, 0, 2\beta^9, 0, \beta^{11}, 0)), \quad (3.24)$$

$$\arctan \left(\frac{1}{4} \right) = P(1, \alpha^{12}, 12, (-\beta^3, 0, -2\beta^9, 0, -\beta^{15}, 0, \beta^{21}, 0, 2\beta^{27}, 0, \beta^{33}, 0)), \quad (3.25)$$

$$\arctan \left(\frac{1}{\sqrt{5}} \right) = P(1, \alpha^{12}, 12, (\beta^2, 0, 2\beta^6, 0, \beta^{10}, 0, -\beta^{14}, 0, -2\beta^{18}, 0, -\beta^{22}, 0)), \quad (3.26)$$

$$\arctan \left(\frac{1}{3\sqrt{5}} \right) = P(1, \alpha^{12}, 12, (\beta^4, 0, 2\beta^{12}, 0, \beta^{20}, 0, -\beta^{28}, 0, -2\beta^{36}, 0, -\beta^{44}, 0)). \quad (3.27)$$

Remark. Identity (3.24) is the same golden ratio base expansion of π that was obtained in Theorem 3.5.

Theorem 3.8. *If r is an integer, then,*

$$\sum_{k=0}^{\infty} \frac{1}{\alpha^{4rk}} \left(\frac{2\beta^r}{4k+1} - \frac{2\beta^{3r}}{4k+3} \right) = \begin{cases} -\arctan \left(\frac{2}{L_r} \right), & r \text{ odd}, \\ \arctan \left(\frac{2}{F_r \sqrt{5}} \right), & r \text{ even}; \end{cases} \quad (3.28)$$

that is

$$P(1, \alpha^{4r}, 4, (2\beta^r, 0, -2\beta^{3r}, 0)) = \begin{cases} -\arctan \left(\frac{2}{L_r} \right), & r \text{ odd}, \\ \arctan \left(\frac{2}{F_r \sqrt{5}} \right), & r \text{ even}. \end{cases} \quad (3.29)$$

Proof. Setting $x = \beta^r$ in the identity

$$2 \arctan x = \arctan \left(\frac{2x}{1-x^2} \right)$$

and using (3.5) and (3.6), we have

$$2 \arctan \frac{1}{\alpha^r} = \begin{cases} \arctan \left(\frac{2}{L_r} \right), & r \text{ odd,} \\ \arctan \left(\frac{2}{F_r \sqrt{5}} \right), & r \text{ even.} \end{cases} \quad (3.30)$$

Setting $n = \alpha^{2r}$ in (3.10) and comparing with (3.30), we obtain (3.28). \square

Examples.

$$\arctan \frac{2}{L_3} = \arctan \frac{1}{2} = P(1, \alpha^{12}, 4, (-2\beta^3, 0, 2\beta^9, 0)), \quad (3.31)$$

$$\arctan \left(\frac{2}{F_2 \sqrt{5}} \right) = \arctan \frac{2}{\sqrt{5}} = P(1, \alpha^8, 4, (2\beta^2, 0, -2\beta^6, 0)). \quad (3.32)$$

4 Zero relations

Zero relations are expansion formulas that evaluate to zero. They are useful in the determination and classification of new expansion formulas. A base α expansion is not considered new if it can be written as a linear combination of existing formulas and known zero relations.

4.1 Zero relations arising from the logarithm formulas

Zero relation from $\log(F_3^2/L_3) = 0$

Theorem 4.1. *We have*

$$\sum_{k=0}^{\infty} \frac{1}{\alpha^{12k}} \left(\frac{1}{6k+1} - \frac{3\beta^2}{6k+2} - \frac{8\beta^4}{6k+3} - \frac{3\beta^6}{6k+4} + \frac{\beta^8}{6k+5} \right) = 0;$$

that is,

$$0 = P(1, \alpha^{12}, 6, (1, -3\beta^2, -8\beta^4, -3\beta^6, \beta^8, 0)).$$

Proof. We have

$$2 \log F_3 - \log L_3 = 0. \quad (4.1)$$

The expansion of $\log L_3$ given in (2.32) has the following base α^{12} , length 12 version:

$$\log L_3 = P(1, \alpha^{12}, 6, (3\beta^2, 3\beta^4, 0, 3\beta^8, 3\beta^{10}, 0)). \quad (4.2)$$

Use of (2.20) and (4.2) in (4.1) yields the zero relation stated in Theorem 4.1. \square

Zero relation from $\log(L_6/(F_4^2 F_3)) = 0$

Theorem 4.2. *We have*

$$0 = P(1, \alpha^{48}, 24, (1, -5\beta^2, -2\beta^4, 3\beta^6, \beta^8, 4\beta^{10}, \beta^{12}, 3\beta^{14}, \\ -2\beta^{16}, -5\beta^{18}, \beta^{20}, 0, \beta^{24}, -5\beta^{26}, -2\beta^{28}, 3\beta^{30}, \beta^{32}, \\ 4\beta^{34}, \beta^{36}, 3\beta^{38}, -2\beta^{40}, -5\beta^{42}, \beta^{44}, 0)).$$

Proof. Write $\log F_3$, $\log F_4$ and $\log L_6$, that is, identities (2.20), (2.22) and (2.34), respectively, in the common base α^{48} and common length 24 and use

$$\log L_6 - 2 \log F_4 - \log F_3 = 0.$$

□

Zero relation from $\log(F_{12}/(F_3^4 L_2^2)) = 0$

Theorem 4.3. *We have*

$$0 = P(1, \alpha^{48}, 24, (1, -4\beta^2, -5\beta^4, 0, \beta^8, 2\beta^{10}, \beta^{12}, 0, -5\beta^{16}, -4\beta^{18}, \beta^{20}, 0, \\ \beta^{24}, -4\beta^{26}, -5\beta^{28}, 0, \beta^{32}, 2\beta^{34}, \beta^{36}, 0, -5\beta^{40}, -4\beta^{42}, \beta^{44}, 0)).$$

Proof. Write $\log F_3$, $\log F_{12}$ and $\log L_2$ from (2.20), (2.24) and (2.31), in common base α^{48} and consider

$$\log F_{12} - 4 \log F_3 - 2 \log L_2 = 0.$$

□

4.2 Zero relations arising from the inverse tangent formulas

Zero relation from $2 \arctan(2/L_3) + \arctan(2/L_5) - \arctan(2/L_1) = 0$

Theorem 4.4. *We have*

$$0 = P(1, \alpha^{60}, 60, (1, 0, -7\beta^2, 0, -4\beta^4, 0, -\beta^6, 0, 7\beta^8, 0, -\beta^{10}, 0, b^{12}, 0, -2\beta^{14}, \\ 0, \beta^{16}, 0, -\beta^{18}, 0, 7\beta^{20}, 0, -\beta^{22}, 0, -4\beta^{24}, 0, -7\beta^{26}, 0, \beta^{28}, 0, -\beta^{30}, 0, \\ 7\beta^{32}, 0, 4\beta^{34}, 0, \beta^{36}, 0, -7\beta^{38}, 0, \beta^{40}, 0, -\beta^{42}, 0, 2\beta^{44}, 0, -\beta^{46}, 0, \\ \beta^{48}, 0, -7\beta^{50}, 0, \beta^{52}, 0, 4\beta^{54}, 0, 7\beta^{56}, 0, -\beta^{58}, 0)).$$

Proof. Using the addition and subtraction formulas for inverse tangents, it is readily verified that

$$\arctan\left(\frac{2}{L_1}\right) - \arctan\left(\frac{2}{L_3}\right) = \arctan\left(\frac{3}{4}\right)$$

and

$$\arctan\left(\frac{2}{L_5}\right) + \arctan\left(\frac{2}{L_3}\right) = \arctan\left(\frac{3}{4}\right);$$

so that

$$2 \arctan\left(\frac{2}{L_3}\right) + \arctan\left(\frac{2}{L_5}\right) - \arctan\left(\frac{2}{L_1}\right) = 0,$$

from which the zero relation follows upon use of (3.29). \square

Remark. The zero relation stated in Theorem 4.4 can also be obtained directly from

$$\arctan \frac{1}{F_3} = \arctan \frac{2}{L_3},$$

by writing (3.12) and (3.31) in the common base α^{60} and common length 60; or from

$$\arctan \frac{1}{F_4} = \arctan \frac{1}{L_2},$$

using (3.17) and (3.19).

Zero relation from

$$2 \arctan(1/L_1) - 2 \arctan(2/(F_2\sqrt{5})) - \arctan(2/(F_6\sqrt{5})) = 0$$

Theorem 4.5. *We have*

$$0 = P(1, \alpha^{24}, 24, (1, 4\beta, 2\beta^2, 0, \beta^4, 2\beta^5, -\beta^6, 0, -2\beta^8, 4\beta^9, -\beta^{10}, 0, \beta^{12}, -4\beta^{13}, 2\beta^{14}, 0, \beta^{16}, -2\beta^{17}, -\beta^{18}, 0, -2\beta^{20}, -4\beta^{21}, -\beta^{22}, 0)).$$

Proof. The identity

$$\begin{aligned} \frac{\pi}{2} - \arctan\left(\frac{2}{\sqrt{5}}\right) &= \arctan\left(\frac{1}{4\sqrt{5}}\right) + \arctan\left(\frac{2}{\sqrt{5}}\right) \\ &= \arctan\left(\frac{\sqrt{5}}{2}\right) \end{aligned}$$

can be arranged as

$$2 \arctan\left(\frac{1}{L_1}\right) - 2 \arctan\left(\frac{2}{F_2\sqrt{5}}\right) - \arctan\left(\frac{2}{F_6\sqrt{5}}\right) = 0$$

which, on account of (3.29), gives the stated zero relation. \square

5 Other degree 1 base α expansions and zero relations

Base α expansions of $\log \alpha$

Theorem 5.1.

$$\log \alpha = P(1, \alpha, 2, (0, -\beta)). \quad (5.1)$$

Proof. We have

$$\log \alpha = \frac{1}{2} \operatorname{Li}_1 \left(\frac{1}{\alpha} \right) = \frac{1}{2} \sum_{k=0}^{\infty} \frac{1}{\alpha^k} \frac{1/\alpha}{k+1} = \sum_{k=0}^{\infty} \frac{1}{\alpha^k} \frac{-\beta}{2k+2}.$$

□

Theorem 5.2.

$$\log \alpha = P(1, \alpha^2, 2, (0, 2\beta^2)). \quad (5.2)$$

Proof. We have

$$\log \alpha = \operatorname{Li}_1 \left(\frac{1}{\alpha^2} \right) = \sum_{k=0}^{\infty} \frac{1}{\alpha^{2k}} \frac{1/\alpha^2}{k+1} = \sum_{k=0}^{\infty} \frac{1}{\alpha^{2k}} \frac{2\beta^2}{2k+2}.$$

□

Another base α expansion of $\log 2$

Theorem 5.3.

$$\log 2 = P(1, \alpha^3, 3, (-\beta, \beta^2, 2\beta^3)). \quad (5.3)$$

Proof. A straightforward consequence of the identity

$$\log 2 = \operatorname{Li}_1 \left(\frac{1}{\alpha} \right) - \operatorname{Li}_1 \left(\frac{1}{\alpha^3} \right).$$

□

Another base α expansion of $\log 5$

Theorem 5.4.

$$\log 5 = P(1, \alpha^4, 2, (4\beta^2, 0)). \quad (5.4)$$

Proof. A consequence of the identity

$$\log 5 = 2 \operatorname{Li}_1 \left(\frac{1}{\alpha^2} \right) - 2 \operatorname{Li}_1 \left(-\frac{1}{\alpha^2} \right).$$

□

A length 2, base α zero relation

Theorem 5.5.

$$P(1, \alpha^2, 2, (1, 3\beta)) = 0. \quad (5.5)$$

Proof. Follows from

$$\operatorname{Li}_1\left(\frac{1}{\alpha^2}\right) + \operatorname{Li}_1\left(-\frac{1}{\alpha}\right) = 0.$$

□

Remark. Relation (5.5) also follows from (5.1) and (5.2).

A length 12, base α zero relation

Theorem 5.6.

$$P(1, \alpha^{12}, 12, (1, \beta, -2\beta^2, 5\beta^3, \beta^4, 10\beta^5, \beta^6, 5\beta^7, -2\beta^8, \beta^9, \beta^{10}, 2\beta^{11})) = 0.$$

Proof. Follows from (2.20) and (5.3).

□

A length 10, base α zero relation

Theorem 5.7.

$$P(1, \alpha^{20}, 10, (1, -5\beta^2, \beta^4, -5\beta^6, -4\beta^8, -5\beta^{10}, \beta^{12}, -5\beta^{14}, \beta^{16}, 0)) = 0.$$

Proof. Ensues from (2.21) and (5.4).

□

A length 5, base α zero relation

Theorem 5.8.

$$P(1, \alpha^5, 5, (\beta, 1, -\beta, -\beta^4, -2\beta^4)) = 0. \quad (5.6)$$

Proof. Setting $p = 2 \cos x$ in the identity

$$\sum_{k=1}^{\infty} \frac{p^k \cos(kx)}{k} = -\frac{1}{2} \log(1 - 2p \cos x + p^2)$$

produces

$$\sum_{k=1}^{\infty} \frac{(2 \cos x)^k \cos(kx)}{k} = 0. \quad (5.7)$$

Now $2 \cos(2\pi/5) = -\beta$.

□

Thus, setting $x = 2\pi/5$ in (5.7) gives

$$\sum_{k=0}^{\infty} \frac{1}{\alpha^{5k}} \left(\frac{\beta}{5k+1} + \frac{1}{5k+2} - \frac{\beta}{5k+3} - \frac{\beta^4}{5k+4} - \frac{2\beta^4}{5k+5} \right) = 0;$$

since

$$\cos\left(\frac{2\pi}{5}(5j-4)\right) = \frac{-\beta}{2} = \cos\left(\frac{2\pi}{5}(5j-1)\right), \quad j = 1, 2, \dots$$

and

$$\cos\left(\frac{2\pi}{5}(5j-2)\right) = \frac{1}{2\beta} = \cos\left(\frac{2\pi}{5}(5j-5)\right), \quad j = 1, 2, \dots$$

References

- [1] K. Adegoke, A non-PSLQ route to BBP-type formulas, *Journal of Mathematics Research* **2**:2 (2010), 56–54.
- [2] K. Adegoke, The golden ratio, Fibonacci numbers and BBP-type formulas, *The Fibonacci Quarterly* **52**:2 (2014), 129–138.
- [3] D. H. Bailey, A compendium of BBP-type formulas for mathematical constants, Preprint available at <http://www.davidhbailey.com/dhbpapers/bbp-formulas.pdf> (2017).
- [4] D. H. Bailey and P. B. Borwein and S. Plouffe, On the rapid computation of various polylogarithmic constants, *Mathematics of Computation* **66**:218 (1997), 903–913.
- [5] D. H. Bailey and R. E. Crandall, On the random character of fundamental constant expansions, *Experimental Mathematics* **10**:2 (2001), 175–190.
- [6] J. Borwein and M. Chamberland, A Golden Example Solved. Undated manuscript.
- [7] H. C. Chan, π in terms of ϕ , *The Fibonacci Quarterly* **44**:2 (2006), 141–144.
- [8] H. C. Chan, Machin-type formulas expressing π in terms of ϕ , *The Fibonacci Quarterly* **46/47**:1 (2008/2009), 32–37.
- [9] B. Cloitre, A BBP formula for π^2 , Undated manuscript, last seen at <https://les-mathematiques.net/phorum/file.php/2/3190/BBPbasePHI.pdf>
- [10] S. Kristensen and O. Mathiasen, BBP-type formulas—an elementary approach, *Journal of Number Theory* **244** (2023), 251–263.
- [11] C. Wei, Several BBP-type formulas for π , *Integral Transforms and Special Functions* **26**:5 (2015), 315–324.
- [12] W. Zhang, New formulae of BBP-type with different moduli, *Journal of Mathematical Analysis and Applications* **398** (2013), 46–60.