

PAC-Bayes Bounds for High-Dimensional Multi-Index Models with Unknown Active Dimension

Maximilian F. Steffen*

Karlsruhe Institute of Technology

Abstract

The multi-index model with sparse dimension reduction matrix is a popular approach to circumvent the curse of dimensionality in a high-dimensional regression setting. Building on the single-index analysis by Alquier, P. & Biau, G. (Journal of Machine Learning Research 14 (2013) 243–280), we develop a PAC-Bayesian estimation method for a possibly misspecified multi-index model with unknown active dimension and an orthogonal dimension reduction matrix. Our main result is a non-asymptotic oracle inequality, which shows that the estimation method adapts to the active dimension of the model, the sparsity of the dimension reduction matrix and the regularity of the link function. Under a Sobolev regularity assumption on the link function the estimator achieves the minimax rate of convergence (up to a logarithmic factor) and no additional price is paid for the unknown active dimension.

Keywords: multi-index model, PAC-Bayesian, adaptive nonparametric estimation, sparsity, oracle inequality, dimension reduction

1 Introduction

A standard task in supervised learning is to estimate or learn, respectively, the conditional expectation of a label $Y \in \mathbb{R}$ given a large vector $\mathbf{X} \in \mathbb{R}^p$ of explanatory random variables based on i.i.d. data $(\mathbf{X}_i, Y_i)_{i=1, \dots, n}$ which are distributed as (\mathbf{X}, Y) . The corresponding nonparametric regression model reads as

$$Y = F(\mathbf{X}) + \varepsilon \quad (1.1)$$

with an observation error ε satisfying $\mathbb{E}[\varepsilon|\mathbf{X}] = 0$ a.s. and the unknown regression function $F: \mathbb{R}^p \rightarrow \mathbb{R}$ given by $F = \mathbb{E}[Y|\mathbf{X} = \cdot]$. If the dimension p is large, the estimation problem suffers from the well-known curse of dimensionality. This is of particular importance in numerous recent applications where p may exceed the sample size n . In this paper, we provide a fully data driven complete calibration of the high-dimensional multi-index model with unknown active dimension.

A popular approach to reduce the effective dimension of the model is to impose a multi-index structure on F (Li, 1991). While we do not assume that the observations exactly follow a multi-index model, our method builds upon an approximation of the regression function of the form

$$F(\mathbf{x}) \approx f^*(\Theta^* \mathbf{x}), \quad \forall \mathbf{x} \in \mathbb{R}^p, \quad (1.2)$$

for some *active dimension* $d^* \ll p$, a sparse *dimension reduction matrix* $\Theta^* \in \mathbb{R}^{d^* \times p}$ and a (measurable) *link function* $f^*: \mathbb{R}^{d^*} \rightarrow \mathbb{R}$. Following the aforementioned Li (1991), the estimation of the space spanned by the rows of Θ^* has been studied extensively in the literature, see e.g. Hristache et al. (2001), Xia (2007) and Dalalyan et al. (2008), but under the assumption of a known active dimension d^* . While

*Financial support of the DFG through project TR 1349/3-1 is gratefully acknowledged. Large parts of this research were carried out while the author was affiliated to Universität Hamburg. The author would like to thank Mathias Trabs for helpful comments.

some research has been done on the estimation of d^* itself, see Xia et al. (2002) and Zhu et al. (2006), the estimation of the overall model has relied on estimating Θ^* and f^* separately to then analyse the propagation error, see Klock et al. (2021). The analysis of high-dimensional multi-index models, where $p \gg n$, is rather limited.

We use a PAC-Bayesian estimation approach, see Guedj (2019) and Alquier (2021) for an overview, which was originally developed by Catoni (2004, 2007) and has been adapted to the single-index model (i.e. $d^* = 1$) without misspecification by Alquier & Biau (2013). In this paper we generalise the PAC-Bayes method for single index models to the more flexible class of multi-index models. In particular, we aim for a method which adapts to the unknown active dimension d^* , the sparsity of Θ^* and the regularity of f^* to achieve a good approximation (1.2) based on the given data.

As a standard assumption in the theory of multi-index models, we suppose that the dimension reduction matrix is (semi-)orthogonal, i.e. $\Theta^*(\Theta^*)^\top = I_{d^* \times d^*}$ is the identity matrix, see Xia (2008, Proposition 1.1). Indeed, this allows for the interpretation of $\Theta^*\mathbf{X}$ as a rotation of the covariates, projected onto the first d^* coordinates followed by another rotation.

The PAC-Bayes approach relies on the following principle: With a prior π for the parameters (d, Θ, f) we consider the Gibbs-posterior probability distribution $\hat{\rho}_\lambda$ whose π -density is (up to normalisation) given by

$$\frac{d\hat{\rho}_\lambda}{d\pi}(d, \Theta, f) \propto \exp(-\lambda R_n(d, \Theta, f)) \quad (1.3)$$

with a tuning parameter $\lambda > 0$ and empirical prediction risk

$$R_n(d^*, \Theta^*, f^*) = \frac{1}{n} \sum_{i=1}^n (Y_i - f^*(\Theta^* \mathbf{X}_i))^2.$$

The estimator for (d^*, Θ^*, f^*) is obtained by simulating a random variable

$$(\hat{d}_\lambda, \hat{\Theta}_\lambda, \hat{f}_\lambda) \sim \hat{\rho}_\lambda. \quad (1.4)$$

While (1.3) coincides with the classical Bayesian posterior distribution only if $Y_i = f(\Theta \mathbf{X}_i) + \varepsilon_i$ with i.i.d. $\varepsilon_i \sim N(0, n/(2\lambda))$, the estimator $\hat{F}_{\hat{d}} := \hat{f}_\lambda(\hat{\Theta}_\lambda \cdot)$ will achieve a small prediction error under quite mild model assumptions.

We will choose a sieve prior that prefers models with a low active dimension, sparse dimension reduction matrices and regular link functions. Let π be supported on $\bigcup_{d=1}^p \{d\} \times \mathcal{S}_d \times \mathcal{F}_d$ for some classes \mathcal{S}_d and \mathcal{F}_d for Θ and f , respectively. For \mathcal{S}_d we will study a class of sparse matrices while \mathcal{F}_d will be given by finite dimensional wavelet approximations. The prior is uniform for a given sparsity and a wavelet projection level. The posterior weighs each triplet of parameters (d, Θ, f) based on its empirical performance (with respect to the empirical loss function) on the data, where the tuning parameter λ determines the impact of $R_n(d, \Theta, f)$ in comparison to the prior beliefs.

We will quantify the accuracy of the estimation procedure in terms of the excess risk

$$\mathcal{E}(d^*, \Theta^*, f^*) := R(d^*, \Theta^*, f^*) - \min_{d, \Theta, f} R(d, \Theta, f) = \mathbb{E}[(f^*(\Theta^* \mathbf{X}) - F(\mathbf{X}))^2],$$

where

$$R(d^*, \Theta^*, f^*) := \mathbb{E}[(Y - f^*(\Theta^* \mathbf{X}))^2]$$

is the prediction risk and the minimum is attained at $\Theta^* = I_{p \times p}$ and $f^* = F$. We will prove an oracle inequality verifying that the PAC-Bayes estimator is not worse than the optimal choices for $\Theta \in \mathcal{S}_d$ and $f \in \mathcal{F}_d$ for any d . In particular, the overall quality of the method depends on the approximation properties of the spaces \mathcal{S}_d and \mathcal{F}_d .

The paper is organised as follows: In Section 2, we explain our estimation method. In Section 3, the main results are stated. The proofs have been postponed to Section 4.

2 Construction of the prior

To construct the prior, we will introduce for any dimension $d = 1, \dots, p$ classes \mathcal{S}_d and \mathcal{F}_d together with priors μ_d and ν_d for the dimension reduction matrix Θ and the link function f , respectively. Based on that we can then define the prior π on $\bigcup_{d=1}^p \{d\} \times \mathcal{S}_d \times \mathcal{F}_d$.

We start for a fixed active dimension $d \in \{1, \dots, p\}$. While we have mentioned above that an optimal dimension reduction matrix Θ^* should be orthogonal, we will not impose this restriction for the estimation method. Instead, we only require that the candidate matrices have $\|\cdot\|_2$ -normed rows, i.e. for $\Theta = (\vartheta_1, \dots, \vartheta_d)^\top \in \mathbb{R}^{d \times p}$ with row vectors $\vartheta_i = (\vartheta_{i,1}, \dots, \vartheta_{i,p}) \in \mathbb{R}^p$ we impose $\|\vartheta_i\|_2 := (\sum_{j=1}^p \vartheta_{i,j}^2)^{1/2} = 1$. To encode sparsity, let

$$\mathcal{I}_d := \{I \mid \emptyset \neq I := I_1 \times \dots \times I_d, I_1, \dots, I_d \subseteq \{1, \dots, p\}\}$$

contain all potential sets of *active coordinates*, that is I_i describes the active coordinates in the i -th argument of the link function. For $I = I_1 \times \dots \times I_d \in \mathcal{I}_d$ the number of active coordinates is $\|I\| := \sum_{i=1}^d |I_i|$, where $|I_i|$ denotes the cardinality of I_i . Note that $\emptyset \neq I = I_1 \times \dots \times I_d$ already implies $I_1, \dots, I_d \neq \emptyset$. The parameter set $\mathcal{S}_d(I)$ of sparse dimension reduction matrices is given by

$$\begin{aligned} \mathcal{S}_d(I) &:= \{\Theta = (\vartheta_1, \dots, \vartheta_d)^\top \in \mathbb{R}^{d \times p} \mid \vartheta_i \in \mathcal{S}(I_i), i = 1, \dots, d\}, \quad \text{where} \\ \mathcal{S}(I_i) &:= \{\vartheta_i = (\vartheta_{i,1}, \dots, \vartheta_{i,p}) \in \mathbb{R}^p \mid \|\vartheta_i\|_2 = 1, \forall j \notin I_i : \vartheta_{i,j} = 0\}. \end{aligned}$$

Finally, we define $\mathcal{S}_d = \bigcup_{I \in \mathcal{I}_d} \mathcal{S}_d(I)$.

Note that $\mathcal{S}_d(I) \supseteq \tilde{\mathcal{S}}_d(I)$ for

$$\tilde{\mathcal{S}}_d(I) := \{\Theta = (\vartheta_1, \dots, \vartheta_d)^\top \in \mathbb{R}^{d \times p} \mid \forall i = 1, \dots, d : \|\vartheta_i\|_2 = 1, \vartheta_{i,j} \neq 0 \text{ iff } \vartheta_{i,j} \in I_i, j = 1, \dots, p\}.$$

In $\tilde{\mathcal{S}}_d(I)$ the index set I exactly describes the sparsity of Θ . However, we consider the prior on the compact set $\mathcal{S}_d(I)$ to ensure the existence of solutions to minimisation problems over $\mathcal{S}_d(I)$ and thus the existence of an oracle dimension reduction matrix as a benchmark for $\hat{\Theta}$.

To construct a prior measure μ_d on \mathcal{S}_d , we use the uniform distribution on the set of dimension reduction matrices with a given active dimension d and with sparsity $i = \|I\|$. These uniform distributions are then weighted geometrically such that sparse dimension reduction matrices are preferred by the prior. Denoting the uniform distribution on $\mathcal{S}_d(I)$ by $\mu_{d,I}$, the prior measure on \mathcal{S}_d is thus given by the mixture

$$\mu_d := C_{\mu,d} \sum_{i=d}^{dp} 10^{-i+d-1} \frac{1}{|\mathcal{I}_{d,i}|} \sum_{I \in \mathcal{I}_{d,i}} \mu_{d,I} \quad \text{where} \quad \mathcal{I}_{d,i} := \{I \in \mathcal{I}_d \mid \|I\| = i\}$$

and with normalisation constant $C_{\mu,d} := 9(1 - 10^{(1-p)d-1})^{-1}$. Here and in the following two analogous constructions the geometric decay 10^{-i+d+1} can be replaced by a^{-i+d+1} for an arbitrary fixed $a > 1$, but we choose 10 for convenience.

To define a class \mathcal{F}_d and a prior ν_d for the link function, we will use a multivariate tensor product wavelet basis on \mathbb{R}^d , see e.g. Daubechies (1992); Giné & Nickl (2016). Let ϕ and ψ be a continuously differentiable scaling and wavelet function on \mathbb{R} , respectively, and write $\psi_0 := \phi$, $\psi_1 := \psi$. We will use compactly supported regular Daubechies wavelets. For $M, N \in \mathbb{N}$ we define the index set

$$\begin{aligned} \mathcal{Z}_{M,N}^d &:= \{l = (0, l_2, 0) \mid l_2 \in \mathbb{Z}^d, \|l_2\|_\infty \leq N\} \\ &\cup \{l = (l_1, l_2, l_3) \in \mathbb{N}_0 \times \mathbb{Z}^d \times \{0, 1\}^d \mid l_1 \leq M, \|l_2\|_\infty \leq 2^{l_1} N, l_3 \neq 0\}, \end{aligned}$$

where l_1 is the approximation level, l_2 is a shift parameter and l_3 is due to the tensor structure. The system $(\Psi_l)_{l \in \mathcal{Z}_{\infty,\infty}^d}$ with

$$\Psi_l(x) := 2^{l_1 d/2} \prod_{i=1}^d \psi_{l_{3,i}}(2^{l_1} x_i - l_{2,i}), \quad x \in \mathbb{R}^d, l = (l_1, l_2, l_3) \in \mathcal{Z}_{\infty,\infty}^d,$$

is an orthonormal basis of $L^2(\mathbb{R}^d)$. In particular, each $f \in L^2(\mathbb{R}^d)$ admits a wavelet series representation $f = \sum_{l \in \mathcal{Z}_{\infty, \infty}^d} \langle f, \Psi_l \rangle \Psi_l$. Throughout, we fix a sufficiently large constant $N \in \mathbb{N}_0$ and abbreviate $\mathcal{Z}_M^d := \mathcal{Z}_{M, N}^d$. For $\xi > 0$ we define the compact wavelet coefficient ball

$$\mathcal{B}_{d, M}(\xi) := \{\beta \in \mathbb{R}^{\mathcal{Z}_M^d} \mid \|\beta\|_{\mathcal{B}} \leq \xi\}, \quad \text{where}$$

$$\|\beta\|_{\mathcal{B}} := L^d \sum_{l \in \mathcal{Z}_M^d} 2^{l_1(d/2+1)} |\beta_l|, \quad L := \|\psi\|_{\infty} \vee \|\phi\|_{\infty} \vee \|\psi'\|_{\infty} \vee \|\phi'\|_{\infty},$$

which determines the finite dimensional approximation space

$$\mathcal{F}_{d, M}(\xi) := \{f = \Phi_{d, M}(\beta) \mid \beta \in \mathcal{B}_{d, M}(\xi)\} \quad \text{via} \quad \Phi_{d, M}(\beta) := \sum_{l \in \mathcal{Z}_M^d} \beta_l \Psi_l, \beta \in \mathbb{R}^{\mathcal{Z}_M^d}.$$

For any $f = \Phi_{d, M}(\beta)$ we write $\|f\|_{\mathcal{B}} := \|\beta\|_{\mathcal{B}}$ which corresponds to the Besov norm with regularity $1 + d$ and integrability parameter 1 on $\text{span}\{\Psi_l : l \in \mathcal{Z}_M^d\}$. In particular, we have for any $f \in \mathcal{F}_{d, M}(\xi)$

$$\|f\|_{\infty} \leq \|f\|_{\mathcal{B}} \leq \xi \quad \text{and} \quad \|(\nabla f)_i\|_{\infty} \leq \|f\|_{\mathcal{B}} \leq \xi, \quad \forall i \in \{1, \dots, d\}. \quad (2.1)$$

For $C > 0$ we set $\mathcal{F}_d := \bigcup_{M=0}^n \mathcal{F}_{d, M}(C + 1)$.

The prior ν_d on \mathcal{F}_d is defined as a random coefficient prior with uniformly distributed coefficients on $\mathcal{F}_{d, M}(C + 1)$ and geometrically decreasing weights in the approximation level M . To this end, let $\tilde{\nu}_{d, M}$ be the uniform distribution on $\mathcal{B}_{d, M}(C + 1)$ and let $\nu_{d, M} := \tilde{\nu}_{d, M}(\Phi_{d, M}^{-1}(\cdot))$ denote the push-forward measure of $\tilde{\nu}_{d, M}$ under $\Phi_{d, M}$. Then

$$\nu_d := C_{\nu, d} \sum_{M=0}^n 10^{-M} \nu_{d, M}, \quad C_{\nu, d} := \frac{9}{10 - 10^{-n}}.$$

We can now define the prior for a fixed active dimension d as the product measure $\pi_d := \delta_d \otimes \mu_d \otimes \nu_d$ with the Dirac measure δ_d in d . Finally, we mix over all possible active dimensions to account for the fact that d^* is unknown. Encoding a preference for simple models, i.e. small active dimensions, via weights 10^{-d} , the final prior on $\bigcup_{d=1}^p \{d\} \times \mathcal{S}_d \times \mathcal{F}_d$ is given by

$$\pi = C_{\pi} \sum_{d=1}^p 10^{-d} \pi_d, \quad C_{\pi} := \frac{9}{1 - 10^{-p}}.$$

The product measure with the Dirac measure in d ensures that a simulation of π will yield a link function and a dimension reduction matrix with matching active dimension.

3 Oracle inequalities

For an active dimension $d \in \{1, \dots, p\}$, an active index set $I \in \mathcal{I}_d$ of the dimension reduction matrix and an approximation level $M \in \{0, \dots, n\}$ of the link function, we define the *oracle choice* on $\mathcal{S}_d(I) \times \mathcal{F}_{d, M}(C)$ as

$$(\Theta_{d, I}^*, f_{d, M}^*) := \arg \min_{(\Theta, f) \in \mathcal{S}_d(I) \times \mathcal{F}_{d, M}(C)} R(d, \Theta, f) \quad (3.1)$$

which is not accessible to the practitioner since $R(d, \Theta, f)$ depends on the unknown distribution of (\mathbf{X}, Y) . Note that the minimisation in f is over $\mathcal{F}_{d, M}(C)$, whereas the prior is defined on $\mathcal{F}_{d, M}(C + 1)$ which ensures that a small neighbourhood of $f_{d, M}^*$ is contained in the support of the prior. A solution to the minimisation problem in (3.1) always exists since we have equivalently

$$(\Theta_{d, I}^*, \beta_{d, M}^*) = \arg \min_{(\Theta, \beta) \in \mathcal{S}_d(I) \times \mathcal{B}_{d, M}(C)} \mathbb{E}[(Y - \Phi_{d, M}(\beta)(\Theta \mathbf{X}))^2]$$

with compact $\mathcal{S}_d(I) \times \mathcal{B}_{d,M}(C)$ and continuous $(\Theta, \beta) \mapsto \mathbb{E}[(Y - \Phi_{d,M}(\beta)(\Theta \mathbf{X}))^2]$. If there is more than one solution, we choose one of them. Our main result gives a theoretical guarantee that the PAC-Bayes estimator $(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda)$ from (1.4) is at least as good as the best oracle $(\Theta_{d,I}^*, f_{d,M}^*)_{d,I,M}$ in terms of the excess risk. To this end, we need some mild assumptions on the regression model (1.1).

Assumption 1.

1. For $K, C \geq 1$ we have $\|\mathbf{X}\|_\infty \leq K$ a.s. and $\|F\|_\infty \leq C$.
2. ε is conditionally on \mathbf{X} sub-Gaussian, i.e. there are constants $\sigma, \Gamma > 0$ such that

$$\mathbb{E}[|\varepsilon|^k | \mathbf{X}] \leq \frac{k!}{2} \sigma^2 \Gamma^{k-2} \text{ a.s.}, \quad \forall k \geq 2.$$

We obtain the following non-asymptotic oracle inequality. It generalises Alquier & Biau (2013, Theorem 2) not only with respect to the multi-index approach with unknown active dimension, but also with respect to some technical but practically relevant aspects like the ℓ^2 -normalisation of Θ and the wavelet basis.

Theorem 2. Under Assumption 1 with $Q = 8(2C + 1)(\Gamma \vee (2C + 1))$ set

$$\lambda = \frac{n}{Q + 2((2C + 1)^2 + 4\sigma^2)}.$$

If $n \geq (10C^{-1}) \vee (30\sqrt{2}ep^{-2})$, then we have for any $\delta \in (0, 1)$

$$\mathcal{E}(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda) \leq \inf_{d,I,M} \left(3\mathcal{E}(d, \Theta_{d,I}^*, f_{d,M}^*) + \frac{\Xi}{n} \left(\|I\| \log(pn) + 4^d N^d 2^{dM} \log(Cn) + \log\left(\frac{2}{\delta}\right) \right) \right)$$

with $(\mathbb{P} \otimes \widehat{\rho}_\lambda)$ -probability of at least $1 - \delta$, where Ξ is a constant only depending on C, K, Γ and σ and the infimum is taken over all triplets (d, I, M) with $d \in \{1, \dots, p\}$, $I \in \mathcal{I}_d$ and $M \in \{0, \dots, n\}$.

Some remarks are in order: **1)** The $1 - \delta$ probability with respect to the coupling $\mathbb{P} \otimes \widehat{\rho}_\lambda$ takes into account the randomness of the data and of the estimate. **2)** The right-hand side of the oracle inequality can be interpreted similar to the classical bias-variance decomposition in non-parametric statistics. The first term $\mathcal{E}(d, \Theta_{d,I}^*, f_{d,M}^*) = \mathbb{E}[(f_{d,M}^*(\Theta_{d,I}^* \mathbf{X}) - F(\mathbf{X}))^2]$ quantifies the approximation error while second term is an upper bound for the stochastic error. In particular, we recover $\frac{\|I\|}{n} \log(pn)$ (or $\frac{\|I\|}{n} \log(p)$ if $p \geq n$) as the typical error term for estimating sparse matrices with sparsity $\|I\|$ while $4^d N^d 2^{dM} \frac{\log(n)}{n}$ is due to the estimation of $4^d N^d 2^{dM}$ many wavelet coefficients each with (squared) accuracy $\frac{\log(n)}{n}$ paying a logarithmic price for adaptiveness. **3)** The infimum over all (d, I, M) in the upper bound shows that the estimator adapts to the active dimension, the sparsity of the dimension reduction matrix and the regularity of the link function. **4)** The factor 3 in the upper bound can be improved to $(1 + \tau)$ for any $\tau > 0$ at the cost of a larger constant Ξ . **5)** One can show the same result in a multi-index model with a known active dimension d^* by using π_{d^*} as a prior instead of π . The only difference (up to a constant) in the result is that the infimum in the upper bound is only taken over all pairs $(I, M) \in \mathcal{I}_{d^*} \times \{0, \dots, n\}$. Consequently, no additional price is paid for not knowing the true active dimension of the model.

In the well-specified setting and under assumptions on the distribution of $\Theta^* \mathbf{X}$ as well as a Besov-type regularity assumption on the link function, we can make the upper bound from Theorem 2 more explicit.

Assumption 3.

1. There exist $d^* \in \{1, \dots, p\}$, $\Theta^* \in \mathcal{S}^{d^*}$ and $f^*: \mathbb{R}^{d^*} \rightarrow \mathbb{R}$ such that $F = f^*(\Theta^* \cdot)$.
2. The random variable $\Theta^* \mathbf{X}$ is bounded, i.e. $\|\Theta^* \mathbf{X}\|_\infty \leq B_1$ for some $B_1 > 0$, and $\Theta^* \mathbf{X}$ has a \mathbb{X}^{d^*} -density on \mathbb{R}^{d^*} bounded by a constant $B_2 > 0$.

For the true dimension reduction matrix Θ^* we write $\|\Theta^*\|_0 := \|I^*\|$ for the minimal $I^* \in \mathcal{I}_{d^*}$ with respect to $\|\cdot\|$ such that $\Theta^* \in \mathcal{S}_{d^*}(I^*)$. The regularity of f^* will be measured in terms of its Besov norm. We recover Sobolev balls for $q = 2$, cf. Giné & Nickl (2016, (4.164)).

Definition 4. The Besov ellipsoid in \mathbb{R}^{d^*} with regularity $\alpha > 0$ and integrability parameter $q \in [0, \infty)$ is given by

$$B_{q,d^*}^\alpha(\xi) := \left\{ f \in L^2(\mathbb{R}^{d^*}) \mid \sum_{l \in \mathcal{Z}_{\infty,\infty}^{d^*}} 2^{ql_1\alpha} |\langle f, \Psi_l \rangle|^q \leq \xi^q \right\}$$

for a radius $\xi > 0$.

Corollary 5. Let the assumptions of Theorem 2 be fulfilled in addition to Assumption 3. Suppose that $f^* \in B_{d^*}^\alpha(\xi)$ with $\xi = C(L^{d^*} 2N^{d^*}/2^{8d^*/2})^{-1}$ for some $\alpha > 2 + d^*$ and $n \geq 2$. Then, with $(\mathbb{P} \otimes \hat{\rho}_\lambda)$ -probability of at least $1 - \delta$, we have

$$\mathcal{E}(\hat{d}_\lambda, \hat{\Theta}_\lambda, \hat{f}_\lambda) \leq \Lambda \left(\left(\frac{\log(Cn)}{n} \right)^{\frac{2\alpha}{2\alpha+d^*}} + \frac{\|\Theta^*\|_0 \log(pn)}{n} + \frac{\log(\frac{2}{\delta})}{n} \right),$$

where Λ only depending on $C, K, \Gamma, \sigma, N, B_1, B_2$ and d^* .

Remark. If Θ^* is sparse (i.e. $\|\Theta^*\|_0$ is small), then the dominating term in the upper bound of the excess risk of the PAC-Bayesian estimator is of order

$$\left(\frac{\log(n)}{n} \right)^{\frac{2\alpha}{2\alpha+d^*}},$$

which is the usual minimax rate (up to a logarithmic factor) for such estimation problems. Note that if $d^* \ll p$, then we have successfully circumvented the curse of dimensionality, since the dimension which appears in the rate is now only d^* . As an alternative to the wavelet construction, one can use the multivariate trigonometric system on $[-1, 1]^{d^*}$, assume $\mathbf{X} \in [-1, 1]^p$ and ℓ^1 -standardised rows of Θ^* (which ensures $\Theta^* \mathbf{X} \in [-1, 1]^{d^*}$) leading to a more direct generalisation of Alquier & Biau 2013. However, the orthogonality assumption on Θ^* seems more natural and in line with the literature.

4 Proofs

We begin with a few preliminary results. The first is a classical result on the Kullback-Leibler divergence. A proof can be found in Catoni (2004, p. 159). Afterwards, we present two lemmata of the ‘‘PAC’’-type, which are extensions of Alquier & Biau (2013, Lemmata 7 & 8). The proofs can be carried out analogously.

Let μ, ν be probability measures on a measurable space (E, \mathcal{A}) . The *Kullback-Leibler divergence* of μ with respect to ν is defined via

$$\mathcal{K}(\mu, \nu) := \begin{cases} \int \log\left(\frac{d\mu}{d\nu}\right) d\mu, & \text{if } \mu \ll \nu \\ \infty, & \text{otherwise} \end{cases}.$$

Lemma 6. Let μ be a probability measure on a measurable space (E, \mathcal{A}) and let $h: E \rightarrow \mathbb{R}$ be a measurable function such that $\int \exp \circ h d\mu < \infty$. With the convention $\infty - \infty = -\infty$, it then holds that

$$\log\left(\int \exp \circ h d\mu\right) = \sup_{\nu} \left(\int h d\nu - \mathcal{K}(\nu, \mu)\right) = -\inf_{\nu} \left(\mathcal{K}(\nu, \mu) - \int h d\nu\right), \quad (4.1)$$

where the supremum and the infimum are taken over all probability measures ν on (E, \mathcal{A}) , but no generality is lost by considering only those probability measures ν on (E, \mathcal{A}) such that $\nu \ll \mu$. If additionally, h is bounded from above on the support of μ , then the supremum and the infimum in (4.1) are both attained by taking $\nu = g$ with the Gibbs-posterior probability distribution g , i.e. $\frac{dg}{d\mu} \propto \exp \circ h$.

Lemma 7. Under Assumption 1, let $Q = 8(2C + 1)(\Gamma \vee (2C + 1))$, $\mathcal{E}_n(d, \Theta, f) := R_n(d, \Theta, f) - R_n(d^*, \Theta^*, f^*)$ and take

$$\lambda \in \left(0, \frac{n}{Q + ((2C + 1)^2 + 4\sigma^2)}\right).$$

Then, for all $\delta \in (0, 1)$ and any probability measure $\rho \ll \pi$, we have that

$$\mathcal{E}(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda) \leq \frac{1}{1 - \frac{\lambda((2C+1)^2 + 4\sigma^2)}{n - Q\lambda}} \left(\int \mathcal{E}_n d\rho + \frac{\mathcal{K}(\rho, \pi) + \log(\frac{1}{\delta})}{\lambda} \right)$$

with $(\mathbb{P} \otimes \widehat{\rho}_\lambda)$ -probability of at least $1 - \delta$, where $(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda) \sim \widehat{\rho}_\lambda$.

Lemma 8. In the situation of Lemma 7, we have

$$\int \mathcal{E}_n d\rho \leq \left(1 + \frac{\lambda((2C + 1)^2 + 4\sigma^2)}{n - Q\lambda}\right) \int \mathcal{E} d\rho + \frac{\mathcal{K}(\rho, \pi) + \log(\frac{1}{\delta})}{\lambda}$$

with $(\mathbb{P} \otimes \widehat{\rho}_\lambda)$ -probability of at least $1 - \delta$.

4.1 Proof of Theorem 2

We extend the proof strategy by Alquier & Biau (2013) to the multi-index setting with unknown active dimension.

We begin by fixing some triplet (d, I, M) with $d \in \{1, \dots, p\}$, $I = I_1 \times \dots \times I_d \in \mathcal{I}_d$, $M \in \{0, \dots, n\}$ as well as $\eta, \gamma \in (0, 1]$ and introduce the probability measure

$$\rho := \rho_{d, I, M, \eta, \gamma} := \kappa_d \otimes \rho_{d, I, \eta}^1 \otimes \rho_{d, M, \gamma}^2, \quad (4.2)$$

where $\rho_{d, I, \eta}^1$ and $\rho_{d, M, \gamma}^2$ are the uniform distribution with respect to $\mu_{d, I}$ and $\nu_{d, M}$ on a ball of radius η and γ around the oracle $\Theta_{d, I}^*$ and $f_{d, M}^*$, respectively. Specifically, we set

$$\begin{aligned} \frac{d\rho_{d, I, \eta}^1}{d\mu_{d, I}}(\Theta) &:= \prod_{i=1}^d \frac{d\rho_{d, I, \eta}^{1, i}}{d\mu_{I_i}}(\vartheta_i), \quad \forall \Theta = (\vartheta_1, \dots, \vartheta_d)^\top, \quad \text{where} \\ \frac{d\rho_{d, I, \eta}^{1, i}}{d\mu_{I_i}}(\vartheta_i) &\propto \mathbb{1}_{\{\|\vartheta_i - \vartheta_{d, I}^*\|_2 \leq \eta\}} \quad \text{and} \\ \frac{d\rho_{d, M, \gamma}^2}{d\nu_{d, M}}(f) &\propto \mathbb{1}_{\{\|f - f_{d, M}^*\|_\psi \leq \gamma\}}, \end{aligned} \quad (4.3)$$

where μ_{I_i} denotes the uniform distribution on $\mathcal{S}(I_i)$. Applying Lemmata 7 and 8, we have that

$$\begin{aligned} \mathcal{E}(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda) &\leq \frac{1}{1 - \frac{\lambda((2C+1)^2 + 4\sigma^2)}{n - Q\lambda}} \left(\int \mathcal{E}_n d\rho + \frac{\mathcal{K}(\rho, \pi) + \log(\frac{2}{\delta})}{\lambda} \right), \quad \text{and} \\ \int \mathcal{E}_n d\rho &\leq \left(1 + \frac{\lambda((2C + 1)^2 + 4\sigma^2)}{n - Q\lambda}\right) \int \mathcal{E} d\rho + \frac{\mathcal{K}(\rho, \pi) + \log(\frac{2}{\delta})}{\lambda} \end{aligned}$$

with a probability of at least $1 - \frac{\delta}{2}$, respectively. Therefore,

$$\mathcal{E}(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda) \leq \frac{1}{1 - \frac{\lambda((2C+1)^2 + 4\sigma^2)}{n - Q\lambda}} \left(\left(1 + \frac{\lambda((2C + 1)^2 + 4\sigma^2)}{n - Q\lambda}\right) \int \mathcal{E} d\rho + 2 \frac{\mathcal{K}(\rho, \pi) + \log(\frac{2}{\delta})}{\lambda} \right) \quad (4.4)$$

holds with a probability of at least $1 - \delta$. To complete the proof, we need to bound the terms on the right hand side of (4.4).

First, we deal with the Kullback-Leibler divergence term using the following two lemmata:

Lemma 9. For $\rho = \rho_{d,I,M,\eta,\gamma}$ from (4.2) and with $\pi_{d,I,M} = \kappa_d \otimes \mu_{d,I} \otimes \nu_{d,M}$, we have

$$\mathcal{K}(\rho, \pi) \leq \|I\| \log(ep) + (\|I\| + M + 1) \log(10) + \mathcal{K}(\rho, \pi_{d,I,M}) =: T_1 + \mathcal{K}(\rho, \pi_{d,I,M}).$$

Lemma 10. For $\rho = \rho_{d,I,M,\eta,\gamma}$ from (4.2) and with $\pi_{d,I,M} = \kappa_d \otimes \mu_{d,I} \otimes \nu_{d,M}$, we have

$$\mathcal{K}(\rho, \pi_{d,I,M}) \leq \|I\| \log\left(\frac{3\sqrt{2}}{\eta}\right) + 4^d N^d 2^{dM+1} \log\left(\frac{C+1}{\gamma}\right) =: T_2.$$

Thus,

$$\mathcal{E}(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda) \leq \frac{1}{1 - \frac{\lambda((2C+1)^2 + 4\sigma^2)}{n - Q\lambda}} \left(\left(1 + \frac{\lambda((2C+1)^2 + 4\sigma^2)}{n - Q\lambda}\right) \int \mathcal{E} d\rho + 2 \frac{T_1 + T_2 + \log\left(\frac{2}{\delta}\right)}{\lambda} \right)$$

with a probability of at least $1 - \delta$.

Second, we control the integral term in (4.4) by splitting it into

$$\begin{aligned} \int \mathcal{E}(c, \Theta, f) d\rho(c, \Theta, f) &= \mathcal{E}(d, \Theta_{d,I}^*, f_{d,M}^*) \\ &\quad + \int \mathbb{E}[(f_{c,M}^*(\Theta_{c,I}^* \mathbf{X}) - f(\Theta_{c,M}^* \mathbf{X}))^2] d\rho(c, \Theta, f) \\ &\quad + \int \mathbb{E}[(f(\Theta_{c,I}^* \mathbf{X}) - f(\Theta \mathbf{X}))^2] d\rho(c, \Theta, f) \\ &\quad + \int \mathbb{E}[2(Y - f_{c,M}^*(\Theta_{c,I}^* \mathbf{X}))(f_{c,M}^*(\Theta_{c,I}^* \mathbf{X}) - f(\Theta_{c,I}^* \mathbf{X}))] d\rho(c, \Theta, f) \\ &\quad + \int \mathbb{E}[2(Y - f_{c,M}^*(\Theta_{c,I}^* \mathbf{X}))(f(\Theta_{c,I}^* \mathbf{X}) - f(\Theta \mathbf{X}))] d\rho(c, \Theta, f) \\ &\quad + \int \mathbb{E}[2(f_{c,M}^*(\Theta_{c,I}^* \mathbf{X}) - f(\Theta_{c,I}^* \mathbf{X}))(f(\Theta_{c,I}^* \mathbf{X}) - f(\Theta \mathbf{X}))] d\rho(c, \Theta, f) \\ &=: \mathcal{E}(d, \Theta_{d,I}^*, f_{d,M}^*) + U_1 + U_2 + U_3 + U_4 + U_5 \end{aligned} \tag{4.5}$$

and treating the terms U_1, \dots, U_5 sequentially. Note, that integrating with respect to $\rho = \kappa_d \otimes \rho_{d,I,\eta}^1 \otimes \rho_{d,M,\gamma}^2$ amounts to fixing the first argument at d and then integrating with respect to $\rho_{d,I,\eta}^1 \otimes \rho_{d,M,\gamma}^2$.

Similarly to (2.1), $f = \Phi_{d,M}(\beta) \in \mathcal{F}_{d,M}(C+1)$ with $\|\beta - \beta_{d,M}^*\|_{\mathcal{B}} \leq \gamma$ implies

$$\|f - f_{d,M}^*\|_\infty \leq \|f - f_{d,M}^*\|_\psi = \|\beta - \beta_{d,M}^*\|_{\mathcal{B}} \leq \gamma.$$

Therefore,

$$U_1 = \int \mathbb{E}[(f_{d,M}^*(\Theta_{d,I}^* \mathbf{X}) - f(\Theta_{d,I}^* \mathbf{X}))^2] d\rho_{d,M,\gamma}^2(f) \leq \int \sup_{x \in \mathbb{R}^d} (f_{d,M}^*(x) - f(x))^2 d\rho_{d,M,\gamma}^2(f) \leq \gamma^2. \tag{4.6}$$

Any $f \in \mathcal{F}_{d,M}(C+1)$ is differentiable as a linear combination of only finitely many basis elements. Therefore, applying the fundamental theorem of calculus to the mapping

$$h: [-1, 1] \rightarrow \mathbb{R}, s \mapsto f((\Theta + s(\Theta_{d,I}^* - \Theta)) \mathbf{X}(\omega))$$

with a fixed $\omega \in \Omega$ (which we will omit from here on) yields

$$f(\Theta_{d,I}^* \mathbf{X}) - f(\Theta \mathbf{X}) = \int_0^1 g'(s) ds = \left\langle (\Theta_{d,I}^* - \Theta) \mathbf{X}, \int_0^1 \nabla f((\Theta + s(\Theta_{d,I}^* - \Theta)) \mathbf{X}) ds \right\rangle$$

and combined with (2.1) we obtain

$$\begin{aligned}
|f(\Theta_{d,M}^* \mathbf{X}) - f(\Theta \mathbf{X})| &= \left| \left\langle (\Theta_{d,I}^* - \Theta) \mathbf{X}, \int_0^1 \nabla f((\Theta + s(\Theta_{d,I}^* - \Theta)) \mathbf{X}) ds \right\rangle \right| \\
&\leq \|(\Theta_{d,I}^* - \Theta) \mathbf{X}\|_2 \left\| \int_0^1 \nabla f((\Theta + s(\Theta_{d,I}^* - \Theta)) \mathbf{X}) ds \right\|_2 \\
&\leq \|\mathbf{X}\|_2 \left(\sum_{i=1}^d \|\vartheta_{d,I,i}^* - \vartheta_i\|_2^2 \right)^{\frac{1}{2}} \sqrt{d} C \\
&\leq pK \left(\sum_{i=1}^d \|\vartheta_{d,I,i}^* - \vartheta_i\|_2^2 \right)^{\frac{1}{2}} \sqrt{d} C \mathbb{P}\text{-a.s.}
\end{aligned} \tag{4.7}$$

Using the above, we deduce

$$\begin{aligned}
U_2 &= \int \mathbb{E}[(f(\Theta_{d,I}^* \mathbf{X}) - f(\Theta \mathbf{X}))^2] d\rho_{d,I,\eta}^1 \otimes \rho_{d,M,\gamma}^2(\Theta, f) \\
&\leq d(pKC)^2 \int \cdots \int \sum_{i=1}^d \|\vartheta_{d,I,i}^* - \vartheta_i\|_2^2 d\rho_{d,I,\eta}^{1,1}(\vartheta_1) \cdots d\rho_{d,I,\eta}^{1,d}(\vartheta_d) \\
&\leq (dpKC\eta)^2.
\end{aligned} \tag{4.8}$$

By construction, $\rho_{d,M,\gamma}^2$ is centered around $f_{d,M}^*$ and thus

$$\int f(x) d\rho_{d,M,\gamma}^2(f) = f_{d,M}^*(x), \quad \forall x \in \mathbb{R}^p. \tag{4.9}$$

In particular, we have

$$U_3 = 0. \tag{4.10}$$

Using Fubini's theorem together with (4.9), we have

$$\begin{aligned}
|U_4| &= 2 \left| \mathbb{E}[(Y - f_{d,M}^*(\Theta_{d,I}^* \mathbf{X})) \int \int f(\Theta_{d,I}^* \mathbf{X}) - f(\Theta \mathbf{X}) d\rho_{d,M,\gamma}^2(f) d\rho_{d,I,\eta}^1(\Theta)] \right| \\
&= 2 \left| \mathbb{E}[(Y - f_{d,M}^*(\Theta_{d,I}^* \mathbf{X})) \int f(\Theta_{d,I}^* \mathbf{X}) - f_{d,M}^*(\Theta \mathbf{X}) d\rho_{d,I,\eta}^1(\Theta)] \right| \\
&\leq 2 \sqrt{R(d, \Theta_{d,I}^*, f_{d,M}^*)} \sqrt{\mathbb{E}[(\int f_{d,M}^*(\Theta_{d,I}^* \mathbf{X}) - f_{d,M}^*(\Theta \mathbf{X}) d\rho_{d,I,\eta}^1(\Theta))^2]},
\end{aligned} \tag{4.11}$$

where the Cauchy-Schwarz inequality has been used in the final step. Repeating the argument from treating U_2 , but now with $f = f_{d,M}^*$, we obtain

$$\mathbb{E}[(\int f_{d,M}^*(\Theta_{d,I}^* \mathbf{X}) - f_{d,M}^*(\Theta \mathbf{X}) d\rho_{d,I,\eta}^1(\Theta))^2] \leq (dpKC\eta)^2. \tag{4.12}$$

Clearly, $f = 0 \in \mathcal{F}_{d,M}(C)$ and thus we have by definition of $(\Theta_{d,I}^*, f_{d,M}^*)$ that

$$R(d, \Theta_{d,I}^*, f_{d,M}^*) \leq R(d, \Theta_{d,I}^*, f) = \mathbb{E}[Y^2] = \mathbb{E}[F(\mathbf{X})^2] + 2\mathbb{E}[F(\mathbf{X})\mathbb{E}[\varepsilon|\mathbf{X}]] + \mathbb{E}[\varepsilon^2] \leq C^2 + \sigma^2. \tag{4.13}$$

Plugging (4.12) and (4.13) into (4.11), we have

$$|U_4| \leq 2dpKC\eta \sqrt{C^2 + \sigma^2}. \tag{4.14}$$

Finally, applying (4.7) again yields

$$\begin{aligned}
|U_5| &\leq 2 \int \mathbb{E}[|f_{d,M}^*(\Theta_{d,I}^* \mathbf{X}) - f(\Theta_{d,I}^* \mathbf{X})| |f(\Theta_{d,I}^* \mathbf{X}) - f(\Theta \mathbf{X})|] d\rho_{d,I,\eta}^1 \otimes \rho_{d,M,\gamma}^2(\Theta, f) \\
&\leq 2\sqrt{dp}KC \mathbb{E} \left[\int |f_{d,M}^*(\Theta_{d,I}^* \mathbf{X}) - f(\Theta_{d,I}^* \mathbf{X})| \left(\sum_{i=1}^d \|\vartheta_{d,I,i}^* - \vartheta_i\|_2^2 \right)^{\frac{1}{2}} d\rho_{d,I,\eta}^1 \otimes \rho_{d,M,\gamma}^2((\vartheta_1, \dots, \vartheta_d)^\top, f) \right] \\
&= 2\sqrt{dp}KC \int |f_{d,M}^*(\Theta_{d,I}^* \mathbf{X}(\omega)) - f(\Theta_{d,I}^* \mathbf{X}(\omega))| \\
&\quad \cdot \left(\sum_{i=1}^d \|\vartheta_{d,I,i}^* - \vartheta_i\|_2^2 \right)^{1/2} d\mathbb{P} \otimes \rho_{d,I,\eta}^1 \otimes \rho_{d,M,\gamma}^2(\omega, (\vartheta_1, \dots, \vartheta_d)^\top, f) \\
&\leq 2\sqrt{dp}KC \sqrt{\int (f_{d,M}^*(\Theta_{d,I}^* \mathbf{X}(\omega)) - f(\Theta_{d,I}^* \mathbf{X}(\omega)))^2 d\mathbb{P} \otimes \rho_{d,I,\eta}^1 \otimes \rho_{d,M,\gamma}^2(\omega, (\vartheta_1, \dots, \vartheta_d)^\top, f)} \\
&\quad \cdot \sqrt{\int \sum_{i=1}^d \|\vartheta_{d,I,i}^* - \vartheta_i\|_2^2 d\mathbb{P} \otimes \rho_{d,I,\eta}^1 \otimes \rho_{d,M,\gamma}^2(\omega, (\vartheta_1, \dots, \vartheta_d)^\top, f)} \tag{4.15} \\
&\leq 2\sqrt{dp}KC \sqrt{\int \mathbb{E}[(f(\Theta_{d,I}^* \mathbf{X}) - f(\Theta \mathbf{X}))^2] d\rho_{d,M,\gamma}^2(f)} \sqrt{\int \sum_{i=1}^d \|\vartheta_{d,I,i}^* - \vartheta_i\|_2^2 \rho_{d,I,\eta}^1((\vartheta_1, \dots, \vartheta_d)^\top)} \\
&\leq 2dpKC\eta\gamma, \tag{4.16}
\end{aligned}$$

where (4.15) follows from the Cauchy-Schwarz inequality for integration with respect to the product measure $\mathbb{P} \otimes \rho_{d,I,\eta}^1 \otimes \rho_{d,M,\gamma}^2$.

Choosing $\eta = (dpn)^{-1}, \gamma = n^{-1}$ when summarising (4.5), (4.6), (4.8), (4.10), (4.14) and (4.16) we have

$$\begin{aligned}
\int \mathcal{E}(c, \Theta, f) d\rho(c, \Theta, f) &\leq \mathcal{E}(d, \Theta_{d,I}^*, f_{d,M}^*) + \gamma^2 + (dpKC\eta)^2 + 2dpKC\eta\sqrt{C^2 + \sigma^2} + 2dpKC\eta\gamma \\
&\leq \mathcal{E}(d, \Theta_{d,I}^*, f_{d,M}^*) + \frac{\Xi_1}{n},
\end{aligned}$$

where Ξ_1 is a constant only depending on C, K and σ . With these choices for η and γ , we can use the assumption that $n \geq (10C^{-1}) \vee (30\sqrt{2}ep^{-2})$ together with $C \geq 1$ (by Assumption 1) to bound

$$\begin{aligned}
T_1 + T_2 &= \|I\| \log(ep) + (\|I\| + M + 1) \log(10) + \|I\| \log(3\sqrt{2}dpn) + 4^d N^d 2^{dM+1} \log(n(C+1)) \\
&\leq \|I\| \log(ep) + (\|I\| + M + 1) \log(10) + \|I\| \log(3\sqrt{2}dpn) + 4^d N^d 2^{dM+1} \log(n(C+1)) \\
&\leq 4\|I\| \log(pn) + 16 \cdot 4^d N^d 2^{dM} \log(Cn).
\end{aligned}$$

Choosing

$$\lambda = \frac{n}{Q + 2((2C+1)^2 + 4\sigma^2)},$$

we have

$$\frac{1}{1 - \frac{\lambda((2C+1)^2 + 4\sigma^2)}{n - Q\lambda}} = 2 \quad \text{as well as} \quad 1 + \frac{\lambda((2C+1)^2 + 4\sigma^2)}{n - Q\lambda} = \frac{3}{2}$$

and there exists a constant Ξ_2 depending only on C, K, Γ and σ such that $\frac{2}{\lambda} \leq \frac{\Xi_2}{n}$. Summarizing the above, we arrive at

$$\mathcal{E}(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda) \leq 3\mathcal{E}(d, \Theta_{d,I}^*, f_{d,M}^*) + \frac{\Xi}{n} \left(\|I\| \log(pn) + 4^d N^d 2^{dM} \log(Cn) + \log\left(\frac{2}{\delta}\right) \right) \tag{4.17}$$

with a probability of at least $1 - \delta$, where Ξ is a constant only depending on C, K, Γ and σ . Note that the upper bound in (4.17) is deterministic. Choosing a triplet (d, I, M) such that this upper bound is

minimised (which is always possible, since there are only finitely many choices for (d, I, M)), we have shown that

$$\mathcal{E}(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda) \leq \inf_{d, I, M} \left(3\mathcal{E}(d, \Theta_{d, I}^*, f_{d, M}^*) + \frac{\Xi}{n} \left(\|I\| \log(pn) + 4^d N^d 2^{dM} \log(Cn) + \log\left(\frac{2}{\delta}\right) \right) \right),$$

with a probability of at least $1 - \delta$. This completes the proof of Theorem 2. \square

4.2 Proof of Corollary 5

Plugging in d^* and I^* in the infimum in Theorem 2, we obtain that

$$\begin{aligned} \mathcal{E}(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda) &\leq \inf_{0 \leq M \leq n} \left(3\mathcal{E}(d^*, \Theta_{d^*, I^*}^*, f_{d^*, M}^*) + \frac{\Xi}{n} \left(\|I\| \log(pn) + 4^{d^*} N^{d^*} 2^{d^* M} \log(Cn) + \log\left(\frac{2}{\delta}\right) \right) \right), \\ &\leq \inf_{\substack{0 \leq M \leq n, \\ f \in \mathcal{F}_{d^*, M}(C)}} \left(3\mathcal{E}(d^*, \Theta^*, f) + \frac{\Xi}{n} \left(\|\Theta^*\|_0 \log(pn) + 4^{d^*} N^{d^*} 2^{d^* M} \log(Cn) + \log\left(\frac{2}{\delta}\right) \right) \right), \end{aligned} \quad (4.18)$$

with a probability of at least $1 - \delta$. The rest of the proof consists of choosing M to balance the terms on the right hand side of (4.18) by using an approximation of f^* , namely

$$f_M = \sum_{(l, k) \in \mathcal{Z}_M^{d^*}} \langle f^*, \Psi_{l, k} \rangle \Psi_{l, k}$$

and then determining the projection level M . To do this, we have to verify that f_M is a valid choice for f in the sense that $f_M \in \mathcal{F}_{d^*, M}(C)$. Indeed, the Cauchy-Schwarz inequality ensures that

$$\begin{aligned} L^{d^*} \sum_{(l, k) \in \mathcal{Z}_M^{d^*}} 2^{l(d/2+1)} |\langle f^*, \Psi_{l, k} \rangle| &\leq L^{d^*} \left(\sum_{(l, k) \in \mathcal{Z}_M^{d^*}} 2^{2l(1-\alpha+d^*/2)} \right)^{\frac{1}{2}} \left(\sum_{(l, k) \in \mathcal{Z}_M^{d^*}} 2^{2l\alpha} |\langle f^*, \Psi_{l, k} \rangle|^2 \right)^{\frac{1}{2}} \\ &\leq L^{d^*} 2N^{d^*} / 2^{8^{d^*/2}} \left(\sum_{(l, k) \in \mathcal{Z}_{\infty, \infty}^{d^*}} 2^{2l\alpha} |\langle f^*, \Psi_{l, k} \rangle|^2 \right)^{\frac{1}{2}} \\ &\leq C. \end{aligned}$$

Using Assumption 3, we see that f_M admits an excess risk of

$$\begin{aligned} \mathcal{E}(d^*, \Theta^*, f_M) &= \mathbb{E}[(f_M(\Theta^* \mathbf{X}) - f^*(\Theta^* \mathbf{X}))^2] \\ &= \int_{[-B_1, B_1]^{d^*}} \varrho(x) (f_M(x) - f^*(x))^2 \mathfrak{K}^{d^*}(dx) \\ &\leq B_2 \int_{[-B_1, B_1]^{d^*}} (f_M(x) - f^*(x))^2 \mathfrak{K}^{d^*}(dx) \\ &\leq B_2 \sum_{(l, k) \in \mathcal{Z}_{\infty, N}^{d^*} \setminus \mathcal{Z}_M^{d^*}} |\langle f^*, \Psi_{l, k} \rangle|^2 \\ &\leq B_2 2^{-2\alpha M} \sum_{(l, k) \in \mathcal{Z}_{\infty, N}^{d^*} \setminus \mathcal{Z}_M^{d^*}} 2^{2l\alpha} |\langle f^*, \Psi_{l, k} \rangle|^2 \\ &\leq B_2 2^{-2\alpha M} (2N^{d^*} / 2^{8^{d^*/2}})^{-1} C L^{-d^*}. \end{aligned} \quad (4.19)$$

Applying (4.19) to (4.18), we see that there exists a constant Λ_1 only depending on $C, K, \Gamma, \sigma, N, B_1, B_2$ and d^* such that

$$\mathcal{E}(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda) \leq \Lambda_1 \inf_{0 \leq M \leq n} \left(2^{-2\alpha M} + 2^{d^* M} \frac{\log(Cn)}{n} + \frac{\|\Theta^*\|_0 \log(pn)}{n} + \frac{\log\left(\frac{2}{\delta}\right)}{n} \right)$$

with a probability of at least $1 - \delta$. To balance the order of the terms depending on M , we choose

$$M = \left\lceil \log \left(\frac{n}{\log(Cn)} \right) / ((2\alpha + d^*) \log(2)) \right\rceil$$

and obtain that for some constant Λ depending only on $C, K, \Gamma, \sigma, N, B_1, B_2$ and d^* , we have

$$\mathcal{E}(\widehat{d}_\lambda, \widehat{\Theta}_\lambda, \widehat{f}_\lambda) \leq \Lambda \left(\left(\frac{\log(Cn)}{n} \right)^{\frac{2\alpha}{2\alpha + d^*}} + \frac{\|\Theta^*\|_0 \log(pn)}{n} + \frac{\log\left(\frac{2}{\delta}\right)}{n} \right)$$

with a probability of at least $1 - \delta$. This completes the proof of Corollary 5. \square

4.3 Proofs of auxiliary lemmata

Proof of Lemma 9. We employ another auxiliary lemma:

Lemma 11. *It holds that*

$$\begin{aligned} \mathcal{K}(\rho_{d,I,M,\eta,\gamma}, \pi) &= \log(G(d, I, M)) + \mathcal{K}(\rho_{d,I,M,\eta,\gamma}, \pi_{d,I,M}), \quad \text{where} \\ G(d, I, M) &:= \frac{1}{721} (1 - 10^{-p})(1 - 10^{(1-p)d-1})(10 - 10^{-n}) 10^{\|I\|+M+1} |\mathcal{I}_{d,\|I\|}|. \end{aligned}$$

Now, we can combine

$$|\mathcal{I}_{d,\|I\|}| = |\{J \mid \emptyset \neq J = J_1 \times \cdots \times J_d, J_1, \dots, J_d \subseteq \{1, \dots, p\}\}| \leq \binom{dp}{\|I\|}$$

with the basic inequality $\binom{dp}{\|I\|} \leq \left(\frac{dpe}{\|I\|}\right)^{\|I\|}$ and the fact that $\|I\| \geq d$, to obtain

$$\begin{aligned} \mathcal{K}(\rho_{d,I,M,\eta,\gamma}, \pi) &= \log(G(d, I, M)) + \mathcal{K}(\rho_{d,I,M,\eta,\gamma}, \pi_{d,I,M}) \\ &\leq \log \left(10^{\|I\|+M+1} \binom{dp}{\|I\|} \right) + \mathcal{K}(\rho_{d,I,M,\eta,\gamma}, \pi_{d,I,M}) \\ &\leq \|I\| \log(ep) + (\|I\| + M + 1) \log(10) + \mathcal{K}(\rho_{d,I,M,\eta,\gamma}, \pi_{d,I,M}). \end{aligned} \quad \square$$

Proof of Lemma 10. We split the proof into two further auxiliary lemmata:

Lemma 12. *For $\rho_{d,I,\eta}^1$ from (4.3), we have*

$$\mathcal{K}(\rho_{d,I,\eta}^1, \mu_{d,I}) \leq \|I\| \log \left(\frac{3\sqrt{2}}{\eta} \right), \quad \forall \eta \in (0, 1].$$

Lemma 13. *For $\rho_{d,M,\gamma}^2$ from (4.3), we have*

$$\mathcal{K}(\rho_{d,M,\gamma}^2, \nu_{d,M}) \leq 2^{3d} N^d 2^{M+1} \log \left(\frac{C+1}{\gamma} \right), \quad \forall \gamma \in (0, 1].$$

The assertion then follows directly via

$$\begin{aligned} \mathcal{K}(\rho, \pi_{d,I,M}) &= \mathcal{K}(\kappa_d \otimes \rho_{d,I,\eta}^1 \otimes \rho_{d,M,\gamma}^2, \kappa_d \otimes \mu_{d,I} \otimes \nu_{d,M}) \\ &= \mathcal{K}(\kappa_d, \kappa_d) + \mathcal{K}(\rho_{d,I,\eta}^1, \mu_{d,I}) + \mathcal{K}(\rho_{d,M,\gamma}^2, \nu_{d,M}) \\ &\leq \|I\| \log \left(\frac{3\sqrt{2}}{\eta} \right) + 2^{3d} N^d 2^{M+1} \log \left(\frac{C+1}{\gamma} \right) =: T_2. \end{aligned} \quad \square$$

Proof of Lemma 11. To simplify the notation we write $\rho = \rho_{d,I,M,\eta,\gamma}$ and $\pi_{d,I,M} = \kappa_d \otimes \mu_{d,I} \otimes \nu_{d,M}$. We will show that

$$\frac{d\rho}{d\pi} = G(d, I, M) \frac{d\rho}{d\pi_{d,I,M}} \quad (4.20)$$

from which we can deduce

$$\mathcal{K}(\rho, \pi) = \int \log\left(\frac{d\rho}{d\pi}\right) d\rho = \log(G(d, I, M)) + \int \log\left(\frac{d\rho}{d\pi_{d,I,M}}\right) d\rho = \log(G(d, I, M)) + \mathcal{K}(\rho, \pi_{d,I,M}).$$

For (4.20), we need to show that

$$\rho(A) = \int_A G(d, I, M)^{-1} \frac{d\rho}{d\pi} d\pi_{d,I,M} \quad (4.21)$$

holds for all $A = A_1 \times A_2 \times A_3$ with $A_1 \in 2^{\{1, \dots, p\}}$, $A_2 \in \mathcal{B}_{\mathcal{S}}$ and $A_3 \in \mathcal{B}_{\mathcal{F}(C+1)}$. Observe that for the sets

$$\begin{aligned} \mathcal{S}_{d,\Leftrightarrow}(J) &:= \{\Theta = (\vartheta_1, \dots, \vartheta_d)^\top \in \mathcal{S}_d \mid (\vartheta_{i,j} \neq 0 \Leftrightarrow j \in J_i) \forall i \in \{1, \dots, d\}, j \in \{1, \dots, p\}\} \\ \mathcal{F}_{d,\widetilde{M},\neq}(C+1) &:= \left\{ f = \Phi_{d,\widetilde{M}}(\beta) \in \mathcal{F}_{d,\widetilde{M}}(C+1) \right. \\ &\quad \left. \mid \exists (l, (k_1, k_2)) \in \mathcal{Z}_{\widetilde{M}}^d : ((\|k_1\|_\infty = 2^l \widetilde{M}) \wedge (\beta_{l,k} \neq 0)) \wedge (k_2 \neq 0) \right\} \end{aligned}$$

with $J = J_1 \times \dots \times J_d \in \mathcal{I}_d$ and $\widetilde{M} \in \{0, \dots, n\}$, we have

$$\kappa_d(\{d\}) = \mu_{d,J}(\mathcal{S}_{d,\Leftrightarrow}(J)) = \nu_{d,\widetilde{M}}(\mathcal{F}_{d,\widetilde{M},\neq}(C+1)) = 1. \quad (4.22)$$

In particular, (4.22) holds for $J = I$ and $\widetilde{M} = M$. Since also $\rho(\{d\} \times \mathcal{S}_{d,\Leftrightarrow}(I) \times \mathcal{F}_{d,M,\neq}(C+1)) = 1$, no generality is lost in additionally assuming that

$$A_1 = \{d\}, \quad A_2 \subseteq \mathcal{S}_{d,\Leftrightarrow}(I) \quad \text{and} \quad A_3 \subseteq \mathcal{F}_{d,M,\neq}(C+1).$$

Now note that

$$\{c\} \cap \{d\} = \emptyset \forall c \neq d, \quad \mathcal{S}_{d,\Leftrightarrow}(J) \cap \mathcal{S}_{d,\Leftrightarrow}(I) = \emptyset \forall J \neq I \quad \text{and} \quad \mathcal{F}_{d,\widetilde{M},\neq}(C+1) \cap \mathcal{F}_{d,M,\neq}(C+1) = \emptyset \forall \widetilde{M} \neq M. \quad (4.23)$$

Combining (4.22) with (4.23), we see that

$$\int_{A_2} \frac{d\rho}{d\pi} d\mu_{d,J} = 0 \forall J \neq I \quad \text{and} \quad \int_{A_3} \frac{d\rho}{d\pi} d\nu_{d,M} = 0 \forall \widetilde{M} \neq M.$$

Therefore, repeated application of Fubini's theorem yields

$$\begin{aligned} \rho(A) &= \int_A \frac{d\rho}{d\pi} d\pi \\ &= \sum_{c=1}^p 10^{-c} \int_A \frac{d\rho}{d\pi} d\pi_c / \left(\frac{1}{9} (1 - 10^{-p}) \right) \\ &= \int_A \frac{d\rho}{d\pi} d\kappa_d \otimes \mu_d \otimes \nu_d / \left(\frac{10^d}{9} (1 - 10^{-p}) \right) \\ &= \int_{A_2 \times A_3} \frac{d\rho}{d\pi}(\Theta, f) d\mu_d \otimes \nu_d(\Theta, f) / \left(\frac{10^d}{9} (1 - 10^{-p}) \right) \\ &= G(d, I, M)^{-1} \int_{A_2 \times A_3} \frac{d\rho}{d\pi}(\Theta, f) d\mu_d \otimes \nu_d(\Theta, f) \\ &= G(d, I, M)^{-1} \int_A \frac{d\rho}{d\pi} d\pi_{d,I,M}. \end{aligned}$$

Thus, we have shown (4.21). \square

Proof of Lemma 12. We will show that

$$\mathcal{K}(\rho_{d,I,\eta}^{1,i}, \mu_{I_i}) \leq |I_i| \log \left(\frac{3\sqrt{2}}{\eta} \right), \forall \eta \in (0, 1], i = 1, \dots, d. \quad (4.24)$$

The assertion follows immediately via

$$\mathcal{K}(\rho, \mu_{d,I}) = \mathcal{K} \left(\bigotimes_{i=1}^d \rho_{d,I,\eta}^{1,i}, \bigotimes_{i=1}^d \mu_{I_i} \right) = \sum_{i=1}^d \mathcal{K}(\rho_{d,I,\eta}^{1,i}, \mu_{I_i}) \leq \sum_{i=1}^d |I_i| \log \left(\frac{3\sqrt{2}}{\eta} \right) = \|I\| \log \left(\frac{3\sqrt{2}}{\eta} \right),$$

where the first equality holds barring a slight breach of conventions for product measures. To show (4.24), fix $i \in \{1, \dots, d\}$ and for simplicity of the notation, set $\rho := \rho_{d,I,\eta}^{1,i}$ and $J = I_i$. Plugging the μ_J -density of ρ into the definition of the Kullback-Leibler divergence, we easily obtain

$$\mathcal{K}(\rho, \mu_J) = -\log \left(\int \mathbb{1}_{\{\|\vartheta - \vartheta_{d,I,i}^*\|_2 \leq \eta\}} \mu_J(d\vartheta) \right) = -\log(\tilde{\mu}(\{\vartheta \in \mathbb{R}^{|J|} \mid \|\vartheta - \tilde{\vartheta}^*\|_2 \leq \eta\})) \quad (4.25)$$

where $\tilde{\vartheta}^*$ is the projection of $\vartheta_{d,I,i}^*$ onto the coordinates whose indices are elements of J and $\tilde{\mu}_J$ denotes the uniform distribution on the unit sphere in $\mathbb{R}^{|J|}$.

We want to show a lower bound for $\tilde{\mu}(\{\vartheta \in \mathbb{R}^{|J|} \mid \|\vartheta - \tilde{\vartheta}^*\|_2 \leq \eta\})$, which is the proportion of the surface of the unit sphere in $\mathbb{R}^{|J|}$ which is covered by the η -ball around $\tilde{\vartheta}^*$ to the surface of the entire unit sphere. By rotational symmetry of the uniform distribution on this sphere, no generality is lost by assuming $\tilde{\vartheta}^* = (1, 0, \dots, 0)^\top \in \mathbb{R}^{|J|}$.

For any $\tilde{\vartheta} = (\tilde{\vartheta}_1, \dots, \tilde{\vartheta}_{|J|})^\top \neq 0$ with $\|\tilde{\vartheta}\|_2 \leq 1$, the η -ball around $\tilde{\vartheta}^*$ covers the same part of the surface of the unit sphere as the $\tilde{\eta}$ -ball around $\tilde{\vartheta}$, where $\tilde{\eta} = \sqrt{\eta^2 \|\tilde{\vartheta}\|_2 - 2\|\tilde{\vartheta}\|_2 + \|\tilde{\vartheta}\|_2^2 + 1}$. Using this dependence between $\tilde{\vartheta}, \eta$ and $\tilde{\eta}$, it is easily checked that $\|\tilde{\vartheta}\|_2 = \sqrt{1 - \tilde{\eta}^2}$ is solved for $0 < \tilde{\eta} = \sqrt{\eta^2 - \frac{1}{4}\eta^4} < 1$. Henceforth, fix this $\tilde{\eta}$. Suppose that $N_{\tilde{\eta}}$ is the smallest number of $\tilde{\eta}$ -balls with centers in the unit ball that suffice to cover the entire unit ball. Denote their centers by $t_1, \dots, t_{N_{\tilde{\eta}}}$. In particular, these balls cover the entire unit sphere and therefore, at least one of them covers at least $N_{\tilde{\eta}}^{-1}$ of the surface of the unit sphere. This can only be the case for $t_j \neq 0$, because otherwise $\tilde{\eta} < 1$ implies $\{\vartheta \in \mathbb{R}^{|J|} \mid \|\vartheta - t_j\|_2 \leq \tilde{\eta}\} \cap \{\vartheta \in \mathbb{R}^{|J|} \mid \|\vartheta\|_2 = 1\} = \emptyset$. If we now change the length of such a $\vartheta := t_j \neq 0$ (without changing its orientation and without changing $\tilde{\eta}$), the coverage of the unit sphere provided by the corresponding $\tilde{\eta}$ -ball also changes. In particular, we will show that if $\|\vartheta\|_2 \neq \sqrt{1 - \tilde{\eta}^2}$, then decreasing (or increasing) $\|\vartheta\|_2$ towards $\sqrt{1 - \tilde{\eta}^2}$, enlarges the coverage of the corresponding ball on the unit sphere. Thus, the proportional coverage of the $\tilde{\eta}$ -ball around $\|\vartheta\|_2^{-1} \sqrt{1 - \tilde{\eta}^2} \vartheta$ (which has, as we showed above, the same proportional coverage of the unit sphere as the η -ball around $\tilde{\vartheta}^*$ that we are actually trying to control) is bounded from below by the proportional coverage of the $\tilde{\eta}$ -ball around $\tilde{\vartheta}$, which in turn is bounded from below by $N_{\tilde{\eta}}^{-1}$. Using the fact that $N_{\tilde{\eta}} \leq (\frac{3}{\tilde{\eta}})^{|J|}$ combined with $\tilde{\eta} \geq \eta/\sqrt{2}$, we have

$$N_{\tilde{\eta}}^{-1} \geq \left(\frac{3}{\tilde{\eta}} \right)^{-|J|} \geq \left(\frac{3\sqrt{2}}{\eta} \right)^{-|J|}$$

and therefore (4.24) follows from (4.25) with

$$\mathcal{K}(\rho, \mu_J) = -\log(\tilde{\mu}(\{\vartheta \in \mathbb{R}^{|J|} \mid \|\vartheta - \tilde{\vartheta}^*\|_2 \leq \eta\})) \leq -\log(N_{\tilde{\eta}}^{-1}) \leq |J| \log \left(\frac{3\sqrt{2}}{\eta} \right).$$

It remains to show that changing the length of $\vartheta \neq 0$ towards $\sqrt{1 - \tilde{\eta}^2}$ increases the proportional coverage of the corresponding $\tilde{\eta}$ -ball. By rotational symmetry, we can assume $\vartheta = (\vartheta_1, 0, \dots, 0)^\top \in \mathbb{R}^{|J|}$ with some $0 < \vartheta_1 \leq 1$ at no loss of generality. Now, it is sufficient to show that

$$\{y \in \mathbb{R}^{|J|} \mid (\|y - \vartheta\|_2 \leq \tilde{\eta}) \wedge (\|y\|_2 = 1)\} \subseteq \{y \in \mathbb{R}^{|J|} \mid (\|y - \tilde{\vartheta}\|_2 \leq \tilde{\eta}) \wedge (\|y\|_2 = 1)\}$$

where $\tilde{\vartheta} = (\sqrt{1 - \tilde{\eta}^2}, 0, \dots, 0)^\top \in \mathbb{R}^{|\mathcal{J}|}$. In this setting, and as $\eta, \tilde{\eta} > 0$, the relationship

$$\tilde{\eta}^2 = \eta^2 \|\tilde{\vartheta}\|_2 - 2\|\tilde{\vartheta}\|_2 + \|\tilde{\vartheta}\|_2^2 + 1$$

is equivalent to

$$\eta^2 = \frac{2\tilde{\vartheta}_1 - \tilde{\vartheta}_1^2 - 1 + \tilde{\eta}^2}{\tilde{\vartheta}_1}.$$

Using elementary calculus techniques together with the fact that $\tilde{\eta} < 1$, it is easy to see that

$$\frac{2\tilde{\vartheta}_1 - \tilde{\vartheta}_1^2 - 1 + \tilde{\eta}^2}{\tilde{\vartheta}_1} \leq 2(1 - \sqrt{1 - \tilde{\eta}^2}).$$

Combining this with the relationship between η and $\tilde{\eta}$, we obtain

$$\begin{aligned} \{y \in \mathbb{R}^{|\mathcal{J}|} \mid (\|y - \vartheta\|_2 \leq \tilde{\eta}) \wedge (\|y\|_2 = 1)\} &= \{y \in \mathbb{R}^{|\mathcal{J}|} \mid (\|y - \tilde{\vartheta}^*\|_2^2 \leq \frac{2\tilde{\vartheta}_1 - \tilde{\vartheta}_1^2 - 1 + \tilde{\eta}^2}{\tilde{\vartheta}_1}) \wedge (\|y\|_2 = 1)\} \\ &\subseteq \{y \in \mathbb{R}^{|\mathcal{J}|} \mid (\|y - \tilde{\vartheta}^*\|_2^2 \leq 2(1 - \sqrt{1 - \tilde{\eta}^2})) \wedge (\|y\|_2 = 1)\} \\ &= \{y \in \mathbb{R}^{|\mathcal{J}|} \mid (\|y - \tilde{\vartheta}^*\|_2^2 \leq \eta^2) \wedge (\|y\|_2 = 1)\} \\ &= \{y \in \mathbb{R}^{|\mathcal{J}|} \mid (\|y - \tilde{\vartheta}\|_2 \leq \tilde{\eta}) \wedge (\|y\|_2 = 1)\}. \quad \square \end{aligned}$$

Proof of Lemma 13. To simplify the notation, we write $\tilde{f}^* = f_{d,M}^*$ and $\tilde{\beta}^* = \beta_{d,M}^*$. We will show that

$$\int \mathbb{1}_{\{\|f - \tilde{f}^*\|_\psi \leq \gamma\}} d\nu_{d,M}(f) = \left(\frac{C+1}{\gamma}\right)^{-|\mathcal{Z}_M^d|}. \quad (4.26)$$

The assertion follows directly with

$$\mathcal{K}(\rho_{d,M,\gamma}^2, \nu_{d,M}) = -\log \left(\int \mathbb{1}_{\{\|f - f^*\|_\psi \leq \gamma\}} d\nu_{d,M}(f) \right) = |\mathcal{Z}_M^d| \log \left(\frac{C+1}{\gamma} \right) \leq 4^d N^d 2^{dM+1} \log \left(\frac{C+1}{\gamma} \right).$$

We now show (4.26) using the definition of $\nu_{d,M}$. If we let $\mathfrak{X}^{\mathcal{Z}_M^d}$ denote the Lebesgue measure on $\mathcal{B}_{d,M}(C+1)$, we obtain

$$\begin{aligned} \int \mathbb{1}_{\{\|f - \tilde{f}^*\|_\psi \leq \gamma\}} d\nu_{d,M}(f) &= \int \mathbb{1}_{\{\|\Phi_{d,M}(\beta) - \tilde{f}^*\|_\psi \leq \gamma\}} d\tilde{\nu}_{d,M}(\beta) \\ &= \int \mathbb{1}_{\{\|\beta - \tilde{\beta}^*\|_{\mathcal{B}} \leq \gamma\}} d\tilde{\nu}_{d,M}(\beta) \\ &= \frac{\int \mathbb{1}_{\{\|\beta\|_{\mathcal{B}} \leq C+1\}} \mathbb{1}_{\{\|\beta - \tilde{\beta}^*\|_{\mathcal{B}} \leq \gamma\}} d\mathfrak{X}^{\mathcal{Z}_M^d}(\beta)}{\int \mathbb{1}_{\{\|\beta - \tilde{\beta}^*\|_{\mathcal{B}} \leq C+1\}} d\mathfrak{X}^{\mathcal{Z}_M^d}(\beta)} \\ &= \frac{\int \mathbb{1}_{\{\|\beta - \tilde{\beta}^*\|_{\mathcal{B}} \leq \gamma\}} d\mathfrak{X}^{\mathcal{Z}_M^d}(\beta)}{\int \mathbb{1}_{\{\|\beta - \tilde{\beta}^*\|_{\mathcal{B}} \leq C+1\}} d\mathfrak{X}^{\mathcal{Z}_M^d}(\beta)} \\ &= \left(\frac{\gamma}{C+1}\right)^{|\mathcal{Z}_M^d|} \frac{\int \mathbb{1}_{\{\|\beta - \tilde{\beta}^*\|_{\mathcal{B}} \leq 1\}} d\mathfrak{X}^{\mathcal{Z}_M^d}(\beta)}{\int \mathbb{1}_{\{\|\beta - \tilde{\beta}^*\|_{\mathcal{B}} \leq 1\}} d\mathfrak{X}^{\mathcal{Z}_M^d}(\beta)} \\ &= \left(\frac{C+1}{\gamma}\right)^{-|\mathcal{Z}_M^d|}, \end{aligned}$$

where we have used that

$$\|\beta\|_{\mathcal{B}} \leq \|\tilde{\beta}^*\|_{\mathcal{B}} + \|\beta - \tilde{\beta}^*\|_{\mathcal{B}} \leq C + \gamma \leq C + 1$$

on $\{\|\beta - \tilde{\beta}^*\|_{\mathcal{B}} \leq \gamma\}$ in the fourth equality. This implies that if the first indicator in the integral in the numerator is 1, so is the second. \square

References

- Alquier, P. (2021). User-friendly introduction to PAC-Bayes bounds. *arXiv preprint arXiv:2110.11216*.
- Alquier, P. & Biau, G. (2013). Sparse single-index model. *Journal of Machine Learning Research*, 14, 243–280.
- Catoni, O. (2004). *Statistical learning theory and stochastic optimization*. Springer.
- Catoni, O. (2007). *PAC-Bayesian supervised classification: The thermodynamics of statistical learning*, volume 56 of *Lecture Notes-Monograph Series*. Institute of Mathematical Statistics.
- Dalalyan, A. S., Juditsky, A., & Spokoiny, V. (2008). A new algorithm for estimating the effective dimension-reduction subspace. *Journal of Machine Learning Research*, 9, 1648–1678.
- Daubechies, I. (1992). *Ten lectures on wavelets*, volume 61 of *CBMS-NSF Regional Conference Series in Applied Mathematics*. Society for Industrial and Applied Mathematics.
- Giné, E. & Nickl, R. (2016). *Mathematical foundations of infinite-dimensional statistical models*. Cambridge University Press.
- Guedj, B. (2019). A primer on PAC-Bayesian learning. *arXiv preprint arXiv:1901.05353*.
- Hristache, M., Juditsky, A., Polzehl, J., & Spokoiny, V. (2001). Structure adaptive approach for dimension reduction. *The Annals of Statistics*, 29(6), 1537–1566.
- Klock, T., Lanteri, A., & Vigogna, S. (2021). Estimating multi-index models with response-conditional least squares. *Electronic Journal of Statistics*, 15(1), 589–629.
- Li, K.-C. (1991). Sliced inverse regression for dimension reduction. *Journal of the American Statistical Association*, 86(414), 316–327.
- Xia, Y. (2007). A constructive approach to the estimation of dimension reduction directions. *The Annals of Statistics*, 35(6), 2654–2690.
- Xia, Y. (2008). A multiple-index model and dimension reduction. *Journal of the American Statistical Association*, 103(484), 1631–1640.
- Xia, Y., Tong, H., Li, W. K., & Zhu, L.-X. (2002). An adaptive estimation of dimension reduction space. *Journal of the Royal Statistical Society. Series B. Statistical Methodology*, 64(3), 363–410.
- Zhu, L., Miao, B., & Peng, H. (2006). On sliced inverse regression with high-dimensional covariates. *Journal of the American Statistical Association*, 101(474), 630–643.