

Dedekind sums and class numbers of imaginary abelian number fields

Stéphane R. LOUBOUTIN
 Aix Marseille Université, CNRS, Centrale Marseille, I2M,
 Marseille, France
 stephane.louboutin@univ-amu.fr

March 27, 2023

Abstract

As a consequence of their work, Bruce C. Berndt and Ronald J. Evans in 1977 and Larry Joel Goldstein and Michael Razar in 1976 obtained a formula for the square of the class number of an imaginary quadratic number field in terms of Dedekind sums. Here we focus only on this formula and give a new short and simple proof of it. We finally express the relative class numbers of imaginary abelian number fields in terms of Dedekind sums.

For $c, d \in \mathbb{Z}$ and $d > 1$ such that $\gcd(c, d) = 1$, the *Dedekind sum* is defined by

$$s(c, d) := \frac{1}{4d} \sum_{n=1}^{|d|-1} \cot\left(\frac{\pi n}{d}\right) \cot\left(\frac{\pi nc}{d}\right) \quad (1)$$

(see [Apo, Chapter 3, Exercise 11] or [RG, (26)]). It depends only on c modulo d .

Let $3 < p \equiv 3 \pmod{4}$ be a prime rational integer. The class number formula applied to the imaginary quadratic field $K = \mathbb{Q}(\sqrt{-p})$ of discriminant $-p$ yields

$$h_K = \frac{\sqrt{p}}{\pi} L(1, \chi_K),$$

where $n \mapsto \chi_K(n) = \left(\frac{n}{p}\right) \in \{-1, 0, +1\}$ is the odd an primitive Dirichlet character modulo p associated with K given by the Legendre symbol and $L(s, \chi_K)$ is the Dirichlet L -function associated with the Dirichlet character χ_K . By [Lou94, Proposition 1] we have

$$L(1, \chi_K) = \frac{\pi}{2p} \sum_{n=1}^{p-1} \chi_K(n) \cot\left(\frac{\pi n}{p}\right).$$

Consequently, we have

$$\begin{aligned} h_K^2 = \frac{p}{\pi^2} L(1, \chi_K)^2 &= \frac{\pi}{4p} \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi_K(ab) \cot\left(\frac{\pi a}{p}\right) \cot\left(\frac{\pi b}{p}\right) \\ &= \frac{\pi}{4p} \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi_K(b) \cot\left(\frac{\pi a}{p}\right) \cot\left(\frac{\pi ab}{p}\right) = \sum_{b=1}^{p-1} \chi_K(b) s(c, p), \end{aligned}$$

⁰2010 Mathematics Subject Classification. Primary. 11F20, 11M20, 11R11, 11R20, 11R29.

Key words and phrases. Dedekind sums. Dirichlet character. L -function. Imaginary quadratic field. Class number.

by the bijective change of variables $b \mapsto ab$ modulo p and by noticing that $\chi_K(a)\chi_K(ab) = \chi_K(a)^2\chi_K(b) = \chi_K(b)$ for $\gcd(b, p) = 1$. We have finally proved a special case of [BE, Corollary 9] for prime discriminants. Since $S := \sum_{c=1}^{p-1} s(c, p) = \sum_{c=1}^{p-1} s(p-c, p) = -S$, we have $S = 0$ and

$$h_K^2 = \sum_{b=1}^{p-1} (1 + \chi_K(b))s(c, p) = 2 \sum_{\substack{b=1 \\ \chi_K(b)=+1}}^{p-1} s(c, p) = 2S(H_2, p),$$

with the notation of [Lou19, Proposition 2]. By [Lou19, Theorem 6] we recover the well known fact that the class number of the imaginary quadratic number field $\mathbb{Q}(\sqrt{-p})$ is odd for $3 < p \equiv 3 \pmod{4}$ a prime integer. We now give a short proof of [BE, Corollary 9] and [GR, p. 358]:

Theorem 1. *Let $-D < 0$ be the discriminant of an imaginary quadratic field K . Let $w_K \in \{2, 4, 6\}$ and h_K be the number of complex roots of unity contained in K and the class number of K . Let χ_K be the real, odd and primitive character modulo D associated with K . Then*

$$h_K = \frac{w_K^2}{4} \sum_{\substack{n=1 \\ \gcd(n, D)=1}}^{D-1} \chi_K(n)s(n, D).$$

Proof. Recalling the class number formula

$$h_K = \frac{w_K \sqrt{D}}{2\pi} L(1, \chi_K),$$

e.g. see [Was, p. 38], the result follows from the following Theorem 2. □

Theorem 2. *Let χ be an odd and primitive character modulo $f > 2$. Then*

$$|L(1, \chi)|^2 = \frac{\pi^2}{f} \sum_{\substack{n=1 \\ \gcd(n, f)=1}}^{f-1} \chi(n)s(n, f). \quad (2)$$

Proof. Using [Lou94, Proposition 1] we get

$$|L(1, \chi)|^2 = \frac{\pi^2}{4f} \sum_{\substack{a=1 \\ \gcd(a, f)=1}}^{f-1} \sum_{\substack{b=1 \\ \gcd(b, f)=1}}^{f-1} \overline{\chi(a)}\chi(b) \cot\left(\frac{\pi a}{f}\right) \cot\left(\frac{\pi b}{f}\right).$$

Making the bijective change of variables $b \mapsto ab \pmod{f}$ we get

$$|L(1, \chi)|^2 = \frac{\pi^2}{4f} \sum_{\substack{b=1 \\ \gcd(b, f)=1}}^{f-1} \chi(b) \sum_{\substack{a=1 \\ \gcd(a, f)=1}}^{f-1} \cot\left(\frac{\pi a}{f}\right) \cot\left(\frac{\pi ab}{f}\right).$$

Noticing that

$$\sum_{\substack{\delta|a \\ \delta|f}} \mu(\delta) = \begin{cases} 1 & \text{if } \gcd(a, f) = 1 \\ 0 & \text{if } \gcd(a, f) > 1 \end{cases}$$

we obtain

$$|L(1, \chi)|^2 = \frac{\pi^2}{4f} \sum_{\delta|f} \frac{\mu(\delta)}{\delta} \sum_{\substack{b=1 \\ \gcd(b, f)=1}}^{f-1} \chi_K(b)s(b, f/\delta).$$

To arrive at the desired result it suffices to prove that for any $d < f$ dividing f and for any primitive modulo f Dirichlet character χ we have $S(\chi, f, d) = 0$, where

$$S(\chi, f, d) := \sum_{b \in (\mathbb{Z}/f\mathbb{Z})^*} \chi(b)s(b, d).$$

To this end we consider the canonical surjective morphism $\phi : (\mathbb{Z}/f\mathbb{Z})^* \rightarrow (\mathbb{Z}/d\mathbb{Z})^*$ and notice that $s(b, d)$ depends only on b modulo d . We obtain

$$S(\chi, f, d) = \sum_{B \in (\mathbb{Z}/d\mathbb{Z})^*} \left(\sum_{\phi(b)=B} \chi(b) \right) s(B, d)$$

and we explain why each sum $\sum_{\phi(b)=B} \chi(b)$ is equal to 0. Indeed, for a given $B \in (\mathbb{Z}/d\mathbb{Z})^*$, we pick up any $b_0 \in (\mathbb{Z}/f\mathbb{Z})^*$ for which $\phi(b_0) = B$. Then $\phi(b) = B$ if and only if $b = b_0x$ with $x \in \ker \phi$. Hence, $\sum_{\phi(b)=B} \chi(b) = \chi(b_0) \sum_{x \in \ker \phi} \chi(x)$. Finally, since χ is primitive modulo f , there exists $x_0 \in \ker \phi$ with $\chi(x_0) \neq 1$ (otherwise χ would be induced by a character modulo d). Therefore the restriction of χ to the multiplicative group $\ker \phi$ is a non trivial character and it follows that $\sum_{x \in \ker \phi} \chi(x) = 0$ (make the change of variables $x \mapsto x_0x$ in this sum). \square

Since $\chi(n^*) = \overline{\chi(n)}$ and $s(n^*, f) = s(n, f)$, where n^* is the inverse in the multiplicative group $(\mathbb{Z}/f\mathbb{Z})^*$ of n with $\gcd(n, f) = 1$, the bijective change of variables $n \mapsto n^*$ shows that the right hand side of (2) being equal to its complex conjugate is indeed a real number. Notice also that the oddness of χ gives $\chi(f-n)s(f-n, f) = \chi(n)s(n, f)$ and

$$|L(1, \chi)|^2 = \frac{2\pi^2}{f} \sum_{\substack{n=1 \\ \gcd(n, f)=1}}^{f/2-1} \Re(\chi(n))s(n, f). \quad (3)$$

Let $Q_K \in \{1, 2\}$, w_K and h_K^- be the Hasse unit index, the number of complex roots of unity in K and class number of an imaginary abelian number field K . Let X_K^- be the set of primitive and odd Dirichlet characters associated with K . The class number formula and the conductor-discriminant formula (see [Was, Theorem 3.11]) give

$$h_K^- = Q_K w_K \prod_{\chi \in X_K^-} \frac{\sqrt{f_\chi}}{2\pi} L(1, \chi),$$

where f_χ denote the conductor of the primitive character χ . Now, instead of using [Was, Theorem 4.17] to compute h_K^- we can use this relative class number formula and (3). For example, if K is an imaginary cyclic quartic field of conductor f and χ is any one of the two conjugate odd and primitive modulo f quartic characters associated with K we have $Q_K = 1$ and

$$h_K^- = w_K \frac{f^2}{4\pi^2} |L(1, \chi)|^2 = \frac{w_K}{2} \sum_{\substack{n=1 \\ \gcd(n, f)=1}}^{f/2-1} \Re(\chi(n))s(n, f).$$

For example, for $K = \mathbb{Q}(\zeta_5)$ we have $Q_K = 1$ by [Was, Corollary 4.13], $w_K = 10$ and since $s(1, 5) = 1/5$, $s(2, 5) = s(3, 5) = 0$ and $s(4, 5) = -1/5$ we get

$$h_K^- = 5 \sum_{n=1}^2 \Re(\chi(n))s(n, 5) = 5 \left(1 \cdot s(1, 5) + 0 \cdot s(2, 5) \right) = 1,$$

where χ is the odd, primitive and quartic character modulo 5 given by $\chi(1) = 1$, $\chi(2) = \zeta_4$, $\chi(3) = -\zeta_4$ and $\chi(4) = -1$ (2 being a generator of the multiplicative group $(\mathbb{Z}/5\mathbb{Z})^*$ we may chose

for χ the character defined by $\chi(2^k) = \zeta_4^k$.

In the same way, for K the imaginary quartic subfield of $\mathbb{Q}(\zeta_{13})$ we have $Q_K = 1$, $w_K = 2$ and since $s(1, 13) = 11/13$, $s(2, 13) = 4/13$, $s(3, 13) = 1/13$, $s(4, 13) = -1/13$ and $s(5, 13) = 0$ we get

$$h_K^- = \sum_{n=1}^5 \Re(\chi(n))s(n, 13) = 11/13 + 1/13 + 1/13 = 1 = 1,$$

where χ is the odd, primitive and quartic character modulo 13 such that $\chi(1) = 1$, $\chi(2) = \zeta_4$, $\chi(3) = 1$ and $\chi(4) = -1$ (2 being a generator of the multiplicative group $(\mathbb{Z}/13\mathbb{Z})^*$ we may chose for χ the character defined by $\chi(2^k) = \zeta_4^k$).

We came up with [BE] and [GR] while working on [LM21] and [LM2?], where the interested reader will find new results and conjectures on Dedekind sums.

References

- [Apo] M. T. Apostol. *Modular functions and Dirichlet series in number theory*. Graduate Texts in Mathematics **41**. Springer-Verlag, New York, 1976.
- [BE] Bruce C. Berndt and Ronald J. Evans. Dedekind sums and class numbers. *Monatsh. Math.* **84** (1977), 265–273.
- [GR] Larry Joel Goldstein and Michael Razar. A generalization of Dirichlet’s class number formula. *Duke Math. J.* **43** (1976), 349–358.
- [LM21] S. Louboutin and M. Munsch. Second moment of Dirichlet L -functions, character sums over subgroups and upper bounds on relative class numbers. *Quart. J. Math.* **72** (2021), 1379–1399.
- [LM2?] S. Louboutin and M. Munsch. Mean square values of L -functions over subgroups for non primitive characters, Dedekind sums and bounds on relative class numbers. *Canad. J. Math.*, to appear.
- [Lou94] S. Louboutin. Quelques formules exactes pour des moyennes de fonctions L de Dirichlet. *Canad. Math. Bull.* **36** (1993), 190–196. Addendum. *Canad. Math. Bull.* **37** (1994), p. 89.
- [Lou19] S. Louboutin. On the denominator of Dedekind sums. *Bull. Korean Math. Soc.* **56** (2019), 815–827.
- [RG] H. Rademacher and E. Grosswald. Dedekind sums. *The Carus Mathematical Monographs*, **16**. The Mathematical Association of America, Washington, D.C., 1972.
- [Was] L. C. Washington. *Introduction to Cyclotomic Fields*. Second Edition. Graduate Texts in Mathematics **83**. Springer-Verlag, New York, 1997.