Convergence analysis of an inexact Forward-Backward algorithm for problems involving weakly convex functions

Ewa Bednarczuk^{1, 2}, Giovanni Bruccola¹, Gabriele Scrivanti³, and The Hung Tran¹

¹Systems Research Institute, PAS, 01-447 Warsaw, Newelska 6, Poland ²Warsaw University of Technology, 00-662 Warsaw, Koszykowa 75, Poland ³Université Paris-Saclay, Inria, CentraleSupélec, CVN, 3 Rue Joliot Curie, 91190, Gif-Sur-Yvette, France.

Abstract

We investigate the convergence properties of exact and inexact forward-backward (FB) algorithms for the minimisation of the two functions f+g defined on a Hilbert space, where f is weakly convex and g is convex and has a Lipschitz-continuous gradient. The main condition ensuring convergence is the sharpness of the objective f+g around the solution set S. We show that the exact (FB) algorithm converges to a global solution provided that at a certain iteration the sequence is sufficiently close to the solution set. The inexact (FB) iterations converge to a ball around S with radius depending upon the inexactness parameter ε . As an application of the analysed algorithm, we consider a feasibility problem involving a sphere and a closed convex set.

KEYWORDS: weakly convex functions, sharpness condition, forward-backward algorithm, inexact forward-backward algorithm, ρ -criticality, proximal subgradient, proximal operator

Mathematical classification index: 90C30 Nonlinear programming - 90C26 Nonconvex programming, global optimization - 90C51 Interior-point methods - 65K10 Numerical optimization and variational techniques - 52A01 Axiomatic and generalized convexity

1 Scope of this Work

Given a Hilbert space \mathcal{X} , we consider a problem of the form

$$\underset{x \in \mathcal{X}}{\text{minimize}} \ f(x) + g(x) \tag{1}$$

where function $f: \mathcal{X} \to (-\infty, +\infty]$ is weakly convex with modulus $\rho \geq 0$ (ρ -weakly convex) and function $g: \mathcal{X} \to \mathbb{R}$ is convex and differentiable with a L_g -Lipschitz continuous gradient. For a function $h: \mathcal{X} \to (-\infty, +\infty]$, the notation dom h will indicate the set

$$dom h := \{x \in \mathcal{X} \mid h(x) < +\infty\}$$

and in the remainder of the paper we will say that function h is *proper* whenever dom $h \neq \emptyset$.

Our standing assumption is that the solution set $S := \operatorname{argmin}_{x \in \mathcal{X}}(f+g)(x)$ is nonempty and that the objective function (f+g) is sharp in the sense of Definition 7 below. Under these assumptions, we analyse the convergence properties of an inexact FB splitting algorithm, where for given inexactness parameters $\varepsilon_t \geq 0$ and step sizes $\alpha_t > 0$, starting from a point $x_0 \in \operatorname{dom}(f+g)$, for any $t \in \mathbb{N}$ the update reads as

$$x_{t+1} \in \varepsilon_t \text{-prox}_{\alpha_t f}(x_t - \alpha_t \nabla g(x_t)).$$

The concept of inexact proximal operator and its computational tractability lie at the core of any proximal method referring to fully convex case, where both f and g are convex,[3, 41, 49, 54, 57]. In our analysis we are using the concept of ε -proximal operator introduced in our paper [7], see also Definition 4 below. Its properties, expressed in terms of global proximal ε -subdifferentials, are investigated in Proposition 4 and Proposition 5 of our paper [7], where we also provide a comparison with other concepts of existing ε -proximal operator in the convex case.

Motivation Weakly convex problems appear, for example, in signal processing [35], image processing, see e.g [44], [6], [56], [42], and machine learning [48]. Moreover, in this paper, we show that it is possible to model a feasibility problem between a sphere and a closed convex set as Problem (1). In the literature, many algorithms are devoted to solve different types of weakly convex problems, see e.g. [24], [26], [25]. Most of these algorithms can be considered special cases of the unified scheme for solving weakly convex problems, recently proposed in [1]. The inexact Sum Rule for weakly convex functions (see [7, Theorem 2]) represents a useful tool for the convergence analysis of algorithms such as the inexact forward-backward algorithm described in this work.

1.1 Related Works

Convergence results for FB The Forward-Backward algorithm (FB) or Proximal gradient algorithm belongs to the class of splitting methods [28, 34, 45] whose aim is to minimise the sum of a smooth function and a non-smooth one. By taking a gradient step on the smooth function and the proximal step on the non-smooth one, it is only needed to access each function separately. This kind of algorithm has been well studied and understood in the case when all the functions are convex [28]. In fact, many variants of FB have been proposed recently. For examples, [22] developed variable metric version of FB algorithm to improve convergence of the algorithm. In [43], and [38], the authors propose an inertial scheme for FB based on the work of Polyak [47]. While in [17], followed by Tseng's Forward-Backward-Forward splitting, the authors design an inertial scheme of FBF algorithm which can be used to obtain inertial primal-dual algorithm.

One crucial requirement in the works above is the ability to compute proximal operators in a closed form. When no such expression is available for the non-smooth term, the computation of the proximal point needs to be addressed/approximated as an independent optimisation problem. If not cautiously treated, the introduced approximation might represent a hindrance to the convergence of the method. One can overcome this issue by using an inexact proximal computation, which allows us to estimate the proximal point of a function at a certain location with a specified level of accuracy [54].

When the hypothesis of convexity is relaxed, two main drawbacks arise. Firstly, the convergence to global minimiser cannot be easily guaranteed anymore. Secondly, the possible lack of convexity of the non-smooth function invalidates the fact that the corresponding proximal operator is single valued.

Convergence under the Kurdyka-Łojasiewicz inequality. In the general non-convex case, the seminal work [2] illustrates that the convergence to a local minimum can be ensured provided that the objective function satisfies the Kurdyka-Łojasiewicz (KL) property [32, 36, 37]. This result, that holds in finite dimension, has an extension to general Hilbert spaces, provided that the objective function is weakly convex and it satisfies the infinite-dimensional version of the Kurdyka-Łojasiewicz property [12]. The use of this property allows us to overcome the fact that in the general non-convex case it is not possible to ensure that the sequence generated by the algorithm satisfies the so-called Fejér-monotonicity property, which is at the core of the convergence proof in the convex case [5]. One can relate the KL property to other conditions concerning the function values and the set of the critical points. For examples, the Luo-Tseng error bound condition combined with some assumption on the separation of the function values at the critical points can lead to the KL property [33].

Convergence under sharpness condition. In [25] the authors discuss the local convergence of standard (projected) subgradient methods for the minimisation of sharp functions that are weakly-convex in a finite-dimensional setting. In [24] the authors propose a stationarity measure for proximal stochastic subgradient methods for an objective function that is expressed as the sum of a closed convex function and an expectation function that is assumed to be weakly convex. The algorithms analysed in [25, 24] perform the (sub)gradient step on the weakly convex term and the proximal step with respect to the convex term.

FB for weakly convex functions Our analysis encompasses the case $\varepsilon_t = 0$, where instead of the approximate (inexact) proximal operator ε_t -prox_{$\alpha_t f$}, exact proximity calculation $\operatorname{prox}_{\alpha_t f}$ is performed for any $t \in \mathbb{N}$. This is studied in [10], where the authors exploit a variable smoothing approach and infer a complexity bound in finite dimension showing convergence to the set of critical points, under the assumption that a closed-form proximal operator is available for the weakly convex function.

Proximal Subdifferential Focusing on the class of weakly convex functions allows us to adopt a specific form of the general Fréchet Subdifferential, which is known as *proximal subdifferential*. There exists a vast literature devoted to proximal subdifferentials, see *e.g.* in the finite dimensional case, the monograph by Rockafellar and Wets [50], in Hilbert spaces the work by Bernard and Thibault [9]. This concept will be the underlying tool in our developments: in particular, make use of the fact that weakly convex functions (among others) enjoy a so-called *globalisation property* [31, 52], which states that the proximal subgradient inequality holds globally (see Definition 1 and Definition 2). Finally, we highlight that for ρ -weakly convex functions the proximal subdifferential coincides with the Clarke subdifferential (see [31, Theorem 3.1])

Contribution. We prove the strong local convergence of the inexat FB iterates in a Hilbert space, in both the exact and the inexact case. Our detailed contribution is as

follows:

- We discuss the notions of ε -critical points and ε -proximal point, which are crucial in some applications when the exact computation of the proximal point is not available. For ρ -weakly convex functions, we relate the proximal operator to the proximal subdifferential, Lemma 2.
- We investigate the relations between sharpness and criticality for weakly convex functions. We determine the location/position of the $(\rho/2)$ -critical points and the solution set S for problem (1) (see Proposition 3).
- We provide conditions for the boundedness and the decrease of the sequence $\operatorname{dist}(x_t, S)$ for $(x_t)_{t\in\mathbb{N}}$ constructed by Algorithm 1 (see Theorem 2).
- We investigate the strong convergence of the sequence generated by Algorithm 1 (see Corollary 4 in the inexact case and Proposition 6 in the exact case).
- We propose a new mathematical model for the feasibility problem, (FP), of a sphere and a closed convex set involving weakly convex functions. We show that our forward-backward scheme for weakly convex functions can be applied to (FP), since all the assumptions required for the convergence to a global solution are satisfied.

Paper Outline. The paper is organised as follows: in section 2 we provide some preliminary facts regarding (global) proximal ε -subdifferentials (where ε is understood as an inexactness parameter) and calculus rules for the computation of the ε -subdifferential of a sum of functions, along with the concept of ε -criticality; in section 3 we show the relation between the proximal operator and the proximal subdifferential of ρ -weakly convex functions (Lemma 2); in section 4 we introduce the concept of μ -sharpness and we prove that for a function satisfying it there exists a neighbourhood of the set of its global minima which does not include critical points; in section 5 we present in Algorithm 1 the inexact FB algorithm and provide a result regarding the descending behaviour of the function value for a sequence generated by Algorithm 1; in section 6 we illustrate some convergence properties related to possibly varying inexactness parameters ε while in section 7 we prove the main convergence results in the case ε is fixed. To conclude, in section 8 we discuss the application of Algorithm 1 to the problem (FP).

2 Preliminaries

We start with the definition of ρ -weak convexity (also known as ρ -paraconvexity, see [31, 51] or ρ -semiconvexity, see [19]).

Definition 1 (ρ -weak convexity). Let \mathcal{X} be a Hilbert space. A function $h: \mathcal{X} \to (-\infty, +\infty]$ is said to be ρ -weakly convex if there exists a constant $\rho \geq 0$ such that for $\lambda \in [0, 1]$ the following inequality holds:

$$(\forall (x,y) \in \mathcal{X}^2) \qquad h(\lambda x + (1-\lambda)y) \leqslant \lambda h(x) + (1-\lambda)h(y) + \lambda (1-\lambda)\frac{\rho}{2}||x-y||^2$$

We refer to ρ as the modulus of weak convexity of the function h.

Weakly convex functions can be equivalently characterised in the following manner:

Proposition 1 ([19, Proposition 1.1.3]). Let \mathcal{X} be a Hilbert space. Function $h: \mathcal{X} \to (-\infty, +\infty]$ is ρ -weakly convex if and only if function $h(\cdot) + \frac{\rho}{2} \| \cdot \|^2$ is convex.

A more detailed discussion about equivalent formulations of weak convexity can be found in [7] and in the references therein.

Example 1. Consider the following function:

$$f: \mathbb{R} \longrightarrow \mathbb{R}$$

$$x \longmapsto |(x-a)(x-b)|$$
(2)

where $(a,b) \in \{(a,b) \in \mathbb{R}^2 \mid a < b\}$. Function f is ρ -weakly convex with $\rho = 2$. As a matter of fact, for every $x \in \mathbb{R}$

$$f(x) + x^{2} = \begin{cases} x^{2} - (a+b)x + ab + x^{2} & \text{if } x \leq a \text{ and } x \geqslant b \\ -x^{2} + (a+b)x - ab + x^{2} & \text{if } a \leq x \leq b \end{cases}$$

$$= \begin{cases} 2x^{2} - (a+b)x + ab & \text{if } x \leq a \text{ and } x \geqslant b \\ (a+b)x - ab & \text{if } a \leq x \leq b \end{cases}$$

$$= \max\{2x^{2} - (a+b)x + ab, (a+b)x - ab\}.$$

In other words, function $x \mapsto f(x) + x^2$ is expressed as the point-wise maximum of convex functions, hence it is itself convex. In conclusion, function f is ρ -weakly convex with $\rho = 2$ by virtue of Proposition 1.

Example 2. Let $Q \in \mathbb{R}^{n \times n}$ be a symmetric matrix and let $f : \mathbb{R}^n \to \mathbb{R}$ be the associated bilinear form, $f(x) = \langle x, Qx \rangle$. It can be shown that f is ρ -weakly convex, where $\rho := |\lambda_0|$, and λ_0 is the smallest negative eigenvalue of Q. Since Q is a real symmetric matrix, it can be decomposed as $Q = PDP^{\top}$, where D is a diagonal matrix of eigenvalues of Q and P is an orthogonal matrix consisting of eigenvectors of Q. Hence

$$(\forall x \in \mathcal{X}) \qquad f(x) = \langle x, Qx \rangle = \langle x, PDP^{\top}x \rangle = \langle P^{\top}x, DP^{\top}x \rangle.$$

If all the eigenvalues of Q are non-negative, then D is a positive semi-definite matrix and $f(x) \ge 0$ for all $x \in \mathbb{R}^n$, which is equivalent to the fact that f is a convex function. On the other hand, if there exists $\lambda_0 < 0$ then we can add a constant $a_0 \ge |\lambda_0|$ so that the diagonal matrix $(D + a_0 Id)$ only admits non-negative entries and we have the following equalities for every $x \in \mathcal{X}$

$$\tilde{f}(x) = \langle P^{\top}x, (D + a_0 Id) P^{\top}x \rangle = \langle x, PDP^{\top}x \rangle + \langle x, P(a_0 Id) P^{\top}x \rangle = f(x) + a_0 ||x||^2,$$

which shows that $f(\cdot) + a_0 \|\cdot\|^2$ is convex because $\tilde{f}(\cdot)$ is convex.

For a general function $h: \mathcal{X} \to (-\infty, +\infty]$, we define the global proximal ε -subdifferential as follows.

Definition 2 (Global proximal ε -subdifferential). Let $\varepsilon \geq 0$. Let \mathcal{X} be a Hilbert space. The global proximal ε -subdifferential of a function $h: \mathcal{X} \to (-\infty, +\infty]$ at $x_0 \in \text{dom } h$ for $C \geq 0$ is defined as follows:

$$\partial_C^{\varepsilon} h(x_0) = \left\{ v \in \mathcal{X} \mid h(x) - h(x_0) \geqslant \langle v, x - x_0 \rangle - C \|x - x_0\|^2 - \varepsilon \quad \forall x \in \mathcal{X} \right\}. \tag{3}$$

For $\varepsilon = 0$, we have

$$\partial_C h(x_0) = \{ v \in \mathcal{X} \mid h(x) \geqslant h(x_0) + \langle v, x - x_0 \rangle - C \|x - x_0\|^2, \ \forall x \in \mathcal{X} \}$$
 (4)

In view of (4), ∂_0 denotes the subdifferential in the sense of convex analysis. The elements of $\partial_C^{\varepsilon} h(x_0)$ are called proximal ε -subgradients.

The notation $\partial^{\varepsilon}h(x)$ is used when the constant in (3) is inessential, i.e. $v \in \partial^{\varepsilon}h(x)$ means that there exists $C \ge 0$ such that $v \in \partial_C^{\varepsilon}h(x)$.

By [31, Proposition 3.1], for a ρ -weakly convex function h, the global proximal subdifferential $\partial_{\rho/2}h(x_0)$ defined by (4) coincides with the set of local proximal subgradients which satisfy the proximal subgradient inequality (4) locally in a neighbourhood of x_0 . Moreover, when h is ρ -weakly convex, by [31, Theorem 3.1], the proximal subdifferential $\partial_{\rho/2}h(x_0)$ coincides with the Clarke subdifferential $\partial_c h(x_0)$ (see also [1]). An extensive study of local proximal subdifferentials in Hilbert spaces can be found in [21, Chapter 2] and in [50, 9].

For any set set-valued mapping $M:\mathcal{X} \rightrightarrows \mathcal{X}$, we will use the notation dom M to indicate the set

$$\operatorname{dom} M := \{ x \in \mathcal{X} \mid M(x) \neq \emptyset \}.$$

Proposition 2 ([7, Proposition 2]). Let \mathcal{X} be a Hilbert space. Let $h: \mathcal{X} \to (-\infty, +\infty]$ be a lower semi-continuous and ρ -weakly convex function with $\rho \geq 0$. Then

$$\operatorname{dom} \partial_{\rho/2} h \subset \operatorname{dom} h$$

and for every $\varepsilon > 0$

$$\operatorname{dom} \partial_{\rho/2}^{\varepsilon} h = \operatorname{dom} h$$

The sum rule for proximal ε -subdifferentials presented below takes into account the moduli of weak convexity of the involved functions.

Theorem 1 ([7, Sum Rule for ε -subdifferential, Theorem 2]). Let \mathcal{X} be a Hilbert space. For i = 0, 1, let function $h_i : \mathcal{X} \to (-\infty, +\infty]$ be proper lower semi-continuous and ρ_i -weakly convex on \mathcal{X} with $\rho_i \geq 0$. Then, for all $x \in \text{dom } h_0 \cap \text{dom } h_1$ and for all $\varepsilon_0, \varepsilon_1 \geq 0$ we have

$$\partial_{\rho_0/2}^{\varepsilon_0} h_0(x) + \partial_{\rho_1/2}^{\varepsilon_1} h_1(x) \subseteq \partial_{\rho/2}^{\varepsilon} (h_0 + h_1)(x)$$

for all $\varepsilon \geqslant \varepsilon_0 + \varepsilon_1$ and for all $\rho \geqslant \rho_0 + \rho_1$. The equality

$$\hat{\mathcal{C}}^{\varepsilon}_{(\rho_0+\rho_1)/2}(h_0+h_1)(x) = \bigcup_{\varepsilon_0,\varepsilon_1,\varepsilon=\varepsilon_0+\varepsilon_1} \hat{\mathcal{C}}^{\varepsilon_0}_{\rho_0/2}h_0(x) + \hat{\mathcal{C}}^{\varepsilon_1}_{\rho_1/2}h_1(x)$$

holds when dom $h_0 \cap \text{dom } h_1$ contains a point at which either $h_0 + \rho_0/2 \|\cdot\|^2$ or $h_1(\cdot) + \rho_1/2 \|\cdot\|^2$ is continuous.

To conclude, we report the following result, commonly known as *Descent Lemma*, that holds for sufficiently regular functions defined on a Hilbert space.

Lemma 1 ([5, Lemma 2.64]). Let \mathcal{X} be a Hilbert space, let U be a nonempty convex subset of \mathcal{X} . Let $h: U \to \mathbb{R}$ be a Fréchet differentiable function such that ∇h is L_h -Lipschitz continuous on U. Let x and y be in U. Then the following holds:

(i)
$$|h(x) - h(y) - \langle x - y | \nabla h(y) \rangle| \le \frac{L_h}{2} ||x - y||^2$$

(ii)
$$|\langle x - y | \nabla h(x) - \nabla h(y) \rangle| \le L_h ||x - y||^2$$

In this work, we will consider ρ -weakly convex functions which are proper lower-semicontinuous. For any $\rho \geq 0$, in analogy to the convex case, we will use the notation $\Gamma_{\rho}(\mathcal{X})$ to indicate the class of proper lower-semicontinuous ρ -weakly convex functions defined on \mathcal{X} . In particular, $\Gamma_0(\mathcal{X})$ denotes the class of proper lower-semicontinuous convex functions defined on \mathcal{X} .

3 Inexact Proximal Operators

In practical context it is not always possible to rely on a closed-form expression for the proximity operator of a function and the computation of the proximal map needs to be addressed as an independent optimisation problem. Examples of such functions are the non-convex ℓ_p -seminorms (i.e. $p \in (0,1)$) or the convex ℓ_p -norms (i.e. $p \geq 1$) – unless p takes some specific values [20]. Another example is given by the combination of a sparsity-promoting functions with a non-orthogonal linear operator, as in the case of the popular discrete Total Variation functional [53] (and its non-convex modifications), which has been extensively used in the context of image and signal processing. In these cases, at each point, the proximal map is defined up to a certain degree of accuracy and in the framework of proximal splitting methods, it is important to carry out a convergence analysis that takes this fact into account. In order to do so, we consider the concept of ε -solution for an optimisation problem and the related notion of ε -proximal point. We illustrate its interaction with the proximal ε -subdifferential from Definition 2.

Definition 3 (ε -solution). Let \mathcal{X} be a Hilbert space. Let $h: \mathcal{X} \to (-\infty, +\infty]$ be a proper function that is bounded from below. Then, for any $\varepsilon > 0$, the element x_{ε} is said to be an ε -solution to the minimisation problem

$$\underset{x \in \mathcal{X}}{\text{minimize}} \ h(x)$$

if the following condition is satisfied:

$$(\forall x \in \mathcal{X}) \qquad \qquad h(x_{\varepsilon}) \leqslant h(x) + \varepsilon.$$

Definition 4 (ε -proximal point). Let \mathcal{X} be a Hilbert space. Let function $h: \mathcal{X} \to (-\infty, +\infty]$ be proper and bounded from below and let $x_0 \in \mathcal{X}$. Any ε -solution to the proximal minimisation problem

minimize
$$h(x) + \frac{1}{2} ||x - x_0||^2$$
, (5)

for $\varepsilon > 0$ is said to be a ε -proximal point for h at x_0 . The set of all ε -proximal point of h at x_0 is denoted as

$$\varepsilon$$
-prox_h $(x_0) := \{x \in \mathcal{X} \mid x \text{ is a } \varepsilon$ -solution of (5)}

The above definition of inexact proximal point is rather general and coincides, in the convex case, with the definition in [57, Equation (2.15)]. In the convex case, the convergence analysis of an inexact Forward Backward algorithm with respect to [57, Equation (2.15)] is proposed in [57, Appendix A].

To proceed with our analysis we consider the following notion of ε -critical point:

Definition 5 (ε -critical point). Let \mathcal{X} be a Hilbert space and $\varepsilon \geq 0$. Let $h: \mathcal{X} \to (-\infty, +\infty]$ be a proper function. A point $x \in \mathcal{X}$ is said to be a ε -critical point for h if $0 \in \partial^{\varepsilon} h(x)$. The set of ε -critical points is identified as

$$\varepsilon$$
- crit $h := \{x \in \mathcal{X} \mid 0 \in \partial^{\varepsilon} h(x)\}$

that is, if $x \in \varepsilon$ -crit h, there exists $a \ge 0$ such that $0 \in \partial_a^\varepsilon h(x)$. For a fixed $a \ge 0$, we say that x is a a- ε -critical point and we write

$$x \in \varepsilon$$
-crit_a h .

In order to study the inexact proximal operator of a function h, we apply Theorem 1 to express the subdifferential of the function in (5) as the sum of the subdifferential of h and of the quadratic term. Then, an explicit form for the ε -subdifferentials of quadratic functions allows us to express the inexact proximal operator in terms of $\partial_{\rho/2}^{\varepsilon}h$. This plays an important role in section 5, where we investigate the convergence of the inexact scheme.

Remark 1 (ε -subdifferential of a convex quadratic function [29, Formula 1.2.5]). Let \mathcal{X} be a Hilbert space. For a function of the form $h_0(\cdot) = \frac{1}{2\alpha} \|\cdot -y\|^2$ for some $y \in \mathcal{X}$ and $\alpha \geq 0$, we have

$$(\forall x \in \mathcal{X}) \qquad \qquad \partial_0^{\varepsilon} h_0(x) = \{ \frac{x - y}{\alpha} + \frac{e}{\alpha} \mid \frac{1}{2\alpha} ||e||^2 \leqslant \varepsilon \}.$$

Corollary 1 ([7, Proposition 5]). Let \mathcal{X} be a Hilbert space and let $h \in \Gamma_{\rho}(\mathcal{X})$. Let $\varepsilon \geq 0$, $\alpha > 0$ and $(x_{\varepsilon}, y) \in \mathcal{X}^2$ such that $x_{\varepsilon} \in \varepsilon$ -prox_{αh}(y). Then there exist $\varepsilon_0, \varepsilon_1 \geq 0$ with $\varepsilon_0 + \varepsilon_1 \leq \varepsilon$ such that, for any $e \in \mathcal{X}$ with $\frac{\|e\|^2}{2\alpha} \leq \varepsilon_0$, we have

$$\frac{y - x_{\varepsilon} - e}{\alpha} \in \partial_{\rho/2}^{\varepsilon_1} h(x_{\varepsilon}) \tag{6}$$

The latter formula (6) leads to the following lemma, which is a generalisation of [55, Lemma 2] and [49, Lemma 2] to proximal ε -subdifferentials.

Corollary 2 ([7, Corollary 1]). Let \mathcal{X} be a Hilbert space. Let $h \in \Gamma_{\rho}(\mathcal{X})$ and let $\varepsilon \geq 0$, $\alpha > 0$. If $x_{\varepsilon} \in \varepsilon$ -prox_{αh}(y), then for any $e \in \mathcal{X}$ with $\frac{\|e\|^2}{2\alpha} \leq \varepsilon$, we have

$$\frac{y - x_{\varepsilon} - e}{\alpha} \in \partial_{\rho/2}^{\varepsilon} h(x_{\varepsilon})$$

3.1 Proximal operators

The main result of the present subsection is Lemma 2 which, for ρ -weakly convex functions, relates the proximal operator to the proximal subdifferential as defined in Definition 2 when $\varepsilon = 0$.

Definition 6 (Proximal Map). Let \mathcal{X} be a Hilbert space. Let $h: \mathcal{X} \to (-\infty, +\infty]$ be a proper ρ -weakly convex function on \mathcal{X} . For any $\alpha > 0$, the set-valued mapping $\operatorname{prox}_{\alpha h}: \mathcal{X} \rightrightarrows \mathcal{X}$ defined as

$$(\forall x \in \mathcal{X}) \qquad \operatorname{prox}_{\alpha h}(x) := \underset{y \in \mathcal{X}}{\operatorname{argmin}} \left\{ h(y) + \frac{1}{2\alpha} \|x - y\|^2 \right\}$$

is called *proximal map* of h at x with respect to parameter α . If $1/\alpha > \rho$, then $\operatorname{prox}_{\alpha h}$ is a single-valued map and $\operatorname{dom} \operatorname{prox}_{\alpha h} := \{x \in \mathcal{X} \mid \operatorname{prox}_{\alpha h}(x) \neq \emptyset\} = \mathcal{X}$. In general, $\operatorname{dom} \operatorname{prox}_{\alpha h}$ could be empty.

Lemma 2. Let \mathcal{X} be a Hilbert space. Let $h: \mathcal{X} \to (-\infty, +\infty]$ be a proper ρ -weakly convex function on \mathcal{X} . Then for $x \in \mathcal{X}$, $p \in \text{dom } h$ and $\alpha > 0$ we have

$$p \in \operatorname{prox}_{\alpha h}(x) \implies \frac{x-p}{\alpha} \in \partial_{\rho/2} h(p).$$
 (7)

If $1/\alpha \geqslant \rho$ then we have the equivalence in (7).

Proof. Take $p \in \text{prox}_{\alpha h}(x)$. For any $y \in \mathcal{X}$ we have

$$h(y) - h(p) + \frac{1}{2\alpha} (\|y - x\|^2 - \|p - x\|^2) \ge 0.$$

Let $z \in \mathcal{X}$ and $\lambda \in (0,1)$, taking $y = \lambda z + (1-\lambda) p$. Then

$$h(\lambda z + (1 - \lambda) p) - h(p) + \frac{1}{2\alpha} (\|\lambda z + (1 - \lambda) p - x\|^2 - \|p - x\|^2) \ge 0.$$

By Definition 1, we obtain

$$\lambda h\left(z\right) + \left(1 - \lambda\right) h\left(p\right) - h\left(p\right) + \frac{\rho}{2} \lambda \left(1 - \lambda\right) \left\|z - p\right\|^2 + \frac{1}{2\alpha} \left(\lambda^2 \left\|z - p\right\|^2 + 2\lambda \left\langle z - p, p - x\right\rangle\right) \geqslant 0.$$

Dividing both sides by λ and letting $\lambda \to 0$, we get

$$h(z) - h(p) + \frac{\rho}{2} ||z - p||^2 \geqslant \frac{1}{\alpha} \langle x - p, z - p \rangle.$$

This holds for all $z \in \mathcal{X}$ as we do not have any restriction when setting y. From the definition of proximal subdifferential, we have $(x-p)/\alpha \in \partial_{\rho/2}h(p)$ which concludes the proof. When $\frac{1}{\alpha}$, $> \rho$ then $h(\cdot) + \frac{1}{2\alpha} ||\cdot -x||^2$ is strongly convex which implies $\partial_{\rho/2}h(p) = \{\frac{x-p}{\alpha}\}$. Hence, the equivalence in (7) holds.

Remark 2. Notice that if function h in Lemma 2 is also assumed to be lower-semicontinuous, by directly applying the definition of critical point (see Definition 5) and the result in Theorem 1, we obtain

$$0 \in \partial_{\rho/2}(h + \frac{1}{2\alpha} \| \cdot -x \|^2)(p) = \partial_{\rho/2}h(p) + \frac{p-x}{\alpha}$$

hence

$$\frac{x-p}{\alpha} \in \partial_{\rho/2}h(p).$$

According to Definition 5, for a ρ -weakly convex function $h: \mathcal{X} \to (-\infty, +\infty]$, a point $x \in \text{dom } h$ is a $\rho/2$ -critical point if $0 \in \partial_{\rho/2}h(x)$, that is

$$(\forall y \in \mathcal{X})$$
 $h(y) - h(x) \ge \langle 0, y - x \rangle - \frac{\rho}{2} ||y - x||^2 = -\frac{\rho}{2} ||y - x||^2$

and this is a necessary (but not sufficient) condition for global minimality. Then, the following equivalence applies for every $p \in \text{dom}(h)$

$$\frac{x-p}{\alpha} \in \partial_{\rho/2}h(p) \iff 0 \in \partial_{\rho/2}h(p) + \frac{1}{2\alpha}\nabla_p\left(\|p-x\|^2\right).$$

4 Sharpness and criticality

In this section we analyse mutual relationship between the solution set S and ε -critical points of the minimisation problem

$$\underset{x \in \mathcal{X}}{\text{minimize}} \ h(x)$$

in the case the objective function h satisfies the following sharpness condition:

Definition 7 (Sharpness, [46, 18]). Let \mathcal{X} be a Hilbert space. Let us consider a function $h: \mathcal{X} \to (-\infty, +\infty]$ and let the set $S = \operatorname{argmin}_{x \in \mathcal{X}} h(x)$ be nonempty. Function h is said to satisfy the sharpness condition locally if there exists $\mu > 0$ and $\delta > 0$ such that

$$(\forall x \in B(S, \delta))$$
 $h(x) - \inf_{x \in \mathcal{X}} h(x) \ge \mu \operatorname{dist}(x, S),$

where we use the convention that if $S = \emptyset$, then

$$(\forall x \in \mathcal{X}) \qquad \operatorname{dist}(x, S) = +\infty.$$

If the condition above is satisfied for every $\delta > 0$ then the sharpness condition is said to hold globally and it reads as

$$(\forall x \in \mathcal{X}) \qquad h(x) - \inf_{x \in \mathcal{X}} h(x) \geqslant \mu \operatorname{dist}(x, S), \tag{8}$$

Example 3. Function $x \mapsto f(x)$ from Example 1 has minima at x = a and x = b with f(a) = f(b) = 0. Take $\delta = \left|\frac{a-b}{2}\right|$ so that $\overline{B(a,\delta)} \cap \overline{B(b,\delta)} = \left\{\frac{a+b}{2}\right\}$, where $B(x_0,\delta)$ denotes the open ball with centre in $x_0 \in \mathbb{R}$ and radius δ and $\overline{B(x_0,\delta)}$ its closure. We can prove that f is globally sharp with respect to the set $S = \{a,b\}$. If $x = \frac{a+b}{2}$ – which corresponds to the midpoint of the interval [a,b] –, then

$$f(x) = \left| -\left(\frac{a-b}{2}\right) \left(\frac{a-b}{2}\right) \right| = \left| \frac{a-b}{2} \right| \operatorname{dist}(x,S)$$

Let us now consider three cases for $x \neq \frac{a+b}{2}$:

- $x \in B(a, \delta)$, so $x \notin B(b, \delta)$ then $f(x) \geqslant \delta |x a| = \delta \operatorname{dist}(x, S)$;
- $x \in B(b, \delta)$, so $x \notin B(a, \delta)$ then $f(x) \ge \delta |x b| = \delta \operatorname{dist}(x, S)$;
- $x \notin B(a, \delta)$ and $x \notin B(b, \delta)$, then $f(x) \ge \delta |x a|$ and $f(x) \ge \delta |x b|$, hence

$$f(x) \ge \delta \min\{|x - a|, |x - b|\} = \delta \operatorname{dist}(x, S).$$

In conclusion, $x \mapsto f(x)$ is sharp for all $x \in \mathbb{R}$ with constant δ , for every $\delta \leqslant \left| \frac{a-b}{2} \right|$.

In the following proposition, we investigate the position of all the ε -critical points of f which do not belong to S.

Proposition 3 (Position of solutions relative to a- ε -critical points). Let \mathcal{X} be a Hilbert space. Let $h: \mathcal{X} \to (-\infty, +\infty]$ be a proper function on \mathcal{X} . Let $S = \operatorname{argmin}_{x \in \mathcal{X}} h(x)$ be non-empty and let h satisfy the sharpness condition (8) with a constant $\mu > 0$. Let $x \in \varepsilon$ -crit_a $(h) \setminus S$ for some $a \ge 0$. Then, for any $0 \le \varepsilon < \mu^2/4a$, the quantities

$$\tau_1(\varepsilon) := \frac{\mu - \sqrt{\mu^2 - 4a\varepsilon}}{2a} \quad and \quad \tau_2(\varepsilon) := \frac{\mu + \sqrt{\mu^2 - 4a\varepsilon}}{2a}$$

are well-defined and, for any $x \in \varepsilon$ -crit_a $(h) \setminus S$, either $\operatorname{dist}(x, S) \ge \tau_2$ or $0 < \operatorname{dist}(x, S) \le \tau_1(\varepsilon)$.

Proof. Let $x \in \mathcal{X}$ be a a- ε -critical point that does not belong to S. By the definition, $0 \in \partial_a^{\varepsilon} h(x)$, which is

$$(\forall y \in \mathcal{X}) \qquad \qquad h(x) - h(y) \leqslant a \|x - y\|^2 + \varepsilon.$$

By the global sharpness of h, this implies that

$$(\forall \overline{x} \in S) \qquad \qquad \mu \operatorname{dist}(x, S) \leqslant a \|x - \overline{x}\|^2 + \varepsilon$$

By taking the infimum in the right-hand side over all $\overline{x} \in S$ we obtain the inequality

$$\mu \operatorname{dist}(x, S) \leq a \operatorname{dist}^{2}(x, S) + \varepsilon.$$
 (9)

which is quadratic with respect to $\operatorname{dist}(x, S)$ and either $\operatorname{dist}(x, S) \ge \tau_2(\varepsilon)$ or $0 < \operatorname{dist}(x, S) \le \tau_1(\varepsilon)$.

From Proposition 3, given $0 \le \varepsilon < \mu^2/a$, for any a- ε -critical point of h which does not belong to S, either the point is included in a neighbourhood of S of radius $\tau_1(\varepsilon)$ or its distance from S is bounded from below by a positive quantity $\tau_2(\varepsilon)$. The result of Proposition 3 is in the spirit of the condition of proper separation of isocost surfaces [40], which is taken as an assumption in [1], where criticality is understood in terms of Clarke's subdifferential.

Definition 8 (Definition 1.1 [1]). A closed function $h : \mathbb{R}^n \to \mathbb{R} \cup \{+\infty\}$ has properly separated isocost surfaces if there exists $\varepsilon > 0$ such that

$$\bar{x} \in \operatorname{crit} h, \bar{y} \in \operatorname{crit} h, h(\bar{x}) \neq h(\bar{y}) \Longrightarrow \|\bar{x} - \bar{y}\| > \varepsilon.$$
 (10)

As reported in [40], the condition in (10) is satisfied for a function h that takes only a finite number of values on crit h or whenever the connected components of crit h are properly separated. In particular, the condition is automatically satisfied when h is convex. In the non-convex case a sufficient condition is that h is a quadratic function defined on a polyhedral set. Moreover, according to [11, Theorem 13], when h is a globally subanalytic function such that $h|_{\text{dom }h}$ is continuous with dom h closed, there exists a finite number of connected components of the set crit h and h is constant on each connected component. In this context criticality is understood in relation to the limiting subdifferential.

Remark 3. When $\varepsilon = 0$, for a a-critical point x that does not belong to S, inequality (9) reads as

$$\mu \operatorname{dist}(x, S) \leq a \operatorname{dist}^2(x, S).$$

and, by dividing both sides by dist(x, S) > 0 we obtain

$$\frac{\mu}{a} \leqslant \operatorname{dist}(x, S) \tag{11}$$

or equivalently

$$a$$
-crit $\backslash S \subset \{x \in \mathcal{X} \mid \frac{\mu}{a} \leq \operatorname{dist}(x,S)\}$

which is the same result as in [25, Lemma 3.1], that is stated for ρ -weakly convex functions and $a = \rho/2$.

Proposition 3, in the form considered in Remark 3, is illustrated in the following example, where inequality (11) is attained:

Example 4. Let us consider function $x \mapsto f(x)$ from Example 1. We notice that $x = \frac{a+b}{2}$ is a $(\rho/2)$ - ε -critical point in the sense of Definition 5 for $\rho = 2$ and $\varepsilon = 0$. We start by rewriting f in the following manner: for every $x \in \mathbb{R}$

$$f(x) = \max\{(x-a)(x-b), -(x-a)(x-b)\}\$$

= \text{max}\{x^2 - (a+b)x + ab, -x^2 - ab + (a+b)x\}

which implies the following inequality:

$$(\forall x \in \mathbb{R}) \qquad f(x) \geqslant -x^2 - ab + (a+b)x.$$

We now notice that for any $(a,b) \in \mathbb{R}^2$ the following identity holds

$$\frac{(a-b)^2}{4} - \frac{(a+b)^2}{4} = \left(\frac{a-b}{2} + \frac{a+b}{2}\right) \left(\frac{a-b}{2} - \frac{a+b}{2}\right) = -ab$$

hence

$$(\forall x \in \mathbb{R}) \quad f(x) \geqslant -x^2 + (a+b)x + \frac{(a-b)^2}{4} - \frac{(a+b)^2}{4} = -\left(x - \frac{a+b}{2}\right)^2 + \frac{(a-b)^2}{4}.$$

Given the fact that $f\left(\frac{a+b}{2}\right) = \frac{(a-b)^2}{4}$, the inequality above reads as

$$(\forall x \in \mathbb{R}) \qquad f(x) - f\left(\frac{a+b}{2}\right) \geqslant -\left(x - \frac{a+b}{2}\right)^2$$

which, according to Definition 4, implies that $0 \in \partial_{\rho/2}^{\varepsilon} f\left(\frac{a+b}{2}\right)$ for every $\varepsilon \geqslant 0$ and for $\rho = 2$. In addition, considering the fact that function f satisfies the sharpness condition with constant $\mu = \left|\frac{a-b}{2}\right|$ (as shown in Example 3), inequality (11) is satisfied at $\overline{x} = \frac{a+b}{2}$:

$$\frac{2\mu}{\rho} = \frac{2\left|\frac{a-b}{2}\right|}{2} = \left|\frac{a-b}{2}\right| = \operatorname{dist}(\overline{x}, S).$$

5 The Inexact FB Algorithm

Assumption 1 summarizes the standing assumptions on functions f and g, which are made in all the results of the subsequent sections.

Assumption 1. Let the following facts hold:

- i) function $f \in \Gamma_{\rho}(\mathcal{X})$ is bounded from below;
- ii) function $g \in \Gamma_0(\mathcal{X})$ is differentiable on its domain with a L_g -Lipschitz continuous gradient;
- iii) dom $f \cap \text{dom } g \neq \emptyset$;
- iv) the set $S = \operatorname{argmin}_{x \in \mathcal{X}} (f + g)(x)$ is non-empty.

We discuss the following forward-backward scheme.

Algorithm 1: Inexact Forward-Backward

```
Initialise: x_0 \in \text{dom}(f+g);

Set: (\varepsilon_t)_{t \in \mathbb{N}} non-negative, (\alpha_t)_{t \in \mathbb{N}} positive;

for t=1,2,\ldots until convergence do
y_t = x_t - \alpha_t \nabla g(x_t)
x_{t+1} \in \varepsilon_t\text{-prox}_{\alpha_t f}(y_t)
end
```

In our analysis below, we do not tackle the problem of computing the ε -proximal point of function f at any given y. Instead we assume that there exists an oracle which provides us with an ε -proximal point. Recall that we defined ε -proximal points as ε -solutions of the corresponding optimisation problem, see Definition 4. An example of an estimation procedure has been proposed in [54, 57] for convex functions.

Inexact Forward-Backward schemes in the general non-convex setting have been proposed in [2, Algorithm 3, Theorem 5.1] and in [14]. In the first work, the scheme relies on a pair of error conditions, namely the sufficient decrease and the relative error condition, which allow to theoretically handle the problems arising from the approximation of the proximal map, but are difficult to treat and implement in practical situations. In the second work, a linesearch-based inexact proximal gradient scheme is designed for problems where function g is not necessarily convex, whereas f is convex and its proximal map can be approximated by means of the strategy proposed in [54, 57]. We therefore highlight that in Algorithm 1 we take the proximal step on the weakly convex function f, while the convex function g is also differentiable with a L_g -Lipschitz continuous gradient.

By applying the result of the sum rule for proximal ε -subdifferentials (Theorem 1) in the form of Corollary 2, we investigate the decreasing behaviour of the objective function in (1) for a sequence generated by the Inexact Forward-Backward scheme in Algorithm 1.

Proposition 4. Let \mathcal{X} be a Hilbert space. Let $f, g : \mathcal{X} \to (-\infty, +\infty]$ satisfy Assumption 1. Let $(x_t)_{t \in \mathbb{N}}$ be the sequence generated by Algorithm 1. Then for any $x \in \mathcal{X}$, the following estimate holds:

$$(f+g)(x) - (f+g)(x_{t+1}) \geqslant \left\langle \frac{x_t - x_{t+1}}{\alpha_t}, x - x_{t+1} \right\rangle - \frac{\rho}{2} \|x - x_{t+1}\|^2 - \frac{L_g}{2} \|x_t - x_{t+1}\|^2 - \varepsilon_t - \sqrt{\frac{2\varepsilon_t}{\alpha_t}} \|x - x_{t+1}\|$$
(12)

Moreover, $(f+g)(x_t)$ satisfies

$$(f+g)(x_t) - (f+g)(x_{t+1}) > -\varepsilon_t - \sqrt{\frac{2\varepsilon_t}{\alpha_t}} \|x_t - x_{t+1}\|$$
(13)

when $2/\alpha_t > \rho + L_a$.

Proof. By Algorithm 1 and Corollary 2, for every $t \in \mathbb{N}$ there exists $e_t \in \mathcal{X}$ with $\frac{\|e_t\|^2}{2\alpha_t} \leq \varepsilon_t$ such that

$$\frac{x_t - \alpha_t \nabla g(x_t) - x_{t+1} - e_t}{\alpha_t} \in \partial_{\rho/2}^{\varepsilon_t} f(x_{t+1}).$$

By definition of proximal ε -subdifferential, for any $x \in \mathcal{X}$

$$f(x) - f(x_{t+1}) \geqslant \left\langle \frac{x_t - x_{t+1}}{\alpha_t} - \nabla g(x_t), x - x_{t+1} \right\rangle - \frac{\rho}{2} \left\| x - x_{t+1} \right\|^2 - \left\langle \frac{e_t}{\alpha_t}, x - x_{t+1} \right\rangle - \varepsilon_t.$$

$$\tag{14}$$

We notice that

$$\langle -\nabla g(x_t), x - x_{t+1} \rangle = \langle -\nabla g(x_t), x - x_t + x_t - x_{t+1} \rangle$$

$$= \langle -\nabla g(x_t), x - x_t \rangle + \langle -\nabla g(x_t), x_t - x_{t+1} \rangle$$

$$\geqslant g(x_t) - g(x) + \langle \nabla g(x_t), x_{t+1} - x_t \rangle$$
(15)

where the last inequality stems from the convexity of function g

$$(\forall x \in X) \qquad q(x) - q(x_t) \geqslant \langle \nabla q(x_t), x - x_t \rangle.$$

By combining (14) and (15) we obtain

$$f(x) - f(x_{t+1}) \ge \left\langle \frac{x_t - x_{t+1}}{\alpha_t}, x - x_{t+1} \right\rangle - \frac{\rho}{2} \|x - x_{t+1}\|^2 - \left\langle \frac{e_t}{\alpha_t}, x - x_{t+1} \right\rangle$$
 (16)
+ $g(x_t) - g(x) + \left\langle \nabla g(x_t), x_{t+1} - x_t \right\rangle - \varepsilon_t,$

As a consequence, for any $x \in \mathcal{X}$

$$(f+g)(x) - (f+g)(x_{t+1}) + g(x_{t+1}) - g(x_{t}) \geqslant \left\langle \frac{x_{t} - x_{t+1}}{\alpha_{t}}, x - x_{t+1} \right\rangle - \frac{\rho}{2} \|x - x_{t+1}\|^{2} + \left\langle \nabla g(x_{t}), x_{t+1} - x_{t} \right\rangle - \left\langle \frac{e_{t}}{\alpha_{t}}, x - x_{t+1} \right\rangle - \varepsilon_{t}$$
(17)

Since g has a Lipschitz continuous gradient, we can use Lemma 1, i.e.

$$\langle \nabla g(x_t), x_{t+1} - x_t \rangle \geqslant g(x_{t+1}) - g(x_t) - \frac{L_g}{2} ||x_t - x_{t+1}||^2.$$

On the other side, by the Cauchy-Schwarz inequality, for any $x \in \mathcal{X}$

$$\left\langle \frac{e_t}{\alpha_t}, x - x_{t+1} \right\rangle \leqslant \frac{1}{\alpha_t} \|e_t\| \|x - x_{t+1}\| \leqslant \frac{1}{\alpha_t} \sqrt{2\alpha_t \varepsilon_t} \|x - x_{t+1}\| \leqslant \sqrt{\frac{2\varepsilon_t}{\alpha_t}} \|x - x_{t+1}\|$$
 (18)

By plugging (17) and (18) into (16) we obtain that for any $x \in \mathcal{X}$

$$(f+g)(x) - (f+g)(x_{t+1})$$

$$\geqslant \left\langle \frac{x_t - x_{t+1}}{\alpha_t}, x - x_{t+1} \right\rangle - \frac{\rho}{2} \|x - x_{t+1}\|^2 - \frac{L_g}{2} \|x_t - x_{t+1}\|^2 - \varepsilon_t - \sqrt{\frac{2\varepsilon_t}{\alpha_t}} \|x - x_{t+1}\|.$$

which is the first assertion (12). To get the second assertion, observe that substituting $x = x_t$ in (12) yields

$$(f+g)(x_t) - (f+g)(x_{t+1}) \ge \left(\frac{1}{\alpha_t} - \frac{\rho}{2} - \frac{L_g}{2}\right) \|x_t - x_{t+1}\|^2 - \varepsilon_t - \sqrt{\frac{2\varepsilon_t}{\alpha_t}} \|x_t - x_{t+1}\|.$$

Requiring $2/\alpha_t > \rho + L_g$ yields

$$(f+g)(x_t) - (f+g)(x_{t+1}) > -\varepsilon_t - \sqrt{\frac{2\varepsilon_t}{\alpha_t}} ||x_t - x_{t+1}||.$$

We highlight that, at this stage of the analysis, we do not assume that the objective function in (1) satisfies the sharpness condition (8), which implies that the result of Proposition 4 applies to a broader class of problems than the one we focus on in this work.

Remark 4. When $\varepsilon_t = 0$ for every $t \in \mathbb{N}$, inequality (13) from Proposition 4 reduces to

$$(f+g)(x_t) - (f+g)(x_{t+1}) > 0,$$

which implies that the objective value diminishes in a strictly monotone way. If the objective function is lower bounded, i.e. $\inf(f+g) > -\infty$, then $(f+g)(x_t)$ is convergent as t goes to infinity. In addition, we observe that the restriction that we have on the step size

$$\alpha_t < \frac{2}{L_g + \rho}$$

generalises the one of the convex case (i.e. when $\rho = 0$) which is $\alpha_t < 2/L_q$ (see [23]).

6 Convergence properties: the general case

Before starting our analysis, we point out that in a recent work [1], the authors propose a unified analysis of descent sequences in the context of weakly convex optimisation, considering a general optimisation scheme (see [1, Equation (1)]), which also accounts for inexact computations. The difference between our scheme and the one proposed in [1, Section 5.1.3] lies in the assumptions imposed on the functions f and g. Namely, instead of sharpness, the authors rely on the assumptions that their objective function satisfies (i) the proper separation of isocost surfaces condition (see Definition 8), and (ii) the sub-differential error bound [1, Definition 3.1], to achieve a convergence results.

In the remainder of the paper we will consider the following assumption.

Assumption 2. Let $(\varepsilon_t)_{t\in\mathbb{N}}$ be a sequence of non-negative values and $(\alpha_t)_{t\in\mathbb{N}}$ and $(\eta_t)_{t\in\mathbb{N}}$ be two sequences of positive values. We assume that the following inequalities are satisfied:

$$1 - \alpha_t \rho - \eta_t > 0 \tag{19}$$

$$\alpha_t \mu^2 \geqslant 2 \left(\varepsilon_t + \frac{\varepsilon_t}{\eta_t} \right) (\alpha_t \rho + \eta_t)$$
 (20)

$$\frac{1}{L_a} \geqslant \alpha_t. \tag{21}$$

In addition, to ease our presentation, especially when discussing the connection between the exact case (that is when $\varepsilon_t = 0$, $\eta_t = 0$ for every t) and the inexact case, we assume that $1/\eta_t = 0$ whenever $\eta_t = 0$.

The existence of α_t , ε_t and η_t satisfying Assumption 1 is discussed in Lemma 8 in Appendix 4. For any sequences $(\alpha_t)_{t\in\mathbb{N}}$, $(\varepsilon_t)_{t\in\mathbb{N}}$ and $(\eta_t)_{t\in\mathbb{N}}$ satisfying Assumption 2 and any $t \in \mathbb{N}$ the following quantities will be used in the sequel,

$$E_{t}^{-} := \frac{\alpha_{t}\mu - \sqrt{\alpha_{t}^{2}\mu^{2} - 2\alpha_{t}\varepsilon_{t}\left(1 + \frac{1}{\eta_{t}}\right)\left(\alpha_{t}\rho + \eta_{t}\right)}}{\alpha_{t}\rho + \eta_{t}}$$

$$E_{t}^{+} := \frac{\alpha_{t}\mu + \sqrt{\alpha_{t}^{2}\mu^{2} - 2\alpha_{t}\varepsilon_{t}\left(1 + \frac{1}{\eta_{t}}\right)\left(\alpha_{t}\rho + \eta_{t}\right)}}{\alpha_{t}\rho + \eta_{t}}.$$
(22)

$$E_t^+ := \frac{\alpha_t \mu + \sqrt{\alpha_t^2 \mu^2 - 2\alpha_t \varepsilon_t \left(1 + \frac{1}{\eta_t}\right) \left(\alpha_t \rho + \eta_t\right)}}{\alpha_t \rho + \eta_t}.$$
 (23)

Remark 5. We notice that the following inequalities hold for every $t \in \mathbb{N}$:

$$E_{t}^{+} = \frac{\alpha_{t}\mu + \sqrt{\alpha_{t}^{2}\mu^{2} - 2\alpha_{t}^{2}\left(\varepsilon_{t} + \frac{\varepsilon_{t}}{\eta_{t}}\right)\left(\rho + \frac{\eta_{t}}{\alpha_{t}}\right)}}{\alpha_{t}\left(\rho + \frac{\eta_{t}}{\alpha_{t}}\right)} = \frac{\mu + \sqrt{\mu^{2} - 2\left(\varepsilon_{t} + \frac{\varepsilon_{t}}{\eta_{t}}\right)\left(\rho + \frac{\eta_{t}}{\alpha_{t}}\right)}}{\left(\rho + \frac{\eta_{t}}{\alpha_{t}}\right)}$$

$$\leq \frac{\mu + \sqrt{\mu^{2} - 2\left(\varepsilon_{t} + \frac{\varepsilon_{t}}{\eta_{t}}\right)\rho}}{\rho} \leq \frac{\mu + \sqrt{\mu^{2} - 2\varepsilon_{t}\rho}}{\rho} = \tau_{2}(\varepsilon_{t})$$

which implies that if $\operatorname{dist}(x_t, S) \leq E_t^+$, then we also have $\operatorname{dist}(x_t, S) \leq \tau_2(\varepsilon_t)$.

We start by analysing a single step of Algorithm 1. Namely, for any $t \in \mathbb{N}$, if condition (26) below holds, it ensures that $\operatorname{dist}(x_{t+1}, S) \leq \frac{1}{\xi_{t+1}} \operatorname{dist}(x_t, S)$, where $\xi_{t+1} > 1$.

Proposition 5. Let \mathcal{X} be a Hilbert space. Let $f, g : \mathcal{X} \to (-\infty, +\infty]$ satisfy Assumption 1 and the function f + g satisfy the sharpness condition with constant $\mu > 0$. Let $(x_t)_{t\in\mathbb{N}}$ be the sequence generated by Algorithm 1 with $(\eta_t)_{t\in\mathbb{N}}$, $(\alpha_t)_{t\in\mathbb{N}}$ and $(\varepsilon_t)_{t\in\mathbb{N}}$ satisfying Assumption 2. Then, at iteration $t \in \mathbb{N}$, we have either dist $(x_{t+1}, S) = 0$ or the following inequality

$$\xi_{t+1}\operatorname{dist}^{2}(x_{t+1},S) \leqslant \operatorname{dist}^{2}(x_{t},S). \tag{24}$$

where

$$\xi_{t+1} := 1 - \alpha_t \rho - \eta_t + \frac{2\alpha_t \mu}{\operatorname{dist}(x_{t+1}, S)} - \frac{2\alpha_t \varepsilon_t}{\operatorname{dist}^2(x_{t+1}, S)} \left(1 + \frac{1}{\eta_t}\right). \tag{25}$$

Moreover, if the following condition is satisfied

$$E_t^- < \operatorname{dist}(x_{t+1}, S) < E_t^+,$$
 (26)

then $\xi_{t+1} > 1$.

Proof. Take an arbitrary but fixed $t \in \mathbb{N}$ and let $\operatorname{dist}(x_{t+1}, S) > 0$. We take any $\overline{x} \in S$ and apply (12) from Proposition 4 for $x = \overline{x}$, which yields the following inequality:

$$(f+g)(\overline{x}) - (f+g)(x_{t+1}) \geqslant \left\langle \frac{x_t - x_{t+1}}{\alpha_t}, \overline{x} - x_{t+1} \right\rangle$$
$$- \frac{\rho}{2} \|\overline{x} - x_{t+1}\|^2 - \frac{L_g}{2} \|x_t - x_{t+1}\|^2 - \varepsilon_t - \sqrt{\frac{2\varepsilon_t}{\alpha_t}} \|\overline{x} - x_{t+1}\|.$$

By using the identity

$$\left(\forall (w,y,z) \in \mathcal{X}^{3}\right) \qquad \qquad \langle w-y,y-z \rangle = \frac{1}{2}\|w-z\|^{2} - \frac{1}{2}\|w-y\|^{2} - \frac{1}{2}\|y-z\|^{2},$$

which follows from [5, Lemma 2.12], we obtain

$$(f+g)(\overline{x}) - (f+g)(x_{t+1})$$

$$\geqslant \frac{1}{2\alpha_{t}} \left(-\|x_{t} - \overline{x}\|^{2} + \|x_{t+1} - \overline{x}\|^{2} + \|x_{t} - x_{t+1}\|^{2} \right)$$

$$- \frac{\rho}{2} \|\overline{x} - x_{t+1}\|^{2} - \frac{L_{g}}{2} \|x_{t} - x_{t+1}\|^{2} - \varepsilon_{t} - \sqrt{\frac{2\varepsilon_{t}}{\alpha_{t}}} \|\overline{x} - x_{t+1}\|$$

$$= \left(\frac{1}{2\alpha_{t}} - \frac{L_{g}}{2} \right) \|x_{t} - x_{t+1}\|^{2} + \left(\frac{1}{2\alpha_{t}} - \frac{\rho}{2} \right) \|\overline{x} - x_{t+1}\|^{2}$$

$$- \frac{1}{2\alpha_{t}} \|x_{t} - \overline{x}\|^{2} - \varepsilon_{t} - \sqrt{\frac{2\varepsilon_{t}}{\alpha_{t}}} \|\overline{x} - x_{t+1}\|.$$
(27)

Notice that for every $\eta > 0$, the following inequality (known as Young's Inequality) holds

$$(\forall (u,v) \in [0,+\infty)^2) \qquad uv \leqslant \frac{1}{2} \left(\eta u^2 + \frac{1}{\eta} v^2 \right). \tag{28}$$

Hence, at iteration t, we use the constant $\eta_t > 0$ to estimate the last term in (27) according to (28) with $v = \sqrt{\frac{1}{\alpha_t}} \|\overline{x} - x_{t+1}\|$ and $u = \sqrt{2\varepsilon_t}$, which yields

$$-\sqrt{\frac{2\varepsilon_t}{\alpha_t}} \|\overline{x} - x_{t+1}\| \ge -\frac{\eta_t}{2\alpha_t} \|\overline{x} - x_{t+1}\|^2 - \frac{\varepsilon_t}{\eta_t}.$$
 (29)

In conclusion, by combining (27) and (29), we obtain

$$(f+g)(\overline{x}) - (f+g)(x_{t+1}) \geqslant \left(\frac{1}{2\alpha_t} - \frac{L_g}{2}\right) \|x_t - x_{t+1}\|^2 + \left(\frac{1}{2\alpha_t} - \frac{\rho}{2} - \frac{\eta_t}{2\alpha_t}\right) \|\overline{x} - x_{t+1}\|^2 - \frac{1}{2\alpha_t} \|x_t - \overline{x}\|^2 - \varepsilon_t - \frac{\varepsilon_t}{\eta_t}.$$
(30)

We then estimate the LHS by the assumption of sharpness on f + g

$$-\mu \operatorname{dist}(x_{t+1}, S) \geqslant \left(\frac{1}{2\alpha_t} - \frac{\rho}{2} - \frac{\eta_t}{2\alpha_t}\right) \|\overline{x} - x_{t+1}\|^2 - \frac{1}{2\alpha_t} \|x_t - \overline{x}\|^2 + \left(\frac{1}{2\alpha_t} - \frac{L_g}{2}\right) \|x_t - x_{t+1}\|^2 - \varepsilon_t - \frac{\varepsilon_t}{\eta_t}.$$

By bringing $\frac{1}{2\alpha_t} \|x_t - \overline{x}\|^2$ to the LHS of the inequality, we observe that both terms involving $\overline{x} \in S$ have a positive sign

$$\frac{1}{2\alpha_t} \|x_t - \overline{x}\|^2 - \mu \operatorname{dist}(x_{t+1}, S) \geqslant \left(\frac{1}{2\alpha_t} - \frac{\rho}{2} - \frac{\eta_t}{2\alpha_t}\right) \|\overline{x} - x_{t+1}\|^2 + \left(\frac{1}{2\alpha_t} - \frac{L_g}{2}\right) \|x_t - x_{t+1}\|^2 - \varepsilon_t - \frac{\varepsilon_t}{\eta_t}.$$

This allows us to take the infimum over $\overline{x} \in S$ in both sides of the inequality, yielding the following inequality involving the distance to the solution set

$$\frac{1}{2\alpha_t} \operatorname{dist}^2(x_t, S) - \mu \operatorname{dist}(x_{t+1}, S) \geqslant \left(\frac{1}{2\alpha_t} - \frac{\rho}{2} - \frac{\eta_t}{2\alpha_t}\right) \operatorname{dist}^2(x_{t+1}, S) + \left(\frac{1}{2\alpha_t} - \frac{L_g}{2}\right) \|x_t - x_{t+1}\|^2 - \varepsilon_t - \frac{\varepsilon_t}{\eta_t} \tag{31}$$

Rearranging the terms in the inequality above, we obtain

$$\operatorname{dist}^{2}(x_{t}, S) \geq (1 - \alpha_{t}\rho - \eta_{t}) \operatorname{dist}^{2}(x_{t+1}, S) + 2\alpha_{t}\mu \operatorname{dist}(x_{t+1}, S) - 2\alpha_{t}(\varepsilon_{t} + \frac{\varepsilon_{t}}{\eta_{t}}) + (1 - \alpha L_{g}) \|x_{t} - x_{t+1}\|^{2}.$$

$$(32)$$

By (21), we can skip the term $(1 - \alpha L_g) \|x_t - x_{t+1}\|^2 \ge 0$ and rewrite (32) as

$$dist^{2}(x_{t}, S) \geqslant \xi_{t+1} dist^{2}(x_{t+1}, S),$$

where

$$\xi_{t+1} = 1 - \alpha_t \rho - \eta_t + \frac{2\alpha_t \mu}{\operatorname{dist}(x_{t+1}, S)} - \frac{2\alpha_t \varepsilon_t}{\operatorname{dist}^2(x_{t+1}, S)} \left(1 + \frac{1}{\eta_t}\right).$$

The above inequality proves (24).

To have $\xi_{t+1} > 1$, by (25) the following inequality must hold

$$(\alpha_t \rho + \eta_t) \operatorname{dist}^2(x_{t+1}, S) - 2\alpha_t \mu \operatorname{dist}(x_{t+1}, S) + 2\alpha_t \varepsilon_t \left(1 + \frac{1}{\eta_t}\right) < 0.$$
 (33)

This is a quadratic inequality (the quadratic term does not vanish as $\eta_t > 0$) with respect to dist (x_{t+1}, S) and its discriminant satisfies

$$\Delta = \alpha_t^2 \mu^2 - 2\alpha_t \varepsilon_t \left(1 + \frac{1}{\eta_t} \right) (\alpha_t \rho + \eta_t) > 0,$$

thanks to condition (20). Thus, for (33) to hold, $\operatorname{dist}(x_{t+1}, S)$ has to be bounded by the solutions of the quadratic equation associated with (33) which is (26)

$$E_t^- < \operatorname{dist}(x_{t+1}, S) < E_t^+$$

Remark 6. When $\varepsilon_t > 0$, in (27) we use Young's inequality (28) on the term $\sqrt{\frac{2\varepsilon_t}{\alpha_t}} \| \overline{x} - x_{t+1} \|$ with $v = \sqrt{\frac{1}{\alpha_t}} \| \overline{x} - x_{t+1} \|$ and $u = \sqrt{2\varepsilon_t}$, as illustrated in (29), to obtain an estimation where all the norms appear with a quadratic exponent. More precisely, we use a modified version of Young's inequality which relies on the constant $\eta_t > 0$. This choice allows the introduction of a positivity constraint on the term $\left(\frac{1}{2\alpha_t} - \frac{\rho}{2} - \frac{\eta_t}{2\alpha_t}\right)$ appearing in (30) – see condition (20) in Assumption 2 – which would otherwise simplify to the negative term $-\frac{\rho}{2}$ in the case of the elementary Young's Inequality (that is when $\eta_t = 1$). The idea behind this strategy is to obtain a result in Proposition 5 for $\varepsilon_t > 0$ that is strongly related to what can be inferred for $\varepsilon_t = 0$, which we will illustrate in Corollary 3. We highlight that the sequence $(\eta_t)_{t\in\mathbb{N}}$ is not related to the model nor to the optimisation method and its presence (which is subordinated to the fact that $\varepsilon_t > 0$ for every $t \in \mathbb{N}$ there exists an explicit choice for α_t , η_t , and ε_t satisfying conditions (19), (20), and (21).

Remark 7. When $\varepsilon_t = 0$ for every $t \in \mathbb{N}$ (and consequently $\eta_t = 0$ for every $t \in \mathbb{N}$), conditions (19), (20), and (21) in Assumption 2 boil down to

$$(\forall t \in \mathbb{N}) \qquad \qquad \alpha_t < \min\left\{\frac{1}{L_g}, \frac{1}{\rho}\right\}.$$

This implies

$$(\forall t \in \mathbb{N}) \qquad \qquad 2 > \rho \alpha_t + L_q \alpha_t$$

which is exactly what is required in Proposition 4 to ensure that the sequence of objective values $((f+g)(x_t))_{t\in\mathbb{N}}$ is monotonically decreasing. To conclude, we also remark that in the case of exact proximal computations, the quantities E_t^- and E_t^+ take the constant values

$$E_t^- = \tau_1(0) = 0, \qquad E_t^+ = \tau_2(0) = \frac{2\mu}{\rho}.$$

for every $t \in \mathbb{N}$.

In case of exact proximal computations, Proposition 5 takes the form presented in the following corollary.

Corollary 3. Let \mathcal{X} be a Hilbert space. Let $f, g: \mathcal{X} \to (-\infty, +\infty]$ satisfy Assumption 1 and the function f + g satisfy the sharpness condition with constant $\mu > 0$. Let $\varepsilon_t = 0$ and $\eta_t = 0$ for every $t \in \mathbb{N}$. Let $(x_t)_{t \in \mathbb{N}}$ be the sequence generated by Algorithm 1 with the stepsize $(\alpha_t)_{t \in \mathbb{N}}$ satisfying the condition

$$(\forall t \in \mathbb{N}) \qquad \qquad \alpha_t < \min\left\{\frac{1}{L_g}, \frac{1}{\rho}\right\}$$

Then, we have either $dist(x_{t+1}, S) = 0$ or the following inequality

$$\xi_{t+1} \operatorname{dist}^{2}(x_{t+1}, S) \leq \operatorname{dist}^{2}(x_{t}, S).$$

where

$$\xi_{t+1} = 1 - \alpha_t \rho + \frac{2\alpha_t \mu}{\operatorname{dist}(x_{t+1}, S)}.$$

Moreover, if for some $t_0 \in \mathbb{N}$ such that $\operatorname{dist}(x_{t_0}, S) < \frac{2\mu}{\rho}$ then following condition is satisfied

$$\operatorname{dist}(x_{t+1}, S) < \frac{2\mu}{\rho},$$

and $\xi_{t+1} > 1$ for all $t \ge t_0$.

As we can see in Proposition 5, for the quantity $dist(x_t, S)$ to diminish, the distance from the next iterate to S has to be bounded from below by E_t^- and from above by E_t^+ .

7 Convergence analysis: all the parameters fixed

In this section we pursue our analysis when all the concerned parameters $\alpha_t, \varepsilon_t, \eta_t$ are fixed for every $t \in \mathbb{N}$ and satisfy Assumption 2. Then, the quantities E_t^+ and E_t^- from (23) and (22) become

$$E^{-} := \frac{\alpha\mu - \sqrt{\alpha^{2}\mu^{2} - 2\alpha\varepsilon\left(1 + \frac{1}{\eta}\right)(\alpha\rho + \eta)}}{(\alpha\rho + \eta)}$$
$$E^{+} := \frac{\alpha\mu + \sqrt{\alpha^{2}\mu^{2} - 2\alpha\varepsilon\left(1 + \frac{1}{\eta}\right)(\alpha\rho + \eta)}}{(\alpha\rho + \eta)}.$$

Lemma 3. Let \mathcal{X} be a Hilbert space. Let $f, g : \mathcal{X} \to (-\infty, +\infty]$ satisfy Assumption 1 and the function f + g satisfy the sharpness condition with constant $\mu > 0$. Let $(x_t)_{t \in \mathbb{N}}$ be the sequence generated by Algorithm 1 with ε , α and η chosen according to Assumption 2. Assuming that for some $t_0 \in \mathbb{N}$ we have

$$dist(x_{t_0}, S) \leqslant E^+ \tag{34}$$

then for every $t \ge t_0$ we have

$$dist(x_t, S) \leqslant E^+. \tag{35}$$

Moreover, if for a certain t_0 we have

$$dist(x_{t_0}, S) \leqslant E^- \tag{36}$$

then, for every $t \ge t_0$ we have

$$dist(x_t, S) \leqslant E^- \tag{37}$$

Proof. Starting from (32) in the proof of Proposition 5, we recall that, for any $t \in \mathbb{N}$, the inequality holds

$$\operatorname{dist}^{2}(x_{t}, S) \geqslant (1 - \alpha \rho - \eta) \operatorname{dist}^{2}(x_{t+1}, S) + 2\alpha \mu \operatorname{dist}(x_{t+1}, S) - 2\alpha(\varepsilon + \frac{\varepsilon}{\eta}).$$

Let now $t = t_0$. From (34), by taking into account the definition of E^+ , we have

$$(1 - \alpha \rho - \eta) \operatorname{dist}^{2}(x_{t+1}, S) + 2\alpha \mu \operatorname{dist}(x_{t+1}, S) - 2\alpha \left(\varepsilon + \frac{\varepsilon}{\eta}\right)$$

$$\leq \left(\frac{\alpha \mu + \sqrt{\alpha^{2} \mu^{2} - 2\alpha \left(\varepsilon + \frac{\varepsilon}{\eta}\right) (\alpha \rho + \eta)}}{\alpha \rho + \eta}\right)^{2}.$$
(38)

which is a second degree inequality with respect $dist(x_{t+1}, S)$. It can be shown that the discriminant has the following form (see equation (60) in Appendix B):

$$\Delta = \left(\frac{\alpha\mu + (1 - \alpha\rho - \eta)\sqrt{\alpha^2\mu^2 - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}}{\alpha\rho + \eta}\right)^2 \geqslant 0.$$

Since $\Delta \ge 0$ we have that the solutions are included in the following range:

$$\frac{-\alpha\mu - \sqrt{\Delta}}{1 - \alpha\rho - \eta} \leqslant \operatorname{dist}(x_{t+1}, S) \leqslant \frac{-\alpha\mu + \sqrt{\Delta}}{1 - \alpha\rho - \eta}$$

where the lower bound is clearly negative, while the upper bound reads as

$$\frac{-\alpha\mu + \sqrt{\Delta}}{1 - \alpha\rho - \eta} = \frac{\alpha\mu + \sqrt{\alpha^2\mu^2 - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}}{(\alpha\rho + \eta)}$$
(39)

(see equation (61) in Appendix B). In conclusion, since the distance from a set is a non-negative value, we obtain

$$0 \leqslant \operatorname{dist}(x_{t+1}, S) \leqslant \frac{\alpha \mu + \sqrt{\alpha^2 \mu^2 - 2\alpha \left(\varepsilon + \frac{\varepsilon}{\eta}\right) (\alpha \rho + \eta)}}{(\alpha \rho + \eta)} = E^+.$$

which by induction proves (35).

The same reasoning can be applied for (36) to obtain (37): see equations (62) and (63) in Appendix B.

Lemma 4. Under the setting of Lemma 3, condition (34) implies that for every $t \ge t_0$ there exists $\zeta_{t+1} \ge 1$ such that

$$\zeta_{t+1} \left(\operatorname{dist}^2(x_{t+1}, S) - (E^-)^2 \right) \le \operatorname{dist}^2(x_t, S) - (E^-)^2.$$
(40)

Proof. We notice that the quantities E^+ and E^- are roots of the quadratic equation associated to inequality (33), so in particular we have

$$(E^{-})^{2} = (1 - \alpha \rho - \eta)(E^{-})^{2} + 2\alpha \mu E^{-} - 2\alpha \varepsilon \left(1 + \frac{1}{\eta}\right). \tag{41}$$

Subtracting (41) from (32), we obtain the following inequality that holds for every $t \in \mathbb{N}$ $\operatorname{dist}^2(x_t, S) - (E^-)^2 \geqslant$

$$(1 - \alpha \rho - \eta)(\operatorname{dist}^{2}(x_{t+1}, S) - (E^{-})^{2}) + 2\alpha \mu(\operatorname{dist}(x_{t+1}, S) - E^{-}) + (1 - \alpha L_{g})\|x_{t} - x_{t+1}\|^{2}$$

$$= (1 - \alpha \rho - \eta + \frac{2\alpha \mu}{\operatorname{dist}(x_{t+1}, S) + E^{-}}) \left(\operatorname{dist}^{2}(x_{t+1}, S) - (E^{-})^{2} \right) + (1 - \alpha L_{g}) \|x_{t} - x_{t+1}\|^{2}.$$

Then,

$$\operatorname{dist}^{2}(x_{t}, S) - (E^{-})^{2} \geqslant \zeta_{t+1} \left(\operatorname{dist}^{2}(x_{t+1}, S) - (E^{-})^{2} \right) + (1 - \alpha L_{g}) \|x_{t} - x_{t+1}\|^{2}, \quad (42)$$

where $\zeta_{t+1} := 1 - \alpha \rho - \eta + \frac{2\alpha\mu}{\operatorname{dist}(x_{t+1},S) + E^-} \geqslant 0$ is a non-negative quantity by virtue of Assumption 2. By skipping the term $(1 - \alpha L_g) \|x_t - x_{t+1}\|^2$, which is positive by (21), we have

$$\operatorname{dist}^{2}(x_{t}, S) - (E^{-})^{2} \geqslant \zeta_{t+1} \left(\operatorname{dist}^{2}(x_{t+1}, S) - (E^{-})^{2} \right).$$

We now need to show that at iteration $t \ge t_0$, we have $\zeta_{t+1} \ge 1$: we notice that the inequality $\zeta_{t+1} \ge 1$ holds whenever

$$\frac{2\alpha\mu}{\operatorname{dist}(x_{t+1},S) + E^{-}} \geqslant \alpha\rho + \eta$$

This is equivalent to the following condition

$$\operatorname{dist}(x_{t+1}, S) \leqslant \frac{2\alpha\mu}{\alpha\rho + \eta} - E^{-} = \frac{\alpha\mu + \sqrt{\alpha^{2}\mu^{2} - 2\alpha\varepsilon\left(1 + \frac{1}{\eta}\right)(\alpha\rho + \eta)}}{(\alpha\rho + \eta)} = E^{+}.$$

By assumption, there exists t_0 such that $\operatorname{dist}(x_{t_0}, S) \leq E^+$ and by Lemma 3 this implies that $\operatorname{dist}(x_{t+1}, S) \leq E^+$. In conclusion, for every $t \geq t_0$ the condition $\zeta_{t+1} \geq 1$ holds. \square

With an additional restriction on the distance function $\operatorname{dist}(x_{t+1}, S)$, *i.e.* (45), then $\operatorname{dist}(x_t, S)$ converges to E^- .

Theorem 2. Let \mathcal{X} be a Hilbert space. Let $f, g: \mathcal{X} \to (-\infty, +\infty]$ satisfy Assumption 1 and the function f+g satisfy the sharpness condition with constant $\mu > 0$. Assume that $\varepsilon_t = \varepsilon \geqslant 0$ and $\alpha_t = \alpha$ for all $t \in \mathbb{N}$ such that we can choose a constant $\eta > 0$ for which Assumption 2 holds. Let $(x_t)_{t \in \mathbb{N}}$ be the sequence generated by Algorithm 1 with respect to $(\varepsilon_t)_{t \in \mathbb{N}}$ and $(\alpha_t)_{t \in \mathbb{N}}$. If there exists $t_0 \in \mathbb{N}$ such that the following holds

$$E^- < \operatorname{dist}(x_{t_0}, S) < E^+ \tag{43}$$

then there exists a constant $\zeta > 1$ such that

$$(\forall t \ge t_0) \qquad \text{dist}^2(x_t, S) - (E^-)^2 \le \left(\frac{1}{\zeta}\right)^{t-t_0} (\text{dist}^2(x_{t_0}, S) - (E^-)^2). \tag{44}$$

implying

$$\limsup_{t\to\infty} \operatorname{dist}(x_t, S) \leqslant E^-.$$

Moreover, if

$$(\forall t \geqslant t_0) \qquad E^- < \operatorname{dist}(x_t, S) < E^+, \tag{45}$$

then

$$\lim_{t \to +\infty} \operatorname{dist}(x_t, S) = E^-.$$

Proof. From Lemma 4 (40) and since E^- is constant for every t, we know that

$$(\forall t \ge t_0)$$
 $\zeta_{t+1} \left(\text{dist}^2(x_{t+1}, S) - (E^-)^2 \right) \le \text{dist}^2(x_t, S) - (E^-)^2,$

where

$$\zeta_{t+1} := 1 - \alpha \rho - \eta + \frac{2\alpha \mu}{\operatorname{dist}(x_{t+1}, S) + E^{-}} \ge 1.$$

From this we infer that for every $t \ge t_0$ the following holds:

$$\operatorname{dist}^{2}(x_{t+1}, S) - (E^{-})^{2} \leq \frac{1}{\zeta_{t+1}} \left(\operatorname{dist}^{2}(x_{t}, S) - (E^{-})^{2} \right)$$

$$\leq \left(\prod_{s=t_{0}}^{t} \frac{1}{\zeta_{s+1}} \right) \left(\operatorname{dist}^{2}(x_{t_{0}}, S) - (E^{-})^{2} \right).$$

$$(46)$$

In order to proceed, we need to discuss the sign of the LHS in (46). In order to do so, we denote by t_1 the first iterate $t_1 > t_0$ such that $\operatorname{dist}(x_{t_1}, S) < E^-$ (without excluding the possibility that such a t_1 does not exist).

For every $t \ge t_1$, by Lemma 3, we have $\operatorname{dist}(x_t, S) \le E^-$, which implies that for every $t \ge t_1$ the LHS in (46) is non-positive:

$$(t \ge t_1)$$
 $\operatorname{dist}^2(x_t, S) - (E^-)^2 \le 0.$ (47)

On the other side, for every t such that $t_0 \le t < t_1$, we have $\operatorname{dist}(x_t, S) > E^-$. By Proposition 5, for $t_0 < t < t_1$ there exists $\xi_t > 1$ such that

$$\sqrt{\xi_t} \operatorname{dist}(x_t, S) \leq \operatorname{dist}(x_{t-1}, S)$$

implying that

$$\operatorname{dist}(x_t, S) < \operatorname{dist}(x_{t-1}, S) < \dots < \operatorname{dist}(x_{t_0}, S). \tag{48}$$

We can therefore define a lower-bound for ζ_t as follows, for $t_0 \leq t < t_1$

$$\zeta_{t} = 1 - \alpha \rho - \eta + \frac{2\alpha \mu}{\operatorname{dist}(x_{t}, S) + E^{-}}$$

$$\geqslant 1 - \alpha \rho - \eta + \frac{2\alpha \mu}{\operatorname{dist}(x_{t_{0}}, S) + E^{-}} = \zeta_{t_{0}}$$

$$> 1 - \alpha \rho - \eta + \frac{2\alpha \mu}{E^{+} + E^{-}} = 1$$
(49)

where the first inequality stems from (48) (we include the equality to take into account the case $t = t_0$), the second from (43) and the last equality from the defintion of E^+ and E^- . As a consequence, the product in the RHS of (46) can be estimated using ζ_{t_0} as follows

$$\left(\prod_{s=t_0}^t \frac{1}{\zeta_{s+1}}\right) \leqslant \left(\frac{1}{\zeta_{t_0}}\right)^{t-t_0+1}$$

implying that

$$(t_0 \le t < t_1)$$
 $\operatorname{dist}^2(x_t, S) - (E^-)^2 \le \left(\frac{1}{\zeta_{t_0}}\right)^{t-t_0} \left(\operatorname{dist}^2(x_{t_0}, S) - (E^-)^2\right).$ (50)

To conclude, we can combine (47) (which holds for $t > t_1$) and (50) (which holds for $t_0 \le t < t_1$) into the following inequality that holds for every $t \ge t_0$ by taking as an upper bound the positive upper bound in (50):

$$(\forall t \ge t_0)$$
 $\operatorname{dist}^2(x_t, S) - (E^-)^2 \le \left(\frac{1}{\zeta_{t_0}}\right)^{t-t_0} \left(\operatorname{dist}^2(x_{t_0}, S) - (E^-)^2\right).$

Taking the $\limsup for t \to \infty$ we conclude that

$$\limsup_{t \to \infty} \left(\operatorname{dist}^{2}(x_{t}, S) - (E^{-})^{2} \right) \leq \limsup_{t \to \infty} \left(\frac{1}{\zeta_{t_{0}}} \right)^{t-t_{0}} \left(\operatorname{dist}^{2}(x_{t_{0}}, S) - (E^{-})^{2} \right) \\
= \left(\operatorname{dist}^{2}(x_{t_{0}}, S) - (E^{-})^{2} \right) \lim_{t \to \infty} \left(\frac{1}{\zeta_{t_{0}}} \right)^{t-t_{0}} \\
= 0$$

where the last equality is due to the fact that $\zeta_{t_0} > 1$ as shown in (49). We then conclude that

$$\limsup_{t \to \infty} \operatorname{dist}(x_t, S) \leqslant E^-.$$

To show the second part of the statement, we notice that if (45) holds for all $t \ge t_0$, then combining it with (44), yields

$$(\forall t \ge t_0) \qquad 0 < \operatorname{dist}^2(x_t, S) - (E^-)^2 \le \left(\frac{1}{\zeta}\right)^{t - t_0} \left(\operatorname{dist}^2(x_{t_0}, S) - (E^-)^2\right) \tag{51}$$

where the RHS is non-negative and monotonically decreases to 0 as $t \to \infty$. Since according to (45), the LHS is positive, (51) implies

$$\lim_{t \to \infty} \operatorname{dist}(x_t, S) = E^-$$

which concludes the proof.

Remark 8. Inequality (44) from Theorem 2 is reminiscent of the results in [30, Theorem 1](in the convex case) and [25, Theorem 5.2] (in the weakly convex case) for subgradient methods. Notice that in these works, exactly as in (44), the LHS might be negative.

From Theorem 2, we know more about $(\operatorname{dist}(x_t, S))_{t \in \mathbb{N}}$ when (45) holds. In fact, we can have a convergence result for the sequence $(x_t)_{t \in \mathbb{N}}$ under the assumption (45) of Theorem 2.

Corollary 4. Under the assumptions of Theorem 2, if (45) holds and (21) from Assumption 2 is satisfied with a strict inequality, that is

$$\alpha < \frac{1}{L_q},$$

then $(x_t)_{t\in\mathbb{N}}$ converges.

Proof. By (42), there exists $\zeta_{t+1} > 1$ such that

$$\zeta_{t+1} \left(\operatorname{dist}^{2}(x_{t+1}, S) - (E^{-})^{2} \right) + (1 - \alpha L_{g}) \|x_{t} - x_{t+1}\|^{2} \leq \operatorname{dist}^{2}(x_{t}, S) - (E^{-})^{2}.$$
 (52)

Thanks to condition (45), the first term on the LHS of (52) is positive, which implies

$$(1 - \alpha L_q) \|x_t - x_{t+1}\|^2 \le \text{dist}^2(x_t, S) - (E^-)^2.$$

By the result of Theorem 2, the RHS can be estimated with a constant $\zeta > 1$

$$(1 - \alpha L_g) \|x_t - x_{t+1}\|^2 \le \left(\frac{1}{\zeta}\right)^{t-t_0} \left[\operatorname{dist}^2(x_{t_0}, S) - (E^-)^2\right]. \tag{53}$$

As both sides of (53) are non-negative for all $t \ge t_0$, taking the square root and summing over t to some $T > t_0$ yields

$$\sqrt{1 - \alpha L_g} \sum_{t \ge t_0}^T \|x_t - x_{t+1}\| \le \sqrt{\operatorname{dist}^2(x_{t_0}, S) - (E^-)^2} \sum_{t \ge t_0}^T \left(\frac{1}{\sqrt{\zeta}}\right)^{t - t_0}.$$
 (54)

Since we have $\zeta > 1$, we also have $\sqrt{\zeta} > 1$. In addition, by assumption, $\sqrt{1 - \alpha L_g}$ is positive. When letting $T \to \infty$, the RHS of (54) is finite, which implies that

$$\sum_{t \ge t_0}^{\infty} \|x_t - x_{t+1}\| < +\infty.$$

By virtue of [13, Theorem 1], we conclude that $(x_t)_{t\in\mathbb{N}}$ is a Cauchy sequence so it converges.

In the exact case $(\varepsilon = 0)$, it is possible to consider a step size which varies in a range $\alpha_t \in [\underline{\alpha}, \overline{\alpha}] \subset \left(0, \min\left\{\frac{1}{\rho}, \frac{1}{L_g}\right\}\right)$ and obtain a similar result as the one in Theorem 2:

Proposition 6 (Strong convergence of the sequence in the exact case). Let \mathcal{X} be a Hilbert space. Let $f, g: \mathcal{X} \to (-\infty, +\infty]$ satisfy Assumption 1 and the function f+g satisfy the sharpness condition with constant $\mu > 0$. Let $(x_t)_{t \in \mathbb{N}}$ be the sequence generated by Algorithm 1 with $\varepsilon = 0$ and step sizes $\alpha_t \in [\underline{\alpha}, \overline{\alpha}] \subset (0, \min\{\frac{1}{\rho}, \frac{1}{L_g}\})$. Assume that there exists a $t_0 \in \mathbb{N}$ such that

$$\operatorname{dist}(x_{t_0}, S) < \frac{2\mu}{\rho}.$$

Then the following holds

$$\lim_{t \to \infty} \operatorname{dist}(x_{t+1}, S) = 0 \tag{55}$$

and the sequence $(x_t)_{t\in\mathbb{N}}$ converges strongly to a point $x^*\in S$.

Proof. From Corollary 3, we have that for every $t \ge t_0$ there exists $\xi_{t+1} = 1 - \alpha_t \rho + \frac{2\alpha_t \mu}{\operatorname{dist}(x_{t+1},S)} > 1$ such that

$$\xi_{t+1} \operatorname{dist}^2(x_{t+1}, S) \le \operatorname{dist}^2(x_t, S).$$
 (56)

Given the lower-bound for α_t , we now define

$$\xi := 1 + \underline{\alpha} \left(\frac{2\mu}{\operatorname{dist}(x_{t_0}, S)} - \rho \right) > 1, \tag{57}$$

which yields the following inequality for every $t \ge t_0$

$$\xi_{t+1} = 1 + \alpha_t \left(\frac{2\mu}{\operatorname{dist}(x_{t+1}, S)} - \rho \right) > 1 + \alpha_t \left(\frac{2\mu}{\operatorname{dist}(x_{t_0}, S)} - \rho \right)$$

$$\geqslant 1 + \underline{\alpha} \left(\frac{2\mu}{\operatorname{dist}(x_{t_0}, S)} - \rho \right) = \xi.$$

Combining the latter with (56), we have that for every $t \ge t_0$

$$\xi \operatorname{dist}^{2}(x_{t+1}, S) < \xi_{t+1} \operatorname{dist}^{2}(x_{t+1}, S) \leq \operatorname{dist}^{2}(x_{t}, S)$$

hence the following "contraction" inequality holds

$$\operatorname{dist}^{2}(x_{t+1}, S) < \frac{1}{\xi}\operatorname{dist}^{2}(x_{t}, S)$$

which by recursiveness yields that for every $t \ge t_0$

$$\operatorname{dist}^{2}(x_{t+1}, S) < \left(\frac{1}{\xi}\right)^{t+1-t_{0}} \operatorname{dist}^{2}(x_{t_{0}}, S).$$
(58)

For $t \to \infty$ we obtain (55) and this concludes the proof of the first part of the statement.

Let us now show the strong convergence of the sequence $(x_t)_{t\in\mathbb{N}}$. We recall inequality (31) from the proof of Proposition 5 with $\varepsilon_t = 0$ and $\eta_t = 0$ for every $t \in \mathbb{N}$ (remembering our technical assumption that $1/\eta_t = 0$ whenever $\eta_t = 0$, see Assumption 2):

$$-\mu \operatorname{dist}(x_{t+1}, S) + \frac{1}{2\alpha_t} \operatorname{dist}^2(x_t, S) \geqslant \left(\frac{1}{2\alpha_t} - \frac{\rho}{2}\right) \operatorname{dist}^2(x_{t+1}, S) + \left(\frac{1}{2\alpha_t} - \frac{L_g}{2}\right) \|x_t - x_{t+1}\|^2.$$

Multiplying by $2\alpha_t$ we get

$$-2\alpha_t \mu \operatorname{dist}(x_{t+1}, S) + \operatorname{dist}^2(x_t, S) \ge (1 - \rho \alpha_t) \operatorname{dist}^2(x_{t+1}, S) + (1 - L_g \alpha_t) \|x_t - x_{t+1}\|^2.$$

Let now $t \ge t_0$. From the inequality above we infer

$$(1 - \alpha_t L_g) \|x_t - x_{t+1}\|^2 \le \operatorname{dist}^2(x_t, S) - \operatorname{dist}^2(x_{t+1}, S) + \underbrace{\alpha_t \rho \operatorname{dist}^2(x_{t+1}, S) - 2\mu \alpha_t \operatorname{dist}(x_{t+1}, S)}_{\le 0}$$

where for every $t \ge t_0$, by virtue of Corollary 3 we have

$$\alpha_t \operatorname{dist}(x_{t+1}, S) (\rho \operatorname{dist}(x_{t+1}, S) - 2\mu) < 0.$$

Since all the terms depending on x_{t+1} are negative we infer the following upper bound for $(1 - \alpha_t L_q) \|x_t - x_{t+1}\|^2$:

$$(t \ge t_0)$$
 $(1 - \alpha_t L_g) \|x_t - x_{t+1}\|^2 \le \text{dist}^2(x_t, S)$

On the other side, given the upper-bound for α_t , we obtain the following lower bound for $(1 - \alpha_t L_g) \|x_t - x_{t+1}\|^2$:

$$(\forall t \in \mathbb{N}) \qquad (1 - \overline{\alpha} L_g) \|x_t - x_{t+1}\|^2 \le (1 - \alpha_t L_g) \|x_t - x_{t+1}\|^2.$$

In conclusion, by taking $\xi > 1$ from (57), for every $t > t_0$ we have

$$||x_t - x_{t+1}|| \leq \frac{\operatorname{dist}(x_t, S)}{\sqrt{1 - \alpha_t L_g}} \leq \frac{\operatorname{dist}(x_t, S)}{\sqrt{1 - \overline{\alpha} L_g}} < \frac{1}{\sqrt{1 - \overline{\alpha} L_g}} \left(\frac{1}{\sqrt{\xi}}\right)^{t - t_0} \operatorname{dist}(x_{t_0}, S),$$

where the last inequality is a consequence of (58). In conclusion, we obtain

$$\sum_{t=0}^{\infty} ||x_t - x_{t+1}|| < +\infty$$

from which we infer that the sequence $(x_t)_{t\in\mathbb{N}}$ is a Cauchy sequence, hence it converges to some $x^* \in \mathcal{X}$ (see [13, Theorem 1]). Eventually, we infer that $x^* \in S$ due to the fact that S is closed (as it is the level set of a lower semi-continuous function) and $\operatorname{dist}(x_t, S) \to 0$ as $t \to +\infty$.

Remark 9 (Strong Convergence in the exact convex case). Under the sharpness condition, for convex function ($\rho = 0$) with exact proximal calculation ($\varepsilon = 0$), starting from (32) one can arrive at

$$2\alpha\mu\operatorname{dist}(x_{t+1},S) \leqslant \operatorname{dist}^2(x_t,S) - \operatorname{dist}^2(x_{t+1},S).$$

Taking the sum for t = 0, ..., T yields

$$2\alpha\mu \sum_{t=0}^{T} \operatorname{dist}(x_{t+1}, S) \leqslant \sum_{t=0}^{T} \left(\operatorname{dist}^{2}(x_{t}, S) - \operatorname{dist}^{2}(x_{t+1}, S) \right) = \operatorname{dist}^{2}(x_{0}, S)$$

hence for $T \to \infty$

$$2\alpha\mu \sum_{t=0}^{\infty} \operatorname{dist}(x_{t+1}, S) \leqslant \operatorname{dist}^{2}(x_{0}, S) < +\infty$$

which implies $\operatorname{dist}(x_t, S) \to 0$ as $t \to \infty$. Hence, we obtain the conclusion in [23, Theorem 3.4(d)] which is the strong convergence of the sequence $(x_t)_{t\in\mathbb{N}}$ to a solution.

8 Feasibility Problem

We are interested in finding a point in the intersection of a closed convex set and the unit sphere centered in the origin (feasibility problem). Feasibility problems (FP) between spheres and one cone appear in source localization and phase retrieval, [39]. (FP) between a sphere and line in Hilbert spaces have been considered in [16], [8], [15] where a non convex Douglas-Rachford algorithm is proposed to solve (FP). The objective of [4] is to derive formula for the projection operator of the intersection of a sphere and a cone, a closed

form is available in some rare cases. In [27], the authors proposed four different schemes to model the feasibility problem between two sets Q and C, possibly non convex. One of these schemes is

$$\underset{x \in \mathcal{X}}{\text{minimize}} \quad \frac{\text{dist}^2(x, Q)}{2} + \frac{\text{dist}^2(x, C)}{2}.$$

In this section, we show that there exists a weakly convex function f which can replace $\operatorname{dist}^2(x,Q)/2$ when Q is the unit sphere. Then, the feasibility problem between a sphere and a closed convex set can be modelled as

$$\underset{x \in \mathcal{X}}{\text{minimize}} \quad f(x) + \frac{\text{dist}^2(x, C)}{2}. \tag{FP}$$

Moreover, we show that problem (FP) is a particular case of (1) and it satisfies all the assumptions that guarantee the convergence of the weakly convex forward-backward algorithm proposed in this paper. In particular, the objective function of (FP) is sharp.

We denote the unit sphere and the unit ball centered in the origin respectively with S and B. We can define a non-negative weakly convex function $f: \mathcal{X} \to (-\infty, +\infty]$, such that

$$\min_{x \in \mathcal{X}} f(x) = 0 \text{ and } \underset{x \in \mathcal{X}}{\operatorname{argmin}} f(x) = \mathcal{S}.$$
 (59)

Let $x \in \mathcal{X}\setminus\{0\}$, the projection of x on the unit sphere \mathcal{S} is $x/\|x\|$. We define $f: \mathcal{X} \to (-\infty, +\infty]$ as

$$(\forall x \in \mathcal{X}) \qquad f(x) := \begin{cases} |\|x\|^2 - \|\frac{x}{\|x\|}\|^2| & x \neq 0 \\ 1 & x = 0 \end{cases} \Rightarrow f(x) := |\|x\|^2 - 1|$$

Hence, $f: \mathcal{X} \to (-\infty, +\infty]$, $x \mapsto |||x||^2 - 1|$ has the optimal value in 0 and satisfies (59). In fact, for $x \in \mathcal{X} \setminus \{0\}$, f(x) is the absolute value of the difference of the norms of x and its projection on \mathcal{S} ,

$$(\forall x \in \mathcal{X} \setminus \{0\})$$
 $f(x) = |\|x\|^2 - \|\frac{x}{\|x\|}\|^2| = |\|x\|^2 - 1|.$

Moreover, $f(0) = 1 = \text{dist}(0, \mathcal{S})$. Then,

$$\underset{x \in \mathcal{X}}{\operatorname{argmin}} \ f(x) = \{ x \in \mathcal{X} \mid ||x|| = 1 \} = \mathcal{S}.$$

In the following, we prove that f(x) is weakly convex and sharp.

Lemma 5. Function f is 2-weakly convex.

Proof. We can rewrite f as

$$(\forall x \in \mathcal{X}) \qquad f(x) := \begin{cases} \|x\|^2 - 1 & x \notin \mathcal{B} \\ 0 & x \in \mathcal{S} \\ 1 - \|x\|^2 & x \in \mathcal{B} \setminus \mathcal{S}. \end{cases}$$

Hence, $f(\cdot) + ||\cdot||^2$ reads as

$$(\forall x \in \mathcal{X}) f(x) + ||x||^2 := \begin{cases} 2||x||^2 - 1 & x \notin \mathcal{B} \\ 1 & x \in \mathcal{S} \\ 1 & x \in \mathcal{B} \setminus \mathcal{S}. \end{cases} = \max\{2||x||^2 - 1, 1\}.$$

In conclusions $f(\cdot) + \|\cdot\|^2$ is convex as it is the point-wise maximum of convex functions.




Figure 1: Comparison between function f (continuous line) and the distance from S (dashed line) for $\mathcal{X} = \mathbb{R}$.

Lemma 6. Function f satisfies the sharpness condition with constant $\mu = 1$.

Proof. We need to show that for $\mu = 1$ the following holds:

$$(\forall x \in \mathcal{X}) \qquad \qquad |\|x\|^2 - 1| \geqslant \mu \operatorname{dist}(x, S).$$

If x = 0, then $\operatorname{dist}(x, \mathcal{S}) = 1$ and the inequality is satisfied for $\mu = 1$. Otherwise, for every $x \in \mathcal{X} \setminus 0$ we have

$$\min_{s \in \mathcal{S}} \|x - s\| = \|x - \frac{x}{\|x\|}\| = \|x\left(1 - \frac{1}{\|x\|}\right)\| = \|x\| \left|1 - \frac{1}{\|x\|}\right| = \|x\| \frac{1}{\|x\|} |\|x\| - 1| = |\|x\| - 1|$$

where the third equality stems from the absolute homogeneity of a norm. We consider the function $f: \mathbb{R} \to [0, +\infty)$ defined as $f(\sigma) := |\sigma^2 - 1|$ which corresponds to the function in (2) with (a, b) = (-1, 1). In this particular setting, it is easy to show that the following inequality is satisfied for every $\delta \leq 1$ (as also illustrated in Figure 1)

$$(\forall \sigma \in \mathbb{R}) \qquad \qquad \delta||\sigma| - 1| = \delta \min\{|\sigma - 1|, |\sigma + 1|\} \leqslant |\sigma^2 - 1|.$$

where the last inequality stems from the sharpness of (2). Choosing $\sigma = ||x||$ we obtain

$$(\forall x \in \mathcal{X} \setminus 0) \qquad \delta|\|x\| - 1| \leqslant |\|x\|^2 - 1|.$$

To conclude, we showed that for every $\mu \in (0,1]$

$$(\forall x \in \mathcal{X}) \qquad \qquad \mu \operatorname{dist}(x, \mathcal{S}) \leqslant f(x)$$

which implies the global sharpness of function f with constant $\mu = 1$.

In the previous part of this section, we proved that f(x) can be a weakly convex function modelling the constraint related to the unit sphere in the feasibility problem between a sphere and a closed convex set C. To model the constraint with respect to the closed convex set C we consider the differentiable function

$$(\forall x \in \mathcal{X})$$

$$g(x) = \frac{\operatorname{dist}^2(x, C)}{2},$$

whose properties are summarised in the following lemma:




Figure 2: Comparison.

Lemma 7 ([5, Corollary 12.30, Example 13.5]). Let $C \subseteq \mathcal{X}$ be a convex set. Then

$$(\forall x \in \mathcal{X}) \qquad \frac{\operatorname{dist}^{2}(x, C)}{2} = \frac{\|x\|^{2}}{2} - \left(\frac{\|\cdot\|^{2}}{2} + \iota_{C}\right)^{*}(x)$$

where $x \mapsto \left(\frac{\|\cdot\|^2}{2} + \iota_C\right)^*(x)$ denotes the Fenchel conjugate of function $x \mapsto \left(\frac{\|\cdot\|^2}{2} + \iota_C\right)(x)$ [5, Definition 13.1]. If in addition C is closed, then function $\operatorname{dist}^2(\cdot, C)$ is Fréchet differentiable on \mathcal{X} and

$$(\forall x \in \mathcal{X})$$

$$\nabla \frac{\operatorname{dist}^{2}(x, C)}{2} = x - \operatorname{proj}_{C}(x).$$

In particular, we highlight that the gradient of $x \mapsto \frac{\text{dist}^2(x,C)}{2}$ is Lipschitz continuous as being the sum of two Lipschitz continuous functions.

The feasibility problem becomes

minimize
$$|||x||^2 - 1| + \frac{\operatorname{dist}^2(x, C)}{2}$$

where the objective function is non-convex. As a matter of facts, according to Lemma 7, the problem can be recast as

minimize
$$|||x||^2 - 1| + \frac{||x||^2}{2} - \left(\frac{||\cdot||^2}{2} + \iota_C\right)^* (x).$$

where function $x \mapsto |\|x\|^2 - 1| + \frac{\|x\|^2}{2}$ is 1-weakly convex by virtue of Lemma 5 and $x \mapsto -\left(\frac{\|\cdot\|^2}{2} + \iota_C\right)^*(x)$ is the negative of a conjugate function, hence it is concave since a Fenchel conjugate function is always convex [5, Proposition 13.13]).

Eventually, the sharpness of f + g, which is the core of our convergence result, holds by virtue of the following theorem, where $h_1 = f$ and $h_2 = g$.

Theorem 3. Let functions $h_1, h_2 : \mathcal{X} \to \mathbb{R}$ be proper and let the following assumptions be satisfied

(i) The global minimiser set S of $h_1 + h_2$ is equal to (or contained in) $S_1 \cap S_1$, if both nonempty, or $S \subset S_1$ where S_1, S_2 are the sets of the minimisers of h_1 and h_2 respectively.

- (ii) The optimal value of $h_1 + h_2$ is zero, $v_{opt} = 0$.
- (iii) $h_1(x) \ge 0, h_2(x) \ge 0$ for all x in a neighborhood of S.
- (iv) h_1 is sharp with respect to S locally or globally with constant $\mu > 0$ and $\inf_{x \in \mathcal{X}} h_1(x) = 0$.

Then $h_1 + h_2$ is locally (globally) sharp.

Proof. By (iv) we have that there exists $\delta_1 > 0$ such that

$$(\forall x \in B(S, \delta_1))$$
 $h_1(x) \ge \mu \operatorname{dist}(x, S).$

By (iii) we know that there exists $\delta_2 > 0$ such that for every $x \in B(S, \delta_2)$ function h_2 is non-negative, hence by taking $\delta = \min\{\delta_1, \delta_2\}$ we have

$$(\forall x \in B(S, \delta)) \qquad h_1(x) + h_2(x) \geqslant \mu \operatorname{dist}(x, S).$$

By (i) and (ii), the above inequality corresponds to the local sharpness of function $h_1 + h_2$.

9 Conclusions

In this paper we investigate the convergence properties of the exact and the inexact forward-backward (FB) algorithms for the minimisation of a function that is expressed as the sum of a weakly convex function and a convex function with Lipschitz continuous gradient. In the inexact case we inferred a convergence result that relies on the hypothesis that the accuracy level $\varepsilon > 0$ for the inexact proximal computation is kept constant throughout all the iterations. In order to carry out our analysis, we exploited the notion of proximal subdifferential and a calculus rule to compute the inexact proximal subdifferential of the sum two functions which controls the modulus of weak convexity ρ . It will be interesting in a future work to extend our results so as to take into account non-constant accuracy levels and non-convex smooth function.

Acknowledgments This work has been supported by the ITN-ETN project TraDE-OPT funded by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 861137. This work represents only the author's view and the European Commission is not responsible for any use that may be made of the information it contains.

References

- [1] ATENAS, F., SAGASTIZÀBAL, C., SILVA, P. J. S., AND SOLODOV, M. A unified analysis of descent sequences in weakly convex optimization, including convergence rates for bundle methods. *SIAM Journal on Optimization* 33, 1 (2023), 89–115.
- [2] Attouch, H., Bolte, J., and Svaiter, B. F. Convergence of descent methods for semi-algebraic and tame problems: proximal algorithms, forward–backward splitting, and regularized Gauss–Seidel methods. *Mathematical Programming* 137, 1 (2013), 91–129.

- [3] Barré, M., Taylor, A. B., and Bach, F. Principled analyses and design of first-order methods with inexact proximal operators. *Mathematical Programming* (2022), 1–46.
- [4] BAUSCHKE, H. H., BUI, M. N., AND WANG, X. Projecting onto the intersection of a cone and a sphere. SIAM Journal on Optimization 28, 3 (2018), 2158–2188.
- [5] BAUSCHKE, H. H., AND COMBETTES, P. L. Convex Analysis and Monotone Operator Theory in Hilbert Spaces, 2nd ed. Springer Publishing Company, Incorporated, 2017.
- [6] BAYRAM, I. On the convergence of the iterative shrinkage/thresholding algorithm with a weakly convex penalty. *IEEE Transactions on Signal Processing* 64, 6 (2015), 1597–1608.
- [7] BEDNARCZUK, E., BRUCCOLA, G., SCRIVANTI, G., AND TRAN, T. H. Calculus rules for proximal ε -subdifferentials and inexact proximity operators for weakly convex functions. $arXiv\ preprint\ arXiv:2211.14525\ (2022)$.
- [8] Benoist, J. The douglas—rachford algorithm for the case of the sphere and the line. Journal of Global Optimization 63 (2015), 363–380.
- [9] BERNARD, F., AND THIBAULT, L. Prox-regular functions in Hilbert spaces. *Journal of Mathematical Analysis and Applications* 303, 1 (2005), 1–14.
- [10] BÖHM, A., AND WRIGHT, S. J. Variable smoothing for weakly convex composite functions. *Journal of optimization theory and applications* 188, 3 (2021), 628–649.
- [11] BOLTE, J., DANIILIDIS, A., AND LEWIS, A. A nonsmooth morse–sard theorem for subanalytic functions. *Journal of mathematical analysis and applications 321*, 2 (2006), 729–740.
- [12] Bolte, J., Daniilidis, A., Ley, O., and Mazet, L. Characterizations of Łojasiewicz inequalities and applications. *Transactions of the American Mathematical Society 362*, 6 (2010), 3319–3363.
- [13] BOLTE, J., SABACH, S., AND TEBOULLE, M. Proximal alternating linearized minimization for nonconvex and nonsmooth problems. *Mathematical Programming* 146, 1-2 (2014), 459–494.
- [14] BONETTINI, S., LORIS, I., PORTA, F., PRATO, M., AND REBEGOLDI, S. On the convergence of a linesearch based proximal-gradient method for nonconvex optimization. *Inverse Problems* 33, 5 (2017), 055005.
- [15] BORWEIN, J. M., LINDSTROM, S. B., SIMS, B., SCHNEIDER, A., AND SKERRITT, M. P. Dynamics of the Douglas–Rachford method for ellipses and *p*-spheres. *Set-Valued and Variational Analysis* 26 (2018), 385–403.
- [16] BORWEIN, J. M., AND SIMS, B. The Douglas–Rachford algorithm in the absence of convexity. Fixed-point algorithms for inverse problems in science and engineering (2011), 93–109.

- [17] BOŢ, R. I., AND CSETNEK, E. R. An inertial forward-backward-forward primal-dual splitting algorithm for solving monotone inclusion problems. *Numerical Algorithms* 71, 3 (2016), 519–540.
- [18] Burke, J. V., and Ferris, M. C. Weak sharp minima in mathematical programming. SIAM Journal on Control and Optimization 31, 5 (1993), 1340–1359.
- [19] CANNARSA, P., AND SINESTRARI, C. Semiconcave functions, Hamilton-Jacobi equations, and optimal control, vol. 58. Springer Science & Business Media, 2004.
- [20] CHAUX, C., COMBETTES, P. L., PESQUET, J.-C., AND WAJS, V. R. A variational formulation for frame-based inverse problems. *Inverse Problems* 23, 4 (June 2007), 1495–1518.
- [21] CLARKE, F. H., LEDYAEV, Y. S., STERN, R. J., AND WOLENSKI, P. R. Non-smooth analysis and control theory, vol. 178. Springer Science & Business Media, 2008.
- [22] Combettes, P. L., and Vũ, B. C. Variable metric forward–backward splitting with applications to monotone inclusions in duality. *Optimization 63*, 9 (2014), 1289–1318.
- [23] COMBETTES, P. L., AND WAJS, V. R. Signal recovery by proximal forward-backward splitting. *Multiscale modeling & simulation* 4, 4 (2005), 1168–1200.
- [24] DAVIS, D., AND DRUSVYATSKIY, D. Stochastic model-based minimization of weakly convex functions. SIAM Journal on Optimization 29, 1 (2019), 207–239.
- [25] DAVIS, D., DRUSVYATSKIY, D., MACPHEE, K. J., AND PAQUETTE, C. Subgradient methods for sharp weakly convex functions. *Journal of Optimization Theory and Applications* 179, 3 (2018), 962–982.
- [26] DAVIS, D., AND GRIMMER, B. Proximally guided stochastic subgradient method for nonsmooth, nonconvex problems. *SIAM Journal on Optimization* 29, 3 (2019), 1908–1930.
- [27] GIBALI, A., SABACH, S., AND VOLDMAN, S. Non-convex split feasibility problems: models, algorithms and theory. *Open Journal of Mathematical Optimization* 1 (2020), 1–15.
- [28] HAN, S.-P., AND LOU, G. A parallel algorithm for a class of convex programs. SIAM Journal on Control and Optimization 26, 2 (1988), 345–355.
- [29] HIRIART-URRUTY, J.-B., AND LEMARÉCHAL, C. Convex analysis and minimization algorithms II: Advanced Theory and Bundle Methods, vol. 306. Springer science & business media, 2013.
- [30] JOHNSTONE, P. R., AND MOULIN, P. Faster subgradient methods for functions with hölderian growth. *Mathematical Programming* 180 (2017), 417–450.
- [31] JOURANI, A. Subdifferentiability and subdifferential monotonicity of γ -paraconvex functions. Control and Cybernetics 25 (1996).

- [32] Kurdyka, K. On gradients of functions definable in o-minimal structures. *Annales de l'Institut Fourier* 48, 3 (1998), 769–783.
- [33] Li, G., and Pong, T. K. Calculus of the exponent of Kurdyka–Łojasiewicz inequality and its applications to linear convergence of first-order methods. *Foundations of computational mathematics* 18, 5 (2018), 1199–1232.
- [34] LIONS, P.-L., AND MERCIER, B. Splitting algorithms for the sum of two nonlinear operators. SIAM Journal on Numerical Analysis 16, 6 (1979), 964–979.
- [35] Liu, Q., Gu, Y., and So, H. C. D.O.A. estimation in impulsive noise via low-rank matrix approximation and weakly convex optimization. *IEEE Transactions on Aerospace and Electronic Systems* 55, 6 (2019), 3603–3616.
- [36] ŁOJASIEWICZ, S. Une propriété topologique des sous-ensembles analytiques réels. Equ. Derivees partielles, Paris 1962, Colloques internat. Centre nat. Rech. sci. 117, 87-89 (1963)., 1963.
- [37] ŁOJASIEWICZ, S. Sur la géométrie semi- et sous- analytique. Annales de l'Institut Fourier 43, 5 (1993), 1575–1595.
- [38] LORENZ, D. A., AND POCK, T. An inertial forward-backward algorithm for monotone inclusions. *Journal of Mathematical Imaging and Vision* 51, 2 (2015), 311–325.
- [39] Luke, D. R., Sabach, S., and Teboulle, M. Optimization on spheres: models and proximal algorithms with computational performance comparisons. *SIAM Journal on Mathematics of Data Science* 1, 3 (2019), 408–445.
- [40] Luo, Z.-Q. T., and Tseng, P. Error bounds and convergence analysis of feasible descent methods: a general approach. *Annals of Operations Research* 46-47 (1993), 157–178.
- [41] MILLÁN, R., AND PENTON, M. Inexact proximal ε -subgradient methods for composite convex optimization problems. *Journal of Global Optimization* 75 (2019).
- [42] MÖLLENHOFF, T., STREKALOVSKIY, E., MOELLER, M., AND CREMERS, D. The primal-dual hybrid gradient method for semiconvex splittings. *SIAM Journal on Imaging Sciences* 8, 2 (2015), 827–857.
- [43] MOUDAFI, A., AND OLINY, M. Convergence of a splitting inertial proximal method for monotone operators. *Journal of Computational and Applied Mathematics* 155, 2 (2003), 447–454.
- [44] NIKOLOVA, M. Estimation of binary images by minimizing convex criteria. In *Proceedings 1998 International Conference on Image Processing. ICIP98 (Cat. No. 98CB36269)* (1998), vol. 2, IEEE, pp. 108–112.
- [45] PASSTY, G. B. Ergodic convergence to a zero of the sum of monotone operators in Hilbert space. *Journal of Mathematical Analysis and Applications* 72, 2 (1979), 383–390.
- [46] POLYAK, B. Sharp minima, 1979. Institute of Control Sciences lecture notes, Moscow, USSR, In IIASA workshop on generalized Lagrangians and their applications, IIASA, Laxenburg, Austria, 1979.

- [47] POLYAK, B. T. Some methods of speeding up the convergence of iteration methods. User computational mathematics and mathematical physics 4, 5 (1964), 1–17.
- [48] RAFIQUE, H., LIU, M., LIN, Q., AND YANG, T. Weakly-convex—concave min—max optimization: provable algorithms and applications in machine learning. *Optimization Methods and Software 37*, 3 (2022), 1087–1121.
- [49] RASCH, J., AND CHAMBOLLE, A. Inexact first-order Primal–Dual algorithms. Computational Optimization and Applications 76, 2 (2020), 381–430.
- [50] ROCKAFELLAR, R., AND WETS, R. J.-B. Variational Analysis. Springer Verlag, Heidelberg, Berlin, New York, 1998.
- [51] ROLEWICZ, S. On paraconvex multifunctions. Oper. Res.-Verf. 31 (1979), 539–546.
- [52] ROLEWICZ, S. On a globalization property. Applicationes Mathematicae 22, 1 (1993), 69–73.
- [53] RUDIN, L. I., OSHER, S., AND FATEMI, E. Nonlinear total variation based noise removal algorithms. *Physica D: nonlinear phenomena 60*, 1-4 (1992), 259–268.
- [54] Salzo, S., and Villa, S. Inexact and accelerated proximal point algorithms. *Journal of Convex Analysis* 19, 4 (2012), 1167–1192.
- [55] SCHMIDT, M., ROUX, N., AND BACH, F. Convergence rates of inexact proximal-gradient methods for convex optimization. *Advances in neural information processing systems* 24 (2011).
- [56] SELESNICK, I., LANZA, A., MORIGI, S., AND SGALLARI, F. Non-convex total variation regularization for convex denoising of signals. *Journal of Mathematical Imaging and Vision 62*, 6 (2020), 825–841.
- [57] VILLA, S., SALZO, S., BALDASSARRE, L., AND VERRI, A. Accelerated and inexact forward-backward algorithms. SIAM Journal on Optimization 23, 3 (2013), 1607– 1633.

A Feasibility of the parameters

Lemma 8. Given ρ , μ and L_g , we denote $\gamma := \max\{\rho, L_g\}$. Then for every $t \in \mathbb{N}$, the following choice of ε_t , η_t and α_t ,

$$\begin{split} &\varepsilon_t \in (0, \frac{\mu^2}{2\gamma} \sqrt{\sqrt{2} - 1}) \\ &\tau_t \in (\frac{2\gamma \varepsilon_t}{\mu^2}, \min\left\{1, \frac{\mu^2}{2\gamma \varepsilon_t} \sqrt{\sqrt{2} - 1}\right\}) \\ &\Delta_{\eta} = \frac{\mu^4 \tau_t^2}{4\gamma_t^2} - \frac{\mu^2 \tau_t}{\gamma_t} \left(\varepsilon_t + 2\varepsilon_t\right) + \varepsilon_t^2 \\ &\underline{\eta}_t = \left(\frac{1}{2} - \frac{\gamma_t \varepsilon_t}{\mu^2 \tau_t}\right) - \frac{\gamma_t}{\mu^2 \tau_t} \sqrt{\Delta_{\eta}} \\ &\overline{\eta}_t = \min\left\{1, \left(\frac{1}{2} - \frac{\gamma_t \varepsilon_t}{\mu^2 \tau_t}\right) + \frac{\rho}{\mu^2 \tau_t} \sqrt{\Delta_{\eta}}\right\} \end{split}$$

$$\eta_t \in (\underline{\eta}_t, \overline{\eta}_t)$$

$$\alpha_t = \tau_t (1 - \eta_t) \frac{1}{\gamma_t}$$

satisfies the inequalities (19), (20) and (21) in Assumption 2.

Proof. For condition (19) to be valid, we need to have that for every $t \in \mathbb{N}$, $\eta_t \in (0,1)$. This implies the following condition on the stepsizes in $(\alpha_t)_{t\in\mathbb{N}}$

$$\frac{1}{\rho} > \frac{1 - \eta_t}{\rho} > \alpha_t$$

which combined with condition (21) implies that

$$\frac{2}{\alpha_t} > \rho + L_g,$$

that is the condition ensuring inequality (13) in Proposition 4.

On the other side, let us focus on condition (20) which reads as

$$\alpha_t \mu^2 \geqslant 2 \left(\varepsilon_t + \frac{\varepsilon_t}{\eta_t} \right) (\alpha_t \rho + \eta_t)$$

If $\varepsilon_t = 0$, then the condition is trivially satisfied. Let us then consider the case when $\varepsilon_t > 0$. From condition (19) we have that $\alpha_t \rho + \eta_t < 1$, so the LHS in (20) satisfies

$$2\left(\varepsilon_t + \frac{\varepsilon_t}{\eta_t}\right)\left(\alpha_t \rho + \eta_t\right) < 2\left(\varepsilon_t + \frac{\varepsilon_t}{\eta_t}\right)$$

Let us set $\gamma = \max\{\rho, L_g\}$ and choose $\alpha_t = \tau_t(1 - \eta_t)\frac{1}{\gamma}$ with $\tau_t \in (0, 1)$ so that (13), (19) and (21) are satisfied. In the following lines we will drop the dependency from t for simplicity. For condition (20) to be satisfied, it is sufficient that the following holds:

$$\eta \varepsilon + \varepsilon < \frac{\mu^2}{2} \alpha \eta$$

Since $\alpha = \tau \frac{1-\eta}{\gamma}$ we have

$$\begin{split} \eta \varepsilon + \varepsilon &< \frac{\mu^2}{2} \tau \left(\frac{1 - \eta}{\gamma} \right) \eta \\ &\iff \frac{\mu^2}{2} \tau \frac{\eta}{\gamma} - \frac{\mu^2}{2} \tau \frac{\eta^2}{\gamma} - \eta \varepsilon - \varepsilon > 0 \\ &\iff \frac{\mu^2}{2} \tau \frac{\eta^2}{\gamma} - \frac{\mu^2}{2} \tau \frac{\eta}{\gamma} + \varepsilon \eta + \varepsilon < 0 \\ &\iff \frac{\mu^2}{2} \tau \frac{\eta^2}{\gamma} + \left(\varepsilon - \frac{\mu^2 \tau}{2 \gamma} \right) \eta + \varepsilon < 0 \end{split}$$

This is a second order inequality in η and the possible solutions lie inside an interval having extrema that are defined by

$$\eta_{1,2} = \frac{-\left(\varepsilon - \frac{\mu^2 \tau}{2\gamma}\right) \pm \sqrt{\left(\varepsilon - \frac{\mu^2 \tau}{2\gamma}\right)^2 - 4\left(\frac{\mu^2 \tau}{2\gamma}\right)\varepsilon}}{\frac{\mu^2 \tau}{\gamma}}$$

In order to have positive values for η_1 and η_2 we need to require that τ is such that

$$\left(\varepsilon - \frac{\mu^2 \tau}{2\gamma}\right) \leqslant 0 \iff \frac{2\gamma \varepsilon}{\mu^2} \leqslant \tau$$

and $\frac{2\gamma\varepsilon}{\mu^2}$ can be ensured to be smaller than 1 if by assumption we take $\varepsilon < \frac{\mu^2}{2\gamma}\sqrt{\sqrt{2}-1}$. On the other side, we also need to choose τ so that Δ_{η} is non-negative, that is

$$\varepsilon^{2} - 2\frac{\mu^{2}\tau}{2\gamma}\varepsilon + \frac{\mu^{4}\tau^{2}}{4\gamma^{2}} - \frac{4\mu^{2}\tau}{2\gamma}\varepsilon \geqslant 0$$

$$\iff \frac{\mu^{4}\tau^{2}}{4\gamma^{2}} - \frac{\mu^{2}\tau}{\gamma}\left(\varepsilon + 2\varepsilon\right) + \varepsilon^{2} \geqslant 0.$$

We obtain a new inequality, this time in τ , which is related to a second order equation having solutions that are defined by

$$\tau_{1,2} = \frac{\frac{\mu^2(3\varepsilon)}{\gamma} \pm \sqrt{\left(\frac{\mu^2}{\gamma} 3\varepsilon\right)^2 - 4\varepsilon^2 \frac{\mu^4}{4\gamma^2}}}{2\varepsilon^2}$$

and we have that $\frac{\mu^2(3\varepsilon)}{\gamma} > 0$ and $\Delta_{\tau} = 9\frac{\mu^4}{\gamma^2}\varepsilon^2 - \frac{\mu^4}{\gamma^2}\varepsilon^2 = 8\frac{\mu^4}{\gamma^2}\varepsilon^2$. We then obtain

$$\tau \leqslant \tau_2 = \frac{\mu^2 3\varepsilon - 2\sqrt{2}\mu^2\varepsilon}{2\varepsilon^2\gamma} = \frac{\mu^2\varepsilon(\sqrt{2}-1)}{2\varepsilon^2\gamma} = \frac{\mu^2}{2\varepsilon\gamma}(\sqrt{2}-1)$$

In conclusion, to wrap up all the conditions we inferred on τ we have

$$\frac{2\gamma\varepsilon}{\mu^2} \leqslant \tau \leqslant \frac{\mu^2}{2\varepsilon\gamma}(\sqrt{2} - 1)$$

where we know that the lower bound is strictly smaller than 1. In order to have admissible values for τ we need the following to hold

$$\frac{2\gamma\varepsilon}{\mu^2} < \frac{\mu^2}{2\gamma\varepsilon}(\sqrt{2} - 1) \iff \varepsilon^2 < \frac{\mu^4}{\gamma^2 4}(\sqrt{2} - 1) \iff \varepsilon < \frac{\mu^2}{2\gamma}\sqrt{\sqrt{2} - 1}$$

which is satisfied by the hypothesis we mentioned above. In conclusion, we need to choose $\tau \in \left(\frac{2\gamma\varepsilon_t}{\mu^2}, \min\left\{1, \frac{\mu^2}{2\gamma\varepsilon_t}\sqrt{\sqrt{2}-1}\right\}\right)$.

Coming back to η , the condition we found for τ allows to state that

$$\frac{-\left(\varepsilon - \frac{\mu^2 \tau}{2\gamma}\right) - \sqrt{\Delta_{\eta}}}{\frac{\mu^2 \tau}{\gamma}} \leqslant \eta \leqslant \frac{-\left(\varepsilon - \frac{\mu^2 \tau}{2\gamma}\right) + \sqrt{\Delta_{\eta}}}{\frac{\mu^2 \tau}{\gamma}}$$

that is

$$\frac{\gamma}{\mu^2 \tau} \left(\frac{\mu^2 \tau}{2\gamma} - \varepsilon \right) - \frac{\gamma}{\mu^2 \tau} \sqrt{\Delta_{\eta}} \leqslant \eta \leqslant \frac{\gamma}{\mu^2 \tau} \left(\frac{\mu^2 \tau}{2\gamma} - \varepsilon \right) + \frac{\gamma}{\mu^2 \tau} \sqrt{\Delta_{\eta}}$$
$$\left(\frac{1}{2} - \frac{\gamma \varepsilon}{\mu^2 \tau} \right) - \frac{\gamma}{\mu^2 \tau} \sqrt{\Delta_{\eta}} \leqslant \eta \leqslant \left(\frac{1}{2} - \frac{\gamma \varepsilon}{\mu^2 \tau} \right) + \frac{\gamma}{\mu^2 \tau} \sqrt{\Delta_{\eta}}$$

and we have $0 < \left(\frac{1}{2} - \frac{\gamma \varepsilon}{\mu^2 \tau}\right) \leq 1$. We conclude that for the choice of τ we found above, we have $[\eta_1, \eta_2] \cap (0, 1) \neq \emptyset$ and we can define

$$\underline{\eta} = \left(\frac{1}{2} - \frac{\gamma \varepsilon}{\mu^2 \tau}\right) - \frac{\gamma}{\mu^2 \tau} \sqrt{\Delta_{\eta}} = \eta_1$$

$$\overline{\eta} = \min \left\{1, \left(\frac{1}{2} - \frac{\gamma \varepsilon}{\mu^2 \tau}\right) + \frac{\rho}{\mu^2 \tau} \sqrt{\Delta_{\eta}}\right\} = \min\{1, \eta_2\}$$

and choose $\eta \in (\eta, \overline{\eta}) \subset (0, 1)$.

To sum up, in order to satisfy the conditions (19), (20) and (21), we need to have for every $t \in \mathbb{N}$

$$\begin{split} \gamma &= \max\{\rho, L_g\} \\ \varepsilon_t &\in (0, \frac{\mu^2}{2\gamma} \sqrt{\sqrt{2} - 1}) \\ \tau_t &\in (\frac{2\gamma \varepsilon_t}{\mu^2}, \min\left\{1, \frac{\mu^2}{2\gamma \varepsilon_t} \sqrt{\sqrt{2} - 1}\right\}) \\ \eta_t &\in (\underline{\eta}_t, \overline{\eta}_t) \\ \alpha_t &= \tau_t (1 - \eta_t) \frac{1}{\gamma} \end{split}$$

B Discriminant from Lemma 3

Let us calculate the discriminant related to inequality (38), in order to define the existence of solutions.

$$\begin{split} &\Delta^{+} = \alpha^{2}\mu^{2} + (1 - \alpha\rho - \eta) \left(\left(\frac{\alpha\mu + \sqrt{\alpha^{2}\mu^{2} - 2\alpha \left(\varepsilon + \frac{\varepsilon}{\eta}\right) (\alpha\rho + \eta)}}{\alpha\rho + \eta} \right)^{2} + 2\alpha \left(\varepsilon + \frac{\varepsilon}{\eta}\right) 60) \right) \\ &= \frac{\alpha^{2}\mu^{2} (\alpha\rho + \eta)^{2}}{(\alpha\rho + \eta)^{2}} \\ &\quad + \frac{(1 - \alpha\rho - \eta) \left(\left(\alpha\mu + \sqrt{\alpha^{2}\mu^{2} - 2\alpha \left(\varepsilon + \frac{\varepsilon}{\eta}\right) (\alpha\rho + \eta)}\right)^{2} + 2\alpha \left(\varepsilon + \frac{\varepsilon}{\eta}\right) (\alpha\rho + \eta)^{2} \right)}{(\alpha\rho + \eta)^{2}} \\ &= \frac{\alpha^{2}\mu^{2} (\alpha\rho + \eta)^{2} + 2\alpha^{2}\mu^{2} (1 - \alpha\rho - \eta)}{(\alpha\rho + \eta)^{2}} - \frac{2\alpha \left(\varepsilon + \frac{\varepsilon}{\eta}\right) (\alpha\rho + \eta) (1 - \alpha\rho - \eta)}{(\alpha\rho + \eta)^{2}} \\ &\quad + \frac{2\alpha\mu\sqrt{\alpha^{2}\mu^{2} - 2\alpha \left(\varepsilon + \frac{\varepsilon}{\eta}\right) (\alpha\rho + \eta)} (1 - \alpha\rho - \eta)}{(\alpha\rho + \eta)^{2}} \\ &\quad + \frac{2\alpha \left(\varepsilon + \frac{\varepsilon}{\eta}\right) (\alpha\rho + \eta)^{2} (1 - \alpha\rho - \eta)}{(\alpha\rho + \eta)^{2}} \end{split}$$

$$= \frac{\alpha^{2}\mu^{2} + (1 - \alpha\rho - \eta)^{2} \left(\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)\right)}{(\alpha\rho + \eta)^{2}} + \frac{2\alpha\mu\left(1 - \alpha\rho - \eta\right)\sqrt{\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}}{(\alpha\rho + \eta)^{2}} = \left(\frac{\alpha\mu + (1 - \alpha\rho - \eta)\sqrt{\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}}{\alpha\rho + \eta}\right)^{2}.$$

Let us calculate the upper bound in equation (39)

$$\frac{-\alpha\mu + \sqrt{\Delta^{+}}}{1 - \alpha\rho - \eta} = \frac{-\alpha\mu + \frac{\alpha\mu + (1 - \alpha\rho - \eta)\sqrt{\alpha^{2}\mu^{2} - 2\alpha(\varepsilon + \frac{\varepsilon}{\eta})(\alpha\rho + \eta)}}{1 - \alpha\rho - \eta}}{1 - \alpha\rho - \eta}$$

$$= \frac{-\alpha\mu(\alpha\rho + \eta) + \alpha\mu + (1 - \alpha\rho - \eta)\sqrt{\alpha^{2}\mu^{2} - 2\alpha(\varepsilon + \frac{\varepsilon}{\eta})(\alpha\rho + \eta)}}{(1 - \alpha\rho - \eta)(\alpha\rho + \eta)}$$

$$= \frac{\alpha\mu(1 - \alpha\rho - \eta) + (1 - \alpha\rho - \eta)\sqrt{\alpha^{2}\mu^{2} - 2\alpha(\varepsilon + \frac{\varepsilon}{\eta})(\alpha\rho + \eta)}}{(1 - \alpha\rho - \eta)(\alpha\rho + \eta)}$$

$$= \frac{\alpha\mu + \sqrt{\alpha^{2}\mu^{2} - 2\alpha(\varepsilon + \frac{\varepsilon}{\eta})(\alpha\rho + \eta)}}{(\alpha\rho + \eta)}.$$
(61)

Let us calculate the discriminant (Δ^{-}) and the upper bound for the second part of

the statement, that is when $dist(x_{t_0}, S) < E^-$.

$$\Delta^{-} = \alpha^{2}\mu^{2} + (1 - \alpha\rho - \eta) \left(\left(\frac{\alpha\mu - \sqrt{\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}}{\alpha\rho + \eta} \right)^{2} + 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right) \right)$$

$$= \frac{\alpha^{2}\mu^{2}(\alpha\rho + \eta)^{2}}{(\alpha\rho + \eta)^{2}}$$

$$+ \frac{(1 - \alpha\rho - \eta)\left(\left(\alpha\mu - \sqrt{\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}\right)^{2} + 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)^{2} \right)}{(\alpha\rho + \eta)^{2}}$$

$$= \frac{\alpha^{2}\mu^{2}(\alpha\rho + \eta)^{2} + 2\alpha^{2}\mu^{2}(1 - \alpha\rho - \eta)}{(\alpha\rho + \eta)^{2}} - \frac{2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)(1 - \alpha\rho - \eta)}{(\alpha\rho + \eta)^{2}}$$

$$- \frac{2\alpha\mu\sqrt{\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)(1 - \alpha\rho - \eta)}}{(\alpha\rho + \eta)^{2}}$$

$$+ \frac{2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)^{2}(1 - \alpha\rho - \eta)}{(\alpha\rho + \eta)^{2}}$$

$$= \frac{\alpha^{2}\mu^{2} + (1 - \alpha\rho - \eta)^{2}\left(\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)\right)}{(\alpha\rho + \eta)^{2}}$$

$$- \frac{2\alpha\mu(1 - \alpha\rho - \eta)\sqrt{\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}}{(\alpha\rho + \eta)^{2}}$$

$$= \left(\frac{\alpha\mu - (1 - \alpha\rho - \eta)\sqrt{\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}}{\alpha\rho + \eta}\right)^{2}.$$
(62)

Let us calculate the upper bound

$$\frac{-\alpha\mu + \sqrt{\Delta^{-}}}{1 - \alpha\rho - \eta} = \frac{-\alpha\mu + \frac{\alpha\mu - (1 - \alpha\rho - \eta)\sqrt{\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}}{1 - \alpha\rho - \eta}}{1 - \alpha\rho - \eta}$$

$$= \frac{-\alpha\mu(\alpha\rho + \eta) + \alpha\mu - (1 - \alpha\rho - \eta)\sqrt{\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}}{(1 - \alpha\rho - \eta)(\alpha\rho + \eta)}$$

$$= \frac{\alpha\mu(1 - \alpha\rho - \eta) - (1 - \alpha\rho - \eta)\sqrt{\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}}{(1 - \alpha\rho - \eta)(\alpha\rho + \eta)}$$

$$= \frac{\alpha\mu - \sqrt{\alpha^{2}\mu^{2} - 2\alpha\left(\varepsilon + \frac{\varepsilon}{\eta}\right)(\alpha\rho + \eta)}}{(\alpha\rho + \eta)}.$$
(63)