

# Newton's methods for solving linear inverse problems with neural network coders

Otmar Scherzer<sup>1,2,3</sup>  
otmar.scherzer@univie.ac.at

Bernd Hofmann<sup>4</sup>  
hofmamb@mathematik.tu-chemnitz.de

Zuhair Nashed<sup>5</sup>  
zuhair.nashed@ucf.edu

March 27, 2023

<sup>1</sup>Faculty of Mathematics  
University of Vienna  
Oskar-Morgenstern-Platz 1  
A-1090 Vienna, Austria

<sup>2</sup>Johann Radon Institute for Computational  
and Applied Mathematics (RICAM)  
Altenbergerstraße 69  
A-4040 Linz, Austria

<sup>3</sup>Christian Doppler Laboratory for  
Mathematical Modeling and Simulation  
of Next Generations of Ultrasound  
Devices (MaMSi)  
Oskar-Morgenstern-Platz 1  
A-1090 Vienna, Austria

<sup>4</sup>Faculty of Mathematics  
Chemnitz University of Technology  
Reichenhainer Str. 39/41  
D-09107 Chemnitz, Germany

<sup>5</sup>College of Sciences  
University of Central Florida  
4393 Andromeda Loop N  
Orlando, FL 32816, USA

## Abstract

Neural networks functions are supposed to be able to encode the desired solution of an inverse problem very efficiently. In this paper, we consider the problem of solving linear inverse problems with neural network coders. First we establish some correspondences of this formulation with existing concepts in regularization theory, in particular with state space regularization, operator decomposition and iterative regularization methods. A Gauss-Newton's method is suitable for solving encoded linear inverse problems, which is supported by a local convergence result. The convergence studies, however, are not complete, and are based on a conjecture on linear independence of activation functions and its derivatives.

This paper is dedicated to Heinz Engl's 70th birthday.

## 1. INTRODUCTION

We start the discussion by considering a *general nonlinear operator equation*

$$N(\vec{p}) = \mathbf{y}, \quad (1.1)$$

where  $N : \vec{P} \rightarrow \mathbf{Y}$  is a nonlinear operator between Hilbert spaces  $\vec{P}$  and  $\mathbf{Y}$ . Particular emphasis is placed on the case when the numerical solution of [Equation 1.1](#) is ill-posed or ill-conditioned. In this situation regularization methods need to be implemented (see [\[14\]](#)) to guarantee stable solvability. Essentially two classes of methods exist as the basis for solving inverse problems numerically, which are *variational methods* (see for instance [\[14, 44, 22, 45, 3, 32\]](#)) and *iterative methods* (see for instance [\[29, 4\]](#)).

Modern *coding theory* (see for instance [\[6, 13, 7, 43, 6\]](#)) assumes that *natural images* can be represented efficiently by a combination of *neural network* functions. Therefore the *set of natural images* is given by the range of a nonlinear mapping  $\Psi$ , which maps neural network coefficients to images. In coding theory very often the considered operator equation is assumed to be *linear* and *ill-posed*. Let us consider therefore the equation

$$F\mathbf{x} = \mathbf{y}, \quad (1.2)$$

where  $F : \mathbf{X} \rightarrow \mathbf{Y}$  is a bounded linear operator with non-closed range mapping between the Hilbert spaces  $\mathbf{X}$  and  $\mathbf{Y}$  such that  $\mathbf{x}$  and  $\mathbf{y}$  are elements of  $\mathbf{X}$  and  $\mathbf{Y}$ , respectively. Combining this with the assumption

---

2000 *Mathematics Subject Classification*: 65J22, 47J07

*Keywords*: Gauss-Newton's method, inverse problems, neural networks

that the solution of Equation 1.2 is a natural image, or in other words that it can be represented as a combination of neural network functions, we get the operator equation

$$N(\vec{p}) = F\Psi(\vec{p}) = \mathbf{y}, \tag{1.3}$$

where  $\Psi : \vec{P} \rightarrow \mathbf{X}$  is the before mentioned nonlinear operator that maps neural network parameters to image functions. We call

- $\mathbf{X}$  the *image space* and
- $\mathbf{Y}$  the *data space*, in accordance with a terminology, which we generated in [2, 1].
- $\vec{P}$  is called the *parameter space*. We make a different notation for  $\vec{P}$ , because it represents parametrizations, and is often considered a space of vectors below.

The advantage of this ansatz is that the solution of Equation 1.2 is sparsely coded. However, the price to pay is that the reformulated Equation 1.3 is nonlinear. Operator equations of the form of Equation 1.3 are not new: They have been studied in abstract settings for instance in the context of

- *state space regularization* [8] and
- in the context of the *degree of ill-posedness* [15, 20, 21, 24, 25] as well as of the *degree of nonlinearity* [24] of nonlinear ill-posed operator equations.
- Another related approach is *finite dimensional approximation* of regularization in Banach spaces (see for instance [42]). Particularly, finite dimensional approximations of regularization methods with neural network functions (in the context of frames and *deep synthesis regularization*) have been studied in [38].

In this paper we aim to link the general regularization of the degree of ill-posedness and nonlinearity with coding theory. We investigate generalized Gauss-Newton’s methods for solving Equation 1.3; Such methods replace the inverse of standard Newton’s method by approximations of outer inverses (see [37]).

The outline of the paper is as follows: In Section 2 we review two decomposition cases as stated in [21] first, one of them is Equation 1.3. The study of decomposition cases follows the work on classifying inverse problems and regularization (see [33]). For operators associated to Equation 1.3, Newton’s methods seem to be better suited than gradient descent methods, which we support by a convergence analysis (see Section 3). Section 3.4 is devoted to solving Equation 1.3, where  $\Psi$  is a shallow neural network synthesis operator.

## 2. DECOMPOSITION CASES

We start with a definition for nonlinear operator equations possessing forward operators that are compositions of a linear and a nonlinear operator. Precisely, we distinguish between a first decomposition case (i), where the linear operator is the inner operator and the nonlinear is the outer one, and a second decomposition case (ii), where the nonlinear operator is the inner operator and the linear is the outer operator.

**Definition 2.1 (Decomposition cases)** Let  $\vec{P}, \mathbf{X}, \mathbf{Y}$  be Hilbert-spaces.

- (i) An operator  $N$  is said to satisfy the *first decomposition case* in an open, non-empty neighborhood  $\mathcal{B}(\vec{p}^\dagger; \rho) \subseteq \vec{P}$  of some point  $\vec{p}^\dagger$  if there exists a linear operator  $F : \vec{P} \rightarrow \mathbf{X}$  and a nonlinear operator  $\Psi : \mathbf{X} \rightarrow \mathbf{Y}$  such that

$$N(\vec{p}) = \Psi(F\vec{p}) \text{ for } \vec{p} \in \mathcal{B}(\vec{p}^\dagger; \rho).$$

- (ii)  $N$  is said to satisfy the *second decomposition case* in a neighborhood  $\mathcal{B}(\vec{p}^\dagger; \rho) \subseteq \vec{P}$  of some point  $\vec{p}^\dagger$  if there exists a linear operator  $F : \mathbf{X} \rightarrow \mathbf{Y}$  and a nonlinear operator  $\Psi : \vec{P} \rightarrow \mathbf{X}$  such that

$$N(\vec{p}) = F\Psi(\vec{p}) \text{ for } \vec{p} \in \mathcal{B}(\vec{p}^\dagger; \rho). \tag{2.1}$$

Typically it is assumed that the nonlinear operator  $\Psi$  is well-posed.

**Remark 2.2 (First decomposition case)** In [21], this decomposition case has been studied under structural conditions, relating the second derivative of  $N$  with the first derivative. Under such assumptions convergence rates conditions (see [21, Lemma 4.1]) could be proven. The first decomposition case also arises in inverse option pricing problems in math finance (see [19] and [23, Sect.4]), where the ill-posed compact linear integration operator occurs as inner operator and a well-posed Nemytskii operator as outer operator.

**Remark 2.3 (Second decomposition case)** Regularization methods for solving operator equations with operators satisfying the second order decomposition case, see Equation 2.1, were probably first analyzed in [8] under the name of *state space regularization*. They considered for instance Tikhonov-type regularization methods, consisting in minimization of

$$J_\lambda(\vec{p}) = \|F\Psi(\vec{p}) - \mathbf{y}\|_{\mathbf{Y}}^2 + \lambda \|\Psi(\vec{p}) - \tilde{\mathbf{x}}\|_{\mathbf{X}}^2, \quad (2.2)$$

where  $\tilde{\mathbf{x}}$  is a prior and  $\lambda > 0$  is a regularization parameter. In [8] they derived estimates for the second derivative of  $J_\lambda(\vec{p}_\lambda)(\mathbf{h}, \mathbf{h})$ , where  $\mathbf{h} \in \vec{P}$ , that is for the curvature of  $J_\lambda$ . If the curvature can be bounded from below by some terms  $\|\mathbf{h}\|_{\vec{P}}^2$  then, for instance, a locally unique minimizer of  $J_\lambda$  can be guaranteed and also domains can be specified where the functional is convex. Conditions, which guarantee convexity are called *curvature to size conditions*. Subsequently, these decomposition cases have been studied exemplarily in [21]. The theory developed there directly applies to Equation 1.3.

Instead of  $J_\lambda$  researchers often study direct regularization with respect to  $\vec{p}$ . For instance in [13] functionals of the form

$$J_\lambda(\vec{p}) = \|F\Psi(\vec{p}) - \mathbf{y}\|_{\mathbf{Y}}^2 + \lambda\mathcal{L}(\vec{p}), \quad (2.3)$$

where  $\mathcal{L}$  is some functional directly regularizing the parameter space. Typically  $\mathcal{L}$  is chosen to penalize for sparsity of parameters. The main difference between Equation 2.2 and Equation 2.3 is that in the prior regularization is performed with respect to the image space  $\mathbf{X}$  and in the later with respect to the parameter space  $\vec{P}$ . Well-posedness of the functional  $J_\lambda$  in Equation 2.2 follows if  $F \circ \Psi$  is lower-semicontinuous, which in turn follows if  $\Psi$  is invertible.

In the following we study the solution of decomposable operator equations, such as Equation 1.3, with Gauss-Newton's methods. Decomposition cases have been used in the analysis of iterative regularization methods as well (see [29]):

**Definition 2.4 (Strong tangential cone condition)** Let  $N : \mathcal{D}(N) \subset \vec{P} \rightarrow \mathbf{Y}$  with  $\mathcal{D}(N)$  its domain be a nonlinear operator.

(i) Then  $N$  is said to satisfy the strong tangential cone condition, originally introduced in [17], if

$$N'(\vec{p}_2) = R_{\vec{p}_2, \vec{p}_1} N'(\vec{p}_1) \text{ for all } \vec{p}_1, \vec{p}_2 \in \mathcal{D}(N). \quad (2.4)$$

where

$$\|R_{\vec{p}_2, \vec{p}_1} - I\| \leq C_T \|\vec{p}_2 - \vec{p}_1\|_{\vec{P}}. \quad (2.5)$$

(ii) In [5] the *order reversed* tangential cone condition,

$$N'(\vec{p}_2) = N'(\vec{p}_1) R_{\vec{p}_2, \vec{p}_1} \text{ for all } \vec{p}_1, \vec{p}_2 \in \mathcal{D}(N), \quad (2.6)$$

together with Equation 2.5, has been introduced.

**Remark 2.5** Equation 2.4 has been used for analyzing *gradient descent methods* (see for instance [17, 29]). For the analysis of Newton's methods Equation 2.6 has been used (see [5, 29]).

The relation to the decomposition cases is as follows:

**Lemma 2.6** Let  $N : \mathcal{D}(N) \subseteq \vec{P} \rightarrow \mathbf{Y}$  with  $\mathcal{D}(N) = \mathcal{B}(\vec{p}^\dagger; \rho)$  satisfy the second decomposition case and assume that  $\Psi'(\vec{p})$  is invertible for  $\vec{p} \in \mathcal{D}(N)$ . Then  $N$  satisfies Equation 2.6.

*Proof:* If  $N$  satisfies the second decomposition case, Equation 2.1, then  $N'(\vec{p}) = F\Psi'(\vec{p})$  for all  $\vec{p} \in \mathcal{D}(N)$ . Note, that because  $F$  is defined on the whole space  $\mathbf{X}$ ,  $\mathcal{D}(N) = \mathcal{D}(\Psi)$ . By using the invertibility assumption on  $\Psi$  we get

$$N'(\vec{p}_2) = F\Psi'(\vec{p}_2) = F\Psi'(\vec{p}_1) \underbrace{\Psi'(\vec{p}_1)^{-1}\Psi'(\vec{p}_2)}_{=: R_{\vec{p}_2, \vec{p}_1}} = N'(\vec{p}_1) R_{\vec{p}_2, \vec{p}_1},$$

which gives the assertion.  $\square$

As we have shown, decomposition cases have been extensively studied in the regularization literature. One conclusion out of these studies is that the order reversed tangential cone condition Equation 2.6 is suitable for analyzing Newton's methods [5, 29] and thus in turn for the coded linear operator Equation 1.3 because of Lemma 2.6. The standard tool for analyzing Newton's methods is the Newton-Mysovskii condition as discussed below.

### 3. THE NEWTON-MYSOVSKII CONDITIONS

In this section we put abstract convergence conditions for Newton type methods in context with decoding. We consider first Newton's methods for solving the *general* operator Equation 1.1. Decomposition cases of the operator  $N$  will be considered afterwards.

**3.1. Newton's method with invertible linearizations.** For Newton's methods *local convergence* is guaranteed under *Newton-Mysovskii* conditions. For comparison reasons, we first recall a simple Newton's method analysis in finite dimensional spaces if the nonlinear operator has derivatives which are invertible. The proof of more general results, such as Theorem 3.6 below, applies here as well, and thus here the proof is omitted. Several variants of Newton-Mysovskii conditions have been proposed in the literature (see for instance [11, 12, 37]). Analysis of Newton's method was an active research area in the last century, see for instance [39, 46].

**Theorem 3.1 (Finite dimensional Newton's method)** *Let  $N : \mathcal{D}(N) \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^n$  be continuously Fréchet-differentiable on a non-empty, open and convex set  $\mathcal{D}(N)$ . Let  $\vec{p}^\dagger \in \mathcal{D}(N)$  be a solution of Equation 1.1. Moreover, we assume that*

- (i)  $N'(\vec{p})$  is invertible for all  $\vec{p} \in \mathcal{D}(N)$  and that
- (ii) the Newton-Mysovskii condition holds: That is, there exist some  $C_N > 0$  such that

$$\|N'(\vec{q})^{-1}(N'(\vec{p} + s(\vec{q} - \vec{p})) - N'(\vec{p}))(\vec{q} - \vec{p})\|_{\vec{p}} \leq sC_N \|\vec{p} - \vec{q}\|_{\vec{p}}^2 \quad (3.1)$$

for all  $\vec{p}, \vec{q} \in \mathcal{D}(N), s \in [0, 1]$ .

Let  $\vec{p}^0 \in \mathcal{D}(N)$  which satisfies

$$\overline{\mathcal{B}(\vec{p}^0; \rho)} \subseteq \mathcal{D}(N) \text{ with } \rho := \|\vec{p}^\dagger - \vec{p}^0\|_{\vec{p}} \text{ and } h := \frac{\rho C_I C_L}{2} < 1. \quad (3.2)$$

Then the Newton's iteration with starting point  $\vec{p}^0$ ,

$$\vec{p}^{k+1} = \vec{p}^k - N'(\vec{p}^k)^{-1}(N(\vec{p}^k) - \mathbf{y}) \quad k \in \mathbb{N}_0, \quad (3.3)$$

satisfies that the iterates  $\{\vec{p}^k : k = 0, 1, 2, \dots\}$  belong to  $\overline{\mathcal{B}(\vec{p}^0; \rho)}$  and converge quadratically to  $\vec{p}^\dagger \in \overline{\mathcal{B}(\vec{p}^0; \rho)}$ .

Now, we turn to the case that  $N$  is a decomposition operator.

**3.2. Newton-Mysovskii conditions with composed operator.** Now, we study the case of convergence of Gauss-Newton's methods where  $N : \vec{P} \rightarrow \mathbf{Y}$  with  $\vec{P} = \mathbb{R}^{n^*}$  and  $\mathbf{Y}$  is an infinite dimensional Hilbert space, where  $F : \mathbf{X} \rightarrow \mathbf{Y}$  is linear and bounded and  $\Psi : \vec{P} = \mathbb{R}^{n^*} \rightarrow \mathbf{X}$ . In this case the Moore-Penrose inverse, or even more general the outer inverse, replaces the inverse in a classical Newton's method (see Equation 3.3), because linearizations of  $N$  will not be invertible anymore as a simple count of dimensions show. We refer now to Gauss-Newton's methods if the linearizations might not be invertible to distinguish between classical Newton's methods also by name.

Before we phrase a convergence result for Gauss-Newton's methods we recall and introduce some definitions:

**Notation 3.2 (Inner, outer and Moore-Penrose inverse)** (see [36, 34]) Let  $L : \vec{P} \rightarrow \mathbf{Y}$  be a linear and bounded operator mapping between two vector spaces  $\vec{P}$  and  $\mathbf{Y}$ . Then

(i) the operator  $B : \mathbf{Y} \rightarrow \vec{P}$  is called a *left inverse* to  $L$  if

$$BL = I .$$

(ii)  $B : \mathbf{Y} \rightarrow \vec{P}$  is called a *right inverse* to  $L$  if

$$LB = I .$$

Left and right inverses are used in different context:

- For a left inverse the nullspace of  $L$  has to be trivial, in contrast to  $B$ .
- For a right inverse the nullspace of  $B$  has to be trivial.

(iii)  $B : \vec{P} \rightarrow \vec{P}$  is called a *inverse* to  $L$  if  $B$  is a right and a left inverse.

(iv)  $B : \vec{P} \rightarrow \mathbf{Y}$  is an *outer inverse* to  $L$  if

$$BLB = B. \quad (3.4)$$

(v) Let  $\vec{P}$  and  $\mathbf{Y}$  be Hilbert-spaces,  $L : \vec{P} \rightarrow \mathbf{Y}$  be a linear bounded operator. We denote the orthogonal projections  $P$  and  $Q$  onto  $\mathcal{N}(L)$ , the nullspace of  $L$  (which is closed), and  $\overline{\mathcal{R}(L)}$ , the closure of the range of  $L$ : That is for all  $\vec{p} \in \vec{P}$  and  $\mathbf{y} \in \mathbf{Y}$  we have

$$P\vec{p} = \operatorname{argmin} \{ \|\vec{p}_1 - \vec{p}\|_{\vec{P}} : \vec{p}_1 \in \mathcal{N}(L) \} \quad \text{and} \quad Q\mathbf{y} = \operatorname{argmin} \{ \|\mathbf{y}_1 - \mathbf{y}\|_{\mathbf{Y}} : \mathbf{y}_1 \in \overline{\mathcal{R}(L)} \}. \quad (3.5)$$

We therefore have

$$\begin{aligned} P : \vec{P} &\rightarrow \mathcal{N}(L) \dot{+} \mathcal{N}(L)^\perp & \text{and} & \quad Q : \mathbf{Y} \rightarrow \overline{\mathcal{R}(L)} \dot{+} \mathcal{R}(L)^\perp \\ \vec{p} &\mapsto P\vec{p} + 0 & & \quad \mathbf{y} \mapsto Q\mathbf{y} + 0 \end{aligned}$$

$B : \mathcal{D}(B) \subseteq \mathbf{Y} \rightarrow \vec{P}$  with  $\mathcal{D}(B) := \overline{\mathcal{R}(L)} \dot{+} \mathcal{R}(L)^\perp$  is called the *Moore-Penrose inverse* of  $L$  if the following identities hold

$$\begin{aligned} LBL &= L, \\ BLB &= B, \\ BL &= I - P, \\ LB &= Q|_{\mathcal{D}(B)}. \end{aligned} \quad (3.6)$$

In coding theory it is often stated that the range of a neural network operator  $\Psi$  forms a manifold in  $\mathbf{X}$ , a space, which contains the natural images. This is the basis of the following definition making use of the Moore-Penrose inverse.

**Definition 3.3 (Lipschitz-differentiable immersion)** Let  $\Psi : \mathcal{D}(\Psi) \subseteq \vec{P} = \mathbb{R}^{n_*} \rightarrow \mathbf{X}$  where  $\mathcal{D}(\Psi)$  is open, non-empty, convex and  $\mathbf{X}$  is a separable (potentially infinite dimensional) Hilbert-space.

(i) We assume that  $\mathcal{M} := \Psi(\mathcal{D}(\Psi))$  is a  $n_*$ -dimensional *submanifold* in  $\mathbf{X}$ :

- Let for all  $\vec{p} = (p_i)_{i=1}^{n_*} \in \mathcal{D}(\Psi)$  denote with  $\Psi'(\vec{p})$  the Fréchet-derivative of  $\Psi$ :

$$\begin{aligned} \Psi'(\vec{p}) &: \vec{P} \rightarrow \mathbf{X}, \\ \vec{q} = (q_i)_{i=1}^{n_*} &\mapsto (\partial_{p_i} \Psi(\vec{p}))_{i=1, \dots, n_*} \vec{q}. \end{aligned}$$

Here  $(\partial_{p_i} \Psi(\vec{p}))_{i=1, \dots, n_*}$  denotes the vector of functions consisting of all partial derivatives with respect to  $\vec{p}$ . In differential geometry notation this coincides with the *tangential mapping*  $T_{\vec{p}}\Psi$ . However, the situation is slightly different here because  $\mathbf{X}$  can be infinite dimensional.

- The *representation mapping* of the derivative

$$\begin{aligned} \Psi' &: \mathcal{D}(\Psi) \rightarrow \mathbf{X}^{n_*}, \\ \vec{p} &\mapsto (\partial_{p_i} \Psi(\vec{p}))_{i=1, \dots, n_*}. \end{aligned}$$

has always the same rank  $n_*$  in  $\mathcal{D}(\Psi)$ , meaning that all elements of  $\partial_{\vec{p}}\Psi(\vec{p})$  are linearly independent. This assumption means, in particular, that  $\Psi$  is an *immersion* and  $\mathcal{M}$  is a submanifold.

(ii) We define

$$\begin{aligned} P_{\vec{p}} : \mathbf{X} &\rightarrow \mathbf{X}_{\vec{p}} := \text{span} \{ \partial_{p_i} \Psi(\vec{p}) : i = 1, \dots, n_* \}, \\ \mathbf{x} &\mapsto P_{\vec{p}} \mathbf{x} := \text{argmin} \{ \| \mathbf{x}_1 - \mathbf{x} \|_{\mathbf{X}} : \mathbf{x}_1 \in \mathbf{X}_{\vec{p}} \} \end{aligned} \quad (3.7)$$

as the projection from

$$\mathbf{X} = \mathbf{X}_{\vec{p}} \dot{+} \mathbf{X}_{\vec{p}}^{\perp}$$

onto  $\mathbf{X}_{\vec{p}}$ , which is well-defined by the closedness of the finite dimensional subspace  $\mathbf{X}_{\vec{p}}$ .

Next we define the inverse of  $\Psi'(\vec{p})$  on  $\mathbf{X}_{\vec{p}}$ :

$$\begin{aligned} \Psi'(\vec{p})^{-1} : \text{span} \{ \partial_{p_i} \Psi(\vec{p}) : i = 1, \dots, n_* \} &\rightarrow \vec{P}, \\ \mathbf{x} = \sum_{i=1}^{n_*} x_i \partial_{p_i} \Psi(\vec{p}) &\mapsto (x_i)_{i=1}^{n_*} \end{aligned}$$

and consequently on  $\mathbf{X}$

$$\begin{aligned} \Psi'(\vec{p})^{\dagger} : \mathbf{X} = \mathbf{X}_{\vec{p}} \dot{+} \mathbf{X}_{\vec{p}}^{\perp} &\rightarrow \vec{P}, \\ \mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2) &\mapsto \Psi'(\vec{p})^{-1} \mathbf{x}_1 \end{aligned} \quad (3.8)$$

which are both well-defined because we assume that  $\Psi$  is an immersion. Note that  $x_i, i = 1, \dots, n_*$  are not necessarily the coordinates with respect to an orthonormal system in  $\text{span} \{ \partial_{p_i} \Psi(\vec{p}) : i = 1, \dots, n_* \}$ .

(iii) Finally, we assume that the operators  $\Psi'(\vec{p})$  are locally bounded and locally Lipschitz-continuous in  $\mathcal{D}(\Psi)$ . That is

$$\| \Psi'(\vec{p}) - \Psi'(\vec{q}) \|_{\vec{P} \rightarrow \mathbf{X}} \leq C_L \| \vec{p} - \vec{q} \|_{\vec{P}} \quad \| \Psi'(\vec{p}) \|_{\vec{P} \rightarrow \mathbf{X}} \leq C_I \text{ for } \vec{p}, \vec{q} \in \mathcal{D}(\Psi). \quad (3.9)$$

If  $\Psi$  satisfies these three properties we call it a *Lipschitz-differentiable immersion*.

The following lemma is proved by standard means:

**Lemma 3.4** *For a Lipschitz-differentiable immersion*

- the function  $\Psi'(\vec{p})^{\dagger} : \mathbf{X} \rightarrow \vec{P}$  is in fact the Moore-Penrose inverse of  $\Psi'(\vec{p})$  and
- for every point  $\vec{p} \in \mathcal{D}(\Psi) \subseteq \vec{P}$  there exists a non-empty closed neighborhood where  $\Psi'(\vec{p})^{\dagger}$  is uniformly bounded and it is Lipschitz-continuous; That is

$$\| \Psi'(\vec{p})^{\dagger} - \Psi'(\vec{q})^{\dagger} \|_{\mathbf{X} \rightarrow \vec{P}} \leq C_L \| \vec{p} - \vec{q} \|_{\vec{P}}, \quad \| \Psi'(\vec{p})^{\dagger} \|_{\mathbf{X} \rightarrow \vec{P}} \leq C_I \text{ for } \vec{p}, \vec{q} \in \mathcal{D}(\Psi). \quad (3.10)$$

- Moreover, the operator  $P_{\vec{p}}$  from Equation 3.7 is bounded.

*Proof:* • We verify the four conditions Equation 3.6 with

- $L = \Psi'(\vec{p}) : \vec{P} = \mathbb{R}^{n_*} \rightarrow \mathbf{X}$ ,  $B = \Psi'(\vec{p})^{\dagger} : \mathbf{X} \rightarrow \vec{P}$ , with  $\mathcal{D}(B) = \mathcal{D}(\Psi'(\vec{p})^{\dagger}) = \mathbf{X}$  and
- $P : \vec{P} \rightarrow \vec{P}$  the zero-operator and  $Q = P_{\vec{p}} : \mathbf{X} \rightarrow \mathbf{X}_{\vec{p}}$ , the projection operator onto  $\mathbf{X}_{\vec{p}}$  (see Equation 3.7).
- First we prove the third identity with  $P = 0$  in Equation 3.6: This follows from the fact that for all  $\vec{q} = (q_i)_{i=1}^{n_*} \in \vec{P}$  we have

$$\Psi'(\vec{p})^{\dagger} \Psi'(\vec{p}) \vec{q} = \Psi'(\vec{p})^{-1} \left( \sum_{i=1}^{n_*} q_i \partial_{p_i} \Psi(\vec{p}) \right) = (q_i)_{i=1}^{n_*} = \vec{q}. \quad (3.11)$$

- For the fourth identity we see that for all  $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2) \in \mathbf{X}$  there exists  $x_i, i = 1, \dots, n_*$  (because  $\partial_{p_i} \Psi(\vec{p}), i = 1, \dots, n_*$  is a basis) such that

$$\mathbf{x} = \sum_{i=1}^{n_*} x_i \partial_{p_i} \Psi(\vec{p}) + \mathbf{x}_2 \text{ with } \mathbf{x}_2 \in \mathbf{X}_{\vec{p}}^{\perp}$$

and thus

$$P_{\vec{p}} \mathbf{x} = \sum_{i=1}^{n_*} x_i \partial_{p_i} \Psi(\vec{p})$$

and therefore

$$\Psi'(\vec{p})^{\dagger} \mathbf{x} = (x_i)_{i=1}^{n_*} = \vec{x}. \quad (3.12)$$

Consequently, we have

$$\Psi'(\vec{p}) \Psi'(\vec{p})^{\dagger} \mathbf{x} = \Psi'(\vec{p}) \vec{x} = P_{\vec{p}} \mathbf{x}. \quad (3.13)$$

- For the second identity we use that for all  $\mathbf{x} \in \mathbf{X}$

$$\underbrace{\Psi'(\vec{p})^\dagger \Psi'(\vec{p}) \Psi'(\vec{p})^\dagger \mathbf{x}}_{\text{Equation 3.12}} \stackrel{=}{=} \underbrace{\Psi'(\vec{p})^\dagger \Psi'(\vec{p}) (\vec{x})}_{\text{Equation 3.11}} \stackrel{=}{=} \underbrace{\vec{x}}_{\text{Equation 3.12}} \stackrel{=}{=} \Psi(\vec{p})^\dagger \mathbf{x}. \quad (3.14)$$

- The first identity is proven analogously.

Thus the Moore-Penrose inverse exists.

- For the proof of the boundedness we argue as follows: First note that

$$\begin{aligned} \|\Psi'(\vec{p})^\dagger\|_{\mathbf{X} \rightarrow \bar{\mathbf{P}}} &= \sup_{\{\mathbf{x} \in \mathbf{X} : \|\mathbf{x}\|_{\mathbf{X}}=1\}} \|\Psi'(\vec{p})^\dagger \mathbf{x}\|_{\bar{\mathbf{P}}} = \sup_{\{\mathbf{x}_1 \in \mathbf{X}_{\bar{\mathbf{P}}} : \|\mathbf{x}_1\|=1\}} \|\Psi'(\vec{p})^{-1} \mathbf{x}_1\|_{\bar{\mathbf{P}}} \\ &= \sup_{\{\mathbf{x}_1 \in \mathbf{X}_{\bar{\mathbf{P}}} : \|\mathbf{x}_1\|=1\}} \|\vec{x}_1\|_{\bar{\mathbf{P}}}, \end{aligned}$$

where we used the representation

$$\mathbf{x}_1 = \sum_{i=1}^{n_*} x_i \mathbf{e}_i \in \mathbf{X}_{\bar{\mathbf{P}}} \text{ with } \mathbf{E} := (\mathbf{e}_1 = \partial_{p_1} \Psi(\vec{p}), \dots, \mathbf{e}_{n_*} = \partial_{p_{n_*}} \Psi(\vec{p})). \quad (3.15)$$

It remains to derive a uniform bound and the continuity for  $\|\vec{x}_1\|_{\bar{\mathbf{P}}}$  with respect to variations of  $\vec{p}$ . For this purpose we use the Gram-Schmidt inductive procedure to obtain the QR-decomposition of the function valued matrix  $\mathbf{E}$ .

- Because of the assumption [Equation 3.9](#), each vector  $\mathbf{e}_j$ ,  $j = 1, \dots, n_*$ , depends continuously of  $\vec{p}$ .
- The Gram-Schmidt procedure performs only additions, multiplications and division and inner products of vectors of functions, which are all locally Lipschitz-continuous with respect to perturbations as long as the vectors are linearly independent. Thus by chain-rule Gram-Schmidt is Lipschitz-continuous in the matrix entries as long as the vectors  $\mathbf{e}_j$ ,  $j = 1, \dots, n_*$  are linearly dependent. Gram-Schmidt produces an orthogonal family  $\hat{\mathbf{e}}_j$ ,  $j = 1, \dots, n_*$ , which therefore are also Lipschitz-continuous with respect to  $\vec{p}$ .
- Calculating the QR-decomposition of  $\mathbf{E}$  we get

$$\mathbf{x}_1 = \vec{x}^T \mathbf{E} = \vec{x}^T Q R \text{ where } R = \begin{pmatrix} \langle \hat{\mathbf{e}}_1, \mathbf{e}_1 \rangle_{\mathbf{Y}} & \cdots & \cdots & \cdots & \langle \hat{\mathbf{e}}_1, \mathbf{e}_{n_*} \rangle_{\mathbf{Y}} \\ 0 & \langle \hat{\mathbf{e}}_2, \mathbf{e}_2 \rangle_{\mathbf{Y}} & \cdots & \cdots & \langle \hat{\mathbf{e}}_2, \mathbf{e}_{n_*} \rangle_{\mathbf{Y}} \\ \vdots & \ddots & \ddots & & \\ \vdots & & \ddots & \ddots & \\ 0 & \cdots & \cdots & 0 & \langle \hat{\mathbf{e}}_{n_*}, \mathbf{e}_{n_*} \rangle_{\mathbf{Y}} \end{pmatrix},$$

or in other word

$$\mathbf{x}_1 R^{-1} Q^T = \vec{x}^T.$$

The QR-algorithm is also Lipschitz-continuous with respect to perturbations in the matrix entries, that with respect to  $\vec{p}$ , as long as the single vectors do not get linearly dependent. Therefore  $Q$  and  $R$  are Lipschitz-continuous as well, and therefore, since  $R$  is a triangular matrix, also  $R^{-1}$ . This guarantees a local uniformly bound and local Lipschitz-continuity for  $\Psi$ .

- The boundedness of  $P_{\bar{\mathbf{P}}}$  is obvious because it is an orthogonal projector.  $\square$

In the following we study a Gauss-Newton's method for solving [Equation 1.3](#), where  $\Psi : \mathcal{D}(\Psi) \subseteq \mathbb{R}^{n_*} \rightarrow \mathbf{X}$  is a Lipschitz-continuous immersion (see [Definition 3.3](#)),  $F : \mathbf{X} \rightarrow \mathbf{Y}$  is bounded and  $N = F \circ \Psi$ .

**Lemma 3.5** *Let  $F : \mathbf{X} \rightarrow \mathbf{Y}$  be linear, bounded, with trivial nullspace and dense range. Moreover, let  $\Psi : \mathcal{D}(\Psi) \subseteq \mathbb{R}^{n_*} \rightarrow \mathbf{X}$  be a Lipschitz-differentiable immersion; see [Definition 3.3](#), and let  $N = F \circ \Psi$ .*

*Then  $\mathcal{D}(N) = \mathcal{D}(\Psi)$  and for every  $\vec{p} \in \mathcal{D}(N)$  the derivative of the operator  $N$  at a point  $\vec{p}$  has a Moore-Penrose inverse  $N'(\vec{p})^\dagger$ , which satisfies:*

- *Decomposition property of the Moore-Penrose inverse:*

$$N'(\vec{p})^\dagger \mathbf{z} = \Psi'(\vec{p})^\dagger F^{-1} \mathbf{z} \text{ for all } \vec{p} \in \mathcal{D}(N), \mathbf{z} \in \mathcal{R}(F) \subseteq \mathbf{Y}. \quad (3.16)$$

In particular this means that

$$N'(\vec{p})^\dagger N'(\vec{p}) = I \text{ on } \mathbb{R}^{n^*} \text{ and } N'(\vec{p})N'(\vec{p})^\dagger = Q|_{\mathcal{R}(FP_{\vec{p}})}, \quad (3.17)$$

where  $I$  denotes the identity operator on  $\mathbb{R}^{n^*}$  and  $Q : \mathbf{Y} \rightarrow \overline{\mathcal{R}(FP_{\vec{p}})} \dot{+} \mathcal{R}(FP_{\vec{p}})^\perp$ , respectively.

- *Generalized Newton-Mysovskii condition:*

$$\begin{aligned} \|N'(\vec{p})^\dagger(N'(\vec{q} + s(\vec{p} - \vec{q}) - N'(\vec{q}))(\vec{p} - \vec{q}))\|_{\vec{p}} &\leq sC_I C_L \|\vec{p} - \vec{q}\|_{\vec{p}}^2 \\ \vec{p}, \vec{q} &\in \mathcal{D}(N), s \in [0, 1]. \end{aligned} \quad (3.18)$$

We recall that the Lipschitz-constants  $C_I$  and  $C_L$  are defined in Equation 3.9.

*Proof:* First of all, we note that

$$N'(\vec{p}) = F\Psi'(\vec{p}) \text{ on } \mathcal{D}(\Psi) = \mathcal{D}(N).$$

To prove Equation 3.16 we have to verify Equation 3.6 with

$$L := N'(\vec{p}) = F\psi'(\vec{p}) : \vec{P} \rightarrow \mathbf{Y} \text{ and } B := \Psi'(\vec{p})^\dagger F^{-1} : \mathcal{R}(F) \subseteq \mathbf{Y} \rightarrow \vec{P}.$$

Note that since we assume that  $F$  has dense range we do not need to define and consider  $B$  on  $\mathcal{R}(F) \dot{+} \underbrace{\mathcal{R}(F)^\perp}_{=\{0\}}$ .

Let us first state that with the notation of Equation 3.6 we have for fixed  $\vec{p}$ :

$$\mathcal{D}(B) = \mathcal{D}(\Psi'(\vec{p})^\dagger F^{-1}) = \mathcal{R}(F) \text{ and } \mathcal{R}(L) = \{F\Psi'(\vec{p})\vec{q} : \vec{q} \in \mathbb{R}^{n^*}\} = \mathcal{R}(FP_{\vec{p}}).$$

We use  $P \equiv 0$  in Equation 3.6.

In particular the first item shows that for  $\mathbf{z} = F\mathbf{x} = FP_{\vec{p}}\mathbf{x} + F(I - P_{\vec{p}})\mathbf{x}$  we have

$$Q\mathbf{z} = Q(FP_{\vec{p}}\mathbf{x} + F(I - P_{\vec{p}})(\mathbf{x})) = FP_{\vec{p}}\mathbf{x}. \quad (3.19)$$

Applying Lemma 3.4 and the invertability of  $F$  on the range of  $F$  shows that

$$\begin{aligned} LBL &= F\Psi'(\vec{p})\Psi'(\vec{p})^\dagger F^{-1}F\Psi'(\vec{p}) = F\Psi'(\vec{p})\underbrace{\Psi'(\vec{p})^\dagger \Psi'(\vec{p})}_{\text{Equation 3.11}} = F\Psi'(\vec{p}) = L, \\ BLB &= \Psi'(\vec{p})^\dagger F^{-1}F\Psi'(\vec{p})\Psi'(\vec{p})^\dagger F^{-1} = \Psi'(\vec{p})^\dagger \underbrace{\Psi'(\vec{p})\Psi'(\vec{p})^\dagger}_{\text{Equation 3.14}} F^{-1} = B, \\ BL &= \Psi'(\vec{p})^\dagger F^{-1}F\Psi'(\vec{p}) \underbrace{=}_{\text{Equation 3.11}} I - P = I \text{ on } \mathbb{R}^{n^*}. \end{aligned}$$

For the forth identity we take some  $\mathbf{z} \in \mathcal{R}(F)$ . Therefore, there exists some  $\mathbf{x} \in \mathbf{X}$  such that  $F\mathbf{x} = \mathbf{z}$  and thus

$$LB\mathbf{z} = F\Psi'(\vec{p})\Psi'(\vec{p})^\dagger F^{-1}\mathbf{z} = F\Psi'(\vec{p})\underbrace{\Psi'(\vec{p})^\dagger \mathbf{z}}_{\text{Equation 3.13}} \underbrace{=}_{\text{Equation 3.19}} Q\mathbf{z}.$$

Thus Equation 3.6 holds and we know that  $N'(\vec{p})^\dagger$  from Equation 3.16 is the Moore-Penrose of  $N'(\vec{p})$ . The last two identities in fact show Equation 3.17.

From Equation 3.9 it follows that

$$\begin{aligned} \|N'(\vec{p})^\dagger(N'(\vec{q} + s(\vec{p} - \vec{q}) - N'(\vec{q}))(\vec{p} - \vec{q}))\|_{\vec{p}} &= \|\Psi'(\vec{p})^\dagger F^{-1}(F\Psi'(\vec{q} + s(\vec{p} - \vec{q}) - F\Psi'(\vec{q}))(\vec{p} - \vec{q}))\|_{\vec{p}} \\ &= \|\Psi'(\vec{p})^\dagger(\Psi'(\vec{q} + s(\vec{p} - \vec{q}) - \Psi'(\vec{q}))(\vec{p} - \vec{q}))\|_{\vec{p}} \\ &\leq C_I C_L s \|\vec{p} - \vec{q}\|_{\vec{p}}^2 \text{ for all } \vec{p}, \vec{q} \in \mathcal{D}(\Psi) = \mathcal{D}(N), \end{aligned}$$

thus Equation 3.18.  $\square$



We have now all ingredients to prove a local convergence rates result for a Gauss-Newton's method, where the operator  $N$  is the composition of a linear bounded operator and a Lipschitz-differentiable immersions:

**Theorem 3.6** *Let  $F : \mathbf{X} \rightarrow \mathbf{Y}$  be linear, bounded, with trivial nullspace and dense range. Moreover, let  $\Psi : \mathcal{D}(\Psi) \subseteq \bar{P} \rightarrow \mathbf{X}$  be a Lipschitz-differentiable immersion with  $\mathcal{D}(\Psi)$  open, non-empty, and convex. Moreover,  $N = F \circ \Psi : \mathcal{D}(\Psi) \rightarrow \mathbf{Y}$ . We assume that there exist  $\bar{p}^\dagger \in \mathcal{D}(\Psi)$  that satisfies*

$$N(\bar{p}^\dagger) = \mathbf{y}. \quad (3.20)$$

Moreover, we assume that there exists  $\bar{p}^0 \in \mathcal{D}(\Psi)$ , which satisfies Equation 3.2. Then, the iterates of the Gauss-Newton's iteration,

$$\bar{p}^{k+1} = \bar{p}^k - N'(\bar{p}^k)^\dagger(N(\bar{p}^k) - \mathbf{y}) \quad k \in \mathbb{N}_0 \quad (3.21)$$

are well-defined elements in  $\overline{\mathcal{B}(\bar{p}^0, \rho)}$  and converge quadratically to  $\bar{p}^\dagger$ .

*Proof:* First of all note, that  $\mathcal{D}(\Psi) = \mathcal{D}(N)$  since  $F$  is defined all over  $\mathbf{X}$ .

Let  $\rho = \|\bar{p}^\dagger - \bar{p}^0\|_{\bar{P}}$ : We prove by induction that  $\bar{p}^k \in \overline{\mathcal{B}(\bar{p}^\dagger; \rho)}$  for all  $k \in \mathbb{N}_0$ .

- For  $k = 0$  the assertion is satisfied by assumption Equation 3.2.
- Let  $\bar{p}^k \in \overline{\mathcal{B}(\bar{p}^\dagger; \rho)}$ . Using the first condition of Equation 3.6, which a Moore-Penrose inverse satisfies, we see that

$$N'(\bar{p}^k)N'(\bar{p}^k)^\dagger N'(\bar{p}^k)(\bar{p}^{k+1} - \bar{p}^\dagger) = N'(\bar{p}^k)(\bar{p}^{k+1} - \bar{p}^\dagger).$$

The definition of Gauss-Newton's method, Equation 3.21, and Equation 3.20 then imply that

$$N'(\bar{p}^k)(\bar{p}^{k+1} - \bar{p}^\dagger) = N'(\bar{p}^k)N'(\bar{p}^k)^\dagger(N(\bar{p}^\dagger) - N(\bar{p}^k) - N'(\bar{p}^k)(\bar{p}^\dagger - \bar{p}^k)),$$

and consequently, using the third identity of Equation 3.6 (note that under the assumptions of this theorem  $P = 0$ , see the proof prior to Equation 3.11), the second identity of Equation 3.6 and that  $F$  is injective, we get

$$\begin{aligned} \bar{p}^{k+1} - \bar{p}^\dagger &= N'(\bar{p}^k)^\dagger N'(\bar{p}^k)(\bar{p}^{k+1} - \bar{p}^\dagger) = N'(\bar{p}^k)^\dagger(N(\bar{p}^\dagger) - N(\bar{p}^k) - N'(\bar{p}^k)(\bar{p}^\dagger - \bar{p}^k)) \\ &= \Psi'(\bar{p}^k)^\dagger(\Psi(\bar{p}^\dagger) - \Psi(\bar{p}^k) - \Psi'(\bar{p}^k)(\bar{p}^\dagger - \bar{p}^k)). \end{aligned}$$

From the Newton-Mysovskii condition Equation 3.18 and Equation 3.2 it then follows that

$$\begin{aligned} \|\bar{p}^{k+1} - \bar{p}^\dagger\|_{\bar{P}} &\leq \frac{C_I C_L}{2} \|\bar{p}^k - \bar{p}^\dagger\|_{\bar{P}}^2 \leq \frac{C_I C_L \rho}{2} \|\bar{p}^k - \bar{p}^\dagger\|_{\bar{P}} < \|\bar{p}^k - \bar{p}^\dagger\|_{\bar{P}} \\ \text{or } \|\bar{p}^{k+1} - \bar{p}^\dagger\|_{\bar{P}} &= \|\bar{p}^k - \bar{p}^\dagger\|_{\bar{P}} = 0. \end{aligned} \quad (3.22)$$

This, in particular shows that  $\bar{p}^{k+1} \in \mathcal{B}(\bar{p}^\dagger; \rho)$ , thus the well-definedness of the Gauss-Newton's iterations in the closed ball.

- Using Equation 3.22 we then get, since  $h = C_I C_L \rho / 2 < 1$ , that

$$\|\bar{p}^{k+1} - \bar{p}^\dagger\|_{\bar{P}} \leq h^{k+1} \|\bar{p}^0 - \bar{p}^\dagger\|_{\bar{P}} \leq h^{k+1} \rho,$$

which converges to 0 for  $k \rightarrow \infty$ .

- Convergence and the first inequality of Equation 3.22 imply quadratic convergence.  $\square$

**Remark 3.7** Based on the assumption of an immersion we have shown in Lemma 3.5 that  $\Psi(\bar{p})^\dagger F^{-1}$  is the Moore-Penrose inverse of  $N = F\Psi(\bar{p})$ . In order to prove (quadratic) convergence of Gauss-Newton's methods one only requires an *outer inverse* (see Notation 3.2). Following [37] (see also [18]) the analysis of Gauss-Newton's method could be based on *outer inverses*, which is more general than for the Moore-Penrose inverse (compare Equation 3.4 and Equation 3.6). However, it is nice to actually see that  $N'(\bar{p})^\dagger$  is a Moore-Penrose inverse, which is the novelty compared to the analysis of [37]. For excellent expositions on Kantorovich and Mysovskii theory see [30, 40, 46] - here we replace the Newton-Mysovskii conditions by properties of an immersion. For aspects related to Newton's methods for singular points see [10, 16]. For applications of generalized inverses in nonlinear analysis see [34, 35].

**3.3. Neural networks.** We want to apply the decomposition theory to Gauss-Newton's methods for solving Equation 1.3, where  $\Psi$  is a *shallow neural network operator*.

**Definition 3.8 (Shallow neural network operator)** Let  $N \in \mathbb{N}$  be fixed. We consider the operator

$$\Psi : \vec{P} := \mathbb{R}^N \times \mathbb{R}^{n \times N} \times \mathbb{R}^N \rightarrow C^1([0, 1]^n) \subseteq \mathbf{X} := L^2([0, 1]^n),$$

$$(\vec{\alpha}, \mathbf{w}, \vec{\theta}) \mapsto \left( \vec{x} \mapsto \sum_{j=1}^N \alpha_j \sigma(\mathbf{w}_j^T \vec{x} + \theta_j) \right) \quad (3.23)$$

where  $\alpha_j, \theta_j \in \mathbb{R}$  and  $\vec{x}, \mathbf{w}_j \in \mathbb{R}^n$ .

Note, that with our previous notation, for instance in Definition 3.3, we have  $n_* = (n + 2) * N$ .

We summarize the notation, because it is quite heavy:

- (i)  $\vec{\cdot}$  denotes a vector in  $\mathbb{R}^n$  or  $\mathbb{R}^N$ ,
- (ii)  $\mathbf{w}$  denotes a matrix: The only exception is Definition 3.10, where it is a tensor.  $\mathbf{w}_j$  denotes a vector, aside from Definition 3.10, where it is again a tensor.

**Example 3.9 (Examples of activation functions)**  $\sigma$  is called the *activation function*, such as

- the *sigmoid function*, defined by

$$\sigma(t) = \frac{1}{1 + e^{-\frac{1}{\varepsilon}t}} \text{ for all } t \in \mathbb{R}. \quad (3.24)$$

Note, we omit the  $\varepsilon$  dependence for notational convenience.

- The *hyperbolic tangent*

$$t \rightarrow \tanh(t) = \frac{e^{2t} - 1}{e^{2t} + 1}. \quad (3.25)$$

- The *ReLU* activation function,

$$\sigma(t) = \max\{0, t\} \text{ for all } t \in \mathbb{R}. \quad (3.26)$$

- The *step function*, which is the pointwise limit of the sigmoid function, with respect to  $\varepsilon \rightarrow 0$ ,

$$\sigma(t) = \begin{cases} 0 & \text{for } t < 0, \\ \frac{1}{2} & \text{for } t = 0, \\ 1 & \text{for } t > 0. \end{cases} \quad (3.27)$$




FIGURE 1. Three different activation functions: Sigmoid, tanh and ReLU. 0-th derivative green, first derivative pink with circles and with pink  $\times$  is the derivative multiplied by  $x$ , 2nd derivative blue. The ReLU function is scaled by a factor 1/10 and the derivative is plotted in original form, derivative times  $x$  is again scaled by 1/10 for visualization purposes.

We only consider shallow neural networks in contrast to *deep neural networks*, which consist of several layers of shallow neural networks (see for instance [26]):

**Definition 3.10 (Deep neural networks)** Let

$$\vec{P}_l := \mathbb{R}^{N_l} \times \mathbb{R}^{n \times N_l} \times \mathbb{R}^{N_l} \text{ for } l = 1, \dots, L \text{ and } \vec{P} := \prod_{l=1}^L \vec{P}_l.$$

Then a deep neural network consisting of  $L$  layers is written as

$$\begin{aligned} \Psi : \vec{P} &\rightarrow L^2([0, 1]^n), \\ (\vec{\alpha}_l, \mathbf{w}_l, \vec{\theta}_l)_{l=1}^L &\mapsto \left( \vec{x} \rightarrow \sum_{j_L=1}^{N_L} \alpha_{j_L}^L \sigma_{\varepsilon_L}^L \left( p_{j_L, L} \left( \sum_{j_{L-1}=1}^{N_{L-1}} \cdots \left( \sum_{j_1=1}^{N_1} \alpha_{j_1, 1} \sigma_{\varepsilon_1}^1 (p_{j_1}^1(\vec{x})) \right) \right) \right) \right), \end{aligned} \quad (3.28)$$

where

$$p_j^i(\vec{x}) = \mathbf{w}_{j,i}^T \vec{x} + \theta_j^i \text{ with } \alpha_j^i, \theta_j^i \in \mathbb{R} \text{ and } \vec{x}, \mathbf{w}_{j,i} \in \mathbb{R}^n \text{ for all } i = 1, \dots, L.$$

Note that the values  $\varepsilon_k$ ,  $k = 1, \dots, L$  can be chosen differently for activation functions at different levels (cf. Equation 3.24).

The success of neural network is due to the universal approximation properties, proven for the first time in [9, 27]. The universal approximation result states that shallow neural networks are universal, that is, that each continuous function can be approximated arbitrarily well by a neural network function. We review this result now.

**Theorem 3.11 ([26])** *In dependence of the smoothness of the activation function  $\sigma$  there exist two classes of results.*

- *Theorem 2 from [26]: Let  $\sigma : \mathbb{R} \rightarrow \mathbb{R}^+$  be a **continuous, bounded and nonconstant** function. Then, for every function  $g \in C(\mathbb{R}^n)$  and every  $\nu > 0$ , there exists a function*

$$\vec{x} \rightarrow G(\vec{x}) = \sum_{j=1}^N \alpha_j \sigma(\mathbf{w}_j^T \vec{x} + \theta_j) \quad \text{with } N \in \mathbb{N}, \alpha_j, \theta_j \in \mathbb{R}, \mathbf{w}_j \in \mathbb{R}^n, \quad (3.29)$$

satisfying

$$|G(\vec{x}) - g(\vec{x})| < \nu \text{ uniformly for all compact subsets } K \subseteq \mathbb{R}^n.$$

- *Theorem 1 from [26]: Let  $\sigma : \mathbb{R} \rightarrow \mathbb{R}^+$  be **unbounded and nonconstant**. Then for every measure  $\mu$  on  $\mathbb{R}^n$  and every constant  $\nu > 0$  and  $p \geq 1$  there exists a function  $G$  of the form Equation 3.29 that satisfies*

$$\int_{\mathbb{R}^n} |G(\vec{x}) - g(\vec{x})|^p d\vec{x} < \nu.$$

The first result applies for instance to the sigmoid and hyperbolic tangent function (see Equation 3.24 and Equation 3.25). The second result applies to the ReLU function (see Equation 3.26). In particular all approximation properties also hold on the compact set  $[0, 1]^n$ , which we are considering.

**3.4. Newton-Mysovskii condition with neural networks.** In the following we verify Newton-Mysovskii conditions for  $\Psi$  being the encoder of Equation 3.23. First we calculate the first and second derivatives of  $\Psi$  with respect to  $\vec{\alpha}$ ,  $\mathbf{w}$  and  $\vec{\theta}$ . The computations can be performed for deep neural network encoders as defined in Equation 3.28 in principle analogously, but are technically and notationally more complicated. To make the notation consistent we define

$$\vec{p} := (\vec{\alpha}, \mathbf{w}, \vec{\theta}) \in \mathbb{R}^N \times \mathbb{R}^{n \times N} \times \mathbb{R}^N = \mathbb{R}^{n*}.$$

**Lemma 3.12** *Let  $\sigma : \mathbb{R} \rightarrow \mathbb{R}^+$  be a two times differentiable function with uniformly bounded function values and first, second order derivatives, such as the sigmoid, hyperbolic tangent functions (see Figure 1)<sup>1</sup>. Then, the derivatives of  $\Psi$  with respect to the coefficients  $\vec{p}$  are given by the following formulas:*

<sup>1</sup>This assumption is actually too restrictive, and only used to see that  $\Psi \in L^2([0, 1]^n)$ .

- Derivative with respect to  $\alpha_s$ ,  $s = 1, \dots, N$ :

$$\frac{\partial \Psi}{\partial \alpha_s}[\vec{p}](\vec{x}) = \sigma \left( \sum_{i=1}^n w_s^i x_i + \theta_s \right) \text{ for } s = 1, \dots, N. \quad (3.30)$$

- Derivative with respect to  $w_s^t$  where  $s = 1, \dots, N$ ,  $t = 1, \dots, n$ :

$$\frac{\partial \Psi}{\partial w_s^t}[\vec{p}](\vec{x}) = \sum_{j=1}^N \alpha_j \sigma' \left( \sum_{i=1}^n w_j^i x_i + \theta_j \right) \delta_{s=j} x_t = \alpha_s \sigma' \left( \sum_{i=1}^n w_s^i x_i + \theta_s \right) x_t \quad (3.31)$$

- Derivative with respect to  $\theta_s$  where  $s = 1, \dots, N$ :

$$\frac{\partial \Psi}{\partial \theta_s}[\vec{p}](\vec{x}) = \sum_{j=1}^N \alpha_j \sigma' \left( \sum_{i=1}^n w_j^i x_i + \theta_j \right) \delta_{s=j} = \alpha_s \sigma' \left( \sum_{i=1}^n w_s^i x_i + \theta_s \right). \quad (3.32)$$

Note, that all the derivatives above are functions in  $\mathbf{X} = L^2([0, 1]^n)$ . In particular, maybe in a more intuitive way, we have

$$D\Psi[\vec{p}](\vec{x})\vec{h} = \left( \frac{\partial \Psi}{\partial \vec{\alpha}}[\vec{p}](\vec{x}) \quad \frac{\partial \Psi}{\partial \mathbf{w}}[\vec{p}](\vec{x}) \quad \frac{\partial \Psi}{\partial \vec{\theta}}[\vec{p}](\vec{x}) \right)^T \vec{h} \text{ for all } \vec{h} = \begin{pmatrix} \vec{h}_{\vec{\alpha}} \\ \mathbf{h}_{\mathbf{w}} \\ \vec{h}_{\vec{\theta}} \end{pmatrix} \in \mathbb{R}^{n_*} \text{ and } \vec{x} \in \mathbb{R}^n. \quad (3.33)$$

Moreover, let  $s_1, s_2 = 1, \dots, N$ ,  $t_1, t_2 = 1, \dots, n$ , then we have in a formal way:

$$\begin{aligned} \frac{\partial^2 \Psi}{\partial \alpha_{s_1} \partial \alpha_{s_2}}(\vec{x}) &= 0, \\ \frac{\partial^2 \Psi}{\partial \alpha_{s_1} \partial w_{s_2}^{t_1}}(\vec{x}) &= \sigma' \left( \sum_{i=1}^n w_{s_1}^i x_i + \theta_{s_1} \right) x_{t_1} \delta_{s_1=s_2}, \\ \frac{\partial^2 \Psi}{\partial \alpha_{s_1} \partial \theta_{s_2}}(\vec{x}) &= \sigma' \left( \sum_{i=1}^n w_{s_1}^i x_i + \theta_{s_1} \right) \delta_{s_1=s_2}, \\ \frac{\partial^2 \Psi}{\partial w_{s_1}^{t_1} \partial w_{s_2}^{t_2}}(\vec{x}) &= \alpha_{s_1} \sigma'' \left( \sum_{i=1}^n w_{s_1}^i x_i + \theta_{s_1} \right) x_{t_1} x_{t_2} \delta_{s_1=s_2}, \\ \frac{\partial^2 \Psi}{\partial w_{s_1}^{t_1} \partial \theta_{s_2}}(\vec{x}) &= \alpha_{s_1} \sigma'' \left( \sum_{i=1}^n w_{s_1}^i x_i + \theta_{s_1} \right) x_{t_1} \delta_{s_1=s_2}, \\ \frac{\partial^2 \Psi}{\partial \theta_{s_1} \partial \theta_{s_2}}(\vec{x}) &= \alpha_{s_1} \sigma'' \left( \sum_{i=1}^n w_{s_1}^i x_i + \theta_{s_1} \right) \delta_{s_1=s_2}, \end{aligned} \quad (3.34)$$

where  $\delta_{a=b} = 1$  if  $a = b$  and 0 else, that is the Kronecker-delta.

The notation of directional derivatives with respect to parameters might be confusing. Note, that for instance  $\frac{\partial \Psi}{\partial \theta_s}[\vec{p}](\vec{x})$  denotes a direction derivative of the functional  $\Psi$  with respect to the variable  $\theta_s$  and this derivative is a function, which depends on  $\vec{x}$ . The argument, where the derivative is evaluated is a vector. So in such a formula  $\theta_s$  has two different meanings. Notationally differentiating between them would be exact but becomes quite unreadable.

**Remark 3.13** • In particular Equation 3.34 shows that

$$\begin{pmatrix} \vec{h}_{\vec{\alpha}} & \mathbf{h}_{\mathbf{w}} & \vec{h}_{\vec{\theta}} \end{pmatrix} D^2\Psi[\vec{p}](\vec{x}) \begin{pmatrix} \vec{h}_{\vec{\alpha}} \\ \mathbf{h}_{\mathbf{w}} \\ \vec{h}_{\vec{\theta}} \end{pmatrix} \text{ is continuous (for fixed } \vec{x}) \text{ with respect to } \vec{p}. \quad (3.35)$$

- We emphasize that under the assumptions of Lemma 3.12 the linear space (for fixed  $\vec{p}$ )

$$\mathcal{R}(D\Psi[\vec{p}]) = \left\{ D\Psi[\vec{p}]\vec{h} : \vec{h} = (\vec{h}_{\vec{\alpha}}, \mathbf{h}_{\mathbf{w}}, \vec{h}_{\vec{\theta}}) \in \mathbb{R}^{N \times (n+2)} \right\} \subseteq L^2([0, 1]^n).$$

- In order to prove convergence of the Gauss-Newton's method, Equation 3.21, by applying Theorem 3.1, we have to prove that  $\Psi$  is a Lipschitz-continuous immersion. Below we lack proving one important property so far, namely, that

$$\partial_k \Psi[\vec{p}], \quad k = 1, \dots, n_* = N(n+2) \quad (3.36)$$

are linearly independent functions. In this paper, this will remain open as a conjecture, and the following statements are valid modulo this conjecture.

In the following we survey some results on linear independence with respect to the coefficients  $\vec{\alpha}, \mathbf{w}, \vec{\theta}$  of the functions  $\vec{x} \rightarrow \sigma(\sum_{i=1}^n w_s^i x_i + \theta_s)$ , which match the functions  $\vec{x} \rightarrow \frac{\partial \Psi}{\partial \alpha_s}[\vec{p}](\vec{x})$ , that is with respect to the first  $N$  variables.

**3.5. Linear independence of activation functions and its derivatives.** The universal approximation results from for instance [9, 27, 26] do not allow to conclude that neural networks function as in Equation 3.23 are linearly independent. Linear independence is a non-trivial research question: We recall a result from [31] from which linear independence of a shallow neural network operator, as defined in Equation 3.23, can be deduced for a variety of activator functions. Similar results on linear independence of shallow network functions based on sigmoid activation functions have been stated in [47, 28], but the discussion in [31] raises questions on the completeness of the proofs. In [31] it is stated that all activation functions from the *Pytorch library* [41] are linearly independent with respect to almost all parameters  $\mathbf{w}$  and  $\theta$ .

**Theorem 3.14 ([31])** *For all activation functions HardShrink, HardSigmoid, HardTanh, HardSwish, LeakyReLU, PReLU, ReLU, ReLU6, RReLU, SoftShrink, Threshold, LogSigmoid, Sigmoid, SoftPlus, Tanh, and TanShrink and the PyTorch functions CELU, ELU, SELU the shallow neural network functions Equation 3.23 formed by randomly generated vectors  $(\mathbf{w}, \vec{\theta})$  are linearly independent.*

**Remark 3.15** (i) Theorem 3.14 states that the functions  $\frac{\partial \Psi}{\partial \alpha_s}$  (taking into account Equation 3.30) are linearly independent for *almost all* parameters  $(\mathbf{w}, \vec{\theta}) \in \mathbb{R}^{n \times N} \times \mathbb{R}^N$ . In other words, the first block of the matrix is  $D\Psi$  in Equation 3.33 consists of functions, which are linearly independent for almost all parameters  $(\mathbf{w}, \vec{\theta})$ . For our results to hold we need on top that the functions  $\frac{\partial \Psi}{\partial w_s^t}$  and  $\frac{\partial \Psi}{\partial \theta_s}$  from the second and third block (see Equation 3.34) are linearly independent within the blocks, respectively, and also across the blocks. So far this has not been proven but can be conjectured already from Figure 1.

(ii) For the sigmoid function we have *obvious symmetries* because

$$\sigma'(\mathbf{w}_j^T \vec{x} + \theta_j) = \sigma'(-\mathbf{w}_j^T \vec{x} - \theta_j) \quad \text{for every } \mathbf{w}_j \in \mathbb{R}^n, \vec{\theta} \in \mathbb{R}^N, \quad (3.37)$$

or in other words for the function  $\Psi$  from Equation 3.23 we have according to Equation 3.32 that

$$\frac{\partial \Psi}{\partial \theta_s}[\vec{\alpha}, \mathbf{w}, \vec{\theta}](\vec{x}) = \alpha_s \sigma'(\mathbf{w}_j^T \vec{x} + \theta_j) = \alpha_s \sigma'(-\mathbf{w}_j^T \vec{x} - \theta_j) = \frac{\partial \Psi}{\partial \theta_s}[\vec{\alpha}, -\mathbf{w}, -\vec{\theta}](\vec{x}) \quad (3.38)$$

or in other words  $\frac{\partial \Psi}{\partial \theta_s}[\vec{\alpha}, \mathbf{w}, -\vec{\theta}]$  and  $\frac{\partial \Psi}{\partial \theta_s}[\vec{\alpha}, -\mathbf{w}, -\vec{\theta}]$  are linearly dependent.

**Conjecture 3.16** We define by  $\mathcal{D}(\Psi)$  a *maximal set of vectors*  $(\vec{\alpha}, \mathbf{w}, \vec{\theta})$  such that the  $n_* = N \times (n+2)$  functions in  $\vec{x}$

$$\vec{x} \rightarrow \frac{\partial \Psi}{\partial \alpha_s}[\vec{\alpha}, \mathbf{w}, \vec{\theta}](\vec{x}), \quad \vec{x} \rightarrow \frac{\partial \Psi}{\partial w_s^t}[\vec{\alpha}, \mathbf{w}, \vec{\theta}](\vec{x}), \quad \vec{x} \rightarrow \frac{\partial \Psi}{\partial \theta_s}[\vec{\alpha}, \mathbf{w}, \vec{\theta}](\vec{x}), \quad s = 1, \dots, N, t = 1, \dots, n,$$

are linearly independent. We assume that  $\mathcal{D}(\Psi)$  is open and dense in  $\mathbf{X} \in L^2([0, 1])^2$  The later is guaranteed by Theorem 3.11. Recall the discussion above: The differentiation variables and the arguments coincide notationally, but are different objects.

**Remark 3.17** • It can be conjectured that for every element from  $\mathcal{D}(\Psi)$  only one element in  $\mathbb{R}^{n_*}$  exists, which satisfies *obvious symmetries* such as formulated in Equation 3.38. These ‘‘mirrored’’ elements are a set of measure zero in  $\vec{P}$ . We conjecture that this corresponds to the set of measure zero as stated in [31], which is derived with Fourier methods.

- Equation 3.34 requires that all components of the vector  $\vec{\alpha}$  are non-zero. This means in particular that for “sparse solutions”, with less than  $n_* = N * (n + 2)$  coefficients, convergence is not guaranteed, because of a locally degenerating submanifold. We consider the manifold given by the function

$$F : \mathbb{R}^2 \rightarrow \mathbb{R}^2.$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} xy \\ x^2 + y^2 \end{pmatrix} \quad (3.39)$$

Then

$$\nabla F(x, y) = \begin{pmatrix} y & x \\ 2x & 2y \end{pmatrix}.$$

We have  $\det \nabla F(x, y) = 2(y^2 - x^2)$ , which vanishes along the diagonals in  $(x, y)$ -space. That is of the diagonals the function is locally a submanifold (see Figure 2):




FIGURE 2. The function  $F$  from Equation 3.39. We have plotted  $F(x, y)$  via its polar coordinates, i.e.  $r = |F(x, y)|$  and  $\theta = \tan^{-1} \left( \frac{xy}{x^2 + y^2} \right)$ . The colors correspond to identical angles.

**3.6. Local convergence of Gauss-Newton’s method with coding networks.** In the following we prove a local convergence result for a Gauss-Newton’s method, for solving operator equations Equation 1.3 where  $F$  is complemented by a shallow neural network coder  $\Psi$ . In order to apply Theorem 3.1 we have to verify that the shallow neural network operator (see Equation 3.23) is a Lipschitz-differentiable immersion.

**Lemma 3.18** *Let  $F : \mathbf{X} = L^2([0, 1]^n) \rightarrow \mathbf{Y}$  be linear, bounded, with trivial nullspace and closed range, and let  $\sigma$  be strictly monotonic (like sigmoid or hyperbolic tangent) and satisfy the assumptions of Lemma 3.12. Moreover, assume that Conjecture 3.16 holds. Then*

- (i) *For every element  $\vec{p} = (\vec{\alpha}, \mathbf{w}, \vec{\theta}) \in \mathbb{R}^{n_*}$  in the maximal set  $\mathcal{D}(\Psi)$  (see Conjecture 3.16),  $\mathcal{R}(D\Psi[\vec{p}])$  is a linear subspace of the space  $\mathbf{X}$  of dimension  $n_* = N \times (n + 2)$ .*
- (ii) *There exists an open neighborhood  $\mathcal{U} \subseteq \mathbb{R}^{N \times (n+2)}$  of vectors  $(\vec{\alpha}, \mathbf{w}, \vec{\theta})$  such that  $\Psi$  is a Lipschitz-differentiable immersion in  $\mathcal{U}$ .*

*Proof:* • It is clear that for each fixed  $\vec{p}$ ,  $D\Psi[\vec{p}] \in L^2([0, 1]^n)$  because of the differentiability assumptions of  $\sigma$ , see Equation 3.35. Conjecture 3.16 implies that  $\mathcal{R}(D\Psi[\vec{p}])$  is a linear subspaces of  $\mathbf{X}$  of dimension  $N \times (n + 2)$  (note the elements are functions).

- $D^2\Psi[\vec{p}] : \mathbb{R}^{N \times (n+2)} \rightarrow L^2([0, 1]^n)$  is continuous (see Equation 3.35) since we assume that the activation function  $\sigma$  is twice differentiable. Now we consider a non-empty open neighborhood  $\mathcal{U}$  of a vector  $\vec{p}$ , with a compact closure.

- Then, from the continuity of  $D^2\Psi$  with respect to  $\vec{p}$ , it follows that  $D\Psi$  is a Fréchet-differentiable with Lipschitz-continuous derivative on  $\mathcal{U}$ . In particular this means that *item (i)* in [Definition 3.3](#) holds. Moreover, [Equation 3.9](#) holds for  $\Psi'$ . That is, there exists constants  $C_L$  and  $C_I$  such that

$$\|\Psi'(\vec{p}) - \Psi'(\vec{q})\|_{\vec{p} \rightarrow \mathbf{Y}} \leq C_L \|\vec{p} - \vec{q}\|_{\vec{p}} \quad \text{and} \quad \|\Psi'(\vec{p})\|_{\vec{p} \rightarrow \mathbf{Y}} \leq C_I \quad \text{for } \vec{p}, \vec{q} \in \mathcal{D}(\Psi). \quad (3.40)$$

□

Note that  $\Psi'(p)^\dagger$  as defined in [Equation 3.8](#) is also uniformly bounded and Lipschitz-continuous as a consequence of [Lemma 3.4](#).

**Theorem 3.19 (Local convergence of Gauss-Newton’s method)** *Let  $F : \mathbf{X} = L^2([0, 1]^n) \rightarrow \mathbf{Y}$  be a linear, bounded operator with trivial nullspace and dense range and let  $N = F \circ \Psi$ , where  $\Psi : \mathcal{D}(\Psi) \subseteq \mathbb{R}^{N \times (n+2)} \rightarrow \mathbf{X}$  is a shallow neural network operator generated by an activation function  $\sigma$  which satisfies the assumptions of [Lemma 3.18](#) and [Conjecture 3.16](#). Let  $\vec{p}^0 \in \mathcal{D}(\Psi)$  be the starting point of the Gauss-Newton’s iteration [Equation 3.21](#) and let  $\vec{p}^\dagger \in \mathcal{D}(\Psi)$  be a solution of [Equation 3.20](#), which satisfy [Equation 3.2](#). Then the Gauss-Newton’s iterations are locally, that is if  $\vec{p}^0$  is sufficiently close to  $\vec{p}^\dagger$ , and quadratically converging.*

*Proof:* The proof is an immediate application of [Lemma 3.18](#) to [Theorem 3.6](#). □

**Remark 3.20** We have shown that a nonlinear operator equation, where the operator is a composition of a linear compact operator and a shallow neural network operator, can be solved with a Gauss-Newton’s method with guaranteed local convergence in the parameter space.

**Conclusion.** We have shown that Gauss-Newton’s methods are efficient algorithms for solving linear inverse problems, where the solution can be encoded with a neural network. The convergence studies, however, are not complete, and are based on a conjecture on linear independence of activation functions and its derivatives.

**Acknowledgements.** This research was funded in whole, or in part, by the Austrian Science Fund (FWF) P 34981 – New Inverse Problems of Super-Resolved Microscopy (NIPSUM). For the purpose of open access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. Moreover, OS is supported by the Austrian Science Fund (FWF), with SFB F68 “Tomography Across the Scales”, project F6807-N36 (Tomography with Uncertainties). The financial support by the Austrian Federal Ministry for Digital and Economic Affairs, the National Foundation for Research, Technology and Development and the Christian Doppler Research Association is gratefully acknowledged. BH is supported by the German Science Foundation (DFG) under the grant HO 1454/13-1 (Project No. 453804957).

## REFERENCES

- [1] A. Aspri, L. Frischauf, Y. Korolev, and O. Scherzer. “Data Driven Reconstruction Using Frames and Riesz Bases”. In: *Deterministic and Stochastic Optimal Control and Inverse Problems*. Ed. by B. Jadamba, A. A. Khan, S. Migórski, and M. Sama. CRC Press, 2021, pp. 303–318. DOI: [10.1201/9781003050575-13](https://doi.org/10.1201/9781003050575-13) (cited on page 2).
- [2] A. Aspri, Y. Korolev, and O. Scherzer. “Data driven regularization by projection”. In: *Inverse Problems* 36.12 (Dec. 2020), p. 125009. ISSN: 0266-5611. DOI: [10.1088/1361-6420/abb61b](https://doi.org/10.1088/1361-6420/abb61b) (cited on page 2).
- [3] A. Bakushinskii and A. Goncharskii. “Ill-Posed Problems: Theory and Applications”. Dordrecht, Boston, London: Kluwer Academic Publishers, 1994 (cited on page 1).
- [4] A. Bakushinskii and A. Goncharskii. “Iterative Methods for the Solution of Incorrect Problems”. Moscow: Nauka, 1989 (cited on page 1).
- [5] B. Blaschke. “Some Newton Type Methods for the Regularization of Nonlinear Ill-Posed Problems”. Linz, Austria: Universitätsverlag Rudolf Trauner, 1996 (cited on pages 3, 4).

- 
- [6] A. Bora, A. Jalal, E. Price, and A. G. Dimakis. “Compressed Sensing using Generative Models”. In: *Proceedings of the 34th International Conference on Machine Learning*. Ed. by D. Precup and Y. W. Teh. Vol. 70. Proceedings of Machine Learning Research. PMLR, Aug. 2017, pp. 537–546 (cited on page 1).
- [7] J. Braunsman, M. Rajković, M. Rumpf, and B. Wirth. “Learning Low Bending and Low Distortion Manifold Embeddings: Theory and Applications”. Preprint on ArXiv 2208.10193. 2022. DOI: [10.48550/arxiv.2208.10193](https://doi.org/10.48550/arxiv.2208.10193) (cited on page 1).
- [8] G. Chavent and K. Kunisch. “Regularization in state space”. In: *Mathematical Modelling and Numerical Analysis* 27 (1993), pp. 535–564. ISSN: 0764-583X (cited on pages 2, 3).
- [9] G. Cybenko. “Approximation by superpositions of a sigmoidal function”. In: *Mathematics of Control, Signals, and Systems* 2.4 (1989), pp. 303–314. DOI: [10.1007/bf02551274](https://doi.org/10.1007/bf02551274) (cited on pages 11, 13).
- [10] D. W. Decker, H. B. Keller, and C. T. Kelley. “Convergence Rates for Newton’s Method at Singular Points”. In: *SIAM Journal on Numerical Analysis* 20.2 (1983), pp. 296–314. ISSN: 0036-1429. DOI: [10.1137/0720020](https://doi.org/10.1137/0720020) (cited on page 9).
- [11] P. Deuffhard and G. Heindl. “Affine invariant convergence theorems for Newton’s method and extensions to related methods”. In: *SIAM Journal on Numerical Analysis* 16.1 (1979), pp. 1–10. ISSN: 0036-1429 (cited on page 4).
- [12] P. Deuffhard and F. A. Potra. “Asymptotic Mesh Independence of Newton–Galerkin Methods via a Refined Mysovskii Theorem”. In: *SIAM Journal on Numerical Analysis* 29.5 (1992), pp. 1395–1412. ISSN: 0036-1429. DOI: [10.1137/0729080](https://doi.org/10.1137/0729080) (cited on page 4).
- [13] M. Duff, N. D. F. Campbell, and M. J. Ehrhardt. “Regularising Inverse Problems with Generative Machine Learning Models”. Preprint on ArXiv 2107.11191. 2021. DOI: [10.48550/arxiv.2107.11191](https://doi.org/10.48550/arxiv.2107.11191) (cited on pages 1, 3).
- [14] H. W. Engl, M. Hanke, and A. Neubauer. “Regularization of inverse problems”. Mathematics and its Applications 375. Dordrecht: Kluwer Academic Publishers Group, 1996. viii+321. ISBN: 0-7923-4157-0 (cited on page 1).
- [15] R. Gorenflo and B. Hofmann. “On autoconvolution and regularization”. In: *Inverse Problems* 10.2 (1994), pp. 353–373. ISSN: 0266-5611. DOI: [10.1088/0266-5611/10/2/011](https://doi.org/10.1088/0266-5611/10/2/011) (cited on page 2).
- [16] P. Grisvard. “Elliptic Problems in Nonsmooth Domains”. Boston: Pitman, 1985 (cited on page 9).
- [17] M. Hanke, A. Neubauer, and O. Scherzer. “A convergence analysis of the Landweber iteration for nonlinear ill-posed problems”. In: *Numerische Mathematik* 72.1 (1995), pp. 21–37. ISSN: 0029-599X. DOI: [10.1007/s002110050158](https://doi.org/10.1007/s002110050158) (cited on page 3).
- [18] W. Häussler. “A Kantorovich-type convergence analysis for the Gauss-Newton-method”. In: *Numerische Mathematik* 48 (1986), pp. 119–125. ISSN: 0029-599X (cited on page 9).
- [19] T. Hein and B. Hofmann. “On the nature of ill-posedness of an inverse problem arising in option pricing”. In: *Inverse Problems* 19.6 (2003), pp. 1319–1338. ISSN: 0266-5611. DOI: [10.1088/0266-5611/19/6/006](https://doi.org/10.1088/0266-5611/19/6/006) (cited on page 3).
- [20] B. Hofmann. “Mathematik Inverser Probleme”. Stuttgart: Teubner, 1999 (cited on page 2).
- [21] B. Hofmann. “On the degree of ill-posedness for nonlinear problems”. In: *Journal of Inverse and Ill-Posed Problems* 2 (1994), pp. 61–76. ISSN: 0928-0219 (cited on pages 2, 3).
- [22] B. Hofmann, ed. “Regularization for Applied Inverse and Ill-Posed Problems”. Leipzig, 1986. DOI: [10.1007/978-3-322-93034-7](https://doi.org/10.1007/978-3-322-93034-7) (cited on page 1).
- [23] B. Hofmann, B. Kaltenbacher, C. Pöschl, and O. Scherzer. “A convergence rates result for Tikhonov regularization in Banach spaces with non-smooth operators”. In: *Inverse Problems* 23.3 (2007), pp. 987–1010. ISSN: 0266-5611. DOI: [10.1088/0266-5611/23/3/009](https://doi.org/10.1088/0266-5611/23/3/009) (cited on page 3).
- [24] B. Hofmann and O. Scherzer. “Factors influencing the ill-posedness of nonlinear problems”. In: *Inverse Problems* 10.6 (1994), pp. 1277–1297. ISSN: 0266-5611. DOI: [10.1088/0266-5611/10/6/007](https://doi.org/10.1088/0266-5611/10/6/007) (cited on page 2).
- [25] B. Hofmann and U. Tautenhahn. “On Ill-Posedness Measures and Space Change in Sobolev Scales”. In: *Zeitschrift für Analysis und ihre Anwendungen. Journal of Analysis and its Applications* 16.4 (1997), pp. 979–1000. DOI: [10.4171/zaa/800](https://doi.org/10.4171/zaa/800) (cited on page 2).



- [26] K. Hornik. “Approximation capabilities of multilayer feedforward networks”. In: *Neural Networks* 4.2 (1991), pp. 251–257. DOI: [10.1016/0893-6080\(91\)90009-t](https://doi.org/10.1016/0893-6080(91)90009-t) (cited on pages 10, 11, 13).
- [27] K. Hornik, M. Stinchcombe, and H. White. “Multilayer Feedforward Networks are Universal Approximators”. In: *Neural Networks* 2 (1989), pp. 359–366 (cited on pages 11, 13).
- [28] G.-B. Huang. “Learning capability and storage capacity of two-hidden-layer feedforward networks”. In: *IEEE Transactions on Neural Networks* 14.2 (2003), pp. 274–281. DOI: [doi:10.1109/tnn.2003.809401](https://doi.org/10.1109/tnn.2003.809401) (cited on page 13).
- [29] B. Kaltenbacher, A. Neubauer, and O. Scherzer. “Iterative regularization methods for nonlinear ill-posed problems”. Vol. 6. Radon Series on Computational and Applied Mathematics. Berlin: Walter de Gruyter, 2008. ISBN: 978-3-11-020420-9. DOI: [10.1515/9783110208276](https://doi.org/10.1515/9783110208276) (cited on pages 1, 3, 4).
- [30] L. Kantorowitsch and G. Akilow. “Funktionalanalysis in normierten Räumen”. Berlin: Akademie Verlag, 1964 (cited on page 9).
- [31] A. Lamperski. “Neural Network Independence Properties with Applications to Adaptive Control”. In: *2022 IEEE 61st Conference on Decision and Control (CDC)*. 2022. DOI: [10.1109/cdc51059.2022.9992994](https://doi.org/10.1109/cdc51059.2022.9992994) (cited on page 13).
- [32] V. A. Morozov. “Methods for Solving Incorrectly Posed Problems”. New York, Berlin, Heidelberg: Springer, 1984 (cited on page 1).
- [33] M. Z. Nashed. “A new approach to classification and regularization of ill-posed operator equations”. In: *Inverse and Ill-Posed Problems*. Vol. 4. 1987, pp. 53–75. DOI: [10.1016/b978-0-12-239040-1.50009-0](https://doi.org/10.1016/b978-0-12-239040-1.50009-0) (cited on page 2).
- [34] M. Z. Nashed. “Inner, outer, and generalized inverses in Banach and Hilbert spaces”. In: *Numerical Functional Analysis and Optimization* 9.3-4 (1987), pp. 261–325. ISSN: 0163-0563. DOI: [10.1080/01630568708816235](https://doi.org/10.1080/01630568708816235) (cited on pages 4, 9).
- [35] M. Z. Nashed. “On the perturbation theory of generalized inverse operators in Banach spaces”. In: *Functional Analysis methods in numerical analysis*. Ed. by M. Z. Nashed. Springer, New York, 1979, pp. 180–195 (cited on page 9).
- [36] M. Nashed, ed. *Generalized inverses and applications*. New York: Academic Press [Harcourt Brace Jovanovich Publishers], 1976, pp. xiv+1054 (cited on page 4).
- [37] M. Nashed and X. Chen. “Convergence of Newton-like methods for singular operator equations using outer inverses”. In: *Numerische Mathematik* 66 (1993), pp. 235–257. ISSN: 0029-599X (cited on pages 2, 4, 9).
- [38] D. Obmann, J. Schwab, and M. Haltmeier. “Deep synthesis network for regularizing inverse problems”. In: *Inverse Problems* 37.1 (2021), p. 015005. ISSN: 0266-5611. DOI: [10.1088/1361-6420/abc7cd](https://doi.org/10.1088/1361-6420/abc7cd) (cited on page 2).
- [39] J. M. Ortega. “The Newton-Kantorovich Theorem”. In: *The American Mathematical Monthly* 75.6 (1968), p. 658. ISSN: 0002-9890. DOI: [10.2307/2313800](https://doi.org/10.2307/2313800) (cited on page 4).
- [40] J. Ortega and W. Rheinboldt. “Iterative Solution of Nonlinear Equations in Several Variables”. New York: Academic Press, 1970 (cited on page 9).
- [41] A. Paszke et al. “PyTorch: An Imperative Style, High-Performance Deep Learning Library”. In: *Proceedings of the 33rd International Conference on Neural Information Processing Systems*. 721. Curran Associates Inc., 2019 (cited on page 13).
- [42] C. Pöschl, E. Resmerita, and O. Scherzer. “Discretization of variational regularization in Banach spaces”. In: *Inverse Problems* 26.10 (2010), p. 105017. ISSN: 0266-5611. DOI: [10.1088/0266-5611/26/10/105017](https://doi.org/10.1088/0266-5611/26/10/105017) (cited on page 2).
- [43] S. Rifai, G. Mesnil, P. Vincent, X. Muller, Y. Bengio, Y. Dauphin, and X. Glorot. “Higher Order Contractive Auto-Encoder”. In: *Machine Learning and Knowledge Discovery in Databases*. 2011, pp. 645–660. DOI: [10.1007/978-3-642-23783-6\\_41](https://doi.org/10.1007/978-3-642-23783-6_41) (cited on page 1).
- [44] O. Scherzer, M. Grasmair, H. Grossauer, M. Haltmeier, and F. Lenzen. “Variational methods in imaging”. Applied Mathematical Sciences 167. New York: Springer, 2009. ISBN: 978-0-387-30931-6. DOI: [10.1007/978-0-387-69277-7](https://doi.org/10.1007/978-0-387-69277-7) (cited on page 1).
- [45] T. Schuster, B. Kaltenbacher, B. Hofmann, and K. S. Kazimierski. “Regularization methods in Banach spaces”. Radon Series on Computational and Applied Mathematics 10. Berlin, Boston: De Gruyter, 2012. xii+283. DOI: [10.1515/9783110255720](https://doi.org/10.1515/9783110255720) (cited on page 1).

- 
- [46] H. Schwetlick. “Numerische Lösung nichtlinearer Gleichungen”. Vol. 17. *Mathematik für Naturwissenschaft und Technik [Mathematics for Science and Technology]*. VEB Deutscher Verlag der Wissenschaften, Berlin, 1979, p. 346 (cited on pages 4, 9).
- [47] S. Tamura and M. Tateishi. “Capabilities of a four-layered feedforward neural network: four layers versus three”. In: *IEEE Transactions on Neural Networks* 8.2 (1997), pp. 251–255. DOI: [doi : 10.1109/72.557662](https://doi.org/10.1109/72.557662) (cited on page 13).