

On the failure of the bootstrap for Chatterjee's rank correlation

Zhexiao Lin* and Fang Han†

April 6, 2023

Abstract

While researchers commonly use the bootstrap for statistical inference, many of us have realized that the standard bootstrap, in general, does not work for Chatterjee's rank correlation. In this paper, we provide proof of this issue under an additional independence assumption, and complement our theory with simulation evidence for general settings. Chatterjee's rank correlation thus falls into a category of statistics that are asymptotically normal but bootstrap inconsistent. Valid inferential methods in this case are Chatterjee's original proposal (for testing independence) and [Lin and Han \(2022\)](#)'s analytic asymptotic variance estimator (for more general purposes).

Keywords: bootstrap, rank correlation, tied data.

1 Introduction

Rank correlation is an essential tool for measuring the association between random variables. Its development is closely linked to the history of statistics as a discipline and has involved many notable figures, including Charles Spearman, Sir Maurice Kendall, Wassily Hoeffding, Jack Kiefer, Murray Rosenblatt, Jaroslav Hájek, Eric Lehmann, Herman Chernoff, and Richard Savage ([Spearman, 1904, 1906](#); [Kendall, 1938, 1970](#); [Hoeffding, 1940, 1948, 1994](#); [Blum et al., 1961](#); [Hájek et al., 1999](#); [Hodges and Lehmann, 1956](#); [Chernoff and Savage, 1958](#)). Unlike other correlation coefficients, a rank correlation relies solely on the rankings of the original data, making it (1) exactly distribution-free when testing independence of continuous random variables; (2) invariant to marginal monotonic transformations (cf. copulas); and (3) robust in the face of outliers and heavy-tailedness. Its usefulness, therefore, is self-explanatory.

Given the remarkable progress made in this area over the past century, it is impressive that, recently, Sourav Chatterjee devised a new rank correlation that is both new and appealing from multiple perspectives. Specifically, consider an independent and identically distributed (i.i.d.) sample $\{X_i, Y_i\}_{i=1, \dots, n}$ from a pair of scalars, (X, Y) , with joint and marginal distribution functions $F_{X,Y}$ and F_X, F_Y , respectively. Let

$$R_i := \sum_{j=1}^n \mathbb{1}(Y_j \leq Y_i) \quad \text{and} \quad L_i := \sum_{j=1}^n \mathbb{1}(Y_j \geq Y_i)$$

*Department of Statistics, University of California, Berkeley, CA 94720, USA; e-mail: zhexiaolin@berkeley.edu

†Department of Statistics, University of Washington, Seattle, WA 98195, USA; e-mail: fanghan@uw.edu

be the rank and “reversed” rank of Y_i with $\mathbf{1}(\cdot)$ representing the indicator function, and let $\{[i], i = 1, \dots, n\}$ be a rearrangement of $\{1, \dots, n\}$ such that $X_{[1]} \leq \dots \leq X_{[n]}$ with ties broken at random. Chatterjee (2021) introduced the following statistic

$$\xi_n := 1 - \frac{n}{2 \sum_{i=1}^n L_i(n - L_i)} \sum_{i=1}^{n-1} |R_{[i+1]} - R_{[i]}|, \quad (1.1)$$

which he showed to be a strongly consistent estimator of Dette-Siburg-Stoimenov’s dependence measure (Dette et al., 2013),

$$\xi = \xi(X, Y) := \frac{\int \text{Var} \{E[\mathbf{1}(Y \geq y) | X]\} dF_Y(y)}{\int \text{Var} \{\mathbf{1}(Y \geq y)\} dF_Y(y)}, \quad (1.2)$$

as long as Y is not almost surely a constant.

Why is ξ_n in (1.1) appealing? There are three reasons Sourav Chatterjee outlines. First, it has a simple form. Second, it has a normal limiting null distribution. Finally, it measures a dependence measure, ξ in (1.2), that satisfies Rényi’s criteria (Rényi, 1959) and Bickel’s definition of a measure of functional dependence (Bickel, 2022): ξ is zero if and only if Y is independent of X and one if and only if Y is a measurable function of X . Therefore, ξ_n is a rank correlation that can accurately quantify both independence and functional dependence. This is something that all aforementioned rank correlations, including Spearman’s ρ , Kendall’s τ , Hoeffding’s D , Blum-Kiefer-Rosenblatt’s r , and Bergsma-Dassios-Yanagimoto’s τ^* (Bergsma and Dassios, 2014; Yanagimoto, 1970), fail to achieve.

Due to these appealing properties, Chatterjee’s rank correlation has gained significant interest and a wave of research has emerged exploring its applications and extensions. Notable recent works include Azadkia and Chatterjee (2021), Cao and Bickel (2020), Shi et al. (2022), Gamboa et al. (2022), Deb et al. (2020), Huang et al. (2020), Auddy et al. (2021), Shi et al. (2023), Lin and Han (2023), Fuchs (2023), Azadkia et al. (2021), Griessenberger et al. (2022), Strothmann et al. (2022), Zhang (2023b), Bickel (2022), Holma (2022), Chatterjee and Vidyasagar (2022), Lin and Han (2022), Han and Huang (2022), Ansari and Fuchs (2022), and Zhang (2023a). Additionally, brief surveys on recent progress of (Chatterjee’s and more) rank correlation methods have been conducted in Han (2021) and Chatterjee (2022).

This paper aims to investigate the validity of the standard bootstrap (Efron, 1979, 1981) when applied to *fixed and continuous* $F_{X,Y}$, with ξ_n taking the form (1.1) to handle ties in resampled data. We prove, in the simple independence case with $F_{X,Y} = F_X F_Y$, that the standard bootstrap results in an inconsistent estimator of ξ_n ’s asymptotic variance, and the bootstrap distribution fails to converge to the limiting distribution of $\sqrt{n}(\xi_n - \xi)$. Simulations further complement the theory, indicating that the standard bootstrap will likely also fail in general settings with $F_{X,Y} \neq F_X F_Y$, even though $\sqrt{n}(\xi_n - \xi)$ still weakly converges to a normal distribution (Lin and Han, 2022). Chatterjee’s rank correlation thus falls into a class of statistics that are root- n consistent, asymptotically normal, but bootstrap inconsistent — a class including Bickel and Freedman’s U-statistics (Bickel and Freedman, 1981, Section 6), Hodges estimator (Beran, 1982, Pages 213-214), and Abadie and Imbens’s matching estimator (Abadie and Imbens, 2008).

It is worth mentioning that there are valid alternatives to using the standard bootstrap for inferring ξ from ξ_n . Sourav Chatterjee derived the limiting null distribution of ξ_n for testing independence between X and Y in Chatterjee (2021). In Lin and Han (2022), conditions were proven under which ξ_n is root- n consistent and asymptotically normal, and an analytic estimator of its asymptotic variance was proposed. This paper thus helps to justify the value of these derivations by demonstrating the inconsistency of an otherwise attractive alternative.

2 Main results

2.1 Setup

In this paper we consider the standard model of $(X_1, Y_1), \dots, (X_n, Y_n)$ to be n independent copies of (X, Y) drawn from a fixed and continuous $F_{X,Y}$ of support in the two-dimensional real space. In this case, with probability one there is no tie in the observation, and hence ξ_n admits the following simpler form,

$$\xi_n := 1 - \frac{3}{n^2 - 1} \sum_{i=1}^{n-1} |R_{[i+1]} - R_{[i]}|. \quad (2.1)$$

To implement the standard bootstrap, consider $(\mathbf{X}_b, \mathbf{Y}_b) = \{(X_{b,i}, Y_{b,i})\}_{i=1}^n$ to be the bootstrap sample, of size n and likely embracing ties, by sampling with replacement from $(\mathbf{X}, \mathbf{Y}) := \{(X_i, Y_i)\}_{i=1}^n$. Let $\tilde{\xi}_b$ be Chatterjee's rank correlation calculated using the bootstrap sample $(\mathbf{X}_b, \mathbf{Y}_b)$. More specifically, write

$$\{[i]_b, i = 1, \dots, n\} \quad (2.2)$$

to be a rearrangement of $\{1, \dots, n\}$ such that $X_{b,[1]_b} \leq \dots \leq X_{b,[n]_b}$, with ties in $\{X_{b,i}\}_{i=1}^n$ broken *in arbitrary way*. Let

$$R_{b,i} := \sum_{j=1}^n \mathbb{1}(Y_{b,j} \leq Y_{b,i}) \quad \text{and} \quad L_{b,i} := \sum_{j=1}^n \mathbb{1}(Y_{b,j} \geq Y_{b,i})$$

be the ranks and reversed ranks of the bootstrap sample. The bootstrapped rank correlation $\tilde{\xi}_b$ is then defined to be

$$\tilde{\xi}_b := 1 - \frac{n}{2 \sum_{i=1}^n L_{b,i}(n - L_{b,i})} \sum_{i=1}^{n-1} |R_{b,[i+1]_b} - R_{b,[i]_b}|, \quad (2.3)$$

namely, plugging (1.1) to the bootstrap sample.

In this paper, we investigate two commonly used versions of the bootstrap in empirical research. The first version involves centering the bootstrap sample at ξ_n , which is calculated using the original sample. The second version involves centering the bootstrap sample at the mean of the bootstrap distribution $E[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}]$. One can then estimate the asymptotic variance using

$$\text{either } nE[(\tilde{\xi}_b - \xi_n)^2 | \mathbf{X}, \mathbf{Y}] \quad \text{or} \quad n \text{Var}[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}],$$

assuming an infinite number of replications for bootstrap. In addition, it is of interest to evaluate

the closeness of the bootstrap distributions of

$$\sqrt{n}(\tilde{\xi}_b - \xi_n) \mid \mathbf{X}, \mathbf{Y} \quad \text{and} \quad \sqrt{n}(\tilde{\xi}_b - \mathbb{E}[\tilde{\xi}_b \mid \mathbf{X}, \mathbf{Y}]) \mid \mathbf{X}, \mathbf{Y}$$

to that of $\sqrt{n}(\xi_n - \xi)$.

2.2 Theory

Our theory section has to be focused on the simple independence case with $F_{X,Y} = F_X F_Y$, only under which we are able to provide the otherwise formidable calculation of the limits of $n\mathbb{E}[(\tilde{\xi}_b - \xi_n)^2]$ and $n \text{Var}[\tilde{\xi}_b \mid \mathbf{X}, \mathbf{Y}]$.

The following result of Chatterjee establishes the limiting distribution of ξ_n under independence.

Proposition 2.1 (Theorem 2.1, Chatterjee (2021)). *Assume $F_{X,Y} = F_X F_Y$ is fixed and continuous. We then have $\sqrt{n}\xi_n$ weakly converges to $N(0, 2/5)$.*

Below is the main theorem of this paper.

Theorem 2.1 (Bootstrap inconsistency). *Assuming the same conditions of Proposition 2.1, the following two statements then hold.*

- (i) *(variance inconsistency) $n\mathbb{E}[(\tilde{\xi}_b - \xi_n)^2 \mid \mathbf{X}, \mathbf{Y}]$ and $n \text{Var}[\tilde{\xi}_b \mid \mathbf{X}, \mathbf{Y}]$ do not converge to $2/5$ in probability.*
- (ii) *(distribution inconsistency) There exists a sequence of measurable events $[\mathcal{E}_i]_{i=1}^\infty$, satisfying*

$$\liminf_{n \rightarrow \infty} \mathbb{P}((X_1, Y_1, \dots, X_n, Y_n) \in \mathcal{E}_n) > 0,$$

such that the distributions of $\sqrt{n}(\tilde{\xi}_b - \xi_n)$ and $\sqrt{n}(\tilde{\xi}_b - \mathbb{E}[\tilde{\xi}_b \mid \mathbf{X}, \mathbf{Y}])$ do not converge to $N(0, 2/5)$ conditional on $(X_1, Y_1, \dots, X_n, Y_n) \in \mathcal{E}_n$.

The authors are intrigued by Theorem 2.1 and believe its significance is best appreciated in the context of mathematical statistics history, where the bootstrap method's validity has been a central topic, with establishing/disproving its consistency being particularly imperative.

For an i.i.d. sample, bootstrap consistency is often linked to the studied statistic's root- n consistency and asymptotic normality. According to Shao and Tu (1995, Page 128), the conventional wisdom seems to suggest that “[u]sually the consistency of the bootstrap distribution estimator requires some smoothness conditions that are *almost the same as* those required for the asymptotic normality of the given statistic and certain moment conditions.” As a matter of fact, this insight has been partly formalized in Mammen (1991, Theorem 1), where Enno Mammen demonstrated, elegantly, that bootstrap consistency is equivalent to asymptotic normality when applied to linear functionals.

Indeed, the majority of theoretical results on bootstrap inconsistency are centered on statistics that do not exhibit a regular pattern of being root- n consistent and asymptotically normal. In this regard, Athreya (1987), Knight (1989), and Hall (1990) focused on the sample mean with a sample drawn from heavy-tailed distributions, Beran and Srivastava (1985) on eigenvalues, Hall et al. (1993) on ranked parameters, and Andrews (2000) and Drton and Williams (2011) on parameters

at the boundary. Furthermore, [Abrevaya and Huang \(2005\)](#), [Kosorok \(2008\)](#), and [Sen et al. \(2010\)](#) examined cubic-root consistent estimators, [Bretagnolle \(1983\)](#) and [Arcones and Gine \(1992\)](#) investigated degenerate U- and V-statistics, and [Dümbgen \(1993\)](#) and [Fang and Santos \(2019\)](#) explored a general class of non-smooth plug-in estimators.

For statistics that exhibit root- n consistency and asymptotic normality, we categorize the cases where bootstrap inconsistency arises to three groups: (1) those that fail due to moment condition, e.g., Bickel and Freedman’s U-statistics ([Bickel and Freedman, 1981](#), Section 6); (2) those that fail at superefficiency points, e.g., Hodges and Stein estimators ([Beran, 1997](#); [Samworth, 2003](#)); and (3) those that do not belong to the previous two groups, including, notably, Abadie and Imbens’s nearest neighbor matching estimator of the average treatment effect ([Abadie and Imbens, 2008](#)).

Abadie and Imbens’s case is particularly relevant to our work on Chatterjee’s rank correlation, as both can be perceived as a type of nearest neighbor graph-based statistics with a *fixed* number of nearest neighbors. To the best of the authors’ knowledge, however, no work has established a general relationship between the inconsistency of the bootstrap and the “irregularity” of graph-based statistics; this would be an interesting future question for mathematical statisticians.

Finally, it should be noted that the issue of bootstrap inconsistency is not a universal problem affecting all rank correlations or rank-based statistics. For example, the bootstrap consistency of Spearman’s ρ and Kendall’s τ can be easily established based on the works of [Bickel and Freedman \(1981\)](#) and [Arcones and Gine \(1992\)](#). On the other hand, the bootstrap inconsistency of Hoeffding’s D , Blum-Kiefer-Rosenblatt’s r , and Bergsma-Dassios-Yanagimoto’s τ^* under independence between X and Y is caused by the non-normal convergence of the degenerate U-statistics, but not by the ranking.

2.3 Simulations

One might be tempted to speculate that the bootstrap inconsistency observed in [Theorem 2.1](#) is solely due to the “degeneracy” property of the null point of independence. While independence does play a crucial role in the cases of Hoeffding’s D , Blum-Kiefer-Rosenblatt’s r , and Bergsma-Dassios-Yanagimoto’s τ^* — they only become degenerate when X is independent of Y — the case of Chatterjee’s rank correlation appears to be different.

As tracking the limits of $nE[(\tilde{\xi}_b - \xi_n)^2]$ and $n \text{Var}[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}]$ under dependence between Y and X is technically intimidating, this paper relies on simulations to illustrate this point. To this end, we investigate the following six methods.

- (i) (V-LH) The asymptotic variance estimator described in [Lin and Han \(2022, Theorem 1.2\)](#);
- (ii) (V-B1) the bootstrap asymptotic variance estimator using $nE[(\tilde{\xi}_b - \xi_n)^2 | \mathbf{X}, \mathbf{Y}]$;
- (iii) (V-B2) the bootstrap asymptotic variance estimator using $n \text{Var}[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}]$;
- (iv) (D-LH) constructing the confidence interval using the idea described in [Lin and Han \(2022, Remark 1.4\)](#);
- (v) (D-HB1) constructing the confidence interval using the hybrid bootstrap ([Shao and Tu, 1995, Section 4.1.5](#)) based on $\sqrt{n}(\tilde{\xi}_b - \xi_n) | \mathbf{X}, \mathbf{Y}$;

Table 1: Variance estimation and empirical coverage probability

| ρ | n | Variance, RMSE | | | Coverage, $\alpha = 0.05$ | | | Coverage, $\alpha = 0.1$ | | |
|--------|-------|----------------|---------|------|---------------------------|-------|-------|--------------------------|-------|-------|
| | | V-LH | V-B1 | V-B2 | D-LH | D-HB1 | D-HB2 | D-LH | D-HB1 | D-HB2 |
| 0 | 1000 | 0.18 | 135.65 | 0.09 | 0.90 | 0.00 | 0.92 | 0.85 | 0.00 | 0.86 |
| | 5000 | 0.08 | 676.92 | 0.09 | 0.94 | 0.00 | 0.91 | 0.89 | 0.00 | 0.85 |
| | 10000 | 0.06 | 1353.32 | 0.09 | 0.95 | 0.00 | 0.92 | 0.90 | 0.00 | 0.85 |
| 0.3 | 1000 | 0.18 | 122.19 | 0.16 | 0.90 | 0.00 | 0.89 | 0.85 | 0.00 | 0.82 |
| | 5000 | 0.07 | 610.14 | 0.16 | 0.94 | 0.00 | 0.88 | 0.89 | 0.00 | 0.81 |
| | 10000 | 0.05 | 1220.34 | 0.16 | 0.95 | 0.00 | 0.89 | 0.90 | 0.00 | 0.82 |
| 0.5 | 1000 | 0.17 | 98.94 | 0.23 | 0.90 | 0.00 | 0.84 | 0.84 | 0.00 | 0.76 |
| | 5000 | 0.07 | 495.33 | 0.24 | 0.95 | 0.00 | 0.85 | 0.89 | 0.00 | 0.77 |
| | 10000 | 0.05 | 990.39 | 0.24 | 0.95 | 0.00 | 0.85 | 0.90 | 0.00 | 0.77 |
| 0.7 | 1000 | 0.15 | 65.81 | 0.26 | 0.91 | 0.00 | 0.81 | 0.84 | 0.00 | 0.72 |
| | 5000 | 0.06 | 329.11 | 0.27 | 0.95 | 0.00 | 0.81 | 0.89 | 0.00 | 0.73 |
| | 10000 | 0.04 | 657.98 | 0.26 | 0.95 | 0.00 | 0.82 | 0.91 | 0.00 | 0.74 |
| 0.9 | 1000 | 0.12 | 23.81 | 0.15 | 0.82 | 0.00 | 0.78 | 0.76 | 0.00 | 0.69 |
| | 5000 | 0.04 | 119.01 | 0.14 | 0.93 | 0.00 | 0.78 | 0.88 | 0.00 | 0.69 |
| | 10000 | 0.03 | 238.42 | 0.15 | 0.94 | 0.00 | 0.77 | 0.89 | 0.00 | 0.68 |

(vi) (D-HB2) constructing the confidence interval using the hybrid bootstrap based on $\sqrt{n}(\tilde{\xi}_b - \mathbb{E}[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}]) | \mathbf{X}, \mathbf{Y}$.

The simulation studies were conducted based on the Gaussian rotation model, where (X, Y) are bivariate Gaussian with mean 0 and covariance matrix Σ , defined as

$$\Sigma = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}, \quad \text{with } \rho \in (-1, 1).$$

We investigate the performance of different methods for estimating ξ_n 's variance and inferring ξ using various sample sizes $n = 1,000, 5,000, 10,000$ and population correlations $\rho = 0, 0.3, 0.5, 0.7, 0.9$. For the bootstrap procedure, we adopt a bootstrap size of 5,000 and simulate 5,000 replications to compute the square roots of the mean squared errors (RMSEs) in estimating $n \text{Var}(\xi_n)$ (of limits 0.4, 0.46, 0.51, 0.47, and 0.24 as ρ changes from 0 to 0.9), as well as the empirical coverage probabilities with the nominal level $\alpha = 0.05$ or 0.1.

Table 1 presents the simulation results, demonstrating that regardless of the strength of dependence characterized by ρ , the bootstrap methods consistently produce erroneous variance estimators and inaccurate confidence intervals. On the other hand, the method proposed by [Lin and Han \(2022\)](#) performs well for large n .

3 Proof of Theorem 2.1

We first introduce some necessary notation. For any integers $n \geq 1$, let $\llbracket n \rrbracket := \{1, 2, \dots, n\}$. A set consisting of distinct elements x_1, \dots, x_n is written as either $\{x_1, \dots, x_n\}$ or $\{x_i\}_{i=1}^n$, and its cardinality is written by $|\{x_i\}_{i=1}^n|$. The corresponding sequence is denoted by $[x_1, \dots, x_n]$ or $[x_i]_{i=1}^n$. For any $a, b \in \mathbb{R}$, write $a \vee b := \max\{a, b\}$ and $a \wedge b := \min\{a, b\}$. For any two real sequences $\{a_n\}$ and $\{b_n\}$, write $a_n = O(b_n)$ if $\limsup |a_n/b_n|$ is bounded.

Given a sample $\{(X_i, Y_i)\}_{i=1}^n$, let $W_i = W_{b,i}$ be the number of times (X_i, Y_i) appears in the bootstrap sample $(\mathbf{X}_b, \mathbf{Y}_b)$; (W_1, \dots, W_n) then follows a multinomial distribution with parameter $(n, 1/n, \dots, 1/n)$, which we denote by $M_n(n; 1/n, \dots, 1/n)$. Introduce

$$\tilde{R}_i := \mathbf{1}(W_i > 0) \cdot \sum_{j=1}^n W_j \mathbf{1}(Y_j \leq Y_i)$$

to represent the rank of Y_i in the bootstrap sample if (X_i, Y_i) appears in the bootstrap sample; note that \tilde{R}_i is different from $R_{b,i}$ as they use different indexing system.

Recall that $\{[i], i = 1, \dots, n\}$ is defined to be the unique rearrangement of $\{1, \dots, n\}$ such that

$$X_{[1]} < \dots < X_{[n]}$$

as F_X has been assumed to be continuous. One could then define

$$\hat{\xi}_b := 1 - \frac{3}{n^2 - 1} \sum_{i=1}^{n-1} \mathbf{1}(W_{[i]} > 0) \left| \tilde{R}_{[i+k(i)]} - \tilde{R}_{[i]} \right|, \quad (3.1)$$

where

$$k(i) := \begin{cases} \min\{k : W_{[i+k]} > 0\}, & \text{if the left set is nonempty,} \\ 0, & \text{otherwise.} \end{cases}$$

Comparing (3.1) with (2.3), we have the following key identity,

$$\sum_{i=1}^{n-1} \mathbf{1}(W_{[i]} > 0) \left| \tilde{R}_{[i+k(i)]} - \tilde{R}_{[i]} \right| = \sum_{i=1}^{n-1} \left| R_{b,[i+1]_b} - R_{b,[i]_b} \right|, \quad (3.2)$$

where $\{[i]_b, i = 1, \dots, n\}$, introduced in (2.2), is defined to be a rearrangement of $\{1, \dots, n\}$ such that $X_{b,[1]_b} \leq \dots \leq X_{b,[n]_b}$, with ties in $\{X_{b,i}\}_{i=1}^n$ broken *in arbitrary way*.

Why does (3.2) hold? The reason is that, no matter how we adjust for the ties on $\{X_{b,i}\}_{i=1}^n$, it is always true that

$$Y_{b,i} = Y_{b,j} \text{ whenever } X_{b,i} = X_{b,j}$$

since they are generated by the same unit from the bootstrap procedure. Then

$$R_{b,[i+1]_b} - R_{b,[i]_b} \text{ is nonzero only if } X_{b,[i+1]_b} \neq X_{b,[i]_b},$$

or equivalently,

$$X_{b,[i+1]_b} \text{ is the right nearest neighbor (NN) of } X_{b,[i]_b}$$

in the bootstrap sample. Note that the bootstrap procedure will not change the order of X_1, \dots, X_n . In order to find the right NN of a unit in the bootstrap sample, it is then equivalent to finding the

right NN in the original sample among those that also appear in the bootstrap sample.

The difference between (3.1) and (2.3) is, then, fully due to the replacement of a random quantity $n/[2\sum_{i=1}^n L_{b,[i]_b}(n - L_{b,[i]_b})]$ by a fixed quantity $3/(n^2 - 1)$, corresponding to the simple form of ξ_n in (2.1). The following lemma shows that this difference is \sqrt{n} -negligible. All lemmas stated in the sequel are under the conditions of Theorem 2.1.

Lemma 3.1. *It holds true that*

$$\lim_{n \rightarrow \infty} n\mathbb{E}[\widehat{\xi}_b - \widetilde{\xi}_b]^2 = 0. \quad (3.3)$$

With lemma 3.1, it then suffices to consider the asymptotic behavior of $\widehat{\xi}_b$. However, (3.1) cannot be further simplified; for instance, one cannot replace \widetilde{R}_i 's by R_i 's. This places more difficulty to analyzing (3.1). Our route to proving Theorem 2.1 is instead built on a seemingly more complicated reformulation of $\widehat{\xi}_b$:

$$\begin{aligned} \widehat{\xi}_b = 1 - \frac{3}{n^2 - 1} \sum_{i=1}^{n-1} \mathbb{1}(W_{[i]} > 0) \sum_{k=1}^{n-i} \left\{ \left[\sum_{j=1}^n W_j \mathbb{1}(Y_{[i]} \wedge Y_{[i+k]} < Y_j < Y_{[i]} \vee Y_{[i+k]}) \right] \right. \\ \left. \mathbb{1}(W_{[i+\ell]} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{[i+k]} > 0) \right\}. \end{aligned}$$

The following lemma establishes the asymptotic equivalence between $\widehat{\xi}_b$ and $\bar{\xi}_b$.

Lemma 3.2. *We have*

$$|\bar{\xi}_b - \widehat{\xi}_b| \leq \frac{6n}{n^2 - 1} = O(n^{-1}).$$

With Lemma 3.2, it then suffices to consider $\bar{\xi}_b$. Conditional on \mathbf{X}, \mathbf{Y} , without loss of generality, we can assume $[X_i]_{i=1}^n$ is strictly increasing. One could then simply write

$$\begin{aligned} \bar{\xi}_b = 1 - \frac{3}{n^2 - 1} \sum_{i=1}^{n-1} \mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left\{ \left[\sum_{j=1}^n W_j \mathbb{1}(Y_i \wedge Y_{i+k} < Y_j < Y_i \vee Y_{i+k}) \right] \right. \\ \left. \mathbb{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \right\}. \end{aligned}$$

For any $i \in \llbracket n-1 \rrbracket$ and $k \in \llbracket n-i \rrbracket$, let

$$S_{i,k} := \left\{ j \in \llbracket n \rrbracket : Y_i \wedge Y_{i+k} < Y_j < Y_i \vee Y_{i+k} \right\} \setminus \left\{ j \in \llbracket n \rrbracket : i < j < i+k \right\}.$$

Then by the event $\{W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket\}$, we have

$$\bar{\xi}_b = 1 - \frac{3}{n^2 - 1} \sum_{i=1}^{n-1} \mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{j \in S_{i,k}} W_j \right] \mathbb{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0).$$

To continue the proof, we introduce the following lemma on some properties of the multinomial distribution.

Lemma 3.3. *Recall that $(W_1, \dots, W_n) \sim M_n(n; 1/n, \dots, 1/n)$. Let $\mathcal{S}, \mathcal{S}', \mathcal{T} \subset \llbracket n \rrbracket$ be mutually disjoint. Then*

$$\mathbb{E} \left[\left(\sum_{s \in \mathcal{S}} W_s \right) \mathbb{1} \left(\sum_{t \in \mathcal{T}} W_t = 0 \right) \right] = |\mathcal{S}| \left(1 - \frac{|\mathcal{T}|}{n} \right)^{n-1};$$

$$\begin{aligned} \mathbb{E}\left[\left(\sum_{s \in \mathcal{S}} W_s\right)^2 \mathbf{1}\left(\sum_{t \in \mathcal{T}} W_t = 0\right)\right] &= |\mathcal{S}| \left(1 - \frac{|\mathcal{T}|}{n}\right)^{n-1} + |\mathcal{S}|^2 \left(1 - \frac{1}{n}\right) \left(1 - \frac{|\mathcal{T}|}{n}\right)^{n-2}; \\ \text{and } \mathbb{E}\left[\left(\sum_{s \in \mathcal{S}} W_s\right) \left(\sum_{s \in \mathcal{S}'} W_s\right) \mathbf{1}\left(\sum_{t \in \mathcal{T}} W_t = 0\right)\right] &= |\mathcal{S}| |\mathcal{S}'| \left(1 - \frac{1}{n}\right) \left(1 - \frac{|\mathcal{T}|}{n}\right)^{n-2}. \end{aligned}$$

Note that $S_{i,k}$ is disjoint with $\{i, i+1, \dots, i+k\}$ almost surely since $[Y_i]_{i=1}^n$ are distinct with probability one. Then we can use Lemma 3.3 to establish the following lemma on the expectation of $\bar{\xi}_b$ conditional on the original sample.

Lemma 3.4. *We have*

$$\mathbb{E}[\bar{\xi}_b | \mathbf{X}, \mathbf{Y}] = 1 - \frac{3}{n^2 - 1} \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} |S_{i,k}| \left[\left(1 - \frac{k-1}{n}\right)^{n-1} - 2 \left(1 - \frac{k}{n}\right)^{n-1} + \left(1 - \frac{k+1}{n}\right)^{n-1} \right].$$

For any $j \in \llbracket n \rrbracket \setminus \{i, i+1, \dots, i+k\}$, by independence of X and Y ,

$$\mathbb{P}(Y_i \wedge Y_{i+k} < Y_j < Y_i \vee Y_{i+k}) = 1/3,$$

and thus

$$\mathbb{E}[|S_{i,k}|] = (n - k - 1)/3.$$

Some calculations then yield the following lemma on the expectation of $\bar{\xi}_b$.

Lemma 3.5. *We have*

$$\mathbb{E}[\bar{\xi}_b] = \frac{1}{e} + O\left(\frac{1}{n}\right).$$

The variance of $\bar{\xi}_b$ conditional on the original sample can be established in a similar way by using Lemma 3.3. For any $i, j \in \llbracket n-1 \rrbracket$, $k \in \llbracket n-i \rrbracket$ and $\ell \in \llbracket n-j \rrbracket$, we write

$$\begin{aligned} S_{i,k \cap j, \ell} &= \left\{ t \in \llbracket n \rrbracket : (Y_i \wedge Y_{i+k}) \vee (Y_j \wedge Y_{j+\ell}) < Y_t < (Y_i \vee Y_{i+k}) \wedge (Y_j \vee Y_{j+\ell}) \right\} \\ &\quad \setminus \left\{ t \in \llbracket n \rrbracket : i < t < i+k, j < t < j+\ell \right\}; \\ S_{i,k \setminus j, \ell} &= \left\{ t \in \llbracket n \rrbracket : Y_i \wedge Y_{i+k} < Y_t < Y_i \vee Y_{i+k}, Y_t > Y_j \vee Y_{j+\ell} \text{ or } Y_t < Y_j \wedge Y_{j+\ell} \right\} \\ &\quad \setminus \left\{ t \in \llbracket n \rrbracket : i < t < i+k, j < t < j+\ell \right\}. \end{aligned}$$

For any four integers n, m, k, ℓ , define

$$\begin{aligned} a_{n,m,k,\ell} &:= \left(1 - \frac{k+\ell-2}{n}\right)^m - 3 \left(1 - \frac{k+\ell-1}{n}\right)^m + 3 \left(1 - \frac{k+\ell}{n}\right)^m - \left(1 - \frac{k+\ell+1}{n}\right)^m; \\ b_{n,m,k,\ell} &:= \left(1 - \frac{k+\ell-2}{n}\right)^m - 4 \left(1 - \frac{k+\ell-1}{n}\right)^m + 6 \left(1 - \frac{k+\ell}{n}\right)^m - 4 \left(1 - \frac{k+\ell+1}{n}\right)^m \\ &\quad + \left(1 - \frac{k+\ell+2}{n}\right)^m; \\ c_{n,m,k} &:= \left(1 - \frac{k-1}{n}\right)^m - 2 \left(1 - \frac{k}{n}\right)^m + \left(1 - \frac{k+1}{n}\right)^m. \end{aligned}$$

We can then use Lemma 3.3 to establish the following result on the variance of $\bar{\xi}_b$ conditional on the original sample.

Lemma 3.6.

$$\begin{aligned}
& \text{Var}[\bar{\xi}_b | \mathbf{X}, \mathbf{Y}] \\
&= \frac{9}{(n^2 - 1)^2} \left[\sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \left(|S_{i,k}| c_{n,n-1,k} + |S_{i,k}|^2 c_{n,n-2,k} (1 - 1/n) \right) \right. \\
&+ 2 \sum_{i,j=1}^{n-1} \sum_{i < j} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-j} \left(|S_{i,k \cap j, \ell}| [a_{n,n-1,k,\ell} \mathbf{1}(k = j - i) + b_{n,n-1,k,\ell} \mathbf{1}(k < j - i)] \right. \\
&+ (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|) (|S_{i,k \cap j, \ell}| + |S_{j,\ell \setminus i,k}|) [a_{n,n-2,k,\ell} \mathbf{1}(k = j - i) + b_{n,n-2,k,\ell} \mathbf{1}(k < j - i)] (1 - 1/n) \\
&+ [\mathbf{1}(j \in S_{i,k}) + \mathbf{1}(j + \ell \in S_{i,k})] (|S_{i,k \cap j, \ell}| + |S_{j,\ell \setminus i,k}|) a_{n,n-2,k,\ell} \mathbf{1}(k < j - i) (1 - 1/n) \\
&+ [\mathbf{1}(i \in S_{j,\ell}) + \mathbf{1}(i + k \in S_{j,\ell})] (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|) a_{n,n-2,k,\ell} \mathbf{1}(k < j - i) (1 - 1/n) \left. \right) \\
&- \left(\sum_{i=1}^{n-1} \sum_{k=1}^{n-i} |S_{i,k}| c_{n,n-1,k} \right)^2 \left. \right] + O\left(\frac{1}{n^2}\right).
\end{aligned}$$

The following lemma further calculates the expectations of those cardinalities present in Lemma 3.6.

Lemma 3.7. Assume $Z_1, \dots, Z_n, Y_1, Y_2, Y_3, Y_4$ are i.i.d. from a continuous distribution. Define

$$\begin{aligned}
S_{12} &:= \sum_{i=1}^n \mathbf{1}\left(Y_1 \wedge Y_2 < Z_i < Y_1 \vee Y_2\right), \\
S_{1234} &:= \sum_{i=1}^n \mathbf{1}\left((Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4)\right), \\
S_{12 \setminus 34} &:= \sum_{i=1}^n \mathbf{1}\left(Y_1 \wedge Y_2 < Z_i < Y_1 \vee Y_2, \left\{Z_i > Y_3 \vee Y_4 \text{ or } Z_i < Y_3 \wedge Y_4\right\}\right), \\
S_{34 \setminus 12} &:= \sum_{i=1}^n \mathbf{1}\left(Y_3 \wedge Y_4 < Z_i < Y_3 \vee Y_4, \left\{Z_i > Y_1 \vee Y_2 \text{ or } Z_i < Y_1 \wedge Y_2\right\}\right), \\
S_{1223} &:= \sum_{i=1}^n \mathbf{1}\left((Y_1 \wedge Y_2) \vee (Y_2 \wedge Y_3) < Z_i < (Y_1 \vee Y_2) \wedge (Y_2 \vee Y_3)\right), \\
S_{12 \setminus 23} &:= \sum_{i=1}^n \mathbf{1}\left(Y_1 \wedge Y_2 < Z_i < Y_1 \vee Y_2, \left\{Z_i > Y_2 \vee Y_3 \text{ or } Z_i < Y_2 \wedge Y_3\right\}\right), \\
S_{23 \setminus 12} &:= \sum_{i=1}^n \mathbf{1}\left(Y_2 \wedge Y_3 < Z_i < Y_2 \vee Y_3, \left\{Z_i > Y_1 \vee Y_2 \text{ or } Z_i < Y_1 \wedge Y_2\right\}\right).
\end{aligned}$$

We then have

$$\begin{aligned}
\mathbb{E}[S_{12}] &= \frac{1}{3}n, \quad \mathbb{E}[S_{12}^2] = \frac{1}{6}n^2 + \frac{1}{6}n, \quad \mathbb{E}[S_{1234}] = \frac{2}{15}n, \quad \mathbb{E}[S_{1234}^2] = \frac{2}{45}n^2 + \frac{4}{45}n, \\
\mathbb{E}[S_{1234}S_{12 \setminus 34}] &= \mathbb{E}[S_{1234}S_{34 \setminus 12}] = \mathbb{E}[S_{12 \setminus 34}S_{34 \setminus 12}] = \frac{1}{45}n^2 - \frac{1}{45}n, \quad \mathbb{E}[S_{1223}] = \frac{1}{6}n, \\
\mathbb{E}[S_{1223}^2] &= \frac{1}{15}n^2 + \frac{1}{10}n, \quad \mathbb{E}[S_{1223}S_{12 \setminus 23}] = \mathbb{E}[S_{1223}S_{23 \setminus 12}] = \mathbb{E}[S_{12 \setminus 23}S_{23 \setminus 12}] = \frac{1}{60}n^2 - \frac{1}{60}n, \\
\mathbb{E}[\mathbf{1}(Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2)S_{1234}] &= \frac{1}{15}n, \quad \text{and } \mathbb{E}[\mathbf{1}(Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2)S_{34 \setminus 12}] = \frac{1}{30}n.
\end{aligned}$$

Applying Lemma 3.7 to Lemma 3.6, we have the following lemma on the expectation of the conditional variance of $\bar{\xi}_b$.

Lemma 3.8. *We have*

$$\limsup_{n \rightarrow \infty} nE[\text{Var}[\bar{\xi}_b | \mathbf{X}, \mathbf{Y}]] \leq \frac{3}{5} - \frac{8}{5} \frac{1}{e^2} \approx 0.3835 < \frac{2}{5}.$$

A more involved analysis of $\text{Var}[\sum_{i=1}^{n-1} \sum_{k=1}^{n-i} |S_{i,k}| c_{n,n-1,k}]$ in Lemma 3.6 can lead to the exact limit value of $nE[\text{Var}[\bar{\xi}_b | \mathbf{X}, \mathbf{Y}]]$ following the same proof techniques as in Lemma 3.8. However, a limit superior of $nE[\text{Var}[\bar{\xi}_b | \mathbf{X}, \mathbf{Y}]]$ smaller than $2/5$ appears to be sufficient for establishing the bootstrap inconsistency.

Combining Lemmas 3.5 and 3.8, a proof of Theorem 2.1 is finally ready.

Proof of Theorem 2.1. By Lemmas 3.1, 3.5, and 3.8, one has

$$\lim_{n \rightarrow \infty} E[\tilde{\xi}_b] = \frac{1}{e}, \quad (3.4)$$

and

$$\limsup_{n \rightarrow \infty} nE[\text{Var}[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}]] \leq \frac{3}{5} - \frac{8}{5} \frac{1}{e^2} \approx 0.3835 < \frac{2}{5}. \quad (3.5)$$

Proposition 2.1, on the other hand, shows

$$\sqrt{n}\xi_n \xrightarrow{d} N(0, 2/5),$$

where \xrightarrow{d} stands for convergence in distribution.

Proof of the first statement. For any $\epsilon > 0$, since $\sqrt{n}\xi_n$ is bounded in probability by the above central limit theorem, one can find $C_1 = C_1(\epsilon) > 0$ such that $P(|\sqrt{n}\xi_n| > C_1) < \epsilon$ for all n sufficiently large.

Note that for any constant $C_2 > 0$,

$$\begin{aligned} E[\tilde{\xi}_b] &= E[E[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}]] \\ &= E[E[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}] \mathbf{1}(E[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}] > C_2)] + E[E[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}] \mathbf{1}(E[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}] \leq C_2)] \\ &\leq E[E[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}] \mathbf{1}(E[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}] > C_2)] + C_2. \end{aligned}$$

By (3.4) and the fact that $E[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}]$ is universally bounded for all \mathbf{X}, \mathbf{Y} , we can take $C_2 < e^{-1}$ and then for all n sufficiently large,

$$P(E[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}] > C_2) \geq 2\epsilon,$$

for some $\epsilon > 0$.

Then for all n sufficiently large, with probability at least ϵ ,

$$nE[(\tilde{\xi}_b - \xi_n)^2 | \mathbf{X}, \mathbf{Y}] \geq n(C_2^2 - 2C_2C_1/\sqrt{n}).$$

This implies that $nE[(\tilde{\xi}_b - \xi_n)^2 | \mathbf{X}, \mathbf{Y}]$ does not converge in probability to any constant.

If, on the other hand, $n \text{Var}[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}]$ converges to $2/5$ in probability, then by the Portmanteau lemma,

$$\liminf_{n \rightarrow \infty} nE[\text{Var}[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}]] \geq \frac{2}{5},$$

which contradicts (3.5). Therefore $n \text{Var}[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}]$ does not converge to 2/5 in probability.

Proof of the second statement. We adopt the argument of [Abadie and Imbens \(2008\)](#). For all sufficiently large n , we can establish in a similar way as above that, with probability at least ϵ ,

$$\sqrt{n}(\tilde{\xi}_b - \xi_n) \geq \sqrt{n}(C_2 - C_1/\sqrt{n}).$$

This shows that $\sqrt{n}(\tilde{\xi}_b - \xi_n)$ cannot converge in distribution to $N(0, 2/5)$.

If $\sqrt{n}(\tilde{\xi}_b - \mathbb{E}[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}])$ converges in distribution to $N(0, 2/5)$, then

$$\liminf_{n \rightarrow \infty} n \text{Var}[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}] \geq 2/5$$

by the Portmanteau lemma. If the convergence in distribution holds for almost all sequences X_1, X_2, \dots and Y_1, Y_2, \dots , then

$$\liminf_{n \rightarrow \infty} n \mathbb{E}[\text{Var}[\tilde{\xi}_b | \mathbf{X}, \mathbf{Y}]] \geq 2/5,$$

which contradicts (3.5). □

4 Auxiliary proofs

Notation. For any two real sequences $\{a_n\}$ and $\{b_n\}$, write $a_n \lesssim b_n$ (or equivalently, $b_n \gtrsim a_n$) if there exists a universal constant $C > 0$ such that $|a_n/b_n| \leq C$ for all sufficiently large n .

4.1 Proof of Lemma 3.1

Proof of Lemma 3.1. Note that for any $i \in \llbracket n-1 \rrbracket$, if $X_{b, [i]_b} = X_{b, [i+1]_b}$, we have $Y_{b, [i]_b} = Y_{b, [i+1]_b}$ and then $R_{b, [i]_b} = R_{b, [i+1]_b}$ from the properties of bootstrap samples. Then $|R_{b, [i+1]_b} - R_{b, [i]_b}|$ is nonzero if and only if $X_{b, [i]_b} < X_{b, [i+1]_b}$. For any $i \in \llbracket n \rrbracket$ such that $W_{[i]} = 0$, $(X_{[i]}, Y_{[i]})$ will not appear in the bootstrap sample. For any $i \in \llbracket n \rrbracket$ such that $W_{[i]} > 0$, the right nearest neighbor of $X_{[i]}$ in the bootstrap sample with strictly larger value is $X_{[i+k]}$, where k is the smallest positive interger such that $W_{[i+k]} > 0$. If such k does not exist, $X_{[i]}$ is the largest value in the bootstrap sample. Therefore we can write $\tilde{\xi}_b$ equivalently as

$$\tilde{\xi}_b := 1 - \frac{n}{2 \sum_{i=1}^n W_i \tilde{L}_i (n - \tilde{L}_i)} \sum_{i=1}^{n-1} \mathbf{1}(W_{[i]} > 0) \left| \tilde{R}_{[i+k(i)]} - \tilde{R}_{[i]} \right|,$$

where

$$\tilde{L}_i = \sum_{j=1}^n W_j \mathbf{1}(Y_j \geq Y_i).$$

Now since $\tilde{R}_i \leq n$ for any $i \in \llbracket n \rrbracket$, we have

$$\begin{aligned} |\tilde{\xi}_b - \hat{\xi}_b| &= \left| \left(\frac{n}{2 \sum_{i=1}^n W_i \tilde{L}_i (n - \tilde{L}_i)} - \frac{3}{n^2 - 1} \right) \sum_{i=1}^{n-1} \mathbf{1}(W_{[i]} > 0) \left| \tilde{R}_{[i+k(i)]} - \tilde{R}_{[i]} \right| \right| \\ &\leq 2n(n-1) \left| \frac{n}{2 \sum_{i=1}^n W_i \tilde{L}_i (n - \tilde{L}_i)} - \frac{3}{n^2 - 1} \right|. \end{aligned}$$

By a careful use of the Efron-Stein inequality, we have the following lemma.

Lemma 4.1. For almost all $\mathbf{X} = (X_1, \dots, X_n)$ and $\mathbf{Y} = (Y_1, \dots, Y_n)$, for any $\epsilon = \epsilon(n) > 0$,

$$\mathbb{P}\left(\left|\sum_{i=1}^n W_i \tilde{L}_i (n - \tilde{L}_i) - \frac{n^3}{6}\right| > \epsilon \mid \mathbf{X}, \mathbf{Y}\right) \lesssim \epsilon^{-2} n^3.$$

Note that if $|\sum_{i=1}^n W_i \tilde{L}_i (n - \tilde{L}_i) - n^3/6| \leq \epsilon$ for some $\epsilon/n^3 \rightarrow 0$, then

$$\left|\frac{1}{\sum_{i=1}^n W_i \tilde{L}_i (n - \tilde{L}_i)} - \frac{6}{n(n^2 - 1)}\right| \lesssim \frac{1}{n^6} \left|n(n^2 - 1) - 6 \sum_{i=1}^n W_i \tilde{L}_i (n - \tilde{L}_i)\right| \lesssim \frac{1}{n^6} (n + \epsilon),$$

and then

$$|\tilde{\xi}_b - \hat{\xi}_b| \lesssim \frac{1}{n^3} \epsilon + \frac{1}{n^2}.$$

Also note that we always have $|\tilde{\xi}_b - \hat{\xi}_b| \lesssim 1$. By Lemma 4.1, if $\epsilon/n^3 \rightarrow 0$, then for almost all \mathbf{X}, \mathbf{Y} ,

$$\begin{aligned} \mathbb{E}[|\tilde{\xi}_b - \hat{\xi}_b|^2 \mid \mathbf{X}, \mathbf{Y}] &\lesssim \mathbb{P}\left(\left|\sum_{i=1}^n W_i \tilde{L}_i (n - \tilde{L}_i) - \frac{n^3}{6}\right| > \epsilon \mid \mathbf{X}, \mathbf{Y}\right) + \frac{1}{n^6} \epsilon^2 + \frac{1}{n^4} \\ &\lesssim n^3 \frac{1}{\epsilon^2} + \frac{1}{n^6} \epsilon^2 + \frac{1}{n^4}. \end{aligned}$$

Taking $\epsilon \asymp n^{9/4}$, we obtain

$$\mathbb{E}[|\tilde{\xi}_b - \hat{\xi}_b|^2 \mid \mathbf{X}, \mathbf{Y}] \lesssim n^{-3/2},$$

and then the proof is complete. \square

4.2 Proof of Lemma 3.2

Proof of Lemma 3.2. By the definition of $k(i)$ for $i \in \llbracket n \rrbracket$, we have for any $i \in \llbracket n \rrbracket$,

$$\left|\tilde{R}_{[i+k(i)]} - \tilde{R}_{[i]}\right| = \sum_{k=1}^{n-i} \left|\tilde{R}_{[i+k]} - \tilde{R}_{[i]}\right| \mathbf{1}\left(W_{[i+\ell]} = 0, \forall \ell \in \llbracket k-1 \rrbracket\right) \mathbf{1}\left(W_{[i+k]} > 0\right).$$

From the definition of $[\tilde{R}_i]_{i=1}^n$, for any $i, k \in \llbracket n \rrbracket$ with $i \neq k$,

$$\left|\tilde{R}_k - \tilde{R}_i\right| = \left|\sum_{j=1}^n W_j \mathbf{1}(Y_j \leq Y_k) - \sum_{j=1}^n W_j \mathbf{1}(Y_j \leq Y_i)\right| = \sum_{j=1}^n W_j \mathbf{1}(Y_i \wedge Y_k < Y_j \leq Y_i \vee Y_k).$$

Combining the above two with (3.1) yields

$$\begin{aligned} \hat{\xi}_b &= 1 - \frac{3}{n^2 - 1} \sum_{i=1}^{n-1} \mathbf{1}(W_{[i]} > 0) \sum_{k=1}^{n-i} \left[\sum_{j=1}^n W_j \mathbf{1}(Y_{[i]} \wedge Y_{[i+k]} < Y_j \leq Y_{[i]} \vee Y_{[i+k]})\right] \\ &\quad \mathbf{1}\left(W_{[i+\ell]} = 0, \forall \ell \in \llbracket k-1 \rrbracket\right) \mathbf{1}\left(W_{[i+k]} > 0\right). \end{aligned}$$

We then have

$$|\bar{\xi}_b - \hat{\xi}_b| \leq \frac{3}{n^2 - 1} \sum_{i=1}^{n-1} \mathbf{1}(W_{[i]} > 0) \sum_{k=1}^{n-i} (W_{[i]} + W_{[i+k]}) \mathbf{1}\left(W_{[i+\ell]} = 0, \forall \ell \in \llbracket k-1 \rrbracket\right) \mathbf{1}\left(W_{[i+k]} > 0\right)$$

$$\leq \frac{6}{n^2 - 1} \sum_{i=1}^n W_i = \frac{6n}{n^2 - 1} = O\left(\frac{1}{n}\right),$$

since among all the units with positive weights, each unit only has at most one right NN and can be the right NN of at most one unit. \square

4.3 Proof of Lemma 3.3

Proof of Lemma 3.3. Note that $(W_1, \dots, W_n) \sim M_n(n; 1/n, \dots, 1/n)$, then since \mathcal{S} and \mathcal{T} are disjoint,

$$\sum_{s \in \mathcal{S}} W_s \mid \sum_{t \in \mathcal{T}} W_t \sim \text{Bin}\left(n - \sum_{t \in \mathcal{T}} W_t, \frac{|\mathcal{S}|/n}{1 - |\mathcal{T}|/n}\right), \quad \sum_{t \in \mathcal{T}} W_t \sim \text{Bin}\left(n, \frac{|\mathcal{T}|}{n}\right).$$

We then obtain

$$\begin{aligned} \mathbb{E}\left[\left(\sum_{s \in \mathcal{S}} W_s\right) \mathbb{1}\left(\sum_{t \in \mathcal{T}} W_t = 0\right)\right] &= \mathbb{E}\left[\sum_{s \in \mathcal{S}} W_s \mid \sum_{t \in \mathcal{T}} W_t = 0\right] \mathbb{P}\left(\sum_{t \in \mathcal{T}} W_t = 0\right) \\ &= n \frac{|\mathcal{S}|/n}{1 - |\mathcal{T}|/n} \left(1 - \frac{|\mathcal{T}|}{n}\right)^n = |\mathcal{S}| \left(1 - \frac{|\mathcal{T}|}{n}\right)^{n-1}, \end{aligned}$$

and

$$\begin{aligned} \mathbb{E}\left[\left(\sum_{s \in \mathcal{S}} W_s\right)^2 \mathbb{1}\left(\sum_{t \in \mathcal{T}} W_t = 0\right)\right] &= \mathbb{E}\left[\left(\sum_{s \in \mathcal{S}} W_s\right)^2 \mid \sum_{t \in \mathcal{T}} W_t = 0\right] \mathbb{P}\left(\sum_{t \in \mathcal{T}} W_t = 0\right) \\ &= \left[n \frac{|\mathcal{S}|/n}{1 - |\mathcal{T}|/n} + n(n-1) \left(\frac{|\mathcal{S}|/n}{1 - |\mathcal{T}|/n}\right)^2\right] \left(1 - \frac{|\mathcal{T}|}{n}\right)^n \\ &= |\mathcal{S}| \left(1 - \frac{|\mathcal{T}|}{n}\right)^{n-1} + |\mathcal{S}|^2 \left(1 - \frac{1}{n}\right) \left(1 - \frac{|\mathcal{T}|}{n}\right)^{n-2}. \end{aligned}$$

Note that from the properties of the multinomial distribution,

$$\sum_{s \in \mathcal{S}} W_s, \sum_{s \in \mathcal{S}'} W_s \mid \sum_{s \in \mathcal{S} \cup \mathcal{S}'} W_s \sim M_2\left(\sum_{s \in \mathcal{S} \cup \mathcal{S}'} W_s; \frac{|\mathcal{S}|}{|\mathcal{S}| + |\mathcal{S}'|}, \frac{|\mathcal{S}'|}{|\mathcal{S}| + |\mathcal{S}'|}\right).$$

It then holds that

$$\begin{aligned} \mathbb{E}\left[\left(\sum_{s \in \mathcal{S}} W_s\right) \left(\sum_{s \in \mathcal{S}'} W_s\right) \mathbb{1}\left(\sum_{t \in \mathcal{T}} W_t = 0\right)\right] &= \mathbb{E}\left[\left(\sum_{s \in \mathcal{S}} W_s\right) \left(\sum_{s \in \mathcal{S}'} W_s\right) \mid \sum_{t \in \mathcal{T}} W_t = 0\right] \mathbb{P}\left(\sum_{t \in \mathcal{T}} W_t = 0\right) \\ &= \mathbb{E}\left[\mathbb{E}\left[\left(\sum_{s \in \mathcal{S}} W_s\right) \left(\sum_{s \in \mathcal{S}'} W_s\right) \mid \sum_{s \in \mathcal{S} \cup \mathcal{S}'} W_s, \sum_{t \in \mathcal{T}} W_t = 0\right] \mid \sum_{t \in \mathcal{T}} W_t = 0\right] \mathbb{P}\left(\sum_{t \in \mathcal{T}} W_t = 0\right) \\ &= \mathbb{E}\left[\left[\left(\sum_{s \in \mathcal{S} \cup \mathcal{S}'} W_s\right)^2 - \left(\sum_{s \in \mathcal{S} \cup \mathcal{S}'} W_s\right)\right] \frac{|\mathcal{S}||\mathcal{S}'|}{(|\mathcal{S}| + |\mathcal{S}'|)^2} \mid \sum_{t \in \mathcal{T}} W_t = 0\right] \mathbb{P}\left(\sum_{t \in \mathcal{T}} W_t = 0\right) \\ &= n(n-1) \left(\frac{(|\mathcal{S}| + |\mathcal{S}'|)/n}{1 - |\mathcal{T}|/n}\right)^2 \frac{|\mathcal{S}||\mathcal{S}'|}{(|\mathcal{S}| + |\mathcal{S}'|)^2} \left(1 - \frac{|\mathcal{T}|}{n}\right)^n \\ &= |\mathcal{S}||\mathcal{S}'| \left(1 - \frac{1}{n}\right) \left(1 - \frac{|\mathcal{T}|}{n}\right)^{n-2}. \end{aligned}$$

The proof is thus complete. \square

4.4 Proof of Lemma 3.4

Proof of Lemma 3.4. For any $i \in \llbracket n-1 \rrbracket$ and $k \in \llbracket n-i \rrbracket$, consider

$$\left(\sum_{j \in S_{i,k}} W_j \right) \mathbf{1}(W_i > 0) \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbf{1}(W_{i+k} > 0).$$

We decompose the above term as

$$\begin{aligned} & \mathbf{1}(W_i > 0) \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbf{1}(W_{i+k} > 0) \\ &= \left[1 - \mathbf{1}(W_i = 0) \right] \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \left[1 - \mathbf{1}(W_{i+k} = 0) \right] \\ &= \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) - \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket \cup \{0\}) \\ &\quad - \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k \rrbracket) + \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k \rrbracket \cup \{0\}). \end{aligned}$$

Note that for any $i \in \llbracket n-1 \rrbracket$ and $k \in \llbracket n-i \rrbracket$, $S_{i,k}$ is disjoint with $\{i, i+1, \dots, i+k\}$. Then, by Lemma 3.3,

$$\begin{aligned} \mathbb{E} \left[\left(\sum_{j \in S_{i,k}} W_j \right) \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k \rrbracket) \right] &= |S_{i,k}| \left(1 - \frac{k}{n} \right)^{n-1}, \\ \mathbb{E} \left[\left(\sum_{j \in S_{i,k}} W_j \right) \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \right] &= |S_{i,k}| \left(1 - \frac{k-1}{n} \right)^{n-1}, \\ \mathbb{E} \left[\left(\sum_{j \in S_{i,k}} W_j \right) \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket \cup \{0\}) \right] &= |S_{i,k}| \left(1 - \frac{k}{n} \right)^{n-1}, \\ \mathbb{E} \left[\left(\sum_{j \in S_{i,k}} W_j \right) \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k \rrbracket \cup \{0\}) \right] &= |S_{i,k}| \left(1 - \frac{k+1}{n} \right)^{n-1}, \end{aligned}$$

and accordingly

$$\begin{aligned} & \mathbb{E}[\bar{\xi}_b \mid \mathbf{X}, \mathbf{Y}] \\ &= 1 - \frac{3}{n^2-1} \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \mathbb{E} \left[\left(\sum_{j \in S_{i,k}} W_j \right) \mathbf{1}(W_i > 0) \mathbf{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbf{1}(W_{i+k} > 0) \mid \mathbf{X}, \mathbf{Y} \right] \\ &= 1 - \frac{3}{n^2-1} \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} |S_{i,k}| \left[\left(1 - \frac{k-1}{n} \right)^{n-1} - 2 \left(1 - \frac{k}{n} \right)^{n-1} + \left(1 - \frac{k+1}{n} \right)^{n-1} \right]. \end{aligned}$$

This completes the proof. \square

4.5 Proof of Lemma 3.5

Proof of Lemma 3.5. We have

$$\begin{aligned} \mathbb{E}[\bar{\xi}_b] &= 1 - \frac{3}{n^2-1} \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \mathbb{E}[|S_{i,k}|] \left[\left(1 - \frac{k-1}{n} \right)^{n-1} - 2 \left(1 - \frac{k}{n} \right)^{n-1} + \left(1 - \frac{k+1}{n} \right)^{n-1} \right] \\ &= 1 - \frac{1}{n^2-1} \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} (n-k-1) \left[\left(1 - \frac{k-1}{n} \right)^{n-1} - 2 \left(1 - \frac{k}{n} \right)^{n-1} + \left(1 - \frac{k+1}{n} \right)^{n-1} \right]. \end{aligned}$$

Note that

$$\begin{aligned}
& \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \left[\left(1 - \frac{k-1}{n}\right)^{n-1} - 2\left(1 - \frac{k}{n}\right)^{n-1} + \left(1 - \frac{k+1}{n}\right)^{n-1} \right] \\
&= \sum_{i=1}^{n-1} \left[1 - \left(1 - \frac{1}{n}\right)^{n-1} - \left(\frac{i}{n}\right)^{n-1} + \left(\frac{i-1}{n}\right)^{n-1} \right] \\
&= (n-1) \left[1 - \left(1 - \frac{1}{n}\right)^{n-1} \right] - \left(1 - \frac{1}{n}\right)^{n-1},
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} k \left[\left(1 - \frac{k-1}{n}\right)^{n-1} - 2\left(1 - \frac{k}{n}\right)^{n-1} + \left(1 - \frac{k+1}{n}\right)^{n-1} \right] \\
&= \sum_{i=1}^{n-1} \left[1 - (n-i+1) \left(\frac{i}{n}\right)^{n-1} + (n-i) \left(\frac{i-1}{n}\right)^{n-1} \right] = n-1 - 2 \frac{\sum_{i=1}^{n-1} i^{n-1}}{n^{n-1}}.
\end{aligned}$$

Note that

$$(1 - 1/n)^n = e^{-1} - (2e)^{-1}n^{-1} + O(n^{-2})$$

and

$$\sum_{i=1}^n i^n / n^n \rightarrow e/(e-1) \quad \text{as } n \rightarrow \infty.$$

We then obtain

$$\begin{aligned}
\mathbb{E}[\bar{\xi}_b] &= 1 - \frac{1}{n+1} \left[(n-1) \left[1 - \left(1 - \frac{1}{n}\right)^{n-1} \right] - \left(1 - \frac{1}{n}\right)^{n-1} \right] + \frac{1}{n^2-1} \left[n-1 - 2 \frac{\sum_{i=1}^{n-1} i^{n-1}}{n^{n-1}} \right] \\
&= 1 - \left[1 - \frac{1}{e} - \left(3 - \frac{1}{2e}\right) \frac{1}{n} + O\left(\frac{1}{n^2}\right) \right] = \frac{1}{e} + \left(3 - \frac{1}{2e}\right) \frac{1}{n} + O\left(\frac{1}{n^2}\right).
\end{aligned}$$

The proof is thus complete. \square

4.6 Proof of Lemma 3.6

Proof of Lemma 3.6. We have

$$\begin{aligned}
& \text{Var}[\bar{\xi}_b | \mathbf{X}, \mathbf{Y}] \\
&= \frac{9}{(n^2-1)^2} \text{Var} \left[\sum_{i=1}^{n-1} \mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{j \in S_{i,k}} W_j \right] \mathbb{1}(W_{i+l} = 0, \forall l \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \mid \mathbf{X}, \mathbf{Y} \right].
\end{aligned} \tag{4.1}$$

In the same way as Lemma 3.4,

$$\begin{aligned}
& \left\{ \mathbb{E} \left[\sum_{i=1}^{n-1} \mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{j \in S_{i,k}} W_j \right] \mathbb{1}(W_{i+l} = 0, \forall l \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \mid \mathbf{X}, \mathbf{Y} \right] \right\}^2 \\
&= \left\{ \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} |S_{i,k}| \left[\left(1 - \frac{k-1}{n}\right)^{n-1} - 2\left(1 - \frac{k}{n}\right)^{n-1} + \left(1 - \frac{k+1}{n}\right)^{n-1} \right] \right\}^2
\end{aligned}$$

$$= \left(\sum_{i=1}^{n-1} \sum_{k=1}^{n-i} |S_{i,k}| c_{n,n-1,k} \right)^2. \quad (4.2)$$

It then suffices to consider

$$\begin{aligned} & \mathbb{E} \left[\left\{ \sum_{i=1}^{n-1} \mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{j \in S_{i,k}} W_j \right] \mathbb{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \right\}^2 \mid \mathbf{X}, \mathbf{Y} \right] \\ &= \sum_{i=1}^{n-1} \mathbb{E} \left[\left\{ \mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{j \in S_{i,k}} W_j \right] \mathbb{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \right\}^2 \mid \mathbf{X}, \mathbf{Y} \right] \\ &+ 2 \sum_{i,j=1, i < j}^{n-1} \mathbb{E} \left[\mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{t \in S_{i,k}} W_t \right] \mathbb{1}(W_{i+s} = 0, \forall s \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \right. \\ &\quad \left. \mathbb{1}(W_j > 0) \sum_{\ell=1}^{n-j} \left[\sum_{t \in S_{j,\ell}} W_t \right] \mathbb{1}(W_{j+s} = 0, \forall s \in \llbracket \ell-1 \rrbracket) \mathbb{1}(W_{j+\ell} > 0) \mid \mathbf{X}, \mathbf{Y} \right]. \end{aligned} \quad (4.3)$$

For the first term in (4.3),

$$\begin{aligned} & \mathbb{E} \left[\left\{ \mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{j \in S_{i,k}} W_j \right] \mathbb{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \right\}^2 \mid \mathbf{X}, \mathbf{Y} \right] \\ &= \mathbb{E} \left[\mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{j \in S_{i,k}} W_j \right]^2 \mathbb{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \mid \mathbf{X}, \mathbf{Y} \right]. \end{aligned}$$

By Lemma 3.3 and the fact that $S_{i,k}$ is disjoint with $\{i, i+1, \dots, i+k\}$,

$$\begin{aligned} & \mathbb{E} \left[\left\{ \mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{j \in S_{i,k}} W_j \right] \mathbb{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \right\}^2 \mid \mathbf{X}, \mathbf{Y} \right] \\ &= \sum_{k=1}^{n-i} |S_{i,k}| \left[\left(1 - \frac{k-1}{n}\right)^{n-1} - 2 \left(1 - \frac{k}{n}\right)^{n-1} + \left(1 - \frac{k+1}{n}\right)^{n-1} \right] \\ &\quad + \sum_{k=1}^{n-i} |S_{i,k}|^2 \left(1 - \frac{1}{n}\right) \left[\left(1 - \frac{k-1}{n}\right)^{n-2} - 2 \left(1 - \frac{k}{n}\right)^{n-2} + \left(1 - \frac{k+1}{n}\right)^{n-2} \right] \\ &= \sum_{k=1}^{n-i} \left(|S_{i,k}| c_{n,n-1,k} + |S_{i,k}|^2 c_{n,n-2,k} (1 - 1/n) \right), \end{aligned}$$

and then

$$\begin{aligned} & \sum_{i=1}^{n-1} \mathbb{E} \left[\left\{ \mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{j \in S_{i,k}} W_j \right] \mathbb{1}(W_{i+\ell} = 0, \forall \ell \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \right\}^2 \mid \mathbf{X}, \mathbf{Y} \right] \\ &= \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \left(|S_{i,k}| c_{n,n-1,k} + |S_{i,k}|^2 c_{n,n-2,k} (1 - 1/n) \right). \end{aligned} \quad (4.4)$$

For the second term in (4.3), for any $i, j \in \llbracket n-1 \rrbracket$ with $i < j$,

$$\begin{aligned}
& \mathbb{E} \left[\mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{t \in S_{i,k}} W_t \right] \mathbb{1}(W_{i+s} = 0, \forall s \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \right. \\
& \left. \mathbb{1}(W_j > 0) \sum_{\ell=1}^{n-j} \left[\sum_{t \in S_{j,\ell}} W_t \right] \mathbb{1}(W_{j+s} = 0, \forall s \in \llbracket \ell-1 \rrbracket) \mathbb{1}(W_{j+\ell} > 0) \mid \mathbf{X}, \mathbf{Y} \right] \\
&= \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-j} \mathbb{E} \left[\mathbb{1}(W_i > 0) \left[\sum_{t \in S_{i,k}} W_t \right] \mathbb{1}(W_{i+s} = 0, \forall s \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \right. \\
& \left. \mathbb{1}(W_j > 0) \left[\sum_{t \in S_{j,\ell}} W_t \right] \mathbb{1}(W_{j+s} = 0, \forall s \in \llbracket \ell-1 \rrbracket) \mathbb{1}(W_{j+\ell} > 0) \mid \mathbf{X}, \mathbf{Y} \right] \\
&:= \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-j} T_{i,j,k,\ell}.
\end{aligned}$$

(a) If $k > j - i$, $T_{i,j,k,\ell}$ is zero for any ℓ since $\mathbb{1}(W_{i+s} = 0, \forall s \in \llbracket k-1 \rrbracket) \mathbb{1}(W_j > 0) = 0$.

(b) If $k = j - i$, for any ℓ ,

$$\begin{aligned}
T_{i,j,k,\ell} &= \mathbb{E} \left[\left[\sum_{t \in S_{i,k}} W_t \right] \left[\sum_{t \in S_{j,\ell}} W_t \right] \mathbb{1}(W_i > 0) \mathbb{1}(W_{i+s} = 0, \forall s \in \llbracket k-1 \rrbracket) \right. \\
& \left. \mathbb{1}(W_j > 0) \mathbb{1}(W_{j+s} = 0, \forall s \in \llbracket \ell-1 \rrbracket) \mathbb{1}(W_{j+\ell} > 0) \mid \mathbf{X}, \mathbf{Y} \right].
\end{aligned}$$

Note that under the event $\{W_{i+s} = 0, \forall s \in \llbracket k-1 \rrbracket\}$ and $\{W_{j+s} = 0, \forall s \in \llbracket \ell-1 \rrbracket\}$, we could decompose the above term as

$$\begin{aligned}
\sum_{t \in S_{i,k}} W_t &= \sum_{t \in S_{i,k} \cap j, \ell} W_t + \sum_{t \in S_{i,k} \setminus j, \ell} W_t + W_j \mathbb{1}(j \in S_{i,k}) + W_{j+\ell} \mathbb{1}(j+\ell \in S_{i,k}), \\
\sum_{t \in S_{j,\ell}} W_t &= \sum_{t \in S_{i,k} \cap j, \ell} W_t + \sum_{t \in S_{j,\ell} \setminus i, k} W_t + W_i \mathbb{1}(i \in S_{j,\ell}) + W_{i+k} \mathbb{1}(i+k \in S_{j,\ell}),
\end{aligned}$$

by noticing that $S_{i,k \cap j, \ell}$, $S_{i,k \setminus j, \ell}$, $S_{j, \ell \setminus i, k}$ are mutually disjoint, and these three sets are disjoint with $\{i, i+1, \dots, i+s\}$ and $\{j, j+1, \dots, j+\ell\}$. The above decomposition also holds for $k \leq j - i$.

If $k = j - i$, then $W_j \mathbb{1}(j \in S_{i,k}) = W_{i+k} \mathbb{1}(i+k \in S_{j,\ell}) = 0$.

We also have for the product of indicator functions in $T_{i,j,k,\ell}$,

$$\begin{aligned}
& \mathbb{1}(W_i > 0) \mathbb{1}(W_{i+s} = 0, \forall s \in \llbracket k-1 \rrbracket) \mathbb{1}(W_j > 0) \mathbb{1}(W_{j+s} = 0, \forall s \in \llbracket \ell-1 \rrbracket) \mathbb{1}(W_{j+\ell} > 0) \\
&= \left[1 - \mathbb{1}(W_i = 0) \right] \left[1 - \mathbb{1}(W_j = 0) \right] \left[1 - \mathbb{1}(W_{j+\ell} = 0) \right] \mathbb{1}(W_{i+s} = 0, \forall s \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{j+s} = 0, \forall s \in \llbracket \ell-1 \rrbracket).
\end{aligned}$$

By Lemma 3.3,

$$\begin{aligned}
& T_{i,j,k,\ell} \\
&= |S_{i,k \cap j, \ell}| a_{n,n-1,k,\ell} + |S_{i,k \cap j, \ell}|^2 a_{n,n-2,k,\ell} (1 - 1/n) \\
& \quad + (|S_{i,k \cap j, \ell}| |S_{i,k \setminus j, \ell}| + |S_{i,k \cap j, \ell}| |S_{j, \ell \setminus i, k}| + |S_{i,k \setminus j, \ell}| |S_{j, \ell \setminus i, k}|) a_{n,n-2,k,\ell} (1 - 1/n) + O(n a_{n,n-2,k,\ell}) \\
&= |S_{i,k \cap j, \ell}| a_{n,n-1,k,\ell} + (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|) (|S_{i,k \cap j, \ell}| + |S_{j, \ell \setminus i, k}|) a_{n,n-2,k,\ell} (1 - 1/n) + O(n a_{n,n-2,k,\ell}).
\end{aligned} \tag{4.5}$$

(c) If $k < j - i$, for any ℓ , the only difference with the case where $k = j - i$ is that we now have to decompose the product of indicator functions in $T_{i,j,k,\ell}$ as

$$\begin{aligned} & \mathbb{1}(W_i > 0) \mathbb{1}(W_{i+s} = 0, \forall s \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \mathbb{1}(W_j > 0) \mathbb{1}(W_{j+s} = 0, \forall s \in \llbracket \ell-1 \rrbracket) \mathbb{1}(W_{j+\ell} > 0) \\ = & \left[1 - \mathbb{1}(W_i = 0)\right] \left[1 - \mathbb{1}(W_{i+k} = 0)\right] \left[1 - \mathbb{1}(W_j = 0)\right] \left[1 - \mathbb{1}(W_{j+\ell} = 0)\right] \\ & \mathbb{1}(W_{i+s} = 0, \forall s \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{j+s} = 0, \forall s \in \llbracket \ell-1 \rrbracket). \end{aligned}$$

By Lemma 3.3,

$$\begin{aligned} & T_{i,j,k,\ell} \\ = & |S_{i,k \cap j, \ell}| b_{n,n-1,k,\ell} + (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|)(|S_{i,k \cap j, \ell}| + |S_{j,\ell \setminus i,k}|) b_{n,n-2,k,\ell} (1 - 1/n) \\ & + [\mathbb{1}(j \in S_{i,k}) + \mathbb{1}(j + \ell \in S_{i,k})] (|S_{i,k \cap j, \ell}| + |S_{j,\ell \setminus i,k}|) a_{n,n-2,k,\ell} (1 - 1/n) \\ & + [\mathbb{1}(i \in S_{j,\ell}) + \mathbb{1}(i + k \in S_{j,\ell})] (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|) a_{n,n-2,k,\ell} (1 - 1/n) + O(a_{n,n-2,k,\ell}). \end{aligned} \quad (4.6)$$

Combining (4.5) and (4.6),

$$\begin{aligned} & \sum_{i,j=1, i < j}^{n-1} \mathbb{E} \left[\mathbb{1}(W_i > 0) \sum_{k=1}^{n-i} \left[\sum_{t \in S_{i,k}} W_t \right] \mathbb{1}(W_{i+s} = 0, \forall s \in \llbracket k-1 \rrbracket) \mathbb{1}(W_{i+k} > 0) \right. \\ & \left. \mathbb{1}(W_j > 0) \sum_{\ell=1}^{n-j} \left[\sum_{t \in S_{j,\ell}} W_t \right] \mathbb{1}(W_{j+s} = 0, \forall s \in \llbracket \ell-1 \rrbracket) \mathbb{1}(W_{j+\ell} > 0) \mid \mathbf{X}, \mathbf{Y} \right] \\ = & \sum_{i,j=1, i < j}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-j} \left(|S_{i,k \cap j, \ell}| [a_{n,n-1,k,\ell} \mathbb{1}(k = j - i) + b_{n,n-1,k,\ell} \mathbb{1}(k < j - i)] \right. \\ & + (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|)(|S_{i,k \cap j, \ell}| + |S_{j,\ell \setminus i,k}|) [a_{n,n-2,k,\ell} \mathbb{1}(k = j - i) + b_{n,n-2,k,\ell} \mathbb{1}(k < j - i)] (1 - 1/n) \\ & + [\mathbb{1}(j \in S_{i,k}) + \mathbb{1}(j + \ell \in S_{i,k})] (|S_{i,k \cap j, \ell}| + |S_{j,\ell \setminus i,k}|) a_{n,n-2,k,\ell} \mathbb{1}(k < j - i) (1 - 1/n) \\ & \left. + [\mathbb{1}(i \in S_{j,\ell}) + \mathbb{1}(i + k \in S_{j,\ell})] (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|) a_{n,n-2,k,\ell} \mathbb{1}(k < j - i) (1 - 1/n) \right) + O(n^2). \end{aligned} \quad (4.7)$$

Plugging (4.2), (4.4), (4.7) into (4.1) completes the proof. \square

4.7 Proof of Lemma 3.7

Proof of Lemma 3.7. Note that all the quantities are invariant under probability transformation. By the continuity of the distribution and the probability integral transformation, we can assume without loss of generality that the distribution is the uniform distribution on $[0, 1]$.

(a) For S_{12} , it is easy to see $P(Y_1 \wedge Y_2 < Z_i < Y_1 \vee Y_2) = 1/3$ for any $i \in \llbracket n \rrbracket$, and then $E[S_{12}] = n/3$. For $i, j \in \llbracket n \rrbracket$ and $i \neq j$, let R_i, R_j be the rank of Z_i, Z_j in $\{Y_1, Y_2, Z_i, Z_j\}$. Then

$$P(Y_1 \wedge Y_2 < Z_i, Z_j < Y_1 \vee Y_2) = P(R_i = 2, R_j = 3) + P(R_i = 3, R_j = 2) = 2 \times 2/4! = 1/6.$$

Then

$$E[S_{12}^2] = n(n-1)/6 + n/3 = n^2/6 + n/6.$$

(b) For S_{1234} , for any $i \in \llbracket n \rrbracket$, to have $(Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4)$, we have

Y_1, Y_2 on different side of Z_i , as well as Y_3, Y_4 . Then

$$P((Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4) \mid Z_i) = 4Z_i^2(1 - Z_i)^2,$$

and then

$$P((Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4)) = 2/15.$$

Then $E[S_{1234}] = 2n/15$. In the same way, for any $i, j \in \llbracket n \rrbracket$ and $i \neq j$,

$$P((Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i, Z_j < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4) \mid Z_i, Z_j) = 4(Z_i \wedge Z_j)^2(1 - Z_i \vee Z_j)^2,$$

and then

$$P((Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i, Z_j < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4)) = 2/45.$$

Then

$$E[S_{1234}^2] = 2n(n-1)/45 + 2n/15 = 2n^2/45 + 4n/45.$$

(c) For $S_{1234}S_{12\setminus 34}$, for any $i, j \in \llbracket n \rrbracket$ and $i \neq j$, to have $(Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4)$ and $Y_1 \wedge Y_2 < Z_j < Y_1 \vee Y_2, Z_j > Y_3 \vee Y_4$ or $Z_j < Y_3 \wedge Y_4$, we consider two cases $Z_i < Z_j$ and $Z_i > Z_j$ separately. If $Z_i < Z_j$, we need one of Y_1, Y_2 smaller than Z_i , and the other larger than Z_j . We also need one of Y_3, Y_4 smaller than Z_i , and the other between Z_i and Z_j . If $Z_i > Z_j$, we need one of Y_1, Y_2 smaller than Z_j , and the other larger than Z_i . We also need one of Y_3, Y_4 larger than Z_i , and the other between Z_i and Z_j . Combining the two cases, we have

$$\begin{aligned} & P((Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4), \\ & Y_1 \wedge Y_2 < Z_j < Y_1 \vee Y_2, Z_j > Y_3 \vee Y_4 \text{ or } Z_j < Y_3 \wedge Y_4 \mid Z_i, Z_j) \\ & = 4[\mathbb{1}(Z_i < Z_j)Z_i^2(Z_j - Z_i)(1 - Z_j) + \mathbb{1}(Z_j < Z_i)Z_j(Z_i - Z_j)(1 - Z_i)^2], \end{aligned}$$

and then

$$\begin{aligned} & P((Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4), \\ & Y_1 \wedge Y_2 < Z_j < Y_1 \vee Y_2, Z_j > Y_3 \vee Y_4 \text{ or } Z_j < Y_3 \wedge Y_4) = 1/45. \end{aligned}$$

Then

$$E[S_{1234}S_{12\setminus 34}] = n(n-1)/45.$$

(d) From the symmetry, $E[S_{1234}S_{34\setminus 12}] = n(n-1)/45$.

(e) For $S_{12\setminus 34}S_{34\setminus 12}$, for any $i, j \in \llbracket n \rrbracket$ and $i \neq j$, to have $Y_1 \wedge Y_2 < Z_i < Y_1 \vee Y_2, Z_i > Y_3 \vee Y_4$ or $Z_i < Y_3 \wedge Y_4$ and $Y_3 \wedge Y_4 < Z_j < Y_3 \vee Y_4, Z_j > Y_1 \vee Y_2$ or $Z_j < Y_1 \wedge Y_2$, we consider $Z_i < Z_j$ and $Z_i > Z_j$ separately. If $Z_i < Z_j$, we need one of Y_1, Y_2 smaller than Z_i , and the other between Z_i and Z_j . We also need one of Y_3, Y_4 larger than Z_j , and the other between Z_i and Z_j . The case when $Z_i > Z_j$ can be analyzed in the same way. Then

$$\begin{aligned} & P(Y_1 \wedge Y_2 < Z_i < Y_1 \vee Y_2, Z_i > Y_3 \vee Y_4 \text{ or } Z_i < Y_3 \wedge Y_4, \\ & Y_3 \wedge Y_4 < Z_j < Y_3 \vee Y_4, Z_j > Y_1 \vee Y_2 \text{ or } Z_j < Y_1 \wedge Y_2 \mid Z_i, Z_j) \\ & = 4[\mathbb{1}(Z_i < Z_j)Z_i(Z_j - Z_i)^2(1 - Z_j) + \mathbb{1}(Z_j < Z_i)Z_j(Z_i - Z_j)^2(1 - Z_i)], \end{aligned}$$

and then

$$\begin{aligned} & \mathbb{P}(Y_1 \wedge Y_2 < Z_i < Y_1 \vee Y_2, Z_i > Y_3 \vee Y_4 \text{ or } Z_i < Y_3 \wedge Y_4, \\ & Y_3 \wedge Y_4 < Z_j < Y_3 \vee Y_4, Z_j > Y_1 \vee Y_2 \text{ or } Z_j < Y_1 \wedge Y_2) = 1/45. \end{aligned}$$

Then

$$\mathbb{E}[S_{12\setminus 34}S_{34\setminus 12}] = n(n-1)/45.$$

(f) For S_{1223} , to have $(Y_1 \wedge Y_2) \vee (Y_2 \wedge Y_3) < Z_i < (Y_1 \vee Y_2) \wedge (Y_2 \vee Y_3)$, we have Y_1, Y_3 on the same side of Z_i , and different side of Y_2 . Then

$$\mathbb{P}((Y_1 \wedge Y_2) \vee (Y_2 \wedge Y_3) < Z_i < (Y_1 \vee Y_2) \wedge (Y_2 \vee Y_3) \mid Z_i) = Z_i(1 - Z_i),$$

and then

$$\mathbb{P}((Y_1 \wedge Y_2) \vee (Y_2 \wedge Y_3) < Z_i < (Y_1 \vee Y_2) \wedge (Y_2 \vee Y_3)) = 1/6.$$

Then $\mathbb{E}[S_{1223}] = n/6$. We also have

$$\mathbb{P}((Y_1 \wedge Y_2) \vee (Y_2 \wedge Y_3) < Z_i, Z_j < (Y_1 \vee Y_2) \wedge (Y_2 \vee Y_3) \mid Z_i, Z_j) = (Z_i \wedge Z_j)(1 - Z_i \vee Z_j)(Z_i \wedge Z_j + 1 - Z_i \vee Z_j),$$

and then

$$\mathbb{P}((Y_1 \wedge Y_2) \vee (Y_2 \wedge Y_3) < Z_i, Z_j < (Y_1 \vee Y_2) \wedge (Y_2 \vee Y_3)) = 1/15.$$

Then

$$\mathbb{E}[S_{1223}^2] = n(n-1)/15 + n/6 = n^2/15 + n/10.$$

(g) For $S_{1223}S_{12\setminus 23}$, by performing similar analysis,

$$\begin{aligned} & \mathbb{P}((Y_1 \wedge Y_2) \vee (Y_2 \wedge Y_3) < Z_i < (Y_1 \vee Y_2) \wedge (Y_2 \vee Y_3), \\ & Y_1 \wedge Y_2 < Z_j < Y_1 \vee Y_2, Z_j > Y_2 \vee Y_3 \text{ or } Z_j < Y_2 \wedge Y_3 \mid Z_i, Z_j) \\ & = \mathbb{1}(Z_i < Z_j)Z_i(Z_j - Z_i)(1 - Z_j) + \mathbb{1}(Z_j < Z_i)Z_j(Z_i - Z_j)(1 - Z_i), \end{aligned}$$

and then

$$\begin{aligned} & \mathbb{P}((Y_1 \wedge Y_2) \vee (Y_2 \wedge Y_3) < Z_i < (Y_1 \vee Y_2) \wedge (Y_2 \vee Y_3), \\ & Y_1 \wedge Y_2 < Z_j < Y_1 \vee Y_2, Z_j > Y_2 \vee Y_3 \text{ or } Z_j < Y_2 \wedge Y_3) = 1/60. \end{aligned}$$

Then

$$\mathbb{E}[S_{1223}S_{12\setminus 23}] = n(n-1)/60.$$

(h) We can establish in the same way that

$$\mathbb{E}[S_{1223}S_{23\setminus 12}] = \mathbb{E}[S_{12\setminus 23}S_{23\setminus 12}] = n(n-1)/60.$$

(i) For $\mathbb{1}(Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2)S_{1234}$, to have $Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2$ and $(Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4)$, we condition on Y_3 and Z_i . If $Y_3 < Z_i$, then we need one of Y_1, Y_2 smaller than Y_3 , and the other larger than Z_i . We also need Y_4 larger than Z_i . Then

$$\begin{aligned} & \mathbb{P}(Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2, (Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4) \mid Z_i, Y_3) \\ & = 2Y_3(1 - Z_i)^2 \mathbb{1}(Y_3 < Z_i) + 2Z_i^2(1 - Y_3) \mathbb{1}(Y_3 > Z_i), \end{aligned}$$

and then

$$P(Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2, (Y_1 \wedge Y_2) \vee (Y_3 \wedge Y_4) < Z_i < (Y_1 \vee Y_2) \wedge (Y_3 \vee Y_4)) = \frac{1}{15}.$$

Then

$$E[\mathbf{1}(Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2)S_{1234}] = n/15.$$

(j) For $\mathbf{1}(Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2)S_{34\setminus 12}$, to have $Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2$ and $Y_3 \wedge Y_4 < Z_i < Y_3 \vee Y_4, Z_i > Y_1 \vee Y_2$ or $Z_i < Y_1 \wedge Y_2$. Conditional on Y_3 and Z_i , if $Y_3 < Z_i$, we need one of Y_1, Y_2 is smaller than Y_3 , and the other between Y_3 and Z_1 . We also need Y_4 larger than Z_1 . Then

$$\begin{aligned} & P(Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2, Y_3 \wedge Y_4 < Z_i < Y_3 \vee Y_4, Z_i > Y_1 \vee Y_2 \text{ or } Z_i < Y_1 \wedge Y_2 \mid Z_i, Y_3) \\ & = 2Y_3(Z_i - Y_3)(1 - Z_i)\mathbf{1}(Y_3 < Z_i) + 2Z_i(Y_3 - Z_i)(1 - Y_3)\mathbf{1}(Y_3 > Z_i), \end{aligned}$$

and then

$$P(Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2, Y_3 \wedge Y_4 < Z_i < Y_3 \vee Y_4, Z_i > Y_1 \vee Y_2 \text{ or } Z_i < Y_1 \wedge Y_2) = \frac{1}{30}.$$

Then

$$E[\mathbf{1}(Y_1 \wedge Y_2 < Y_3 < Y_1 \vee Y_2)S_{34\setminus 12}] = n/30.$$

The proof is thus complete. \square

4.8 Proof of Lemma 3.8

Proof of Lemma 3.8. By Lemma 3.7, we have

$$E[|S_{i,k}|] = (n - k - 1)/3 \quad \text{and} \quad E[|S_{i,k}|^2] = (n - k - 1)^2/6 + (n - k - 1)/6.$$

One could then obtain

$$E\left[\sum_{i=1}^{n-1} \sum_{k=1}^{n-i} |S_{i,k}| c_{n,n-1,k}\right] = \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \left(\frac{1}{3}(n - k - 1)\right) c_{n,n-1,k} = O(n^2), \quad (4.8)$$

and

$$E\left[\sum_{i=1}^{n-1} \sum_{k=1}^{n-i} |S_{i,k}|^2 c_{n,n-2,k}\right] = \frac{1}{6} \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \left((n - k - 1)(n - k)\right) c_{n,n-2,k} = \frac{1}{6} \left(1 - \frac{1}{e}\right) n^3 + O(n^2). \quad (4.9)$$

When $k < j - i$, i, j, k, ℓ are distinct. Again by Lemma 3.7, in this case we have

$$E[|S_{i,k \cap j, \ell}|] = \frac{2}{15}(n - k - \ell - 2),$$

and

$$E[(|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|)(|S_{i,k \cap j, \ell}| + |S_{j, \ell \setminus i, k}|)] = \frac{1}{9}(n - k - \ell - 2)^2 + \frac{1}{45}(n - k - \ell - 2),$$

and

$$\begin{aligned} & E\{\mathbf{1}(j \in S_{i,k}) + \mathbf{1}(j + \ell \in S_{i,k})\}(|S_{i,k \cap j, \ell}| + |S_{j, \ell \setminus i, k}|) + [\mathbf{1}(i \in S_{j, \ell}) + \mathbf{1}(i + k \in S_{j, \ell})](|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|) \\ & = 4\left(\frac{1}{15} + \frac{1}{30}\right)(n - k - \ell - 2) = \frac{2}{5}(n - k - \ell - 2). \end{aligned}$$

Then

$$\begin{aligned}
& \mathbb{E} \left[\sum_{i,j=1, i < j}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-j} |S_{i,k \cap j, \ell}| b_{n,n-1,k,\ell} \mathbb{1}(k < j - i) \right] = \mathbb{E} \left[\sum_{i=1}^{n-3} \sum_{j=i+2}^{n-1} \sum_{k=1}^{j-i-1} \sum_{\ell=1}^{n-j} |S_{i,k \cap j, \ell}| b_{n,n-1,k,\ell} \right] \\
&= \frac{2}{15} \sum_{i=1}^{n-3} \sum_{j=i+2}^{n-1} \sum_{k=1}^{j-i-1} \sum_{\ell=1}^{n-j} (n - k - \ell - 2) b_{n,n-1,k,\ell} \\
&= \frac{2}{15} \sum_{k=1}^{n-4} \sum_{\ell=1}^{n-k-3} \sum_{j=k+2}^{n-\ell} \sum_{i=1}^{j-k-1} (n - k - \ell - 2) b_{n,n-1,k,\ell} \\
&= \frac{1}{15} \sum_{k=1}^{n-4} \sum_{\ell=1}^{n-k-3} (n - k - \ell)(n - k - \ell - 1)(n - k - \ell - 2) b_{n,n-1,k,\ell} \\
&= \frac{1}{15} \left(1 - \frac{2}{e} + \frac{1}{e^2} \right) n^3 + O(n^2), \tag{4.10}
\end{aligned}$$

and

$$\begin{aligned}
& \mathbb{E} \left[\sum_{i,j=1, i < j}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-j} (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|) (|S_{i,k \cap j, \ell}| + |S_{j,\ell \setminus i, k}|) b_{n,n-2,k,\ell} \mathbb{1}(k < j - i) \right] \\
&= \mathbb{E} \left[\sum_{i=1}^{n-3} \sum_{j=i+2}^{n-1} \sum_{k=1}^{j-i-1} \sum_{\ell=1}^{n-j} (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|) (|S_{i,k \cap j, \ell}| + |S_{j,\ell \setminus i, k}|) b_{n,n-2,k,\ell} \right] \\
&= \frac{1}{9} \sum_{i=1}^{n-3} \sum_{j=i+2}^{n-1} \sum_{k=1}^{j-i-1} \sum_{\ell=1}^{n-j} (n - k - \ell - 2)^2 b_{n,n-2,k,\ell} + \frac{1}{45} \sum_{i=1}^{n-3} \sum_{j=i+2}^{n-1} \sum_{k=1}^{j-i-1} \sum_{\ell=1}^{n-j} (n - k - \ell - 2) b_{n,n-2,k,\ell} \\
&= \frac{1}{18} \sum_{k=1}^{n-4} \sum_{\ell=1}^{n-k-3} (n - k - \ell)(n - k - \ell - 1)(n - k - \ell - 2)^2 b_{n,n-2,k,\ell} \\
&\quad + \frac{1}{90} \sum_{k=1}^{n-4} \sum_{\ell=1}^{n-k-3} (n - k - \ell)(n - k - \ell - 1)(n - k - \ell - 2) b_{n,n-2,k,\ell} \\
&= \frac{1}{18} \left[\left(1 - \frac{2}{e} + \frac{1}{e^2} \right) n^4 + \left(-13 + \frac{15}{e} - \frac{3}{e^2} \right) n^3 \right] + \frac{1}{90} \left(1 - \frac{2}{e} + \frac{1}{e^2} \right) n^3 + O(n^2), \tag{4.11}
\end{aligned}$$

and

$$\begin{aligned}
& \mathbb{E} \left[\sum_{i,j=1, i < j}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-j} \left([\mathbb{1}(j \in S_{i,k}) + \mathbb{1}(j + \ell \in S_{i,k})] (|S_{i,k \cap j, \ell}| + |S_{j,\ell \setminus i, k}|) a_{n,n-2,k,\ell} \mathbb{1}(k < j - i) \right. \right. \\
&\quad \left. \left. + [\mathbb{1}(i \in S_{j,\ell}) + \mathbb{1}(i + k \in S_{j,\ell})] (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|) a_{n,n-2,k,\ell} \mathbb{1}(k < j - i) \right) \right] \\
&= \frac{2}{5} \sum_{i=1}^{n-3} \sum_{j=i+2}^{n-1} \sum_{k=1}^{j-i-1} \sum_{\ell=1}^{n-j} (n - k - \ell - 2) a_{n,n-2,k,\ell} \\
&= \frac{1}{5} \sum_{k=1}^{n-4} \sum_{\ell=1}^{n-k-3} (n - k - \ell)(n - k - \ell - 1)(n - k - \ell - 2) a_{n,n-2,k,\ell} \\
&= \frac{1}{5} \left(1 - \frac{1}{e} \right) n^3 + O(n^2). \tag{4.12}
\end{aligned}$$

When $k = j - i$, we have $j = i + k$ and then by Lemma 3.7,

$$\mathbb{E}[|S_{i,k \cap j, \ell}|] = \frac{1}{6}(n - k - \ell - 1),$$

and

$$\mathbb{E}[(|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|)(|S_{i,k \cap j, \ell}| + |S_{j, \ell \setminus i, k}|)] = \frac{7}{60}(n - k - \ell - 1)^2 + \frac{1}{20}(n - k - \ell - 1).$$

Then

$$\begin{aligned} \mathbb{E}\left[\sum_{i,j=1, i < j}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-j} |S_{i,k \cap j, \ell}| a_{n,n-1,k,\ell} \mathbf{1}(k = j - i)\right] &= \mathbb{E}\left[\sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-i-k} |S_{i,k \cap i+k, \ell}| a_{n,n-1,k,\ell}\right] \\ &= \frac{1}{6} \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-i-k} (n - k - \ell - 1) a_{n,n-1,k,\ell} = \frac{1}{6} \sum_{k=1}^{n-2} \sum_{\ell=1}^{n-k-1} \sum_{i=1}^{n-k-\ell} (n - k - \ell - 1) a_{n,n-1,k,\ell} \\ &= \frac{1}{6} \sum_{k=1}^{n-2} \sum_{\ell=1}^{n-k-1} (n - k - \ell)(n - k - \ell - 1) a_{n,n-1,k,\ell} = O(n^2), \end{aligned} \quad (4.13)$$

and

$$\begin{aligned} \mathbb{E}\left[\sum_{i,j=1, i < j}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-j} (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|)(|S_{i,k \cap j, \ell}| + |S_{j, \ell \setminus i, k}|) a_{n,n-2,k,\ell} \mathbf{1}(k = j - i)\right] \\ &= \mathbb{E}\left[\sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-i-k} (|S_{i,k \cap i+k, \ell}| + |S_{i,k \setminus i+k, \ell}|)(|S_{i,k \cap i+k, \ell}| + |S_{i+k, \ell \setminus i, k}|) a_{n,n-2,k,\ell}\right] \\ &= \frac{7}{60} \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-i-k} (n - k - \ell - 1)^2 a_{n,n-2,k,\ell} + \frac{1}{20} \sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-i-k} (n - k - \ell - 1) a_{n,n-2,k,\ell} \\ &= \frac{7}{60} \sum_{k=1}^{n-2} \sum_{\ell=1}^{n-k-1} (n - k - \ell)(n - k - \ell - 1)^2 a_{n,n-2,k,\ell} \\ &\quad + \frac{1}{20} \sum_{k=1}^{n-2} \sum_{\ell=1}^{n-k-1} (n - k - \ell)(n - k - \ell - 1) a_{n,n-2,k,\ell} \\ &= \frac{7}{60} \left(1 - \frac{1}{e}\right) n^3 + O(n^2). \end{aligned} \quad (4.14)$$

Combining all the pieces from (4.8) to (4.14), we have

$$\begin{aligned} &\sum_{i=1}^{n-1} \sum_{k=1}^{n-i} \left(|S_{i,k}| c_{n,n-1,k} + |S_{i,k}|^2 c_{n,n-2,k} (1 - 1/n) \right) \\ &+ 2 \sum_{i,j=1, i < j}^{n-1} \sum_{k=1}^{n-i} \sum_{\ell=1}^{n-j} \left(|S_{i,k \cap j, \ell}| [a_{n,n-1,k,\ell} \mathbf{1}(k = j - i) + b_{n,n-1,k,\ell} \mathbf{1}(k < j - i)] \right. \\ &+ (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|)(|S_{i,k \cap j, \ell}| + |S_{j, \ell \setminus i, k}|) [a_{n,n-2,k,\ell} \mathbf{1}(k = j - i) + b_{n,n-2,k,\ell} \mathbf{1}(k < j - i)] (1 - 1/n) \\ &+ [\mathbf{1}(j \in S_{i,k}) + \mathbf{1}(j + \ell \in S_{i,k})] (|S_{i,k \cap j, \ell}| + |S_{j, \ell \setminus i, k}|) a_{n,n-2,k,\ell} \mathbf{1}(k < j - i) (1 - 1/n) \\ &\left. + [\mathbf{1}(i \in S_{j,\ell}) + \mathbf{1}(i + k \in S_{j,\ell})] (|S_{i,k \cap j, \ell}| + |S_{i,k \setminus j, \ell}|) a_{n,n-2,k,\ell} \mathbf{1}(k < j - i) (1 - 1/n) \right) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{6} \left(1 - \frac{1}{e}\right) n^3 + \frac{2}{15} \left(1 - \frac{2}{e} + \frac{1}{e^2}\right) n^3 + \frac{7}{30} \left(1 - \frac{1}{e}\right) n^3 + \frac{1}{9} \left(1 - \frac{2}{e} + \frac{1}{e^2}\right) n^4 - \frac{1}{9} \left(1 - \frac{2}{e} + \frac{1}{e^2}\right) n^3 \\
&\quad + \frac{1}{9} \left(-13 + \frac{15}{e} - \frac{3}{e^2}\right) n^3 + \frac{1}{45} \left(1 - \frac{2}{e} + \frac{1}{e^2}\right) n^3 + \frac{2}{5} \left(1 - \frac{1}{e}\right) n^3 + O(n^2) \\
&= \frac{1}{9} \left(1 - \frac{2}{e} + \frac{1}{e^2}\right) n^4 + \left(-\frac{3}{5} + \frac{7}{9e} - \frac{13}{45e^2}\right) n^3 + O(n^2). \tag{4.15}
\end{aligned}$$

By the proof of Lemma 3.5,

$$\begin{aligned}
&\mathbb{E} \left[\left(\sum_{i=1}^{n-1} \sum_{k=1}^{n-i} |S_{i,k}| c_{n,n-1,k} \right)^2 \right] \geq \left[\mathbb{E} \left(\sum_{i=1}^{n-1} \sum_{k=1}^{n-i} |S_{i,k}| c_{n,n-1,k} \right) \right]^2 \\
&= \frac{1}{9} \left[\left(1 - \frac{1}{e}\right) n^2 - \left(3 - \frac{1}{2e}\right) n + O(1) \right]^2 = \frac{1}{9} \left(1 - \frac{1}{e}\right)^2 n^4 - \frac{2}{9} \left(1 - \frac{1}{e}\right) \left(3 - \frac{1}{2e}\right) n^3 + O(n^2). \tag{4.16}
\end{aligned}$$

Plugging (4.15) and (4.16) to Lemma 3.6 then yields

$$\begin{aligned}
&\mathbb{E}[\text{Var}[\bar{\xi}_b | \mathbf{X}, \mathbf{Y}]] \leq \frac{9}{n^4} \left(1 + O(n^{-2})\right) \\
&\quad \left[\frac{1}{9} \left(1 - \frac{2}{e} + \frac{1}{e^2}\right) n^4 + \left(-\frac{3}{5} + \frac{7}{9e} - \frac{13}{45e^2}\right) n^3 - \frac{1}{9} \left(1 - \frac{1}{e}\right)^2 n^4 + \frac{2}{9} \left(1 - \frac{1}{e}\right) \left(3 - \frac{1}{2e}\right) n^3 + O(n^2) \right] + O\left(\frac{1}{n^2}\right) \\
&= \left(\frac{3}{5} - \frac{8}{5e^2}\right) \frac{1}{n} + O\left(\frac{1}{n^2}\right),
\end{aligned}$$

and the proof is thus complete. \square

4.9 Proof of Lemma 4.1

Proof of Lemma 4.1. From the definition of $[\tilde{L}_i]_{i=1}^n$, we have

$$\begin{aligned}
\sum_{i=1}^n W_i \tilde{L}_i (n - \tilde{L}_i) &= \sum_{i=1}^n W_i \left(\sum_{j=1}^n W_j \mathbb{1}(Y_j \geq Y_i) \right) \left(n - \sum_{j=1}^n W_j \mathbb{1}(Y_j \geq Y_i) \right) \\
&= \sum_{i=1}^n W_i \left(\sum_{j=1}^n W_j \mathbb{1}(Y_j \geq Y_i) \right) \left(\sum_{j=1}^n W_j \mathbb{1}(Y_j < Y_i) \right).
\end{aligned}$$

We first study the mean. Note that (W_1, \dots, W_n) is independent of \mathbf{X} and \mathbf{Y} . We can assume $[Y_i]_{i=1}^n$ is strictly increasing with probability one without loss of generality conditional on \mathbf{X} and \mathbf{Y} . Then

$$\begin{aligned}
&\mathbb{E} \left[\sum_{i=1}^n W_i \left(\sum_{j=1}^n W_j \mathbb{1}(Y_j \geq Y_i) \right) \left(\sum_{j=1}^n W_j \mathbb{1}(Y_j < Y_i) \right) \mid \mathbf{X}, \mathbf{Y} \right] \\
&= \mathbb{E} \left[\sum_{i=1}^n W_i \left(n - \sum_{j=1}^{i-1} W_j \right) \left(\sum_{j=1}^{i-1} W_j \right) \right] \\
&= n \sum_{i=1}^n \mathbb{E} \left[W_i \left(\sum_{j=1}^{i-1} W_j \right) \right] - \sum_{i=1}^n \mathbb{E} \left[W_i \left(\sum_{j=1}^{i-1} W_j \right)^2 \right].
\end{aligned}$$

Since $(W_1, \dots, W_n) \sim M_n(n; 1/n, \dots, 1/n)$, for any $i \in \llbracket n \rrbracket$, we have

$$\sum_{j=1}^{i-1} W_j \mid W_i \sim \text{Bin}\left(n - W_i, \frac{i-1}{n-1}\right) \quad \text{and} \quad W_i \sim \text{Bin}\left(n, \frac{1}{n}\right).$$

Then for any $i \in \llbracket n \rrbracket$,

$$\mathbb{E} \left[W_i \left(\sum_{j=1}^{i-1} W_j \right) \right] = \mathbb{E} \left[\frac{i-1}{n-1} W_i (n - W_i) \right] = (i-1)(1 - 1/n),$$

and

$$\begin{aligned} \mathbb{E} \left[W_i \left(\sum_{j=1}^{i-1} W_j \right)^2 \right] &= \mathbb{E} \left[\frac{(i-1)(n-i)}{(n-1)^2} W_i (n - W_i) \right] + \mathbb{E} \left[\frac{(i-1)^2}{(n-1)^2} W_i (n - W_i)^2 \right] \\ &= \frac{(i-1)(n-i)}{n} + \frac{(i-1)^2(n^2 - 2n + 2)}{n^2} = (i-1)(1 - 1/n) + (i-1)^2(1 - 1/n)(1 - 2/n). \end{aligned}$$

It then holds that

$$\begin{aligned} &\mathbb{E} \left[\sum_{i=1}^n W_i \left(\sum_{j=1}^n W_j \mathbf{1}(Y_j \geq Y_i) \right) \left(\sum_{j=1}^n W_j \mathbf{1}(Y_j < Y_i) \right) \mid \mathbf{X}, \mathbf{Y} \right] \\ &= \frac{(n-1)^2(n^2 + 2n - 2)}{6n} = \frac{1}{6}(n^3 - 5n + 6 - 2n^{-1}). \end{aligned} \quad (4.17)$$

We next consider the variance. Note that

$$\begin{aligned} &\text{Var} \left[\sum_{i=1}^n W_i \left(\sum_{j=1}^n W_j \mathbf{1}(Y_j \geq Y_i) \right) \left(\sum_{j=1}^n W_j \mathbf{1}(Y_j < Y_i) \right) \mid \mathbf{X}, \mathbf{Y} \right] \\ &= \text{Var} \left[\sum_{i=1}^n W_i \left(n - \sum_{j=1}^{i-1} W_j \right) \left(\sum_{j=1}^{i-1} W_j \right) \right] \\ &= \text{Var} \left[\sum_{i=1}^n W_i \left(\sum_{j=i}^n W_j \right) \left(\sum_{j=1}^{i-1} W_j \right) \right]. \end{aligned}$$

We can consider Z_1, Z_2, \dots, Z_n i.i.d. with distribution $\mathbb{P}(Z_1 = k) = 1/n$ for any $k \in \llbracket n \rrbracket$. Then (W_1, \dots, W_n) has the same law with $(\sum_{k=1}^n \mathbf{1}(Z_k = 1), \dots, \sum_{k=1}^n \mathbf{1}(Z_k = n))$.

To apply the Efron-Stein inequality, consider \tilde{Z}_1 with the same distribution as Z_1 and is independent of Z_1, \dots, Z_n . Let $(\tilde{W}_1, \dots, \tilde{W}_n)$ be the new values by replacing Z_1 with \tilde{Z}_1 . Assume $Z_1 = k$ and $\tilde{Z}_1 = \ell$ for some $k, \ell \in \llbracket n \rrbracket$. If $k = \ell$, then $W_i = \tilde{W}_i$ for all $i \in \llbracket n \rrbracket$, and then the difference is zero. Then we assume $k < \ell$ without loss of generality. In this case, $\tilde{W}_k = W_k - 1$, $\tilde{W}_\ell = W_\ell + 1$, and $\tilde{W}_i = W_i$ for other $i \in \llbracket n \rrbracket$. Thus we have the following results for different $i \in \llbracket n \rrbracket$,

- $i = k$,

$$\begin{aligned} &\tilde{W}_i \left(\sum_{j=i}^n \tilde{W}_j \right) \left(\sum_{j=1}^{i-1} \tilde{W}_j \right) - W_i \left(\sum_{j=i}^n W_j \right) \left(\sum_{j=1}^{i-1} W_j \right) \\ &= (W_k - 1) \left(\sum_{j=k}^n W_j \right) \left(\sum_{j=1}^{k-1} W_j \right) - W_k \left(\sum_{j=k}^n W_j \right) \left(\sum_{j=1}^{k-1} W_j \right) \\ &= - \left(\sum_{j=k}^n W_j \right) \left(\sum_{j=1}^{k-1} W_j \right). \end{aligned}$$

- $i = \ell$,

$$\begin{aligned}
& \widetilde{W}_i \left(\sum_{j=i}^n \widetilde{W}_j \right) \left(\sum_{j=1}^{i-1} \widetilde{W}_j \right) - W_i \left(\sum_{j=i}^n W_j \right) \left(\sum_{j=1}^{i-1} W_j \right) \\
&= (W_\ell + 1) \left(\sum_{j=\ell}^n W_j + 1 \right) \left(\sum_{j=1}^{\ell-1} W_j - 1 \right) - W_\ell \left(\sum_{j=\ell}^n W_j \right) \left(\sum_{j=1}^{\ell-1} W_j \right) \\
&= \left(\sum_{j=\ell}^n W_j + 1 \right) \left(\sum_{j=1}^{\ell-1} W_j - 1 \right) + W_\ell \left(\sum_{j=1}^{\ell-1} W_j - \sum_{j=\ell}^n W_j - 1 \right).
\end{aligned}$$

- $k < i < \ell$,

$$\begin{aligned}
& \widetilde{W}_i \left(\sum_{j=i}^n \widetilde{W}_j \right) \left(\sum_{j=1}^{i-1} \widetilde{W}_j \right) - W_i \left(\sum_{j=i}^n W_j \right) \left(\sum_{j=1}^{i-1} W_j \right) \\
&= W_i \left(\sum_{j=i}^n W_j + 1 \right) \left(\sum_{j=1}^{i-1} W_j - 1 \right) - W_i \left(\sum_{j=i}^n W_j \right) \left(\sum_{j=1}^{i-1} W_j \right) \\
&= W_i \left(\sum_{j=1}^{i-1} W_j - \sum_{j=i}^n W_j - 1 \right).
\end{aligned}$$

For other $i \in \llbracket n \rrbracket$, the difference is zero.

We thus have

$$\begin{aligned}
& \sum_{i=1}^n \widetilde{W}_i \left(\sum_{j=i}^n \widetilde{W}_j \right) \left(\sum_{j=1}^{i-1} \widetilde{W}_j \right) - \sum_{i=1}^n W_i \left(\sum_{j=i}^n W_j \right) \left(\sum_{j=1}^{i-1} W_j \right) \\
&= - \left(\sum_{j=k}^n W_j \right) \left(\sum_{j=1}^{k-1} W_j \right) + \left(\sum_{j=\ell}^n W_j + 1 \right) \left(\sum_{j=1}^{\ell-1} W_j - 1 \right) + \sum_{i=k+1}^{\ell} W_i \left(\sum_{j=1}^{i-1} W_j - \sum_{j=i}^n W_j - 1 \right) \\
&= \left[- \left(\sum_{j=k}^{\ell-1} W_j \right) \left(\sum_{j=1}^{k-1} W_j \right) - \left(\sum_{j=\ell}^n W_j \right) \left(\sum_{j=1}^{k-1} W_j \right) \right] + \left[\left(\sum_{j=\ell}^n W_j \right) \left(\sum_{j=1}^{k-1} W_j \right) \right. \\
&\quad \left. + \left(\sum_{j=\ell}^n W_j \right) \left(\sum_{j=k}^{\ell-1} W_j \right) - \sum_{j=\ell}^n W_j + \sum_{j=1}^{k-1} W_j + \sum_{j=k}^{\ell-1} W_j - 1 \right] + \left[\sum_{i=k+1}^{\ell} W_i \left(\sum_{j=1}^{k-1} W_j \right) \right. \\
&\quad \left. + \sum_{i=k+1}^{\ell} W_i \left(\sum_{j=k}^{i-1} W_j \right) - \sum_{i=k+1}^{\ell} W_i \left(\sum_{j=i}^{\ell-1} W_j \right) - \sum_{i=k+1}^{\ell} W_i \left(\sum_{j=\ell}^n W_j \right) - \sum_{i=k+1}^{\ell} W_i \right].
\end{aligned}$$

Note that

$$\sum_{i=k+1}^{\ell} W_i \left(\sum_{j=k}^{i-1} W_j \right) = \sum_{i=k}^{\ell-1} W_i \left(\sum_{j=i+1}^{\ell} W_j \right), \quad \sum_{i=1}^n W_i = n.$$

By some algebra,

$$\begin{aligned} & \sum_{i=1}^n \widetilde{W}_i \left(\sum_{j=i}^n \widetilde{W}_j \right) \left(\sum_{j=1}^{i-1} \widetilde{W}_j \right) - \sum_{i=1}^n W_i \left(\sum_{j=i}^n W_j \right) \left(\sum_{j=1}^{i-1} W_j \right) \\ &= (W_\ell - W_k + 1) \left(\sum_{j=1}^{k-1} W_j - \sum_{j=\ell}^n W_j - 1 \right) + (W_k + W_\ell) \left(\sum_{j=k}^{\ell-1} W_j \right) - \sum_{j=k}^{\ell-1} W_j^2. \end{aligned}$$

Using the fact that $\sum_{i=1}^n W_i = n$ and all W_i 's are nonnegative, we have

$$\begin{aligned} & \left| \sum_{i=1}^n \widetilde{W}_i \left(\sum_{j=i}^n \widetilde{W}_j \right) \left(\sum_{j=1}^{i-1} \widetilde{W}_j \right) - \sum_{i=1}^n W_i \left(\sum_{j=i}^n W_j \right) \left(\sum_{j=1}^{i-1} W_j \right) \right| \\ & \leq (W_\ell + W_k + 1)(n + 1) + (W_\ell + W_k)n + \sum_{i=1}^n W_i^2 \end{aligned}$$

Note that $Z_1, \dots, Z_n, \widetilde{Z}_1$ are i.i.d, and $W_k \sim \text{Bin}(n, \frac{1}{n})$ for any $k \in \llbracket n \rrbracket$. Then from the Efron-Stein inequality,

$$\text{Var} \left[\sum_{i=1}^n W_i \left(\sum_{j=1}^n W_j \mathbf{1}(Y_j \geq Y_i) \right) \left(\sum_{j=1}^n W_j \mathbf{1}(Y_j < Y_i) \right) \mid \mathbf{X}, \mathbf{Y} \right] \lesssim n^3. \quad (4.18)$$

From the Chebyshev inequality, and the bias-variance decomposition using (4.17) and (4.18), for any $\epsilon > 0$, for almost all \mathbf{X}, \mathbf{Y} ,

$$\begin{aligned} & \mathbb{P} \left(\left| \sum_{i=1}^n W_i \widetilde{L}_i (n - \widetilde{L}_i) - \frac{n^3}{6} \right| \geq \epsilon \mid \mathbf{X}, \mathbf{Y} \right) \\ & \leq \epsilon^{-2} \left[\text{Var} \left[\sum_{i=1}^n W_i \widetilde{L}_i (n - \widetilde{L}_i) \mid \mathbf{X}, \mathbf{Y} \right] + \mathbb{E} \left[\left(\sum_{i=1}^n W_i \widetilde{L}_i (n - \widetilde{L}_i) - \frac{n^3}{6} \right)^2 \mid \mathbf{X}, \mathbf{Y} \right] \right] \\ & \lesssim \epsilon^{-2} (n^3 + n^2) \lesssim \epsilon^{-2} n^3, \end{aligned}$$

and then the proof is complete. \square

Acknowledgement

The authors would like to thank Sourav Chatterjee for confirming the failure of the bootstrap for his rank correlation and for generously sharing his insights with the authors. The authors would also like to thank Peter Bickel for pointing out the works of Beran and for discussing the relation between bootstrap inconsistency and adaptive estimators, to thank Mathias Drton for sharing his work (Drton and Williams, 2011) that led the authors to Beran (1997) and Samworth (2003), and to thank Andres Santos for explaining his findings in Fang and Santos (2019). The authors also benefited from discussions with Lihua Lei, Bodhisattva Sen, and Jon Wellner.

This work was partly motivated by discussions with Mona Azadkia, David Childers, Peng Ding, Andreas Hagemann, and Mauricio Olivares. In particular, Peng posed a question to the second author regarding the practical relevance of the results presented in Lin and Han (2022) and noted that bootstrap is commonly used in practice. This paper serves as a response to Peng's inquiry.

References

- Abadie, A. and Imbens, G. W. (2008). On the failure of the bootstrap for matching estimators. *Econometrica*, 76(6):1537–1557.
- Abrevaya, J. and Huang, J. (2005). On the bootstrap of the maximum score estimator. *Econometrica*, 73(4):1175–1204.
- Andrews, D. W. (2000). Inconsistency of the bootstrap when a parameter is on the boundary of the parameter space. *Econometrica*, 68(2):399–405.
- Ansari, J. and Fuchs, S. (2022). A simple extension of Azadkia & Chatterjee’s rank correlation to a vector of endogenous variables. Available at [arXiv:2212.01621](https://arxiv.org/abs/2212.01621).
- Arcones, M. A. and Gine, E. (1992). On the bootstrap of U and V statistics. *The Annals of Statistics*, 20(2):655–674.
- Athreya, K. (1987). Bootstrap of the mean in the infinite variance case. *The Annals of Statistics*, 15(2):724–731.
- Auddy, A., Deb, N., and Nandy, S. (2021). Exact detection thresholds for Chatterjee’s correlation. Available at [arXiv:2104.15140v1](https://arxiv.org/abs/2104.15140v1).
- Azadkia, M. and Chatterjee, S. (2021). A simple measure of conditional dependence. *The Annals of Statistics*, 49(6):3070–3102.
- Azadkia, M., Taeb, A., and Bühlmann, P. (2021). A fast non-parametric approach for causal structure learning in polytrees. Available at [arXiv:2111.14969](https://arxiv.org/abs/2111.14969).
- Beran, R. (1982). Estimated sampling distributions: the bootstrap and competitors. *The Annals of Statistics*, 10(1):212–225.
- Beran, R. (1997). Diagnosing bootstrap success. *Annals of the Institute of Statistical Mathematics*, 49:1–24.
- Beran, R. and Srivastava, M. S. (1985). Bootstrap tests and confidence regions for functions of a covariance matrix. *The Annals of Statistics*, 13(1):95–115.
- Bergsma, W. and Dassios, A. (2014). A consistent test of independence based on a sign covariance related to Kendall’s tau. *Bernoulli*, 20(2):1006–1028.
- Bickel, P. J. (2022). Measures of independence and functional dependence. Available at [arXiv:2206.13663](https://arxiv.org/abs/2206.13663).
- Bickel, P. J. and Freedman, D. A. (1981). Some asymptotic theory for the bootstrap. *The Annals of Statistics*, 9(6):1196–1217.

- Blum, J. R., Kiefer, J., and Rosenblatt, M. (1961). Distribution free tests of independence based on the sample distribution function. *The Annals of Mathematical Statistics*, 32(2):485–498.
- Bretagnolle, J. (1983). Lois limites du bootstrap de certaines fonctionnelles. *Annales de l'I.H.P. Probabilités et Statistiques*, 19(3):281–296.
- Cao, S. and Bickel, P. J. (2020). Correlations with tailored extremal properties. Available at [arXiv:2008.10177v2](https://arxiv.org/abs/2008.10177v2).
- Chatterjee, S. (2021). A new coefficient of correlation. *Journal of the American Statistical Association*, 116(535):2009–2022.
- Chatterjee, S. (2022). A survey of some recent developments in measures of association. Available at [arXiv:2211.04702](https://arxiv.org/abs/2211.04702).
- Chatterjee, S. and Vidyasagar, M. (2022). Estimating large causal polytree skeletons from small samples. Available at [arXiv:2209.07028](https://arxiv.org/abs/2209.07028).
- Chernoff, H. and Savage, I. R. (1958). Asymptotic normality and efficiency of certain nonparametric test statistics. *The Annals of Mathematical Statistics*, 29(4):972–994.
- Deb, N., Ghosal, P., and Sen, B. (2020). Measuring association on topological spaces using kernels and geometric graphs. Available at [arXiv:2010.01768v2](https://arxiv.org/abs/2010.01768v2).
- Dette, H., Siburg, K. F., and Stoimenov, P. A. (2013). A copula-based non-parametric measure of regression dependence. *Scandinavian Journal of Statistics*, 40(1):21–41.
- Drton, M. and Williams, B. (2011). Quantifying the failure of bootstrap likelihood ratio tests. *Biometrika*, 98(4):919–934.
- Dümbgen, L. (1993). On nondifferentiable functions and the bootstrap. *Probability Theory and Related Fields*, 95:125–140.
- Efron, B. (1979). Bootstrap methods: Another look at the jackknife. *The Annals of Statistics*, 7(1):1–26.
- Efron, B. (1981). Nonparametric standard errors and confidence intervals. *Canadian Journal of Statistics*, 9(2):139–158.
- Fang, Z. and Santos, A. (2019). Inference on directionally differentiable functions. *The Review of Economic Studies*, 86(1):377–412.
- Fuchs, S. (2023). Quantifying directed dependence via dimension reduction. *Journal of Multivariate Analysis (in press)*.
- Gamboa, F., Gremaud, P., Klein, T., and Lagnoux, A. (2022). Global sensitivity analysis: a new generation of mighty estimators based on rank statistics. *Bernoulli*, 28(4):2345–2374.

- Griessenberger, F., Junker, R. R., and Trutschnig, W. (2022). On a multivariate copula-based dependence measure and its estimation. *Electronic Journal of Statistics*, 16(1):2206–2251.
- Hájek, J., Šidák, Z., and Sen, P. K. (1999). *Theory of Rank Tests* (2nd ed.). Probability and Mathematical Statistics. Academic Press, Inc., San Diego, CA.
- Hall, P. (1990). Asymptotic properties of the bootstrap for heavy-tailed distributions. *The Annals of Probability*, 18(3):1342–1360.
- Hall, P., Härdle, W., and Simar, L. (1993). On the inconsistency of bootstrap distribution estimators. *Computational Statistics and Data Analysis*, 16(1):11–18.
- Han, F. (2021). On extensions of rank correlation coefficients to multivariate spaces. *Bernoulli News*, 28:7–11.
- Han, F. and Huang, Z. (2022). Azadkia-Chatterjee’s correlation coefficient adapts to manifold data. Available at [arXiv:2209.11156](https://arxiv.org/abs/2209.11156).
- Hodges, Jr., J. L. and Lehmann, E. L. (1956). The efficiency of some nonparametric competitors of the t -test. *The Annals of Mathematical Statistics*, 27(2):324–335.
- Hoeffding, W. (1940). Maszstabinvariante Korrelationstheorie. *Schr. Math. Inst. u. Inst. Angew. Math. Univ. Berlin*, 5:181–233.
- Hoeffding, W. (1948). A non-parametric test of independence. *The Annals of Mathematical Statistics*, 19(4):546–557.
- Hoeffding, W. (1994). *Scale-Invariant Correlation Theory*, pages 57–107. Springer New York.
- Holma, A. (2022). Correlation coefficient based feature screening: With applications to microarray data. Master’s thesis, Umeå University.
- Huang, Z., Deb, N., and Sen, B. (2020). Kernel partial correlation coefficient – a measure of conditional dependence. Available at [arXiv:2012.14804v1](https://arxiv.org/abs/2012.14804v1).
- Kendall, M. G. (1938). A new measure of rank correlation. *Biometrika*, 30(1/2):81–93.
- Kendall, M. G. (1970). *Rank Correlation Methods* (4th ed.). Griffin.
- Knight, K. (1989). On the bootstrap of the sample mean in the infinite variance case. *The Annals of Statistics*, 15(2):1168–1175.
- Kosorok, M. R. (2008). Bootstrapping the Grenander estimator. *Beyond parametrics in interdisciplinary research: Festschrift in honor of Professor Pranab K. Sen*, 1:282–292.
- Lin, Z. and Han, F. (2022). Limit theorems of Chatterjee’s rank correlation. Available at [arXiv:2204.08031](https://arxiv.org/abs/2204.08031).

- Lin, Z. and Han, F. (2023). On boosting the power of Chatterjee’s rank correlation. *Biometrika (in press)*.
- Mammen, E. (1991). *When does Bootstrap Work?: Asymptotic Results and Simulations*. Springer.
- Rényi, A. (1959). On measures of dependence. *Acta Math. Acad. Sci. Hungar.*, 10(3–4):441–451.
- Samworth, R. (2003). A note on methods of restoring consistency to the bootstrap. *Biometrika*, 90(4):985–990.
- Sen, B., Banerjee, M., and Woodroffe, M. (2010). Inconsistency of bootstrap: The Grenander estimator. *The Annals of Statistics*, 38(4):1953–1977.
- Shao, J. and Tu, D. (1995). *The Jackknife and Bootstrap*. Springer.
- Shi, H., Drton, M., and Han, F. (2022). On the power of Chatterjee’s rank correlation. *Biometrika*, 109(2):317–333.
- Shi, H., Drton, M., and Han, F. (2023). On Azadkia-Chatterjee’s conditional dependence coefficient. *Bernoulli (in press)*.
- Spearman, C. (1904). The proof and measurement of association between two things. *The American Journal of Psychology*, 15(1):72–101.
- Spearman, C. (1906). ‘Footrule’ for measuring correlation. *British Journal of Psychology*, 2(1):89–108.
- Strothmann, C., Dette, H., and Siburg, K. F. (2022). Rearranged dependence measures. Available at [arXiv:2201.03329](https://arxiv.org/abs/2201.03329).
- Yanagimoto, T. (1970). On measures of association and a related problem. *Annals of the Institute of Statistical Mathematics*, 22(1):57–63.
- Zhang, Q. (2023a). On relationships between Chatterjee’s and Spearman’s correlation coefficients. Available at [arXiv:2302.10131](https://arxiv.org/abs/2302.10131).
- Zhang, Q. (2023b). On the asymptotic null distribution of the symmetrized Chatterjee’s correlation coefficient. *Statistics and Probability Letters (in press)*.