

Provably well-founded strict partial orders

Toshiyasu Arai
 Graduate School of Mathematical Sciences
 University of Tokyo
 3-8-1 Komaba, Meguro-ku, Tokyo 153-8914, JAPAN
 tosarai@ms.u-tokyo.ac.jp

Abstract

In this note we show through infinitary derivations that each provably well-founded strict partial order in ACA_0 admits an embedding to an ordinal $< \varepsilon_0$.

1 Provably well-founded relations

A *strict partial order* \prec is an irreflexive $\forall n(n \not\prec n)$ and transitive $\forall n, m, k(n \prec m \prec k \rightarrow n \prec k)$, relation on ω . Let ‘ \prec is a strict partial order’ denotes the formula $\forall n(n \not\prec n) \wedge \forall n, m, k(n \prec m \prec k \rightarrow n \prec k)$. $<_{\varepsilon_0}$ denotes a standard ε_0 -order, while $<_{\omega}$ the usual order on ω .

Theorem 1.1 *Assume $\text{ACA}_0 \vdash \text{TI}(\prec)$ for a primitive recursive relation \prec . Then there exist an ordinal $\alpha_1 < \varepsilon_0$ and a primitive recursive function f such that $\text{I}\Sigma_1$ proves*

$$\prec \text{ is a strict partial order } \rightarrow \forall n, m (n \prec m \rightarrow f(n) <_{\varepsilon_0} f(m) <_{\varepsilon_0} \alpha_1).$$

Theorem 1.1 is shown in [1] by modifying Takeuti’s proof in [4,5] in terms of Gentzen’s finitary proof [3]. In this note we show Theorem 1.1 through infinitary derivations.

Corollary 1.2 *Assume $\text{ACA}_0 \vdash \text{TI}(\prec)$ for a primitive recursive relation \prec . Then there exists an extension \prec' of \prec such that \prec' is primitive recursive, a well order, and $\text{ACA}_0 \vdash \text{TI}(\prec')$.*

Proof. Let $n \prec' m \Leftrightarrow f(n) <_{\varepsilon_0} f(m) \vee (f(n) = f(m) \wedge n <_{\omega} m)$. □

2 Proof

Assume for a primitive recursive relation \prec , $\text{ACA}_0 \vdash \text{TI}(\prec)$. In what follows argue in $\text{I}\Sigma_1$, and assume that \prec is a strict partial order. There exists an ordinal $\alpha_0 < \varepsilon_0$ such that, cf. [2]

$$\forall n [\vdash_0^{\alpha_0} E(n)] \tag{1}$$

where $\vdash_c^\alpha \Gamma$ designates that ‘there exists a (primitive recursive) infinitary derivation of Γ with ω -rule and the following inferences (*prg*) and (*Rep*)

$$\frac{\{\vdash_c^\beta \Gamma, E(m)\}_{m < n}}{\vdash_c^\alpha \Gamma} \text{ (prg)}$$

where $\beta <_{\varepsilon_0} \alpha$, E is a fresh predicate symbol and $(E(n)) \in \Gamma$. The subscript 0 in $\vdash_0^{\alpha_0} \Gamma$ indicates that a witnessed derivation is cut-free.

$$\frac{\vdash_c^\beta \Gamma}{\vdash_c^\alpha \Gamma} \text{ (Rep)}$$

where $\beta <_{\varepsilon_0} \alpha$.

Formally we understand by (1) the following fact. There exist a primitive recursive tree $T \subset <^\omega \omega$ and a primitive recursive function H such that to each node $\sigma \in T$, a five data $H(\sigma) = (seq(\sigma), ord(\sigma), rul(\sigma), crk(\sigma), num(\sigma))$ are assigned by H . Let $\Gamma = seq(\sigma)$, $\alpha = ord(\sigma)$, $c = crk(\sigma)$ and $n = num(\sigma)$. Then $H(\sigma)$ indicates that a sequent Γ is derived by a derivation in depth at most α with cut rank c . $J = rul(\sigma)$ is the last inference.

$$\frac{\{\sigma_i \vdash_c^{\beta_i} \Gamma_i\}_{i \in I}}{\sigma \vdash_c^\alpha \Gamma} \text{ (J)}$$

has to be locally correct with respect to inferences $(\vee), (\wedge), (\exists), (\forall), (cut), (prg)$ and (*Rep*), and $\beta_i <_{\varepsilon_0} \alpha$ for each i . Moreover when $J = rul(\sigma) = (prg)$, $(E(n)) \in \Gamma = seq(\sigma)$ with $n = num(\sigma)$ is the main formula of the (*prg*). Then¹ $H(\langle n \rangle) = (\{E(n)\}, \alpha_0, rul(\langle n \rangle), 0)$ for each n . Although T is not assumed to be well-founded, $rul(\sigma)$ is either (*prg*) or (*Rep*) for each $\sigma \in T$. Therefore $seq(\sigma) \subset \{E(n) : n \in \omega\}$. Let us assume that

$$\frac{\{\sigma * \langle m \rangle \vdash_c^\beta \Gamma, E(m)\}_{m < n}}{\sigma \vdash_c^\alpha \Gamma} \text{ (prg)} \quad \frac{\sigma * \langle 0 \rangle \vdash_c^\beta \Gamma}{\sigma \vdash_c^\alpha \Gamma} \text{ (Rep)}$$

First we define nodes $\sigma_m \in T$ by induction on m as follows. Let $\beta_m = ord(\sigma_m)$ and $\Gamma_m = seq(\sigma_m)$ and $J_m = rul(\sigma_m)$. Namely $\sigma_m \vdash_0^{\beta_m} \Gamma_m$. It enjoys

$$\forall n ((E(n)) \in \Gamma_m \Rightarrow m \preceq n) \tag{2}$$

Case 1. $\neg \exists n <_\omega m (m < n)$: Then let $\sigma_m = \langle m \rangle$. This means that $\beta_m = \alpha_0$ and $\Gamma_m = \{E(m)\}$.

Case 2. $\exists n <_\omega m (m < n)$: Let $n_0 <_\omega m$ be the $<_\omega$ -least number such that $m < n_0$ and $\beta_{n_0} = \min_{<_{\varepsilon_0}} \{\beta_n : n <_\omega m, m < n\}$. Consider the last inference $J_{n_0} = rul(\sigma_{n_0})$ in the derivation of $\sigma_{n_0} \vdash_0^{\beta_{n_0}} \Gamma_{n_0}$.

Case 2.1. The last inference J_{n_0} is a (*prg*):

$$\frac{\{\sigma_{n_0} * \langle n \rangle \vdash_0^\beta \Gamma_{n_0}, E(n)\}_{n < n_1}}{\sigma_{n_0} \vdash_0^{\beta_{n_0}} \Gamma_{n_0}} \text{ (prg)}$$

¹ $H(\langle \rangle)$ is arbitrary for the root $\langle \rangle$ of the tree.

where $\beta <_{\varepsilon_0} \beta_{n_0}$ and $(E(n_1)) \in \Gamma_{n_0}$ with $n_1 = \text{num}(\sigma_{n_0})$. We have $m \prec n_0 \preceq n_1$ by (2). Then let $\sigma_m = \sigma_{n_0} * \langle m \rangle$. Let $\beta_m = \beta$ and $\Gamma_m = \Gamma_{n_0} \cup \{E(m)\}$. If $(E(n)) \in \Gamma_{n_0}$, then $m \prec n_0 \preceq n$ by (2). Hence (2) is enjoyed for σ_m since \prec is assumed to be transitive.

Case 2.2. The last inference J_{n_0} is a (*Red*):

$$\frac{\sigma_{n_0} * \langle 0 \rangle \vdash_0^\beta \Gamma_{n_0}}{\sigma_{n_0} \vdash_0^{\beta_{n_0}} \Gamma_{n_0}} \text{ (Rep)}$$

where $\beta < \beta_{n_0}$. Then let $\sigma_m = \sigma_{n_0} * \langle 0 \rangle$. This means $\beta_m = \beta$ and $\Gamma_m = \Gamma_{n_0}$. Again (2) is enjoyed for σ_m by the transitivity of \prec .

Lemma 2.1 $\forall m \forall n <_\omega m [m \prec n \Rightarrow \beta_m <_{\varepsilon_0} \beta_n]$.

Proof. In **Case 2**, if $n <_\omega m$ and $m \prec n$, then $\beta_m <_{\varepsilon_0} \beta_{n_0} \leq_{\varepsilon_0} \beta_n$. \square

Now let us define $\alpha_1 = \omega^{\alpha_0}$ and f as follows.

$$f(n) = \max_{<_{\varepsilon_0}} \{ \omega^{\beta_{n_0}} \# \dots \# \omega^{\beta_{n_{\ell-1}}} \# \omega^{\beta_{n_\ell}} : \forall i < \ell (n_i \prec n_{i+1} \ \& \ n_i <_\omega n_\ell = n) \}$$

where $\#$ denotes the natural sum. Note that $n_i \neq n_j$ for $i < j \leq \ell$ since \prec is assumed to be a strict partial order. The following Lemma 2.2 shows Theorem 1.1.

Lemma 2.2 $\forall n, m [n \prec m \Rightarrow f(n) <_{\varepsilon_0} f(m) < \omega^{\alpha_0+1} = \alpha_1]$.

Proof. Let $n_0, \dots, n_{\ell-1} <_\omega n_\ell = n \prec m$ be such that $n_0 \prec \dots \prec n_{\ell-1} \prec n_\ell$ and

$$f(n) = \omega^{\beta_{n_0}} \# \dots \# \omega^{\beta_{n_{\ell-1}}} \# \omega^{\beta_{n_\ell}}.$$

Then $n_i \prec m$ and $n_i \neq m$. Let $A = \{i \leq \ell : m <_\omega n_i\}$ and $B = \{i \leq \ell : n_i <_\omega m\}$. Then $A \cup B = \{0, \dots, \ell\}$ and $A \cap B = \emptyset$. By Lemma 2.1 we obtain $\forall i \in A (\beta_{n_i} <_{\varepsilon_0} \beta_m)$, and hence

$$\sum \{ \omega^{\beta_{n_i}} : i \in A \} <_{\varepsilon_0} \omega^{\beta_m} \tag{3}$$

where $\sum \{ \alpha_0, \dots, \alpha_n \} = \alpha_0 \# \dots \# \alpha_n$. On the other side let

$$\gamma := \max_{<_{\varepsilon_0}} \{ \omega^{\beta_{m_0}} \# \dots \# \omega^{\beta_{m_{k-1}}} : \forall i < k (m_i \prec m_{i+1} \ \& \ m_i <_\omega m_k = m) \}$$

and $B = \{n_{i_0} \prec \dots \prec n_{i_{\ell-1}}\}$. Then $n_{i_0} \prec \dots \prec n_{i_{\ell-1}} \prec m$ and $n_j <_\omega m$ for each $n_j \in B$ since \prec is assumed to be transitive. Therefore

$$\sum \{ \omega^{\beta_{n_i}} : i \in B \} \leq_{\varepsilon_0} \gamma \tag{4}$$

By (4) and (3) we conclude

$$f(n) = \sum \{ \omega^{\beta_{n_i}} : i \in B \} \# \sum \{ \omega^{\beta_{n_i}} : i \in A \} <_{\varepsilon_0} \gamma \# \omega^{\beta_m} = f(m).$$

□

When \prec is elementary recursive, then so is f . For almost all theories T , Theorem 1.1 holds if the ordinal ε_0 is replaced by the proof-theoretic ordinal of T provided that a reasonable ordinal analysis of T is given.

References

- [1] T. Arai, Some results on cut-elimination, provable well-orderings, induction and reflection. *Ann. Pure Appl. Logic* 95 (1998) 93-184.
- [2] T. Arai, *Ordinal analysis with an introduction to proof theory*, Springer, 2020.
- [3] G. Gentzen, Beweibarkeit und Unbeweisbarkeit von Anfangsfällen der transfiniten Induktion in der reinen Zahlentheorie, *Math. Ann.* 119 (1943) 140-161.
- [4] G. Takeuti, A remark on Gentzen's paper "Beweibarkeit und Unbeweisbarkeit von Anfangsfällen der transfiniten Induktion in der reinen Zahlentheorie", *Proc. Japan Acad.* 39 (1963) 263-269.
- [5] G. Takeuti, *Proof Theory*, second edition, North-Holland, Amsterdam (1987) reprinted from Dover Publications (2013)