

A weighted Adams' inequality in \mathbb{R}^4 and applications

Wenjing Chen^{*†} and Shiqi Zhang

School of Mathematics and Statistics, Southwest University, Chongqing, 400715, P.R. China

Abstract

In this paper, we establish a weighted Adams' inequality in some appropriate weighted Sobolev space in \mathbb{R}^4 . Then we give an improvement inequality by proving the concentration-compactness result. In the last part, we consider an application to elliptic equation involving new exponential growth at infinity in \mathbb{R}^4 .

Keywords: Adams' inequality; Exponential growth; Radial function.

1 Introduction and statement of main results

Let $\Omega \subset \mathbb{R}^N$, $N \geq 2$ be a bounded domain. By Sobolev embedding theorem, we know that $W_0^{k,p}(\Omega) \hookrightarrow L^q(\Omega)$, for $1 \leq q \leq \frac{Np}{N-kp}$, $kp < N$ and $W_0^{k,p}(\Omega) \hookrightarrow L^q(\Omega)$, for $1 \leq q < \infty$, $kp = N$. However, it is known that $W_0^{k,p}(\Omega) \hookrightarrow L^\infty(\Omega)$ does not hold. In this case, it was provided independently by Yudovich [43], Peetre [33], Pohazaev [34], Trudinger [40] that $W_0^{1,N}(\Omega) \hookrightarrow L_\phi(\Omega)$ where $L_\phi(\Omega)$ is the Orlicz space with the N-function $\phi(t) = e^{|t|^{\frac{N}{N-1}}} - 1$. To be precise, Moser [30] proved the following inequality:

$$\sup_{u \in W_0^{1,N}(\Omega), |\nabla u|_{L^N(\Omega)} \leq 1} \int_{\Omega} e^{\alpha u^{\frac{N}{N-1}}} dx < +\infty \iff \alpha \leq \alpha_N = N\omega_{\frac{1}{N-1}},$$

where ω_{N-1} denotes the area of the unit sphere in \mathbb{R}^N .

This result has led to many related results. For instance, the attainability of the supremum has been proved by Carleson and Chang [12] when the domain is a unit ball in \mathbb{R}^N , by Flucher [19] on an open and bounded domain of \mathbb{R}^2 with smooth boundary and by Lin [27] on bounded smooth domain in any dimension. For the unbounded domain, several extensions of Trudinger-Moser type inequalities have been established, see [11, 26, 29, 37]. In particular, Ruf [37] proved the following sharp inequality:

$$\sup_{u \in H^1(\mathbb{R}^2), \int_{\mathbb{R}^2} (|u|^2 + |\nabla u|^2) dx \leq 1} \int_{\mathbb{R}^2} (e^{\alpha u^2} - 1) dx < +\infty \iff \alpha \leq 4\pi.$$

Li and Ruf [26] generalized the inequality to the N-dimensional ($N \geq 2$) case as follows:

$$\sup_{u \in W^{1,N}(\mathbb{R}^N), \int_{\mathbb{R}^N} (|u|^N + |\nabla u|^N) dx \leq 1} \int_{\mathbb{R}^N} \Phi_N \left(\alpha |u|^{\frac{N}{N-1}} \right) dx < +\infty \iff \alpha \leq \alpha_N,$$

^{*}Corresponding author.

[†]E-mail address: wjchen@swu.edu.cn (W. Chen), sqzhangmaths@163.com (S. Zhang).

where $\Phi_N(t) = e^t - \sum_{j=0}^{N-2} \frac{t^j}{j!}$. Trudinger-Moser inequalities were also given on manifolds [20, 25, 41], on Heisenberg groups [17, 23], without boundary conditions [15, 16] and references therein.

Adams [1] studied the similar result to higher order Sobolev spaces. Indeed, Adams proved that

$$\sup_{u \in W_0^{2,2}(\Omega), \int_{\Omega} |\Delta u|^2 dx \leq 1} \int_{\Omega} e^{32\pi^2 u^2} dx < \infty. \quad (1.1)$$

When Ω is replaced by the whole space \mathbb{R}^4 , the integrals in (1.1) are infinite. But Ruf and Sani [38] established the corresponding Adams type inequality as follows:

$$\sup_{u \in W_{rad}^{2,2}(\mathbb{R}^4), \|u\|_{W^{2,2}} \leq 1} \int_{\mathbb{R}^4} (e^{32\pi^2 u^2} - 1) dx < +\infty,$$

where $\|u\|_{W^{2,2}}^2 = \int_{\mathbb{R}^4} (|\Delta u|^2 + |\nabla u|^2 + u^2) dx$. More related results can see [14, 24] and references therein.

Recently, many Trudinger-Moser type and Adams' type inequalities on weighted Sobolev spaces have been studied. Calanchi and Ruf [9] first studied the case of weighted Sobolev norms with weights of logarithmic type. In fact, let $B = B_1(0)$ be the unit open ball in \mathbb{R}^2 , $\omega_0(x) = \left(\log \frac{e}{|x|}\right)^\beta$, $0 \leq \beta \leq 1$. Denote by $H_{0,rad}^1(B, \omega_0)$ the standard weighted radial Sobolev space defined as the space of all radial functions of the completion of C_0^∞ with respect to the norm

$$\|u\|_{\omega_0}^2 = \int_B |\nabla u|^2 \omega_0(x) dx.$$

Then, for $0 \leq \beta < 1$,

$$\sup_{u \in H_{0,rad}^1(B, \omega_0), \|u\|_{\omega_0} \leq 1} \int_B e^{\alpha |u|^{\frac{2}{1-\beta}}} dx < +\infty \iff \alpha \leq \alpha_{2,\beta} = 2 [2\pi(1-\beta)]^{\frac{1}{1-\beta}},$$

for $\beta = 1$,

$$\sup_{u \in H_{0,rad}^1(B, \omega_0), \|u\|_{\omega_0} \leq 1} \int_B e^{\alpha e^{2\pi u^2}} dx < +\infty \iff \alpha \leq 2.$$

Moreover, Calanchi and Ruf [10] generalized the similar results to the N-dimensional case. Here, we have to mention the work where Nguyen [32] provided a simple proof which based on a suitable change of function and obtained the existence of maximizers for this inequality when β is small enough. The attainability of the supremum of this inequality has been given by Roy [35, 36]. For the related results on unbounded domains, we can cite [2, 3, 5, 6]. In particular, Aouaoui and Jlel [5] extended the work of Calanchi and Ruf to the whole space \mathbb{R}^2 , considering the following weight:

$$\omega_1(x) = \begin{cases} \left(\log\left(\frac{e}{|x|}\right)\right)^\beta, & \text{if } |x| < 1, \\ \chi_0(|x|), & \text{if } |x| \geq 1, \end{cases} \quad (1.2)$$

where $0 < \beta \leq 1$ and $\chi_0 : [1, +\infty) \rightarrow (0, +\infty)$ is a continuous function such that $\chi_0(1) = 1$ and $\inf_{t \geq 1} \chi_0(t) > 0$. Denote by F_β as the space of all radial functions of the completion of $C_0^\infty(\mathbb{R}^2)$ with respect to the norm

$$\|u\|_{\omega_1}^2 = \int_{\mathbb{R}^2} (|\nabla u|^2 \omega_1(x) + |u|^2) dx.$$

More precisely,

Theorem A (i) Let $0 < \beta < 1$ and ω_1 be defined by (1.2). For all $u \in F_\beta$, we have

$$\int_{\mathbb{R}^2} \left(e^{|u|^{\frac{2}{1-\beta}}} - 1 \right) dx < +\infty.$$

Moreover,

$$\sup_{u \in F_\beta, \|u\|_{\omega_1} \leq 1} \int_{\mathbb{R}^2} \left(e^{\alpha|u|^{\frac{2}{1-\beta}}} - 1 \right) dx < +\infty,$$

if and only if $\alpha < \alpha_{2,\beta}$.

(ii) Let $\beta = 1$ and ω_1 be defined by (1.2). For all $u \in F_1$, we have

$$\int_{\mathbb{R}^2} \left(e^{\alpha(e^{u^2}-1)} - 1 \right) dx < +\infty.$$

Moreover, if $\alpha \leq 2e^{-\left(\inf_{s \geq 1} \chi(s)\right)^{-\frac{1}{2}}}$, then

$$\sup_{u \in F_1, \|u\|_{\omega_1} \leq 1} \int_{\mathbb{R}^2} \left(e^{\alpha(e^{2\pi u^2}-1)} - 1 \right) dx < +\infty.$$

If $\alpha > 2e^{\int_0^{+\infty} \log^2(1+t)e^{-2t} dt}$, then

$$\sup_{u \in F_1, \|u\|_{\omega_1} \leq 1} \int_{\mathbb{R}^2} \left(e^{\alpha(e^{2\pi u^2}-1)} - 1 \right) dx = +\infty.$$

The similar results in $\mathbb{R}^N (N \geq 2)$ was provided by Aouaoui and Jlel [6], considering the following weight:

$$\omega_2(x) = \begin{cases} \left(\log\left(\frac{e}{|x|}\right)\right)^{\beta(N-1)}, & \text{if } |x| < 1, \\ \chi_1(|x|), & \text{if } |x| \geq 1, \end{cases} \quad (1.3)$$

where $0 < \beta \leq 1$ and $\chi_1 : [1, +\infty) \rightarrow (0, +\infty)$ is a continuous function such that $\chi_1(1) = 1$ and $\inf_{t \geq 1} \chi_1(t) > 0$. Denote by E_β as the space of all radial functions of the completion of $C_0^\infty(\mathbb{R}^N)$ with respect to the norm

$$\|u\|_{\omega_2}^N = \int_{\mathbb{R}^N} |\nabla u|^N \omega_2(x) dx.$$

More details see [6], here we just mention the main result of the case $0 < \beta < 1$ as follows.

Theorem B Let $0 < \beta < 1$ and ω_2 be defined by (1.3). Assume that there exists a positive $C > 0$ such that

$$\int_1^{+\infty} \left(\int_r^{+\infty} \frac{ds}{s(\chi_1(s))^{\frac{1}{N-1}}} \right)^{\frac{k}{1-\beta}} r^{N-1} dr \leq C^k, \quad \text{for all } k \in \mathbb{N}^*.$$

For all $\alpha > 0$ and $u \in E_\beta$, we have

$$\int_{\mathbb{R}^N} \left(e^{\alpha|u|^{\frac{N}{(1-\beta)(N-1)}}} - 1 \right) dx < +\infty.$$

Moreover,

$$\sup_{u \in E_\beta, \|u\|_{\omega_2} \leq 1} \int_{\mathbb{R}^N} \left(e^{\alpha|u|^{\frac{N}{(1-\beta)(N-1)}}} - 1 \right) dx < +\infty,$$

if and only if

$$\alpha \leq \alpha_{N,\beta} = N \left[\omega_{\frac{N-1}{N-1}}^{\frac{1}{1-\beta}} (1-\beta) \right]^{\frac{1}{1-\beta}}.$$

Extension of the Trudinger-Moser inequality to higher order Sobolev spaces can see [44]. Zhu and Wang [44] proved the following Adams' inequality when the domain is a unit ball in \mathbb{R}^4 .

Theorem C Let $B = B_1(0) \subset \mathbb{R}^4$ denote the unit ball in \mathbb{R}^4 and $\beta \in (0, 1)$. Then

$$\sup_{u \in W_{0,rad}^{2,2}(B,\omega), \|\Delta u\|_\omega \leq 1} \int_B e^{\alpha|u|^{\frac{2}{1-\beta}}} dx < +\infty, \quad (1.4)$$

if and only if

$$\alpha \leq \alpha_\beta = 4 [8\pi^2(1-\beta)]^{\frac{1}{1-\beta}},$$

where $\omega(x) = \left(\log \frac{e}{|x|}\right)^\beta$, and

$$W_{0,rad}^{2,2}(B,\omega) = cl \left\{ u \in C_{0,rad}^\infty(B); \|\Delta u\|_\omega^2 = \int_B |\Delta u|^2 \omega(x) dx < +\infty \right\}.$$

A nature question is whether the similar result as (1.4) holds on unbounded domain. In this paper, we focus on the bi-Laplacian operator in \mathbb{R}^4 . We define the radial weight

$$\omega_\beta(x) = \begin{cases} \left(\log \left(\frac{e}{|x|}\right)\right)^\beta, & \text{if } |x| < 1, \\ \chi(|x|), & \text{if } |x| \geq 1, \end{cases} \quad (1.5)$$

where $0 < \beta < 1$, $\chi : [1, +\infty) \rightarrow (0, +\infty)$ is a continuous function such that $\chi(1) = 1$, $\inf_{t \geq 1} \chi(t) > 0$ and satisfies

$$\int_1^{+\infty} \chi(y)^{-1} y^{-1} dy < +\infty, \quad (1.6)$$

and

$$\int_1^{+\infty} \chi(y)^{-1} y dy < +\infty. \quad (1.7)$$

We define the weighted Sobolev space

$$X_\beta = \left\{ u \in L_{rad}^{\frac{2}{1-\beta}}(\mathbb{R}^4), \int_{\mathbb{R}^4} \omega_\beta(x) |\Delta u|^2 dx < +\infty \right\}, \quad \text{for all } 0 < \beta < 1,$$

with the norm

$$\|u\|_\beta^2 = \int_{\mathbb{R}^4} \omega_\beta(x) |\Delta u|^2 dx.$$

For the function χ , we make the following assumptions: there exists a constant $M > 0$ such that

$$\frac{1}{r^8} \left(\int_1^r t^3 \chi(t) dt \right) \left(\int_1^r \frac{t^3}{\chi(t)} dt \right) \leq M, \quad \text{for all } r \geq 1; \quad (1.8)$$

$$\frac{1}{r^8} \int_1^r t^3 \chi(t) dt \leq M, \quad \text{for all } r \geq 1; \quad (1.9)$$

$$\frac{\max_{r \leq t \leq 4r} \chi(t)}{\min_{r \leq t \leq 4r} \chi(t)} \leq M, \quad \text{for all } r \geq 1. \quad (1.10)$$

The conditions (1.8), (1.9) and (1.10) are used to guarantee the weight ω_β is in Muckenhoupt's A_2 -class, that is, ω_β satisfies

$$\sup_{B \subset \mathbb{R}^4} \left(\frac{1}{|B|} \int_B \omega_\beta(x) dx \right) \left(\frac{1}{|B|} \int_B \omega_\beta(x)^{-1} dx \right) < +\infty, \quad (1.11)$$

see [13, 21, 31] for more details.

The main results are as follows.

Theorem 1.1. *Let $0 < \beta < 1$ and ω_β be defined by (1.5). We suppose that*

(D₁)

$$\int_1^{+\infty} r^3 \left(\int_r^{+\infty} \frac{1}{\chi(y)y} dy \right)^{\frac{k}{1-\beta}} dr < C_1^k, \quad C_1 > 0.$$

(D₂)

$$\int_1^{+\infty} r^{3-\frac{2k}{1-\beta}} \left(\int_1^r \frac{y}{\chi(y)} dy \right)^{\frac{k}{1-\beta}} dr < C_2^k, \quad C_2 > 0.$$

For $u \in X_\beta$, we have

$$\int_{\mathbb{R}^4} \left(e^{|u|^{\frac{2}{1-\beta}}} - 1 \right) dx < +\infty. \quad (1.12)$$

Moreover, if $\alpha \leq \alpha_\beta$, then

$$\sup_{u \in X_\beta, \|u\|_\beta \leq 1} \int_{\mathbb{R}^4} \left(e^{\alpha|u|^{\frac{2}{1-\beta}}} - 1 \right) dx < +\infty. \quad (1.13)$$

If $\alpha > \alpha_\beta$, then

$$\sup_{u \in X_\beta, \|u\|_\beta \leq 1} \int_{\mathbb{R}^4} \left(e^{\alpha|u|^{\frac{2}{1-\beta}}} - 1 \right) dx = +\infty.$$

Remark 1.1. An example of function χ satisfying (1.6), (1.7), (1.8), (1.9), (1.10), (D₁) and (D₂) is given by

$$\chi(y) = y^\alpha, \quad \max\{2, 4(1-\beta)\} < \alpha < 4.$$

Notice that there is already the result of Adams' inequality in the $B \subset \mathbb{R}^4$, that is Theorem C. The key point and difficulty of the proof in Theorem 1.1 is that we need a radial lemma to deal with $u(x)$ if $|x|$ large enough. Inspired by [39, 44], we make a suitable change of variable. The more details see Lemma 2.1. Then, we establish an Lions [28] type result about an improvement of (1.13) for weakly convergent sequences in E_β with a constant larger than that found in (1.13) which is used to treat nonlinear equations with exponential growth at infinity. More precisely, we prove the theorem as follows.

Theorem 1.2. *Let $0 < \beta < 1$. Assume that (D_1) and (D_2) hold. Let $\{u_n\} \subset X_\beta$ and $u \in X_\beta \setminus \{0\}$ be such that $\|u_n\|_\beta = 1$ and $u_n \rightharpoonup u$ weakly in X_β . Then*

$$\sup_n \int_{\mathbb{R}^4} \left(e^{p\alpha_\beta |u_n|^{\frac{2}{1-\beta}}} - 1 \right) dx < +\infty, \quad \text{for all } 0 < p < P_\beta(u), \quad (1.14)$$

where

$$P_\beta(u) = \begin{cases} \frac{1}{(1-\|u\|_\beta^2)^{\frac{1}{1-\beta}}}, & \text{if } \|u\|_\beta < 1, \\ +\infty, & \text{if } \|u\|_\beta = 1. \end{cases} \quad (1.15)$$

Next, we will use the Adams' inequalities established in Theorem 1.1 and the improved inequality given in Theorem 1.2 to study the weighted elliptic problem involving nonlinearities which satisfy some conditions including the exponential growth. For $0 < \beta < 1$, we consider the problem

$$\Delta(\omega_\beta(x)\Delta u) = f(u) \quad \text{in } \mathbb{R}^4. \quad (1.16)$$

We assume

(F_1) $f : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function such that $f(s) = 0$, for all $s \leq 0$ and

$$|f(s)| \leq c_0 \left(|s|^{p-1} + |s|^{q-1} \left(e^{\alpha|s|^{\frac{2}{1-\beta}}} - 1 \right) \right), \quad \text{for all } s \in \mathbb{R},$$

with $c_0 > 0$, $\min\{p, q\} > \frac{2}{1-\beta}$, $\alpha > 0$.

(F_2) There exist $M_0 > 0$ and $s_0 > 0$ such that

$$F(s) = \int_0^s f(t)dt \leq M_0 f(s), \quad \text{for all } s \geq s_0.$$

(F_3) There exists $\kappa > \frac{2}{\pi^2 e^4} \left(\frac{\alpha_\beta}{\alpha} \right)^{1-\beta}$ such that $\liminf_{s \rightarrow \infty} \frac{f(s)s}{e^{\alpha s \frac{2}{1-\beta}}} \geq \kappa$.

Theorem 1.3. *Let $0 < \beta < 1$. Assume that (D_1) , (D_2) and $(F_1) - (F_3)$ hold. Then, problem (1.16) has at least one nontrivial and nonnegative weak solution.*

Remark 1.2. An example of function f satisfying the conditions $(F_1) - (F_3)$ is given by

$$f(s) = \begin{cases} C s^{p-1} \left(\frac{2\alpha}{1-\beta} s^{\frac{2}{1-\beta}} + 1 \right) \left(e^{\alpha s \frac{2}{1-\beta}} - 1 \right), & \text{if } s \geq 0, \\ 0, & \text{if } s < 0, \end{cases}$$

where $p > \frac{2}{1-\beta}$, $\alpha > 0$ and $C > 0$ is chosen large enough.

Indeed, we have

$$F(s) = \int_0^s f(t)dt = \begin{cases} C s^p \left(e^{\alpha s \frac{2}{1-\beta}} - 1 \right), & \text{if } s \geq 0, \\ 0, & \text{if } s < 0. \end{cases}$$

Remark 1.3. In general, we often use the following (AR) condition to guarantee the Mountain Pass geometry and the boundedness of the (PS) sequences.

(AR) There exists $\theta > 2$ such that $0 < \theta F(t) \leq f(t)t$ for all $t \geq 0$.

In this paper, we only use the conditions (F_1) and (F_3) to prove the Mountain Pass geometry of I_β and use the condition (F_2) to guarantee the boundedness of (PS) sequences. The main difficulty of the proof is the lack of compactness. In fact, we can prove that the (PS) condition is satisfied at a certain level of energy using Theorem 1.2 and (F_1) mainly. Inspired by [42], by considering the family of Adams type function, we will estimate the level of the energy I_β using the condition (F_3) . It is worth mentioning that the condition (F_3) we give in this paper is weaker than the corresponding condition (H_5) in [42]. The more details see Section 5.

Remark 1.4. The conditions (1.6) and (1.7) do not work in the proof of Theorem 1.1 and Theorem 1.2, but are crucial to the proof of Theorem 1.3. More specifically, the conditions play an important part in Proposition 5.1. Indeed, the condition (1.6) is to guarantee the boundedness of $u(e_1)$. The condition (1.7) is to guarantee $u(x) \rightarrow 0$ when $|x| \rightarrow \infty$ which is sufficient to prove the compactness of the embedding. More details see Proposition 5.1.

This paper is organized as follows: In Section 2, we prove some results which will be used in the next sections. Theorem 1.1 will be proved in Section 3. Section 4 is devoted for a concentration compactness result of Lions type which will be used to prove the application in Section 5. The constant C in this paper will change according to different situations.

2 Preliminaries

In this section, we establish an estimate of $u(x)$ for $|x|$ large enough which plays an important role in the proof of Theorem 1.1. First, we prove the following density property.

Proposition 2.1. *If χ satisfies the conditions (1.8), (1.9) and (1.10), then $C_{0,rad}^\infty(\mathbb{R}^4)$ is dense in X_β .*

Proof. Similar to [5, 6], we only need to prove $\omega_\beta \in A_2$, that is, we show that (1.11) holds.

Let $r > 0$ and $x_0 \in \mathbb{R}^4$. Denote by $B(x_0, r)$ the open ball of \mathbb{R}^4 of center x_0 and radius r .

Case 1. $B(x_0, r) \cap B(0, r) \neq \emptyset$. In this case $B(x_0, r) \subset B(0, 3r)$ which implies that

$$\begin{aligned} & \frac{1}{|B(x_0, r)|} \left(\int_{B(x_0, r)} \omega_\beta(x) dx \right) \left(\frac{1}{|B(x_0, r)|} \int_{B(x_0, r)} \omega_\beta(x)^{-1} dx \right) \\ & \leq \frac{c}{r^8} \left(\int_0^{3r} \omega_\beta(t) t^3 dt \right) \left(\int_0^{3r} \frac{t^3}{\omega_\beta(t)} dt \right). \end{aligned} \quad (2.1)$$

If $3r < 1$, after a simple calculation, we have

$$\limsup_{r \rightarrow 0^+} \frac{c}{r^8} \left(\int_0^{3r} t^3 (1 - \log t)^\beta dt \right) \left(\int_0^{3r} \frac{t^3}{(1 - \log t)^\beta} dt \right) < +\infty. \quad (2.2)$$

If $3r \geq 1$, then

$$\begin{aligned} & \frac{c}{r^8} \left(\int_0^{3r} \omega_\beta(t) t^3 dt \right) \left(\int_0^{3r} \frac{t^3}{\omega_\beta(t)} dt \right) \\ & = \frac{c}{r^8} \left(\int_0^1 t^3 (1 - \log t)^\beta dt + \int_1^{3r} t^3 \chi(t) dt \right) \\ & \times \left(\int_0^1 \frac{t^3}{(1 - \log t)^\beta} dt + \int_1^{3r} \frac{t^3}{\chi(t)} dt \right). \end{aligned} \quad (2.3)$$

Since $\inf_{t \geq 1} \chi(t) > 0$, then

$$\limsup_{r \rightarrow +\infty} \frac{1}{r^8} \left(\int_1^{3r} \frac{t^3}{\chi(t)} dt \right) = 0 < +\infty. \quad (2.4)$$

Combining (1.8), (1.9), (2.2), (2.3) and (2.4), we deduce from (2.1) that

$$\frac{1}{|B(x_0, r)|} \left(\int_{B(x_0, r)} \omega_\beta(x) dx \right) \left(\frac{1}{|B(x_0, r)|} \int_{B(x_0, r)} \omega_\beta(x)^{-1} dx \right) < \infty. \quad (2.5)$$

Case 2. $B(x_0, r) \cap B(0, r) = \emptyset$. In this case, we have

$$\frac{|x_0|}{2} \leq |x| \leq 2|x_0|, \quad \text{for all } x \in B(x_0, r).$$

Hence,

$$\begin{aligned} & \left(\frac{1}{|B(x_0, r)|} \int_{B(x_0, r)} \omega_\beta(x) dx \right) \left(\frac{1}{|B(x_0, r)|} \int_{B(x_0, r)} \omega_\beta(x)^{-1} dx \right) \\ & \leq \frac{\sup_{\frac{|x_0|}{2} \leq |x| \leq 2|x_0|} \omega_\beta(x)}{\inf_{\frac{|x_0|}{2} \leq |x| \leq 2|x_0|} \omega_\beta(x)} \\ & \leq \sup_{\alpha > 0} \left(\frac{\sup_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)}{\inf_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)} \right). \end{aligned} \quad (2.6)$$

If $4\alpha < 1$, then

$$\frac{\sup_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)}{\inf_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)} = \frac{(1 - \log \alpha)^\beta}{(1 - \log(4\alpha))^\beta},$$

and consequently,

$$\sup_{0 < \alpha < \frac{1}{4}} \left(\frac{\sup_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)}{\inf_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)} \right) = \sup_{0 < \alpha < \frac{1}{4}} \left(\frac{1 - \log \alpha}{1 - \log(4\alpha)} \right)^\beta < +\infty. \quad (2.7)$$

If $\frac{1}{4} \leq \alpha < 1$, by the continuity of the function ω_β and the boundedness theorem, we get

$$\frac{\sup_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)}{\inf_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)} \leq \frac{\sup_{\frac{1}{4} \leq t \leq 4} \omega_\beta(t)}{\inf_{\frac{1}{4} \leq t \leq 4} \omega_\beta(t)} < +\infty,$$

and consequently,

$$\sup_{\frac{1}{4} \leq \alpha < 1} \left(\frac{\sup_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)}{\inf_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)} \right) < +\infty. \quad (2.8)$$

If $\alpha \geq 1$, then by (1.10) we have

$$\frac{\sup_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)}{\inf_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)} \leq \frac{\sup_{\alpha \leq t \leq 4\alpha} \chi(t)}{\inf_{\alpha \leq t \leq 4\alpha} \chi(t)} < +\infty,$$

and consequently,

$$\sup_{\alpha \geq 1} \left(\frac{\sup_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)}{\inf_{\alpha \leq t \leq 4\alpha} \omega_\beta(t)} \right) < +\infty. \quad (2.9)$$

Combining (2.7), (2.8) and (2.9), we deduce from (2.6) that

$$\frac{1}{|B(x_0, r)|} \left(\int_{B(x_0, r)} \omega_\beta(x) dx \right) \left(\frac{1}{|B(x_0, r)|} \int_{B(x_0, r)} \omega_\beta(x)^{-1} dx \right) < \infty. \quad (2.10)$$

From (2.5) and (2.10), we have $\omega_\beta \in A_2$. Then, $C_{0,rad}^\infty(\mathbb{R}^4)$ is dense in X_β . \square

Now, using the density of $C_{0,rad}^\infty(\mathbb{R}^4)$ in X_β , we can prove the following radial lemma which is based on a suitable change of variable, see [39, 44].

Lemma 2.1. *Fix $0 < \beta < 1$ and let $u \in X_\beta$. Then one has*

$$|u(x)| \leq \frac{\sqrt{2}c_1}{4\pi} \left(\int_{|x| \geq 1} |\Delta u|^2 \chi(|x|) dx \right)^{\frac{1}{2}} \left[\left(\int_{|x|}^{+\infty} \chi(y)^{-1} y^{-1} dy \right)^{\frac{1}{2}} + \left(\int_1^{|x|} \chi(y)^{-1} y |x|^{-2} dy \right)^{\frac{1}{2}} \right] + \frac{c_2}{|x|^2},$$

for $|x| \geq 1$.

(2.11)

Proof. Since $C_{0,rad}^\infty(\mathbb{R}^4)$ is dense in X_β , it is enough to prove (2.11) when $u \in C_{0,rad}^\infty(\mathbb{R}^4)$. Setting

$$\omega(s) = 4\pi u(x), \quad r = |x| = s^{-\frac{1}{2}}, \quad \text{for } 0 < s \leq 1.$$

Then we have

$$\begin{aligned} \lim_{s \rightarrow 0^+} \omega(0) &= 0, \quad \lim_{s \rightarrow 1^-} \omega'(s) = \omega'(1) = -2\pi u'(1), \quad c_2 \triangleq |-2\pi u'(1)| \\ \omega''(s) &= \pi \cdot s^{-3} \Delta u \Big|_{|x|=s^{-\frac{1}{2}}}, \end{aligned}$$

and

$$\begin{aligned} \int_{|x| \geq 1} |\Delta u|^2 \chi(|x|) dx &= 2\pi^2 \int_1^{+\infty} |\Delta u|^2 \chi(r) r^3 dr \\ &= \pi^2 \int_0^1 \left| \Delta u \left(s^{-\frac{1}{2}} \right) \right|^2 \chi \left(s^{-\frac{1}{2}} \right) s^{-3} ds \\ &= \int_0^1 |\omega''(s)|^2 \chi \left(s^{-\frac{1}{2}} \right) s^3 ds. \end{aligned}$$

Let us now evaluate $\omega(s)$, we have

$$\begin{aligned}
|\omega(s)| &= \left| \int_0^s \omega'(z) dz \right| \\
&= \left| \int_0^s \left(\int_z^1 \omega''(t) dt - \omega'(1) \right) dz \right| \\
&\leq \int_0^s \left| \int_z^1 \omega''(t) dt \right| dz + \left| \int_0^s c_2 dz \right| \\
&\leq \left(\int_0^s \int_z^1 |\omega''(t)|^2 \chi(t^{-\frac{1}{2}}) t^2 dt dz \right)^{\frac{1}{2}} \cdot \left(\int_0^s \int_z^1 \chi(t^{-\frac{1}{2}})^{-1} t^{-2} dt dz \right)^{\frac{1}{2}} + |c_2 s| \\
&\leq \left(\int_0^s |\omega''(t)|^2 \chi(t^{-\frac{1}{2}}) t^2 dt \int_0^t dz + \int_s^1 |\omega''(t)|^2 \chi(t^{-\frac{1}{2}}) t^2 dt \int_0^s dz \right)^{\frac{1}{2}} \\
&\quad \cdot \left(\int_0^s \int_z^1 \chi(t^{-\frac{1}{2}})^{-1} t^{-2} dt dz \right)^{\frac{1}{2}} + |c_2 s| \\
&\leq \left(\int_0^s |\omega''(t)|^2 \chi(t^{-\frac{1}{2}}) t^3 dt + \int_s^1 |\omega''(t)|^2 \chi(t^{-\frac{1}{2}}) t^3 dt \right)^{\frac{1}{2}} \\
&\quad \cdot \left(\int_0^s \int_z^1 \chi(t^{-\frac{1}{2}})^{-1} t^{-2} dt dz \right)^{\frac{1}{2}} + |c_2 s| \\
&= \left(\int_0^1 |\omega''(t)|^2 \chi(t^{-\frac{1}{2}}) t^3 dt \right)^{\frac{1}{2}} \left(\int_0^s \int_z^1 \chi(t^{-\frac{1}{2}})^{-1} t^{-2} dt dz \right)^{\frac{1}{2}} + |c_2 s| \\
&= \left(\int_{|x| \geq 1} |\Delta u|^2 \chi(|x|) dx \right)^{\frac{1}{2}} \left(\int_0^s \int_z^1 \chi(t^{-\frac{1}{2}})^{-1} t^{-2} dt dz \right)^{\frac{1}{2}} + |c_2 s|,
\end{aligned}$$

and

$$\begin{aligned}
\int_0^s \int_z^1 \chi(t^{-\frac{1}{2}})^{-1} t^{-2} dt dz &= \int_0^s \chi(t^{-\frac{1}{2}})^{-1} t^{-2} dt \int_0^t dz + \int_s^1 \chi(t^{-\frac{1}{2}})^{-1} t^{-2} dt \int_0^s dz \\
&= \int_0^s \chi(t^{-\frac{1}{2}})^{-1} t^{-1} dt + \int_s^1 \chi(t^{-\frac{1}{2}})^{-1} t^{-2} s dt.
\end{aligned}$$

Thus we have

$$|\omega(s)| \leq \left(\int_{|x| \geq 1} |\Delta u|^2 \chi(|x|) dx \right)^{\frac{1}{2}} \left(\int_0^s \chi(t^{-\frac{1}{2}})^{-1} t^{-1} dt + \int_s^1 \chi(t^{-\frac{1}{2}})^{-1} t^{-2} s dt \right)^{\frac{1}{2}} + |c_2 s|.$$

Consequently,

$$\begin{aligned}
|u(x)| &\leq \frac{c_1}{4\pi} \left(\int_{|x|\geq 1} |\Delta u|^2 \chi(|x|) dx \right)^{\frac{1}{2}} \left[\left(\int_0^{\frac{1}{|x|^2}} \chi \left(t^{-\frac{1}{2}} \right)^{-1} t^{-1} dt \right)^{\frac{1}{2}} + \left(\int_{\frac{1}{|x|^2}}^1 \chi \left(t^{-\frac{1}{2}} \right)^{-1} t^{-2} |x|^{-2} dt \right)^{\frac{1}{2}} \right] \\
&\quad + \frac{c_2}{|x|^2} \\
&= \frac{\sqrt{2}c_1}{4\pi} \left(\int_{|x|\geq 1} |\Delta u|^2 \chi(|x|) dx \right)^{\frac{1}{2}} \left[\left(\int_{|x|}^{+\infty} \chi(y)^{-1} y^{-1} dy \right)^{\frac{1}{2}} + \left(\int_1^{|x|} \chi(y)^{-1} y |x|^{-2} dy \right)^{\frac{1}{2}} \right] + \frac{c_2}{|x|^2}.
\end{aligned}$$

□

3 Proof of Theorem 1.1

In this section, we prove the capital Adams' inequality in detail. We start by proving the first conclusion of Theorem 1.1. For $u \in X_\beta$, we have

$$\int_{\mathbb{R}^4} \left(e^{|u|^{\frac{2}{1-\beta}}} - 1 \right) dx = \int_{|x|\geq 1} \left(e^{|u|^{\frac{2}{1-\beta}}} - 1 \right) dx + \int_{|x|<1} \left(e^{|u|^{\frac{2}{1-\beta}}} - 1 \right) dx. \quad (3.1)$$

On the one hand,

$$\int_{|x|\geq 1} \left(e^{|u|^{\frac{2}{1-\beta}}} - 1 \right) dx = \sum_{k=1}^{+\infty} \frac{1}{k!} \int_{|x|\geq 1} |u|^{\frac{2k}{1-\beta}} dx. \quad (3.2)$$

From (2.11) and (D_1) , (D_2) , we get

$$\begin{aligned}
\int_{|x|\geq 1} |u|^{\frac{2k}{1-\beta}} dx &\leq \int_{|x|\geq 1} \left\{ \frac{\sqrt{2}c_1}{4\pi} \|u\|_\beta \left[\left(\int_{|x|}^{+\infty} \chi(y)^{-1} y^{-1} dy \right)^{\frac{1}{2}} + \left(\int_1^{|x|} \chi(y)^{-1} y |x|^{-2} dy \right)^{\frac{1}{2}} \right] + \frac{c_2}{|x|^2} \right\}^{\frac{2k}{1-\beta}} dx \\
&\leq C \|u\|_\beta^{\frac{2k}{1-\beta}} \left[\int_{|x|\geq 1} \left(\int_{|x|}^{+\infty} \chi(y)^{-1} y^{-1} dy \right)^{\frac{k}{1-\beta}} dx + \int_{|x|\geq 1} \left(\int_1^{|x|} \chi(y)^{-1} y |x|^{-2} dy \right)^{\frac{k}{1-\beta}} dx \right] \\
&\quad + C \int_{|x|\geq 1} \frac{1}{|x|^{\frac{4k}{1-\beta}}} dx \\
&= C \|u\|_\beta^{\frac{2k}{1-\beta}} 2\pi^2 \left[\int_1^{+\infty} r^3 \left(\int_r^{+\infty} \chi(y)^{-1} y^{-1} dy \right)^{\frac{k}{1-\beta}} dr + \int_1^{+\infty} r^{3-\frac{2k}{1-\beta}} \left(\int_1^r \frac{y}{\chi(y)} dy \right)^{\frac{k}{1-\beta}} dr \right] \\
&\quad + 2\pi^2 \int_1^{+\infty} \frac{1}{r^{\frac{4k}{1-\beta}-3}} dr < C^k, \quad \text{for all } k \geq 1.
\end{aligned} \quad (3.3)$$

Combining (3.2) and (3.3), we have

$$\int_{|x|\geq 1} \left(e^{|u|^{\frac{2}{1-\beta}}} - 1 \right) dx \leq \sum_{k=1}^{+\infty} \frac{C^k}{k!} = e^C - 1 < +\infty. \quad (3.4)$$

Now in order to estimate the second integral in (3.1), set

$$v(x) = \begin{cases} u(x) - u(e_1), & 0 \leq |x| < 1, \\ 0, & |x| \geq 1, \end{cases} \quad (3.5)$$

where $e_1 = (1, 0, 0, 0) \in \mathbb{R}^4$. Clearly $v \in W_{0,rad}^{2,2}(B, \omega)$. After an elementary calculation, we have

$$|u|^{\frac{2}{1-\beta}} = |v + u(e_1)|^{\frac{2}{1-\beta}} \leq (1 + \epsilon)|v|^{\frac{2}{1-\beta}} + \left(1 - \frac{1}{(1 + \epsilon)^{\frac{1-\beta}{1+\beta}}}\right)^{\frac{1+\beta}{\beta-1}} |u(e_1)|^{\frac{2}{1-\beta}},$$

for fix $\epsilon > 0$.

Then, using [7, Lemma 2(ii)], we have

$$\begin{aligned} \int_{|x|<1} e^{|u|^{\frac{2}{1-\beta}}} dx &\leq \int_{|x|<1} e^{(1+\epsilon)|v|^{\frac{2}{1-\beta}}} \cdot e^{\left(1 - \frac{1}{(1+\epsilon)^{\frac{1-\beta}{1+\beta}}}\right)^{\frac{1+\beta}{\beta-1}} |u(e_1)|^{\frac{2}{1-\beta}}} dx \\ &\leq e^{\left(1 - \frac{1}{(1+\epsilon)^{\frac{1-\beta}{1+\beta}}}\right)^{\frac{1+\beta}{\beta-1}} |u(e_1)|^{\frac{2}{1-\beta}}} \int_{|x|<1} e^{(1+\epsilon)|v|^{\frac{2}{1-\beta}}} dx < +\infty. \end{aligned} \quad (3.6)$$

Combining (3.1), (3.4) and (3.6), we conclude that

$$\int_{\mathbb{R}^4} \left(e^{|u|^{\frac{2}{1-\beta}}} - 1 \right) dx < +\infty, \quad \text{for all } u \in X_\beta.$$

This ends the proof of (1.12).

On the other hand, for any $u \in X_\beta$, $\|u\|_\beta \leq 1$, notice that

$$\|\Delta v\|_\omega^2 = \int_B |\Delta v|^2 \omega(x) dx = \int_B |\Delta u|^2 \omega_\beta(x) dx \leq \|u\|_\beta^2 \leq 1. \quad (3.7)$$

Let $\alpha < \alpha_\beta$. Plainly, there exists $\epsilon > 0$ such that $\alpha(1 + \epsilon) < \alpha_\beta$.

Combining (3.7) and (1.4), we have

$$\begin{aligned} \sup_{u \in X_\beta, \|u\|_\beta \leq 1} \int_{|x|<1} e^{\alpha|u|^{\frac{2}{1-\beta}}} dx &\leq C \sup_{u \in X_\beta, \|u\|_\beta \leq 1} \int_{|x|<1} e^{\alpha(1+\epsilon)|v|^{\frac{2}{1-\beta}}} dx \\ &\leq C \sup_{v \in W_{0,rad}^{2,2}(B, \omega), \|v\|_\beta \leq 1} \int_{|x|<1} e^{\alpha_\beta|v|^{\frac{2}{1-\beta}}} dx < +\infty. \end{aligned} \quad (3.8)$$

Furthermore

$$\int_{|x|\geq 1} \left(e^{\alpha|u|^{\frac{2}{1-\beta}}} - 1 \right) dx = \sum_{k=1}^{+\infty} \frac{\alpha^k}{k!} \int_{|x|\geq 1} |u|^{\frac{2k}{1-\beta}} dx. \quad (3.9)$$

Combining (3.3) and (3.9), we infer

$$\sup_{u \in X_\beta, \|u\|_\beta \leq 1} \int_{|x|\geq 1} \left(e^{\alpha|u|^{\frac{2}{1-\beta}}} - 1 \right) dx < +\infty. \quad (3.10)$$

It follows from (3.8) and (3.10) that

$$\sup_{u \in X_\beta, \|u\|_\beta \leq 1} \int_{\mathbb{R}^4} \left(e^{\alpha|u|^{\frac{2}{1-\beta}}} - 1 \right) dx < +\infty, \quad \text{for all } \alpha < \alpha_\beta.$$

Now we prove that

$$\sup_{u \in X_\beta, \|u\|_\beta \leq 1} \int_{\mathbb{R}^4} \left(e^{\alpha_\beta|u|^{\frac{2}{1-\beta}}} - 1 \right) dx < +\infty.$$

For any $u \in X_\beta$, $\|u\|_\beta = \int_{\mathbb{R}^4} |\Delta u|^2 \omega_\beta(x) dx \leq 1$, we discuss it in three cases.

Case 1. Assume that

$$\int_{|x|<1} |\Delta u|^2 \omega_\beta(x) dx = 1.$$

Then

$$\int_{|x|\geq 1} |\Delta u|^2 \omega_\beta(x) dx = 0. \quad (3.11)$$

From (2.11), we have

$$|u(x)| \leq c_2, \quad \text{for all } |x| \geq 1. \quad (3.12)$$

Having in mind that $u \in C_{0,rad}^\infty(\mathbb{R}^4)$. Then combining (3.11) and (3.12), we deduce that for any $|x| \geq 1$, $u(x) = 0$. In particular, we have $u(1) = 0$. So, $u|_{B \in W_{0,rad}^{2,2}(B, \omega)}$. Therefore, by Theorem C, we get

$$\int_{|x|<1} \left(e^{\alpha_\beta|u|^{\frac{2}{1-\beta}}} - 1 \right) dx \leq \int_{|x|<1} e^{\alpha_\beta|u|^{\frac{2}{1-\beta}}} dx \leq \sup_{v \in W_{0,rad}^{2,2}(B, \omega), \|\Delta v\|_\omega \leq 1} \int_B e^{\alpha_\beta|v|^{\frac{2}{1-\beta}}} dx < +\infty.$$

Case 2. Assume that

$$\int_{|x|<1} |\Delta u|^2 \omega_\beta(x) dx = 0.$$

We set $v(x)$ as (3.5), similar to (3.6), we have

$$\int_{|x|<1} \left(e^{\alpha_\beta|u|^{\frac{2}{1-\beta}}} - 1 \right) dx \leq \int_{|x|<1} e^{\alpha_\beta|u|^{\frac{2}{1-\beta}}} dx \leq C \int_{|x|<1} e^{\alpha_\beta(1+\epsilon)|v|^{\frac{2}{1-\beta}}} dx. \quad (3.13)$$

Notice that $(1+\epsilon)^{\frac{1-\beta}{2}} v \in W_{0,rad}^{2,2}(B, \omega)$ and

$$\left\| \Delta \left((1+\epsilon)^{\frac{1-\beta}{2}} v \right) \right\|_\omega^2 = (1+\epsilon)^{1-\beta} \int_{|x|<1} |\Delta v|^2 \omega(x) dx = (1+\epsilon)^{1-\beta} \int_{|x|<1} |\Delta u|^2 \omega_\beta(x) dx = 0 < 1.$$

Then, by Theorem C, we get

$$\int_{|x|<1} \left(e^{\alpha_\beta|u|^{\frac{2}{1-\beta}}} - 1 \right) dx \leq C \int_{|x|<1} e^{\alpha_\beta \left| (1+\epsilon)^{\frac{1-\beta}{2}} v \right|^{\frac{2}{1-\beta}}} dx \leq \sup_{v \in W_{0,rad}^{2,2}(B, \omega), \|\Delta v\|_\omega \leq 1} \int_B e^{\alpha_\beta|v|^{\frac{2}{1-\beta}}} dx < +\infty.$$

Case 3. Assume that

$$0 < \int_{|x|<1} |\Delta u|^2 \omega_\beta(x) dx < 1.$$

Similar to the previous case, we also have (3.13).

Let $1 + \epsilon = \left(\int_{|x|<1} |\Delta u|^2 \omega_\beta(x) dx \right)^{-\frac{1}{1-\beta}}$, we have

$$\left\| \Delta \left((1 + \epsilon)^{\frac{1-\beta}{2}} v \right) \right\|_\omega^2 = (1 + \epsilon)^{1-\beta} \int_{|x|<1} |\Delta u|^2 \omega_\beta(x) dx = 1.$$

Then, by Theorem C, we get

$$\int_{|x|<1} \left(e^{\alpha_\beta |u|^{\frac{2}{1-\beta}}} - 1 \right) dx \leq C \int_{|x|<1} e^{\alpha_\beta \left| (1+\epsilon)^{\frac{1-\beta}{2}} v \right|^{\frac{2}{1-\beta}}} dx \leq \sup_{v \in W_{0,rad}^{2,2}(B,\omega), \|\Delta v\|_\omega \leq 1} \int_B e^{\alpha_\beta |v|^{\frac{2}{1-\beta}}} dx < +\infty.$$

This ends the proof of (1.13).

Next, we show that if $\alpha > \alpha_\beta$, then the supremum is infinite.

We consider the family of Adams type function, see [4, 44]

$$u_\epsilon = \begin{cases} \left(\frac{\log \frac{e^4}{\epsilon}}{\alpha_\beta} \right)^{\frac{1-\beta}{2}} - \frac{|x|^{2(1-\beta)}}{2 \left(\frac{\alpha_\beta}{4} \epsilon \right)^{\frac{1-\beta}{2}} \left(\frac{1}{4} \log \frac{e^4}{\epsilon} \right)^{\frac{1+\beta}{2}}} + \frac{1}{2 \left(\frac{\alpha_\beta}{4} \right)^{\frac{1-\beta}{2}} \left(\frac{1}{4} \log \frac{e^4}{\epsilon} \right)^{\frac{1+\beta}{2}}}, & |x| \leq \sqrt[4]{\epsilon}, \\ \frac{\left(\log \frac{e}{|x|} \right)^{1-\beta}}{\left(\frac{\alpha_\beta}{16} \log \frac{e^4}{\epsilon} \right)^{\frac{1-\beta}{2}}}, & \sqrt[4]{\epsilon} < |x| \leq \frac{1}{2}, \\ \eta_\epsilon, & |x| > \frac{1}{2}, \end{cases}$$

where $\eta \in C_0^\infty(B_1)$ is such that

$$\eta_\epsilon|_{x=\frac{1}{2}} = \frac{1}{\left(\frac{\alpha_\beta}{16} \log \frac{e^4}{\epsilon} \right)^{\frac{1-\beta}{2}}} (\log 2e)^{1-\beta}, \quad \frac{\partial \eta_\epsilon}{\partial x} \Big|_{x=\frac{1}{2}} = -\frac{2(1-\beta)}{\left(\frac{\alpha_\beta}{16} \log \frac{e^4}{\epsilon} \right)^{\frac{1-\beta}{2}}} (\log 2e)^{-\beta},$$

and $\eta_\epsilon, \nabla \eta_\epsilon, \Delta \eta_\epsilon$ are all $O\left(\frac{1}{\left(\log \frac{e^4}{\epsilon}\right)^{\frac{1-\beta}{2}}}\right)$ as $\epsilon \rightarrow 0^+$.

By direct computation, we get

$$\Delta u_\epsilon = \begin{cases} -\frac{(1-\beta)(4-2\beta)|x|^{-2\beta}}{\left(\frac{\alpha_\beta}{4} \epsilon \right)^{\frac{1-\beta}{2}} \left(\frac{1}{4} \log \frac{1}{\epsilon} \right)^{\frac{1+\beta}{2}}}, & |x| \leq \sqrt[4]{\epsilon}, \\ \frac{1-\beta}{\left(\frac{\alpha_\beta}{16} \log \frac{1}{\epsilon} \right)^{\frac{1-\beta}{2}}} \left(\log \frac{1}{|x|} \right)^{-\beta} \frac{1}{|x|^2} \left(\frac{-\beta}{\log \frac{1}{|x|}} - 2 \right), & \sqrt[4]{\epsilon} < |x| \leq \frac{1}{2}, \\ \Delta \eta_\epsilon, & |x| > \frac{1}{2}. \end{cases}$$

Now, we evaluate the weighted Sobolev norm of u_ϵ as follows

$$\begin{aligned} \|u_\epsilon\|_\beta^2 &= 2\pi^2 \underbrace{\int_0^{\sqrt[4]{\epsilon}} |\Delta u_\epsilon|^2 \left(\log \frac{e}{|x|} \right)^\beta |x|^3 dx}_{A_1} + 2\pi^2 \underbrace{\int_{\sqrt[4]{\epsilon}}^{\frac{1}{2}} |\Delta u_\epsilon|^2 \left(\log \frac{e}{|x|} \right)^\beta |x|^3 dx}_{A_2} \\ &\quad + 2\pi^2 \underbrace{\int_{\frac{1}{2}}^1 |\Delta \eta_\epsilon|^2 \left(\log \frac{e}{|x|} \right)^\beta |x|^3 dx + 2\pi^2 \int_1^{+\infty} |\Delta \eta_\epsilon|^2 \chi(|x|) |x|^3 dx}_{A_3}. \end{aligned}$$

Similar to [4, 44], an easy computation gives

$$A_1 = O\left(\frac{1}{\log \frac{\epsilon}{\sqrt[4]{\epsilon}}}\right), A_2 = 1 + O\left(\frac{1}{\left(\log \frac{\epsilon}{\sqrt[4]{\epsilon}}\right)^{1-\beta}}\right), A_3 = O\left(\frac{1}{\left(\log \frac{\epsilon}{\sqrt[4]{\epsilon}}\right)^{1-\beta}}\right).$$

Then

$$\|u_\epsilon\|_\beta^2 \rightarrow 1, \quad \text{as } \epsilon \rightarrow 0^+.$$

Now we normalize u_ϵ , set

$$\tilde{u}_\epsilon = \frac{u_\epsilon}{\|u_\epsilon\|_\beta} \in X_\beta.$$

Moreover, we have

$$\tilde{u}_\epsilon \geq \frac{\left(\frac{\log \frac{\epsilon^4}{\epsilon}}{\alpha_\beta}\right)^{\frac{1-\beta}{2}}}{\|u_\epsilon\|_\beta}, \quad \text{for all } x \in \mathbb{R}^4, |x| \leq \sqrt[4]{\epsilon}.$$

So when $\alpha > \alpha_\beta$, for any $u \in X_\beta$, $\|u\|_\beta \leq 1$, we have

$$\begin{aligned} \int_{\mathbb{R}^4} \left(e^{\alpha|u|^{\frac{2}{1-\beta}}} - 1 \right) dx &\geq \lim_{\epsilon \rightarrow 0^+} \int_{|x| \leq \sqrt[4]{\epsilon}} \left(e^{\alpha|\tilde{u}_\epsilon|^{\frac{2}{1-\beta}}} - 1 \right) dx \\ &\geq \lim_{\epsilon \rightarrow 0^+} \frac{\pi^2}{2} \epsilon \left(\left(\frac{\epsilon^4}{\epsilon} \right)^{\frac{\alpha}{\alpha_\beta}} - 1 \right) = +\infty. \end{aligned}$$

It follows that

$$\sup_{u \in X_\beta, \|u\|_\beta \leq 1} \int_{\mathbb{R}^4} \left(e^{\alpha|u|^{\frac{2}{1-\beta}}} - 1 \right) dx = +\infty.$$

This ends the proof of Theorem 1.1.

4 Proof of Theorem 1.2

In this section, we prove the improvement of the Adams' inequality using the inequality we have proved in the previous section, that is the proof of Theorem 1.2.

Let $\{u_n\} \subset X_\beta$ and $u \in X_\beta \setminus \{0\}$ be such that $\|u_n\|_\beta = 1$ and $u_n \rightharpoonup u$ weakly in X_β . For $a, b \in \mathbb{R}$, $q > 1$. If q' its conjugate i.e. $\frac{1}{q} + \frac{1}{q'} = 1$ and by Young inequality, we have

$$e^{a+b} \leq \frac{1}{q} e^{qa} + \frac{1}{q'} e^{q'b}.$$

Also we have

$$\begin{aligned} |u_n|^{\frac{2}{1-\beta}} &= |u_n - u + u|^{\frac{2}{1-\beta}} \leq (|u_n - u| + |u|)^{\frac{2}{1-\beta}} \\ &\leq (1 + \epsilon)|u_n - u|^{\frac{2}{1-\beta}} + \left(1 - \frac{1}{(1 + \epsilon)^{\frac{1-\beta}{1+\beta}}}\right)^{\frac{\beta+1}{\beta-1}} |u|^{\frac{2}{1-\beta}}, \end{aligned}$$

which implies that

$$\begin{aligned} \int_{\mathbb{R}^4} \left(e^{p\alpha_\beta |u_n|^{\frac{2}{1-\beta}}} - 1 \right) dx &\leq \frac{1}{q} \int_{\mathbb{R}^4} \left(e^{(1+\epsilon)pq\alpha_\beta |u_n - u|^{\frac{2}{1-\beta}}} - 1 \right) dx \\ &\quad + \frac{1}{q'} \int_{\mathbb{R}^4} \left(e^{pq'\alpha_\beta \left(1 - \frac{1}{(1+\epsilon)^{\frac{1-\beta}{1+\beta}}}\right)^{\frac{\beta+1}{\beta-1}} |u|^{\frac{2}{1-\beta}}} - 1 \right) dx. \end{aligned} \quad (4.1)$$

By (1.12), we have

$$\begin{aligned} &\int_{\mathbb{R}^4} \left(e^{pq'\alpha_\beta \left(1 - \frac{1}{(1+\epsilon)^{\frac{1-\beta}{1+\beta}}}\right)^{\frac{\beta+1}{\beta-1}} |u|^{\frac{2}{1-\beta}}} - 1 \right) dx \\ &\leq \int_{\mathbb{R}^4} \left(e^{\left| \left(pq'\alpha_\beta \left(1 - \frac{1}{(1+\epsilon)^{\frac{1-\beta}{1+\beta}}}\right)^{\frac{\beta+1}{\beta-1}} \right)^{\frac{1-\beta}{2}} |u|^{\frac{2}{1-\beta}} \right|} - 1 \right) dx < +\infty. \end{aligned} \quad (4.2)$$

To complete the proof, we have to prove that for every p such that $0 < p < P_\beta(u)$,

$$\sup_n \int_{\mathbb{R}^4} \left(e^{(1+\epsilon)pq\alpha_\beta |u_n - u|^{\frac{2}{1-\beta}}} - 1 \right) dx < +\infty, \quad (4.3)$$

for some $q > 1$ and $\epsilon > 0$.

If $\|u\|_\beta < 1$ and $0 < p < P_\beta(u) = \frac{1}{(1 - \|u\|_\beta^2)^{\frac{1}{1-\beta}}}$, there exists $\nu > 0$ such that

$$p(1 - \|u\|_\beta^2)^{\frac{1}{1-\beta}}(1 + \nu) < 1.$$

Moreover, by Brezis-Lieb Lemma [8], we can easily get

$$\|u_n - u\|_\beta^2 = \|u_n\|_\beta^2 - \|u\|_\beta^2 + o(1) = 1 - \|u\|_\beta^2 + o(1),$$

where $o(1) \rightarrow 0$ as $n \rightarrow +\infty$.

Then

$$\lim_{n \rightarrow +\infty} \|u_n - u\|_\beta^{\frac{2}{1-\beta}} = \left(\lim_{n \rightarrow +\infty} \|u_n - u\|_\beta^2 \right)^{\frac{1}{1-\beta}} = (1 - \|u\|_\beta^2)^{\frac{1}{1-\beta}}.$$

So for all $\epsilon > 0$, there exists $N_\epsilon \geq 1$ such that

$$\|u_n - u\|_\beta^{\frac{2}{1-\beta}} \leq (1 - \|u\|_\beta^2)^{\frac{1}{1-\beta}}(1 + \epsilon), \quad \text{for all } n \geq N_\epsilon.$$

Choose $q = 1 + \epsilon$, $\epsilon = \sqrt[3]{1 + \nu} - 1$, one has

$$pq(1 + \epsilon)\|u_n - u\|_\beta^{\frac{2}{1-\beta}} \leq p(1 + \nu) (1 - \|u\|_\beta^2)^{\frac{1}{1-\beta}} < 1.$$

Consequently, using (1.13), we get

$$\begin{aligned} \int_{\mathbb{R}^4} \left(e^{(1+\epsilon)pq\alpha_\beta|u_n-u|^{\frac{2}{1-\beta}}} - 1 \right) dx &= \int_{\mathbb{R}^4} \left(e^{pq(1+\epsilon)\|u_n-u\|_\beta^{\frac{2}{1-\beta}} \alpha_\beta \left(\frac{|u_n-u|}{\|u_n-u\|_\beta} \right)^{\frac{2}{1-\beta}}} - 1 \right) dx \\ &< +\infty. \end{aligned} \quad (4.4)$$

If $\|u\|_\beta = 1$ and $0 < p < P_\beta(u) = +\infty$, we also have

$$\lim_{n \rightarrow +\infty} \|u_n - u\|_\beta^{\frac{2}{1-\beta}} = \left(\lim_{n \rightarrow +\infty} \|u_n - u\|_\beta^2 \right)^{\frac{1}{1-\beta}} = 0.$$

So for all $\epsilon > 0$, there exists $N_\epsilon \geq 1$ such that

$$\|u_n - u\|_\beta^{\frac{2}{1-\beta}} < \frac{1}{p(1 + \epsilon)^2}, \quad \text{for all } n \geq N_\epsilon.$$

Choose $q = 1 + \epsilon$, one has

$$pq(1 + \epsilon)\|u_n - u\|_\beta^{\frac{2}{1-\beta}} < p(1 + \epsilon)^2 \cdot \frac{1}{p(1 + \epsilon)^2} = 1.$$

Consequently, using (1.13), we also get

$$\begin{aligned} \int_{\mathbb{R}^4} \left(e^{(1+\epsilon)pq\alpha_\beta|u_n-u|^{\frac{2}{1-\beta}}} - 1 \right) dx &= \int_{\mathbb{R}^4} \left(e^{pq(1+\epsilon)\|u_n-u\|_\beta^{\frac{2}{1-\beta}} \alpha_\beta \left(\frac{|u_n-u|}{\|u_n-u\|_\beta} \right)^{\frac{2}{1-\beta}}} - 1 \right) dx \\ &< +\infty. \end{aligned} \quad (4.5)$$

Now (4.3) follows from (4.4) and (4.5).

Combining (4.1), (4.2) and (4.3), the proof is complete.

5 Application

In this section, we introduce an application of the previous inequalities and give the details of the proof of Theorem 1.3.

Notice that problem (1.16) has a variational structure, so we use the variational methods to deal with the problem. The energy function I_β associated with problem (1.16) is given by

$$I_\beta(v) = \frac{1}{2}\|v\|_\beta^2 - \int_{\mathbb{R}^4} F(v)dx, \quad \text{for all } v \in X_\beta.$$

We can easily see that I_β is well defined and $I_\beta \in C^1(X_\beta, \mathbb{R})$.

A weak solution of problem (1.16) is a function $u \in X_\beta$ which satisfies

$$\int_{\mathbb{R}^4} \omega_\beta(x) \Delta u \Delta v dx = \int_{\mathbb{R}^4} f(u) v dx, \quad \text{for all } v \in X_\beta.$$

It is well-known that finding a weak solution of problem (1.16) is equivalent to finding a critical point of I_β .

First, we give the embedding properties which will be useful in the rest of our work.

Proposition 5.1. *Let $0 < \beta < 1$. Assume that (D_1) and (D_2) holds. Then for all $q \geq \frac{2}{1-\beta}$, $X_\beta \hookrightarrow L^q(\mathbb{R}^4)$ with continuous embedding. Furthermore, for all $q > \frac{2}{1-\beta}$, the embedding $X_\beta \hookrightarrow L^q(\mathbb{R}^4)$ is compact.*

Proof. Let $u \in X_\beta$, by Lemma 2.1, we have

$$|u(e_1)| \leq \frac{\sqrt{2}c_1}{4\pi} \|u\|_\beta \left(\int_1^{+\infty} \chi(y)^{-1} y^{-1} dy \right)^{\frac{1}{2}} + c_2.$$

Set

$$v(x) = \begin{cases} u(x) - u(e_1), & 0 \leq |x| < 1, \\ 0, & |x| \geq 1, \end{cases}$$

where $e_1 = (1, 0, 0, 0) \in \mathbb{R}^4$. Clearly $v \in W_{0,rad}^{2,2}(B, \omega)$.

From [44, Lemma 2.1], we have

$$|v(x)| \leq \frac{1}{2\sqrt{2}\pi} \left(\frac{\left(\log \frac{e}{|x|} \right)^{1-\beta} - 1}{1-\beta} \right)^{\frac{1}{2}} \|u\|_\beta.$$

Then

$$\begin{aligned} |u(x)| &\leq |v(x)| + |u(e_1)| \\ &\leq \frac{1}{2\sqrt{2}\pi} \left(\frac{\left(\log \frac{e}{|x|} \right)^{1-\beta} - 1}{1-\beta} \right)^{\frac{1}{2}} \|u\|_\beta + \frac{\sqrt{2}c_1}{4\pi} \|u\|_\beta \left(\int_1^{+\infty} \chi(y)^{-1} y^{-1} dy \right)^{\frac{1}{2}} + c_2, \quad \text{for } |x| < 1. \end{aligned}$$

By (1.6), it is easy to infer that

$$\int_{|x|<1} |u(x)|^{\frac{2k}{1-\beta}} dx \leq c_3 \|u\|_\beta^{\frac{2k}{1-\beta}} + c_4 \leq (c_3 + c_5) \|u\|_\beta^{\frac{2k}{1-\beta}}. \quad (5.1)$$

On the other hand, we can infer from (3.3) that

$$\int_{|x|\geq 1} |u|^{\frac{2k}{1-\beta}} dx \leq c_6 \|u\|_\beta^{\frac{2k}{1-\beta}} + c_7 \leq (c_6 + c_8) \|u\|_\beta^{\frac{2k}{1-\beta}}. \quad (5.2)$$

Combining (5.1) and (5.2), one has

$$\int_{\mathbb{R}^4} |u|^{\frac{2k}{1-\beta}} dx \leq c_9 \|u\|_\beta^{\frac{2k}{1-\beta}}.$$

Then we deduce that the continuous embedding $X_\beta \hookrightarrow L^{\frac{2k}{1-\beta}}(\mathbb{R}^4)$ holds for all $k \in N^*$. By simple interpolation, it follows that X_β is continuously embedded into $L^q(\mathbb{R}^4)$, for all $q \geq \frac{2}{1-\beta}$. Proceeding as the proof of [6, proposition 2], we deduce that the embedding $X_\beta \hookrightarrow L^q(\mathbb{R}^4)$ is compact for all $q > \frac{2}{1-\beta}$. \square

Now we consider problem (1.16) and verify that the corresponding energy function I_β has the Mountain Pass geometry structure.

Lemma 5.1. *Assume that $(F_1) - (F_3)$ hold. Then*

(i) *There exist $\rho_0, \tau > 0$ such that $I_\beta(u) \geq \tau$, for $u \in X_\beta$, $\|u\|_\beta = \rho_0$.*

(ii) *There exists $\|e_0\|_\beta > \rho_0$ such that $I_\beta(e_0) < 0$.*

Proof. (i) Given $u \in X_\beta$, by (F_1) , we get

$$\begin{aligned} \int_{\mathbb{R}^4} |F(u)| dx &\leq c_0 \left(\int_{\mathbb{R}^4} |u|^p dx + \int_{\mathbb{R}^4} |u|^q \left(e^{\alpha|u|^{\frac{2}{1-\beta}}} - 1 \right) dx \right) \\ &\leq c_0 \left(\|u\|_{L^p(\mathbb{R}^4)}^p + \|u\|_{L^{q+1}(\mathbb{R}^4)}^q \left(\int_{\mathbb{R}^4} \left(e^{\alpha(q+1)|u|^{\frac{2}{1-\beta}}} - 1 \right) dx \right)^{\frac{1}{q+1}} \right). \end{aligned} \quad (5.3)$$

For $u \in X_\beta$ such that

$$\|u\|_\beta = \rho_0 < \left(\frac{\alpha_\beta}{\alpha(1+q)} \right)^{\frac{1-\beta}{2}}.$$

By (1.13), there exists a constant $C_3 > 0$ such that

$$\int_{\mathbb{R}^4} \left(e^{\alpha(q+1)|u|^{\frac{2}{1-\beta}}} - 1 \right) dx = \int_{\mathbb{R}^4} \left(e^{\alpha(q+1)\|u\|_\beta^{\frac{2}{1-\beta}} \left| \frac{u}{\|u\|_\beta} \right|^{\frac{2}{1-\beta}}} - 1 \right) dx \leq C_3.$$

Since $X_\beta \hookrightarrow L^p(\mathbb{R}^4)$, $X_\beta \hookrightarrow L^{q+1}(\mathbb{R}^4)$ and combine with (5.3), we have

$$\int_{\mathbb{R}^4} |F(u)| dx \leq C_4 \left(\|u\|_\beta^p + \|u\|_\beta^q \right)$$

where $C_4 > 0$. Then

$$I_\beta(u) = \frac{1}{2} \|u\|_\beta^2 - \int_{\mathbb{R}^4} F(u) dx \geq \frac{1}{2} \|u\|_\beta^2 - C_4 \left(\|u\|_\beta^p + \|u\|_\beta^q \right), \quad \text{for } u \in X_\beta, \|u\|_\beta = \rho_0.$$

One can choose $\rho_0 > 0$ small enough such that

$$\rho_0 < \min \left\{ \left(\frac{\alpha_\beta}{\alpha(1+q)} \right)^{\frac{1-\beta}{2}}, \left(\frac{1}{4C_4} \right)^{\frac{1}{\min\{p,q\}-2}} \right\}.$$

Then we deduce that

$$I_\beta(u) \geq \rho_0^2 \left(\frac{1}{2} - 2C_4 \rho_0^{\min\{p,q\}-2} \right) = \tau > 0, \quad \text{for } u \in X_\beta, \|u\|_\beta = \rho_0.$$

(ii) From (F_3) , we have

$$\lim_{s \rightarrow \infty} \frac{f(s)s}{e^{\alpha s} s^{\frac{2}{1-\beta}}} = \lim_{s \rightarrow \infty} \frac{f(s)s}{s^{\frac{2}{1-\beta}}} \cdot \frac{s^{\frac{2}{1-\beta}}}{e^{\alpha s} s^{\frac{2}{1-\beta}}} \geq \kappa > 0.$$

Since $\lim_{s \rightarrow \infty} \frac{s^{\frac{2}{1-\beta}}}{e^{\alpha s} s^{\frac{2}{1-\beta}}} = 0$, we infer

$$\lim_{s \rightarrow \infty} \frac{f(s)}{s^{\frac{2}{1-\beta}-1}} = \infty.$$

Then, there exist $s_1 > 0$ and $M > 0$ such that for any $s \geq s_1$,

$$f(s) \geq M s^{\frac{2}{1-\beta}-1}.$$

It follows that

$$F(s) \geq \frac{1-\beta}{2} M s^{\frac{2}{1-\beta}}.$$

By (F_1) , we have

$$\begin{aligned} I_\beta(tu) &= \frac{1}{2} t^2 \|u\|_\beta^2 - \int_{\mathbb{R}^4} F(tu) dx \\ &\leq \frac{1}{2} t^2 \|u\|_\beta^2 - \int_{|tu| \geq s_1} F(tu) dx \\ &\leq \frac{1}{2} t^2 \|u\|_\beta^2 - \frac{1-\beta}{2} M t^{\frac{2}{1-\beta}} \int_{|tu| \geq s_1} u^{\frac{2}{1-\beta}} dx. \end{aligned}$$

Then, we obtain

$$I_\beta(tu) \rightarrow -\infty, \quad \text{when } t \rightarrow +\infty.$$

Thus, there exists $e_0 \in X_\beta$, $\|e_0\|_\beta > \rho_0$ such that $I_\beta(e_0) < 0$. \square

By Lemma 5.1 and the Mountain Pass Theorem, we know that there exists a $(PS)_m$ sequence of the functional I_β at the mountain pass level

$$m = \inf_{\gamma \in \Gamma} \sup_{0 \leq t \leq 1} I_\beta(\gamma(t)),$$

where

$$\Gamma = \{\gamma \in C([0, 1], X_\beta), \gamma(0) = 0, I_\beta(\gamma(1)) < 0\}.$$

Next, we establish an important estimate involving the level m .

Lemma 5.2. *Assume that $(F_1) - (F_3)$ hold. Then there holds*

$$0 < m < m^* \triangleq \frac{1}{2} \left(\frac{\alpha_\beta}{\alpha} \right)^{1-\beta}.$$

Proof. By the previous lemma, clearly, $m \geq \rho_1 > 0$.

Next, we consider the family of Adams type function

$$\omega_n(x) = \begin{cases} \left(\frac{\log(e^4 n)}{\alpha_\beta} \right)^{\frac{1-\beta}{2}} - \frac{|x|^{2(1-\beta)}}{2 \left(\frac{\alpha_\beta}{4n} \right)^{\frac{1-\beta}{2}} \left(\frac{1}{4} \log(e^4 n) \right)^{\frac{1+\beta}{2}}} + \frac{1}{2 \left(\frac{\alpha_\beta}{4} \right)^{\frac{1-\beta}{2}} \left(\frac{1}{4} \log(e^4 n) \right)^{\frac{1+\beta}{2}}}, & 0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}, \\ \frac{\left(\log \frac{e}{|x|} \right)^{1-\beta}}{\left(\frac{\alpha_\beta}{16} \log(e^4 n) \right)^{\frac{1-\beta}{2}}}, & \frac{1}{\sqrt[4]{n}} < |x| \leq \frac{1}{2}, \\ \eta_n, & |x| > \frac{1}{2}, \end{cases}$$

where $\eta_n \in C_0^\infty(B_1)$ is such that

$$\eta_n|_{x=\frac{1}{2}} = \frac{1}{\left(\frac{\alpha_\beta}{16} \log(e^4 n) \right)^{\frac{1-\beta}{2}}} (\log 2e)^{1-\beta}, \quad \frac{\partial \eta_n}{\partial x}|_{x=\frac{1}{2}} = -\frac{2(1-\beta)}{\left(\frac{\alpha_\beta}{16} \log(e^4 n) \right)^{\frac{1-\beta}{2}}} (\log 2e)^{-\beta},$$

and $\eta_n, \nabla \eta_n, \Delta \eta_n$ are all $O\left(\frac{1}{(\log(e^4 n))^{\frac{1-\beta}{2}}}\right)$ as $n \rightarrow \infty$.

As in the proof of Section 3, after a series of computation, we get

$$\|\omega_n\|_\beta \rightarrow 1, \quad \text{as } n \rightarrow \infty.$$

Let

$$v_n = \frac{\omega_n}{\|\omega_n\|_\beta}.$$

We also have

$$\|v_n\|_\beta = 1,$$

and

$$v_n(x) \geq \left(\frac{\log(e^4 n)}{\alpha_\beta} \right)^{\frac{1-\beta}{2}} + o(1), \quad \text{for all } x \in \mathbb{R}^4, \quad 0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}. \quad (5.4)$$

In order to prove $m < m^*$, we only need to prove that

$$\max_{t \geq 0} I_\beta(tv_n) < m^*.$$

By contradiction, we suppose that

$$\max_{t \geq 0} I_\beta(tv_n) \geq m^*.$$

Then, there exists $t_n > 0$ such that

$$\max_{t \geq 0} I_\beta(tv_n) = I_\beta(t_n v_n) \geq m^*,$$

that is

$$\frac{1}{2} t_n^2 \|v_n\|_\beta^2 - \int_{\mathbb{R}^4} F(t_n v_n) dx \geq \frac{1}{2} \left(\frac{\alpha_\beta}{\alpha} \right)^{1-\beta}.$$

Combining with (F_1) , we get

$$t_n^2 \geq \left(\frac{\alpha_\beta}{\alpha} \right)^{1-\beta}. \quad (5.5)$$

Moreover, we have

$$\frac{d}{dt}I_\beta(tv_n)|_{t=t_n} = t_n - \int_{\mathbb{R}^4} f(t_nv_n)v_n dx = 0,$$

that is

$$t_n^2 = \int_{\mathbb{R}^4} f(t_nv_n)t_nv_n dx. \quad (5.6)$$

Now we claim that $\{t_n\}$ is bounded.

In fact, it follows from (F_3) that for any $\epsilon > 0$, there exists $s(\epsilon)$ such that

$$f(s)s \geq (\kappa - \epsilon)e^{\alpha s^{\frac{2}{1-\beta}}}, \quad \text{for all } s \geq s(\epsilon). \quad (5.7)$$

Notice that $t_nv_n \geq s(\epsilon)$ for $0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}$ and for large $n \in \mathbb{N}$. Then it follows from (5.4), (5.6) and (5.7) that for large n ,

$$\begin{aligned} t_n^2 &= \int_{\mathbb{R}^4} f(t_nv_n)t_nv_n dx \\ &\geq \int_{0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}} f(t_nv_n)t_nv_n dx \\ &\geq (\kappa - \epsilon) \int_{0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}} e^{\alpha t_n^{\frac{2}{1-\beta}} v_n^{\frac{2}{1-\beta}}} dx \\ &\geq 2\pi^2(\kappa - \epsilon) \int_0^{\frac{1}{\sqrt[4]{n}}} r^3 e^{\alpha t_n^{\frac{2}{1-\beta}} \frac{\log(e^4 n)}{\alpha\beta}} dr, \end{aligned} \quad (5.8)$$

that is

$$1 \geq 2\pi^2(\kappa - \epsilon) e^{\alpha \frac{\log(e^4 n)}{\alpha\beta} t_n^{\frac{2}{1-\beta}} - \log(4n) - \log t_n^2}.$$

Therefore, $\{t_n\}$ is bounded.

From (5.5), we have

$$\lim_{n \rightarrow +\infty} t_n^2 \geq \left(\frac{\alpha\beta}{\alpha}\right)^{1-\beta}.$$

In fact, we can prove that

$$\lim_{n \rightarrow +\infty} t_n^2 = \left(\frac{\alpha\beta}{\alpha}\right)^{1-\beta}. \quad (5.9)$$

Indeed, if $\lim_{n \rightarrow +\infty} t_n^2 > \left(\frac{\alpha\beta}{\alpha}\right)^{1-\beta}$, it follows that for large n there exists $\delta > 0$ such that

$$t_n^{\frac{2}{1-\beta}} \geq \frac{\alpha\beta}{\alpha} + \delta. \quad (5.10)$$

Combining (5.8) and (5.10), we infer $t_n \rightarrow \infty$, as $n \rightarrow \infty$ which contradicts the boundedness of $\{t_n\}$. Therefore, (5.9) follows immediately.

Combining (5.8) and (5.9), we have

$$\begin{aligned}
\lim_{n \rightarrow \infty} t_n^2 &= \left(\frac{\alpha_\beta}{\alpha} \right)^{1-\beta} \geq (\kappa - \epsilon) \lim_{n \rightarrow \infty} \int_{0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}} e^{\alpha t_n^{\frac{2}{1-\beta}} v_n^{\frac{2}{1-\beta}}} dx \\
&\geq 2\pi^2(\kappa - \epsilon) \lim_{n \rightarrow \infty} \int_0^{\frac{1}{\sqrt[4]{n}}} r^3 e^{\log(e^4 n)} dr \\
&= \frac{\pi^2}{2}(\kappa - \epsilon)e^4.
\end{aligned}$$

It follows that

$$\kappa \leq \frac{2}{\pi^2 e^4} \left(\frac{\alpha_\beta}{\alpha} \right)^{1-\beta},$$

which contradicts with (F_3) . \square

Lemma 5.3. *Assume that $(F_1) - (F_3)$ hold. Let $\{u_n\}$ be a $(PS)_m$ sequence of I_β with $m \in (0, m^*)$. Then $\{u_n\}$ is bounded in X_β .*

Proof. Let $\{u_n\} \subset X_\beta$ be a $(PS)_m$ sequence of I_β with $m \in (0, m^*)$. It follows that

$$I_\beta(u_n) = \frac{1}{2} \|u_n\|_\beta^2 - \int_{\mathbb{R}^4} F(u_n) dx \rightarrow m, \quad n \rightarrow +\infty, \quad (5.11)$$

and

$$|I'_\beta(u_n)u_n| = \left| \|u_n\|_\beta^2 - \int_{\mathbb{R}^4} f(u_n)u_n dx \right| \leq \epsilon_n \|u_n\|_\beta, \quad (5.12)$$

where $\epsilon_n \rightarrow 0$, as $n \rightarrow +\infty$.

From (F_2) , we have for any $\epsilon > 0$ there exists $s_\epsilon > 0$ such that

$$F(s) \leq \epsilon s f(s), \quad \text{for all } s \geq s_\epsilon. \quad (5.13)$$

Combining (5.11), (5.12) and (5.13), for fix $0 < \epsilon < \frac{1}{2}$, we have

$$\begin{aligned}
\frac{1}{2} \|u_n\|_\beta^2 &\leq C + \int_{\mathbb{R}^4} F(u_n) dx \\
&\leq C + \int_{|u_n| \leq s_\epsilon} F(u_n) dx + \int_{|u_n| \geq s_\epsilon} F(u_n) dx \\
&\leq C' + \epsilon \int_{\mathbb{R}^4} f(u_n)u_n dx \\
&\leq C' + \epsilon \|u_n\|_\beta^2 + \epsilon \epsilon_n \|u_n\|_\beta.
\end{aligned}$$

Therefore, we can deduce that $\{u_n\}$ is bounded in X_β . \square

Now we turn to the compactness properties of $(PS)_m$ sequence.

Lemma 5.4. *Assume that $(F_1) - (F_3)$ hold. Let $\{u_n\}$ be a $(PS)_m$ sequence of I_β such that*

$$\liminf_{n \rightarrow +\infty} \|u_n\|_\beta < \left(\frac{\alpha_\beta}{\alpha} \right)^{\frac{1-\beta}{2}}, \quad (5.14)$$

then $\{u_n\}$ has a convergence subsequence in X_β .

Proof. By Lemma 5.3, up to a subsequence, one has

$$u_n \rightharpoonup u \text{ weakly in } X_\beta \text{ and } u_n(x) \rightarrow u(x) \text{ a.e. } x \in \mathbb{R}^4.$$

By (5.14), up to a subsequence, there exist $r_1 > 1$ and $0 < \gamma_0 < 1$ such that

$$\|u_n\|_\beta < \left(\frac{\gamma_0 \alpha \beta}{r_1 \alpha} \right)^{\frac{1-\beta}{2}}. \quad (5.15)$$

By (F_1) , we get

$$\left| \int_{\mathbb{R}^4} f(u_n)(u_n - u) dx \right| \leq c_1 \int_{\mathbb{R}^4} |u_n|^{p-1} |u_n - u| dx + c_1 \int_{\mathbb{R}^4} |u_n|^{q-1} \left(e^{\alpha|u_n|^{\frac{2}{1-\beta}}} - 1 \right) |u_n - u| dx. \quad (5.16)$$

Using Hölder inequality, we have

$$\begin{aligned} \int_{\mathbb{R}^4} |u_n|^{p-1} |u_n - u| dx &\leq \left(\int_{\mathbb{R}^4} |u_n|^p \right)^{\frac{p-1}{p}} \left(\int_{\mathbb{R}^4} |u_n - u|^p \right)^{\frac{1}{p}} \\ &= \|u_n\|_{L^p(\mathbb{R}^4)}^{p-1} \|u_n - u\|_{L^p(\mathbb{R}^4)}. \end{aligned} \quad (5.17)$$

Let $r_2 > 1$ and $r_3 > 1$ be such that

$$\begin{cases} \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} = 1, \\ r_2 > \frac{r_1}{r_1 - 1}, \\ r_2 \geq \frac{2}{(1-\beta)(q-1)}, \\ r_3 > \frac{2}{1-\beta}. \end{cases}$$

From (1.13), (5.15) and Hölder inequality, we have

$$\begin{aligned} &\int_{\mathbb{R}^4} |u_n|^{q-1} \left(e^{\alpha|u_n|^{\frac{2}{1-\beta}}} - 1 \right) |u_n - u| dx \\ &\leq \|u_n\|_{L^{r_2(q-1)}(\mathbb{R}^4)}^{q-1} \|u_n - u\|_{L^{r_3}(\mathbb{R}^4)} \left(\int_{\mathbb{R}^4} \left(e^{r_1 \alpha |u_n|^{\frac{2}{1-\beta}}} - 1 \right) dx \right)^{\frac{1}{r_1}} \\ &\leq \|u_n\|_{L^{r_2(q-1)}(\mathbb{R}^4)}^{q-1} \|u_n - u\|_{L^{r_3}(\mathbb{R}^4)} \left(\int_{\mathbb{R}^4} \left(e^{r_1 \alpha \|u_n\|_\beta^{\frac{2}{1-\beta}} \left(\frac{u_n}{\|u_n\|_\beta} \right)^{\frac{2}{1-\beta}}} - 1 \right) dx \right)^{\frac{1}{r_1}} \\ &\leq \|u_n\|_{L^{r_2(q-1)}(\mathbb{R}^4)}^{q-1} \|u_n - u\|_{L^{r_3}(\mathbb{R}^4)} \left(\int_{\mathbb{R}^4} \left(e^{\gamma_0 \alpha \beta \left(\frac{u_n}{\|u_n\|_\beta} \right)^{\frac{2}{1-\beta}}} - 1 \right) dx \right)^{\frac{1}{r_1}} \\ &\leq c_2 \|u_n\|_{L^{r_2(q-1)}(\mathbb{R}^4)}^{q-1} \|u_n - u\|_{L^{r_3}(\mathbb{R}^4)}. \end{aligned} \quad (5.18)$$

Since the embeddings $X_\beta \hookrightarrow L^p(\mathbb{R}^4)$ and $X_\beta \hookrightarrow L^{r_3}(\mathbb{R}^4)$ are compact, we can immediately deduce from (5.16), (5.17) and (5.18) that

$$\int_{\mathbb{R}^4} f(u_n)(u_n - u)dx \rightarrow 0.$$

On the other hand, since $\langle I'_\beta(u_n), u_n - u \rangle \rightarrow 0$, as $n \rightarrow +\infty$, then we infer

$$\int_{\mathbb{R}^4} \omega_\beta \Delta u_n \Delta (u_n - u) \rightarrow 0, \quad n \rightarrow +\infty.$$

Thus,

$$\|u_n - u\|_\beta^2 = \langle u_n, u_n - u \rangle - \langle u, u_n - u \rangle = \langle u_n, u_n - u \rangle + o_n(1) \rightarrow 0, \quad n \rightarrow +\infty.$$

That is, $u_n \rightarrow u$ strongly in X_β . □

Lemma 5.5. *Assume that $(F_1) - (F_3)$ hold. Then I_β satisfies the $(PS)_m$ condition if $m \in (0, m^*)$.*

Proof. Let $\{u_n\} \subset X_\beta$ be a $(PS)_m$ sequence of I_β with $m \in (0, m^*)$. By Lemma 5.3, up to a subsequence, one has

$$u_n \rightharpoonup u \text{ weakly in } E_\beta \text{ and } u_n(x) \rightarrow u(x) \text{ a.e. } x \in \mathbb{R}^4,$$

and

$$|\langle I'_\beta(u_n), u_n \rangle| = \left| \|u_n\|_\beta^2 - \int_{\mathbb{R}^4} f(u_n)u_n dx \right| \leq C, \quad \text{for all } n.$$

Since u_n is bounded in X_β , then

$$\int_{\mathbb{R}^4} f(u_n)u_n dx \leq C, \quad \text{for all } n.$$

By [18, Lemma 2.1], we get $f(u_n) \rightarrow f(u)$ in $L^1_{loc}(\mathbb{R}^4)$. Then, it follows from (F_2) and Generalized Lebesgue Dominated Convergence Theorem that $F(u_n) \rightarrow F(u)$ in $L^1_{loc}(\mathbb{R}^4)$. Moreover, inspired by [22], we can deduce that

$$F(u_n) \rightarrow F(u) \text{ in } L^1(\mathbb{R}^4) \text{ as } n \rightarrow +\infty.$$

In fact, by (5.13), we have

$$\int_{|u_n| \geq s_\epsilon} |F(u_n)| dx \leq \epsilon \int_{|u_n| \geq s_\epsilon} f(u_n)u_n dx \leq C\epsilon. \quad (5.19)$$

By (F_1) , let $\lambda \triangleq \max\{p, q\}$, we can choose a constant $C_\epsilon > 0$ such that

$$F(s) \leq C_\epsilon |s|^\lambda, \quad \text{for all } s \leq s_\epsilon.$$

Then, for $R > 0$, we have

$$\begin{aligned} \int_{\{|x| > R\} \cap \{|u_n| \leq s_\epsilon\}} F(u_n) dx &\leq C_\epsilon \int_{\{|x| > R\} \cap \{|u_n| \leq s_\epsilon\}} |u_n|^\lambda dx \\ &\leq C_\epsilon \int_{\{|x| > R\} \cap \{|u_n| \leq s_\epsilon\}} |u_n - u|^\lambda dx + C_\epsilon \int_{\{|x| > R\} \cap \{|u_n| \leq s_\epsilon\}} |u|^\lambda dx. \end{aligned}$$

Now using the compactness of embedding $X_\beta \hookrightarrow L^\lambda(\mathbb{R}^4)$, we can choose $R > 0$ such that

$$\int_{\{|x|>R\} \cap \{|u_n| \leq s_\epsilon\}} |F(u_n)| dx \leq C\epsilon. \quad (5.20)$$

Similarly, one has

$$\int_{|u_n| \geq s_\epsilon} |F(u)| dx \leq C\epsilon \quad \text{and} \quad \int_{\{|x|>R\} \cap \{|u_n| \leq s_\epsilon\}} |F(u)| dx \leq C\epsilon. \quad (5.21)$$

Noticing that

$$\int_{\{|x|<R\} \cap \{|u_n| \leq s_\epsilon\}} |F(u_n) - F(u)| \leq \int_{|x|<R} |F(u_n) - F(u)| < \epsilon. \quad (5.22)$$

Combining (5.19), (5.20), (5.21) and (5.22), we have the fact that $F(u_n) \rightarrow F(u)$ in $L^1(\mathbb{R}^4)$. Furthermore, we get $I'_\beta(u) = 0$, namely, u is a weak solution of (1.16).

If $u = 0$, since

$$\int_{\mathbb{R}^4} F(u_n) dx \rightarrow \int_{\mathbb{R}^4} F(0) dx = 0 \quad \text{and} \quad I_\beta(u_n) \rightarrow m,$$

then

$$\|u_n\|_\beta \rightarrow (2m)^{\frac{1}{2}} < \left(\frac{\alpha_\beta}{\alpha}\right)^{\frac{1-\beta}{2}}.$$

By Lemma 5.4, we deduce that $u_n \rightarrow 0$ strongly in X_β which is contradiction with the fact that $m > 0$.

Thus, $u \neq 0$.

Clearly, up to a subsequence, we have

$$\liminf_{n \rightarrow +\infty} \|u_n\|_\beta \geq \|u\|_\beta > 0.$$

Set

$$v_n = \frac{u_n}{\|u_n\|_\beta}, \quad v = \frac{u}{\lim_{n \rightarrow +\infty} \|u_n\|_\beta}.$$

Then $\|v\|_\beta < 1$, $v_n \rightharpoonup v$ weakly in X_β , $\|v_n\|_\beta = 1$.

We have

$$\begin{aligned} \lim_{n \rightarrow +\infty} \|u_n\|_\beta^2 &= 2 \left(m + \int_{\mathbb{R}^4} F(u) dx \right) \\ &= \frac{2 \left(m + \int_{\mathbb{R}^4} F(u) dx \right) (m - I_\beta(u))}{(m - I_\beta(u))} \\ &= 2(m - I_\beta(u)) \frac{m + \int_{\mathbb{R}^4} F(u) dx}{\int_{\mathbb{R}^4} F(u) dx - \frac{1}{2}\|u\|_\beta^2} \\ &= 2(m - I_\beta(u)) \frac{1}{1 - \|v\|_\beta^2} \\ &\leq \frac{2m}{1 - \|v\|_\beta^2} < \frac{\left(\frac{\alpha_\beta}{\alpha}\right)^{1-\beta}}{1 - \|v\|_\beta^2}. \end{aligned}$$

It follows that there exist $r_1 > 1$ and $0 < \gamma_1 < 1$ such that, up to a subsequence,

$$\alpha r_1 \|u_n\|_{\beta}^{\frac{2}{1-\beta}} < \frac{\gamma_1 \alpha_{\beta}}{\left(1 - \|v\|_{\beta}^2\right)^{\frac{1}{1-\beta}}} = \gamma_1 P_{\beta}(v) \alpha_{\beta}, \quad \text{for all } n,$$

where P_{β} is defined in (1.15).

By Theorem 1.2, we get

$$\begin{aligned} \sup_n \int_{\mathbb{R}^4} \left(e^{\alpha r_1 |u_n|^{\frac{2}{1-\beta}}} - 1 \right) dx &= \sup_n \int_{\mathbb{R}^4} \left(e^{\alpha r_1 \|u_n\|^{\frac{2}{1-\beta}} |v_n|^{\frac{2}{1-\beta}}} - 1 \right) dx \\ &\leq \sup_n \int_{\mathbb{R}^4} \left(e^{\gamma_1 P_{\beta}(v) \alpha_{\beta} |v_n|^{\frac{2}{1-\beta}}} - 1 \right) dx < +\infty. \end{aligned}$$

Continuing exactly as in the proof of Lemma 5.4, we can show that $u_n \rightarrow u$ strongly in X_{β} . □

Proof of Theorem 1.3 In view of Lemma 5.1–Lemma 5.5, we can apply the Mountain-Pass Theorem to deduce that I_{β} admits a critical point u such that $I_{\beta}(u) = m > 0$. The nonnegativity of u is immediate. From the proof of Lemma 5.5, we can infer that u is nontrivial.

Acknowledgments

The authors have been supported by National Natural Science Foundation of China 11971392 and Natural Science Foundation of Chongqing, China cstc2021ycjh-bgzxm0115.

References

- [1] D.R. Adams, A sharp inequality of J. Moser for higher order derivatives, *Ann. of Math.(2)*, **128** (1988), no. 2, 385-398.
- [2] S. Aouaoui, A new Trudinger-Moser type inequality and an application to some elliptic equation with doubly exponential nonlinearity in the whole space \mathbb{R}^2 , *Arch. Math. (Basel)*, **114** (2020), no. 2, 199-214.
- [3] S. Aouaoui, A new Singular Trudinger-Moser Type Inequality with Logarithmic Weights and Applications, *Adv. Nonlinear Stud.*, **20** (2020), no. 1, 113-139.
- [4] S. Aouaoui, F.S.B. Albuquerque, Adams' type inequality and application to a quasilinear nonhomogeneous equation with singular and vanishing radial potentials in \mathbb{R}^4 , *Ann. Mat. Pura Appl.(4)*, **198** (2019), no. 4, 1331-1349.
- [5] S. Aouaoui, R. Jlel, On some elliptic equation in the whole Euclidean space \mathbb{R}^2 with nonlinearities having new exponential growth condition, *Commun. Pure Appl. Anal.*, **19** (2020), no. 10, 4771-4796.
- [6] S. Aouaoui, R. Jlel, New weighted sharp Trudinger-Moser inequalities defined on the whole euclidean space \mathbb{R}^N and applications, *Calc. Var. Partial Differential Equations*, **60** (2021), no. 1, Paper No. 50, 44 pp.

- [7] D. Brahim, J. Rached, Forth order weighted elliptic problem under exponential nonlinear growth, <https://www.researchgate.net/publication/358144431> (2022).
- [8] H. Brézis, E. Lieb, A relation between pointwise convergence of functions and convergence of functionals, *Proc. Amer. Math. Soc.*, **88** (1983), no. 3, 486-490.
- [9] M. Calanchi, B. Ruf, On Trudinger-Moser type inequalities with logarithmic weights, *J. Differential Equations*, **258** (2015), no. 6, 1967-1989.
- [10] M. Calanchi, B. Ruf, Trudinger-Moser type inequalities with logarithmic weights in dimension N, *Nonlinear Anal.*, **121** (2015), 403-411.
- [11] D.M. Cao, Nontrivial solution of semilinear elliptic equation with critical exponent in \mathbb{R}^2 , *Comm. Partial Differential Equations*, **17** (1992), no. 3-4, 407-435.
- [12] L. Carleson, S.-Y.A. Chang, On the existence of an extremal function for an inequality of J. Moser, *Bull. Sci. Math.(2)*, **110** (1986), no. 2, 113-127.
- [13] A.C. Cavalheiro, Weighted Sobolev Spaces and Degenerate Elliptic Equations, *Bol. Soc. Paran. Mat.(3)*, **26** (2008), no. 1-2, 117-132.
- [14] L. Chen, G.Z. Lu, M.C. Zhu, Existence and nonexistence of extremals for critical Adams inequalities in \mathbb{R}^4 and Trudinger-Moser inequalities in \mathbb{R}^2 , *Adv. Math.*, **368** (2020), 107143, 61 pp.
- [15] A. Cianchi, Moser-Trudinger inequalities without boundary conditions and isoperimetric problems, *Indiana Univ. Math. J.*, **54** (2005), no. 3, 669-705.
- [16] A. Cianchi, Moser-Trudinger trace inequalities, *Adv. Math.*, **217** (2008), no. 5, 2005-2044.
- [17] W.S. Cohn, G.Z. Lu, Best constants for Moser-Trudinger inequalities on the Heisenberg group, *Indiana Univ. Math. J.*, **50** (2001), no. 4, 1567-1591.
- [18] D.G. de Figueiredo, O.H. Miyagaki, B. Ruf, Elliptic equations in \mathbb{R}^2 with nonlinearities in the critical growth range, *Calc. Var. Partial Differential Equations*, **3** (1995), no. 2, 139-153.
- [19] M. Flucher, Extremal functions for Trudinger-Moser inequality in 2 dimensions, *Comment. Math. Helv.*, **67** (1992), no. 3, 471-497.
- [20] L. Fontana, Sharp borderline Sobolev inequalities on compact Riemannian manifolds, *Comment. Math. Helv.*, **68** (1993), no. 3, 415-454.
- [21] T. Kilpeläinen, Weighted Sobolev spaces and capacity, *Ann. Acad. Sci. Fenn. Ser. A I Math.*, **19** (1994), no. 1, 95-113.
- [22] N. Lam, G.Z. Lu, Existence and multiplicity of solutions to equations of N-Laplacian type with critical exponential growth in \mathbb{R}^N , *J. Funct. Anal.*, **262** (2012), no. 3, 1132-1165.
- [23] N. Lam, G.Z. Lu, Sharp Moser-Trudinger inequality on the Heisenberg group at the critical case and applications, *Adv. Math.*, **231** (2012), no. 6, 3259-3287.
- [24] N. Lam, G.Z. Lu, A new approach to sharp Moser-Trudinger and Adams type inequalities: a rearrangement-free argument, *J. Differential Equations*, **255** (2013), no. 3, 298-325.

- [25] Y.X. Li, Moser-Trudinger inequality on compact Riemannian manifolds of dimension two, *J. Partial Differential Equations*, **14** (2001), no. 2, 163-192.
- [26] Y.X. Li, B. Ruf, A sharp Trudinger-Moser type inequality for unbounded domains in \mathbb{R}^n , *Indiana Univ. Math. J.*, **57** (2008), no. 1, 451-480.
- [27] K.-C. Lin, Extremal functions for Moser's inequality, *Trans. Amer. Math. Soc.*, **348** (1996), no. 7, 2663-2671.
- [28] P.L. Lions, The concentration-compactness principle in the Calculus of variations. The limit case. I., *Rev. Mat. Iberoamericana*, **1** (1985), no. 1, 145-201.
- [29] G.Z. Lu, M.C. Zhu, A sharp Trudinger-Moser type inequality involving L^n norm in the entire space \mathbb{R}^n , *J. Differential Equations*, **267** (2019), no. 5, 3046-3082.
- [30] J. Moser, A sharp form of an inequality by N. Trudinger, *Indiana Univ. Math. J.*, **20** (1970/71), 1077-1092.
- [31] E. Nakai, N. Tomita, K. Yabuta, Density of the set of all infinitely differentiable functions with compact support in weighted Sobolev spaces, *Sci. Math. Jpn.*, **60** (2004), no. 1, 121-127.
- [32] V.H. Nguyen, Remarks on the Moser-Trudinger type inequality with logarithmic weights in dimension N , *Proc. Am. Math. Soc.*, **147** (2019), no. 12, 5183-5193.
- [33] J. Peetre, Espaces d'interpolation et théorème de Soboleff (French), *Ann. Inst. Fourier (Grenoble)*, **16** (1966), no. fasc. 1, 279-317.
- [34] S. I. Pohozaev, The Sobolev embedding in the case $pl = n$, Proc. of the Technical Scientific Conference on Advances of Scientific Research 1964-1965, Mathematics Section, Energet. Inst., Moscow, Moskov, 1965, pp. 158-170.
- [35] P. Roy, Extremal function for Moser-Trudinger type inequality with logarithmic weight, *Nonlinear Anal.*, **135** (2016), 194-204.
- [36] P. Roy, On attainability of Moser Trudinger inequality with logarithmic weights in higher dimensions, *Discrete Contin. Dyn. Syst.*, **39** (2019), no. 9, 5207-5222.
- [37] B. Ruf, A sharp Trudinger-Moser type inequality for unbounded domains in \mathbb{R}^2 , *J. Funct. Anal.*, **219** (2005), no. 2, 340-367.
- [38] B. Ruf, F. Sani, Sharp Adams-type inequalities in \mathbb{R}^n , *Trans. Amer. Math. Soc.*, **365** (2013), no. 2, 645-670.
- [39] C. Tarsi, Adams' inequality and limiting Sobolev embeddings into Zygmund spaces, *Potential Anal.*, **37** (2012), no. 4, 353-385.
- [40] N. S. Trudinger, On imbeddings into Orlicz spaces and some applications, *J. Math. Mech.*, **17** (1967), 473-483.
- [41] Y.Y. Yang, A sharp form of the Moser-Trudinger inequality on a compact Riemannian surface, *Trans. Amer. Math. Soc.*, **359** (2007), no. 12, 5761-5776.

- [42] Y.Y. Yang, Adams type inequalities and related elliptic partial differential equations in dimension four, *J. Differential Equations*, **252** (2012), no. 3, 2266-2295.
- [43] V.I. Yudovich, Some estimates connected with integral operators and with solutions of elliptic equations (Russian), *Dokl. Akad. Nauk SSSR*, **138** (1961), 805-808.
- [44] M.C. Zhu, L.F. Wang, Adams' inequality with logarithmic weights in \mathbb{R}^4 , *Proc. Amer. Math. Soc.*, **149** (2021), no. 8, 3463-3472.