

SIGN CHANGES OF THE PARTIAL SUMS OF A RANDOM MULTIPLICATIVE FUNCTION II

MARCO AYMONE

ABSTRACT. We study two models of random multiplicative functions: Rademacher random multiplicative functions supported on the squarefree integers f , and Rademacher random completely multiplicative functions f^* . We prove that the partial sums $\sum_{n \leq x} f^*(n)$ and $\sum_{n \leq x} \frac{f(n)}{\sqrt{n}}$ change sign infinitely often as $x \rightarrow \infty$, almost surely. The case $\sum_{n \leq x} \frac{f^*(n)}{\sqrt{n}}$ is leaved as an open question and we stress the possibility of only a finite number of sign changes, almost surely.

1. INTRODUCTION

1.1. **Main results.** A Rademacher random multiplicative function f is defined as follows: Over the primes, the values $f(p)$ are *i.i.d.* random variables taking ± 1 values with probability $1/2$ in each instance (Rademacher random variables), and at the other values of n ,

$$f(n) := \mu^2(n) \prod_{p|n} f(p),$$

where μ is the Möbius function.

By a Rademacher random completely multiplicative function f^* , we mean that

$$f^*(n) = \prod_{p^a || n} f(p)^a.$$

In a previous paper, Aymone, Heap and Zhao [3] proved that the partial sums $\sum_{n \leq x} f(n)$ of a Rademacher random multiplicative function change sign infinitely often¹ as $x \rightarrow \infty$, almost surely. The key idea was to show that, almost surely

$$\lim_{\sigma \rightarrow 1/2^+} \int_1^\infty \left(\sum_{n \leq x} f(n) \right) \frac{dx}{x^{1+\sigma}} = 0,$$

$$\limsup_{\sigma \rightarrow 1/2^+} \int_1^\infty \left| \sum_{n \leq x} f(n) \right| \frac{dx}{x^{1+\sigma}} = \infty,$$

and these visibly capture an infinite number of sign changes.

¹Here we say that a function $M(x)$ changes sign infinitely often as $x \rightarrow \infty$ if at a certain point x_0 , $M(x)$ is not always non-negative or non-positive for all $x \geq x_0$.

Our task here is to study sign changes for weighted sums of f and f^* :

$$\sum_{n \leq x} \frac{f(n)}{n^\alpha}, \sum_{n \leq x} \frac{f^*(n)}{n^\alpha}, \alpha \geq 0.$$

In both cases, we have that for $\alpha > 1/2$, the partial sums converge to an Euler product, and hence these partial sums become positive for all x sufficiently large, almost surely. Therefore, we restrict ourselves to the cases $0 \leq \alpha \leq 1/2$.

Theorem 1.1. *Let f be a Rademacher random multiplicative function. Then for each $0 \leq \alpha \leq 1/2$, $\sum_{n \leq x} \frac{f(n)}{n^\alpha}$ changes sign infinitely often as $x \rightarrow \infty$, almost surely.*

Theorem 1.2. *Let f^* be a Rademacher random completely multiplicative function. Then for each $0 \leq \alpha < 1/2$ we have that $\sum_{n \leq x} \frac{f^*(n)}{n^\alpha}$ changes sign infinitely often as $x \rightarrow \infty$, almost surely.*

The case $\sum_{n \leq x} \frac{f^*(n)}{\sqrt{n}}$ is leaved as an open question, and it is possible that these sums change sign only a finite number of times, here we give an heuristic reason. In a recent paper, Angelo and Xu [1] (and later improvements by Kerr and Klurman [20]) studied the weighted sums $\sum_{n \leq x} \frac{f^*(n)}{n}$ in the context of a “*random Turán conjecture*”. They proved that the probability of $\sum_{n \leq x} \frac{f^*(n)}{n}$ being positive for all $x \geq 1$ is at least $1 - 10^{-45}$. This implies a certain positivity for harmonic sums of f^* and it may be possible that such result may be extended for $\sum_{n \leq x} \frac{f^*(n)}{n^\sigma}$, where $\sigma > 1/2$. Further, after a convolution argument, $f^* = f * \mathbf{1}_{PS}$, where $\mathbf{1}_{PS}(n)$ is the indicator that n is a perfect square. Therefore,

$$\sum_{n \leq x} \frac{f^*(n)}{\sqrt{n}} \approx \sum_{n \leq x} \frac{f(n)}{\sqrt{n}} \log(x/n).$$

The later is known as a Riesz mean, and at some instances these averages stabilizes oscillatory properties of an oscillating arithmetic function providing a finite number of sign changes for these averages.

In the deterministic case of the Liouville function λ , in two interesting papers [24] by Mossinghoff and Trudgian and other by Humphries [19] is discussed sign changes of $\sum_{n \leq x} \frac{\lambda(n)}{n^\alpha}$. In [24] is leaved as an open problem to prove or disprove that $\sum_{n \leq x} \frac{\lambda(n)}{\sqrt{n}}$ is negative for all $x \geq 17$. A remarkable consequence of this fact is that this implies that all non-trivial zeros of the Riemann zeta function have multiplicity at most 2, see Theorem 2.4 of [24]. In [19] it is showed that under standard assumptions on the non-trivial zeros of the Riemann zeta function, the set of x for which $\sum_{n \leq x} \frac{\lambda(n)}{\sqrt{n}}$ is negative has total logarithmic asymptotic density.

1.2. Proof method. Our proof follows closely the lines of [3]. The difference here is that, in contrast with unweighted sums of a Rademacher random multiplicative function, we have the almost sure limits

$$\lim_{\sigma \rightarrow 1/2^+} \int_1^\infty \left(\sum_{n \leq x} \frac{f(n)}{\sqrt{n}} \right) \frac{dx}{x^{1/2+\sigma}} = \infty,$$

$$\limsup_{\sigma \rightarrow 1/2^+} \int_1^\infty \left| \sum_{n \leq x} \frac{f(n)}{\sqrt{n}} \right| \frac{dx}{x^{1/2+\sigma}} = \infty,$$

and this alone does not capture an infinite number of sign changes, unless that we make it quantitative the two divergences above, and that these divergences differ in size. This can, indeed, be achieved by using the following quantitative statement by Harper:

Theorem 1.3 (Harper [11], page 25). *Let p run over the primes and $(f(p))_p$ be a sequence of independent Rademacher random variables. For fixed $A > 3$, almost surely, there exists a sequence $\sigma_k \rightarrow 1/2^+$ such that*

$$\sup_{1 \leq t \leq 2 \left(\log \left(\frac{1}{\sigma_k - 1/2} \right) \right)^2} \left(\sum_p \frac{f(p) \cos(t \log p)}{p^{\sigma_k}} - 2 \log \log \left(\frac{1}{\sigma_k - 1/2} \right) \right) \geq \log \left(\frac{1}{\sigma_k - 1/2} \right) - A \log \log \left(\frac{1}{\sigma_k - 1/2} \right).$$

1.3. Background. Here we are intended to give a list of results in the literature (perhaps non-complete) on random multiplicative functions. The first result on partial sums of random multiplicative functions is due to Wintner [27], where he considered a question by Levy [23] and showed that $\sum_{n \leq x} f(n)$ is almost surely $\Omega(x^{1/2-\epsilon})$ and $\ll x^{1/2+\epsilon}$, for any $\epsilon > 0$.

Later, these results have been improved by Erdős [9] and subsequently by Halász [10]. The current best upper bound is due to Basquin [5] and to Lau, Tenenbaum and Wu [22], and states that, almost surely,

$$\sum_{n \leq x} f(n) \ll \sqrt{x} (\log \log x)^{2+\epsilon},$$

for any $\epsilon > 0$.

The current best Omega bound is due to Harper [15]:

$$\sum_{n \leq x} f(n) = \Omega(\sqrt{x} (\log \log x)^{1/4-\epsilon}),$$

for any $\epsilon > 0$, almost surely.

Central limit Theorems also have been studied in various settings: [8], [12], [18], [21], [26].

The problem of moments also have been extensively studied, with the remarkable result of Harper that shows that the first moment exhibits better than squareroot cancellation [14]. Other related works (including related models): [2], [6], [7], [13], [16], [17], [28].

And another viewpoints of investigation [1], [3], [4], [20], [25].

2. PRELIMINARIES

2.1. Notation. Here p stands for a generic prime number. We use the standard Vinogradov notation $f(x) \ll g(x)$ or Landau's $f(x) = O(g(x))$ whenever there exists a constant $c > 0$ such that $|f(x)| \leq c|g(x)|$, for all x in a set of parameters. When not specified, this set of parameters is an infinite interval (a, ∞) for sufficiently large $a > 0$. The standard $f(x) = o(g(x))$ means that $f(x)/g(x) \rightarrow 0$ when $x \rightarrow a$, where a could be a complex number or $\pm\infty$. For $g(x) > 0$ for all x , we say that $f(x) = \Omega(g(x))$ if $\limsup_{x \rightarrow \infty} \frac{|f(x)|}{g(x)} > 0$.

2.2. Some Lemmas. By the pioneering work of Wintner [27], we have that $F(s) := \sum_{n \leq x} \frac{f(n)}{n^s}$ converges for all $Re(s) > 1/2$, almost surely. His proof can be modified to show that the same is true for $F^*(s) = \sum_{n \leq x} \frac{f^*(n)}{n^s}$. Moreover we have:

Lemma 2.1. *For all $Re(s) > 1/2$, almost surely we have*

$$F(s) = \prod_p \left(1 + \frac{f(p)}{p^s} \right),$$

$$F^*(s) = \prod_p \left(1 - \frac{f(p)}{p^s} \right)^{-1}.$$

This leads to the almost sure identities

$$(1) \quad F(s) = \exp \left(\sum_p \frac{f(p)}{p^s} - \frac{1}{2} \log \zeta(2s) + O(1) \right),$$

$$(2) \quad F^*(s) = \exp \left(\sum_p \frac{f(p)}{p^s} + \frac{1}{2} \log \zeta(2s) + O(1) \right),$$

where the $O(1)$ term above is actually an analytic function that is uniformly bounded in $Re(s) \geq 1/3 + \epsilon$, for small fixed $\epsilon > 0$. For a proof of these results we refer reader to Lemma 2.4 of [3] and the references therein. To conclude this section, we have

Lemma 2.2. *Let $(f(p))_p$ be independent Rademacher random variables. Then, for all $\epsilon > 0$, we have almost surely as $\sigma \rightarrow 1/2^+$*

$$\sum_p \frac{f(p)}{p^\sigma} \ll \left(\log \left(\frac{1}{\sigma - 1/2} \right) \right)^{1/2+\epsilon}.$$

For a proof we refer reader to Lemma 2.1 and Remark 2.1 of [3].

3. PROOF OF THE MAIN RESULTS

3.1. The case $\sum_{n \leq x} \frac{f(n)}{\sqrt{n}}$. By the current best almost sure upper bound for the partial sums of a Rademacher random multiplicative function [5] and [22],

$$\sum_{n \leq x} f(n) \ll \sqrt{x} (\log \log x)^{2+\epsilon},$$

we have that by partial summation

$$\sum_{n \leq x} \frac{f(n)}{\sqrt{n}} \ll (\log x)^2, \text{ a.s.}$$

Further refinements of this last upper bound could, perhaps, be achieved, by following the lines of [2]. Now, for $Re(s) = \sigma > 1/2$,

$$F(s) := \sum_{n=1}^{\infty} \frac{f(n)}{n^s} = \sum_{n=1}^{\infty} \frac{f(n)}{\sqrt{n}} \frac{1}{n^{s-1/2}} = (s - 1/2) \int_1^{\infty} \left(\sum_{n \leq x} \frac{f(n)}{\sqrt{n}} \right) \frac{dx}{x^{s+1/2}}.$$

By equation (1) and Lemma 2.2 combined with the estimate $\log \zeta(2\sigma) = \log(1/(2\sigma - 1)) + O(1)$ valid for $\sigma \rightarrow 1^+$, we obtain that as $\sigma \rightarrow 1/2^+$, almost surely we have $F(\sigma) = (2\sigma - 1)^{1/2+o(1)}$. Hence, we almost surely have as $\sigma \rightarrow 1/2^+$

$$(3) \quad \int_1^{\infty} \left(\sum_{n \leq x} \frac{f(n)}{\sqrt{n}} \right) \frac{dx}{x^{\sigma+1/2}} = \frac{1}{(2\sigma - 1)^{1/2+o(1)}}.$$

On the other hand, for $t \geq 1$

$$\left| \frac{F(\sigma + it)}{t} \right| \ll \int_1^{\infty} \left| \sum_{n \leq x} \frac{f(n)}{\sqrt{n}} \right| \frac{dx}{x^{\sigma+1/2}}.$$

Now, by equation (1),

$$|F(s)| = \exp \left(\sum_p \frac{f(p) \cos(t \log p)}{p^\sigma} - \frac{1}{2} Re(\log \zeta(2s)) + O(1) \right).$$

We recall that $|\log \zeta(\sigma + it)| \leq \log \log |t| + O(1)$, for $\sigma \geq 1$ and $t \geq 2$. By Theorem 1.3, almost surely we have an infinite sequence of points $\sigma_k \rightarrow 1/2^+$ and $t_k \geq 1$ such that

$$\left| \frac{F(\sigma_k + it_k)}{t_k} \right| \geq \exp \left((1 + o(1)) \log \left(\frac{1}{\sigma_k - 1/2} \right) \right) \gg \frac{1}{(\sigma_k - 1/2)^{0.9}}.$$

Therefore, almost surely, there exists a sequence $\sigma_k \rightarrow 1/2^+$ such that

$$(4) \quad \int_1^\infty \left| \sum_{n \leq x} \frac{f(n)}{\sqrt{n}} \right| \frac{dx}{x^{\sigma_k + 1/2}} \gg \frac{1}{(\sigma_k - 1/2)^{0.9}}.$$

To complete the argument, we see that the quantities at the left of (3) and (4) have different size, and so the partial sums $\sum_{n \leq x} \frac{f(n)}{\sqrt{n}}$ cannot be always non-negative or non-positive, there always must be infinite sign changes, almost surely.

3.2. The case $\sum_{n \leq x} \frac{f^*(n)}{n^\alpha}$. Let then $0 \leq \alpha < 1/2$. Let $F^*(s)$ be the Dirichlet series of f^* . Similarly as in the previous case, by equation (2) and Lemma 2.2, we have that

$$F^*(\sigma) = \frac{1}{(2\sigma - 1)^{1/2 + o(1)}},$$

almost surely as $\sigma \rightarrow 1/2^+$.

By partial summation and integration,

$$F^*(s) = (s - \alpha) \int_1^\infty \left(\sum_{n \leq x} \frac{f^*(n)}{n^\alpha} \right) \frac{dx}{x^{s+1-\alpha}}.$$

Therefore, almost surely as $\sigma \rightarrow 1/2^+$

$$(5) \quad \int_1^\infty \left(\sum_{n \leq x} \frac{f^*(n)}{n^\alpha} \right) \frac{dx}{x^{\sigma+1-\alpha}} = \frac{1 + o(1)}{(1/2 - \alpha)} \frac{1}{(2\sigma - 1)^{1/2 + o(1)}}.$$

And similarly to (4), almost surely there exists a sequence $\sigma_k \rightarrow 1/2^+$ such that

$$(6) \quad \int_1^\infty \left| \sum_{n \leq x} \frac{f^*(n)}{n^\alpha} \right| \frac{dx}{x^{\sigma+1-\alpha}} \gg \frac{1}{(2\sigma_k - 1)^{0.99}}.$$

Just as before, (5) and (6) captures infinite sign changes, almost surely.

3.3. The case $\sum_{n \leq x} \frac{f(n)}{n^\alpha}$. This case can be treated similarly as the previous one, although we can proceed in a more elementary way as in [3], that is, Theorem 1.3 is not needed here.

The proof of Theorems 1.1 and 1.2 are now complete.

Acknowledgements This project started while I am as a visiting Professor at Aix-Marseille Université, and I am warmly thankful for the kind hospitality here, and special thanks to Prof. Olivier Ramaré. This project is funded by CNPq grant PDE no. 400010/2022-4 (200121/2022-7) and by CNPq grant Universal no. 403037/2021-2.

REFERENCES

- [1] R. ANGELO AND M. XU, *On a Turán conjecture and random multiplicative functions*, Quart. J. Math., online first (2022).
- [2] M. AYMONE, W. HEAP, AND J. ZHAO, *Partial sums of random multiplicative functions and extreme values of a model for the Riemann zeta function*, J. Lond. Math. Soc. (2), 103 (2021), pp. 1618–1642.
- [3] ———, *Sign changes of the partial sums of a random multiplicative function*, Bull. Lond. Math. Soc., 55 (2023), pp. 78–89.
- [4] M. AYMONE AND V. SIDORAVICIUS, *Partial sums of biased random multiplicative functions*, J. Number Theory, 172 (2017), pp. 343–382.
- [5] J. BASQUIN, *Sommes friables de fonctions multiplicatives aléatoires*, Acta Arith., 152 (2012), pp. 243–266.
- [6] J. BENATAR, A. NISHRY, AND B. RODGERS, *Moments of polynomials with random multiplicative coefficients*, Mathematika, 68 (2022), pp. 191–216.
- [7] A. BONDARENKO AND K. SEIP, *Helson’s problem for sums of a random multiplicative function*, Mathematika, 62 (2016), pp. 101–110.
- [8] S. CHATTERJEE AND K. SOUNDARARAJAN, *Random multiplicative functions in short intervals*, Int. Math. Res. Not. IMRN, (2012), pp. 479–492.
- [9] P. ERDŐS, *Some unsolved problems*, Magyar Tud. Akad. Mat. Kutató Int. Közl., 6 (1961), pp. 221–254.
- [10] G. HALÁSZ, *On random multiplicative functions*, in Hubert Delange colloquium (Orsay, 1982), vol. 83 of Publ. Math. Orsay, Univ. Paris XI, Orsay, 1983, pp. 74–96.
- [11] A. J. HARPER, *Bounds on the suprema of Gaussian processes, and omega results for the sum of a random multiplicative function*, Ann. Appl. Probab., 23 (2013), pp. 584–616.
- [12] ———, *On the limit distributions of some sums of a random multiplicative function*, J. Reine Angew. Math., 678 (2013), pp. 95–124.
- [13] ———, *Moments of random multiplicative functions, II: High moments*, Algebra Number Theory, 13 (2019), pp. 2277–2321.
- [14] ———, *Moments of random multiplicative functions, I: Low moments, better than squareroot cancellation, and critical multiplicative chaos*, Forum Math. Pi, 8 (2020), pp. e1, 95.
- [15] ———, *Almost sure large fluctuations of random multiplicative functions*, Int. Math. Res. Not., (2022).
- [16] A. J. HARPER, A. NIKEGHBALI, AND M. RADZIWIŁ Ł, *A note on Helson’s conjecture on moments of random multiplicative functions*, in Analytic number theory, Springer, Cham, 2015, pp. 145–169.
- [17] W. P. HEAP AND S. LINDQVIST, *Moments of random multiplicative functions and truncated characteristic polynomials*, Q. J. Math., 67 (2016), pp. 683–714.
- [18] B. HOUGH, *Summation of a random multiplicative function on numbers having few prime factors*, Math. Proc. Cambridge Philos. Soc., 150 (2011), pp. 193–214.
- [19] P. HUMPHRIES, *The distribution of weighted sums of the Liouville function and Pólya’s conjecture*, J. Number Theory, 133 (2013), pp. 545–582.
- [20] B. KERR AND O. KLURMAN, *How negative can $\sum_{n \leq x} \frac{f(n)}{n}$ be?*, arXiv:2211.05540, (2022).
- [21] O. KLURMAN, I. D. SHKREDOV, AND M. W. XU, *On the random Chowla conjecture*, GAFA (to appear), (2023).

- [22] Y.-K. LAU, G. TENENBAUM, AND J. WU, *On mean values of random multiplicative functions*, Proc. Amer. Math. Soc., 141 (2013), pp. 409–420.
- [23] P. LÉVY, *Sur les séries dont les termes sont des variables éventuelles indépendents*, Studia Mathematica, 03 (1931), pp. 119–155.
- [24] M. J. MOSSINGHOFF AND T. S. TRUDGIAN, *Between the problems of Pólya and Turán*, J. Aust. Math. Soc., 93 (2012), pp. 157–171.
- [25] J. NAJNUDEL, *On consecutive values of random completely multiplicative functions*, Electron. J. Probab., 25 (2020), pp. Paper No. 59, 28.
- [26] K. SOUNDARARAJAN AND M. XU, *Central limit theorems for random multiplicative functions*, arXiv:2212.06098, (2022).
- [27] A. WINTNER, *Random factorizations and Riemann's hypothesis*, Duke Math. J., 11 (1944), pp. 267–275.
- [28] M. W. XU, *Better than square-root cancellation for random multiplicative functions*, arXiv:2303.06774, (2023).

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE MINAS GERAIS (UFMG), Av. ANTÔNIO CARLOS, 6627, CEP 31270-901, BELO HORIZONTE, MG, BRAZIL.

Email address: `aymone.marco@gmail.com`