

FLEXIBILITY OF GENERALIZED ENTROPY FOR WANDERING DYNAMICS

JAVIER CORREA AND HELLEN DE PAULA

ABSTRACT. We show a flexibility result in the context of generalized entropy. The space of dynamical systems we work with is, homeomorphisms on the sphere whose non-wandering set consist in only one fixed point.

Keywords: Generalized entropy, Polynomial entropy, Flexibility.

2020 Mathematics subject classification: 37A35, 37B40 and 37E30.

1. INTRODUCTION

An important problem in dynamical systems is to measure the complexity of a map in terms of its orbits. The topological entropy of a system studies the exponential growth rate at which orbits are separated and has become a crucial tool for classifying highly chaotic dynamical systems. However, there are many interesting families of dynamical systems, where every system has vanishing entropy, and therefore, another object is needed. The object that studies the polynomial growth rate is called polynomial entropy and was introduced by J. P. Marco in [8]. S. Galatolo introduced in [5] the concept of generalized topological entropy that extends classical and polynomial entropy for one-parameter families of orders of growth. This concept is a translation of A. Katok and J. P. Thouvenot notion of slow entropy defined in [7]. For a more detailed exposition on the works related to the previous quantities check [4], [3] and [1]. In this article, we are going to work with the notion of generalized entropy introduced by the first author and E. Pujals in [4]. This object, instead of quantifying the complexity of a system with a single number, works directly in the space of the orders of growth \mathbb{O} . In subsection 2.1 we show how is constructed $o(f)$, the generalized entropy of a map f .

In this text we are interested in the families of dynamical systems on the sphere whose non-wandering set consist in only one fixed point. We shall denote by S^2 the sphere and $\Omega(f)$ the non-wandering set of a map $f : S^2 \rightarrow S^2$. We define \mathcal{H} as the family of homeomorphisms $f : S^2 \rightarrow S^2$ such that $\#\Omega(f) = 1$. A simple consequence of the variational principle is that any map in \mathcal{H} has vanishing classical entropy. The family \mathcal{H} is naturally interesting because the orientation preserving ones are in a direct bijection with Brouwer homeomorphisms of the plane (through compactifying the plane with one point).

For the family \mathcal{H} , in the context of polynomial entropy we have the result of L. Haseux and F. Le Roux in [6]. We shall call the polynomial entropy of a map by $h_{pol}(f)$.

Theorem 1.1 (L. Haseux and F. Le Roux). *The following two happens:*

- (1) For any $f \in \mathcal{H}$, $h_{pol}(f) \in \{1\} \cup [2, \infty]$.
- (2) For any $t \in \{1\} \cup [2, \infty]$, there exists $f \in \mathcal{H}$ such that $h_{pol}(f) = t$. Moreover, $h_{pol}(f) = 1$ if and only if f is the compactification of a map conjugate to translation in \mathbb{R}^2 .

The first part of the theorem gives natural boundaries and the second part is a flexibility result in the sense introduced by J. Bochi, A. Katok and F. Rodriguez-Hertz in [2]. The article [6] has two parts. First, a way to code the orbits of a dynamical system with $\#\Omega(f) = 1$, that allows the computation of $h_{pol}(f)$ and second, the construction of maps in the point (2) of theorem 1.1. We have extended the first part in [3] to generalized entropy and also to maps with finite non-wandering set. With this tool, we computed the generalized entropy of Morse-Smale diffeomorphisms on surfaces and some maps in the boundary of chaos. Our main result in this article is an extension of the second part of the work by L. Haseux and F. Le Roux.

In order to enunciate our theorem, we need to fix some terminology associated to \mathbb{O} the space of orders of growth. To us, an order of growth is the class of non-decreasing sequences in $(0, \infty)$ that have the same asymptotic speed at ∞ . We represent this classes by $[a(n)]$ for the general case and the formula between brackets if it is defined by one, for instance $[n^2]$. The family of polynomial orders of growth is $\mathbb{P} = \{[n^t] : 0 < t < \infty\}$. We say that an order of growth $[a(n)]$ verifies the linearly invariant property (LIP) if for some $m \geq 2$, $[a(mn)] = [a(n)]$. The set of orders of growth that verifies LIP is represented by \mathbb{L} (observe that $\mathbb{P} \subset \mathbb{L}$). The abstract completion of \mathbb{O} by its partial order is the complete set of orders of growth $\overline{\mathbb{O}}$ and the generalized entropy of a map $o(f)$ is defined in $\overline{\mathbb{O}}$. General elements of $\overline{\mathbb{O}}$ are represented by o . We define $\overline{\mathbb{L}}$ as the countable completion of \mathbb{L} in $\overline{\mathbb{O}}$ (Check subsections 2.1, 2.3 and 2.4 for more details).

By our results in [3], is immediate that for any $f \in \mathcal{H}$, either $o(f) = [n]$ or $[n^2] \leq o(f) \leq \sup(\mathbb{P})$. The main result in this article is the following.

Theorem 1. *For any $o \in \overline{\mathbb{L}}$ verifying $o = [n]$ or $[n^2] \leq o \leq \sup(\mathbb{P})$, there exists $f \in \mathcal{H}$ such that $o(f) = o$.*

We observe that the order of growth $[n^2 \log(n)]$ belongs to $\overline{\mathbb{L}}$ and therefore the following corollary holds.

Corollary 1. *There exists $f_1, f_2 \in \mathcal{H}$ such that $o(f_1) = [n^2]$ and $o(f_2) = [n^2 \log(n)]$. In particular, $h_{pol}(f_1) = h_{pol}(f_2)$ yet $o(f_1) \neq o(f_2)$.*

The first part of theorem 1.1 implies that polynomial orders of growth is the right scope to analyze the dispersion of orbits for maps in \mathcal{H} . However, we argue, in the light corollary 1,

that generalized entropy allows to appreciate on a deeper level how rich the universe of possible complexities actually is.

Let us comment on the hypothesis $\overline{\mathbb{L}}$. There is a weaker property an order of growth may verify that we call the bounded jump property (BJP) and was introduced in [3]. We shall use \mathbb{B} to represent the elements in \mathbb{O} that verifies BJP and $\overline{\mathbb{B}}$ to represent the countable completion of \mathbb{B} . The BJP is in fact necessary: for any $f : M \rightarrow M$ with M a compact metric space, $o(f) \in \overline{\mathbb{B}}$ (see proposition 2.4). Therefore, the best version of theorem 1 that may be true is for any $o \in \overline{\mathbb{B}}$. The linearly invariant property implies the bounded jump property ($\mathbb{L} \subset \mathbb{B}$) but the converse is false. For example any exponential order of growth $[e^{tn}]$ fails to verify LIP. The LIP is in fact associated to sub-exponential orders of growth (see proposition 2.7), but even in this context we believe to be possible the construction of elements in $[a(n)] \in \mathbb{B} \setminus \mathbb{L}$ such that $[a(n)] < \sup(\mathbb{P})$. On the other hand, any simple example of $[a(n)] < \sup(\mathbb{P})$ constructed by formulas verify LIP. For instance $[n^t]$, $[\log(n)]$, $[\log(\log(n))]$ belong to \mathbb{L} . Moreover, we consider it an nice hypothesis to work with. Since it is naturally associated to sub-exponential orders of growth, any proof involving LIP, will use arguments intrinsic to the world of vanishing entropy that will not be an adaptation from the positive entropy setting.

We would like to finish this introduction with some comments and open questions. Consider M a compact metric space and $f : M \rightarrow M$ a homeomorphism. A simple consequence of the variational principle is that $h(f|_{\Omega(f)}) = h(f)$. However, for generalized entropy it is only true $o(f|_{\Omega(f)}) \leq o(f)$. In order to have a clean notation we call $o(f, \Omega(f)) = o(f|_{\Omega(f)})$. As it was observed in [4] the generalized entropy may “spike” from $o(f, \Omega(f))$ to $o(f)$. Moreover, for the class of homeomorphisms worked in this article \mathcal{H} , every map verifies $o(f, \Omega(f))$ is the class of the constant sequence $[c]$ and $[n] \leq o(f) \leq \sup(\mathbb{P})$. Therefore, $o(f, \Omega(f)) < o(f)$. We consider that a possible path to understand this jumps in the general case is the following. Define $\hat{M} = M/\Omega(f)$, this is, collapse the non-wandering set of f to a single point. The map f , induces a homeomorphism $\hat{f} : \hat{M} \rightarrow \hat{M}$ whose non-wandering set contains only one point. Now, with the examples we have worked so far, the inequalities

$$\max\{o(f, \Omega(f)), o(\hat{f})\} \leq o(f) \leq o(f, \Omega(f))o(\hat{f}),$$

always seems to hold and we ask if this is true in general. It is simple to construct examples where one inequality is strict. If the answer is positive and $o(f, \Omega(f)) = [c]$, then $o(f) = o(\hat{f})$ because $[c]$ is the identity for the product in $\overline{\mathbb{O}}$. Therefore, a simpler version of our previous question is the following: If $o(f, \Omega(f)) = [c]$, is $o(f) = o(\hat{f})$?

This work is structured as follows:

- In section 2, we show the construction of the set of orders growth and generalized entropy. We also explain the coding of orbits for maps in \mathcal{H} , and prove some simple results about the bounded jump and linearly invariant properties.
- In section 3, we prove theorem 1. We split the proof in two cases. We address first when $o \in \mathbb{B}$ and then when $o \in \overline{\mathbb{B}}$.

2. PRELIMINARIES

2.1. Orders of growth and generalized entropy. Let us briefly recall how the complete set of the orders of growth and the generalized entropy of a map are defined in [4]. First, we consider the space of non-decreasing sequences in $[0, \infty)$:

$$\mathcal{O} = \{a : \mathbb{N} \rightarrow [0, \infty) : a(n) \leq a(n+1), \forall n \in \mathbb{N}\}.$$

Next, we define the equivalence relationship \approx in \mathcal{O} by $a(n) \approx b(n)$ if and only if there exist $c_1, c_2 \in (0, \infty)$ such that $c_1 a(n) \leq b(n) \leq c_2 a(n)$ for all $n \in \mathbb{N}$. Since two sequences are related, if both have the same order of growth, we call the quotient space $\mathbb{O} = \mathcal{O}/\approx$ as the space of the orders of growth. If $a(n)$ belongs to \mathcal{O} , we are going to denote $[a(n)]$ as the associated class in \mathbb{O} . If a sequence is defined by a formula (for example, n^2), then the order of growth associated will be represented by the formula between the brackets ($[n^2] \in \mathbb{O}$).

We define in \mathbb{O} a natural partial order. We say that $[a(n)] \leq [b(n)]$ if there exists $C > 0$ such that $a(n) \leq Cb(n)$, for all $n \in \mathbb{N}$. Observe that \leq is well defined. We consider $\overline{\mathbb{O}}$ the Dedekind-MacNeille completion. This is the smallest complete lattice that contains \mathbb{O} . In particular, it is uniquely defined and we will consider that $\mathbb{O} \subset \overline{\mathbb{O}}$. We will also call $\overline{\mathbb{O}}$ the complete set of the orders of growth.

Now, we proceed to define the generalized entropy of a dynamical system in the complete space of the orders of growth. Given (M, d) , a compact metric space and $f : M \rightarrow M$ a continuous map. We define in M the distance

$$d_n^f(x, y) = \max\{d(f^k(x), f^k(y)); 0 \leq k \leq n-1\},$$

and we denote the dynamical ball as $B(x, n, \varepsilon) = \{y \in M; d_n^f(x, y) < \varepsilon\}$. A set $G \subset M$ is a (n, ε) -generator if $M = \cup_{x \in G} B(x, n, \varepsilon)$. We define $g_{f, \varepsilon}(n)$ as the smallest possible cardinality of a finite (n, ε) -generator. If we fix $\varepsilon > 0$, then $g_{f, \varepsilon}(n)$ is a non-decreasing sequence of natural numbers, thus, $g_{f, \varepsilon}(n) \in \mathcal{O}$. For a fixed n , if $\varepsilon_1 < \varepsilon_2$, then $g_{f, \varepsilon_1}(n) \geq g_{f, \varepsilon_2}(n)$, and therefore $[g_{f, \varepsilon_1}(n)] \geq [g_{f, \varepsilon_2}(n)]$ in \mathbb{O} . We define the generalized entropy of f as

$$o(f) = \text{“lim”}_{\varepsilon \rightarrow 0} [g_{f, \varepsilon}(n)] = \sup\{[g_{f, \varepsilon}(n)] \in \mathbb{O} : \varepsilon > 0\} \in \overline{\mathbb{O}}.$$

This object is a dynamical invariant.

Theorem 2.1 (Correa-Pujals [4]). *Let M and N be two compact metric spaces and $f : M \rightarrow M$, $g : N \rightarrow N$, two continuous maps. Suppose there exists $h : M \rightarrow N$, a homeomorphism, such that $h \circ f = g \circ h$. Then, $o(f) = o(g)$.*

We also define the generalized entropy through the point of view of (n, ε) -separated. We say that $E \subset M$ is (n, ε) -separated if $B(x, n, \varepsilon) \cap E = \{x\}$, for all $x \in E$. We define $s_{f, \varepsilon}(n)$ as the maximal cardinality of a (n, ε) -separated set. Analogously, if we fix $\varepsilon > 0$, then $s_{f, \varepsilon}(n)$ is a non-decreasing sequence of natural numbers. Again, for a fixed n , if $\varepsilon_1 < \varepsilon_2$, then $s_{f, \varepsilon_1}(n) \geq s_{f, \varepsilon_2}(n)$, and therefore, $[s_{f, \varepsilon_1}(n)] \geq [s_{f, \varepsilon_2}(n)]$. By standard arguments it is verified

$$o(f) = \sup\{[s_{f, \varepsilon}(n)] \in \mathbb{O} : \varepsilon > 0\}.$$

The order of growth defined through open coverings is also equivalent. Consider $\mathcal{U} = \{U_1, \dots, U_k\}$ an open covering of M and define

$$\mathcal{U}^n = \{U_{i_0} \cap f^{-1}(U_{i_1}) \cap \dots \cap f^{-n}(U_{i_n}) \neq \emptyset : i_0, \dots, i_n \in \{1, \dots, k\}\},$$

which is also an open covering of M . Next, we define the sequence $a_{f,\mathcal{U}}(n)$ as the cardinal of the smallest sub-covering of \mathcal{U}^n . Then, it holds

$$o(f) = \sup\{[a_{f,\mathcal{U}}(n)] : \mathcal{U} \text{ is a finite open covering of } M\}.$$

Now, we shall explain how the generalized topological entropy is related to the classical notion of topological entropy and polynomial entropy. Given a dynamical system f , we recall that the topological entropy of f is

$$h(f) = \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{\log(g_{f,\varepsilon}(n))}{n},$$

and the polynomial entropy of f is

$$h_{pol}(f) = \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{\log(g_{f,\varepsilon}(n))}{\log(n)}.$$

We define the family of exponential orders of growth as the set $\mathbb{E} = \{[\exp(tn)]; t \in (0, \infty)\} \subset \mathbb{O}$ and the family of polynomials orders of growth as the set $\mathbb{P} = \{[n^t]; t \in (0, \infty)\}$. In [4], the authors defined two natural projections, $\pi_{\mathbb{E}} : \mathbb{O} \rightarrow [0, \infty]$ and $\pi_{\mathbb{P}} : \mathbb{O} \rightarrow [0, \infty]$, for which the following theorem holds.

Theorem 2.2 (Correa-Pujals [4]). *Let M be a compact metric space and $f : M \rightarrow M$, a continuous map. Then, $\pi_{\mathbb{E}}(o(f)) = h(f)$ and $\pi_{\mathbb{P}}(o(f)) = h_{pol}(f)$.*

The projection $\pi_{\mathbb{E}}$ verifies for elements in \mathbb{O} , $\pi_{\mathbb{E}}([a(n)]) = \limsup_{n \rightarrow \infty} \frac{\log(a(n))}{n}$.

2.2. Coding of orbits in wandering dynamics. Let us consider M a compact metric space and $f : M \rightarrow M$ a homeomorphism such that $\Omega(f) = \{p_1, \dots, p_k\}$. Let \mathcal{F} be a finite family of non-empty subsets of $M \setminus \Omega(f)$. We denote by $\cup \mathcal{F}$ the union of all the elements of \mathcal{F} and by $\infty_{\mathcal{F}}$ the complement of $\cup \mathcal{F}$. Let us fix a positive integer n and consider $\underline{x} = (x_0, \dots, x_{n-1})$ a finite sequence of points in M and $\underline{w} = (w_0, \dots, w_{n-1})$ a finite sequence of elements of $\mathcal{F} \cup \{\infty_{\mathcal{F}}\}$. We say that \underline{w} is a coding of \underline{x} , relative to \mathcal{F} , if for every $i = 0, \dots, n-1$, holds $x_i \in w_i$. Whenever the family \mathcal{F} is fixed, we simplify the notation by using ∞ instead of $\infty_{\mathcal{F}}$. Note that if the sets of \mathcal{F} are not disjoint, we can have more than one coding for a given sequence. We denote the set of all the codings of all orbits $(x, f(x), \dots, f^{n-1}(x))$ of length n by $\mathcal{A}_n(f, \mathcal{F})$. We define the sequence $c_{f,\mathcal{F}}(n) = \#\mathcal{A}_n(f, \mathcal{F})$, and it is easy to see that $c_{f,\mathcal{F}}(n) \in \mathbb{O}$.

We say that:

- a set Y is wandering if $f^n(Y) \cap Y = \emptyset$ for every $n \geq 1$.
- Y is a compact neighborhood if it is compact and is the closure of an open set.

- the subsets Y_1, \dots, Y_L of $M \setminus \{\Omega(f)\}$ are mutually singular if, for every $N > 0$, there exists a point x and times n_1, \dots, n_L such that $f^{n_i}(x) \in Y_i$ for every $i = 1, \dots, L$, and $|n_i - n_j| > N$ for every $i \neq j$.

Let us call Σ as the family of finite families of wandering compact neighborhoods that are mutually singular. Given $\delta > 0$, we define Σ_δ as the subset of Σ formed by every family whose every element has a diameter smaller than δ .

Theorem 2.3 (Correa - de Paula [3]). *Let M be a compact metric space and $f : M \rightarrow M$ a homeomorphism such that $\Omega(f)$ is finite. Then,*

$$o(f) = \sup\{[c_{f,\mathcal{F}}(n)] \in \mathbb{O} : \mathcal{F} \in \Sigma\}.$$

In addition, the equation also holds if we switch Σ by Σ_δ .

2.3. Bounded Jump property. We say that a class of orders of growth $[a(n)]$ verifies the bounded jump property (BJP) if there exists a constant $C > 0$ such that $a(n+1) \leq Ca(n)$. Note that this definition does not depend on the choice of the class representative. We call $\mathbb{B} \subset \mathbb{O}$ as the set of orders of growth that verify the BJP and $\overline{\mathbb{B}}$ the subset in $\overline{\mathbb{O}}$ defined by:

$$\overline{\mathbb{B}} = \{sup(\Gamma) \in \overline{\mathbb{O}} : \Gamma \subset \mathbb{B} \text{ and } \Gamma \text{ is countable}\}.$$

The set $\overline{\mathbb{B}}$ is the countable completion of \mathbb{B} .

Proposition 2.4. *Let us consider M a compact metric space and $f : M \rightarrow M$ a homeomorphism. Then, it is verified*

$$(1) \text{ For } \mathcal{U} \text{ a finite covering of } M, [a_{f,\mathcal{U}}(n)] \in \mathbb{B}.$$

$$(2) o(f) \in \overline{\mathbb{B}}.$$

Proof. We shall begin with proving the result for open coverings. Consider $\mathcal{U} = \{U_1, \dots, U_k\}$ a finite open covering of M and fix $n \in \mathbb{N}$. Suppose that $\mathcal{V} \subset \mathcal{U}^n$ is a sub-covering such that $a_{f,\mathcal{U}}(n) = \#\mathcal{V}$. Then, $\mathcal{V}' = \{V \cap f^{-(n+1)}(U_i) : V \in \mathcal{V}, U_i \in \mathcal{U}\} \subset \mathcal{U}^{n+1}$ is also a sub-covering with at most $\#U\#V$ elements. Therefore, $a_{f,\mathcal{U}}(n+1) \leq \#U a_{f,\mathcal{U}}(n)$.

Now that we know $[a_{f,\mathcal{U}}(n)] \in \mathbb{B}$, we only need to show that $o(f)$ is obtained as the supremum of a countable family of open coverings. By lemma A.1 in [4], given $\varepsilon > 0$ if $diam(\mathcal{U}) \leq \varepsilon$, then $[a_{f,\mathcal{U}}(n)] \geq [s_{f,\varepsilon}(n)]$. Therefore, if $\{\mathcal{U}_k\}_{k \in \mathbb{N}}$ is a family of open coverings such that $diam(\mathcal{U}_k) \leq \frac{1}{k}$, then $o(f) = \sup\{[a_{f,\mathcal{U}_k}(n)] : k \in \mathbb{N}\}$. \square

In [3] we proved an analogous result for $[c_{f,\mathcal{F}}(n)]$.

Lemma 2.5 (Correa - de Paula [3]). *Let us consider M a compact metric space and $f : M \rightarrow M$ a homeomorphism whose non-wandering set is finite. Given $\mathcal{F} \in \Sigma$, the class $[c_{f,\mathcal{F}}(n)] \in \mathbb{B}$.*

The next result show how we use the BJP. We recall that a set $\mathcal{S} \subset \mathbb{N}$ is syndetic if there exists $N \in \mathbb{N}$ such that for all n , the interval $[n, n+N]$ contains at least one point of \mathcal{S} .

Lemma 2.6 (Correa - de Paula [3]). *Let us consider $[a(n)] \in \mathbb{B}$, a syndetic set \mathcal{S} and a sequence $b(n) \in \mathcal{O}$. If there exist two constants c_1 and c_2 such that $c_1 b(n) \leq a(n) \leq c_2 b(n)$ for all $n \in \mathcal{S}$, then $[a(n)] = [b(n)]$.*

This lemma tells us that if we know an order of growth in a syndetic set and said order of growth verifies the BJP, then we understand the order of growth in \mathbb{N} . For example, if we prove that $c_1 n^k \leq c_{f,\mathcal{F}}(n) \leq c_2 n^k$ for every $n \in \mathcal{S}$, then $[c_{f,\mathcal{F}}(n)] = [n^k]$.

2.4. Linearly invariant property. We say that a class of orders of growth $[a(n)]$ verifies the linearly invariant property if there exists an integer $m \geq 2$ such that $[a(n)] = [a(mn)]$. We call $\mathbb{L} \subset \mathbb{O}$ as the set of orders of growth that verify the LIP and $\overline{\mathbb{L}}$ the subset in $\overline{\mathbb{O}}$ defined by:

$$\overline{\mathbb{L}} = \{\sup(\Gamma) \in \overline{\mathbb{O}} : \Gamma \subset \mathbb{L} \text{ and } \Gamma \text{ is countable}\}.$$

The set $\overline{\mathbb{L}}$ is the countable completion of \mathbb{L} .

Orders of growth in \mathbb{L} verify the following properties

Proposition 2.7. *Let $[a(n)] \in \mathbb{L}$. Then, it is verified:*

- (1) $[a(n)] \in \mathbb{B}$.
- (2) For all integer $m \geq 2$ it is verified $[a(mn)] = [a(n)]$.
- (3) $\pi_{\mathbb{E}}([a(n)]) = 0$.

Proof. (1). The BJP is equivalent to $[a(n)] = [a(n+1)]$. Suppose that $[a(mn)] = [a(n)]$ for some $m \geq 2$. Since $a(n) \leq a(n+1) \leq a(mn)$ for $n \geq 1$, we infer $[a(n)] = [a(n+1)]$.

(2). Let us suppose $[a(2n)] = [a(n)]$ and consider $c_2 > c_1 > 0$ such that $c_1 \leq \frac{a(2n)}{a(n)} \leq c_2$. Observe that

$$c_1^2 \leq \frac{a(4n)}{a(2n)} \frac{a(2n)}{a(n)} \leq c_2^2 \quad \forall n \in \mathbb{N}.$$

From this, we deduce that $[a(4n)] = [a(n)]$ and by induction we conclude $[a(2^k n)] = [a(n)]$. Now, we fix $m \in \mathbb{N}$ and choose $2^k > m$. Since $a(n) \leq a(mn) \leq a(2^k n)$ and $[a(2^k n)] = [a(n)]$ we see $[a(mn)] = [a(n)]$.

(3). It is simple to observe that

$$\pi_{\mathbb{E}}([a(n)]) = \pi_{\mathbb{E}}([a(mn)]) = \limsup_n \frac{\log(a(mn))}{n} = m \limsup_n \frac{\log(a(mn))}{mn} \geq m \pi_{\mathbb{E}}([a(n)]).$$

However, $m \geq 2$ and therefore $\pi_{\mathbb{E}}([a(n)]) = 0$. □

A similar reasoning as the one done in proposition 2.4 and the third property of the previous proposition implies the following result.

Proposition 2.8. *Consider $f : M \rightarrow M$ a dynamical system such that $[a_{f,\mathcal{U}}(n)] \in \mathbb{L}$ for all finite open covering \mathcal{U} . Then:*

- (1) $o(f) \in \overline{\mathbb{L}}$,
(2) and $h(f) = 0$.

3. PROOF OF THEOREM 1

In this section we shall prove theorem 1. For $o = [n]$, the result holds because for any translation in the plane, the compactification f verifies $o(f) = [n]$. For this type of maps, all the mutually singular sets consist of a single set and by theorem 2.3 we conclude.

Now, we are only interested in the case $[n^2] \leq o \leq \sup(\mathbb{P})$. We are going to prove first the result for the particular case $o \in \mathbb{L}$, and then, adapt to obtain the general result ($o \in \overline{\mathbb{L}}$).

3.1. Construction of the map. We are going to recall the construction done in [6], that will provide us a homeomorphism $f : S^2 \rightarrow S^2$. Let us fix $[a(n)] \in \mathbb{L}$ such that $[n^2] < [a(n)] < \sup(\mathbb{P})$. For said $[a(n)]$ there must exist $L \geq 3$ such that $[a(n)] \leq [n^L]$. From now on, L is going to be fixed.

Let us consider P_1, \dots, P_L copies of the plane \mathbb{R}^2 . For each P_i we denote by $O_i = \{(x, y) \in P_i : y > 0\}$ the upper half plane. For each $i = 1, \dots, L - 1$ we are going to select maps $\varphi_i : (0, \infty) \rightarrow \mathbb{R}$ and with them define $\Phi_i : O_i \rightarrow O_{i+1}$ by $\Phi_i(x, y) = (x + \varphi_i(y), y)$.

The first property we want for φ_i is that the limit in 0 is $-\infty$. Now, we define \sim the equivalence relation in $\cup P_i$ generated by $(x, y) \sim \Phi_i(x, y)$ if $(x, y) \in O_i$. The quotient space $P = \cup P_i / \sim$ is a Hausdorff simply connected non-compact surface, and thus is homeomorphic to the plane. Figure 3.1 represents the quotient space P .




FIGURE 3.1. Gluing of the plane P for $L = 4$.

Let $T : \cup P_i \rightarrow \cup P_i$ be defined as the translation $T(x, y) = (x + 1, y)$ on each P_i . The map T commutes with each Φ_i and therefore it defines $F : P \rightarrow P$ an orientation preserving homeomorphism on the plane without fixed points. The map F induces a homeomorphism f in the compactification of the plane $\mathbb{R}^2 \cup \{\infty\} = S^2$ that verifies $\Omega(f) = \{\infty\}$. In particular, we can apply theorem 2.3 to compute its generalized entropy. Figure 3.2 represents the dynamics of f .

To compute $o(f)$ we need first to understand the location of the mutually singular sets. Let us call $\pi : \cup P_i \rightarrow P$ the projection and for each i , define $Y_i = \pi([-1/3, 1/3] \times [0, 1])$ with




FIGURE 3.2. Dynamics of f for $L = 4$.

$[-1/3, 1/3] \times [0, 1] \subset P_i$. For the map constructed f , the family of sets $\{Y_1, \dots, Y_L\}$ is mutually singular. Moreover, any other one, up to iteration of its elements, with diameter small enough, is equivalent to this one. Figure 3.3 represents the location of the sets Y_i .




FIGURE 3.3. Locations of sets Y_i for $L = 4$.

The key observation to be made, is that for a point $(x, y) \in P_i$ such that its projection $\pi(x, y) \in Y_i$ needs for $-\varphi_i(y) > 0$ iterates to reach Y_{i+1} . This means that with the height y we can control the amount of time the point $\pi(x, y) \in Y_1$ is going to need to reach each one of the sets Y_i .

3.2. proof of the theorem for the particular case. Up to this point we followed the construction done [6]. Our next step will be to modify the choice of the maps φ_i done in [6] such that $[c_{f, \mathcal{F}}(n)] = [a(n)]$.

Let us construct φ_i by steps. We shall chose φ_1 such that it is negative, increasing, and, for each positive integer k_1 , assumes the value $-k_1$ on a non trivial interval I_{k_1} . This collection of intervals tends to 0 when k_1 tends to $+\infty$. For convenience, we assume that all these intervals

are included in the interval $(0, 1]$. The number k_1 is going to be our main variable, and we would like to stress that any point of the form $\pi(x, y) \in Y_1$ with $y \in I_{k_1}$ is going to need k_1 iterates of f to reach Y_2 .

Now we look at k_1 iterations of the points in the previous step that are in Y_2 . These points do have the same height y when looked in P_2 . Now, the restriction of φ_2 to I_{k_1} tell us how many iterates are needed for these points in Y_2 to reach Y_3 . We are going to ask that φ_2 restricted to I_{k_1} is continuous, increasing and varies from $-(k_1 + a_2(k_1))$ to $-k_1$ (we postpone for later the choice of the integer $0 < a_2(k_1) \leq k_1$). Moreover, we need that for each integer $-k_2$ in the interval $[-(k_1 + a_2(k_1)), -k_1]$, φ_2 assumes the value $-k_2$ on a non trivial sub-interval $I_{k_1, k_2} \subset I_{k_1}$. Now, any point with height in I_{k_1, k_2} is going to need k_1 iterations to go from Y_1 to Y_2 and k_2 iterations to go from Y_2 to Y_3 . We would like to observe that for k_1 fixed, there are $a_2(k_1)$ possible iterations need to reach from Y_2 to Y_3 . Finally we would like remark that we chose the interval $[-(k_1 + a_2(k_1)), -k_1]$ instead of $[-a_2(k_1), 0]$ because we also need that φ_2 tends to $-\infty$ on 0.

Likewise, we define φ_3 to be increasing on each interval I_{k_1, k_2} and assumes each integer $-k_3$ between $-(3k_1 - k_2 + a_3(k_1))$ and $-(3k_1 - k_2)$ on a non trivial sub-interval I_{k_1, k_2, k_3} of I_{k_1, k_2} . To explain why we choose the interval like this, observe first that $2k_1 \leq k_1 + k_2 \leq 3k_1$. By the choice of the interval, if $0 < a_3(k_1) \leq k_1$, then

$$4k_1 \leq k_1 + k_2 + k_3 \leq 5k_1.$$

This give us the following control: A point in Y_1 with height in I_{k_1} is going to reach Y_2 in k_1 iterates, reach Y_3 in between $2k_1$ to $3k_1$ iterates and Y_4 in between $4k_1$ to $5k_1$ iterates. Similar to the previous step, for k_1 fixed, there are $a_3(k_1)$ possible iterations needed to reach from Y_3 to Y_4 .

We define φ_i inductively until φ_{L-1} . The map φ_{i+1} is chosen such that for k_1, \dots, k_i fixed, it assumes each integer value $-k_{i+1}$ in a sub-interval $I_{k_1, \dots, k_{i+1}}$ of I_{k_1, \dots, k_i} . The value of $-k_{i+1}$ ranges in an interval of length $a_{i+1}(k_1) \leq k_1$ and such that

$$2ik_1 \leq k_1 + \dots + k_{i+1} \leq (2i + 1)k_1.$$

For $i + 1 = L - 1$ we see

$$(2L - 4)k_1 \leq k_1 + \dots + k_{L-1} \leq (2L - 3)k_1.$$

Let us finally address the choice of $a_i(n)$. We choose them such that

$$\left[n \sum_{k_1=1}^n a_2(k_1) \cdots a_{L-1}(k_1) \right] = [a(n)].$$

In order to see why this is possible, consider the case $a_i(k_1) = k_1$ for all i and observe that

$$n \sum_{k_1=1}^n a_2(k_1) \cdots a_{L-1}(k_1) = n \sum_{k_1=1}^n k_1^{L-2},$$

that is a polynomial of degree L on n . Since $[a(n)] \leq [n^L]$, we just lower the values of $a_i(k_1)$ to obtain the order of growth of $a(n)$.

Our task now is to compute $[c_{f, \mathcal{F}}(n)]$.

Consider a word $w \in \mathcal{A}_n(f, \mathcal{F})$ and $x \in S^2$ a point such that w is the coding of the itinerary of x . We call the displacement of w , k units to the right, to the word of length n that codes the itinerary of $f^{-k}(x)$. The displacement of w , k units to the left, is the word of length n that codes the itinerary of $f^k(x)$.

Consider the syndetic set $\mathcal{S} = (2L - 2)\mathbb{N}$. In light of lemma 2.6 if we prove that $d_1 a(n) \leq c_{f, \mathcal{F}}(n) \leq d_2 a(n)$ for $n \in \mathcal{S}$, then $[c_{f, \mathcal{F}}(n)] = [a(n)]$.

Let us consider $b_2 > b_1 > 0$ such that

$$b_1 a(n) \leq n \sum_{k_1=1}^n a_2(k_1) \cdots a_{L-1}(k_1) \leq b_2 a(n) \quad \forall n \in \mathbb{N}.$$

By linear invariance of $[a(n)]$ there exists $c_2 > c_1 > 0$ such that

$$c_1 a(k(2L - 2)) \leq a(k) \leq c_2 a(k(2L - 2)) \quad \forall k \in \mathbb{N}.$$

We fix now some $n \in \mathcal{S}$ and define $k \in \mathbb{N}$ such that $n = k(2L - 2)$. If a word in $\mathcal{A}_n(f, \mathcal{F})$ begins with Y_1 and is associated to $k_1 \leq k$, then all the symbols Y_2, \dots, Y_L appear in w . By our construction, there are $a_2(k_1) \cdots a_{L-1}(k_1)$ of those. Now this words can be displaced at least k to the right creating new unique words in $\mathcal{A}_n(f, \mathcal{F})$. This is true because all symbols Y_i appear before $(2L - 3)k_1 \leq (2L - 3)k$. From this we infer that in $\mathcal{A}_n(f, \mathcal{F})$ there are at least $k \sum_{k_1=1}^k a_2(k_1) \cdots a_{L-1}(k_1)$ distinct words. Then,

$$c_{f, \mathcal{F}}(n) \geq k \sum_{k_1=1}^k a_2(k_1) \cdots a_{L-1}(k_1) \geq b_1 a(k) \geq b_1 c_1 a(n).$$

On the other hand, every word in $\mathcal{A}_n(f, \mathcal{F})$ is obtained from displacing a word that starts with Y_1 . We can displace up to n spaces to the right or $(2L - 2)n$ to the left. We only need to count up to $k_1 = n$ and therefore,

$$c_{f, \mathcal{F}}(n) \leq (2L - 1)n \sum_{k_1=1}^n a_2(k_1) \cdots a_{L-1}(k_1) \leq (2L - 1)c_2 a(n).$$

In conclusion, if $d_1 = b_1 c_1$ and $d_2 = (2L - 1)c_2$, we have proved

$$d_1 a(n) \leq c_{f, \mathcal{F}}(n) \leq d_2 a(n) \quad \forall n \in \mathcal{S}.$$

Thus,

$$[c_{f, \mathcal{F}}(n)] = [a(n)]$$

and with this, we have finished the proof of theorem 1 for the case $o = [a(n)] \in \mathbb{L}$.

3.3. Lemmas of orders of growth. To prove the general case of theorem 1 the following lemma will simplify our reasoning.

Lemma 3.1. *Let $\Gamma \subset \mathbb{O}$ a countable subset of orders of growth, then there exists a countable and ordered subset $\hat{\Gamma} \subset \mathbb{O}$ such that $\sup(A) = \sup(B)$. Moreover, if $\Gamma \subset \mathbb{B}$ or $\Gamma \subset \mathbb{L}$, then so does $\hat{\Gamma}$.*

In order to prove lemma 3.1 the following is necessary.

Lemma 3.2. *Suppose that $[a(n)], [b(n)] \in \mathbb{O}$. Then:*

$$(1) \sup\{[a(n)], [b(n)]\} = [\max\{a(n), b(n)\}].$$

(2) *If $[a(n)], [b(n)] \in \mathbb{B}$ or $[a(n)], [b(n)] \in \mathbb{L}$, so does $\sup\{[a(n)], [b(n)]\}$.*

Proof. (1) We know that $[a(n)] \leq [\max\{a(n), b(n)\}]$ and $[b(n)] \leq [\max\{a(n), b(n)\}]$, then $\sup\{[a(n)], [b(n)]\} \leq [\max\{a(n), b(n)\}]$. But, if we suppose

$$\sup\{[a(n)], [b(n)]\} < [\max\{a(n), b(n)\}],$$

then there exists $[c(n)] \in \mathbb{O}$ such that

$$\sup\{[a(n)], [b(n)]\} < [c(n)] < [\max\{a(n), b(n)\}].$$

By definition, we know $[a(n)] \leq \sup\{[a(n)], [b(n)]\}$, and then $[a(n)] \leq [c(n)]$. Analogously $[b(n)] \leq [c(n)]$. Thus, there must exist constants $d_1, d_2 > 0$ such that $a(n) \leq d_1 c(n)$ and $b(n) \leq d_2 c(n)$. Let $d = \max\{d_1, d_2\}$, and observe $\max\{a(n), b(n)\} \leq d c(n)$, which implies $[\max\{a(n), b(n)\}] \leq [c(n)]$, a contradiction. Therefore,

$$\sup\{[a(n)], [b(n)]\} = [\max\{a(n), b(n)\}],$$

as we wanted.

(2) For elements in \mathbb{B} is an immediate conclusion from part (1) of this lemma. For elements in \mathbb{L} is necessary to use point (2) of proposition 2.7. \square

Proof of lemma 3.1. Let $\Gamma = \{[b_1(n)], [b_2(n)], \dots, [b_k(n)], \dots\} \subset \mathbb{O}$, we construct the subset $\hat{\Gamma}$ as follows:

$$\begin{aligned} [a_1(n)] &= [b_1(n)], \\ [a_2(n)] &= \sup\{[b_1(n)], [b_2(n)]\} = [\max\{b_1(n), b_2(n)\}], \\ [a_3(n)] &= \sup\{[a_1(n)], [b_3(n)]\} = \sup\{\sup\{[b_1(n)], [b_2(n)]\}, [b_3(n)]\} \\ &= \sup\{[b_1(n)], [b_2(n)], [b_3(n)]\} = [\max\{b_1(n), b_2(n), b_3(n)\}], \\ &\vdots \\ [a_k(n)] &= \sup\{[a_{k-1}(n)], [b_k(n)]\} = \dots = [\max\{b_1(n), b_2(n), \dots, b_k(n)\}], \\ &\vdots \end{aligned}$$

The previous identities hold by lemma 3.2 and an inductive argument. A is clearly a countable set and it is easy see that it is an ordered set, in fact

$$\begin{aligned} [a_1(n)] &= \sup\{[b_1(n)], [b_2(n)]\} \leq \sup\{[b_1(n)], [b_2(n)], [b_3(n)]\} \\ &= [a_2(n)] \leq \dots \leq [a_k(n)] \leq [a_{k+1}(n)] \leq \dots \end{aligned}$$

If $\Gamma \subset \mathbb{B}$ or $\Gamma \subset \mathbb{L}$, by the second item of 3.2 so does $\hat{\Gamma}$.

We shall prove now that $\sup(\hat{\Gamma}) = \sup(\Gamma)$. Since $\sup(\hat{\Gamma}) \geq [a_k(n)] \geq [b_k(n)]$, we deduce $\sup(\hat{\Gamma}) \geq \sup(\Gamma)$. On the other hand, $\sup(\Gamma) \geq [b_k(n)]$, for all $k \in \mathbb{N}$, and by definition of $[a_k(n)]$ we infer $\sup(\Gamma) \geq [a_k(n)]$ for all $k \in \mathbb{N}$. Therefore, $\sup(\Gamma) \leq \sup(\hat{\Gamma})$. \square

3.4. Proof of the general case. Let us consider $\Gamma \subset \mathbb{L}$ countable such that $[n^2] \leq o = \sup(\Gamma) \leq \sup(\mathbb{P})$. We want to construct f such that $o(f) = o$.

By lemma 3.1 we might suppose that Γ is ordered. Moreover, if $\Gamma = \{[a_k(n)] : k \in \mathbb{N}\}$, we may also assume that $[a_1(n)] \geq [n^2]$.

We construct first $f_1 \in \mathcal{H}$ such that $o(f_1) = [a_1(n)]$ using the technique of subsection 3.2. We shall call ∞ to the only fixed point of f_1 and D_1 as the disk associated to the lower semi-plane in P_1 . We also call S_1 to the closure of its complement. The dynamics constructed in subsection 3.2 lay entirely in the region S_1 . Therefore, $o(f_1|_{S_1}) = [a_1(n)]$.




FIGURE 3.4. $f_2 : S^2 \rightarrow S^2$.

We now define $f_2 \in \mathcal{H}$ such that $f_2|_{S_1} = f_1|_{S_1}$ and $f_2|_{D_1}$ verifies $o(f_2|_{D_1}) = [a_2(n)]$. The construction is made in an analogous way by adding new planes P_i^2 and using the lower semi-plane of P_1 . Figure 3.4 represents the construction of f_2 .

We choose one of the new planes, and call D_2 to the region associated to its lower half-plane. We call S_2 to the closure of the complement of D_2 and observe that dynamics created so far lay entirely in S_2 . In particular,

$$o(f_2|_{S_2}) = o(f_2) = \sup\{o(f_2|_{S_1}), o(f_2|_{D_1})\} = \sup\{[a_1(n)], [a_2(n)]\} = [a_2(n)].$$

Inductively, we define a family of regions D_k and $S_k = \overline{D_k^c}$ where D_k is a disk, contains ∞ in its border, $D_{k+1} \subset D_k$, and we may also ask that $\text{diam}(D_k) \rightarrow 0$. By induction we also have a family of maps $f_k \in \mathcal{H}$ such that:

- $f_k(S_i) = S_i$ for all $i \leq k$.
- $f_{k+1}|_{S_k} = f_k|_{S_k}$.
- $o(f_k) = o(f_k|_{S_k}) = [a_k(n)]$.

This family of dynamical systems in \mathcal{H} , define a unique map $f \in \mathcal{H}$ with $f(x) = f_k(x)$ if $x \in S_k$.

Let us check that $o(f) = \sup(\Gamma)$. We observe first that

$$o(f) \geq o(f|_{S_k}) = o(f_{k|S_k}) = [a_k(n)],$$

thus $o(f) \geq \sup(\Gamma)$. To see the other inequality, we chose $\varepsilon > 0$ and k such that $\text{diam}(D_k) \leq \varepsilon/2$. Since $f(D_k) = D_k$, for any $x_k \in D_k$, the dynamical ball $B(x_k, n, \varepsilon)$ contains D_k for every n . If G_n is an (n, ε) generator set of $f|_{S_k}$ with $g_{f|_{S_k}, \varepsilon}(n) = \#G_n$, then $G_n \cup \{x_k\}$ is an (n, ε) -generator set of f . Therefore, $g_{f, \varepsilon}(n) \leq g_{f|_{S_k}, \varepsilon}(n) + 1$ and then

$$[g_{f, \varepsilon}(n)] \leq [g_{f|_{S_k}, \varepsilon}(n)] \leq o(f|_{S_k}) = [a_k(n)] \leq \sup(\Gamma).$$

From this, we conclude $o(f) \leq \sup(\Gamma)$. This proves $o(f) = \sup(\Gamma)$ and finishes the proof of theorem 1.

REFERENCES

- [1] A. K. ADAM KANIGOWSKI AND D. WEI, *Survey on entropy-type invariants of sub-exponential growth in dynamical systems*, arXiv:2004.04655, (2020).
- [2] J. BOCHI, A. KATOK, AND F. R. HERTZ, *Flexibility of Lyapunov exponents*, arXiv:1908.07891, (2019).
- [3] J. CORREA AND H. DE PAULA, *Polynomial entropy of Morse-Smale diffeomorphisms on surfaces*, Bull. Sci. Math., 182 (2023).
- [4] J. CORREA AND E. R. PUJALS, *Orders of growth and generalized entropy*, J. Inst. Math. Jussieu, (2021), p. 1–33.
- [5] S. GALATOLO, *Global and local complexity in weakly chaotic dynamical systems*, Discrete Contin. Dyn. Syst., 9 (2003), pp. 1607–1624.
- [6] L. HAUSEUX AND F. LE ROUX, *Entropie polynomiale des homéomorphismes de Brouwer*, Annales Henri Lebesgue, 2 (2019), pp. 39–57.
- [7] A. KATOK AND J. P. THOUVENOT, *Slow entropy type invariants and smooth realization of commuting measure-preserving transformations*, Ann. Inst. Henri Poincaré Probab. Stat., 33 (1997), pp. 323–338.
- [8] J. P. MARCO, *Polynomial entropies and integrable hamiltonian systems*, Regul. Chaotic Dyn., 18 (2013), pp. 623–655.

UNIVERSIDADE FEDERAL DE MINAS GERAIS

Email address: jcorrea@mat.ufmg.br

CENTRO FEDERAL DE EDUCAÇÃO TECNOLÓGICA DE MINAS GERAIS

Email address: hellenlimadepaula@ufmg.br