

On the self-similarity of the norm one group of p -adic division algebras

Francesco Nosedà

Illir Snopce

Abstract

Let p be a prime, K a finite extension of \mathbb{Q}_p , D a finite dimensional central division K -algebra, and $SL_1(D)$ the group of elements of D of reduced norm 1. When $p \geq \dim(SL_1(D))$, we provide an infinite family of congruence subgroups of $SL_1(D)$ that do not admit self-similar actions on regular rooted p -ary trees. The proof requires results on \mathbb{Z}_p -Lie lattices that also lead to the classification of the torsion-free p -adic analytic pro- p groups G of dimension less than p with the property that all the nontrivial closed subgroups of G admit a self-similar action on a p -ary tree. As a consequence we obtain that a nontrivial torsion-free p -adic analytic pro- p group G of dimension less than p is isomorphic to the maximal pro- p Galois group of a field that contains a primitive p -th root of unity if and only if all the nontrivial closed subgroups of G admit a self-similar action on a regular rooted p -ary tree.

1 Introduction

Let K be a p -adic field, that is, a finite field extension of \mathbb{Q}_p for some prime p . For D a finite dimensional central division algebra over K , the group $SL_1(D)$ of elements of D of reduced norm 1 is algebraic, in the sense that it is the group of K -rational points of an algebraic group, and it has the structure of compact p -adic analytic group. The groups of type $SL_1(D)$ and their associated \mathbb{Q}_p -Lie algebras $sl_1(D)$ play an important role in the classification of, respectively, simple algebraic groups and simple Lie algebras defined over p -adic fields. More precisely, if G is the group of K -rational points of an anisotropic absolutely simple simply connected algebraic group defined over K then G is isomorphic to the group $SL_1(D)$ for some D [Bru67, Proposition 6]. Analogously, if \mathcal{L} is an anisotropic absolutely simple K -Lie algebra then \mathcal{L} is isomorphic to $sl_1(D)$ for some D ; see, for instance, [Sch14, Theorem (2), page 6]. Due to their prominent role, these objects have been studied thoroughly; see, for instance, [Rie70, Pra88, Ers07, Ers22, Sch14]; see also [Klo03]. Moreover, we point out that the groups $SL_1(D)$ have strong connections with the Morava stabilizer groups in stable homotopy theory; see, for instance, [Rav86, Appendix 2] and [Hen98]. The main goal of the present paper is to study the groups $SL_1(D)$ from the point of view of their self-similar actions on regular rooted trees.

Groups that admit a faithful self-similar action on some regular rooted tree form an interesting class that contains many important examples such as the Grigorchuk 2-group [Gri80] and the Gupta-Sidki p -groups [Gup83]. More recently, there has been an intensive study on the self-similar actions of other families of groups including iterated monodromy groups [Nek05], finitely

Mathematics Subject Classification (2020): primary 20E08, 20E18; secondary 11E95, 11S20, 16K20, 17B20.

Key words: self-similar group, division algebra over p -adic field, p -adic analytic group, pro- p group, p -adic Lie lattice, maximal pro- p Galois group.

generated nilpotent groups [Ber07], arithmetic groups [Kap12], finite p -groups [Sun11, Bab20], groups of type FP_n [Koc20], and p -adic analytic pro- p groups [Nos19, Nos22a, Nos22b].

We say that a group G is self-similar of index d , where $d \geq 1$ is an integer, if G admits a faithful self-similar action on a regular rooted d -ary tree in such a way that the action is transitive on the first level. We say that G is self-similar if it is self-similar of some index d ; see, for instance, [Nek05] for a general treatment of self-similar actions. It is known that a group is self-similar of index d if and only if it admits a simple virtual endomorphism of index d . A virtual endomorphism of G of index D is a group homomorphism $\varphi : D \rightarrow G$, where $D \leq G$ is a subgroup of index d ; the virtual endomorphism φ is said to be simple if there are no nontrivial normal subgroups N of G that are φ -invariant, that is, such that $N \subseteq D$ and $\varphi(N) \subseteq N$.

Throughout the paper we let p be a prime. In [Nos19] we initiated the study of self-similar actions of p -adic analytic pro- p groups by exploiting the Lazard's correspondence, an isomorphism of categories between the category of saturable finitely generated pro- p groups and the category of saturable \mathbb{Z}_p -Lie lattices; see [Laz65, IV (3.2.6)] and, for instance, [Gon07] and [Gon09] for more details. We recall that the Lazard's correspondence restricts to an isomorphism between the category of uniform pro- p groups and the category of powerful \mathbb{Z}_p -Lie lattices; see [Dix99]. A \mathbb{Z}_p -Lie lattice is a finitely generated free \mathbb{Z}_p -module endowed with a structure of Lie algebra over \mathbb{Z}_p . Let L be a \mathbb{Z}_p -Lie lattice and $k \in \mathbb{N}$. A virtual endomorphism of L of index p^k is an algebra homomorphism $\varphi : M \rightarrow L$ where M is a subalgebra of L of index p^k , where the index is taken with respect to the additive group structure. An ideal I of L is said to be φ -invariant if $I \subseteq M$ and $\varphi(I) \subseteq I$. We say that a virtual endomorphism φ is simple if there are no nonzero ideals I of L that are φ -invariant. Finally, we say that L is self-similar of index p^k if there exists a simple virtual endomorphism of L of index p^k , and we say that L is self-similar if it is self-similar of index p^k for some k .

We introduce some more notation in order to state our results. As above, let D be a finite dimensional central division K -algebra, where K is a finite field extension of \mathbb{Q}_p . Let d and f be the degree and inertia degree of K/\mathbb{Q}_p , respectively, and let n be the index of D over K , that is $\dim_K(D) = n^2$. Let \mathcal{O}_K be the integral closure of \mathbb{Z}_p in K , Δ be the unique maximal \mathcal{O}_K -order in D , and $sl_1(\Delta)$ be the set of elements of Δ of reduced trace equal to 0. Observe that $sl_1(\Delta)$ is a \mathbb{Z}_p -Lie lattice of dimension $d(n^2 - 1)$.

The following two theorems are proved in Section 3.

Theorem A *Assume that $f \geq 2$ holds or both $n \geq 3$ and $(p, n) \neq (3, 3)$ hold. Let N be a maximal proper \mathbb{Z}_p -submodule of $sl_1(\Delta)$. Then $[N, N] = [sl_1(\Delta), sl_1(\Delta)]$.*

Theorem B *Assume that $f \geq 2$ or $(p, n) \notin \{(2, 2), (3, 3)\}$, and let $m \in \mathbb{N}$. Then $p^m sl_1(\Delta)$ is not self-similar of index p .*

Let $SL_1(D)$ be the set of elements of D (equivalently, of Δ) of reduced norm equal to 1, and observe that $SL_1(D)$ is a compact p -adic analytic group of dimension $d(n^2 - 1)$. Let \mathfrak{p} be the unique maximal two-sided ideal of Δ . For $l \in \mathbb{N}$, define $SL_1^l(D) = SL_1(D) \cap (1 + \mathfrak{p}^l)$ and $sl_1^l(\Delta) = sl_1(\Delta) \cap \mathfrak{p}^l$. For $p \geq 5$ and $l \geq ne$, where e is the ramification index of K/\mathbb{Q}_p , $SL_1^l(D)$ is a uniform pro- p open subgroup of $SL_1(D)$, and the powerful \mathbb{Z}_p -Lie lattice associated with $SL_1^l(D)$ is $sl_1^l(\Delta)$ [Ers22, Proposition 4.4]. In particular, for $p \geq 5$ and all integers $m \geq 1$, the \mathbb{Z}_p -Lie lattice associated with $SL_1^{mne}(D)$ is $p^m sl_1(\Delta)$.

The following theorem, proved in Section 5, provides a family of unsolvable torsion free p -adic analytic pro- p groups of arbitrarily high dimension (for sufficiently large p) that do not

admit a self-similar action on a p -ary tree. We point out that, to the extent of our knowledge, such families were only known for solvable groups (cf. [Nos22a, Proposition 2.25] and [Nos19, Proposition A]).

Theorem C *Assume that $p \geq d(n^2 - 1)$, and let $m \geq 1$ be an integer. Then $SL_1^{mne}(D)$ is a uniform pro- p group and the following holds.*

- (1) $SL_1^{mne}(D)$ is not self-similar of index p .
- (2) Assume that $n \neq 2$, and let H be a subgroup of $SL_1^{mne}(D)$ of index p . Then $[H, H] = [SL_1^{mne}(D), SL_1^{mne}(D)]$.

Theorem C, together with [Nos19, Theorem D] and [Nos22b, Theorem 1], provides evidence for the following generalization of [Nos19, Conjecture E].

Conjecture D *Assume that $n \geq 2$ and $p \geq d(n^2 - 1)$. Then no open subgroup of $SL_1^{ne}(D)$ is self-similar.*

We recall that in order to prove the conjecture it is enough to prove that $SL_1^{ne}(D)$ is not self-similar; see [Nos19, Corollary 1.4].

A finitely generated pro- p group G is said to be strongly hereditarily self-similar of index p if G and all of its nontrivial finitely generated closed subgroups are self-similar of index p . The groups that are strongly hereditarily self-similar of index p were classified inside the class of solvable torsion-free p -adic analytic pro- p groups of dimension strictly less than p [Nos22a, Theorem A]. In an analogous fashion to what is done for groups, we say that a \mathbb{Z}_p -Lie lattice L is strongly hereditarily self-similar of index p if L and all of its nonzero subalgebras are self-similar of index p . For $p \geq 3$, the Lie lattices that are strongly hereditarily self-similar of index p were classified inside the class of solvable \mathbb{Z}_p -Lie lattices, without restriction on the dimension [Nos22a, Theorem 2.34]. In this paper we show that the aforementioned classifications work over much wider classes, by not restricting to solvable groups and Lie lattices.

Theorem E *Assume $p \geq 5$, let L be a \mathbb{Z}_p -Lie lattice, and define $d = \dim(L)$. Then L is strongly hereditarily self-similar of index p if and only if L is isomorphic to one of the following Lie lattices:*

- (1) the abelian Lie lattice \mathbb{Z}_p^d , with $d \geq 1$;
- (2) the metabelian Lie lattice $\mathbb{Z}_p \times \mathbb{Z}_p^{d-1}$, where the canonical generator of \mathbb{Z}_p acts on \mathbb{Z}_p^{d-1} by multiplication by the scalar p^s , with $d \geq 2$ and $s \in \mathbb{N}$.

The following theorem proves [Nos22a, Conjecture E].

Theorem F *Let p be a prime, and let G be a torsion-free p -adic analytic pro- p group. Define $d = \dim(G)$, and assume that $p > d$. Then G is strongly hereditarily self-similar of index p if and only if G is isomorphic to one of the following groups:*

- (1) the abelian pro- p group \mathbb{Z}_p^d , with $d \geq 1$;

- (2) the metabelian pro- p group $\mathbb{Z}_p \rtimes \mathbb{Z}_p^{d-1}$, where the canonical generator of \mathbb{Z}_p acts on \mathbb{Z}_p^{d-1} by multiplication by the scalar $1 + p^s$, with $d \geq 2$ and integer $s \geq 1$.

The proof of Theorem [E](#) is given in Section [4](#), while the proof of Theorem [F](#) is given in Section [5](#) and relies on Theorem [E](#) and [[Nos22a](#), Proposition 3.1], the latter being the key to exploit Lie methods. We point out that, for the proof of Theorem [E](#), one first applies classical results on Lie algebras over fields of characteristic zero to reduce the problem to the study of the \mathbb{Z}_p -Lie lattices associated with the finite dimensional division \mathbb{Q}_p -algebras introduced above; then, one can apply Theorem [B](#).

The groups that appear in Theorem [F](#) have been object of study in the past few decades, since they are connected with several topics, such as, maximal pro- p Galois groups of fields [[War92](#)], pro- p groups that have constant generating number on open subgroups [[Klo11](#)], Bloch-Kato pro- p groups [[Qua14](#)], hereditarily uniform pro- p groups [[Klo14](#)], and Frattini-injective pro- p groups [[Sno20](#)]. With the aid of Theorem [F](#), we can prove a rather strong generalization of [[Nos22a](#), Theorem 3.9]. The proof of Theorem [G](#) is given in Section [5](#).

Theorem [G](#) *Let G be a nontrivial torsion-free p -adic analytic pro- p group, and assume that $p > \dim(G)$. Then the following are equivalent.*

- (1) G is strongly hereditarily self-similar of index p .
- (2) G is isomorphic to the maximal pro- p Galois group of some field that contains a primitive p -th root of unity.
- (3) G has constant generating number on open subgroups.
- (4) G is a Bloch-Kato pro- p group.
- (5) G is a hereditarily uniform pro- p group.
- (6) G is a Frattini-injective pro- p group.

Notation. The set $\mathbb{N} = \{0, 1, 2, \dots\}$ of natural numbers is assumed to contain 0. Throughout the paper we let p be a prime, and we denote by \mathbb{Z}_p the ring of p -adic integers, by \mathbb{Q}_p the field of p -adic numbers, by \mathbb{F}_p the residue field of \mathbb{Z}_p , and by $v_p : \mathbb{Q}_p \rightarrow \mathbb{Z} \cup \{\infty\}$ the p -adic valuation. The expression $x \equiv_a y$ means that x is equivalent to y modulo a , where the context where the equivalence is applied will always be clear. The characteristic of a field K is denoted by $\text{ch}(K)$. When R is a commutative ring, we denote by R^* the associated multiplicative group; moreover, when L is an R -Lie algebra and M is an R -submodule of L , $[M, M]$ denotes the set of finite sums of brackets of elements of M , and we observe that $[M, M]$ coincides with the R -submodule of L generated by such brackets.

Acknowledgments. The authors thank Gopal Prasad for feedback on a question on division algebras. The first author also thanks José J. Ramón Marí for a technical suggestion.

2 Preliminary results

We collect both general and technical facts that will be used in Section [3](#).

2.1 Lemmas in cyclic extensions

For an arbitrary field K , let F/K be a finite cyclic Galois field extension, let n be the degree of F/K , and let ϑ be a generator of $\text{Gal}(F/K)$. Let $T : F \rightarrow K$ be the trace map, and let $F^0 = \ker(T)$. Since F/K is finite and separable, the trace map is surjective, and the trace form, $(\alpha, \beta) \in F \times F \mapsto T(\alpha\beta)$, is nondegenerate. In particular, F^0 is a K -vector subspace of F of dimension $n - 1$. We will prove some results that will be applied in Section 3.2 to the residue field extension κ_F/κ_K introduced at the beginning of Section 3. The range of a function f will be denoted by $\text{ran}(f)$.

Lemma 2.1 *Let $0 < j < n$, and assume that $(\text{ch}(K), n) \neq (2, 2)$. Then there exists $\alpha \in F^0$ such that $\vartheta^j(\alpha) \neq \alpha$.*

PROOF: Observe that $\vartheta^j \neq \text{id}$. For $\text{ch}(K) \nmid n$ the lemma is clear, since $F = K \oplus F^0$ and ϑ^j fixes the elements of K , so that ϑ^j does not fix all the elements of F^0 . Below we will prove the lemma for $n \geq 3$. Since, under the assumptions, $\text{ch}(K) \mid n$ implies $n \geq 3$, the lemma is proven in all the cases.

Assume that $n \geq 3$. Let $S = \vartheta^j - \text{id}$, a K -linear map $F \rightarrow F$. Let $F_1 = \ker(S)$, and observe that F_1 is the fixed field of ϑ^j . Hence, F_1 is an intermediate field of the extension F/K , and $F_1 \neq F$. Assume by contradiction that ϑ^j fixes the elements of F^0 , in other words, $F^0 \subseteq F_1$. Then $F^0 = F_1$, which implies that $n - 1$ divides n , a contradiction since $n \geq 3$. \square

Lemma 2.2 *Let $\alpha \in F$ and $\alpha \neq 0$. Then $\{\alpha\vartheta(\beta) - \beta\vartheta^{-1}(\alpha) : \beta \in F\} = F^0$.*

PROOF: Let $V = \{\alpha\vartheta(\beta) - \beta\vartheta^{-1}(\alpha) : \beta \in F\}$. Since ϑ preserves the trace, one can see that $V \subseteq F^0$. Let $f : F \rightarrow F$ be the K -linear map defined by $f(\beta) = \alpha\vartheta(\beta) - \beta\vartheta^{-1}(\alpha)$, and observe that $V = \text{ran}(f)$. Since the range of f has dimension less than n , there exists $\beta_0 \in \ker(f)$ such that $\beta_0 \neq 0$. An easy computation shows that, for all $\beta \in F$, $\beta \in \ker(f)$ if and only if $\vartheta(\beta/\beta_0) = \beta/\beta_0$. Since the fixed field of ϑ is K , we have $\ker(f) = \beta_0 K$. Hence, V has dimension $n - 1$ over K , and the lemma follows. \square

Remark 2.3 By applying Lemma 2.2 with the generator ϑ^{-1} in place of ϑ , one sees that if $\beta \in F$ and $\beta \neq 0$ then $\{\alpha\vartheta(\beta) - \beta\vartheta^{-1}(\alpha) : \alpha \in F\} = F^0$.

The following is the additive form Hilbert's Theorem 90.

Lemma 2.4 *We have $\{\vartheta(\alpha) - \alpha : \alpha \in F\} = F^0$.*

PROOF: The K -linear map $\vartheta - \text{id} : F \rightarrow F$ has kernel of dimension 1 and range included in F^0 . \square

Lemma 2.5 *Let k be an integer such that $k + 1 \not\equiv_n 0$. Then $\text{span}_K(\alpha\vartheta(\beta) - \beta\vartheta^k(\alpha) : \alpha, \beta \in F) = F$.*

PROOF: Define $V = \text{span}_K(\alpha\vartheta(\beta) - \beta\vartheta^k(\alpha) : \alpha, \beta \in F)$, and observe that $\{\vartheta(\beta) - \beta : \beta \in F\} = F^0$, by Lemma 2.4. By taking $\alpha = 1$ one sees that $F^0 \subseteq V$. It is enough to show that there exists $\gamma \in V$ such that $T(\gamma) \neq 0$. Assume by contradiction that $T(\alpha\vartheta(\beta) - \beta\vartheta^k(\alpha)) = 0$ for all $\alpha, \beta \in F$. By substituting α with $\vartheta(\alpha)$, we get $T(\vartheta(\alpha\beta) - \beta\vartheta^{k+1}(\alpha)) = 0$ for all α, β . Hence, for all α, β , we have $T(\alpha\beta) = T(\vartheta(\alpha\beta)) = T(\beta\vartheta^{k+1}(\alpha))$, so that $T([\vartheta^{k+1}(\alpha) - \alpha]\beta) = 0$.

Since the trace form is nondegenerate, we get $\vartheta^{k+1}(\alpha) = \alpha$ for all $\alpha \in F$. This is a contradiction since $\vartheta^{k+1} \neq \text{id}$. \square

For the applications of Section 3.2 we need a basis of F over K with the properties given in the following lemma.

Lemma 2.6 *There exists a basis $(\tau_i)_{0 \leq i < n}$ of F over K such that the following holds.*

- (1) $(\tau_i)_{1 \leq i < n}$ is a basis of F^0 over K .
- (2) If $\text{ch}(K) \nmid n$ then $\tau_0 = 1$.
- (3) If $\text{ch}(K) \mid n$ then $\tau_1 = 1$.
- (4) If $n \geq 2$ and $(\text{ch}(K), n) \neq (2, 2)$ then $\tau_{n-1} \notin K$.

PROOF: There exists a basis $(\gamma_i)_{0 \leq i < n}$ of F over K such that $\gamma_0 = 1$. There exists i_0 such that $T(\gamma_{i_0}) \neq 0$. In case $\text{ch}(K) \nmid n$, take $i_0 = 0$. We define

$$\begin{aligned} \tau_0 &= \gamma_{i_0} \\ \tau_i &= \gamma_{i-1} - T(\gamma_{i-1})T(\gamma_{i_0})^{-1}\gamma_{i_0} & \text{if } 0 < i \leq i_0 \\ \tau_i &= \gamma_i - T(\gamma_i)T(\gamma_{i_0})^{-1}\gamma_{i_0} & \text{if } i > i_0, \end{aligned}$$

and one can see that $(\tau_i)_{0 \leq i < n}$ satisfies the desired properties. \square

Remark 2.7 Let $(\tau_i)_{0 \leq i < n}$ be a basis of F over K with the properties stated in Lemma 2.6. Define $\xi_i = \tau_i - \vartheta(\tau_i)$ for $0 \leq i < n$, and recall Lemma 2.4. Define $i_0 = 0$ when $\text{ch}(K) \nmid n$, and $i_0 = 1$ when $\text{ch}(K) \mid n$. Observe that $\xi_{i_0} = 0$. Then the set $(\xi_i)_{0 \leq i < n, i \neq i_0}$ is linearly independent over K .

Remark 2.8 Let V be a K -vector subspace of F of dimension $n - 1$. Observe that if $\alpha \in F$ and $\alpha V = V$ then $\alpha \in K$. In fact, V is a vector space over the subfield $K(\alpha)$ generated by α over K , and we have $\dim_K(V) = \dim_{K(\alpha)}(V) \cdot [K(\alpha) : K]$, from which we get $[K(\alpha) : K] = 1$. Now, let (ξ_0, ξ_1) be an ordered set of elements of F that is linearly independent over K . Since $\xi_0/\xi_1 \notin K$, by the above observation we have $\xi_0 V \neq \xi_1 V$. Hence, $\xi_0 V + \xi_1 V = F$.

2.2 Lemmas over discrete valuation rings

This section is divided into three parts. In the first, we discuss the notion of just-infinite Lie-lattice; in the second, we introduce the convenient terminology of relative invariant exponents and s -invariants; and in the third, we introduce a notation for the parametrization of maximal proper submodules of a lattice.

Let R be a commutative ring. An R -lattice is a finitely generated free R -module. When an R -lattice is endowed with the structure of an R -Lie algebra, we call it an R -Lie lattice. For our purposes, even when proving or recalling general properties, it is natural to work over a principal ideal domain, so that a submodule of a lattice is a lattice. So, we assume that R is a principal ideal domain, and we denote by F the field of fractions of R . When V is a finite dimensional F -vector space, a full R -lattice in V is a finitely generated R -submodule of V whose span over F is V . Observe that, since R is a principal ideal domain, a full R -lattice is an R -lattice. Finally, if L is an R -lattice then one can identify L with a full R -lattice in $L \otimes_R F$.

For the first part of the section, we say that an R -Lie lattice L is just infinite if it is nonzero and, for all nonzero ideals I of L , we have $\dim_R(I) = \dim_R(L)$. Also, L is said to be hereditarily just infinite if all subalgebras M of L with $\dim_R(M) = \dim_R(L)$ are just infinite.

Remark 2.9 Let L be a \mathbb{Z}_p -lattice and M be a \mathbb{Z}_p -submodule of L . Then $\dim_{\mathbb{Z}_p}(M) = \dim_{\mathbb{Z}_p}(L)$ if and only if $[L : M]$ is finite, where the index is taken with respect to the additive group structure. Also, M is a maximal proper \mathbb{Z}_p -submodule of L if and only if $[L : M] = p$.

Lemma 2.10 Let L be an R -Lie lattice, and let $\mathcal{L} := L \otimes_R F$ be the associated F -Lie algebra. Then the following are equivalent.

- (1) L is just infinite.
- (2) \mathcal{L} is nonzero and it has no nonzero proper ideals.
- (3) L is hereditarily just infinite.

PROOF: By identifying $L \subseteq \mathcal{L}$, one can prove that $\mathcal{I} = \mathcal{L}$ for all nonzero ideals \mathcal{I} of \mathcal{L} if and only if $\dim_R(I) = \dim_R(L)$ for all nonzero ideals I of L . This proves the equivalence between (1) and (2). The proof may be completed after observing that if M is an R -submodule of L such that $\dim_R(M) = \dim_R(L)$ then the F -Lie algebras $M \otimes_R F$ and \mathcal{L} may be identified. \square

The following lemma is a version of [Nos22b, Lemma 10].

Lemma 2.11 Let L be a just-infinite \mathbb{Z}_p -Lie lattice, and let φ be a virtual endomorphism of L that is not injective. Then φ is not simple.

Lemma 2.12 Let L be a just-infinite \mathbb{Z}_p -Lie lattice with $\dim_{\mathbb{Z}_p}(L) \geq 2$, and let $k \in \mathbb{N}$. Assume that for all \mathbb{Z}_p -submodules M of L of index p^k we have $[M, M] = [L, L]$. Then $p^m L$ is not self-similar of index p^k for all $m \in \mathbb{N}$.

PROOF: Let $m \in \mathbb{N}$ and $\varphi : M \rightarrow p^m L$ be a virtual endomorphism of $p^m L$ of index p^k . We have to show that φ is not simple. Observe that $p^m L$ is just infinite (Lemma 2.10) and $\dim_{\mathbb{Z}_p}(p^m L) \geq 2$; as a consequence, $p^m L$ is not abelian. By Lemma 2.11, we can assume that φ is injective. Let $M' = \varphi(M)$, so that φ induces an isomorphism of \mathbb{Z}_p -Lie algebras $M \simeq M'$. By [Nos21, Corollary 8], M' has index p^k in $p^m L$. Hence, $p^{-m} M$ and $p^{-m} M'$ are submodules of L of index p^k . By assumption, $[p^{-m} M, p^{-m} M] = [L, L] = [p^{-m} M', p^{-m} M']$. Hence, $[M, M] = [p^m L, p^m L] = [M', M']$. It follows that $[p^m L, p^m L]$ is a nonzero ideal of $p^m L$ that is φ -invariant, and we conclude that φ is not simple. \square

For the second part of the section, we assume that R is a discrete valuation ring, and we let $\varrho \in R$ be a uniformizing parameter. For the basic properties of the Hermite and Smith canonical forms, the reader may consult, for instance, [Adk92, Section 5.2]. Recall that F denotes the field of fractions of R .

Let V be a finite dimensional F -vector space, and let $n = \dim_F(V)$. Also, let L and M be two full R -lattices in V . We will define the invariant exponents $s_R(L, M)$ of M relative to L over R as a multiset of integers. We abuse the notation by denoting multisets with the same notation as for ordered sets, for instance (s_0, \dots, s_{n-1}) , possibly adding exponents to indicate multiplicities, for instance $(s_0^{f_0}, \dots, s_{n-1}^{f_{n-1}})$. The definition of $s_R(L, M)$ goes as follows. There exist a basis (e_0, \dots, e_{n-1}) of L over R and $s_0, \dots, s_{n-1} \in \mathbb{Z}$ such that $(\varrho^{s_0} e_0, \dots, \varrho^{s_{n-1}} e_{n-1})$ is a

basis of M over R . We define $s_R(L, M) = (s_0, \dots, s_{n-1})$, where one can prove that the multiset (s_0, \dots, s_{n-1}) only depends on the pair (L, M) and not on the chosen basis.

Let V be endowed with the structure of F -Lie algebra, and assume that $[V, V] = V$. Let L be a full R -lattice in V . Then $[L, L]$ is a full R -lattice in V . We define the s -invariants $s_R(L)$ of L over R to be the invariant exponents $s_R(L, [L, L])$ of $[L, L]$ relative to L over R .

For the third part of the section, we keep the assumptions of the second part, and we introduce some notation for the classification of maximal proper submodules. Let L be an R -lattice, and let $(x_\lambda)_{0 \leq \lambda < n}$ be a basis of L over R , where $n = \dim_R(L)$. We consider pairs (μ, b) where μ is an integer with $0 \leq \mu < n$ and $b = (b_\lambda)_{\lambda \neq \mu}$ is a sequence of elements of R . With any such pair we associate a maximal proper R -submodule $N_{\mu, b}$ of L defined to be the span over R of (y_λ) , where

$$\begin{aligned} y_\mu &= \varrho x_\mu \\ y_\lambda &= x_\lambda + b_\lambda x_\mu \quad \text{if } \lambda \neq \mu. \end{aligned}$$

Any maximal proper R -submodule of L is equal to $N_{\mu, b}$ for some (μ, b) . The proof of the following lemma is left to the reader.

Lemma 2.13 *Consider pairs (μ, b) and (ν, c) . The following holds.*

- (1) *Assume that $\mu = \nu$. Then $N_{\mu, b} = N_{\mu, c}$ if and only if $b_\lambda \equiv_\varrho c_\lambda$ for all $\lambda \neq \mu$.*
- (2) *Assume that $\mu \neq \nu$. Then $N_{\mu, b} = N_{\nu, c}$ if and only if $b_\nu \not\equiv_\varrho 0$, $c_\mu \equiv_\varrho b_\nu^{-1}$, and $c_\lambda \equiv_\varrho -b_\nu^{-1} b_\lambda$ for all $\lambda \neq \mu, \nu$.*

2.3 Lemmas over the ring of integers of a p -adic field

Let K be a finite extension of \mathbb{Q}_p , and let \mathcal{O}_K be the integral closure of \mathbb{Z}_p in K . Let d , e , and f be the degree, ramification index, and inertia degree of K/\mathbb{Q}_p , respectively. Let $\pi \in \mathcal{O}_K$ be a uniformizing parameter.

Remark 2.14 Let B be maximal proper \mathbb{Z}_p -submodule of \mathcal{O}_K . Since $[\mathcal{O}_K : B] = p$ and $[\mathcal{O}_K : \pi\mathcal{O}_K] = p^f$, we see that the condition $B \subseteq \pi\mathcal{O}_K$ implies $f = 1$. Hence, the conditions $B \subseteq \pi\mathcal{O}_K$ and $B = \pi\mathcal{O}_K$ are equivalent.

We recall the following known property; see, for instance, [Rei03, Theorem 5.6, page 71].

Lemma 2.15 *Let $(\alpha_i)_{0 \leq i < f}$ be a sequence of elements of \mathcal{O}_K whose reduction modulo π is a basis of $\mathcal{O}_K/\pi\mathcal{O}_K$ over \mathbb{F}_p . Let $(\beta_j)_{0 \leq j < e}$ be a sequence of elements of \mathcal{O}_K such that $v_K(\beta_j) = j$ for all j , where $v_K : K \rightarrow \mathbb{Z} \cup \{\infty\}$ is the valuation of K . Then $(\alpha_i \beta_j)_{0 \leq i < f, 0 \leq j < e}$ is a basis of \mathcal{O}_K over \mathbb{Z}_p .*

We fix a sequence $(\alpha_i)_{0 \leq i < f}$ of elements of \mathcal{O}_K whose reduction modulo π is a basis of $\mathcal{O}_K/\pi\mathcal{O}_K$ over \mathbb{F}_p . Moreover, we assume that $\alpha_0 = 1$. We define $\varepsilon_{ij} = \alpha_i \pi^j$ for $0 \leq i < f$ and $0 \leq j < e$, so that (ε_{ij}) is a basis of \mathcal{O}_K over \mathbb{Z}_p . Observe that $\varepsilon_{00} = 1$.

Lemma 2.16 *Let $s \in \mathbb{Z}$, and let $s = qe + r$, where $q, r \in \mathbb{Z}$ and $0 \leq r < e$. Then $s_{\mathbb{Z}_p}(\mathcal{O}_K, \pi^s \mathcal{O}_K) = (q^{d-rf}, (q+1)^{rf})$.*

PROOF: Since $v_K(p) = e$, by Lemma 2.15 we have that $(\alpha_i p^q \pi^{j-s} : 0 \leq i < f, r \leq j < e; \alpha_i p^{q+1} \pi^{j-s} : 0 \leq i < f, 0 \leq j < r)$ is a basis of \mathcal{O}_K over \mathbb{Z}_p . By multiplying these basis elements by π^s , we get that $(p^q \alpha_i \pi^j : 0 \leq i < f, r \leq j < e; p^{q+1} \alpha_i \pi^j : 0 \leq i < f, 0 \leq j < r)$ is a basis of $\pi^s \mathcal{O}_K$ over \mathbb{Z}_p . The result follows. \square

Remark 2.17 For $0 \leq i < f$ and $0 \leq j < e$, define

$$\gamma_{ij} = \begin{cases} p\varepsilon_{ij} & \text{if } j = 0 \\ \varepsilon_{ij} & \text{if } j \neq 0. \end{cases}$$

By applying the argument in the proof of Lemma 2.16 with $s = 1$, it follows that (γ_{ij}) is a basis of $\pi \mathcal{O}_K$ over \mathbb{Z}_p .

Corollary 2.18 *Let V be a finite dimensional K -vector space, and let $n = \dim_K(V)$. Let L and M be full \mathcal{O}_K -lattices in V , and let $s_{\mathcal{O}_K}(L, M) = (s_0, \dots, s_{n-1})$. For $0 \leq l < n$, let $s_l = q_l e + r_l$, where $q_l, r_l \in \mathbb{Z}$ and $0 \leq r_l < e$. Then $s_{\mathbb{Z}_p}(L, M) = (q_0^{d-r_0 f}, (q_0 + 1)^{r_0 f}, \dots, q_{n-1}^{d-r_{n-1} f}, (q_{n-1} + 1)^{r_{n-1} f})$.*

PROOF: There exist a basis e_0, \dots, e_{n-1} of L over \mathcal{O}_K and integers s_0, \dots, s_{n-1} such that $(\pi^{s_l} e_l)$ is a basis of M over \mathcal{O}_K . By definition, $S_{\mathcal{O}_K}(L, M) = (s_0, \dots, s_{n-1})$. Now the corollary follows from Lemma 2.16. \square

Lemma 2.19 *Let B be a maximal proper \mathbb{Z}_p -submodule of \mathcal{O}_K . Then the following holds.*

(1) *If $\pi \mathcal{O}_K \subseteq B$ then $s_{\mathbb{Z}_p}(B, \pi \mathcal{O}_K) = (0^{d-f+1}, 1^{f-1})$.*

(2) *If $\pi \mathcal{O}_K \not\subseteq B$ then $s_{\mathbb{Z}_p}(B, \pi \mathcal{O}_K) = (-1, 0^{d-f-1}, 1^f)$.*

PROOF: We will show that the relative invariant exponents $s_{\mathbb{Z}_p}(B, \pi \mathcal{O}_K)$ are either equal to $(0^{d-f+1}, 1^{f-1})$ or to $(-1, 0^{d-f-1}, 1^f)$. Since the invariant exponents are all nonnegative if and only if $\pi \mathcal{O}_K \subseteq B$, the result follows. We use the letters λ and μ for pairs of integers such as $\lambda = (\lambda_0, \lambda_1)$, where $0 \leq \lambda_0 < f$ and $0 \leq \lambda_1 < e$. Recall the basis (ε_λ) of \mathcal{O}_K over \mathbb{Z}_p introduced above, and consider the basis (γ_λ) of $\pi \mathcal{O}_K$ over \mathbb{Z}_p given in Remark 2.17. There exist μ and a sequence $(b_\lambda)_{\lambda \neq \mu}$ of elements of \mathbb{Z}_p such that (β_λ) is a basis of B over \mathbb{Z}_p , where

$$\beta_\lambda = \begin{cases} p\varepsilon_\mu & \text{if } \lambda = \mu \\ \varepsilon_\lambda + b_\lambda \varepsilon_\mu & \text{if } \lambda \neq \mu. \end{cases}$$

The idea of the proof is to write the basis (γ_λ) in coordinates with respect to (β_λ) over \mathbb{Q}_p , and then compute the Smith canonical form of the resulting change-of-basis matrix. We divide the proof into two cases.

(1) Assume that $\mu_1 = 0$ and, for all λ , if $\lambda \neq \mu$ and $\lambda_1 \neq 0$ then $b_\lambda \equiv_p 0$. By defining

$$\delta_\lambda = \begin{cases} \gamma_\mu & \text{if } \lambda = \mu \\ \gamma_\lambda + b_\lambda \gamma_\mu & \text{if } \lambda \neq \mu \text{ and } \lambda_1 = 0 \\ \gamma_\lambda + p^{-1} b_\lambda \gamma_\mu & \text{if } \lambda \neq \mu \text{ and } \lambda_1 \neq 0, \end{cases}$$

one proves that (δ_λ) is a basis of $\pi \mathcal{O}_K$ over \mathbb{Z}_p and

$$\delta_\lambda = \begin{cases} \beta_\mu & \text{if } \lambda = \mu \\ p\beta_\lambda & \text{if } \lambda \neq \mu \text{ and } \lambda_1 = 0 \\ \beta_\lambda & \text{if } \lambda \neq \mu \text{ and } \lambda_1 \neq 0, \end{cases}$$

from which the invariant exponents are computed to be $(0^{d-f+1}, 1^{f-1})$.

(2) Assume that $\mu_1 \neq 0$ or there exists λ such that $\lambda \neq \mu$, $\lambda_1 \neq 0$, and $b_\lambda \not\equiv_p 0$. The proof is similar to the one of the first case after defining

$$\delta_\lambda = \begin{cases} \gamma_\mu & \text{if } \lambda = \mu \\ \gamma_\lambda + pb_\lambda\gamma_\mu & \text{if } \lambda \neq \mu \text{ and } \lambda_1 = 0 \\ \gamma_\lambda + b_\lambda\gamma_\mu & \text{if } \lambda \neq \mu \text{ and } \lambda_1 \neq 0, \end{cases}$$

and the invariant exponents are $(-1, 0^{d-f-1}, 1^f)$. \square

In the following lemma, the indices take values $0 \leq \lambda_0, \mu_0 < f$ and $0 \leq \lambda_1, \mu_1 < e$.

Lemma 2.20 *Given (μ_0, μ_1) and a sequence $(b_{\lambda_0\lambda_1})_{(\lambda_0, \lambda_1) \neq (\mu_0, \mu_1)}$ of elements of \mathbb{Z}_p , define $\beta_{\mu_0\mu_1} = p\varepsilon_{\mu_0\mu_1}$ and $\beta_{\lambda_0\lambda_1} = \varepsilon_{\lambda_0\lambda_1} + b_{\lambda_0\lambda_1}\varepsilon_{\mu_0\mu_1}$ if $(\lambda_0, \lambda_1) \neq (\mu_0, \mu_1)$. Let B be the span of $(\beta_{\lambda_0\lambda_1})$ over \mathbb{Z}_p . Then the following are equivalent.*

(1) $B \subseteq \pi\mathcal{O}_K$.

(2) $f = 1$, $\mu_1 = 0$, and for all (λ_0, λ_1) with $(\lambda_0, \lambda_1) \neq (\mu_0, \mu_1)$ and $\lambda_1 \neq 0$ we have $b_{\lambda_0\lambda_1} \equiv_p 0$.

PROOF: Observe that when $f = 1$ the index λ_0 only takes the value 0. We first prove that (2) implies (1). Assume (2). Since $f = 1$, we can write a generic element γ of B as $\sum_{\lambda_1} c_{\lambda_1}\beta_{0\lambda_1}$ for some sequence (c_{λ_1}) of elements of \mathbb{Z}_p . Since $\mu_1 = 0$, we have

$$\gamma = c_0\beta_{00} + \sum_{\lambda_1 \neq 0} c_{\lambda_1}\beta_{0\lambda_1} = c_0p + \sum_{\lambda_1 \neq 0} c_{\lambda_1}(\pi^{\lambda_1} + b_{0\lambda_1}).$$

Since $b_{0\lambda_1} \equiv_p 0$ for $\lambda_1 \neq 0$ and since π divides p in \mathcal{O}_K , we see that γ is a multiple of π in \mathcal{O}_K .

We now prove that (2) implies (1) by contrapositive. By Remark 2.14, if $f \neq 1$ we have $B \not\subseteq \pi\mathcal{O}_K$. Assume that $f = 1$. If $\mu_1 \neq 0$ then $\beta_{00} = 1 + b_{00}\pi^{\mu_1}$ is in B but not in $\pi\mathcal{O}_K$. If there exists (λ_0, λ_1) such that $(\lambda_0, \lambda_1) \neq (\mu_0, \mu_1)$, $\lambda_1 \neq 0$, and $b_{\lambda_0\lambda_1} \not\equiv_p 0$ then, by Lemma 2.13, one can reduce the proof to the case where $\mu_1 \neq 0$. \square

Let L be an \mathcal{O}_K -lattice, and let $n = \dim_{\mathcal{O}_K}(L)$.

Remark 2.21 If M is a proper \mathcal{O}_K -submodule of L then $[L : M] \geq p^f$. It follows that, for $f \geq 2$, no maximal proper \mathbb{Z}_p -submodule of L is an \mathcal{O}_K -submodule of L . Moreover, let N be a maximal proper \mathbb{Z}_p -submodule of L . If N is an \mathcal{O}_K -submodule of L then $f = 1$ and N is a maximal proper \mathcal{O}_K -submodule of L .

We adapt the general notation presented in Section 2.2 in order to treat the maximal proper \mathbb{Z}_p -submodules of L . We consider indices $\lambda = (\lambda_0, \lambda_1, \lambda_2)$ with $0 \leq \lambda_0 < f$, $0 \leq \lambda_1 < e$, and $0 \leq \lambda_2 < n$, and similarly for μ and ν . Let (e_{λ_2}) be a basis of L over \mathcal{O}_K . We define $x_\lambda = \varepsilon_{\lambda_0\lambda_1}e_{\lambda_2}$, so that (x_λ) is a basis of L over \mathbb{Z}_p . Given an index μ and a sequence $b = (b_\lambda)_{\lambda \neq \mu}$ of elements of \mathbb{Z}_p , we define

$$\begin{aligned} y_\mu &= px_\mu \\ y_\lambda &= x_\lambda + b_\lambda x_\mu \quad \text{if } \lambda \neq \mu, \end{aligned}$$

and $N_{\mu,b}$ to be the span of (y_λ) over \mathbb{Z}_p . Hence, $N_{\mu,b}$ is a maximal proper \mathbb{Z}_p -submodule of L , and any maximal proper \mathbb{Z}_p -submodule of L is equal to $N_{\mu,b}$ for some pair (μ, b) . Given a pair (μ, b) , we define

$$\begin{aligned}\beta_{\mu_0\mu_1} &= p\varepsilon_{\mu_0\mu_1} \\ \beta_{\lambda_0\lambda_1} &= \varepsilon_{\lambda_0\lambda_1} + b_{\lambda_0\lambda_1\mu_2}\varepsilon_{\mu_0\mu_1} \quad \text{if } (\lambda_0, \lambda_1) \neq (\mu_0, \mu_1),\end{aligned}$$

and $B_{\mu,b}$ to be the span over \mathbb{Z}_p of $(\beta_{\lambda_0\lambda_1})$. Hence, $B_{\mu,b}$ is a maximal proper \mathbb{Z}_p -submodule of \mathcal{O}_K .

Lemma 2.22 *Let N be a maximal proper \mathbb{Z}_p -submodule of L . Consider the following conditions.*

- (1) *For all (μ, b) such that $N = N_{\mu,b}$ we have $B_{\mu,b} \subseteq \pi\mathcal{O}_K$.*
- (2) *$f = 1$ and for all (μ, b) such that $N = N_{\mu,b}$ the following holds: $\mu_1 = 0$ and for all λ with $\lambda_1 \neq 0$ we have $b_\lambda \equiv_p 0$.*
- (3) *$f = 1$ and there exists (μ, b) such that $N = N_{\mu,b}$, $\mu_1 = 0$, and for all λ with $\lambda_1 \neq 0$ we have $b_\lambda \equiv_p 0$.*
- (4) *N is an \mathcal{O}_K -submodule of L .*

Then (1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4).

PROOF: Observe that (2) \Rightarrow (3) is immediate. We start by proving (1) \Rightarrow (2). The claim that $f = 1$, under assumption (1), follows from Remark 2.14. Assume by contrapositive that there exists (μ, b) such that $N = N_{\mu,b}$ and either $\mu_1 \neq 0$ or there exists ν with $\nu_1 \neq 0$ and $b_\nu \not\equiv_p 0$. In case $\mu_1 \neq 0$, Lemma 2.20 implies that $B_{\mu,b} \not\subseteq \pi\mathcal{O}_K$. In case there exists ν_1 with $\nu_1 \neq 0$ and $b_\nu \not\equiv_p 0$, Lemma 2.13 implies that $N = N_{\nu,c}$ for some c , and we have $B_{\nu,c} \not\subseteq \pi\mathcal{O}_K$ again by Lemma 2.20.

We now prove that (3) \Rightarrow (4). Assume (3), and observe that, since $f = 1$, we have $\lambda_0 = 0$ for all λ . Given (μ, b) with the properties of assumption (3), we define $z_{\mu_2} = \pi e_{\mu_2}$ and $z_{\lambda_2} = e_{\lambda_2} + b_{00\lambda_2}e_{\mu_2}$ for $\lambda_2 \neq \mu_2$. Moreover, we define M to be the span of (z_{λ_2}) over \mathcal{O}_K , so that M is a maximal proper \mathcal{O}_K -submodule of L . We will show that $N = M$. Since M is a proper \mathbb{Z}_p -submodule of L , it is enough to show that the generators y_λ of N are in M . We have

$$\begin{aligned}y_\mu &= pe_{\mu_2} \\ y_\lambda &= z_{\lambda_2} \quad \text{if } \lambda_1 = 0 \text{ and } \lambda_2 \neq \mu_2 \\ y_\lambda &= \pi^{\lambda_1}e_{\lambda_2} + b_{0\lambda_1\lambda_2}e_{\mu_2} \quad \text{if } \lambda_1 \neq 0.\end{aligned}$$

By assumption, π divides $b_{0\lambda_1\lambda_2}$ for $\lambda_1 \neq 0$; also, observe that π divides p in \mathcal{O}_K and $\pi L \subseteq M$. Hence, any generator of N is in M , since it is either a generator of M as well, or a multiple of π in L . \square

Remark 2.23 The conditions in the statment of Lemma 2.22 are equivalent. Since we will not need this fact, we leave the proof to the reader.

Lemma 2.24 *Let L be an \mathcal{O}_K -Lie lattice, and let N be a maximal proper \mathbb{Z}_p -submodule of L . If N is not an \mathcal{O}_K -submodule of L then $[N, N] = [L, L]$.*

PROOF: Assume that N is not an \mathcal{O}_K -submodule of L , and let (e_{λ_2}) be a basis of L over \mathcal{O}_K . By Lemma 2.22, there exists (μ, b) such that $N = N_{\mu, b}$ and $B := B_{\mu, b} \not\subseteq \pi\mathcal{O}_K$. We have $[L, L] = \sum_{\lambda_2, \nu_2: \lambda_2 \neq \nu_2} \mathcal{O}_K[e_{\lambda_2}, e_{\nu_2}]$. Since it is clear that $[N, N] \subseteq [L, L]$, it is enough to prove that $\mathcal{O}_K[e_{\lambda_2}, e_{\nu_2}] \subseteq [N, N]$ for all λ_2 and ν_2 with $\lambda_2 \neq \nu_2$. We take λ_2 and ν_2 with $\lambda_2 \neq \nu_2$, and we divide the proof into two cases.

- (1) Assume that $\lambda_2 = \mu_2$ or $\nu_2 = \mu_2$. Without loss of generality, we can assume that $\lambda_2 = \mu_2$. Let $z \in \mathcal{O}_K[e_{\mu_2}, e_{\nu_2}]$, and let $\delta \in B \cap \mathcal{O}_K^*$. Then there exists $\gamma \in \mathcal{O}_K$ such that $z = [\delta e_{\mu_2}, \gamma e_{\nu_2}]$. We have $\gamma = \sum_{\nu_0, \nu_1} c_{\nu_0 \nu_1} \varepsilon_{\nu_0 \nu_1}$ and $\delta = \sum_{\lambda_0, \lambda_1} d_{\lambda_0 \lambda_1} \beta_{\lambda_0 \lambda_1}$ for some indexed sets $(c_{\nu_0 \nu_1})$ and $(d_{\lambda_0 \lambda_1})$ of elements of \mathbb{Z}_p . Hence,

$$z = \sum_{\lambda_0, \lambda_1, \nu_0, \nu_1} d_{\lambda_0 \lambda_1} c_{\nu_0 \nu_1} [\beta_{\lambda_0 \lambda_1} e_{\mu_2}, \varepsilon_{\nu_0 \nu_1} e_{\nu_2}].$$

Since $[\beta_{\lambda_0 \lambda_1} e_{\mu_2}, \varepsilon_{\nu_0 \nu_1} e_{\nu_2}] = [\beta_{\lambda_0 \lambda_1} e_{\mu_2}, \varepsilon_{\nu_0 \nu_1} e_{\nu_2} + b_{\nu_0 \nu_1 \nu_2} \varepsilon_{\mu_0 \mu_1} e_{\mu_2}] = [y_{\lambda_0 \lambda_1 \mu_2}, y_{\nu_0 \nu_1 \nu_2}]$, we have $z \in [N, N]$.

- (2) Assume that $\lambda_2 \neq \mu_2$ and $\nu_2 \neq \mu_2$. Let $z \in \mathcal{O}_K[e_{\lambda_2}, e_{\nu_2}]$, so that $z = \gamma[e_{\lambda_2}, e_{\nu_2}]$ for some $\gamma \in \mathcal{O}_K$. We have $\gamma = \sum_{\lambda_0, \lambda_1} c_{\lambda_0 \lambda_1} \varepsilon_{\lambda_0 \lambda_1}$ for some indexed set $(c_{\lambda_0 \lambda_1})$ of elements of \mathbb{Z}_p . Observe that $[y_{\lambda_0 \lambda_1 \lambda_2}, y_{00 \nu_2}] = \varepsilon_{\lambda_0 \lambda_1} [e_{\lambda_2}, e_{\nu_2}] + b_{00 \nu_2} \varepsilon_{\lambda_0 \lambda_1} \varepsilon_{\mu_0 \mu_1} [e_{\lambda_2}, e_{\mu_2}] + b_{\lambda_0 \lambda_1 \lambda_2} \varepsilon_{\mu_0 \mu_1} [e_{\mu_2}, e_{\nu_2}]$. Hence,

$$\begin{aligned} \sum_{\lambda_0, \lambda_1} c_{\lambda_0 \lambda_1} [y_{\lambda_0 \lambda_1 \lambda_2}, y_{00 \nu_2}] &= \gamma[e_{\lambda_2}, e_{\nu_2}] + b_{00 \nu_2} \gamma \varepsilon_{\mu_0 \mu_1} [e_{\lambda_2}, e_{\mu_2}] + \\ &\quad \left(\sum_{\lambda_0, \lambda_1} c_{\lambda_0 \lambda_1} b_{\lambda_0 \lambda_1 \lambda_2} \right) \varepsilon_{\mu_0 \mu_1} [e_{\mu_2}, e_{\nu_2}]. \end{aligned}$$

By definition, the sum on the left-hand side is in $[N, N]$; also, by the previous item, the last two terms on the right-hand side are in $[N, N]$. Since the first term on the right-hand side is z , it follows that $z \in [N, N]$. \square

3 Self-similarity in division algebras

The proofs of Theorem A and Theorem B are given in Section 3.3. In order to prove the desired results, we need some of the structure properties of D , which we state below, and several preliminary results. Specific results for $n = 2$ and $n \geq 3$ are given in Section 3.1 and Section 3.2, respectively.

Let $\pi \in \mathcal{O}_K$ be a uniformizing parameter. From the general theory we know that there exist F , θ , and Π with the following properties; see, for instance, [Rei03, Section 14].

- (1) $K \subseteq F \subseteq D$ and F is a unital K -subalgebra of D and an unramified field extension of K of degree n ; moreover, F/K is Galois and cyclic.
- (2) θ is a generator of $\text{Gal}(F/K)$, $\Pi \in D$, $\Pi^n = \pi$, and $\Pi\alpha = \theta(\alpha)\Pi$ for all $\alpha \in F$.
- (3) D is generated by F and Π as a unital K -algebra. More precisely,

$$D = \bigoplus_{0 \leq j < n} F\Pi^j.$$

(4) Let \mathcal{O}_F be the integral closure of \mathcal{O}_K in F . Then

$$\Delta = \bigoplus_{0 \leq j < n} \mathcal{O}_F \Pi^j.$$

(5) Let $t_{D/K} : D \rightarrow K$ be the reduced-trace map, and let $T_{F/K} : F \rightarrow K$ be the trace map. Let $x = \sum_j \alpha_j \Pi^j$, where $\alpha_j \in F$, be an element of D . Then $t_{D/K}(x) = T_{F/K}(\alpha_0)$.

We denote the residue fields of \mathcal{O}_K and \mathcal{O}_F by $\kappa_K = \mathcal{O}_K/\pi\mathcal{O}_K$ and $\kappa_F = \mathcal{O}_F/\pi\mathcal{O}_F$, respectively, and by $v_K : K \rightarrow \mathbb{Z} \cup \{\infty\}$ and $v_F : F \rightarrow \mathbb{Z} \cup \{\infty\}$ the respective valuations. We denote by $\text{pr} : \mathcal{O}_F \rightarrow \kappa_F$ the projection, and we adopt the notation $\bar{\alpha} = \text{pr}(\alpha)$ when convenient. The extension κ_F/κ_K has degree n . Let F^0 be the kernel of the trace map $F \rightarrow K$, and define $\mathcal{O}_F^0 = \mathcal{O}_F \cap F^0$. Let κ_F^0 be the kernel of the trace map $\kappa_F \rightarrow \kappa_K$. Observe that F^0 and κ_F^0 have dimension $n - 1$ over K and κ_K , respectively.

The following lemma is an immediate consequence of the properties listed above, and it will be used constantly in Section 3.2. From the lemma we see that D is graded over $\mathbb{Z}/n\mathbb{Z}$ as a K -Lie algebra.

Lemma 3.1 *For all $j, k \in \mathbb{Z}$ and $\alpha, \beta \in F$ we have $[\alpha \Pi^j, \beta \Pi^k] = (\alpha \theta^j(\beta) - \beta \theta^k(\alpha)) \Pi^{j+k}$, where $[\cdot, \cdot]$ denotes the commutator in D .*

Let $sl_1(D)$ be the kernel of the reduced-trace map $D \rightarrow K$, so that $sl_1(\Delta) = \Delta \cap sl_1(D)$. We have

$$sl_1(D) = F^0 \oplus \left(\bigoplus_{1 \leq j < n} F \Pi^j \right) \quad \text{and} \quad sl_1(\Delta) = \mathcal{O}_F^0 \oplus \left(\bigoplus_{1 \leq j < n} \mathcal{O}_F \Pi^j \right).$$

Remark 3.2 Assume that $n \geq 2$. Since $sl_1(D)$ is simple both as a K -Lie algebra and as a \mathbb{Q}_p -Lie algebra, $sl_1(\Delta)$ is just infinite both as an \mathcal{O}_K -Lie lattice and as a \mathbb{Z}_p -Lie lattice; see Lemma 2.10.

We point out some properties that will be used in Section 3.1 and Section 3.2.

Remark 3.3 There exists $\tau_0 \in \mathcal{O}_F$ such that $v_K(T_{F/K}(\tau_0)) = 0$, otherwise the trace map of the extension κ_F/κ_K would be identically zero. We have $\mathcal{O}_F = \mathcal{O}_K \tau_0 \oplus \mathcal{O}_F^0$. In particular, there exists a basis $(\tau_i)_{0 \leq i < n}$ of \mathcal{O}_F over \mathcal{O}_K such that $(\tau_i)_{1 \leq i < n}$ is a basis of \mathcal{O}_F^0 over \mathcal{O}_K .

Lemma 3.4 $\pi(\mathcal{O}_F^0) = \kappa_F^0$.

PROOF: The inclusion ' \subseteq ' is clear. For ' \supseteq ', let $(\tau_i)_{0 \leq i < n}$ be a basis of \mathcal{O}_F over \mathcal{O}_K such that $(\tau_i)_{1 \leq i < n}$ is a basis of \mathcal{O}_F^0 over \mathcal{O}_K . Observe that $(\bar{\tau}_i)_{0 \leq i < n}$ is a basis of κ_F over κ_K and that $(\bar{\tau}_i)_{1 \leq i < n}$ is a basis of κ_F^0 over κ_K . Also, note that $T_{F/K}(\tau_0) \not\equiv_{\pi} 0$. Let $x \in \kappa_F^0$. There exists $\alpha \in \mathcal{O}_F$ such that $\text{pr}(\alpha) = x$. We have $\alpha = \sum_{0 \leq i < n} a_i \tau_i$ for some $a_i \in \mathcal{O}_K$. We have $\pi \mid a_0$, since $a_0 T_{F/K}(\tau_0) = T_{F/K}(\alpha) \equiv_{\pi} 0$. Let $\beta = \alpha - a_0 \tau_0$. Then $\beta \in \mathcal{O}_F^0$ and $\text{pr}(\beta) = x$. \square

Lemma 3.5 *Let U be an \mathcal{O}_K -submodule of \mathcal{O}_F . Then the following holds.*

- (1) If $\text{pr}(U) = \kappa_F$ then $U = \mathcal{O}_F$.
- (2) If $U \subseteq \mathcal{O}_F^0$ and $\text{pr}(U) = \kappa_F^0$ then $U = \mathcal{O}_F^0$.

PROOF: The statement is a consequence of Nakayama's Lemma and Lemma 3.4. \square

3.1 The case $n = 2$

We aim to prove the following theorem.

Theorem 3.6 *Assume that $n = 2$ and $p \neq 2$, and let $m \in \mathbb{N}$. Then $p^m sl_1(\Delta)$, regarded as a \mathbb{Z}_p -Lie lattice, is not self-similar of index p .*

The theorem is a consequence of Lemma 3.8 and Lemma 3.11 proven below. For the expository convenience of this section, we introduce the terminology “standard basis” in the following definition, which the reader should compare with [Nos19, Lemma 2.20]. We point out that, in general, standard bases do not exist, even for 3-dimensional unsolvable Lie lattices.

Definition 3.7 *Let L be a 3-dimensional \mathcal{O}_K -Lie lattice. A standard basis of L is a basis (e_0, e_1, e_2) of L over \mathcal{O}_K with the property that $[e_i, e_{i+1}] = u_{i+2}\pi^{s_{i+2}}e_{i+2}$ for all $i \in \mathbb{Z}/3\mathbb{Z}$, where $u_i \in \mathcal{O}_K^*$ for all i , $s_0 = 1$, $s_1 = s_2 = 0$, and $-u_1u_2$ is not a square modulo π in \mathcal{O}_K , that is, there is no $\alpha \in \mathcal{O}_K$ such that $-u_1u_2 \equiv_\pi \alpha^2$.*

Lemma 3.8 *Assume that $n = 2$ and $p \neq 2$. Then there exists a standard basis of $sl_1(\Delta)$.*

PROOF: Since p is odd, $2 \in \mathcal{O}_K^*$. Hence, since $T_{F/K}(1) = 2$, there exists $\xi \in \mathcal{O}_F$ such that $(1, \xi)$ is a basis of \mathcal{O}_F over \mathcal{O}_K and $T_{F/K}(\xi) = 0$ (see Remark 3.3). By observing that $\theta(\xi) = -\xi$, so that $N_{F/K}(\xi) = -\xi^2$, we see that $\xi^2 \in \mathcal{O}_K$; moreover, since $\xi \in \mathcal{O}_F^*$, we have $\xi^2 \in \mathcal{O}_K^*$. By defining $e_0 = \xi$, $e_1 = \Pi$, and $e_2 = \xi\Pi$, we get a presentation of $sl_1(\Delta)$ of the form $sl_1(\Delta) = \mathcal{O}_K e_0 \oplus \mathcal{O}_K e_1 \oplus \mathcal{O}_K e_2$ with $[e_i, e_{i+1}] = u_{i+2}\pi^{s_{i+2}}e_{i+2}$ for $i \in \mathbb{Z}/3\mathbb{Z}$, where $s_0 = 1$, $s_1 = s_2 = 0$, $u_0 = -2\pi$, $u_1 = -2\xi^2$, and $u_2 = 2$. Since $-u_1u_2 = 4\xi^2$, it remains to show that ξ^2 is not a square modulo π in \mathcal{O}_K , which follows from the fact that the image of ξ in κ_F is not in κ_K . \square

For the proof of Lemma 3.11 we need two lemmas, for which we introduce the following context. Let L be a 3-dimensional \mathcal{O}_K -Lie lattice that admits a standard basis (e_0, e_1, e_2) . We define $M_0 = [L, L] = \text{Span}_{\mathcal{O}_K}(\pi e_0, e_1, e_2)$, which is a maximal proper \mathcal{O}_K -submodule of L . The next lemma is a version of [Nos19, Lemma 2.25, Lemma 2.26] where the ring of coefficients is \mathcal{O}_K in place of \mathbb{Z}_p , and where M is not assumed to be a subalgebra. We leave to the reader to check that the new lemma has the same proof as its original counterparts.

Lemma 3.9 *Let M be a maximal proper \mathcal{O}_K -submodule of L . Then the following holds.*

- (1) *If $M = M_0$ then $s_{\mathcal{O}_K}(M) = (0, 1^2)$ and $M = [L, L]$.*
- (2) *If $M \neq M_0$ then $s_{\mathcal{O}_K}(M) = (-1, 1, 2)$ and $M + [M, M] = L$.*

Lemma 3.10 *Let N be a maximal proper \mathbb{Z}_p -submodule of L . Then the following holds.*

- (1) *If N is not an \mathcal{O}_K -submodule of L then:*
 - (a) *if $[N, N] \subseteq N$ then $s_{\mathbb{Z}_p}(N) = (0^{3d-f+1}, 1^{f-1})$;*
 - (b) *if $[N, N] \not\subseteq N$ then $s_{\mathbb{Z}_p}(N) = (-1, 0^{3d-f-1}, 1^f)$.*
- (2) *If $N = M_0$ then $s_{\mathbb{Z}_p}(N) = (0^{3d-2f}, 1^{2f}) = (0^{3d-2}, 1^2)$.*
- (3) *If N is an \mathcal{O}_K -submodule of L and $N \neq M_0$ then:*

- (a) if $d = 1$ then $s_{\mathbb{Z}_p}(N) = (-1, 1, 2)$;
(b) if $d \geq 2$ then $s_{\mathbb{Z}_p}(N) = (-1^f, 0^{3d-4f}, 1^{3f}) = (-1, 0^{3d-4}, 1^3)$.

PROOF: When N is an \mathcal{O}_K -submodule of L , the result follows from a straightforward computation by applying Lemma 3.9 and Corollary 2.18; one has also to observe that we have $f = 1$ in this case (see Remark 2.21). Assume that N is not an \mathcal{O}_K -submodule of L . We will show that the s -invariants $s_{\mathbb{Z}_p}(N) = s_{\mathbb{Z}_p}(N, [N, N])$ are either $(0^{3d-f+1}, 1^{f-1})$ or $(-1, 0^{3d-f-1}, 1^f)$. Since the s -invariants are all nonnegative if and only if $[N, N] \subseteq N$, the result follows. Let (e_0, e_1, e_2) be a standard basis of L , and recall that $[L, L] = \mathcal{O}_K \pi e_0 \oplus \mathcal{O}_K e_1 \oplus \mathcal{O}_K e_2$. By Lemma 2.24, $[N, N] = [L, L]$. We use the notation of Section 2.3; in particular, from the paragraph below Remark 2.21, the reader should recall the domains where the indices $\lambda = (\lambda_0, \lambda_1, \lambda_2)$ and the pairs (μ, b) take values, and the definition of x_λ , y_λ , $N_{\mu, b}$, and $B_{\mu, b}$. We divide the proof into two cases.

- (1) Assume that there exists (μ, b) such that $N = N_{\mu, b}$ and $\mu_2 \neq 0$. We define

$$z_\lambda = \begin{cases} x_\lambda & \text{if } \lambda_2 \neq 0 \\ x_\lambda & \text{if } \lambda_2 = 0 \text{ and } \lambda_1 \neq 0 \\ px_\lambda & \text{if } \lambda_2 = 0 \text{ and } \lambda_1 = 0, \end{cases}$$

and from Remark 2.17 one can see that (z_λ) is a basis of $[L, L]$ over \mathbb{Z}_p . We define a new basis (w_λ) of $[L, L]$ over \mathbb{Z}_p by

$$w_\lambda = \begin{cases} z_\mu & \text{if } \lambda = \mu \\ z_\lambda + b_\lambda z_\mu & \text{if } \lambda \neq \mu \text{ and } \lambda_2 \neq 0 \\ z_\lambda + b_\lambda z_\mu & \text{if } \lambda \neq \mu \text{ and } \lambda_2 = 0 \text{ and } \lambda_1 \neq 0 \\ z_\lambda + pb_\lambda z_\mu & \text{if } \lambda \neq \mu \text{ and } \lambda_2 = 0 \text{ and } \lambda_1 = 0. \end{cases}$$

One can compute that

$$w_\lambda = \begin{cases} p^{-1}y_\mu & \text{if } \lambda = \mu \\ y_\lambda & \text{if } \lambda \neq \mu \text{ and } \lambda_2 \neq 0 \\ y_\lambda & \text{if } \lambda \neq \mu \text{ and } \lambda_2 = 0 \text{ and } \lambda_1 \neq 0 \\ py_\lambda & \text{if } \lambda \neq \mu \text{ and } \lambda_2 = 0 \text{ and } \lambda_1 = 0, \end{cases}$$

from which the s -invariants are computed to be $(-1, 0^{3d-f-1}, 1^f)$.

- (2) Assume that for all (μ, b) with $N = N_{\mu, b}$ we have $\mu_2 = 0$. We take a pair (μ, b) with $N = N_{\mu, b}$. For all λ with $\lambda_2 \neq 0$, we have $b_\lambda \equiv_p 0$, otherwise, we could apply Lemma 2.13 (with $R = \mathbb{Z}_p$ and $\varrho = p$) to get a pair (ν, c) such that $N = N_{\nu, c}$ and $\nu_2 \neq 0$. We claim that $N = Be_0 \oplus \mathcal{O}_K e_1 \oplus \mathcal{O}_K e_2$, where $B = B_{\mu, b}$ (the claim is proven below). It follows that $s_{\mathbb{Z}_p}(N) = s_{\mathbb{Z}_p}(N, [L, L]) = S_{\mathbb{Z}_p}(B, \pi\mathcal{O}_K) \cup (0^{2d})$. Lemma 2.19 gives the value of $S_{\mathbb{Z}_p}(B, \pi\mathcal{O}_K)$, and the s -invariants of N turn out to be $(0^{3d-f+1}, 1^{f-1})$ or $(-1, 0^{3d-f-1}, 1^f)$, as desired. For the proof of the claim, let $z \in Be_0 \oplus \mathcal{O}_K e_1 \oplus \mathcal{O}_K e_2$. Then, for some sequence (c_λ) of elements of \mathbb{Z}_p , we have

$$\begin{aligned} z &= \left(\sum_{\lambda: \lambda_2=0} c_\lambda \beta_{\lambda_0 \lambda_1} \right) e_0 + \left(\sum_{\lambda: \lambda_2 \neq 0} c_\lambda \varepsilon_{\lambda_0 \lambda_1} \right) e_{\lambda_2} \\ &= \sum_{\lambda: \lambda_2=0} c_\lambda y_\lambda + \sum_{\lambda: \lambda_2 \neq 0} c_\lambda (y_\lambda - p^{-1} b_\lambda y_\mu), \end{aligned}$$

where $\varepsilon_{\lambda_0\lambda_1}$ and $\beta_{\lambda_0\lambda_1}$ are defined in Section 2.3. Since $b_\lambda \equiv_p 0$ when $\lambda_2 \neq 0$, we see that $z \in N_{\mu,b} = N$. Hence, $Be_0 \oplus \mathcal{O}_Ke_1 \oplus \mathcal{O}_Ke_2 \subseteq N$. Since $Be_0 \oplus \mathcal{O}_Ke_1 \oplus \mathcal{O}_Ke_2$ is a maximal proper \mathbb{Z}_p -submodule of L and N is proper, we get the equality. \square

Now, we are ready to prove the lemma that completes the proof of Theorem 3.6.

Lemma 3.11 *Let L be a 3-dimensional \mathcal{O}_K -Lie lattice that admits a standard basis, and let $m \in \mathbb{N}$. Then p^mL , regarded as a \mathbb{Z}_p -Lie lattice, is not self-similar of index p .*

PROOF: Let $\varphi : N \rightarrow p^mL$ be a virtual endomorphism of p^mL of index p . We have to show that φ is not simple. We start by proving that p^mL is just infinite as \mathbb{Z}_p -Lie lattice. With the same proof as for [Nos22b, Lemma 9], one can see that L is just infinite as an \mathcal{O}_K -Lie lattice. Then, being nonabelian, the K -Lie algebra $L \otimes_{\mathcal{O}_K} K$ is simple (Lemma 2.10). By [Bou75, I.6.10], $L \otimes_{\mathcal{O}_K} K$ is simple as a \mathbb{Q}_p -Lie algebra. Observe that L is a full \mathbb{Z}_p -lattice in the \mathbb{Q}_p -vector space $L \otimes_{\mathcal{O}_K} K$, so that we can identify the \mathbb{Q}_p -Lie algebras $L \otimes_{\mathcal{O}_K} K$ and $L \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$. Hence, by a second application of Lemma 2.10, L is just infinite as a \mathbb{Z}_p -Lie lattice. Finally, again by Lemma 2.10, p^mL is just infinite as a \mathbb{Z}_p -Lie lattice, as desired.

Now, by Lemma 2.11, we can assume that φ is injective. We define $N' = \varphi(N)$, so that φ induces an isomorphism $N \simeq N'$ of \mathbb{Z}_p -Lie algebras, from which we get $s_{\mathbb{Z}_p}(N) = s_{\mathbb{Z}_p}(N')$. By index stability [Nos21, Corollary 8], the index of N' in p^mL is p . It follows that $p^{-m}N$ and $p^{-m}N'$ are maximal proper \mathbb{Z}_p -submodules of L . Since N and N' have the same s -invariants, since the lists of s -invariants that appear in Lemma 3.10 are different among the (sub)items (by looking at the multiplicities of 1 and 2, for instance), and since the items of the lemma exhaust all the cases, we see that one of the following possibilities holds:

- none of $p^{-m}N$ and $p^{-m}N'$ is an \mathcal{O}_K -submodule of L ;
- $p^{-m}N = M_0 = p^{-m}N'$;
- both $p^{-m}N$ and $p^{-m}N'$ are \mathcal{O}_K -submodules of L different from M_0 .

By Lemma 2.24 and Lemma 3.9, one of the following possibilities holds:

- $[p^{-m}N, p^{-m}N] = [L, L] = [p^{-m}N', p^{-m}N']$;
- $p^{-m}N = [L, L] = p^{-m}N'$;
- $p^{-m}N + [p^{-m}N, p^{-m}N] = L = p^{-m}N' + [p^{-m}N', p^{-m}N']$.

Hence, one of the following possibilities holds:

- $[N, N] = [p^mL, p^mL] = [N', N']$;
- $p^mN = [p^mL, p^mL] = p^mN'$;
- $p^mN + [N, N] = p^m(p^mL) = p^mN' + [N', N']$.

Finally, one of $[p^mL, p^mL]$ and $p^m(p^mL)$ is a nonzero ideal of p^mL that is φ -invariant. Hence, φ is not simple. \square

3.2 The case $n \geq 3$

In this section we work over \mathcal{O}_K , and we aim to prove the following theorem.

Theorem 3.12 *Assume that $n \geq 3$ and $(p, n) \neq (3, 3)$. Let M be a maximal proper \mathcal{O}_K -submodule of $sl_1(\Delta)$. Then $[M, M] = [sl_1(\Delta), sl_1(\Delta)]$.*

The theorem follows from Lemma 3.14, Corollary 3.28, and Corollary 3.29. The corollaries themselves are a consequence of a rather long list of technical lemmas. We do not make the overall assumption that $n \geq 3$. Notably, Lemma 3.16 works for $n = 2$ and $p \neq 2$ as well, and it provides a computation of $[sl_1(\Delta), sl_1(\Delta)]$ alternative to the one given in [Nos19], for instance. On the other hand, the reader may assume that $n \geq 3$ from the beginning and not lose anything fundamental.

Recall the realization of $sl_1(\Delta)$ given in the introduction to Section 3. We denote $L = sl_1(\Delta)$, $L_0 = \mathcal{O}_F^0$, and $L_j = \mathcal{O}_F \Pi^j$ for $0 < j < n$. Hence,

$$L = \bigoplus_{0 \leq j < n} L_j,$$

and we observe that L is graded over $\mathbb{Z}/n\mathbb{Z}$ as an \mathcal{O}_K -Lie algebra. We define Λ to be the set of pairs (i, j) of integers such that $0 \leq i, j < n$ and $(i, j) \neq (0, 0)$. We endow Λ with the binary relation

$$(i, j) \leq (l, k) \quad : \iff \quad j < k \text{ or } (j = k \text{ and } i \leq l),$$

which is a linear order. If we denote an element of Λ by λ , say, then we denote its components by (λ_0, λ_1) . For $\lambda \in \Lambda$, we define $\Lambda_\lambda = \{\eta \in \Lambda : \eta < \lambda\}$. Recall that we denote by $\text{pr} : \mathcal{O}_F \rightarrow \kappa_F$ the projection, and that we adopt the notation $\bar{\alpha} = \text{pr}(\alpha)$ when convenient.

Lemma 3.13 *There exists a basis $(\tau_i)_{0 \leq i < n}$ of \mathcal{O}_F over \mathcal{O}_K such that the following conditions hold.*

- (1) $(\tau_i)_{1 \leq i < n}$ is a basis of \mathcal{O}_F^0 over \mathcal{O}_K .
- (2) If $p \nmid n$ then $\tau_0 = 1$.
- (3) If $p \mid n$ then $\bar{\tau}_1 = 1$.
- (4) If $n \geq 2$ and $(p, n) \neq (2, 2)$ then $\bar{\tau}_{n-1} \notin \kappa_K$.

PROOF: Let $(t_i)_{0 \leq i < n}$ be a basis of κ_F over κ_K with the properties stated in Lemma 2.6. From Nakayama's Lemma and Lemma 3.4, it follows that there exists a basis $(\tau_i)_{0 \leq i < n}$ of \mathcal{O}_F over \mathcal{O}_K such that $(\tau_i)_{1 \leq i < n}$ is a basis of \mathcal{O}_F^0 over \mathcal{O}_K , $\tau_0 = 1$ if $p \nmid n$, and $\bar{\tau}_i = t_i$ for $0 \leq i < n$. The other required properties are immediate. \square

We fix a basis $(\tau_i)_{0 \leq i < n}$ of \mathcal{O}_F over \mathcal{O}_K with the properties stated in Lemma 3.13. For $\eta \in \Lambda$, we define $x_\eta = \tau_{\eta_0} \Pi^{\eta_1}$. Then $(x_\eta)_{\eta \in \Lambda}$ is a basis of L over \mathcal{O}_K . For $\lambda \in \Lambda$, $e : \Lambda_\lambda \rightarrow \mathcal{O}_K$, and $\eta \in \Lambda$ we define

$$y_\eta = \begin{cases} x_\eta + e_\eta x_\lambda & \text{if } \eta < \lambda \\ \pi x_\lambda & \text{if } \eta = \lambda \\ x_\eta & \text{if } \eta > \lambda, \end{cases}$$

and $M_{\lambda, e} = \text{span}_{\mathcal{O}_K}(y_\eta : \eta \in \Lambda)$, which is a maximal proper \mathcal{O}_K -submodule of L . For the next lemma see, for instance, [Adk92, Section 5.2].

Lemma 3.14 *Let M be a maximal proper \mathcal{O}_K -submodule of $sl_1(\Delta)$. Then there exist $\lambda \in \Lambda$ and $e : \Lambda_\lambda \rightarrow \mathcal{O}_K$ such that $M = M_{\lambda,e}$.*

Remark 3.15 In most proofs of the statements below we construct an \mathcal{O}_K -submodule U of \mathcal{O}_F (respectively, of \mathcal{O}_F^0), usually through a set of generators, say $(\alpha_i)_{i \in I}$, where I is a set of indices. In these situations, the aim is to prove that $U = \mathcal{O}_F$ (respectively, $U = \mathcal{O}_F^0$). This is achieved by proving that $\text{pr}(U) = \kappa_F$ (respectively, $\text{pr}(U) = \kappa_F^0$), and by applying Corollary 3.5. Observe that $\text{pr}(U)$ is generated by $(\bar{\alpha}_i)_{i \in I}$ over κ_K . The proof of $\text{pr}(U) = \kappa_F$ (respectively, $\text{pr}(U) = \kappa_F^0$) is achieved by applying the relevant results of Section 2.1 to the residue field extension κ_F/κ_K and the generator ϑ given by the reduction modulo π of the restriction of θ to \mathcal{O}_F . We observe that $(\bar{\tau}_i)_{0 \leq i < n}$ is a basis of κ_F over κ_K that satisfies the properties of Lemma 2.6, and we note that $\text{ch}(\kappa_K) = p$. We will make constant use of Lemma 3.1 without further mention. Also, recall that $\Pi^n = \pi$.

Lemma 3.16 *Assume that $(p, n) \neq (2, 2)$. Then*

$$[L, L] = \pi L_0 \oplus \left(\bigoplus_{1 \leq j < n} L_j \right).$$

PROOF: Since the lemma is trivial for $n = 1$, we assume that $n \geq 2$. We denote $L' = [L, L]$ and $L'_j = L' \cap L_j$, for $0 \leq j < n$. Observe that L'_j is generated over \mathcal{O}_K by the brackets $[x_\eta, x_\mu]$ where $\eta, \mu \in \Lambda$ and $\eta_1 + \mu_1 \equiv_n j$; also, L' is homogeneous, that is, $L' = \bigoplus_{0 \leq j < n} L'_j$. Observe that if $\eta_1 + \mu_1 = 0$ then $[x_\eta, x_\mu] = 0$. Hence, $L'_0 \subseteq \pi L_0$. It is enough to show that $\pi L_0 \subseteq L'_0$ and $L_j \subseteq L'_j$ for $0 < j < n$.

We first argue for L_j . Let $0 < j < n$. The brackets $[x_{i0}, x_{lj}] = (\tau_i - \theta^j(\tau_i))\tau_l \Pi^j$ belong to L'_j for $1 \leq i < n$ and $0 \leq l < n$. We will show that $N_j := \text{span}_{\mathcal{O}_K}([x_{i0}, x_{lj}] : 1 \leq i < n \text{ and } 0 \leq l < n)$ is equal to L_j , from which $L_j \subseteq L'_j$ follows. We have $N_j = U_j \Pi^j$, where $U_j = \text{span}_{\mathcal{O}_K}([\tau_i - \theta^j(\tau_i)]\tau_l : 1 \leq i < n \text{ and } 0 \leq l < n)$. Now the desired result follows from Remark 3.15 and Lemma 2.1.

Finally we argue for πL_0 . The brackets $[x_{01}, x_{l,n-1}] = \pi(\tau_0 \theta(\tau_l) - \tau_l \theta^{-1}(\tau_0))$ belong to L'_0 for $0 \leq l < n$. We will show that $N_0 := \text{span}_{\mathcal{O}_K}([x_{01}, x_{l,n-1}] : 0 \leq l < n)$ is equal to πL_0 , from which $\pi L_0 \subseteq L'_0$ follows. We have $N_0 = \pi U_0$, where $U_0 = \text{span}_{\mathcal{O}_K}(\tau_0 \theta(\tau_l) - \tau_l \theta^{-1}(\tau_0) : 0 \leq l < n)$. Now the desired result follows from Remark 3.15 and Lemma 2.2. \square

Remark 3.17 Let M be a maximal proper \mathcal{O}_K -submodule of L . Clearly, $[M, M] \subseteq [L, L]$; hence, in order to prove that $[L, L] = [M, M]$, it is enough to prove that $[L, L] \subseteq [M, M]$. For $(p, n) \neq (2, 2)$, it is enough to prove that $\pi L_0 \subseteq [M, M]$ and $L_m \subseteq [M, M]$ for $0 < m < n$; see Lemma 3.16. Observe that $\pi L \subseteq M$, hence, $\pi^2[L, L] \subseteq [M, M]$.

For the next sequence of lemmas and corollaries, we take $\lambda \in \Lambda$ and $e : \Lambda_\lambda \rightarrow \mathcal{O}_K$, and we define $M = M_{\lambda,e}$. We order the lemmas according to the order in which they are applied in the corollaries.

Lemma 3.18 *Assume that $\lambda_1 = 0$, and let m be an integer such that $2 \leq m < n$. Then $L_m \subseteq [M, M]$.*

PROOF: Observe that $n \geq 3$. Denote $\eta = (i, 1)$ and $\mu = (l, m - 1)$, for $0 \leq i < n$ and $0 \leq l < n$. Observe that $\lambda < \eta$ and $\lambda < \mu$, so that $[y_\eta, y_\mu] = [x_\eta, x_\mu]$, which belongs to $[M, M]$. We will show that $N := \text{span}_{\mathcal{O}_K}([x_{i1}, x_{l, m-1}] : 0 \leq i < n \text{ and } 0 \leq l < n)$ is equal to L_m , from which $L_m \subseteq [M, M]$ follows. Since $[x_{i1}, x_{l, m-1}] = (\tau_i \theta(\tau_l) - \tau_l \theta^{m-1}(\tau_i)) \Pi^m$, we have $N = U \Pi^m$, where $U = \text{span}_{\mathcal{O}_K}(\tau_i \theta(\tau_l) - \tau_l \theta^{m-1}(\tau_i) : 0 \leq i < n \text{ and } 0 \leq l < n)$. Now the desired result follows from Remark 3.15 and Lemma 2.5. \square

Lemma 3.19 *Assume that $n \geq 3$, $\lambda_1 = 0$, and $\lambda_0 < n - 1$. Then $L_1 \subseteq [M, M]$.*

PROOF: Denote $\eta = (n-1, 0)$ and $\mu = (l, 1)$, for $0 \leq l < n$. Observe that $\lambda < \eta$ and $\lambda < \mu$, so that $[y_\eta, y_\mu] = [x_\eta, x_\mu]$, which belongs to $[M, M]$. We will show that $N := \text{span}_{\mathcal{O}_K}([x_{n-1,0}, x_{l1}] : 0 \leq l < n)$ is equal to L_1 , from which $L_1 \subseteq [M, M]$ follows. Since, $[x_{n-1,0}, x_{l1}] = (\tau_{n-1} - \theta(\tau_{n-1})) \tau_l \Pi$, we have $N = U \Pi$, where $U = \text{span}_{\mathcal{O}_K}([\tau_{n-1} - \theta(\tau_{n-1})] \tau_l : 0 \leq l < n)$. Now the desired result follows from Remark 3.15 and the fact that $\bar{\tau}_{n-1} \notin \kappa_K$. \square

Lemma 3.20 *Assume that $n \geq 3$, $(p, n) \neq (3, 3)$, $\lambda_1 = 0$, and $\lambda_0 = n - 1$. Then $L_1 \subseteq [M, M]$.*

PROOF: Let $i_0 = 1$ if $p \nmid n$, and $i_0 = 2$ if $p \mid n$. Observe that $i_0 < n - 1$. Denote $\eta = (i_0, 0)$ and $\mu = (l, 1)$, for $0 \leq l < n$. Observe that $\eta < \lambda < \mu$, so that $[y_\eta, y_\mu] = [x_\eta, x_\mu] + e_\eta [x_\lambda, x_\mu]$, which belongs to $[M, M]$. By denoting $\xi_j = \tau_j - \theta(\tau_j)$ for $j \in \{i_0, n-1\}$, we have $[y_\eta, y_\mu] = \tau_l (\xi_{i_0} + e_{i_0} \xi_{n-1}) \Pi$. We will show that $N := \text{span}_{\mathcal{O}_K}([y_{i_0 0}, y_{l1}] : 0 \leq l < n)$ is equal to L_1 , from which $L_1 \subseteq [M, M]$ follows. We have $N = U \Pi$, where $U = \text{span}_{\mathcal{O}_K}(\tau_l (\xi_{i_0} + e_{i_0} \xi_{n-1}) : 0 \leq l < n)$. Observe that $\xi_{i_0} + e_{i_0} \xi_{n-1} \not\equiv_\pi 0$; see Remark 2.7. Now the desired result follows from Remark 3.15. \square

Lemma 3.21 *Assume that $\lambda_1 = 0$. Then $\pi L_0 \subseteq [M, M]$.*

PROOF: For $n = 1$ the result is trivial. Assume $n \geq 2$. Denote $\eta = (0, 1)$ and $\mu = (l, n - 1)$, for $0 \leq l < n$. Observe that $\lambda < \eta$ and $\lambda < \mu$, so that $[y_\eta, y_\mu] = [x_\eta, x_\mu]$, which belongs to $[M, M]$. We will show that $N := \text{span}_{\mathcal{O}_K}([x_{01}, x_{l, n-1}] : 0 \leq l < n)$ is equal to πL_0 , from which $\pi L_0 \subseteq [M, M]$ follows. Since $[x_{01}, x_{l, n-1}] = \pi(\tau_0 \theta(\tau_l) - \tau_l \theta^{-1}(\tau_0))$, we have $N = \pi U$, where $U = \text{span}_{\mathcal{O}_K}(\tau_0 \theta(\tau_l) - \tau_l \theta^{-1}(\tau_0) : 0 \leq l < n)$. Now the desired result follows from Remark 3.15 and Lemma 2.2. \square

Lemma 3.22 *Assume that $n \geq 3$, $\lambda_1 = 1$, and $\lambda_0 < n - 1$. Then $\pi L_0 \subseteq [M, M]$.*

PROOF: By considering the indices $\eta = (n-1, 1)$ and $\mu = (l, n-1)$, for $0 \leq l < n$, the proof is similar to the one of Lemma 3.21. \square

Lemma 3.23 *Assume that $n \geq 3$, $\lambda_1 = 1$, and $\lambda_0 = n - 1$. Then $\pi L_0 \subseteq [M, M]$.*

PROOF: Denote $\eta = (0, 1)$ and $\mu = (l, n - 1)$, for $0 \leq l < n$, and observe that $\eta < \lambda < \mu$. Denote $\sigma_0 = \tau_0 + e_\eta \tau_{n-1}$, and observe that $\sigma_0 \not\equiv_\pi 0$. After noticing that $[y_\eta, y_\mu] = [x_\eta, x_\mu] + e_\eta [x_\lambda, x_\mu] = \pi(\sigma_0 \theta(\tau_l) - \tau_l \theta^{-1}(\sigma_0))$, the proof proceeds as in the case of Lemma 3.21. \square

Lemma 3.24 *Assume that $\lambda_1 \geq 1$. Let k and m be integers such that $k \geq 0$ and $1 \leq m < n$, and assume that $(\lambda_1, m) \neq (1, 1)$. Assume that $\pi^{k+1} L \subseteq [M, M]$ and $\pi^k L_j \subseteq [M, M]$ for $m < j < n$. Then $\pi^k L_m \subseteq [M, M]$.*

PROOF: Observe that we have $n \geq 3$. We divide the proof into three cases.

- (1) Case $\lambda_1 < m$. Denote $\eta = (i, 0)$ and $\mu = (l, m)$, for $1 \leq i < n$ and $0 \leq l < n$. Observe that $\eta < \lambda < \mu$, so that $[y_\eta, y_\mu] = [x_\eta, x_\mu] + e_\eta[x_\lambda, x_\mu]$. We also have $[x_\lambda, x_\mu] \in L_{\lambda_1+m}$ when $\lambda_1 + m < n$, and $[x_\lambda, x_\mu] \in \pi L$ when $\lambda_1 + m \geq n$. From the assumptions, since $\lambda_1 + m > m$, one can deduce that $\pi^k[x_\eta, x_\mu] \in [M, M]$. Observe that $[x_\eta, x_\mu] = [x_{i0}, x_{lm}] = (\tau_i - \theta^m(\tau_i))\tau_l\Pi^m$. We will show that $N_1 := \text{span}_{\mathcal{O}_K}([x_{i0}, x_{lm}] : 1 \leq i < n \text{ and } 0 \leq l < n)$ is equal to L_m , from which $\pi^k L_m \subseteq [M, M]$ follows. We have $N_1 = U_1\Pi^m$, where $U_1 = \text{span}_{\mathcal{O}_K}([\tau_i - \theta^m(\tau_i)]\tau_l : 1 \leq i < n \text{ and } 0 \leq l < n)$. Now the desired result follows from Remark 3.15 and Lemma 2.1.
- (2) Case $\lambda_1 > m$. Denote $\eta = (i, 0)$ and $\mu = (l, m)$, for $1 \leq i < n$ and $0 \leq l < n$. Observe that $\eta, \mu < \lambda$, so that $[y_\eta, y_\mu] = [x_\eta, x_\mu] + e_\eta[x_\lambda, x_\mu] + e_\mu[x_\eta, x_\lambda]$. We have $[x_\lambda, x_\mu] \in L_{\lambda_1+m}$ when $\lambda_1 + m < n$, and $[x_\lambda, x_\mu] \in \pi L$ when $\lambda_1 + m \geq n$; also, $[x_\eta, x_\lambda] \in L_{\lambda_1}$. From the assumptions, since $\lambda_1 + m > \lambda_1 > m$, one can deduce that $\pi^k[x_\eta, x_\mu] \in [M, M]$. From here, the argument for this case is completed in exactly the same way as for case (1).
- (3) Case $\lambda_1 = m$. Observe that $m \geq 2$.
Denote $\eta = (i, 1)$ and $\mu = (l, m-1)$, for $0 \leq i < n$ and $0 \leq l < n$. Observe that $\eta, \mu < \lambda$, so that $[y_\eta, y_\mu] = [x_\eta, x_\mu] + e_\eta[x_\lambda, x_\mu] + e_\mu[x_\eta, x_\lambda]$. We have $[x_\lambda, x_\mu] \in L_{2m-1}$ when $2m-1 < n$, and $[x_\lambda, x_\mu] \in \pi L$ when $2m-1 \geq n$; also, $[x_\eta, x_\lambda] \in L_{m+1}$ when $m+1 < n$, and $[x_\eta, x_\lambda] \in \pi L$ when $m+1 \geq n$. From the assumptions, since $2m-1 \geq m+1 > m$, one can deduce that $\pi^k[x_\eta, x_\mu] \in [M, M]$. After noticing that $[x_\eta, x_\mu] = [x_{i1}, x_{l,m-1}] = (\tau_i\theta(\tau_l) - \tau_l\theta^{m-1}(\tau_i))\Pi^m$, the proof proceeds as in the case of Lemma 3.18. \square

Lemma 3.25 *Assume that $n \geq 3$, $(p, n) \neq (3, 3)$, and $\lambda_1 = 1$. Let k be an integer such that $k \geq 0$, and assume that $\pi^k L_2 \subseteq [M, M]$. Then $\pi^k L_1 \subseteq [M, M]$.*

PROOF: Denote $\eta = (i, 0)$ and $\mu = (l, 1)$, for $1 \leq i < n$, $0 \leq l < n$, and $l \neq \lambda_0$. Observe that $\eta < \lambda$; also, $\lambda < \mu$ when $l > \lambda_0$, and $\mu < \lambda$ when $l < \lambda_0$. Define $z_{il} = [x_{i0}, x_{l1}]$ when $l > \lambda_0$, and $z_{il} = [x_{i0}, x_{l1}] + e_l[x_{i0}, x_\lambda]$ when $l < \lambda_0$, where $e_l = e_{l1}$. Hence, $[y_\eta, y_\mu] = z_{il} + e_\eta[x_\lambda, x_\mu]$. We have $[x_\lambda, x_\mu] \in L_2$. From the assumptions one can deduce that $\pi^k z_{il} \in [M, M]$. For $1 \leq i < n$ we define $\xi_i = \tau_i - \theta(\tau_i)$, while for $0 \leq l < n$ and $l \neq \lambda_0$ we define $\sigma_l = \tau_l$ when $l > \lambda_0$, and $\sigma_l = \tau_l + e_l\tau_{\lambda_0}$ when $l < \lambda_0$. Observe that $z_{il} = \xi_i\sigma_l\Pi$. We will show that $N := \text{span}_{\mathcal{O}_K}(z_{il} : 1 \leq i < n, 0 \leq l < n, \text{ and } l \neq \lambda_0)$ is equal to L_1 , from which $\pi^k L_1 \subseteq [M, M]$ follows. We have $N = U\Pi$, where $U = \text{span}_{\mathcal{O}_K}(\xi_i\sigma_l : 1 \leq i < n, 0 \leq l < n, \text{ and } l \neq \lambda_0)$. Now we want to apply Remark 3.15. In order to do so, we observe that $(\bar{\sigma}_l)_{l \neq \lambda_0}$ spans a subspace of dimension $n-1$ over κ_K . Hence, the desired result follows from Remark 2.7 and Remark 2.8. \square

Lemma 3.26 *Assume that $2 \leq \lambda_1 \leq n-2$, and let k be an integer such that $k \geq 1$. Assume that $\pi^k L_j \subseteq [M, M]$ for $0 < j < n$. Then $\pi^k L_0 \subseteq [M, M]$.*

PROOF: Observe that $n \geq 4$. Denote $\eta = (0, 1)$ and $\mu = (l, n-1)$, for $0 \leq l < n$. Observe that $\eta < \lambda < \mu$, so that $[y_\eta, y_\mu] = [x_\eta, x_\mu] + e_\eta[x_\lambda, x_\mu]$. Since $n < \lambda_1 + n - 1 < 2n$, we have $[x_\lambda, x_\mu] \in \pi L_{\lambda_1-1}$. From the assumptions, since $0 < \lambda_1 - 1 < n$, one can deduce that $\pi^{k-1}[x_\eta, x_\mu] \in [M, M]$. After noticing that $[x_\eta, x_\mu] = [x_{01}, x_{l,n-1}] = \pi(\tau_0\theta(\tau_l) - \tau_l\theta^{-1}(\tau_0))$, the proof proceeds as in the case of Lemma 3.21. \square

Lemma 3.27 *Assume that $n \geq 3$ and $\lambda_1 = n-1$. Let k be an integer such that $k \geq 1$, and assume that $\pi^k L_j \subseteq [M, M]$ for $0 < j < n$. Then $\pi^k L_0 \subseteq [M, M]$.*

PROOF: We divide the proof into two parts.

- (1) Case $\lambda_0 < n - 1$. Denote $\eta = (i, 1)$ and $\mu = (n - 1, n - 1)$, for $0 \leq i < n$. Observe that $\eta < \lambda < \mu$, so that $[y_\eta, y_\mu] = [x_\eta, x_\mu] + e_\eta[x_\lambda, x_\mu]$. Since $\lambda_1 + \mu_1 = 2n - 2$ and in particular $n < \lambda_1 + \mu_1 < 2n$, we have $[x_\lambda, x_\mu] \in \pi L_{n-2}$. From the assumptions, since $0 < n - 2 < n$, one can deduce that $\pi^{k-1}[x_\eta, x_\mu] \in [M, M]$. Observe that $[x_\eta, x_\mu] = [x_{i1}, x_{n-1, n-1}] = \pi(\tau_i\theta(\tau_{n-1}) - \tau_{n-1}\theta^{-1}(\tau_i))$. We will show that $N := \text{span}_{\mathcal{O}_K}([x_{i1}, x_{n-1, n-1}] : 0 \leq i < n)$ is equal to πL_0 , from which $\pi^k L_0 \subseteq [M, M]$ follows. We have $N = \pi U$, where $U = \text{span}_{\mathcal{O}_K}(\tau_i\theta(\tau_{n-1}) - \tau_{n-1}\theta^{-1}(\tau_i) : 0 \leq i < n)$. Now the desired result follows from Remark 3.15 and Remark 2.3.
- (2) Case $\lambda_0 = n - 1$. Denote $\eta = (i, 1)$ and $\mu = (0, n - 1)$, for $0 \leq i < n$. Observe that $\eta < \lambda$ and $\mu < \lambda$, so that $[y_\eta, y_\mu] = [x_\eta, x_\mu] + e_\eta[x_\lambda, x_\mu] + e_\mu[x_\eta, x_\lambda]$. Since $\lambda_1 + \mu_1 = 2n - 2$ and in particular $n < \lambda_1 + \mu_1 < 2n$, we have $[x_\lambda, x_\mu] \in \pi L_{n-2}$. Define $z_i = [x_\eta, x_\mu] + e_\mu[x_\eta, x_\lambda]$ for $0 \leq i < n$, and define $\sigma_0 = \tau_0 + e_\mu\tau_{n-1}$. Observe that $\sigma_0 \not\equiv_\pi 0$. From the assumptions, since $0 < n - 2 < n$, one can deduce that $\pi^{k-1}z_i \in [M, M]$. After noticing that $z_i = \pi(\tau_i\theta(\sigma_0) - \sigma_0\theta^{-1}(\tau_i))$, the proof proceeds as in the case of item (1). \square

Corollary 3.28 *Assume that $n \geq 3$, $(p, n) \neq (3, 3)$, and $\lambda_1 \leq 1$. Then $[M, M] = [L, L]$.*

PROOF: We divide the proof into two parts.

- (1) Case $\lambda_1 = 0$. The result follows from Remark 3.17 and Lemmas 3.18, 3.19, 3.20, and 3.21.
- (2) Case $\lambda_1 = 1$. We will prove that if $k \geq 0$ is an integer such that $\pi^{k+1}[L, L] \subseteq [M, M]$ then $\pi^k[L, L] \subseteq [M, M]$. Since $\pi^2[L, L] \subseteq [M, M]$, see Remark 3.17, the corollary follows by descending induction on k . Let $k \geq 0$ be an integer such that $\pi^{k+1}[L, L] \subseteq [M, M]$. We have $\pi^{k+1}L_j \subseteq [M, M]$ for $0 < j < n$. By applying Lemma 3.22 in case $\lambda_0 < n - 1$, and by applying Lemma 3.23 in case $\lambda_0 = n - 1$, we see that $\pi^{k+1}L_0 \subseteq [M, M]$. Hence, $\pi^{k+1}L \subseteq [M, M]$. By Lemma 3.24 and descending induction on m , $\pi^k L_m \subseteq [M, M]$ for $2 \leq m < n$. By Lemma 3.25, $\pi^k L_1 \subseteq [M, M]$. It follows that $\pi^k[L, L] \subseteq [M, M]$. \square

Corollary 3.29 *Assume that $\lambda_1 \geq 2$. Then $[M, M] = [L, L]$.*

PROOF: Observe that $n \geq 3$. We will prove that if $k \geq 0$ is an integer such that $\pi^{k+1}[L, L] \subseteq [M, M]$ then $\pi^k[L, L] \subseteq [M, M]$. Since $\pi^2[L, L] \subseteq [M, M]$, see Remark 3.17, the corollary follows by descending induction on k . Let $k \geq 0$ be an integer such that $\pi^{k+1}[L, L] \subseteq [M, M]$. We have $\pi^{k+1}L_j \subseteq [M, M]$ for $0 < j < n$. By applying Lemma 3.26 in case $\lambda_1 < n - 1$, and by applying Lemma 3.27 in case $\lambda_1 = n - 1$, we see that $\pi^{k+1}L_0 \subseteq [M, M]$. Hence, $\pi^{k+1}L \subseteq [M, M]$. By Lemma 3.24 and descending induction on m , $\pi^k L_m \subseteq [M, M]$ for $1 \leq m < n$. It follows that $\pi^k[L, L] \subseteq [M, M]$. \square

3.3 Proof of Theorem A and Theorem B

Proof of Theorem A. The theorem follows from Remark 2.21, Lemma 2.24, and Theorem 3.12. \square

Proof of Theorem B. Denote $L = sl_1(\Delta)$. In case $n = 1$ we have $L = \{0\}$, and the result is trivial. Assume that $n \geq 2$, in which case L is just infinite as a \mathbb{Z}_p -Lie lattice (Remark 3.2) and $\dim_{\mathbb{Z}_p}(L) \geq 3$. In case $f \geq 2$, the result follows from Theorem A and Lemma 2.12. Assume that

$(p, n) \notin \{(2, 2), (3, 3)\}$. In case $n = 2$ we have $p \neq 2$, and the result follows from Theorem 3.6. In case $n \geq 3$ we have $(p, n) \neq (3, 3)$, and the result follows from Theorem A and Lemma 2.12. \square

4 Proof of Theorem E

The ‘only if’ part of Theorem E follows from [Nos22a, Theorem 2.34] and Theorem 4.1 below. The ‘if’ part follows from [Nos22a, Remark 2.35, Proposition 2.40].

Theorem 4.1 *Assume that $p \geq 5$, and let L be an unsolvable \mathbb{Z}_p -Lie lattice. Then L is not strongly hereditarily self-similar of index p .*

PROOF: Observe that $\dim(L) \geq 3$. We will prove that there exists a nonzero \mathbb{Z}_p -subalgebra L_0 of L that is not self-similar of index p . Let $\mathcal{L} = L \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$, which is an unsolvable \mathbb{Q}_p -Lie algebra of dimension d . By the Levi decomposition theorem [Jac62, page 91] and by the unsolvability of \mathcal{L} , there exists a nonzero semisimple \mathbb{Q}_p -subalgebra \mathcal{L}_1 of \mathcal{L} . By the structure theorem of semisimple Lie algebras [Jac62, page 71], there exists a simple \mathbb{Q}_p -subalgebra \mathcal{M} of \mathcal{L}_1 . We divide the proof into two cases according whether \mathcal{M} contains a nonzero nilpotent element or not.

First, we assume that \mathcal{M} contains a nonzero nilpotent element e . By the Jacobson-Morozov theorem (see [Bou05, Chapter VIII, Section 11, Proposition 2]), the element e may be completed to an sl_2 -triple, that is, there exist $f, h \in \mathcal{M}$ such that the set $\{e, f, h\}$ is linearly independent over \mathbb{Q}_p , $[h, e] = 2e$, $[h, f] = -2f$, and $[e, f] = h$. Let M be the \mathbb{Z}_p -subalgebra of \mathcal{L} generated by e, f , and h , and observe that M is a \mathbb{Z}_p -Lie lattice isomorphic to $sl_2(\mathbb{Z}_p)$; moreover, there exists $m \in \mathbb{N}$ such that $p^m M \subseteq L$. By [Nos22a, Proposition 2.41], there exists a nonzero \mathbb{Z}_p -subalgebra L_0 of $p^m M$ which is not self-similar of index p .

Second, we assume that \mathcal{M} contains no nonzero nilpotent elements. By [Sch14, Lemma 3.2.1], \mathcal{M} is anisotropic; hence, by [Sch14, Theorem (2), page 6], \mathcal{M} is isomorphic, as \mathbb{Q}_p -Lie algebra, to the derived algebra $[D, D]$ of a finite dimensional division \mathbb{Q}_p -algebra D . Let K be the center of D . Hence, K is a finite field extension of \mathbb{Q}_p , and D is a central division K -algebra. Let \mathcal{O}_K be the integral closure of \mathbb{Z}_p in K , and let Δ be the unique maximal \mathcal{O}_K -order in D . Observe that, since $sl_1(D) = [D, D]$, $sl_1(\Delta)$ is a \mathbb{Z}_p -Lie subalgebra of $[D, D]$. Let M be the \mathbb{Z}_p -Lie lattice inside \mathcal{M} that corresponds to $sl_1(\Delta)$ via an isomorphism $\mathcal{M} \rightarrow sl_1(D)$. There exists $m \in \mathbb{N}$ such that $p^m M \subseteq L$. By Theorem B, we can take $L_0 = p^m M$. \square

5 Proof of Theorem C, Theorem F, and Theorem G

Proof of Theorem C. For $n = 1$, the group $G := SL_1^{mne}(D)$ is trivial, and the statements are true. Assume that $n \geq 2$, and note that $p \geq 3$. We first argue that G is a uniform pro- p group whose associated powerful \mathbb{Z}_p -Lie lattice is $L_G = p^m sl_1(\Delta)$. For $p \geq 5$, this has already been observed, while when $p = 3$ we have $n = 2$ and $d = 1$, and one can consult, for instance, [Klo03, Section 2]. Now, item (1) follows from [Nos19, Proposition A] and Theorem B. For item (2), observe that $n \geq 3$, hence $p \geq 11$. By [Nos19, Theorem 3.1], H is saturable and the index of L_H in L_G is p , where L_H is the \mathbb{Z}_p -Lie lattice associated with H . Also, by [Gon07, Theorem B], $[L_G, L_G] = [G, G]$ and $[L_H, L_H] = [H, H]$. Observe that $p^{-m} L_H$ is a submodule of $p^{-m} L_G = sl_1(\Delta)$ of index p . By Theorem A, $[p^{-m} L_H, p^{-m} L_H] = [sl_1(\Delta), sl_1(\Delta)]$, hence, $[L_H, L_H] = [L_G, L_G]$. The desired $[H, H] = [G, G]$ follows. \square

Proof of Theorem F. We denote by $L^d(s)$ and $G^d(s)$ the Lie lattices and groups appearing in item (2) of the statements of Theorem E and Theorem F, respectively. We proceed with the proof by first assuming that G is isomorphic to one of the groups listed in the statement. Then G is solvable, and we can apply [Nos22a, Theorem A] to deduce the desired result (see also [Nos22a, Proposition 3.7]). Now assume that G is strongly hereditarily self-similar of index p . If $d \leq 2$ then G is solvable, and again we can apply [Nos22a, Theorem A]. We assume that $d \geq 3$ and observe that in this case $p \geq 5$. By [Nos22a, Remark 3.2], we can apply [Nos22a, Proposition 3.1] and deduce that G is saturable and the \mathbb{Z}_p -Lie lattice L associated with G is strongly hereditarily self-similar of index p . By Theorem E, L is isomorphic to either \mathbb{Z}_p^d or to $L^d(s)$ for some $s \in \mathbb{N}$; in fact, in the latter case we have $s \geq 1$, since L is residually nilpotent by [Gon09, Theorem B]. Then, since the group associated with \mathbb{Z}_p^d (respectively, $L^d(s)$) is \mathbb{Z}_p^d (respectively, $G^d(s)$), see [Nos22a, Remark 3.6], G is isomorphic to either \mathbb{Z}_p^d or $G^d(s)$, as desired. \square

Proof of Theorem G. Denote $d = \dim(G)$, and consider the following property.

- (0) $G \simeq \mathbb{Z}_p^d$, or $d \geq 2$ and there exists an integer $s \geq 1$ such that $G \simeq \mathbb{Z}_p \times \mathbb{Z}_p^{d-1}$, where \mathbb{Z}_p acts on \mathbb{Z}_p^{d-1} by multiplication by $1 + p^s$.

The equivalences (0) \Leftrightarrow (1), (0) \Leftrightarrow (3), (0) \Leftrightarrow (5), and (0) \Leftrightarrow (6) follow from Theorem F, [Klo11, Corollary 2.4], [Klo14, Corollary 1.13], and [Sno20, Theorem 1.2], respectively. The implication (0) \Rightarrow (2) is a consequence of [Nos22a, Theorem A, Theorem B] and the fact that the groups appearing in item (0) are solvable. The implication (2) \Rightarrow (4) follows from the Norm Residue Theorem (cf. [Hae19]). For $p = 2$, the implication (4) \Rightarrow (0) follows from the assumption $p > d$, so that $d = 1$ and $G \simeq \mathbb{Z}_2$; instead, for $p \neq 2$, the implication follows from [Qua14, Theorem B, Proposition 3.4] and the fact that a p -adic analytic pro- p group does not admit a nonabelian free pro- p subgroup. \square

References

- [Adk92] Adkins W. A., Weintraub S. H., *Algebra - An approach via module theory*, Graduate Texts in Mathematics, vol. 136, Springer-Verlag, 1992.
- [Ati69] Atiyah M. F.; Macdonald I. G., *Introduction to commutative algebra*, Addison-Wesley, 1969.
- [Bab20] Babai A., Fathalikhani K., Fernández Alcober G. A., Vannacci M., *On self-similar finite p -groups*, Groups Geom. Dyn. **14** (2020), 337–348.
- [Ber07] Berlatto A., Sidki S. N., *Virtual endomorphisms of nilpotent groups*, Groups Geom. Dyn. **1** (2007), 21–46.
- [Bou75] Bourbaki N., *Lie groups and Lie algebras - Chapters 1-3*, Springer, 1975.
- [Bou05] ———, *Lie groups and Lie algebras - Chapters 7-9*, Springer, 2005.
- [Bru67] Bruhat F., Tits J., *Groupes algébriques simples sur un corps local*, Proc. Conf. on Local Fields - Driebergen (Springer T.A., ed.), Springer, 1967, pp. 23–36.
- [Dix99] Dixon J. D., Sautoy M. P. F. du, Mann A., Segal D., *Analytic pro- p groups*, 2nd ed., Cambridge Studies in Advanced Mathematics, vol. 61, Cambridge University Press, 1999.
- [Ers07] Ershov M., *Finite presentability of $SL_1(D)$* , Israel Journal of Mathematics **158** (2007), 297–347.
- [Ers22] ———, *On the second cohomology of the norm one group of a p -adic division algebra*, arXiv:2206.12775v1, with an appendix by Ershov M. and Weigel T., 2022.
- [Gon07] González-Sánchez J., *On p -saturable groups*, J. Algebra **315** (2007), 809–823.
- [Gon09] González-Sánchez J., Klopsch B., *Analytic pro- p groups of small dimensions*, J. Group Theory **12** (2009), 711–734.

- [Gri80] Grigorchuk R. I., *On Burnside's problem on periodic groups*, Funktsional. Anal. i Prilozhen. **14** (1980), 53–54.
- [Gup83] Gupta N., Sidki S. N., *On the Burnside problem for periodic groups*, Math. Z. **182** (1983), 385–388.
- [Hae19] Haesemeyer C., Weibel C. A., *The norm residue theorem in motivic cohomology*, Annals of Mathematics Studies, vol. 200, Princeton University Press, 2019.
- [Hen98] Henn H.-W., *Centralizers of elementary abelian p -subgroups and mod- p cohomology of profinite groups*, Duke Mathematical Journal **91** (1998), no. 3, 561–585.
- [Jac62] Jacobson N., *Lie Algebras*, Dover Publications, 1962.
- [Kap12] Kapovich M., *Arithmetic aspects of self-similar groups*, Groups Geom. Dyn. **6** (2012), 737–754.
- [Klo03] Klopsch B., *Zeta functions related to the pro- p group $SL_1(\Delta_p)$* , Math. Proc. Camb. Phil. Soc. **135** (2003), 45–57.
- [Klo11] Klopsch B., Snopce I., *Pro- p groups with constant generating number on open subgroups*, Journal of Algebra **331** (2011), 263–270.
- [Klo14] ———, *A characterization of uniform pro- p groups*, Quart. J. Math. **65** (2014), 1277–1291.
- [Koc20] Kochloukova D. H., Sidki S. N., *Self-similar groups of type FP_n* , Geometriae Dedicata **204** (2020), 241–264.
- [Lam91] Lam T. Y., *A first course in noncommutative rings*, Springer-Verlag, 1991.
- [Laz65] Lazard M., *Groupes analytiques p -adiques*, Publ. Math. Inst. Hautes Études Sci. **26** (1965), 5–219.
- [Nek05] Nekrashevych V., *Self-similar groups*, Mathematical Surveys and Monographs 117, American Mathematical Society, 2005.
- [Nos19] Noseda F., Snopce I., *On self-similarity of p -adic analytic pro- p groups of small dimension*, Journal of Algebra **540** (2019), 317–345.
- [Nos21] ———, *Index-stable compact p -adic analytic groups*, Arch. Math. **116** (2021), 153–160, with an appendix by Jean-Pierre Serre.
- [Nos22a] ———, *On hereditarily self-similar p -adic analytic pro- p groups*, Groups Geom. Dyn. **16** (2022), no. 1, 85–114.
- [Nos22b] ———, *On the self-similarity index of p -adic analytic pro- p groups*, J. Group Theory **25** (2022), no. 3, 433–445.
- [Pra88] Prasad G., Raghunathan M. S., *Topological central extensions of $SL_1(D)$* , Inventiones Mathematicae **92** (1988), 645–689.
- [Qua14] Quadrelli C., *Bloch-Kato pro- p groups and locally powerful groups*, Forum Math. **26** (2014), 793–814.
- [Rav86] Ravenel D. C., *Complex cobordism and stable homotopy groups of spheres*, Academic Press, 1986.
- [Rei03] Reiner, I., *Maximal Orders*, London Mathematical Society Monographs New Series, Oxford University Press, USA, 2003.
- [Rie70] Riehm C., *The norm 1 group of p -adic division algebra*, American Journal of Mathematics **92** (1970), no. 2, 499–523.
- [Sch14] Schoeneberg, T., *Semisimple Lie algebras and their classification over p -adic fields*, Société Mathématique de France, 2014.
- [Sno20] Snopce I., Tanushevski S., *Frattini-injectivity and Maximal pro- p Galois groups*, arXiv:2009.09297, 2020.
- [Sun11] Šunic Z., *Finite self-similar p -groups with abelian first level stabilizers*, Internat. J. Algebra Comput. **21** (2011), 355–364.
- [War92] Ware R., *Galois groups of maximal p -extensions*, Transactions of the American Mathematical Society **333** (1992), no. 2, 721–728.

Francesco Nosedà
Mathematics Institute
Federal University of Rio de Janeiro
Av. Athos da Silveira Ramos, 149
21941-909, Rio de Janeiro, RJ
Brazil
nosedà@im.ufrj.br

Iliir Snopce
Mathematics Institute
Federal University of Rio de Janeiro
Av. Athos da Silveira Ramos, 149
21941-909, Rio de Janeiro, RJ
Brazil
ilir@im.ufrj.br

March 28, 2023