

WEIGHTED $L^p \rightarrow L^q$ -BOUNDEDNESS OF COMMUTATORS AND PARAPRODUCTS IN THE BLOOM SETTING

TIMO S. HÄNNINEN, EMIEL LORIST, AND JAAKKO SINKO

ABSTRACT. We characterize the $L^p(\mu) \rightarrow L^q(\lambda)$ boundedness of commutators of Calderón–Zygmund operators and pointwise multiplication in the upper triangular Bloom setting:

Let $1 < q < p < \infty$, $\mu \in A_p$, $\lambda \in A_q$ and define the Bloom weight ν by $\nu^{1/p+1/q'} := \mu^{1/p}\lambda^{-1/q}$. For a locally integrable function b and a non-degenerate Calderón–Zygmund operator T , the commutator $[b, T] : L^p(\mu) \rightarrow L^q(\lambda)$ is bounded if and only if $M_\nu^\# b \in L^r(\nu)$, where $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$ and $M_\nu^\# b$ is the weighted sharp maximal function defined by

$$M_\nu^\# b := \sup_Q \frac{1_Q}{\nu(Q)} \int_{\mathbb{R}^d} |b - \langle b \rangle_Q| dx.$$

In parallel to commutators, we also characterize the boundedness of dyadic paraproducts Π_b in the Bloom setting for all $p, q \in (1, \infty)$. This completes the characterisation of the $L^p(\mu) \rightarrow L^q(\lambda)$ -boundedness of both commutators of Calderón–Zygmund operators and pointwise multiplication and of paraproducts in the Bloom setting for all $p, q \in (1, \infty)$.

CONTENTS

1. Introduction	1
2. Preliminaries	6
3. Upper bounds	16
4. Lower bounds	24
5. Multiplier condition	31
Acknowledgements	36
References	36

1. INTRODUCTION

1.1. Commutators. The boundedness of the commutator $[b, T]$ of a Calderón–Zygmund singular integral operator T and pointwise multiplication by $b \in L^1_{\text{loc}}$ plays an important role in harmonic analysis, which dates back to the work of Nehari [Neh57] on the boundedness of $[b, H]$ on L^p , where H denotes the Hilbert transform. This result was later extended by Coifman, Rochberg, and Weis [CRW76], who showed that $[b, T] : L^p \rightarrow L^p$ is bounded for a wide class of Calderón–Zygmund operators T with convolution kernels, $b \in \text{BMO}$ and $p \in (1, \infty)$. Subsequent works have yielded a full characterization of those $b \in L^1_{\text{loc}}$ for which $[b, T] : L^p \rightarrow L^q$ is bounded for non-degenerate Calderón–Zygmund operators. The state of the art result can be formulated as follows: For $p, q \in (1, \infty)$, T a non-degenerate Calderón–Zygmund

2020 *Mathematics Subject Classification.* Primary: 42B20; Secondary: 47B47.

Key words and phrases. commutator, paraproduct, singular integral, Muckenhoupt weight.

operator and $b \in L^1_{\text{loc}}$, there holds

$$\| [b, T] \|_{L^p \rightarrow L^q} \approx \begin{cases} \|b\|_{\text{BMO}} & p = q, & [\text{CRW76}], \\ \|b\|_{\dot{C}^{0,\alpha}} & \frac{\alpha}{d} = \frac{1}{p} - \frac{1}{q} & p < q, & [\text{Jan78}], \\ \|b\|_{\dot{L}^r} & \frac{1}{r} = \frac{1}{q} - \frac{1}{p} & p > q, & [\text{Hyt21a}], \end{cases}$$

where $\dot{C}^{0,\alpha}$ denotes the space of all α -Hölder continuous functions and \dot{L}^r denotes the Lebesgue space L^r modulo constants. We refer to [Hyt21a] and [HOS22, Section 1.1] for a survey of all results leading up to this full characterization. An analogous characterization in a different context, namely on the boundedness of a Toeplitz type operator between Hardy spaces in the unit ball, was obtained by Pau and Perälä [PP20].

Weighted versions of commutator estimates in the case $p = q$ date back to the work of Strömberg, who gave an alternative proof of the boundedness of $[b, H]$ on L^p using the Fefferman-Stein maximal function, which also showed the boundedness of $[b, H]$ on $L^p(w)$ for any Muckenhoupt weight $w \in A_p$ (cf. [Tor86, p. 417]). A sharp version of this result in terms of the weight characteristic $[w]_{A_p}$ was obtained by Chung [Chu11], which was afterwards extended to any Calderón-Zygmund operator by Chung, Pereyra, and Perez [CPP12]. In particular, they showed that any Calderón-Zygmund operator T satisfies

$$(1.1) \quad \| [b, T] \|_{L^p(w) \rightarrow L^p(w)} \lesssim [w]_{A_p}^{2 \max\{1, \frac{1}{p-1}\}} \|b\|_{\text{BMO}}.$$

for all $w \in A_p$ and $p \in (1, \infty)$ and the dependence on $[w]_{A_p}$ is sharp.

These results have been generalized to a two-weight setting by taking weights $\mu, \lambda \in A_p$ and asking for a characterization of those $b \in L^1_{\text{loc}}$ such that $[b, T]$ is bounded from $L^p(\mu)$ to $L^p(\lambda)$. In this setting, long before even the one-weight works [Chu11, CPP12], Bloom [Blo85] proved that for $\mu, \lambda \in A_p$ and $p \in (1, \infty)$, it holds that $[b, H]: L^p(\mu) \rightarrow L^p(\lambda)$ is bounded if and only if $b \in \text{BMO}_\nu$, where $\nu := \mu^{1/p} \lambda^{-1/p}$, and BMO_ν is defined as the space of all $f \in L^1_{\text{loc}}$ such that

$$\|f\|_{\text{BMO}_\nu} := \sup_Q \frac{1}{\nu(Q)} \int_Q |f - \langle f \rangle_Q| dx < \infty,$$

where $\langle f \rangle_Q = \frac{1}{|Q|} \int_Q f$ and the supremum is taken over all cubes Q in \mathbb{R}^d . Note that $\text{BMO}_\nu = \text{BMO}$ when $\mu = \lambda$. The sufficiency of $b \in \text{BMO}_\nu$ for the boundedness of $[b, T]$ from $L^p(\mu)$ to $L^p(\lambda)$ was later extended to general Calderón-Zygmund operators by Segovia and Torrea [ST93]. We refer to [LOR21, Section 6] for a discussion of the necessity of the conditions $\mu, \lambda \in A_p$ for the boundedness of the commutator $[b, T]$ from $L^p(\mu)$ to $L^p(\lambda)$.

The study of characterizing the boundedness $[b, T]: L^p(\mu) \rightarrow L^q(\lambda)$ of commutators (or other operators) for $p, q \in (1, \infty)$ by means of conditions on the Bloom weight ν , which is connected to the two-weights μ and λ by the relation $\nu^{\frac{1}{p} + \frac{1}{q}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$, is called the *Bloom setting*. In this setting, the weights μ and λ are typically satisfying the a priori assumption $\mu \in A_p$ and $\lambda \in A_q$.

The Bloom setting has in recent years attracted renewed attention, starting with the works [HLW16, HLW17] of Holmes, Lacey and Wick. This has led to the following current state of the art result: For $1 < p \leq q < \infty$, $\mu \in A_p$, $\lambda \in A_q$, T a non-degenerate Calderón-Zygmund operator and $b \in L^1_{\text{loc}}$, there holds

$$\| [b, T] \|_{L^p(\mu) \rightarrow L^q(\lambda)} \lesssim \begin{cases} ([\mu]_{A_p} [\lambda]_{A_q})^{\max\{1, \frac{1}{p-1}\}} \|b\|_{\text{BMO}_\nu} & p = q, & [\text{LOR17}], \\ C_{\mu, \lambda} \|b\|_{\text{BMO}_\nu^\alpha} & \frac{\alpha}{d} = \frac{1}{p} - \frac{1}{q} & p < q, & [\text{HOS22}], \end{cases}$$

and

$$\|[b, T]\|_{L^p(\mu) \rightarrow L^q(\lambda)} \gtrsim C_{\mu, \lambda} \begin{cases} \|b\|_{\text{BMO}_\nu} & p = q, & [\text{Hyt21a}], \\ \|b\|_{\text{BMO}_\nu^\alpha} & \frac{\alpha}{d} = \frac{1}{p} - \frac{1}{q} \quad p < q, & [\text{HOS22}], \end{cases}$$

where $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$ and BMO_ν^α is defined as the space of all $f \in L^1_{\text{loc}}$ such that

$$\|f\|_{\text{BMO}_\nu^\alpha} := \sup_Q \frac{1}{\nu(Q)^{1+\frac{\alpha}{d}}} \int_Q |f - \langle f \rangle_Q| dx < \infty.$$

where the supremum is taken over all cubes Q in \mathbb{R}^d . We note that these results recover the unweighted setting for $p \leq q$, since $\text{BMO}_1^\alpha = \dot{C}^{0, \alpha}$. Moreover, we note that $\text{BMO}_\nu^0 = \text{BMO}_\nu$. We refer [HOS22, Section 1.2] for a survey of the results leading up to this characterization.

Any characterization for the off-diagonal (upper triangular) case $p > q$ in the Bloom setting has been missing so far. In this paper, we will characterize the boundedness of $[b, T]$ for non-degenerate Calderón–Zygmund operators in this case $1 < q < p < \infty$. This completes the characterisation of the boundedness of $[b, T]$ from $L^p(\mu)$ to $L^q(\lambda)$ for all $p, q \in (1, \infty)$ in the Bloom setting. The case $q < p$ is important, for example, for questions related to the Jacobian problem (see, e.g., [Hyt21b, Lin17, Lin22]).

Fix $1 < q < p < \infty$ and $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$. In the unweighted case, the sufficiency of the condition $b \in L^r$ for the boundedness of $[b, T]$ is very simple. Indeed, this is a direct consequence of the boundedness of T on L^p and L^q and Hölder's inequality:

$$\begin{aligned} \|[b, T]f\|_{L^q} &\leq \|bTf\|_{L^q} + \|T(bf)\|_{L^q} \leq \|b\|_{L^r} \|Tf\|_{L^p} + \|T\|_{L^q \rightarrow L^q} \|bf\|_{L^q} \\ &\leq (\|T\|_{L^p \rightarrow L^p} + \|T\|_{L^q \rightarrow L^q}) \|b\|_{L^r} \|f\|_{L^p}. \end{aligned}$$

The surprising fact from [Hyt21a] is that $b \in \dot{L}^r$ is actually also necessary for the boundedness of $[b, T]$ from L^p to L^q , proving that, after factoring out constants, there is no mutual cancellation between the two terms of the commutator. This is in stark contrast with the case $p = q$ and thus $r = \infty$, in which case the two individual terms are only bounded if $b \in L^\infty$ but combined one can allow $b \in \text{BMO}$ by the seminal result of Coifman, Rochberg and Weis [CRW76].

In the weighted setting it is well known that a Calderón–Zygmund operator T is bounded on $L^p(\mu)$ for $\mu \in A_p$. Translating above observations to the Bloom setting, one may therefore conjecture that the boundedness of $[b, T]$ from $L^p(\mu)$ to $L^q(\lambda)$ for $\mu \in A_p$ and $\lambda \in A_q$ can be characterized by the boundedness of the multiplication map $f \mapsto bf$, modulo constants. By Hölder's inequality, one easily checks that

$$\|f \mapsto bf\|_{L^p(\mu) \rightarrow L^q(\lambda)} = \|b\nu^{-1}\|_{L^r(\nu)},$$

which would suggest the right-hand side, modulo constants, as the canonical condition on b for the boundedness of $[b, T]$ from $L^p(\mu)$ to $L^q(\lambda)$. However, although this condition is clearly sufficient for the boundedness of $[b, T]$, it turns out to be non-necessary, as we shall prove in Section 5. Thus, in the Bloom setting, $[b, T]$ can be bounded from $L^p(\mu)$ to $L^q(\lambda)$ for more $b \in L^1_{\text{loc}}$ than those b for which the individual terms of the commutator are bounded, i.e. cancellation plays a role in the Bloom setting.

To state our condition on b , let us introduce a weighted sharp maximal function. For $f \in L^1_{\text{loc}}$ and a weight w , we define

$$M_w^\# f := \sup_Q \frac{\mathbf{1}_Q}{w(Q)} \int |f - \langle f \rangle_Q| dx.$$

We will show that boundedness of $[b, T]$ from $L^p(\mu)$ to $L^q(\lambda)$ for $\mu \in A_p$ and $\lambda \in A_q$ for $p > q$ can be characterized by the assumption $M_\nu^\# b \in L^r(\nu)$. Note that when

$\nu = 1$, we have

$$\|M_\nu^\# b\|_{L^r(\nu)} \approx \inf_c \|(b-c)\nu^{-1}\|_{L^r(\nu)},$$

by the classical result of Fefferman and Stein, which actually extends to the case $\nu \in A_{r'}$ (see Proposition 5.1). However, the assumptions on μ and λ only yield $\nu \in A_{2r'}$ (see Lemma 2.23), which explains why we cannot characterize the boundedness of $[b, T]$ from $L^p(\mu)$ to $L^q(\lambda)$ by

$$\inf_c \|(b-c)\nu^{-1}\|_{L^r(\nu)} < \infty.$$

Our main result for commutators reads as follows. We refer to Subsection 2.6 for a definition of non-degenerate ω -Calderón-Zygmund operators.

Theorem A. *Let $1 < q < p < \infty$, $\mu \in A_p$ and $\lambda \in A_q$. Set $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$ and $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. Let T be a non-degenerate ω -Calderón-Zygmund operator with Dini continuous kernel. For $b \in L_{\text{loc}}^1$, we have*

$$\|[b, T]\|_{L^p(\mu) \rightarrow L^q(\lambda)} \lesssim [\mu]_{A_p}^{\max\{1, \frac{1}{p-1}\}} [\lambda]_{A_q}^{\max\{1, \frac{1}{q-1}\}} \|M_\nu^\# b\|_{L^r(\nu)},$$

and

$$\|[b, T]\|_{L^p(\mu) \rightarrow L^q(\lambda)} \gtrsim C_{\mu, \lambda} \|M_\nu^\# b\|_{L^r(\nu)}.$$

The proof of the upper estimate in Theorem A can be found in Theorem 3.4 and the lower estimate in Corollary 4.3. These theorems actually prove more general statements:

- In Theorem 3.4 we also revisit the upper bound in the case $1 < p < q < \infty$ from [HOS22]. Indeed, we obtain a quantitative bound on $\|[b, T]\|_{L^p(\mu) \rightarrow L^q(\lambda)}$ in terms of $[\mu]_{A_p}$ and $[\lambda]_{A_q}$, which recovers the known sharp one-weight estimate in (1.1) for $\mu = \lambda$ and $q \rightarrow p$. Tracking this dependence in the proof of [HOS22, Theorem 2.12] would yield a worse dependence on $[\mu]_{A_p}$ and $[\lambda]_{A_q}$.
- In Theorem 4.1 we replace the conditions $\mu \in A_p$ and $\lambda \in A_q$ in the lower bound by a weaker condition on the pair of weights (μ, λ) .

In a follow-up paper, we will characterize when $[b, T]$ is compact from $L^p(\mu)$ to $L^q(\lambda)$ in the case $1 < q < p < \infty$, which in the unweighted setting was recently characterized by Hytönen, Li, Tao and Yang in [HLTY22]. Combined with the case $1 < p < q < \infty$ from [HOS22], this will also complete the characterization of the compactness of commutators in the Bloom setting.

1.2. Paraproducts. Paraproducts also play a vital role in harmonic analysis, for example in the celebrated dyadic representation theorem for Calderón-Zygmund operators by Hytönen [Hyt12, Hyt17]. A paraproduct Π_b with a $b \in L_{\text{loc}}^1$ is typically less singular than the commutator $[b, T]$ for a Calderón-Zygmund operator T . Indeed, using the aforementioned dyadic representation theorem, one can write the commutator of $[b, T]$ as the sum of compositions of paraproducts and Haar shift operators (cf. [HLW17, Section 5]). Consequently, the sharp exponent in the dependence on a weight characteristic is typically smaller for paraproducts than for commutators. For example, in the one-weight setting, it is well-known (see e.g. [Per19]) that for $p \in (1, \infty)$, $w \in A_p$ and $b \in L_{\text{loc}}^1$ the paraproduct Π_b is bounded on $L^p(w)$ with sharp dependence on $[w]_{A_p}$ given by

$$\|\Pi_b\|_{L^p(w) \rightarrow L^p(w)} \lesssim [w]_{A_p}^{\max\{1, \frac{1}{p-1}\}} \|b\|_{\text{BMO}},$$

which should be compared to (1.1).

In the recent work [FH22] by Fragkiadaki and Holmes Fay, the Bloom setting has also been investigated for paraproducts in the case $p = q$. They showed for $p \in (1, \infty)$, $\mu, \lambda \in A_p$ and $\nu := \mu^{1/p} \lambda^{-1/p}$ that

$$\|\Pi_b\|_{L^p(\mu) \rightarrow L^p(\lambda)} \lesssim [\mu]_{A_p}^{\frac{1}{p-1}} [\lambda]_{A_p} \|b\|_{\text{BMO}_\nu},$$

which should be compared to the result for commutators in [LOR17] mentioned before. Moreover, they showed that in case $p = 2$ and $\lambda = \mu^{-1}$ their estimate is sharp.

In this paper, we will give a full characterization of the boundedness of Π_b from $L^p(\lambda)$ to $L^q(\mu)$ for $p, q \in (1, \infty)$ and $b \in L^1_{\text{loc}}$ in the Bloom setting.

Our main result for paraproducts reads as follows. We refer to Subsection 2.8 for a definition of the paraproduct Π_b .

Theorem B. *Let $1 < p, q < \infty$, $\mu \in A_p$ and $\lambda \in A_q$. Set $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$, $\frac{\alpha}{d} := \frac{1}{p} - \frac{1}{q}$ and $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. For $b \in L^1_{\text{loc}}$ we have*

$$\|\Pi_b\|_{L^p(\mu) \rightarrow L^q(\lambda)} \lesssim [\mu]_{A_p}^{\frac{1}{p-1}} [\lambda]_{A_q} \begin{cases} \|b\|_{\text{BMO}_\nu^c} & p \leq q, \\ [\nu]_{A_\infty}^{1/r} \|M_\nu^\# b\|_{L^r(\nu)} & p > q, \end{cases}$$

and

$$\|\Pi_b\|_{L^p(\mu) \rightarrow L^q(\lambda)} \gtrsim C_{\mu, \lambda} \begin{cases} \|b\|_{\text{BMO}_\nu^c} & p \leq q, \\ \|M_\nu^\# b\|_{L^r(\nu)} & p > q. \end{cases}$$

We note that, in the setting of Theorem B, we always have $\nu \in A_{2r'}$ and hence $\nu \in A_\infty$. The proof of the upper estimate in Theorem B can be found in Theorem 3.10 (see also Remark 3.11) and the lower estimate in Theorem 4.4. Once again, these theorems actually prove more general statements:

- Viewing a paraproduct as a bilinear operator, one would expect the upper bound in Theorem B to hold for the pair of weights (μ, λ) in a genuinely multilinear weight class (cf. [ALM22, Section 4.6]). In Theorem 3.10 we will show that it suffices to assume the strictly weaker condition

$$\sup_Q \langle \mu^{1-p'} \rangle_Q^{\frac{1}{p'}} \langle \lambda \rangle_Q^{\frac{1}{q}} \langle \nu \rangle_Q^{\frac{1}{p} + \frac{1}{q'}} < \infty$$

for the upper bound to hold. In fact, this condition on the weights is also necessary for the upper bound (by $\|b\|_{\text{BMO}_\nu^c}$ when $q \geq p$, or by $\|M_\nu^\# b\|_{L^r(\nu)}$ when $q < p$), as we will show in Lemma 3.8.

- For the lower estimate, we again replace the conditions $\mu \in A_p$ and $\lambda \in A_q$ in Theorem 4.4 by a weaker condition on the pair of weights (μ, λ) .

1.3. **Outline.** This paper is organized as follows:

- In Section 2 we will discuss some preliminaries on dyadic analysis, weights, Calderón–Zygmund operators and paraproducts.
- In Section 3 we will prove the upper bounds for both commutators and paraproducts using sparse domination techniques.
- In Section 4 we will prove the lower bounds for both commutators and paraproducts.
- Finally, in Section 5 we will give an example of a $b \in L^1_{\text{loc}}$ which does not satisfy $\inf_c \|(b-c)\nu^{-1}\|_{L^r(\nu)} < \infty$, but for which the commutator $[b, T]$ and the paraproduct Π_b are bounded.

2. PRELIMINARIES

2.1. **Notation.** The standard or most used notation is summarized in the following table:

Q	Cube with sides parallel to the coordinate axes.
$\ell(Q)$	Sidelength of cube Q .
aQ	Cube concentric to cube Q with sidelength $a\ell(Q)$.
\mathcal{D}	Dyadic lattice: A countable collection of cubes such that $Q \cap R \in \{Q, R, \emptyset\}$ for all $Q, R \in \mathcal{D}$.
\widehat{Q}	Dyadic parent: Smallest cube $R \in \mathcal{D}$ such that $Q \subsetneq R$.
$\langle f \rangle_Q^\mu$	Average: $\langle f \rangle_Q^\mu := \frac{1}{\mu(Q)} \int_Q f \, d\mu$.
$D_Q f$	Difference of averages: $D_Q f := \sum_{R: \widehat{R}=Q} \langle f \rangle_R 1_R - \langle f \rangle_Q 1_Q$.
p'	Conjugate exponent of p : $\frac{1}{p'} + \frac{1}{p} := 1$.
α	Exponent defined by $\frac{\alpha}{d} := \frac{1}{p} - \frac{1}{q}$.
r	Exponent defined by $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$.
μ'	Dual weight of μ w.r.t. exponent p defined by $\mu' := \mu^{-p'/p}$.
λ'	Dual weight of λ w.r.t. exponent q defined by $\lambda' := \lambda^{-q'/q}$.
ν	Bloom weight defined by $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{1/p} \lambda^{-q}$.
$\ f\ _{\text{BMO}_\nu^\alpha}$	$\ f\ _{\text{BMO}_\nu^\alpha} := \sup_Q \frac{1}{\nu(Q)^{1+\frac{\alpha}{d}}} \int_Q f - \langle f \rangle_Q $
$\ f\ _{L^p(\mu)}$	Lebesgue norm: $\ f\ _{L^p(\mu)} := (\int_{\mathbb{R}^d} f ^p \, d\mu)^{1/p}$.
$\Pi_b f$	Dyadic paraproduct: $\Pi_b f := \sum_{Q \in \mathcal{D}} D_Q b \langle f \rangle_Q$.
$M^\mu f$	Maximal operator: $M^\mu f := \sup_Q \langle f \rangle_Q^\mu 1_Q$.
$M_\nu^\# b$	Weighted sharp maximal operator:

$$M_\nu^\# b := \sup_Q \frac{1}{\nu(Q)} \int |b - \langle b \rangle_Q| \, dx.$$

If μ is the Lebesgue measure, “ μ ” is omitted from the notation; similarly with λ and ν . The supremum \sup_Q is either over all cubes Q or over all dyadic cubes $Q \in \mathcal{D}$ in a dyadic lattice \mathcal{D} , as understood from the context. We will make extensive use of the notation “ \lesssim ” to indicate inequalities up to an implicit multiplicative constant. These constants may depend on p, q, d and also on e.g. the Calderón–Zygmund operator T and properties of its kernel and sparseness constants, but not on any of the functions under consideration. If these implicit constants depend on the weights μ, λ, ν , this will be denoted by “ $\lesssim_{\mu, \lambda, \nu}$ ”.

2.2. **Dyadic lattices.** By a *cube* Q we mean a cube with sides parallel to the coordinate axes.

Definition 2.1 (Dyadic lattice). A collection \mathcal{D} is called a *dyadic lattice* on the Euclidean space \mathbb{R}^d if $\mathcal{D} = \bigcup_{k \in \mathbb{Z}} \mathcal{D}_k$, where \mathcal{D}_k is a partition of the Euclidean space \mathbb{R}^d by cubes of side length 2^{-k} and the partition \mathcal{D}_{k+1} refines the partition \mathcal{D}_k . The cubes $Q \in \mathcal{D}$ of a dyadic lattice \mathcal{D} are called *dyadic cubes*. If \mathcal{D} is a dyadic lattice and $Q_0 \in \mathcal{D}$, we denote

$$\mathcal{D}(Q_0) := \{Q \in \mathcal{D} : Q \subseteq Q_0\}.$$

From the definition it follows that the dyadic lattice \mathcal{D} is a countable collection of cubes with *dyadic nestedness property*: $Q \cap R \in \{Q, R, \emptyset\}$ for all $Q, R \in \mathcal{D}$. For $Q \in \mathcal{D}$ we define its dyadic parent \widehat{Q} to be the smallest $R \in \mathcal{D}$ such that $Q \subsetneq R$.

Example. The dyadic lattice

$$\mathcal{D}^0 := \{2^{-k}([0, 1) + j) : k \in \mathbb{Z}, j \in \mathbb{Z}^d\}$$

is called *the standard dyadic lattice* and the dyadic lattices

$$\mathcal{D}^\alpha := \{2^{-k}([0, 1) + j + (-1)^k \alpha) : k \in \mathbb{Z}, j \in \mathbb{Z}^d\} \quad \text{where } \alpha \in \{0, \frac{1}{3}, \frac{2}{3}\}^d,$$

are examples of *shifted* dyadic lattices.

The geometric observation that generic cubes can be approximated by shifted dyadic cubes is known as *the one-third trick* [Oki92] and can be stated as follows (e.g. [HNVW16, Lemma 3.2.26]):

Lemma 2.2 (Shifted dyadic cubes approximate generic cubes: one-third trick). *For each cube $Q \subseteq \mathbb{R}^d$ there are $\alpha \in \{0, \frac{1}{3}, \frac{2}{3}\}^d$ and $R \in \mathcal{D}^\alpha$ such that $Q \subseteq R$ and $\ell(R) \leq 3\ell(Q)$.*

Thanks to this approximation, dyadic and non-dyadic quantities are often comparable when measures are doubling. As an example that we will use in what follows, we can compare dyadic and non-dyadic sharp maximal functions.

Corollary 2.3 (Dyadic and non-dyadic sharp maximal functions are comparable). *Let ν is a doubling measure and $f \in L^1_{\text{loc}}$. Then*

$$M_\nu^\# f(x) \approx_\nu \sup_{\alpha \in \{0, \frac{1}{3}, \frac{2}{3}\}^d} M_{\nu, \mathcal{D}^\alpha}^\# f(x), \quad x \in \mathbb{R}^d,$$

where in $M_{\nu, \mathcal{D}^\alpha}^\# f$ the supremum is taken over cubes $Q \in \mathcal{D}^\alpha$.

Proof. Let Q be a cube. Then there are $\alpha \in \{0, \frac{1}{3}, \frac{2}{3}\}^d$ and $R \in \mathcal{D}^\alpha$ such that $Q \subseteq R$ and $\ell(R) \leq 3\ell(Q)$. For one term, the observation that $R \subseteq 5Q$ together with the assumption that ν is doubling gives

$$\frac{1}{\nu(Q)} = \frac{\nu(5Q)}{\nu(Q)} \frac{1}{\nu(5Q)} \leq C_\nu \frac{1}{\nu(R)}.$$

For the other term, because $Q \subseteq R$, we obtain

$$\int_Q |b - \langle b \rangle_Q| dx \leq 2 \inf_c \int_Q |b - c| dx \leq 2 \inf_c \int_R |b - c| dx \leq 2 \int_R |b - \langle b \rangle_R| dx.$$

This finishes the proof. \square

2.3. Dyadic analysis. In what follows, we introduce Carleson and sparse collections, the basic estimates for them, and the related theory of the A_∞ measures. The results are well-known but perhaps somewhat scattered, so we hope that the systematic exposition in this section will be convenient for the reader.

Throughout this section an arbitrary dyadic lattice \mathcal{D} is fixed; for brevity “ \mathcal{D} ” is suppressed in the most notation. A basic tool in dyadic analysis is the well-known Hardy–Littlewood maximal inequality. The *dyadic Hardy–Littlewood maximal operator* M^μ is defined by

$$M^\mu f := \sup_{Q \in \mathcal{D}} \langle |f| \rangle_Q^\mu 1_Q.$$

Lemma 2.4 (Dyadic Hardy–Littlewood maximal inequality). *Let μ be a locally finite Borel measure and $p \in (1, \infty)$. Then*

$$\|M^\mu\|_{L^p(\mu)} \leq p' \|f\|_{L^p(\mu)}$$

for all $f \in L^p(\mu)$.

The Hardy–Littlewood maximal inequality and the Carleson embedding theorem are closely tied: each can be derived from the other. A family $\{s_Q\}_{Q \in \mathcal{D}}$ of complex numbers indexed by dyadic cubes satisfies *the μ -Carleson packing condition* if there is a constant $C > 0$ such that

$$\sum_{Q' \in \mathcal{D}: Q' \subseteq Q} |s_{Q'}| \mu(Q') \leq C \mu(Q)$$

for all cubes $Q \in \mathcal{D}$. The least constant in the estimate is called *the Carleson norm* and is denoted by $\|s\|_{\text{Car}(\mu)}$, i.e.

$$\|s\|_{\text{Car}(\mu)} := \sup_{Q \in \mathcal{D}} \frac{1}{\mu(Q)} \sum_{Q' \in \mathcal{D}: Q' \subseteq Q} |s_{Q'}| \mu(Q').$$

Lemma 2.5 (Dyadic Carleson embedding theorem). *Let μ be a locally finite Borel measure and let $p \in (1, \infty)$. Then*

$$\left(\sum_{Q \in \mathcal{D}} |\langle f \rangle_Q^\mu|^p s_Q \mu(Q) \right)^{1/p} \leq p' \|s\|_{\text{Car}(\mu)}^{1/p} \|f\|_{L^p(\mu)}.$$

for all $\{s_Q\}_{Q \in \mathcal{D}} \in \text{Car}(\mu)$ and $f \in L^p(\mu)$.

In this context the A_∞ -class enters because it is precisely the class that preserves the Carleson packing condition. A locally finite Borel measure w is said to satisfy *the dyadic Fujii–Wilson A_∞ -condition* with respect to a locally finite Borel measure μ - and this is denoted by $w \in A_\infty(\mu)$ - if

$$(2.1) \quad \int_Q \sup_{R \in \mathcal{D}(Q): x \in R} \frac{w(R)}{\mu(R)} d\mu(x) \leq C w(Q)$$

for all cubes $Q \in \mathcal{D}$. The least admissible constant C is denoted by $[w]_{A_\infty(\mu)}$ and is called *the dyadic Fujii–Wilson A_∞ -characteristic*.

The following lemma can be found, for example, in [HV21, Proposition 3.7].

Lemma 2.6 (Carleson condition is preserved by A_∞ measures). *Let w, μ be locally finite Borel measures. Then*

$$[w]_{A_\infty(\mu)} = \sup_{s := \{s_Q\}_{Q \in \mathcal{D}}} \frac{\|s\|_{\text{Car}(w)}}{\|s\|_{\text{Car}(\mu)}}.$$

This fact, combined with Lemma 2.5, implies the weighted version of the Carleson embedding theorem:

Lemma 2.7 (Weighted version of the dyadic Carleson embedding theorem). *Let μ be a locally finite Borel measure, $w \in A_\infty(\mu)$ and let $p \in (1, \infty)$. Then*

$$\left(\sum_{Q \in \mathcal{D}} |\langle f \rangle_Q^w|^p s_Q w(Q) \right)^{1/p} \leq p' [w]_{A_\infty(\mu)}^{1/p} \|s\|_{\text{Car}(\mu)}^{1/p} \|f\|_{L^p(w)}.$$

Dyadic cubes are ordered into a binary tree by nestedness. Iterated stopping time arguments along this tree generate Carleson collections of dyadic cubes. A collection \mathcal{S} of dyadic cubes is called *(C, μ)-Carleson* if there exists a constant $C > 0$ such that

$$\sum_{S' \in \mathcal{S}: S' \subseteq S} \mu(S') \leq C \mu(S)$$

for all $S \in \mathcal{S}$. The associated Carleson norm is given by

$$\|\mathcal{S}\|_{\text{Car}(\mu)} := \|1_{\mathcal{S}}\|_{\text{Car}(\mu)} = \sup_{S \in \mathcal{S}} \frac{1}{\mu(S)} \sum_{S' \in \mathcal{S}: S' \subseteq S} \mu(S').$$

Thus, a Carleson collection \mathcal{S} of dyadic cubes is a special case of a Carleson family of complex numbers via the correspondence $s_Q := 1_{\mathcal{S}}(Q)$. In fact, iterated stopping time arguments typically generate sparse collections, which are collections of large disjoint parts and hence in particular satisfy the Carleson packing condition.

Definition 2.8 (Sparseness). Let $\gamma \in (0, 1)$. A collection \mathcal{S} of dyadic cubes is called (γ, μ) -sparse if for each $S \in \mathcal{S}$ there exists $E_S \subseteq S$ such that $\mu(E_S) \geq \gamma\mu(S)$ and such that the sets $\{E_S\}_{S \in \mathcal{S}}$ are disjoint. If μ is the Lebesgue measure, it is omitted from the notation.

Indeed, every γ -sparse collection \mathcal{S} satisfies the γ^{-1} -Carleson packing condition:

$$\|\mathcal{S}\|_{\text{Car}(\mu)} := \sup_{S \in \mathcal{S}} \frac{1}{\mu(S)} \sum_{S' \in \mathcal{S}: S' \subseteq S} \mu(S') \leq \frac{1}{\gamma} \sup_{S \in \mathcal{S}} \frac{1}{\mu(S)} \sum_{S' \in \mathcal{S}: S' \subseteq S} \mu(E_{S'}) \leq \frac{1}{\gamma}.$$

For many purposes Carleson (or in particular sparse) collections behave like disjoint collections; an instance of this is captured by the following lemma:

Lemma 2.9. *Let μ be a locally finite Borel measure. Assume that \mathcal{S} is a μ -Carleson (or in particular μ -sparse) collection of dyadic cubes. Then*

$$\left\| \sum_{S \in \mathcal{S}} a_S 1_S \right\|_{L^p(\mu)} \leq p \|\mathcal{S}\|_{\text{Car}(\mu)}^{1/p'} \left(\sum_{S \in \mathcal{S}} |a_S|^p \mu(S) \right)^{1/p}$$

for all families $\{a_S\}_{S \in \mathcal{S}}$ of complex numbers.

Proof. By Hölder's inequality, we have

$$\int_{\mathbb{R}^d} \left(\sum_{S \in \mathcal{S}} a_S 1_S \right) g \, d\mu \leq \left(\sum_{S \in \mathcal{S}} |a_S|^p \mu(S) \right)^{1/p} \left(\sum_{S \in \mathcal{S}} (|g|_S^\mu)^{p'} \mu(S) \right)^{1/p'}.$$

From this the statement follows by $L^p - L^{p'}$ duality and the dyadic Carleson embedding theorem (Lemma 2.5). \square

As before, the fact that an A_∞ -measure preserves the Carleson packing condition leads to a weighted version of this lemma:

Lemma 2.10. *Let μ be a locally finite Borel measure and $w \in A_\infty(\mu)$. Assume that \mathcal{S} is a μ -Carleson (or, in particular, μ -sparse) collection of dyadic cubes. Then*

$$\left\| \sum_{S \in \mathcal{S}} a_S 1_S \right\|_{L^p(w)} \leq p [w]_{A_\infty(\mu)}^{1/p'} \|\mathcal{S}\|_{\text{Car}^\mathcal{P}(\mu)}^{1/p'} \left(\sum_{S \in \mathcal{S}} |a_S|^p w(S) \right)^{1/p}.$$

Proof. The statement follows by combining Lemma 2.6 and Lemma 2.9. \square

As mentioned above, every γ -sparse collection satisfies the γ^{-1} -Carleson packing condition. An obstruction for the converse is a point mass: a mass point can not be divided between two sets. Excluding this obstruction, the converse also holds (see [Ver96, Corollary 6]):

Lemma 2.11 (Carleson and sparse are equivalent). *Let μ be a locally finite Borel measure with no point masses. Then each (C, μ) -Carleson collection of dyadic cubes is $(\frac{1}{C}, \mu)$ -sparse.*

For different proofs and generalizations, see [LN18, CO17, Hän18, Bar19, Rey22]. As an immediate corollary of the equivalence of being Carleson and being sparse, we see that A_∞ -measures do not only preserve Carleson, but also sparseness:

Corollary 2.12 (A_∞ -measures preserve sparseness). *Let μ and w be measures with no point masses. Assume that $w \in A_\infty(\mu)$. Then every (γ, μ) -sparse collection is $(\gamma [w]_{A_\infty(\mu)}^{-1}, w)$ -sparse.*

Proof. The statement follows by combining Lemma 2.6 and Lemma 2.11. \square

When measures are doubling, the above estimates involving dyadic cubes translate easily into their counterparts involving generic cubes, by using shifted dyadic lattices (Lemma 2.2) in a typical fashion. In particular, we will need the following non-dyadic counterpart:

Lemma 2.13 (Weighted basic lemma for sparse collections of generic cubes). *Let $p \in (1, \infty)$. Assume that w and μ are doubling measures and that $w \in A_\infty(\mu)$. Assume that \mathcal{S} is a (γ, μ) -sparse countable collection of cubes. Then*

$$\left\| \sum_{S \in \mathcal{S}} a_S 1_S \right\|_{L^p(w)} \lesssim_{\mu, w} [w]_{A_\infty(\mu)}^{1/p'} \left(\sum_S |a_S|^p w(S) \right)^{1/p}.$$

The implicit constant depends on the measures μ and w via their doubling constants.

Proof. The lemma follows from its dyadic counterpart in Lemma 2.10 via shifted dyadic cubes (see Lemma 2.2). \square

2.4. Muckenhoupt weights. A locally integrable function $w: \mathbb{R}^d \rightarrow (0, \infty)$ is called a *weight*. By associating the measure $w(E) := \int_E w \, dx$ for measurable $E \subseteq \mathbb{R}^d$ to a weight w , the statements of the previous subsection are applicable to w .

For $p \in (1, \infty)$ the (non-dyadic) A_p -characteristic $[w]_{A_p}$ is defined by

$$[w]_{A_p} := \sup_Q \langle w \rangle_Q \langle w^{-p'/p} \rangle_Q^{p/p'},$$

and for $p = \infty$ the (non-dyadic) Fujii-Wilson A_∞ -characteristic from the previous section takes the form

$$[w]_{A_\infty} := \sup_Q \frac{1}{w(Q)} \int_Q \sup_{R \ni x} \frac{w(Q \cap R)}{|R|} \, dx,$$

where the suprema are over all cubes. We will write $w \in A_p$ if $[w]_{A_p} < \infty$.

Remark 2.14. The dyadic version of the A_p -characteristic is defined similarly, but with the suprema over dyadic cubes instead of generic cubes. The dyadic version is used in the context of dyadic operators (e.g. dyadic paraproducts), while in the context of non-dyadic operators (e.g. commutators) the non-dyadic version is used; this distinction is suppressed in the notation and understood from the context.

Example. The power weight $w_\delta(x) := |x|^\delta$ satisfies $w_\delta \in A_p$ if and only if $-d < \delta < (p-1)d$.

We will only need the following well-known properties for A_p -characteristic:

Lemma 2.15 (Duality and monotonicity of the A_p characteristic). *Let w be a weight.*

- (Duality) $[w]_{A_p}^{1/p} = [w^{-p'/p}]_{A_{p'}}^{1/p'}$ for $p \in (1, \infty)$.
- (Monotonicity) $[w]_{A_p} \geq [w]_{A_q}$ and $[w]_{A_p} \gtrsim [w]_{A_\infty}$ for $1 < p \leq q < \infty$.

Proof. The duality is clear from the definitions. The inequality $[w]_{A_p} \geq [w]_{A_q}$ for $1 < p \leq q < \infty$ follows by Jensen's inequality and for $[w]_{A_p} \gtrsim [w]_{A_\infty}$ we refer to [HP13, Proposition 2.2]. \square

We will use the following bound for the weighted norm of the maximal operator.

Lemma 2.16 ([Buc93]). *Let $p \in (1, \infty)$ and $w \in A_p$. Then*

$$\|Mf\|_{L^p(w)} \lesssim [w]_{A_p}^{\frac{1}{p-1}} \|f\|_{L^p(w)}$$

for all $f \in L^p(w)$.

2.5. Bloom weight and the Bloom–Muckenhoupt joint characteristics. An operator $U: L^p(\mu) \rightarrow L^q(\lambda)$ is bounded if and only if

$$\left| \int_{\mathbb{R}^d} (Uf)g \, dx \right| \leq C \|f\|_{L^p(\mu)} \|g\|_{L^{q'}(\lambda^{-q'/q})}$$

for all $f \in L^p(\mu)$ and $g \in L^{q'}(\lambda^{q'/q})$. The weight $\lambda^{-q'/q}$ appearing in this bilinear estimate is called the *dual weight* of the weight λ with respect to the weighted space $L^q(\lambda)$ and is denoted by $\lambda' := \lambda^{-q'/q}$. Similarly, we write $\mu' := \mu^{-p'/p}$ for the dual weight of the weight μ with respect to the weighted space $L^p(\mu)$.

Definition 2.17 (Bloom weight). Let μ, λ be weights and let $p, q \in (1, \infty)$. The Bloom weight $\nu = \nu_{L^p(\mu) \rightarrow L^q(\lambda)}$ associated with $L^p(\mu) \rightarrow L^q(\lambda)$ -boundedness is defined by

$$(2.2) \quad \nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{1/p} (\lambda')^{1/q'} = \mu^{1/p} \lambda^{-1/q}.$$

Remark 2.18. An operator $U: L^p(\mu) \rightarrow L^q(\lambda)$ is bounded if and only if $U^*: L^{q'}(\lambda') \rightarrow L^{p'}(\mu')$ is bounded, where U^* is the adjoint with respect to the unweighted integral dual pairing. Note that the Bloom weight is invariant under duality in that

$$\nu_{L^{q'}(\lambda') \rightarrow L^{p'}(\mu')} = \nu_{L^p(\mu) \rightarrow L^q(\lambda)}.$$

We will use joint weight characteristics for our upper and lower bounds in the Bloom setting, which we will introduce next. We note that these characteristics are all finite in case $\mu \in A_p$, $\lambda \in A_q$ and $\nu^{\frac{1}{p} + \frac{1}{q'}} = \mu^{1/p} \lambda^{-1/q}$, as we shall show after the definitions.

Definition 2.19 (Joint Bloom–Muckenhoupt characteristic for upper bounds). Let $p, q \in (1, \infty)$ and let μ, λ, ν be weights. We define

$$[\mu', \lambda, \nu]_{B_{p',q}} := \sup_Q \langle \mu' \rangle_Q^{1/p'} \langle \lambda \rangle_Q^{1/q} \langle \nu \rangle_Q^{1/p+1/q'},$$

and write $(\mu', \lambda, \nu) \in B_{p',q}$ if $[\mu', \lambda, \nu]_{B_{p',q}} < \infty$.

The class $B_{p',q}$ is exactly the class that plays an important role in the weighted $L^p \times L^{q'} \rightarrow L^{r'}$ -boundedness of bilinear operators for $\frac{1}{r'} = \frac{1}{p} + \frac{1}{q'}$ (cf. [LOP⁺09]).

Definition 2.20 (Joint Bloom–Muckenhoupt characteristic for lower bounds). Let $p, q \in (1, \infty)$ and let μ, λ, ν be weights. We define

$$[\mu, \lambda']_{B_{p,q'}(\nu)} := \sup_Q \left(\frac{\mu(Q)}{\nu(Q)} \right)^{1/p} \left(\frac{\lambda'(Q)}{\nu(Q)} \right)^{1/q'}$$

and write $(\mu, \lambda') \in B_{p,q'}(\nu)$ if $[\mu, \lambda']_{B_{p,q'}(\nu)} < \infty$.

We note that the class of weights $B_{p,q'}(\nu)$ is closely related to the Muckenhoupt classes used in weighted $L^{p'} \rightarrow L^{q'}$ -boundedness of linear operators, using ν as the base measure on \mathbb{R}^d (cf. [FH18]).

Remark 2.21.

- (i) Interpreting the lower bound Bloom characteristic as an assumption on the triplet of weights (μ, λ', ν) through

$$[\mu, \lambda', \nu]_{B_{p,q'}(\nu)} := \sup_Q \left(\frac{\mu(Q)}{\nu(Q)} \right)^{1/p} \left(\frac{\lambda'(Q)}{\nu(Q)} \right)^{1/q'} \left(\frac{\nu(Q)}{\nu(Q)} \right)^{1/p'+1/q'},$$

we see that the notation for the lower and upper bound Bloom characteristics could be unified, identifying the upper bound Bloom weight class with $B_{p',q}(1)$.

- (ii) Fix $b \in L^1_{\text{loc}}$. Note that the norms $\|b\|_{\text{BMO}_\nu^\alpha}$ and $\|M_\nu^\# b\|_{L^r(\nu)}$ are each decreasing as ν increases.
- (iii) For weights $(\mu, \lambda, \nu) \in B_{p',q}$ we have by the Lebesgue differentiation theorem that

$$\mu'(x)^{1/p'} \lambda(x)^{1/q} \nu(x)^{1/p+1/q'} \leq [\mu', \lambda, \nu]_{B_{p',q}}, \quad x \in \mathbb{R}^d$$

and thus

$$\nu^{\frac{1}{p} + \frac{1}{q'}} \leq [\mu', \lambda, \nu]_{B_{p',q}} \cdot \mu^{1/p} \lambda^{-1/q}.$$

Therefore, in view of (ii), we will always assume to be in the Bloom setting $\nu^{\frac{1}{p} + \frac{1}{q'}} = \mu^{1/p} \lambda^{-1/q}$ when proving upper bounds for weights in the class $B_{p',q}$.

- (iv) Similarly, for weights $(\mu, \lambda) \in B_{p,q'}(\nu)$ we have

$$\mu^{1/p} \lambda^{-1/q} \leq [\mu, \lambda']_{B_{p,q'}(\nu)} \cdot \nu^{\frac{1}{p} + \frac{1}{q'}}$$

and we will assume to be in the Bloom setting when proving lower bounds for weights in the class $[\mu, \lambda']_{B_{p,q'}(\nu)}$.

As already noted, the assumption that these joint characteristics are finite relaxes the typical Bloom-setting assumption that $\mu \in A_p$ and $\lambda \in A_q$ separately.

Lemma 2.22 (Separate Muckenhoupt conditions imply the joint conditions). *Let $p, q \in (1, \infty)$, $\mu \in A_p$, $\lambda \in A_q$ and let $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{1/p} \lambda^{-1/q}$. Then*

$$\begin{aligned} [\mu, \lambda']_{B_{p,q'}(\nu)} &\leq [\mu]_{A_p}^{1/p} [\lambda]_{A_q}^{1/q}, \\ [\mu', \lambda, \nu]_{B_{p',q}} &\leq [\mu]_{A_p}^{1/p} [\lambda]_{A_q}^{1/q}. \end{aligned}$$

Proof. The first estimate follows from Jensen's inequality, see [HOS22, Proposition 3.1]. The second estimate is a direct corollary of Hölder's inequality. \square

The joint Bloom characteristic for upper bounds $(\mu', \lambda, \nu) \in B_{p',q}$ with $\nu^{1/r'} := \mu^{1/p} \lambda^{-1/q}$ implies that $\nu \in A_{2r'}$ for $\frac{1}{r'} = \frac{1}{p} + \frac{1}{q'}$. So, in particular, $\nu \in A_{2r'}$ if $\mu \in A_p$ and $\lambda \in A_q$ and this result is sharp:

Lemma 2.23 (Muckenhoupt class for Bloom weight). *Let $p, q \in (1, \infty)$ and set $\frac{1}{r'} := \frac{1}{p} + \frac{1}{q'}$. Let μ, λ be weights and set $\nu^{1/r'} := \mu^{1/p} \lambda^{-1/q}$.*

- (i) $(\mu', \lambda, \nu) \in B_{p',q}$ if and only if $\mu' \in A_{2p'}$, $\lambda \in A_{2q}$ and $\nu \in A_{2r'}$.
- (ii) If $\mu \in A_p$, $\lambda \in A_q$, then $\nu \in A_{2r'}$ with

$$[\nu]_{A_{2r'}} \leq [\mu]_{A_p}^{r'/p} [\lambda]_{A_q}^{r'/q}.$$

- (iii) For each power weight $w \in A_{2r'}$ there exist power weights $\mu \in A_p$ and $\lambda \in A_q$ such that $w^{1/r'} = \mu^{1/p} \lambda^{-1/q} =: \nu^{1/r'}$.

Proof. For (i) we refer to [LOP⁺09, Theorem 3.6], from which the qualitative claim in (ii) follows by Lemma 2.22. The weight characteristic estimate in (ii) is shown in [HOS22, Proposition 3.1]. Finally, the statement in (iii) follows from the fact that $w_\delta := |x|^\delta$ satisfies $w_\delta \in A_s$ if and only if $-d < \delta < (s-1)d$. \square

We end this subsection with a lemma on A_∞ -conditions implied by the Bloom characteristic for lower bounds:

Lemma 2.24 (Weights satisfy the A_∞ -condition with respect to their Bloom weight).

Let $p, q \in (1, \infty)$, let μ, λ be weights and set $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{1/p} \lambda^{-1/q}$. Then

$$\begin{aligned} [\mu]_{A_\infty}(\nu) &\lesssim [\mu, \lambda']_{B_{p,q'}(\nu)}^p, \\ [\lambda']_{A_\infty}(\nu) &\lesssim [\mu, \lambda']_{B_{p,q'}(\nu)}^{q'} \end{aligned}$$

for the dyadic versions of the weight characteristics.

Remark 2.25. Assuming that the weights μ, λ, ν are doubling, these estimates hold also for the non-dyadic versions of the weight characteristics, but with the implicit constant depending on the doubling constants. This follows from the one-third trick.

Proof. Fix a cube Q . By Hölder's inequality and the definition of the dyadic weight characteristic,

$$\begin{aligned} & \int_{\mathbb{R}^d} \sup_{R \subseteq Q} 1_R \frac{\mu(R)}{\nu(R)} \, d\nu \\ & \leq \int_{\mathbb{R}^d} \left(\sup_{R \subseteq Q} 1_R \frac{\mu(R)}{\nu(R)} \right)^{1/p} \left(\sup_{R \subseteq Q} 1_R \frac{\mu(R)}{\nu(R)} \right)^{1/p'} \, d\nu \\ & \leq [\mu, \lambda']_{B_{p,q'}(\nu)} \int_{\mathbb{R}^d} \left(\sup_{R \subseteq Q} 1_R \frac{\lambda'(R)}{\nu(R)} \right)^{-1/q'} \left(\sup_{R \subseteq Q} 1_R \frac{\mu(R)}{\nu(R)} \right)^{1/p'} \, d\nu \\ & \leq [\mu, \lambda']_{B_{p,q'}(\nu)} \left(\int_{\mathbb{R}^d} \left(\sup_{R \subseteq Q} 1_R \frac{\lambda'(R)}{\nu(R)} \right)^{-p/q'} \, d\nu \right)^{1/p} \left(\int_{\mathbb{R}^d} \sup_{R \subseteq Q} 1_R \frac{\mu(R)}{\nu(R)} \, d\nu \right)^{1/p'}. \end{aligned}$$

Therefore, via a monotone convergence argument to ensure finiteness, we have

$$\int_{\mathbb{R}^d} \sup_{R \subseteq Q} 1_R \frac{\mu(R)}{\nu(R)} \, d\nu \leq [\mu, \lambda']_{B_{p,q'}(\nu)}^p \int_{\mathbb{R}^d} \left(\sup_{R \subseteq Q} 1_R \frac{\lambda'(R)}{\nu(R)} \right)^{-p/q'} \, d\nu.$$

Note that $\lambda'(Q) = \int_Q \lambda' \nu^{-1} \, d\nu$ and hence $\frac{\lambda'(Q)}{\nu(Q)} = \langle \lambda' \nu^{-1} \rangle_Q^\nu$. Let $\alpha \in (1, \infty)$ be an auxiliary exponent. Note that $t \mapsto t^{-p/(q'\alpha)}$ is convex. By Jensen's inequality, we have

$$\left(\frac{\lambda'(Q)}{\nu(Q)} \right)^{-p/q'} \leq \left(\langle (\lambda' \nu^{-1})^{-p/(q'\alpha)} \rangle_Q^\nu \right)^\alpha.$$

Therefore, by the dyadic Hardy–Littlewood maximal inequality,

$$\int_{\mathbb{R}^d} \left(\sup_{R \subseteq Q} 1_R \frac{\lambda'(R)}{\nu(R)} \right)^{-p/q'} \, d\nu \leq (\alpha')^\alpha \int_Q (\lambda' \nu^{-1})^{-p/q'} \, d\nu = \int_Q (\lambda' \nu^{-1})^{-p/q'} \nu \, dx.$$

Since $(\lambda' \nu^{-1})^{-p/q'} \nu = \mu$ by assumption, choosing $\alpha = 2$ yields

$$\frac{1}{\mu(Q)} \int_{\mathbb{R}^d} \sup_{R \subseteq Q} 1_R \frac{\mu(R)}{\nu(R)} \, d\nu \leq 4 [\mu, \lambda']_{B_{p,q'}(\nu)}^p$$

Taking the supremum over all dyadic cubes yields the first estimate, the second is proven similarly. \square

2.6. Calderón–Zygmund operators and kernels. We now turn to the definition of the operators under consideration. We start with the definition of Calderón–Zygmund operators and their kernels.

A function $\omega : [0, \infty) \rightarrow [0, \infty)$ is called a *modulus of continuity* if it is increasing and satisfies $\lim_{t \rightarrow 0} \omega(t) = 0$. It is said to satisfy *the Dini condition* if $\int_0^1 \frac{\omega(t)}{t} \, dt < \infty$.

Definition 2.26 (Non-degenerate ω -Calderón–Zygmund kernel). Let $\omega : [0, \infty) \rightarrow [0, \infty)$ be a modulus of continuity. A kernel $K : \mathbb{R}^d \times \mathbb{R}^d \setminus \{x = y\} \rightarrow \mathbb{C}$ is called a (*two-variable*) ω -Calderón–Zygmund kernel if it satisfies the standard size and continuity conditions: for $x \neq y$

$$|K(x, y)| \lesssim_K \frac{1}{|x - y|^d},$$

and for $|x - x'| \leq \frac{1}{2}|x - y|$

$$|K(x, y) - K(x', y)| + |K(y, x) - K(y, x')| \leq \frac{1}{|x - y|^d} \omega \left(\frac{|x - x'|}{|x - y|} \right).$$

The kernel $K : \mathbb{R}^d \times \mathbb{R}^d \setminus \{x = y\} \rightarrow \mathbb{C}$ is called *non-degenerate* if for each $x \in \mathbb{R}^d$ and $r > 0$ there is $y \in \mathbb{R}^d$ with $|x - y| \gtrsim_K r$ and

$$|K(y, x)| \gtrsim_K \frac{1}{r^d}.$$

The notion of non-degeneracy in Definition 2.26 was introduced by Hytönen in [Hyt21a], to which we refer for an overview of earlier non-degeneracy assumptions in the literature.

Definition 2.27 (ω -Calderón–Zygmund operator). Let $p \in (1, \infty)$ and let $T : L^p \rightarrow L^p$ be a bounded linear operator. Then T is called an ω -Calderón–Zygmund operator if there is an ω -Calderón–Zygmund kernel K such that for every $f \in L_c^\infty$ one has the kernel representation

$$Tf(x) = \int_{\mathbb{R}^d} K(x, y)f(y) dy \quad x \notin \text{supp}(f).$$

Estimates involving a Calderón–Zygmund operator will implicitly depend on the operator norm of T on L^p , the modulus of continuity ω , and the implicit constant in the definition of the kernel.

2.7. Commutators. Next, we turn to the definition of commutators between an ω -Calderón–Zygmund operator and pointwise multiplication by $b \in L_{\text{loc}}^1$. In general, the commutator

$$[b, T]f := bT(f) - T(bf)$$

may be undefined for $f \in L_c^\infty$, since for such f we only have $bf \in L^1$. If T is weak L^1 -bounded, this is not an issue, which is for example the case when ω satisfies the Dini condition. In this case we have the following identity:

Lemma 2.28. *Let T be a ω -Calderón–Zygmund operator and $b \in L_{\text{loc}}^1$. Suppose that $T : L^1 \rightarrow L^{1, \infty}$ is bounded. For $f \in L_c^\infty$ and $g \in L_c^\infty$ with supports separated by a positive distance, we have*

$$\langle g, [b, T]f \rangle = \langle b, gTf - fT^*g \rangle.$$

Remark 2.29. By the classical theory of Calderón–Zygmund operators, an ω -Calderón–Zygmund operator T is bounded from L^1 to $L^{1, \infty}$ if ω is Dini continuous.

Proof of Lemma 2.28. We start by noting that, by the support condition on f and g and the kernel estimates, we have $gTf, fT^*g \in L_c^\infty$, so the right-hand side is well-defined. It remains to show that

$$(2.3) \quad \langle g, T(bf) \rangle = \langle b, fT^*g \rangle.$$

Define $h = bf \in L^1$ and note that $T(h) \in L^{1, \infty}$ is well-defined. Moreover $T(h \mathbf{1}_{|h| \leq n}) \rightarrow T(h)$ in $L^{1, \infty}$ and thus $T(h \mathbf{1}_{|h| \leq n_k}) \rightarrow T(h)$ pointwise a.e. for some subsequence $(n_k)_{k \geq 1}$. Therefore, for $x \notin \text{supp}(f)$, we have by the dominated convergence theorem

$$T(bf)(x) = \lim_{k \rightarrow \infty} \int_{\mathbb{R}^d} K(x, y)h(y) \mathbf{1}_{|h| \leq n_k} dy = \int_{\mathbb{R}^d} K(x, y)b(y)f(y) dy.$$

from which (2.3) follows by Fubini's theorem. \square

Note that

$$\langle b, gTf - fT^*g \rangle = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (b(x) - b(y))K(x, y)f(y)g(x) dy dx.$$

for $f \in L_c^\infty$ and $g \in L_c^\infty$ with supports separated by a positive distance, where the integrand is Lebesgue integrable due to the estimates for ω -Calderón–Zygmund kernels.

In our lower bounds for commutators we will avoid well-definedness issues of $[b, T]$ for $b \in L_{\text{loc}}^1$ without assuming the $T : L^1 \rightarrow L^{1, \infty}$ boundedness (e.g. Dini continuity). Indeed, as in [Hyt21a], we will work under the assumption that we study an operator U_b with off-support kernel representation with ω -Calderón–Zygmund kernel,

$$\langle g, U_b f \rangle = \int \int (b(y) - b(x))K(y, x)f(x)g(y) dx dy,$$

for $f \in L_c^\infty$ and $g \in L_c^\infty$ with supports separated by a positive distance. Assumptions are formulated entirely in terms of the boundedness of U_b and this kernel representation, with no reference to T or $[b, T]$. The prototype of such an operator is of course the commutator $U_b := [b, T]$ with T an ω -Calderón–Zygmund operator with a Dini continuous ω .

2.8. Paraproducts. We end this section with the introduction of paraproducts. Let \mathcal{D} be a dyadic lattice and $b \in L_{\text{loc}}^1$. The difference of averages $D_Q b$ for $Q \in \mathcal{D}$ is defined by

$$D_Q b := \sum_{R \in \mathcal{D}: \tilde{R}=Q} \langle b \rangle_{R1_R} - \langle b \rangle_{Q1_Q}.$$

The difference $D_Q b$ can also be written in terms of Haar projections

$$D_Q b = \sum_{i \in \{0,1\}^d \setminus \{0\}} \langle b, h_Q^i \rangle h_Q^i,$$

where $(h_Q^i)_{i \in \{0,1\}^d}$ are the Haar functions associated with Q .

The paraproduct Π_b associated with $b \in L_{\text{loc}}^1$ is formally defined as

$$\Pi_b f := \sum_{Q \in \mathcal{D}} D_Q b \langle f \rangle_Q$$

whenever say $f \in L_c^\infty$. Since the assumption $b \in L_{\text{loc}}^1$ alone is insufficient to make sense of convergence of the sum over all dyadic cubes, we include into the definition an a priori assumption on unconditional convergence:

Definition 2.30 (Paraproduct). Let $b \in L_{\text{loc}}^1$. Assume moreover that b is such that for each $f \in L_c^\infty$ the sum

$$\sum_{Q \in \mathcal{D}} D_Q b \langle f \rangle_Q$$

over all dyadic cubes converges unconditionally in L_{loc}^1 , by which we mean unconditional convergence in $L^1(K)$ for every compact set K . Then the operator $\Pi_b : L_c^\infty \rightarrow L_{\text{loc}}^1$ defined by

$$\Pi_b f := \Pi_{b, \mathcal{D}} f := \sum_{Q \in \mathcal{D}} D_Q b \langle f \rangle_Q$$

is called the *dyadic paraproduct* associated with the function b .

We write $\Pi_{b, \mathcal{D}'} f := \sum_{Q \in \mathcal{D}'} D_Q b \langle f \rangle_Q$ when $\mathcal{D}' \subseteq \mathcal{D}$.

Lemma 2.31. *Let $b \in L^1_{\text{loc}}$. Assume that $\Pi_b : L^\infty_c \rightarrow L^1_{\text{loc}}$ is a paraproduct. Let $\mathcal{D}_n \subseteq \mathcal{D}$ such that $\bigcup_{n=1}^\infty \mathcal{D}_n = \mathcal{D}$. Then*

$$\lim_{n \rightarrow \infty} \int |1_Q \Pi_{b, \mathcal{D}_n} f| dx = \int |1_Q \Pi_{b, \mathcal{D}} f| dx$$

for $f \in L^\infty_c$ and Q a cube.

Proof. The lemma follows by the triangle inequality together with the unconditional convergence in $L^1(Q)$. \square

Stronger conditions on b ensure stronger unconditional convergence. In the unweighted setting, for example, if $b \in \text{BMO}$, then for every $f \in L^p$ the sum $\Pi_{b, \mathcal{D}} f$ converges unconditionally in L^p for every $p \in (1, \infty)$. In the extension of this to the Bloom setting, Burkholder's weak L^1 -inequality is a useful tool:

Lemma 2.32 (Burkholder). *Let \mathcal{D} be a dyadic lattice and let $(v_Q)_{Q \in \mathcal{D}}$ be a finitely nonzero sequence of scalars. We have for $f \in L^1$ and*

$$\left\| \sum_{Q \in \mathcal{D}} v_Q D_Q f \right\|_{L^{1, \infty}} \leq 2 \sup_{Q \in \mathcal{D}} |v_Q| \|f\|_{L^1}.$$

Proof. Viewing \mathcal{D} as a filtration on \mathbb{R}^d , we can view $\sum_{Q \in \mathcal{D}} v_Q D_Q f$ as a martingale transform. The statement then follows from Burkholder's weak L^1 -inequality for martingale transforms [Bur79]. \square

Remark 2.33. In \mathbb{R}^d there are two more paraproducts of interest, which can be treated using similar methods as employed in the current paper. We refer to [FH22] for a discussion of these paraproducts in the context of Bloom boundedness with $p = q$.

3. UPPER BOUNDS

In this section we will give sufficient conditions for the Bloom boundedness of commutators and paraproducts for all $1 < p, q < \infty$. The results in the case $q < p$ are entirely new, whereas in the case $p \leq q$ we obtain sharper quantitative bounds in terms of the weight characteristics than currently available in the literature.

3.1. Commutators. We start with the upper bound for commutators, for which we will use the main sparse domination result of [LOR17]. To state their result, for $f, b \in L^1_{\text{loc}}$ and a sparse family of cubes \mathcal{S} , we define the auxiliary sparse operator

$$\mathcal{A}_{\mathcal{S}, b} f(x) = \sum_{Q \in \mathcal{S}} |b - \langle b \rangle_Q| \langle f \rangle_Q \mathbf{1}_Q(x), \quad x \in \mathbb{R}^d,$$

and its formal adjoint

$$\mathcal{A}_{\mathcal{S}, b}^* f(x) = \sum_{Q \in \mathcal{S}} \langle |b - \langle b \rangle_Q| f \rangle_Q \mathbf{1}_Q(x), \quad x \in \mathbb{R}^d.$$

Theorem 3.1 ([LOR17, Theorem 1.1]). *Let T be an ω -Calderón-Zygmund operator with Dini continuous kernel and let $b \in L^1_{\text{loc}}$. For every $f \in L^\infty_c$, there exist 3^d dyadic lattices \mathcal{D}^k and $\frac{1}{2 \cdot 9^d}$ -sparse families $\mathcal{S}_k \subseteq \mathcal{D}^k$ such that*

$$|[b, T]f(x)| \lesssim \sum_{k=1}^{3^d} (\mathcal{A}_{\mathcal{S}_k, b} |f|(x) + \mathcal{A}_{\mathcal{S}_k, b}^* |f|(x)), \quad x \in \mathbb{R}^d.$$

In view of Theorem 3.1, to prove Bloom upper estimates for the commutators $[b, T]$, it suffices prove Bloom estimates for $\mathcal{A}_{\mathcal{S}, b}$ and $\mathcal{A}_{\mathcal{S}, b}^*$. We will need the following lemma, which is a special case of the main result in [FH18].

Lemma 3.2 ([FH18]). *Let $1 < p \leq q < \infty$, set $\frac{1}{s} := \frac{1}{p} + \frac{1}{q'}$ and let $w \in A_q$. Let \mathcal{D} be a dyadic lattice. For any γ -sparse collection $\mathcal{S} \subseteq \mathcal{D}$ we have*

$$\left\| \sum_{Q \in \mathcal{S}} \left(\frac{1}{|Q|^s} \int_Q |f|^s dx \right)^{1/s} \mathbf{1}_Q \right\|_{L^p(w^{p/q}) \rightarrow L^q(w)} \lesssim [w]_{A_q}^{\max\{\frac{1}{p'} + \frac{1}{q}, \frac{1}{q-1}\}}$$

Proof. Let $f \in L^p(w)$ and write

$$\mathcal{A}_{\mathcal{S},s} f := \sum_{Q \in \mathcal{S}} \left(\frac{1}{|Q|^s} \int_Q |f|^s dx \right)^{1/s} \mathbf{1}_Q.$$

Using [FH18, Theorem 1.1] with parameters $p = p/s$, $q = q/s$, $r = 1/s$, $\alpha = s$, $\omega = w$ and $\sigma = w^{-q'/q} = w'$, we obtain

$$\begin{aligned} \|\mathcal{A}_{\mathcal{S},s} f\|_{L^q(w)} &= \|(\mathcal{A}_{\mathcal{S},s} f)^s\|_{L^{q/s}(w)}^{1/s} \\ &\lesssim \left(\sup_{Q \in \mathcal{S}} |Q|^{-s} w(Q)^{\frac{s}{q}} w'(Q)^{\frac{p-s}{p}} ([w]_{A_\infty}^{\frac{s}{p'}} + [w']_{A_\infty}^{\frac{s}{q}}) \right)^{1/s} \left\| \frac{|f|^s}{w'} \right\|_{L^{p/s}(w')}^{1/s} \\ &\lesssim \sup_{Q \in \mathcal{S}} |Q|^{-1} w(Q)^{1/q} w'(Q)^{\frac{1}{q'}} ([w]_{A_\infty}^{\frac{1}{p'}} + [w']_{A_\infty}^{\frac{1}{q}}) \|f\|_{L^p(w^{p/q})}, \end{aligned}$$

where we used

$$\begin{aligned} (1 - q') \left(1 - \frac{p}{s} \right) &= (1 - q') \frac{p}{q'} = \frac{p}{q} \\ \frac{p-s}{ps} &= \frac{1}{s} - \frac{1}{p} = \frac{1}{q'}, \end{aligned}$$

in the last step. Noting that

$$\sup_{Q \in \mathcal{S}} |Q|^{-1} w(Q)^{1/q} w'(Q)^{\frac{1}{q'}} \leq [w]_{A_q}^{\frac{1}{q}},$$

and by Lemma 2.15

$$[w]_{A_\infty}^{\frac{1}{p'}} + [w']_{A_\infty}^{\frac{1}{q}} \leq [w]_{A_q}^{\frac{1}{p'}} + [w]_{A_q}^{\frac{1}{q(q-1)}} \leq 2 [w]_{A_q}^{\max\{\frac{1}{p'}, \frac{1}{q-1} - \frac{1}{q}\}},$$

finishes the proof. \square

Using Lemma 3.2, we can now prove Bloom estimates for $\mathcal{A}_{\mathcal{S},b}^*$.

Proposition 3.3 (Bloom estimate for sparse operator). *Let $1 < p, q < \infty$ and define $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$ and $\frac{\alpha}{d} := \frac{1}{p} - \frac{1}{q}$. Take $\mu \in A_p$, $\lambda \in A_q$ and set $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. Let \mathcal{D} be a dyadic lattice. For any γ -sparse collection $\mathcal{S} \subseteq \mathcal{D}$ and $b \in L_{\text{loc}}^1$ we have*

$$\|\mathcal{A}_{\mathcal{S},b}^*\|_{L^p(\mu) \rightarrow L^q(\lambda)} \lesssim [\mu]_{A_p}^{\max\{1, \frac{1}{p-1}\}} \begin{cases} [\lambda]_{A_q}^{\max\{\frac{1}{p'} + \frac{1}{q}, \frac{1}{q-1}\}} \cdot \|b\|_{\text{BMO}_\nu^\alpha} & p \leq q, \\ [\lambda]_{A_q}^{\max\{1, \frac{1}{q-1}\}} \cdot \|M_\nu^\sharp b\|_{L^r(\nu)} & p > q. \end{cases}$$

Proof. We start by noting that by [LOR17, Lemma 5.1], there is a $\frac{\gamma}{4}$ -sparse collection $\tilde{\mathcal{S}} \subset \mathcal{D}$ with $\mathcal{S} \subseteq \tilde{\mathcal{S}}$ so that for any $Q \in \mathcal{S}$ we have

$$\begin{aligned} \int_Q |b - \langle b \rangle_Q| |f| dx &\lesssim \int_Q \sum_{R \in \tilde{\mathcal{S}}: R \subseteq Q} \langle |b - \langle b \rangle_R| \rangle_R |f| \mathbf{1}_R dx \\ (3.1) \qquad \qquad \qquad &= \sum_{R \in \tilde{\mathcal{S}}: R \subseteq Q} \int_R |b - \langle b \rangle_R| \cdot \langle |f| \rangle_R dx. \end{aligned}$$

Furthermore, in order to use Proposition 3.2 efficiently, we define for $s \in (0, \infty)$

$$\mathcal{A}_{\mathcal{S},s} f := \sum_{Q \in \mathcal{S}} \left(\frac{1}{|Q|^s} \int_Q |f|^s dx \right)^{1/s} \mathbf{1}_Q.$$

We first consider the case $p \leq q$. Since $\frac{1}{s} := \frac{1}{p} + \frac{1}{q'} \geq 1$, we have by Minkowski's inequality

$$\begin{aligned} \sum_{R \in \tilde{\mathcal{S}}: R \subseteq Q} \int_R |b - \langle b \rangle_R| \cdot \langle |f| \rangle_R dx &\leq \|b\|_{\text{BMO}_\nu^\alpha} \sum_{R \in \tilde{\mathcal{S}}: R \subseteq Q} \nu(R)^{1/s} \langle |f| \rangle_R \\ &= \|b\|_{\text{BMO}_\nu^\alpha} \sum_{R \in \tilde{\mathcal{S}}: R \subseteq Q} \left(\int_Q \langle |f| \rangle_R^s \mathbf{1}_R d\nu \right)^{1/s} \\ &\leq \|b\|_{\text{BMO}_\nu^\alpha} \left(\int_Q \left(\sum_{R \in \tilde{\mathcal{S}}: R \subseteq Q} \langle |f| \rangle_R \mathbf{1}_R \right)^s d\nu \right)^{1/s}. \end{aligned}$$

And thus, combined with (3.1), Proposition 3.2 and the definition of ν , we obtain for $f \in L^p(\mu)$

$$\begin{aligned} \|\mathcal{A}_{\mathcal{S},b}^* f\|_{L^q(\lambda)} &\lesssim \|b\|_{\text{BMO}_\nu^\alpha} \|\mathcal{A}_{\mathcal{S},s}(\mathcal{A}_{\tilde{\mathcal{S}},1} f \cdot \nu^{1/s})\|_{L^q(\lambda)} \\ &\lesssim \|b\|_{\text{BMO}_\nu^\alpha} [\lambda]_{A_q}^{\max\{\frac{1}{p'} + \frac{1}{q}, \frac{1}{q-1}\}} \|\mathcal{A}_{\tilde{\mathcal{S}},1} f \cdot \nu^{1/s}\|_{L^p(\lambda^{p/q})} \\ &= \|b\|_{\text{BMO}_\nu^\alpha} [\lambda]_{A_q}^{\max\{\frac{1}{p'} + \frac{1}{q}, \frac{1}{q-1}\}} \|\mathcal{A}_{\tilde{\mathcal{S}},1} f\|_{L^p(\mu)} \\ &\lesssim \|b\|_{\text{BMO}_\nu^\alpha} [\lambda]_{A_q}^{\max\{\frac{1}{p'} + \frac{1}{q}, \frac{1}{q-1}\}} [\mu]_{A_p}^{\max\{1, \frac{1}{p-1}\}} \|f\|_{L^p(\mu)}, \end{aligned}$$

finishing the case $p \leq q$.

Next, we consider the case $p > q$. We have

$$\begin{aligned} \sum_{R \in \tilde{\mathcal{S}}: R \subseteq Q} \int_R |b - \langle b \rangle_R| \cdot \langle |f| \rangle_R dx &= \int_Q \sum_{R \in \tilde{\mathcal{S}}: R \subseteq Q} \frac{1}{\nu(R)} \int_R |b - \langle b \rangle_R| dx \cdot \langle |f| \rangle_R \mathbf{1}_R d\nu \\ &\leq \int_Q M_\nu^\sharp b \cdot \mathcal{A}_{\tilde{\mathcal{S}},1} f d\nu. \end{aligned}$$

Thus, combined with (3.1) and Proposition 3.2, we obtain for $f \in L^p(\mu)$

$$\begin{aligned} \|\mathcal{A}_{\mathcal{S},b}^* f\|_{L^q(\lambda)} &\lesssim \|\mathcal{A}_{\mathcal{S},1}(M_\nu^\sharp b \cdot \mathcal{A}_{\tilde{\mathcal{S}},1} f \cdot \nu)\|_{L^q(\lambda)} \\ &\lesssim [\lambda]_{A_q}^{\max\{1, \frac{1}{q-1}\}} \|M_\nu^\sharp b \cdot \mathcal{A}_{\tilde{\mathcal{S}},1} f \cdot \nu\|_{L^q(\lambda)}. \end{aligned}$$

Finally, using Hölder's inequality, the definition of ν and Proposition 3.2 once more, we obtain

$$\begin{aligned} \|M_\nu^\sharp b \cdot \mathcal{A}_{\tilde{\mathcal{S}},1} f \cdot \nu\|_{L^q(\lambda)} &= \|M_\nu^\sharp b \cdot \nu^{1/r} \cdot \mathcal{A}_{\tilde{\mathcal{S}},1} f \cdot \nu^{1/r'} \lambda^{1/q}\|_{L^q} \\ &\leq \|M_\nu^\sharp b \cdot \nu^{1/r}\|_{L^r} \|\mathcal{A}_{\tilde{\mathcal{S}},1} f \cdot \nu^{1/r'} \lambda^{1/q}\|_{L^p} \\ &= \|M_\nu^\sharp b\|_{L^r(\nu)} \|\mathcal{A}_{\tilde{\mathcal{S}},1} f\|_{L^p(\mu)} \\ &\lesssim [\mu]_{A_p}^{\max\{1, \frac{1}{p-1}\}} \|M_\nu^\sharp b\|_{L^r(\nu)} \|f\|_{L^p(\mu)}, \end{aligned}$$

finishing the proof. \square

Since $\mathcal{A}_{\mathcal{S},b}^*$ is the formal adjoint of $\mathcal{A}_{\mathcal{S},b}$, we can also deduce upper Bloom estimates for $\mathcal{A}_{\mathcal{S},b}$ from Proposition 3.3. Combining Theorem 3.1 and Proposition 3.3, we therefore obtain our desired result.

Theorem 3.4 (Bloom upper estimate for commutators). *Let $p, q \in (1, \infty)$ and define $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$ and $\frac{\alpha}{d} := \frac{1}{p} - \frac{1}{q}$. Take $\mu \in A_p$, $\lambda \in A_q$ and set $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. Let T be an ω -Calderón-Zygmund operator with Dini continuous ω and let $b \in L_{\text{loc}}^1$. We*

have

$$\| [b, T] \|_{L^p(\mu) \rightarrow L^q(\lambda)} \lesssim [\mu]_{A_p}^{\max\{1, \frac{1}{p-1}\}} [\lambda]_{A_q}^{\max\{1, \frac{1}{q-1}\}} \cdot \begin{cases} \|b\|_{\text{BMO}_\nu^\alpha} & p \leq q, \\ \|M_\nu^\# b\|_{L^r(\nu)} & p > q. \end{cases}$$

Proof. By Proposition 3.3 and duality, using $\frac{1}{p'} + \frac{1}{q} \leq 1$ when $p \leq q$, we have

$$\begin{aligned} \|\mathcal{A}_{\mathcal{S}, b}\|_{L^p(\mu) \rightarrow L^q(\lambda)} &= \|\mathcal{A}_{\mathcal{S}, b}^*\|_{L^{q'}(\lambda') \rightarrow L^{p'}(\mu')} \\ &\lesssim [\mu']_{A_{p'}}^{\max\{1, \frac{1}{p'-1}\}} [\lambda']_{A_{q'}}^{\max\{1, \frac{1}{q'-1}\}} \cdot \begin{cases} \|b\|_{\text{BMO}_\nu^\alpha} & p \leq q \\ \|M_\nu^\# b\|_{L^r(\nu)} & p > q \end{cases} \\ &= [\mu]_{A_p}^{\max\{1, \frac{1}{p-1}\}} [\lambda]_{A_q}^{\max\{1, \frac{1}{q-1}\}} \cdot \begin{cases} \|b\|_{\text{BMO}_\nu^\alpha} & p \leq q \\ \|M_\nu^\# b\|_{L^r(\nu)} & p > q. \end{cases} \end{aligned}$$

Therefore, the theorem follows from Theorem 3.1 and the density of the bounded, compactly supported functions in $L^p(\mu)$. \square

Remark 3.5. For $p = q$, Theorem 3.4 was already obtained in [HLW17]. We followed the proof using sparse domination from [LOR17], who considered the case $p = q$ as well. If $\mu = \lambda = w$ for $w \in A_p$, it is known that the dependence on $[w]_{A_p}$ in Theorem 3.4 is sharp, see [Chu11, Per19] and the references therein.

A qualitative version of Theorem 3.4 for $p > q$ was recently obtained in [HOS22]. Tracking the constants in [HOS22, Theorem 2.12] would yield quantitatively worse behaviour in $[\mu]_{A_p}$ and $[\lambda]_{A_q}$ than Theorem 3.4. Note that the dependence on $[\mu]_{A_p}$ and $[\lambda]_{A_q}$ in Theorem 3.4 can be slightly improved in the case $p > q$, using the full power of Proposition 3.3. Since we do not know if the obtained bound is sharp, we leave this to the interested reader.

3.2. Paraproducts. Next, we consider Bloom upper bounds for paraproducts. We start our analysis with a sparse domination result for finite truncations of Π_b for $b \in L_{\text{loc}}^1$, which generalizes [FH22, Theorem 3.1]. We will employ a stopping time argument that has two innovative features: “coupled stopping conditions” and “uniformity over parts of the input data”. In our further considerations it will be crucial that this sparse domination result still contains the terms $\langle |b - \langle b \rangle_Q| \rangle_Q$ rather than the more typical sparse estimate of paraproducts using $\|b\|_{\text{BMO}}$.

Theorem 3.6 (Sparse domination of paraproducts). *Let $f, b \in L_{\text{loc}}^1(\mathbb{R}^d)$, and let \mathcal{D} be a dyadic lattice. For every $Q_0 \in \mathcal{D}$, there exist a $\frac{1}{2^{d+2}}$ -sparse family $\mathcal{S} \subseteq \mathcal{D}(Q_0)$ such that for all finite collections $\mathcal{F} \subseteq \mathcal{D}(Q_0)$*

$$\left| \sum_{Q \in \mathcal{F}} D_Q b \langle f \rangle_Q \right| \lesssim \sum_{Q \in \mathcal{S}} \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q \mathbf{1}_Q.$$

Proof. We will show that for each $Q \in \mathcal{D}$ there exists a collection of pairwise disjoint cubes $\{P_k\}_k \subseteq \mathcal{D}(Q)$ such that $\sum_k |P_k| \leq \frac{1}{2} |Q|$ and such that

$$(3.2) \quad \left| \sum_{R \in \mathcal{D}(Q) \cap \mathcal{F}} D_R b \langle f \rangle_R \right| \leq 2^6 \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q \mathbf{1}_Q \\ + \sum_k |D_{\widehat{P}_k} b \langle |f| \rangle_{\widehat{P}_k} \mathbf{1}_{P_k}| + \sum_k \left| \sum_{R \in \mathcal{D}(P_k) \cap \mathcal{F}} D_R b \langle f \rangle_R \right|$$

uniformly over all finite collections $\mathcal{F} \subseteq \mathcal{D}$. Note that

$$|D_{\widehat{P}_k} b| \mathbf{1}_{P_k} = \langle b \rangle_{P_k} - \langle b \rangle_{\widehat{P}_k} \mathbf{1}_{P_k}.$$

Iterating the estimate (3.2), starting at the cube Q_0 , we obtain a $\frac{1}{2}$ -sparse collection $\tilde{\mathcal{S}} \subseteq \mathcal{D}(Q_0)$ such that

$$(3.3) \quad \left| \sum_{Q \in \tilde{\mathcal{F}}} D_Q b \langle f \rangle_Q \right| \leq \sum_{Q \in \tilde{\mathcal{S}}} 2^6 \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q \mathbf{1}_Q \\ + \sum_{Q \in \tilde{\mathcal{S}} \setminus \{Q_0\}} |\langle b \rangle_Q - \langle b \rangle_{\hat{Q}}| \langle |f| \rangle_{\hat{Q}} \mathbf{1}_Q$$

uniformly over all finite collections $\mathcal{F} \subseteq \mathcal{D}$. Noting that

$$\mathcal{S} := \tilde{\mathcal{S}} \cup \{\hat{Q} : Q \in \tilde{\mathcal{S}} \setminus \{Q_0\}\} \subseteq \mathcal{D}(Q_0)$$

is $\frac{1}{2^{d+2}}$ -sparse and for every $Q \in \mathcal{D}$ we have

$$\langle b \rangle_Q - \langle b \rangle_{\hat{Q}} \langle |f| \rangle_{\hat{Q}} \mathbf{1}_Q \leq 2^d \langle |b - \langle b \rangle_{\hat{Q}}| \rangle_{\hat{Q}} \langle |f| \rangle_{\hat{Q}} \mathbf{1}_Q$$

then yields the result.

Let us prove (3.2). Let $\{P_k\}_k$ be the collection of maximal cubes $P \in \mathcal{D}(Q)$ such that

$$(3.4) \quad \langle |f| \rangle_P > 4 \langle |f| \rangle_Q$$

or there exists a finite collection \mathcal{F}_P such that on P we have

$$(3.5) \quad \left| \sum_{R \in \mathcal{F}_P : P \supseteq R \subseteq Q} D_R b \langle f \rangle_R \right| > 2^5 \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q =: a_Q.$$

We remark that the left-hand side is constant on P .

For $R \in \mathcal{D}(Q)$ define

$$v_R := \begin{cases} \langle f \rangle_R & R \in \bigcup_k \{S \in \mathcal{F}_{P_k} : P_k \supseteq S \subseteq Q\} \\ 0 & \text{otherwise.} \end{cases}$$

By the stopping condition (3.4) we have $|v_R| \leq 4 \langle |f| \rangle_Q$ for $R \in \mathcal{D}(Q)$. Therefore, using the stopping conditions (3.4) and (3.5), Burkholder's weak L^1 -inequality for martingale differences (see Lemma 2.32) and the weak L^1 -boundedness of the dyadic maximal operator, we have

$$\sum_k |P_k| \leq \left| \left\{ \left| \sum_{R \in \mathcal{D}(Q)} v_R D_R b \right| > a_Q \right\} \right| + \left| \left\{ \sup_{R \in \mathcal{D}(Q)} \langle |f| \rangle_R > 4 \langle |f| \rangle_Q \right\} \right| \\ \leq \frac{2}{a_Q} \cdot \sup_{R \in \mathcal{D}(Q)} |v_R| \cdot \left\| \sum_{R \in \mathcal{D}(Q)} D_R b \right\|_{L^1} + \frac{1}{4 \langle |f| \rangle_Q} \cdot \|f \mathbf{1}_Q\|_{L^1} \\ \leq \frac{2}{2^5 \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q} \cdot 4 \langle |f| \rangle_Q \cdot \|b - \langle b \rangle_Q\|_{L^1(\mathbb{R}^d)} + \frac{|Q|}{4} \\ = \frac{1}{2} |Q|.$$

Now, to show (3.2), we write

$$\left| \sum_{R \in \mathcal{D}(Q) \cap \mathcal{F}} D_R b \langle f \rangle_R \right| \leq \left| \sum_{R \in \mathcal{D}(Q) \cap \mathcal{F}} D_R b \langle f \rangle_R \right| \mathbf{1}_{Q \setminus \bigcup_k P_k} \\ + \sum_k \left| \sum_{R \in \mathcal{F} : \hat{P}_k \supseteq R \subseteq Q} D_R b \langle f \rangle_R \right| \mathbf{1}_{P_k} \\ + \sum_k |D_{\hat{P}_k} b| \langle |f| \rangle_{\hat{P}_k} \mathbf{1}_{P_k} \\ + \sum_k \left| \sum_{R \in \mathcal{D}(P_k) \cap \mathcal{F}} D_R b \langle f \rangle_R \right|.$$

By the stopping condition (3.5), we have

$$\mathbf{1}_{Q \setminus \cup_k P_k} \left| \sum_{R \in \mathcal{D}(Q) \cap \mathcal{F}} D_R b \langle f \rangle_R \right| \leq 2^5 \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q,$$

and

$$\sum_k \mathbf{1}_{P_k} \left| \sum_{R \in \mathcal{F}: \hat{P}_k \subsetneq R \subset Q} D_R b \langle f \rangle_R \right| \leq 2^5 \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q,$$

both uniformly over all finite collections $\mathcal{F} \subseteq \mathcal{D}$. This concludes the proof of (3.2) and thus finishes the proof of the theorem. \square

Remark 3.7. The estimate in (3.3) is interesting in its own right, since it does not use the doubling property of the Lebesgue measure. In fact, it directly generalizes to the setting where one has a locally finite Borel measure μ on \mathbb{R}^d instead of the Lebesgue measure.

Let μ, λ, ν be weights and $f, b \in L^1_{\text{loc}}$. In the case $1 < p \leq q < \infty$ the sparse operator arising in Theorem 3.6 can be estimated as follows

$$\left\| \sum_{Q \in \mathcal{S}} \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q \mathbf{1}_Q \right\|_{L^q(\lambda)} \leq \left\| \sum_{Q \in \mathcal{S}} \frac{\nu(Q)^{\frac{1}{p} + \frac{1}{q'}}}{|Q|} \langle |f| \rangle_Q \mathbf{1}_Q \right\|_{L^q(\lambda)} \|b\|_{\text{BMO}_\nu^\alpha}.$$

Therefore, to prove the Bloom-weighted boundedness of the (finitely truncated) paraproduct $\Pi_{b, \mathcal{F}} f$, one could analyze the Bloom-weighted boundedness of the sparse operator

$$f \mapsto \sum_{Q \in \mathcal{S}} \frac{\nu(Q)^{\frac{1}{p} + \frac{1}{q'}}}{|Q|} \langle |f| \rangle_Q \mathbf{1}_Q.$$

In the case $p = q$ and $\mu, \lambda \in A_p$, this operator was studied in [FH22, Theorem 2.6]. Note that, viewing a paraproduct as a bilinear operator, one expects weighted boundedness to hold for (μ, λ) in a genuinely multilinear weight class (cf. [ALM22, Section 4.6]).

The next lemma identifies the canonical weight class for Bloom estimates of paraproducts for all $p, q \in (1, \infty)$. In Proposition 3.9, we will study Bloom estimates for the sparse operator arising in Theorem 3.6 using this weight class.

Lemma 3.8 (Bloom weight class for paraproducts). *Let $1 < p, q < \infty$ and define $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$ and $\frac{\alpha}{d} := \frac{1}{p} - \frac{1}{q}$. Let μ, λ, ν be weights and $\mu' = \mu^{-p'/p}$. Suppose that for all $b \in L^1_{\text{loc}}$ we have*

$$\|\Pi_b\|_{L^p(\mu) \rightarrow L^q(\lambda)} \lesssim \begin{cases} \|b\|_{\text{BMO}_\nu^\alpha} & p \leq q, \\ \|M_\nu^\sharp b\|_{L^r(\nu)} & p > q. \end{cases}$$

Then we have

$$[(\mu', \lambda, \nu)]_{B_{p', q}} = \sup_Q \langle \mu' \rangle_Q^{\frac{1}{p'}} \langle \lambda \rangle_Q^{\frac{1}{q}} \langle \nu \rangle_Q^{\frac{1}{p} + \frac{1}{q'}} < \infty.$$

Proof. For a fixed cube $Q \in \mathcal{D}$, define $b = \mathbf{1}_{Q_+} - \mathbf{1}_{Q_-}$, where Q_+ and Q_- are the right and left halves of Q along the first coordinate axis respectively. Note that

$$\begin{aligned} \|b\|_{\text{BMO}_\nu^\alpha} &= \frac{|Q|}{\nu(Q)^{\frac{1}{p} + \frac{1}{q'}}} & p \leq q, \\ \|M_\nu^\sharp b\|_{L^r(\nu)} &= \frac{|Q|}{\nu(Q)} \|M^\nu \mathbf{1}_Q\|_{L^r(\nu)} \leq r' \frac{|Q|}{\nu(Q)^{\frac{1}{p} + \frac{1}{q'}}} & p > q. \end{aligned}$$

Applying the boundedness of Π_b to $f = \mathbf{1}_Q \mu'$, we get

$$\langle \mu' \rangle_Q \cdot \|\mathbf{1}_Q\|_{L^q(\lambda)} \lesssim \frac{|Q|}{\nu(Q)^{\frac{1}{p} + \frac{1}{q'}}} \cdot \left(\int_Q \mu^{-\frac{p}{p-1}} d\mu \right)^{\frac{1}{p}}.$$

Rearranging the terms, we obtain

$$\langle \mu' \rangle_Q^{\frac{1}{p}} \langle \lambda \rangle_Q^{\frac{1}{q}} \langle \nu \rangle_Q^{\frac{1}{p} + \frac{1}{q'}} \leq C,$$

for some $C > 0$ independent of Q . \square

By Lemma 3.8 and Remark 2.21(iii) it is canonical to assume the Bloom relation $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$ when studying upper bounds for paraproducts. In this setting, the assumption $(\mu', \lambda, \nu) \in B_{p',q}$ is exactly the same as the assumption that $(\mu, \lambda') \in A_{(p,q')}$, where $A_{(p,q')}$ denotes a multilinear weight class (see [LOP⁺09]). This assumption is strictly weaker than $\mu \in A_p$ and $\lambda \in A_q$ (see Lemma 2.23).

In the following proposition we will prove the Bloom estimates for the sparse operator arising in Theorem 3.6 for weights $(\mu', \lambda, \nu) \in B_{p',q}$ satisfying the Bloom relation $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. Note that $\mu', \lambda, \nu \in A_\infty$ by Lemma 2.23.

Proposition 3.9. *Let $1 < p, q < \infty$ and define $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$ and $\frac{\alpha}{d} := \frac{1}{p} - \frac{1}{q}$. Let $(\mu', \lambda, \nu) \in B_{p',q}$ with $\mu' = \mu^{-p'/p}$ and $\nu^{\frac{1}{p} + \frac{1}{q'}} = \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. Let \mathcal{D} be a dyadic lattice and $\gamma \in (0, 1)$. For any γ -sparse family $\mathcal{S} \subseteq \mathcal{D}$ and $b \in L^1_{\text{loc}}$ we have*

$$\|f \mapsto \sum_{Q \in \mathcal{S}} \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q \mathbf{1}_Q\|_{L^p(\mu) \rightarrow L^q(\lambda)} \lesssim C_{\mu, \lambda, \nu} \cdot \begin{cases} \|b\|_{\text{BMO}_\nu^\alpha} & p \leq q, \\ \|M_\nu^\alpha b\|_{L^r(\nu)} & p > q, \end{cases}$$

where

$$C_{\mu, \lambda} = [(\mu', \lambda, \nu)]_{B_{p',q}} \cdot [\mu']_{A_\infty}^{1/p} [\lambda]_{A_\infty}^{1/q'} [\nu]_{A_\infty}^{(1/r)_+}.$$

Proof. Throughout the proof we will write and $r_+ = \frac{1}{(1/r)_+}$. Note that

$$\|h\|_{L^p(\mu)} = \|h/\mu'\|_{L^p(\mu')}, \quad h \in L^p(\mu),$$

so, by duality, it suffices to show

$$\sum_{Q \in \mathcal{S}} \int_Q |b - \langle b \rangle_Q| \langle |f| \mu' \rangle_Q \langle |g| \lambda \rangle_Q dx \lesssim C_{\mu, \lambda} \|f\|_{L^p(\mu')} \|g\|_{L^{q'}(\lambda)} \cdot \begin{cases} \|b\|_{\text{BMO}_\nu^\alpha} & p \leq q, \\ \|M_\nu^\alpha b\|_{L^r(\nu)} & p > q. \end{cases}$$

for $f \in L^p(\mu')$ and $g \in L^{q'}(\lambda)$. Since $\frac{1}{p} + \frac{1}{q'} + \frac{1}{r_+} \geq 1$, we have by Hölder's inequality and Lemma 2.7

$$\begin{aligned} & \sum_{Q \in \mathcal{S}} \int_Q |b - \langle b \rangle_Q| \cdot \langle |f| \mu' \rangle_Q \langle |g| \lambda \rangle_Q dx \\ &= \sum_{Q \in \mathcal{S}} \frac{1}{\nu(Q)^{\frac{1}{p} + \frac{1}{q'}}} \int_Q |b - \langle b \rangle_Q| \cdot \langle |f| \rangle_Q^{\mu'} \langle |g| \rangle_Q^\lambda \langle \mu' \rangle_Q \langle \lambda \rangle_Q \cdot \nu(Q)^{\frac{1}{p} + \frac{1}{q'}} dx \\ &\leq [(\mu', \lambda, \nu)]_{B_{p',q}} \sum_{Q \in \mathcal{S}} \frac{1}{\nu(Q)^{\frac{1}{p} + \frac{1}{q'}}} \int_Q |b - \langle b \rangle_Q| \cdot \langle |f| \rangle_Q^{\mu'} \mu'(Q)^{\frac{1}{p}} \cdot \langle |g| \rangle_Q^\lambda \lambda(Q)^{\frac{1}{q'}} dz \end{aligned}$$

$$\begin{aligned}
&\leq [(\mu', \lambda, \nu)]_{B_{p',q}} \left(\sum_{Q \in \mathcal{S}} (\langle |f| \rangle_Q^{\mu'})^p \mu'(Q) \right)^{1/p} \left(\sum_{Q \in \mathcal{S}} (\langle |g| \rangle_Q^\lambda)^{q'} \lambda(Q) \right)^{1/q'} \\
&\quad \cdot \left(\sum_{Q \in \mathcal{S}} \left(\frac{1}{\nu(Q)^{\frac{1}{p} + \frac{1}{q'}}} \int_Q |b - \langle b \rangle_Q| \, dx \right)^{r_+} \right)^{1/r_+} \\
&\leq \frac{pq'}{\gamma^{\frac{1}{p} + \frac{1}{q'}}} [(\mu', \lambda, \nu)]_{p,q} [\mu']_{A_\infty}^{1/p} [\lambda]_{A_\infty}^{1/q'} \|f\|_{L^p(\mu')} \|g\|_{L^{q'}(\lambda)} \\
&\quad \cdot \left(\sum_{Q \in \mathcal{S}} \left(\frac{1}{\nu(Q)^{1/r'}} \int_Q |b - \langle b \rangle_Q| \, dx \right)^{r_+} \right)^{1/r_+}.
\end{aligned}$$

Now if $p \leq q$, we have $r_+ = \infty$, in which case we see directly that

$$\left(\sum_{Q \in \mathcal{S}} \left(\frac{1}{\nu(Q)^{1/r'}} \int_Q |b - \langle b \rangle_Q| \, dx \right)^{r_+} \right)^{1/r_+} \leq \|b\|_{\text{BMO}_\nu^\alpha}.$$

For the case that $p > q$, we have $r_+ = r$ and by (2.12)

$$\begin{aligned}
&\left(\sum_{Q \in \mathcal{S}} \left(\frac{1}{\nu(Q)^{1/r'}} \int_Q |b - \langle b \rangle_Q| \, dx \right)^{r_+} \right)^{1/r_+} \\
&\leq \frac{[\nu]_{A_\infty}^{1/r}}{\gamma^{1/r}} \left(\sum_{Q \in \mathcal{S}} \left(\frac{1}{\nu(Q)} \int_Q |b - \langle b \rangle_Q| \, dx \right)^r \nu(E_Q) \right)^{1/r} \\
&\leq \frac{[\nu]_{A_\infty}^{1/r}}{\gamma^{1/r}} \left(\sum_{Q \in \mathcal{S}} \int_{E_Q} (M_\nu^\# b)^r \, d\nu \right)^{1/r} \leq \frac{[\nu]_{A_\infty}^{1/r}}{\gamma^{1/r}} \|M_\nu^\# b\|_{L^r(\nu)},
\end{aligned}$$

finishing the proof. \square

Combining Theorem 3.6 and Proposition 3.9, we now obtain a Bloom upper bound for paraproducts for weights $(\mu', \lambda, \nu) \in B_{p',q}$ with $\nu^{\frac{1}{p} + \frac{1}{q'}} = \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. The case $p = q$, $\mu \in A_p$ and $\lambda \in A_q$ has previously been obtained in [FH22].

Theorem 3.10 (Bloom upper estimate for paraproducts). *Let $1 < p, q < \infty$ and define $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$ and $\frac{\alpha}{d} := \frac{1}{p} - \frac{1}{q}$. Let $(\mu', \lambda, \nu) \in B_{p',q}$ with $\mu' = \mu^{-p'/p}$ and $\nu^{\frac{1}{p} + \frac{1}{q'}} = \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. Let \mathcal{D} be a dyadic lattice. For any $b \in L_{\text{loc}}^1$ we have*

$$\|\Pi_b\|_{L^p(\mu) \rightarrow L^q(\lambda)} \lesssim [(\mu', \lambda, \nu)]_{B_{p',q}} [\mu']_{A_\infty}^{1/p} [\lambda]_{A_\infty}^{1/q'} \cdot \begin{cases} \|b\|_{\text{BMO}_\nu^\alpha} & p \leq q, \\ [\nu]_{A_\infty}^{1/r} \|M_\nu^\# b\|_{L^r(\nu)} & p > q, \end{cases}$$

where

$$\Pi_b f := \sum_{Q \in \mathcal{D}} D_Q b \langle f \rangle_Q, \quad f \in L^p(\mu),$$

converges unconditionally in $L^q(\lambda)$ if $b \in \text{BMO}_\nu^\alpha$ when $p \leq q$ and if $M_\nu^\# b \in L^r(\nu)$ when $p > q$.

Proof. Let $f \in L^p(\mu)$. By density we may assume without loss of generality that f has compact support. Moreover, by considering quadrants separately if needed, we may assume that there is a $Q_0 \in \mathcal{D}$ such that $\text{supp } f \subseteq Q_0$. By Theorem 3.6, we can find a $\frac{1}{2^{d+2}}$ -sparse family of cubes $\mathcal{S} \subseteq \mathcal{D}(Q_0)$ such that for all finite collections $\mathcal{F} \subseteq \mathcal{D}(Q_0)$ we have

$$\left| \sum_{Q \in \mathcal{F}} D_Q b \langle f \rangle_Q \right| \leq 2^{d+5} \sum_{Q \in \mathcal{S}} \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q \mathbf{1}_Q,$$

Now set $\tilde{\mathcal{S}} = \mathcal{S} \cup \{Q \in \mathcal{D} : Q_0 \subsetneq Q\}$, which is $\frac{1}{2^{d+2}}$ -sparse as well. Since, for any $Q \in \mathcal{D}$, we have

$$\begin{aligned} |D_Q b| &= \left| \sum_{P \in \mathcal{D}: \tilde{P}=Q} \langle b \rangle_P \mathbf{1}_P - \langle b \rangle_Q \mathbf{1}_Q \right| \\ &\leq \sum_{P \in \mathcal{D}: \tilde{P}=Q} \langle |b - \langle b \rangle_Q| \rangle_P \mathbf{1}_P \leq 2^d \langle |b - \langle b \rangle_Q| \rangle_Q \mathbf{1}_Q, \end{aligned}$$

it follows that for all finite collections $\mathcal{F} \subseteq \mathcal{D}$ we have

$$\begin{aligned} \left| \sum_{Q \in \mathcal{F}} D_Q b \langle f \rangle_Q \right| &= \left| \sum_{Q \in \mathcal{F} \cap \mathcal{D}(Q_0)} D_Q b \langle f \rangle_Q \right| + \sum_{Q \in \mathcal{F}: Q_0 \subsetneq Q} |D_Q b \langle f \rangle_Q| \\ &\lesssim \sum_{Q \in \tilde{\mathcal{S}}} \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q \mathbf{1}_Q. \end{aligned}$$

Since the right-hand side is finite a.e. and does not depend on \mathcal{F} , this implies that $\sum_{Q \in \mathcal{D}} |D_Q b \langle f \rangle_Q|$ converges pointwise a.e. and for any choice of $\epsilon_Q \in \{0, 1\}$ we have

$$\left| \sum_{Q \in \mathcal{D}} \epsilon_Q D_Q b \langle f \rangle_Q \right| \lesssim \sum_{Q \in \tilde{\mathcal{S}}} \langle |b - \langle b \rangle_Q| \rangle_Q \langle |f| \rangle_Q \mathbf{1}_Q.$$

The norm estimate now follows from Proposition 3.9 and the unconditional convergence follows from the dominated convergence theorem. \square

Remark 3.11. Let $p, q \in (1, \infty)$, $\mu \in A_p$ and $\lambda \in A_p$ and set $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. By Lemma 2.15 and Lemma 2.22 we have

$$[(\mu', \lambda, \nu)]_{B_{p', q}} [\mu']_{A_\infty}^{1/p} [\lambda]_{A_\infty}^{1/q'} \lesssim [\mu]_{A_p}^{\frac{1}{p-1}} [\lambda]_{A_q},$$

so, in particular, Theorem 3.10 proves the upper bound in Theorem B.

Let $w \in A_2$. In the case $p = q = 2$, $\mu = w$, $\lambda = w^{-1}$ and thus $\nu = w$, it was shown in [FH22, Section 3.1] that there is a $b \in \text{BMO}_w$ such that

$$\|\Pi_b\|_{L^2(w) \rightarrow L^2(w^{-1})} \gtrsim \|b\|_{\text{BMO}_w} [w]_{A_2}^2$$

In this setting we have

$$[\mu]_{A_2} [\lambda]_{A_2} = [w]_{A_2}^2,$$

so our result, as well as [FH22, Theorem 2.6], is sharp. We do not know if Theorem 3.10 is also sharp for other parameters.

Remark 3.12. We note that the claimed unconditional convergence in 3.10 is not automatic, since we do not have $\lambda \in A_q$ in general (see Lemma 2.23). As a consequence, we note that the functions with finite Haar expansion are not necessarily dense in $L^q(\lambda)$.

4. LOWER BOUNDS

We next turn to lower bounds for commutators and paraproducts. Throughout the section, recall the following for weights μ, λ, ν :

- We set $\lambda' := \lambda^{-q'/q}$ and

$$[\mu, \lambda']_{B_{p, q'}(\nu)} := \sup_Q \left(\frac{\mu(Q)}{\nu(Q)} \right)^{1/p} \left(\frac{\lambda'(Q)}{\nu(Q)} \right)^{1/q'}.$$

- Suppose $\nu^{\frac{1}{p} + \frac{1}{q'}} = \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. Then the condition $(\mu, \lambda') \in B_{p, q'}(\nu)$ implies the conditions $\mu \in A_\infty(\nu)$ and $\lambda' \in A_\infty(\nu)$ by Lemma 2.24.

4.1. Statement and overview of proof for commutators. We start by proving that, in the Bloom setting, the sharp maximal function condition

$$\|M_\nu^\# b\|_{L^r(\nu)} < +\infty$$

is necessary for the $L^p(\mu) \rightarrow L^q(\lambda)$ boundedness of commutators $[b, T]$ in the range $1 < q < p < \infty$. We will prove this under weaker assumptions on the weights than $\mu \in A_p$, $\lambda \in A_q$.

The main challenge in the Bloom setting for $1 < q < p < \infty$ is finding a condition that is not only sufficient but also necessary. The proof of the lower bound builds upon techniques from [HHL16, Hyt21a, HOS22], and upon weighted sparse analysis.

Theorem 4.1 (Sharp maximal function condition is necessary in the Bloom setting). *Let $1 < q < p < \infty$ and set $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$. Let $b \in L_{\text{loc}}^1$ and K be a non-degenerate ω -Calderón-Zygmund kernel. Assume the following:*

- (Weights) *Let μ and λ be weights and set $\lambda' := \lambda^{-q'/q}$ and $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. Assume that μ, λ', ν are doubling and $(\mu, \lambda') \in B_{p, q'}(\nu)$.*
- (Boundedness) *Assume that $U_b : L^p(\mu) \rightarrow L^q(\lambda)$ is a bounded linear operator.*
- (Off-support bilinear integral form representation) *Assume that*

$$\int_{\mathbb{R}^d} g U_b f \, dx = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (b(y) - b(x)) K(y, x) f(x) g(y) \, dx \, dy$$

whenever the functions $f \in L_c^\infty$ and $g \in L_c^\infty$ have supports separated by a positive distance.

Then

$$\|M_\nu^\# b\|_{L^r(\nu)} \lesssim_{\mu, \lambda', \nu} [\mu, \lambda']_{B_{p, q'}(\nu)} [\mu]_{A_\infty(\nu)}^{1/p'} [\lambda']_{A_\infty(\nu)}^{1/q} \|U_b\|_{L^p(\mu) \rightarrow L^q(\nu)},$$

where the implicit constant depends on the doubling constants of μ, λ', ν .

Remark 4.2 (Unweighted setting). In the unweighted case $\mu = \lambda = \nu = 1$ the theorem recovers [Hyt21a, Theorem 2.5.1] because of the well-known comparison

$$\|M^\# b\|_{L^r} \approx \inf_c \|b - c\|_{L^r}.$$

From Theorem 4.1 we can easily deduce the lower bound in Theorem A.

Corollary 4.3 (Lower bound for commutator). *Let $1 < q < p < \infty$, $\mu \in A_p$ and $\lambda \in A_q$. Set $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$ and $\nu^{1/p + 1/q'} := \mu^{1/p} \lambda^{-1/q}$. Let T be a non-degenerate ω -Calderón-Zygmund operator with Dini continuous ω , and let $b \in L_{\text{loc}}^1$. Then*

$$\|M_\nu^\# b\|_{L^r(\nu)} \lesssim C_{\mu, \lambda} \|[b, T]\|_{L^p(\mu) \rightarrow L^q(\lambda)}.$$

Proof of Corollary 4.3. This follows from observing that the assumptions of Corollary 4.3 imply the assumptions of Theorem 4.1, as follows. From $\mu \in A_p$ and $\lambda \in A_q$ (or equivalently $\lambda' \in A_{q'}$) it follows that $(\mu, \lambda') \in B_{p, q'}(\nu)$ by Lemma 2.22 and that $\nu \in A_{2r'}$ by Lemma 2.23. In particular, μ, λ', ν are doubling since they are A_∞ -weights. Since ω is Dini continuous, T is bounded from L^1 to $L^{1, \infty}$. Therefore, by Lemma 2.28, we have the kernel representation

$$\int f [b, T] g = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (b(x) - b(y)) K(x, y) f(y) g(x) \, dy \, dx$$

whenever the functions $f \in L_c^\infty$ and $g \in L_c^\infty$ have supports separated by a positive distance. \square

Proof of Theorem 4.1. The proof proceeds in three main steps: (1) discretization of the Lebesgue norm of the sharp maximal function, (2) control of mean oscillations by bilinear forms, and (3) use of a sequential-type testing condition on bilinear forms for general abstract operators. Each of these steps is stated as a separate lemma. How the lemmas are combined to yield the theorem is detailed in what follows.

First, we discretize. Fix a parameter $\gamma \in (0, 1)$. By combining Corollary 2.3 and Lemma 4.7, we obtain

$$\|M_\nu^\#\|_{L^r(\nu)} \approx_\nu \sup_{\mathcal{D}} \sup_{S \subseteq \mathcal{D}} \left(\sum_{S \in \mathcal{S}} \left(\frac{1}{\nu(S)} \int_S |b - \langle b \rangle_S| dx \right)^r \nu(S) \right)^{1/r},$$

where the supremum is taken over all dyadic lattices \mathcal{D} and over all collections $S \subseteq \mathcal{D}$ that are (γ, ν) -sparse in the lattice \mathcal{D} .

Second, we control each oscillation by testing a bilinear form against a pair of test functions. By Lemma 4.8, for each cube $S \in \mathcal{S}$ there exist a cube \tilde{S} with $\ell(\tilde{S}) \sim \ell(S)$ and $\text{dist}(\tilde{S}, S) \sim \ell(S)$ and functions $f_S^i, g_{\tilde{S}}^i$ with $|f_S^i| \lesssim 1_S$, $|g_{\tilde{S}}^i| \lesssim 1_{\tilde{S}}$ for $i = 1, 2$ such that

$$\int_S |b - \langle b \rangle_S| dx \lesssim \sum_{i=1}^2 \left| \int_{\mathbb{R}^d} f_S^i U_b g_{\tilde{S}}^i dx \right|$$

Therefore,

$$\left(\sum_{S \in \mathcal{S}} \left(\frac{1}{\nu(S)} \int_S |b - \langle b \rangle_S| dx \right)^r \nu(S) \right)^{1/r} \leq \sum_{i=1}^2 \left(\sum_{S \in \mathcal{S}} \left| \frac{1}{\nu(S)} \int_{\mathbb{R}^d} f_S^i U_b g_{\tilde{S}}^i dx \right|^r \nu(S) \right)^{1/r}.$$

Finally, the right hand-side, the so-called sequential testing condition on bilinear forms, is bounded by the operator norm by Lemma 4.10. The proof is completed. \square

4.2. Statement and overview of proof for paraproducts. The lower bound for dyadic paraproducts is stated and proved as for commutators, except that the structure of dyadic paraproducts - in particular the use of dyadic cubes instead of generic cubes - simplifies estimations.

Theorem 4.4 (Necessity in the Bloom setting for paraproducts). *Let $p, q \in (1, \infty)$, set $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$ and $\frac{\alpha}{d} := \frac{1}{p} - \frac{1}{q}$. Let μ and λ be weights and set $\lambda' := \lambda^{-q'/q}$ and $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. Assume that $\Pi_b : L^p(\mu) \rightarrow L^q(\lambda)$ is bounded and that $(\mu, \lambda') \in B_{p, q'}(\nu)$. Then the following holds:*

$$\begin{aligned} \|b\|_{\text{BMO}_\nu^\alpha} &\lesssim [\mu, \lambda']_{B_{p, q'}(\nu)} \|\Pi_b\|_{L^p(\mu) \rightarrow L^q(\lambda)} & p \leq q, \\ \|M_\nu^\# b\|_{L^r(\nu)} &\lesssim [\mu, \lambda']_{B_{p, q'}(\nu)} [\mu]_{A_\infty(\nu)}^{1/p'} [\lambda']_{A_\infty(\nu)}^{1/q} \|\Pi_b\|_{L^p(\mu) \rightarrow L^q(\lambda)} & p > q. \end{aligned}$$

Remark 4.5. In this dyadic context the suprema in the weighted sharp maximal function, the space BMO_ν^α and in all the weight characteristics are taken over the dyadic lattice associated with the paraproduct.

Remark 4.6. Note that the assumption $(\mu, \lambda') \in B_{p, q'}(\nu)$ is weaker than the assumption $\mu \in A_p$ and $\lambda \in A_q$ by Lemma 2.22.

Proof of Theorem 4.4. The case $q < p$ follows from combining Lemmas 4.7, 4.9, and 4.10. Note that no dependence on doubling constants occurs in this dyadic context because of the following: (1) The dyadic version of the weighted dyadic sharp maximal function is used. (2) The pairs of test functions that Lemma 4.9 gives to Lemma 4.10 are of the form $(f_{Q_S}, g_{Q_S}) = (1_S, 1_S)$. Therefore, inside the proof of Lemma 4.10, we can use Lemma 2.10 (dyadic estimate) instead of Lemma

2.13 (non-dyadic estimate). The case $p \leq q$ follows by combining Lemma 4.9 and Lemma 4.13. \square

4.3. Discretizing the norm of the weighted sharp maximal function. The first step in our proofs for both commutators and paraproducts is the discretization of the $L^r(\nu)$ -norm of the weighted sharp maximal function.

Lemma 4.7 (Discretized norm of the weighted sharp maximal function). *Let $b \in L^1_{\text{loc}}$, $\gamma \in (0, 1)$, and \mathcal{D} a dyadic lattice. Let ν be a locally finite Borel measure and $r \in (0, \infty)$. Then*

$$\left\| \sup_{Q \in \mathcal{D}} \frac{1_Q}{\nu(Q)} \int_Q |b - \langle b \rangle_Q| dx \right\|_{L^r(\nu)} \approx \sup_{\mathcal{S} \subseteq \mathcal{D}} \left(\sum_{S \in \mathcal{S}} \left(\frac{1}{\nu(S)} \int_S |b - \langle b \rangle_S| dx \right)^r \nu(S) \right)^{1/r},$$

where the supremum is taken over all (γ, ν) -sparse collections $\mathcal{S} \subseteq \mathcal{D}$.

Proof. Since for every cube Q

$$\inf_c \int_Q |b - c| dx \approx \int_Q |b - \langle b \rangle_Q| dx,$$

the claimed conclusion is equivalent to the comparison

$$\left\| \sup_{Q \in \mathcal{D}} \frac{1_Q}{\nu(Q)} \inf_c \int_Q |b - c| dx \right\|_{L^r(\nu)} \approx \sup_{\mathcal{S} \subseteq \mathcal{D}} \left(\sum_{S \in \mathcal{S}} \left(\frac{1}{\nu(S)} \inf_c \int_Q |b - c| dx \right)^r \nu(S) \right)^{1/r}.$$

This comparison follows from a standard stopping time argument. Indeed, the classical stopping time argument of principal cubes is abstracted in [HHL16, Lemma 2.4], whose particular case $\tau_Q := \inf_c \int_Q |b - c| dx$ recovers the comparison. \square

4.4. Controlling oscillations by testing bilinear forms against pairs of test functions. Mean oscillation can be controlled by testing the bilinear form of operator against pairs of test functions. This is only step in the proof of lower bounds that relies on the concrete structure of operator.

For commutators $[b, T]$, the argument for the Beurling transform $T = S$ is classical [CRW76]. The argument for a very general class of Calderón–Zygmund singular kernels, together with a discussion on previous results, can be found in [Hyt21a] (cf. [HOS22, Proposition 4.2]):

Lemma 4.8 (Oscillations are dominated by testing the commutator against test functions; [Hyt21a]). *Let $b \in L^1_{\text{loc}}(\mathbb{R}^d, \mathbb{C})$ and let K be a non-degenerate ω -Calderón–Zygmund kernel. Let $\langle g, [b, T]f \rangle$ denote (as convenient self-explanatory abbreviation) the off-support bilinear form*

$$\langle g, [b, T]f \rangle := \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (b(y) - b(x))K(y, x)f(x)g(y) dx dy$$

for functions $f \in L^\infty_c$ and $g \in L^\infty_c$ with supports separated by a positive distance.

Then, for each cube Q , there exist a cube \tilde{Q} with $\ell(\tilde{Q}) \sim \ell(Q)$ and $\text{dist}(\tilde{Q}, Q) \sim \ell(Q)$ and functions $f^i_Q, g^i_{\tilde{Q}}$ with $|f^i_Q| \leq 1_Q$, $|g^i_{\tilde{Q}}| \leq 1_{\tilde{Q}}$ for $i = 1, 2$ such that

$$\int_Q |b - \langle b \rangle_Q| dx \lesssim \sum_{i=1}^2 \left| \langle g^i_{\tilde{Q}}, [b, T]f^i_Q \rangle \right|.$$

In the case of paraproducts, the argument and test functions are particularly simple.

Lemma 4.9 (Oscillations are dominated by testing paraproduct against indicators of cubes). *Let $b \in L^1_{\text{loc}}$ and let \mathcal{D} be a dyadic lattice. Then for every $Q \in \mathcal{D}$ we have*

$$\int_Q |b - \langle b \rangle_Q| dx \lesssim \int_{\mathbb{R}^d} |1_Q \Pi_{b, \mathcal{D}}(1_Q)| dx.$$

Proof. Let $Q \in \mathcal{D}$. Using the triangle inequality to replace $\langle b \rangle_Q$ by a term more appropriate for a paraproduct, we obtain formally

$$\begin{aligned} \int_Q |b - \langle b \rangle_Q| dx &\leq 2 \inf_c \int_Q |b - c| dx \\ &\leq 2 \int_Q |b - \langle b \rangle_Q| + \sum_{R \supseteq Q} D_R b \langle 1_Q \rangle_R dx \\ &= 2 \int_Q \left| \sum_{R \subseteq Q} D_R b \langle 1_Q \rangle_R + \sum_{R \supseteq Q} D_R b \langle 1_Q \rangle_R \right| dx \\ &= 2 \int_{\mathbb{R}^d} |1_Q \Pi_b(1_Q)| dx. \end{aligned}$$

Rigorously, we consider the truncations

$$\mathcal{D}_{M,N} := \{Q \in \mathcal{D} : 2^{-M} \leq \ell(Q) \leq 2^N\},$$

so that the term

$$\sum_{R \in \mathcal{D}_{M,N} : R \supseteq Q} D_R b \langle 1_Q \rangle_R$$

is finite. Recall that for each $f \in L_c^\infty$ the sum $\Pi_{b,\mathcal{D}} f$ converges unconditionally in L_{loc}^1 by the definition of dyadic paraproducts. Then, by a similar calculation as above together with Lemma 2.31, we obtain

$$\int_Q |b - \langle b \rangle_Q| dx \leq \lim_{N \rightarrow \infty} \lim_{M \rightarrow \infty} \int_{\mathbb{R}^d} |1_Q \Pi_{b,\mathcal{D}_{M,N}} 1_Q| dx = \int_{\mathbb{R}^d} |1_Q \Pi_b 1_Q| dx.$$

□

4.5. Sequential testing condition for general operators. The final steps in our arguments for the lower bounds for paraproducts and commutators is the necessity of a sequential testing condition for general operators. We consider the cases $p < q$ and $p \geq q$ separately.

Lemma 4.10 (Sequential testing condition on bilinear form is necessary). *Let $1 < q < p < \infty$ and set $\frac{1}{r} := \frac{1}{p} - \frac{1}{q}$. Let μ and λ be weights and $U : L^p(\mu) \rightarrow L^q(\lambda)$ a bounded linear operator. Assume the following:*

- (Weights) Set $\lambda' := \lambda^{-q'/q}$ and $\nu^{\frac{1}{p} + \frac{1}{q'}} := \mu^{\frac{1}{p}} \lambda^{-\frac{1}{q}}$. Assume that μ, λ', ν are doubling and $(\mu, \lambda') \in B_{p,q'}(\nu)$.
- (Test functions) To each cube P assign a cube Q_P with $\ell(Q_P) \lesssim \ell(P)$ and $\text{dist}(Q_P, P) \lesssim \ell(P)$ and a function f_{Q_P} with $|f_{Q_P}| \lesssim 1_{Q_P}$. Similarly, to each cube P assign a cube R_P with $\ell(R_P) \lesssim \ell(P)$ and $\text{dist}(R_P, P) \lesssim \ell(P)$ and a function g_{R_P} with $|g_{R_P}| \lesssim 1_{R_P}$.
- (Sparse collection) Assume that \mathcal{S} is a (δ, ν) -sparse collection of cubes.

Then

$$\begin{aligned} &\left(\sum_{S \in \mathcal{S}} \left| \frac{1}{\nu(S)} \int_{\mathbb{R}^d} g_{R_S} U f_{Q_S} dx \right|^r \nu(S) \right)^{1/r} \\ &\lesssim_{\mu, \lambda', \nu} [\mu, \lambda', \nu]_{B_{p,q'}(\nu)} [\mu]_{A_\infty(\nu)}^{1/p'} [\lambda']_{A_\infty(\nu)}^{1/q} \|U\|_{L^p(\mu) \rightarrow L^q(\lambda)}. \end{aligned}$$

The implicit constant depends on the doubling constants of the weights μ, λ', ν .

Remark 4.11 (Formulation in terms of bilinear form and weak-form boundedness). As in [Hyt21a, Theorem 2.5.1], Lemma 4.10 can be formulated entirely in terms

of a bilinear form $I : L_c^\infty \times L_c^\infty \rightarrow \mathbb{C}$. The boundedness then takes the following weaker form:

$$\sum_{i=1}^N |I(f_{Q_i}, g_{R_i})| \lesssim \left\| \sum_{i=1}^N \|f_{Q_i}\|_{L^\infty(Q_i)} 1_{Q_i} \right\|_{L^p(\mu)} \left\| \sum_{i=1}^N \|g_{R_i}\|_{L^\infty(R_i)} 1_{R_i} \right\|_{L^{q'}(\lambda')}$$

whenever $(f_{Q_i}, g_{R_i}) \in L^\infty(Q_i) \times L^\infty(R_i)$ for $i = 1, \dots, N$ and $N \in \mathbb{N}$. In the case of commutators, the pairs of test functions $(f_{Q_i}, g_{R_i}) \in L^\infty(Q_i) \times L^\infty(R_i)$ are such that $\text{dist}(Q_i, R_i) \sim \ell(Q_i) \sim \ell(R_i)$, as in Lemma 4.8.

Remark 4.12 (Weak-form boundedness in case $q \geq p$). When $q \geq p$, the weak-form boundedness becomes

$$|I(f_Q, g_R)| \lesssim \|f_Q\|_{L^\infty(Q)} \|g_R\|_{L^\infty(R)} \mu(Q)^{1/p} \lambda'(Q)^{1/q'}.$$

For instances of uses of such L^∞ -type testing-conditions in the context of stopping time arguments, see for example [Scu10, HH16, Hän17].

Proof of Lemma 4.10. The argument builds upon ideas from [Hyt21a] and [HOS22]. By the $\ell^r(\nu)$ - $\ell^{r'}(\nu)$ duality, the estimate of the conclusion is equivalent to the estimate

$$\sum_{S \in \mathcal{S}} \left(\frac{1}{\nu(S)} \int_{\mathbb{R}^d} g_{R_S} U f_{Q_S} dx \right) \gamma_S \nu(S) \lesssim \left(\sum_{S \in \mathcal{S}} \gamma_S^{r'} \nu(S) \right)^{1/r'}.$$

By assumption we have $1 = \frac{r'}{q'} + \frac{r'}{p}$, so

$$\gamma_S \leq [\mu, \lambda']_{B_{p, q'}(\nu)} \left(\gamma_S^{r'} \frac{\nu(S)}{\lambda^{-q'/q}(S)} \right)^{1/q'} \cdot \left(\gamma_S^{r'} \frac{\nu(S)}{\mu(S)} \right)^{1/p} =: [\mu, \lambda']_{B_{p, q'}(\nu)} \beta_S \cdot \alpha_S.$$

Therefore,

$$\sum_{S \in \mathcal{S}} \int_{\mathbb{R}^d} g_{R_S} U f_{Q_S} dx \gamma_S \lesssim_{\mu, \lambda', \nu} \int_{\mathbb{R}^d} \sum_{S \in \mathcal{S}} (\beta_S g_{R_S}) (U \alpha_S f_{Q_S}) dx.$$

Randomization is used to decouple the factors in the summands. Let $(\epsilon_S)_{S \in \mathcal{S}}$ be independent Rademacher variables on a probability space, i.e. $\mathbb{E}(\epsilon_S \epsilon_T) = \delta_{S, T}$ for all $S, T \in \mathcal{D}$ and $|\epsilon_S| = 1$. Now,

$$\begin{aligned} \int_{\mathbb{R}^d} \sum_{S \in \mathcal{S}} \beta_S g_{R_S} U \alpha_S f_{Q_S} dx &= \mathbb{E} \int_{\mathbb{R}^d} \left(\sum_{T \in \mathcal{S}} \epsilon_T \beta_T g_{R_T} \right) U \left(\sum_{S \in \mathcal{S}} \epsilon_S \alpha_S f_{Q_S} \right) dx \\ &=: \mathbb{E} \int \tilde{g}_\epsilon U \tilde{f}_\epsilon dx. \end{aligned}$$

By Hölder's inequality and boundedness of U from $L^p(\lambda)$ to $L^q(\mu)$, we have

$$\begin{aligned} \int \tilde{g}_\epsilon U \tilde{f}_\epsilon dx &= \int (\tilde{g}_\epsilon(\lambda')^{-1/q'}) U (\tilde{f}_\epsilon \lambda^{1/q}) dx \\ &\leq \|\tilde{g}_\epsilon\|_{L^{q'}(\lambda')} \|U \tilde{f}_\epsilon\|_{L^q(\lambda)} \\ &\leq \|U\|_{L^p(\mu) \rightarrow L^q(\lambda)} \|\tilde{g}_\epsilon\|_{L^{q'}(\lambda')} \|\tilde{f}_\epsilon\|_{L^p(\mu)}. \end{aligned}$$

The proof is completed by checking the estimates

$$\begin{aligned} \|\tilde{f}_\epsilon\|_{L^p(\mu)} &\leq \left(\sum_S \gamma_S^{r'} \nu(S) \right)^{1/p}, \\ \|\tilde{g}_\epsilon\|_{L^{q'}(\lambda')} &\leq \left(\sum_S \gamma_S^{r'} \nu(S) \right)^{1/q'}. \end{aligned}$$

and taking expectations.

We tackle the estimate for \tilde{f}_ϵ ; the estimate for \tilde{g}_ϵ is tackled similarly. Let us first write out what suffices to be checked. On the one hand, recalling that $\tilde{f}_\epsilon := \sum_S \epsilon_S \alpha_S f_{Q_S}$ and that by assumption $f_{Q_P} \lesssim 1_{Q_P}$, we see that

$$\|\tilde{f}_\epsilon\|_{L^p(\mu)} \lesssim \left\| \sum_{S \in \mathcal{S}} \alpha_S 1_{Q_S} \right\|_{L^p(\mu)}.$$

On the other hand, recalling that $\alpha_S := \left(\gamma_S^{r'} \frac{\nu(S)}{\mu(S)} \right)^{1/p}$, we see that

$$\sum_S \alpha_S^p \mu(S) \leq \sum_S \gamma_S^{r'} \nu(S).$$

Thus, it suffices to check that

$$\left\| \sum_{S \in \mathcal{S}} \alpha_S 1_{Q_S} \right\|_{L^p(\mu)} \lesssim_{\mu, \nu} \left(\sum_S \alpha_S^p \mu(S) \right)^{1/p}.$$

This estimate is checked as follows. By assumption $\text{dist}(Q_S, S) \lesssim \ell(S)$ and $\ell(Q_S) \lesssim \ell(S)$, so there is an $a \geq 1$ such that $aS \supseteq Q_S$. Therefore,

$$\left\| \sum_{S \in \mathcal{S}} \alpha_S 1_{Q_S} \right\|_{L^p(\mu)} \leq \left\| \sum_S \alpha_S 1_{aS} \right\|_{L^p(\mu)}.$$

Since \mathcal{S} is (δ, ν) -sparse by assumption, for each $S \in \mathcal{S}$ there is a $E_S \subseteq S$ such $\nu(E_S) \geq \delta \nu(S)$ and such that the sets $\{E_S\}_{S \in \mathcal{S}}$ are disjoint. Since ν is doubling, we have

$$\nu(E_S) \geq \delta \nu(S) \geq \delta c_{a, \nu} \nu(aS).$$

Since $a \geq 1$, we also have $E_S \subseteq S \subseteq aS$, so $\{aS\}_{S \in \mathcal{S}}$ is $(\delta c_{a, \nu}, \nu)$ -sparse.

Now, since by the preceding $\{aS\}_{S \in \mathcal{S}}$ is ν -sparse and by assumption $\mu \in A_\infty(\nu)$, it follows by Lemma 2.13 that

$$\left\| \sum_{S \in \mathcal{S}} \alpha_S 1_{aS} \right\|_{L^p(\mu)} \lesssim_{\mu, \nu} [\mu]_{A_\infty(\nu)}^{1/p'} \left(\sum_{S \in \mathcal{S}} \alpha_S^p \mu(aS) \right)^{1/p}.$$

Since μ is doubling, we have $\mu(S) \lesssim_{a, \mu} \mu(aS)$, finishing the proof. \square

We end this section with the necessity of testing condition for general operators in the case $p \geq q$, which is much simpler than the case $p < q$.

Lemma 4.13 (Testing condition in case $q \geq p$, cf. [HOS22]). *Let $1 < p \leq q < \infty$. Let μ, λ and ν be weights and set $\lambda' := \lambda^{-q'/q}$. Assume that $U : L^p(\mu) \rightarrow L^q(\lambda)$ is bounded and $(\mu, \lambda') \in B_{p, q'}(\nu)$. Then*

$$\sup_Q \frac{1}{\nu(Q)^{1/p+1/q'}} \int_{\mathbb{R}^d} 1_Q |U 1_Q| \, dx \leq [\mu, \lambda']_{B_{p, q'}(\nu)} \|U\|_{L^p(\mu) \rightarrow L^q(\lambda)}.$$

Proof. By the relation $(\lambda')^{1/q'} \lambda^{1/q} = 1$ and Hölder's inequality,

$$\begin{aligned} \int_{\mathbb{R}^d} 1_Q |U 1_Q| \, dx &= \int_{\mathbb{R}^d} ((\lambda')^{1/q'} 1_Q) (|U 1_Q| \lambda^{1/p}) \, dx \\ &\leq \|1_Q\|_{L^{q'}(\lambda')} \|U 1_Q\|_{L^q(\lambda)} \\ &\leq \|U\|_{L^p(\mu) \rightarrow L^q(\lambda)} \|1_Q\|_{L^{q'}(\lambda')} \|1_Q\|_{L^p(\mu)}. \end{aligned}$$

The claim now follows from the definition of $[\mu, \lambda']_{B_{p, q'}(\nu)}$. \square

5. MULTIPLIER CONDITION

Let $1 < q < p < \infty$ and $1/r := 1/q - 1/p$. In this section we will compare the multiplier condition

$$\inf_c \|f \mapsto (b-c)f\|_{L^p(\mu) \rightarrow L^q(\lambda)} = \inf_c \|(b-c)\nu^{-1}\|_{L^r(\nu)} < \infty,$$

and sharp maximal condition $M_\nu^\# b \in L^r(\nu)$ and prove that, unlike in the unweighted setting, the multiplier condition is not necessary for the boundedness of the commutator or for the boundedness of the paraproduct when $p > q$.

5.1. Conditions under which the multiplier and sharp maximal conditions are equivalent. In this subsection we show that under the assumption that $\nu \in A_{r'}$, the multiplier norm and the $L^r(\nu)$ -norm of the sharp maximal function are comparable. In the next subsection we will see that this is not possible in general. In particular, the comparability result of this subsection shows that in the proof of Theorem 5.3(ii) in the next subsection it will be crucial to use a weight $\nu \notin A_{r'}$.

Proposition 5.1. *Let $b \in L_{\text{loc}}^1$, $r \in (1, \infty)$ and let ν be a weight.*

(i) *If ν is doubling, we have*

$$\|M_\nu^\# b\|_{L^r(\nu)} \lesssim_\nu \inf_c \|(b-c)\nu^{-1}\|_{L^r(\nu)}.$$

(ii) *If $\nu \in A_{r'}$, we have*

$$\inf_c \|(b-c)\nu^{-1}\|_{L^r(\nu)} \lesssim [\nu]_{A_{r'}}^{r-1} [\nu]_{A_\infty} \|M_\nu^\# b\|_{L^r(\nu)}.$$

To prove Proposition 5.1, we will need the following weighted version of Lemma 3.6 in [Hyt21a]. Its proof is almost identical.

Lemma 5.2. *Let $b \in L_{\text{loc}}^1$, $1 < r < \infty$ and $\nu \in A_{r'}$. If we have for all cubes Q that*

$$\|(b - \langle b \rangle_Q)\nu^{-1}\|_{L^r(\nu|_Q)} \leq C,$$

then there is a constant c such that

$$\|(b-c)\nu^{-1}\|_{L^r(\nu)} \leq C.$$

Proof. Let us consider a sequence of cubes $Q_1 \subseteq Q_2 \subseteq \dots$ with $\bigcup_{j=1}^\infty Q_j = \mathbb{R}^d$. For $j \leq k$, we have

$$|\langle b \rangle_{Q_k} - \langle b \rangle_{Q_j}| = |(b(x) - \langle b \rangle_{Q_j}) - (b(x) - \langle b \rangle_{Q_k})|$$

and hence, multiplying by $\nu^{-1/r'}$, taking the L^r -average over Q_j and using Minkowski's inequality, we have

$$\begin{aligned} & \left(\frac{1}{|Q_j|} \int_{Q_j} |\langle b \rangle_{Q_k} - \langle b \rangle_{Q_j}|^r \nu^{1-r}(x) dx \right)^{1/r} \\ & \leq |Q_j|^{-1/r} \|(b - \langle b \rangle_{Q_j})\nu^{-1}\|_{L^r(\nu|_{Q_j})} + |Q_j|^{-1/r} \|(b - \langle b \rangle_{Q_k})\nu^{-1}\|_{L^r(\nu|_{Q_j})} \\ & \leq C|Q_j|^{-1/r} + |Q_j|^{-1/r} \|(b - \langle b \rangle_{Q_k})\nu^{-1}\|_{L^r(\nu|_{Q_k})} \\ & \leq 2C|Q_j|^{-1/r}. \end{aligned}$$

Rearranging the terms in this estimate, we obtain

$$(5.1) \quad |\langle b \rangle_{Q_k} - \langle b \rangle_{Q_j}| \leq 2C \cdot \nu^{1-r}(Q_j)^{-1/r}.$$

Note that because $\nu \in A_{r'}$, we know that $\nu^{1-r} \in A_r$. Furthermore, any A_∞ weight gives infinite measure to \mathbb{R}^d and thus $\nu^{1-r}(Q_j)^{-1/r} \rightarrow 0$ as $j \rightarrow \infty$. Thus inequality

(5.1) implies that $(\langle b \rangle_{Q_j})_{j=1}^\infty$ is a Cauchy sequence and thus converges to some c . We conclude by Fatou's lemma that

$$\begin{aligned} \int_{\mathbb{R}^d} |b - c|^r \nu^{1-r} dx &= \int_{\mathbb{R}^d} \lim_{j \rightarrow \infty} 1_{Q_j} |b - \langle b \rangle_{Q_j}|^r \nu^{1-r} dx \\ &\leq \liminf_{j \rightarrow \infty} \int_{Q_j} |b - \langle b \rangle_{Q_j}|^r \nu^{1-r} dx \leq C^r. \end{aligned}$$

This finishes the proof. \square

We are now ready to prove Proposition 5.1.

Proof of Proposition 5.1. (i): Let c be a constant. Note that for $x \in \mathbb{R}^d$ we have

$$M_\nu^\# b(x) \leq 2 \sup_{Q \ni x} \frac{1}{\nu(Q)} \int_Q (b - c) dx = 2 M^\nu((b - c)\nu^{-1})(x).$$

So the claim follows from the Hardy–Littlewood maximal inequality. The non-dyadic versions of the maximal functions are used in this context. Therefore, due to the one-third trick, the constant in the Hardy–Littlewood maximal inequality depends on the doubling constant of the measure ν .

For (ii) it suffices to estimate

$$\|(b - \langle b \rangle_Q)\nu^{-1}\|_{L^r(\nu \mathbf{1}_Q)} = \|(b - \langle b \rangle_Q)\mathbf{1}_Q\|_{L^r(\nu^{1-r})}$$

uniformly over Q by Lemma 5.2. Fix a cube Q . By [Hyt21a, Lemma 3.1.2] there exists a $\frac{1}{2}$ -sparse family \mathcal{S} such that

$$\mathbf{1}_Q |b - \langle b \rangle_Q| \lesssim \sum_{S \in \mathcal{S}} \frac{1}{|S|} \int_S |b - \langle b \rangle_S| dx.$$

Let $g \in L^{r'}(\nu)$ be positive. By Hölder's inequality, we estimate

$$\begin{aligned} &\int_Q |b(x) - \langle b \rangle_Q| g(x) dx \\ &\lesssim \sum_{S \in \mathcal{S}} \frac{1}{\nu(S)} \left(\int_S |b - \langle b \rangle_S| dx \right) \langle g \rangle_S \nu(S) \\ &\leq \left(\sum_{S \in \mathcal{S}} \left(\frac{1}{\nu(S)} \int_S |b - \langle b \rangle_S| dx \right)^r \nu(S) \right)^{1/r} \left(\sum_{S \in \mathcal{S}} \langle g \rangle_S^{r'} \nu(S) \right)^{1/r'}. \end{aligned}$$

For the first term, we estimate using Corollary 2.12

$$\begin{aligned} \left(\sum_{S \in \mathcal{S}} \left(\frac{1}{\nu(S)} \int_S |b - \langle b \rangle_S| dx \right)^r \nu(S) \right)^{1/r} &\lesssim [\nu]_{A_\infty}^{1/r} \left(\sum_{S \in \mathcal{S}} \left(\frac{1}{\nu(S)} \int_S |b - \langle b \rangle_S| dx \right)^r \nu(E_S) \right)^{1/r} \\ &= [\nu]_{A_\infty}^{1/r} \left(\sum_{S \in \mathcal{S}} \int_{E_S} \left(\frac{1}{\nu(S)} \int_S |b - \langle b \rangle_S| dx \right)^r d\nu \right)^{1/r} \\ &\leq [\nu]_{A_\infty}^{1/r} \|M_\nu^\# b\|_{L^r(\nu)}. \end{aligned}$$

For the second term, we have using Corollary 2.12 and Lemma 2.16 that

$$\begin{aligned} \left(\sum_{S \in \mathcal{S}} \langle g \rangle_S^{r'} \nu(S) \right)^{1/r'} &\lesssim [\nu]_{A_\infty}^{1/r'} \left(\sum_{S \in \mathcal{S}} \langle g \rangle_S^{r'} \nu(E_S) \right)^{1/r'} = [\nu]_{A_\infty}^{1/r'} \left(\sum_S \int_{E_S} \langle g \rangle_S^{r'} d\nu \right)^{1/r'} \\ &\leq [\nu]_{A_\infty}^{1/r'} \|Mg\|_{L^{r'}(\nu)} \\ &\lesssim [\nu]_{A_\infty}^{1/r'} [\nu]_{A_{r'}}^{\frac{1}{r'-1}} \|g\|_{L^{r'}(\nu)}. \end{aligned}$$

Combining these estimates and using duality, we get

$$\|(b - \langle b \rangle_Q)\mathbf{1}_Q\|_{L^r(\nu^{1-r})} \lesssim [\nu]_{A_{r'}}^{r-1} [\nu]_{A_\infty} \|M_\nu^\# b\|_{L^r(\nu)},$$

finishing the proof. \square

5.2. Multiplier condition is in general non-necessary for boundedness. In the previous subsection, we have seen that the pointwise multiplication condition and the sharp maximal condition are equivalent for $\nu \in A_{r'}$, which combined with Theorem A yields the following. We have

$$(5.2) \quad \|[b, T]\|_{L^p(\mu) \rightarrow L^q(\lambda)} \underset{c}{\sim}_{\mu, \lambda} \inf_c \|(b-c)\nu^{-1}\|_{L^r(\nu)}, \quad \text{assuming } \nu \in A_{r'},$$

for $1 < q < p < \infty$, $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$, $\mu \in A_p$, $\lambda \in A_q$ and $\nu^{\frac{1}{p} + \frac{1}{q'}}$:= $\mu^{1/p} \lambda^{-1/q}$, which is a direct analog of the unweighted setting in [Hyt21a]. In this subsection we will prove that the multiplier condition

$$\inf_c \|(b-c)\nu^{-1}\|_{L^r(\nu)} < \infty$$

is in general not necessary for the $L^p(\mu) \rightarrow L^q(\lambda)$ -boundedness of commutators or paraproducts. From (5.2), we know that a counterexample must satisfy $\nu \in A_{2r'} \setminus A_{r'}$.

Theorem 5.3. *Let T be an ω -Calderón-Zygmund operator with ω satisfying the Dini condition.*

(i) (Sufficiency) *Suppose $1 < p, q < \infty$, $\mu \in A_p$ and $\lambda \in A_q$. For every $b \in L^1_{\text{loc}}$,*

$$\begin{aligned} \|[b, T]\|_{L^p(\mu) \rightarrow L^q(\lambda)} &\leq (\|T\|_{L^p(\mu) \rightarrow L^p(\mu)} + \|T\|_{L^q(\lambda) \rightarrow L^q(\lambda)}) \\ &\quad \cdot \inf_c \|f \mapsto (b-c)f\|_{L^p(\mu) \rightarrow L^q(\lambda)}. \end{aligned}$$

(ii) (Non-necessity) *Suppose $1 < q < p < \infty$. Then there are $\mu \in A_p$, $\lambda \in A_q$ and $b \in L^1_{\text{loc}}$ such that*

$$\begin{aligned} \inf_c \|f \mapsto (b-c)f\|_{L^p(\mu) \rightarrow L^q(\lambda)} &= \infty, \\ \|[b, T]\|_{L^p(\mu) \rightarrow L^q(\lambda)} &< \infty. \end{aligned}$$

Proof. The proof of (i) is straightforward and well-known. Let c be a constant and denote $\Theta(c) = \|f \mapsto (b-c)f\|_{L^p(\mu) \rightarrow L^q(\lambda)}$. Then

$$\begin{aligned} \|[b, T]f\|_{L^q(\lambda)} &= \|[b-c, T]f\|_{L^q(\lambda)} \\ &\leq \|(b-c)Tf\|_{L^q(\lambda)} + \|T((b-c)f)\|_{L^q(\lambda)} \\ &\leq \Theta(c)\|Tf\|_{L^p(\mu)} + \|T\|_{L^q(\lambda) \rightarrow L^q(\lambda)}\|(b-c)f\|_{L^q(\lambda)} \\ &\leq \Theta(c)(\|Tf\|_{L^p(\mu)} + \|T\|_{L^q(\lambda) \rightarrow L^q(\lambda)}\|f\|_{L^p(\mu)}) \\ &\leq \Theta(c)(\|T\|_{L^p(\mu) \rightarrow L^p(\mu)} + \|T\|_{L^q(\lambda) \rightarrow L^q(\lambda)})\|f\|_{L^p(\mu)}. \end{aligned}$$

For (ii) the idea is to use Lemma 5.5 below. Let $b \in L^\infty$ be given by $b := \mathbf{1}_{B(0,1)}$, where $B(0,1)$ is the unit ball. Set $\frac{1}{r} := \frac{1}{q} - \frac{1}{p}$, let $\gamma := d(r' - 1)$ and define $\nu(x) := |x|^\gamma$. Then $\nu \in A_{2r'}$ since $-d < \gamma < d(2r' - 1)$. By Lemma 2.23, there exist power weights $\mu \in A_p$ and $\lambda \in A_q$ such that $\nu^{1/r'} = \mu^{1/p} \lambda^{-1/q}$. By Theorem 3.4 and Lemma 5.5(i) we deduce that

$$\|[b, T]\|_{L^p(\mu) \rightarrow L^q(\lambda)} \lesssim_{\mu, \lambda} \|M_\nu^\# b\|_{L^r(\nu)} < \infty.$$

Also, because for all constants c we have by Hölder's inequality that

$$(5.3) \quad \|f \mapsto (b-c)f\|_{L^p(\mu) \rightarrow L^q(\lambda)} = \|(b-c)\nu^{-1}\|_{L^r(\nu)},$$

Lemma 5.5(ii) shows that

$$\inf_c \|f \mapsto (b-c)f\|_{L^p(\mu) \rightarrow L^q(\lambda)} = \infty. \quad \square$$

We get a similar result to Theorem 5.3 for paraproducts instead of commutators. The sufficiency is an immediate consequence of Theorem 3.10, Proposition 5.1(i) and equality (5.3). The non-necessity part follows by using Theorem 3.10 instead of Theorem 3.4, in an otherwise identical proof to the non-necessity part of Theorem 5.3. We write this result as follows:

Theorem 5.4. *Let $1 < q < p < \infty$.*

(i) (Sufficiency) *Suppose $\mu \in A_p$ and $\lambda \in A_q$. For every $b \in L^1_{\text{loc}}$,*

$$\|\Pi_b\|_{L^p(\mu) \rightarrow L^q(\lambda)} \lesssim [\mu]_{A_p}^{\frac{1}{p-1}} [\lambda]_{A_q} [\nu]_{A_\infty}^{1/q-1/p} \cdot \inf_c \|f \mapsto (b-c)f\|_{L^p(\mu) \rightarrow L^q(\lambda)}.$$

(ii) (Non-necessity) *There are $\mu \in A_p$, $\lambda \in A_q$ and $b \in L^1_{\text{loc}}$ such that*

$$\begin{aligned} \inf_c \|f \mapsto (b-c)f\|_{L^p(\mu) \rightarrow L^q(\lambda)} &= \infty, \\ \|\Pi_b\|_{L^p(\mu) \rightarrow L^q(\lambda)} &< \infty. \end{aligned}$$

We illustrate the non-necessity using power weights. The following lemma is key in the proof of non-necessity. We use the notation $B(a, s) = \{x \in \mathbb{R}^d : |x - a| \leq s\}$.

Lemma 5.5. *Let $r \in (1, \infty)$ and set $b := \mathbf{1}_{B(0,1)}$.*

(i) *Let $\nu(x) := |x|^\gamma$ for $\gamma \geq 0$. Then*

$$\|M_\nu^\sharp b\|_{L^r(\nu)} < \infty.$$

(ii) *Let $\nu(x) := |x|^{d(r'-1)}$. Then*

$$\inf_c \|(b-c)\nu^{-1}\|_{L^r(\nu)} = \infty.$$

Before proving Lemma 5.5, and thus finalising the proof of Theorem 5.3, we document a useful property of power weights with a non-negative power:

Lemma 5.6. *Suppose $\gamma \geq 0$ and $\nu(x) := |x|^\gamma$. Then we have for all cubes Q*

$$\ell(Q)^{\gamma+d} \lesssim \nu(Q)$$

In particular, for all $\alpha > 0$ we have

$$\sup_{Q: \ell(Q) \geq \alpha} \frac{|Q|}{\nu(Q)} \lesssim \alpha^{-\gamma}.$$

Proof. Let Q be a cube. Note that, since $\gamma \geq 0$, we have $\nu(Q) \geq \nu(Q')$, where Q' is the cube with center 0 and side length $\ell(Q)$. Integrating in polar coordinates shows that $\nu(Q') \gtrsim \ell(Q)^{\gamma+d}$ proving the first estimate. The second estimate is an immediate corollary of the first. \square

Proof of Lemma 5.5. For (i) we write

$$\int_{\mathbb{R}^d} (M_\nu^\sharp b)^r \nu \, dx = \left(\int_{B(0, \frac{1}{2})} + \int_{B(0,2) \setminus B(0, \frac{1}{2})} + \int_{\mathbb{R}^d \setminus B(0,2)} \right) (M_\nu^\sharp b)^r \nu \, dx,$$

and estimate $M_\nu^\sharp b$ separately in each of the three domains.

For $|x| \leq 1/2$: Suppose $Q \ni x$. If $\ell(Q) < \frac{1}{2\sqrt{d}}$, then $Q \subseteq B(0, 1)$ and thus

$$\int_Q |b - \langle b \rangle_Q| \, dx = 0.$$

For cubes Q with $\ell(Q) \geq \frac{1}{2\sqrt{d}}$, we use the estimate

$$\int_Q |b - \langle b \rangle_Q| \, dx \leq 2|Q|,$$

which yields

$$M_\nu^\# b(x) \leq 2 \sup_{Q \ni x: \ell(Q) \geq \frac{1}{2\sqrt{d}}} \frac{|Q|}{\nu(Q)}.$$

An application of Lemma 5.6 shows that $M_\nu^\# b(x) \lesssim 1$.

For $1/2 < |x| \leq 2$: We again estimate

$$M_\nu^\# b(x) \leq 2 \sup_{Q \ni x} \frac{|Q|}{\nu(Q)}.$$

We split into two cases: $\ell(Q) \leq \frac{1}{4\sqrt{d}}$ and $\ell(Q) \geq \frac{1}{4\sqrt{d}}$. In the first case, we have $Q \subseteq \{y \in \mathbb{R}^d: |y| \geq 1/4\}$ and therefore

$$|Q| \leq 4^\gamma \nu(Q).$$

In the second case, we use Lemma 5.6 to get

$$|Q| \lesssim \ell(Q)^{\gamma+d} \lesssim \nu(Q).$$

Combining these estimates, we again obtain $M_\nu^\# b(x) \lesssim 1$.

For $|x| > 2$: Note first that for any cube Q ,

$$\int_Q |b - \langle b \rangle_Q| dx \leq 2 \int_Q |b| dx \leq 2 \int_{\mathbb{R}^d} |b| dx \lesssim 1.$$

If $Q \cap B(0, 1) = \emptyset$, we trivially have

$$\int_Q |b - \langle b \rangle_Q| dx = 0.$$

Thus, by Lemma 5.6, we get

$$M_\nu^\# b(x) \lesssim \sup_{Q \ni x, Q \cap B(0,1) \neq \emptyset} \frac{1}{\nu(Q)} \lesssim \sup_{Q \ni x, Q \cap B(0,1) \neq \emptyset} \frac{1}{\ell(Q)^{\gamma+d}}.$$

Note that for all cubes Q in the supremum we have

$$\ell(Q)\sqrt{d} \geq |x| - 1 \geq \frac{1}{2}|x|$$

and thus $\ell(Q) \gtrsim |x|$. This yields

$$M_\nu^\# b(x) \lesssim \frac{1}{|x|^{\gamma+d}}.$$

The three domains have been considered and using the achieved point-wise estimates respectively, we get

$$\int_{\mathbb{R}^d} (M_\nu^\# b)^r \nu dx \lesssim \int_{B(0,2)} |x|^\gamma dx + \int_{\mathbb{R}^d \setminus B(0,2)} |x|^{-r(\gamma+d)+\gamma} dx$$

Because $\gamma \geq 0$ and $r > 1$, we have $-r(\gamma+d)+\gamma = -\gamma(r-1) - dr < -d$. Therefore both the first and the second integral is finite.

For (ii) suppose first that $c = 0$. Then

$$\|(b-c)\nu^{-1}\|_{L^r(\nu)}^r \geq \int_{B(0,1)} |x|^{d(r'-1)(1-r)} dx = \infty,$$

because $d(r'-1)(1-r) = -d$. Next, suppose that $c \neq 0$. Then

$$\|(b-c)\nu^{-1}\|_{L^r(\nu)}^r \geq |c|^r \int_{\mathbb{R}^d \setminus B(0,1)} |x|^{d(r'-1)(1-r)} dx = \infty,$$

again because $d(r'-1)(1-r) = -d$. \square

ACKNOWLEDGEMENTS

The authors thank Tuomas Hytönen and Tuomas Oikari for helpful and inspiring discussions. All the authors are supported by the Academy of Finland (through Projects 332740 and 336323).

REFERENCES

- [ALM22] E. Airta, K. Li, and H. Martikainen. Two-weight inequalities for multilinear commutators in product spaces. *Potential Anal.*, (Online First), 2022.
- [Bar19] A. Barron. *Sparse bounds in harmonic analysis and semiperiodic estimates*. PhD thesis, Brown University, 2019.
- [Blo85] S. Bloom. A commutator theorem and weighted BMO. *Trans. Amer. Math. Soc.*, 292(1):103–122, 1985.
- [Buc93] S. Buckley. Estimates for operator norms on weighted spaces and reverse Jensen inequalities. *Trans. Amer. Math. Soc.*, 340(1):253–272, 1993.
- [Bur79] D. L. Burkholder. A sharp inequality for martingale transforms. *Ann. Probab.*, 7(5):858–863, 1979.
- [Chu11] D. Chung. Sharp estimates for the commutators of the Hilbert, Riesz transforms and the Beurling-Ahlfors operator on weighted Lebesgue spaces. *Indiana Univ. Math. J.*, 60(5):1543–1588, 2011.
- [CO17] C. Cascante and J.M. Ortega. Two-weight norm inequalities for vector-valued operators. *Mathematika*, 63(1):72–91, 2017.
- [CPP12] D. Chung, M.C. Pereyra, and C. Perez. Sharp bounds for general commutators on weighted Lebesgue spaces. *Trans. Amer. Math. Soc.*, 364(3):1163–1177, 2012.
- [CRW76] R. R. Coifman, R. Rochberg, and G. Weiss. Factorization theorems for Hardy spaces in several variables. *Ann. of Math. (2)*, 103(3):611–635, 1976.
- [FH18] S. Fackler and T.P. Hytönen. Off-diagonal sharp two-weight estimates for sparse operators. *New York J. Math.*, 24:21–42, 2018.
- [FH22] V. Fragkiadaki and I. Holmes Fay. Paraproducts, Bloom BMO and sparse BMO functions. *Rev. Mat. Iberoam.*, (Online first), 2022.
- [Hän17] T.S. Hänninen. Two-weight inequality for operator-valued positive dyadic operators by parallel stopping cubes. *Israel J. Math.*, 219(1):71–114, 2017.
- [Hän18] T.S. Hänninen. Equivalence of sparse and Carleson coefficients for general sets. *Ark. Mat.*, 56(2):333–339, 2018.
- [HH16] Timo S. Hänninen and Tuomas P. Hytönen. Operator-valued dyadic shifts and the $T(1)$ theorem. *Monatsh. Math.*, 180(2):213–253, 2016.
- [HHL16] T.S. Hänninen, T.P. Hytönen, and K. Li. Two-weight L^p - L^q bounds for positive dyadic operators: unified approach to $p \leq q$ and $p > q$. *Potential Anal.*, 45(3):579–608, 2016.
- [HLTY22] T.P. Hytönen, K. Li, J. Tao, and D. Yang. The L^p -to- L^q compactness of commutators with $p > q$. arXiv:2208.10016, 2022.
- [HLW16] I. Holmes, M.T. Lacey, and B.D. Wick. Bloom’s inequality: commutators in a two-weight setting. *Arch. Math. (Basel)*, 106(1):53–63, 2016.
- [HLW17] I. Holmes, M.T. Lacey, and B.D. Wick. Commutators in the two-weight setting. *Math. Ann.*, 367(1-2):51–80, 2017.
- [HNVW16] T.P. Hytönen, J.M.A.M. van Neerven, M.C. Veraar, and L. Weis. *Analysis in Banach spaces. Vol. I. Martingales and Littlewood-Paley theory*, volume 63 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer, Cham, 2016.
- [HOS22] T.P. Hytönen, T. Oikari, and J. Sinko. Fractional Bloom boundedness and compactness of commutators. arXiv:2207.01385, 2022.
- [HP13] T.P. Hytönen and C. Pérez. Sharp weighted bounds involving A_∞ . *Anal. PDE*, 6(4):777–818, 2013.
- [HV21] T.S. Hänninen and I.E. Verbitsky. On two-weight norm inequalities for positive dyadic operators. *Potential Anal.*, 55(2):229–249, 2021.
- [Hyt12] T.P. Hytönen. The sharp weighted bound for general Calderón-Zygmund operators. *Ann. of Math.*, 175(3):1473–1506, 2012.
- [Hyt17] T.P. Hytönen. Representation of singular integrals by dyadic operators, and the A_2 theorem. *Expo. Math.*, 35(2):166–205, 2017.
- [Hyt21a] T.P. Hytönen. The L^p -to- L^q boundedness of commutators with applications to the Jacobian operator. *J. Math. Pures Appl. (9)*, 156:351–391, 2021.
- [Hyt21b] T.P. Hytönen. Of commutators and Jacobians. In *Geometric aspects of harmonic analysis*, volume 45 of *Springer INdAM Ser.*, pages 455–466. Springer, Cham, 2021.

- [Jan78] S. Janson. Mean oscillation and commutators of singular integral operators. *Ark. Mat.*, 16:263–270, 1978.
- [Lin17] S. Lindberg. On the Hardy space theory of compensated compactness quantities. *Arch. Ration. Mech. Anal.*, 224(2):709–742, 2017.
- [Lin22] S. Lindberg. A note on the jacobian problem of coifman, lions, meyer and semmes. arXiv:2208.13576, 2022.
- [LN18] A.K. Lerner and F. Nazarov. Intuitive dyadic calculus: The basics. *Expositiones Mathematicae*, 2018.
- [LOP⁺09] A.K. Lerner, S. Ombrosi, C. Pérez, R.H. Torres, and R. Trujillo-González. New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory. *Adv. Math.*, 220(4):1222–1264, 2009.
- [LOR17] A.K. Lerner, S. Ombrosi, and I.P. Rivera-Ríos. On pointwise and weighted estimates for commutators of Calderón-Zygmund operators. *Adv. Math.*, 319:153–181, 2017.
- [LOR21] A.K. Lerner, S. Ombrosi, and I.P. Rivera-Ríos. On two weight estimates for iterated commutators. *J. Funct. Anal.*, 281(8):Paper No. 109153, 46, 2021.
- [Neh57] Z. Nehari. On bounded bilinear forms. *Ann. of Math.*, 65:153–162, 1957.
- [Oki92] K. Okikiolu. Characterization of subsets of rectifiable curves in \mathbf{R}^n . *J. London Math. Soc. (2)*, 46(2):336–348, 1992.
- [Per19] M.C. Pereyra. Dyadic harmonic analysis and weighted inequalities: the sparse revolution. In *New trends in applied harmonic analysis. Vol. 2—harmonic analysis, geometric measure theory, and applications*, Appl. Numer. Harmon. Anal., pages 159–239. Birkhäuser/Springer, Cham, 2019.
- [PP20] Jordi Pau and Antti Perälä. A Toeplitz-type operator on Hardy spaces in the unit ball. *Trans. Amer. Math. Soc.*, 373(5):3031–3062, 2020.
- [Rey22] G. Rey. Greedy approximation algorithms for sparse collections. arXiv:2202.10267, 2022.
- [Scu10] J. Scurry. A characterization of two-weight inequalities for a vector-valued operator. arXiv:1007.3089, 2010.
- [ST93] C. Segovia and J.L. Torrea. Higher order commutators for vector-valued Calderón-Zygmund operators. *Trans. Amer. Math. Soc.*, 336(2):537–556, 1993.
- [Tor86] A. Torchinsky. *Real-variable methods in harmonic analysis*, volume 123 of *Pure and Applied Mathematics*. Academic Press, Inc., Orlando, FL, 1986.
- [Ver96] I.E. Verbitsky. Imbedding and multiplier theorems for discrete Littlewood-Paley spaces. *Pacific J. Math.*, 176(2):529–556, 1996.

TIMO S. HÄNNINEN, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF HELSINKI,
P.O. Box 68, FI-00014 HELSINKI, FINLAND
Email address: timo.s.hanninen@helsinki.fi

EMIEL LORIST, DELFT INSTITUTE OF APPLIED MATHEMATICS, DELFT UNIVERSITY OF TECHNOLOGY,
P.O. BOX 5031, 2600 GA DELFT, THE NETHERLANDS
Email address: e.lorist@tudelft.nl

JAAKKO SINKO, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF HELSINKI,
P.O. Box 68, FI-00014 HELSINKI, FINLAND
Email address: jaakko.sinko@helsinki.fi