

Inference Rules for Binary Predicates in a Multigranular Framework

Stephen J. Hegner
DBMS Research of New Hampshire
PO Box, 2153, New London, NH 03257, USA
dbmsnh@gmx.com

M. Andrea Rodríguez*
Millennium Institute for Foundational Research on Data
Departamento Ingeniería Informática y Ciencias de la Computación
Edmundo Larenas 219, Universidad de Concepción
4070409 Concepción, Chile
andrea@udec.cl

Abstract

In a multigranular framework, the two most important binary predicates are those for subsumption and disjointness. In the first part of this work, a sound and complete inference system for assertions using these predicates is developed. It is customized for the granular framework; particularly, it models both bottom and top granules correctly, and it requires all granules other than the bottom to be nonempty. Furthermore, it is single use, in the sense that no assertion is used more than once as an antecedent in a proof.

In the second part of this work, a method is developed for extending a sound and complete inference system on a framework which admits Armstrong models to one which provides sound and complete inference on all assertions, both positive and negative. This method is then applied to the binary granule predicates, to obtain a sound and complete inference system for subsumption and disjointness, as well as their negations.

1 Introduction

In a multigranular relational database system such as MGDB [HR17] [HR18], the ordinary flat attributes are replaced with ones which admit a hierarchical structure. Specifically, a

*Author to whom all correspondence regarding this paper should be directed.

constrained granulated attribute schema, or *CGAS* includes a set $\text{Granules}(\mathfrak{S})$, called the *granules* of \mathfrak{S} , together with a set of constraints $\text{Constr}(\mathfrak{S})$ which relate the granules to one another. The knowledge model is *open world* in the sense that an assertion φ may be true (if $\text{Constr}(\mathfrak{S}) \models \varphi$), false (if $\text{Constr}(\mathfrak{S}) \models \neg\varphi$), or unknown (if neither $\text{Constr}(\mathfrak{S}) \models \varphi$ nor $\text{Constr}(\mathfrak{S}) \models \neg\varphi$) holds. Although it can be shown that inference is decidable, if the nature of the constraints is not restricted carefully, it becomes highly intractable. To address this issue, the constraints which are permitted are required to lie in one of two groups. The first group, which is the main topic of this paper, consists of the binary forms of subsumption ($\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_2$) (equivalent to containment, both proper and improper, in the underlying semantics) as well as of disjointness ($\sqcap\{\mathfrak{g}_1, \mathfrak{g}_2\} = \perp$), together with their negations ($\mathfrak{g}_1 \not\sqsubseteq \mathfrak{g}_2$) and ($\sqcap\{\mathfrak{g}_1, \mathfrak{g}_2\} \neq \perp$). In the context of geographic entities of Chile, simple examples of positive binary statements include the statements that the province of Llanquihue lies within the region of Los Lagos, and that the provinces of Llanquihue and Osorno are disjoint, represented in (1-1) below.

$$(\text{Llanquihue_prv} \sqsubseteq \text{Los_Lagos_rgn}) \quad (\sqcap\{\text{Llanquihue_prv}, \text{Osorno_prv}\} = \perp) \quad (1-1)$$

Similarly, simple examples of negative binary statements include that Puyehue National Park is neither contained in nor disjoint from the region of Los Lagos, represented in (1-2) below.

$$(\text{Puyehue_npr} \not\sqsubseteq \text{Los_Lagos_rgn}) \quad (\sqcap\{\text{Puyehue_npr}, \text{Los_Lagos_rgn}\} \neq \perp) \quad (1-2)$$

The second group consists of assertions which express that a single granule is either subsumed by or else equal to the join (or union) of a (finite) set of other granules. A typical example, shown in (1-3) below, asserts that the region of Los Lagos is the disjoint union of its constituent provinces.

$$\text{Los_Lagos_rgn} = \sqcup\{\text{Osorno_prv}, \text{Llanquihue_prv}, \text{Chiloé_prv}, \text{Palena_prv}\} \quad (1-3)$$

Management of complexity of inference for this type of constraint requires that the granules be grouped into *granularities*, and that such assertions be *bigranular*. For details on this, including the meaning of bigranular, the reader is referred to [HR18]. Despite their importance, such assertions will not be considered further in this paper.

The focus of this paper is the first group of constraints; more precisely, to provide a consistent and complete inference system for the two forms of positive binary constraints ($\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_2$) and ($\sqcap\{\mathfrak{g}_1, \mathfrak{g}_2\} = \perp$), together with the associated negative binary constraints ($\mathfrak{g}_1 \not\sqsubseteq \mathfrak{g}_2$) and ($\sqcap\{\mathfrak{g}_1, \mathfrak{g}_2\} \neq \perp$). The construction of this inference system is divided into two steps. In the first step, an inference system for positive binary constraints only is developed, and in the second, that system is extended to include negative binary constraints.

For the first step, the inference rules shown in Figs. 1.1 and 1.2, taken together, are shown to be *sat-conditionally complete*; that is, they are complete when applied to a consistent set of constraints. In these rules, \mathfrak{g} , \mathfrak{g}_1 , \mathfrak{g}_2 , \mathfrak{g}_3 , \mathfrak{g}'_1 , and \mathfrak{g}'_2 are variables which may take on the

$$\frac{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_2) \quad (\mathfrak{g}_2 \sqsubseteq \mathfrak{g}_3)}{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_3)} \qquad \frac{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}'_1) \quad (\prod\{\mathfrak{g}'_1, \mathfrak{g}_2\} = \perp)}{(\prod\{\mathfrak{g}_1, \mathfrak{g}_2\} = \perp)}$$

Figure 1.1: Main inference rules for positive binary constraints

$$\overline{(\mathfrak{g} \sqsubseteq \mathfrak{g})} \qquad \overline{(\mathfrak{g} \sqsubseteq \top)} \qquad \overline{(\perp \sqsubseteq \mathfrak{g})} \qquad \overline{(\prod\{\perp, \mathfrak{g}\} = \perp)}$$

Figure 1.2: Tautological inference rules for positive binary constraints

value of any granule, while \perp and \top represent the special bottom (least) and top (greatest) granules.

The rules of Fig. 1.1 are the main ones — asserting respectively that subsumption is closed under transitivity, and that if two granules are disjoint, so too are any of their subgranules under subsumption. The four rules of Fig. 1.2 represent tautologies — statements which are always true. The conclusion holds without any premises. Although these tautological rules may seem trivial, they are necessary to ensure complete sat-conditional inference.

In order to derive unsatisfiability of a set of positive assertions, the rule of Fig. 1.3 must be included as well. It requires that no granule, other than \perp , may have empty semantics, so such a granule may never be disjoint from itself.

$$\frac{(\prod\{\mathfrak{g}, \mathfrak{g}\} = \perp)}{\mathbf{false}} \langle!(\mathfrak{g} \neq \perp)\rangle$$

Figure 1.3: Unsatisfiability inference rule on positive binary constraints

This first part of the paper is based substantially upon the earlier work reported in [AP88], which provides overlapping but not identical rules in a context in which neither \perp nor \top exist as special granules, and in which any granule may have empty semantics. Since these differences are substantial, the full development of this inference system for positive assertions requires considerable development.

In the second part of this paper, it is shown that a consistent and complete set of inference rules for both positive and negative constraints may be obtained by augmenting the rules for positive constraints with *swapped* versions, in which a single antecedent is swapped with the consequent, and each negated. Shown in Fig. 1.4 are the main rules so obtained from Fig. 1.1. Similarly, Fig. 1.5 shows the swapped versions of the rules of Fig. 1.3. Informally speaking, the **false** conclusion becomes a **true** premise under a swap, which is trivial and so may be omitted. The resulting rule thus expresses a tautology. Together, the rules of Figs. 1.1–1.2 and 1.4–1.5 provide a sat-conditionally complete inference system for all binary constraints, positive and negative. (For technical reasons to be explained later, the tautological rules of Fig. 1.2 do not need to be swapped.)

$$\frac{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_2) \quad (\mathfrak{g}_1 \not\sqsubseteq \mathfrak{g}_3)}{(\mathfrak{g}_2 \not\sqsubseteq \mathfrak{g}_3)} \quad \frac{(\mathfrak{g}_2 \sqsubseteq \mathfrak{g}_3) \quad (\mathfrak{g}_1 \not\sqsubseteq \mathfrak{g}_3)}{(\mathfrak{g}_1 \not\sqsubseteq \mathfrak{g}_2)} \quad \frac{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}'_1) \quad (\prod\{\mathfrak{g}_1 \perp, \mathfrak{g}_2\} \neq \perp)}{(\mathfrak{g}'_1 \not\sqsubseteq \mathfrak{g}'_2)}$$

Figure 1.4: Swapped main inference rules for binary constraints

$$\overline{(\prod\{\mathfrak{g}, \mathfrak{g}\} \neq \perp)} \quad \langle!(\mathfrak{g} \neq \perp)\rangle$$

Figure 1.5: Tautological inference rule for negative binary constraints

In order to derive unsatisfiability for a mix of negative and positive constraints, a simple augmentation of Figs. 1.1–1.5 suffices. Namely, for every positive assertion α , a rule which states that α and $\neg\alpha$ cannot both hold is added. Since there are only two types of assertions in the binary logic of granules, the two rules of Fig. 1.6 suffice.

$$\frac{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_2) \quad (\mathfrak{g}_1 \not\sqsubseteq \mathfrak{g}_2)}{\mathbf{false}} \quad \frac{(\prod\{\mathfrak{g}_1, \mathfrak{g}_2\} = \perp) \quad (\prod\{\mathfrak{g}'_1, \mathfrak{g}'_2\} \neq \perp)}{\mathbf{false}}$$

Figure 1.6: Unsatisfiable-pair rules for binary constraints

The combined rules given in Figs. 1.1–1.6 are consistent and complete for inference on all binary constraints, positive and negative.

An important aspect of the augmentation procedure which is developed is that it is not limited to binary constraints of granule logic. Rather, it applies to any *Armstrong context* in which the positive constraints satisfy certain properties. This opens the door to the application of these results to other contexts.

This report is organized as follows. Sec. 2 provides a summary of granule spaces, which retain the essential features of CGASs which are necessary for this paper, while eliminating the unnecessary details. In Sec. 3, essential ideas for proof systems are reviewed and/or developed. In Sec. 4, the consistency and completeness of the binary inference rules of Figs. 1.1–1.3 for positive binary constraints is established. In Sec. 5, the semantics of negation in the setting in an Armstrong setting, which includes the logic of binary predicates on granules, is developed. In Sec. 6, the results of Sec. 5 are used to develop an associated proof system, establishing that the rules of Figs. 1.4–1.6 provide a sufficient augmentation of those of Figs. 1.1–1.3 to obtain a complete set of inference rules for both positive and negative granule constraints. Sec. 7 contains information the relationship of this work to other publications. Sec. 8 contains conclusions and further directions.

2 The Semantics of Granules

In the framework reported [HR17] and [HR18], a central notion is that of *granularity*, which provides a classification system for the underlying granules. For example, relative to the examples of Chilean spatial entities described in Sec. 1, three principal granularities are *Region*, *Province*, and *NatIPark*, with *Los_Lagos_rgn* a granule of granularity *Region*, *Osorno_prv*, *Llanquihue_prv*, *Chiloé_prv*, and *Palena_prv* granules of granularity *Province*, and *Puyehue_npk* a granule of granularity *NatIPark*. From the perspective of a multigranular DBMS, such classification of granules into granularities is essential not only for the purposes of organizing concepts, but also for the systematic expression and tractable inference of constraints of the form (1-3). Nevertheless, within the context of inference on binary constraints of the forms illustrated in Figs. 1.1–1.6, they are of secondary importance at best. Therefore, in this section, a simplified version of the CGAS (constrained granulated attribute schema) of [HR18] is developed, in which granularities are ignored, with only binary constraints considered. The result is termed a *simple monogranular attribute schema*, or *SMAS*. This approach has the advantage of allowing a much simpler development and presentation of the main results of this report. In addition, it allows this report to be read and understood without first gaining a thorough understanding of the concepts developed in [HR17] and [HR18].

2.1 Granule spaces A *granule space* is a triple $\mathcal{G} = (\text{Granules}\langle\mathcal{G}\rangle, \perp_{\mathcal{G}}, \top_{\mathcal{G}})$ in which $\text{Granules}\langle\mathcal{G}\rangle$ is a set, called the set of *granules*. with $\perp_{\mathcal{G}}, \top_{\mathcal{G}} \in \text{Granules}\langle\mathcal{G}\rangle$. $\perp_{\mathcal{G}}$ is called the *bottom granule* of \mathcal{G} while $\top_{\mathcal{G}}$ is called the *top granule* of \mathcal{G} ; they are always distinct. The special notation $\text{Granules}_{\perp}\langle\mathcal{G}\rangle$ is used to denote the set $\text{Granules}\langle\mathcal{G}\rangle \setminus \{\perp_{\mathcal{G}}\}$, while $\text{Granules}_{\perp\top}\langle\mathcal{G}\rangle$ is used to denote the set $\text{Granules}\langle\mathcal{G}\rangle \setminus \{\perp_{\mathcal{G}}, \top_{\mathcal{G}}\}$.

2.2 Granule structures and binary constraints A *granule structure* over \mathcal{G} is a pair $\sigma = (\text{Dom}\langle\sigma\rangle, \text{GnletoDom}_{\sigma})$ in which $\text{Dom}\langle\sigma\rangle$ is a nonempty set and $\text{GnletoDom}_{\sigma} : \text{Granules}\langle\mathcal{G}\rangle \rightarrow \mathbf{2}^{\text{Dom}\langle\sigma\rangle}$ is a function which assigns a subset of $\text{Dom}\langle\sigma\rangle$ to each granule, subject to the conditions that $\text{GnletoDom}_{\sigma}(\perp_{\mathcal{G}}) = \emptyset$, $\text{GnletoDom}_{\sigma}(\top_{\mathcal{G}}) = \text{Dom}\langle\sigma\rangle$, and for every $g \in \text{Granules}_{\perp}\langle\mathcal{G}\rangle$, $\text{GnletoDom}_{\sigma}(g) \neq \emptyset$.

The *positive binary constraints* over \mathcal{G} are of two forms. For $g_1, g_2 \in \text{Granules}\langle\mathcal{G}\rangle$, the *subsumption rule* $(g_1 \sqsubseteq g_2)$ holds in the granule structure σ iff $\text{GnletoDom}_{\sigma}(g_1) \subseteq \text{GnletoDom}_{\sigma}(g_2)$. Likewise, the *disjointness rule* $(\sqcap\{g_1, g_2\} = \perp)$ holds in σ iff $\text{GnletoDom}_{\sigma}(g_1) \cap \text{GnletoDom}_{\sigma}(g_2) = \emptyset$. The set of all such positive subsumption rules (resp. positive disjointness rules) over \mathcal{G} is denoted $\text{SubConstr}^+\langle\mathcal{G}\rangle$ (resp. $\text{DisjConstr}^+\langle\mathcal{G}\rangle$). Combining these, the set of all positive binary constraints over \mathcal{G} is

$$\text{BinConstr}^+\langle\mathcal{G}\rangle = \text{SubConstr}^+\langle\mathcal{G}\rangle \cup \text{DisjConstr}^+\langle\mathcal{G}\rangle.$$

The *negative binary constraints* over \mathcal{G} are the negations of the positive constraints. For $g_1, g_2 \in \text{Granules}\langle\mathcal{G}\rangle$, $(g_1 \not\sqsubseteq g_2)$ denotes that $(g_1 \sqsubseteq g_2)$ does not hold; *i.e.*, that $\text{GnletoDom}_{\sigma}(g_1) \not\subseteq \text{GnletoDom}_{\sigma}(g_2)$, with $\text{SubConstr}^-\langle\mathcal{G}\rangle$ denoting the set of all such negative subsumption rules. Similarly, $(\sqcap\{g_1, g_2\} \neq \perp)$ denotes that $(\sqcap\{g_1, g_2\} = \perp)$ does not hold; *i.e.*, that $\text{GnletoDom}_{\sigma}(g_1) \cap \text{GnletoDom}_{\sigma}(g_2) \neq \emptyset$. The set of all negative binary constraints over \mathcal{G} is $\text{BinConstr}^-\langle\mathcal{G}\rangle = \text{SubConstr}^-\langle\mathcal{G}\rangle \cup \text{DisjConstr}^-\langle\mathcal{G}\rangle$, with $\text{DisjConstr}^-\langle\mathcal{G}\rangle$

denoting the set of all such negative subsumption rules.

$\text{BinConstr}^\pm(\mathcal{G}) = \text{BinConstr}^+(\mathcal{G}) \cup \text{BinConstr}^-(\mathcal{G})$ is the set of *all binary constraints* over \mathcal{G} .

Following standard logical terminology, the granule structure σ is a *model* of $\varphi \in \text{BinConstr}^\pm(\mathcal{G})$ if φ holds in σ , and σ is a *model* of $\Phi \subseteq \text{BinConstr}^\pm(\mathcal{G})$ if it is a model of every $\varphi \in \Phi$. φ (resp. Φ) is *satisfiable* if it has at least one model. The set of all models of φ (resp. Φ) is denoted $\text{Mod}(\varphi)$ (resp. $\text{Mod}(\Phi)$). The constraint φ (resp. the set Φ of constraints) is *satisfiable* if $\text{Mod}(\varphi) \neq \emptyset$ (resp. $\text{Mod}(\Phi) \neq \emptyset$).

For $\varphi_1, \varphi_2 \in \text{BinConstr}^\pm(\mathcal{G})$, φ_1 *semantically entails* φ_2 , written $\varphi_1 \models \varphi_2$, just in case $\text{Mod}(\varphi_1) \subseteq \text{Mod}(\varphi_2)$. Likewise, for $\Phi \subseteq \text{BinConstr}^\pm(\mathcal{G})$ and $\varphi \in \text{BinConstr}^\pm(\mathcal{G})$ (resp. $\Phi' \subseteq \text{BinConstr}^\pm(\mathcal{G})$), write $\Phi \models \varphi$ (resp. $\Phi \models \Phi'$) just in case $\text{Mod}(\Phi) \subseteq \text{Mod}(\varphi)$. (resp. $\text{Mod}(\Phi) \subseteq \text{Mod}(\Phi')$).

2.3 Positive binary tautologies In standard logic terminology, a *tautology* is a statement which is true in every model. Define the *positive binary tautologies* of \mathcal{G} as follows.

$$\begin{aligned} \text{BinTaut}^+(\mathcal{G}) = & \{(g \sqsubseteq g) \mid g \in \text{Granules}(\mathcal{G})\} \\ & \cup \{(\perp_{\mathcal{G}} \sqsubseteq g) \mid g \in \text{Granules}(\mathcal{G})\} \\ & \cup \{(g \sqsubseteq \top_{\mathcal{G}}) \mid g \in \text{Granules}(\mathcal{G})\} \\ & \cup \{(\prod\{\perp_{\mathcal{G}}, g\} = \perp) \mid g \in \text{Granules}(\mathcal{G})\} \end{aligned}$$

That $\text{BinTaut}^+(\mathcal{G})$ consists of exactly those members of $\text{BinConstr}^+(\mathcal{G})$ which are tautologies is established next.

2.4 Proposition — Characterization of positive binary tautologies *Let \mathcal{G} be a granule space. Then $\text{BinTaut}^+(\mathcal{G})$ is precisely the subset of $\text{BinConstr}^+(\mathcal{G})$ consisting of tautologies. More precisely, if $\varphi \in \text{BinConstr}^+(\mathcal{G})$, then $\varphi \in \text{BinTaut}^+(\mathcal{G})$ iff $\sigma \in \text{Mod}(\varphi)$ for every granule structure σ over \mathcal{G} .*

PROOF: First of all, it is immediate from the definition of constraint satisfaction in 2.2 that every $\varphi \in \text{BinTaut}^+(\mathcal{G})$ is a tautology; that $\sigma \in \text{Mod}(\varphi)$ for every granule structure σ over \mathcal{G} . To show they are the only tautologies, choose any $\varphi \in \text{BinConstr}^+(\mathcal{G}) \setminus \text{BinTaut}^+(\mathcal{G})$. If $\varphi \in \text{SubConstr}^+(\mathcal{G})$, then it must be of the form $(g_1 \sqsubseteq g_2)$ with $g_1 \neq \perp_{\mathcal{G}}$, $g_2 \neq \top_{\mathcal{G}}$, and $g_1 \not\sqsubseteq g_2$. Define $\sigma' = (\text{Dom}(\sigma'), \text{GnletoDom}_{\sigma'})$ to have $\text{Dom}(\sigma') = \{x_1, x_2\}$ (any two-element set) with $\text{GnletoDom}_{\sigma'}(g_1) = \{x_1, x_2\}$ and $\text{GnletoDom}_{\sigma'}(g_2) = \{x_2\}$. Clearly, σ' is a granule structure which is not a model of φ , so the latter cannot be a tautology. On the other hand, if $\varphi \in \text{DisjConstr}^+(\mathcal{G})$, then it must be of the form $(\prod\{g_1, g_2\} = \perp)$ with $g_1 \neq \perp_{\mathcal{G}}$ and $g_2 \neq \perp_{\mathcal{G}}$. Then σ' , as defined above, is not a model of φ , so again the latter is not a tautology. \square

2.5 Unsatisfiable positive binary constraints Define the *unsatisfiable positive binary constraints* of \mathcal{G} as follows.

$$\begin{aligned} \text{BinUnsat}^+(\mathcal{G}) = & \{(g \sqsubseteq \perp_{\mathcal{G}}) \mid g \in \text{Granules}_{\neq}(\mathcal{G})\} \\ & \cup \{(\prod\{g, g\} = \perp) \mid g \in \text{Granules}_{\neq}(\mathcal{G})\} \end{aligned}$$

That $\text{BinUnsat}^+(\mathcal{G})$ consists of exactly those members of $\text{BinConstr}^+(\mathcal{G})$ which are unsatisfiable is established next.

2.6 Proposition — Characterization of unsatisfiable positive binary constraints

Let \mathcal{G} be a granule space. Then $\text{BinUnsat}^+(\mathcal{G})$ is precisely the subset of $\text{BinConstr}^+(\mathcal{G})$ consisting of unsatisfiable assertions. More precisely, if $\varphi \in \text{BinConstr}^+(\mathcal{G})$, then $\varphi \in \text{BinUnsat}^+(\mathcal{G})$ iff for no granule structure σ over \mathcal{G} is it the case that $\sigma \in \text{Mod}(\varphi)$.

PROOF: First of all, it is immediate from the definition of constraint satisfaction in 2.2 that every $\varphi \in \text{BinUnsat}^+(\mathcal{G})$ is unsatisfiable; that for no granule structure σ over \mathcal{G} is it the case that $\sigma \in \text{Mod}(\varphi)$. To show they are the only unsatisfiable members of $\text{BinConstr}^+(\mathcal{G})$, choose any $\varphi \in \text{BinConstr}^+(\mathcal{G}) \setminus \text{BinUnsat}^+(\mathcal{G})$. If $\varphi \in \text{SubConstr}^+(\mathcal{G})$, then it must be of the form $(g_1 \sqsubseteq g_2)$ with $g_2 \neq \perp_{\mathcal{G}}$. If $g_1 = \perp_{\mathcal{G}}$, then φ is a tautology, and so trivially satisfiable. Otherwise, define $\sigma' = (\text{Dom}\langle\sigma'\rangle, \text{GnletoDom}_{\sigma'})$ to have $\text{Dom}\langle\sigma'\rangle = \{x_1, x_2\}$ (any two-element set) with $\text{GnletoDom}_{\sigma'}(g_1) = \{x_1\}$. and $\text{GnletoDom}_{\sigma'}(g_2) = \{x_1, x_2\}$. Clearly, σ' is a granule structure which is a model of φ , so the latter is satisfiable. On the other hand, if $\varphi \in \text{DisjConstr}^+(\mathcal{G})$, then it must be of the form $(\bigvee\{g_1, g_2\} = \perp)$ with either $g_1 \not\stackrel{\text{id}}{=} g_2$ or else $g_1 = g_2 = \perp_{\mathcal{G}}$. If $g_1 = g_2 = \perp_{\mathcal{G}}$, then $(\bigvee\{g_1, g_2\} = \perp)$ is a tautology and there is nothing further to prove. Otherwise, define $\sigma'' = (\text{Dom}\langle\sigma''\rangle, \text{GnletoDom}_{\sigma''})$ to have $\text{Dom}\langle\sigma''\rangle = \{x_1, x_2\}$ (any two-element set) with $\text{GnletoDom}_{\sigma''}(g_1) = \{x_1\}$ and $\text{GnletoDom}_{\sigma''}(g_2) = \{x_2\}$. Clearly, σ'' is a granule structure which is a model of φ , rendering it satisfiable. \square

2.7 Simple monogranular attribute schemata A *simple monogranular attribute schema*, or *SMAS*, is a pair $\mathfrak{G} = (\text{GrSpace}\langle\mathfrak{G}\rangle, \text{Constr}(\mathfrak{G}))$ in which $\text{GrSpace}\langle\mathfrak{G}\rangle$ is a granule space and $\text{Constr}(\mathfrak{G}) \subseteq \text{BinConstr}^+(\text{GrSpace}\langle\mathfrak{G}\rangle)$. To understand the concepts of 2.2 in the context of this SMAS, it is only necessary to replace \mathcal{G} with $\text{GrSpace}\langle\mathfrak{G}\rangle$ everywhere. However, as they occur frequently, $\perp_{\mathfrak{G}}$ (resp. $\top_{\mathfrak{G}}$) will often be used as synonyms for the more cumbersome $\perp_{\text{GrSpace}\langle\mathfrak{G}\rangle}$ (resp. $\top_{\text{GrSpace}\langle\mathfrak{G}\rangle}$).

Define $\text{Constr}^+(\mathfrak{G}) = \text{Constr}(\mathfrak{G}) \cap \text{BinConstr}^+(\mathfrak{G})$. Thus, $\text{Constr}^+(\mathfrak{G})$ is the set of all positive constraints in $\text{Constr}(\mathfrak{G})$. Similarly, define $\text{Constr}^-(\mathfrak{G}) = \text{Constr}(\mathfrak{G}) \cap \text{BinConstr}^-(\mathfrak{G})$.

Define the *closure* of $\text{Constr}(\mathfrak{G})$ to be

$$\overline{\text{Constr}(\mathfrak{G})} = \{\varphi \in \text{BinConstr}^+(\mathfrak{G}) \mid \text{Constr}(\mathfrak{G}) \models \varphi\}.$$

A granular structure σ is a *model* of \mathfrak{G} precisely in the case that it is a model of $\text{Constr}(\mathfrak{G})$. Call an SMAS \mathfrak{G} *satisfiable* if $\text{Constr}(\mathfrak{G})$ has that property, and *unsatisfiable* otherwise.

2.8 Equivalence and identity of granules It is important to distinguish two distinct notions of equality between granules. Relative to \mathfrak{G} , two granules $g_1, g_2 \in \text{Granules}\langle\mathfrak{G}\rangle$ are *equivalent* if $\text{Constr}(\mathfrak{G}) \models \{(g_1 \sqsubseteq g_2), (g_2 \sqsubseteq g_1)\}$. Thus, two granules are equivalent if they have the same value under every granule structure which is a model. On the other hand, the two granules $g_1, g_2 \in \text{Granules}\langle\mathfrak{G}\rangle$ are *identical* if they are the same granule; that is, if they have the same name. Clearly, two distinct granules may be equivalent without being identical. In order to distinguish these two, the expression $(g_1 = g_2)$ will always mean that g_1 and g_2 are equivalent. To express identity, the notation $(g_1 \stackrel{\text{id}}{=} g_2)$ is always used.

2.9 Implicit symmetry of disjunction Throughout this work, for any $g_1, g_2 \in \text{Granules}\langle\mathfrak{G}\rangle$, $(\prod\{g_1, g_2\} = \perp)$ and $(\prod\{g_2, g_1\} = \perp)$ will always be regarded as the same constraint. Clearly, the semantics does not depend upon any order on g_1 and g_2 . This will eliminate the need for largely trivial inference rules which would otherwise be required to assure that $(\prod\{g_1, g_2\} = \perp)$ and $(\prod\{g_2, g_1\} = \perp)$, as well as $(\prod\{g_1, g_2\} \neq \perp)$ and $(\prod\{g_2, g_1\} \neq \perp)$, are the same.

It will also permit writing constraints of the form $(\prod S = \perp)$ and $(\prod S \neq \perp)$, in which S is a nonempty set consisting of at most two granules. If $S = \{g_1, g_2\}$, then $(\prod S = \perp)$ represents indifferently both $(\prod\{g_1, g_2\} = \perp)$ and $(\prod\{g_2, g_1\} = \perp)$, while if $S = \{g\}$, then $(\prod S = \perp)$ represents $(\prod\{g, g\} = \perp)$. The meaning of $(\prod S \neq \perp)$ is analogous.

2.10 Relationship of SMASs to CGASs This section provides a brief comparison of the SMAS, as defined in 2.7 and the CGAS, as defined in [HR18, Sec. 2] and the earlier *granularity schema*, as defined in [HR17, Sec. 3]. The reader who is not familiar with these latter concepts, or who is not interested in the comparison, may safely skip this discussion.

A CGAS $\mathfrak{G} = (\mathbf{Glt}\langle\mathfrak{G}\rangle, \text{GrAsgn}\langle\mathfrak{G}\rangle, \text{Constr}^\pm\langle\mathfrak{G}\rangle)$ consists of three components, a *granularity poset* $\mathbf{Glt}\langle\mathfrak{G}\rangle$, a *granule assignment* $\text{GrAsgn}\langle\mathfrak{G}\rangle$, and a set $\text{Constr}^\pm\langle\mathfrak{G}\rangle$ of constraints. Relative to such a CGAS, an SMAS $\mathfrak{G} = (\text{GrSpace}\langle\mathfrak{G}\rangle, \text{Constr}\langle\mathfrak{G}\rangle)$ differs in three fundamental ways.

- (a) In an SMAS, there is no granularity poset, since there is no classification of granules into granularities.
- (b) The granule space $\text{GrSpace}\langle\mathfrak{G}\rangle$ of the SMAS corresponds roughly to the granule assignment $\text{GrAsgn}\langle\mathfrak{G}\rangle$ of the CGAS. However, since there is no granularity hierarchy in an SMAS, the classification of granules into granularities, which is embedded in $\text{GrAsgn}\langle\mathfrak{G}\rangle$ in the CGAS, is absent in $\text{GrSpace}\langle\mathfrak{G}\rangle$.
- (c) In a CGAS, some binary constraints, are already embedded in $\text{GrAsgn}\langle\mathfrak{G}\rangle$, including in particular those which arise from the classification of granules into granularities. More specifically, the granule preorder $\overline{\sqsubseteq}_{\mathfrak{G}}$, as well as the constraints arising from the requirement that granules of distinct granularities be disjoint, are already embedded in the granule assignment. On the other hand, in a SMAS, no constraints are embedded in the granule space; constraints are only embedded in $\text{Constr}\langle\mathfrak{G}\rangle$.

3 Background on Proof Systems

The results of this paper are based upon simple notions of mathematical logic in general, and of propositional logic in particular. There are many excellent presentations of those subjects, at various levels and from various perspectives, including, but not limited to, [Mon76], [Men97], [End01], [BA12], and [Gal15]. It is assumed that the reader is familiar with the basic ideas presented in the appropriate chapters of those books. The purpose of this section is to establish notation and terminology for proof systems.

3.1 Notation \mathbb{N} denotes the set $\{0, 1, 2, \dots\}$ of natural numbers. Unless stated specifically to the contrary, an expression of the form $[n_1, n_2]$, with $n_1, n_2 \in \mathbb{N}$, denotes the interval $\{n \in \mathbb{N} \mid n_1 \leq n \leq n_2\}$.

$\text{Card}\langle S \rangle$ denotes the cardinality of the set S .

3.2 Context — SMAS Unless stated specifically to the contrary, for the rest of this paper, $\mathfrak{G} = (\text{GrSpace}\langle \mathfrak{G} \rangle, \text{Constr}\langle \mathfrak{G} \rangle)$ will denote an arbitrary SMAS.

3.3 WFFs for granule logic At first glance, it might seem that $\text{BinConstr}^\pm\langle \mathfrak{G} \rangle$ forms a suitable set over which to express inference rules for binary granule constraints. However, that choice is too limiting. For example, to express that the subsumption relation \sqsubseteq is transitive, it would be necessary to have a separate rule of the form

$$\frac{(g_1 \sqsubseteq g_2) \quad (g_2 \sqsubseteq g_3)}{(g_1 \sqsubseteq g_3)} \quad (3.3-1)$$

for each triple $\langle g_1, g_2, g_3 \rangle$ of granules. It would be far better to be able to assert, with a single rule, that that such a relationship holds for any triple of granules. To this end, it is appropriate to introduce a set $\{\mathfrak{g}_1, \mathfrak{g}_2, \dots, \mathfrak{g}_i, \dots\}$ of granule variables, with the understanding that each such variable, by default, may be bound to any granule. The general transitivity rule then becomes

$$\frac{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_2) \quad (\mathfrak{g}_2 \sqsubseteq \mathfrak{g}_3)}{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_3)} \quad (3.3-2)$$

with $\langle \mathfrak{g}_1, \mathfrak{g}_2, \mathfrak{g}_3 \rangle$ representing any triple of granules, as already illustrated in Fig. 1.1. In the case that variable may only be bound to certain granules, an explicit qualifier may be used, as already illustrated in the rules of Figs. 1.3 and 1.5 by the use of $|\mathfrak{g} \neq \perp$.

To formalize this, begin by associating with \mathfrak{G} a countable set $\text{VarSet}\langle \mathfrak{G} \rangle = \{\mathfrak{g}_1, \mathfrak{g}_2, \dots, \mathfrak{g}_i, \dots\}$ of *granule variables*, with $\text{VarSet}\langle \mathfrak{G} \rangle \cap \text{Granules}\langle \mathfrak{G} \rangle = \emptyset$. Relative to this set, the *positive binary well-formed formulas* (or *positive binary WFFs*) of \mathfrak{G} , denoted $\text{BinWFF}^+\langle \mathfrak{G} \rangle$, is the set consisting of all expressions of the forms $(\gamma_1 \sqsubseteq \gamma_2)$ and $(\prod\{\gamma_1, \gamma_2\} = \perp)$ with $\gamma_1, \gamma_2 \in \text{Granules}\langle \mathfrak{G} \rangle \cup \text{VarSet}\langle \mathfrak{G} \rangle$. In other words, $\text{BinWFF}^+\langle \mathfrak{G} \rangle$ is similar to $\text{BinConstr}^+\langle \mathfrak{G} \rangle$, except that both granule names and granule variables may be used in the expressions. Thus, in addition to the members of $\text{BinConstr}^+\langle \mathfrak{G} \rangle$, $\text{BinWFF}^+\langle \mathfrak{G} \rangle$ includes all expressions of the forms $(\mathfrak{g}_1 \sqsubseteq g_2)$, $(g_1 \sqsubseteq \mathfrak{g}_2)$, $(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_2)$, $(\prod\{\mathfrak{g}_1, g_2\} = \perp)$, $(\prod\{g_1, \mathfrak{g}_2\} = \perp)$, and $(\prod\{\mathfrak{g}_1, \mathfrak{g}_2\} = \perp)$, for $\mathfrak{g}_1, \mathfrak{g}_2 \in \text{VarSet}\langle \mathfrak{G} \rangle$ and $g_1, g_2 \in \text{Granules}\langle \mathfrak{G} \rangle$. Similarly, the *negative binary WFFs* of \mathfrak{G} , denoted $\text{BinWFF}^-\langle \mathfrak{G} \rangle$, is the set consisting of all expressions of the forms $(\gamma_1 \not\sqsubseteq \gamma_2)$ and $(\prod\{\gamma_1, \gamma_2\} \neq \perp)$ with $\gamma_1, \gamma_2 \in \text{Granules}\langle \mathfrak{G} \rangle \cup \text{VarSet}\langle \mathfrak{G} \rangle$. Finally, the set of *all binary WFFs* of \mathfrak{G} , denoted $\text{BinWFF}^\pm\langle \mathfrak{G} \rangle$, is $\text{BinWFF}^+\langle \mathfrak{G} \rangle \cup \text{BinWFF}^-\langle \mathfrak{G} \rangle$.

A more formal and complete development of inference rules is found in 3.9 and that which follows.

3.4 Context — Granule variables Unless stated specifically to the contrary, for the rest of this paper, $\text{VarSet}\langle \mathfrak{G} \rangle = \{\mathfrak{g}_1, \mathfrak{g}_2, \dots, \mathfrak{g}_i, \dots\}$ will denote a countable set of granule variables for the SMAS \mathfrak{G} identified in 3.2.

3.5 Semantics of WFFs To extend the notions of semantic entailment and model of 2.2 to WFFs involving variables, it is necessary to bind those variables to granules. More precisely, define a *substitution* for \mathfrak{G} to be a possibly partial function $s : \text{VarSet}\langle\mathfrak{G}\rangle \rightarrow \text{VarSet}\langle\mathfrak{G}\rangle \cup \text{Granules}\langle\mathfrak{G}\rangle$. Define $\text{Subst}\langle\varphi, s\rangle$ to be the WFF in which each variable g in the domain of s is replaced by $s(g)$. For example, if $s_0 = \{g_1 \mapsto g_3, g_2 \mapsto g_4, g_3 \mapsto g_3\}$, then $\text{Subst}\langle(g_1 \sqsubseteq g_2), s_0\rangle = (g_3 \sqsubseteq g_4)$. A *ground* substitution s is one for which $s(\text{VarSet}\langle\mathfrak{G}\rangle) \subseteq \text{Granules}\langle\mathfrak{G}\rangle$; in other words, for every g for which $s(g)$ is defined, $s(g)$ is a granule, never a variable. A substitution *complete* for $\varphi \in \text{BinWFF}^\pm\langle\mathfrak{G}\rangle$ if it is defined for every $g \in \text{VarSet}\langle\mathfrak{G}\rangle$ which occurs in φ . If s is ground and complete for φ , say that φ *holds for* s in a granule structure σ if $\text{Subst}\langle\varphi, s\rangle$ holds in σ . Thus, a WFF is assigned a truth value once all of its variables have been bound to granules. Now, for $\varphi_1, \varphi_2 \in \text{BinWFF}^\pm\langle\mathfrak{G}\rangle$, $\varphi_1 \models \varphi_2$, precisely in the case that $\text{Subst}\langle\varphi_1, s\rangle \models \text{Subst}\langle\varphi_2, s\rangle$ for every ground substitution s which is complete for both φ_1 and φ_2 .

These ideas extend easily to $\Phi \subseteq \text{BinWFF}^\pm\langle\mathfrak{G}\rangle$. Define $\text{Subst}\langle\Phi, s\rangle = \{\text{Subst}\langle\psi, s\rangle \mid \psi \in \Phi\}$, say that s is *complete* for Φ if it is complete for each $\psi \in \Phi$, and for $\varphi \in \text{BinWFF}^\pm\langle\mathfrak{G}\rangle$, define $\Phi \models \varphi$ to hold iff $\text{Subst}\langle\Phi, s\rangle \models \text{Subst}\langle\varphi, s\rangle$ for every ground substitution s which is complete for both Φ and φ . Similarly, for $\Phi' \subseteq \text{BinWFF}^\pm\langle\mathfrak{G}\rangle$, $\Phi \models \Phi'$ holds iff $\text{Subst}\langle\Phi, s\rangle \models \text{Subst}\langle\Phi', s\rangle$ for every ground substitution s which is complete for $\Phi \cup \Phi'$.

Finally, satisfiability is always defined with respect to ground substitutions. More precisely, $\varphi \in \text{BinWFF}^\pm\langle\mathfrak{G}\rangle$ (resp. $\Phi \subseteq \text{BinWFF}^\pm\langle\mathfrak{G}\rangle$) is *satisfiable* if there is a ground substitution s which is complete for φ (resp. for Φ) with $\text{Mod}(\text{Subst}\langle\varphi, s\rangle) \neq \emptyset$ (resp. $\text{Mod}(\text{Subst}\langle\Phi, s\rangle) \neq \emptyset$).

It is easy to see that these notions for $\text{BinWFF}^\pm\langle\mathfrak{G}\rangle$ generalize those of 2.2 for $\text{BinConstr}^\pm\langle\mathfrak{G}\rangle$.

3.6 General logical systems and negation Many of the concepts introduced in the remainder of this section, as well as in Secs. 5 and 6, apply in a more general setting than just $\text{BinWFF}^\pm\langle\mathfrak{G}\rangle$. In this spirit, the framework of a general logical system \mathcal{L} will be assumed as a context, as needed. This context includes notions of WFFs (denoted $\text{WFF}\langle\mathcal{L}\rangle$), structure, model, and semantic entailment (denoted \models). It is assumed that \mathcal{L} is *nontrivial* in that there is at least one $\varphi \in \text{WFF}\langle\mathcal{L}\rangle$ which is satisfiable, and that it contains at least one tautology **true** and one unsatisfiable assertion **false**. It is furthermore assumed that \mathcal{L} has a negation operator, with $\neg\varphi$ denoting the negation of $\varphi \in \text{WFF}\langle\mathcal{L}\rangle$, $\neg\neg\varphi$ equivalent to φ , and excluded middle (φ and $\neg\varphi$ cannot both be satisfiable). Further detailed properties of \mathcal{L} will not be developed here, except as necessary in for a particular issue. However, it is worthwhile noting that there is no assumption that \mathcal{L} be closed under conjunction or disjunction. In fact, the main example of \mathcal{L} in this paper, $\text{BinWFF}^\pm\langle\mathfrak{G}\rangle$, is closed under neither.

For readers not comfortable with this generalization, it suffices to fix $\text{WFF}\langle\mathcal{L}\rangle$ to be $\text{BinWFF}^\pm\langle\mathfrak{G}\rangle$ and \models to be the semantic entailment relation for that context. The assertion **true** (resp. **false**) may be taken to be (a synonym for) any element of $\text{BinTaut}^+\langle\mathfrak{G}\rangle$ (resp. $\text{BinUnsat}^+\langle\mathfrak{G}\rangle$). In $\text{BinWFF}^\pm\langle\mathfrak{G}\rangle$, to avoid unnecessary pedantry, $(g_1 \not\sqsubseteq g_2)$ (resp. $(\prod\{g_1, g_2\} \neq \perp)$), will be regarded as the same WFF as $\neg(g_1 \sqsubseteq g_2)$ (resp. $\neg(\prod\{g_1, g_2\} = \perp)$).

3.7 Proof systems Semantic entailment is a purely model-based means of relating WFFs. In order to deduce that a set of formulas implies another; that is, to determine whether $\Phi \models \varphi$ holds, a computational procedure is necessary. This is the rôle of a proof system.

More precisely, a proof system \mathcal{A} for a set $\mathfrak{A} \subseteq \text{WFF}\langle\mathcal{L}\rangle$ consists of a set of inference rules, denoted $\text{InfRules}\langle\mathcal{A}\rangle$, by which new WFFs may be obtained from others. Let $\Phi \subseteq \mathfrak{A}$ and let $\varphi \in \mathfrak{A}$. Write $\Phi \vdash_{\mathcal{A}} \varphi$ just in case φ may be obtained from Φ by repeated application of the inference rules of \mathcal{A} . Write $\Phi \vdash_{\mathcal{A}}^{(1)} \varphi$ just in case φ may be obtained from Φ by a single application of a single inference rule of \mathcal{A} .

The proof system \mathcal{A} is *sound* if whenever $\Phi \vdash_{\mathcal{A}} \varphi$, then $\Phi \models \varphi$. In words, only conclusions which are true (in the sense of \models) may be proven. Soundness is always an essential property of a proof system.

The proof system \mathcal{A} is *complete* if whenever $\Phi \models \varphi$, then $\Phi \vdash_{\mathcal{A}} \varphi$. In words, all inferences which are true may be proven using \mathcal{A} . While not essential, completeness is nevertheless a highly desirable property.

The proof system \mathcal{A} is *decidable* if there is an algorithm (which always terminates) for determining whether or not $\Phi \vdash_{\mathcal{A}} \varphi$ holds. All proof systems considered in this report are decidable, since the set \mathfrak{A} will always be finite.

3.8 Sat-conditional completeness A proof system \mathcal{A} for \mathfrak{A} is said to be *sat-complete* if for any satisfiable $\Phi \subseteq \mathfrak{A}$ and any $\varphi \in \mathfrak{A}$, $\Phi \models \varphi$ implies $\Phi \vdash_{\mathcal{A}} \varphi$.

Thus, \mathcal{A} is complete as long as the input set Φ is satisfiable, but it need not be complete in the case that Φ is not satisfiable. (Of course, if Φ is not satisfiable, then $\Phi \models \varphi$ holds trivially, so sat-completeness is not a serious limitation. It does, require, however, that an alternate means of detecting unsatisfiability be used.)

3.9 Notation for rules and proofs The representation of proof rules employed in this paper is widely used; the purpose here is to clarify terminology and notation. A generic rule is illustrated in (3.9-1) below.

$$\frac{\alpha_1 \ \alpha_2 \ \dots \ \alpha_k}{\beta} \quad (r) \tag{3.9-1}$$

The set $\{\alpha_i \mid i \in [1, k]\} \subseteq \text{WFF}\langle\mathcal{L}\rangle$ consists of the *antecedents* of the rule and is denoted $\text{Antecedents}\langle r \rangle$. Similarly, $\beta \in \text{WFF}\langle\mathcal{L}\rangle$ is the *consequent*, and is denoted $\text{Consequent}\langle r \rangle$. This rule denotes exactly that $\{\alpha_i \mid i \in [1, k]\} \vdash_{\mathcal{A}} \beta$ using a single application of the rule named r . If the rule name is clear from context, it may be omitted. If $k = 0$; *i.e.*, if $\{\alpha_i \mid i \in [1, k]\} = \emptyset$, then the rule expresses that β is a tautology. If β is **false**, the rule expresses that $\alpha_1 \wedge \alpha_2 \wedge \dots \wedge \alpha_k$ is unsatisfiable.

As already suggested in 3.3, substitutions may be applied to rules (as well as formulas). For example, (3.3-1) is the result of applying the substitution $s_1 = \{\mathfrak{g}_1 \mapsto g_1, \mathfrak{g}_2 \mapsto g_2, \mathfrak{g}_3 \mapsto g_3\}$, to (3.3-2). If the set of substitutions is restricted, this is expressed via an appropriate expression in angle brackets, as illustrated in Figs. 1.2 and 1.5.

The result of applying a substitution to a rule is called an *instance* of that rule. It will sometimes be useful to label a rule instance in a proof. For example, (3.9-2) below shows a

single instance labelled with d . In order to distinguish rule names from labels, the former will always be written in parentheses (as illustrated in (3.9-1), while labels will never be enclosed in parentheses. Note that while a rule will in general have only one name, there may be several instances of it (and hence labels for those instances) in a single proof.

$$\frac{\alpha_1 \ \alpha_2 \ \dots \ \alpha_k}{\beta} \ d \quad (3.9-2)$$

As examples, Figs. 1.1–1.6 provide the rules on granules which will be studied in this paper. Later in the paper, 4.13, 6.5, and 6.16 provide the same rules, with names.

A great advantage of the notation of 3.9 is that proofs involving the application of several rules may be presented in a clear fashion visually. For example, formulas (3.9-3) and (3.9-4) below show two ways of proving $(g_1 \sqsubseteq g_5)$ from the set

$$S_{3.9} = \{(g_1 \sqsubseteq g_2), (g_2 \sqsubseteq g_3), (g_3 \sqsubseteq g_4), (g_4 \sqsubseteq g_5)\},$$

using the rule to the left in Fig. 1.1. The various deduction instances have been labelled, with d_1 – d_3 in (3.9-3) and with d'_1 – d'_3 in (3.9-4). All are instances of the rule (3.3-2).

$$\frac{\frac{(g_1 \sqsubseteq g_2) \ (g_2 \sqsubseteq g_3)}{(g_1 \sqsubseteq g_3)} \ d_1 \quad (g_3 \sqsubseteq g_4)}{(g_1 \sqsubseteq g_4)} \ d_2 \quad (g_4 \sqsubseteq g_5)}{(g_1 \sqsubseteq g_5)} \ d_3 \quad (3.9-3)$$

$$\frac{\frac{(g_1 \sqsubseteq g_2) \ (g_2 \sqsubseteq g_3)}{(g_1 \sqsubseteq g_3)} \ d'_1 \quad \frac{(g_3 \sqsubseteq g_4) \ (g_4 \sqsubseteq g_5)}{(g_3 \sqsubseteq g_5)} \ d'_2}{(g_1 \sqsubseteq g_5)} \ d'_3 \quad (3.9-4)$$

Diagrams such as (3.9-3) and (3.9-4) are called *proof trees* (for the inference system under consideration).

The *antecedents* of a combined rule are the formulas which are at the top level. In both (3.9-3) and (3.9-4), the set of antecedents is $S_{3.9}$. Each combined rule has a single *consequent*, which is at the bottom. In both examples, it is $(g_1 \sqsubseteq g_5)$.

A formula which occurs in a proof tree, other than as an antecedent or consequent, is called an *internal formula*. In (3.9-3), the set of internal formulas is $\{(g_1 \sqsubseteq g_3), (g_1 \sqsubseteq g_4)\}$, while in (3.9-4), it is $\{(g_1 \sqsubseteq g_3), (g_3 \sqsubseteq g_5)\}$,

Finally, a formula φ which is regarded as an axiom will always be regarded as a proof of itself.

3.10 Soundness and minimality of rules and proof systems Let \mathcal{A} be a proof system for \mathfrak{A} , and let r be a rule, as depicted in (3.10-1) below.

$$\frac{\alpha_1 \ \alpha_2 \ \dots \ \alpha_k}{\beta} \ (r) \quad (3.10-1)$$

Call r *sound* if it produces consistent inference by itself; *i.e.*, if $\{\alpha_1, \alpha_2, \dots, \alpha_k\} \models \beta$.

Call a sound r *minimal* if whenever any α_i is removed from the antecedents, the resulting rule r' , as shown in (3.10-2) below, is no longer sound.

$$\frac{\alpha_1 \ \alpha_2 \ \dots \ \alpha_{i-1} \ \alpha_{i+1} \ \dots \ \alpha_k}{\beta} \quad (r') \quad (3.10-2)$$

Put another way, the rule r is minimal if for any $j \in [1, k]$, $\{\alpha_i \mid i \in [1, k] \setminus \{i\}\} \not\models \beta$.

Say that \mathcal{A} is *sound* (resp. *minimal*) if all members of $\text{InfRules}\langle\mathcal{A}\rangle$ have that property.

3.11 Single-use vs. multiple-use proofs It may be the case that an antecedent of a proof is used several times. For example, given the set

$S_{3.11} = \{(g_1 \sqsubseteq g_2), (g_2 \sqsubseteq g_3), (g_3 \sqsubseteq g_2), (g_2 \sqsubseteq g_3), (g_3 \sqsubseteq g_4)\}$, of constraints, in the proof shown in (3.11-1) below, each of $(g_1 \sqsubseteq g_2)$, $(g_1 \sqsubseteq g_3)$, and $(g_1 \sqsubseteq g_3)$ is used more than once.

$$\frac{\frac{\frac{(g_1 \sqsubseteq g_2) \ (g_2 \sqsubseteq g_3)}{(g_1 \sqsubseteq g_3)} \quad d_1 \quad (g_3 \sqsubseteq g_2)}{(g_1 \sqsubseteq g_2)} \quad d_2 \quad (g_2 \sqsubseteq g_3)}{(g_1 \sqsubseteq g_3)} \quad d_3 \quad (g_3 \sqsubseteq g_4)}{(g_1 \sqsubseteq g_4)} \quad d_4 \quad (3.11-1)$$

Call such a proof *multiple use*. On the other hand, call proofs in which no formula occurs as an antecedent more than once *single use*. The proofs of (3.9-1) and (3.9-2) are single use.

Call a proof system *single use* if for any set $\{\alpha_1, \alpha_2, \dots, \alpha_k\}$ of formulas, and any single formula β , if there is a proof of β from $\{\alpha_1, \alpha_2, \dots, \alpha_k\}$, then there is a single-use proof.

It is worth noting now that even though the proof of (3.11-1) is multiple use, a single-use proof is also possible, as shown in (3.11-2) below.

$$\frac{\frac{(g_1 \sqsubseteq g_2) \ (g_2 \sqsubseteq g_3)}{(g_1 \sqsubseteq g_3)} \quad d_1 \quad (g_3 \sqsubseteq g_4)}{(g_1 \sqsubseteq g_4)} \quad d_4 \quad (3.11-2)$$

The proof systems developed in this paper for granule rules are all single use, as will be established later.

The idea of “using up” formulas, without permitting their re-use, was introduced by Girard in the context of *linear logic* [Gir95]. Since it is not central to the results of this paper, linear logic will not be considered further.

3.12 Armstrong models Let \mathcal{C} be any set of sentences in a framework for logic. Given $\Phi \subseteq \mathcal{C}$, an *Armstrong model* M of Φ , relative to \mathcal{C} , has the property that for every $\varphi \in \mathcal{C}$, it is the case that $M \in \text{Mod}(\varphi)$ iff $\Phi \models \varphi$. In other words, an Armstrong model M is minimal in the sense that it satisfies only those constraints of \mathcal{C} which must be satisfied in order for M to be a model of Φ . Say that \mathcal{C} *admits Armstrong models*, or that it is an *Armstrong context*,

if every satisfiable subset of \mathcal{C} has such a model. See [Fag82] for a comprehensive discussion of Armstrong models, including a succinct characterization of the conditions under which they exist [Fag82, Thm. 3.1].

In the context of this paper, the important case is $\mathcal{C} = \text{BinConstr}^+(\mathfrak{G})$. While it follows easily from the more general result of [HR17, 3.18] that $\text{BinConstr}^+(\mathfrak{G})$ admits Armstrong models, a much stronger result is established in this section. Specifically, it is shown how to construct an Armstrong model, called the canonical model, for $\text{Constr}^+(\mathfrak{G})$. Since \mathfrak{G} is arbitrary, this is essentially equivalent to showing how to construct a canonical model for any $\Phi \subseteq \text{BinConstr}^+(\mathfrak{G})$. Using that model, it is shown that the positive inference rules identified in Sec. 1 are both sound and complete.

4 Inference for Positive Binary Granular Constraints

In this section, a complete inference system is developed for $\text{BinConstr}^+(\mathfrak{G})$; *i.e.*, for binary rules of the form $(g_1 \sqsubseteq g_2)$ and $(\sqcap\{g_1, g_2\} = \perp)$. It is based substantially upon [AP88], although significant differences in the two frameworks (particularly, the absence of the granules $\perp_{\mathfrak{G}}$ and $\top_{\mathfrak{G}}$ in [AP88]), as well as the need to ensure that proofs are single use (a property not considered in [AP88]), render the task substantial.

While the presentation here is designed to be independent of [AP88], referring to the latter only for proofs, readers may nevertheless be interested in consulting that work as a reference point. To that end, a brief summary of the terminology and notation of [AP88]¹ is provided below.

4.1 Summary of terminology, notation, and context of [AP88] The work of [AP88] has its roots in general knowledge bases, and so uses different terminology and notation than does this report, whose roots are in the spatial domain. In [AP88], the equivalent of an SMAS is called a *scheme*, with *types* corresponding to the granules of an SMAS, and *objects* corresponding to the elements of the domain $\text{Dom}\langle\sigma\rangle$ of a granule structure σ . A *potential state* assigns objects to types, much as the function $\text{GnletoDom}_{\sigma}$ assigns domain elements to granules. The subsumption constraint $(g_1 \sqsubseteq g_2)$ of this paper is written $g_1 \textit{ isa } g_2$, while the disjointness constraint $(\sqcap\{g_1, g_2\} = \perp)$ is written $g_1 \textit{ dis } g_2$.

From the perspective of semantics, there are two significant points of difference with the SMAS framework of this paper. First, there are no specially identified types corresponding to $\perp_{\mathfrak{G}}$ and $\top_{\mathfrak{G}}$, although it is possible to require that a type g be assigned the empty set of objects via the constraint $g \textit{ dis } g$. Such a type is termed *inconsistent*, with a scheme *inconsistent* if it contains at least one inconsistent type. Second, the set of objects associated with a type is not required to be nonempty. This is in sharp contrast to the SMAS framework, in which a granule structure assigns to each granule, other than $\perp_{\mathfrak{G}}$, a *nonempty* set of domain elements.

¹[AP86] is a preliminary conference version of the journal article [AP88]. Readers who have difficulty accessing [AP88] may find [AP86] a useful alternative.

4.2 Granule quasi-structures and quasi-models In [AP88], a more relaxed version of (the notion corresponding to) a granule structure σ than employed in an SMAS (see 2.2) is used, in which there are no restrictions on the mapping GnletoDom_σ . To keep the distinction clear, a structure in the sense of [AP88] will be termed a quasi-structure. Formally, a *granule quasi-structure* over \mathfrak{G} is a pair $\sigma = (\text{Dom}\langle\sigma\rangle, \text{GnletoDom}_\sigma)$ in which $\text{Dom}\langle\sigma\rangle$ is a nonempty set and $\text{GnletoDom}_\sigma : \text{Granules}\langle\mathfrak{G}\rangle \rightarrow \mathbf{2}^{\text{Dom}\langle\sigma\rangle}$ is a function which assigns a subset of $\text{Dom}\langle\sigma\rangle$ to each granule, subject to the further property that $\bigcup\{\text{GnletoDom}_\sigma(g) \mid g \in \text{Granules}\langle\mathfrak{G}\rangle\} = \text{Dom}\langle\sigma\rangle$. Note in particular that there is no requirement that granules other than $\perp_{\mathfrak{G}}$ map to nonempty sets, and there are no restrictions on the images of $\perp_{\mathfrak{G}}$ or $\top_{\mathfrak{G}}$. There is, however, the weaker property that every domain element occur in the image of some granule under GnletoDom_σ .

A granule quasi-structure σ is a *quasi-model* of the subsumption constraint $(g_1 \sqsubseteq g_2)$ if $\text{GnletoDom}_\sigma(g_1) \subseteq \text{GnletoDom}_\sigma(g_2)$, and it is a quasi-model of the disjointness constraint $(\prod\{g_1, g_2\} = \perp)$ if $\text{GnletoDom}_\sigma(g_1) \cap \text{GnletoDom}_\sigma(g_2) = \emptyset$. For $\Phi \subseteq \text{BinConstr}^+\langle\mathfrak{G}\rangle$, σ is a quasi-model of Φ if it is a quasi-model of each $\varphi \in \Phi$. Finally, it is a quasi-model for \mathfrak{G} if it is a quasi-model of $\text{Constr}\langle\mathfrak{G}\rangle$ (but not necessarily of $\text{BinTaut}^+\langle\mathfrak{G}\rangle$).

The *empty quasi-structure* over \mathfrak{G} , denoted $\text{Empty}\langle\mathfrak{G}\rangle$, has $\text{Dom}\langle\text{Empty}\langle\mathfrak{G}\rangle\rangle = \emptyset$ with $\text{GnletoDom}_{\text{Empty}\langle\mathfrak{G}\rangle}$ the function which maps every granule to \emptyset . This quasi-structure is interesting primarily because it is always a quasi-model, thus showing that any SMAS is satisfiable in the context of quasi-models.

To avoid confusion with quasi-models, the granule structures (resp. models) of this paper will sometimes be called *full structures* (resp. *full models*).

4.3 Inference rules on quasi-models The inference rules of [AP88, Lem. 3] are shown in Fig. 4.1. Define the inference system $\text{QBinInfPos}\langle\mathfrak{G}\rangle$ to have

$$\text{InfRules}\langle\text{QBinInfPos}\langle\mathfrak{G}\rangle\rangle = \{(I1), (I2), (D1), (M1), (M2)\}.$$

The main rules are (I2) and (M1), which assert that subsumption is transitive, and that

$$\begin{array}{ccc} (I2) \frac{(g_1 \sqsubseteq g_2) \quad (g_2 \sqsubseteq g_3)}{(g_1 \sqsubseteq g_3)} & (M1) \frac{(g_1 \sqsubseteq g'_1) \quad (\prod\{g'_1, g_2\} = \perp)}{(\prod\{g_1, g_2\} = \perp)} & \\ \\ (I1) \frac{}{(g \sqsubseteq g)} & (D1) \frac{(\prod\{g', g'\} = \perp)}{(\prod\{g', g\} = \perp)} & (M2) \frac{(\prod\{g', g'\} = \perp)}{(g' \sqsubseteq g)} \end{array}$$

Figure 4.1: Inference rules of $\text{QBinInfPos}\langle\mathfrak{G}\rangle$

disjointness is preserved under reduction of granule size via subsumption, respectively. The rule (I1) asserts the tautology that every granule subsumes itself. These three rules make sense equally in the contexts of full models and quasi-models.

The remaining two rules, (D1) and (M2), assert that if a granule is disjoint from itself, then it is also disjoint from every other granule as well, and is also subsumed by every

granule, respectively. This makes sense in the context of quasi-models, since any granule in a quasi-model may be required to be empty. However, in the context of full models, they rules apply only to the case that g' is bound to $\perp_{\mathfrak{G}}$. This is explained in more detail in 4.13 and 4.14.

4.4 Proposition — Soundness and completeness of $\text{QBinInfPos}\langle\mathfrak{G}\rangle$ *The inference rules of $\text{QBinInfPos}\langle\mathfrak{G}\rangle$, shown in Fig. 4.1 are sound and complete in the context of quasi-models. More precisely, they are complete when the concept of model used is that of quasi-model.*

PROOF: See [AP88, Thm. 1]. \square

The strategy of augmenting $\text{QBinInfPos}\langle\mathfrak{G}\rangle$ to obtain an inference system which is complete for full models is a straightforward, and is executed in two steps. In the first, additional rules which enforce all members of $\text{BinTaut}^+\langle\mathfrak{G}\rangle$ as constraints are added, and in the second, additional rules which require that the members of $\text{BinUnsat}^+\langle\mathfrak{G}\rangle$ can never be satisfied are added. The details follow.

4.5 Strong quasi-structures and quasi-models Call quasi-structure $\sigma = (\text{Dom}\langle\sigma\rangle, \text{GnletoDom}_\sigma)$ over \mathfrak{G} *strong* if $\text{GnletoDom}_\sigma(\perp_{\mathfrak{G}}) = \emptyset$ and $\text{GnletoDom}_\sigma(\top_{\mathfrak{G}}) = \text{Dom}\langle\sigma\rangle$. A quasi-model of \mathfrak{G} which is also a strong quasi-structure is called a *strong quasi-model* (of \mathfrak{G}).

Clearly, every full structure (resp. full model) is also a strong quasi-structure (resp. strong quasi-model). The only difference is that in a strong quasi-structure σ , there is no requirement that $\text{GnletoDom}_\sigma(g) \neq \emptyset$ for $g \in \text{Granules}_{\neq}\langle\mathfrak{G}\rangle$.

Also, note that $\text{Empty}\langle\mathfrak{G}\rangle$ is always a strong quasi-model, whatever be $\text{Constr}\langle\mathfrak{G}\rangle$. Thus, just as is the case for ordinary quasi-structures, any SMAS is satisfiable in the context of strong quasi-models.

4.6 Proposition — Tautologies and strong quasi-structures *If σ is a quasi-structure over \mathfrak{G} which furthermore satisfies $\text{BinTaut}^+\langle\mathfrak{G}\rangle$, then it is a strong quasi-structure over \mathfrak{G} . Consequently, if σ is a quasi-model of \mathfrak{G} which satisfies $\text{BinTaut}^+\langle\mathfrak{G}\rangle$, then it is a strong quasi-model.*

PROOF: The two differences between $\sigma = (\text{Dom}\langle\sigma\rangle, \text{GnletoDom}_\sigma)$ being an ordinary quasi-structure and a strong quasi-structure are that, in the latter, it is additionally required that (a) $\text{GnletoDom}_\sigma(\perp_{\mathfrak{G}}) = \emptyset$; and (b) $\text{GnletoDom}_\sigma(\top_{\mathfrak{G}}) = \text{Dom}\langle\sigma\rangle$. In a quasi-structure, neither of these conditions is required to hold. To prove the assertion, it suffices to show that requiring that all constraints in $\text{BinTaut}^+\langle\mathfrak{G}\rangle$ hold implies that (a) and (b) hold. This is easily verified, as follows.

For (a), letting $g \stackrel{\text{id}}{=} \perp_{\mathfrak{G}}$ in $(\prod\{\perp_{\mathfrak{G}}, g\} = \perp) \in \text{BinTaut}^+\langle\mathfrak{G}\rangle$, it follows that $\text{GnletoDom}_\sigma(\perp_{\mathfrak{G}}) \cap \text{GnletoDom}_\sigma(\perp_{\mathfrak{G}}) = \emptyset$, which can only hold if $\text{GnletoDom}_\sigma(\perp_{\mathfrak{G}}) = \emptyset$.

For (b), the tautology $(g \sqsubseteq \top_{\mathfrak{G}}) \in \text{BinTaut}^+\langle\mathfrak{G}\rangle$ guarantees that $\text{GnletoDom}_\sigma(g) \subseteq \text{GnletoDom}_\sigma(\top_{\mathfrak{G}})$ for every $g \in \text{Granules}\langle\mathfrak{G}\rangle$, which with the additional condition

$\bigcup\{\text{GnletoDom}_\sigma(g) \mid g \in \text{Granules}\langle\mathfrak{G}\rangle\} = \text{Dom}\langle\sigma\rangle$ (see 4.2) guarantees that $\text{GnletoDom}_\sigma(\top_{\mathfrak{G}}) = \text{Dom}\langle\sigma\rangle$, completing the proof. \square

4.7 Extending $\text{QBinInfPos}\langle\mathfrak{G}\rangle$ to strong quasi-models Define the inference system $\text{SQBinInfPos}\langle\mathfrak{G}\rangle$ to have

$\text{InfRules}\langle\text{SQBinInfPos}\langle\mathfrak{G}\rangle\rangle = \text{InfRules}\langle\text{QBinInfPos}\langle\mathfrak{G}\rangle\rangle \cup \{(T1), (T2), (T3)\}$,
with the rules (T1), (T2), and (T3) as defined in Fig. 4.2.

$$(T1) \frac{}{(\prod\{\perp_{\mathfrak{G}}, \mathfrak{g}\} = \perp)} \quad (T2) \frac{}{(\perp_{\mathfrak{G}} \sqsubseteq \mathfrak{g})} \quad (T3) \frac{}{(\mathfrak{g} \sqsubseteq \top_{\mathfrak{G}})}$$

Figure 4.2: Additional tautological rules $\text{SQBinInfPos}\langle\mathfrak{G}\rangle$

4.8 Proposition — Soundness and completeness of $\text{SQBinInfPos}\langle\mathfrak{G}\rangle$ in the context of strong quasi-models *The inference rules of $\text{SQBinInfPos}\langle\mathfrak{G}\rangle$ are sound and complete in the context of strong quasi-models. More precisely, they are complete when the concept of model used is that of strong quasi-model.*

PROOF: The result follows from 4.6, together with the fact that the added rules (T1), (T2), and (T3) of $\text{SQBinInfPos}\langle\mathfrak{G}\rangle$, together with (I1) of $\text{QBinInfPos}\langle\mathfrak{G}\rangle$, force all constraints in $\text{BinTaut}^+\langle\mathfrak{G}\rangle$ to hold. \square

4.9 Proposition — Strong quasi-structures and $\text{BinUnsat}^+\langle\mathfrak{G}\rangle$ *If σ is a strong quasi-structure over \mathfrak{G} which has the further property that for no $\varphi \in \text{BinUnsat}^+\langle\mathfrak{G}\rangle$ is it the case that $\text{Constr}\langle\mathfrak{G}\rangle \cup \text{BinTaut}^+\langle\mathfrak{G}\rangle \models \varphi$, then σ is a (full) structure over \mathfrak{G} .*

Consequently, if σ is a strong quasi-model of \mathfrak{G} which does not satisfy any member of $\text{BinUnsat}^+\langle\mathfrak{G}\rangle$, then it is a (full) model of \mathfrak{G} .

PROOF: The difference between $\sigma = (\text{Dom}\langle\sigma\rangle, \text{GnletoDom}_\sigma)$ being a strong quasi-structure and a full structure is that, in the latter, it is additionally required that, for $g \in \text{Granules}_\neq\langle\mathfrak{G}\rangle$, $\text{GnletoDom}_\sigma(g) \neq \emptyset$. Now, for any $g \in \text{Granules}_\neq\langle\mathfrak{G}\rangle$, it is the case that $(\prod\{g, g\} = \perp) \in \text{BinUnsat}^+\langle\mathfrak{G}\rangle$, which forbids $\text{GnletoDom}_\sigma(g) = \text{GnletoDom}_\sigma(g) \cap \text{GnletoDom}_\sigma(g) = \emptyset$. \square

4.10 Theorem — Sat-completeness of $\text{SQBinInfPos}\langle\mathfrak{G}\rangle$ *The inference rules of $\text{SQBinInfPos}\langle\mathfrak{G}\rangle$ are sound and sat-complete in the context of full models.*

PROOF: The result follows from 4.8 and 4.9, since for sat-completeness, the possibility of a structure satisfying any member of $\text{BinUnsat}^+\langle\mathfrak{G}\rangle$ is ruled out. \square

4.11 Extending $\text{SQBinInfPos}\langle\mathfrak{G}\rangle$ to models Define the inference system $\text{SQBinInfPosExt}\langle\mathfrak{G}\rangle$ to have

$\text{InfRules}\langle\text{SQBinInfPosExt}\langle\mathfrak{G}\rangle\rangle = \text{InfRules}\langle\text{SQBinInfPos}\langle\mathfrak{G}\rangle\rangle \cup \{(U1), (U2)\}$,
with the rules (U1) and (U2) as defined in Fig. 4.3.

$$(U1) \frac{(\prod\{\mathfrak{g}, \mathfrak{g}\} = \perp)}{\mathbf{false}} \ll(\mathfrak{g} \neq \perp_{\mathfrak{G}})\qquad (U2) \frac{(g \sqsubseteq \perp_{\mathfrak{G}})}{\mathbf{false}} \ll(\mathfrak{g} \neq \perp_{\mathfrak{G}})$$

Figure 4.3: Additional rules of $\text{SQBinInfPosExt}\langle\mathfrak{G}\rangle$

4.12 Proposition — Soundness and completeness of $\text{QBinInfPosExt}\langle\mathfrak{G}\rangle$ in the context of full models *The inference rules of $\text{SQBinInfPosExt}\langle\mathfrak{G}\rangle$ are sound and complete in the context of full models.*

PROOF: Soundness and sat-completeness follow from 4.10, while (U1) and (U2) ensure that unsatisfiability will be detected. \square

4.13 Unnecessary rules of $\text{BinInfPosSat}\langle\mathfrak{G}\rangle$ While the inference system $\text{QBinInfPosExt}\langle\mathfrak{G}\rangle$ is sound and complete for full models, it contains some rules which are not necessary. Specifically, the rules (D1), (M2), and (U2) may be removed. Define the inference system $\text{BinInfPos}\langle\mathfrak{G}\rangle$ to have

$$\text{InfRules}\langle\text{BinInfPos}\langle\mathfrak{G}\rangle\rangle = (\text{InfRules}\langle\text{SQBinInfPosExt}\langle\mathfrak{G}\rangle\rangle) \setminus \{(D1), (M2), (U2)\}.$$

Due to its importance, this inference system is shown in full in Fig. 4.4. That it is sound

$$(I2) \frac{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_2) \quad (\mathfrak{g}_2 \sqsubseteq \mathfrak{g}_3)}{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_3)} \qquad (M1) \frac{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}'_1) \quad (\prod\{\mathfrak{g}'_1, \mathfrak{g}_2\} = \perp)}{(\prod\{\mathfrak{g}_1, \mathfrak{g}_2\} = \perp)}$$

$$(I1) \frac{}{(\mathfrak{g} \sqsubseteq \mathfrak{g})} \qquad (T1) \frac{}{(\prod\{\perp_{\mathfrak{G}}, \mathfrak{g}\} = \perp)} \qquad (T2) \frac{}{(\perp_{\mathfrak{G}} \sqsubseteq \mathfrak{g})} \qquad (T3) \frac{}{(\mathfrak{g} \sqsubseteq \top_{\mathfrak{G}})}$$

$$(U1) \frac{(\prod\{\mathfrak{g}, \mathfrak{g}\} = \perp)}{\mathbf{false}} \ll(\mathfrak{g} \neq \perp_{\mathfrak{G}})$$

Figure 4.4: Inference rules of $\text{BinInfPos}\langle\mathfrak{G}\rangle$

and complete for full model is established in 4.14 below.

4.14 Proposition — Soundness and completeness of $\text{BinInfPos}\langle\mathfrak{G}\rangle$ *The inference rules of $\text{BinInfPos}\langle\mathfrak{G}\rangle$ are sound and complete in the context of full models.*

PROOF: First of all, the premise $(\prod\{\mathfrak{g}', \mathfrak{g}'\} = \perp)$ in rules (D1) and (M2) can only hold in the case that \mathfrak{g}' is bound to $\perp_{\mathfrak{G}}$, in view of rule (U1). Since $(\prod\{\perp_{\mathfrak{G}}, \perp_{\mathfrak{G}}\} = \perp)$ is a tautology by (T1), the premise for \mathfrak{g} bound to $\perp_{\mathfrak{G}}$ becomes **true** in both (D1) and (M2). The resulting rules are precisely (T1) and (T2), respectively, rendering the original (D1) and (M2) unnecessary.

To justify removing (U2), it suffices that to note that the unsatisfiability of $(g \sqsubseteq \perp_{\mathfrak{G}})$ may be deduced from the unsatisfiability of $(\prod\{g, g\} = \perp)$, using additionally (T2) and (M1), as shown in Fig. 4.5. \square

$$\frac{\frac{(g \sqsubseteq \perp_{\mathfrak{G}}) \quad \overline{(\prod\{\perp_{\mathfrak{G}}, g\} = \perp)}}{(\prod\{g, g\} = \perp)} \quad \langle\langle g \neq \perp_{\mathfrak{G}} \rangle\rangle}{\text{false}}$$

Figure 4.5: Deducing the unsatisfiability of $(g \sqsubseteq \perp_{\mathfrak{G}})$ from that of $(\prod\{g, g\} = \perp)$

4.15 Armstrong context strategy The approach to extending inference systems such as $\text{BinInfPos}\langle\mathfrak{G}\rangle$ to include negative constraints as well requires that the context (in this case $\text{BinConstr}^+\langle\mathfrak{G}\rangle$) be Armstrong (see 3.12). A straightforward way to show that a context is Armstrong is to use the characterization given in [Fag82, Thm. 3.1]. The proof has already been developed in [HR17, 3.15-3.18] for a much more general class of granular constraints than $\text{BinConstr}^+\langle\mathfrak{G}\rangle$, and so will not be repeated here. The reader is referred to that article for details. For completeness, the formal result is recorded below.

4.16 Theorem — Armstrong context $\text{BinConstr}^+\langle\mathfrak{G}\rangle$ is an Armstrong context. More precisely, if \mathfrak{G} is satisfiable, then $\text{Constr}^+(\mathfrak{G}) \cup \text{BinTaut}^+\langle\mathfrak{G}\rangle$ admits an Armstrong model.

PROOF: See [HR17, 3.18]. \square

Attention is now turned to the task of showing that proofs in $\text{BinInfPos}\langle\mathfrak{G}\rangle$ are single use. For this task, it is useful to begin by characterizing properties of \mathfrak{G} via an associated graph.

4.17 The graph of an SMAS The *graph* of \mathfrak{G} , denoted $\text{Graph}\langle\mathfrak{G}\rangle$, has as vertices the elements of $\text{Granules}\langle\mathfrak{G}\rangle$. It has two types of edges, one directed and one undirected, defined as follows.

(s-edge) For $g_1, g_2 \in \text{Granules}\langle\mathfrak{G}\rangle$, there is a directed *subsumption edge* from g_1 to g_2 just in case $(g_1 \sqsubseteq g_2) \in \text{Constr}^+(\mathfrak{G}) \cup \text{BinTaut}^+\langle\mathfrak{G}\rangle$. Formally, this edge is regarded as the ordered pair $\langle g_1, g_2 \rangle$. The set of all subsumption edges of $\text{Graph}\langle\mathfrak{G}\rangle$ is denoted $\text{SubEdges}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$.

(d-edge) For $g_1, g_2 \in \text{Granules}\langle\mathfrak{G}\rangle$, there is an undirected *disjointness edge* between g_1 and g_2 just in case $(\prod\{g_1, g_2\} = \perp) \in \text{Constr}^+(\mathfrak{G}) \cup \text{BinTaut}^+\langle\mathfrak{G}\rangle$. Formally, if $g_1 \stackrel{\text{id}}{\neq} g_2$, this edge is regarded as the unordered pair $\{g_1, g_2\}$. In the case that $g_1 \stackrel{\text{id}}{=} g_2$, it is regarded as the singleton $\{g_1\} = \{g_2\} = \{g_1, g_2\}$. The set of all disjointness edges of $\text{Graph}\langle\mathfrak{G}\rangle$ is denoted $\text{DisjEdges}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$.

The set of *all edges* of $\text{Graph}\langle\mathfrak{G}\rangle$ is

$$\text{Edges}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle = \text{SubEdges}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle \cup \text{DisjEdges}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle.$$

In the equivalent of this graph in [AP88], subsumption edges are called *black edges* and disjointness edges are called *red edges*.

A *subsumption path* in $\text{Graph}\langle\mathfrak{G}\rangle$ is a sequence $p = \langle g_1, g_2, \dots, g_k \rangle$, with $k \geq 2$, of elements of $\text{Granules}\langle\mathfrak{G}\rangle$ with the property that for each $i \in [1, k-1]$, $\langle g_i, g_{i+1} \rangle \in \text{SubEdges}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$. Such a path p is said to be *from* g_1 *to* g_k , and to *underlie* $\langle g_1, g_k \rangle$. The set of all subsumption paths of $\text{Graph}\langle\mathfrak{G}\rangle$ is denoted $\text{SubPaths}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$. The length k of p is denoted $\text{Length}\langle p \rangle$.

The *underlying constraint sequence* of p is $\text{ConstrSeq}\langle p \rangle = \langle (g'_1 \sqsubseteq g'_2), (g'_2 \sqsubseteq g'_3), \dots, (g'_i \sqsubseteq g'_{i+1}), \dots, (g'_{k-1} \sqsubseteq g'_k) \rangle$, with $(g'_i \sqsubseteq g'_{i+1})$ the i^{th} -constraint of p . The *underlying constraint set* of p is the underlying constraint sequence with the order removed; more precisely, $\text{ConstrSet}\langle p \rangle = \{(g'_1 \sqsubseteq g'_2), (g'_2 \sqsubseteq g'_3), \dots, (g'_i \sqsubseteq g'_{i+1}), \dots, (g'_{k-1} \sqsubseteq g'_k)\}$,

The relation $\text{Sub}^*\text{Edges}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$ on $\text{Granules}\langle\mathfrak{G}\rangle$ is defined by $\langle g_1, g_2 \rangle \in \text{Sub}^*\text{Edges}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$ iff there is a $\langle g'_1, g'_2, \dots, g'_k \rangle \in \text{SubPaths}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$ with $g'_1 \stackrel{\text{id}}{=} g_1$ and $g'_k \stackrel{\text{id}}{=} g_2$. In this case, the path $p = \langle g'_1, g'_2, \dots, g'_k \rangle$ is said to *underlie* $\langle g_1, g_2 \rangle$.

4.18 Proposition — Existence of underlying paths *Let $g_1, g_2 \in \text{Granules}\langle\mathfrak{G}\rangle$, and assume further that \mathfrak{G} is satisfiable. Then the entailment $\text{Constr}^+\langle\mathfrak{G}\rangle \cup \text{BinTaut}^+\langle\mathfrak{G}\rangle \models (g_1 \sqsubseteq g_2)$ holds iff there is a subsumption path p which underlies $\langle g_1, g_2 \rangle$.*

PROOF: That the result then holds in the context of quasi-models is shown in [AP88, Thm. 2]. Upon adding the tautologies and taking into account the results of 4.6, it is easy to see that it holds in the context of strong quasi-models as well. Finally, with the requirement that \mathfrak{G} be satisfiable, in view of 4.9, the result applies to the context of full models also. \square

Since a subsumption constraint holds in \mathfrak{G} iff it has an underlying path, it also has a simple proof based upon that path. The details follow.

4.19 Linear proofs for subsumption constraints Let $p' = \langle g'_1, g'_2, \dots, g'_k \rangle \in \text{SubPaths}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$. The *left-linear proof tree* (in $\text{BinInfPos}\langle\mathfrak{G}\rangle$) associated with p , denoted $\text{LLinear}\langle p \rangle$, is defined as follows.

- (a) If $k = 2$ and $(g'_1 \sqsubseteq g'_2) \in \text{Constr}\langle\mathfrak{G}\rangle$, then $\text{LLinear}\langle p \rangle$ is simply $(g'_1 \sqsubseteq g'_2)$. In this case, the proof uses no rules at all, since the desired consequence is in the set of antecedents.
- (b) If $k = 2$ and $(g'_1 \sqsubseteq g'_2) \in \text{BinTaut}^+\langle\mathfrak{G}\rangle \setminus \text{Constr}\langle\mathfrak{G}\rangle$, then $\text{LLinear}\langle p \rangle$ is the tree $\overline{(g'_1 \sqsubseteq g'_2)}$. In this case, the proof uses at most the rules in $\{(I1), (T1), (T2), (T3)\}$.
- (c) If $k > 2$ and for every $i \in [1, k-1]$ it is the case that $(g'_i \sqsubseteq g'_{i+1}) \in \text{Constr}\langle\mathfrak{G}\rangle$, then $\text{LLinear}\langle p \rangle$ is defined as shown in Fig. 4.6. In this case, only rule (I2) is used.
- (d) If $k > 2$ and for one or more $i \in [1, k-1]$ it is the case that $(g'_i \sqsubseteq g'_{i+1}) \in \text{BinTaut}^+\langle\mathfrak{G}\rangle \setminus \text{Constr}\langle\mathfrak{G}\rangle$, then $\text{LLinear}\langle p \rangle$ is as shown in Fig. 4.6, but with $(g'_i \sqsubseteq g'_{i+1})$ replaced by the tree $\overline{(g'_i \sqsubseteq g'_{i+1})}$. In this case, at most the rules in $\{(I1), (I1), (T1), (T2), (T3)\}$ are used.

4.20 Lemma — Existence of left-linear proofs *Let $g_1, g_2 \in \text{Granules}\langle\mathfrak{G}\rangle$. If \mathfrak{G} is satisfiable and $\text{BinConstr}^+\langle\mathfrak{G}\rangle \cup \text{BinTaut}^+\langle\mathfrak{G}\rangle \models (g_1 \sqsubseteq g_2)$, then there is a left-linear proof of*

$$\begin{array}{c}
\frac{(g'_1 \sqsubseteq g'_2) \quad (g'_2 \sqsubseteq g'_3)}{(g'_1 \sqsubseteq g'_3)} \quad (g'_3 \sqsubseteq g'_4) \\
\hline
(g'_1 \sqsubseteq g'_4) \\
\vdots \\
(g'_1 \sqsubseteq g'_i) \quad (g'_i \sqsubseteq g'_{i+1}) \\
\hline
(g'_1 \sqsubseteq g'_{i+1}) \\
\vdots \\
(g'_1 \sqsubseteq g'_{k-1}) \quad (g'_k \sqsubseteq g'_{k-1}) \\
\hline
(g'_1 \sqsubseteq g'_k)
\end{array}$$

Figure 4.6: A left-linear proof tree for subsumption

this in $\text{BinInfPos}\langle\mathfrak{G}\rangle$.

PROOF: Use 4.18 to establish the existence of an underlying subsumption path, and then 4.19 to show that a proof of the given form is possible. \square

4.21 Reduced subsumption paths The subsumption path $p = \{g'_1, g'_2, \dots, g'_k\} \in \text{SubPaths}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$ is *reduced* if either (a) $\text{Length}\langle p \rangle = 2$, or else (b) no granule in $\text{Granules}_\neq\langle\mathfrak{G}\rangle$ occurs more than once in p , while $\perp_{\mathfrak{G}}$ does not occur at all.

4.22 Lemma — Existence of reduced subsumption paths Let $\langle g_1, g_2 \rangle \in \text{Granules}\langle\mathfrak{G}\rangle \times \text{Granules}\langle\mathfrak{G}\rangle$. If there is a subsumption path in $\text{Graph}\langle\mathfrak{G}\rangle$ which underlies $\langle g_1, g_2 \rangle$, then there is such a path which is reduced.

PROOF: Let $p = \langle g'_1, g'_2, \dots, g'_k \rangle$ be a subsumption path which underlies $\langle g_1, g_2 \rangle$. If $\text{Length}\langle p \rangle = 2$, then there is nothing further to prove, since all such paths are reduced. So, assume that $\text{Length}\langle p \rangle > 2$ and that some $g \in \text{Granules}\langle\mathfrak{G}\rangle$ occurs more than once in p . If these multiple occurrences are the endpoints of p ; that is, if $\stackrel{\text{id}}{=} g'_1 \stackrel{\text{id}}{=} g'_k$, then $\langle g_1, g_2 \rangle$ is of the form $\langle g, g \rangle$ for some $g \in \text{Granules}\langle\mathfrak{G}\rangle$. In that case, replace p with $\langle g, g \rangle$, which also underlies $\langle g_1, g_2 \rangle$ and is furthermore reduced since it is of length two.

Otherwise, writing $p = \langle g'_1, g'_2, \dots, g'_k \rangle$, suppose that for $j_1, j_2 \in [1, k]$ with $j_1 < j_2$ and either $j_1 > 1$ or $j_2 < k$, it is the case that $g'_{j_1} \stackrel{\text{id}}{=} g'_{j_2}$. Remove the elements in the subsequence $\langle g'_{j_1+1}, \dots, g'_{j_2} \rangle$ from p , with the resulting sequence being $p' = \langle g'_1, g'_2, \dots, g'_{j_1}, g'_{j_2+1}, \dots, g'_k \rangle$. It is easy to see that $p' \in \text{SubPaths}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$, and that it also underlies $\langle g_1, g_2 \rangle$, but with $\text{Length}\langle p' \rangle < \text{Length}\langle p \rangle$. If another inessential cycle remains in p' , this process may be repeated. Since each iteration reduces the length of the path, the process must terminate eventually, resulting in a path which is free of internal cycles.

The only satisfiable subsumption constraints which involve $\perp_{\mathfrak{G}}$, are of the form $(\perp_{\mathfrak{G}} \sqsubseteq g)$ for some $g \in \text{Granules}\langle\mathfrak{G}\rangle$, and these are already in $\text{BinTaut}^+\langle\mathfrak{G}\rangle$. Thus, there is no need to use $\perp_{\mathfrak{G}}$ in any path of length greater than two. \square

4.23 Lemma — Single-use left-linear proofs *If p is a reduced subsumption path in $\text{Graph}\langle\mathfrak{G}\rangle$, then the proof $\text{LLinear}\langle p\rangle$ (see 4.19) is single use.*

PROOF: This is immediate, since in any reduced path of length greater than two, no granule can occur more than once. \square

4.24 Proposition — Deducing satisfiability of subsumption constraints *If $\text{Constr}^+(\mathfrak{G})$ is satisfiable, then for any $g_1, g_2 \in \text{Granules}\langle\mathfrak{G}\rangle$ with $\text{Constr}^+(\mathfrak{G}) \models (g_1 \sqsubseteq g_2)$, there is a single-use left-linear proof in $\text{BinInfPos}\langle\mathfrak{G}\rangle$ of $(g_1 \sqsubseteq g_2)$, using only subsumption constraints in $\text{Constr}^+(\mathfrak{G})$ as premises, and using at most the rules which lie in $\{(I2), (I1), (T1), (T2), (T3)\}$. If $g_1, g_2 \in \text{Granules}_{\neq} \langle\mathfrak{G}\rangle$ with $g_1 \stackrel{\text{id}}{\neq} g_2$, then at most rules (I2) and (T3) need be used.*

PROOF: It suffices to combine 4.20 and 4.23. \square

It is also the case that proofs of disjointness have a special form, but in the general case, based upon two subsumption paths rather than one. The details follow.

4.25 Protected pairs Define the set of *doublets* of \mathfrak{G} to be

$$\text{Doublets}\langle\mathfrak{G}\rangle = \{\{g_1, g_2\} \mid g_1, g_2 \in \text{Granules}\langle\mathfrak{G}\rangle \text{ and } g_1 \stackrel{\text{id}}{\neq} g_2\}.$$

Say that a pair $S = \{g_1, g_2\} \in \text{Doublets}\langle\mathfrak{G}\rangle$ is *protected for \mathfrak{G}* if there is a pair $S' = \{g'_1, g'_2\} \in \text{Granules}\langle\mathfrak{G}\rangle$ for which $\langle g_1, g'_1 \rangle, \langle g_2, g'_2 \rangle \in \text{Sub}^*\text{Edges}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$ and $S' \in \text{DisjEdges}\langle\text{Graph}\langle\mathfrak{G}\rangle\rangle$. See Fig. 4.7. Thus, S is protected if its two granules are (not




Figure 4.7: Visualization of the protected pair $\{g_1, g_2\}$

necessarily proper) subgranules of corresponding ones in a pair S' which is declared explicitly to be disjoint, either as a tautology or else as a constraint of \mathfrak{G} . In this case, S' is called a *protector* of S . A pair $S \in \text{Doublets}\langle\mathfrak{G}\rangle$ is *unprotected (for \mathfrak{G})* if it is not protected. The set of all protected pairs for \mathfrak{G} is denoted $\text{ProtectedPr}\langle\mathfrak{G}\rangle$, while $\text{UnProtectedPr}\langle\mathfrak{G}\rangle = \text{Doublets}\langle\mathfrak{G}\rangle \setminus \text{ProtectedPr}\langle\mathfrak{G}\rangle$ is the set of unprotected pairs for \mathfrak{G} .

4.26 Lemma — Protection implies disjointness *For any $S \in \text{Doublets}\langle\mathfrak{G}\rangle$, $S \in \text{ProtectedPr}\langle\mathfrak{G}\rangle$ iff $\text{Constr}^+(\mathfrak{G}) \models (\sqcap S = \perp)$.*

PROOF: See [AP88, Thm. 3]. \square

4.27 One- and two-path left-linear proofs for disjointness constraints Suppose that $(\prod\{g_1, g_2\} = \perp)$ is the constraint to prove. There are four cases to consider. First of all, if $(\prod\{g_1, g_2\} = \perp) \in \text{Constr}^+(\mathfrak{G})$, then the constraint itself forms its proof tree. Second, if

$(\prod\{g_1, g_2\} = \perp)$ is a tautology; that is, if $g_1 \stackrel{\text{id}}{=} \perp_{\mathfrak{G}}$, then the proof tree is $\overline{(\prod\{\perp_{\mathfrak{G}}, g_2\} = \perp)}$ (and analogously if $g_2 \stackrel{\text{id}}{=} \perp_{\mathfrak{G}}$). Third, neither of these hold, but for some $g'_1 \in \text{Granules}\langle\mathfrak{G}\rangle$, $(\prod\{g'_1, g_2\} = \perp) \in \text{Constr}^+(\mathfrak{G})$, then the proof is of the form shown in Fig. 4.8. An analogous result holds $(\prod\{g_1, g'_2\} = \perp) \in \text{Constr}^+(\mathfrak{G})$ for some $g'_2 \in \text{Granules}\langle\mathfrak{G}\rangle$. Finally, if none of

$$\frac{\begin{array}{c} \Sigma_1 \\ \vdots \\ \text{left-linear proof} \\ \vdots \\ (g_1 \sqsubseteq g'_1) \end{array} \quad \begin{array}{c} \Sigma_2 \\ \vdots \\ \text{left-linear proof} \\ \vdots \\ (g_2 \sqsubseteq g'_2) \end{array} \quad (\prod\{g'_1, g_2\} = \perp)}{(\prod\{g_1, g_2\} = \perp)}$$

Figure 4.8: A one-path linear proof of disjointness

these holds, then the proof is as shown in Fig. 4.9.

$$\frac{\begin{array}{c} \Sigma_1 \\ \vdots \\ \text{left-linear proof} \\ \vdots \\ (g_1 \sqsubseteq g'_1) \end{array} \quad \begin{array}{c} \Sigma_2 \\ \vdots \\ \text{left-linear proof} \\ \vdots \\ (g_2 \sqsubseteq g'_2) \end{array} \quad (\prod\{g'_1, g'_2\} = \perp)}{(\prod\{g'_1, g_2\} = \perp)} \quad \frac{\quad}{(\prod\{g_1, g_2\} = \perp)}$$

Figure 4.9: A two-path linear proof of disjointness

4.28 Proposition — Deducing satisfiability of disjointness constraints *If $\text{Constr}^+(\mathfrak{G})$ is satisfiable, then for any $g_1, g_2 \in \text{Granules}\langle\mathfrak{G}\rangle$ with $\text{Constr}^+(\mathfrak{G}) \models (\prod\{g_1, g_2\} = \perp)$, there is a single-use left-linear proof in $\text{BinInfPos}\langle\mathfrak{G}\rangle$ of at most two paths of this and using at most those rules which lie in $\{(I2), (M1), (I1), (T1), (T2), (T3)\}$.*

PROOF: The proof follows directly from the constructions of 4.27. It only remains to show single usage. In all but the last case, it is trivial that the proof is single use. For the last case, it suffice to note that since g'_1 and g'_2 are disjoint, every granule in the path underlying $\langle g_1, g'_1 \rangle$ must be disjoint from every granule in the path underlying $\langle g_2, g'_2 \rangle$, so there is no possibility of a constraint being used in both chains. \square

4.29 Example For a simple example of the proof technique of 4.28, let $\mathfrak{G}_{4.29}$ be an SMAS with $\{g_i \mid i \in [1, 8]\} \subseteq \text{Granules}\langle\mathfrak{G}_{4.29}\rangle$ and $\{(g_1 \sqsubseteq g_2), (g_2 \sqsubseteq g_3), (g_3 \sqsubseteq g_4), (g_5 \sqsubseteq g_6), (g_6 \sqsubseteq g_7), (g_7 \sqsubseteq g_8), (\bigwedge\{g_4, g_8\} = \perp)\} \subseteq \text{BinConstr}^+\langle\mathfrak{G}_{4.29}\rangle$. Assume further that $\text{BinConstr}^+\langle\mathfrak{G}_{4.29}\rangle$ is satisfiable. In Fig. 4.10 is shown a two-path left-linear proof that $\text{BinConstr}^+\langle\mathfrak{G}_{4.29}\rangle \models (\bigwedge\{g_1, g_5\} = \perp)$ using the proof rules of $\text{BinInfPos}\langle\mathfrak{G}_{4.29}\rangle$.

$$\begin{array}{c}
\frac{(g_1 \sqsubseteq g_2) \quad (g_2 \sqsubseteq g_3)}{(g_1 \sqsubseteq g_3)} \quad (g_3 \sqsubseteq g_4) \quad \frac{(g_5 \sqsubseteq g_6) \quad (g_6 \sqsubseteq g_7)}{(g_5 \sqsubseteq g_7)} \quad (g_7 \sqsubseteq g_8) \\
\hline
(g_1 \sqsubseteq g_4) \quad (g_5 \sqsubseteq g_8) \quad (\bigwedge\{g_4, g_8\} = \perp) \\
\hline
(\bigwedge\{g_1, g_5\} = \perp)
\end{array}$$

Figure 4.10: Example proof of disjointness

4.30 Proposition — Deducing unsatisfiability using $\text{BinInfPos}\langle\mathfrak{G}\rangle$ *If \mathfrak{G} is not satisfiable, then there is a single-use proof in $\text{BinInfPos}\langle\mathfrak{G}\rangle$ whose last step is the application of rule (U1).*

PROOF: The completeness of $\text{BinInfPos}\langle\mathfrak{G}\rangle$ guarantees that there is a proof of the unsatisfiability of \mathfrak{G} whose last step is an instantiation of (U1), since (U1) is the only rule which allows the deduction of **false**. However, such an argument alone does not guarantee the the proof will be single use. To ensure that condition, additional work is required.

For the most general case, the idea of the proof is to build upon previous results which ensure single-use proofs, specifically 4.28. The strategy is to identify a granule $g \in \text{Granules}_{\neq}\langle\mathfrak{G}\rangle$ for which $(\bigwedge\{g, g\} = \perp)$ is provable from $\text{Constr}^+\langle\mathfrak{G}\rangle$, in a form similar to that illustrated in Fig. 4.9, and then append to the end of it an application of (U1) to deduce **false**, as illustrated in Fig. 4.11. There are, however, two details which require attention. First of all,

$$\begin{array}{c}
\begin{array}{c} \Sigma_1 \\ \vdots \\ \text{left-linear proof} \\ \vdots \\ (g \sqsubseteq g'_1) \end{array} \quad \begin{array}{c} \Sigma_2 \\ \vdots \\ \text{left-linear proof} \\ \vdots \\ (g \sqsubseteq g'_2) \quad (\bigwedge\{g'_1, g'_2\} = \perp) \end{array} \\
\hline
(g \sqsubseteq g'_1) \quad (\bigwedge\{g'_1, g\} = \perp) \\
\hline
(\bigwedge\{g, g\} = \perp) \\
\hline
\text{false}
\end{array}$$

Figure 4.11: A two-path linear proof of unsatisfiability

since $\text{Constr}^+(\mathfrak{G})$ is unsatisfiable, care must be taken in identifying what a proof is, since **false** entails anything and everything. Second, since g is used in both linear paths, a situation which does not occur in Fig. 4.9, it must be ensured via other means that no rule is used more than once.

The strategy is to switch to the proof system $\text{SQBinInfPos}\langle\mathfrak{G}\rangle$ for all deductions of Fig. 4.11 except the last one which deduces **false**. As established in 4.8, any set of constraints is satisfiable in the context of strong quasi-models. Furthermore, every rule of $\text{BinInfPos}\langle\mathfrak{G}\rangle$, save for (U1), is also a rule of $\text{SQBinInfPos}\langle\mathfrak{G}\rangle$. Therefore, all of the tree of Fig. 4.11 except the lowest level, in which (U1) is applied, is also a proof tree in $\text{SQBinInfPos}\langle\mathfrak{G}\rangle$. Hence, the entire tree is a proof tree in $\text{BinInfPos}\langle\mathfrak{G}\rangle$.

Now, let $p_1 = \langle g_{11}, g_{12}, \dots, g_{1k_1} \rangle$ (resp. $p_2 = \langle g_{21}, g_{22}, \dots, g_{2k_2} \rangle$) be the path which underlies the proof of $(g \sqsubseteq g'_1)$ from Σ_1 (resp. of $(g \sqsubseteq g'_2)$ from resp. Σ_2). As in 4.27, it is assumed that both p_1 and p_2 are reduced. If $\Sigma_1 \cap \Sigma_2 \neq \emptyset$, then some granule other than g must occur in both p_1 and p_2 . In that case, let $g' \stackrel{\text{id}}{=} g_{1i_1} \stackrel{\text{id}}{=} g_{2i_2}$ be the granule with the highest index in p_1 which also occurs in p_2 , and replace p_1 with $p'_1 = \langle g_{1i_1}, g_{1i_2}, \dots, g_{1k_1} \rangle$, p_2 with $p'_2 = \langle g_{2i_1}, g_{2i_2}, \dots, g_{2k_1} \rangle$, and g with g' throughout the proof. The result is clearly single use, since for Σ_1 and Σ_2 to have a constraint in common would require that p_1 and p_2 have at least two granules in common.

In the less general case, in which a final proof of false is appended to a tree of the form of Fig. 4.8, the proof similar but easier, since there is only one left-linear proof of subsumption. For the remaining possibilities (corresponding to the first two cases of 4.27), the proof is trivial. \square

Finally, the main result of this section may be established.

4.31 Theorem — $\text{BinInfPos}\langle\mathfrak{G}\rangle$ and single use *The proof system $\text{BinInfPos}\langle\mathfrak{G}\rangle$ is sound, complete, and single use.*

PROOF: This follows from 4.24, 4.28, and 4.30. \square

4.32 Example For a simple example of the proof technique of 4.30(b), let $\mathfrak{G}_{4.32}$ be an SMAS with $\{g_i \mid i \in [1, 7]\} \subseteq \text{Granules}\langle\mathfrak{G}_{4.32}\rangle$ and $\{(g_1 \sqsubseteq g_2), (g_2 \sqsubseteq g_3), (g_3 \sqsubseteq g_4), (g_1 \sqsubseteq g_5), (g_5 \sqsubseteq g_6), (g_6 \sqsubseteq g_7), (\bigcap\{g_4, g_7\} = \perp)\} \subseteq \text{BinConstr}^+\langle\mathfrak{G}_{4.32}\rangle$.

Assume further that $g_1 \stackrel{\text{id}}{\neq} \perp_{\mathfrak{G}_{4.32}}$. In Fig. 4.12 is shown a proof that $\text{BinConstr}^+\langle\mathfrak{G}_{4.32}\rangle$ is unsatisfiable, using the proof rules of $\text{BinInfPos}\langle\mathfrak{G}_{4.32}\rangle$.

5 Negation in the Armstrong Context

In the remainder of this paper, the goal is to show how to augment the inference system $\text{BinInfPos}\langle\mathfrak{G}\rangle$ of 4.13 to include all assertions in $\text{BinConstr}^\pm\langle\mathfrak{G}\rangle$; that is, to include those in $\text{BinConstr}^-\langle\mathfrak{G}\rangle$ as well as those of $\text{BinConstr}^+\langle\mathfrak{G}\rangle$ which are already covered by $\text{BinInfPos}\langle\mathfrak{G}\rangle$. The approach taken is a general one, not limited to $\text{BinConstr}^+\langle\mathfrak{G}\rangle$. Rather, it applies to

$$\begin{array}{c}
\frac{(g_1 \sqsubseteq g_2) \quad (g_2 \sqsubseteq g_3)}{(g_1 \sqsubseteq g_3)} \quad \frac{(g_3 \sqsubseteq g_4)}{(g_1 \sqsubseteq g_4)} \quad \frac{(g_1 \sqsubseteq g_5) \quad (g_5 \sqsubseteq g_6)}{(g_1 \sqsubseteq g_6)} \quad \frac{(g_6 \sqsubseteq g_7)}{(g_1 \sqsubseteq g_7)} \quad (\bigcap\{g_4, g_7\} = \perp) \\
\hline
\frac{(g_1 \sqsubseteq g_4) \quad (\bigcap\{g_1, g_4\} = \perp)}{(\bigcap\{g_1\} = \perp)} \\
\hline
\mathbf{false}
\end{array}$$

Figure 4.12: Example proof of unsatisfiability

any Armstrong context \mathcal{C} . There is absolutely no added complexity in taking this more general approach. On the contrary, it is easier to develop the ideas independently of a specific context. This approach not only avoids involving irrelevant properties specific to $\text{BinConstr}^\pm(\mathfrak{G})$, but also opens the door for these results to be applied in other contexts. Nevertheless, explicit guidance on how the general results apply to $\text{BinConstr}^\pm(\mathfrak{G})$ will always be kept at the forefront.

In this section, some essential properties of negation in an Armstrong Context are developed. In the next section, Sec. 6, the construction of a set of inference rules which are complete for both positive and negative assertions is provided.

5.1 Context Unless stated specifically to the contrary, for the rest of this paper, \mathcal{L} will denote a general logical system (see 3.6), with $\mathcal{C} \subseteq \text{WFF}\langle\mathcal{L}\rangle$ an Armstrong context (see 3.12). In other words, every satisfiable subset of \mathcal{C} will admit an Armstrong model. For readers not interested in any generalization, it suffices to take $\mathcal{C} = \text{BinConstr}^+(\mathfrak{G})$, with \mathcal{L} the logic which includes all assertions in $\text{BinConstr}^\pm(\mathfrak{G})$.

5.2 Armstrong constraint sets It is useful to have an explicit notation for the negations of the assertions in \mathcal{C} . To this end, for any $\Phi \subseteq \mathcal{C}$, define $\text{Not}\langle\Phi\rangle = \{\neg\varphi \mid \varphi \in \Phi\}$, and define $\text{All}^\pm(\mathcal{C}) = \mathcal{C} \cup \text{Not}\langle\mathcal{C}\rangle$. For the specific case of $\mathcal{C} = \text{BinConstr}^+(\mathfrak{G})$, it is immediate that $\text{Not}\langle\text{BinConstr}^+(\mathfrak{G})\rangle = \text{BinConstr}^-(\mathfrak{G})$ and $\text{All}^\pm(\text{BinConstr}^+(\mathfrak{G})) = \text{BinConstr}^\pm(\mathfrak{G})$.

An *extended Armstrong constraint set* over \mathcal{C} is a satisfiable subset of $\text{All}^\pm(\mathcal{C})$. In particular, for $\mathcal{C} = \text{BinConstr}^+(\mathfrak{G})$, an extended Armstrong constraint set is a subset of $\text{BinConstr}^\pm(\mathfrak{G})$.

For $\Phi \subseteq \text{All}^\pm(\mathcal{C})$, define $\text{Only}^+(\Phi) = \Phi \cap \mathcal{C}$, and define $\text{Only}^-(\Phi) = \Phi \cap \text{Not}\langle\mathcal{C}\rangle$. Thus, if $\mathcal{C} = \text{BinConstr}^+(\mathfrak{G})$, then $\text{Only}^+(\Phi) = \Phi \cap \text{BinConstr}^+(\mathfrak{G})$ and $\text{Only}^-(\Phi) = \Phi \cap \text{BinConstr}^-(\mathfrak{G})$.

5.3 Equivalent and complementary pairs Two WFFs may be equivalent without being identical. For example, in $\text{BinConstr}^+(\mathfrak{G})$, $(\perp_{\mathfrak{G}} \sqsubseteq g_1)$ and $(g_2 \sqsubseteq \top_{\mathfrak{G}})$ are equivalent for any $g_1, g_2 \in \text{Granules}\langle\mathfrak{G}\rangle$, since both are tautologies and so true in any granular structure.

More generally, $\varphi_1, \varphi_2 \in \text{WFF}\langle\mathcal{L}\rangle$ are *equivalent* if they have the same models. In this case, write $\varphi_1 \equiv \varphi_2$, and say that $\{\varphi_1, \varphi_2\}$ forms an *equivalent pair*.

A two-element set $\{\varphi_1, \varphi_2\} \subseteq \text{WFF}\langle\mathcal{L}\rangle$ forms a *complementary pair* if $\{\varphi_1, \neg\varphi_2\}$ forms an equivalent pair, with φ_1 and φ_2 called *complements* of one another.

5.4 Example — Complementary pairs in $\text{BinConstr}^+\langle\mathfrak{G}\rangle$ For any $g_1, g_2 \in \text{Granules}_\perp\langle\mathfrak{G}\rangle$, $\{(\perp_{\mathfrak{G}} \sqsubseteq g_1), (\prod\{g_2, g_2\} = \perp)\}$ forms a complementary pair of elements in $\mathcal{C} = \text{BinConstr}^+\langle\mathfrak{G}\rangle$, with $(\perp_{\mathfrak{G}} \sqsubseteq g_1)$ a tautology and $(\prod\{g_2, g_2\} = \perp)$ unsatisfiable.

It holds generally that a complementary pair in \mathcal{C} must consist of a tautology and an unsatisfiable assertion. Central to proving this is to note that the empty set admits an Armstrong model, as established below.

5.5 Lemma — Armstrong model for \emptyset *In the context identified in 5.1, the empty set $\emptyset \subseteq \mathcal{C}$ admits an Armstrong model.*

PROOF: \mathcal{L} is assumed to be nontrivial in the sense that there is a $\varphi \in \text{WFF}\langle\mathcal{L}\rangle$ which admits a model (see 3.6). Since any model of φ is also a model of \emptyset , it follows that the empty set is satisfiable. Since \mathcal{C} is an Armstrong context and $\emptyset \subseteq \mathcal{C}$, \emptyset must have an Armstrong model, as required. \square

5.6 Lemma — Complementary pairs in the Armstrong context *Let $\{\varphi_1, \varphi_2\} \subseteq \mathcal{C}$. If $\{\varphi_1, \varphi_2\}$ is a complementary pair, then one must be a tautology, with the other unsatisfiable.*

PROOF: Let M_\emptyset be an Armstrong model for \emptyset , which must exist in view of 5.5, and let $\{\varphi_1, \varphi_2\} \subseteq \mathcal{C}$ be a complementary pair. M_\emptyset is a model of φ_1 iff it is a tautology (just by definition of tautology; see, [Mon76, Def. 8.22] or [End01, p. 23]), and similarly for φ_2 . Thus, if neither is a tautology, then M_\emptyset is a model of neither, and so it must be a model of both $\neg\varphi_1$ and $\neg\varphi_2$, which is impossible, since they are complements of one another. Thus, one of φ_1 or φ_2 must be a tautology, completing the proof. \square

The characterization of overlap of \mathcal{C} and $\text{Not}\langle\mathcal{C}\rangle$ now follows easily.

5.7 Proposition — Overlap of \mathcal{C} and $\text{Not}\langle\mathcal{C}\rangle$ in the Armstrong context *If $\varphi_1 \in \mathcal{C}$ and $\varphi_2 \in \text{Not}\langle\mathcal{C}\rangle$ with $\varphi_1 \equiv \varphi_2$, then φ_1 must be either a tautology or else unsatisfiable.*

PROOF: Assume that $\varphi_1 \in \mathcal{C}$, $\varphi_2 \in \text{Not}\langle\mathcal{C}\rangle$ with $\varphi_1 \equiv \varphi_2$. Then φ_2 is of the form $\neg\varphi_3$ for some $\varphi_3 \in \mathcal{C}$, whence $\varphi_1 \equiv \neg\varphi_3$. It then follows that $\{\varphi_1, \varphi_3\}$ is a complementary pair, whence by 5.6, one of φ_1, φ_3 must be a tautology, with the other unsatisfiable. \square

With the preceding results in hand, the focus is now turned to the more general topic of incorporating negation in inference. Although the idea of contraposition is well-known in logic, it is worthwhile to recall it precisely here, since the idea is central to results of this section.

5.8 Lemma — General swapping of premise and conclusion *Let $\mathfrak{A} \subseteq \text{WFF}\langle\mathcal{L}\rangle$ be closed under logical negation; i.e., $\text{Not}\langle\mathfrak{A}\rangle \subseteq \mathfrak{A}$. (Note that this holds in particular for $\mathfrak{A} = \text{All}^\pm\langle\mathcal{C}\rangle$.) In addition, let $\Phi \subseteq \mathfrak{A}$ be finite with $\varphi, \psi \in \mathfrak{A}$,*

(a) $\Phi \cup \{\varphi\} \models \psi$ *iff* $\Phi \cup \{\neg\psi\} \models \neg\varphi$.

(b) *If $\Phi \cup \{\varphi\} \models \psi$ and $\Phi \cup \{\varphi\}$ is satisfiable, with φ is essential in the sense that $\Phi \not\models \psi$, then $\Phi \cup \{\neg\varphi\}$ is also satisfiable.*

In particular, taking ψ to be unsatisfiable, the following special cases are obtained.

(a') $\Phi \cup \{\varphi\}$ *is unsatisfiable iff* $\Phi \models \neg\varphi$,

(b') *If $\Phi \cup \{\varphi\}$ is unsatisfiable and φ is essential to that unsatisfiability in the sense that Φ is satisfiable, then $\Phi \cup \{\neg\varphi\}$ is also satisfiable.*

PROOF: Part (a) is just *contraposition* [End01, Cor. 24D]. For part (b), since $\Phi \not\models \psi$, there is a model M of $\Phi \cup \{\neg\psi\}$, which must also be a model of $\neg\varphi$ by (a). Hence M is also a model of $\Phi \cup \{\neg\varphi\}$. Parts (a') and (b') follow by taking ψ to be the identically false assertion **false** in (a) and (b). \square

5.9 Remark — Swapping in the presence of unsatisfiability In the first part of 5.8, it is important to note that if $\Phi \models \psi$; that is, if φ is superfluous in $\Phi \cup \{\varphi\} \models \psi$, then $\Phi \cup \{\neg\psi\}$ must be unsatisfiable, so that $\Phi \cup \{\neg\psi\} \models \neg\varphi$ holds in a rather trivial way, as **false** $\models \neg\varphi$. Just take φ to be the identically true assertion **true** to see this.

Similarly, in the second part, if Φ is unsatisfiable, so that $\Phi \cup \{\varphi\}$ is unsatisfiable regardless of φ , then $\Phi \models \neg\varphi$ holds again in a rather trivial way, since false implies anything.

The point is that while the result holds in general, it is, for the most part, interesting only in the case that φ is essential on the left-hand side for the entailment to hold.

When applied within an Armstrong context, the general idea of swapping premise and conclusion leads to remarkable results. Specifically, negative premises are of no consequence in deriving positive conclusions, and in the derivation of a negative conclusion, at most one negative premise is necessary. These ideas are formalized in the next two results.

5.10 Lemma — Satisfiability and unsatisfiability in extended Armstrong constraint sets *Let Φ be an extended Armstrong constraint set over \mathcal{C} ,*

(a) *If Φ satisfiable, then any Armstrong model of $\text{Only}^+\langle\Phi\rangle$ relative to \mathcal{C} is also a model of all of Φ .*

(b) *If Φ is not satisfiable, then either $\text{Only}^+\langle\Phi\rangle$ is unsatisfiable or else there is some $\neg\varphi \in \text{Only}^-\langle\Phi\rangle$ with $\text{Only}^+\langle\Phi\rangle \cup \{\neg\varphi\}$ unsatisfiable.*

PROOF: For part (a), if $\Phi = \text{Only}^+\langle\Phi\rangle$, then there is nothing further to prove. Otherwise, let M be an Armstrong model of Φ relative to \mathcal{C} , and let $\neg\varphi \in \text{Only}^-\langle\Phi\rangle$. Since $\Phi \cup \{\neg\varphi\}$ is satisfiable, it follows from 5.8(a) that $\Phi \not\models \varphi$. Thus, since M is an Armstrong model of Φ , it

cannot be a model of φ , whence it is a model of $\neg\varphi$. Since $\neg\varphi$ was chosen arbitrarily from $\text{Only}^-\langle\Phi\rangle$, it follows that M is a model of all of Φ , as required.

For part (b), assume that Φ is not satisfiable. If $\text{Only}^+\langle\Phi\rangle$ is not satisfiable, there is nothing further to prove. So, assume that $\text{Only}^+\langle\Phi\rangle$ is satisfiable, and let M be an Armstrong model of $\text{Only}^+\langle\Phi\rangle$ relative to \mathcal{C} . Since Φ is not satisfiable, there must be some $\neg\varphi \in \text{Only}^-\langle\Phi\rangle$ with $M \notin \text{Mod}(\neg\varphi)$, whence $M \in \text{Mod}(\varphi)$. Thus, by the definition of Armstrong model, $\text{Only}^+\langle\Phi\rangle \models \varphi$. Finally, applying 5.8(a'), $\text{Only}^+\langle\Phi\rangle \cup \{\neg\varphi\}$ is unsatisfiable, completing the proof. \square

5.11 Proposition — Entailment in extended Armstrong constraint sets *Let Φ be an extended Armstrong constraint set over \mathcal{C} , and let $\beta \in \mathcal{C}$.*

- (a) *If $\Phi \models \beta$, then it must be the case that $\text{Only}^+\langle\Phi\rangle \models \beta$.*
- (b) *If $\Phi \models \neg\beta$, then either $\text{Only}^+\langle\Phi\rangle \models \neg\beta$, or else there is a $\neg\varphi \in \text{Only}^-\langle\Phi\rangle$ with the property that $\text{Only}^+\langle\Phi\rangle \cup \{\neg\varphi\} \models \neg\beta$ (and hence $\text{Only}^+\langle\Phi\rangle \cup \{\beta\} \models \varphi$) holds, with both $\text{Only}^+\langle\Phi\rangle \cup \{\neg\varphi\}$ and $\text{Only}^+\langle\Phi\rangle \cup \{\beta\}$ satisfiable.*

PROOF: Part (a): If Φ is not satisfiable, the result holds trivially. So, assume that Φ is satisfiable with $\Phi \models \beta$, and let M be an Armstrong model for $\text{Only}^+\langle\Phi\rangle$. By 5.10(a), M is a model of all of Φ , and hence a model of β as well. Finally, by the definition of Armstrong model, the only constraints in \mathcal{C} which hold in M are those implied by $\text{Only}^+\langle\Phi\rangle$, whence $\text{Only}^+\langle\Phi\rangle \models \beta$.

Part (b): If Φ is not satisfiable, then the result is immediate. So, assume that Φ is satisfiable with $\Phi \models \neg\beta$. In view of 5.8(a'), $\Phi \cup \{\beta\}$ must be unsatisfiable. Then, applying 5.10(b), it must be the case that either $\text{Only}^+\langle\Phi\rangle \cup \{\beta\}$ is unsatisfiable or else there is $\neg\varphi \in \text{Only}^-\langle\Phi\rangle$ with $\text{Only}^+\langle\Phi\rangle \cup \{\beta\} \cup \{\neg\varphi\}$ unsatisfiable. If $\text{Only}^+\langle\Phi\rangle \cup \{\beta\}$ is unsatisfiable, a second application of 5.8(a') yields $\text{Only}^+\langle\Phi\rangle \models \neg\beta$. If $\text{Only}^+\langle\Phi\rangle \cup \{\beta\}$ is satisfiable but there is $\neg\varphi \in \text{Only}^-\langle\Phi\rangle$ with $\text{Only}^+\langle\Phi\rangle \cup \{\beta\} \cup \{\neg\varphi\}$ unsatisfiable, an application of 5.8(a) yields $\text{Only}^+\langle\Phi\rangle \cup \{\neg\varphi\} \models \neg\beta$, completing the proof. \square

6 Inference on Binary Rules Involving Negation

In this section, it is shown how to extend a sound and complete proof system \mathcal{A} for an Armstrong context \mathcal{C} to one which applies to all of $\text{All}^\pm(\mathcal{C})$. It builds upon the theoretical ideas developed in Sec. 5, together with the fundamental operation of *swapping* to each rule in \mathcal{A} , in which the conclusion of a rule is swapped with a premise, and each negated. It is shown that augmenting \mathcal{A} with the set of all such swapped rules results in a sound and complete inference system for $\text{All}^\pm(\mathcal{C})$.

6.1 Context Throughout this section, unless stated specifically to the contrary, \mathcal{A} will be taken to be a sound but not necessarily complete inference system for \mathcal{C} . Note that the context introduced in 5.1 continues to hold.

As in the previous section, the reader who is not interested in the general case may fix \mathcal{C} to be $\text{BinConstr}^+(\mathfrak{G})$.

While the following result is straightforward, it is recorded formally because it provides the key connection between the results of Sec. 5 and the type of proof system developed in this section.

6.2 Proposition — Satisfiable inference in the Armstrong setting *Let Φ be a satisfiable subset of \mathcal{C} , and let $\beta \in \mathcal{C}$ also be satisfiable. Assume furthermore that \mathcal{A} complete.*

- (a) $\Phi \models \beta$ holds iff $\text{Only}^+(\Phi) \vdash_{\mathcal{A}} \beta$.
- (b) $\Phi \models \neg\beta$ holds iff either $\text{Only}^+(\Phi) \cup \{\beta\} \vdash_{\mathcal{A}} \mathbf{false}$ or else there is a $\neg\varphi \in \text{Only}^-(\Phi)$ for which $\text{Only}^+(\Phi) \cup \{\beta\} \vdash_{\mathcal{A}} \varphi$.

PROOF: The proof follows immediately from 5.11 and the fact that \mathcal{A} is sound and complete for \mathcal{C} . \square

6.3 Swapping and swap closure Let \mathcal{A} be an inference system for \mathcal{C} . Say that a rule $r \in \text{InfRules}\langle\mathcal{A}\rangle$ has nonempty antecedent set if $\text{Antecedents}\langle r \rangle \neq \emptyset$.

Let $r \in \text{InfRules}\langle\mathcal{A}\rangle$ have nonempty antecedent set $\text{Antecedents}\langle r \rangle = \{\alpha_i \mid i \in [1, k]\}$ and $\text{Consequent}\langle r \rangle = \{\beta\}$. The α_i -swap of r , denoted $\text{Swap}\langle r: \alpha_i \rangle$, is the rule obtained by replacing α_i with $\neg\beta$ as an antecedent, and replacing β with $\neg\alpha_i$ as the consequent. Visually, the transformation is

$$\frac{\alpha_1 \ \alpha_2 \ \dots \ \alpha_{i-1} \ \alpha_i \ \alpha_{i+1} \ \dots \ \alpha_{k-1} \ \alpha_k}{\beta} \rightsquigarrow \frac{\alpha_1 \ \alpha_2 \ \dots \ \alpha_{i-1} \ \neg\beta \ \alpha_{i+1} \ \dots \ \alpha_{k-1} \ \alpha_k}{\neg\alpha_i} \quad (6.3-1)$$

If β is unsatisfiable; *i.e.*, $\beta \equiv \mathbf{false}$, then $\neg\beta$ is simply omitted as a premise. In this case, the transformation is

$$\frac{\alpha_1 \ \alpha_2 \ \dots \ \alpha_{i-1} \ \alpha_i \ \alpha_{i+1} \ \dots \ \alpha_{k-1} \ \alpha_k}{\mathbf{false}} \rightsquigarrow \frac{\alpha_1 \ \alpha_2 \ \dots \ \alpha_{i-1} \ \alpha_{i+1} \ \dots \ \alpha_{k-1} \ \alpha_k}{\neg\alpha_i} \quad (6.3-2)$$

Define the *swap closure* $\text{SwapCl}\langle\mathcal{A}\rangle$ of \mathcal{A} to be the inference system on $\text{All}^\pm(\mathcal{C})$ given by $\text{InfRules}\langle\text{SwapCl}\langle\mathcal{A}\rangle\rangle = \text{InfRules}\langle\mathcal{A}\rangle \cup \{\text{Swap}\langle r: \alpha \rangle \mid r \in \text{InfRules}\langle\mathcal{A}\rangle \text{ and } \text{Antecedents}\langle r \rangle \neq \emptyset \text{ and } \alpha \in \text{Antecedents}\langle r \rangle\}$.

6.4 Proposition — Soundness of swapping

- (a) *If r is a sound and minimal inference rule of \mathcal{A} , then $\text{Swap}\langle r: \alpha_i \rangle$ is a sound and minimal inference rule on $\text{All}^\pm(\mathcal{C})$.*
- (b) *If every rule of \mathcal{A} is sound and minimal, then $\text{SwapCl}\langle\mathcal{A}\rangle$ is sound as well.*

PROOF: The proof of (a) follows directly from 5.8. The proof of (b) then follows immediately from the definition of swap closure. \square

6.5 The swap closure of $\text{BinInfPos}\langle\mathfrak{G}\rangle$ Upon applying the construction of 6.3(b) to the rules of $\text{BinInfPos}\langle\mathfrak{G}\rangle$, a new inference system, denoted $\text{BinInf}\langle\mathfrak{G}\rangle$, is obtained. Its rules are defined by

$\text{InfRules}\langle\text{BinInf}\langle\mathfrak{G}\rangle\rangle = \text{InfRules}\langle\text{SwapCl}\langle\text{BinInfPos}\langle\mathfrak{G}\rangle\rangle\rangle$
 $= \{(I2), (M1), (I1), (T1), (T2), (T3), (U1), (I2\text{-sa}), (I2\text{-sb}), (M2\text{-sa}), (M2\text{-sb}), (U1\text{-s})\}$
 with five additional rules not found in $\text{InfRules}\langle\text{BinInfPos}\langle\mathfrak{G}\rangle\rangle$ shown in Fig. 6.1, with the way in which each rule was obtained via swapping shown to its far right. Note that rules (I1) and (T1)–(T3) are not swapped, since they all have empty antecedent set.

$$\begin{array}{l}
 (I2\text{-sa}) \frac{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_2) (\mathfrak{g}_1 \not\sqsubseteq \mathfrak{g}_3)}{(\mathfrak{g}_2 \not\sqsubseteq \mathfrak{g}_3)} \quad (\text{Swap}\langle(I2):(\mathfrak{g}_2 \sqsubseteq \mathfrak{g}_3)\rangle) \\
 \\
 (I2\text{-sb}) \frac{(\mathfrak{g}_1 \not\sqsubseteq \mathfrak{g}_3) (\mathfrak{g}_2 \sqsubseteq \mathfrak{g}_3)}{(\mathfrak{g}_1 \not\sqsubseteq \mathfrak{g}_2)} \quad (\text{Swap}\langle(I2):(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_2)\rangle) \\
 \\
 (M1\text{-sa}) \frac{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}'_1) (\prod\{\mathfrak{g}_1, \mathfrak{g}_2\} \neq \perp)}{(\prod\{\mathfrak{g}'_1, \mathfrak{g}'_2\} \neq \perp)} \quad (\text{Swap}\langle(M1):(\prod\{\mathfrak{g}'_1, \mathfrak{g}'_2\} = \perp)\rangle) \\
 \\
 (M1\text{-sb}) \frac{(\prod\{\mathfrak{g}_1, \mathfrak{g}_2\} \neq \perp) (\prod\{\mathfrak{g}'_1, \mathfrak{g}_2\} = \perp)}{(\mathfrak{g}_1 \not\sqsubseteq \mathfrak{g}'_1)} \quad (\text{Swap}\langle(M1):(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}'_1)\rangle) \\
 \\
 (U1\text{-s}) \frac{}{(\prod\{\mathfrak{g}, \mathfrak{g}\} \neq \perp)} \langle!(\mathfrak{g} \neq \perp)\rangle \quad (\text{Swap}\langle(U1):(\prod\{\mathfrak{g}, \mathfrak{g}\} = \perp)\rangle)
 \end{array}$$

Figure 6.1: Swapped rules for $\text{BinInfPos}\langle\mathfrak{G}\rangle$

The goal is to show that $\text{BinInf}\langle\mathfrak{G}\rangle$ is both sound and complete. The first part is easy.

6.6 Proposition — Soundness of $\text{BinInf}\langle\mathfrak{G}\rangle$ *The inference system $\text{BinInf}\langle\mathfrak{G}\rangle$ is sound.*

PROOF: The proof follows immediately from 6.4. \square

6.7 Conversion to positive-only proofs Let the setting be as in 6.2(b), and suppose that $\Phi \models \neg\beta$. Of course, there is no proof of this using \mathcal{A} , since that proof system does not include members of $\text{Not}\langle\mathcal{C}\rangle$ for either antecedents or consequents of rules. However, using 6.2(b) and the completeness of \mathcal{A} , it must be the case that either $\text{Only}^+\langle\Phi\rangle \cup \{\beta\} \vdash_{\mathcal{A}} \mathbf{false}$ or else $\text{Only}^+\langle\Phi\rangle \cup \{\beta\} \vdash_{\mathcal{A}} \varphi$ for some $\neg\varphi \in \text{Only}^-\langle\Phi\rangle$. The idea is to take a proof in \mathcal{A} for $\text{Only}^+\langle\Phi\rangle \cup \{\beta\} \vdash_{\mathcal{A}} \mathbf{false}$ or $\text{Only}^+\langle\Phi\rangle \cup \{\beta\} \vdash_{\mathcal{A}} \varphi$, as the case may be, and convert it to a proof $\Phi \vdash_{\text{SwapCl}(\mathcal{A})} \beta$ by *contrapositioning* certain rules in the original proof, which is developed

in detail in 6.13 below. In support of this, it is first useful to introduce an alternative representation for proof trees, using trees represented in vertex-edge format..

6.8 The tree of an inference rule The notation for proof rules in which the premises are separated from the conclusion by a single horizontal line is something of a standard in logic. Implicitly, it represents a tree structure. However, for the purposes of formalizing the process of contrapositioning, it is advantageous to make this tree structure explicit, as illustrated in (6.8-1) and (6.8-2).

$$\frac{\alpha_1 \ \alpha_2 \ \dots \ \alpha_{i-1} \ \alpha_i \ \alpha_{i+1} \ \dots \ \alpha_{k-1} \ \alpha_k}{\beta} \ d \quad \equiv \quad \begin{array}{c} \alpha_1 \ \alpha_2 \ \dots \ \alpha_{i-1} \ \alpha_i \ \alpha_{i+1} \ \dots \ \alpha_{k-1} \ \alpha_k \\ \swarrow \quad \nwarrow \quad \quad \quad \nearrow \quad \nwarrow \quad \quad \quad \nearrow \quad \nwarrow \quad \quad \quad \nearrow \\ d \quad d \quad \quad \quad d \quad d \quad \quad \quad d \quad d \quad \quad \quad d \quad d \end{array} \beta \quad (6.8-1)$$

$$\frac{}{\beta} \ e \quad \equiv \quad \begin{array}{c} \mathbf{true} \\ \uparrow \\ e \\ \beta \end{array} \quad (6.8-2)$$

Each antecedent, as well as the consequent, is a vertex. The consequent is the root of the tree, with a directed edge from it to each antecedent, as illustrated in (6.8-1). If the antecedent set is empty, an artificial antecedent vertex **true** is added, as illustrated in (6.8-2). Formally, each edge is labelled with the instance name for that rule. However, those labels may be omitted if the name of the instance is clear from context.

6.9 Proof trees in vertex-edge format Proof trees are represented in vertex-edge format in the obvious way, by connecting the individual trees together. Fig. 6.2 shows the proof trees for the proofs of (3.9-3) and (3.9-4).






Figure 6.2: Vertex-edge representations for (3.9-3) and (3.9-4)

6.10 Swapped inference rules in vertex-edge format Given an inference rule r , the representation of $\text{Swap}\langle r:\alpha_i \rangle$ is, in principle, straightforward, as illustrated in (6.10-1). All that is necessary is to swap the identities of the vertex labelled α_i with that of the conclusion β , and negate each. The edges remain fixed; thus, this representation is called *fixed-edge positioning*.



While this works fine for an isolated inference rule, it leads to problems when such rules are interconnected to form proofs because vertices must be moved around, creating a representational maze. It is more suitable to keep the position of the vertices fixed, and move the edges as necessary. This *fixed-vertex positioning* is shown in (6.10-2).



Although the vertices α_i and β have been negated, all retain their original positions. On the other hand, the edges must be redrawn. In addition to reversing the original arrow from β to α_i , each edge from β to α_j for $j \neq i$ is replaced by an edge from α_i to α_j .

6.11 Swapping of rule instances The notion of swapping introduced in 6.3 extends naturally to rule instances. Specifically, if d is an instance of rule r obtained via substitution s (see 3.5 and 3.9), then for $\alpha_i \in \text{Antecedents}\langle r \rangle$, $\text{Swap}\langle d:\text{Subst}\langle \alpha_i, s \rangle \rangle$ is the rule instance obtained by applying s to all entries of $\text{Swap}\langle r:\alpha_i \rangle$.

6.12 Proof trees in multiple-use contexts In order to describe the process of proof contrapositioning (see 6.13), it is necessary to have a means of identifying the vertices of the proof tree unambiguously. For the case of single-use proofs, this is an easy task, since the WFF associated with that vertex identifies it uniquely. However, if the proof is not single use, there may be several vertices labelled with the same WFF. In that case, the solution is to identify a non-root vertex as a pair $\langle \alpha, d \rangle$ in which α is the WFF and d is the rule for which $\alpha \in \text{Antecedents}\langle d \rangle$. The root vertex is identified specially as $\langle \beta, \text{root} \rangle$. As a concrete example, consider the proof tree of (3.11-1), which is shown in Fig. 6.3. There are two vertices labelled $(g_1 \sqsubseteq g_2)$; one is a leaf, an antecedent of rule d_1 , while the second is an interior vertex, an antecedent of rule d_3 . These are represented as $\langle (g_1 \sqsubseteq g_2), d_1 \rangle$ and $\langle (g_1 \sqsubseteq g_2), d_3 \rangle$, respectively. Note that the root vertex is denoted $\langle (g_1 \sqsubseteq g_4), \text{root} \rangle$.

Given a proof tree τ , notation such as $\langle \alpha, d \rangle \in \text{Antecedents}\langle \tau \rangle$ will be used with the obvious meaning — that $\alpha \in \text{Antecedents}\langle \tau \rangle$.




Figure 6.3: Vertex-edge representation for (3.11-1)

It should be observed that this convention assumes that a single rule instance does not have multiple instances of the same antecedent. In most situations, this is a reasonable assumption. To work with multiple instances of the same antecedent, it is necessary to employ trees which have ordered vertices, and to tag the instances with an index based upon that order. The details are straightforward, but will not be used in this work, because they would render the presentation much more complex.

6.13 Proof contrapositioning Figs. 6.4, 6.5, and 6.6 illustrate the strategy of *proof contrapositioning*, in which rule swapping is applied to an entire proof tree. Figure 6.4 shows a *source tree*, call it τ , in vertex-edge format. It is an arbitrary proof tree using the rules

\mathcal{A} ; no special properties are assumed. Each entry of the form $\nabla_{ij}^{T_{ij}}$ represents a subtree of the entire proof tree which remains unchanged during the transformation. The most important vertex in the tree is $\langle \alpha_{1j_1}, d_1 \rangle$, called the (*designated*) *swap leaf* of τ . It is the antecedent which will be swapped with the consequent β to obtain a proof of $\neg \alpha_{1j_1}$ from $(\text{Antecedents}\langle \tau \rangle \setminus \{\alpha_{1j_1}\}) \cup \{\neg \beta\}$. The *swap path* is

$$\langle \langle \alpha_{1j_1}, d_1 \rangle, \langle \alpha_{2j_2}, d_2 \rangle, \dots, \langle \alpha_{ij_i}, d_i \rangle, \dots, \langle \alpha_{(k-1)j_{k-1}}, d_{i-1} \rangle, \langle \alpha_{kj_k}, d_k \rangle, \langle \beta, \text{root} \rangle \rangle,$$

the (unique) path from the swap leaf to the root. The members of this sequence are precisely the vertices whose assertions will be negated in the contrapositioned proof. In Fig. 6.4, this path goes right down the middle of the diagram. For $i \in [1, k-1]$, $\text{Consequent}\langle d_i \rangle = \alpha_{(i+1)j_{i+1}} \in \text{Antecedents}\langle d_{i+1} \rangle$, with $\text{Consequent}\langle d_k \rangle = \beta$.

In the result, called the *contrapositioned tree*, each rule instance d_i in the swap-rule sequence is replaced by $\text{Swap}\langle d_i; \alpha_{ij_i} \rangle$. This tree is illustrated in Fig. 6.5, using fixed-vertex positioning as described in 6.10. For compactness of notation, this replacement rule is labelled d'_i in Fig. 6.5. Notice that the changes described in 6.10 are followed. In particular, in the transition from d_i to d'_i , the antecedent α_{ij_i} and the consequent $\alpha_{(i+1)j_{i+1}}$ (β for d_k) are negated, and the arrow connecting them is reversed. Furthermore, the source of all arrows of d_i is changed to $\neg \alpha_{ij_i}$ in d'_i .




Figure 6.4: The source proof to undergo contrapositioning

In Fig. 6.6, this same contrapositioned tree is shown in fixed-edge format. Although the result may seem more aesthetically pleasing, note that there has been substantial rearrangement — the swapped rules occur in the flipped tree in the opposite order from that in the source tree, since the swapping operation exchanges an antecedent of the rule with the




Figure 6.5: The contrapositioned proof in fixed-position format

consequent (and then complements both). Thus, while fixed-edge positioning may result in a tree which looks more natural, it is fixed-vertex positioning which illustrates, in a far more understandable manner, the relationship between the source tree and its contrapositioned counterpart.




Figure 6.6: The contrapositioned proof in hierarchical format

The contrapositioning of the tree τ is denoted $\text{Contraposn}\langle\tau:\langle\alpha_{1j_i}, d_1\rangle\rangle$. The swap leaf determines the entire swap path.

For examples of contrapositioning, the reader is encouraged to look ahead to 6.20 and 6.21. The immediate topic is to establish the key completeness result surrounding contrapositioning.

6.14 Lemma — Contrapositioning preserves proof *If τ is a proof tree over \mathcal{A} with $\langle \alpha, d \rangle \in \text{Antecedents}\langle \tau \rangle$, then $\text{Contraposn}\langle \tau: \langle \alpha, d \rangle \rangle$ is a proof of $\neg\alpha$ from $(\text{Antecedents}\langle \tau \rangle \setminus \{\alpha\}) \cup \{\neg\text{Consequent}\langle \tau \rangle\}$ in the proof system $\text{SwapCl}\langle \mathcal{A} \rangle$.*

PROOF: The result follows directly from the construction given in 6.13. It is easy to verify that $\text{Contraposn}\langle \tau: \langle \alpha, d \rangle \rangle$ is a proof tree constructed from the rules of $\text{SwapCl}\langle \mathcal{A} \rangle$, with the required antecedents and consequent. \square

6.15 Theorem — Sat-completeness of Armstrong inference systems *If \mathcal{A} is sound, complete and single use, then $\text{SwapCl}\langle \mathcal{A} \rangle$ is sound, sat-complete, and single use also.*

PROOF: Consistency of $\text{SwapCl}\langle \mathcal{A} \rangle$ follows from 6.4(b). To show sat-completeness, assume that \mathcal{A} is complete, and let $\Phi \subseteq \text{All}^\pm(\mathcal{C})$ be satisfiable, and let $\psi \in \mathcal{C}$. There are two cases to consider: $\Phi \models \psi$ and $\Phi \models \neg\psi$.

For the first case, $\Phi \models \psi$, it suffices to invoke 6.2(a), which ensures that $\text{Only}^+(\Phi) \vdash_{\mathcal{A}} \psi$, whence $\Phi \vdash_{\mathcal{A}} \psi$ holds *a fortiori*.

For the second case, if $\Phi \models \neg\psi$, then by 6.2(b), either $\text{Only}^+(\Phi) \cup \{\psi\} \vdash_{\mathcal{A}} \mathbf{false}$, or else there is a single $\neg\varphi \in \text{Only}^-(\Phi)$ with the property that $\text{Only}^+(\Phi) \cup \{\psi\} \models \varphi$. For the first subcase, $\text{Only}^+(\Phi) \cup \{\psi\} \vdash_{\mathcal{A}} \mathbf{false}$, let τ be the proof tree in \mathcal{A} modelled after that of Fig. 6.4, with $\alpha_{1j_1} = \psi$ and $\beta = \mathbf{false}$. For the second subcase, $\text{Only}^+(\Phi) \cup \{\psi\} \models \varphi$ for some $\neg\varphi \in \text{Only}^-(\Phi)$, let τ be the proof tree in \mathcal{A} modelled after that of Fig. 6.4, with $\alpha_{1j_1} = \psi$ and $\beta = \varphi$. In either subcase, the completeness of \mathcal{A} guarantees that such a tree exists. If ψ is used more than once as an antecedent, pick any occurrence. (It will be shown shortly that such multiple occurrences are impossible.)

Now, consider the tree $\tau' = \text{Contraposn}\langle \tau: \langle \psi, d_1 \rangle \rangle$, with d_1 the rule which has $\psi = \alpha_{1j_1}$ as antecedent, as in Fig. 6.4. In both subcases identified above, the contrapositioned tree τ' will have $\neg\psi$ as the conclusion. In the first, $\neg\beta = \neg\mathbf{false} = \mathbf{true}$, so $\neg\beta$ is simply omitted as an antecedent of τ' . In the second, $\neg\beta = \neg\varphi$, which is exactly the single negative sentence which must be added as an antecedent of the entire proof. It cannot be the case that there are multiple occurrences of ψ in the original tree τ . If there were, the resulting tree would exhibit a proof of $\neg\psi$ from a set of premises which includes ψ . Since all of the rules are sound, this is impossible.

From this, it follows that $\text{Contraposn}\langle \tau: \langle \psi, d_1 \rangle \rangle$ is a proof of ψ from Φ in $\text{SwapCl}\langle \mathcal{A} \rangle$, so the latter is sat-complete. The single-use property follows from the argument of the previous paragraph. \square

To obtain a system which is complete even for unsatisfiable formulas of $\text{All}^\pm(\mathcal{C})$, the notion of complement-pair rules is introduced.

6.16 Complement-pair rules *A complement-pair rule is one which asserts that a WFF and its negation cannot both be true. A generic such rule is illustrated in (6.16-1).*

$$\frac{\alpha \quad \neg\alpha}{\mathbf{false}} \quad (6.16-1)$$

The *complement-pair set* of \mathcal{C} , denoted $\text{CompPrSet}\langle\mathcal{C}\rangle$, includes, for each $\alpha \in \mathcal{C}$, a complement-pair rule of the form shown (6.16-1). In the case that the rules of \mathcal{A} are represented generically using variables, it is only necessary to add rules which instantiate to all complement-pair rules. For the main example $\text{BinConstr}^+\langle\mathfrak{G}\rangle$, $\text{CompPrSet}\langle\text{BinConstr}^+\langle\mathfrak{G}\rangle\rangle$ is given by the two rules shown in Fig. 6.7.

$$(C1) \frac{(\mathfrak{g}_1 \sqsubseteq \mathfrak{g}_2) \quad (\mathfrak{g}_1 \not\sqsubseteq \mathfrak{g}_2)}{\mathbf{false}} \qquad (C2) \frac{(\prod\{\mathfrak{g}_1, \mathfrak{g}_2\} = \perp) \quad (\prod\{\mathfrak{g}_1, \mathfrak{g}_2\} \neq \perp)}{\mathbf{false}}$$

Figure 6.7: Complement-pair rules for $\text{BinConstr}^+\langle\mathfrak{G}\rangle$

The *full swap closure* of \mathcal{A} , denoted $\text{FullSwapCl}\langle\mathcal{A}\rangle$, is $\text{SwapCl}\langle\mathcal{A}\rangle \cup \text{CompPrSet}\langle\mathcal{C}\rangle$. Thus, $\text{FullSwapCl}\langle\text{BinInfPos}\langle\mathfrak{G}\rangle\rangle = \text{SwapCl}\langle\text{BinInfPos}\langle\mathfrak{G}\rangle\rangle \cup \text{CompPrSet}\langle\text{BinConstr}^+\langle\mathfrak{G}\rangle\rangle = \{(\text{I2}), (\text{M1}), (\text{I1}), (\text{T1}), (\text{T2}), (\text{T3}), (\text{U1}), (\text{I2-sa}), (\text{I2-sb}), (\text{M1-sa}), (\text{M1-sb}), (\text{U1-s}), (\text{C1}), (\text{C2})\}$.

The justification for this terminology is that a complement-pair rule may be regarded as the swap of an identity rule, as illustrated in (6.16-2). The middle inference rule is equivalent to the leftmost, adding only a **true** premise, while the rightmost rule results from the swap of the antecedent **true** and the consequent α .

$$\frac{\alpha}{\alpha} \quad \equiv \quad \frac{\alpha \quad \mathbf{true}}{\alpha} \quad \overset{\text{swap}}{\rightsquigarrow} \quad \frac{\alpha \quad \neg\alpha}{\mathbf{false}} \quad (6.16-2)$$

Thus, if identity rules, such as the leftmost rule of (6.16-2), were added to \mathcal{A} , then the complement-pair rules would be included in the swap closure.

6.17 Lemma — Characterization of unsatisfiability in $\text{All}^\pm\langle\mathcal{C}\rangle$ *Let $\Phi \subseteq \text{All}^\pm\langle\mathcal{C}\rangle$. If Φ is unsatisfiable, then either $\text{Only}^+\langle\Phi\rangle$ is unsatisfiable or else there is a $\neg\varphi \in \text{Only}^-\langle\Phi\rangle$ with the property that $\text{Only}^+\langle\Phi\rangle \models \varphi$.*

PROOF: If $\text{Only}^+\langle\Phi\rangle$ is unsatisfiable, then applying 5.11(b) with $\beta \equiv \mathbf{true}$ (so $\neg\beta \equiv \mathbf{false}$), it follows that either $\text{Only}^+\langle\Phi\rangle \models \mathbf{false}$; *i.e.*, $\text{Only}^+\langle\Phi\rangle$ is unsatisfiable, or else there is a $\neg\varphi \in \text{Only}^-\langle\Phi\rangle$ with the property that $\text{Only}^+\langle\Phi\rangle \cup \{\neg\varphi\} \models \mathbf{false}$; *i.e.*, that $\text{Only}^+\langle\Phi\rangle \cup \{\neg\varphi\}$ is unsatisfiable. From this, it follows from 5.8(a') that $\text{Only}^+\langle\Phi\rangle \models \varphi$, as required. \square

6.18 Theorem — Completeness of $\text{FullSwapCl}\langle\mathcal{A}\rangle$ *If \mathcal{A} is complete, then the proof system $\text{FullSwapCl}\langle\mathcal{A}\rangle$ is both sound and complete.*

PROOF: Assume that \mathcal{A} is complete. Soundness and sat-completeness of $\text{FullSwapCl}\langle\mathcal{A}\rangle$ follow from 6.15. Indeed, the rules of $\text{CompPrSet}\langle\mathcal{A}\rangle$ are clearly sound, and adding them to $\text{SwapCl}\langle\mathcal{A}\rangle$ has no effect on sat-completeness.

To show that completeness extends to unsatisfiable inferences, let $\Phi \subseteq \text{All}^\pm\langle\mathcal{C}\rangle$ be unsatisfiable. If $\text{Only}^+\langle\Phi\rangle$ is unsatisfiable, then this is already provable using only the rules of \mathcal{A} , since it is assumed to be complete. Trivially, a proof that $\text{Only}^+\langle\Phi\rangle$ is unsatisfiable is also a proof that all of Φ is unsatisfiable.

If $\text{Only}^+(\Phi)$ is satisfiable, then in view of 6.17, there is a $\neg\varphi \in \text{Only}^-(\Phi)$ with the property that $\text{Only}^+(\Phi) \models \varphi$. A proof tree establishing that Φ is unsatisfiable is shown in (6.18-1).

$$\begin{array}{c}
 \Sigma \\
 \vdots \\
 \text{proof using } \mathcal{A} \\
 \vdots \\
 \varphi \qquad \qquad \neg\varphi \\
 \hline
 \text{false}
 \end{array} \tag{6.18-1}$$

In that proof, Σ is a minimal subset of $\text{Only}^+(\Phi)$ for which $\Sigma \models \varphi$. The fact that \mathcal{A} is complete guarantees that the associated proof exists. Thus, $\Sigma \cup \{\neg\varphi\}$ is provably unsatisfiable, whence Φ has this property as well. \square

To close this section, two examples are provided which illustrate the process of proof using contrapositioning.

6.19 Corollary — Completeness of $\text{BinInf}(\mathfrak{G})$ *The inference system $\text{BinInf}(\mathfrak{G})$ is sound and complete.*

PROOF: Soundness follows from 6.6, while completeness follows from 4.31 and 6.18 \square

6.20 Example — Inference involving contrapositioning In this example, a proof which involves contrapositioning is presented. Let $\mathfrak{G}_{6.20}$ be an SMAS with $\{g_i \mid i \in [1, 6]\} \subseteq \text{Granules}(\mathfrak{G}_{6.20})$ and

$$\{(g_1 \sqsubseteq_{\mathfrak{G}} g_2), (g_2 \sqsubseteq_{\mathfrak{G}} g_3), (g_4 \sqsubseteq_{\mathfrak{G}} g_5), (g_5 \sqsubseteq_{\mathfrak{G}} g_6), (g_1 \not\sqsubseteq_{\mathfrak{G}} g_6)\} \subseteq \text{BinConstr}^{\pm}(\mathfrak{G}_{6.20}).$$

It will be shown that $\text{BinConstr}^{\pm}(\mathfrak{G}_{6.20}) \models (g_3 \not\sqsubseteq_{\mathfrak{G}} g_4)$ using the proof rules of $\text{BinInf}(\mathfrak{G}_{6.20})$. As the first step, it is necessary to identify the proof of positive assertions which will be contrapositioned. Using 5.8(a) with $\Phi \leftarrow \text{BinConstr}^{\pm}(\mathfrak{G}_{6.20}) \setminus \{(g_1 \not\sqsubseteq_{\mathfrak{G}} g_6)\}$, $\varphi \leftarrow (g_3 \sqsubseteq_{\mathfrak{G}} g_4)$, and $\psi \leftarrow (g_1 \sqsubseteq_{\mathfrak{G}} g_6)$, it suffices to show that

$$(\text{BinConstr}^{\pm}(\mathfrak{G}_{6.20}) \setminus \{(g_1 \not\sqsubseteq_{\mathfrak{G}} g_6)\}) \cup \{(g_3 \sqsubseteq_{\mathfrak{G}} g_4)\} \models (g_1 \sqsubseteq_{\mathfrak{G}} g_6).$$

The various trees which illustrate the proof process are shown in Fig. 6.8. The first tree, at the top of the figure, represents the proof of

$$(\text{BinConstr}^{\pm}(\mathfrak{G}_{6.20}) \setminus \{(g_1 \not\sqsubseteq_{\mathfrak{G}} g_6)\}) \cup \{(g_3 \sqsubseteq_{\mathfrak{G}} g_4)\} \models (g_1 \sqsubseteq_{\mathfrak{G}} g_6)$$

in standard proof-tree format. Below it, to the left, is this same tree in vertex-edge format, with the swap path highlighted using double-line arrows. This tree is the instantiation, for this example, of the generic tree of Fig. 6.4. To its right is the contrapositioning of this tree using fixed vertex positioning, corresponding to the generic tree of Fig. 6.5. The swap path is again highlighted using double-line arrows, while edges whose source vertex has been altered as a result of this transformation are show using dashed lines. Just below is the same tree in fixed-edge positioning, corresponding to Fig. 6.6, while at the bottom of the figure is the same result proof in standard inference format. In all cases, the the proof rules used have not been identified explicitly, but these are very easy to determine.




Figure 6.8: Proof trees for 6.20

6.21 Example — Inference using a complement-pair rule In this example, a proof of unsatisfiability which involves the use of a complement-pair rule is developed. Let $\mathfrak{G}_{6.21}$ be an SMAS with $\{g_i \mid i \in [1, 6]\} \subseteq \text{Granules}(\mathfrak{G}_{6.21})$ and $\{(g_1 \sqsubseteq_{\mathfrak{E}} g_2), (g_2 \sqsubseteq_{\mathfrak{E}} g_3), (g_4 \sqsubseteq_{\mathfrak{E}} g_5), (g_5 \sqsubseteq_{\mathfrak{E}} g_6), (\prod_{\mathfrak{E}} \{g_3, g_6\} = \perp_{\mathfrak{E}}), (\prod_{\mathfrak{E}} \{g_1, g_4\} \neq \perp_{\mathfrak{E}})\} \subseteq \text{BinConstr}^{\pm}(\mathfrak{G}_{6.21})$. It will be shown that $\text{BinConstr}^{\pm}(\mathfrak{G}_{6.21})$ is unsatisfiable, using the proof rules of $\text{BinInf}(\mathfrak{G}_{6.20})$.

The strategy is simple. First, develop a proof that $\text{BinConstr}^+(\mathfrak{G}_{6.21}) \setminus \{(\sqcap\{g_1, g_4\} \neq \perp)\} \models$




Figure 6.9: Proof trees for 6.21

$(\sqcap\{g_1, g_4\} = \perp)$, and then apply the complement-pair rule which states that both elements of the pair $\{(\sqcap\{g_1, g_4\} = \perp), (\sqcap\{g_1, g_4\} \neq \perp)\}$ cannot be true. The associated proof trees, in both vertex-edge format and traditional logic format, are shown in Fig. 6.9.

6.22 Relationship to RCC5 Given two granules $g_1, g_2 \in \text{Granules}(\mathfrak{G})$, the truth values of the three sentences $(g_1 \sqsubseteq g_2)$, $(g_2 \sqsubseteq g_1)$, and $(\sqcap\{g_1, g_2\} = \perp)$ together represent what is known about their relationship. To formalize this, for $\varphi \in \text{BinConstr}^+(\mathfrak{G})$, define

$$\text{State}_{\mathfrak{G}}\langle\varphi\rangle = \begin{cases} \text{true} & \text{if } \text{Constr}(\mathfrak{G}) \models \varphi, \\ \text{false} & \text{if } \text{Constr}(\mathfrak{G}) \models \neg\varphi, \\ \text{unknown} & \text{otherwise,} \end{cases}$$

and then define the *state vector of* $\langle g_1, g_2 \rangle$ in \mathfrak{G} to be the triple

$$\text{State}_{\mathfrak{G}}\langle g_1, g_2 \rangle = \langle \text{State}_{\mathfrak{G}}\langle (g_1 \sqsubseteq g_2) \rangle, \text{State}_{\mathfrak{G}}\langle (g_2 \sqsubseteq g_1) \rangle, \text{State}_{\mathfrak{G}}\langle (\sqcap\{g_1, g_2\} = \perp) \rangle \rangle.$$

Call $\text{State}_{\mathfrak{G}}\langle g_1, g_2 \rangle$ *complete* if no entry has the value **unknown**.

The *region connection calculus*, or *RCC*, and its variants, have complete state vectors, rather than individual predicates, as their starting point. The papers [RC89], [RCC92], [CBGG97], [LY03], and [Ste04] form a small sample of the vast work in this area. The most commonly investigated variant is RCC8, whose associated state vectors have four entries, since boundaries of regions are also modelled. For a comparison to the work of this paper, the more appropriate variant is RCC5, which focuses on regions without boundaries, so the state vector has only three entries. Table 6.1 shows the state vectors for the version of

RCC5 described in [Ste04, Table 1], which is particularly appropriate for comparison to the granule logic of this paper, since it supports empty granules, unlike most variants of RCC.² To avoid any confusion, the variant of RCC described in Table 6.1 will be called *RCC5+*.

	$(g_1 \sqsubseteq g_2)$	$(g_2 \sqsubseteq g_1)$	$(\prod\{g_2, g_1\} = \perp)$	Symmetric
$DC\langle g_1, g_2 \rangle$	false	false	true	✓
$PO\langle g_1, g_2 \rangle$	false	false	false	✓
$EQ\langle g_1, g_2 \rangle$	true	true	false	✓
$PP\langle g_1, g_2 \rangle$	true	false	false	
$PP^{-1}\langle g_1, g_2 \rangle$	false	true	false	
$EQ_{\emptyset}\langle g_1, g_2 \rangle$	true	true	true	✓
$PP_{\emptyset}\langle g_1, g_2 \rangle$	true	false	true	
$PP_{\emptyset}^{-1}\langle g_1, g_2 \rangle$	false	true	true	

Table 6.1: Predicates of RCC5+

The predicate names which are used are those which have become standard in the field. DC stands for *disconnected* (i.e., disjoint), PO for *partial overlap*, EQ for *equal*, PP for *proper part*, and PP^{-1} for *proper part inverse*. The relations DC, PO, and EQ are *symmetric* in that they are their own inverses. On the other hand PP is not symmetric; hence the need for PP^{-1} . These five relations constitute ordinary RCC5, and are fine as long as the regions are nonempty. However, they fail on empty regions. The assertions $EQ\langle \perp_{\mathfrak{G}}, \perp_{\mathfrak{G}} \rangle$, $PP\langle \perp_{\mathfrak{G}}, g_2 \rangle$, and $PP^{-1}\langle \perp_{\mathfrak{G}}, g_2 \rangle$ all fail to hold, the latter two for any $g \in \text{Granules}_{\neq \perp}(\mathfrak{G})$, counter to the expected meaning of these predicates. The solution is to introduce three additional relation names. The statements $EQ\langle g_1, g_2 \rangle$, $PP\langle g_1, g_2 \rangle$, and $PP^{-1}\langle g_1, g_2 \rangle$ cover the cases for which $g_1 \stackrel{id}{\neq} \perp_{\mathfrak{G}}$, while $EQ_{\emptyset}\langle g_1, g_2 \rangle$, $PP_{\emptyset}\langle g_1, g_2 \rangle$, and $PP_{\emptyset}^{-1}\langle g_1, g_2 \rangle$ cover the cases for which $g_1 \stackrel{id}{=} \perp_{\mathfrak{G}}$.

A key feature of this framework is that for any pair $\langle g_1, g_2 \rangle$ of granules, $R\langle g_1, g_2 \rangle$ holds for exactly one $R \in \{DC, PO, EQ, PP, PP^{-1}, EQ_{\emptyset}, PP_{\emptyset}, PP_{\emptyset}^{-1}\}$. This can be seen from the fact that every possible state vector is represented by one of the predicates. This property is sometimes called *JEPD*, short for *jointly exclusive and pairwise disjoint*.

Inference within RCC is often performed via the use of a *composition table* [LY03]. Such a table presents, for $R_1, R_2 \in \{DC, PO, EQ, PP, PP^{-1}, EQ_{\emptyset}, PP_{\emptyset}, PP_{\emptyset}^{-1}\}$, the relations $R_3 \in \{DC, PO, EQ, PP, PP^{-1}, EQ_{\emptyset}, PP_{\emptyset}, PP_{\emptyset}^{-1}\}$ for which $R_3 \subseteq R_1 \circ R_2$, with \circ denoting ordinary relational composition. Less formally, the table identifies the relations for which $R_3\langle g_1, g_3 \rangle$ could hold, given that $R_1\langle g_1, g_2 \rangle$ and $R_2\langle g_2, g_3 \rangle$ hold.

Presenting the entire composition table for RCC5+ is beyond the scope of this discussion. However, a simple example of inference is appropriate, in order to compare RCC5+ to the framework of this paper. Let $\mathfrak{G}_{6.22}$ have $\text{Granules}_{\neq \perp}(\mathfrak{G}_{6.22}) = \{g_1, g_2, g_3\}$, with

$$\text{Constr}(\mathfrak{G}) = \{(g_1 \not\sqsubseteq g_2), (g_2 \not\sqsubseteq g_1), (\prod\{g_1, g_2\} \neq \perp), (g_2 \sqsubseteq g_3), (g_3 \not\sqsubseteq g_2), (\prod\{g_2, g_3\} \neq \perp)\}.$$

²In [Ste04, Table 1], rather than $(\prod\{g_1, g_2\} = \perp)$, a predicate which is best described as $(g_1 \sqsubseteq \overline{g_2})$ is used for the third state parameter, with $\overline{g_2}$ the complement of g_2 . Since complements are not part of the framework of an SMAS, $(\prod\{g_1, g_2\} = \perp)$ is used instead. It is easy to verify that the results are identical.

It is immediate that these constraints correspond exactly to the RCC5+ conditions $\{\text{PO}\langle g_1, g_2 \rangle, \text{PP}\langle g_2, g_3 \rangle\}$. Even without constructing the entire composition table for RCC5+, it is straightforward to see that $\text{PO}\langle g_1, g_3 \rangle$ and $\text{PP}\langle g_1, g_3 \rangle$ are the only possibilities for the relationship on $\langle g_1, g_3 \rangle$. This can be done by looking at a composition table for RCC8, such as [LY03, Table 2], and collapsing TPP and NTPP into PP, or just by direct inspection, drawing a Venn diagram.

Now, consider applying the proof rules of $\text{BinInf}\langle \mathfrak{G}_{6.22} \rangle$. The only possible deductions are shown in Fig. 6.10. For both $(g_3 \sqsubseteq g_1)$ and $(\prod\{g_1, g_3\} = \perp)$ false, with the truth value

$$\frac{(g_2 \sqsubseteq g_3) \quad (g_2 \not\sqsubseteq g_1)}{(g_3 \not\sqsubseteq g_1)} \qquad \frac{(g_2 \sqsubseteq g_3) \quad (\prod\{g_1, g_2\} \neq \perp)}{(\prod\{g_1, g_3\} \neq \perp)}$$

Figure 6.10: Proofs for 6.22

of $(g_1 \sqsubseteq g_3)$ unknown, a check of Table 6.1 shows that the only possibilities are $\text{PO}\langle g_1, g_3 \rangle$ and $\text{PP}\langle g_1, g_3 \rangle$, in agreement with the result from a composition table.

Whether RCC is preferable to the framework of this paper depends entirely upon the type of knowledge to be modelled. If almost all relationships between granules are known completely, so that one of the eight predicates of Table 6.1 applies to each pair $\langle g_1, g_2 \rangle$, then RCC will likely be superior. However, in the context of substantial incomplete information, RCC will be overwhelmed by the need for disjunctive representation, rendering the approach of this paper the preferred one.

6.23 Relationship to AFDs In [DeB87] and [PDGV89, Sec. 5.2], a framework for dependencies on relational databases which includes both functional dependencies (FDs) and *afunctional dependencies* (AFDs) is developed, with a sound and complete inference system. Since FDs form the original Armstrong context [Arm74], it is worth asking how this work fits into the framework developed here.

AFDs are not true logical negations of FDs. The relation R satisfies the AFD $X \not\rightarrow Y$ if for **every** tuple $t \in R$, there is a second tuple $t' \in R$ with $t[X] = t'[X]$ and $t[Y] \neq t'[Y]$. On the other hand, the true logical negation of the FD $X \rightarrow Y$ has the property that for **some** tuple $t \in R$, there is a second tuple $t' \in R$ with $t[X] = t'[X]$ and $t[Y] \neq t'[Y]$. Thus, despite the utility of AFDs, it is not possible to define the combined inference system for FDs and AFDs by applying the swap approach of this paper to constraints on FDs.

6.24 Other contexts It should perhaps be mentioned that the techniques of this section can also be applied to the inference systems $\text{QBinInfPos}\langle \mathfrak{G} \rangle$ (see 4.3) and $\text{SQBinInfPos}\langle \mathfrak{G} \rangle$ (see 4.7), since their underlying frameworks (consisting of quasi-models and strong quasi-models, respectively), are easily shown to be Armstrong contexts. However, this direction will not be pursued here, since there is no immediate need for the result.

7 Relationship to Other Work

XXXXXX

8 Conclusions and Further Directions

In this paper, two main contributions have been made.

SINGLE-USE PROOF SYSTEM FOR BINARY GRANULE CONSTRAINTS: A sound and complete proof system for the predicates of subsumption and disjunction of granules, $\text{BinInfPos}\langle\mathfrak{G}\rangle$, has been developed. This result builds upon the earlier work in [AP88], adding critical new features, such as modelling of the bottom granule $\perp_{\mathfrak{G}}$ and top granule $\top_{\mathfrak{G}}$, and the requirement that all granules other than $\perp_{\mathfrak{G}}$ have nonempty semantics. It furthermore provides single-use proofs, which are necessary for extension to a proof system that involves the negations of these predicates as well.

PROOF SYSTEM AUGMENTATION TO NEGATIVE CONSTRAINTS A general method has been developed to extend any sound and complete single-use inference system over an Armstrong context \mathcal{C} to one which is sound and complete on the assertions in \mathcal{C} together with their negations. This method is then applied to the inference system $\text{BinInfPos}\langle\mathfrak{G}\rangle$ to obtain the sound and complete inference $\text{BinInf}\langle\mathfrak{G}\rangle$ system on all binary predicates of an SMAS, both positive and negative.

The immediate further direction is the following.

EXTEND TO GRANULES WITH BOUNDARY: In spatial modelling, it is often advantageous or even necessary to be able to work with granules with boundary. Work is currently underway to extend the notion of SMAS of this paper to such granules. This requires replacing the single disjunction predicate ($\sqcap\{g_1, g_2\} = \perp$), with two, the first ($\sqcap\{g_1^{\circ}, g_2^{\circ}\} = \perp$) for interiors and the second ($\sqcap\{g_1^{\partial}, g_2^{\partial}\} = \perp$) for boundaries. The resulting framework can be shown to be an Armstrong context, with a sound and complete single-use inference system which extends $\text{BinInfPos}\langle\mathfrak{G}\rangle$ naturally. The methods of Sec. 6 of this paper apply to this new framework as well, providing an inference system for both positive and negative constraints for binary predicates with boundary. The modelling power of this framework is equivalent to that of the four-intersection model [EF91] and RCC8 [Ste04].

References

- [AP86] Paolo Atzeni and D. Stott Parker Jr. Formal properties of net-based knowledge representation schemes. In *Proceedings of the Second International Conference on Data Engineering, February 5-7, 1986, Los Angeles, California, USA*, pages 700–706. IEEE Computer Society, 1986.

- [AP88] Paolo Atzeni and D. Stott Parker Jr. Formal properties of net-based knowledge representation schemes. *Data Knowl. Eng.*, 3:137–147, 1988.
- [Arm74] William Ward Armstrong. Dependency structures of data base relationships. In *IFIP Congress*, pages 580–583, 1974.
- [BA12] Mordechai Ben-Ari. *Mathematical Logic for Computer Science*. Springer, third edition, 2012.
- [CBGG97] Anthony G. Cohn, Brandon Bennett, John Gooday, and Nicholas Mark Gotts. Qualitative spatial representation and reasoning with the region connection calculus. *GeoInformatica*, 1(3):275–316, 1997.
- [DeB87] Paul De Bra. Functional dependency implications, inducing horizontal decompositions. In Joachim Biskup, János Demetrovics, Jan Paredaens, and Bernhard Thalheim, editors, *MFDBS 87, 1st Symposium on Mathematical Fundamentals of Database Systems, Dresden, GDR, January 19-23, 1987, Proceedings*, volume 305 of *Lecture Notes in Computer Science*, pages 80–98. Springer, 1987.
- [EF91] Max J. Egenhofer and Robert D. Franzosa. Point-set topological spatial relations. *Int. J. Geogr. Inf. Sci.*, 5(2):161–174, 1991.
- [End01] Herbert B. Enderton. *A Mathematical Introduction to Logic*. Harcourt, second edition, 2001.
- [Fag82] Ronald Fagin. Horn clauses and database dependencies. *J. Assoc. Comp. Mach.*, 29(4):952–985, 1982.
- [Gal15] Jean H. Gallier. *Logic for Computer Science*. Dover, second edition, 2015.
- [Gir95] Jean-Yves Girard. Linear logic: Its syntax and semantics. In Jean-Yves Girard, Yves Lafont, and Laurent Regnier, editors, *Advances in Linear Logic*, volume 222 of *London Mathematical Society Lecture Notes Series*, pages 1–42. Cambridge University Press, 1995.
- [HR17] Stephen J. Hegner and M. Andrea Rodríguez. A model for multigranular data and its integrity. *Informatica, Lith. Acad. Sci.*, 28:45–78, 2017.
- [HR18] Stephen J. Hegner and M. Andrea Rodríguez. Implicit representation of bi-granular rules for multigranular data. In Sven Hartmann, Hui Ma, Abdelkader Hameurlain, Günther Permul, and Roland R. Wagner, editors, *Database and Expert Systems Applications — 29th International Conference, DEXA 2018, Regensburg, Germany, September 3-6, 2018, Proceedings, Part I*, volume 11029 of *Lecture Notes in Computer Science*, pages 372–389. Springer, 2018.
- [LY03] Sanjiang Li and Mingsheng Ying. Region connection calculus: Its models and composition table. *Artif. Intell.*, 145(1-2):121–146, 2003.

- [Men97] Elliott Mendelson. *Introduction to Mathematical Logic*. Chapman and Hall, fourth edition, 1997.
- [Mon76] J. Donald Monk. *Mathematical Logic*. Springer-Verlag, 1976.
- [PDGV89] Jan Paredaens, Paul De Bra, Marc Gyssens, and Dirk Van Gucht. *The Structure of the Relational Database Model*. Springer-Verlag, 1989.
- [RC89] David A. Randell and Anthony G. Cohn. Modelling topological and metrical properties in physical processes. In Ronald J. Brachman, Hector J. Levesque, and Raymond Reiter, editors, *Proceedings of the 1st International Conference on Principles of Knowledge Representation and Reasoning (KR'89)*. Toronto, Canada, May 15-18 1989, pages 357–368. Morgan Kaufmann, 1989.
- [RCC92] David A. Randell, Zhan Cui, and Anthony G. Cohn. A spatial logic based on regions and connection. In Bernhard Nebel, Charles Rich, and William R. Swartout, editors, *Proceedings of the 3rd International Conference on Principles of Knowledge Representation and Reasoning (KR'92)*. Cambridge, MA, October 25-29, 1992., pages 165–176. Morgan Kaufmann, 1992.
- [Ste04] John G. Stell. Part and complement: Fundamental concepts in spatial relations. *Ann. Math. Artif. Intell.*, 41(1):1–17, 2004.