

# DIFFUSIVE LIMIT OF THE UNSTEADY NEUTRON TRANSPORT EQUATION IN BOUNDED DOMAINS

ZHIMENG OUYANG

ABSTRACT. The justification of hydrodynamic limits in non-convex domains has long been an open problem due to the singularity at the grazing set. In this paper, we investigate the unsteady neutron transport equation in a general bounded domain with the in-flow, diffuse-reflection, or specular-reflection boundary condition. Using a novel kernel estimate, we demonstrate the optimal  $L^2$  diffusive limit in the presence of both initial and boundary layers. Previously, this result was only proved for convex domains when the time variable is involved. Our approach is highly robust, making it applicable to all basic types of physical boundary conditions.

## CONTENTS

1.	Introduction	1
2.	Asymptotic Analysis	6
3.	Remainder Equation	11
4.	Remainder Estimate	13
5.	Diffusive Limit	16
6.	Diffuse-Reflection Boundary Problem	17
7.	Specular-Reflection Boundary Problem	19
	References	21

## 1. INTRODUCTION

The neutron transport equation is a fundamental model for studying particle dynamics in confined spaces, such as nuclear reactors and medical imaging devices (e.g., CT, MRI). Its investigation dates back to the dawn of the atomic age and has motivated significant developments in applied mathematics, such as the discontinuous Galerkin (DG) method [9, 7, 8] and the inverse transport problem [1, 2].

We consider the unsteady neutron transport equation in a three-dimensional smooth bounded domain  $\Omega$  without assuming any convexity. Denote the time variable  $t \in \mathbb{R}^+$ , the space variable  $x = (x_1, x_2, x_3) \in \Omega$ , and the velocity variable  $w = (w_1, w_2, w_3) \in \mathbb{S}^2$ . Let the neutron density  $u^\varepsilon(t, x, w)$  satisfy the in-flow boundary problem

$$\begin{cases} \varepsilon \partial_t u^\varepsilon + w \cdot \nabla_x u^\varepsilon + \varepsilon^{-1} (u^\varepsilon - \bar{u}^\varepsilon) = 0 & \text{in } \mathbb{R}^+ \times \Omega \times \mathbb{S}^2, \\ u^\varepsilon(0, x, w) = u_o(x, w) & \text{in } \Omega \times \mathbb{S}^2, \\ u^\varepsilon(t, x_0, w) = g(t, x_0, w) & \text{for } x_0 \in \partial\Omega \text{ and } w \cdot n(x_0) < 0. \end{cases} \quad (1.1)$$

Here,  $\bar{u}^\varepsilon$  is the velocity average of  $u^\varepsilon$ , i.e.,

$$\bar{u}^\varepsilon(t, x) = \frac{1}{4\pi} \int_{\mathbb{S}^2} u^\varepsilon(t, x, w) dw. \quad (1.2)$$

---

2020 *Mathematics Subject Classification*. Primary 35Q49, 82D75; Secondary 35Q62, 35Q20.

*Key words and phrases*. non-convex domain, boundary layer, initial layer, diffusive limit.

Z. Ouyang is supported by NSF Grant DMS-2202824.

Throughout the paper, we will use  $\bar{f}(t, x)$  to denote the velocity average of a function  $f(t, x, w)$ .  $u_o$  and  $g$  are the given initial data and boundary data, respectively, and  $n(x_0)$  is the outward unit normal vector at boundary point  $x_0 \in \partial\Omega$ . Our goal is to study the asymptotic behavior of the solution  $u^\varepsilon$  as the Knudsen number  $\varepsilon \rightarrow 0^+$ .

When the incoming particles are distributed like a probability average of the outgoing particles (possibly plus a small perturbation), then the boundary condition in (1.1) is replaced by

$$u^\varepsilon(t, x_0, w) = \mathcal{P}_\gamma[u^\varepsilon](t, x_0) + \varepsilon h(t, x_0, w) \quad \text{for } x_0 \in \partial\Omega \text{ and } w \cdot n(x_0) < 0, \quad (1.3)$$

where

$$\mathcal{P}_\gamma[u^\varepsilon](t, x_0) := c_\gamma \int_{w \cdot n > 0} u^\varepsilon(t, x_0, w)(w \cdot n) dw \quad (1.4)$$

normalized with the constant  $c_\gamma = \frac{1}{\pi}$  satisfying  $c_\gamma \int_{w \cdot n > 0} (w \cdot n) dw = 1$ . This is called the diffuse-reflection boundary condition.

In the specular-reflection boundary problem, where the particles bounce back specularly (with a small perturbation), the boundary condition in (1.1) is replaced by

$$u^\varepsilon(t, x_0, w) = u^\varepsilon(t, x_0, \mathcal{R}w) + \varepsilon h(t, x_0, w) \quad \text{for } x_0 \in \partial\Omega \text{ and } w \cdot n(x_0) < 0 \quad (1.5)$$

with  $\mathcal{R}w := w - 2(w \cdot n)n$ .

For both reflection type boundary conditions, we additionally assume the compatibility condition

$$\int_{w \cdot n < 0} h(t, x_0, w)(w \cdot n) dw = 0 \quad \text{for all } x_0 \in \partial\Omega \text{ and } t \in \mathbb{R}^+ \quad (1.6)$$

on the given perturbative boundary data  $h$ , so that the solution enjoys the null flux condition at the boundary:

$$\int_{\mathbb{S}^2} u^\varepsilon(t, x_0, w)(w \cdot n) dw = 0 \quad \text{for all } x_0 \in \partial\Omega \text{ and } t \in \mathbb{R}^+. \quad (1.7)$$

**1.1. Notation.** Based on the flow direction, we can divide the phase boundary  $\gamma := \partial\Omega \times \mathbb{S}^2$  into the incoming boundary  $\gamma_- := \{(x_0, w) \in \gamma : w \cdot n(x_0) < 0\}$ , the outgoing boundary  $\gamma_+ := \{(x_0, w) \in \gamma : w \cdot n(x_0) > 0\}$ , and the grazing set  $\gamma_0 := \{(x_0, w) \in \gamma : w \cdot n(x_0) = 0\}$ . In particular, the boundary conditions of (1.1), (1.3) and (1.5) are only prescribed on  $\gamma_-$ . Denote  $\Gamma := [0, T] \times \gamma$  and we may also split  $\Gamma = \Gamma_- \cup \Gamma_+ \cup \Gamma_0$  in a parallel manner.

Let  $\langle \cdot, \cdot \rangle_\sigma$  denote the inner product with respect to variable(s)  $\sigma$  which can be  $t, x, w$ , their combinations, or the domain specified, and let  $\langle\langle \cdot, \cdot \rangle\rangle$  be the inner product over  $[0, t] \times \Omega \times \mathbb{S}^2$  for  $t \in [0, T]$ , i.e.,

$$\langle\langle f, g \rangle\rangle := \int_0^t \iint_{\Omega \times \mathbb{S}^2} fg \, dw dx dt'. \quad (1.8)$$

Also, let  $\langle \cdot, \cdot \rangle_{\gamma_\pm}$  denote the inner product on  $\gamma_\pm$  with measure  $d\gamma := |w \cdot n| dw dS_x$ , where  $dS_x$  is the differential of the boundary surface. When the time variable is involved, we use  $\langle \cdot, \cdot \rangle_{\Gamma_\pm}$  to denote

$$\langle f, g \rangle_{\Gamma_\pm} := \int_0^t \iint_{\partial\Omega \times \mathbb{S}^2} fg |w \cdot n| dw dS_x dt'. \quad (1.9)$$

Define the bulk and boundary  $L^2$  norms

$$\|f\|_{L^2} := \left( \iint_{\Omega \times \mathbb{S}^2} |f(x, w)|^2 dw dx \right)^{\frac{1}{2}}, \quad |f|_{L^2_{\gamma_\pm}} := \left( \int_{\gamma_\pm} |f(x, w)|^2 d\gamma \right)^{\frac{1}{2}}, \quad (1.10)$$

and the  $L^\infty$  norms

$$\|f\|_{L^\infty} := \operatorname{ess\,sup}_{(x, w) \in \Omega \times \mathbb{S}^2} |f(x, w)|, \quad |f|_{L^\infty_{\gamma_\pm}} := \operatorname{ess\,sup}_{(x, w) \in \gamma_\pm} |f(x, w)|. \quad (1.11)$$

When the  $t$  variable is involved, we use  $\|f\|_{L^2}$ ,  $\|f\|_{L^2_{\Gamma_\pm}}$ ,  $\|f\|_{L^\infty}$  and  $\|f\|_{L^\infty_{\Gamma_\pm}}$  to denote the corresponding norms.

For  $k, m, l \in \mathbb{N}$ ,  $1 \leq p, q, r \leq \infty$ , let  $W^{k,p}$  be the Sobolev space, and let  $\|\cdot\|_{W_t^{l,r} W_x^{k,p} W_w^{m,q}}$  denote the usual Sobolev norm for  $t \in \mathbb{R}^+$ ,  $x \in \Omega$ , and  $w \in \mathbb{S}^2$ . If we use mixed norms on the boundary, then denote by  $|\cdot|_{W_x^{k,p} W_w^{m,q}(\gamma_\pm)}$  the mixed Sobolev norm for  $(x, w) \in \gamma_\pm$  with measure  $d\gamma = |w \cdot n| dw dS_x$ . The same

convention also applies to  $\|\cdot\|_{W_t^{l,r}W_x^{k,p}W_w^{m,q}(\Gamma_{\pm})}$  when  $t$  is involved. Sometimes we will omit the variables in the subscript when there is no possibility of confusion.

We will abuse notation for the functions under change of variables: for example, if  $\mathcal{T} : (t, x, w) \mapsto (\tau, y, v)$ , and let  $f(t, x, w) = f \circ \mathcal{T}^{-1}(\tau, y, v) =: \tilde{f}(\tau, y, v)$ , then we just write  $f(\tau, y, v)$  instead of  $\tilde{f}(\tau, y, v)$  for notational simplicity. However, unless otherwise specified, the integrals in the  $L^p$ -norms and the inner products are still with respect to the original variables  $t, x, w$ .

Throughout this paper,  $C > 0$  denotes a constant that only depends on the spatial domain  $\Omega$  or temporal domain  $[0, T]$  for some given  $T > 0$ , but does not depend on the data or  $\varepsilon$ . It is referred as universal constant and can change from one inequality to another. We write  $a \lesssim b$  to denote  $a \leq Cb$  and  $a \gtrsim b$  to denote  $a \geq Cb$ . Also, we write  $a \simeq b$  if  $a \lesssim b$  and  $a \gtrsim b$ . We will use  $0 < \delta \ll 1$  to denote a sufficiently small constant independent of the data or  $\varepsilon$ .

**1.2. Main Results.** Our main theorems establish the diffusive limit of the unsteady neutron transport equation in a bounded domain, under three different types of boundary conditions: in-flow, diffuse-reflection, and specular-reflection. As the assumption on data and the limit problem differ among the three cases, it is better to state our results separately:

**Theorem 1.1** (In-flow case). *Given  $u_o$  and  $g$  satisfying*

$$\|u_o\|_{W_x^{3,\infty}L_w^\infty} + \|g\|_{W_t^{2,\infty}W_x^{3,\infty}W_w^{1,\infty}(\Gamma_-)} \lesssim 1 \quad (1.12)$$

and the compatibility condition

$$u_o(x_0, w) = g(0, x_0, w) = C(x_0) \quad \text{for every } x_0 \in \partial\Omega, \quad (1.13)$$

there exists a unique solution  $u^\varepsilon(t, x, w) \in L^\infty([0, T] \times \Omega \times \mathbb{S}^2)$  to (1.1) for any  $T > 0$ . Moreover, the solution obeys the estimate

$$\|u^\varepsilon - U_0\|_{L^2([0, T] \times \Omega \times \mathbb{S}^2)} \lesssim \varepsilon^{\frac{1}{2}}, \quad (1.14)$$

where  $U_0(t, x, w) = \bar{U}_0(t, x)$  satisfies the heat equation with Dirichlet boundary condition:

$$\begin{cases} \partial_t \bar{U}_0 - \Delta_x \bar{U}_0 = 0 & \text{in } [0, T] \times \Omega, \\ \bar{U}_0(0, x) = \bar{u}_o(x) & \text{in } \Omega, \\ \bar{U}_0(t, x_0) = \Phi_{0,\infty}(t; \nu_1, \nu_2) & \text{on } [0, T] \times \partial\Omega, \end{cases} \quad (1.15)$$

in which  $\Phi_{0,\infty}$  is determined by solving the boundary layer equation (Milne problem) for  $\Phi_0$ :

$$\begin{cases} \sin \phi \frac{\partial \Phi_0}{\partial \eta} + \Phi_0 - \bar{\Phi}_0 = 0, \\ \Phi_0(t; 0, \nu_1, \nu_2; \phi, \psi) = g(t; \nu_1, \nu_2; \phi, \psi) \quad \text{for } \sin \phi > 0, \\ \lim_{\eta \rightarrow \infty} \Phi_0(t; \eta, \nu_1, \nu_2; \phi, \psi) = \Phi_{0,\infty}(t; \nu_1, \nu_2). \end{cases} \quad (1.16)$$

Here, the change of variables  $(x; w) \mapsto (\eta, \nu_1, \nu_2; \phi, \psi)$  near the boundary are defined in Section 2.3.

*Remark 1.1.* If we make the stronger assumption on the boundary data that for some  $\alpha > 0$

$$\|e^{\alpha t} g\|_{W_t^{2,\infty}W_x^{3,\infty}W_w^{1,\infty}(\Gamma_-)} \lesssim 1, \quad (1.17)$$

then based on Theorem 4.1, we have the approximation estimate with time decay for some  $0 < \beta < \alpha$

$$\|e^{\beta t} (u^\varepsilon - U_0)\|_{L^2(\mathbb{R}^+ \times \Omega \times \mathbb{S}^2)} \lesssim \varepsilon^{\frac{1}{2}}. \quad (1.18)$$

*Remark 1.2.* The estimate (1.14) achieves the optimal convergence rate in  $L_{t,x,w}^2$ . In [27, 28, 29] for 2D/3D convex domains, it is justified that

$$\|u^\varepsilon - \tilde{U}_0 - \tilde{U}_0^B - U_0^I\|_{L^2} \lesssim \varepsilon^{\frac{5}{6}-}, \quad (1.19)$$

where  $\tilde{U}_0^B$  is the boundary layer with geometric correction,  $U_0^I$  is the initial layer, and  $\tilde{U}_0$  is the corresponding interior solution. [26, Theorem 2.1] reveals that the two interior solutions differ by

$$\|\tilde{U}_0 - U_0\|_{L^2} \lesssim \varepsilon^{\frac{2}{3}}. \quad (1.20)$$

Due to the rescaling in the normal spatial variable  $\eta = \varepsilon^{-1}\mu$ , for general non-constant in-flow boundary data, the boundary layer  $\tilde{U}_0^B \neq 0$  satisfies

$$\|\tilde{U}_0^B\|_{L^2} \simeq \varepsilon^{\frac{1}{2}}. \quad (1.21)$$

Also, due to the rescaling in the temporal variable  $\tau = \varepsilon^{-2}t$ , we have

$$\|U_0^I\|_{L^2} \simeq \varepsilon. \quad (1.22)$$

Hence, we conclude that

$$\|u^\varepsilon - U_0\|_{L^2} \simeq \varepsilon^{\frac{1}{2}}. \quad (1.23)$$

This suggests the optimality of our diffusive approximation (1.14).

In the case of the diffuse and specular reflection boundary conditions, we have the following analogues:

**Theorem 1.2** (Diffuse-reflection case). *If we consider (1.1) with the boundary condition replaced by (1.3), then under the assumption*

$$\|u_o\|_{W_x^{3,\infty}L_w^\infty} + \|h\|_{W_t^{1,\infty}W_x^{2,\infty}L_w^\infty(\Gamma_-)} \lesssim 1 \quad (1.24)$$

with the compatibility conditions (1.6) and

$$u_o(x_0, w) = C(x_0), \quad h(0, x_0, w) = 0 \quad \text{for every } x_0 \in \partial\Omega, \quad (1.25)$$

there exists a unique solution  $u^\varepsilon(t, x, w) \in L^\infty([0, T] \times \Omega \times \mathbb{S}^2)$  for any  $T > 0$ . Moreover, the solution obeys the estimate

$$\|u^\varepsilon - U_0\|_{L^2([0, T] \times \Omega \times \mathbb{S}^2)} \lesssim \varepsilon^{\frac{1}{2}}, \quad (1.26)$$

where  $U_0(t, x, w) = \bar{U}_0(t, x)$  satisfies the heat equation with Neumann boundary condition:

$$\begin{cases} \partial_t \bar{U}_0 - \Delta_x \bar{U}_0 = 0 & \text{in } [0, T] \times \Omega, \\ \bar{U}_0(0, x) = \bar{u}_o(x) & \text{in } \Omega, \\ \frac{\partial \bar{U}_0}{\partial n}(t, x_0) = 0 & \text{on } [0, T] \times \partial\Omega. \end{cases} \quad (1.27)$$

**Theorem 1.3** (Specular-reflection case). *If we consider (1.1) with the boundary condition replaced by (1.5), then under the assumption*

$$\|u_o\|_{W_x^{3,\infty}L_w^\infty} + \|h\|_{W_t^{1,\infty}W_x^{2,\infty}W_w^{1,\infty}(\Gamma_-)} \lesssim 1 \quad (1.28)$$

with the compatibility conditions (1.6) and  $h|_{\Gamma_0} = 0$ , as well as

$$u_o(x_0, w) = C(x_0), \quad \nabla_x u_o(x_0, w) = 0, \quad h(0, x_0, w) = 0 \quad \text{for every } x_0 \in \partial\Omega, \quad (1.29)$$

there exists a unique solution  $u^\varepsilon(t, x, w) \in L^\infty([0, T] \times \Omega \times \mathbb{S}^2)$  for any  $T > 0$ . Moreover, the solution obeys the estimate

$$\|u^\varepsilon - U_0\|_{L^2([0, T] \times \Omega \times \mathbb{S}^2)} \lesssim \varepsilon^{\frac{1}{2}}, \quad (1.30)$$

where  $U_0(t, x, w) = \bar{U}_0(t, x)$  satisfies the heat equation with Neumann boundary condition:

$$\begin{cases} \partial_t \bar{U}_0 - \Delta_x \bar{U}_0 = 0 & \text{in } [0, T] \times \Omega, \\ \bar{U}_0(0, x) = \bar{u}_o(x) & \text{in } \Omega, \\ \frac{\partial \bar{U}_0}{\partial n}(t, x_0) = 0 & \text{on } [0, T] \times \partial\Omega. \end{cases} \quad (1.31)$$

*Remark 1.3.* To compare the three boundary problems: In all the three cases, we achieve the  $L^2$  diffusive limit with the same convergence rate

$$\|u^\varepsilon - U_0\|_{L^2([0, T] \times \Omega \times \mathbb{S}^2)} \lesssim \varepsilon^{\frac{1}{2}}, \quad (1.32)$$

and in a similar fashion as Remark 1.1, the corresponding time decay if assuming decay on the boundary data. For the in-flow case, the interior solution  $U_0$  satisfies the heat equation with Dirichlet boundary determined by boundary layers. For both diffuse and specular cases,  $U_0$  does not depend on boundary layers due to the vanishing leading-order boundary layer. However, the proof of the specular case still requires introducing a

next-order (artificial) “boundary layer”, in order to enforce the perfect specular boundary condition for the remainder (because we cannot control the boundary trace of remainder in the energy or kernel estimate); while the diffuse case allows small perturbation that can be controlled by  $(1 - \mathcal{P}_\gamma)[\mathcal{R}]$  in the energy, and thus no boundary layer is needed in the analysis.

**1.3. Overview of the Method.** The neutron transport equation has been investigated from various perspectives since the 1960s. For the physical modeling and formal asymptotic expansion, we refer to [15, 16, 17, 18, 19, 20, 21, 22, 23]. For theoretical analysis, we refer to [6, 5, 3, 4]. For recent developments on the diffusive limit and the half-space problem, we refer to [30, 29, 13, 24, 25, 26] and the references therein.

In bounded domains, the mismatch of boundary conditions for the asymptotic expansion calls for a kinetic correction – the so-called boundary layer (Knudsen layer). As far as we are aware of, there are two main approaches:

In flat domains (e.g. half space or  $\mathbb{T}^2 \times [0, 1]$ ), the boundary layer  $U_0^B(t; \eta, \iota_1, \iota_2; \phi, \psi)$  relies on the Milne problem (cf. [6, 5])

$$\sin \phi \frac{\partial U_0^B}{\partial \eta} + U_0^B - \overline{U_0^B} = 0. \quad (1.33)$$

This formulation is consistent with the intuitive derivation from the transport operator under change of coordinates and rescaling (see (2.26)).

Unfortunately, in curved domains, the non-vanishing curvature and geometric effects are non-negligible due to the grazing set singularity. A surprising counter-example is constructed in [30] so that

$$\lim_{\varepsilon \rightarrow 0} \left\| \|u^\varepsilon - U_0 - U_0^B - U_0^I\| \right\|_{L^\infty} \neq 0. \quad (1.34)$$

As [30, 29] pointed out, in curved convex domains, the boundary layer  $\tilde{U}_0^B(t; \eta, \iota_1, \iota_2; \phi, \psi)$  relies on the so-called  $\varepsilon$ -Milne problem with geometric correction:

$$\sin \phi \frac{\partial \tilde{U}_0^B}{\partial \eta} - \frac{\varepsilon}{1 - \varepsilon \eta} \cos \phi \frac{\partial \tilde{U}_0^B}{\partial \phi} + \tilde{U}_0^B - \overline{\tilde{U}_0^B} = 0, \quad (1.35)$$

where the extra correction term  $-\frac{\varepsilon}{1 - \varepsilon \eta} \cos \phi \frac{\partial \tilde{U}_0^B}{\partial \phi}$  helps provide the weighted  $W^{1, \infty}$  bound of  $\tilde{U}_0^B$  and eventually leads to the desired diffusive limit.

In both flat and convex domains, the proof relies on expanding the boundary layer to higher order which provides sufficient  $\varepsilon$  power to close the remainder estimate. However, in non-convex domains, as [31, 13] reveals, the Milne problem with or without geometric correction does not guarantee  $W^{1, \infty}$  regularity of the boundary layer. As a consequence, we cannot even bound the remainder induced by the leading-order boundary layer.

In this paper, we will employ a fresh approach to justify the  $L^2$  diffusive expansion in general bounded domains with a cutoff boundary layer and the novel remainder estimate. We intend to show that

$$\lim_{\varepsilon \rightarrow 0} \left\| \|u^\varepsilon - U_0 - U_0^B - U_0^I\| \right\|_{L^2} = 0 \quad (1.36)$$

for the leading-order interior solution  $U_0$ , boundary layer  $U_0^B$ , and initial layer  $U_0^I$ .

It is well-known that the classical bounds for the remainder  $\mathcal{R} := u^\varepsilon - U_0 - U_0^B - U_0^I$  read (see [27, 28, 29])

$$\|\mathcal{R}(t)\|_{L^2} + \varepsilon^{-\frac{1}{2}} \|\mathcal{R}\|_{L_{\Gamma_+}^2} + \varepsilon^{-1} \|\|\mathcal{R} - \overline{\mathcal{R}}\|\|_{L^2} \lesssim \delta \|\|\mathcal{R}\|\|_{L^2} + 1, \quad (1.37)$$

$$\|\|\mathcal{R}\|\|_{L^2} \lesssim \varepsilon^{-1} \|\|\mathcal{R} - \overline{\mathcal{R}}\|\|_{L^2} + \|\mathcal{R}\|_{L_{\Gamma_+}^2} + \varepsilon^{\frac{1}{2}} \|\mathcal{R}(t)\|_{L^2} + 1, \quad (1.38)$$

which yield

$$\|\mathcal{R}(t)\|_{L^2} + \varepsilon^{-\frac{1}{2}} \|\mathcal{R}\|_{L_{\Gamma_+}^2} + \varepsilon^{-1} \|\|\mathcal{R} - \overline{\mathcal{R}}\|\|_{L^2} + \|\|\mathcal{R}\|\|_{L^2} \lesssim 1. \quad (1.39)$$

Clearly, without expanding to higher-order boundary layers, this is insufficient for the validity of (1.36).

The bottleneck of (1.39) lies in the  $\overline{\mathcal{R}}$  bound (1.38). Initiated from [13] for the steady problem with the in-flow boundary, we design several delicate test functions in the kernel estimate to obtain

$$\|\|\mathcal{R}\|\|_{L^2} \lesssim \|\|\mathcal{R} - \overline{\mathcal{R}}\|\|_{L^2} + \|\mathcal{R}\|_{L_{\Gamma_+}^2} + \varepsilon^{\frac{1}{2}} \|\mathcal{R}(t)\|_{L^2} + \varepsilon^{\frac{1}{2}}, \quad (1.40)$$

which will lead to the desired remainder estimate

$$\|\mathcal{R}(t)\|_{L^2} + \varepsilon^{-\frac{1}{2}} \|\mathcal{R}\|_{L^2_{\Gamma_+}} + \varepsilon^{-1} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2} + \varepsilon^{-\frac{1}{2}} \|\overline{\mathcal{R}}\|_{L^2} \lesssim 1. \quad (1.41)$$

The key to achieve (1.40) is a tricky combination of three conservation laws. We introduce the auxiliary function  $\xi(t, x)$  satisfying  $-\Delta_x \xi = \overline{\mathcal{R}}$  and  $\xi|_{\partial\Omega} = 0$ . For the remainder equation

$$\varepsilon \partial_t \mathcal{R} + w \cdot \nabla_x \mathcal{R} + \varepsilon^{-1} (\mathcal{R} - \overline{\mathcal{R}}) = \mathcal{S}, \quad (1.42)$$

testing against  $\varepsilon^{-1} \xi$  yields

$$\langle \partial_t \overline{\mathcal{R}}, \xi \rangle - \varepsilon^{-1} \langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x \xi \rangle = \varepsilon^{-1} \langle \mathcal{S}, \xi \rangle, \quad (1.43)$$

the choice of test function  $w \cdot \nabla_x \xi$  yields

$$\begin{aligned} \varepsilon \langle \partial_t \mathcal{R}, w \cdot \nabla_x \xi \rangle + \langle \mathcal{R}, w \cdot \nabla_x \xi \rangle_{\Gamma_+} - \langle \mathcal{G}, w \cdot \nabla_x \xi \rangle_{\Gamma_-} \\ - \langle \mathcal{R}, w \cdot \nabla_x (w \cdot \nabla_x \xi) \rangle + \varepsilon^{-1} \langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x \xi \rangle = \langle \mathcal{S}, w \cdot \nabla_x \xi \rangle, \end{aligned} \quad (1.44)$$

while the test function  $\varepsilon \partial_t \xi$  leads to

$$\varepsilon^2 \langle \partial_t \overline{\mathcal{R}}, \partial_t \xi \rangle - \varepsilon \langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x \partial_t \xi \rangle = \varepsilon \langle \mathcal{S}, \partial_t \xi \rangle. \quad (1.45)$$

On the one hand, the linear combination of (1.43) and (1.44) crucially cancels  $\varepsilon^{-1} \langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x \xi \rangle$  so that it kills the worst contribution of  $\varepsilon^{-1} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}$ , and also provides the control of  $-\langle \overline{\mathcal{R}}, w \cdot \nabla_x (w \cdot \nabla_x \xi) \rangle \simeq \|\overline{\mathcal{R}}\|_{L^2}^2$ . On the other hand, the estimate of  $\varepsilon \langle \partial_t \mathcal{R}, w \cdot \nabla_x \xi \rangle \sim \varepsilon \langle \mathcal{R}, w \cdot \nabla_x \partial_t \xi \rangle$  (up to some good terms) calls for the control of  $\partial_t \nabla_x \xi$ , which is in turn provided by (1.45) as  $\varepsilon^2 \langle \partial_t \overline{\mathcal{R}}, \partial_t \xi \rangle = \varepsilon^2 \|\nabla_x \partial_t \xi\|_{L^2}^2$ .

Combined with the key favorable sign of  $\langle \partial_t \overline{\mathcal{R}}, \xi \rangle \sim \|\nabla_x \xi(t)\|_{L^2}^2$  and a careful analysis of the source terms  $\varepsilon^{-1} \langle \mathcal{S}, \xi \rangle$ ,  $\langle \mathcal{S}, w \cdot \nabla_x \xi \rangle$ , and  $\varepsilon \langle \mathcal{S}, \partial_t \xi \rangle$  using Hardy's inequality and delicately chosen norms, we are able to create an extra gain of  $\varepsilon^{\frac{1}{2}}$  for the kernel bound. We then conclude the remainder estimate without any further expansion of higher-order boundary layers.

The remaining of this paper is structured as follows: Section 2 focuses on the asymptotic expansion for the interior solution, boundary and initial layers. Section 3 presents the setup of the remainder equation, and Section 4 elaborates the estimates of  $\mathcal{R} - \overline{\mathcal{R}}$  and  $\overline{\mathcal{R}}$ . Finally, the main theorem for the inflow case is proved in Section 5, while the diffuse and specular boundary problems are discussed in Section 6 and Section 7, respectively.

## 2. ASYMPTOTIC ANALYSIS

We seek a solution to (1.1) in the form of

$$u^\varepsilon = U + U^I + U^B + \mathcal{R} = (U_0 + \varepsilon U_1 + \varepsilon^2 U_2) + (U_0^I + \varepsilon U_1^I) + U_0^B + \mathcal{R}, \quad (2.1)$$

where the interior solution

$$U(t, x, w) := U_0(t, x, w) + \varepsilon U_1(t, x, w) + \varepsilon^2 U_2(t, x, w), \quad (2.2)$$

the initial layer

$$U^I(\tau, x, w) := U_0^I(\tau, x, w) + \varepsilon U_1^I(\tau, x, w), \quad (2.3)$$

and the boundary layer

$$U^B(t; \eta, \iota_1, \iota_2; \phi, \psi) := U_0^B(t; \eta, \iota_1, \iota_2; \phi, \psi). \quad (2.4)$$

Here,  $U_0$ ,  $U_1$ ,  $U_2$ ,  $U_0^I$ ,  $U_1^I$ , and  $U_0^B$  will be constructed in the following subsections, and  $\mathcal{R}(t, x, w)$  is the remainder.

**2.1. Interior Solution.** Inserting the asymptotic expansion ansatz  $u^\varepsilon \sim \sum_{k=0}^{\infty} U_k \varepsilon^k$  into (1.1), we get a hierarchy of equations by looking at each order of  $\varepsilon$ . Following the analysis in [29], we deduce that

$$U_0 = \bar{U}_0, \quad \partial_t \bar{U}_0 - \Delta_x \bar{U}_0 = 0, \quad (2.5)$$

$$U_1 = \bar{U}_1 - w \cdot \nabla_x U_0, \quad \partial_t \bar{U}_1 - \Delta_x \bar{U}_1 = 0, \quad (2.6)$$

$$U_2 = \bar{U}_2 - w \cdot \nabla_x U_1 - \partial_t U_0, \quad \partial_t \bar{U}_2 - \Delta_x \bar{U}_2 = 0. \quad (2.7)$$

The initial and boundary conditions for  $\bar{U}_0$ ,  $\bar{U}_1$ , and  $\bar{U}_2$  are determined by the initial and boundary layers.

**2.2. Initial Layer.** Based on the principle of dominant balance, we find the correct scaling for the time variable is  $\tau = \varepsilon^{-2}t$ , and so  $\partial_t = \varepsilon^{-2}\partial_\tau$ . Under the substitution  $t \mapsto \tau$ , the rescaled equation of (1.1) for the initial layer reads

$$\partial_\tau U^I + \varepsilon w \cdot \nabla_x U^I + (U^I - \bar{U}^I) = 0. \quad (2.8)$$

Inserting the expansion  $U^I \sim \sum_{k=0}^{\infty} U_k^I \varepsilon^k$  into (2.8) and comparing the order of  $\varepsilon$ , we find that

$$\partial_\tau U_0^I + U_0^I - \bar{U}_0^I = 0, \quad (2.9)$$

$$\partial_\tau U_1^I + U_1^I - \bar{U}_1^I = -w \cdot \nabla_x U_0^I. \quad (2.10)$$

Let us consider the general initial layer problem

$$\begin{cases} \frac{d\Theta}{d\tau} + \Theta - \bar{\Theta} = S, \\ \Theta(0, x, w) = \Theta_o(x, w). \end{cases} \quad (2.11)$$

Here,  $S(\tau, x, w)$  is the forcing term, and  $\Theta_o(x, w)$  is the given initial data. We are interested in the solution  $\Theta(\tau, x, w)$  that satisfies

$$\lim_{\tau \rightarrow \infty} \Theta(\tau, x, w) = \Theta_\infty(x) \quad (2.12)$$

for some  $\Theta_\infty(x)$  that does not depend on  $w$ .

The following proposition guarantees the solvability and regularity of the above initial layer problem:

**Proposition 2.1** (Initial layer problem). *Let  $k \in \mathbb{N}$ . Assume*

$$\|\Theta_o\|_{W_x^{k,\infty} L_w^\infty} + \|e^{\alpha\tau} S\|_{L_\tau^\infty W_x^{k,\infty} L_w^\infty} \lesssim 1 \quad (2.13)$$

for some  $\alpha > 0$ . Then there exist a unique solution  $\Theta(\tau, x, w) \in L_\tau^\infty W_x^{k,\infty} L_w^\infty$  to (2.11) and a  $\Theta_\infty(x) \in W_x^{k,\infty}$  such that (2.12) holds in the sense of

$$\|\Theta_\infty\|_{W_x^{k,\infty}} + \|e^{\beta\tau}(\Theta - \Theta_\infty)\|_{L_\tau^\infty W_x^{k,\infty} L_w^\infty} \lesssim 1 \quad (2.14)$$

for any  $0 < \beta \leq \min\{1, \alpha\}$ .

*Proof.* Based on ODE theory, the initial value problem (2.11) admits a unique solution. We first integrate the equation in  $w$  to find  $\partial_\tau \bar{\Theta} = \bar{S}$  and so  $\bar{\Theta} = \bar{\Theta}_o + \int_0^\tau \bar{S}$ . Then subtracting this from (2.11), we get

$$\begin{cases} \frac{d}{d\tau}(\Theta - \bar{\Theta}) + (\Theta - \bar{\Theta}) = S - \bar{S}, \\ (\Theta - \bar{\Theta})|_{\tau=0} = \Theta_o - \bar{\Theta}_o, \end{cases} \quad (2.15)$$

which can be solved using the integrating factor  $e^\tau$ . We have

$$\Theta - \bar{\Theta} = e^{-\tau}(\Theta_o - \bar{\Theta}_o) + \int_0^\tau e^{\tau'-\tau}(S - \bar{S}) d\tau', \quad (2.16)$$

and thus

$$\Theta = \bar{\Theta}_o + e^{-\tau}(\Theta_o - \bar{\Theta}_o) + \int_0^\tau \left\{ \bar{S} + e^{\tau'-\tau}(S - \bar{S}) \right\} d\tau'. \quad (2.17)$$

Let

$$\Theta_\infty := \overline{\Theta}_\infty = \lim_{\tau \rightarrow \infty} \overline{\Theta} = \overline{\Theta}_o + \int_0^\infty \overline{S} \, d\tau. \quad (2.18)$$

Then it is straightforward to verify (2.14).  $\square$

Let  $\Theta_0$  be the solution to (2.11) with  $S = 0$  and the initial data  $\Theta_{0,o} = u_o$  satisfying (1.12). Then by (2.18) we have  $\Theta_{0,\infty} = \overline{u}_o$ . Define the leading-order initial layer to be

$$U_0^I(\tau, x, w) := \Theta_0(\tau, x, w) - \Theta_{0,\infty}(x), \quad (2.19)$$

so that it satisfies

$$\begin{cases} \partial_\tau U_0^I + U_0^I - \overline{U}_0^I = 0, \\ U_0^I(0, x, w) = u_o(x, w) - \overline{u}_o(x), \\ \lim_{\tau \rightarrow \infty} U_0^I(\tau, x, w) = 0, \end{cases} \quad (2.20)$$

as well as

$$\|e^\tau U_0^I\|_{L_\tau^\infty W_x^{3,\infty} L_w^\infty} \lesssim 1. \quad (2.21)$$

Next, we define the second-order initial layer to be

$$U_1^I(\tau, x, w) := \Theta_1(\tau, x, w) - \Theta_{1,\infty}(x), \quad (2.22)$$

where  $\Theta_1$  is determined by solving (2.11) with  $S = -w \cdot \nabla_x U_0^I$  and  $\Theta_{1,o} = w \cdot \nabla_x U_0(0, x, w)$ , and so  $\Theta_{1,\infty}(x) = \lim_{\tau \rightarrow \infty} \overline{\Theta}_1(\tau, x)$  whose explicit formula is given by (2.18). Then  $U_1^I$  satisfies

$$\begin{cases} \partial_\tau U_1^I + U_1^I - \overline{U}_1^I = -w \cdot \nabla_x U_0^I, \\ U_1^I(0, x, w) = w \cdot \nabla_x U_0(0, x, w) - \Theta_{1,\infty}(x), \\ \lim_{\tau \rightarrow \infty} U_1^I(\tau, x, w) = 0, \end{cases} \quad (2.23)$$

as well as

$$\|\Theta_{1,\infty}\|_{W_x^{2,\infty}} + \|e^\tau U_1^I\|_{L_\tau^\infty W_x^{2,\infty} L_w^\infty} \lesssim 1. \quad (2.24)$$

**2.3. Boundary Layer.** We first perform change of coordinates depending on the geometry of the boundary. For the smooth surface  $\partial\Omega$ , there exists an orthogonal curvilinear coordinates system  $(\iota_1, \iota_2)$  such that the coordinate lines coincide with the principal directions at any  $x_0 \in \partial\Omega$ . Assume that  $\partial\Omega$  is parameterized by  $\mathbf{r} = \mathbf{r}(\iota_1, \iota_2)$ , and let  $\varsigma_i := L_i^{-1} \partial_{\iota_i} \mathbf{r}$  ( $i = 1, 2$ ) be the two orthogonal unit tangential vectors, where  $L_i := |\partial_{\iota_i} \mathbf{r}|$ . Then  $\{\varsigma_1, \varsigma_2, n\}$  form an orthonormal basis at each point in a neighborhood of the boundary. Hence, we can define a new coordinate system  $(\mu, \iota_1, \iota_2)$  near the boundary, where  $\mu$  denotes the normal distance to the boundary surface  $\partial\Omega$  so that  $x = \mathbf{r} - \mu n$ . We then define the rescaled normal spatial variable  $\eta = \varepsilon^{-1} \mu$  again due to the principle of dominant balance, and so  $\partial_\mu = \varepsilon^{-1} \partial_\eta$ .

Define also the spherical substitution  $w \mapsto (\phi, \psi)$  for the velocity  $w \in \mathbb{S}^2$  via

$$-w \cdot n = \sin \phi, \quad w \cdot \varsigma_1 = \cos \phi \sin \psi, \quad w \cdot \varsigma_2 = \cos \phi \cos \psi, \quad \text{for } \phi \in [-\frac{\pi}{2}, \frac{\pi}{2}], \quad \psi \in [0, 2\pi). \quad (2.25)$$

Under the change of variables  $(x, w) \mapsto (\eta, \iota_1, \iota_2; \phi, \psi)$  in the phase space, the transport operator becomes

$$\begin{aligned} \varepsilon \partial_t + w \cdot \nabla_x &= \varepsilon \partial_t + \varepsilon^{-1} \sin \phi \frac{\partial}{\partial \eta} + \frac{R_1 \cos \phi \sin \psi}{L_1(R_1 - \varepsilon \eta)} \frac{\partial}{\partial \iota_1} + \frac{R_2 \cos \phi \cos \psi}{L_2(R_2 - \varepsilon \eta)} \frac{\partial}{\partial \iota_2} \\ &\quad - \left( \frac{\sin^2 \psi}{R_1 - \varepsilon \eta} + \frac{\cos^2 \psi}{R_2 - \varepsilon \eta} \right) \cos \phi \frac{\partial}{\partial \phi} \\ &\quad + \frac{\sin \psi}{R_1 - \varepsilon \eta} \left\{ \frac{R_1 \cos \phi}{L_1 L_2} \left[ \varsigma_1 \cdot \left( \varsigma_2 \times (\partial_{\iota_1 \iota_2} \mathbf{r} \times \varsigma_2) \right) \right] - \sin \phi \cos \psi \right\} \frac{\partial}{\partial \psi} \\ &\quad - \frac{\cos \psi}{R_2 - \varepsilon \eta} \left\{ \frac{R_2 \cos \phi}{L_1 L_2} \left[ \varsigma_2 \cdot \left( \varsigma_1 \times (\partial_{\iota_1 \iota_2} \mathbf{r} \times \varsigma_1) \right) \right] - \sin \phi \sin \psi \right\} \frac{\partial}{\partial \psi}, \end{aligned} \quad (2.26)$$

where  $R_i$  ( $i = 1, 2$ ) is the radius of curvature.



Inserting the expansion  $U^B \sim \sum_{k=0}^{\infty} U_k^B \varepsilon^k$  into the rescaled equation and comparing the order of  $\varepsilon$  (see (2.26)), we expect that the standard leading-order boundary layer  $U_0^B(t; \eta, \iota_1, \iota_2; \phi, \psi)$  should satisfy

$$\sin \phi \frac{\partial U_0^B}{\partial \eta} + U_0^B - \overline{U_0^B} = 0. \quad (2.27)$$

Consider the general boundary layer problem (Milne problem)

$$\sin \phi \frac{\partial \Phi}{\partial \eta} + \Phi - \overline{\Phi} = S, \quad \overline{\Phi}(t; \eta, \iota_1, \iota_2) := \frac{1}{4\pi} \int_{-\pi}^{\pi} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \Phi(t; \eta, \iota_1, \iota_2; \phi, \psi) \cos \phi \, d\phi \, d\psi, \quad (2.28)$$

with boundary condition

$$\Phi(t; 0, \iota_1, \iota_2; \phi, \psi) = \rho(t; \iota_1, \iota_2; \phi, \psi) \quad \text{for } \sin \phi > 0. \quad (2.29)$$

We are interested in the solution  $\Phi(t; \eta, \iota_1, \iota_2; \phi, \psi)$  that satisfies

$$\lim_{\eta \rightarrow \infty} \Phi(t; \eta, \iota_1, \iota_2; \phi, \psi) = \Phi_{\infty}(t; \iota_1, \iota_2) \quad (2.30)$$

for some  $\Phi_{\infty}(t; \iota_1, \iota_2)$  that does not depend on the velocity variables  $(\phi, \psi)$ .

According to [12, Section 4], we have the well-posedness and regularity of the above Milne problem:

**Proposition 2.2** (Milne problem). *Assume*

$$\|\rho\|_{W_t^{2,\infty} W_{\iota_1, \iota_2}^{3,\infty} W_w^{1,\infty}} + \|e^{\alpha \eta} S\|_{W_t^{2,\infty} W_x^{3,\infty} W_w^{1,\infty}} \lesssim 1 \quad (2.31)$$

for some  $\alpha > 0$ . Then there exist a unique solution  $\Phi \in W_t^{2,\infty} W_{\iota_1, \iota_2}^{3,\infty} W_{\psi}^{1,\infty} L_{\eta, \phi}^{\infty}$  to (2.28)(2.29) and a  $\Phi_{\infty} \in W_t^{2,\infty} W_{\iota_1, \iota_2}^{3,\infty}$  such that (2.30) holds in the sense of

$$\|\Phi_{\infty}\|_{W_t^{2,\infty} W_{\iota_1, \iota_2}^{3,\infty}} + \|e^{\beta \eta} (\Phi - \Phi_{\infty})\|_{W_t^{2,\infty} W_{\iota_1, \iota_2}^{3,\infty} W_{\psi}^{1,\infty} L_{\eta, \phi}^{\infty}} \lesssim 1, \quad (2.32)$$

$$\|e^{\beta \eta} \sin \phi \partial_{\eta} (\Phi - \Phi_{\infty})\|_{L^{\infty}} + \|e^{\beta \eta} \sin \phi \partial_{\phi} (\Phi - \Phi_{\infty})\|_{L^{\infty}} \lesssim 1 \quad (2.33)$$

for any  $0 < \beta < \min\{1, \alpha\}$ .

Let  $\Phi_0$  be the solution to (2.28) with  $S = 0$  and the boundary data  $\rho(t; \iota_1, \iota_2; \phi, \psi) = g(t, x_0, w)$  satisfying (1.12), and so there exists  $\Phi_{0,\infty}(t; \iota_1, \iota_2) = \lim_{\eta \rightarrow \infty} \Phi_0(t; \eta, \iota_1, \iota_2; \phi, \psi)$ . Then  $\Psi_0 := \Phi_0 - \Phi_{0,\infty}$  satisfies

$$\begin{cases} \sin \phi \frac{\partial \Psi_0}{\partial \eta} + \Psi_0 - \overline{\Psi_0} = 0, \\ \Psi_0|_{\{\eta=0, \sin \phi > 0\}} = g - \Phi_{0,\infty}, \\ \lim_{\eta \rightarrow \infty} \Psi_0 = 0, \end{cases} \quad (2.34)$$

as well as

$$\|\Phi_{0,\infty}\|_{W_t^{2,\infty} W_{\iota_1, \iota_2}^{3,\infty}} + \|e^{\beta \eta} \Psi_0\|_{W_t^{2,\infty} W_{\iota_1, \iota_2}^{3,\infty} W_{\psi}^{1,\infty} L_{\eta, \phi}^{\infty}} \lesssim 1, \quad (2.35)$$

$$\|e^{\beta \eta} \sin \phi \partial_{\eta} \Psi_0\|_{L^{\infty}} + \|e^{\beta \eta} \sin \phi \partial_{\phi} \Psi_0\|_{L^{\infty}} \lesssim 1 \quad (2.36)$$

for any  $0 < \beta < 1$ .

Additionally, let  $\chi(r) \in C^{\infty}(\mathbb{R})$  and  $\tilde{\chi}(r) = 1 - \chi(r)$  be smooth cutoff functions satisfying  $\chi(r) = 1$  for  $|r| \leq 1$  and  $\chi(r) = 0$  for  $|r| \geq 2$ . We define the cutoff boundary layer

$$U_0^B(t; \eta, \iota_1, \iota_2; \phi, \psi) := \chi(\varepsilon^{\frac{1}{2}} \eta) \tilde{\chi}(\varepsilon^{-1} \phi) \Psi_0(t; \eta, \iota_1, \iota_2; \phi, \psi). \quad (2.37)$$

Here, the space cutoff  $\chi(\varepsilon^{\frac{1}{2}} \eta)$  restricts the boundary layer effect within a thin layer near the boundary and helps avoid self-interactions, while the velocity cutoff  $\tilde{\chi}(\varepsilon^{-1} \phi)$  truncates the grazing singularity. With these

cutoffs, the modified boundary layer  $U_0^B$  satisfies

$$\begin{cases} \sin \phi \frac{\partial U_0^B}{\partial \eta} + U_0^B - \overline{U_0^B} = \sin \phi \cdot \varepsilon^{\frac{1}{2}} \chi'(\varepsilon^{\frac{1}{2}} \eta) \tilde{\chi}(\varepsilon^{-1} \phi) \Psi_0 + \chi(\varepsilon^{\frac{1}{2}} \eta) \left[ \overline{\chi(\varepsilon^{-1} \phi) \Psi_0} - \chi(\varepsilon^{-1} \phi) \overline{\Psi_0} \right], \\ U_0^B(t; 0, \iota_1, \iota_2; \phi, \psi) = \tilde{\chi}(\varepsilon^{-1} \phi) \left[ g(t; \iota_1, \iota_2; \phi, \psi) - \Phi_{0,\infty}(t; \iota_1, \iota_2) \right] \quad \text{for } \sin \phi > 0, \\ \lim_{\eta \rightarrow \infty} U_0^B(t; \eta, \iota_1, \iota_2; \phi, \psi) = 0. \end{cases} \quad (2.38)$$

Moreover, it holds that

$$\| \| e^{\beta \eta} U_0^B \| \|_{W_t^{2,\infty} W_{\iota_1, \iota_2}^{3,\infty} W_\psi^{1,\infty} L_{\eta, \phi}^\infty} \lesssim 1 \quad (2.39)$$

for any  $0 < \beta < 1$ .

**2.4. Matching Procedure.** Now we construct the initial layer, boundary layer, and interior solution for each level of the asymptotic expansion in (2.1) via a matching procedure.

Construction of  $U_0^I$ ,  $U_0^B$ , and  $U_0$ . For the leading-order expansion, we want to enforce the matching initial condition

$$(U_0 + U_0^I)|_{t=0} = u_o, \quad (2.40)$$

and the matching boundary condition

$$(U_0 + U_0^B)|_{\Gamma_-} = g + O(\varepsilon^{0+}). \quad (2.41)$$

With the initial layer  $U_0^I$  given in (2.19) that satisfies (2.20) and the boundary layer  $U_0^B$  given in (2.37) that satisfies (2.38), we require the interior solution  $U_0$  to satisfy the following initial-boundary value problem (combining (2.5)):

$$\begin{cases} U_0 = \overline{U_0}, \quad \partial_t \overline{U_0} - \Delta_x \overline{U_0} = 0, \\ \overline{U_0}(0, x) = \overline{u}_o(x), \\ \overline{U_0}(t, x_0) = \Phi_{0,\infty}(t; \iota_1, \iota_2) \quad \text{for } x_0 \in \partial\Omega, \end{cases} \quad (2.42)$$

which uniquely determines  $U_0(t, x, w)$  due to classical theory for the heat equation. Also, with (1.12), (2.35), and by the standard parabolic estimate (cf. [14]), we have for any  $2 \leq p < \infty$

$$\| \| U_0 \| \|_{W_t^{2,\infty} W_x^{3,p}} + \| U_0 \|_{W_t^{2,\infty} W_{\iota_1, \iota_2}^{3,p}} \lesssim 1. \quad (2.43)$$

Then the joint boundary value is

$$(U_0 + U_0^B)|_{\Gamma_-} = g - \chi(\varepsilon^{-1} \phi) (g - \Phi_{0,\infty}). \quad (2.44)$$

Here the additional term  $\chi(\varepsilon^{-1} \phi) (g - \Phi_{0,\infty}) \sim O(\varepsilon^{\frac{1}{p}})$  in  $L_w^p$ -norm due to the velocity cutoff.

Construction of  $U_1^I$  and  $U_1$ . For the next-order expansion, we want to enforce the matching initial condition

$$(U_1 + U_1^I)|_{t=0} = 0. \quad (2.45)$$

With the initial layer  $U_1^I$  given in (2.22) that satisfies (2.23), we require the interior solution  $U_1$  to satisfy (combining (2.6)):

$$\begin{cases} U_1 = \overline{U_1} - w \cdot \nabla_x U_0, \quad \partial_t \overline{U_1} - \Delta_x \overline{U_1} = 0, \\ \overline{U_1}(0, x) = \Theta_{1,\infty}(x), \\ \overline{U_1}(t, x_0) = 0 \quad \text{for } x_0 \in \partial\Omega, \end{cases} \quad (2.46)$$

which uniquely determines  $U_1(t, x, w)$ . Also, with (2.43) and (2.24), we have for any  $2 \leq p < \infty$

$$\| \| U_1 \| \|_{W_t^{2,\infty} W_x^{2,p} L_w^\infty} + \| U_1 \|_{W_t^{2,\infty} W_{\iota_1, \iota_2}^{2,p} L_w^\infty} \lesssim 1. \quad (2.47)$$

Construction of  $U_2$ . Lastly, we solve for a higher-order expansion  $U_2(t, x, w)$  from (see (2.7))

$$\begin{cases} U_2 = \bar{U}_2 - w \cdot \nabla_x U_1 - \partial_t U_0, & \partial_t \bar{U}_2 - \Delta_x \bar{U}_2 = 0, \\ \bar{U}_2(0, x) = 0, \\ \bar{U}_2(t, x_0) = 0 & \text{for } x_0 \in \partial\Omega. \end{cases} \quad (2.48)$$

In view of (2.43) and (2.47), we have for any  $2 \leq p < \infty$

$$\|U_2\|_{W_t^{1,\infty} W_x^{1,p} L_w^\infty} + \|U_2\|_{W_t^{1,\infty} W_{t_1,t_2}^{1,p} L_w^\infty} \lesssim 1. \quad (2.49)$$

To summarize, we have obtained the well-posedness and regularity estimates of the interior solution, initial layer, and boundary layer:

**Proposition 2.3.** *Assume (1.12) holds for the initial data  $u_o$  and boundary data  $g$ . Let  $U_0, U_1, U_2$  be constructed via (2.42)(2.46)(2.48),  $U_0^I, U_1^I$  in Subsection 2.2, and  $U_0^B$  in Subsection 2.3. Then we have for any  $2 \leq p < \infty$*

$$\|U_0\|_{W_t^{2,\infty} W_x^{3,p}} + \|U_0\|_{W_t^{2,\infty} W_{t_1,t_2}^{3,p}} \lesssim 1, \quad (2.50)$$

$$\|U_1\|_{W_t^{2,\infty} W_x^{2,p} L_w^\infty} + \|U_1\|_{W_t^{2,\infty} W_{t_1,t_2}^{2,p} L_w^\infty} \lesssim 1, \quad (2.51)$$

$$\|U_2\|_{W_t^{1,\infty} W_x^{1,p} L_w^\infty} + \|U_2\|_{W_t^{1,\infty} W_{t_1,t_2}^{1,p} L_w^\infty} \lesssim 1, \quad (2.52)$$

and

$$\|e^\tau U_0^I\|_{L_\tau^\infty W_x^{3,\infty} L_w^\infty} \lesssim 1, \quad (2.53)$$

$$\|e^\tau U_1^I\|_{L_\tau^\infty W_x^{2,\infty} L_w^\infty} \lesssim 1, \quad (2.54)$$

$$\|e^{\beta\eta} U_0^B\|_{W_t^{2,\infty} W_{t_1,t_2}^{3,\infty} W_\psi^{1,\infty} L_{n,\phi}^\infty} \lesssim 1 \quad (2.55)$$

for any  $0 < \beta < 1$ .

### 3. REMAINDER EQUATION

Denote the approximate solution

$$u_a := (U_0 + \varepsilon U_1 + \varepsilon^2 U_2) + (U_0^I + \varepsilon U_1^I) + U_0^B, \quad (3.1)$$

and so the remainder

$$\mathcal{R} := u^\varepsilon - u_a. \quad (3.2)$$

Inserting (3.2) into (1.1), we have

$$\begin{aligned} \varepsilon \partial_t (u_a + \mathcal{R}) + w \cdot \nabla_x (u_a + \mathcal{R}) + \varepsilon^{-1} (u_a + \mathcal{R}) - \varepsilon^{-1} (\bar{u}_a + \bar{\mathcal{R}}) &= 0, \\ (u_a + \mathcal{R})|_{t=0} &= u_o, \quad (u_a + \mathcal{R})|_{\Gamma_-} = g, \end{aligned} \quad (3.3)$$

which yields

$$\begin{aligned} \varepsilon \partial_t \mathcal{R} + w \cdot \nabla_x \mathcal{R} + \varepsilon^{-1} (\mathcal{R} - \bar{\mathcal{R}}) &= -\varepsilon \partial_t u_a - w \cdot \nabla_x u_a - \varepsilon^{-1} (u_a - \bar{u}_a), \\ \mathcal{R}|_{t=0} &= u_o - u_a|_{t=0}, \quad \mathcal{R}|_{\Gamma_-} = g - u_a|_{\Gamma_-}. \end{aligned} \quad (3.4)$$

Consider the initial-boundary value problem for the remainder  $\mathcal{R}(t, x, w)$ :

$$\begin{cases} \varepsilon \partial_t \mathcal{R} + w \cdot \nabla_x \mathcal{R} + \varepsilon^{-1} (\mathcal{R} - \bar{\mathcal{R}}) = \mathcal{S} & \text{in } \mathbb{R}^+ \times \Omega \times \mathbb{S}^2, \\ \mathcal{R}(0, x, w) = \mathcal{I}(x, w) & \text{in } \Omega \times \mathbb{S}^2, \\ \mathcal{R}(t, x_0, w) = \mathcal{G}(t, x_0, w) & \text{for } x_0 \in \partial\Omega \text{ and } w \cdot n(x_0) < 0, \end{cases} \quad (3.5)$$

where

$$\bar{\mathcal{R}}(t, x) = \frac{1}{4\pi} \int_{\mathbb{S}^2} \mathcal{R}(t, x, w) dw. \quad (3.6)$$

In particular, we know the initial data is given by

$$\mathcal{I} := \varepsilon^2 w \cdot \nabla_x U_1|_{t=0} + \varepsilon^2 \partial_t U_0|_{t=0}, \quad (3.7)$$

and the boundary data is given by

$$\mathcal{G} := \chi(\varepsilon^{-1}\phi)(g - \Phi_{0,\infty}) - \varepsilon U_1^I|_{\Gamma_-} + \varepsilon(w \cdot \nabla_x U_0)|_{\Gamma_-} + \varepsilon^2(w \cdot \nabla_x U_1)|_{\Gamma_-} + \varepsilon^2 \partial_t \Phi_{0,\infty}. \quad (3.8)$$

Note that with the compatibility condition  $u_o|_\gamma = g|_{t=0} = C(x_0)$  (see (1.13)), we actually have  $U_0^B|_{t=0} = 0$  from the Milne problem and  $U_0^I|_{\Gamma_-} = 0$  from the initial layer problem.

The source term can be split as

$$\begin{aligned} \mathcal{S} &:= -\varepsilon \partial_t u_a - w \cdot \nabla_x u_a - \varepsilon^{-1}(u_a - \bar{u}_a) \\ &=: \mathcal{S}^{IS} + \mathcal{S}^{IL} + \mathcal{S}_1^{BL} + \mathcal{S}_2^{BL} + \mathcal{S}_3^{BL} \end{aligned} \quad (3.9)$$

with the expressions

$$\mathcal{S}^{IS} := -\varepsilon^2 w \cdot \nabla_x U_2 - \varepsilon^2 \partial_t U_1 - \varepsilon^3 \partial_t U_2, \quad (3.10)$$

$$\mathcal{S}^{IL} := -\varepsilon w \cdot \nabla_x U_1^I, \quad (3.11)$$

$$\mathcal{S}_1^{BL} := \left( \frac{\sin^2 \psi}{R_1 - \varepsilon \eta} + \frac{\cos^2 \psi}{R_2 - \varepsilon \eta} \right) \cos \phi \frac{\partial U_0^B}{\partial \phi}, \quad (3.12)$$

$$\mathcal{S}_2^{BL} := -\varepsilon \partial_t U_0^B - \frac{R_1 \cos \phi \sin \psi}{L_1(R_1 - \varepsilon \eta)} \frac{\partial U_0^B}{\partial \iota_1} - \frac{R_2 \cos \phi \cos \psi}{L_2(R_2 - \varepsilon \eta)} \frac{\partial U_0^B}{\partial \iota_2} \quad (3.13)$$

$$\begin{aligned} & - \frac{\sin \psi}{R_1 - \varepsilon \eta} \left\{ \frac{R_1 \cos \phi}{L_1 L_2} \left[ \varsigma_1 \cdot \left( \partial_{\iota_1 \iota_2} \mathbf{r} \times \varsigma_2 \right) \right] - \sin \phi \cos \psi \right\} \frac{\partial U_0^B}{\partial \psi} \\ & + \frac{\cos \psi}{R_2 - \varepsilon \eta} \left\{ \frac{R_2 \cos \phi}{L_1 L_2} \left[ \varsigma_2 \cdot \left( \partial_{\iota_1 \iota_2} \mathbf{r} \times \varsigma_1 \right) \right] - \sin \phi \sin \psi \right\} \frac{\partial U_0^B}{\partial \psi}, \\ \mathcal{S}_3^{BL} &:= -\varepsilon^{-1} \left\{ \sin \phi \cdot \varepsilon^{\frac{1}{2}} \chi'(\varepsilon^{\frac{1}{2}} \eta) \tilde{\chi}(\varepsilon^{-1} \phi) \Psi_0 + \chi(\varepsilon^{\frac{1}{2}} \eta) \left[ \overline{\chi(\varepsilon^{-1} \phi) \Psi_0} - \chi(\varepsilon^{-1} \phi) \bar{\Psi}_0 \right] \right\}. \end{aligned} \quad (3.14)$$

We now give preliminary estimates of the initial, boundary, and source terms specified above. The proof is straightforward and largely based on Proposition 2.3 and the certain definitions with rescaling and cutoffs (cf. [13, Section 3.3]).

**Lemma 3.1.** *Assume (1.12)(1.13) hold for  $u_o$  and  $g$ . For the initial term  $\mathcal{I}(x, w)$  given in (3.7), we have*

$$\|\mathcal{I}\|_{L^2} \lesssim \varepsilon^2. \quad (3.15)$$

For the boundary term  $\mathcal{G}(t, x_0, w)$  given in (3.8), we have

$$\|\mathcal{G}\|_{L_{\Gamma_-}^2} \lesssim \varepsilon. \quad (3.16)$$

For the source term  $\mathcal{S}(t, x, w)$  given in (3.9)–(3.14), we have

$$\|\|\mathcal{S}^{IS}\|\|_{L^2} \lesssim \varepsilon^2, \quad \|\|\mathcal{S}^{IL}\|\|_{L^2} \lesssim \varepsilon^2, \quad (3.17)$$

$$\|\|(1 + \eta) \mathcal{S}_1^{BL}\|\|_{L^2} \lesssim 1, \quad \|\|(1 + \eta) \mathcal{S}_2^{BL}\|\|_{L^2} \lesssim \varepsilon^{\frac{1}{2}}, \quad (3.18)$$

$$\|\|(1 + \eta) \mathcal{S}_3^{BL}\|\|_{L^2} \lesssim 1, \quad \|\|(1 + \eta) \mathcal{S}_3^{BL}\|\|_{L_t^2 L_x^2 L_w^1} \lesssim \varepsilon^{\frac{1}{2}}. \quad (3.19)$$

Moreover, the boundary layer  $U_0^B$  defined in (2.37) satisfies

$$\|\|(1 + \eta) U_0^B\|\|_{L^2} \lesssim \varepsilon^{\frac{1}{2}}. \quad (3.20)$$

In preparation for the a priori estimate of the remainder, we need to write the weak formulation of the remainder equation, which depends essentially on the validity of integration by parts for the transport operator (cf. [10, Lemma 2.2]):

**Lemma 3.2** (Green's identity). *Assume  $f(t, x, w), g(t, x, w) \in L^\infty([0, T]; L^2(\Omega \times \mathbb{S}^2))$  with  $\partial_t f + w \cdot \nabla_x f, \partial_t g + w \cdot \nabla_x g \in L^2([0, T] \times \Omega \times \mathbb{S}^2)$  and  $f, g \in L^2_{\Gamma_+}$ . Then for almost all  $t, s \in [0, T]$ , we have*

$$\begin{aligned} & \int_s^t \iint_{\Omega \times \mathbb{S}^2} \left\{ (\partial_t f + w \cdot \nabla_x f)g + (\partial_t g + w \cdot \nabla_x g)f \right\} \\ &= \iint_{\Omega \times \mathbb{S}^2} f(t)g(t) - \iint_{\Omega \times \mathbb{S}^2} f(s)g(s) + \int_s^t \int_{\gamma} fg(w \cdot n). \end{aligned} \quad (3.21)$$

With this result, the weak formulation of (3.5) takes the following form: for any test function  $\varphi(t, x, w) \in L^\infty([0, T]; L^2(\Omega \times \mathbb{S}^2))$  with  $\partial_t \varphi + w \cdot \nabla_x \varphi \in L^2([0, T] \times \Omega \times \mathbb{S}^2)$  and  $\varphi \in L^2_{\Gamma_+}$ , it holds for any  $t \in [0, T]$  that

$$\begin{aligned} & \varepsilon \langle \mathcal{R}(t), \varphi(t) \rangle_{x, w} - \varepsilon \langle \mathcal{I}, \varphi(0) \rangle_{x, w} + \langle \mathcal{R}, \varphi \rangle_{\Gamma_+} - \langle \mathcal{G}, \varphi \rangle_{\Gamma_-} \\ &= \varepsilon \langle \mathcal{R}, \partial_t \varphi \rangle + \langle \mathcal{R}, w \cdot \nabla_x \varphi \rangle - \varepsilon^{-1} \langle \mathcal{R} - \overline{\mathcal{R}}, \varphi \rangle + \langle \mathcal{S}, \varphi \rangle. \end{aligned} \quad (3.22)$$

Here, the time integral is taken over  $[0, t]$ .

#### 4. REMAINDER ESTIMATE

Our goal of this section is to prove the remainder estimate for

$$\mathcal{R} := u^\varepsilon - \left[ (U_0 + \varepsilon U_1 + \varepsilon^2 U_2) + (U_0^I + \varepsilon U_1^I) + U_0^B \right]. \quad (4.1)$$

**Theorem 4.1** (Remainder estimate). *Under the assumption (1.12) with (1.13), we have*

$$\|\mathcal{R}\|_{L^\infty_t L^2_{x, w}} + \varepsilon^{-\frac{1}{2}} \|\mathcal{R}\|_{L^2_{\Gamma_+}} + \varepsilon^{-\frac{1}{2}} \|\overline{\mathcal{R}}\|_{L^2} + \varepsilon^{-1} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2} \lesssim 1, \quad (4.2)$$

where the implicit constant is independent of  $\varepsilon$ .

The combination of the following two lemmas, which will be presented in the rest of this section, will lead immediately to the desired estimate above.

**Lemma 4.2** (Energy estimate). *Under the assumption (1.12) with (1.13), we have*

$$\|\mathcal{R}\|_{L^\infty_t L^2_{x, w}}^2 + \varepsilon^{-1} \|\mathcal{R}\|_{L^2_{\Gamma_+}}^2 + \varepsilon^{-2} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}^2 \lesssim \delta \varepsilon^{-1} \|\overline{\mathcal{R}}\|_{L^2}^2 + \delta^{-1}. \quad (4.3)$$

Here  $0 < \delta \ll 1$  is a constant that can be taken sufficiently small.

*Proof.* Taking  $\varphi = \varepsilon^{-1} \mathcal{R}$  in the weak formulation (3.22) and using the fundamental theorem of calculus and divergence theorem, we obtain

$$\frac{1}{2} \|\mathcal{R}(t)\|_{L^2}^2 - \frac{1}{2} \|\mathcal{I}\|_{L^2}^2 + \frac{1}{2} \varepsilon^{-1} \|\mathcal{R}\|_{L^2_{\Gamma_+}}^2 - \frac{1}{2} \varepsilon^{-1} \|\mathcal{G}\|_{L^2_{\Gamma_-}}^2 = -\varepsilon^{-2} \langle \mathcal{R} - \overline{\mathcal{R}}, \mathcal{R} \rangle + \varepsilon^{-1} \langle \mathcal{S}, \mathcal{R} \rangle. \quad (4.4)$$

Observing the orthogonality  $\langle \overline{\mathcal{R}}, \mathcal{R} - \overline{\mathcal{R}} \rangle_w = 0$ , we then rearrange the terms to arrive at

$$\frac{1}{2} \|\mathcal{R}(t)\|_{L^2}^2 + \frac{1}{2} \varepsilon^{-1} \|\mathcal{R}\|_{L^2_{\Gamma_+}}^2 + \varepsilon^{-2} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}^2 = \frac{1}{2} \|\mathcal{I}\|_{L^2}^2 + \frac{1}{2} \varepsilon^{-1} \|\mathcal{G}\|_{L^2_{\Gamma_-}}^2 + \varepsilon^{-1} \langle \mathcal{S}, \mathcal{R} \rangle. \quad (4.5)$$

By (3.15) and (3.16), we know

$$\|\mathcal{I}\|_{L^2}^2 + \varepsilon^{-1} \|\mathcal{G}\|_{L^2_{\Gamma_-}}^2 \lesssim \varepsilon. \quad (4.6)$$

Now we split the last term in (4.5) and estimate each part separately. Using Young's inequality and by (3.17), we have

$$\varepsilon^{-1} \left| \langle \mathcal{S}^{IS} + \mathcal{S}^{IL}, \mathcal{R} \rangle \right| \lesssim \delta \|\mathcal{R}\|_{L^2}^2 + \delta^{-1} \varepsilon^{-2} \|\mathcal{S}^{IS} + \mathcal{S}^{IL}\|_{L^2}^2 \lesssim \delta \|\mathcal{R}\|_{L^2}^2 + \delta^{-1} \varepsilon^2. \quad (4.7)$$

Likewise, by (3.18) and (3.19), we have

$$\begin{aligned} \varepsilon^{-1} \left| \langle \mathcal{S}_1^{BL} + \mathcal{S}_2^{BL} + \mathcal{S}_3^{BL}, \mathcal{R} - \overline{\mathcal{R}} \rangle \right| &\lesssim \delta \varepsilon^{-2} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}^2 + \delta^{-1} \|\mathcal{S}_1^{BL} + \mathcal{S}_2^{BL} + \mathcal{S}_3^{BL}\|_{L^2}^2 \\ &\lesssim \delta \varepsilon^{-2} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}^2 + \delta^{-1}. \end{aligned} \quad (4.8)$$

For  $\varepsilon^{-1} \langle \mathcal{S}_1^{BL}, \overline{\mathcal{R}} \rangle$ , we use integration by parts in  $\phi$  (note that the velocity integral contains a Jacobian  $\cos \phi$ ) followed by (3.20) and get

$$\begin{aligned} \varepsilon^{-1} \left| \langle \mathcal{S}_1^{BL}, \overline{\mathcal{R}} \rangle \right| &= \varepsilon^{-1} \left| \left\langle \left( \frac{\sin^2 \psi}{R_1 - \varepsilon \eta} + \frac{\cos^2 \psi}{R_2 - \varepsilon \eta} \right) \cos \phi \partial_\phi U_0^B, \overline{\mathcal{R}} \right\rangle \right| \\ &= \varepsilon^{-1} \left| \left\langle \left( \frac{\sin^2 \psi}{R_1 - \varepsilon \eta} + \frac{\cos^2 \psi}{R_2 - \varepsilon \eta} \right) 2 \sin \phi U_0^B, \overline{\mathcal{R}} \right\rangle \right| \\ &\lesssim \varepsilon^{-1} \left\| U_0^B \right\|_{L_t^2 L_x^2 L_w^1} \left\| \overline{\mathcal{R}} \right\|_{L^2} \lesssim \varepsilon^{-\frac{1}{2}} \left\| \overline{\mathcal{R}} \right\|_{L^2} \lesssim \delta \varepsilon^{-1} \left\| \overline{\mathcal{R}} \right\|_{L^2}^2 + \delta^{-1}. \end{aligned} \quad (4.9)$$

Turning to the remaining terms, we use (3.18)(3.19) to estimate

$$\begin{aligned} \varepsilon^{-1} \left| \langle \mathcal{S}_2^{BL} + \mathcal{S}_3^{BL}, \overline{\mathcal{R}} \rangle \right| &\lesssim \varepsilon^{-1} \left\| \overline{\mathcal{R}} \right\|_{L^2} \left( \left\| \mathcal{S}_2^{BL} \right\|_{L_t^2 L_x^2 L_w^1} + \left\| \mathcal{S}_3^{BL} \right\|_{L_t^2 L_x^2 L_w^1} \right) \\ &\lesssim \varepsilon^{-\frac{1}{2}} \left\| \overline{\mathcal{R}} \right\|_{L^2} \lesssim \delta \varepsilon^{-1} \left\| \overline{\mathcal{R}} \right\|_{L^2}^2 + \delta^{-1}. \end{aligned} \quad (4.10)$$

Collecting (4.7) through (4.10), we deduce that

$$\varepsilon^{-1} \left| \langle \mathcal{S}, \mathcal{R} \rangle \right| \lesssim \delta \varepsilon^{-2} \left\| \mathcal{R} - \overline{\mathcal{R}} \right\|_{L^2}^2 + \delta \varepsilon^{-1} \left\| \overline{\mathcal{R}} \right\|_{L^2}^2 + \delta^{-1}. \quad (4.11)$$

Finally, combining (4.5) with (4.6) and (4.11) settles (4.3).  $\square$

**Lemma 4.3** (Kernel estimate). *Under the assumption (1.12) with (1.13), we have*

$$\left\| \overline{\mathcal{R}} \right\|_{L^2}^2 \lesssim \varepsilon \left\| \mathcal{R} \right\|_{L_t^\infty L_x^2 L_w}^2 + \left\| \mathcal{R} - \overline{\mathcal{R}} \right\|_{L^2}^2 + \left\| \mathcal{R} \right\|_{L_{\Gamma^+}^2}^2 + \varepsilon. \quad (4.12)$$

*Proof.* We will use an auxiliary Dirichlet problem for the Poisson equation to construct our test functions. Let  $\xi(t, x) := (-\Delta_x)^{-1} \overline{\mathcal{R}}(t, x)$  be the solution to (for fixed  $t$ )

$$\begin{cases} -\Delta_x \xi = \overline{\mathcal{R}} & \text{in } \Omega, \\ \xi = 0 & \text{on } \partial\Omega. \end{cases} \quad (4.13)$$

Based on the standard elliptic estimate and the trace theorem, for every  $t \in \mathbb{R}^+$  we have

$$\left\| \xi(t) \right\|_{H^2(\Omega)} + \left\| \nabla_x \xi(t) \right\|_{H^{\frac{1}{2}}(\partial\Omega)} \lesssim \left\| \overline{\mathcal{R}}(t) \right\|_{L^2(\Omega)}. \quad (4.14)$$

We first test (3.5) with  $\varphi = \xi$  (or equivalently, take  $\varphi = \xi$  in (3.22)) to get

$$\varepsilon \langle \partial_t \mathcal{R}, \xi \rangle + \langle \mathcal{R}, \xi \rangle_{\Gamma^+} - \langle \mathcal{G}, \xi \rangle_{\Gamma^-} - \langle \mathcal{R}, w \cdot \nabla_x \xi \rangle + \varepsilon^{-1} \langle \mathcal{R} - \overline{\mathcal{R}}, \xi \rangle = \langle \mathcal{S}, \xi \rangle. \quad (4.15)$$

Notice that  $\langle \mathcal{R} - \overline{\mathcal{R}}, \xi \rangle_w = 0$  and  $\langle \partial_t \mathcal{R} - \partial_t \overline{\mathcal{R}}, \xi \rangle_w = 0$  by orthogonality,  $\langle \overline{\mathcal{R}}, w \cdot \nabla_x \xi \rangle_w = 0$  by symmetry, and that the boundary terms vanish due to  $\xi|_{\partial\Omega} = 0$ . Hence, we are left with

$$\varepsilon \langle \partial_t \overline{\mathcal{R}}, \xi \rangle - \langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x \xi \rangle = \langle \mathcal{S}, \xi \rangle. \quad (4.16)$$

We then test (3.5) with  $\varphi = w \cdot \nabla_x \xi$  to get

$$\begin{aligned} \varepsilon \langle \partial_t \mathcal{R}, w \cdot \nabla_x \xi \rangle + \langle \mathcal{R}, w \cdot \nabla_x \xi \rangle_{\Gamma^+} - \langle \mathcal{G}, w \cdot \nabla_x \xi \rangle_{\Gamma^-} \\ - \langle \mathcal{R}, w \cdot \nabla_x (w \cdot \nabla_x \xi) \rangle + \varepsilon^{-1} \langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x \xi \rangle = \langle \mathcal{S}, w \cdot \nabla_x \xi \rangle. \end{aligned} \quad (4.17)$$

Adding  $\varepsilon^{-1} \times (4.16)$  and (4.17) to eliminate  $\varepsilon^{-1} \langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x \xi \rangle$ , we obtain

$$\begin{aligned} \langle \partial_t \overline{\mathcal{R}}, \xi \rangle + \varepsilon \langle \partial_t \mathcal{R}, w \cdot \nabla_x \xi \rangle + \langle \mathcal{R}, w \cdot \nabla_x \xi \rangle_{\Gamma^+} - \langle \mathcal{G}, w \cdot \nabla_x \xi \rangle_{\Gamma^-} \\ - \langle \mathcal{R}, w \cdot \nabla_x (w \cdot \nabla_x \xi) \rangle = \varepsilon^{-1} \langle \mathcal{S}, \xi \rangle + \langle \mathcal{S}, w \cdot \nabla_x \xi \rangle. \end{aligned} \quad (4.18)$$

On the one hand, we may split

$$- \langle \mathcal{R}, w \cdot \nabla_x (w \cdot \nabla_x \xi) \rangle = - \langle \overline{\mathcal{R}}, w \cdot \nabla_x (w \cdot \nabla_x \xi) \rangle - \langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x (w \cdot \nabla_x \xi) \rangle, \quad (4.19)$$

where the kernel part has the key positivity:

$$- \langle \overline{\mathcal{R}}, w \cdot \nabla_x (w \cdot \nabla_x \xi) \rangle \simeq - \langle \overline{\mathcal{R}}, \Delta_x \xi \rangle = \left\| \overline{\mathcal{R}} \right\|_{L^2}^2, \quad (4.20)$$

while the non-kernel part can be bounded as

$$\left| \langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x (w \cdot \nabla_x \xi) \rangle \right| \lesssim \left\| \mathcal{R} - \overline{\mathcal{R}} \right\|_{L^2} \left\| \xi \right\|_{L_t^2 H_x^2} \lesssim \delta \left\| \overline{\mathcal{R}} \right\|_{L^2}^2 + \delta^{-1} \left\| \mathcal{R} - \overline{\mathcal{R}} \right\|_{L^2}^2. \quad (4.21)$$

On the other hand, integration by parts reveals that

$$\langle\langle \partial_t \overline{\mathcal{R}}, \xi \rangle\rangle = -\langle\langle \partial_t \Delta_x \xi, \xi \rangle\rangle = \langle\langle \partial_t \nabla_x \xi, \nabla_x \xi \rangle\rangle = \frac{1}{2} \|\nabla_x \xi(t)\|_{L^2}^2 - \frac{1}{2} \|\nabla_x \xi(0)\|_{L^2}^2, \quad (4.22)$$

where (3.15) yields

$$\frac{1}{2} \|\nabla_x \xi(0)\|_{L^2}^2 \lesssim \|\overline{\mathcal{R}}(0)\|_{L^2}^2 = \|\overline{\mathcal{I}}\|_{L^2}^2 \lesssim \|\mathcal{I}\|_{L^2}^2 \lesssim \varepsilon^4. \quad (4.23)$$

In addition, we bound

$$\begin{aligned} \varepsilon \langle\langle \partial_t \mathcal{R}, w \cdot \nabla_x \xi \rangle\rangle &= \varepsilon \langle\langle \partial_t (\mathcal{R} - \overline{\mathcal{R}}), w \cdot \nabla_x \xi \rangle\rangle \\ &= -\varepsilon \langle\langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x \partial_t \xi \rangle\rangle + \varepsilon \langle (\mathcal{R} - \overline{\mathcal{R}})(t), w \cdot \nabla_x \xi(t) \rangle_{x,w} - \varepsilon \langle (\mathcal{R} - \overline{\mathcal{R}})(0), w \cdot \nabla_x \xi(0) \rangle_{x,w}, \end{aligned} \quad (4.24)$$

by

$$\varepsilon \left| \langle\langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x \partial_t \xi \rangle\rangle \right| \lesssim \varepsilon \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2} \|\nabla_x \partial_t \xi\|_{L^2} \lesssim \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}^2 + \varepsilon^2 \|\nabla_x \partial_t \xi\|_{L^2}^2, \quad (4.25)$$

$$\varepsilon \left| \langle (\mathcal{R} - \overline{\mathcal{R}})(t), w \cdot \nabla_x \xi(t) \rangle_{x,w} \right| \lesssim \varepsilon \|(\mathcal{R} - \overline{\mathcal{R}})(t)\|_{L^2} \|\xi(t)\|_{H^1} \lesssim \varepsilon \|\mathcal{R}(t)\|_{L^2}^2, \quad (4.26)$$

$$\varepsilon \left| \langle (\mathcal{R} - \overline{\mathcal{R}})(0), w \cdot \nabla_x \xi(0) \rangle_{x,w} \right| \lesssim \varepsilon \|\mathcal{R}(0)\|_{L^2}^2 = \varepsilon \|\mathcal{I}\|_{L^2}^2 \lesssim \varepsilon^5. \quad (4.27)$$

Also, the boundary terms can be controlled using (4.14) and (3.16):

$$\begin{aligned} \left| \langle \mathcal{R}, w \cdot \nabla_x \xi \rangle_{\Gamma_+} - \langle \mathcal{G}, w \cdot \nabla_x \xi \rangle_{\Gamma_-} \right| &\lesssim \left( \|\mathcal{R}\|_{L^2_{\Gamma_+}} + \|\mathcal{G}\|_{L^2_{\Gamma_-}} \right) \|\nabla_x \xi\|_{L^2_{t,\partial\Omega}} \\ &\lesssim \delta \|\overline{\mathcal{R}}\|_{L^2}^2 + \delta^{-1} \|\mathcal{R}\|_{L^2_{\Gamma_+}}^2 + \delta^{-1} \varepsilon^2. \end{aligned} \quad (4.28)$$

Collecting (4.18) through (4.28), we deduce that

$$\begin{aligned} \|\nabla_x \xi(t)\|_{L^2}^2 + \|\overline{\mathcal{R}}\|_{L^2}^2 &\lesssim \varepsilon^2 \|\nabla_x \partial_t \xi\|_{L^2}^2 + \varepsilon \|\mathcal{R}(t)\|_{L^2}^2 + \delta^{-1} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}^2 + \delta^{-1} \|\mathcal{R}\|_{L^2_{\Gamma_+}}^2 + \delta^{-1} \varepsilon^2 \\ &\quad + \varepsilon^{-1} |\langle\langle \mathcal{S}, \xi \rangle\rangle| + |\langle\langle \mathcal{S}, w \cdot \nabla_x \xi \rangle\rangle|. \end{aligned} \quad (4.29)$$

Next, we estimate the term  $\varepsilon^2 \|\nabla_x \partial_t \xi\|_{L^2}^2$  on the RHS above. Let  $\zeta := \partial_t \xi = (-\Delta_x)^{-1} \partial_t \overline{\mathcal{R}}$  denote the solution to

$$\begin{cases} -\Delta_x \zeta = \partial_t \overline{\mathcal{R}} & \text{in } \Omega, \\ \zeta = 0 & \text{on } \partial\Omega. \end{cases} \quad (4.30)$$

Poincaré's inequality indicates that

$$\|\zeta(t)\|_{L^2(\Omega)} \lesssim \|\nabla_x \zeta(t)\|_{L^2(\Omega)} = \|\nabla_x \partial_t \xi(t)\|_{L^2(\Omega)}. \quad (4.31)$$

Testing (3.5) against  $\varphi = \varepsilon \zeta$  and similar to (4.16) we get

$$\varepsilon^2 \langle\langle \partial_t \overline{\mathcal{R}}, \zeta \rangle\rangle - \varepsilon \langle\langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x \zeta \rangle\rangle = \varepsilon \langle\langle \mathcal{S}, \zeta \rangle\rangle. \quad (4.32)$$

Again, integration by parts reveals that

$$\varepsilon^2 \langle\langle \partial_t \overline{\mathcal{R}}, \zeta \rangle\rangle = -\varepsilon^2 \langle\langle \Delta_x \zeta, \zeta \rangle\rangle = \varepsilon^2 \|\nabla_x \zeta\|_{L^2}^2 = \varepsilon^2 \|\nabla_x \partial_t \xi\|_{L^2}^2. \quad (4.33)$$

Also, we have

$$\varepsilon \left| \langle\langle \mathcal{R} - \overline{\mathcal{R}}, w \cdot \nabla_x \zeta \rangle\rangle \right| \lesssim \varepsilon \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2} \|\nabla_x \zeta\|_{L^2} \lesssim \delta \varepsilon^2 \|\nabla_x \partial_t \xi\|_{L^2}^2 + \delta^{-1} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}^2. \quad (4.34)$$

Hence, (4.32) together with (4.33) and (4.34) yields

$$\varepsilon^2 \|\nabla_x \partial_t \xi\|_{L^2}^2 \lesssim \delta^{-1} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}^2 + \varepsilon |\langle\langle \mathcal{S}, \zeta \rangle\rangle|. \quad (4.35)$$

We then insert (4.35) into (4.29) and obtain

$$\begin{aligned} \|\nabla_x \xi(t)\|_{L^2}^2 + \varepsilon^2 \|\nabla_x \partial_t \xi\|_{L^2}^2 + \|\overline{\mathcal{R}}\|_{L^2}^2 &\lesssim \varepsilon \|\mathcal{R}(t)\|_{L^2}^2 + \delta^{-1} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}^2 + \delta^{-1} \|\mathcal{R}\|_{L^2_{\Gamma_+}}^2 + \delta^{-1} \varepsilon^2 \\ &\quad + \varepsilon^{-1} |\langle\langle \mathcal{S}, \xi \rangle\rangle| + |\langle\langle \mathcal{S}, w \cdot \nabla_x \xi \rangle\rangle| + \varepsilon |\langle\langle \mathcal{S}, \zeta \rangle\rangle|. \end{aligned} \quad (4.36)$$

Now it remains to handle the last three terms above involving the source term  $\mathcal{S}$ . By using (4.14), (4.31), and (3.17), we see that

$$\begin{aligned} & \varepsilon^{-1} \left| \langle \mathcal{S}^{IS} + \mathcal{S}^{IL}, \xi \rangle \right| + \left| \langle \mathcal{S}^{IS} + \mathcal{S}^{IL}, w \cdot \nabla_x \xi \rangle \right| + \varepsilon \left| \langle \mathcal{S}^{IS} + \mathcal{S}^{IL}, \zeta \rangle \right| \\ & \lesssim \varepsilon^{-1} \left( \left\| \mathcal{S}^{IS} \right\|_{L^2} + \left\| \mathcal{S}^{IL} \right\|_{L^2} \right) \left\| \xi \right\|_{L_t^2 H_x^1} + \varepsilon \left( \left\| \mathcal{S}^{IS} \right\|_{L^2} + \left\| \mathcal{S}^{IL} \right\|_{L^2} \right) \left\| \zeta \right\|_{L^2} \\ & \lesssim \varepsilon \left\| \overline{\mathcal{R}} \right\|_{L^2} + \varepsilon^3 \left\| \nabla_x \partial_t \xi \right\|_{L^2} \lesssim \delta \left\| \overline{\mathcal{R}} \right\|_{L^2}^2 + \delta \varepsilon^2 \left\| \nabla_x \partial_t \xi \right\|_{L^2}^2 + \delta^{-1} \varepsilon^2. \end{aligned} \quad (4.37)$$

Analogous to (4.9), we may transfer  $\mathcal{S}_1^{BL}$  to  $U_0^B$  via integration by parts in  $\phi$ . Noting that  $\xi|_{\partial\Omega} = 0$ , we then invoke Hardy's inequality and (3.18)(3.19)(3.20) followed by (4.14) to bound

$$\begin{aligned} & \varepsilon^{-1} \left| \langle \mathcal{S}_1^{BL} + \mathcal{S}_2^{BL} + \mathcal{S}_3^{BL}, \xi \rangle \right| \\ & \lesssim \varepsilon^{-1} \left\langle \left| U_0^B \right| + \left| \mathcal{S}_2^{BL} \right| + \left| \mathcal{S}_3^{BL} \right|, \left| \int_0^\mu \partial_\mu \xi \right| \right\rangle = \left\langle \left| \eta U_0^B \right| + \left| \eta \mathcal{S}_2^{BL} \right| + \left| \eta \mathcal{S}_3^{BL} \right|, \left| \frac{1}{\mu} \int_0^\mu \partial_\mu \xi \right| \right\rangle \\ & \lesssim \left( \left\| \eta U_0^B \right\|_{L_t^2 L_x^2 L_w^1} + \left\| \eta \mathcal{S}_2^{BL} \right\|_{L_t^2 L_x^2 L_w^1} + \left\| \eta \mathcal{S}_3^{BL} \right\|_{L_t^2 L_x^2 L_w^1} \right) \left\| \frac{1}{\mu} \int_0^\mu \partial_\mu \xi \right\|_{L^2} \\ & \lesssim \varepsilon^{\frac{1}{2}} \left\| \partial_\mu \xi \right\|_{L^2} \lesssim \varepsilon^{\frac{1}{2}} \left\| \xi \right\|_{L_t^2 H_x^1} \lesssim \varepsilon^{\frac{1}{2}} \left\| \overline{\mathcal{R}} \right\|_{L^2} \lesssim \delta \left\| \overline{\mathcal{R}} \right\|_{L^2}^2 + \delta^{-1} \varepsilon. \end{aligned} \quad (4.38)$$

In a parallel fashion, we have

$$\begin{aligned} & \left| \langle \mathcal{S}_1^{BL} + \mathcal{S}_2^{BL} + \mathcal{S}_3^{BL}, w \cdot \nabla_x \xi \rangle \right| \\ & \lesssim \left\langle \left| U_0^B \right| + \left| \mathcal{S}_2^{BL} \right| + \left| \mathcal{S}_3^{BL} \right|, \left| (\nabla_x \xi) \Big|_{\mu=0} + \int_0^\mu \partial_\mu (\nabla_x \xi) \right| \right\rangle \\ & \lesssim \left\langle \left| U_0^B \right| + \left| \mathcal{S}_2^{BL} \right| + \left| \mathcal{S}_3^{BL} \right|, \left| \nabla_x \xi \Big|_{\mu=0} \right\rangle + \varepsilon \left\langle \left| \eta U_0^B \right| + \left| \eta \mathcal{S}_2^{BL} \right| + \left| \eta \mathcal{S}_3^{BL} \right|, \left| \frac{1}{\mu} \int_0^\mu \partial_\mu (\nabla_x \xi) \right| \right\rangle \\ & \lesssim \varepsilon^{\frac{1}{2}} \left\| \nabla_x \xi \Big|_{L_{t,\partial\Omega}^2} \right\| + \varepsilon^{\frac{3}{2}} \left\| \partial_\mu (\nabla_x \xi) \right\|_{L^2} \lesssim \varepsilon^{\frac{1}{2}} \left\| \nabla_x \xi \Big|_{L_{t,\partial\Omega}^2} \right\| + \varepsilon^{\frac{3}{2}} \left\| \xi \right\|_{L_t^2 H_x^1} \lesssim \varepsilon^{\frac{1}{2}} \left\| \overline{\mathcal{R}} \right\|_{L^2} \lesssim \delta \left\| \overline{\mathcal{R}} \right\|_{L^2}^2 + \delta^{-1} \varepsilon, \end{aligned} \quad (4.39)$$

and

$$\begin{aligned} & \varepsilon \left| \langle \mathcal{S}_1^{BL} + \mathcal{S}_2^{BL} + \mathcal{S}_3^{BL}, \zeta \rangle \right| \\ & \lesssim \varepsilon \left\langle \left| U_0^B \right| + \left| \mathcal{S}_2^{BL} \right| + \left| \mathcal{S}_3^{BL} \right|, \left| \int_0^\mu \partial_\mu \zeta \right| \right\rangle = \varepsilon^2 \left\langle \left| \eta U_0^B \right| + \left| \eta \mathcal{S}_2^{BL} \right| + \left| \eta \mathcal{S}_3^{BL} \right|, \left| \frac{1}{\mu} \int_0^\mu \partial_\mu \zeta \right| \right\rangle \\ & \lesssim \varepsilon^2 \left( \left\| \eta U_0^B \right\|_{L_t^2 L_x^2 L_w^1} + \left\| \eta \mathcal{S}_2^{BL} \right\|_{L_t^2 L_x^2 L_w^1} + \left\| \eta \mathcal{S}_3^{BL} \right\|_{L_t^2 L_x^2 L_w^1} \right) \left\| \frac{1}{\mu} \int_0^\mu \partial_\mu \zeta \right\|_{L^2} \\ & \lesssim \varepsilon^{\frac{5}{2}} \left\| \partial_\mu \zeta \right\|_{L^2} \lesssim \varepsilon^{\frac{5}{2}} \left\| \nabla_x \partial_t \xi \right\|_{L^2} \lesssim \delta \varepsilon^2 \left\| \nabla_x \partial_t \xi \right\|_{L^2}^2 + \delta^{-1} \varepsilon^3. \end{aligned} \quad (4.40)$$

Collecting (4.37) through (4.40), we find

$$\varepsilon^{-1} \left| \langle \mathcal{S}, \xi \rangle \right| + \left| \langle \mathcal{S}, w \cdot \nabla_x \xi \rangle \right| + \varepsilon \left| \langle \mathcal{S}, \zeta \rangle \right| \lesssim \delta \left\| \overline{\mathcal{R}} \right\|_{L^2}^2 + \delta \varepsilon^2 \left\| \nabla_x \partial_t \xi \right\|_{L^2}^2 + \delta^{-1} \varepsilon. \quad (4.41)$$

Combined with (4.36), this yields

$$\left\| \nabla_x \xi(t) \right\|_{L^2}^2 + \varepsilon^2 \left\| \nabla_x \partial_t \xi \right\|_{L^2}^2 + \left\| \overline{\mathcal{R}} \right\|_{L^2}^2 \lesssim \varepsilon \left\| \mathcal{R}(t) \right\|_{L^2}^2 + \delta^{-1} \left\| \mathcal{R} - \overline{\mathcal{R}} \right\|_{L^2}^2 + \delta^{-1} \left\| \mathcal{R} \right\|_{L_{\Gamma_+}^2}^2 + \delta^{-1} \varepsilon, \quad (4.42)$$

and thus (4.12) follows.  $\square$

## 5. DIFFUSIVE LIMIT

We are now ready to prove our main result for the in-flow boundary problem:

*Proof of Theorem 1.1.* The well-posedness of (1.1) is rather standard and was addressed already in [6, 5, 29]. The existence of  $U_0$  is guaranteed in Section 2.4. This leaves us to demonstrate the approximation estimate (1.14).



Given the interior solutions  $U_0, U_1, U_2$ , the initial layers  $U_0^I, U_1^I$ , and the boundary layer  $U_0^B$  constructed in Section 2, the remainder estimate (Theorem 4.1) implies

$$\| \| u^\varepsilon - U_0 - \varepsilon U_1 - \varepsilon^2 U_2 - U_0^I - \varepsilon U_1^I - U_0^B \| \|_{L^2} \lesssim \varepsilon^{\frac{1}{2}}. \quad (5.1)$$

On the other hand, using the estimates recalled in Proposition 2.3, we see that

$$\| \| \varepsilon U_1 + \varepsilon^2 U_2 \| \|_{L^2} \lesssim \varepsilon, \quad \| \| U_0^I + \varepsilon U_1^I \| \|_{L^2} \lesssim \varepsilon, \quad \| \| U_0^B \| \|_{L^2} \lesssim \varepsilon^{\frac{1}{2}}, \quad (5.2)$$

where we take into account the temporal rescaling  $\tau = \varepsilon^{-2}t$  for the initial layer and the normal spatial rescaling  $\eta = \varepsilon^{-1}\mu$  for the boundary layer. Consequently, (1.14) follows by combining (5.1) with (5.2). This completes the proof of the theorem.  $\square$

## 6. DIFFUSE-REFLECTION BOUNDARY PROBLEM

Consider the unsteady neutron transport equation with the diffuse-reflection boundary condition:

$$\begin{cases} \varepsilon \partial_t u^\varepsilon + w \cdot \nabla_x u^\varepsilon + \varepsilon^{-1} (u^\varepsilon - \bar{u}^\varepsilon) = 0 & \text{in } \mathbb{R}^+ \times \Omega \times \mathbb{S}^2, \\ u^\varepsilon(0, x, w) = u_o(x, w) & \text{in } \Omega \times \mathbb{S}^2, \\ u^\varepsilon(t, x_0, w) = \mathcal{P}_\gamma[u^\varepsilon](t, x_0) + \varepsilon h(t, x_0, w) & \text{for } x_0 \in \partial\Omega \text{ and } w \cdot n(x_0) < 0. \end{cases} \quad (6.1)$$

Here,  $\mathcal{P}_\gamma[u^\varepsilon]$  represents the weighted average over the outgoing boundary, i.e.,

$$\mathcal{P}_\gamma[u^\varepsilon](t, x_0) := c_\gamma \int_{w \cdot n > 0} u^\varepsilon(t, x_0, w) (w \cdot n) dw, \quad (6.2)$$

where the constant  $c_\gamma = \frac{1}{\pi}$  satisfies the normalization condition  $c_\gamma \int_{w \cdot n > 0} (w \cdot n) dw = 1$ . In addition, for the perturbative boundary data  $h$ , we assume the compatibility condition (1.6) to make sure  $u^\varepsilon$  has zero flux (1.7) at the boundary.

**6.1. Asymptotic Analysis.** By analogy with what has gone in Section 2, we expand the exact solution as

$$u^\varepsilon = U + U^I + \mathcal{R} = (U_0 + \varepsilon U_1 + \varepsilon^2 U_2) + (U_0^I + \varepsilon U_1^I) + \mathcal{R}. \quad (6.3)$$

Here, the initial layer  $U^I := U_0^I + \varepsilon U_1^I$  and the corresponding  $\Theta_{0,\infty}, \Theta_{1,\infty}$  can be taken the same as the in-flow case. To determine the interior solutions, as noted in [11], we should instead enforce the Neumann boundary condition here. Specifically,  $U_0$  is solved from

$$\begin{cases} U_0 = \bar{U}_0, \quad \partial_t \bar{U}_0 - \Delta_x \bar{U}_0 = 0, \\ \bar{U}_0(0, x) = \bar{u}_o(x), \\ \frac{\partial \bar{U}_0}{\partial n}(t, x_0) = 0 \quad \text{for } x_0 \in \partial\Omega. \end{cases} \quad (6.4)$$

Then  $U_1$  is solved from

$$\begin{cases} U_1 = \bar{U}_1 - w \cdot \nabla_x U_0, \quad \partial_t \bar{U}_1 - \Delta_x \bar{U}_1 = 0, \\ \bar{U}_1(0, x) = \Theta_{1,\infty}(x), \\ \frac{\partial \bar{U}_1}{\partial n}(t, x_0) = 0 \quad \text{for } x_0 \in \partial\Omega, \end{cases} \quad (6.5)$$

and  $U_2$  from

$$\begin{cases} U_2 = \bar{U}_2 - w \cdot \nabla_x U_1 - \partial_t U_0, \quad \partial_t \bar{U}_2 - \Delta_x \bar{U}_2 = 0, \\ \bar{U}_2(0, x) = 0, \\ \frac{\partial \bar{U}_2}{\partial n}(t, x_0) = 0 \quad \text{for } x_0 \in \partial\Omega. \end{cases} \quad (6.6)$$

Note that we do not need to introduce boundary layers here because the leading-order boundary condition is automatically satisfied by the interior solution  $U_0$  and the next-order boundary trace can be controlled in the remainder estimate.

Moreover, under the assumption (1.24), the well-posedness and corresponding regularity estimates as (2.50)–(2.54) in Proposition 2.3 still hold.

**6.2. Remainder Estimate.** With a view to achieving the same approximation estimate (1.26), we look at the remainder

$$\mathcal{R} := u^\varepsilon - \left[ (U_0 + \varepsilon U_1 + \varepsilon^2 U_2) + (U_0^I + \varepsilon U_1^I) \right], \quad (6.7)$$

and derive the initial-boundary value problem

$$\begin{cases} \varepsilon \partial_t \mathcal{R} + w \cdot \nabla_x \mathcal{R} + \varepsilon^{-1} (\mathcal{R} - \overline{\mathcal{R}}) = \mathcal{S} & \text{in } \mathbb{R}^+ \times \Omega \times \mathbb{S}^2, \\ \mathcal{R}(0, x, w) = \mathcal{I}(x, w) & \text{in } \Omega \times \mathbb{S}^2, \\ \mathcal{R}(t, x_0, w) = \mathcal{P}_\gamma[\mathcal{R}](t, x_0) + \mathcal{H}(t, x_0, w) & \text{for } x_0 \in \partial\Omega \text{ and } w \cdot n(x_0) < 0, \end{cases} \quad (6.8)$$

where

$$\mathcal{S} := -\varepsilon w \cdot \nabla_x U_1^I - \varepsilon^2 w \cdot \nabla_x U_2 - \varepsilon^2 \partial_t U_1 - \varepsilon^3 \partial_t U_2, \quad (6.9)$$

$$\mathcal{I} := \varepsilon^2 w \cdot \nabla_x U_1|_{t=0} + \varepsilon^2 \partial_t U_0|_{t=0}, \quad (6.10)$$

$$\mathcal{H} := \varepsilon h - \varepsilon(1 - \mathcal{P}_\gamma)[U_1^I]|_{\Gamma_-} + \varepsilon(1 - \mathcal{P}_\gamma)[w \cdot \nabla_x U_0]|_{\Gamma_-} + \varepsilon^2(1 - \mathcal{P}_\gamma)[w \cdot \nabla_x U_1]|_{\Gamma_-}, \quad (6.11)$$

with the parallel estimates as in Lemma 3.1:

$$\|\mathcal{S}\|_{L^2} \lesssim \varepsilon^2, \quad \|\mathcal{I}\|_{L^2} \lesssim \varepsilon^2, \quad \|\mathcal{H}\|_{L_{\mathbb{F}_-}^2} \lesssim \varepsilon. \quad (6.12)$$

Also, it can be verified that

$$\iint_{\Omega \times \mathbb{S}^2} \mathcal{I}(x, w) dw dx = 0, \quad \iint_{\Omega \times \mathbb{S}^2} \mathcal{S}(t, x, w) dw dx = 0 \quad \text{for all } t \in \mathbb{R}^+, \quad (6.13)$$

$$\int_{w \cdot n < 0} \mathcal{H}(t, x_0, w)(w \cdot n) dw = 0 \quad \text{for all } x_0 \in \partial\Omega \text{ and } t \in \mathbb{R}^+, \quad (6.14)$$

which yields

$$\int_{\Omega} \overline{\mathcal{R}}(t, x) dx = 0 \quad \text{for all } t \in \mathbb{R}^+ \quad (6.15)$$

and

$$\int_{\mathbb{S}^2} \mathcal{R}(t, x_0, w)(w \cdot n) dw = 0 \quad \text{for all } x_0 \in \partial\Omega \text{ and } t \in \mathbb{R}^+. \quad (6.16)$$

Observe that by the orthogonality  $\langle \mathcal{P}_\gamma[\mathcal{R}], (1 - \mathcal{P}_\gamma)[\mathcal{R}] \rangle_{\gamma_+} = 0$  and the compatibility (6.14), the boundary integral equals

$$\int_{\Gamma} \mathcal{R}^2(w \cdot n) = \|\mathcal{R}\|_{L_{\mathbb{F}_+}^2}^2 - \|\mathcal{P}_\gamma[\mathcal{R}] + \mathcal{H}\|_{L_{\mathbb{F}_-}^2}^2 = \|(1 - \mathcal{P}_\gamma)[\mathcal{R}]\|_{L_{\mathbb{F}_+}^2}^2 - \|\mathcal{H}\|_{L_{\mathbb{F}_-}^2}^2. \quad (6.17)$$

Correspondingly, we have the energy bound

$$\|\mathcal{R}\|_{L_t^\infty L_{x,w}^2}^2 + \varepsilon^{-1} \|(1 - \mathcal{P}_\gamma)[\mathcal{R}]\|_{L_{\mathbb{F}_+}^2}^2 + \varepsilon^{-2} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}^2 \lesssim \delta \varepsilon^{-1} \|\overline{\mathcal{R}}\|_{L^2}^2 + \delta^{-1} \quad (6.18)$$

with a small constant  $0 < \delta \ll 1$ .

Paralleling Lemma 4.3, we may deduce the kernel estimate

$$\|\overline{\mathcal{R}}\|_{L^2}^2 \lesssim \varepsilon \|\mathcal{R}\|_{L_t^\infty L_{x,w}^2}^2 + \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2}^2 + \|(1 - \mathcal{P}_\gamma)[\mathcal{R}]\|_{L_{\mathbb{F}_+}^2}^2 + \varepsilon, \quad (6.19)$$

by using test functions built from  $\xi(t, x) := (-\Delta_x)^{-1} \overline{\mathcal{R}}(t, x)$  with the Neumann boundary condition instead. To be specific, (6.15) allows us to construct  $\xi$  via

$$\begin{cases} -\Delta_x \xi = \overline{\mathcal{R}} & \text{in } \Omega, \\ \frac{\partial \xi}{\partial n} = 0 & \text{on } \partial\Omega, \\ \int_{\Omega} \xi = 0. \end{cases} \quad (6.20)$$

Consequently, we see the boundary terms

$$\int_{\Gamma} \mathcal{R} \xi (w \cdot n) = \langle \mathcal{R}, \xi \rangle_{\Gamma_+} - \langle \mathcal{P}_\gamma[\mathcal{R}] + \mathcal{H}, \xi \rangle_{\Gamma_-} = 0 \quad (6.21)$$

from (6.16), and

$$\begin{aligned} \int_{\Gamma} \mathcal{R} (w \cdot \nabla_x \xi) (w \cdot n) &= \langle \mathcal{R}, w \cdot \nabla_x \xi \rangle_{\Gamma_+} - \langle \mathcal{P}_\gamma[\mathcal{R}] + \mathcal{H}, w \cdot \nabla_x \xi \rangle_{\Gamma_-} \\ &= \langle (1 - \mathcal{P}_\gamma)[\mathcal{R}], w \cdot \nabla_x \xi \rangle_{\Gamma_+} - \langle \mathcal{H}, w \cdot \nabla_x \xi \rangle_{\Gamma_-}, \end{aligned} \quad (6.22)$$

thanks to  $\frac{\partial \xi}{\partial n} \Big|_{\partial \Omega} = (n \cdot \nabla_x \xi) \Big|_{\partial \Omega} = 0$ . The remaining proof of (6.19) then follows from analogous arguments.

Combining (6.18) with (6.19) yields the remainder estimate

$$\|\mathcal{R}\|_{L_t^\infty L_{x,w}^2} + \varepsilon^{-\frac{1}{2}} \|(1 - \mathcal{P}_\gamma)[\mathcal{R}]\|_{L_{\Gamma_+}^2} + \varepsilon^{-\frac{1}{2}} \|\overline{\mathcal{R}}\|_{L^2} + \varepsilon^{-1} \|\mathcal{R} - \overline{\mathcal{R}}\|_{L^2} \lesssim 1, \quad (6.23)$$

and Theorem 1.2 can thus be proved accordingly.

## 7. SPECULAR-REFLECTION BOUNDARY PROBLEM

Consider the unsteady neutron transport equation with the specular-reflection boundary condition:

$$\begin{cases} \varepsilon \partial_t u^\varepsilon + w \cdot \nabla_x u^\varepsilon + \varepsilon^{-1} (u^\varepsilon - \overline{u}^\varepsilon) = 0 & \text{in } \mathbb{R}^+ \times \Omega \times \mathbb{S}^2, \\ u^\varepsilon(0, x, w) = u_o(x, w) & \text{in } \Omega \times \mathbb{S}^2, \\ u^\varepsilon(t, x_0, w) = \mathcal{R}_\gamma[u^\varepsilon](t, x_0, w) + \varepsilon h(t, x_0, w) & \text{for } x_0 \in \partial \Omega \text{ and } w \cdot n(x_0) < 0, \end{cases} \quad (7.1)$$

where the specular-reflection operator

$$\mathcal{R}_\gamma[u^\varepsilon](t, x_0, w) := u^\varepsilon(t, x_0, \mathcal{R}w) \quad \text{with} \quad \mathcal{R}w := w - 2(w \cdot n)n, \quad (7.2)$$

and  $h$  is a given perturbation satisfying the compatibility condition (1.6), so that  $u^\varepsilon$  has zero flux at each boundary point.

**7.1. Asymptotic Analysis.** Let us expand the exact solution as

$$u^\varepsilon = U + U^I + U^B + \mathcal{R} = (U_0 + \varepsilon U_1 + \varepsilon^2 U_2) + (U_0^I + \varepsilon U_1^I) + (U_0^B + \varepsilon U_1^B) + \mathcal{R}. \quad (7.3)$$

Here, the initial layer  $U^I := U_0^I + \varepsilon U_1^I$  and the interior solution  $U := U_0 + \varepsilon U_1 + \varepsilon^2 U_2$  can be constructed identically to the diffuse-reflection case.

However, unlike the previous scenario, some kind of ‘‘boundary layer’’ need to be introduced (if  $h \neq 0$ ) in order to guarantee the perfect specular boundary condition for the remainder. Specifically, we let  $U_0^B = 0$  because the leading-order interior solution  $U_0$  already satisfies the specular boundary condition. We then define  $U_1^B$  to be an extension of  $h$  into a thin layer of the interior of domain:

$$U_1^B = E[h](t; \eta, \iota_1, \iota_2; \phi, \psi) := \chi(\eta) \mathbb{1}_{\{\sin \phi > 0\}} h(t; \iota_1, \iota_2; \phi, \psi), \quad (7.4)$$

and let it act as the next-order boundary layer.

**7.2. Remainder Estimate.** Proceeding in a parallel fashion, we look at the remainder

$$\mathcal{R} := u^\varepsilon - \left[ (U_0 + \varepsilon U_1 + \varepsilon^2 U_2) + (U_0^I + \varepsilon U_1^I) + \varepsilon U_1^B \right], \quad (7.5)$$

and derive the initial-boundary value problem

$$\begin{cases} \varepsilon \partial_t \mathcal{R} + w \cdot \nabla_x \mathcal{R} + \varepsilon^{-1} (\mathcal{R} - \overline{\mathcal{R}}) = \mathcal{S} & \text{in } \mathbb{R}^+ \times \Omega \times \mathbb{S}^2, \\ \mathcal{R}(0, x, w) = \mathcal{I}(x, w) & \text{in } \Omega \times \mathbb{S}^2, \\ \mathcal{R}(t, x_0, w) = \mathcal{R}(t, x_0, \mathcal{R}w) & \text{for } x_0 \in \partial \Omega \text{ and } w \cdot n(x_0) < 0. \end{cases} \quad (7.6)$$

In particular,  $\mathcal{R}$  satisfies the perfect specular boundary condition without any perturbation, due to the introduction of the ‘‘boundary layer’’  $U_1^B$  and the observation that

$$U_0 = \mathcal{R}_\gamma[U_0], \quad U_1 = \mathcal{R}_\gamma[U_1], \quad U_2 = \mathcal{R}_\gamma[U_2], \quad U_0^I = \mathcal{R}_\gamma[U_0^I], \quad U_1^I = \mathcal{R}_\gamma[U_1^I] \quad (7.7)$$

from the construction of  $U_0, U_1, U_2, U_0^I, U_1^I$  and the compatibility conditions (1.29).

Compared to the diffuse-reflection case, the expression of  $\mathcal{I}$  remains the same, while the source term  $\mathcal{S}$  now contains additional terms coming from  $U_1^B$  (see (2.26) for the transport operator under the boundary coordinates):

$$\mathcal{S}^{BL} := -\varepsilon^2 \partial_t U_1^B - \varepsilon w \cdot \nabla_x U_1^B - (U_1^B - \overline{U_1^B}) =: \mathcal{S}_a^{BL} + \mathcal{S}_b^{BL}. \quad (7.8)$$

Here, let  $\mathcal{S}_a^{BL} := \varepsilon(\mathcal{S}_1^{BL} + \mathcal{S}_2^{BL})[U_1^B]$  denote the terms corresponding to  $\mathcal{S}_1^{BL} + \mathcal{S}_2^{BL}$  in the inflow case, which now raise by one order of  $\varepsilon$  and so  $\| (1 + \eta) \mathcal{S}_a^{BL} \|_{L^2} \lesssim \varepsilon^{\frac{3}{2}}$ . In particular, the term containing  $\partial_\phi U_1^B$  is still good due to the continuity of  $E[h]$  across the grazing set where  $w \cdot n = 0$  under assumption (1.28) with the compatibility condition  $h|_{\Gamma_0} = 0$ . The remaining part

$$\mathcal{S}_b^{BL} := -\sin \phi \partial_\eta U_1^B - (U_1^B - \overline{U_1^B}) = -\chi'(\eta) \sin \phi \mathbf{1}_{\{\sin \phi > 0\}} h - \chi(\eta) \left( \mathbf{1}_{\{\sin \phi > 0\}} h - \overline{\mathbf{1}_{\{\sin \phi > 0\}} h} \right) \quad (7.9)$$

which corresponds to  $\mathcal{S}_3^{BL}$  in the inflow case now has no cancellation to exploit since it does not satisfy the Milne problem. In view of the scaling  $\eta = \varepsilon^{-1} \mu$ , we see  $\| U_1^B \|_{L^2} \lesssim \varepsilon^{\frac{1}{2}}$  and those terms in  $\mathcal{S}_b^{BL}$  contribute to

$$\| (1 + \eta) \mathcal{S}^{BL} \|_{L^2} \lesssim \varepsilon^{\frac{1}{2}}, \quad (7.10)$$

which is acceptable.

Besides, due to the presence of  $U_1^B$ , we have

$$\iint_{\Omega \times \mathbb{S}^2} \mathcal{S} = -\varepsilon^2 \iint_{\Omega \times \mathbb{S}^2} \partial_t U_1^B, \quad \text{and so} \quad \int_{\Omega} \overline{\mathcal{R}} = -\varepsilon \iint_{\Omega \times \mathbb{S}^2} U_1^B, \quad (7.11)$$

considering the null flux of  $\mathcal{R}$  and  $U_1^B$  under (1.6). Despite the fact that  $\overline{\mathcal{R}}$  does not necessarily have zero average, we may redefine

$$\tilde{\mathcal{R}} := \mathcal{R} - \frac{1}{|\Omega|} \int_{\Omega} \overline{\mathcal{R}}, \quad \tilde{\mathcal{S}} := \mathcal{S} - \frac{1}{4\pi|\Omega|} \iint_{\Omega \times \mathbb{S}^2} \mathcal{S}, \quad (7.12)$$

so that  $\int_{\Omega} \tilde{\mathcal{R}} = 0$ ,  $\iint_{\Omega \times \mathbb{S}^2} \tilde{\mathcal{S}} = 0$ , and that  $\tilde{\mathcal{R}}$  solves (7.6) with  $\mathcal{S}$  replaced by  $\tilde{\mathcal{S}}$ .

Noticing that the difference  $|\int_{\Omega} \overline{\mathcal{R}}| \lesssim \varepsilon^2$  has higher order of smallness, we only need to obtain the remainder estimate for  $\tilde{\mathcal{R}}$ . For simplicity, we will abuse notation  $\mathcal{R}$  for  $\tilde{\mathcal{R}}$  in what follows.

The specular boundary condition on  $\mathcal{R}$  yields

$$\int_{\mathbb{S}^2} \mathcal{R}(t, x_0, w)(w \cdot n) dw = 0 \quad \text{for all } x_0 \in \partial\Omega \text{ and } t \in \mathbb{R}^+, \quad (7.13)$$

as well as

$$\int_{\Gamma} \mathcal{R}^2(w \cdot n) = \|\mathcal{R}\|_{L^2_{\Gamma_+}}^2 - \|\mathcal{R}\|_{L^2_{\Gamma_-}}^2 = 0. \quad (7.14)$$

Also, with the same test functions as in the diffuse-reflection case where  $\frac{\partial \xi}{\partial n} = 0$  on  $\partial\Omega$ , the corresponding boundary terms will vanish:

$$\int_{\Gamma} \mathcal{R} \xi(w \cdot n) = \langle \mathcal{R}, \xi \rangle_{\Gamma_+} - \langle \mathcal{R}, \xi \rangle_{\Gamma_-} = 0, \quad (7.15)$$

$$\int_{\Gamma} \mathcal{R}(w \cdot \nabla_x \xi)(w \cdot n) = \langle \mathcal{R}, w \cdot \nabla_x \xi \rangle_{\Gamma_+} - \langle \mathcal{R}, w \cdot \nabla_x \xi \rangle_{\Gamma_-} = 0. \quad (7.16)$$

Arguing analogously, we thus find the energy bound

$$\|\mathcal{R}\|_{L_t^\infty L_{x,w}^2}^2 + \varepsilon^{-2} \|\|\mathcal{R} - \overline{\mathcal{R}}\|\|_{L^2}^2 \lesssim \delta \varepsilon^{-1} \|\|\overline{\mathcal{R}}\|\|_{L^2}^2 + \delta^{-1}, \quad (7.17)$$

and the kernel bound

$$\|\|\overline{\mathcal{R}}\|\|_{L^2}^2 \lesssim \varepsilon \|\|\mathcal{R}\|\|_{L_t^\infty L_{x,w}^2}^2 + \|\|\mathcal{R} - \overline{\mathcal{R}}\|\|_{L^2}^2 + \varepsilon. \quad (7.18)$$

For the kernel estimate, however, the fresh terms  $\varepsilon^{-1} |\langle \mathcal{S}^{BL}, \xi \rangle| + |\langle \mathcal{S}^{BL}, w \cdot \nabla_x \xi \rangle| + \varepsilon |\langle \mathcal{S}^{BL}, \zeta \rangle|$  in (4.36) require a different treatment, especially for the worst part  $\varepsilon^{-1} |\langle \mathcal{S}_b^{BL}, \xi \rangle|$ : note that the estimate like (4.38) no longer applies since now we do not have  $\xi|_{\partial\Omega} = 0$ . Instead, using integration by parts in  $\mu$  and noticing

that the boundary term vanishes due to  $\int_{\mathbb{S}^2} \sin \phi U_1^B \Big|_{\mu=0} dw = \int_{w \cdot n < 0} h(t, x_0, w)(w \cdot n) dw = 0$  for each  $x_0 \in \partial\Omega$  and all  $t \in \mathbb{R}^+$  (which is precisely the compatibility condition (1.6)), we find

$$\begin{aligned} \varepsilon^{-1} \left| \langle \mathcal{S}_b^{BL}, \xi \rangle \right| &= \varepsilon^{-1} \left| \langle \sin \phi \partial_\eta U_1^B + (U_1^B - \overline{U_1^B}), \xi \rangle \right| = \left| \langle \sin \phi \partial_\mu U_1^B, \xi \rangle \right| = \left| \langle \sin \phi U_1^B, \partial_\mu \xi \rangle \right| \quad (7.19) \\ &\lesssim \left\| U_1^B \right\|_{L_t^2 L_x^2 L_w^1} \left\| \partial_\mu \xi \right\|_{L^2} \lesssim \varepsilon^{\frac{1}{2}} \left\| \xi \right\|_{L_t^2 H_x^1} \lesssim \varepsilon^{\frac{1}{2}} \left\| \overline{\mathcal{R}} \right\|_{L^2} \lesssim \delta \left\| \overline{\mathcal{R}} \right\|_{L^2}^2 + \delta^{-1} \varepsilon. \end{aligned}$$

The rest of the terms can be bounded either in a similar way as (4.38)–(4.40), or even more directly considering the higher order of  $\varepsilon$  they possess.

Finally, by combining (7.17) and (7.18) we have the remainder estimate

$$\left\| \mathcal{R} \right\|_{L_t^\infty L_{x,w}^2} + \varepsilon^{-\frac{1}{2}} \left\| \overline{\mathcal{R}} \right\|_{L^2} + \varepsilon^{-1} \left\| \mathcal{R} - \overline{\mathcal{R}} \right\|_{L^2} \lesssim 1. \quad (7.20)$$

This leads to the proof of (1.30) and so that of Theorem 1.3.

## REFERENCES

- [1] G. BAL, *Inverse transport theory and applications*, Inverse Problems, 25 (2009), p. 053001.
- [2] G. BAL AND K. REN, *On multi-spectral quantitative photoacoustic tomography in diffusive regime*, Inverse Problems, 28 (2012), p. 025010.
- [3] C. BARDOS, F. GOLSE, AND B. PERTHAME, *The Rosseland approximation for the radiative transfer equations*, Comm. Pure Appl. Math., 40 (1987), pp. 69–721.
- [4] C. BARDOS, F. GOLSE, B. PERTHAME, AND R. SENTIS, *The nonaccretive radiative transfer equations: existence of solutions and Rosseland approximation*, J. Funct. Anal., 77 (1988), pp. 434–460.
- [5] C. BARDOS, R. SANTOS, AND R. SENTIS, *Diffusion approximation and computation of the critical size*, Trans. Amer. Math. Soc., 284 (1984), pp. 617–649.
- [6] A. BENSOUSSAN, J.-L. LIONS, AND G. C. PAPANICOLAOU, *Boundary layers and homogenization of transport processes*, Publ. Res. Inst. Math. Sci., 15 (1979), pp. 53–157.
- [7] B. COCKBURN, S. HOU, AND C.-W. SHU, *The Runge-Kutta local projection discontinuous Galerkin finite element method for conservation laws IV. the multidimensional case*, Math. Comp., 54 (1990), pp. 545–581.
- [8] B. COCKBURN, G. E. KARNIADAKIS, AND C.-W. SHU, *The development of discontinuous galerkin methods*, in Discontinuous Galerkin methods (Newport, RI, 1999), Berlin, 2000, Springer.
- [9] B. COCKBURN AND C.-W. SHU, *TVB Runge-Kutta local projection discontinuous Galerkin finite element method for conservation laws II. general framework*, Math. Comp., 52 (1989), pp. 411–435.
- [10] R. ESPOSITO, Y. GUO, C. KIM, AND R. MARRA, *Non-isothermal boundary in the Boltzmann theory and Fourier law*, Comm. Math. Phys., 323 (2013), pp. 177–239.
- [11] Y. GUO AND L. WU, *Geometric correction in diffusive limit of neutron transport equation in 2D convex domains*, Arch. Rational Mech. Anal., 226 (2017), pp. 321–403.
- [12] ———, *Regularity of Milne problem with geometric correction in 3D*, Math. Models Methods Appl. Sci., 27 (2017), pp. 453–524.
- [13] ———,  *$L^2$  diffusive expansion for neutron transport equation*, arXiv:2301.11996, (2023).
- [14] N. V. KRYLOV, *Lectures on elliptic and parabolic equations in Sobolev spaces.*, American Mathematical Society, Providence, RI, 2008.
- [15] E. W. LARSEN, *A functional-analytic approach to the steady, one-speed neutron transport equation with anisotropic scattering*, Comm. Pure Appl. Math., 27 (1974), pp. 523–545.
- [16] ———, *Solutions of the steady, one-speed neutron transport equation for small mean free paths*, J. Mathematical Phys., 15 (1974), pp. 299–305.
- [17] ———, *Neutron transport and diffusion in inhomogeneous media I*, J. Mathematical Phys., 16 (1975), pp. 1421–1427.
- [18] ———, *Asymptotic theory of the linear transport equation for small mean free paths II*, SIAM J. Appl. Math., 33 (1977), pp. 427–445.
- [19] E. W. LARSEN AND J. D’ARRUDA, *Asymptotic theory of the linear transport equation for small mean free paths I*, Phys. Rev., 13 (1976), pp. 1933–1939.
- [20] E. W. LARSEN AND G. J. HABETLER, *A functional-analytic derivation of Case’s full and half-range formulas*, Comm. Pure Appl. Math., 26 (1973), pp. 525–537.
- [21] E. W. LARSEN AND J. B. KELLER, *Asymptotic solution of neutron transport problems for small mean free paths*, J. Mathematical Phys., 15 (1974), pp. 75–81.
- [22] E. W. LARSEN AND P. F. ZWEIFEL, *On the spectrum of the linear transport operator*, J. Mathematical Phys., 15 (1974), pp. 1987–1997.
- [23] ———, *Steady, one-dimensional multigroup neutron transport with anisotropic scattering*, J. Mathematical Phys., 17 (1976), pp. 1812–1820.
- [24] Q. LI, J. LU, AND W. SUN, *Diffusion approximations and domain decomposition method of linear transport equations: asymptotics and numerics*, J. Comput. Phys., 292 (2015), pp. 141–167.
- [25] ———, *Half-space kinetic equations with general boundary conditions*, Math. Comp., 86 (2017), pp. 1269–1301.

- [26] ———, *Validity and regularization of classical half-space equations*, J. Stat. Phys., 166 (2017), pp. 398–433.
- [27] L. WU, *Diffusive limit with geometric correction of unsteady neutron transport equation*, Kinet. Relat. Models, 10 (2017), pp. 1163–1203.
- [28] ———, *Asymptotic analysis of unsteady neutron transport equation*, Math. Methods Appl. Sci., 42 (2019), pp. 2544–2585.
- [29] ———, *Diffusive limit of transport equation in 3D convex domains*, Peking Math. J., 4 (2021), pp. 203–284.
- [30] L. WU AND Y. GUO, *Geometric correction for diffusive expansion of steady neutron transport equation*, Comm. Math. Phys., 336 (2015), pp. 1473–1553.
- [31] L. WU, X. YANG, AND Y. GUO, *Asymptotic analysis of transport equation in annulus*, J. Stat. Phys., 165 (2016), pp. 585–644.

(Z. Ouyang)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO

*Email address:* `ouyangzm9386@uchicago.edu`