

A groupoid approach to the Wodzicki residue

Nathan Couchet¹ and Robert Yuncken^{* 2}

¹Université Clermont Auvergne, CNRS, LMBP, F-63000 Clermont-Ferrand, France, nathan.couchet@uca.fr

²Institut Élie Cartan de Lorraine, Université de Lorraine, CNRS, IECL, F-57000 Metz, France, robert.yuncken@univ-lorraine.fr

KEYWORDS: Wodzicki residue, pseudodifferential calculus, classical symbol, tangent groupoid, filtered manifold, non-commutative geometry.

MSC: Primary: 47G30; secondary: 22A22, 35S05, 58H05, 58J40.

Abstract

Originally, the noncommutative residue was studied in the 80's by Wodzicki in his thesis [31] and Guillemin [16]. In this article we give a definition of the Wodzicki residue, using the language of r -fibered distributions from [22], [28], in the context of filtered manifolds. We show that this groupoidal residue behaves like a trace on the algebra of pseudodifferential operators on filtered manifolds and coincides with the usual residue Wodzicki in the case where the manifold is trivially filtered. Moreover, in the context of Heisenberg calculus, we show that the groupoidal residue coincides with Ponge's definition [23] for contact and codimension 1 foliation Heisenberg manifolds.

1 Introduction.

One of the remarkable features of the theory of pseudodifferential operators is the noncommutative residue of Wodzicki, which he defined in 1984 in his thesis [32]. The noncommutative residue was also defined in 1985 in a article of Guillemin, [16, Definition 6 p 151], in which he proposes to associate to an operator P a zeta function:

$$\zeta(P, s) = \sum_k \lambda_k^s, \quad (1.1)$$

where λ_k are the eigenvalues of P , and then shows that it admits a meromorphic continuation. The residues of this zeta function are linked to the number of eigenvalues of P denoted by $N(\lambda) = \text{card}\{\lambda_k, \lambda_k \leq \lambda\}$, leading Guillemin to a Weyl-type formula, as in [30].

Let P be a classical pseudodifferential operator of order $m \in \mathbb{Z}$ on a manifold M of dimension d . This means that in any chart the symbol of P admits an asymptotic expansion:

$$a(x, \xi) \sim \sum_k a_{m-k}(x, \xi). \quad (1.2)$$

In the following discussion, we fix a chart (U, ϕ) and we identify U with its image in \mathbb{R}^d .

Definition 1.1. [29, p 58] , [21, 1.8 Formule locale], [31, § 7]

*This article/publication is based upon work from COST Action CaLISTA CA21109 supported by COST (European Cooperation in Science and Technology). www.cost.eu.

We define the residue at x of a classical pseudodifferential operator P on M of order m to be:

$$Res_x^W(P) = \frac{1}{(2\pi)^d} \left(\int_{\mathbb{S}^{d-1}} a_{-d}(x, \xi) d\sigma(\xi) \right) dx, \quad (1.3)$$

where $a_{-d}(x, \xi)$ is the homogeneous part of order $-d$ in the variable ξ coming from the asymptotic expansion of P in any chart. Moreover, if M is a compact riemannian manifold, we define also the (global) residue of P :

$$Res^W(P) = \int_M Res_x^W(P) dx, \quad (1.4)$$

where dx is the smooth measure coming from the riemannian structure.

It is a profound theorem that these quantities are independent of the choice of chart. In this paper, we propose to give another definition of this residue for operators of order $-dim(M)$, which extends naturally to filtered manifolds, using the groupoidal calculus from [28] — that is using tangent groupoids $\mathbb{T}M, \mathbb{T}_H M$. We therefore manage to circumvent the chart machinery. Moreover our definition extends essentially without change to define non commutative residue on pseudodifferential operator on any filtered manifold, including that of Ponge on a Heisenberg manifold [23], using the filtered tangent groupoid of [5],[27]. Note that the compact hypothesis is to make (1.4) valid, we can define Res_x^W for M non-compact.

The noncommutative residue has proved to be connected with other geometrical objects. For instance :

1. There is link between this residue for a differential operator on a compact manifold and the asymptotic expansion of the trace of the heat operator e^{-tP} , see [1].
2. Connes showed in 1988 - see [7, Proposition 5 p313], [4, section 2.6 p17], [20, section 7.6], [25, Proposition 4.11 p 16] - that when P is a pseudodifferential operator of order $-dim(M)$ and M is compact, then the Dixmier trace of P coincides to this residue up to a constant.
3. Ponge showed in [23, Proposition 6.3 p 454] the link between the residue of the Kohn-Laplacian \square_b on a CR compact manifold, see [23, Equation (6.9)], and the volume of M , defined at [23, Equation (6.4)].

Let us very briefly recall the groupoid approach to pseudodifferential operators, first observed by Debord-Skandalis [11] in 2014 and developed by van Erp and the second author some years later [28]. The tangent groupoid of Connes is:

$$\mathbb{T}M = M \times M \times \mathbb{R}^* \bigcup TM \times \{0\}, \quad (1.5)$$

which is seen as a smooth glueing of the tangent bundle TM with a family of pair groupoids $M \times M$ over \mathbb{R}^* . In the filtered case the appropriate substitute of the tangent bundle TM is a bundle of nilpotent osculating groups $\mathcal{T}_H M$ whose fibers are denoted by $\mathcal{T}_H M_x$ and we define the filtered tangent groupoid by:

$$\mathbb{T}_H M = M \times M \times \mathbb{R}^* \bigcup \mathcal{T}_H M \times \{0\}. \quad (1.6)$$

The bundle of osculating groups admits a family of automorphism $(\delta_s)_{s>0}$ called dilations which generalises the homotheties on $\mathbb{T}M$ in the trivially-filtered case.

Using these dilations we obtain a smooth \mathbb{R}_+^* -action on $\mathbb{T}_H M$, namely the Debord-Skandalis action [11], see also [28] where it is called the “zoom action” and [9]:

Definition 1.2. Let M be a filtered manifold. We define the Debord-Skandalis action of \mathbb{R}_+^* on $\mathbb{T}_H M$, $s \in \mathbb{R}_+^* \mapsto \alpha_s \in \text{Aut}(\mathbb{T}_H M)$ by:

$$\begin{cases} \alpha_s(y, x, t) = (y, x, s^{-1}t) & (x, y) \in M, \\ \alpha_s(x, \xi, 0) = (x, \delta_s(\xi), 0) & x \in M, \xi \in \mathcal{T}_H M_x. \end{cases} \quad (1.7)$$

The key point in the groupoidal approach to pseudodifferential operators [28] is to consider the set of distributions on the tangent groupoid $\mathbb{T}_H M$ which are essentially homogeneous for the Debord-Skandalis action in the following sense. A r -fibered distribution on $\mathbb{T}_H M$ is a continuous $C^\infty(M \times \mathbb{R})$ -linear map $\mathbb{k} : C^\infty(\mathbb{T}_H M) \rightarrow C^\infty(M \times \mathbb{R})$. This definition implies that the support of such a distribution is r -proper and we denote them by $\mathcal{E}'_r(\mathbb{T}_H M)$. They were first studied by Androulidakis-Skandalis [2] and Lescure-Manchon-Vassout [22] in the general case of submersions. Thanks to a result from [22], r -fibered distributions can be seen as smooth maps \mathbb{k} from $M \times \mathbb{R}$ to compactly supported distributions in the r -fibers, $\mathbb{k}(x, t) \in \mathcal{E}'(r^{-1}(x, t))$, and whose support is r -proper. We say that a r -fibered distribution is properly supported if its support is also s -proper.

Recall that $C_p^\infty(\mathbb{T}_H M, \Omega_r)$ denotes the proper smooth sections of the 1-density bundle tangent to the r -fibers, see [9, Definition 5.9]. We say that a properly supported r -fibered distribution \mathbb{k} is essentially homogeneous of order m for the Debord-Skandalis if:

$$s \in \mathbb{R}_+^* \mapsto s^{-m} \alpha_{s*} \mathbb{k} - \mathbb{k} \in C_p^\infty(\mathbb{T}_H M, \Omega_r). \quad (1.8)$$

We refer to the function appearing in (1.8) as the co-cycles of \mathbb{k} . The set of essentially homogeneous distributions of this kind is denoted $\Psi_{vEY}^m(\mathbb{T}_H M)$. We will refer to notations and concepts from [28] and [9, section 5.3]. Thanks to this, van Erp and the second author define a pseudodifferential operator as the restriction at $t = 1$ of an element $\mathbb{k} \in \Psi_{vEY}^m(\mathbb{T}_H M)$, see [28, Theorem 2 p3]. These elements are the H -pseudodifferential operators on the filtered manifold M and are denoted by $\Psi_H^m(M)$.

One of the main theorems of the article [28] is that the groupoidal calculus coincides, in the case of a trivially filtered manifold, with the classical calculus of Kohn-Nirenberg and Hörmander, namely:

$$\Psi_{\text{Hör}}^m(M) = \Psi_H^m(M). \quad (1.9)$$

A simple but important first observation of the present paper is the following. We will consider $m \in \mathbb{Z}$ in the whole paper. If M is a filtered manifold, we denote d_H the homogeneous dimension of M . Note that, if we fix a smooth measure dx on M , we obtain a canonical smooth family of 1-densities $d\lambda_x$ on the tangent fibers $T_x M$ and hence on the osculating groups $\mathcal{T}_H M_x$.

Lemma 1.3. *Let M be a filtered manifold and $\mathbb{k} \in \Psi_{vEY}^{-d_H}(\mathbb{T}_H M)$. For every $x \in M$, the function defined by:*

$$s \mapsto \left(s^{d_H} \alpha_{s*} \mathbb{k} - \mathbb{k} \right) |_{(x,0,0)}, \quad (1.10)$$

is a group homomorphism from (\mathbb{R}_+^, \times) to $(\mathbb{C}, +)$. More precisely, there exists a constant $r_x \in \mathbb{C}$ such that for all $s > 0$:*

$$\left(s^{d_H} \alpha_{s*} \mathbb{k} - \mathbb{k} \right) |_{(x,0,0)} = r_x \log(s) d\lambda_x. \quad (1.11)$$

We shall give the proof in the next section.

We may define the quantity $r_x dx$ to be the groupoidal residue of the pseudodifferential operator with kernel $\mathbb{k}|_{t=1}$, which is indeed a polyhomogeneous pseudodifferential operator by (1.9). As we shall see in Lemma 2.1, $r_x dx$ does not depend on the r -fibered distribution \mathbb{k} representing the pseudo-differential operator P at $t = 1$. Therefore we can define:

Definition 1.4. Let M be a filtered manifold of homogeneous dimension d_H and $P \in \Psi_H^m(M)$ with $m \leq -d_H$. Let $\mathbb{k} \in \Psi_{vEY}^{-d_H}(\mathbb{T}_H M)$ be an element in vEY groupoidal calculus such that $\mathbb{k}|_{t=1}$ is the Schwartz kernel of P . We define the groupoidal residue of P at $x \in M$, denoted $Res_x(P)$, for any $s \in \mathbb{R}_+^* \setminus \{1\}$ to be:

$$Res_x(P) := \frac{1}{\log(s)} \left(s^{d_H} \alpha_{s*} \mathbb{k} - \mathbb{k} \right) |_{(x,0,0)}. \quad (1.12)$$

The above definition is for scalar-valued operators. For operators between vector bundles, we should take, for any $s \in \mathbb{R}_+^* \setminus \{1\}$:

$$Res_x(P) := \frac{1}{\log(s)} \text{Tr} \left(s^{d_H} \alpha_{s*} \mathbb{k} - \mathbb{k} \right) |_{(x,0,0)}. \quad (1.13)$$

For details on the groupoidal calculus with vector-bundle coefficients, see [10]. We will restrict our attention to scalar-valued operators for simplicity.

Now come the main results of this paper:

1. We shall show, see Theorem 2.3, that similarly to the Wodzicki residue Res_x^W , the groupoidal residue Res_x from Definition 1.4 defines a trace on operators of appropriate order. More precisely if $P \in \Psi_H^m(M)$ where M is a filtered manifold and $m \leq -d_H$, $Q \in \Psi_H^0(M)$ then :

$$Res_x([P, Q]) = 0, \quad (1.14)$$

where $[,]$ denotes the commutator of operators.

2. We shall show, see Theorem 4.3, that the groupoidal residue from Definition 1.4 coincides with the Wodzicki residue in the case of a trivially filtered manifold.
3. We shall show, see Theorem 5.1 and Corollary 5.2, that the definition of the non-commutative residue made by Raphaël Ponge [23] in 2007 coincides with groupoidal residue from Definition 1.4 for \mathcal{V} -pseudo-differential operators in the calculus of BG, [3, section § 10] where M is a contact manifold or a foliation of codimension 1.

Definition 1.4 only applies to pseudo-differential operators of order $\leq -d_H$. It should be possible to extend this definition to operators of arbitrary order. In the case of the classical unfiltered calculus, this will be treated in a forthcoming article by Higson, Sukochev and Zanin [17]. We learnt of their work while writing this article. We refer to that article for details.

2 Basic properties of the Wodzicki residue on a filtered manifold.

Maintaining the notation of the introduction, let us start by proving Lemmas 1.3 and 2.1, which show the well-definedness of our residue. Our conventions for the filtered tangent groupoid are such that the range and source maps are given by :

$$\begin{cases} r(y, x, t) = y, & s(y, x, t) = x, & t \neq 0 \\ r(x, v, 0) = x, & s(x, v, 0) = x. \end{cases} \quad (2.1)$$

Proof of Lemma (1.3). Set:

$$F : s \in \mathbb{R}_+^* \mapsto F_s := \left(s^{d_H} \alpha_{s^*} \mathbb{k} - \mathbb{k} \right) \in C_p^\infty(\mathbb{T}_H M, \Omega_r). \quad (2.2)$$

We will show that it is a group morphism when restricted at $(x, 0, 0)$. First, the reader can easily check that:

$$F_{st} = s^{d_H} \alpha_{s^*} F_t + F_s, \quad s, t > 0. \quad (2.3)$$

Thus we need to show:

$$s^{d_H} \alpha_{s^*} F_t|_{(x,0,0)} = F_t|_{(x,0,0)}. \quad (2.4)$$

We are therefore going to prove (2.4). In the fiber $(x, \cdot, 0)$, we can write:

$$F_s|_{(x, \cdot, 0)} = f_x d\lambda_x, \quad (2.5)$$

for some $f_x \in C_c^\infty(\mathcal{T}_H M_x)$ and where $d\lambda_x$ is a Haar measure on the nilpotent (graded) osculating group $\mathcal{T}_H M_x$ such that $\delta_{s^*}(d\lambda_x) = s^{-d_H} d\lambda_x$. The Debord-Skandalis action at $t = 0$ acts by the dilations δ_s . So for the fiber $(x, \cdot, 0)$ we have:

$$\alpha_{s^*}(F_s)|_{(x, \cdot, 0)} = \delta_{s^*}(F_s|_{(x, \cdot, 0)}) \quad (2.6)$$

$$\underbrace{=}_{(2.5)} \left((\delta_s^{-1})^* f_x \right) \delta_{s^*}(d\lambda_x) \quad (2.7)$$

$$= s^{-d_H} \left((\delta_s^{-1})^* f_x \right) d\lambda_x. \quad (2.8)$$

Since the Debord-Skandalis action α_s fixes the points $(x, 0, 0)$, this gives $s^{d_H} \alpha_{s^*} F_t|_{(x,0,0)} = F_t|_{(x,0,0)}$ as desired. Therefore, $s \mapsto F_s|_{(x,0,0)}$ is a morphism from (\mathbb{R}_+^*, \times) to $(\mathbb{C}, +)$. Moreover, note that the map from (2.2) is smooth, see [28, Lemma 21]. \square

Lemma 2.1. *Let M be a filtered manifold and $P \in \Psi_H^m(M)$, with $m \leq -d_H$. The Definition 1.4 of the groupoidal residue does not depend on the r -fibered distribution \mathbb{k} that represents P .*

Proof. Let $\mathbb{k}, \mathbb{k}' \in \Psi_{vEY}^{-d_H}(\mathbb{T}_H M)$ such that:

$$\mathbb{k}|_{t=1} = P, \quad \mathbb{k}'|_{t=1} = P. \quad (2.9)$$

Let $x \in M$ be fixed. We need to show that the two co-cycles (1.12) related to these r -fibered distributions give the same groupoidal residue at x . Thanks to [28, Corollary 33], we have:

$$\mathbb{k}|_{t=0} - \mathbb{k}'|_{t=0} \in C_p^\infty(\mathcal{T}_H M, \Omega_r). \quad (2.10)$$

Using (2.10), there exists $f_x \in C_c^\infty(\mathcal{T}_H M_x)$ such that:

$$(\mathbb{k} - \mathbb{k}')|_{(x, \cdot, 0)} = f_x d\lambda_x, \quad (2.11)$$

where $d\lambda_x$ is a left Haar measure on the nilpotent (graded) Lie group $\mathcal{T}_H M_x$. To conclude, remember that the Debord-Skandalis action fixes $(x, 0, 0)$. Indeed, we compute:

$$\left(s^{d_H} \alpha_{s^*} (\mathbb{k} - \mathbb{k}') - (\mathbb{k} - \mathbb{k}') \right)|_{(x,0,0)} = s^{d_H} \alpha_{s^*} (\mathbb{k} - \mathbb{k}')|_{(x,0,0)} - (\mathbb{k} - \mathbb{k}')|_{(x,0,0)} \quad (2.12)$$

$$= s^{d_H} \delta_{s^*}(f_x d\lambda_x) - f_x d\lambda_x \quad (2.13)$$

$$= 0, \quad (2.14)$$

as in the previous proof. \square

Now we can conclude that the groupoidal residue is well-defined.

Let us now recall a basic fact about convolution Lie groups. In the following Lemma the commutators are in the sense of convolution product. Also recall that every connected nilpotent Lie group is unimodular, see [13, Proposition 2.30] or [12, Proposition 5.5.4 et Corollary 5.5.5].

Lemma 2.2. *Let G be a connected nilpotent Lie group G . Given $f \in C_c^\infty(G)$ and $g \in \mathcal{E}'(G)$ then $[f, g](e) = 0$.*

Proof. Recall that convolution on Lie group is defined by:

$$\underbrace{u \star v}_{\in \mathcal{E}'(G)} = \underbrace{(u \otimes v)}_{\in \mathcal{E}'(G \times G)} \circ m^*, \quad (2.15)$$

where $u \in \mathcal{E}'(G)$, $v \in \mathcal{E}'(G)$, $(m^*f)(x, y) = f(xy)$. We may also take $v \in \mathcal{D}'(G)$ and in this case $u \star v \in \mathcal{D}'(G)$, see [14, Theorem 5.1.1 p51 on \mathbb{R}^d]. The commutator is well-defined because $C_c^\infty(G)$ is a two-sided ideal in $\mathcal{E}'(G)$. Still inspired by the [14, Theorem 5.2.1 p53 on \mathbb{R}^d], one can prove that for $f \in C_c^\infty(G)$ and $g \in \mathcal{E}'(G)$:

$$f \star g(x) = \langle g(y), f(xy^{-1}) \rangle, \quad (2.16)$$

and:

$$g \star f(x) = \langle g(y), f(y^{-1}x) \rangle. \quad (2.17)$$

See that it is in the two previous equations that we use the unimodularity of G . Then we have proved $f \star g(e) = g \star f(e)$, that is $[f, g](e) = 0$. □

Theorem 2.3. *Let M be a filtered manifold of homogeneous dimension d_H . Let $P \in \Psi_H^m(M)$ be a pseudodifferential operator M with $m \leq -d_H$ and $Q \in \Psi_H^0(M)$. Then the groupoidal residue of Definition 1.4 satisfies the trace property, that is for all $x \in M$:*

$$Res_x([P, Q]) = 0. \quad (2.18)$$

Proof. We denote by \mathbb{k}_P and \mathbb{k}_Q the two associated essentially homogeneous r -fibered distributions of order $-d_H$ and 0 respectively. By definition, there exists $f, g \in C_p^\infty(\mathbb{T}_H M, \Omega_r)$ such that:

$$Res_x([P, Q]) = Res_x(PQ) - Res_x(QP) \quad (2.19)$$

$$= \frac{1}{\log(s)} \left((s^{d_H} \alpha_{s^*}(\mathbb{k}_P \star \mathbb{k}_Q) - \mathbb{k}_P \star \mathbb{k}_Q)|_{(x,0,0)} - (s^{d_H} \alpha_{s^*}(\mathbb{k}_Q \star \mathbb{k}_P) - \mathbb{k}_Q \star \mathbb{k}_P)|_{(x,0,0)} \right) \quad (2.20)$$

$$= \frac{1}{\log(s)} \left(s^{d_H} ((s^{-d_H} \mathbb{k}_P + f) \star (\mathbb{k}_Q + g) - \mathbb{k}_P \star \mathbb{k}_Q)|_{(x,0,0)} - s^{d_H} ((\mathbb{k}_Q + g) \star (s^{-d_H} \mathbb{k}_P + f) + \mathbb{k}_Q \star \mathbb{k}_P)|_{(x,0,0)} \right) \quad (2.21)$$

$$= \frac{1}{\log(s)} \left(\mathbb{k}_P \star g + f \star \mathbb{k}_Q - \mathbb{k}_Q \star f - g \star \mathbb{k}_P \right)|_{(x,0,0)} \quad (2.22)$$

$$= \frac{1}{\log(s)} \left([\mathbb{k}_P, g] + [\mathbb{k}_Q, f] \right)|_{(x,0,0)} \quad (2.23)$$

$$= 0. \quad (2.24)$$

The equality in equation (2.21) is true since \mathbb{k}_P and \mathbb{k}_Q are essentially homogeneous r -fibered distributions respectively of order $-d_H$ and 0 respectively and α_s is a groupoid

automorphism for all $s > 0$. The last equality is true by virtue of Lemma 2.2 applied fiberwise to the osculating groups which are by definition connected (graded) nilpotent Lie groups. Indeed the convolution in the brackets $[\mathbb{k}_P, g]$ and $[\mathbb{k}_Q, f]$ are done fiberwise. In the fiber $(x, 0, 0)$ the convolution is done between a distribution with compact support and a function on $\mathcal{T}_H M_x$ with compact support. \square

Remark 1. *Theorem 2.3 is in particular satisfied for $Q \in \Psi_H^{m'}(M)$ with $m' \leq -1$. In this case, we would have PQ and QP in the space $\Psi_H^{-d_H-1}(M)$. Therefore, the two r -fibered distributions $\mathbb{k}_P \star \mathbb{k}_Q$, $\mathbb{k}_Q \star \mathbb{k}_P$ would be two continuous functions, thanks to [28, Theorem 52]. Therefore the two co-cycles in (2.20) would be 0, as again the Debord-Skandalis action fixes $(x, 0, 0)$. So it is only the critical case $m' = 0$ which remained to be proven.*

3 Pseudo-homogeneous functions and kernels

It is well-known, see for instance [19],[3, p4], that classical pseudodifferential operators admit also a kernel expansion in terms of (pseudo)-homogeneous functions which is precisely linked to the asymptotic expansion of the symbol. Let us recall the details.

Consider \mathbb{R}^d equipped with a one-parameter family of dilations $(\delta_s)_{s>0}$. Typically, these will come from a grading on \mathbb{R}^d where δ_s acts on the subspace of graded degree k by multiplication by s^k . We will be particularly interested in the trivial dilation structure:

$$\delta_s(\xi_1, \dots, \xi_d) = (s\xi_1, \dots, s\xi_d), \quad (3.1)$$

and shall also encounter the Heisenberg dilation structure:

$$\delta_s(\xi_0, \xi_1, \dots, \xi_d) = (s^2\xi_0, s\xi_1, \dots, s\xi_d). \quad (3.2)$$

Now we can define:

Definition 3.1. [19, Definition 7.1.1 p353],[3, Definition (15.19)]

Let $U \subset \mathbb{R}^d$ be an open subset. We denote by $\mathcal{H}_G^m(U \times \mathbb{R}^d)$ the space of functions $f \in C^\infty(U \times \mathbb{R}^d \setminus \{0\})$ homogeneous of order m satisfying:

$$f(x, \delta_s(\xi)) = s^m f(x, \xi), \quad s > 0, \quad (3.3)$$

where δ_s is as in (3.1) or (3.2). We define the set $\Psi\text{hf}_G^m(U \times \mathbb{R}^d)$ of smooth pseudo-homogeneous functions of degree m in the second variable as follows. If $m \notin \mathbb{N}$ then :

$$\Psi\text{hf}_G^m(U \times \mathbb{R}^d) = \mathcal{H}_G^m(U \times \mathbb{R}^d). \quad (3.4)$$

If $m \in \mathbb{N}$ then $\Psi\text{hf}_G^m(U \times \mathbb{R}^d)$ is the set of $k \in C^\infty(U \times \mathbb{R}^d \setminus \{0\})$ of the form:

$$k(x, \xi) = f(x, \xi) + \log(|\xi|)p(x, \xi), \quad (3.5)$$

where p is a homogeneous polynomial in ξ of degree m having C^∞ -coefficients in x , where the function $f \in \mathcal{H}_G^m(U \times \mathbb{R}^d)$ and $|\cdot|$ is a homogeneous quasi-norm on \mathbb{R}^d , in the sense of dilations (3.1) or (3.2).

We also recall, see [9]:

Definition 3.2. Let $U \subset \mathbb{R}^d$ be an open subset. We denote by $\mathcal{HS}_G^m(U \times \mathbb{R}^d)$ the space of functions $f \in C^\infty(U \times \mathbb{R}^d)$ which are homogeneous of order m modulo Schwartz-class meaning:

$$f(x, \delta_s(\xi)) - s^m f(x, \xi), \quad (3.6)$$

is a Schwartz-class function of ξ , with smooth dependance on x .

Definition 3.3. [19, Equation (7.1.2) p354] [23, Definition 3.5 p414], [3, Equations (15.40)-(15.41)]

A distribution kernel $k \in \mathcal{D}'(U \times U)$ is said to have a pseudo-homogeneous expansion of degree $m \in \mathbb{R}$ if:

$$k \sim \sum_j k_{m+j}, \quad (3.7)$$

where $k_{m+j} \in \Psi\text{hf}_G^{m+j}(U \times \mathbb{R}^d)$ and where the symbol \sim means that for all $N \in \mathbb{N}$, there exists $J_N \in \mathbb{N}$:

$$k - \sum_{j \in J_N} k_{m+j} \in C^N(U \times U), \quad (3.8)$$

where $C^N(U \times U)$ denotes the space of class C^N functions. The space of kernels having a pseudo-homogeneous expansion of degree m is denoted $\Psi\text{hk}_G^m(U)$.

Remark 2. *Our use of $\mathcal{D}'(U \times U)$ in the above definition follows the standard practice of most works on pseudodifferential operators. Stricly speaking, when we realize these as kernels in the groupoid calculus, we will need to replace these by r -fibred distributions :*

$$k(x, y)dy \in \mathcal{D}'_r(U \times U), \quad (3.9)$$

see [22], [28]. The Lebesgue measure dy is homogeneous of degree $-d$ with respect to the trivial dilation δ_s so this introduces a degree-shift in orders of kernels, see the next theorem. Thus, the kernel of a pseudodifferential operator on U of order $m < 0$ will be given by an element of $\Psi\text{hk}_G^{-m-d}(U)$. This is a well-known technical detail, and we will not remark on it further.

We now state an important theorem, also see [19, thm 7.1.1, 7.1.6, 7.1.7, 7.1.8].

Theorem 3.4. *Seeley 1969 [26, theorem 1 p 209]*

Let $U \subset \mathbb{R}^d$ be an open subset and $m < 0$. Then $P \in \Psi_{\text{Hör}}^m(U)$ if and only if:

$$Pu(x) = \int_U k(x, x-y)u(y)dy, \quad u \in C_c^\infty(U), \quad (3.10)$$

with Schwartz kernel satisfying $k \in \Psi\text{hk}_G^{-m-d}(U)$.

Moreover, in the case of the trivial dilations structure, the asymptotic expansion of the symbol:

$$a \sim \sum_{j=0}^{\infty} a_{m-j}, \quad a_{m-j} \in \mathcal{H}_G^{m-j}(U \times \mathbb{R}^d), \quad (3.11)$$

and the kernel:

$$k \sim \sum_{j=0}^{\infty} k_{m+j}, \quad k_{m+j} \in \Psi\text{hf}_G^{m+j}(U \times \mathbb{R}^d), \quad (3.12)$$

are related by an adapted Fourier transform as follows see [19, Equation (7.1.81) p393]. Take $\psi \in C_c^\infty(\mathbb{R}^d)$ is any cut-off function satisfying:

$$\psi(z) := \begin{cases} 1 & \text{if } |z| \leq \frac{1}{2}, \\ 0 & \text{if } |z| > 1. \end{cases} \quad (3.13)$$

Set $\kappa = -m - d$. Then for $m - j < 0$:

$$a_{m-j}(x, \xi) = \lim_{t \rightarrow +\infty} \int_{\mathbb{R}^d} k_{\kappa+j}(x, z)\psi\left(\frac{z}{t}\right)e^{-i\xi \cdot z} dz, \quad x \in U. \quad (3.14)$$

4 The Wodzicki residue coincides with the groupoidal residue

We shall prove in this section that the groupoidal residue $Res_x(P)$ and the Wodzicki residue $Res_x^W(P)$ coincide when P is a classical pseudodifferential operator of order $\leq -d$ on a trivially filtered manifold, see Theorem 4.3. We begin by recalling exponential coordinates $\mathbb{E}xp^{\overline{X}}$ on $\mathbb{T}M$, see also [28].

Given a vector field X on M and a point $x \in M$, we write $\exp(X).x$ for the time one flow of x along X if defined. If $\overline{X} = (X_1, \dots, X_n)$ is a local frame of vector fields and $v \in \mathbb{R}^n$ then we set $v.\overline{X} = \sum_{k=1}^n v_k X_k$. Also, note that the dilations δ_s on \mathbb{R}^d in this case are given by $\delta_s(v) = sv$. The following Lemma lists the properties of the exponential charts of $\mathbb{T}_H M$ which we will need in the sequel.

Lemma 4.1. [28, Lemma 27 p 14], [9, Proposition 5.13], [8]

Let M be a smooth manifold of dimension d and $\mathbb{k} \in \Psi_{vEY}^m(\mathbb{T}M)$. Given $x_0 \in M$, (U_0, ϕ) a chart on x_0 and $\overline{X} = (X_1, \dots, X_d)$ a local frame on x_0 , we have:

1. There exists an open neighbourhood U of $U_0 \times \{0\}$ with $U \subset U_0 \times \mathbb{R}^d$ such that:

$$Exp^{\overline{X}} : U \rightarrow M \times M, (x, v) \mapsto (x, \exp(v.\overline{X}).x), \quad (4.1)$$

is a diffeomorphism onto its image.

2. The derivative of $Exp^{\overline{X}}$ at $(x_0, 0)$ is :

$$d_{(x_0, 0)} Exp^{\overline{X}} : (w, v) \in T_x U_0 \times \mathbb{R}^d \mapsto (w, v.\overline{X}|_x). \quad (4.2)$$

3. Put $\tilde{U} := \{(x, v, t) \in U_0 \times \mathbb{R}^d \times \mathbb{R}, (x, \delta_t(v)) \in U\}$. Then the map:

$$\mathbb{E}xp^{\overline{X}} : \tilde{U} \rightarrow \mathbb{T}M, (x, v, t) \mapsto \begin{cases} \mathbb{E}xp^{\overline{X}}(x, v, t) = (Exp^{\overline{X}}(x, \delta_t(-v)), t), & t \neq 0 \\ \mathbb{E}xp^{\overline{X}}(x, v, 0) = (x, v.\overline{X}|_x, 0) & t = 0, \end{cases} \quad (4.3)$$

defines the inverse of a smooth chart for the tangent groupoid $\mathbb{T}M$.

4. Let $\mathbb{U} = \mathbb{E}xp^{\overline{X}}(\tilde{U}) \in \mathbb{T}M$ be the domain of this chart. Then:

$$\mathbb{U} = (TM \times \{0\}) \cup (Exp^{\overline{X}}(U) \times \mathbb{R}^*), \quad (4.4)$$

is an open neighbourhood of $(TM \times \{0\}) \cup (\text{diag}(U_0) \times \mathbb{R}^*)$ in $\mathbb{T}M$, where $\text{diag}(U_0) = \{(x, x), x \in U_0\}$. Moreover \mathbb{U} is invariant for the Debord-Skandalis action 1.2, and the pullback of this action under $\mathbb{E}xp^{\overline{X}}$:

$$\tilde{\alpha}_s := (\mathbb{E}xp^{\overline{X}})^{-1} \circ \alpha_s \circ \mathbb{E}xp^{\overline{X}} : \tilde{U} \rightarrow \tilde{U}, \quad (4.5)$$

is given by:

$$\tilde{\alpha}_s(x, v, t) = (x, \delta_s(v), s^{-1}t). \quad (4.6)$$

5. There exists $\chi_{\mathbb{U}} \in C_c^\infty(\mathbb{T}M)$ invariant under the Debord-Skandalis action α_s such that:

$$\chi_{\mathbb{U}} = \begin{cases} 1 & \text{in a neighbourhood of } \{(x_0, x_0)\} \times \mathbb{R}, \\ 0 & \text{outside } \mathbb{U}. \end{cases} \quad (4.7)$$

6. $\mathbb{k}\chi_{\mathbb{U}} \in \Psi_{vEY}^m(\mathbb{T}M)$ has support in \mathbb{U} . Moreover $\tilde{\mathbb{k}} = (\mathbb{E}xp^{\overline{X}})_*^{-1}(\mathbb{k}\chi_{\mathbb{U}})$ has support in \tilde{U} and is essentially homogeneous for the action $\tilde{\alpha}_s$.

Lemma 4.2. *Let $U \subset \mathbb{R}^d$ be an open subset. If $k_0 \in \Psi\text{hf}_G^0(U \times \mathbb{R}^d)$ then $\text{Res}_x(P) = \text{Res}_x^W(P)$ is verified for the operator $P \in \Psi_{\text{Hör}}^{-d}(U)$ with Schwartz kernel $\chi(x-y)k_0(x, x-y)$ where $\chi \in C_c^\infty(\mathbb{R}^d)$ is 1 in a neighbourhood of 0 and 0 at infinity.*

Proof. By definition of $\Psi\text{hf}_G^0(U \times \mathbb{R}^d)$, we can write $k_0(x, z) = f_0(x, z) + \log(|z|)p_0(x)$, where p_0 is a smooth function on U and f_0 is a smooth function homogeneous of order 0 with respect to z . Let χ be as in the statement. Then we set:

1. $l(x, y) = \chi(x-y) \ln(|x-y|)p_0(x)dy$ to be the kernel of the operator $P = Op(l)$ whose kernel's asymptotic expansion is given by $l_0(x, z) = \ln(|z|)p_0(x)dz$ and $l_j(x, z) = 0$ if $j > 0$.
2. $r(x, y) = \chi(x-y)f_0(x, x-y)dy$ to be the kernel of the operator $P = Op(r)$ whose kernel's asymptotic expansion is given by $r_0(x, z) = f_0(x, z)dz$ and $r_j(x, z) = 0$ if $j > 0$.

We compute the Wodzicki residue at $x \in U$ respectively for each of these operators, using Definition 1.3. In both cases, recall that equation (3.14) gives us the link between the asymptotic symbol expansion and the asymptotic kernel expansion.

1. We get, by denoting \mathcal{F}_2 the Fourier transform with respect to the second variable:

$$a_{-d}(x, \xi) \underset{(3.14)}{=} \mathcal{F}_2\left(p_0(x) \log(|\cdot|)\right)(\xi) \quad (4.8)$$

$$= p_0(x) \mathcal{F}_2(\log(|\cdot|))(\xi), \quad (4.9)$$

where $\mathcal{F}_2(\log(|\cdot|))$ is interpreted as the Fourier transform of the tempered distribution $z \mapsto \log(|z|)$. Now, the Fourier transform of the logarithm in \mathbb{R}^d is well known and given, for $\xi \neq 0$ by:

$$\mathcal{F}_2(\log(|\cdot|))(\xi) = -\frac{1}{|\xi|^d} \frac{(2\pi)^d}{\omega_d}, \quad (4.10)$$

where $\omega_d = \frac{(2\pi)^d}{\sqrt{\pi^d} \Gamma(\frac{d}{2}) 2^{d-1}}$ denotes the surface area of the unit $(d-1)$ -sphere $\mathbb{S}^{d-1} := \{\xi \in \mathbb{R}^d, |\xi| = 1\} \subset \mathbb{R}^d$. Then in (4.9) we may now write:

$$a_{-d}(x, \xi) \underset{(4.10)}{=} -\frac{p_0(x) (2\pi)^d}{|\xi|^d \omega_d}. \quad (4.11)$$

Therefore from (1.3) we get:

$$\text{Res}_x^W(Op(l)) = \frac{1}{(2\pi)^d} \int_{\mathbb{S}^{d-1}} -\frac{p_0(x) (2\pi)^d}{|\xi|^d \omega_d} d\sigma(\xi) dx = -p_0(x) dx, \quad (4.12)$$

where $d\sigma$ denotes the usual surface measure on \mathbb{S}^{d-1} .

2. As $f_0 \in \mathcal{H}^0(U \times \mathbb{R}^d)$, we may extend it to $f_0 \in C^\infty(U) \otimes L^\infty(\mathbb{R}^d)$ by attributing any value at $\xi = 0$ for all $x \in U$. The result is a tempered distribution (generalized function) which is homogeneous of degree 0.

We now proceed with $x \in U$ fixed. Thanks to [15, Proposition 2.4.7 p 140], or [6, p86], there exists $b_x \in \mathbb{C}$ and Ω_x a smooth function on the sphere \mathbb{S}^{d-1} with integral 0 on \mathbb{S}^{d-1} such that:

$$\mathcal{F}_2(f_0(x, \cdot))(\xi) = b_x \delta_0 + W_{\Omega_x}(\xi), \quad (4.13)$$

where W_{Ω_x} is the principal value distribution whose restriction to $\mathbb{R}^d \setminus \{0\}$ is:

$$\Omega_x \left(\frac{\xi}{|\xi|} \right) \frac{1}{|\xi|^d}, \quad (4.14)$$

see [15, Equation (2.4.12)]. When $\xi \neq 0$ we have that:

$$a_{-d}(x, \xi) \underbrace{=}_{(3.14)} \mathcal{F}_2(f_0(x, \cdot))(\xi) \quad (4.15)$$

$$\underbrace{=}_{(4.13)} W_{\Omega_x}(\xi), \quad (4.16)$$

is smooth in $\mathbb{R}^d \setminus \{0\}$. It follows that:

$$\int_{\mathbb{S}^{d-1}} a_{-d}(x, \xi) d\sigma(\xi) \underbrace{=}_{(4.16)} \int_{\mathbb{S}^{d-1}} W_{\Omega_x}(\xi) d\sigma(\xi) \quad (4.17)$$

$$\underbrace{=}_{(4.14)} \int_{\mathbb{S}^{d-1}} \Omega_x(\xi) d\sigma(\xi) \quad (4.18)$$

$$= 0, \quad (4.19)$$

where the last equality is true by the assumption on Ω_x . We therefore have:

$$Res_x^W(Op(r)) = 0. \quad (4.20)$$

Now we look at the co-cycles at x of the r -fibred distributions essentially homogeneous associated to the operators $Op(l)$, $Op(r)$ and prove that at $(x, 0, 0)$ we recover the residue values (4.12) and (4.20). First, we can respectively define two elements in $\Psi_{\text{vEY}}^m(\mathbb{T}M)$ such that their restrictions in $t = 1$ give the kernels l and r .

1. Set:

$$\begin{cases} \mathbb{I}(x, y, t) = \frac{1}{t^d} \chi\left(\frac{x-y}{t}\right) \log\left(\frac{|x-y|}{t}\right) p_0(x) dy & \text{if } t \neq 0 \\ \mathbb{I}(x, v, 0) = \chi(v) \log(|v|) p_0(x) d\lambda_x(v) & \text{if } t = 0, \end{cases} \quad (4.21)$$

where $d\lambda_x$ denotes the Haar measure on the tangent space $T_x M$ at x . Writing this in exponential coordinates according to Lemma 4.1 with respect to the standard coordinate frame \bar{X} for \mathbb{R}^d , we get:

$$\tilde{\mathbb{I}}(x, v, t) = \left(\mathbb{E}xp^{\bar{X}}\right)_*^{-1} \mathbb{I}(x, v, t) = \chi(v) \log(|v|) p_0(x) d\lambda_x(v). \quad (4.22)$$

Recalling $\tilde{\alpha}_s$ from Lemma 4.1 and using the fact that $\delta_{s*}(d\lambda_x) = s^{-d} d\lambda_x$, we get:

$$s^d \tilde{\alpha}_{s*} \tilde{\mathbb{I}}(x, v, t) - \tilde{\mathbb{I}}(x, v, t) = s^d \tilde{\mathbb{I}}\left(x, \delta_{s^{-1}}(v), st\right) \delta_{s*}(d\lambda_x(v)) - \tilde{\mathbb{I}}(x, v, t) \quad (4.23)$$

$$\begin{aligned} &= -\log(s) \chi(s^{-1}v) p_0(x) d\lambda_x(v) \\ &\quad + \left(\chi(s^{-1}v) - \chi(v)\right) \log(|v|) p_0(x) d\lambda_x(v). \end{aligned} \quad (4.24)$$

We deduce that :

$$s^d \tilde{\alpha}_{s*} \tilde{\mathbb{I}} - \tilde{\mathbb{I}} \in C_p^\infty(\tilde{U}, \Omega_r). \quad (4.25)$$

2. Set:

$$\begin{cases} \mathbb{r}(x, y, t) = \chi\left(\frac{x-y}{t}\right) f_0\left(x, \frac{x-y}{t}\right) dy & \text{if } t \neq 0 \\ \mathbb{r}(x, v, 0) = \chi(v) f_0(x, v) d\lambda_x(v) & \text{if } t = 0, \end{cases} \quad (4.26)$$

In the same exponential coordinates we get:

$$\tilde{\mathbb{r}}(x, v, t) = \left(\mathbb{E}xp^{\bar{X}}\right)_*^{-1} \mathbb{r}(x, v, t) = \chi(v) f_0(x, v) d\lambda_x(v), \quad (4.27)$$

and the homogeneity of f_0 gives:

$$s^d \tilde{\alpha}_{s*} \tilde{\mathbb{r}} - \tilde{\mathbb{r}} \in C_p^\infty(\tilde{U}, \Omega_r). \quad (4.28)$$

We move now to the computations of the co-cycles restricted in $(x, 0, 0)$.

1. Using (4.24) we get, for all $s \in \mathbb{R}_+^* \setminus \{1\}$:

$$\left(\tilde{\alpha}_{s^*} \tilde{\mathbb{I}} - s^{-d} \tilde{\mathbb{I}}\right)|_{(x,0,0)} = -\frac{1}{s^d} \log(s) p_0(x) d\lambda_x \quad (4.29)$$

$$= \frac{1}{s^d} \log(s) \text{Res}_x^W(\text{Op}(l)), \quad (4.30)$$

where we use the canonical identification of the smooth family of 1-densities $d\lambda_x$ with the smooth measure dx on M . Then:

$$\frac{1}{\log(s)} \left(s^d \tilde{\alpha}_{s^*} \tilde{\mathbb{I}} - \tilde{\mathbb{I}}\right)|_{(x,0,0)} = \text{Res}_x^W(\text{Op}(l)). \quad (4.31)$$

2. Using (4.27) we get, for all $s \in \mathbb{R}_+^* \setminus \{1\}$:

$$\left(\tilde{\alpha}_{s^*} \tilde{\mathfrak{r}} - s^{-d} \tilde{\mathfrak{r}}\right)|_{(x,0,0)} = \left(f_0(x, v) \left[\chi\left(\frac{v}{s}\right) - \chi(v)\right] d\lambda_x(v)\right)|_{(x,0,0)} \quad (4.32)$$

$$= 0 \quad (4.33)$$

$$= \frac{1}{s^d} \log(s) \text{Res}_x^W(\text{Op}(r)). \quad (4.34)$$

Then:

$$\frac{1}{\log(s)} \left(s^d \tilde{\alpha}_{s^*} \tilde{\mathfrak{r}} - \tilde{\mathfrak{r}}\right)|_{(x,0,0)} = \text{Res}_x^W(\text{Op}(r)). \quad (4.35)$$

This completes the proof. \square

Theorem 4.3. *Let M be a (trivially) filtered manifold of dimension d and $P \in \Psi_H^m(M)$ a classical pseudodifferential operator of order m on M , with $m \leq -d$, $m \in \mathbb{Z}$. Let \mathbb{k} be any essentially homogeneous r -fibered distribution of order $-d$ that extends P at $t = 1$. Then:*

$$\text{Res}_x^W(P) = \frac{1}{\log(s)} \left(s^d \alpha_{s^*} \mathbb{k} - \mathbb{k}\right)|_{(x,0,0)}, \quad \forall s \in \mathbb{R}_+^* \setminus \{1\}, \quad \forall x \in M. \quad (4.36)$$

Proof. Consider first the case $m \leq -d - 1$. Thanks to [28, Theorem 52] we already know that:

$$\mathbb{k} \in C^0(\mathbb{T}M, \Omega_r), \quad (4.37)$$

as we have supposed here $m \leq -d - 1$. That means:

$$\mathbb{k}(x, v, 0) = \mathbb{l}_0(x, v) d\lambda_x, \quad (4.38)$$

for some $\mathbb{l}_0 \in C^0(TM)$ and where $d\lambda_x$ is the Haar measure on $T_x M$.

We can now evaluate the co-cycle term to term. Moreover the points $(x, 0, 0)$ are fixed by the Debord-Skandalis action. Thus, using the facts that $\delta_{s^*} d\lambda_x = s^{-d} d\lambda_x$, we get:

$$\frac{1}{\log(s)} \left(s^d \tilde{\alpha}_{s^*} \mathbb{k} - \mathbb{k}\right)|_{(x,0,0)} = \frac{1}{\log(s)} \left(\mathbb{l}_0(x, 0) d\lambda_x - \mathbb{l}_0(x, 0) d\lambda_x\right) = 0. \quad (4.39)$$

This agrees with the Wodzicki residue in this case. Indeed, for an operator of this order the term a_{-d} appearing in the asymptotic expansion of its symbol is always zero.

If $m = -d$ and (U_0, ϕ) is a chart on $x \in M$, then the kernel admits an asymptotic expansion $k \sim \sum_{j=0}^{+\infty} k_j$ in U_0 , thanks to Seeley's Theorem 3.4. Thus we may write:

$$k(x, x - y) - \chi(x - y)k_0(x, x - y) \sim \sum_{j \geq 1} k_j(x, x - y), \quad (4.40)$$

with $\chi \in C_c^\infty(\mathbb{R}^d)$ is such that χ is equal to 1 in a neighbourhood of 0 and 0 at infinity. Let us denote the left hand side of (4.40) by $\bar{k}(x, x - y)$. Then $\bar{P} = Op(\bar{k}) \in \Psi_{\text{Hör}}^{m-1}(M)$ and using what we did just before, $Res_x^W(\bar{P}) = Res_x(\bar{P})$ for all $x \in M$. It suffices to apply Lemma 4.2 to the function $k_0 \in \Psi\text{hf}_G^0$ to conclude, as we have $k(x, z) = \bar{k}(x, z) + \chi(z)k_0(x, z)$. \square

5 Ponge's non commutative residue for an Heisenberg manifold.

In his article [23], Ponge defined a noncommutative residue that fits the context of a Heisenberg manifold. In this section, we will show that Ponge's residue coincides with the groupoidal residue of Definition 1.4 for pseudodifferential operators of order $\leq -d_H$ on a contact manifold or a codimensional one foliation. Again, we will restrict our attention to scalar-valued operators to simplify notation, although one can easily generalise to vector bundles using (1.13).

Let M be a Heisenberg manifold of dimension $d + 1$ with hyperplane bundle $\mathcal{V} \leq TM$. The algebra of Heisenberg pseudodifferential operators of BG [3] is denoted $\Psi_{\mathcal{V}}^\bullet(M)$. It is shown in [9] that this coincides with the groupoidal calculus when $M = \mathbb{H}_n \times \mathbb{R}^m$, \mathbb{H}_n being the $2n + 1$ dimensional Heisenberg group, or M is a contact manifold or a codimensional one foliation. That is:

$$\Psi_{\mathcal{V}}^m(M) = \Psi_H^m(M), \quad (5.1)$$

for M in these cases, though we expect it to be true also for a general Heisenberg manifold.

Ponge's noncommutative residue is defined as follows. Let $(X_j)_{j \in \{0, \dots, d\}}$ be a local H -frame of vector fields on an open subset $U \subset M$ and $\Psi_x : U \rightarrow \mathbb{R}^{d+1}$ be a privileged change of coordinates centered at x , see [23, p 415 and Definitions 2.3 ,2.4]. The latter assertion means that if $(X_j)_{j \in \{0, \dots, d\}}$ is a local H -frame of vector fields, then we have $\Psi_x(x) = 0$ and $(\Psi_x)_* X_j(x) = \partial_j|_x$. In this context, the noncommutative residue of Ponge of a Heisenberg pseudodifferential operator $P \in \Psi_{\mathcal{V}}^{-d_H}(U)$ of degree $-d_H$ on U , is defined as follows.

Let p_{-d-2} be the term of degree $-d_H = -(d + 2)$ from the asymptotic expansion of the symbol of $P \in \Psi_{\mathcal{V}}^{-d_H}(U)$, see [23, 2.7 p409]. Then set the noncommutative residue of P at x :

$$c_P(x) = \frac{|d\Psi_x|}{(2\pi)^{d+1}} \int_{\mathbb{S}^d} p_{-(d+2)}(x, \xi) d\xi, \quad (5.2)$$

where $|d\Psi_x|$ is the jacobian of Ψ_x , see [23, Lemma 3.9].

The reader can compare this definition in contrast with the "non-graded" non commutative residue (1.3). We begin with the case of the model groups $M = \mathbb{H}_n \times \mathbb{R}^m = \mathbb{R}^{d+1}$, where $d = 2n + m$. If $n = 0$ then $\mathbb{H}_0 = \mathbb{R}$ by convention. We equip M with the model vector fields $\bar{X} = (X_0, X_1, \dots, X_d)$ of [3, chapter 1 p12-13], also see [9, section 5.1], so that (X_0, \dots, X_{2n}) generate \mathbb{H}_n , X_0 is central and (X_{2n+1}, \dots, X_d) are the usual vectors fields on \mathbb{R}^{d+1} . Recall that $\mathcal{E}'_r(M \times M)$ denotes the set of r -fibered distributions on the pair groupoid $M \times M$.

Theorem 5.1. *Given a model manifold $M = \mathbb{H}_n \times \mathbb{R}^m$ of homogeneous dimension $d_H = d + 2$, $2n + m = d$, with the standard model structure as in [9] and $P \in \Psi_{\mathcal{V}}^m(M)$, with $m \leq -d_H$, then the residue of Ponge at x and our residue from Definition 1.4 coincide:*

$$\text{Res}_x(P) = c_P(x). \quad (5.3)$$

Proof. We will rely heavily on [3] and [9]. We denote $k \in \mathcal{E}'_r(M \times M)$ the kernel of P . Since the Heisenberg and groupoidal calculi coincide, see [9, Theorem 5.16], there exists $\mathbb{k} \in \Psi_{vEY}^{-d_H}(\mathbb{T}_H M)$ such that $\mathbb{k}|_{t=1} = k$. Moreover we may suppose, see [28, proposition 42] that \mathbb{k} is homogeneous on the nose of order $-d_H$ outside $t \in [-1, 1]$. We pull this back via exponential coordinates, again as in [9, section 5], yielding $\tilde{\mathbb{k}} \in \mathcal{E}'_r(M \times \mathbb{R}^{d+1} \times \mathbb{R})$ with the equality $\mathbb{k} = \mathbb{E}xp_*^{\bar{X}}(\tilde{\mathbb{k}})$. Note that in the case of model manifold, the exponential coordinate chart is globally define.

Next we must consider the symbol of P . By the definition of the Heisenberg calculus, [3, Chapter 3, § 10], P is defined starting from a graded-polyhomogeneous function $f \in S_{phg,G}^{-d_H}(M \times \mathbb{R}^{d+1})$. Letting $\bar{\sigma}(x, \xi) = (x, \sigma_0(x, \xi), \dots, \sigma_d(x, \xi))$ be the coordinate transform [3, Equations (10.14),(10.15)] obtained from the symbols of the model vector fields (X_0, \dots, X_d) , Beals and Greiner define the \mathcal{V} -symbol associated to f by:

$$q(x, \xi) = \bar{\sigma}^* f(x, \xi). \quad (5.4)$$

See also [9, Definition 5.5]. The symbol and kernel are related by fiberwise Fourier transform, after the abovementioned coordinate changes. Explicitly, we have:

$$f = \mathcal{F}_2(\tilde{\mathbb{k}})|_{t=1}, \quad (5.5)$$

where \mathcal{F}_2 is the fiberwise Fourier transform in the second variable. Extending this fiberwise Fourier transform to all $t \in \mathbb{R}$, let us put:

$$u = \mathcal{F}_2(\tilde{\mathbb{k}}) \in C^\infty(M \times \mathbb{R}^{d+1} \times \mathbb{R}). \quad (5.6)$$

Note that the Debord-Skandalis action transforms under the Fourier transform as:

$$\mathcal{F}_2 \circ \tilde{\alpha}_{s*} \circ \mathcal{F}_2^{-1} = \beta_s^*, \quad (5.7)$$

where $\beta_s : M \times \mathbb{R}^{d+1} \times \mathbb{R} \rightarrow M \times \mathbb{R}^{d+1} \times \mathbb{R}$ are the dilations:

$$\beta_s(x, v, t) = (x, \delta_s(v), st), \quad (5.8)$$

see [9, Proposition .13]. The essential homogeneity of \mathbb{k} and consequently of $\tilde{\mathbb{k}} = (\mathbb{E}xp_*^{\bar{X}})^{-1}(\mathbb{k})$, therefore implies that $u \in \mathcal{HS}_G^m(M \times \mathbb{R}^{d+1} \times \mathbb{R})$ where the homogeneity modulo Schwartz is with respect to the dilations β_s (thanks to the hypothesis on the homogeneity of \mathbb{k} outside $t \in [-1, 1]$), see the proof of [9, Theorem 5.16].

Set $u_0 = u|_{t=0}$. By [9, Proposition 3.2] we have $u_0 \in \mathcal{HS}_G^{-d_H}(M \times \mathbb{R}^{d+1})$. Therefore by a well-known Lemma, eg [9, Theorem 2.1], outside a compact neighbourhood containing 0 of ξ we may write:

$$u_0 = u'_0 + u''_0, \quad (5.9)$$

where $u'_0 \in \mathcal{H}_G^{-d_H}(M \times \mathbb{R}^{d+1})$ and $u''_0 \in S_G(M \times \mathbb{R}^{d+1})$.

We may extend the homogeneous function u'_0 as a tempered distribution (not necessarily homogeneous), still denoted u'_0 . Indeed, we may first extend u'_0 to a distribution such as in [3, section § 15], [24, Lemma 3.1], or in [18, Theorem 3.2.4], and it is tempered because it has polynomial growth at infinity. Therefore we can find $u_0''' \in C^\infty(M, \mathcal{E}'(\mathbb{R}^{d+1})) = C^\infty(M) \otimes \mathcal{E}'(\mathbb{R}^{d+1})$ such that the following holds everywhere:

$$u_0 = u'_0 + u''_0 + u_0''' . \quad (5.10)$$

We can compute for all $s \in \mathbb{R}_+^* \setminus \{1\}$:

$$Res_x(P) = \frac{1}{\log(s)} \left(s^{d_H} \alpha_{s*} \mathbb{k} - \mathbb{k} \right) |_{(x,0,0)} \quad (5.11)$$

$$= \frac{1}{\log(s)} \left(s^{d_H} \alpha_{s*} \mathbb{E}xp_*^{\overline{X}}(\tilde{\mathbb{k}}) - \mathbb{E}xp_*^{\overline{X}}(\tilde{\mathbb{k}}) \right) |_{(x,0,0)} \quad (5.12)$$

$$= \frac{1}{\log(s)} \left(s^{d_H} \tilde{\alpha}_{s*} \tilde{\mathbb{k}} - \tilde{\mathbb{k}} \right) \circ \left(\mathbb{E}xp^{\overline{X}} \right)^{-1} |_{(x,0,0)}, \quad (5.13)$$

where we used in (5.12) the equality $\tilde{\alpha}_s = \left(\mathbb{E}xp^{\overline{X}} \right)^{-1} \circ \alpha_s \circ \mathbb{E}xp^{\overline{X}}$, see (4.5). Also recall that $\mathbb{E}xp^{\overline{X}}|_{M \times \{0\} \times \{0\}} = id_{M \times \{0\} \times \{0\}}$. We next continue to compute in (5.13):

$$Res_x(P) = \frac{1}{\log(s)} \left(s^{d_H} \tilde{\alpha}_{s*} \mathcal{F}_2^{-1}(u) - \mathcal{F}_2^{-1}(u) \right) |_{(x,0,0)} \quad (5.14)$$

$$= \frac{1}{\log(s)} \mathcal{F}_2^{-1} \left(s^{d_H} \beta_s^* u - u \right) |_{(x,0,0)}, \quad (5.15)$$

where we use the equality $\mathcal{F}_2 \circ \tilde{\alpha}_{s*} \mathcal{F}_2^{-1} = \beta_s^*$ recalled earlier. Now we use (5.10):

$$Res_x(P) = \frac{1}{\log(s)} \mathcal{F}_2^{-1} \left(s^{d_H} \beta_s^* (u'_0 + u''_0 + u_0''') - (u'_0 + u''_0 + u_0''') \right) |_{(x,0,0)} \quad (5.16)$$

$$= \frac{1}{\log(s)} \mathcal{F}_2^{-1} \left(s^{d_H} \beta_s^* u'_0 - u'_0 + s^{d_H} \beta_s^* (u''_0 + u_0''') - (u''_0 + u_0''') \right) |_{(x,0,0)}. \quad (5.17)$$

Since u''_0 is Schwartz class in ξ and u_0''' is compactly supported in ξ , their fiberwise Fourier transforms are smooth.

Therefore $\mathcal{F}_2^{-1}(u''_0)$ and $\mathcal{F}_2^{-1}(u_0''')$, can be evaluated at $(x, 0, 0)$ and as in the proof of Theorem 4.3, for all $s \in \mathbb{R}_+^* \setminus \{1\}$, we get:

$$\frac{1}{\log(s)} \mathcal{F}_2^{-1} \left(s^{d_H} \beta_s^* (u''_0 + u_0''') - (u''_0 + u_0''') \right) |_{(x,0,0)} = 0. \quad (5.18)$$

Hence:

$$Res_x(P) = \frac{1}{\log(s)} \mathcal{F}_2^{-1} \left(s^{d_H} \beta_s^* u'_0 - u'_0 \right) |_{(x,0,0)}. \quad (5.19)$$

Finally, we use [23, Lemma 3.1 p 414 and Equation (3.2)] which assert that we have for all $s \in \mathbb{R}_+^* \setminus \{1\}$:

$$\beta_s^* u'_0 = s^{-d_H} u'_0 + s^{-d_H} \log(s) c_0(u'_0) \delta_0, \quad (5.20)$$

where:

$$c_0(u'_0) = \int_{|\xi|=1} u'_0(x, \xi) d\sigma(\xi). \quad (5.21)$$

Then, using (5.20), Equation (5.19) becomes:

$$Res_x(P) = \frac{1}{\log(s)} \mathcal{F}_2^{-1} \left(\log(s) c_0(u'_0) \delta_0 \right) |_{(x,0,0)} \quad (5.22)$$

$$= \frac{1}{(2\pi)^{d+1}} \int_{|\xi|=1} u'_0(x, \xi) d\sigma(\xi), \quad (5.23)$$

where the constant $\frac{1}{(2\pi)^{d+1}}$ appears in the inverse Fourier transform formula. Now, thanks to [3, Equation (3.25) p 27] we see that:

$${}^t(d\Psi_x)\xi = \sigma(x, \xi), \quad (5.24)$$

where σ is defined in [9, Equation (5.12)]. The reader can compute the Jacobian of Ψ_x and see that it is triangular and unipotent and so has determinant equal to one, see also [8].

According to [9, the proof p8 of Theorem 1.12], the purely homogeneous component u'_0 of u_0 is the first term in the asymptotic expansion of the polyhomogeneous function $f = u|_{t=1}$, see [9, Equation (3.16)] and the remarks which follow. Therefore:

$$u'_0(x, \xi) = p_{-(d+2)}(x, \xi), \quad (5.25)$$

where $p_{-(d+2)}$ is the term of degree $-(d+2)$ in the asymptotic expansion of $f = u|_{t=1}$. We have shown that :

$$Res_x(P) = \frac{|d\Psi_x|}{(2\pi)^{d+1}} \int_{|\xi|=1} p_{-(d+2)}(x, \xi) d\sigma(\xi). \quad (5.26)$$

□

Corollary 5.2. *If M is a contact manifold or a codimensional one foliation, then the groupoidal residue of Definition 1.4 agree with's Ponge noncommutative residue for operators $P \in \Psi_{\mathcal{V}}^{-d_H}(M)$, meaning that (5.3) still holds.*

Proof. Darboux' Theorem for a contact manifold or Frobenius' Theorem for a codimensional one foliation, implies that around any $x \in M$, there is a local coordinate system which identifies M with the model space \mathbb{H}_n , $d = 2n$, or $\mathbb{H}_0 \times \mathbb{R}^d$. Since both Ponge's residue and the groupoidal residue are independent of (priveleged) coordinates, the result follows. □

References

- [1] Thomas Ackermann. A note on the wodzicki residue. *Journal of Geometry and Physics*, 20(4):404–406, 1996.
- [2] Iakovos Androulidakis and Georges Skandalis. Pseudodifferential calculus on a singular foliation. *Journal of noncommutative geometry*, 5(1):125–152, 2010.
- [3] Richard Beals and Peter Charles Greiner. *Calculus on Heisenberg Manifolds.(AM-119), Volume 119*. Princeton University Press, 1988.
- [4] Duván Cardona, César del Corral, and Vishvesh Kumar. Dixmier traces for discrete pseudo-differential operators. *Journal of Pseudo-Differential Operators and Applications*, 11(2):647–656, 2020.
- [5] Woocheol Choi and Raphael Ponge. Tangent maps and tangent groupoid for carnot manifolds. *Differential Geometry and its Applications*, 62:136–183, 2019. <https://doi.org/10.1016/j.difgeo.2018.11.002>.
- [6] Ronald R Coifman and Yves Meyer. Au dela des opérateurs pseudo-différentiels. 1978.
- [7] Alain Connes. *Noncommutative Geometry*. Gulf Professional Publishing, 1994.
- [8] Nathan Couchet. *Une approche groupoïdale au calcul pseudo-différentiel sur les variétés d'Heisenberg et au résidu de Wodzicki*. PhD thesis, Université Clermont Auvergne, LMBP. In preparation for 2023.

- [9] Nathan Couchet and Robert Yuncken. On polyhomogeneous symbols and the heisenberg pseudodifferential calculus. 2022. <https://arxiv.org/abs/2210.15391>.
- [10] Shantanu Dave and Stefan Haller. Graded hypoellipticity of bgg sequences. *Annals of Global Analysis and Geometry*, 62(4):721–789, 2022.
- [11] Claire Debord and Georges Skandalis. Adiabatic groupoid, crossed product by \mathbb{R}_+^* and pseudodifferential calculus. *Advances in Mathematics*, 257:66–91, 2014. <https://doi.org/10.1016/j.aim.2014.02.012>.
- [12] Jacques Faraut. *Analysis on Lie groups, volume 110 of.* 2008.
- [13] Gerald B Folland. *A course in abstract harmonic analysis*, volume 29. CRC press, 2016.
- [14] Friedrich Gerard Friedlander, Mark Suresh Joshi, and M Joshi. *Introduction to the Theory of Distributions*. Cambridge University Press, 1998.
- [15] Loukas Grafakos. *Classical fourier analysis*, volume 2. Springer, 2008.
- [16] Victor Guillemin. A new proof of weyl’s formula on the asymptotic distribution of eigenvalues. *Advances in Mathematics*, 55(2):131–160, 1985.
- [17] Nigel Higson, Fedor Sukochev, and Dmitriy Zanin. In preparation.
- [18] Lars Hörmander. *The analysis of partial differential operators*. Springer, second edition, 1983.
- [19] George C Hsiao and Wolfgang L Wendland. *Boundary integral equations*. Springer, 2008.
- [20] H. Figueroa J. Varilly, J. Gracia Bondia. *Elements of noncommutative geometry*. Springer, chap 7 :251-318, 2001.
- [21] Christian Kassel. Le résidu non commutatif. *Séminaire Bourbaki*, 708:199–229, 1989. http://www.numdam.org/article/SB_1988-1989__31__199_0.pdf.
- [22] Jean-Marie Lescure, Dominique Manchon, and Stéphane Vassout. About the convolution of distributions on groupoids. *Journal of Noncommutative Geometry*, 11(2):757–789, 2017. <https://doi.org/10.4171/JNCG/11-2-10>.
- [23] Raphaël Ponge. Noncommutative residue for heisenberg manifolds. applications in cr and contact geometry. *Journal of Functional Analysis*, 252(2):399–463, 2007. <https://www.sciencedirect.com/science/article/pii/S0022123607002753>.
- [24] Raphaël Ponge. *Heisenberg calculus and spectral theory of hypoelliptic operators on Heisenberg manifolds*. American Mathematical Soc., 2008. <https://doi.org/10.1090/memo/0906>.
- [25] Raphael Ponge. Connes’ integration and weyl’s laws. *arXiv preprint arXiv:2107.01242*, 2021.
- [26] Robert Seeley. *Topics in pseudo-differential operators*. Springer, 2011.
- [27] Erik van Erp and Robert Yuncken. On the tangent groupoid of a filtered manifold. *Bull. Lond. Math. Soc.*, 49(6):1000–1012, 2017. <https://doi.org/10.1112/blms.12096>.
- [28] Erik Van Erp and Robert Yuncken. A groupoid approach to pseudodifferential calculi. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2019(756):151–182, 2019. <https://doi.org/10.1515/crelle-2017-0035>.

- [29] Stéphane Vassout. *Feuilletages et résidu non commutatif longitudinal*. PhD thesis, Paris 6, 2001.
- [30] Hermann Weyl. Über die asymptotische verteilung der eigenwerte. *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse*, 1911:110–117, 1911.
- [31] Mariusz Wodzicki. Local invariants of spectral asymmetry. *Inventiones mathematicae*, 75(1):143–177, 1984.
- [32] Mariusz Wodzicki. *Spectral asymmetry and noncommutative residue*. PhD thesis, Thesis, Stekhlov Institute of Mathematics, Moscow, 1984.