

Notes on the value function approach to multiobjective bilevel optimization

Daniel Hoff* Patrick Mehlitz†

March 29, 2023

This paper is concerned with the value function approach to multiobjective bilevel optimization which exploits a lower level frontier-type mapping in order to replace the hierarchical model of two interdependent multiobjective optimization problems by a single-level multiobjective optimization problem. As a starting point, different value-function-type reformulations are suggested and their relations are discussed. Here, we focus on the situations where the lower level problem is solved up to efficiency or weak efficiency, and an intermediate solution concept is suggested as well. We study the graph-closedness of the associated efficiency-type and frontier-type mappings. These findings are then used for two purposes. First, we investigate existence results in multiobjective bilevel optimization. Second, for the derivation of necessary optimality conditions via the value function approach, it is inherent to differentiate frontier-type mappings in a generalized way. Here, we are concerned with the computation of upper coderivative estimates for the frontier-type mapping associated with the setting where the lower level problem is solved up to weak efficiency. We proceed in two ways, relying, on the one hand, on a weak domination property and, on the other hand, on a scalarization approach. Throughout the paper, illustrative examples visualize our findings, the necessity of crucial assumptions, and some flaws in the related literature.

Keywords: Bilevel optimization, Existence results, Generalized differentiation, Parametric optimization, Multiobjective optimization, Set-valued and variational analysis

MSC: 49J52, 49J53, 90C29, 90C30

1 Introduction

In this note, we are concerned with multiobjective bilevel optimization problems of type

$$\min_{x,y} \kappa\{F(x,y) \mid x \in X, y \in \widehat{\Psi}(x)\} \quad (\text{BPP})$$

*Technische Universität Ilmenau, Institute of Mathematics, 98684 Ilmenau, Germany, daniel.hoff@tu-ilmenau.de, ORCID: 0009-0001-2725-2692

†Brandenburgische Technische Universität Cottbus–Senftenberg, Institute of Mathematics, 03046 Cottbus, Germany, mehlitz@b-tu.de, ORCID: 0000-0002-9355-850X

where $F: \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^p$ is a continuous vector-valued function, $\mathcal{K} \subset \mathbb{R}^p$ is a nonempty, closed, convex, pointed cone with nonempty interior, $X \subset \mathbb{R}^n$ is a nonempty, closed set, and $\widehat{\Psi}: \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ is a suitable solution mapping associated with the parametric multiobjective optimization problem

$$\min_y \mathcal{C} \{f(x, y) \mid y \in \Gamma(x)\}. \quad (\text{P}(x))$$

Above, $f: \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^q$ is a continuous vector-valued function, $\mathcal{C} \subset \mathbb{R}^q$ is a nonempty, closed, convex, pointed cone with nonempty interior, and $\Gamma: \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ is a set-valued mapping with a closed graph. In our model problem, \mathcal{K} and \mathcal{C} play the role of ordering cones, i.e., they specify what *minimization* means in (BPP) and (P(x)), up to different notions of optimality/efficiency in multiobjective optimization. Indeed, even for a fixed ordering cone, there exist diverse concepts of efficiency which aim to characterize the valuable points in the feasible set when faced with multiple objective functions. In this paper, we focus our attention on so-called efficiency and weak efficiency. Exemplary, we refer the interested reader to the monographs [23, 36] for a detailed introduction to multiobjective optimization and to Section 2.2 where we summarize some essential foundations of this topic.

In case where $p := q := 1$ and $\mathcal{K} := \mathcal{C} := \mathbb{R}_+$, (BPP) reduces to a so-called standard bilevel optimization problem which is used to model situations of interdependent decision making between two parties that cooperate with each other. Let us note that this model is closely related to the so-called optimistic approach to bilevel optimization as shown in [20, Proposition 6.9]. In the literature, (BPP) and (P(x)) are referred to as the upper and lower level problem, respectively. Due to numerous underlying applications from, e.g., finance, economics, data as well as natural sciences, and energy management, bilevel optimization can be found among the most active fields of research within the optimization community, see the survey papers [13, 14]. For an introduction to bilevel optimization, we refer the interested reader to the monographs [2, 12, 18, 52]. We emphasize that bilevel optimization problems are, in general, highly irregular and nonconvex optimization problems which are only implicitly given as a closed-form representation of the solution mapping $\widehat{\Psi}$ associated with (P(x)) is often not available. This causes the model (BPP) to be rather challenging from a theoretical as well as numerical perspective.

Since the set of efficient or weakly efficient points of a multiobjective optimization problem is, usually, not a singleton, the minimization of a scalar function over the set of efficient or weakly efficient points, in order to reduce the number of reasonable and economically utilizable points, is a nearby topic, see e.g. [6, 7, 9, 33], and such problems already possess a hierarchical structure. So-called semivectorial bilevel optimization problems, where only $p := 1$ and $\mathcal{K} := \mathbb{R}_+$ are demanded in (BPP), i.e., only the underlying parametric optimization problem (P(x)) possesses multiple objective functions, provide a much more general model paradigm and have been investigated, e.g., in [8, 10, 17, 19, 39, 57]. The even more general situation where the objective functions of both decision makers in (BPP) are allowed to be vector-valued has been considered, e.g., in [11, 24, 25, 27, 37, 50]. For formal completeness, let us also mention that the setting where $q := 1$ and $\mathcal{C} := \mathbb{R}_+$ hold while the objective function of (BPP) is vector-valued is also reasonable, see e.g. [26, 47, 54, 56]. It is well known from the literature that scalarization techniques can be used to transfer multiobjective optimization problems into scalar ones, see e.g. [36, Section 5]. Hence, whenever the lower level problem (P(x)) possesses multiple objective functions, it seems to be nearby to replace it with a scalarized counterpart and to treat the scalarization parameters as additional upper level variables. This procedure, however, comes with a price. It has been shown in the setting of semivectorial bilevel optimization that this reformulation induces additional local minimizers

which might be attractive for solution algorithms, see [19]. Even worse, the more detailed study [4] revealed that this transformation also induces additional stationary points while the constraint qualifications needed to tackle the reformulated problem might be stronger than the ones applicable to the original model. These observations are likely to address multiobjective bilevel optimization problems, too. Hence, one should not tackle $(P(x))$ with scalarization approaches for the theoretical and numerical treatment of (BPP) in a lightheaded way.

In this note, we are concerned with the so-called value function approach to bilevel optimization, which dates back to [46] in the scalar case. Let us, for a moment, assume that $(P(x))$ is a scalar problem, i.e., $q := 1$ and $\mathcal{C} := \mathbb{R}_+$. With the aid of the so-called lower level optimal value function $\varphi: \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ given by

$$\forall x \in \mathbb{R}^n: \quad \varphi(x) := \inf_y \{f(x, y) \mid y \in \Gamma(x)\}, \quad (1.1)$$

where $\overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, \infty\}$ and $\inf \emptyset := \infty$, one can equivalently reformulate (BPP) by means of

$$\min_{x,y} \mathcal{K} \{F(x, y) \mid x \in X, f(x, y) \leq \varphi(x), y \in \Gamma(x)\}, \quad (1.2)$$

as the feasible sets of (BPP) and (1.2) coincide. Noting that φ is only implicitly available while being inherently nonsmooth, and observing that classical constraint qualifications from nonsmooth programming are likely to fail at all feasible points of (1.2) , see [55, Proposition 3.2], (1.2) is still a rather challenging problem. Nevertheless, the so-called *value function reformulation* (1.2) has turned out to be useful for the construction of necessary optimality conditions and solution algorithms for (BPP) in the setting where, additionally, $p := 1$ and $\mathcal{K} := \mathbb{R}_+$, see e.g. [15, 16, 21, 43, 55] and the references therein.

Based on its successful application in the scalar situation, the paper [37] suggests an extension of the value function approach to multiobjective bilevel optimization. In order to recover the approach of [37], let us introduce the set-valued mapping $\widehat{\Phi}: \mathbb{R}^n \rightrightarrows \mathbb{R}^q$ by means of

$$\forall x \in \mathbb{R}^n: \quad \widehat{\Phi}(x) := f(x, \widehat{\Psi}(x)) := \{f(x, y) \mid y \in \widehat{\Psi}(x)\},$$

and observe that (BPP) is closely related to

$$\min_{x,y} \mathcal{K} \{F(x, y) \mid x \in X, f(x, y) \in \widehat{\Phi}(x), y \in \Gamma(x)\} \quad (VFR)$$

which, again, we will refer to as value function reformulation of (BPP) . Let us note at this point that (BPP) and (VFR) do not need to be equivalent anymore in general, see [Example 3.19](#). In [37], the authors exploit the model problem (VFR) for the derivation of necessary optimality conditions for the associated model (BPP) where the lower level parametric optimization problem is solved up to efficient points. More precisely, the authors of [37] transfer the classical arguments from [55], which address bilevel optimization problems with scalar upper and lower level objective functions, to the multiobjective situation. Among others, this makes the (generalized) differentiation of the so-called frontier mapping $\widehat{\Phi}$ from above necessary. Therefore, Mordukhovich's coderivative construction is exploited, see e.g. [42, 49]. Let us mention that the computation of coderivatives associated with frontier maps has already been studied in the literature before, see e.g. [34, 38, 53].

Now, we would like to mention an important and crucial observation which seemingly has been overseen in most of the aforementioned literature dealing with multiobjective bilevel optimization. If the lower level problem $(P(x))$ is solved up to efficiency, then the associated

solution mapping is likely to possess a graph which is not closed - even under comparatively strict assumptions. This non-closedness also addresses the associated frontier mapping which is used in the definition of the associated value function reformulation. Let us point out two immediate disadvantages which result from this non-closedness. First, it will be difficult to guarantee that (BPP) possesses efficient or at least weakly efficient points in this case. Second, for the computation of coderivatives, the underlying set-valued mapping needs to possess, at least locally around the reference point, a closed graph, so, in turn, this concept formally cannot be applied to the frontier mapping of interest in general.

In this note, we address both of the aforementioned issues and, along the way, comment on related peculiarities of multiobjective bilevel optimization. Throughout the paper, several examples are presented to illustrate our arguments. Furthermore, we compare our results with related findings addressing (1.2) in the scalar case. To start, let us mention that whenever $(P(x))$ is solved up to weak efficiency only, then the associated solution map is likely to possess a closed graph under mild assumptions, and this also extends to the associated frontier mapping. This can be distilled e.g. from [31, Section 3.5], but we provide a self-contained proof here for the purpose of completeness. In Section 3 of the paper, we are not only concerned with some formal properties of the value function reformulation, but also with the derivation of existence results for (BPP) which is a closely related issue. As already mentioned, this is reasonable under mild assumptions if $(P(x))$ is solved up to weak efficiency. We also show that the efficiency and associated frontier mapping of a fully linear parametric multiobjective optimization problem possess closed graphs, giving rise to yet another criterion for the existence of efficient and weakly efficient points for (BPP). Furthermore, we suggest an intermediate solution approach between efficiency and weak efficiency addressing $(P(x))$ which naturally comes along with a closed graph of the associated mappings $\widehat{\Psi}$ as well as $\widehat{\Phi}$ and, thus, the superordinate problem (BPP) is also likely to possess efficient and weakly efficient points. Section 4 is dedicated to the computation of upper coderivative estimates for the weak frontier mapping, i.e., the frontier mapping associated with $(P(x))$ when solved up to weak efficiency. First, we follow the approach promoted in [34], which relies on the so-called domination property, see e.g. [32,41], and point out some associated advantages and disadvantages. Furthermore, we illustrate a crucial error in [37, Theorem 4.1], that is concerned with the computation of an upper estimate for the coderivative of the efficiency-related frontier mapping of $(P(x))$. Unluckily, this shows that the results obtained in [37, Section 4] are not reliable. A second approach to coderivative estimates we discuss here is based on a weighted-sum scalarization of the problem $(P(x))$ in the convex situation. This idea is motivated by considerations from [57].

The remainder of the paper is organized as follows. In Section 2, we comment on the basic notation used in this paper, as well as some preliminaries from set-valued analysis and generalized differentiation. Furthermore, some foundations of multiobjective optimization are presented. Section 3 is dedicated to the detailed study of model problems in multiobjective bilevel optimization with a particular emphasis on value-function-type reformulations and their relations to each other. Based on a discussion of closedness properties of efficiency-type and associated frontier mappings in Section 3.1, some existence results are stated in Section 3.2. Furthermore, an intermediate way between efficiency and weak efficiency on how to interpret $(P(x))$ is suggested in Section 3.3. In Section 4, we are concerned with the derivation of upper coderivative estimates for frontier mappings. Based on a classical approach from the literature, we recover some known results in the novel setting of weak efficiency in Section 4.1. Relying on a scalarization approach, some new findings are obtained in Section 4.2. The paper closes with some concluding remarks in Section 5.

2 Preliminaries

In this section, we comment on the notation used in this paper. Particularly, some foundations of set-valued and variational analysis are provided. Furthermore, we present some necessary basics of multiobjective optimization.

2.1 Basic notation

The notation in this paper is fairly standard and mainly follows [42, 49].

Let \mathbb{R}_+ and \mathbb{R}_- represent the nonnegative and nonpositive real numbers, respectively. Throughout the paper, we equip \mathbb{R}^n , where $n \in \mathbb{N}$ is a positive integer, with the Euclidean inner product $\langle \cdot, \cdot \rangle: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ and the associated Euclidean norm $\|\cdot\|: \mathbb{R}^n \rightarrow \mathbb{R}_+$. We use $\mathbb{S}_1(0)$ to represent the unit sphere around the origin in \mathbb{R}^n where the dimension of the underlying space will be always clear from the context. For some set $\Omega \subset \mathbb{R}^n$, $\text{cl } \Omega$, $\text{int } \Omega$, $\text{conv } \Omega$, and $\text{cone } \Omega$ denote the closure, the interior, the convex hull, and the convex conic hull of Ω , respectively.

Whenever $v: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a mapping which is differentiable at $\bar{x} \in \mathbb{R}^n$, then $v'(\bar{x}) \in \mathbb{R}^{m \times n}$ denotes the Jacobian of v at \bar{x} . Partial derivatives are represented in analogous fashion.

2.1.1 Convexity properties of vector-valued functions

In this paper, we are partially concerned with convexity properties of set-valued mappings with respect to (w.r.t.) a given so-called ordering cone, see e.g. [36]. For a nonempty, closed, convex, pointed cone $\mathcal{K} \subset \mathbb{R}^p$, a mapping $F: \mathbb{R}^m \rightarrow \mathbb{R}^p$ is said to be \mathcal{K} -convex if

$$\forall y, y' \in \mathbb{R}^m \forall \alpha \in [0, 1]: \quad \alpha F(y) + (1 - \alpha)F(y') - F(\alpha y + (1 - \alpha)y') \in \mathcal{K}.$$

We note that whenever F is \mathcal{K} -convex and $Y \subset \mathbb{R}^m$ is convex, then $F(Y) + \mathcal{K}$ is convex, and the function $y \mapsto \langle \lambda, F(y) \rangle$ is convex for each $\lambda \in \mathcal{K}^*$. Here, we made use of

$$\mathcal{K}^* := \{\lambda \mid \forall z \in \mathcal{K}: \langle \lambda, z \rangle \geq 0\}$$

to denote the dual cone of \mathcal{K} which is a closed, convex cone as well. Let us mention that a componentwise convex mapping $F: \mathbb{R}^m \rightarrow \mathbb{R}^p$ is \mathbb{R}_+^p -convex.

The mapping F is said to be \mathcal{K} -strictly-quasiconvex if

$$\begin{aligned} \forall y, y' \in \mathbb{R}^m \forall \alpha \in (0, 1): \\ F(y') - F(y) \in \mathcal{K}, y \neq y' \implies F(y') - F(\alpha y + (1 - \alpha)y') \in \text{int } \mathcal{K}. \end{aligned}$$

Exemplary, in case where $\mathcal{K} := \mathbb{R}_+^p$, F is \mathbb{R}_+^p -strictly-quasiconvex if F_1, \dots, F_p are so-called semi-strictly quasiconvex in the sense that

$$\forall y, y' \in \mathbb{R}^m \forall \alpha \in (0, 1): \quad F_i(y') \geq F_i(y), y \neq y' \implies F_i(y') > F_i(\alpha y + (1 - \alpha)y')$$

has to hold for all $i = 1, \dots, p$. The latter is clearly inherent whenever the functions F_1, \dots, F_p are strictly convex.

2.1.2 Set-valued mappings

In this paper, we are concerned with so-called set-valued mappings which assign to each element of the pre-image space a (potentially empty) subset of the image space. Such objects will be denoted by $\Upsilon: \mathbb{R}^n \rightrightarrows \mathbb{R}^m$. Throughout the paper, we use

$$\text{gph } \Upsilon := \{(x, y) \mid y \in \Upsilon(x)\}, \quad \text{dom } \Upsilon := \{x \mid \Upsilon(x) \neq \emptyset\}$$

to represent the graph and the domain of Υ , respectively. For a set $S \subset \mathbb{R}^m$, we define the set-valued mappings $\Upsilon + S, \Upsilon - S: \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ by means of

$$\forall x \in \mathbb{R}^n: \quad (\Upsilon + S)(x) := \Upsilon(x) + S, \quad (\Upsilon - S)(x) := \Upsilon(x) - S$$

for brevity of notation. A set-valued mapping is referred to as polyhedral whenever its graph can be represented as the union of finitely many convex polyhedral sets.

For fixed $\bar{x} \in \text{dom } \Upsilon$, we call Υ closed at \bar{x} whenever for each sequence $((x^k, y^k))_{k \in \mathbb{N}} \subset \text{gph } \Upsilon$ and any $\bar{y} \in \mathbb{R}^m$ such that $x^k \rightarrow \bar{x}$ and $y^k \rightarrow \bar{y}$, we also have $\bar{y} \in \Upsilon(\bar{x})$. Note that this property is inherent whenever $\text{gph } \Upsilon$ is closed. Additionally, whenever Υ is closed at \bar{x} , $\Upsilon(\bar{x})$ is a closed set. Observe that closedness of $\text{gph } \Upsilon$ does not necessarily give closedness of $\text{dom } \Upsilon$. However, closedness of $\text{dom } \Upsilon$ and closedness of Υ at each point from $\text{dom } \Upsilon$ give closedness of $\text{gph } \Upsilon$. We refer to Υ as locally bounded at \bar{x} if there exist a bounded set $B \subset \mathbb{R}^m$ and some neighborhood $U \subset \mathbb{R}^n$ of \bar{x} such that $\Upsilon(x) \subset B$ for all $x \in U$. Furthermore, Υ is called lower semicontinuous at \bar{x} (in Berge's sense) w.r.t. $\Omega \subset \mathbb{R}^n$ whenever for each open set $V \subset \mathbb{R}^m$ such that $\Upsilon(\bar{x}) \cap V \neq \emptyset$, there is neighborhood $U \subset \mathbb{R}^n$ of \bar{x} such that $\Upsilon(x) \cap V \neq \emptyset$ for all $x \in \Omega \cap U$. For some fixed pair $(\bar{x}, \bar{y}) \in \text{gph } \Upsilon$, we say that Υ is inner semicontinuous at (\bar{x}, \bar{y}) w.r.t. $\Omega \subset \mathbb{R}^n$ whenever for each neighborhood $V \subset \mathbb{R}^m$ of \bar{y} , there exists a neighborhood $U \subset \mathbb{R}^n$ of \bar{x} such that $\Upsilon(x) \cap V \neq \emptyset$ for all $x \in \Omega \cap U$. If $\Omega := \mathbb{R}^n$ can be chosen, Υ is called inner semicontinuous at (\bar{x}, \bar{y}) for brevity. Note that Υ is lower semicontinuous at \bar{x} w.r.t. Ω if and only if it is inner semicontinuous at (\bar{x}, y) w.r.t. Ω for each $y \in \Upsilon(\bar{x})$. Furthermore, observe that Υ is inner semicontinuous at (\bar{x}, \bar{y}) w.r.t. Ω if and only if for each sequence $(x^k)_{k \in \mathbb{N}} \subset \Omega$ such that $x^k \rightarrow \bar{x}$, there exists a sequence $(y^k)_{k \in \mathbb{N}} \subset \mathbb{R}^m$ satisfying $y^k \rightarrow \bar{y}$ and $y^k \in \Upsilon(x^k)$ for all large enough $k \in \mathbb{N}$. Let us also mention that Υ is referred to as inner semicompact at \bar{x} w.r.t. Ω if for each sequence $(x^k)_{k \in \mathbb{N}} \subset \Omega$ with $x^k \rightarrow \bar{x}$, there are a subsequence $(x^{k_\ell})_{\ell \in \mathbb{N}}$ and a convergent sequence $(y^\ell)_{\ell \in \mathbb{N}}$ such that $y^\ell \in \Upsilon(x^{k_\ell})$ holds for all $\ell \in \mathbb{N}$. If $\Omega := \mathbb{R}^n$ can be chosen, Υ is called inner semicompact at \bar{x} . Clearly, if Υ is locally bounded at \bar{x} , then it is inner semicompact at \bar{x} w.r.t. each Ω . Furthermore, whenever there exists $\bar{y} \in \Upsilon(\bar{x})$ such that Υ is inner semicontinuous at (\bar{x}, \bar{y}) w.r.t. Ω , then Υ is inner semicompact at \bar{x} w.r.t. Ω . In the definitions of lower and inner semicontinuity as well as inner semicompactness, special emphasis is laid on settings where $\Omega \in \{\mathbb{R}^n, \text{dom } \Upsilon\}$.

In the subsequently stated lemma, we characterize the closedness of the graph of set-valued mappings of special type.

Lemma 2.1. *For a function $v: \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$, we consider the set-valued mapping $\Upsilon: \mathbb{R}^n \rightrightarrows \mathbb{R}$ given by*

$$\Upsilon(x) := \{v(x)\} - \mathbb{R}_+$$

with the conventions $\{\infty\} - \mathbb{R}_+ := \mathbb{R}$ and $\{-\infty\} - \mathbb{R}_+ := \emptyset$. Then $\text{gph } \Upsilon$ is closed if and only if v is upper semicontinuous.

Proof. We observe that $\text{gph } \Upsilon$ equals $\text{hypo } v := \{(x, \alpha) \in \mathbb{R}^n \times \mathbb{R} \mid \alpha \leq v(x)\}$, the so-called hypograph of v , and $\text{hypo } v$ is closed if and only if v is an upper semicontinuous function, see e.g. [49, Theorem 1.6]. \square

For two given set-valued mappings $\Upsilon_1: \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ and $\Upsilon_2: \mathbb{R}^m \rightrightarrows \mathbb{R}^p$, their composition $\Upsilon_2 \circ \Upsilon_1: \mathbb{R}^n \rightrightarrows \mathbb{R}^p$ is defined by

$$\forall x \in \mathbb{R}^n: \quad (\Upsilon_2 \circ \Upsilon_1)(x) := \bigcup_{y \in \Upsilon_1(x)} \Upsilon_2(y).$$

The associated so-called intermediate mapping $\Xi: \mathbb{R}^n \times \mathbb{R}^p \rightrightarrows \mathbb{R}^m$ is given by

$$\forall x \in \mathbb{R}^n \forall z \in \mathbb{R}^p: \quad \Xi(x, z) := \{y \in \mathbb{R}^m \mid z \in \Upsilon_2(y)\}$$

and plays an important role for the analysis of the underlying composition, see e.g. [5, Section 5.3] or [42, Section 3.1.2].

2.1.3 Variational analysis and generalized differentiation

The following definitions are taken from [42, 49].

For some set $\Omega \subset \mathbb{R}^n$, which is closed locally around some point $\bar{x} \in \Omega$, we call

$$N_\Omega(\bar{x}) := \left\{ \eta \left| \begin{array}{l} \exists (x^k)_{k \in \mathbb{N}}, (y^k)_{k \in \mathbb{N}} \subset \mathbb{R}^n \exists (\alpha_k)_{k \in \mathbb{N}} \subset (0, \infty): \\ x^k \rightarrow \bar{x}, \alpha_k(x^k - y^k) \rightarrow \eta, y^k \in \Pi_\Omega(x^k) \forall k \in \mathbb{N} \end{array} \right. \right\}$$

the limiting (or Mordukhovich) normal cone to Ω at \bar{x} . Above, $\Pi_\Omega: \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ denotes the projector of Ω which is given by

$$\forall x \in \mathbb{R}^n: \quad \Pi_\Omega(x) := \underset{y}{\operatorname{argmin}} \{\|y - x\| \mid y \in \Omega\}.$$

We note that $N_\Omega(\bar{x})$ is a closed cone which reduces to the classical normal cone of convex analysis as soon as Ω is a convex set.

For a set-valued mapping $\Upsilon: \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ and some point $(\bar{x}, \bar{y}) \in \text{gph } \Upsilon$ where $\text{gph } \Upsilon$ is locally closed, the set-valued mapping $D^*\Upsilon(\bar{x}, \bar{y}): \mathbb{R}^m \rightrightarrows \mathbb{R}^n$ given by

$$\forall y^* \in \mathbb{R}^m: \quad D^*\Upsilon(\bar{x}, \bar{y})(y^*) := \{x^* \mid (x^*, -y^*) \in N_{\text{gph } \Upsilon}(\bar{x}, \bar{y})\}$$

is referred to as the (limiting or Mordukhovich) coderivative of Υ at (\bar{x}, \bar{y}) . In case of a single-valued continuous mapping $v: \mathbb{R}^n \rightarrow \mathbb{R}^m$, which can naturally be interpreted as a singleton-valued set-valued mapping, and $\bar{x} \in \mathbb{R}^n$, we make use of $D^*v(\bar{x}) := D^*\Upsilon(\bar{x}, v(\bar{x}))$ for brevity of notation. If the mapping v is continuously differentiable at \bar{x} , then $D^*v(\bar{x})(y^*) = \{v'(\bar{x})^\top y^*\}$ is valid for all $y^* \in \mathbb{R}^m$.

2.2 Foundations of multiobjective optimization

For a continuous vector function $F: \mathbb{R}^m \rightarrow \mathbb{R}^p$, a nonempty, closed set $S \subset \mathbb{R}^m$, and some closed, convex, pointed cone $\mathcal{K} \subset \mathbb{R}^p$ with nonempty interior, we investigate the rather general multiobjective optimization problem

$$\min_{\mathcal{K}} \{F(y) \mid y \in S\}. \tag{MOP}$$

Recall that some feasible point $\bar{y} \in S$ of (MOP) is called efficient (weakly efficient) whenever

$$(\{F(\bar{y})\} - \mathcal{K} \setminus \{0\}) \cap F(S) = \emptyset \quad ((\{F(\bar{y})\} - \text{int } \mathcal{K}) \cap F(S) = \emptyset)$$

holds. The set of all efficient (weakly efficient) points of (MOP) will be denoted by $\mathcal{E}(F, S, \mathcal{K})$ ($\mathcal{E}_w(F, S, \mathcal{K})$). Further, $\mathcal{N}(F, S, \mathcal{K}) := F(\mathcal{E}(F, S, \mathcal{K}))$ ($\mathcal{N}_w(F, S, \mathcal{K}) := F(\mathcal{E}_w(F, S, \mathcal{K}))$) is referred to as the nondominated (weakly nondominated) set of (MOP). Let us mention that the computation of efficient (weakly efficient) points makes knowledge of F , S , and \mathcal{K} necessary, while, in principle, nondominated (weakly nondominated) points can be characterized from knowledge of $F(S)$ and \mathcal{K} only. Clearly, the inclusions $\mathcal{E}(F, S, \mathcal{K}) \subset \mathcal{E}_w(F, S, \mathcal{K})$ and $\mathcal{N}(F, S, \mathcal{K}) \subset \mathcal{N}_w(F, S, \mathcal{K})$ hold by definition. Let us also point out that continuity of F and the openness of $\text{int } \mathcal{K}$ give closedness of the sets $\mathcal{E}_w(F, S, \mathcal{K})$ and $\mathcal{N}_w(F, S, \mathcal{K})$. In contrast, the sets $\mathcal{E}(F, S, \mathcal{K})$ and $\mathcal{N}(F, S, \mathcal{K})$ do not need to be closed, see [Example 2.3](#) below. Let us also note that the closedness of $\mathcal{E}_w(F, S, \mathcal{K})$ and $\mathcal{N}_w(F, S, \mathcal{K})$ guarantees $\text{cl } \mathcal{E}(F, S, \mathcal{K}) \subset \mathcal{E}_w(F, S, \mathcal{K})$ and $\text{cl } \mathcal{N}(F, S, \mathcal{K}) \subset \mathcal{N}_w(F, S, \mathcal{K})$, and these inclusions can be strict. In case where $p := 1$, there is no difference between efficiency and weak efficiency as well as nondominance and weak nondominance, respectively, since the only cones satisfying our requirements would be \mathbb{R}_+ and \mathbb{R}_- in this case.

In this paper, we are concerned with so-called *domination* properties of (MOP), see e.g. [\[32, 40, 41\]](#).

Definition 2.2. *We say that (MOP) possesses the domination property (the weak domination property) if the following inclusion is valid:*

$$F(S) \subset \mathcal{N}(F, S, \mathcal{K}) + \mathcal{K} \quad \left(F(S) \subset \mathcal{N}_w(F, S, \mathcal{K}) + \mathcal{K} \right).$$

It is clear from the definition that validity of the domination property also gives validity of the weak domination property. Observe that, by convexity of \mathcal{K} , (MOP) possesses the domination property (the weak domination property) if and only if

$$F(S) + \mathcal{K} = \mathcal{N}(F, S, \mathcal{K}) + \mathcal{K} \quad \left(F(S) + \mathcal{K} = \mathcal{N}_w(F, S, \mathcal{K}) + \mathcal{K} \right).$$

Thus, the discussion at the beginning of [\[37, Section 4\]](#) is completely superfluous. Sufficient criteria for the domination and weak domination property can be found in [\[32, 40\]](#). Exemplary, let us mention that (MOP) possesses the domination property whenever F is continuous while S is compact. This follows by combining [\[32, Theorem 2\]](#) and [\[36, Theorem 3.38\]](#). Furthermore, we also want to point the reader's attention to the fact that even in the presence of the domination property, $\mathcal{E}(F, S, \mathcal{K})$ and $\mathcal{N}(F, S, \mathcal{K})$ might be non-closed.

Example 2.3. *Let $F: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the identity, let $S \subset \mathbb{R}^2$ be given by*

$$S := \{y \in \mathbb{R}_+^2 \mid y_1^2 + y_2^2 \geq 1\} \cup ([-1, 0] \times [1, \infty)),$$

and let $\mathcal{K} := \mathbb{R}_+^2$. Then we find

$$\mathcal{E}(F, S, \mathcal{K}) = \mathcal{N}(F, S, \mathcal{K}) = \{(\cos t, \sin t) \in \mathbb{R}^2 \mid t \in [0, \pi/2)\} \cup \{(-1, 1)\}.$$

On the one hand, this can be used to see that the associated problem (MOP) possesses the domination property. On the other hand, $\mathcal{E}(F, S, \mathcal{K})$ and $\mathcal{N}(F, S, \mathcal{K})$ are not closed.

It follows from [51, Example 3.4.2] that the efficient set $\mathcal{E}(F, S, \mathcal{K})$ might not be closed even if the set $F(S)$ is compact and convex. However, for instance in [48, Lemma 7.1], it is shown that if $\mathcal{K} := \mathbb{R}_+^p$ and all functions F_1, \dots, F_p are strictly quasiconvex while S is convex, then $\mathcal{E}(F, S, \mathcal{K})$ coincides with $\mathcal{E}_w(F, S, \mathcal{K})$ and so $\mathcal{E}(F, S, \mathcal{K})$ is closed since $\mathcal{E}_w(F, S, \mathcal{K})$ is closed. This result can also be extended to more general ordering cones.

Lemma 2.4. *Let F be \mathcal{K} -strictly-quasiconvex, and let S be convex. Then $\mathcal{E}(F, S, \mathcal{K}) = \mathcal{E}_w(F, S, \mathcal{K})$ and $\mathcal{N}(F, S, \mathcal{K}) = \mathcal{N}_w(F, S, \mathcal{K})$.*

Proof. We only show equivalence of the efficient and weakly efficient set. This automatically gives equivalence of the nondominated and weakly nondominated set. Since $\mathcal{E}(F, S, \mathcal{K}) \subset \mathcal{E}_w(F, S, \mathcal{K})$ holds in general, we only need to prove the converse inclusion. Fix $\bar{y} \in \mathcal{E}_w(F, S, \mathcal{K})$ and suppose that $\bar{y} \notin \mathcal{E}(F, S, \mathcal{K})$. Then we find some $\tilde{y} \in S \setminus \{\bar{y}\}$ satisfying $F(\tilde{y}) \in \{F(\bar{y})\} - \mathcal{K} \setminus \{0\}$. For all $\alpha \in (0, 1)$, $\alpha\tilde{y} + (1 - \alpha)\bar{y} \in S$ follows by convexity of S , and since F is \mathcal{K} -strictly-quasiconvex, we obtain

$$F(\alpha\tilde{y} + (1 - \alpha)\bar{y}) \in (\{F(\bar{y})\} - \text{int } \mathcal{K}) \cap F(S).$$

This is a contradiction to $\bar{y} \in \mathcal{E}_w(F, S, \mathcal{K})$. □

3 Value function models in multiobjective bilevel optimization

In order to make (BPP) explicit, one has to specify the precise meaning behind the optimization goals at the (upper and) lower level stage. Therefore, let us define set-valued mappings $\Psi, \Psi_w: \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ and $\Phi, \Phi_w: \mathbb{R}^n \rightrightarrows \mathbb{R}^q$ by means of

$$\begin{aligned} \Psi(x) &:= \mathcal{E}(f(x, \cdot), \Gamma(x), \mathcal{C}), & \Phi(x) &:= \mathcal{N}(f(x, \cdot), \Gamma(x), \mathcal{C}), \\ \Psi_w(x) &:= \mathcal{E}_w(f(x, \cdot), \Gamma(x), \mathcal{C}), & \Phi_w(x) &:= \mathcal{N}_w(f(x, \cdot), \Gamma(x), \mathcal{C}) \end{aligned}$$

for each $x \in \mathbb{R}^n$. Note that the identities

$$\text{dom } \Psi = \text{dom } \Phi, \quad \text{dom } \Psi_w = \text{dom } \Phi_w$$

hold by definition of these mappings. We will refer to Ψ and Ψ_w as the efficiency and weak efficiency mapping associated with (P(x)), respectively. Similarly, Φ and Φ_w are called the frontier and weak frontier mapping associated with (P(x)). For brevity of notation, we further introduce the set-valued mapping $\Sigma: \mathbb{R}^n \rightrightarrows \mathbb{R}^q$ given by

$$\forall x \in \mathbb{R}^n: \quad \Sigma(x) := f(x, \Gamma(x)). \tag{3.1}$$

In the course of this note, we are mainly concerned with the optimization problems

$$\begin{aligned} \min_{x,y} \mathcal{K} \{F(x, y) \mid x \in X, y \in \Psi(x)\}, & \quad (\mathcal{E}\text{-BPP}) \\ \min_{x,y} \mathcal{K} \{F(x, y) \mid x \in X, y \in \Psi_w(x)\} & \quad (\mathcal{E}_w\text{-BPP}) \end{aligned}$$

as well as their associated respective value function reformulations

$$\begin{aligned} \min_{x,y} \mathcal{K} \{F(x, y) \mid x \in X, f(x, y) \in \Phi(x), y \in \Gamma(x)\}, & \quad (\mathcal{E}\text{-VFR}) \\ \min_{x,y} \mathcal{K} \{F(x, y) \mid x \in X, f(x, y) \in \Phi_w(x), y \in \Gamma(x)\} & \quad (\mathcal{E}_w\text{-VFR}) \end{aligned}$$

which are apparently equivalent.

Proposition 3.1.

- (a) For each $x \in \mathbb{R}^n$, $\Psi(x) = \{y \in \Gamma(x) \mid f(x, y) \in \Phi(x)\}$ holds true.
- (b) For each $x \in \mathbb{R}^n$, $\Psi_w(x) = \{y \in \Gamma(x) \mid f(x, y) \in \Phi_w(x)\}$ holds true.

Proof. Let us start with the proof of the first assertion. Indeed, if $y \in \Psi(x)$ holds for some $x \in \mathbb{R}^n$, then $y \in \Gamma(x)$ and $f(x, y) \in f(x, \Psi(x)) = \Phi(x)$ follow trivially, i.e., the inclusion \subset is obvious. For the proof of the converse inclusion \supset , pick $y \in \Gamma(x)$ such that $f(x, y) \in \Phi(x)$ for some $x \in \mathbb{R}^n$. By definition of Φ , we find $y' \in \Psi(x)$ such that $f(x, y) = f(x, y')$. Since $(\{f(x, y')\} - \mathcal{C} \setminus \{0\}) \cap \Sigma(x) = \emptyset$, we also have $(\{f(x, y)\} - \mathcal{C} \setminus \{0\}) \cap \Sigma(x) = \emptyset$, and this gives $y \in \Psi(x)$.

The proof of the second assertion is completely analogous. □

Corollary 3.2.

- (a) Problems $(\mathcal{E}\text{-BPP})$ and $(\mathcal{E}\text{-VFR})$ possess the same feasible set.
- (b) Problems $(\mathcal{E}_w\text{-BPP})$ and $(\mathcal{E}_w\text{-VFR})$ possess the same feasible set.

Let us note that whenever $q := 1$ and $\mathcal{C} := \mathbb{R}_+$ hold, then both problems $(\mathcal{E}\text{-VFR})$ and $(\mathcal{E}_w\text{-VFR})$ are equivalent to

$$\min_{x,y} \kappa \{F(x, y) \mid x \in X, f(x, y) = \varphi(x), y \in \Gamma(x)\} \quad (3.4)$$

which, up to the relation sign in the constraint involving φ , is the same as the classical value function reformulation (1.2). In scalar bilevel programming, however, it has turned out to be beneficial to work with the constraint $f(x, y) \leq \varphi(x)$ as this opens the door to algorithmic applications and induces a sign condition for the Lagrange multiplier associated with this constraint in the context of nonsmooth optimality conditions. Noting that, for each $x \in \mathbb{R}^n$ and $y \in \Gamma(x)$, $f(x, y) < \varphi(x)$ is not possible, we can reformulate (1.2) and (3.4) as

$$\min_{x,y} \kappa \{F(x, y) \mid x \in X, f(x, y) \in \{\varphi(x)\} - \mathbb{R}_+, y \in \Gamma(x)\}, \quad (3.5)$$

with the conventions $\{\infty\} - \mathbb{R}_+ := \mathbb{R}$ and $\{-\infty\} - \mathbb{R}_+ := \emptyset$. Observing that \mathbb{R}_+ is the ordering cone associated with the lower level problem in the scalar situation, this motivates to investigate

$$\min_{x,y} \kappa \{F(x, y) \mid x \in X, f(x, y) \in \Phi(x) - \mathcal{C}, y \in \Gamma(x)\}, \quad (3.6a)$$

$$\min_{x,y} \kappa \{F(x, y) \mid x \in X, f(x, y) \in \Phi_w(x) - \mathcal{C}, y \in \Gamma(x)\} \quad (3.6b)$$

as potential surrogates of $(\mathcal{E}\text{-VFR})$ and $(\mathcal{E}_w\text{-VFR})$, respectively. As the following results show, the addition of the lower level ordering cone in (3.6a) and (3.6b) does not change the problem.

Proposition 3.3.

- (a) For each $x \in \mathbb{R}^n$, $\Psi(x) = \{y \in \Gamma(x) \mid f(x, y) \in \Phi(x) - \mathcal{C}\}$ holds true.
- (b) For each $x \in \mathbb{R}^n$, $\Psi_w(x) = \{y \in \Gamma(x) \mid f(x, y) \in \Phi_w(x) - \mathcal{C}\}$ holds true.

Proof. Let us start with the proof of the first assertion. Clearly, as $0 \in \mathcal{C}$, the inclusion \subset is clear from [Proposition 3.1](#). Conversely, fix $x \in \mathbb{R}^n$ and $y \in \Gamma(x)$ such that $f(x, y) \in \Phi(x) - \mathcal{C}$. Then there exist $y' \in \Psi(x)$ and $c \in \mathcal{C}$ such that $f(x, y) = f(x, y') - c$. By definition of efficiency, this gives $c = 0$, i.e., $f(x, y) = f(x, y') \in \Phi(x)$, and, hence, $y \in \Psi(x)$ by [Proposition 3.1](#).

In order to prove the second assertion, we observe that the inclusion \subset follows again from $0 \in \mathcal{C}$ and [Proposition 3.1](#). Thus, pick an arbitrary $x \in \mathbb{R}^n$ and $y \in \Gamma(x)$ such that $f(x, y) \in \Phi_w(x) - \mathcal{C}$. Then there exist $y' \in \Psi_w(x)$ and $c \in \mathcal{C}$ such that $f(x, y) = f(x, y') - c$. Suppose that $y \notin \Psi_w(x)$. Then there are $\tilde{y} \in \Gamma(x)$ and $\tilde{c} \in \text{int } \mathcal{C}$ such that $f(x, \tilde{y}) = f(x, y) - \tilde{c}$, and this gives $f(x, \tilde{y}) = f(x, y') - (c + \tilde{c})$. As we have $c + \tilde{c} \in \text{int } \mathcal{C}$, $y' \notin \Psi_w(x)$ follows which is a contradiction. Hence, we find $y \in \Psi_w(x)$ which yields $f(x, y) \in \Phi_w(x)$, i.e., $y \in \Psi_w(x)$ due to [Proposition 3.1](#). \square

Corollary 3.4.

- (a) Problems [\(E-VFR\)](#) and [\(3.6a\)](#) possess the same feasible set.
- (b) Problems [\(E_w-VFR\)](#) and [\(3.6b\)](#) possess the same feasible set.

In order to guarantee the existence of efficient or weakly efficient points for the above model problems [\(E-BPP\)](#), [\(E_w-BPP\)](#), [\(E-VFR\)](#), and [\(E_w-VFR\)](#), and to make their algorithmic treatment reasonable, it is important to guarantee that their respective feasible sets are nonempty and closed. Recalling that the mapping f is continuous, and the sets X and $\text{gph } \Gamma$ are assumed to be closed, this reduces to the closedness of $\text{gph } \Psi$, $\text{gph } \Psi_w$, $\text{gph } \Phi$, and $\text{gph } \Phi_w$, respectively. This is addressed in [Sections 3.1](#) and [3.2](#) below. As we will see, $\text{gph } \Psi$ and $\text{gph } \Phi$ are not likely to be closed in several popular settings. In [Section 3.3](#), we suggest an intermediate approach that bridges the model problems [\(E-BPP\)](#) and [\(E_w-BPP\)](#) (and, similarly, [\(E-VFR\)](#) and [\(E_w-VFR\)](#)) and comes along with a guaranteed closedness of the feasible set.

3.1 Closedness properties of efficiency and frontier mappings

In the following lemma, we show that Ψ_w as well as Φ_w are indeed closed under mild assumptions, see e.g. [\[31, Corollary 3.5.6\]](#) for a related result.

Lemma 3.5. *Fix $\bar{x} \in \text{dom } \Psi_w$ and let Γ be lower semicontinuous at \bar{x} w.r.t. its domain. Then Ψ_w is closed at \bar{x} . If, additionally, Γ is locally bounded at \bar{x} , Φ_w is closed at \bar{x} .*

Proof. Let $(x^k)_{k \in \mathbb{N}} \subset \mathbb{R}^n$ and $(y^k)_{k \in \mathbb{N}} \subset \mathbb{R}^m$ as well as $\bar{y} \in \mathbb{R}^m$ be chosen such that $x^k \rightarrow \bar{x}$, $y^k \rightarrow \bar{y}$, and $y^k \in \Psi_w(x^k)$ for all $k \in \mathbb{N}$. By definition of Ψ_w , we find $y^k \in \Gamma(x^k)$ and

$$(\{f(x^k, y^k)\} - \text{int } \mathcal{C}) \cap \Sigma(x^k) = \emptyset \quad (3.7)$$

for all $k \in \mathbb{N}$. Since Γ is closed at \bar{x} , we obtain $\bar{y} \in \Gamma(\bar{x})$. Suppose that $\bar{y} \notin \Psi_w(\bar{x})$. Then there are $\tilde{y} \in \Gamma(\bar{x})$ and $\bar{z} \in \text{int } \mathcal{C}$ such that $f(\bar{x}, \bar{y}) - \bar{z} = f(\bar{x}, \tilde{y})$. Noting that $(x^k)_{k \in \mathbb{N}} \subset \text{dom } \Psi_w \subset \text{dom } \Gamma$, the assumed lower semicontinuity of Γ yields the existence of a sequence $(\tilde{y}^k)_{k \in \mathbb{N}} \subset \mathbb{R}^m$ such that $\tilde{y}^k \rightarrow \tilde{y}$ and $\tilde{y}^k \in \Gamma(x^k)$ for all $k \in \mathbb{N}$. We set

$$z^k := f(x^k, y^k) - f(\bar{x}, \bar{y}) - f(x^k, \tilde{y}^k) + f(\bar{x}, \tilde{y}) + \bar{z}$$

for each $k \in \mathbb{N}$. Continuity of f gives us $z^k \rightarrow \bar{z}$, and due to $\bar{z} \in \text{int } \mathcal{C}$, $z^k \in \text{int } \mathcal{C}$ holds for all large enough $k \in \mathbb{N}$. Some rearrangements yield

$$f(x^k, y^k) - z^k = f(x^k, \tilde{y}^k) + (f(\bar{x}, \bar{y}) - \bar{z} - f(\bar{x}, \tilde{y})) = f(x^k, \tilde{y}^k),$$

i.e.,

$$(\{f(x^k, y^k)\} - \text{int } \mathcal{C}) \cap \Sigma(x^k) \neq \emptyset$$

for large enough $k \in \mathbb{N}$, contradicting (3.7). Hence, $\bar{y} \in \Psi_w(\bar{x})$ follows, i.e., Ψ_w is closed at \bar{x} .

In order to verify the closedness of Φ_w at \bar{x} , we pick sequences $(x^k)_{k \in \mathbb{N}} \subset \mathbb{R}^n$ and $(z^k)_{k \in \mathbb{N}} \subset \mathbb{R}^p$ such that $x^k \rightarrow \bar{x}$, $z^k \in \Phi_w(x^k)$ for all $k \in \mathbb{N}$, and $z^k \rightarrow \bar{z}$ for some $\bar{z} \in \mathbb{R}^p$. This gives the existence of $(y^k)_{k \in \mathbb{N}} \subset \mathbb{R}^m$ such that $y^k \in \Psi_w(x^k)$ and $f(x^k, y^k) = z^k$ for all $k \in \mathbb{N}$. Local boundedness of Γ at \bar{x} guarantees $y^k \rightarrow \bar{y}$ along a subsequence (without relabeling) for some $\bar{y} \in \mathbb{R}^m$. The first part of the proof shows $\bar{y} \in \Psi_w(\bar{x})$, and continuity of f yields $\bar{z} = f(\bar{x}, \bar{y})$, i.e., $\bar{z} \in \Phi_w(\bar{x})$, showing the closedness of Φ_w at \bar{x} . \square

As a corollary, we can state conditions which guarantee the closedness of $\text{gph } \Psi_w$ and $\text{gph } \Phi_w$.

Corollary 3.6. *Let $\text{dom } \Gamma$ be closed, and let Γ be locally bounded and lower semicontinuous w.r.t. its domain at each point from $\text{dom } \Gamma$. Then $\text{gph } \Psi_w$ and $\text{gph } \Phi_w$ are closed.*

Proof. The closedness of $\text{gph } \Gamma$ and the assumed local boundedness guarantee that, for each $\bar{x} \in \text{dom } \Gamma$, $\Gamma(\bar{x})$ is nonempty and compact. Thus, [36, Theorem 6.3] shows $\text{dom } \Psi_w = \text{dom } \Gamma$, i.e., $\text{dom } \Psi_w$ and $\text{dom } \Phi_w$ are closed. Furthermore, Lemma 3.5 shows that Ψ_w and Φ_w are closed at each point of their domain, respectively. Taking these facts together, the assertion follows. \square

Let us relate these findings to the classical results from [1] which address the scalar situation $q := 1$ and $\mathcal{C} := \mathbb{R}_+$. Then the closedness of Ψ_w corresponds to the closedness of the classical solution mapping while the closedness of Φ_w corresponds to continuity (w.r.t. the domain) of the so-called optimal value function. Consulting [1, Theorems 4.2.1, 4.2.2], these properties are obtained under continuity of the objective function and lower semicontinuity of the constraint mapping Γ . In this regard, Lemma 3.5 amounts to a reasonable generalization to the vector-valued situation.

Remark 3.7. *Recall that the efficient set as well as the associated nondominated set in multiobjective optimization do not need to be closed even in situations where all objective functions are continuous and the feasible set is compact. This issue naturally extends to the parametric setting. Typically, the set-valued mappings Ψ and Φ do not possess closed graphs in many situations.*

Let us inspect some simple examples.

Example 3.8. *We consider $(\mathbf{P}(x))$ with $n := m := 1$, $q := 2$, $\mathcal{C} := \mathbb{R}_+^2$, and*

$$\forall x, y \in \mathbb{R}: \quad f(x, y) := (\sin y, y), \quad \Gamma(x) := [0, 2\pi].$$

The efficiency map is given by

$$\forall x \in \mathbb{R}: \quad \Psi(x) = \{0\} \cup (\pi, 3\pi/2]$$

which is not closed for each $x \in \mathbb{R}$, whereas the weak efficiency map

$$\forall x \in \mathbb{R}: \quad \Psi_w(x) = \{0\} \cup [\pi, 3\pi/2]$$

is closed at each $x \in \mathbb{R}$.

Example 3.9. We consider $(\mathbf{P}(x))$ with $n := 1$, $m := 2$, $q := 2$, $\mathcal{C} := \mathbb{R}_+^2$, and

$$\forall x \in \mathbb{R} \forall y \in \mathbb{R}^2: \quad f(x, y) := y, \quad \Gamma(x) := \{y \mid y_1 + xy_2 \geq 0\}.$$

For $x > 0$, efficient points and weakly efficient points of $(\mathbf{P}(x))$ coincide. In case $x = 0$, there are no efficient point of $(\mathbf{P}(x))$, but all points from $\{0\} \times \mathbb{R}$ are weakly efficient. Moreover, for $x < 0$, there is neither an efficient nor a weakly efficient point. Hence, the efficiency map and frontier map are given by

$$\forall x \in \mathbb{R}: \quad \Psi(x) = \Phi(x) = \begin{cases} \{y \mid y_1 + xy_2 = 0\} & x > 0, \\ \emptyset & x \leq 0, \end{cases}$$

and these mappings do not possess closed graphs. Furthermore, the weak efficiency map and the weak frontier map are given by

$$\forall x \in \mathbb{R}: \quad \Psi_w(x) = \Phi_w(x) = \begin{cases} \{y \mid y_1 + xy_2 = 0\} & x \geq 0, \\ \emptyset & x < 0 \end{cases}$$

and possess closed graphs, see [Corollary 3.6](#) while noting that Γ is not locally bounded at each point $x \geq 0$.

Remark 3.10. Assume that, for each $x \in \text{dom } \Gamma$, $f(x, \cdot): \mathbb{R}^m \rightarrow \mathbb{R}^q$ is \mathcal{C} -strictly-quasiconvex while $\Gamma(x)$ is convex. Then, due to [Lemma 2.4](#), we find $\Psi(x) = \Psi_w(x)$ and $\Phi(x) = \Phi_w(x)$ for all $x \in \text{dom } \Gamma$, i.e., there is no difference between the models $(\mathcal{E}\text{-BPP})$ and $(\mathcal{E}_w\text{-BPP})$, $(\mathcal{E}\text{-VFR})$ and $(\mathcal{E}_w\text{-VFR})$, or [\(3.6a\)](#) and [\(3.6b\)](#), respectively. Particularly, [Lemma 2.4](#) and [Corollary 3.6](#) can be used to infer closedness results on Ψ and Φ .

In the following lemma, we show that the efficiency and frontier mapping of a fully linear multiobjective parametric optimization problem possess a closed graph, respectively.

Lemma 3.11. For matrices $A \in \mathbb{R}^{r \times n}$, $B \in \mathbb{R}^{r \times m}$, $D \in \mathbb{R}^{q \times m}$, and $d \in \mathbb{R}^r$, let $f: \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^q$ and $\Gamma: \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ be given by

$$\forall x \in \mathbb{R}^n \forall y \in \mathbb{R}^m: \quad f(x, y) := Dy, \quad \Gamma(x) := \{y \mid Ax + By \leq d\}. \quad (3.8)$$

Furthermore, let the ordering cone $\mathcal{C} := \mathbb{R}_+^q$ be fixed. Then the associated mappings Ψ and Φ possess closed graphs.

Proof. We prove the closedness of $\text{gph } \Psi$ by combining a classical scalarization argument and results from scalar linear parametric optimization. Afterwards, the closedness of $\text{gph } \Phi$ is shown via Hofmann's lemma.

Fix a sequence $((x^k, y^k))_{k \in \mathbb{N}} \subset \text{gph } \Psi$ and a pair $(\bar{x}, \bar{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ such that $(x^k, y^k) \rightarrow (\bar{x}, \bar{y})$. Let $k \in \mathbb{N}$ be fixed for the moment. Due to [\[23, Theorem 4.7\]](#), y^k is an optimal solution of the scalar linear optimization problem

$$\min_y \left\{ \mathbf{e}^\top Dy \mid Ax^k + By \leq d, Dy \leq Dy^k \right\} \quad (3.9)$$

where $\mathbf{e} \in \mathbb{R}^q$ denotes the all-ones vector. The dual of this problem is given by

$$\max_{u, v} \left\{ (d - Ax^k)^\top u + (Dy^k)^\top v \mid B^\top u + D^\top v = D^\top \mathbf{e}, u \leq 0, v \leq 0 \right\}. \quad (3.10)$$

As (3.9) possesses a solution, (3.10) is solvable as well.

Let us inspect (3.10) which can be interpreted as a linear parametric optimization problem which multiple parameters, namely x^k and y^k , which merely appear in the objective function. Based on [45, Sections 7, 8], the set of parameters where this parametric problem possesses a solution is a convex polyhedron and can be partitioned into finitely many so-called regions of stability, being convex polyhedra as well, such that, in the interior of any such region, precisely one of the vertices associated with the convex polyhedron

$$Q := \{(u, v) \mid B^\top u + D^\top v = D^\top \mathbf{e}, u \leq 0, v \leq 0\}$$

is optimal, while on the boundary, this vertex is still optimal but may not be the unique maximizer. Anyhow, since there exist only finitely many regions of stability, there is an infinite index set $N \subset \mathbb{N}$ such that all points from the subsequence $((x^k, y^k))_{k \in N}$ belong to the same region of stability which is associated with some fixed vertex $(\bar{u}, \bar{v}) \in \mathbb{R}^n \times \mathbb{R}^m$ of Q . As all regions of stability are closed, (\bar{u}, \bar{v}) does not only solve (3.10) for each $k \in N$ but also the problem

$$\max_{u, v} \left\{ (d - A\bar{x})^\top u + (D\bar{y})^\top v \mid (u, v) \in Q \right\}. \quad (3.11)$$

Strong duality of linear programming yields $\mathbf{e}^\top Dy^k = (d - Ax^k)^\top \bar{u} + (Dy^k)^\top \bar{v}$ for each $k \in N$, and taking the limit $k \rightarrow_N \infty$ gives $\mathbf{e}^\top D\bar{y} = (d - A\bar{x})^\top \bar{u} + (D\bar{y})^\top \bar{v}$. As \bar{y} is feasible to

$$\min_y \left\{ \mathbf{e}^\top Dy \mid A\bar{x} + By \leq d, Dy \leq D\bar{y} \right\},$$

which is the dual of (3.11), it already must be a minimizer. Applying [23, Theorem 4.7] once more gives $\bar{y} \in \Psi(\bar{x})$. This shows the closedness of $\text{gph } \Psi$.

In the remainder of the proof, we show that $\text{gph } \Phi$ is closed. Thus, let $(\bar{x}, \bar{z}) \in \mathbb{R}^n \times \mathbb{R}^q$ and $((x^k, z^k))_{k \in \mathbb{N}} \subset \text{gph } \Phi$ be chosen such that $(x^k, z^k) \rightarrow (\bar{x}, \bar{z})$. By definition of Φ , we find a sequence $(y^k)_{k \in \mathbb{N}} \subset \mathbb{R}^m$ such that $y^k \in \Psi(x^k)$ and $z^k = Dy^k$ hold for each $k \in \mathbb{N}$. Applying Hofmann's lemma, see e.g. [22, Lemma 3C.4], for each $k \in \mathbb{N}$ to the linear system

$$Dy \leq z^k, -Dy \leq -z^k, By \leq d - Ax^k$$

yields some $\tilde{y}^k \in \Gamma(x^k)$ which satisfies $D\tilde{y}^k = z^k$ and $\|\tilde{y}^k\| \leq \kappa(\|z^k\| + \|d - Ax^k\|)$ where the constant $\kappa > 0$ does not depend on k . Particularly, for each $k \in \mathbb{N}$, $D\tilde{y}^k = z^k = Dy^k$, i.e., $\tilde{y}^k \in \Psi(x^k)$. Further, by boundedness of $(x^k)_{k \in \mathbb{N}}$ and $(z^k)_{k \in \mathbb{N}}$, $(\tilde{y}^k)_{k \in \mathbb{N}}$ is bounded as well so we may assume $\tilde{y}^k \rightarrow \tilde{y}$ for some $\tilde{y} \in \mathbb{R}^m$. The first part of the proof gives $\tilde{y} \in \Psi(\bar{x})$, so that $D\tilde{y} = \bar{z}$ yields $\bar{z} \in \Phi(\bar{x})$. Hence, $\text{gph } \Phi$ is closed. \square

We note that the first part of the above proof has been inspired by classical arguments of Isermann who showed that all efficient points of a linear multiobjective optimization problem can be found with the aid of the weighted-sum scalarization technique where all weights are positive, see [35].

In the context of Proposition 3.3 and Corollary 3.4, the following result should be taken into account.

Lemma 3.12. *Let Γ be locally bounded at each point of its domain. Then the following assertions hold.*

- (a) *The set $\text{gph } \Phi$ is closed if and only if $\text{gph}(\Phi - \mathcal{C})$ is closed.*
- (b) *The set $\text{gph } \Phi_w$ is closed if and only if $\text{gph}(\Phi_w - \mathcal{C})$ is closed.*

Proof. We only prove the first assertion. The second one can be shown in analogous fashion while incorporating some arguments from the proof of [Proposition 3.3](#).

We show both implications separately.

[\implies] Let $\text{gph } \Phi$ be closed, and choose a sequence $((x^k, z^k))_{k \in \mathbb{N}} \subset \text{gph}(\Phi - \mathcal{C})$ such that $(x^k, z^k) \rightarrow (\bar{x}, \bar{z})$ for some $(\bar{x}, \bar{z}) \in \mathbb{R}^n \times \mathbb{R}^q$. By definition, we find sequences $(c^k)_{k \in \mathbb{N}} \subset \mathcal{C}$ and $(y^k)_{k \in \mathbb{N}} \subset \mathbb{R}^m$ such that $z^k = f(x^k, y^k) - c^k$ and $y^k \in \Psi(x^k)$, i.e., $(x^k, f(x^k, y^k)) \in \text{gph } \Phi$, for each $k \in \mathbb{N}$. As the sequence $((x^k, y^k))_{k \in \mathbb{N}} \subset \text{gph } \Gamma$ is bounded by assumption, it possesses an accumulation point (\bar{x}, \bar{y}) which, by continuity of f and closedness of $\text{gph } \Phi$, satisfies $f(\bar{x}, \bar{y}) \in \Phi(\bar{x})$. Furthermore, $(c^k)_{k \in \mathbb{N}}$ possesses the accumulation point $f(\bar{x}, \bar{y}) - \bar{z}$ which, by closedness of \mathcal{C} , belongs to \mathcal{C} . Hence, we have shown $\bar{z} \in (\Phi - \mathcal{C})(\bar{x})$, and closedness of $\text{gph}(\Phi - \mathcal{C})$ follows.

[\impliedby] Let $\text{gph}(\Phi - \mathcal{C})$ be closed, and choose a sequence $((x^k, z^k))_{k \in \mathbb{N}} \subset \text{gph } \Phi$ such that $(x^k, z^k) \rightarrow (\bar{x}, \bar{z})$ for some $(\bar{x}, \bar{z}) \in \mathbb{R}^n \times \mathbb{R}^q$. On the one hand, since $0 \in \mathcal{C}$, we also have $((x^k, z^k))_{k \in \mathbb{N}} \subset \text{gph}(\Phi - \mathcal{C})$, and $(\bar{x}, \bar{z}) \in \text{gph}(\Phi - \mathcal{C})$ follows, i.e., there are $\bar{y} \in \Psi(\bar{x})$ and $c \in \mathcal{C}$ such that $\bar{z} = f(\bar{x}, \bar{y}) - c$. On the other hand, there is a sequence $(y^k)_{k \in \mathbb{N}} \subset \mathbb{R}^m$ such that $z^k = f(x^k, y^k)$ and $y^k \in \Psi(x^k)$ for all $k \in \mathbb{N}$. As the sequence $((x^k, y^k))_{k \in \mathbb{N}} \subset \text{gph } \Gamma$ is bounded by assumption, it possesses an accumulation point (\bar{x}, \bar{y}) which, by closedness of $\text{gph } \Gamma$, belongs to $\text{gph } \Gamma$. Combining these observations, we have $f(\bar{x}, \bar{y}) = \bar{z} = f(\bar{x}, \bar{y}) - c$. Now, $\bar{y} \in \Psi(\bar{x})$ guarantees $c = 0$, i.e., $\bar{z} = f(\bar{x}, \bar{y}) \in \Phi(\bar{x})$, and closedness of $\text{gph } \Phi$ follows. \square

3.2 Existence results in multiobjective bilevel optimization

The following example shows that the closedness of $\text{gph } \hat{\Psi}$ is, indeed, essential for the existence of efficient and weakly efficient points of [\(BPP\)](#).

Example 3.13. We consider [\(BPP\)](#) with $n := m := 1$, $p := 2$, $\mathcal{K} := \mathbb{R}_+^2$, $X := \mathbb{R}$, and

$$\forall x, y \in \mathbb{R}: \quad F(x, y) := (\cos(y) + 1, (y - \pi)^2).$$

Furthermore, we assume that $\hat{\Psi} \in \{\Psi, \Psi_w\}$ is the efficiency or weak efficiency map of the parametric optimization problem from [Example 3.8](#), i.e.,

$$\forall x \in \mathbb{R}: \quad \Psi(x) = \{0\} \cup (\pi, 3\pi/2], \quad \Psi_w(x) = \{0\} \cup [\pi, 3\pi/2].$$

In case $\hat{\Psi} := \Psi$, there is neither an efficient nor a weakly efficient point of [\(BPP\)](#). However, in case $\hat{\Psi} := \Psi_w$, $\mathbb{R} \times \{\pi\}$ is a feasible subset of [\(BPP\)](#), and these are precisely the efficient and, simultaneously, weakly efficient points of the problem.

The following result addresses the existence of efficient and weakly efficient point of [\(BPP\)](#).

Theorem 3.14. Let $\text{gph } \hat{\Psi}$ be closed and $\text{gph } \hat{\Psi} \cap (X \times \mathbb{R}^m)$ be nonempty and bounded. Then there exist an efficient and a weakly efficient point of [\(BPP\)](#).

Proof. Problem [\(BPP\)](#) is equivalent to

$$\min_{x, y} \kappa \{F(x, y) \mid (x, y) \in X \times \mathbb{R}^m, (x, y) \in \text{gph } \hat{\Psi}\}. \quad (3.12)$$

Due to the postulated assumptions, the set $\text{gph } \hat{\Psi} \cap (X \times \mathbb{R}^m)$ is nonempty, bounded, and closed and, thus, compact. Then $F(\text{gph } \hat{\Psi} \cap (X \times \mathbb{R}^m))$ is compact because F is continuous. The existence of an efficient point of [\(3.12\)](#), which then is efficient and, thus, weakly efficient for [\(BPP\)](#), follows by [\[36, Theorem 6.3\]](#). \square

Let us note that the boundedness assumption on $\text{gph } \widehat{\Psi} \cap (X \times \mathbb{R}^m)$ in [Theorem 3.14](#) is inherently satisfied whenever X and $\bigcup_{x \in X} \Gamma(x)$ are bounded.

Based on [Corollary 3.6](#), we obtain from [Theorem 3.14](#) an existence result for efficient and weakly efficient points of $(\mathcal{E}_w\text{-BPP})$.

Corollary 3.15. *Let $\text{dom } \Gamma$ be closed, and let Γ be locally bounded and lower semicontinuous w.r.t. its domain at each point from $\text{dom } \Gamma$. Let $\text{gph } \Psi_w \cap (X \times \mathbb{R}^m)$ be nonempty and bounded. Then there exist an efficient and a weakly efficient point of $(\mathcal{E}_w\text{-BPP})$.*

We note that [Corollary 3.15](#) generalizes [19, Theorem 5.5] to multiobjective bilevel optimization problems, but does not cover the situation were the lower level variable comes from an infinite-dimensional space.

Similar as in [Corollary 3.15](#), [Theorem 3.14](#) gives an existence result for efficient and weakly efficient points of $(\mathcal{E}\text{-BPP})$ and $(\mathcal{E}_w\text{-BPP})$ whenever the lower level data is fully linear as described in [\(3.8\)](#).

Corollary 3.16. *Let the lower level data be given as in [\(3.8\)](#) and fix $\mathcal{C} := \mathbb{R}_+^q$. Then the following assertions hold.*

- (a) *Let $\text{gph } \Psi \cap (X \times \mathbb{R}^m)$ be nonempty and bounded. Then there exist an efficient and a weakly efficient point of $(\mathcal{E}\text{-BPP})$.*
- (b) *Let $\text{gph } \Psi_w \cap (X \times \mathbb{R}^m)$ be nonempty and bounded. Then there exist an efficient and a weakly efficient point of $(\mathcal{E}_w\text{-BPP})$.*

Proof. The first assertion directly follows from [Lemma 3.11](#) and [Theorem 3.14](#). The second statement can be obtained from [Corollary 3.6](#) and [Theorem 3.14](#) as $\text{gph } \Gamma$ is a convex polyhedral set and, thus, closed, and these properties also ensure that Γ is lower semicontinuous w.r.t. its domain at each point from $\text{dom } \Gamma$, see e.g. [49, Example 9.35]. \square

Remark 3.17. *Taking a look back at the optimal value function reformulation for standard scalar bilevel optimization problems, we know from [Lemma 2.1](#) that the feasible set of [\(3.5\)](#) is closed if and only if φ is upper semicontinuous, and this can also be seen from [\(1.2\)](#) as sublevel sets of extended-real-valued functions are closed if and only if they are lower semicontinuous. The latter property is clearly milder than the continuity of φ on its domain $\{x \mid |\varphi(x)| < \infty\}$ which seems to be necessary for this desirable closedness when focusing on the model problem [\(3.4\)](#).*

Thus, having problems [\(3.6a\)](#) and [\(3.6b\)](#) as well as [Corollary 3.4](#) at hand, it is seemingly reasonable to work out conditions which ensure that the graphs of $\Phi - \mathcal{C}$ and $\Phi_w - \mathcal{C}$ are closed. This is enough to obtain the closedness of the feasible sets of [\(3.6a\)](#) and [\(3.6b\)](#) which, in turn, provides the basis for further existence results in multiobjective bilevel optimization. However, note that due to [Lemma 3.12](#), it turns out that whenever the feasibility mapping Γ is locally bounded, then this apparently milder approach yields no new insights. Observe that this is in accordance with the aforementioned scalar situation as the optimal value function φ is lower semicontinuous on its domain whenever Γ is locally bounded.

3.3 An intermediate approach to multiobjective bilevel optimization

On the one hand, the feasible set of $(\mathcal{E}\text{-BPP})$ is likely to be much smaller than the one of $(\mathcal{E}_w\text{-BPP})$ since the solution criterion behind Ψ is sharper than the one behind Ψ_w . On the other hand, from the viewpoint of existence theory, necessary optimality conditions, and

solution algorithms, a closed feasible set is desirable. The latter is given for $(\mathcal{E}_w\text{-BPP})$ under mild assumptions, see [Corollary 3.6](#), while the feasible set of $(\mathcal{E}\text{-BPP})$ is not likely to be closed apart from selected situations.

In this subsection, we suggest yet another possibility for the choice of $\widehat{\Psi}$ in the general multiobjective bilevel optimization problem (BPP) which induces an intermediate model between $(\mathcal{E}\text{-BPP})$ and $(\mathcal{E}_w\text{-BPP})$. To start, however, we would like to suggest an intermediate value function model which bridges $(\mathcal{E}\text{-VFR})$ and $(\mathcal{E}_w\text{-VFR})$ to some extent. There are two reasons for this approach. First, it allows for a simple definition of an intermediate solution map such that the associated models (BPP) and (VFR) are equivalent. Second, noting that, in order to determine efficient and weakly efficient points of some multiobjective optimization problem, one has to compute its nondominated and weakly nondominated points first before identifying their pre-images in the feasible set, this procedure seems the more natural one.

Therefore, let us introduce $\overline{\Phi}: \mathbb{R}^n \rightrightarrows \mathbb{R}^q$ by means of

$$\text{gph } \overline{\Phi} := \text{cl gph } \Phi$$

and consider

$$\min_{x,y} \kappa \{F(x,y) \mid x \in X, f(x,y) \in \overline{\Phi}(x), y \in \Gamma(x)\}. \quad (\overline{\mathcal{E}}\text{-VFR})$$

As $\text{gph } \Phi_w$ is closed under mild assumptions, see e.g. [Corollary 3.6](#), the inclusions $\text{gph } \Phi \subset \text{gph } \overline{\Phi} \subset \text{gph } \Phi_w$ are likely to hold and would give $\Phi(x) \subset \overline{\Phi}(x) \subset \Phi_w(x)$ for each $x \in \mathbb{R}^n$. Hence, $(\overline{\mathcal{E}}\text{-VFR})$ can be interpreted as an intermediate problem between $(\mathcal{E}\text{-VFR})$ and $(\mathcal{E}_w\text{-VFR})$ since the feasible set of $(\overline{\mathcal{E}}\text{-VFR})$ is not smaller than the one of $(\mathcal{E}\text{-VFR})$ and likely to be not larger than the one of $(\mathcal{E}_w\text{-VFR})$. Furthermore, $(\overline{\mathcal{E}}\text{-VFR})$ possesses a closed feasible set by construction of $\overline{\Phi}$, and, thus, efficient and weakly efficient points of $(\overline{\mathcal{E}}\text{-VFR})$ exist under standard assumptions.

Based on $\overline{\Phi}$, let us introduce $\overline{\Psi}: \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ by means of

$$\forall x \in \mathbb{R}^n: \quad \overline{\Psi}(x) := \{y \in \Gamma(x) \mid f(x,y) \in \overline{\Phi}(x)\}.$$

In the subsequent lemma, we comment on the properties of $\overline{\Psi}$. The simple proof follows by definition of $\overline{\Psi}$ as well as our above discussion on the properties of $\overline{\Psi}$ and is, thus, omitted.

Lemma 3.18.

- (a) *The set $\text{gph } \overline{\Psi}$ is closed.*
- (b) *Whenever $\text{gph } \Phi_w$ is closed, then the inclusions $\text{gph } \Psi \subset \text{gph } \overline{\Psi} \subset \text{gph } \Psi_w$ hold. Particularly, $\Psi(x) \subset \overline{\Psi}(x) \subset \Psi_w(x)$ is valid for each $x \in \mathbb{R}^n$ in this situation.*
- (c) *We have $\text{cl gph } \Psi \subset \text{gph } \overline{\Psi}$.*

By definition of $\overline{\Psi}$, the problem

$$\min_{x,y} \kappa \{F(x,y) \mid x \in X, y \in \overline{\Psi}(x)\} \quad (\overline{\mathcal{E}}\text{-BPP})$$

is equivalent to $(\overline{\mathcal{E}}\text{-VFR})$, and due to [Lemma 3.18 \(b\)](#), can be seen as an intermediate model between $(\mathcal{E}\text{-BPP})$ and $(\mathcal{E}_w\text{-BPP})$. Clearly, $(\overline{\mathcal{E}}\text{-BPP})$ has a closed feasible set and, due to [Theorem 3.14](#), is likely to possess efficient and weakly efficient points which can be interpreted as approximately efficient and approximately weakly efficient for $(\mathcal{E}\text{-BPP})$.

Let us point out that one does not obtain equality in [Lemma 3.18 \(c\)](#) in general situations. Indeed, when defining $\tilde{\Psi}: \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ by $\text{gph } \tilde{\Psi} := \text{cl gph } \Psi$ and $\tilde{\Phi}: \mathbb{R}^n \rightrightarrows \mathbb{R}^q$ by $\tilde{\Phi}(x) :=$

$f(x, \tilde{\Psi}(x))$ for all $x \in \mathbb{R}^n$, then the corresponding problems **(BPP)** and **(VFR)** for $\hat{\Psi} := \tilde{\Psi}$ and $\hat{\Phi} := \tilde{\Phi}$, respectively, are not necessarily equivalent. We omitted this undesirable behavior by starting with the definition of a suitable intermediate frontier mapping before introducing an intermediate efficiency mapping.

Example 3.19. We consider **(P(x))** with $n := 1$, $m := q := 2$, $\mathcal{C} := \mathbb{R}_+^2$, and

$$\forall x \in \mathbb{R} \forall y \in \mathbb{R}^2: \quad f(x, y) := (\sin y_1, y_2), \quad \Gamma(x) := \text{conv}\{(0, 0), (2\pi, 2\pi)\} \cup \{(0, \pi)\}$$

We easily find

$$\begin{aligned} \forall x \in \mathbb{R}: \quad \Psi(x) &= \{y \mid y_1 = y_2, y_2 \in \{0\} \cup (\pi, 3\pi/2]\}, \\ \Phi(x) &= \{(\sin z, z) \mid z \in \{0\} \cup (\pi, 3\pi/2]\} \end{aligned}$$

which gives

$$\begin{aligned} \forall x \in \mathbb{R}: \quad \tilde{\Psi}(x) &= \{y \mid y_1 = y_2, y_2 \in \{0\} \cup [\pi, 3\pi/2]\}, \\ \tilde{\Phi}(x) &= \{(\sin z, z) \mid z \in \{0\} \cup [\pi, 3\pi/2]\}. \end{aligned}$$

Observe, however, that

$$\forall x \in \mathbb{R}: \quad \{y \in \Gamma(x) \mid f(x, y) \in \tilde{\Phi}(x)\} = \tilde{\Psi}(x) \cup \{(0, \pi)\},$$

i.e., the feasible set of any superordinate model **(BPP)** for $\hat{\Psi} := \tilde{\Psi}$ is likely to be smaller than the feasible set of the associated value function reformulation **(VFR)** for $\hat{\Phi} := \tilde{\Phi}$.

Let us also mention that

$$\forall x \in \mathbb{R}: \quad \bar{\Phi}(x) = \{(\sin z, z) \mid z \in \{0\} \cup [\pi, 3\pi/2]\} = \tilde{\Phi}(x), \quad \bar{\Psi}(x) = \tilde{\Psi}(x) \cup \{(0, \pi)\}$$

holds in this example. This also shows $\text{cl gph } \Psi \subsetneq \text{gph } \bar{\Psi}$.

The following example shows that **(E-BPP)**, **(E_w-BPP)**, and **(Ē-BPP)** may possess pairwise different solutions.

Example 3.20. Motivated by **Example 2.3**, for $n := 1$, $m := q := 2$, and $\mathcal{C} := \mathbb{R}_+^2$, we consider the parametric multiobjective optimization problem **(P(x))** with the data

$$\forall x \in \mathbb{R} \forall y \in \mathbb{R}^2: \quad f(x, y) := y, \quad \Gamma(x) := \{y \in \mathbb{R}_+^2 \mid y_1^2 + y_2^2 \geq x^2\} \cup ([-1, 0] \times [|x|, \infty)).$$

A simple calculation reveals

$$\begin{aligned} \forall x \in \mathbb{R}: \quad \Psi(x) &= \begin{cases} \{(|x| \cos t, |x| \sin t) \mid t \in [0, \pi/2)\} \cup \{(-1, |x|)\} & x \neq 0, \\ \{(-1, 0)\} & x = 0, \end{cases} \\ \Psi_w(x) &= \{(|x| \cos t, |x| \sin t) \mid t \in [0, \pi/2]\} \cup ([-1, 0] \times \{|x|\}) \\ &\quad \cup ([|x|, \infty) \times \{0\}) \cup (\{-1\} \times [|x|, \infty)), \\ \bar{\Psi}(x) &= \{(|x| \cos t, |x| \sin t) \mid t \in [0, \pi/2]\} \cup \{(-1, |x|)\}. \end{aligned}$$

Observe that $\text{gph } \bar{\Psi} = \text{gph } \Psi \cup \{(x, 0, |x|) \mid x \in \mathbb{R}\}$ holds true.

Now, we consider the superordinate semivectorial bilevel optimization problem

$$\min_{x, y} \{(y_1 + 1/4)^2 + y_2^2 \mid x \geq 1, y \in \hat{\Psi}(x)\}$$

where $\widehat{\Psi} \in \{\Psi, \Psi_w, \overline{\Psi}\}$. Let us start with the investigation of $\widehat{\Psi} := \Psi$. The associated problem $(\mathcal{E}\text{-BPP})$ does not possess a global minimizer, but there is a local minimizer at $(1, (-1, 1))$. Whenever $\widehat{\Psi} := \Psi_w$, the associated problem $(\mathcal{E}_w\text{-BPP})$ possesses the uniquely determined global minimizer $(1, (-1/4, 1))$, and no other local minimizers exist. Finally, for $\widehat{\Psi} := \overline{\Psi}$, the associated problem $(\overline{\mathcal{E}}\text{-BPP})$ possesses the uniquely determined global minimizer $(1, (0, 1))$ (which is not feasible to $(\mathcal{E}\text{-BPP})$) and an additional local minimizer at $(1, (-1, 1))$.

Finally, we would like to point the reader's attention to the fact that, in contrast to [Corollary 3.4](#), $(\overline{\mathcal{E}}\text{-VFR})$ and

$$\min_{x,y} \kappa\{F(x,y) \mid x \in X, f(x,y) \in \overline{\Phi}(x) - \mathcal{C}, y \in \Gamma(x)\}. \quad (3.13)$$

are not necessarily equivalent as the feasible set of [\(3.13\)](#) is likely to be larger than the one of $(\overline{\mathcal{E}}\text{-VFR})$. However, if $\text{gph } \Phi_w$ is closed, the feasible set of [\(3.13\)](#) is included in the feasible set of [\(3.6b\)](#) which, by [Corollary 3.4](#), equals the feasible set of $(\mathcal{E}_w\text{-VFR})$. Thus, the enlargement of the feasible set in [\(3.13\)](#) in comparison with $(\overline{\mathcal{E}}\text{-VFR})$ is somewhat controllable.

Example 3.21. Let us reinspect the semivectorial bilevel optimization problem from [Example 3.20](#). One can easily check that, for $\bar{x} := 1, \bar{y} := (-1/4, 1) \in \Gamma(\bar{x})$ satisfies

$$f(\bar{x}, \bar{y}) \in \overline{\Phi}(\bar{x}) - \mathbb{R}_+^2 = \overline{\Psi}(\bar{x}) - \mathbb{R}_+^2,$$

but (\bar{x}, \bar{y}) is not feasible for the associated problem $(\overline{\mathcal{E}}\text{-BPP})$ and, thus, $(\overline{\mathcal{E}}\text{-VFR})$. Keeping our above discussion in mind and noticing that $\text{gph } \Phi_w$ is closed, the associated model [\(3.13\)](#) possesses the global minimizer (\bar{x}, \bar{y}) , see [Example 3.20](#) as well.

4 Towards necessary optimality conditions

For the derivation of necessary optimality conditions and constraint qualifications for $(\mathcal{E}\text{-BPP})$ and $(\mathcal{E}_w\text{-BPP})$ via $(\mathcal{E}\text{-VFR})$ and $(\mathcal{E}_w\text{-VFR})$, respectively, in terms of the limiting tools of generalized differentiation, it is important to have formulas or at least upper estimates for the coderivative of Φ and Φ_w available. Here, we focus on the latter and comment on two approaches which can be used to obtain such estimates in terms of initial problem data.

4.1 Coderivative estimates via the domination property

In the literature, one can find several papers dealing with the derivation of upper coderivative estimates for the frontier mapping Φ in cases where $(\mathbf{P}(x))$ possesses the domination property for all $x \in \mathbb{R}^n$ from a neighborhood of the reference point, see e.g. [\[34, 38, 53\]](#). Apart from the fact that these papers completely ignore the fact that $\text{gph } \Phi$ is likely to be non-closed, so that the coderivative is not well-defined, they obtain validity of the upper estimate

$$D^*\Phi(\bar{x}, \bar{z})(z^*) \subset D^*(\Sigma + \mathcal{C})(\bar{x}, \bar{z})(z^*) \quad (4.1)$$

only for points $(\bar{x}, \bar{z}) \in \text{gph } \Phi$ and $z^* \in \mathcal{C}_>$ where

$$\mathcal{C}_> := \{z^* \in \mathbb{R}^q \mid \forall z \in \mathcal{C} \cap \mathbb{S}_1(0): \langle z^*, z \rangle > 0\}.$$

In practice, this means that the estimate [\(4.1\)](#) is, more or less, only reasonable for arguments $z^* \in \mathbb{R}^q$ from $\text{int } \mathcal{C}^*$. For the derivation of necessary optimality conditions for $(\mathcal{E}\text{-BPP})$ via $(\mathcal{E}\text{-VFR})$, however, this additional restriction on the variable z^* might be too restrictive.

In the proof of [37, Theorem 4.1], the authors claim that for $\mathcal{C} := \mathbb{R}_+^q$, one has the estimate

$$D^*\Phi(\bar{x}, \bar{z})(z^*) \subset D^*\Sigma(\bar{x}, \bar{z})(z^*) \quad (4.2)$$

for all $z^* \in \mathbb{R}^q$. This estimate, however, does not depend on the choice of the ordering cone \mathcal{C} which is somewhat dubious. As the following example shows, this estimate is indeed false, and so are [37, Theorems 4.1, 4.4].

Example 4.1. We consider $(\mathbf{P}(x))$ with $n := 1$, $m := q := 2$, $\mathcal{C} := \mathbb{R}_+^2$, and

$$\forall x \in \mathbb{R} \forall y \in \mathbb{R}^2: \quad f(x, y) := y, \quad \Gamma(x) := \{y \mid y_1 + 2y_2 \geq 0, 2y_1 + y_2 \geq 0, y_1, y_2 \leq 2\},$$

i.e., the associated problem $(\mathbf{P}(x))$ does not depend on the parameter. We find

$$\forall x \in \mathbb{R}: \quad \Sigma(x) = \Gamma(x), \quad \Phi(x) = \Psi(x) = \text{conv}\{(-1, 2), (0, 0)\} \cup \text{conv}\{(2, -1), (0, 0)\}$$

and observe that $(\mathbf{P}(x))$ possesses the domination property (for each $x \in \mathbb{R}$). Let us fix $\bar{x} := 0$ and $\bar{z} := (0, 0)$. Then we find

$$\begin{aligned} N_{\text{gph } \Sigma}(\bar{x}, \bar{z}) &= \text{cone}\{(0, -1, -2), (0, -2, -1)\}, \\ N_{\text{gph } \Phi}(\bar{x}, \bar{z}) &= \text{cone}\{(0, -1, -2), (0, -2, -1)\} \cup \text{cone}\{(0, 2, 1)\} \cup \text{cone}\{(0, 1, 2)\}. \end{aligned}$$

This can be easily seen from $\text{gph } \Sigma = \mathbb{R} \times \Sigma(0)$ and $\text{gph } \Phi = \mathbb{R} \times \Phi(0)$, the product rule for the computation of limiting normals, see e.g. [42, Proposition 1.2], and a simple calculation of the normal cones to $\Sigma(0)$ (which is convex) and $\Phi(0)$ in \mathbb{R}^2 . Hence, for $z^* := (-1, -2)$, we find

$$D^*\Phi(\bar{x}, \bar{z})(z^*) = \{0\}, \quad D^*\Sigma(\bar{x}, \bar{z})(z^*) = \emptyset,$$

showing that the estimate (4.2) fails to be true for general arguments.

Our next example indicates that the results from [34, 38, 53] can be directly generalized to weak frontier mappings.

Example 4.2. We consider $(\mathbf{P}(x))$ with $n := m := 1$, $q := 2$, $\mathcal{C} := \mathbb{R}_+^2$, and

$$\forall x, y \in \mathbb{R}: \quad f(x, y) := (xy, y), \quad \Gamma(x) := [0, 1].$$

Then we find $\Sigma(x) = \text{conv}\{(0, 0), (x, 1)\}$ for all $x \in \mathbb{R}$ as well as

$$\forall x \in \mathbb{R}: \quad \Phi_w(x) = \begin{cases} \Sigma(x) & x \leq 0, \\ \{(0, 0)\} & x > 0, \end{cases} \quad \Psi_w(x) = \begin{cases} [0, 1] & x \leq 0, \\ \{0\} & x > 0. \end{cases}$$

Using this, one can easily check that $\Sigma(x) \subset \Phi_w(x) + \mathbb{R}_+^2$ holds for all $x \in \mathbb{R}$, i.e., the associated problem $(\mathbf{P}(x))$ possesses the weak domination property for all $x \in \mathbb{R}$ (actually, it also possesses the domination property for all $x \in \mathbb{R}$). Let us consider $\bar{x} := \bar{y} := 0$ and $\bar{z} := (0, 0)$. It is not hard to see that

$$\begin{aligned} N_{\text{gph } \Sigma}(\bar{x}, \bar{z}) &= \{0\} \times \mathbb{R} \times \mathbb{R}_-, \\ N_{\text{gph } (\Sigma + \mathcal{C})}(\bar{x}, \bar{z}) &= \{0\} \times \mathbb{R}_- \times \mathbb{R}_-, \\ N_{\text{gph } \Phi_w}(\bar{x}, \bar{z}) &= (\{0\} \times \mathbb{R} \times \mathbb{R}) \cup (\mathbb{R}_+ \times \mathbb{R} \times \{0\}), \\ N_{\text{gph } \Psi_w}(\bar{x}, \bar{y}) &= (\{0\} \times \mathbb{R}) \cup (\mathbb{R}_+ \times \{0\}). \end{aligned}$$

This directly shows that the inclusion

$$D^*\Phi_w(\bar{x}, \bar{z})(z^*) \subset D^*(\Sigma + \mathcal{C})(\bar{x}, \bar{z})(z^*) \quad (4.3)$$

holds for all $z^* \in \text{int } \mathbb{R}_+^2$, but is violated for, e.g., $z^* := (1, 0)$ already, where

$$D^*\Phi_w(\bar{x}, \bar{z})(z^*) = \mathbb{R}_+, \quad D^*\Sigma(\bar{x}, \bar{z})(z^*) = \{0\}.$$

Let us now formally prove that the estimate (4.3) holds for all $z^* \in \mathcal{C}_>^*$.

Lemma 4.3. *Let $(\bar{x}, \bar{z}) \in \text{gph } \Phi_w$ be fixed and let $\text{gph } \Phi_w$ and $\text{gph}(\Sigma + \mathcal{C})$ be locally closed around (\bar{x}, \bar{z}) . Assume that $(\mathbf{P}(x))$ possesses the weak domination property for all $x \in U$ where $U \subset \mathbb{R}^n$ is a neighborhood of \bar{x} . Then, for each $z^* \in \mathcal{C}_>^*$, the estimate (4.3) is valid.*

Proof. Fix $z^* \in \mathcal{C}_>^*$. By mimicking the proof of [34, Lemma 3.2], which does not rely on the precise underlying notion of nondominance, we find

$$D^*\Phi_w(\bar{x}, \bar{z})(z^*) \subset D^*(\Phi_w + \mathcal{C})(\bar{x}, \bar{z})(z^*).$$

Furthermore, locally around \bar{x} , the mappings $\Phi_w + \mathcal{C}$ and $\Sigma + \mathcal{C}$ are identical by the weak domination property. Noting that the coderivative is a local construction, the claim follows. \square

We would like to point the reader's attention to the similarities in the estimates (4.1) and (4.3). Indeed, the right-hand side is the same in both estimates which is somewhat unsatisfying as the precise underlying notion of nondominance seems to make no difference.

With the aid of Lemma 4.3, we are in position to obtain an upper estimate for the coderivative of Φ_w in terms of initial problem data. The following result is inspired by [34, Theorem 3.3].

Theorem 4.4. *Let $(\bar{x}, \bar{z}) \in \text{gph } \Phi_w$ be fixed and let $\text{gph } \Phi_w$ be locally closed around (\bar{x}, \bar{z}) . Assume that $(\mathbf{P}(x))$ possesses the weak domination property for all $x \in U$ where $U \subset \mathbb{R}^n$ is a neighborhood of \bar{x} . Furthermore, let Γ be locally bounded at \bar{x} . Let f be locally Lipschitz continuous at all points (\bar{x}, \bar{y}) such that $\bar{y} \in \Psi_w(\bar{x})$ and $f(\bar{x}, \bar{y}) = \bar{z}$ hold. Then, for each $z^* \in \mathcal{C}_>^*$,*

$$D^*\Phi_w(\bar{x}, \bar{z})(z^*) \subset \bigcup_{\substack{\bar{y} \in \Psi_w(\bar{x}) \\ f(\bar{x}, \bar{y}) = \bar{z}}} \bigcup_{(x^*, y^*) \in D^*f(\bar{x}, \bar{y})(z^*)} (\{x^*\} + D^*\Gamma(\bar{x}, \bar{y})(y^*)).$$

If, additionally, f is continuously differentiable at all points (\bar{x}, \bar{y}) such that $\bar{y} \in \Psi_w(\bar{x})$ and $f(\bar{x}, \bar{y}) = \bar{z}$ hold, then, for each $z^ \in \mathcal{C}_>^*$,*

$$D^*\Phi_w(\bar{x}, \bar{z})(z^*) \subset \bigcup_{\substack{\bar{y} \in \Psi_w(\bar{x}) \\ f(\bar{x}, \bar{y}) = \bar{z}}} (\{f'_x(\bar{x}, \bar{y})^\top z^*\} + D^*\Gamma(\bar{x}, \bar{y})(f'_y(\bar{x}, \bar{y})^\top z^*)).$$

Proof. Let us note that closedness of $\text{gph } \Gamma$ and local boundedness of Γ at \bar{x} yield that, locally around (\bar{x}, \bar{z}) , $\text{gph } \Sigma$ and $\text{gph}(\Sigma + \mathcal{C})$ are closed. Combining Lemma 4.3 and the first part of the proof of [34, Theorem 3.3] while taking into account that $-N_{\mathcal{C}}(0) = \mathcal{C}^* \supset \mathcal{C}_>^*$ since \mathcal{C} is a cone gives

$$D^*\Phi_w(\bar{x}, \bar{z})(z^*) \subset D^*\Sigma(\bar{x}, \bar{z})(z^*)$$

for each $z^* \in \mathcal{C}_>^*$.

Defining $\tilde{\Gamma}: \mathbb{R}^n \rightrightarrows \mathbb{R}^n \times \mathbb{R}^m$ by $\tilde{\Gamma}(x) := \{x\} \times \Gamma(x)$ for each $x \in \mathbb{R}^n$ gives $\Sigma = f \circ \tilde{\Gamma}$, and

$$D^*\tilde{\Gamma}(\bar{x}, (\bar{x}, y))(x^*, y^*) = \{x^*\} + D^*\Gamma(\bar{x}, y)(y^*)$$

holds for each $y \in \Gamma(\bar{x})$ and $(x^*, y^*) \in \mathbb{R}^n \times \mathbb{R}^m$, see e.g. [5, Lemma 5.7]. Thus, in order to infer the more general assertion of the theorem, we apply the chain rule from [5, Theorem 5.2]. Therefore, we note that the involved intermediate mapping

$$(x, z) \mapsto \{(x, y) \mid y \in \Gamma(x), f(x, y) = z\} \quad (4.4)$$

is inner semicompact at (\bar{x}, \bar{z}) w.r.t. its domain since Γ is assumed to be locally bounded at \bar{x} , and that f is Lipschitz continuous at (\bar{x}, \bar{y}) which serves as a qualification condition. Then we find

$$D^*\Sigma(\bar{x}, \bar{z})(z^*) \subset \bigcup_{\substack{\bar{y} \in \Gamma(\bar{x}) \\ f(\bar{x}, \bar{y}) = \bar{z}}} \bigcup_{(x^*, y^*) \in D^*f(\bar{x}, \bar{y})(z^*)} D^*\tilde{\Gamma}(\bar{x}, (\bar{x}, \bar{y}))(x^*, y^*),$$

and noting that, due to **Proposition 3.1**, each $\bar{y} \in \Gamma(\bar{x})$ with $f(\bar{x}, \bar{y}) = \bar{z}$ already belongs to $\Psi_w(\bar{x})$ yields the claim.

The second statement of the theorem is an obvious consequence of the first assertion. \square

Let us note that the chain rule from [5, Theorem 5.2] also allows for the derivation of refined upper estimates in terms of *directional* coderivatives.

We would like to point the interested reader to the fact that in the proofs of [34, Theorem 3.3] and [37, Theorem 4.1], the authors rely on the chain rule from [42, Theorem 3.18(i)] which is only possible if the intermediate mapping from (4.4) possesses certain inner semicontinuity properties. These, however, are not inherent from the assumptions stated in [34, 37], and one can easily check that even inner semicontinuity of Γ is not sufficient for these requirements. Thus, the aforementioned results are not reliable. Let us also note that, in situations where f is Lipschitzian, the coderivative of Σ can be computed exploiting a combination of the image and pre-image rule as we have

$$\text{gph } \Sigma = \pi(\{(x, y, z) \mid (x, y, f(x, y) - z) \in \text{gph } \Gamma \times \{0\}\})$$

where $\pi: \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{R}^q \rightarrow \mathbb{R}^n \times \mathbb{R}^q$ is given by $\pi(x, y, z) := (x, z)$ for all $x \in \mathbb{R}^n$, $y \in \mathbb{R}^m$, and $z \in \mathbb{R}^q$, see e.g. [5, Sections 5.1.3, 5.1.4]. This, however, still makes some additional boundedness assumption on Γ necessary, and the resulting upper estimate on the coderivative of Σ would also comprise a union over $\Psi_w(\bar{x})$. More precisely, this approach precisely recovers the estimates from **Theorem 4.4**.

Let us specify **Theorem 4.4** for the scalar case where $q := 1$ and $\mathcal{C} := \mathbb{R}_+$. Then the weak domination property is inherently satisfied. Furthermore, we find $1 \in \mathcal{C}_>^* = (0, \infty)$, and $D^*\Phi_w(\bar{x}, \varphi(\bar{x}))(1)$ equals the so-called limiting subdifferential at \bar{x} of the optimal valued function φ from (1.1) provided the latter is continuous around \bar{x} , see [42, Definition 1.77, Theorem 1.80]. Then the estimates in **Theorem 4.4** recover those obtained in [44, Theorem 7] under slightly different assumptions. Hence, one can interpret **Theorem 4.4** as a generalization of this result to the multiobjective situation. Nevertheless, **Theorem 4.4** is of limited practical use as it provides no information about situations where the argument z^* of the coderivative is chosen in $\mathbb{R}^q \setminus \mathcal{C}_>^*$, and such values of z^* might be of interest when deriving necessary optimality conditions for (\mathcal{E}_w -VFR) as the concept of subdifferentials is not reasonable in vector-valued scenarios whenever scalarization is not taken into account.

4.2 Coderivative estimates via scalarization

In this section, we aim to find upper coderivative estimates for the weak frontier mapping Φ_w via a scalarization approach. This, however, makes additional assumptions necessary.

Assumption 4.5. *For each $x \in \mathbb{R}^n$, $f(x, \cdot)$ is \mathcal{C} -convex and $\Gamma(x)$ is convex.*

We emphasize that [Assumption 4.5](#) guarantees that $\Sigma(x) + \mathcal{C}$ is convex for each $x \in \mathbb{R}^n$. Hence, due to [\[36, Corollary 5.29\]](#), for each $\bar{x} \in \text{dom } \Psi_w$, we find the equivalence

$$y \in \Psi_w(\bar{x}) \iff \exists \lambda \in \mathcal{C}^* \cap \mathbb{S}_1(0): y \in \Psi_w^{\text{sc}}(\bar{x}, \lambda)$$

where $\Psi_w^{\text{sc}}: \mathbb{R}^n \times \mathbb{R}^q \rightrightarrows \mathbb{R}^m$ is the solution mapping associated with the scalar parametric optimization problem

$$\min_y \{ \langle \lambda, f(x, y) \rangle \mid y \in \Gamma(x) \}. \quad (\text{P}(x, \lambda))$$

Based on the chain rule, we obtain a simple upper estimate for the coderivative of Φ_w in terms of the coderivative of Ψ_w , as we have $\Phi_w = f \circ \tilde{\Psi}_w$ where $\tilde{\Psi}_w: \mathbb{R}^n \rightrightarrows \mathbb{R}^n \times \mathbb{R}^m$ is given by $\tilde{\Psi}_w(x) := \{x\} \times \Psi_w(x)$ for each $x \in \mathbb{R}^n$. We omit the proof as it mainly parallels the arguments used to validate [Theorem 4.4](#).

Lemma 4.6. *Fix $(\bar{x}, \bar{z}) \in \text{gph } \Phi_w$ and let $\text{gph } \Phi_w$ be locally closed around (\bar{x}, \bar{z}) . Let Ψ_w be inner semicompact at \bar{x} w.r.t. $\text{dom } \Psi_w$. Furthermore, for each $\bar{y} \in \Psi_w(\bar{x})$ with $f(\bar{x}, \bar{y}) = \bar{z}$, let $\text{gph } \Psi_w$ be closed locally around (\bar{x}, \bar{y}) , and let f be locally Lipschitz continuous at (\bar{x}, \bar{y}) . Then, for each $z^* \in \mathbb{R}^q$,*

$$D^* \Phi_w(\bar{x}, \bar{z})(z^*) \subset \bigcup_{\substack{\bar{y} \in \Psi_w(\bar{x}) \\ f(\bar{x}, \bar{y}) = \bar{z}}} \bigcup_{(x^*, y^*) \in D^* f(\bar{x}, \bar{y})(z^*)} (x^* + D^* \Psi_w(\bar{x}, \bar{y})(y^*)).$$

If, additionally, f is continuously differentiable at all points (\bar{x}, \bar{y}) such that $\bar{y} \in \Psi_w(\bar{x})$ and $f(\bar{x}, \bar{y}) = \bar{z}$ hold, then, for each $z^ \in \mathbb{R}^q$,*

$$D^* \Phi_w(\bar{x}, \bar{z})(z^*) \subset \bigcup_{\substack{\bar{y} \in \Psi_w(\bar{x}) \\ f(\bar{x}, \bar{y}) = \bar{z}}} (f'_x(\bar{x}, \bar{y})^\top z^* + D^* \Psi_w(\bar{x}, \bar{y})(f'_y(\bar{x}, \bar{y})^\top z^*)).$$

Let us emphasize that [Lemma 4.6](#) is correct even in the absence of [Assumption 4.5](#) which is not needed for the proof. We note that the estimates from [Lemma 4.6](#) are closely related to those ones obtained in [Theorem 4.4](#). Indeed, the only difference (apart from the underlying assumptions) is that in [Theorem 4.4](#), all estimates are valid for $z^* \in \mathcal{C}_>^*$ only, while the coderivative of the efficiency mapping Ψ_w appearing in [Lemma 4.6](#) is replaced by the coderivative of the feasibility mapping Γ in [Theorem 4.4](#). Let us also visualize this in exemplary way.

One can easily check that the estimate provided in [Lemma 4.6](#) is actually sharp in the context of [Example 4.2](#). Indeed, one obtains

$$D^* \Phi_w(\bar{x}, \bar{z})(z^*) = D^* \Psi_w(\bar{x}, \bar{y})(z_2^*) = \begin{cases} \{0\} & z_2^* \neq 0, \\ \mathbb{R}_+ & z_2^* = 0 \end{cases}$$

for each $z^* \in \mathbb{R}^2$ in this situation (observe that the only point $y \in \Psi_w(\bar{x})$ which satisfies $f(\bar{x}, y) = \bar{z}$ in the present situation is \bar{y}). We also note that the estimates from [Theorem 4.4](#) do not apply for $z^* \in \mathbb{R} \times \{0\}$ as these vectors do not belong to $\text{int } \mathbb{R}_+^2$.

It is an essential disadvantage of the estimates from [Lemma 4.6](#) that they are not stated in terms of initial problem data as the coderivative of Ψ_w is a fully implicit object. Subsequently, we aim to estimate it from above. In order to deal with this issue, [Assumption 4.5](#) turns out to be helpful as we will demonstrate subsequently. For brevity of notation, we introduce an intermediate mapping $\Xi: \mathbb{R}^n \times \mathbb{R}^m \rightrightarrows \mathbb{R}^q$ given by

$$\forall x \in \mathbb{R}^n \forall y \in \mathbb{R}^m: \quad \Xi(x, y) := \{\lambda \in \mathcal{C}^* \cap \mathbb{S}_1(0) \mid y \in \Psi_w^{\text{sc}}(x, \lambda)\}.$$

For given $(x, y) \in \text{gph } \Psi_w$, $\Xi(x, y)$ comprises all (normalized) scalarization parameters for which y as a minimizer of $(\mathbf{P}(x, \lambda))$. We note that, by compactness of $\mathcal{C}^* \cap \mathbb{S}_1(0)$, Ξ is inner semicompact w.r.t. its domain at each point of its domain.

Lemma 4.7. *Fix $(\bar{x}, \bar{y}) \in \text{gph } \Psi_w$ and let $\text{gph } \Psi_w$ be closed locally around (\bar{x}, \bar{y}) . Furthermore, for each $\lambda \in \Xi(\bar{x}, \bar{y})$, let $\text{gph } \Psi_w^{\text{sc}}$ be closed locally around $((\bar{x}, \lambda), \bar{y})$. Additionally, let one of the following conditions hold:*

- (a) Ψ_w^{sc} is a polyhedral set-valued mapping, \mathcal{C} is a polyhedral cone, and the unit sphere $\mathbb{S}_1(0)$ is taken w.r.t. the 1- or ∞ -norm,
- (b) for each $\lambda \in \Xi(\bar{x}, \bar{y})$, the qualification condition

$$(0, \eta) \in D^* \Psi_w^{\text{sc}}((\bar{x}, \lambda), \bar{y})(0), \eta \in -N_{\mathcal{C}^* \cap \mathbb{S}_1(0)}(\lambda) \implies \eta = 0$$

holds.

Finally, let [Assumption 4.5](#) hold. Then, for each $y^* \in \mathbb{R}^m$, we have

$$D^* \Psi_w(\bar{x}, \bar{y})(y^*) \subset \bigcup_{\lambda \in \Xi(\bar{x}, \bar{y})} \{x^* \mid (x^*, \eta) \in D^* \Psi_w^{\text{sc}}((\bar{x}, \lambda), \bar{y})(y^*), \eta \in -N_{\mathcal{C}^* \cap \mathbb{S}_1(0)}(\lambda)\}.$$

Proof. Defining $\Upsilon: \mathbb{R}^n \rightrightarrows \mathbb{R}^n \times \mathbb{R}^q$ by $\Upsilon(x) := \{x\} \times (\mathcal{C}^* \cap \mathbb{S}_1(0))$ for each $x \in \mathbb{R}^n$, we find $\Psi_w = \Psi_w^{\text{sc}} \circ \Upsilon$ since [Assumption 4.5](#) holds. Since Ξ is inner semicompact at (\bar{x}, \bar{y}) w.r.t. its domain, the intermediate mapping

$$(x, y) \mapsto \{(x, \lambda) \in \Upsilon(x) \mid y \in \Psi_w^{\text{sc}}(x, \lambda)\} = \{(x, \lambda) \mid \lambda \in \Xi(x, y)\}$$

is inner semicompact at (\bar{x}, \bar{y}) w.r.t. its domain as well. Thus, the result follows from the chain rule stated in [[5](#), Theorem 5.2] while observing that

$$D^* \Upsilon(\bar{x}, (\bar{x}, \lambda))(x^*, \eta) = \begin{cases} \{x^*\} & \eta \in -N_{\mathcal{C}^* \cap \mathbb{S}_1(0)}(\lambda), \\ \emptyset & \text{otherwise} \end{cases}$$

holds for each $\lambda \in \Xi(\bar{x}, \bar{y})$ and $(x^*, \eta) \in \mathbb{R}^n \times \mathbb{R}^q$ while each of the additionally postulated assumptions guarantees validity of the qualification condition which is imposed in [[5](#), Theorem 5.2]. \square

Let us note that [Lemma 4.7](#) generalizes [[57](#), Proposition 3.2] to arbitrary ordering cones while coming along with a simpler proof and another type of qualification condition. Furthermore, we would like to mention that [Lemma 4.7](#) can be used to derive necessary optimality conditions for $(\mathcal{E}_w\text{-BPP})$ directly.

In order to obtain a valuable coderivative estimate for the weak frontier mapping from [Lemmas 4.6](#) and [4.7](#), it remains to estimate the appearing coderivative of Ψ_w^{sc} . This, however, is a well-established subject in variational analysis since Ψ_w^{sc} is the solution mapping of a

scalar parametric optimization problem. Exemplary, let us mention that, in the presence of [Assumption 4.5](#), $(\mathbf{P}(x, \lambda))$ is a convex optimization problem for each $x \in \mathbb{R}^n$ and $\lambda \in \mathcal{C}^*$. Assuming, additionally, that f is a smooth function, we find

$$\forall x \in \mathbb{R}^n \forall \lambda \in \mathcal{C}^*: \quad \Psi_{\mathbf{w}}^{\text{sc}}(x, \lambda) = \left\{ y \mid -f'_y(x, y)^\top \lambda \in N_{\Gamma(x)}(y) \right\},$$

i.e., $\Psi_{\mathbf{w}}^{\text{sc}}$ possesses the structure of a so-called *parametric variational system*. Consulting e.g. [42, Section 4.4], coderivative estimates for $\Psi_{\mathbf{w}}^{\text{sc}}$ can be obtained in terms of derivatives of f and coderivatives of the normal cone mapping $(x, y) \mapsto N_{\Gamma(x)}(y)$ under suitable constraint qualifications. The remaining coderivative of the normal cone map may be specified for diverse different types of constraint systems, see e.g. [3, 28–30]. Exemplary, this has been worked out in the proof of [57, Corollary 3.4] in the particular situation where $\mathcal{C} := \mathbb{R}_+^q$ and Γ is induced by smooth parametric inequality constraints.

As the purpose of this note is mainly focused on the value behind the model reformulations $(\mathcal{E}\text{-VFR})$ and $(\mathcal{E}_{\mathbf{w}}\text{-VFR})$, we abstain from the postulation of any more technical results, but simply combine [Lemmas 4.6](#) and [4.7](#) to obtain a coderivative estimate which can be made fully explicit by respecting the above comments by the interested reader.

Theorem 4.8. *Fix $(\bar{x}, \bar{z}) \in \text{gph } \Phi_{\mathbf{w}}$ and let $\text{gph } \Phi_{\mathbf{w}}$ be locally closed around (\bar{x}, \bar{z}) . Furthermore, let [Assumption 4.5](#) be valid. Let $\Psi_{\mathbf{w}}$ be inner semicompact at \bar{x} w.r.t. $\text{dom } \Psi_{\mathbf{w}}$. Furthermore, for each $\bar{y} \in \Psi_{\mathbf{w}}(\bar{x})$ with $f(\bar{x}, \bar{y}) = \bar{z}$, let $\text{gph } \Psi_{\mathbf{w}}$ be closed locally around (\bar{x}, \bar{y}) , let f be locally Lipschitz continuous at (\bar{x}, \bar{y}) , and, for each $\lambda \in \Xi(\bar{x}, \bar{y})$, let $\text{gph } \Psi_{\mathbf{w}}^{\text{sc}}$ be closed locally around $((\bar{x}, \lambda), \bar{y})$, and let one of the conditions (a) or (b) of [Lemma 4.7](#) be valid. Then, for each $z^* \in \mathbb{R}^q$,*

$$D^* \Phi_{\mathbf{w}}(\bar{x}, \bar{z})(z^*) \subset \bigcup_{\substack{\bar{y} \in \Psi_{\mathbf{w}}(\bar{x}) \\ f(\bar{x}, \bar{y}) = \bar{z}}} \left\{ x_1^* + x_2^* \left| \begin{array}{l} (x_1^*, y^*) \in D^* f(\bar{x}, \bar{y})(z^*), \\ (x_2^*, \eta) \in D^* \Psi_{\mathbf{w}}^{\text{sc}}((\bar{x}, \lambda), \bar{y})(y^*), \\ \lambda \in \Xi(\bar{x}, \bar{y}), \eta \in -N_{\mathcal{C}^* \cap \mathbb{S}_1(0)}(\lambda) \end{array} \right. \right\}.$$

If, additionally, f is continuously differentiable at all points (\bar{x}, \bar{y}) such that $\bar{y} \in \Psi_{\mathbf{w}}(\bar{x})$ and $f(\bar{x}, \bar{y}) = \bar{z}$ hold, then, for each $z^* \in \mathbb{R}^q$,

$$D^* \Phi_{\mathbf{w}}(\bar{x}, \bar{z})(z^*) \subset \bigcup_{\substack{\bar{y} \in \Psi_{\mathbf{w}}(\bar{x}) \\ f(\bar{x}, \bar{y}) = \bar{z}}} \left\{ f'_x(\bar{x}, \bar{y})^\top z^* + x^* \left| \begin{array}{l} (x^*, \eta) \in D^* \Psi_{\mathbf{w}}^{\text{sc}}((\bar{x}, \lambda), \bar{y})(f'_y(\bar{x}, \bar{y})^\top z^*), \\ \lambda \in \Xi(\bar{x}, \bar{y}), \eta \in -N_{\mathcal{C}^* \cap \mathbb{S}_1(0)}(\lambda) \end{array} \right. \right\}.$$

Let us note that, based on [Lemma 3.5](#), one can easily check that the numerous closedness assumptions as well as the inner semicompactness assumption on $\Psi_{\mathbf{w}}$ in [Theorem 4.8](#) are inherently satisfied whenever Γ is lower semicontinuous w.r.t. its domain locally around \bar{x} , and locally bounded at \bar{x} . Both assumptions are inherent whenever $\Gamma(x) := \Omega$ holds for some nonempty and compact set $\Omega \subset \mathbb{R}^m$ and all $x \in \mathbb{R}^n$. Then it remains to validate condition (a) or (b) of [Lemma 4.7](#).

Remark 4.9. *Let [Assumption 4.5](#) be valid and $\varphi^{\text{sc}}: \mathbb{R}^n \times \mathbb{R}^q \rightarrow \overline{\mathbb{R}}$ be the optimal value function associated with $(\mathbf{P}(x, \lambda))$, i.e.,*

$$\forall x \in \mathbb{R}^n \forall \lambda \in \mathbb{R}^q: \quad \varphi^{\text{sc}}(x, \lambda) := \inf_y \{ \langle \lambda, f(x, y) \rangle \mid y \in \Gamma(x) \}.$$

Then we find the representation

$$\forall x \in \mathbb{R}^n: \quad \Phi_{\mathbf{w}}(x) = \{ z \in \Sigma(x) \mid \exists \lambda \in \mathcal{C}^* \cap \mathbb{S}_1(0): \langle \lambda, z \rangle = \varphi^{\text{sc}}(x, \lambda) \},$$

where $\Sigma: \mathbb{R}^n \rightrightarrows \mathbb{R}^q$ has been defined in (3.1). Using $\Upsilon: \mathbb{R}^n \rightrightarrows \mathbb{R}^n \times \mathbb{R}^q$ from the proof of Lemma 4.7 as well as $\Upsilon_1: \mathbb{R}^n \times \mathbb{R}^q \rightarrow \mathbb{R}^q \times \mathbb{R}$ and $\Upsilon_2: \mathbb{R}^q \times \mathbb{R} \rightarrow \mathbb{R}^q$ given by

$$\forall x \in \mathbb{R}^n \forall \lambda \in \mathbb{R}^q \forall \alpha \in \mathbb{R}: \quad \Upsilon_1(x, \lambda) := (\lambda, \varphi^{\text{sc}}(x, \lambda)), \quad \Upsilon_2(\lambda, \alpha) := \{z \mid \langle \lambda, z \rangle - \alpha = 0\},$$

we find $\Phi_w(x) = \Sigma(x) \cap (\Upsilon_2 \circ (\Upsilon_1 \circ \Upsilon))(x)$. Seemingly, this representation offers the possibility to obtain an upper estimate for the coderivative of Φ_w via an intersection rule, see e.g. [42, Proposition 3.20], and the chain rule, see e.g. [5, Theorem 5.2]. Indeed, in case where φ^{sc} is a locally Lipschitzian function, which holds under fairly mild assumptions, one can apply the chain rule from [5] twice in order to find an upper estimate of the coderivative of $\Upsilon_2 \circ (\Upsilon_1 \circ \Upsilon)$ which comprises the limiting subdifferential of (a non-specified multiple of) φ^{sc} . The latter, however, cannot be approximated from above by the Cartesian product of the associated subdifferentials w.r.t. the variable x and λ in general which is a first difficulty. A second problem pops up as the constraint qualification from [42, Proposition 3.20] which is sufficient to get the intersection rule working is also likely to fail. Hence, we abstain here from providing the details as this approach does not seem too reasonable.

5 Conclusions

In this paper, we visualized that multiobjective bilevel optimization is, up to a certain degree, only reasonable if the lower level efficiency-type and frontier-type mapping possess closed graphs as this is a fundamental assumption in order to guarantee the existence of solutions (in whatever sense) and to allow for the derivation of optimality conditions. We presented conditions which guarantee validity of these closedness properties, gave existence results, and discussed the computation of upper coderivative estimates for frontier-type mappings. The latter is an essential step on the route to necessary optimality conditions as visualized in the recent paper [37] which, however, has some flaws which we carved out here. Furthermore, we commented on different ways on how to introduce a value function reformulation of a multiobjective bilevel optimization problem which might be important in algorithmic applications.

In the future, it has to be investigated whether the value function reformulation of a multiobjective optimization problem can be exploited beneficially for its algorithmic solution. Whenever the lower level problem is a scalar optimization problem, this has already been demonstrated in several different ways in the literature, see e.g. [14, Section 20.6.4] for an overview.

References

- [1] B. Bank, J. Guffat, D. Klatte, B. Kummer, and K. Tammer. *Non-Linear Parametric Optimization*. Birkhäuser, Basel, 1983.
- [2] J. F. Bard. *Practical Bilevel Optimization: Algorithms and Applications*. Kluwer Academic, Dordrecht, 1998.
- [3] M. Benko, H. Gfrerer, and J. V. Outrata. Stability analysis for parametric variational systems with implicit constraints. *Set-Valued and Variational Analysis*, 28:167–193, 2020. doi:10.1007/s11228-019-00516-1.

- [4] M. Benko and P. Mehlitz. On implicit variables in optimization theory. *Journal of Nonsmooth Analysis and Optimization*, 2:7215, 2021. doi:10.46298/jnsao-2021-7215.
- [5] M. Benko and P. Mehlitz. Calmness and calculus: two basic patterns. *Set-Valued and Variational Analysis*, 30:81–117, 2022. doi:10.1007/s11228-021-00589-x.
- [6] S. Bolintineanu. Necessary conditions for nonlinear suboptimization over the weakly efficient set. *Journal of Optimization Theory and Applications*, 78(3):579–598, 1993. doi:10.1007/BF00939883.
- [7] S. Bolintineanu. Optimality conditions for minimization over the (weakly or properly) efficient set. *Journal of Mathematical Analysis and Applications*, 173(2):523 – 541, 1993. doi:10.1006/jmaa.1993.1085.
- [8] H. Bonnel. Optimality conditions for the semivectorial bilevel optimization problem. *Pacific Journal of Optimization*, 2(3):447–468, 2006.
- [9] H. Bonnel and C. Y. Kaya. Optimization over the efficient set of multi-objective convex optimal control problems. *Journal of Optimization Theory and Applications*, 147(1):93–112, 2010. doi:10.1007/s10957-010-9709-y.
- [10] H. Bonnel and J. Morgan. Semivectorial bilevel optimization problem: penalty approach. *Journal of Optimization Theory and Applications*, 131(3):365–382, 2006. doi:10.1007/s10957-006-9150-4.
- [11] B. Dandurand, P. Guarneri, G. M. Fadel, and M. M. Wiecek. Bilevel multiobjective packing optimization for automotive design. *Structural and Multidisciplinary Optimization*, 50:663–682, 2014. doi:10.1007/s00158-014-1120-0.
- [12] S. Dempe. *Foundations of Bilevel Programming*. Kluwer Academic, Dordrecht, 2002.
- [13] S. Dempe. Annotated bibliography on bilevel programming and mathematical programs with equilibrium constraints. *Optimization*, 52(3):333–359, 2003. doi:10.1080/0233193031000149894.
- [14] S. Dempe. Bilevel optimization: theory, algorithms, applications and a bibliography. In *Bilevel Optimization: Advances and Next Challenges*, pages 581–672. Springer, Cham, 2020. doi:10.1007/978-3-030-52119-6_20.
- [15] S. Dempe, J. Dutta, and B. S. Mordukhovich. New necessary optimality conditions in optimistic bilevel programming. *Optimization*, 56(5-6):577–604, 2007. doi:10.1080/02331930701617551.
- [16] S. Dempe and S. Franke. On the solution of convex bilevel optimization problems. *Computational Optimization and Applications*, 63(3):685–703, 2016. doi:10.1007/s10589-015-9795-8.
- [17] S. Dempe, N. Gadhi, and A. B. Zemkoho. New optimality conditions for the semivectorial bilevel optimization problem. *Journal of Optimization Theory and Applications*, 157(1):54–74, 2013. doi:10.1007/s10957-012-0161-z.
- [18] S. Dempe, V. Kalashnikov, G. Pérez-Valdéz, and N. Kalashnykova. *Bilevel Programming Problems - Theory, Algorithms and Applications to Energy Networks*. Springer, Berlin, 2015.

- [19] S. Dempe and P. Mehrlitz. Semivectorial bilevel programming versus scalar bilevel programming. *Optimization*, 69(4):657–679, 2020. doi:10.1080/02331934.2019.1625900.
- [20] S. Dempe, B. S. Mordukhovich, and A. B. Zemkoho. Sensitivity analysis for two-level value functions with applications to bilevel programming. *SIAM Journal on Optimization*, 22(4):1309–1343, 2012. doi:10.1137/110845197.
- [21] S. Dempe and A. B. Zemkoho. The bilevel programming problem: reformulations, constraint qualifications and optimality conditions. *Mathematical Programming*, 138:447–473, 2013. doi:10.1007/s10107-011-0508-5.
- [22] A. L. Dontchev and R. T. Rockafellar. *Implicit Functions and Solution Mappings*. Springer, Heidelberg, 2014.
- [23] M. Ehrgott. *Multicriteria Optimization*. Springer, Berlin, 2005.
- [24] G. Eichfelder. Multiobjective bilevel optimization. *Mathematical Programming*, 123:419–449, 2010. doi:10.1007/s10107-008-0259-0.
- [25] G. Eichfelder. Methods for multiobjective bilevel optimization. In S. Dempe and A. B. Zemkoho, editors, *Bilevel Optimization: Advances and Next Challenges*, pages 423–449. Springer, Cham, 2020. doi:10.1007/978-3-030-52119-6_15.
- [26] N. Gadhi and S. Dempe. Necessary optimality conditions and a new approach to multiobjective bilevel optimization problems. *Journal of Optimization Theory and Applications*, 155:100–114, 2012. doi:10.1007/s10957-012-0046-1.
- [27] E. Gebhardt and J. Jahn. Global solver for nonlinear bilevel optimization problems. *Pacific Journal of Optimization*, 5(2):387–401, 2009. URL <http://www.ybook.co.jp/pjo.html>.
- [28] H. Gfrerer and J. V. Outrata. On computation of generalized derivatives of the normal-cone mapping and their applications. *Mathematics of Operations Research*, 41(4):1535–1556, 2016. doi:10.1287/moor.2016.0789.
- [29] H. Gfrerer and J. V. Outrata. On computation of limiting coderivatives of the normal-cone mapping to inequality systems and their applications. *Optimization*, 65(4):671–700, 2016. doi:10.1080/02331934.2015.1066372.
- [30] H. Gfrerer and J. V. Outrata. On the Aubin property of a class of parameterized variational systems. *Mathematical Methods of Operations Research*, 86(3):443–467, 2017. doi:10.1007/s00186-017-0596-y.
- [31] A. Göpfert, H. Riahi, C. Tammer, and C. Zălinescu. *Variational Methods in Partically Ordered Spaces*. Springer, New York, 2003.
- [32] M. I. Henig. The domination property in multicriteria optimization. *Journal of Mathematical Analysis and Applications*, 114(1):7–16, 1986. doi:10.1016/0022-247X(86)90061-2.
- [33] R. Horst and N. V. Thoai. Maximizing a concave function over the efficient or weakly-efficient set. *European Journal of Operational Research*, 117(2):239 – 252, 1999. doi:10.1016/S0377-2217(98)00230-6.

- [34] N. Q. Huy, B. S. Mordukhovich, and J.-C. Yao. Coderivatives of frontier and solution maps in parametric multiobjective optimization. *Taiwanese Journal of Mathematics*, 12(8):2083–2111, 2008. doi:10.11650/twjmath/1500405137.
- [35] H. Isermann. Proper efficiency and the linear vector maximization problem. *Operations Research*, 22(1):189–191, 1974. URL <https://www.jstor.org/stable/169229>.
- [36] J. Jahn. *Vector Optimization - Theory, Applications, and Extensions*. Springer, Berlin, 2011.
- [37] L. Lafhim and A. B. Zemkoho. Extension of the value function reformulation to multiobjective bilevel optimization. *Optimization Letters*, 2022. doi:10.1007/s11590-022-01948-9.
- [38] S. J. Li and X. W. Xue. Sensitivity analysis of gap functions for vector variational inequality via coderivatives. *Optimization*, 63(7):1075–1098, 2014. doi:10.1080/02331934.2012.688970.
- [39] B. Liu, Z. Wan, J. Chen, and G. Wang. Optimality conditions for pessimistic semivectorial bilevel programming problems. *Journal of Inequalities and Applications*, 41(1), 2014. doi:10.1186/1029-242X-2014-41.
- [40] C. G. Liu, K. F. Ng, and W. H. Yang. Merit functions in vector optimization. *Mathematical Programming*, 119:215–237, 2009. doi:10.1007/s10107-008-0208-y.
- [41] D. T. Luc. On the domination property in vector optimization. *Journal of Optimization Theory and Applications*, 43:327–330, 1984. doi:10.1007/BF00936168.
- [42] B. S. Mordukhovich. *Variational Analysis and Generalized Differentiation*. Springer, Berlin, 2006.
- [43] B. S. Mordukhovich, N. M. Nam, and H. M. Phan. Variational analysis of marginal functions with applications to bilevel programming. *Journal of Optimization Theory and Applications*, 152(3):557–586, 2012. doi:10.1007/s10957-011-9940-1.
- [44] B. S. Mordukhovich, N. M. Nam, and N. Y. Yen. Subgradients of marginal functions in parametric mathematical programming. *Mathematical Programming*, 116:369–396, 2009. doi:10.1007/s10107-007-0120-x.
- [45] F. Nožička, J. Guddat, H. Hollatz, and B. Bank. *Theorie der linearen parametrischen Optimierung*. Akademie-Verlag, Berlin, 1974.
- [46] J. V. Outrata. On the numerical solution of a class of Stackelberg problems. *Zeitschrift für Operations Research*, 34(4):255–277, 1990. doi:10.1007/BF01416737.
- [47] M. Pilecka. *Set-valued optimization and its application to bilevel optimization*. Phd thesis, Technische Universität Bergakademie Freiberg, 2016.
- [48] F. Plastria. On the structure of the weakly efficient set for quasiconvex vector minimization. *Journal of Optimization Theory and Applications*, 184:547–564, 2020. doi:10.1007/s10957-019-01608-6.
- [49] R. T. Rockafellar and R. J.-B. Wets. *Variational Analysis*. Springer, Berlin, 2009.

- [50] S. Ruuska, K. Miettinen, and M. M. Wiecek. Connections between single-level and bilevel multiobjective optimization. *Journal of Optimization Theory and Applications*, 153:60–74, 2012. doi:10.1007/s10957-011-9943-y.
- [51] Y. Sawaragi, H. Nakayama, and T. Tanino. *Theory of Multiobjective Optimization*. Academic Press, Orlando, 1985.
- [52] K. Shimizu, Y. Ishizuka, and J. F. Bard. *Nondifferentiable and Two-Level Mathematical Programming*. Kluwer Academic, Dordrecht, 1997.
- [53] X. W. Xue, S. J. Li, C. M. Liao, and J.-C. Yao. Sensitivity analysis of parametric vector set-valued optimization problems via coderivatives. *Taiwanese Journal of Mathematics*, 15(4):2533–3554, 2011. URL <https://www.jstor.org/stable/taiwmath.15.6.2533>.
- [54] J. J. Ye. Necessary optimality conditions for multiobjective bilevel programs. *Mathematics of Operations Research*, 36(1):165–184, 2011. doi:10.1287/moor.1100.0480.
- [55] J. J. Ye and D. L. Zhu. Optimality conditions for bilevel programming problems. *Optimization*, 33(1):9–27, 1995. doi:10.1080/02331939508844060.
- [56] Y. Yin. Multiobjective bilevel optimization for transportation planning and management problems. *Journal of Advanced Transportation*, 36(1):93–105, 2002. doi:10.1002/atr.5670360106.
- [57] A. B. Zemkoho. Solving ill-posed bilevel programs. *Set-Valued and Variational Analysis*, 24:423–448, 2016. doi:10.1007/s11228-016-0371-x.