

MODELS OF ABELIAN VARIETIES OVER VALUED FIELDS, USING MODEL THEORY

YATIR HALEVI

ABSTRACT. Given an elliptic curve E over a perfect defectless henselian valued field (F, val) with perfect residue field, there exists an integral separated smooth group scheme \mathcal{E} over \mathcal{O}_F with $\mathcal{E} \times_{\text{Spec } \mathcal{O}_F} \text{Spec } F \cong E$. The definable group $\mathcal{E}(\mathcal{O})$ is the maximal generically stable subgroup of E . We also give some partial results on general Abelian varieties over F .

The construction of \mathcal{E} is by means of generating a birational group law over \mathcal{O}_F by the aid of a generically stable generic type of a definable subgroup of E .

1. INTRODUCTION

Let E be an elliptic curve over the field of fractions $F = \text{Frac}(R)$ of a valuation ring R . A *model* of E is a scheme \mathcal{E} (of finite type) over R for which $\mathcal{E} \times_{\text{Spec } R} \text{Spec } F \cong E$. There are many different models one can consider. For example, if E is given by a Weierstrass equation over R then we can take the projective closure in \mathbb{P}_R^2 ; we get a proper model over R but it might not be smooth. If we now take the smooth locus over R the resulting model is smooth but we might lose properness. We might also desire the model to be a group scheme over R .

When R is a discrete valuation ring, for any abelian variety A over F , Néron constructed a model \mathcal{A} over R which can be seen as a best possible model. Néron relaxed the assumption of properness and concentrated on smoothness and the group structure. The model Néron constructed (*the Néron model of A over R*) is a smooth separated group scheme of finite type over R satisfying that $\mathcal{A} \times_{\text{Spec } R} \text{Spec } F \cong A$ and the *Néron mapping property*. This latter property is extremely important, but will not be important here; we just remark that it implies that $\mathcal{A}(R) = \mathcal{A}_F(F)$.

It is well known that if R is a valuation subring of an algebraically closed field then a Néron model might not necessarily exist, even for an elliptic curve over R (see, e.g., Remark 5.12). Nevertheless, one might hope to find well-behaved models. See [Nér64, BLR90] for more information on models over DVRs.

The main aim of this paper is to give a model theoretic approach for constructing such models by means of finding birational group laws (in the sense of [BLR90]) over R . We build heavily on results from [HRK19] and [Hal19], where generically stable groups in algebraically closed valued fields and stably pointed varieties were studied, respectively. Before stating the main theorems we elaborate on the former.

Let (K, val) be an algebraically closed valued field. A type-definable subgroup $H \subseteq G$ of an algebraic group G over K is said to be a connected generically stable group if there exists a generically stable type p , concentrated on H , satisfying that $gp = p$ for any $g \in H$ (so p is the unique generic type of H). We recall that a type p concentrated on G is generically stable if for any open affine subvariety $V \subseteq G$, on which p is concentrated, $\text{val}(f(c)) \in \Gamma_K$ for any $f \in K[V]$ and $c \models p|_K$.

The author was partially supported by ISF grants No. 555/21 and 290/19 and by the Fields Institute for Research in Mathematical Sciences.

We now give the main theorem for elliptic curves. Let F be a perfect defectless henselian (non-trivially) valued field with a perfect residue field. Let \mathcal{O}_F be its valuation ring and let $\mathcal{O}_{F^{alg}}$ be its extension to the algebraic closure F^{alg} of F . Below the types and definable groups are objects in the algebraically closed valued field F^{alg} .

Theorem (Theorem 5.8). *Let E be an elliptic curve given by a Weierstrass equation over \mathcal{O}_F . There exists a smooth integral separated \mathcal{O}_F -group scheme \mathcal{E} of finite presentation over \mathcal{O}_F with geometrically integral fibers satisfying that $\mathcal{E}_F \cong E$ and*

- (1) $\mathcal{E}(\mathcal{O}_{F^{alg}})$ is a connected generically stable group,
- (2) $\mathcal{E}(\mathcal{O}_{F^{alg}}) = E(F^{alg})$ if and only if \mathcal{E} is an Abelian scheme over \mathcal{O}_F if and only if $\Delta \in \mathcal{O}_F^\times$, where Δ is the discriminant of E ,
- (3) For any integral group scheme \mathcal{G} of finite type over $\mathcal{O}_{F^{alg}}$ with integral special fiber and $\mathcal{G}_{F^{alg}} \cong E_{F^{alg}}$ we have $\mathcal{G}_R(R) \subseteq \mathcal{E}_R(R)$ for any valued field extension $(F^{alg}, \mathcal{O}_{F^{alg}}) \leq (\text{Frac}(R), R)$.

For general Abelian varieties over an algebraically closed valued field K , with valuation ring \mathcal{O}_K , we have the following partial result.

Theorem (Theorem 4.8). *Let A be an Abelian variety over K and assume that A is a connected generically stable group with a unique generic type p . Then the following are equivalent:*

- (1) *There exists an open affine subvariety $V \subseteq A$ such that*

$$\{f \in K[V] : p \vdash \text{val}(f(x)) \geq 0\}$$

is a finitely generated algebra over \mathcal{O}_K .

- (2) *There exists an integral Abelian scheme \mathcal{A} over \mathcal{O}_K with $\mathcal{A}_K \cong A$.*

Our hope is that this theorem may be generalized to the generic type of the maximal generically stable subgroup, in the sense of [HRK19], and that condition (1) always holds.

The key model theoretic input is the following, stated for algebraically closed valued fields just for simplicity; it holds for valued fields as above but requires some more assumptions. It should be seen as an analog of [HRK19, Theorem 6.11] for non-affine algebraic groups. We do not know if the finite generation assumption can be dropped.

Proposition (Proposition 4.7). *Let G be an algebraic group over K and let H be a connected generically stable Zariski dense subgroup with p its generically stable generic. Further assume that there exists an open affine subvariety $V \subseteq G$ with*

$$\{f \in K[V] : p \vdash \text{val}(f(x)) \geq 0\}$$

a finitely generated algebra over \mathcal{O}_K .

Then there exists a smooth integral separated \mathcal{O}_K -group scheme \mathcal{G} of finite presentation over \mathcal{O}_K with geometrically integral fibers satisfying that $\mathcal{G}_K \cong G$ and that under this isomorphism $H \cong \mathcal{G}(\mathcal{O})$.

Acknowledgements I would like to thank Silvain Rideau-Kikuchi for numerous conversations and Martin Hils for going over an early draft. Appendix A is a result of several discussions with Itay Kaplan. Finally, I would like to thank Udi Hrushovski, the seeds (and seedlings) of this paper grew out of our fruitful discussions during my PhD.

2. PRELIMINARIES AND NOTATION

We will usually not distinguish between singletons and sequences, thus we may write $a \in M$ and actually mean $a = (a_1, \dots, a_n) \in M^n$, unless a distinction is necessary. We use juxtaposition ab for concatenation of sequences, or AB for $A \cup B$ if dealing with sets. That being said, since

we will be dealing with groups, we will try to differentiate between concatenation ab and group multiplication $a \cdot b$ by denoting the latter by $a \cdot b$.

We review the necessary model theory of algebraically closed valued fields. See [Hal21, HL16, HHM08] for more information. We will assume basic knowledge of model theory (and specifically the model theory of NIP structures), see [Sim15].

We will also require many results from [Hal19]. We will not be able to fully write all of them here as facts, but will cite them carefully when needed. The published version [Hal19] has several minor mistakes and one major error, see [Hal22] for a full corrigendum. The arxiv version [Hal21] has all the corrections implemented so it will be our main cited version.

Valued fields. Let (F, val) be a non-trivially valued field with value group Γ_F . The valuation ring of F is the ring $\mathcal{O}_F = \{x \in F : \text{val}(x) \geq 0\}$. It is a local ring with maximal ideal $\mathfrak{m}_F = \{x \in F : \text{val}(x) > 0\}$. The residue field is the quotient $\mathbf{k}_F = \mathcal{O}_F/\mathfrak{m}_F$ and the quotient map $\text{res} : \mathcal{O}_F \rightarrow \mathbf{k}_F$ is called the residue map.

Let ACVF be the theory of non trivially valued algebraically closed fields in the three sorted language: the valued field sort which will be denoted by VF, the value group by Γ and the residue field by \mathbf{k} .

We will treat the valued field sort and the model as interchangeable, e.g. when we say that K is a model of ACVF we really mean that $K = \text{VF}(M)$ for some $M \models \text{ACVF}$. For any definable set D and set A , we will write $D(A)$ for $\text{dcl}(A) \cap D$. So $\Gamma(A) := \text{dcl}(A) \cap \Gamma$, and $\mathbf{k}(A) := \text{dcl}(A) \cap \mathbf{k}$.

We work in some large (larger than any set or field in question) saturated model \mathbb{K} of ACVF (technically $\mathbb{K} = \text{VF}(\mathbb{U})$ for some saturated model \mathbb{U}).¹ All valued fields in question will be seen as subfields of \mathbb{K} . We set $\mathcal{O} = \mathcal{O}_{\mathbb{K}}$, $\Gamma = \Gamma_{\mathbb{K}}$ and $\mathbf{k} = \mathbf{k}_{\mathbb{K}}$.

The theory ACVF has NIP; we will mostly deal with *generically stable* types in this theory. The precise definition of such types will not be important here (see Appendix A). What will be important is that in ACVF an invariant type is generically stable if and only if it is stably dominated if and only if it is orthogonal to Γ .

This latter definition will be our most used property of such types. We say that a global type p is *orthogonal to Γ* (over A) if it is A -definable and for every $B \supseteq A$ and B -definable function f into Γ , f_*p is a constant type, where $f_*p = \text{tp}(f(a)/\mathbb{U})$ for $a \models p$. For a type p concentrated on an affine variety V over a model $K \models \text{ACVF}$ this exactly means that $\text{val}(f(c)) \in \Gamma_K$ for any $f \in K[V]$ and $c \models p|K$.

Example 2.1. Let $p_{\mathcal{O}}$ be the global generic type of the closed ball \mathcal{O} (in the sense of [HHM08, Definition 7.17]), i.e $p_{\mathcal{O}}(x)$ says that $x \in \mathcal{O}$ but x is not in any proper sub-ball of \mathcal{O} . One sees that for every polynomial in n variables $f \in K[X]$ and $c \models p_{\mathcal{O}}^{\otimes n}|K$, $\text{val}(f(c)) = \min_i\{\text{val}(b_i)\}$, where $\{b_1, \dots, b_m\}$ are the coefficients of f .

For two generically stable types p and q we denote by $p(x) \otimes q(y)$ the generically stable type $\text{tp}(a, b/\mathbb{U})$ where $b \models q|\mathbb{U}$ and $a \models p|b\mathbb{U}$. It is equal to $q(y) \otimes p(x)$. We shorthand $p^{\otimes n}$ for $p \otimes \dots \otimes p$ (n times).

If p and q are generically stable types concentrated on some (type-)definable group G then we write pq for $f_*(p \otimes q)$, where $f(x, y) = x \cdot y$, and likewise p^n for $h_*p^{\otimes n}$ where $h(x_1, \dots, x_n) = x_1 \cdot \dots \cdot x_n$.

Generically stable groups. We review some definitions from [HRK19].

¹There are standard techniques from set theory that ensure the generalized continuum hypothesis from some point on while fixing a fragment of the universe (so this does not affect the questions asked in this paper), see [HK23].

Let G be a type-definable group, p an A -definable type on G and $g \in G$. The *left translate* of p by g , gp , is the definable type such that for any $A \cup g \subseteq A'$,

$$d \models p|_{A'} \Leftrightarrow gd \models gp|_{A'}.$$

Similarly the right translate pg .

Definition 2.2. Let G be a type-definable group. A definable type p is *left generic* if for any $A = \text{acl}(A)$ over which it is defined and $g \in G$, gp is definable over A . Similarly right generic.

Any generically stable left generic is also right generic.

Definition 2.3. A type-definable group will be called *generically stable* if it has a generically stable generic.

For a generically stable group G , let G^0 be the intersection of all definable subgroups of finite index. It is of bounded index and called the *connected component* of G . We have the following: p is the unique generic type of G if and only if for all $g \in G$, $gp = p$ if and only if $G = G^0$. If $G = G^0$ then we say that G is *connected*.

Algebraic Geometry. We will assume basic knowledge of schemes and group schemes, see [GW10, Sta23] for the relevant definitions.

For a field F , by a variety over F we mean a geometrically integral separated scheme of finite type over F . We will reserve capital letters V, U, W, \dots for varieties and calligraphic letters $\mathcal{V}, \mathcal{U}, \mathcal{W}, \dots$ for general schemes, usually over valuation rings.

For an affine scheme $\mathcal{V} = \text{Spec } A$ we denote by $D_{\mathcal{V}}(f)$ the *principal open affine subscheme* defined by f , i.e. $\text{Spec } A_f$ for some $f \in A$. For a projective variety we denote by $D_{+, \mathcal{V}}(f)$ the projective analog. Similarly for affine or projective varieties.

For any variety V over F and field $L \supseteq F$ we denote by $V_L := V \times_{\text{Spec } F} \text{Spec } L$ the base change of V to L . Since V is geometrically integral over F , V_L is still a variety over L .

Let F be a valued field. Most of our schemes over \mathcal{O}_F will be quasi-compact integral and faithfully flat separated dominant over \mathcal{O}_F . For a valued field extension $L \supseteq F$, the base change $\mathcal{V}_{\mathcal{O}_L} = \mathcal{V} \times_{\text{Spec } \mathcal{O}_F} \text{Spec } \mathcal{O}_L$ is still quasi-compact and faithfully flat separated dominant over \mathcal{O}_L . If we further assume that \mathcal{V}_F is a variety over F then by Fact 3.19, $\mathcal{V}_{\mathcal{O}_L}$ is also an integral scheme. As \mathcal{V} is separated over \mathcal{O}_F , by the valuative criterion for separatedness, we have $\mathcal{V}_{\mathcal{O}_L}(\mathcal{O}_L) \subseteq \mathcal{V}_L(L)$ for any valued field extension $L \supseteq F$.

We will denote by V the definable set $V_{\mathbb{K}}(\mathbb{K})$, and by $\mathcal{V}(\mathcal{O})$ the pro-definable set $\mathcal{V}_{\mathcal{O}_{\mathbb{K}}}(\mathcal{O})$, hopefully without causing too much confusion.

Given such a scheme \mathcal{V} with \mathcal{V}_F a variety, we can view $\mathcal{V}(\mathcal{O})$ as a type-definable subset of the definable set \mathcal{V}_F or as a pro-definable subset of an inverse limit of powers of \mathcal{O} , in (possibly) infinitely many variables.

It will be convenient to view it as a type-definable subset of \mathcal{V}_F when, e.g., \mathcal{V} is a group scheme over \mathcal{O}_F , and then $\mathcal{V}(\mathcal{O})$ is a type-definable subgroup of \mathcal{V}_F . It will be convenient to view it as a pro-definable set when, e.g., we consider the pro-definable reduction map $r : \mathcal{V}(\mathcal{O}) \rightarrow \mathcal{V}_{\mathbf{k}}$. In this situation the reduction map is just given by res . If we view it as a type-definable subset of \mathcal{V}_F then, in general, the reduction map r is not equal to the res map on $\mathcal{V}(\mathcal{O})$.

We refer to [Hal21, Section 2.2] for an explanation how these can be thought of as (pro-)definable sets.

2.1. The Functor. We review some necessary definitions and results from [Hal21]. The results of this section will be tacitly assumed throughout the paper without further reference.

Let K be an algebraically closed valued field. The category (SPVar/K) , of *stably pointed varieties*, is the category of pairs (V, p) , where V is a variety over K and p is a Zariski dense

generically stable type over K , concentrated on V . Maps are morphisms between varieties that map the types accordingly. The category $(\text{SPVar}^{\text{aff}}/K)$ is the restriction of (SPVar/K) to affine varieties.

We define a functor from $(\text{SPVar}^{\text{aff}}/K)$ to the category of affine schemes over \mathcal{O}_K given by $\Phi(V, p) = \text{Spec } K[V]^p$, where

$$K[V]^p = \{f \in K[V] : p \vdash \text{val}(f(x)) \geq 0\}.$$

The scheme $\mathcal{V} := \Phi(V, p)$ is an affine integral scheme flat over \mathcal{O}_K . Note that by definition p is concentrated on $\mathcal{V}(\mathcal{O})$. Moreover, $V \cong \mathcal{V}_K := \mathcal{V} \times_{\mathcal{O}_K} K$ and \mathcal{V} enjoys the maximum modulus principle with respect to p (mmp w.r.t. p):

For every regular function f on $\mathcal{V}_{\mathbb{K}}$ there is some $\gamma_f \in \Gamma$ such that $p \vdash \text{val}(f(x)) = \gamma_f$ and for any $h \in \mathcal{V}(\mathcal{O})$, $\text{val}(f(h)) \geq \gamma_f$. As a result if $\mathcal{U} \subseteq \mathcal{V}$ is an open affine subscheme with $\mathcal{U}(\mathcal{O}) \neq \emptyset$ then p is concentrated on $\mathcal{U}(\mathcal{O})$. For the definition of the maximum modulus principle for general quasi-compact separated schemes over \mathcal{O}_K see [Hal21, Definition 4.3.1].

If either K is sufficiently saturated or \mathcal{V} is of finite type over \mathcal{O}_K then \mathcal{V} has an \mathcal{O}_K -point. Thus $\Phi(V_{\mathbb{K}}, p)$ is faithfully flat and dominant over \mathcal{O} [Hal21, Proposition 3.1.4]. On the other hand, by [Hal21, Proposition 4.2.15] $\Phi(V, p) \times_{\mathcal{O}_K} \mathcal{O} = \Phi(V_{\mathbb{K}}, p)$ and thus by faithfully flat descent ([GW10, Corollary 14.12]) $\mathcal{V} = \Phi(V, p)$ is faithfully flat over \mathcal{O}_K as well (though might not have an \mathcal{O}_K -point). It is dominant over \mathcal{O}_K by [GW10, Exercise 14.14].

We now expand the above to a larger family of valued fields.

Definition 2.4. (1) We say that a non-trivially valued field (F, val) is *gracious* if it is perfect henselian defectless with perfect residue field.

(2) We say that a generically stable F -definable type p is *strictly based on F* if $\Gamma_{F(c)} = \Gamma_F$ for $c \models p|F$.

Remark 2.5. Note that if F has a divisible value group then every generically stable type over F is strictly based on F .

Let $(\text{SPVar}^{\text{aff}}/F)$ be the category of pairs (V, p) , where V is an affine variety over F and p is a generically stable F -definable type, strictly based on F and Zariski dense in V . Likewise, (SPVar/F) for general varieties.

For a gracious valued field F and $(V, p) \in (\text{SPVar}^{\text{aff}}/F)$ we let

$$F[V]^p = \{f \in F[V] : p \vdash \text{val}(f(x)) \geq 0\},$$

and $\Phi_F(V, p) = \text{Spec } F[V]^p$. So $\text{Spec } \Phi_F(V, p)$ is an integral affine scheme. The following is an important result allowing us apply descent arguments.

Fact 2.6. [Hal21, Corollary 4.2.13, Proposition 4.2.15] *Let F be a gracious field, $K \supseteq F$ a model of ACVF and $(V, p) \in (\text{SPVar}^{\text{aff}}/F)$. Then*

$$\Phi_K(V_K, p) = \Phi_F(V, p) \times_{\text{Spec } \mathcal{O}_F} \text{Spec } \mathcal{O}_K.$$

As a result, by faithfully flat descent, $\Phi_F(V, p)$ is faithfully flat over \mathcal{O}_F [GW10, Corollary 14.12] and dominant over \mathcal{O}_F by [GW10, Exercise 14.14]. Due to the above fact we usually omit the subscript from Φ_F , unless a precise distinction is needed. In particular, the definable set $\Phi(V, p)(\mathcal{O})$ is well-defined.

If $\Phi(V, p)$ is of finite type over \mathcal{O}_F then by flatness, [Nag66], it is finitely presented over \mathcal{O}_F .

The function Φ commutes with products in the following sense: If $(V, p), (U, q) \in (\text{SPVar}^{\text{aff}}/F)$ and $p \otimes q$ is strictly based on F then $\Phi(V \times_F U, p \otimes q) = \Phi(V, p) \times_{\text{Spec } \mathcal{O}_F} \Phi(U, q)$, see [Hal21, Proposition 4.2.18].

We do not know if Φ commutes with open immersions, e.g. for $U \subseteq V$ an open subvariety, we do not know if $\Phi(U, p)$ is open subscheme of $\Phi(V, p)$. However, by passing to an open subvariety of U we can always assume that this is the case (taking $W = U$ below):

Lemma 2.7. *Let F be a gracious valued field. Let $(V, p), (U, q) \in (\text{SPVar}^{\text{aff}}/F)$, $W \subseteq U$ an open affine subvariety and $\varphi : W \rightarrow V$ an open immersion with $\varphi_*q = p$. Then there exist $f \in F[U]$ and $g \in F[V]$, with $q \vdash \text{val}(f(x)) = 0$ and $p \vdash \text{val}(g(x)) = 0$, such that*

- (1) $D_U(f) \subseteq W$ and φ restricts to an isomorphism φ' between $D_U(f)$ and $D_V(g)$
- (2) For $\mathcal{V} = \Phi(V, p)$ and $\mathcal{U} = \Phi(U, q)$, $\Phi(D_U(f), q) = D_{\mathcal{U}}(f)$ and $\Phi(D_V(g), p) = D_{\mathcal{V}}(g)$
- (3) $\Phi(\varphi') : D_{\mathcal{U}}(f) \rightarrow D_{\mathcal{V}}(g)$ is an isomorphism.

Proof. By a standard argument, there are $f \in F[U]$ and $g \in F[V]$ such that $D_U(f) \subseteq W$, $D_V(g) \subseteq f(W)$ and that φ restricts to an isomorphism between $D_U(f)$ and $D_V(g)$. This gives (1). As p and q are strictly based on F we may further assume that $q \vdash \text{val}(f(x)) = 0$ and $p \vdash \text{val}(g(x)) = 0$ (since $\text{val}(f(c)) \in \Gamma_F$ for $c \models q|F$, and likewise for g and p). In particular, $(F[U]_f)^q = (F[U]^q)_f$ and likewise for g . Thus if we set $\mathcal{U} = \Phi(U, q)$ and $\mathcal{V} = \Phi(V, p)$ we get (2).

Item (3) follows since Φ obviously preserves isomorphisms. \square

3. SOME MORE ON THE FUNCTOR

For the entirety of this section we let F be a gracious valued field and K an algebraically closed valued field containing F . The aim of this section is to prove some more properties of Φ .

3.1. Integrality of the Fibers. In this subsection we prove that given $(V, p) \in (\text{SPVar}^{\text{aff}}/F)$, the fibers of $\Phi(V, p) \rightarrow \text{Spec } \mathcal{O}_F$ are geometrically integral. We then deduce, assuming $\Phi(V, p)$ is of finite type over \mathcal{O}_F , that $\Phi(V, p)$ is (locally) smooth over \mathcal{O}_F .

We denote by $(K, \text{val})^{\text{sh}}$ the Shelah expansion of (K, val) , see [Sim15, Section 3.3] and Appendix A. We will require the following result whose proof we postpone to the appendix, see Proposition A.4.

Proposition 3.1. *Let T be a complete NIP theory, in a first order language \mathcal{L} and $M \models T$. Let $\mathbb{U} \succ M$ be a large saturated model of T and \mathbb{U}^* an expansion to a saturated model of $\text{Th}(M^{\text{sh}})$ (so $\mathbb{U}^* \upharpoonright \mathcal{L} = \mathbb{U}$).*

For every global type p , if p is generically stable over M then there exists a unique extension $p \subseteq q \in S^{\mathcal{L}^{\text{sh}}}(\mathbb{U}^)$ which is generically stable over M .*

The following observations is simple, yet useful:

Fact 3.2. *Let $R \subseteq S$ be (possibly trivial) valuation rings with fraction field F . Then for every S -algebras A and B*

$$A \otimes_R B = A \otimes_S B.$$

Proof. Let $a \in A$ and $b \in B$ and let $r \in S \setminus R$, so $r^{-1} \in R$.

$$(ra) \otimes b = (ra) \otimes (r^{-1}rb) = (rr^{-1})a \otimes (rb) = a \otimes (rb),$$

as needed. \square

Proposition 3.3. *Let $(V, p) \in (\text{SPVar}^{\text{aff}}/F)$ and let $\mathcal{V} = \Phi(V, p)$. All the fibers of $\mathcal{V} \rightarrow \text{Spec } \mathcal{O}_F$ are geometrically integral.*

Proof. After base change, we may assume that $F = K$.

Let v be the valuation on K , $\mathcal{O}_v := \mathcal{O}_K$ and $\Gamma_v := \Gamma_K$. Let $\mathfrak{p}_w \in \text{Spec } \mathcal{O}_K$ be a prime ideal corresponding to a coarsening w of v with value group $\Gamma_w := \Gamma_v/\Delta_w$ and residue field \mathbf{k}_w ; so $\mathcal{O}_w := (\mathcal{O}_v)_{\mathfrak{p}_w}$ is the corresponding valuation ring.

If \mathfrak{p}_w corresponds to the generic point, the generic fiber is just $\mathcal{V} \times_{\text{Spec } \mathcal{O}_v} \text{Spec } K = V$ so integral. Otherwise, w is a non-trivial coarsening of v .

Let $(K, \text{val})^{sh}$ be the Shelah expansion of (K, val) . Let \tilde{p} be a generically stable extension of p living in the saturated expansion $(\mathbb{K}, \text{val})^*$, as given by Proposition 3.1. Possibly after passing to T^{eq} , which we can do by Fact A.3, the valuation w is now definable in $(K, v)^{sh}$ and thus defines a coarsening of v in \mathbb{K} as well, which we will also denote by w . Let p_w be the restriction of \tilde{p} to (\mathbb{K}, w) ; p is just the restriction of \tilde{p} to (\mathbb{K}, v) . Note that p_w is still K -definable and Zariski dense in V . It is generically stable by Fact A.3. Let $K[V]^{p_w} = \{f \in K[V] : w(f(c)) \geq 0, \text{ for } c \models p_w|K\}$.

Claim. *Viewing $K[V]^{p_v} \otimes_{\mathcal{O}_v} \mathcal{O}_w$ as a subset of $K[V]^{p_w}$,*

$$K[V]^{p_w} = K[V]^{p_v} \otimes_{\mathcal{O}_v} \mathcal{O}_w.$$

Proof. Let $c \models \tilde{p}|K$, then obviously $c \models p_v|K$ and $c \models p_w|K$.

Let $f \in K[V]$ be a regular function satisfying $w(f(c)) \geq 0$. As p_w is generically stable, $w(f(c)) \in \Gamma_w$. Since w is a coarsening of v , there are two possibilities for $v(f(c))$, either $v(f(c)) \geq 0$ or $0 > v(f(c)) \in \Delta_w$. If $v(f(c)) \geq 0$ there is nothing to show. If $0 > v(f(c)) \in \Delta_w$ then let $a \in K$ be such that $v(f(c)) = v(a)$. Such an element exists since p_v is orthogonal to Γ_v . Writing $f(c) = a \cdot \frac{f(c)}{a}$ concludes the proof since $v\left(\frac{f(c)}{a}\right) = 0$ and $v(a) \in \Delta_w$ and hence $w(a) = 0$. \square (claim)

The fiber over \mathfrak{p}_w is by definition $\mathcal{V} \times_{\text{Spec } \mathcal{O}_v} \text{Spec } \mathbf{k}_w$, or in other words

$$\text{Spec}(K[V]^p \otimes_{\mathcal{O}_v} \mathbf{k}_w).$$

As $\mathbf{k}_w = \mathcal{O}_w \otimes_{\mathcal{O}_v} \mathbf{k}_w$, by Fact 3.2 this is equal to

$$\text{Spec}(K[V]^{p_v} \otimes_{\mathcal{O}_v} \mathcal{O}_w \otimes_{\mathcal{O}_v} \mathbf{k}_w).$$

By the claim above and Fact 3.2 again we get that it is equal to

$$\text{Spec}(K[V]^{p_w} \otimes_{\mathcal{O}_v} \mathbf{k}_w).$$

It is thus sufficient to prove the following claim:

Claim. *For any $(V, p) \in (SPVar^{aff}/K)$, $\Phi(V, p)$ has an integral special fiber.*

Proof. We show that the special fiber $K[V]^p \otimes_{\mathcal{O}_K} \mathbf{k}_K \cong K[V]^p / (\mathfrak{m}_K K[V]^p)$ is an integral domain. Let $f, g \in K[V]^p$ and assume that $fg \in \mathfrak{m}_K K[V]^p$. In particular $\text{val}(f(c)g(c)) > 0$ for $c \models p|K$ and thus without loss of generality $\text{val}(f(c)) > 0$. Since p is generically stable, there is some $a \in K$ such that $\text{val}(f(c)) = \text{val}(a) > 0$ so $\text{val}(f(c)/a) = 0$ and hence $f(c)/a \in K[V]^p$. Consequently, $f = a \cdot f/a \in \mathfrak{m}_K K[V]^p$, as needed. \square (claim)

Apply the last claim to (K, w) and p_w and conclude. \square

We now turn to showing smoothness.

By [Nag66, Theorem 3], a flat scheme of finite type over a valuation ring is finitely presented. Actually the proof gives a bit more (see [Nag66, page 159]):

Fact 3.4. *Let R be a valuation ring with maximal ideal \mathfrak{m} and let $R[X]$ be the polynomial ring over R with variables $X = (x_1, \dots, x_n)$. Then for every $I \trianglelefteq R[X]$ such that $R[X]/I$ is flat over R , if $f_1 + \mathfrak{m}[X], \dots, f_r + \mathfrak{m}[X]$ generate $(I + \mathfrak{m}[X])/\mathfrak{m}[X]$ then f_1, \dots, f_r generate I .*

Proposition 3.5. *Let \mathcal{V} be an integral affine scheme of finite type over \mathcal{O}_K with an \mathcal{O}_K -point. If the special fiber $\mathcal{V}_{\mathfrak{k}}$ is reduced then the smooth locus of \mathcal{V} over \mathcal{O}_K is open non-empty and has an \mathcal{O}_K -point.*

Proof. Assume that $\mathcal{V} = \text{Spec } \mathcal{O}_K[X]/I$. Since \mathcal{V} has an \mathcal{O}_K -point it is faithfully flat over \mathcal{O}_K by [Hal21, Proposition 3.1.4], finitely presented over \mathcal{O}_K by [Nag66, Theorem 3] and $\mathcal{V}_{\mathbf{k}}(\mathbf{k}_K)$ is non empty.

The coordinate ring of the special fiber is isomorphic to

$$(\mathcal{O}_K[X]/I) \otimes_{\mathcal{O}_K} \mathbf{k}_K \cong \mathcal{O}_K[X]/(I + \mathfrak{m}_K[X]) \cong (\mathcal{O}_K[X]/\mathfrak{m}_K[X]) / (I + \mathfrak{m}_K[X]/\mathfrak{m}_K[X]);$$

by Fact 3.4 there are generators f_1, \dots, f_r of I such that $\overline{f}_1, \dots, \overline{f}_r$, with $\overline{f}_i = f_i + \mathfrak{m}_K[X]$, are non-zero and generate $I + \mathfrak{m}_K[X]$.

As $\mathcal{V}_{\mathbf{k}}$ is reduced (and of finite type over \mathbf{k}_K) and \mathbf{k}_K is perfect, by [GW10, Theorem 6.19] the singular locus of $\mathcal{V}_{\mathbf{k}}$ is a proper closed subvariety. Thus there is some $a \in \mathcal{V}_{\mathbf{k}}(\mathbf{k}_K)$ and a minor $\det \overline{M}(a)$ of $\left(\frac{\partial \overline{f}_i}{\partial x_j}(a) \right)_{1 \leq i \leq r, 1 \leq j \leq n}$ of order $n - d$ with $d = \dim \mathcal{V}_{\mathbf{k}}$ and $\det \overline{M}(a) \neq 0$.

By [Hal21, Theorem 3.2.4], there is some $b \in \mathcal{V}(\mathcal{O}_K)$ with $r(b) = a$, where $r : \mathcal{V}(\mathcal{O}_K) \rightarrow \mathcal{V}_{\mathbf{k}}(\mathbf{k}_K)$ is the reduction map. Viewing b as an element of \mathcal{O}_K^n we exactly get that $\text{res}(b) = a$.

Hence $\text{val}(\det M(b)) = 0$ where $\det M(b)$ is the corresponding minor of $\left(\frac{\partial f_i}{\partial x_j}(b) \right)_{1 \leq i \leq r, 1 \leq j \leq n}$. This exactly means that the prime ideal $\mathfrak{p}_b = \{f \in \mathcal{O}_K[X]/I : \text{val}(f(b)) > 0\}$ of $\mathcal{O}_K[X]/I$ is a smooth point of \mathcal{V} over \mathcal{O}_K of relative dimension d .

Thus the smooth locus is open non-empty and contains the \mathcal{O}_K -point b . \square

Corollary 3.6. *Let $(V, p) \in (\text{SPVar}^{\text{aff}}/F)$. If $\Phi_F(V, p)$ is of finite type over \mathcal{O}_F then there exists an open affine subvariety $U \subseteq V$ such that $\Phi_F(U, p)$ is an open affine subscheme of $\Phi_F(V, p)$ and $\Phi_F(U, p)$ is smooth over \mathcal{O}_F .*

Proof. The \mathcal{O}_K -scheme $\mathcal{V}_{\mathcal{O}_K} = \Phi_K(V_K, p)$ satisfies the assumptions of Proposition 3.5 (by Proposition 3.3). As \mathcal{V} is affine, the smooth locus of \mathcal{V} over \mathcal{O}_F is a finite union of principal open affine subschemes of \mathcal{V} (given by Jacobian determinants). Base changing this open subscheme to $\text{Spec } \mathcal{O}_K$ we get the smooth locus of $\mathcal{V}_{\mathcal{O}_K}$.

Hence there exists some $f \in F[V]^p$ such that $D_{\mathcal{V}_{\mathcal{O}_K}}(f)$ is smooth over \mathcal{O}_K and $D_{\mathcal{V}_{\mathcal{O}_K}}(f)(\mathcal{O}_K) \neq \emptyset$. As smoothness descends under faithfully flat descent [DG67, IV.17.7.3], $\mathcal{U} := D_{\mathcal{V}}(f)$ is smooth over \mathcal{O}_F .

Since $\mathcal{U}(\mathcal{O})$ is non empty, p is concentrated on $\mathcal{U}(\mathcal{O})$, i.e. $\text{val}(f(c)) = 0$ for $c \models p|F$. Consequently, $(F[V]^p)_f = (F[V]_f)^p$ and thus, setting $U = \text{Spec } F[V]_f$, $\Phi_F(U, p) = \mathcal{U}$. \square

3.2. Strongly Stably dominated types. We recall the definition of a strongly stably dominated type (at least an equivalent definition, see [HL16, Proposition 8.1.2]).

Definition 3.7. Let $L = \text{dcl}(L)$ be a valued field and p an A -definable type on a variety V over L . We say that p is *strongly stably dominated* if there exists some L -definable map g into a variety over the residue field for which $\dim(p) = \dim(g_*p)$.

Every strongly stably dominated type is generically stable; not every generically stable type is strongly stably dominated, but every generically stable type concentrated on an algebraic curve is strongly stably dominated [HL16].

Fact 3.8. *Let L be a valued field with $\text{dcl}(L) = L$, V an affine variety over L , with $n = \dim(V)$, and p a Zariski dense L -definable type concentrated on V . If p is strongly stably dominated then there exists an open affine subvariety $U \subseteq V$ and a finite morphism $f : U \rightarrow \mathbb{A}_L^n$ with $f_*p = p_{\mathcal{O}}^{\otimes n}$.*

Proof. By [HL16, Lemma 8.1.2], we can find such $f : U \rightarrow \mathbb{A}_L^n$ with f quasi-finite. By [Sta23, Tag 02NW], we may further reduce f to achieve finiteness. \square

The purpose of this section is to present some results on strongly stably dominated types. We start with some known results.

Fact 3.9. [HHS21, Lemmas 2.13, 2.14]

- (1) *The generic types of a definable generically stable group are strongly stably dominated.*
- (2) *If the unique generically stable type of a type-definable connected generically stable subgroup H of an algebraic group G is strongly stably dominated then H is definable.*

For the next result we will need the following, which is hidden inside the proof of [Hal21, Theorem 3.2.4].

Fact 3.10. *Let \mathcal{V} be an affine integral scheme flat and dominant over \mathcal{O}_K . If $\mathcal{V}(\mathcal{O}) = \emptyset$ then $\mathcal{V}_{\mathbf{k}} = \emptyset$. If \mathcal{V} is of finite presentation over \mathcal{O}_K then $\mathcal{V}(\mathcal{O}_K) = \emptyset$ implies that $\mathcal{V}_{\mathbf{k}}(\mathbf{k}_K) = \emptyset$.*

Proof. Let $\mathcal{V} = \text{Spec } \mathcal{O}_K[X]/I$ be the given affine scheme. Assume that $\mathcal{V}_{\mathbf{k}} \neq \emptyset$, after base-changing we also assume that $\mathcal{V}_{\mathbf{k}}(\mathbf{k}_K) \neq \emptyset$. By [Hal21, Lemma 3.2.2(2)], there exists $b \in \mathcal{V}(\mathcal{O})$. If \mathcal{V} is finitely presented over \mathcal{O}_K then $\mathcal{V}(\mathcal{O}_K) \neq \emptyset$ as well. \square

In the following, it is convenient to view $\mathcal{V}(\mathcal{O})$ has a pro-definable subset of an inverse limit of (powers of) \mathcal{O} . Thus b might in general be an infinite tuple.

Lemma 3.11. *Let \mathcal{V} be an affine integral scheme flat over \mathcal{O}_K , with \mathcal{V}_K geometrically integral, and assume that it has the mmp w.r.t. p , a generically stable type over K concentrated on $\mathcal{V}(\mathcal{O})$. If $\mathcal{V}_{\mathbf{k}}$ is integral then for any $b \models p|K$, $\mathcal{V}_{\mathbf{k}} = \text{Spec}(\mathbf{k}_K[\text{res}(b)])$. In particular, r_*p is the unique generic type of $\mathcal{V}_{\mathbf{k}}$.*

Proof. Assume that $\mathcal{V} = \text{Spec } \mathcal{O}_K[X]/I$. Since p is concentrated on $\mathcal{V}(\mathcal{O})$, \mathcal{V} has an \mathcal{O} -point. Thus, as $\mathcal{V}_{\mathcal{O}}$ is integral by Fact 3.19, $\mathcal{V}_{\mathcal{O}}$ is dominant and faithfully flat over \mathcal{O} [Hal21, Proposition 3.1.4]; by faithfully flat descent so is \mathcal{V} over \mathcal{O}_K .

Let $b \models p|K$. By [Hal21, Lemma 3.2.2(1)], $\mathcal{O}_K[X]/I \cong \mathcal{O}_K[b]$, given by $X \mapsto b$. Now let $\mathcal{O}_K[b] \otimes_{\mathcal{O}_K} \mathbf{k}_K \rightarrow \mathbf{k}_K[\text{res}(b)]$ be the surjection given by the residue map; we claim it is injective. Note that every element of $\mathcal{O}_K[b] \otimes_{\mathcal{O}_K} \mathbf{k}_K$ is of the form $f(b) \otimes 1$.

Let $f \in \mathcal{O}_K[X]/I$ and assume that $\overline{f}(\text{res}(b)) = 0$, i.e. $\text{val}(f(b)) > 0$. Thus $\text{val}(f(c)) > 0$ for any $c \models p|L$ for any large enough model L containing K . By the maximum modulus principle, $D_{\mathcal{V}}(f)(\mathcal{O}) = \emptyset$, where $D_{\mathcal{V}}(f)$ is the principal open subscheme of \mathcal{V} given by f .

Note that $D_{\mathcal{V}}(f)$ is also integral, flat and dominant over \mathcal{O}_K as an open subscheme of such; so by Fact 3.10 we get that $D_{\mathcal{V}}(f)_{\mathbf{k}}(\mathbf{k}) = \emptyset$ which gives $f = 0$ in $\mathcal{O}_K[b] \otimes_{\mathcal{O}_K} \mathbf{k}_K \subseteq \mathcal{O}_K[b] \otimes_{\mathcal{O}_K} \mathbf{k}$. \square

Lemma 3.12. *Let $(V, p) \in (\text{SPVar}^{\text{aff}}/K)$. The type p is strongly stably dominated if and only if $\dim V = \dim \mathcal{V}_{\mathbf{k}}$, for $\mathcal{V} = \Phi(V, p)$. Furthermore, in this situation p is stably dominated via $r : \mathcal{V}(\mathcal{O}) \rightarrow \mathcal{V}_{\mathbf{k}}$.*

Proof. Assume that $\dim V = \dim \mathcal{V}_{\mathbf{k}}$. Since $\mathcal{V} \rightarrow \text{Spec } \mathcal{O}_K$ has integral fibers (Proposition 3.3), by Lemma 3.11 r_*p is the generic type of $\mathcal{V}_{\mathbf{k}}$, so $\dim(p) = \dim V = \dim \mathcal{V}_{\mathbf{k}} = \dim(r_*p)$, so p is strongly stably dominated. By [Hal21, Proposition 4.1.4], p is stably dominated via r .

For the other direction, by Fact 3.8 there exists an open affine subvariety $U \subseteq V$ and a finite morphism $f : U \rightarrow \mathbb{A}_K^n$ satisfying $f_*p = p_{\mathcal{O}}^{\otimes n}$ for $n = \dim V$. Writing $f = (f_i)$, then $f_i \in K[U]^p$ for all i . Letting $\overline{f}_i = \text{res} \circ f_i$, by Lemma 3.11 we have that for $c \models p|K$

$$\overline{f}_i(c) = \text{res}(f_i(c)) \in \mathbf{k}_K[\text{res}(c)] \cong K[U]^p \otimes_{\mathcal{O}_K} \mathbf{k}_K.$$

Since $f(c) \models p_{\mathcal{O}}^{\otimes n}$ this implies that $\dim \mathbf{k}_K[\text{res}(c)] = n$, i.e. $\dim \mathcal{V}_{\mathbf{k}} = n$. \square

The following two results will not be used in the rest of the paper, but we believe they might be interesting in their own right. The first can be seen as a geometrical analog of [HRK19, Proposition 4.6] and the second as a “strengthening” of [HRK19, Lemma 5.1].

Below we will repeatedly use the fact that Φ commutes with products.

Proposition 3.13. *Let G be an algebraic group over K and p a Zariski dense generically stable over K type concentrated on G . Assume that $p^2 = p$, let $U \subseteq G$ be an open subvariety and let $H = \text{Stab}(p)$. Let $\mathcal{U} = \Phi(U, p)$ and $r : \mathcal{U}(\mathcal{O}) \rightarrow \mathcal{U}_{\mathbf{k}}$ be the reduction map.*

*Then there exists pro- K -definable connected group \mathfrak{h} in \mathbf{k} , a pro- K -definable homomorphism $\phi : H \rightarrow \mathfrak{h}$ and an injective pro- K -definable map $f : r_*p \rightarrow \mathfrak{h}$, with $f_*r_*p = \phi_*p$ the unique generic type of \mathfrak{h} .*

If p is strongly stably dominated then \mathfrak{h} can be taken definable and p is stably dominated via ϕ .

Proof. Let $U \subseteq G$ be an affine open subvariety over K and $W \subseteq G \times G$ an affine open subvariety such that $a \cdot b \in U$ for any $(a, b) \in W$. Since G is irreducible, there is no harm in assuming that $W \subseteq U \times U$. Let $\mathcal{W} = \Phi(W, p \otimes p)$ and $\mathcal{U} = \Phi(U, p)$, by Lemma 2.7 we may further assume that $\mathcal{W} \subseteq \mathcal{U} \times \mathcal{U}$ is an open subscheme. By functoriality of Φ the multiplication map induces a morphism $M : \mathcal{W} \rightarrow \mathcal{U}$ over \mathcal{O}_K and thus also $M_{\mathbf{k}} : \mathcal{W}_{\mathbf{k}} \rightarrow \mathcal{U}_{\mathbf{k}}$ which is now a pro- \mathbf{k}_K -definable map; note that $\mathcal{W}_{\mathbf{k}} \subseteq \mathcal{U}_{\mathbf{k}} \times \mathcal{U}_{\mathbf{k}}$ is an affine open subvariety over \mathbf{k}_K . Furthermore, by further restricting W we may assume that $M(a, M(b, c)) = M(M(a, b), c)$ for all $(a, b, c) \models p^{\otimes 3}|K$. Likewise there exists an affine open subvariety $W' \subseteq W$, where the inverse map $W' \rightarrow W$ is defined. Again, by functoriality, this defines $\iota : \mathcal{W}' \rightarrow \mathcal{W}$ which induces a map $\iota_{\mathbf{k}} : \mathcal{W}'_{\mathbf{k}} \rightarrow \mathcal{W}_{\mathbf{k}}$.

We claim that $(r_*p, M_{\mathbf{k}})$ is an abstract group chunk over K , in the sense of [HRK19, Definition 3.14], where $r : \mathcal{U}(\mathcal{O}) \rightarrow \mathcal{U}_{\mathbf{k}}$.

(1) Let $(a, b) \models p \otimes p|K$, so $(r(a), r(b)) \models r_*p \otimes r_*p|K$. We need to show that $M_{\mathbf{k}}(r(a), r(b)) \models p|K$. Indeed,

$$M_{\mathbf{k}}(r(a), r(b)) = r(M(a, b)) = r(a \cdot b) \models r_*p,$$

where the latter equality follows since $p^2 = p$.

(2) Let $(a, b) \models p \otimes p|K$, as $(a^{-1}, a \cdot b) \models p \otimes p|K$ as well, $(a^{-1}, ab) \in \mathcal{W}(\mathcal{O})$. Following the diagrams we get

$$M_{\mathbf{k}}(\iota_{\mathbf{k}}(r(a)), M_{\mathbf{k}}(r(a), r(b))) = M_{\mathbf{k}}(r(\iota(a)), r(M(a, b))) = r(M(\iota(a), M(a, b))) = r(b),$$

and thus $r(b) \in \text{dcl}(K, r(a), M_{\mathbf{k}}(r(a), r(b)))$ and likewise

$$r(a) \in \text{dcl}(K, r(b), M_{\mathbf{k}}(r(a), r(b))).$$

(3) Generic associativity for $(r_*p, M_{\mathbf{k}})$ follows directly from that of (p, M) as was done above.

By [HRK19, Proposition 3.15], there exists pro- K -definable connected group \mathfrak{h} in \mathbf{k} and an injective pro- K -definable map $f : r_*p \rightarrow \mathfrak{h}$ such that for any $(a, b) \models r_*p^{\otimes 2}|K$, $f(M_{\mathbf{k}}(a, b)) = f(a) \cdot f(b)$ and f_*r_*p is the unique generic type of \mathfrak{h} . In particular, if $(a, b) \models p^{\otimes 2}$ then

$$f(r(a \cdot b)) = f(M_{\mathbf{k}}(r(a), r(b))) = f(r(a)) \cdot f(r(b)),$$

so by [HRK19, Proposition 3.16] there exists a unique pro- K -definable homomorphism $\phi : H \rightarrow \mathfrak{h}$, where $H = \text{Stab}(p)$ such that for any $a \models p|K$, $f(r(a)) = g(a)$.

If p is strongly stably dominated then H is definable by Fact 3.9, so we may assume that \mathfrak{h} is definable as in [HRK19, Proposition 4.6]. We are left in showing that p is stably dominated via ϕ . By Lemma 3.12, p is stably dominated via r . Since f is K -definable and injective, p is also stably dominated via ϕ . \square

Lemma 3.14. *Let A be a commutative algebraic group over K and p a strongly stably dominated type concentrated on A , which is definable over K . If p^n is Zariski dense in A then it is the generic of a coset of a generically stable connected definable subgroup of A , where $n = \dim A$.*

Proof. We may assume that $n > 0$. By [HRK19, Lemma 5.1], there exists a type-definable generically stable connected subgroup H of A with unique generic $p^{\mp 2n}$, where

$$p^{\mp 2n} = f_*p^{\otimes 2n}$$

for $f : A^{2n} \rightarrow A$ sending $(a_1, \dots, a_{2n}) \mapsto a_1^{-1} \cdot a_2 \cdot a_3^{-1} \cdot \dots \cdot a_{2n}$ and, moreover, p is concentrated on a coset of H . Let $a \in A$ be such that ap is concentrated on H , where ap is the pushforward of p under the multiplication-by- a map.

Since $p^{\mp 2n}$ is Zariski dense (because p^n is Zariski dense) and strongly stably dominated, by Fact 3.9 H is definable. By [HRK19, Proposition 4.6], there exists an ω -stable connected group \mathfrak{h} in the residue field sort and a definable surjection $\phi : H \rightarrow \mathfrak{h}$, such that the generic of H is stably dominated by the generic of \mathfrak{h} via ϕ , and $\dim H = \dim(p^{\mp 2n}) = \dim(\phi_* p^{\mp 2n}) = \dim \mathfrak{h} = n$. Since H is Abelian, \mathfrak{h} is necessarily Abelian as well.

Claim. $\varphi_*((ap)^n) = \varphi_* p^{\mp 2n}$.

Proof. Let $q := \varphi_*(ap)$. Since $\dim(\mathfrak{h}) = n$, $\dim(q^n \cdot q^n) = \dim(q^n)$ (e.g., since \dim is equal to the Morley rank). By [Kow00, Lemma 1.3], q^n is a generic of a coset of a type-definable subgroup of \mathfrak{h} with generic $q^n q^{-n} = q^{\mp 2n}$. Since \mathfrak{h} is connected with unique generic $q^{\mp 2n} = \phi_* p^{\mp 2n}$, $q^n = \varphi_*((ap)^n) = \varphi_* p^{\mp 2n}$, as needed. \square (claim)

By [HRK19, Lemma 4.9], $(ap)^n$ is a generic of H so, by connectedness of H , $(ap)^n = a^n p^n = p^{\mp 2n}$, as desired. \square

3.3. A finiteness condition. The purpose of this section is to provide a condition that guarantees that $\Phi(V, p)$ is of finite type over \mathcal{O}_F , for $(V, p) \in (\text{SPVar}^{\text{aff}}/F)$. Then we show that generic types of Abelian schemes over \mathcal{O}_K satisfy this condition.

Definition 3.15. We say that an extension of valued fields $(L, w)/(M, v)$ is *henselian* if w is the unique extension of v to L .

Proposition 3.16. *Let $(V, p) \in (\text{SPVar}^{\text{aff}}/F)$. If there exists a finite morphism $f : U \rightarrow \mathbb{A}_F^n$ satisfying that $f_* p = p_{\mathcal{O}}^{\otimes n}$ and for $c \models p|F$, $(F(c)/F(f(c)), \text{val})$ is a henselian extension then there exists an open subvariety $U \subseteq V$ such that $\Phi(U, p)$ is of finite type over \mathcal{O}_F .*

Proof. Let $c \models p|F$ and let $f : U \rightarrow \mathbb{A}_F^n$ be the finite morphism of varieties over F satisfying the assumption.

Assume that $f(c) = (f_1(c), \dots, f_n(c))$. Identifying $F[U]$ with $F[c]$ we have that $f_i(c) \in F[c]$ for $1 \leq i \leq n$, and as $f(c) \models p_{\mathcal{O}}^{\otimes n}|F$ they constitute a separating transcendence basis to $F(f(c))$ over F , i.e. for any (multi-)index set $I \subseteq \mathbb{N}$ and $g \in F[f(c)]$, if $g = \sum_{\nu \in I} a_{\nu} f(c)^{\nu}$ then

$$\text{val}(g) = \min\{\text{val}(a_{\nu})\}.$$

Since (F, val) is a defectless valued field, by the generalized stability theorem, [Kuh10, Theorem 1.1], $(F(c), \text{val})$ is also defectless. So $F(c)/F(f(c))$ is defectless henselian, and hence by, e.g. [Kuh, Lemma 11.15] this extension has a separating basis (as a valued vector space). I.e., since p is strictly based on F , $\Gamma_{F(c)} = \Gamma_F$, there are basis elements $g_1, \dots, g_d \in F(U)$, with $\text{val}(g_i(c)) = 0$, such that for any $a_1, \dots, a_d \in F[f(c)]$,

$$\text{val}\left(\sum_i a_i g_i(c)\right) = \min_i \{\text{val}(a_i)\}.$$

Let $W \subseteq U$ be an open subvariety for which $g_i \in F[W]$ for all $1 \leq i \leq d$. Note that $g_i, f_j \in F[W]^p$, for $1 \leq i \leq d$, $1 \leq j \leq n$, and by the above they generate $F[W]$ as an F -algebra. On the other hand if $h \in F[W]^p$ then by the choice of the g_i and f_j , it is in the \mathcal{O}_F -algebra generated by them. \square

Corollary 3.17. *Let $(V, p) \in (\text{SPVar}^{\text{aff}}/F)$. Assume that*

- (\dagger) *there exists a finite morphism $f : U \rightarrow \mathbb{A}_F^n$ satisfying*
- (a) *$f_* p = p_{\mathcal{O}}^{\otimes n}$ and*

(b) for any $c \models p|F$, $\text{tp}_{ACF}(c/F(f(c))) \vdash p|F$.

Then there exists an open subvariety $U \subseteq V$ such that $\Phi(U, p)$ is of finite type over \mathcal{O}_F .

Proof. Let $c \models p|F$ and let $f : U \rightarrow \mathbb{A}_F^n$ be the finite morphism of varieties over F satisfying (\dagger) . Henselianity of the extension $(F(c)/F(f(c)), \text{val})$ follows from condition (b) and quantifier elimination in ACVF. \square

We now show that the generic type of an Abelian scheme over \mathcal{O}_K satisfies (\dagger) of Corollary 3.17. We first need the following version of Noether's normalization lemma.

Lemma 3.18. *Let \mathcal{V} be an integral projective scheme of finite type over \mathcal{O}_K with $\mathcal{V}_{\mathbf{k}}$ integral and $n = \dim \mathcal{V}_K = \dim \mathcal{V}_{\mathbf{k}}$. Let p be a generically stable Zariski dense K -definable type concentrated on $\mathcal{V}(\mathcal{O})$ and assume that \mathcal{V} has the mmp w.r.t. p . Then there exists a finite morphism $f : \mathcal{V}_K \rightarrow \mathbb{P}_K^n$, mapping p to $p_{\mathcal{O}^{\otimes n}}$, which descends to \mathbf{k} in the sense that the following diagram commutes*

$$\begin{array}{ccc} \mathcal{V}_K(K) & \xrightarrow{f} & \mathbb{P}_K^n(K) \\ r \downarrow & & \downarrow r \\ \mathcal{V}_{\mathbf{k}}(\mathbf{k}_K) & \xrightarrow{\bar{f}} & \mathbb{P}_{\mathbf{k}}^n(\mathbf{k}_K) \end{array}$$

Proof. Assume that \mathcal{V}_K is embedded inside \mathbb{P}_K^m and likewise $\mathcal{V}_{\mathbf{k}}$ inside $\mathbb{P}_{\mathbf{k}}^m$.

By Noether normalization for projective varieties [Mum95, Corollary 2.29], there exists a linear subspace $\bar{L} \subseteq \mathbb{P}_{\mathbf{k}}^m(\mathbf{k}_K)$ of codimension $n+1$ disjoint from $\mathcal{V}_{\mathbf{k}}(\mathbf{k}_K)$. This linear subspace is defined by $n+1$ linearly independent vectors (forms) $\bar{v}_1, \dots, \bar{v}_{n+1} \in \mathbf{k}_K^{m+1}$. Let $v_1, \dots, v_{n+1} \in \mathcal{O}_K^{m+1}$ be lifts of these vectors. They are linearly independent over K and thus define a linear subspace of $L \subseteq \mathbb{P}_K^m(K)$ of codimension $n+1$. By properness of \mathcal{V} over \mathcal{O}_K , every K -point of \mathcal{V} is also an \mathcal{O}_K -point, so L is also disjoint from $\mathcal{V}_K(K)$.

Let $f : \mathcal{V}_K(K) \rightarrow \mathbb{P}_K^n(K)$ be the projection with center L and let $\bar{f} : \mathcal{V}_{\mathbf{k}}(\mathbf{k}_K) \rightarrow \mathbb{P}_{\mathbf{k}}^n(\mathbf{k}_K)$ be its reduction; it is the projection with center \bar{L} (by definition). By [Mum95, Corollary 2.29], they are both finite surjective morphisms.

We are left with verifying that p maps to $p_{\mathcal{O}^{\otimes n}}$. Let $b \models p|K$. By [Hal21, Proposition 4.3.5], any affine open subscheme $\mathcal{U} \subseteq \mathcal{V}$, with $\mathcal{U}(\mathcal{O}) \neq \emptyset$, has the mmp w.r.t. p . Thus by Lemma 3.11, $\text{res}(b)$ is a \mathbf{k}_K -generic of $\mathcal{V}_{\mathbf{k}}(\mathbf{k}_K)$ and since \bar{f} is dominant $\bar{f}(\text{res}(b)) \models (p_{\mathbf{k}}^{\otimes n})|K$, where $p_{\mathbf{k}}$ is the generic type of $\mathbb{P}_{\mathbf{k}}^1$. Since the diagram commutes, $f_*p = p_{\mathcal{O}^{\otimes n}}$. \square

We recall a few properties of Abelian schemes. By an Abelian scheme over \mathcal{O}_K we mean a smooth proper group scheme over \mathcal{O}_K whose fibers are geometrically connected (in particular \mathcal{G} is of finite presentation over \mathcal{O}_K and \mathcal{G}_K and $\mathcal{G}_{\mathbf{k}}$ are integral). We wish to show that if \mathcal{A} is an Abelian scheme over \mathcal{O}_K then $\mathcal{A}(\mathcal{O})$ is generically stable.

Fact 3.19. *Let \mathcal{V} be a scheme which is flat over \mathcal{O}_F and assume that \mathcal{V}_F is geometrically integral, then \mathcal{V} and $\mathcal{V} \times_{\text{Spec } \mathcal{O}_F} \text{Spec } \mathcal{O}_K$ are integral schemes.*

Proof. We may assume that $\mathcal{V} = \text{Spec } A$ is affine. Since A is flat over \mathcal{O}_F ,

$$A \otimes_{\mathcal{O}_F} \mathcal{O}_K \hookrightarrow A \otimes_{\mathcal{O}_F} K.$$

We also have $A \otimes_{\mathcal{O}_F} K \cong A \otimes_{\mathcal{O}_F} (F \otimes_F K) \cong (A \otimes_{\mathcal{O}_F} F) \otimes_F K$, and since the latter is an integral domain (by assumption), $A \otimes_{\mathcal{O}_F} \mathcal{O}_K$ is an integral domain.

Likewise, since A is flat over \mathcal{O}_F ,

$$A \otimes_{\mathcal{O}_F} \mathcal{O}_F \hookrightarrow A \otimes_{\mathcal{O}_F} \mathcal{O}_K,$$

so A is an integral domain as well. \square

Lemma 3.20. *Let \mathcal{G} be an integral separated group scheme of finite type over \mathcal{O}_F , with \mathcal{G}_F and \mathcal{G}_k geometrically integral. Then*

- (1) $\mathcal{G}(\mathcal{O})$ is a connected generically stable group.
- (2) If p is a K -definable type concentrated on $\mathcal{G}(\mathcal{O})$ and there exists an affine open subscheme $\mathcal{U} \subseteq \mathcal{G}$ for which p is concentrated on $\mathcal{U}(\mathcal{O})$ and $\mathcal{U}_{\mathcal{O}_K} \cong \Phi(\mathcal{U}_K, p)$ then p is the unique generic type of $\mathcal{G}(\mathcal{O})$.

Proof. By Fact 3.19, we may base change to K ; so we assume that $F = K$.

(1) By [Hal21, Proposition 5.1.6], $\mathcal{G}(\mathcal{O})$ is a generically stable group, i.e. has a generically stable K -definable generic type q . Furthermore, by its proof r_*q is a generic of \mathcal{G}_k and q is stably dominated by r_*q via r . The map r is surjective by [Hal21, Theorem 3.2.4]. For any $g \in \mathcal{G}(\mathcal{O})$, $r_*(gq) = r_*(g)r_*q = r_*q$ by integrality of \mathcal{G}_k so by [HRK19, Lemma 4.9] $gq = q$; i.e. $\mathcal{G}(\mathcal{O})$ is connected.

(2). By [Hal21, Theorem 5.2.2], \mathcal{U} has the mmp w.r.t. q ; by the assumption it also has the mmp w.r.t. p . Thus for every open affine subscheme $\mathcal{W} \subseteq \mathcal{U}$ with $\mathcal{W}(\mathcal{O}) \neq \emptyset$, both types are concentrated on $\mathcal{W}(\mathcal{O})$. By [Hal21, Lemma 4.3.7], $p = q$. □

Let \mathcal{A} be an integral Abelian scheme over \mathcal{O}_K . By Lemma 3.20, $\mathcal{A}(\mathcal{O}) = \mathcal{A}_K$ is a connected generically stable group with unique generic type p ; so by [Hal21, Theorem 5.2.2], \mathcal{A} has the mmp w.r.t. p . By [Hal21, Theorem 3.2.4], $r : \mathcal{A}_K \rightarrow \mathcal{A}_k$ is a surjective group homomorphism.

Fact 3.21. *Every Abelian scheme over a valuation ring R is projective over R .*

Proof. The result will follow by [Ray70, Theorem XI.1.4] once we note that the scheme $\text{Spec } R$ is geometrically unibranch [Sta23, Tag 0BQ2] which is straightforward since R is a valuation ring. □

Proposition 3.22. *Let \mathcal{A} be an integral Abelian scheme over \mathcal{O}_K and p its unique generically stable generic type. Then p satisfies (\dagger) of Corollary 3.17.*

Proof. Consider the commutative diagram supplied by Lemma 3.18. Any affine open subscheme $\mathcal{U} \subseteq \mathcal{A}$ with $\mathcal{U}(\mathcal{O}) \neq \emptyset$ has the mmp w.r.t. p , so by Lemma 3.11, r_*p is the generic type of \mathcal{A}_k . Let $c \models p|K$ and let $d \in \mathcal{A}_K = \mathcal{A}(\mathcal{O})$ with $f(c) = f(d)$; so $f(d) \models p_{\mathcal{O}}^{\otimes n}$. Since \bar{f} is a dominant morphism, necessarily $r(d) \models r_*p|K$.

By [Hal21, Lemma 4.1.6], p is stably dominated by r_*p via the surjective group homomorphism r . By [HRK19, Lemma 4.9], and connectedness, we conclude that $d \models p|K$, as needed. □

4. GETTING A GROUP SCHEME

Let F be a gracious valued field and $K \supseteq F$ an algebraically closed valued field.

In [HRK19, Theorem 6.11], Hrushovski and Rideau-Kikuchi proved that given an *affine* algebraic group G over K and a generically stable type p concentrating on G satisfying $p^2 = p$ there exists an affine group scheme \mathcal{G} over \mathcal{O} and a pro-definable isomorphism $G \cong \mathcal{G}_K$ under which $\text{Stab}(p)$ is mapped to $\mathcal{G}(\mathcal{O})$.

We still do not have a full analog for general algebraic groups over K . The purpose of this section is to answer this to the affirmative when the functor Φ (locally) gives a scheme of finite type over \mathcal{O}_K . We then apply this to give a model theoretic criterion for good reduction of (some) Abelian varieties over K .

We review some definitions and results from [BLR90]. Let \mathcal{S} be any scheme.

Definition 4.1. An open subscheme \mathcal{U} of a scheme \mathcal{X} over \mathcal{S} is called \mathcal{S} -dense if $\mathcal{U} \times_{\mathcal{S}} \mathcal{S}'$ is schematically dense in $\mathcal{X} \times_{\mathcal{S}} \mathcal{S}'$ for all morphisms $\mathcal{S}' \rightarrow \mathcal{S}$.

Fact 4.2. [DG67, IV.11.10.10] *If $\mathcal{X} \rightarrow \mathcal{S}$ is flat and of finite presentation then an open subscheme \mathcal{U} of \mathcal{X} is \mathcal{S} -dense if and only if for all $s \in \mathcal{S}$, the fiber $\mathcal{U}_s := \mathcal{U} \times_{\mathcal{S}} \kappa(s)$ is schematically dense in \mathcal{X}_s , where $\kappa(s)$ is the residue class field of s in \mathcal{S} .*

Note that in EGA such denseness is referred to as *universally schematically dense relative to \mathcal{S}* and in SGA3 as *schematically dense in \mathcal{X} relative to \mathcal{S}* . Furthermore, in [BLR90, Section 2.5] \mathcal{S} -denseness is only defined for smooth \mathcal{S} -schemes and in return they only require that \mathcal{U}_s be Zariski dense in \mathcal{X}_s . The reason for this is the following.

Fact 4.3. *Assume that \mathcal{X} is smooth and finitely presented over \mathcal{S} and let $\mathcal{U} \subseteq \mathcal{X}$ be an open subscheme. Then \mathcal{U} is \mathcal{S} -dense if and only if \mathcal{U}_s is Zariski dense in \mathcal{X}_s for all $s \in \mathcal{S}$.*

In particular, if furthermore for every $s \in \mathcal{S}$, \mathcal{X}_s is irreducible and \mathcal{U}_s is not the empty scheme, then \mathcal{U} is \mathcal{S} -dense.

Proof. As $\mathcal{X} \rightarrow \mathcal{S}$ is flat and of finite presentation, if \mathcal{U} is \mathcal{S} -dense then each \mathcal{U}_s is schematically dense in \mathcal{X}_s by Fact 4.2.

Assume that \mathcal{U}_s is Zariski dense in \mathcal{X}_s for all $s \in \mathcal{S}$. In particular, for any such $s \in \mathcal{S}$, \mathcal{X}_s is a smooth scheme over the field $\kappa(s)$ and hence reduced so being Zariski dense is equivalent to being schematically dense by [Sta23, Tag 056D]. \square

By an \mathcal{S} -rational map $\mathcal{X} \dashrightarrow \mathcal{Y}$ we mean an equivalence class of \mathcal{S} -morphisms $\mathcal{U} \rightarrow \mathcal{Y}$ where \mathcal{U} is some \mathcal{S} -dense open subscheme of \mathcal{X} . An \mathcal{S} -rational map $\varphi : \mathcal{X} \dashrightarrow \mathcal{Y}$ is called \mathcal{S} -birational if φ can be defined by an \mathcal{S} -morphism $\mathcal{U} \rightarrow \mathcal{Y}$ which induces an isomorphism from \mathcal{U} onto an \mathcal{S} -dense open subscheme of \mathcal{Y} , see [BLR90, Section 2.5] for more information.

Definition 4.4. Let \mathcal{S} be a scheme and \mathcal{X} a smooth separated \mathcal{S} -scheme finitely presented and faithfully flat over \mathcal{S} . An \mathcal{S} -birational group law on \mathcal{X} is an \mathcal{S} -rational map

$$m : \mathcal{X} \times_{\mathcal{S}} \mathcal{X} \dashrightarrow \mathcal{X}, \quad (x, y) \mapsto xy,$$

such that

(a) the \mathcal{S} -rational maps

$$\Psi_1 : \mathcal{X} \times_{\mathcal{S}} \mathcal{X} \dashrightarrow \mathcal{X} \times_{\mathcal{S}} \mathcal{X}, \quad (x, y) \mapsto (x, xy),$$

$$\Psi_2 : \mathcal{X} \times_{\mathcal{S}} \mathcal{X} \dashrightarrow \mathcal{X} \times_{\mathcal{S}} \mathcal{X}, \quad (x, y) \mapsto (xy, y),$$

are \mathcal{S} -birational, and

(b) m is associative; i.e. $(xy)z = x(yz)$ whenever both sides are defined.

As in Weil's original group chunk theorem an \mathcal{S} -birational group law gives rise to a group scheme:

Fact 4.5. [BLR90, Theorem 6.6.1] *Let \mathcal{S} be a scheme, and let m be an \mathcal{S} -birational group law on a smooth and separated \mathcal{S} -scheme \mathcal{X} which is finitely presented and faithfully flat over \mathcal{S} . Then there exists a smooth and separated \mathcal{S} -group scheme $\overline{\mathcal{X}}$ of finite presentation with a group law \overline{m} together with an \mathcal{S} -dense open subscheme $\mathcal{X}' \subseteq \mathcal{X}$ and an open immersion $\mathcal{X}' \hookrightarrow \overline{\mathcal{X}}$ having \mathcal{S} -dense image such that \overline{m} restricts to m on \mathcal{X}' .*

The group scheme $\overline{\mathcal{X}}$ is unique scheme satisfying the above, up to canonical isomorphism.

Lemma 4.6. *Let $(G, p) \in (\text{SPVar}/F)$ with G an algebraic group over F and assume that $p \otimes p$ and $p \otimes p \otimes p$ are strictly based on F and that $p^2 = p$. For any affine open subvariety $V \subseteq G$, for which $\Phi(V, p)$ is of finite type over \mathcal{O}_F ,*

$$\Phi(m) : \Phi(V, p) \times_{\mathcal{O}_F} \Phi(V, p) \dashrightarrow \Phi(V, p),$$

is an \mathcal{O}_F -birational group law, where $m : V \times V \dashrightarrow V$ is the rational map given by the group multiplication on G .

Proof. By Corollary 3.6, we may assume that $\Phi(V, p)$ is smooth and of finite type over \mathcal{O}_F . Thus $\Phi(V, p)$ is faithfully flat over \mathcal{O}_F and finitely presented over \mathcal{O}_F .

Let $m : G \times_F G \rightarrow G$, $(x, y) \mapsto xy$, be the multiplication map on G and consider the isomorphisms of varieties

$$\psi_1 : G \times_F G \rightarrow G \times_F G, (x, y) \mapsto (x, xy)$$

and

$$\psi_2 : G \times_F G \rightarrow G \times_F G, (x, y) \mapsto (xy, y);$$

note that $m_*(p \otimes p) = p$, $(\psi_1)_*(p \otimes p) = p \otimes p$ and $(\psi_2)_*(p \otimes p) = p \otimes p$ since $p^2 = p$.

Consider $W = \psi_1^{-1}(V \times V) \cap \psi_2^{-1}(V \times V) \cap (V \times V)$, it is an open affine subvariety of $V \times V$ since ψ_1, ψ_2 are isomorphisms and $G \times G$ is separated (by [Sta23, Tag 01KU]). We now restrict ψ_1 and ψ_2 to W : $\psi_1 : W \rightarrow V \times V$ and $\psi_2 : W \rightarrow V \times V$, so they are now open immersions.

As $F[V \times V]^{p \otimes p} \cong F[V]^p \otimes_{\mathcal{O}_F} F[V]^p$ (by [Hal21, Proposition 4.2.18]), we get that $\Phi(V \times V, p \otimes p) = \mathcal{V} \times_{\mathcal{O}_F} \mathcal{V}$, for $\mathcal{V} = \Phi(V, p)$. Thus, by applying Lemma 2.7 twice, we are supplied with $f \in F[V \times V]^{p \otimes p}$ and $g_1, g_2 \in F[V \times V]^{p \otimes p}$, with $p \otimes p \vdash \text{val}(f(x)) = \text{val}(g_1(x)) = \text{val}(g_2(x)) = 0$, such that, for $\mathcal{U} := D_{\mathcal{V} \times \mathcal{V}}(f)$,

$$\Phi(\psi_1) : \mathcal{U} \rightarrow D_{\mathcal{V} \times \mathcal{V}}(g_1) \text{ and } \Phi(\psi_2) : \mathcal{U} \rightarrow D_{\mathcal{V} \times \mathcal{V}}(g_2)$$

are isomorphisms. Since $p \otimes p \vdash \text{val}(f(x)) = 0$, $\Phi(D_{V \times V}(f), p \otimes p) = D_{\mathcal{V} \times \mathcal{V}}(f)$ and so is faithfully flat over \mathcal{O}_F .

Note that $\mathcal{V} \times_{\mathcal{O}_F} \mathcal{V}$ is smooth finitely presented and flat over \mathcal{O}_F , and thus so is \mathcal{U} , and all the fibers of $\mathcal{V} \times_{\mathcal{O}_F} \mathcal{V} \rightarrow \text{Spec } \mathcal{O}_F$ are (geometrically) integral (Proposition 3.3). In order to conclude, that \mathcal{U} is \mathcal{O}_F -dense in $\mathcal{V} \times_{\mathcal{O}_F} \mathcal{V}$ we need it, by Fact 4.3, to meet all the fibers of $\mathcal{V} \times_{\mathcal{O}_K} \mathcal{V} \rightarrow \text{Spec } \mathcal{O}_F$. But this is automatic since \mathcal{U} is faithfully flat over \mathcal{O}_F .

Thus $\Phi(\psi_1) : \mathcal{V} \times_{\mathcal{O}_F} \mathcal{V} \dashrightarrow \mathcal{V} \times_{\mathcal{O}_F} \mathcal{V}$ and $\Phi(\psi_2) : \mathcal{V} \times_{\mathcal{O}_F} \mathcal{V} \dashrightarrow \mathcal{V} \times_{\mathcal{O}_F} \mathcal{V}$ are \mathcal{O}_F -birational by the paragraph after [BLR90, Definition 1, Section 5.1].

Now, by letting π_i be the projection on the i -th coordinate, we get that $\pi_2 \circ \Phi(\psi_1) = \pi_1 \circ \Phi(\psi_2)$. Setting $\tilde{m} = \pi_2 \circ \Phi(\psi_1)$ we get an \mathcal{O}_F -rational map $\mathcal{V} \times_{\mathcal{O}_F} \mathcal{V} \dashrightarrow \mathcal{V}$ which gives an \mathcal{O}_F -birational group law (getting associativity is easy, by further restricting $D_{V \times V}(f)$ in the beginning). \square

Proposition 4.7. *Let $(G, p) \in (SPVar/F)$ with G an algebraic group over F , with p satisfying $p^2 = p$ and $p \otimes p, p \otimes p \otimes p$ strongly based on F . Further assume there exists an open subvariety $V \subseteq G$ for which $\Phi(V, p)$ is of finite type over \mathcal{O}_F . Then there exists a smooth integral separated \mathcal{O}_F -group scheme \mathcal{G} of finite presentation over \mathcal{O}_F with geometrically integral fibers and an isomorphism $G \cong \mathcal{G}_F$ such that under this isomorphism $\text{Stab}(p)$ is isomorphic to $\mathcal{G}(\mathcal{O})$.*

Moreover, there is some $f \in F[V]$, such that $\Phi(D_V(f), p)$ is isomorphic to an open subscheme of \mathcal{G} .

Proof. By Lemma 4.6 and Fact 4.5 there exists a smooth separated \mathcal{O}_F -group scheme \mathcal{G} of finite presentation and faithfully flat over \mathcal{O}_F together with an \mathcal{O}_F -dense open subscheme $\mathcal{U} \subseteq \Phi(V, p)$ and an open immersion $\mathcal{U} \rightarrow \mathcal{G}$ having \mathcal{O}_F -dense image. Let $\mathcal{V} = \Phi(V, p)$.

Since \mathcal{U} is \mathcal{O}_F -dense in \mathcal{V} , and $\mathcal{V} \rightarrow \text{Spec } \mathcal{O}_F$ has geometrically integral fibers, $\mathcal{U}_{\mathbf{k}}$ is Zariski dense in $\mathcal{V}_{\mathbf{k}}$. The latter is non-empty and so $\mathcal{U}_{\mathbf{k}}$ is non-empty as well. Thus there exists an element $f \in F[V]^p$ such that $D_{\mathcal{V}}(f) \subseteq \mathcal{U}$ and $D_{\mathcal{V}}(f)_{\mathbf{k}}$ is non-empty (and thus $D_{\mathcal{V}}(f)_{\mathbf{k}}(\mathbf{k})$ is non empty). Replace \mathcal{U} by $D_{\mathcal{V}}(f)$.

Since $\Phi(V, p)$ is an integral scheme and flat dominant over \mathcal{O}_F so is \mathcal{U} . By Fact 3.10, $\mathcal{U}(\mathcal{O}) \neq \emptyset$ so $\mathcal{U}_{\mathcal{O}_K}$ is faithfully flat over \mathcal{O}_K by [Hal21, Lemma 3.1.4]. By faithfully flat descent \mathcal{U} is faithfully flat over \mathcal{O}_F as well. Furthermore, by the mmp w.r.t. p , p is then concentrated on $\mathcal{U}(\mathcal{O})$ so $p \vdash \text{val}(f(x)) = 0$. Consequently, $(F[V]^p)_f = (F[V]_f)^p$, i.e. $\Phi(\mathcal{U}_F, p) = \mathcal{U}$.

By applying the uniqueness clause of Fact 4.5 to the variety V over F together with the F -birational group law inherited from F , we deduce that G is isomorphic to \mathcal{G}_F since $\mathcal{V}_F = V$. We now identify between \mathcal{G}_F and G .

Identify \mathcal{U} with its image inside \mathcal{G} . Since \mathcal{U} is an integral open subscheme of the group scheme \mathcal{G} , \mathcal{G} is integral as well. Since $\mathcal{U} \rightarrow \text{Spec } \mathcal{O}_F$ has geometrically integral fibers (by Proposition 3.3) and $\mathcal{U}_{\mathfrak{k}} \neq \emptyset$ is an open subvariety of the group scheme $\mathcal{G}_{\mathfrak{k}}$ the latter is an algebraic group (i.e. geometrically integral). Likewise, all the fibers are algebraic groups.

By Lemma 3.20, $\mathcal{G}(\mathcal{O})$ is generically stable with p its unique generic type; so $\text{Stab}(p) = \mathcal{G}(\mathcal{O})$. \square

Theorem 4.8. *Let A be an Abelian variety over K and assume that A is a generically stable group with a unique generic type p . Then the following are equivalent*

- (1) *There exists an open affine subvariety $V \subseteq A$ such that $\Phi(V, p)$ is of finite type over \mathcal{O}_K .*
- (2) *There exists an integral Abelian scheme \mathcal{A} over \mathcal{O}_K with $A \cong \mathcal{A}_K$.²*
- (3) *p satisfies (\dagger) of Corollary 3.17.*

Proof. (1) \implies (2). By Proposition 4.7, there exists a smooth integral separated group scheme \mathcal{A} over \mathcal{O}_K which is of finite presentation over \mathcal{O}_K , with integral fibers, satisfying $A \cong \mathcal{A}_K$. Since $\text{Stab}(p) = A$ it follows that $\mathcal{A}(\mathcal{O}) = \mathcal{A}_K$. As A is an Abelian variety, it is projective and so proper over K , we can thus conclude by [Hal21, Proposition 5.3.4] that \mathcal{A} is universally closed over \mathcal{O}_K so \mathcal{A} is proper over \mathcal{O}_K . Since all the fibers of $\mathcal{A} \rightarrow \text{Spec } \mathcal{O}_K$ are geometrically irreducible, \mathcal{A} is an Abelian scheme over \mathcal{O}_K .

(2) \implies (3). This is Proposition 3.22.

(3) \implies (1). This is Proposition 3.16. \square

In Theorem 5.8, we will show that the generic type of the maximal generically stable subgroup of an elliptic curve over K always satisfies (3).

Question 4.9. For A and p as in Theorem 4.8. Does p always satisfy (3)?

More generally, if a maximal (connected) generically stable subgroup of A exists, does its generic type satisfy (3)?

5. ELLIPTIC CURVES

Let F be a gracious valued field and K a model of ACVF containing F .

Let E be an elliptic curve over F , i.e. a smooth projective curve of genus 1, which we see as a closed subscheme of \mathbb{P}_F^2 given by a Weierstrass equation

$$y^2z + a_1xyz + a_3yz^2 = x^3 + a_2x^2z + a_4xz^2 + a_6z^3.$$

It is automatically geometrically integral and separated [You]. Note that $E = D_{+,E}(z) \cup D_{+,E}(y)$. The group $E(F)$ is Abelian; we denote its multiplication by \cdot and by e its identity element (which is the unique point at infinity).

After a suitable change of coordinates, we may assume that $\text{val}(a_i) \geq 0$ for $i = 1, 2, 3, 4, 6$. As E is a curve, every generically stable type concentrated on E is strongly stably dominated ([HL16, Lemma 8.1.4(1)]) and by Lemma 3.14, every generically stable type concentrated on E is a translate of a generic of a connected generically stable subgroup of E . Recall that by Fact 3.9, every type-definable generically stable subgroup of E is definable.

For the following, recall that we denote by $p_{\mathcal{O}}$ the unique generic of the closed ball \mathcal{O} .

² \mathcal{A} is *probably* the Néron model of A over \mathcal{O}_K , see the unpublished result [CH].

Lemma 5.1. *For any gracious valued field $L \supseteq F$, $x \models p_{\mathcal{O}}|L$ and y an element satisfying that*

$$y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6,$$

we have that

(1) $\text{val}(y) \geq 0$ and $\{x^i y^j\}_{i \geq 0, j=0,1}$ constitutes a separating basis of $L(x, y)$ over L .

As a result,

(i) *The collection of formulas stating that $x \models p_{\mathcal{O}}$ and*

$$y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$$

determines a unique generically stable F -definable type $p(x, y)$.

(ii) *p satisfies (\dagger) of Proposition 3.16.*

(iii) *$p^{\otimes n}$ is strongly based on F , for any $n \geq 1$.*

Proof. The proof of (1) is similar to the proof of Proposition 3.16. As $x \models p_{\mathcal{O}}|L$, x constitutes a separating transcendence basis for $L(x)$ over L ; by [Kuh, Theorem 1.1] $L(x)$ is a defectless valued field. By the choice of y , $L(x, y)/L(x)$ is an extension of valued fields of degree 2. As $\text{val}(a_i) \geq 0$ for $i = 1, 2, 3, 4, 6$ necessarily $\text{val}(y) \geq 0$ (for example, by [Kuh, Lemma 5.6(a)]); by the choice of y , $\mathbf{k}_L(\text{res}(x), \text{res}(y))$ is an extension of degree of 2 of $\mathbf{k}_{L(x)} = \mathbf{k}_L(\text{res}(x))$ and so $\text{val}(y) = 0$. By the fundamental inequality for extension of valued fields we have $\mathbf{k}_L(\text{res}(x), \text{res}(y)) = \mathbf{k}_{L(x,y)}$, and the result follows.

(i) A curve satisfying $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$ must be irreducible [You]; thus applying (1) to a model L containing F we conclude, by quantifier elimination in ACVF, that these formulas determine a unique generically stable type (indeed, it must be orthogonal to Γ). It is clearly F -definable.

(ii) This is a rephrasing of (i) with the finite morphism being to projection on the x -coordinate.

(iii) This follows from (1). \square

Let \mathcal{V} be the closed subscheme of $\mathbb{A}_{\mathcal{O}_F}^2$ (in variables x, y) given by $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$, and let $V = \mathcal{V}_F = D_{+,E}(z)$. By Lemma 5.1, \mathcal{V} has the mmp w.r.t. p , where p is the type from Lemma 5.1.

The following is straightforward (using Lemma 5.1).

Corollary 5.2. *We have $\mathcal{V} \cong \Phi(V, p)$ and for any $\mathfrak{p} \in \text{Spec } \mathcal{O}_F$, the fiber $\Phi(V, p)_{\mathfrak{p}}$ is the reduction of \mathcal{V} to \mathbf{k}_w , where w is the coarsening of $\text{val} \upharpoonright F$ corresponding to \mathfrak{p} .*

The identity element of $E(K)$ is the only K -point not in $V(K)$. The multiplication law, see e.g. [Sil09, Page 59], on points $(x_1, y_1), (x_2, y_2) \in V$ with $(x_2, y_2) \neq (x_1, y_1)^{\pm 1}$ is given by $(x_3, y_3) = (x_1, y_1) \cdot (x_2, y_2)$, where

$$x_3 = \lambda^2 + a_1\lambda - a_2 - x_1 - x_2,$$

$$y_3 = -(\lambda + a_1)x_3 - \nu - a_3$$

and

$$\lambda = \frac{y_2 - y_1}{x_2 - x_1}, \nu = \frac{y_1x_2 - y_2x_1}{x_2 - x_1}.$$

For the inverse,

$$(x_1, y_1)^{-1} = (x_1, -y_1 - a_1x_1 - a_3).$$

We note that this implies that for any type q concentrated on $\mathcal{V}(\mathcal{O})$, q^{-1} is also concentrated on $\mathcal{V}(\mathcal{O})$.

Lemma 5.3. *Let q be any generically stable K -definable type concentrated on V with $pq = q$, then q is concentrated on $\mathcal{V}(\mathcal{O})$.*

Proof. Let $d \models q|K$ and let $c \models p|Kd$. For any element $g \in V$ write $g = (g_x, g_y)$.

We first make some observations. If $\text{val}(d_x) \geq 0$ then d_y is a root of a monic polynomial over \mathcal{O}_K so $\text{val}(d_y) \geq 0$ ([Kuh, Lemma 5.6(a)]) and we are done; so we assume towards a contradiction that $\text{val}(d_x) < 0$.

Next note that $\text{val}(d_y) \leq \text{val}(d_x)$; otherwise since $a_1xy = x^3 + a_2x^2 + a_4x - y^2 - a_3y$ we would have $\text{val}(a_1d_xd_y) = 3\text{val}(d_x)$ and so $\text{val}(a_1) + \text{val}(d_y) = 2\text{val}(d_x)$, contradiction.

Now we note that $\text{val}(d_x^3) \leq \text{val}(d_y^2)$; otherwise as above we would get that $\text{val}(a_1) + \text{val}(d_x) = \text{val}(d_y)$ which contradicts the above.

Thus $\text{val}(d_y) \leq \text{val}(d_x) < 0$ and $\text{val}(d_x^3) \leq \text{val}(d_y^2)$; in particular $\text{val}(d_x^2) < \text{val}(d_y)$.

By the multiplication law, if $g = c \cdot d$ then

$$g_x = \frac{(d_y^2 + a_1d_yd_x - a_2d_x^2 - d_x^3) + (-a_1d_y + 2a_2d_x + d_x^2)c_x + (-2d_y - a_1d_x)c_y}{d_x^2 - 2d_xc_x + c_x^2} + \frac{a_1c_yc_x + (-a_2 + d_x)c_x^2 + c_y^2 - c_x^3}{d_x^2 - 2d_xc_x + c_x^2}$$

By Lemma 5.1(1), and since $c \models p|Kd$ we get

$$\begin{aligned} \text{val}(g_x) &= \min\{\text{val}(d_y^2 + a_1d_yd_x - a_2d_x^2 - d_x^3), \text{val}(-a_1d_y + 2a_2d_x + d_x^2), \\ &\text{val}(-2d_y - a_1d_x), \text{val}(a_1), \text{val}(d_x - a_2), 0\} - \min\{\text{val}(d_x^2), \text{val}(-2d_x), 0\}. \end{aligned}$$

Since (d_x, d_y) is a point on the elliptic curve, $\text{val}(d_y^2 + a_1d_yd_x - a_2d_x^2 - d_x^3) = \text{val}(-a_3d_y + a_4d_x + a_6) \geq \text{val}(d_y)$; obviously $\text{val}(-2d_y - a_1d_x) \geq \text{val}(d_y)$ and as $\text{val}(d_x^2) < \text{val}(d_y)$ we have $\text{val}(-a_1d_y + 2a_2d_x + d_x^2) = \text{val}(d_x^2)$. Hence, $\text{val}(g_x) = \text{val}(d_x^2) - \text{val}(d_x^2) = 0$; but we assumed that $pq = q$ so $\text{val}(g_x) = \text{val}(d_x)$, contradiction.

We conclude that $\text{val}(d_x), \text{val}(d_y) \geq 0$. \square

Lemma 5.4. *Let q be a generically stable K -definable type concentrated on $\mathcal{V}(\mathcal{O})$ then $pq = p$.*

Proof. By base-changing everything to a maximally complete ACVF containing K and since the conclusion descends, there is no harm in assuming K is maximally complete.

Let $d \models q|K$ and $c \models p|Kd$. For any element $g \in V$ we write $g = (g_x, g_y)$. As q is concentrated on $\mathcal{V}(\mathcal{O})$, we have $\text{val}(d_x), \text{val}(d_y) \geq 0$. Moreover, as p is concentrated on $\mathcal{V}(\mathcal{O})$, p^{-1} is also concentrated on $\mathcal{V}(\mathcal{O})$.

Claim. (1) $\text{val}(d_x - c_x) = 0$
(2) $\text{val}(d_x - (cd^{-1})_x) = 0$
(3) pq and pq^{-1} are concentrated on $\mathcal{V}(\mathcal{O})$.

Proof. (1) As $c \models p|Kd$ and $\text{val}(c_x) = 0$, by Lemma 5.1(1) we get $\text{val}(d_x - c_x) = \min\{\text{val}(d_x), 0\} = 0$.

(2) Note that $(cd^{-1})_x = \lambda^2 + a_1\lambda - a_2 - c_x - d_x$, where $\lambda = \frac{-d_y - a_1d_x - a_3 - c_y}{d_x - c_x}$. After taking common denominator, since $\text{val}(a_1), \text{val}(a_3), \text{val}(d_x), \text{val}(d_y) \geq 0$, as in (1) (and using (1)), we conclude.

(3) Since q and p are concentrated on $\mathcal{V}(\mathcal{O})$, so is pq by (1) and the formulas for multiplication. We can also apply the argument for pq^{-1} . \square (claim)

Let $f \in K[V]$ be some regular function. By the mmp of \mathcal{V} w.r.t. p and item (3) of the claim, $\text{val}(f(c)) \leq \text{val}(f(cd))$, we now show the other direction; since f is arbitrary this will give the desired conclusion that $pq = p$.

By the multiplication law on E , there exist $f_i, g_i \in K[V]$ and some integer n such that for any $a, b \in V$, with $a \neq b^{\pm 1}$, we have

$$f(a \cdot b) = \frac{\sum_i f_i(a)g_i(b)}{(b_x - a_x)^n}.$$

As K is maximally complete and $(c, d) \models (p \otimes q)|K$, by [HHM08, Lemma 12.4] (see also [Hal21, Proposition 4.2.17]), f_i and g_i may be chosen such that $\text{val}(\sum_i f_i(c)g_i(d)) = \min_i \{\text{val}(f_i(c))\}$ and $\text{val}(g_i(d)) = 0$.

We can now use the above claim and the fact that $\text{val}(f_i(c)) \leq \text{val}(f_i(cd^{-1}))$ (by the mmp of \mathcal{V} w.r.t. p and the fact that pq^{-1} is concentrated on $\mathcal{V}(\mathcal{O})$) to compute

$$\begin{aligned} \text{val}(f(cd)) &= \text{val}\left(\frac{\sum_i f_i(c)g_i(d)}{(d_x - c_x)^n}\right) = \text{val}\left(\sum_i f_i(c)g_i(d)\right) = \\ &= \min_i \{\text{val}(f_i(c))\} \leq \min_i \{\text{val}(f_i(cd^{-1}))\} = \min_i \{\text{val}(f_i(cd^{-1})g_i(d))\} \\ &\leq \text{val}\left(\sum_i f_i(cd^{-1})g_i(d)\right) = \text{val}\left(\frac{\sum_i f_i(cd^{-1})g_i(d)}{(d_x - (cd^{-1})_x)^n}\right) = \text{val}(f(c)), \end{aligned}$$

as required. \square

Proposition 5.5. *The type p is a generic of a connected definable subgroup of E ; moreover we have $pq = p$ for any generically stable generic q of a connected definable subgroup of E . As a result, for any connected definable generically stable subgroup H of E we have $H \subseteq \text{Stab}(p)$: p is the generic of the maximal generically stable (type-)definable subgroup of E .*

Proof. As p is concentrated on $\mathcal{V}(\mathcal{O})$, by Lemma 5.4 we know that $p^2 = p$, so p is the unique generic type of $\text{Stab}(p)$. The latter is definable since p is strongly stably dominated.

Let q be any generically stable generic of a connected (type-)definable subgroup of E . As E is Abelian $pq = (pq)^2$ is also a generically stable generic (of a type-definable subgroup). Since $p(pq) = pq$, by Lemma 5.3 pq is concentrated on $\mathcal{V}(\mathcal{O})$. Then by Lemma 5.4, $p(pq) = p$ so $pq = p$; thus $\text{Stab}(q) \subseteq \text{Stab}(p)$. \square

Remark 5.6. By [HRK19, Corollary 6.19], there exists a (necessarily unique) K -definable generically stable type p_∞ concentrated on E satisfying that $p_\infty^2 = p_\infty$ and that $qp_\infty = p_\infty$ for any other generically stable K -definable type q concentrated on E satisfying $q^2 = q$, i.e. a generic type of a generically stable subgroup.³ Proposition 5.5 gives a direct proof of this fact in this situation.

We will also require the following folklore result.

Fact 5.7. *Let $f : \mathcal{X} \rightarrow \mathcal{S}$ be a flat, separated of finite presentation morphism. If f has proper and geometrically connected fibers then it is proper.*

Proof. In the case when \mathcal{S} is locally Noetherian and f is of finite type this is [DG67, IV.15.7.10]. However, if we assume that f is of finite presentation then the locally Noetherianity assumption may be dropped by Noetherian approximation. \square

Theorem 5.8. *Let E be an elliptic curve over F and let p be the generically stable generic of the maximal generically stable definable subgroup.*

There exists a smooth integral separated \mathcal{O}_F -group scheme \mathcal{E} of finite presentation over \mathcal{O}_F with geometrically integral fibers, satisfying $E \cong \mathcal{E}_F$ such that under this isomorphism $\text{Stab}(p) = \mathcal{E}(\mathcal{O})$, for which the following are equivalent:

³We have been informed by Rideau-Kikuchi by private communication that there is a problem with the published proof of [HRK19, Corollary 6.19]. However, for elliptic curves this problem can be circumvented.

- (1) $\text{Stab}(p) = \mathcal{E}_F = E$.
- (2) \mathcal{E} is proper over \mathcal{O}_F , and thus an Abelian scheme over \mathcal{O}_F .
- (3) (Assuming E is given by a Weierstrass equation over \mathcal{O}_F) $\text{val}(\Delta) = 0$, where Δ is the discriminant.

As a result, there exists a proper integral group scheme \mathcal{G} over \mathcal{O}_F with $\mathcal{G}_F \cong E$ if and only if $\text{Stab}(p) = \mathcal{G}_F$ if and only if $\text{val}(\Delta) = 0$ (when E is given by a Weierstrass equation over \mathcal{O}_F).

Remark 5.9. In the “as a result”, we can replace “there exists a proper integral group scheme \mathcal{G} over \mathcal{O}_F with $\mathcal{G}_F \cong E$ ” with “there exists a proper irreducible smooth curve with geometrically connected fibers all of genus one, with a given section in $\mathcal{G}(\mathcal{O}_F)$, satisfying $\mathcal{G}_F \cong E$ ”, since any such scheme must necessarily be a group scheme (by [KM85, Theorem 2.1.2]).

Proof. We may assume that E is given by (the projective closure of) the Weierstrass equation $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$, with

$$\text{val}(a_1), \text{val}(a_3), \text{val}(a_2), \text{val}(a_4), \text{val}(a_6) \geq 0.$$

By Lemma 5.1(ii), p satisfies (†) of Corollary 3.17 so the existence of \mathcal{E} is given by Proposition 4.7 (and Lemma 5.1(iii)) and Proposition 3.16.

(1) \iff (2). If $\text{Stab}(p) = E$ then, by [Hal21, Proposition 5.3.4], \mathcal{E} is universally closed over \mathcal{O}_F , so \mathcal{E} is proper over \mathcal{O}_F . As the fibers are geometrically integral, \mathcal{E} is an Abelian scheme over \mathcal{O}_F . For the other direction, if \mathcal{E} is proper over \mathcal{O}_F then $\text{Stab}(p) = \mathcal{E}(\mathcal{O}) = \mathcal{E}_F = E$.

(2) \iff (3). Since the assumption is preserved and the conclusion descends, there is no harm in base changing and assuming that $F = K$. We first make some observations. As \mathcal{E} is smooth over \mathcal{O}_K , for any $\mathfrak{p} \in \text{Spec } \mathcal{O}_K$, the fiber $\mathcal{E}_{\mathfrak{p}}$ is an algebraic group of dimension 1 over $\kappa(\mathfrak{p})$, i.e. is either \mathbb{G}_a , \mathbb{G}_m or an elliptic curve (note that these options are pair-wise not birational).

On the other hand, for the open affine subvariety $V \subseteq E$ from before, we have that $\Phi(V, p)$ is birational with \mathcal{E} (the “moreover” clause of Proposition 4.7). So for any $\mathfrak{p} \in \text{Spec } \mathcal{O}_K$, $\mathcal{E}_{\mathfrak{p}}$ is birational to $\Phi(V, p)_{\mathfrak{p}}$ which is just an open subvariety of the reduction of \mathcal{V} to $\kappa(\mathfrak{p})$ (see Corollary 5.2). Let w be the coarsening of val corresponding to \mathfrak{p} ; so $\mathcal{E}_{\mathfrak{p}}$ is an elliptic curve if and only if $w(\Delta) = 0$.

If \mathcal{E} is an Abelian scheme over \mathcal{O}_K then the special fiber must be an elliptic curve so $\text{val}(\Delta) = 0$. If on the other hand, $\text{val}(\Delta) = 0$ then $w(\Delta) = 0$ for any coarsening w of val , hence $\mathcal{E}_{\mathfrak{p}_w}$ is proper over \mathbf{k}_w for any such coarsening. Since all the fibers are (geometrically) integral they are all Abelian varieties. By Fact 5.7, \mathcal{E} is proper over \mathcal{O}_K , i.e. it is an Abelian scheme over \mathcal{O}_K .

For the final statement. The statement “ $\text{Stab}(p) = E$ is equivalent to $\text{val}(\Delta) = 0$ ” is given by the above arguments, so we show they are equivalent to the middle statement. If $\text{Stab}(p) = E$ then \mathcal{E} is a proper group scheme over \mathcal{O}_F with $\mathcal{E}_F \cong E$ (by (2)). For the other direction, there is no harm in assuming that $F = K$. Let \mathcal{G} be a proper integral group scheme over \mathcal{O}_K for which $\mathcal{G}_K = E$. By [Hal21, Proposition 5.1.6], $\mathcal{G}(\mathcal{O}) = \mathcal{G}_K = E$ is a generically stable group say with principal generic q . As $\text{Stab}(q)$ is an intersection of definable subgroups of E of finite index, [HRK19], and E is a divisible group, $\text{Stab}(q) = E$; thus $q = p$ as needed. \square

Remark 5.10. Let E be an elliptic curve over a fraction field over a DVR R given by a Weierstrass equation over R . This same equation defines a closed subscheme of \mathbb{P}_R^2 . It is known that the smooth locus of this scheme carries the structure of a smooth group scheme whose generic fiber is isomorphic to E [Sil94, Theorem IV.5.3]. It will be interesting to check if similar arguments give a similar result over a general valued field. Our motivation here was to show that the techniques of the previous sections have applications for some Abelian varieties.

Remark 5.11. It is well known that an elliptic curve over a fraction field of a DVR has a Néron model; this model is in general not proper. It is well known that an elliptic curve E over a DVR

has a proper Néron model if and only if it has good reduction (i.e. has some proper model). Given [CH], the theorem above can be seen as a generalization of this latter fact.

Remark 5.12. It is well known that an elliptic curve over an algebraically closed valued field might not have a Néron model, see e.g. [CH, The comment by user gdb].

Another argument is provided by [Hal21, Proposition 5.3.4]. Indeed, over a model of ACVF this cited result shows that a Néron model must be proper. So the existence of a Néron model would imply, using Theorem 5.8, that $\text{val}(\Delta) = 0$. But obviously this does not hold for all elliptic curves.

We finish with the following geometric interpretation of Proposition 5.5.

Proposition 5.13. *For any integral group scheme \mathcal{G} of finite type over \mathcal{O}_K with integral special fiber and satisfying $\mathcal{G}_K \cong E$, we have $\mathcal{G}(\mathcal{O}) \subseteq \mathcal{E}(\mathcal{O})$ (as subsets of E). Where \mathcal{E} is the group scheme supplied by Theorem 5.8.*

As a result, for any valued field extension $(K, \mathcal{O}_K) \leq (L, \mathcal{O}_L)$ we have $\mathcal{G}_{\mathcal{O}_L}(\mathcal{O}_L) \subseteq \mathcal{E}_{\mathcal{O}_L}(\mathcal{O}_L)$.

Proof. By Lemma 3.20, $\mathcal{G}(\mathcal{O})$ is a connected generically stable group with a unique generic type q , i.e. $\text{Stab}(q) = \mathcal{G}(\mathcal{O})$. By Proposition 5.5, $\mathcal{G}(\mathcal{O}) \subseteq \text{Stab}(p) = \mathcal{E}(\mathcal{O})$, where the latter equality is by Theorem 5.8.

The final statement is by model completeness. \square

Question 5.14. Is it true that for every generically stable type q on E there exists some affine open subvariety $V \subseteq E$ for which $\Phi(V, q)$ is of finite type over \mathcal{O}_K ?

APPENDIX A. GENERICALLY STABLE TYPES AND THE SHELAH EXPANSION

Let T be a complete NIP theory in a first order language \mathcal{L} and let \mathbb{U} be a large saturated model.

Definition A.1. A global type p is said to be *generically stable* if it is definable and finitely satisfiable in some small model M . We then say that p is *generically stable over M* .

Remark A.2. (1) For every set B for which a generically stable type p is B -invariant, p is the unique B -invariant extension of $p|_B$.
 (2) A global A -invariant type is generically stable if and only if every Morley sequence of p is totally indiscernible if and only if there exists a Morley sequence of p which is totally indiscernible.

Fact A.3. (1) *Let p be a generically stable type. Then p is still generically stable in any reduct.*

(2) *There is a unique expansion of any type p to T^{eq} . If p is generically stable then this expansion is still generically stable.*

Proof. (1) Assume that p is definable and finitely satisfiable in some small model M . Let \tilde{p} be the reduct of the type. Since \tilde{p} is still finitely satisfiable in M it is M -invariant. Now, we use the fact that every Morley sequence of p is totally indiscernible to show that there exists such a Morley sequence in \tilde{p} .

(2) Straightforward. \square

Let $M \models T$ be a small model and $M \prec N$ an $|M|^+$ -saturated model. The Shelah expansion of M , denoted by M^{sh} , is a structure with universe M and language $\mathcal{L}^{\text{sh}} = \mathcal{L} \cup \{R_{\varphi(x,c)} : \varphi(x,y) \in \mathcal{L}, c \in N\}$, where for every $a \in M$, $M^{\text{sh}} \models R_{\varphi(x,c)}(a) \iff N \models \varphi(a,c)$. The definable sets of M^{sh} do not depend on the $|M|^+$ -saturated model from which we take the parameters for the externally definable sets. Since T is NIP, $\text{Th}(M^{\text{sh}})$ eliminates quantifiers in \mathcal{L}^{sh} [She09].

By resplendency of saturated models, we may expand \mathbb{U} to a saturated model of $\text{Th}(M^{sh})$ in the language \mathcal{L}^{sh} . We denote this expansion by \mathbb{U}^* .

The following is probably well know but we could not find a proof of it anywhere, a version for measures appears in [BK19, Theorem 6.2].

Proposition A.4. *Let p be a global type. If p is definable and finitely satisfiable in $A \subseteq M$. Then there exists a unique extension $p \subseteq q \in S^{\mathcal{L}^{sh}}(\mathbb{U}^*)$ which is definable and finitely satisfiable in A .*

As a result, if p is generically stable over M then so is q .

Proof. Let $M^{sh} \prec L^*$ be an $|M|^+$ -saturated extension and $(L^*, M^{sh}) \prec (\mathbb{V}^*, \mathbb{U}^{**})$ a saturated extension of cardinality $|\mathbb{U}|$. By uniqueness of saturated models we may assume that $\mathbb{U}^{**} = \mathbb{U}^*$. Let $L, \mathbb{U}, \mathbb{V}$ be their reducts to \mathcal{L} , respectively.

As p is definable and finitely satisfiable in A , there exists an extension $\tilde{p} \in S(\mathbb{V})$ which is still definable and finitely satisfiable in A (and with the same definition). Seen as a partial type over \mathbb{V}^* , \tilde{p} is still finitely satisfiable in A . Let \tilde{p}^{sh} be a completion of \tilde{p} in $S^{\mathcal{L}^{sh}}(\mathbb{V}^*)$ which is still finitely satisfiable in A . We will show that $\tilde{p}^{sh}|_{\mathbb{U}^*}$ is definable over A .

For each $c \in N$ choose some $\tilde{c} \in L$ for which $\text{tp}(c/M) = \text{tp}(\tilde{c}/M)$. In particular, for each relation $R_{\varphi(x,c)}(x, y)$, $c \in N$, we have that

$$\varphi(M, c) = \varphi(M, \tilde{c}).$$

We first show the following:

Claim.

$$(\star) \quad R_{\varphi(x,y,c)}(x, b) \in \tilde{p}^{sh}|_{\mathbb{U}^*} \iff \varphi(x, b, \tilde{c}) \in \tilde{p}.$$

Proof. Assume that (\star) is not true. Thus there exists $b \in \mathbb{U}^*$ with

$$\neg R_{\varphi(x,y,c)}(x, b) \wedge \varphi(x, b, \tilde{c}) \in \tilde{p}^{sh}.$$

Since \tilde{p}^{sh} is finitely satisfiable in A there exists $m \in A \subseteq M^{sh}$ with

$$\mathbb{V}^* \models \neg R_{\varphi(x,y,c)}(m, b) \wedge \varphi(m, b, \tilde{c}),$$

but since we also have

$$(\mathbb{V}^*, \mathbb{U}^*) \models \forall (a_1, a_2) \in \mathbb{U}^* (R_{\varphi(x,y,c)}(a_1, a_2) \leftrightarrow \varphi(a_1, a_2, \tilde{c})),$$

we arrive to a contradiction. □ (claim)

We now show that $\tilde{p}^{sh}|_{\mathbb{U}^*}$ is definable over A . Since \tilde{p} is definable over A , we have $\varphi(x, b, \tilde{c}) \in \tilde{p} \iff \mathbb{V} \models \psi(b, \tilde{c})$ for some \mathcal{L} -formula over A . By the choice of \tilde{c} , we have that $\psi(M, \tilde{c}) = \psi(M, \tilde{c})$, thus

$$(L^*, M^{sh}) \models \forall a \in M^{sh} (R_{\psi(x,c)}(a) \leftrightarrow \psi(a, \tilde{c})).$$

Since $(L^*, M^{sh}) \prec (\mathbb{V}^*, \mathbb{U}^*)$, for all $a \in \mathbb{U}^*$

$$\mathbb{U}^* \models R_{\psi(x,c)}(a) \iff \mathbb{V}^* \models \psi(a, \tilde{c}).$$

Together with the claim this combines to

$$R_{\varphi(x,y,c)}(x, b) \in \tilde{p}^{sh}|_{\mathbb{U}^*} \iff \mathbb{U}^* \models R_{\psi(x,c)}(b).$$

Uniqueness follows from (\star) . □

REFERENCES

- [BK19] Gareth Boxall and Charlotte Kestner. Theories with distal shelah expansions. preprint, <https://arxiv.org/abs/1801.02346>, 2019.
- [BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud. *Néron models*, volume 21 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1990.
- [CH] Kestutis Cesnavičius and David Holmes. Do abelian varieties have neron models over arbitrary valuation rings? MathOverflow. URL:<https://mathoverflow.net/q/307819> (version: 2018-08-08).
- [DG67] Jean Dieudonné and Alexander Grothendieck. Éléments de géométrie algébrique. *Inst. Hautes Études Sci. Publ. Math.*, 4, 8, 11, 17, 20, 24, 28, 32, 1961–1967.
- [GW10] Ulrich Görtz and Torsten Wedhorn. *Algebraic geometry I*. Advanced Lectures in Mathematics. Vieweg + Teubner, Wiesbaden, 2010. Schemes with examples and exercises.
- [Hal19] Yatir Halevi. On stably pointed varieties and generically stable groups in ACVF. *Ann. Pure Appl. Logic*, 170(2):180–217, 2019.
- [Hal21] Yatir Halevi. On stably pointed varieties and generically stable groups in ACVF, 2021. preprint, <https://arxiv.org/abs/1611.05422>.
- [Hal22] Yatir Halevi. Corrigendum to “On stably pointed varieties and generically stable groups in ACVF” [Ann. Pure Appl. Log. 170(2) (2019) 180–217]. *Ann. Pure Appl. Logic*, 173(1):Paper No. 103045, 3, 2022.
- [HHM08] Deirdre Haskell, Ehud Hrushovski, and Dugald Macpherson. *Stable domination and independence in algebraically closed valued fields*, volume 30 of *Lecture Notes in Logic*. Association for Symbolic Logic, Chicago, IL; Cambridge University Press, Cambridge, 2008.
- [HHS21] Martin Hils, Ehud Hrushovski, and Pierre Simon. Definable equivariant retractions in non-archimedean geometry, 2021. preprint, <https://arxiv.org/abs/2101.02619>.
- [HK23] Yatir Halevi and Itay Kaplan. Saturated models for the working model theorist. *Bulletin of Symbolic Logic*, page 1–5, 2023.
- [HL16] Ehud Hrushovski and François Loeser. *Non-archimedean tame topology and stably dominated types*, volume 192 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2016.
- [HRK19] Ehud Hrushovski and Silvain Rideau-Kikuchi. Valued fields, metastable groups. *Selecta Math. (N.S.)*, 25(3):Paper No. 47, 58, 2019.
- [KM85] Nicholas M. Katz and Barry Mazur. *Arithmetic moduli of elliptic curves*, volume 108 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1985.
- [Kow00] Piotr Kowalski. Stable groups and algebraic groups. *J. London Math. Soc. (2)*, 61(1):51–57, 2000.
- [Kuh] Franz-Viktor Kuhlmann. *Book on Valuation Theory*. <http://math.usask.ca/~fvk/Fvkbook.htm>.
- [Kuh10] Franz-Viktor Kuhlmann. Elimination of ramification I: the generalized stability theorem. *Trans. Amer. Math. Soc.*, 362(11):5697–5727, 2010.
- [Mum95] David Mumford. *Algebraic geometry. I*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Complex projective varieties, Reprint of the 1976 edition.
- [Nag66] Masayoshi Nagata. Finitely generated rings over a valuation ring. *J. Math. Kyoto Univ.*, 5:163–169, 1966.
- [Nér64] André Néron. Modèles minimaux des variétés abéliennes sur les corps locaux et globaux. *Inst. Hautes Études Sci. Publ. Math.*, (21):128, 1964.
- [Ray70] Michel Raynaud. *Faisceaux amples sur les schémas en groupes et les espaces homogènes*. Lecture Notes in Mathematics, Vol. 119. Springer-Verlag, Berlin-New York, 1970.
- [She09] Saharon Shelah. Dependent first order theories, continued. *Israel J. Math.*, 173:1–60, 2009.
- [Sil94] Joseph H. Silverman. *Advanced topics in the arithmetic of elliptic curves*, volume 151 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994.
- [Sil09] Joseph H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer, Dordrecht, second edition, 2009.
- [Sim15] Pierre Simon. *A guide to NIP theories*, volume 44 of *Lecture Notes in Logic*. Association for Symbolic Logic, Chicago, IL; Cambridge Scientific Publishers, Cambridge, 2015.
- [Sta23] The Stacks Project Authors. Stacks project. <http://stacks.math.columbia.edu>, 2023.
- [You] Alex Youcis. Are elliptic curve irreducible? Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/4001445> (version: 2021-01-27).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, 199 ABBA KHOUSHY AVENUE, HAIFA, ISRAEL

Email address: ybenar@campus.haifa.ac.il