

The q -fractional Euler top system with one control on a fractional Leibniz algebroid

Gheorghe IVAN

Abstract. We determine the fractional almost Poisson realizations for fractional Euler top system with one control. These realizations allow us to introduce a fractional Leibniz algebroid structure on \mathbf{R}^3 and also to define the q -fractional Euler top system with one control on a fractional Leibniz algebroid. Finally, the numerical integration of them are discussed. ¹

1 Introduction

Fractional calculus is an interdisciplinary area of applied mathematics with many applications in several fields of science and engineering. For example, the fractional calculus has been applied to variational problems in: fluid mechanics, geometrical mechanics, chaotic dynamics, quantum mechanics, nanotechnology [34, 1, 32, 10, 31, 11, 4, 20, 17]. The fractional derivatives and integrals describe with great precision the dynamic behavior of various systems of differential equations [38, 29, 30, 39, 21, 12, 16, 26, 27].

The Lie algebroids have been used to geometric formulation of Hamiltonian mechanics. In Hamiltonian mechanics the approach is to consider the cotangent bundle. The cotangent bundle forms a phase space consisting of the configuration and momentum space. A state is a point in the phase space and an observable is given by Hamiltonian function. This function is used to generate the sections of the bundle. The evolution is given by a bundle morphism (named, anchor) and is defined by Hamilton's equations. The concept of Leibniz algebroid is a wakened version of a Lie algebroid, where the bilinear operation on sections of the vector bundle is not necessarily skew-symmetric and the anchor is a homomorphism of Leibniz algebra.

The Lie algebroids and Leibniz algebroids have proven to be powerful tools for geometric formulation of the Hamiltonian mechanics [2, 3, 7, 13, 14, 33, 36]. Also, they have been used in the investigation of many fractional dynamical systems [22, 23, 35, 6, 24].

In this paper we refer to a special class of dynamical systems. This class is formed by a family of differential equations on \mathbf{R}^3 which depends by a triple of real parameters $(\alpha_1, \alpha_2, \alpha_3)$ and one control parameter $k \in \mathbf{R}$, called the Euler top system with one control. It is described by the following nonlinear differential equations on \mathbf{R}^3 :

$$\dot{x}^1 = \alpha_1 x^2 x^3, \quad \dot{x}^2 = \alpha_2 x^1 x^3 - k x^1, \quad \dot{x}^3 = \alpha_3 x^1 x^2, \quad (1.1)$$

where x^1, x^2, x^3 are state variables and $\alpha_i, k \in \mathbf{R}$ for $i = \overline{1, 3}$ such that $\alpha_1 \alpha_2 \alpha_3 \neq 0$.

The family of dynamical systems (1.1) generalizes for example: the Lagrange system [37], a particular case of the Rabinovich system which is determined by the nonlinear terms [6] and the general Euler top system [15, 25].

The goal of our work is to study the fractional-order model associated to Euler top system with one control (1.1).

The paper is structured as follows. In Section 2, fractional Euler top system with one control (2.4) is presented. We prove that the fractional top system (2.4) has two fractional almost Poisson realizations (Proposition 2.1). In Section 3, the Proposition 2.1 is used to give the construction of q -fractional Euler top system with control on a fractional Leibniz algebroid. In Section 4, the numerical integration q -fractional Euler top system with one control (3.8)

¹MSC 2020: 26A33, 17B66, 34A08, 65L12, 70H05.

Key words: Caputo fractional derivative, fractional Leibniz algebroid, fractional Euler top system, numerical integration.

is discussed. As example, the numerical simulation for the q -fractional Lagrange system with one control (3.9) is given.

2 Fractional almost Poisson realizations of fractional Euler top system with one control

We start by recalling of the notion fractional Leibniz structure [19].

Let $f \in C^\infty(\mathbf{R})$ and $q \in \mathbf{R}, q > 0$. The q -order Caputo differential operator [3, 8] is described by $D_t^q f(t) = I^{m-q} f^{(m)}(t)$, $q > 0$, where $f^{(m)}(t)$ is the m -order derivative of the function f , $m \in \mathbf{N}^*$ is an integer such that $m - 1 \leq q \leq m$ and I^q is the q -order Riemann-Liouville integral operator expressed by:

$$I^q f(t) = \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} f(s) ds, \quad q > 0,$$

where Γ is the Euler Gamma function. If $q = 1$, then $D_t^q f(t) = df/dt$. We suppose $q \in (0, 1)$.

Let M be a n -dimensional smooth manifold, $U \subset M$ a local chart, $(x^i), i = \overline{1, n}$ a system of coordinates on U and $D_{x^i}^q f$ the Caputo partial derivatives for $f \in C^\infty(U)$.

We denote by $\mathcal{X}^q(U)$ be the module of the fractional vector fields generated by the operators $D_{x^i}^q, i = \overline{1, n}$. A fractional vector field $\overset{q}{X} \in \mathcal{X}^q(U)$ [19], has the form:

$$\overset{q}{X} = \overset{q}{X}^i D_{x^i}^q, \quad \overset{q}{X}^i \in C^\infty(U), \quad i = \overline{1, n}.$$

Let $\mathcal{D}^q(U)$ the module generated by 1-forms $d(x^i)^q, i = \overline{1, n}$ on U . The fractional exterior derivative $d^q : C^\infty(U) \rightarrow \mathcal{D}^q(U)$ is expressed by:

$$d^q(f) = d(x^i)^q D_{x^i}^q(f), \quad f \in C^\infty(U).$$

Let $\overset{q}{P} \in \mathcal{X}^q(M) \times \mathcal{X}^q(M)$ be a fractional 2-tensor field and $d^q f, d^q g \in \mathcal{D}^q(M)$. The bilinear map $[\cdot, \cdot]^q : C^\infty(M) \times C^\infty(M) \rightarrow C^\infty(M)$ defined by:

$$[f, h]^q = \overset{q}{P}(d^q f, d^q h), \quad (\forall) f, h \in C^\infty(M),$$

is called the *fractional Leibniz bracket*.

The *fractional Leibniz vector field* $\overset{q}{X}_h$ associated to $h \in C^\infty(M)$, is defined by:

$$\overset{q}{X}_h(f) = [f, h]^q, \quad (\forall) f \in C^\infty(M),$$

Locally, if $\overset{q}{P}$ is a skew-symmetric fractional 2-tensor field on M generated by the matrix $\overset{q}{P} = (\overset{q}{P}^{ij})$, then the dynamical system associated to $\overset{q}{X}_h$ is given by:

$$D_t^q x^i(t) = [x^i(t), h(t)]^q, \quad \text{where } [x^i, h]^q = \overset{q}{P}^{ij} \cdot D_{x^j}^q h,$$

or equivalently

$$D_t^q x^i = \overset{q}{P}^{ij} \cdot D_{x^j}^q h. \quad (2.1)$$

The system (2.1) is called the *fractional Leibniz system* associated to matrix $\overset{\alpha}{P}$ with the Hamiltonian $h \in C^\infty(M)$.

Let now $[\cdot, \cdot]^q$ be a fractional Leibniz structure on \mathbf{R}^n with the associated matrix $\overset{q}{P} = (\overset{q}{P}^{ij})$, i.e. $\overset{q}{P}^{ij} = [x^i, x^j]^q, i, j = \overline{1, n}$.

Definition 2.1 ([19]) Let be a fractional dynamical system on \mathbf{R}^n given by:

$$D_t^q x^i(t) = f^i(x^1(t), \dots, x^n(t)), \quad f^i \in C^\infty(\mathbf{R}^n, \mathbf{R}), \quad i = \overline{1, n}. \quad (2.2)$$

We say that the fractional dynamics (2.2) has a *fractional Leibniz realization*, if there exist a Leibniz structure $[\cdot, \cdot]^q$ on \mathbf{R}^n generated by the matrix $\overset{q}{P} = (\overset{q}{P}^{ij})$ and a Hamiltonian $\overset{q}{H} \in C^\infty(\mathbf{R}^n, \mathbf{R})$ such that (2.2) can be written in the following form:

$$D_t^q x(t) = \overset{q}{P}(x(t)) \cdot \nabla^q \overset{q}{H}(x(t)), \quad (2.3)$$

where $x(t) = (x^1(t), \dots, x^n(t))^T$ and $\nabla^q \overset{q}{H} = (D_{x^1}^q \overset{q}{H}, \dots, D_{x^n}^q \overset{q}{H})^T$. \square

A fractional Leibniz realization is denoted by $(\mathbf{R}^n, [\cdot, \cdot]^q, \overset{q}{H})$ or $(\mathbf{R}^n, \overset{q}{P}, \overset{q}{H})$.

If the matrix $\overset{q}{P} = (\overset{q}{P}^{ij})$ is skew-symmetric, then $(\mathbf{R}^n, \overset{q}{P}, \overset{q}{H})$ is called the *fractional almost Poisson realization* of the system (2.2).

A *fractional Casimir* of the configuration $(\mathbf{R}^n, [\cdot, \cdot]^q, \overset{q}{H})$ is a function $\overset{q}{C} \in C^\infty(\mathbf{R}^n, \mathbf{R})$ such that $[\overset{q}{C}, f]^q = 0$ for all $f \in C^\infty(\mathbf{R}^n, \mathbf{R})$, that is $\overset{q}{P}(x(t)) \cdot \nabla^q \overset{q}{C}(x(t)) = 0$.

The *fractional Euler top system with parameters* is defined by the following set of fractional differential equations:

$$\begin{cases} D_t^q x^1(t) = \alpha_1 x^2(t) x^3(t) \\ D_t^q x^2(t) = \alpha_2 x^1(t) x^3(t) - k x^1(t), \quad q \in (0, 1), \\ D_t^q x^3(t) = \alpha_3 x^1(t) x^2(t) \end{cases} \quad (2.4)$$

where $\alpha_i, k \in \mathbf{R}$ for $i = \overline{1, 3}$ such that $\alpha_1 \alpha_2 \alpha_3 \neq 0$.

For $k \neq 0$, the fractional dynamics (2.4) is called the *fractional Euler top system with one control*. If $k = 0$, then (2.4) is the *fractional Euler top system*, [28].

For $q = 1$, (2.4) becomes the *Euler top system with parameters* given by

$$\dot{x}^1 = \alpha_1 x^2 x^3, \quad \dot{x}^2 = \alpha_2 x^1 x^3 - k x^1, \quad \dot{x}^3 = \alpha_3 x^1 x^2, \quad (2.5)$$

where $\alpha_i, k \in \mathbf{R}$ for $i = \overline{1, 3}$ such that $\alpha_1 \alpha_2 \alpha_3 \neq 0$.

For $k \neq 0$, the dynamics (2.5) is called the *Euler top system with one control*. If $k = 0$, then (2.5) is the *Euler top system* [18, 15].

Example 2.1 (i) For $\alpha = (1, 1, 1)$ and $k \neq 0$, the system (2.4) becomes:

$$D_t^q x^1 = x^2 x^3, \quad D_t^q x^2 = x^1 x^3 - k x^1, \quad D_t^q x^3 = x^1 x^2. \quad (2.6)$$

The system (2.6) is called the *fractional Lagrange system with one control*. If $k = 0$ and $q = 1$, the system (2.6) is the *Lagrange system* in the theory of static $SU(2)$ -monopoles [37].

(ii) For $\alpha = (1, -1, 1)$ and $k \neq 0$, the system (2.4) becomes:

$$D_t^q x^1 = x^2 x^3, \quad D_t^q x^2 = -x^1 x^3 - k x^1, \quad D_t^q x^3 = x^1 x^2. \quad (2.7)$$

The system (2.7) is called the *fractional Rabinovich system with one control*. For $k = 0$ and $q = 1$, the system (2.7) is the *Rabinovich system* [6]. \square

Proposition 2.1 Let be $q \in (0, 1]$. The fractional Euler top system with parameters (2.4) has the following fractional almost Poisson realizations:

(i) $(\mathbf{R}^3, P_1^\alpha, \overset{q}{H}_1^\alpha)$ with the fractional Casimir $\overset{q}{C}_1^\alpha$, where

$$P_1^\alpha(x, k) = \begin{pmatrix} 0 & k & -\alpha_3 x^2 \\ -k & 0 & -\frac{\alpha_2 \alpha_3}{\alpha_1} x^1 \\ \alpha_3 x^2 & \frac{\alpha_2 \alpha_3}{\alpha_1} x^1 & 0 \end{pmatrix}, \quad (2.8)$$

$$\overset{q}{H}_1^\alpha(x) = \frac{1}{\Gamma(q+2)} [(x^1)^{1+q} - \frac{\alpha_1}{\alpha_3} (x^3)^{1+q}], \quad (2.9)$$

$$\overset{q}{C}_1^\alpha(x, k) = \frac{1}{\Gamma(q+2)} [\alpha_2 (x^1)^{1+q} - \alpha_1 (x^2)^{1+q}] - \frac{k \alpha_1}{\Gamma(q+1) \alpha_3} (x^3)^q. \quad (2.10)$$

(ii) $(\mathbf{R}^3, P_2^\alpha, \overset{q}{H}_2^\alpha)$ with the fractional Casimir $\overset{q}{C}_2^\alpha$, where

$$P_2^\alpha(x) = \begin{pmatrix} 0 & x^3 & 0 \\ -x^3 & 0 & -\frac{\alpha_3}{\alpha_1} x^1 \\ 0 & \frac{\alpha_3}{\alpha_1} x^1 & 0 \end{pmatrix}, \quad (2.11)$$

$$\overset{q}{H}_2^\alpha(x, k) = \frac{1}{\Gamma(q+2)} [-\alpha_2 (x^1)^{1+q} + \alpha_1 (x^2)^{1+q}] + \frac{k \alpha_1}{\Gamma(q+1) \alpha_3} (x^3)^q, \quad (2.12)$$

$$\overset{q}{C}_2^\alpha(x) = \frac{1}{\Gamma(q+2)} [\alpha_3 (x^1)^{1+q} - \alpha_1 (x^3)^{1+q}]. \quad (2.13)$$

Proof. (i) Using the relations $D_{x^i}^q (x^j)^m = \delta_i^j \frac{\Gamma(1+m)}{\Gamma(1+m-q)} (x^j)^{m-q}$, $i, j = \overline{1, 3}$ follows:

$$D_{x^1}^q \overset{q}{H}_1^\alpha = x^1, \quad D_{x^2}^q \overset{q}{H}_1^\alpha = 0, \quad D_{x^3}^q \overset{q}{H}_1^\alpha = -\frac{\alpha_1}{\alpha_3} x^3, \quad D_{x^1}^q \overset{q}{C}_1^\alpha = \alpha_2 x^1, \quad D_{x^2}^q \overset{q}{C}_1^\alpha = -\alpha_1 x^2,$$

$$D_{x^3}^q \overset{q}{C}_1^\alpha = -\frac{k \alpha_1}{\alpha_3}. \quad \text{Then: } P_1^\alpha \cdot \nabla \overset{q}{H}_1^\alpha =$$

$$= \begin{pmatrix} 0 & k & -\alpha_3 x^2 \\ -k & 0 & -\frac{\alpha_2 \alpha_3}{\alpha_1} x^1 \\ \alpha_3 x^2 & \frac{\alpha_2 \alpha_3}{\alpha_1} x^1 & 0 \end{pmatrix} \begin{pmatrix} x^1 \\ 0 \\ -\frac{\alpha_1}{\alpha_3} x^3 \end{pmatrix} = \begin{pmatrix} \alpha_1 x^2 x^3 \\ \alpha_2 x^1 x^3 - k x^1 \\ \alpha_3 x^1 x^2 \end{pmatrix} = \begin{pmatrix} D_t^q x^1 \\ D_t^q x^2 \\ D_t^q x^3 \end{pmatrix}.$$

Hence $D_t^q x(t) = P_1^\alpha(x(t)) \cdot \nabla \overset{q}{H}_1^\alpha(x(t))$ and (2.4) is a fractional almost Poisson system. Also, $\overset{q}{C}_1^\alpha$ is a fractional Casimir, since:

$$P_1^\alpha \cdot \nabla \overset{q}{C}_1^\alpha = \begin{pmatrix} 0 & k & -\alpha_3 x^2 \\ -k & 0 & -\frac{\alpha_2 \alpha_3}{\alpha_1} x^1 \\ \alpha_3 x^2 & \frac{\alpha_2 \alpha_3}{\alpha_1} x^1 & 0 \end{pmatrix} \begin{pmatrix} \alpha_2 x^1 \\ -\alpha_1 x^2 \\ -\frac{k \alpha_1}{\alpha_3} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

(ii) We have $D_{x^1}^q \overset{q}{H}_2^\alpha = -\alpha_2 x^1$, $D_{x^2}^q \overset{q}{H}_2^\alpha = \alpha_1 x^2$, $D_{x^3}^q \overset{q}{H}_2^\alpha = \frac{k \alpha_1}{\alpha_3}$, $D_{x^1}^q \overset{q}{C}_2^\alpha = \alpha_3 x^1$, $D_{x^2}^q \overset{q}{C}_2^\alpha = 0$, $D_{x^3}^q \overset{q}{C}_2^\alpha = -\alpha_1 x^3$. Similarly, $D_t^q x = P_2^\alpha \cdot \nabla \overset{q}{H}_2^\alpha$ and $P_2^\alpha \cdot \nabla \overset{q}{C}_2^\alpha = 0$. \square

Remark 2.1 If in Proposition 2.1 we replace $\alpha = (\alpha_1, \alpha_2, \alpha_3)$ with $\alpha = (1, 1, 1)$ (resp. $\alpha = (1, -1, 1)$) we obtain two fractional almost Poisson realizations for the fractional Lagrange (resp. Rabinovich) system with parameter (2.6) (resp. (2.7)). \square

3 Fractional Euler top system with one control on a fractional Leibniz algebroid

A $\widehat{\text{fractional}}$ Leibniz algebroid structure [23] on a vector bundle $\pi : E \rightarrow M$ is given by a bracket $[\cdot, \cdot]^q$ on the space of sections $\text{Sec}(\pi)$ and two vector bundle morphisms $\overset{q}{\rho}_1, \overset{q}{\rho}_2 : E \rightarrow T^q M$ (called the *left* and *right fractional anchor*) such that for all $\sigma_1, \sigma_2 \in \text{Sec}(\pi)$ and $f, g \in C^\infty(M)$ we have:

$$\begin{cases} [e_u, e_v]^q = C_{uv}^w e_w, \\ [f\sigma_1, g\sigma_2]^q = f\overset{q}{\rho}_1(\sigma_1)(g)\sigma_2 - g\overset{q}{\rho}_2(\sigma_2)(f)\sigma_1 + fg[\sigma_1, \sigma_2]^q. \end{cases}$$

A vector bundle $\pi : E \rightarrow M$ endowed with a fractional Leibniz algebroid structure $([\cdot, \cdot]^q, \overset{q}{\rho}_1, \overset{q}{\rho}_2)$ on E , is called *fractional Leibniz algebroid* over M and it is denoted by $(E, [\cdot, \cdot]^q, \overset{q}{\rho}_1, \overset{q}{\rho}_2)$.

In the paper [23] has shown that there exists a bijective correspondence between the fractional Leibniz algebroid structures on a vector bundle E and the linear fractional 2-tensor fields on the dual vector bundle E^* .

More precisely, a linear fractional 2-tensor field $\overset{q}{\Lambda}$ on the dual vector bundle $\pi^* : E^* \rightarrow M$ defines a fractional Leibniz structure on the vector bundle $\pi : E \rightarrow M$.

The bracket $[\cdot, \cdot]_{\overset{q}{\Lambda}}$ is defined by:

$$\begin{aligned} [F, G]_{\overset{q}{\Lambda}} &= \overset{q}{\Lambda} (d^{qr} F, d^{qr} G), \quad (\forall) F, G \in C^\infty(E^*), \quad \text{where} \\ d^{qr} F &= d(x^i)^q D_{x^i}^q H + d(\xi_u)^r D_{\xi_u}^r F = d^q(F) + d^r(F). \end{aligned}$$

If (x^i) , (x^i, y^u) resp., (x^i, ξ_u) for $i = \overline{1, n}$, $u = \overline{1, m}$ are coordinates on M, E resp. E^* , then the linear fractional tensor $\overset{q}{\Lambda}$ on E^* has the form:

$$\overset{q}{\Lambda} = C_{uv}^w \xi_w D_{\xi_u}^r \otimes D_{\xi_v}^r + \overset{q}{\rho}_{1u}^i D_{\xi_u}^r \otimes D_{x^i}^q - \overset{q}{\rho}_{2u}^i D_{x^i}^q \otimes D_{\xi_u}^r.$$

The fractional system associated to vector field $\overset{q}{X}_H$ with $H \in C^\infty(E^*)$ given by:

$$\overset{q}{X}_H(F) = \overset{q}{\Lambda} (d^{qr} F, d^{qr} H), \quad \text{for all } F \in C^\infty(E^*) \quad (3.1)$$

is called a *fractional dynamical system on the fractional Leibniz algebroid* $(E, [\cdot, \cdot]^q, \overset{q}{\rho}_1, \overset{q}{\rho}_2)$, [19].

Locally, the fractional dynamical system (3.1) reads:

$$\begin{cases} D_t^q \xi_u = [\xi_u, H]_{\overset{q}{\Lambda}} = C_{uv}^w \xi_w D_{\xi_v}^r H + \overset{q}{\rho}_{1u}^i D_{x^i}^q H \\ D_t^q x^i = [x^i, H]_{\overset{q}{\Lambda}} = -\overset{q}{\rho}_{2u}^i D_{\xi_u}^r H \end{cases}. \quad (3.2)$$

In the following we suppose that $r = q$.

If $\tilde{P} = (C_{uv}^w \xi_w)$, $\overset{q}{\rho}_1 = (\overset{q}{\rho}_{1u}^i)$ and $\overset{q}{\rho}_2 = (\overset{q}{\rho}_{2u}^i)$, for $i = \overline{1, 3}$, $u, v, w = \overline{1, 3}$, then the fractional system (3.2) (with condition $r = q$) can be written in the matrix form:

$$\begin{pmatrix} D_t^q \xi_1 \\ D_t^q \xi_2 \\ D_t^q \xi_3 \end{pmatrix} = \tilde{P} \begin{pmatrix} D_{\xi_1}^q H \\ D_{\xi_2}^q H \\ D_{\xi_3}^q H \end{pmatrix} + \overset{q}{\rho}_1 \begin{pmatrix} D_{x^1}^q H \\ D_{x^2}^q H \\ D_{x^3}^q H \end{pmatrix}, \quad \begin{pmatrix} D_t^q x^1 \\ D_t^q x^2 \\ D_t^q x^3 \end{pmatrix} = -\overset{q}{\rho}_2 \begin{pmatrix} D_{\xi_1}^q H \\ D_{\xi_2}^q H \\ D_{\xi_3}^q H \end{pmatrix}. \quad (3.3)$$

As example we give the *fractional Euler top system with one control on a fractional Leibniz algebroid*.

Let the vector bundle $\pi : E = \mathbf{R}^3 \times \mathbf{R}^3 \rightarrow \mathbf{R}^3$ and $\pi^* : E^* = \mathbf{R}^3 \times (\mathbf{R}^3)^* \rightarrow \mathbf{R}^3$ its dual. We consider on E^* the linear 2-tensor field $\tilde{\Lambda}$, defined by the matrix \tilde{P} , the fractional anchors ${}^q\rho_1, {}^q\rho_2 : \mathbf{R}^3 \times \mathbf{R}^3 \rightarrow T(\mathbf{R}^3)$ and the function H defined by:

$$\tilde{P} = P(x, \xi, k) = \begin{pmatrix} 0 & k & -\alpha_3 x^2 \xi_2 \\ -k & 0 & -\frac{\alpha_2 \alpha_3}{\alpha_1} x^1 \xi_1 \\ \alpha_3 x^2 \xi_2 & \frac{\alpha_2 \alpha_3}{\alpha_1} x^1 \xi_1 & 0 \end{pmatrix}, \quad (3.4)$$

$${}^q\rho_1 = P_1^\alpha(x, k) = \begin{pmatrix} 0 & k & -\alpha_3 x^2 \\ -k & 0 & -\frac{\alpha_2 \alpha_3}{\alpha_1} x^1 \\ \alpha_3 x^2 & \frac{\alpha_2 \alpha_3}{\alpha_1} x^1 & 0 \end{pmatrix}, \quad (3.5)$$

$${}^q\rho_2 = - \begin{pmatrix} 0 & (x^3)^q & 0 \\ -(x^3)^q & 0 & -\frac{\alpha_3}{\alpha_1} (x^1)^q \\ 0 & \frac{\alpha_3}{\alpha_1} (x^1)^q & 0 \end{pmatrix} \quad (3.6)$$

$$H(x, \xi, k) = \frac{1}{\Gamma(1+q)} [-\alpha_2 (x^1)^q (\xi_1)^q + \alpha_1 (x^2)^q (\xi_2)^q + \frac{k\alpha_1}{\alpha_3} (\xi_3)^q]. \quad (3.7)$$

Proposition 3.1 *Let be $k \neq 0$ and $q \in (0, 1]$. The fractional system (3.3) on the fractional Leibniz algebroid $(\mathbf{R}^3 \times \mathbf{R}^3, \tilde{P}, {}^q\rho_1, {}^q\rho_2)$ associated to H , where $\tilde{P}, {}^q\rho_1, {}^q\rho_2, H$ are given by the relations (3.4) – (3.7), is described by:*

$$\begin{cases} D_t^q \xi_1 &= k\alpha_1((x^2)^q - x^2 \xi_2 + (\xi_2)^q) \\ D_t^q \xi_2 &= k\alpha_2((x^1)^q - x^1 \xi_1 + (\xi_1)^q) \\ D_t^q \xi_3 &= \alpha_2 \alpha_3 (-(x^1)^q x^2 \xi_2 + x^1 (x^2)^q \xi_1 + x^1 (\xi_2)^q - x^2 (\xi_1)^q). \\ D_t^q x^1 &= \alpha_1 (x^2)^q (x^3)^q \\ D_t^q x^2 &= \alpha_2 (x^1)^q (x^3)^q - k(x^1)^q \\ D_t^q x^3 &= \alpha_3 (x^1)^q (x^2)^q \end{cases} \quad (3.8)$$

Proof. We have $D_{\xi_1}^q H = -\alpha_2 (x^1)^q$, $D_{\xi_2}^q H = \alpha_1 (x^2)^q$, $D_{\xi_3}^q H = \frac{k\alpha_1}{\alpha_3}$, $D_{x^1}^q H = -\alpha_2 (\xi_1)^q$, $D_{x^2}^q H = \alpha_1 (\xi_2)^q$, $D_{x^3}^q H = 0$. Then (3.3) are written as follows:

$$\begin{pmatrix} D_t^q \xi_1 \\ D_t^q \xi_2 \\ D_t^q \xi_3 \end{pmatrix} = \tilde{P} \begin{pmatrix} -\alpha_2 (x^1)^q \\ \alpha_1 (x^2)^q \\ \frac{k\alpha_1}{\alpha_3} \end{pmatrix} + {}^q\rho_1 \begin{pmatrix} -\alpha_2 (\xi_1)^q \\ \alpha_1 (\xi_2)^q \\ 0 \end{pmatrix}, \quad \begin{pmatrix} D_t^q x^1 \\ D_t^q x^2 \\ D_t^q x^3 \end{pmatrix} = -{}^q\rho_2 \begin{pmatrix} -\alpha_2 (x^1)^q \\ \alpha_1 (x^2)^q \\ \frac{k\alpha_1}{\alpha_3} \end{pmatrix}.$$

Replacing now the matrices $\tilde{P}, {}^q\rho_1$ and ${}^q\rho_2$ one obtains the equations (3.8). \square

The dynamical system (3.8) is called the *q-fractional Euler top system with one control on the fractional Leibniz algebroid $\pi : E = \mathbf{R}^3 \times \mathbf{R}^3 \rightarrow \mathbf{R}^3$ for $k \neq 0$.*

Example 3.1 Taking $k \neq 0, \alpha = (1, 1, 1)$ and $q \in (0, 1)$ in the relations (3.4) – (3.8), the fractional system (3.3) on the fractional Leibniz algebroid $(\mathbf{R}^3 \times \mathbf{R}^3, \tilde{P}, \rho_1^q, \rho_2^q)$ associated to function H , is given by:

$$\left\{ \begin{array}{l} D_t^q \xi_1 = k[(x^2)^q - x^2 \xi_2 + (\xi_2)^q] \\ D_t^q \xi_2 = k[(x^1)^q - x^1 \xi_1 + (\xi_1)^q] \\ D_t^q \xi_3 = -(x^1)^q x^2 \xi_2 + x^1 (x^2)^q \xi_1 + x^1 (\xi_2)^q - x^2 (\xi_1)^q \\ D_t^q x^1 = (x^2)^q (x^3)^q \\ D_t^q x^2 = (x^1)^q (x^3)^q - k(x^1)^q \\ D_t^q x^3 = (x^1)^q (x^2)^q. \end{array} \right. \quad (3.9)$$

The dynamical system (3.9) is called the q -fractional Lagrange system with one control on the fractional Leibniz algebroid $\pi : E = \mathbf{R}^3 \times \mathbf{R}^3 \rightarrow \mathbf{R}^3$. \square

4 Numerical integration of the fractional Euler top with one control (3.8)

Let $f : \mathbf{R} \rightarrow \mathbf{R}$ be an integrable function. For $q \in (0, 1]$ we consider a function of C^1 -class $g^q : \mathbf{R} \times \mathbf{R} \rightarrow \mathbf{R}$, $(t, s) \rightarrow g_t^q(s)$ for $t, s \in \mathbf{R}$ and $t_0 \leq s \leq t$.

A Riemann integral of f with respect to g_t^q is defined by ${}_{t_0}I_{t, g_t^q} f(t) = \int_{t_0}^t f(s) g_t^q(s) ds$.

If the function g_t^q is defined by

$$g_t^q(s) = \frac{1}{\Gamma(q)} (t-s)^{q-1} e^{-\rho(t-s)}, \quad (4.1)$$

where $\Gamma(q)$ is the Euler Gamma function and $\rho > 0$, then the Riemann integral ${}_{t_0}I_{t, g_t^q} f(t)$ becomes the fractional integral, denoted with ${}_{t_0}I_t^q f(t)$, where:

$${}_{t_0}I_t^q f(t) = \frac{1}{\Gamma(q)} \int_{t_0}^t (t-s)^{q-1} e^{-\rho(t-s)} f(s) ds. \quad (4.2)$$

Remark 4.1 For $\rho = 0$, the relation (4.2) is the fractional Riemann-Liouville integral ([31], Section 2.1). Also, the fractional integral defined by (4.2) is a special case of the generalized fractional El-Nabulsi integral [10]. \square

If $\varphi : \mathbf{R} \times \mathbf{R}^n \rightarrow \mathbf{R}^n$ is a deterministic function, we will denote by:

$${}_{t_0}I_{t, g_t^q} \varphi(t, x(t)) = \int_{t_0}^t \varphi(s, x(s)) g_t^q(s) ds,$$

where $x(t) = (x^1(t), \dots, x^n(t))$ and the integral is Riemann integral with respect to g_t^q .

We call *Volterra integral equation with respect to g_t^q* , the functional Volterra type equation given by:

$$x(t) = x(t_0) + {}_{t_0}I_{t, g_t^q} \varphi(t, x(t)). \quad (4.3)$$

The equation (4.3) can be written formally in the following way:

$$dx(s) = \varphi(s, x(s)) g_t^q(s) ds. \quad (4.4)$$

We apply the above considerations for the fractional Euler top system (3.8)

$$\begin{cases} D_t^q x^i(t) &= F_i(x^1(t), x^2(t), x^3(t)), & i = \overline{1, 3} \\ D_t^q \xi_i(t) &= F_{i+3}(x^1(t), x^2(t), x^3(t), \xi_1(t), \xi_2(t), \xi_3(t)), & i = \overline{1, 3} \\ x(t_0) &= (x^1(t_0), x^2(t_0), x^3(t_0)), \quad \xi(t_0) = (\xi_1(t_0), \xi_2(t_0), \xi_3(t_0)) \end{cases} \quad (4.5)$$

where $t \in (t_0, \tau)$, $q \in (0, 1)$ and

$$\begin{cases} F_1(x) &= \alpha_1(x^2)^q(x^3)^q \\ F_2(x) &= \alpha_2(x^1)^q(x^3)^q - k(x^1)^q \\ F_3(x) &= \alpha_3(x^1)^q(x^2)^q \\ F_4(x, \xi) &= k\alpha_1((x^2)^q - x^2\xi_2 + (\xi_2)^q) \\ F_5(x, \xi) &= k\alpha_2((x^1)^q - x^1\xi_1 + (\xi_1)^q) \\ F_6(x, \xi) &= \alpha_2\alpha_3(-(x^1)^qx^2\xi_2 + x^1(x^2)^q\xi_1 + x^1(\xi_2)^q - x^2(\xi_1)^q). \end{cases} \quad (4.6)$$

Since the functions $F_i(x)$ and $F_{i+3}(x, \xi)$ for $i = \overline{1, 3}$ are continuous, the initial value problem (4.5) is equivalent to system of Volterra integral equations [31, 34], which is given as follows:

$$\begin{cases} x^i(t) &= x^i(t_0) + {}_{t_0}I_{t, g_t^q} F_i(x^1(t), x^2(t), x^3(t)), & i = \overline{1, 3}, \\ \xi_i(t) &= \xi_i(t_0) + {}_{t_0}I_{t, g_t^q} F_{i+3}(x^1(t), x^2(t), x^3(t), \xi_1(t), \xi_2(t), \xi_3(t)), & i = \overline{1, 3}. \end{cases} \quad (4.7)$$

The equations (4.7) can be written in the following form:

$$\begin{cases} dx^1(s) &= F_1(x(s))g_t^q(s)ds \\ dx^2(s) &= F_2(x(s))g_t^q(s)ds, \\ dx^3(s) &= F_3(x(s))g_t^q(s)ds \\ d\xi_1(s) &= F_4(x(s), \xi(s))g_t^q(s)ds \\ d\xi_2(s) &= F_5(x(s), \xi(s))g_t^q(s)ds \\ d\xi_3(s) &= F_6(x(s), \xi(s))g_t^q(s)ds, \end{cases} \quad (4.8)$$

where g_t^q and $F_i(x(s))$, $F_{i+3}(x(s), \xi(s))$, $i = \overline{1, 3}$ are given in (4.1) and (4.6).

The system (4.8) is called the *Volterra integral equations associated to Euler top system* (4.6).

The problem for solving the system (4.8) is reduced to one of solving a sequence of systems of fractional equations in increasing dimension on successive intervals $[j, (j+1)]$.

For the numerical integration of the system (4.8) one can use the first Euler scheme [9, 31], which is expressed as follows:

$$\begin{cases} x^i(j+1) &= x^i(j) + hF_i(x^1(j), x^2(j), x^3(j))g_t^q(j), & i = \overline{1, 3}, \\ \xi_i(j+1) &= \xi_i(j) + hF_{i+3}(x^1(j), x^2(j), x^3(j), \xi_1(j), \xi_2(j), \xi_3(j))g_t^q(j), & i = \overline{1, 3}, \end{cases} \quad (4.9)$$

where $j = 0, 1, 2, \dots, N$, $h = \frac{T}{N}$, $T > 0$, $N > 0$.

More precisely, if $t_0 = 0$, $\rho > 0$ and $\varepsilon > 0$, the numerical integration of the system (4.9) is given by:

$$\left\{ \begin{array}{l} x^1(j+1) = x^1(j) + h \frac{1}{\Gamma(q)} (t-j)^{q-1} \varepsilon^{-\rho(t-j)} (\alpha_1 (x^2)^q(j) (x^3)^q(j)) \\ x^2(j+1) = x^2(j) + h \frac{1}{\Gamma(q)} (t-j)^{q-1} \varepsilon^{-\rho(t-j)} (\alpha_2 (x^1)^q(j) (x^3)^q(j) - k (x^1)^q(j)) \\ x^3(j+1) = x^3(j) + h \frac{1}{\Gamma(q)} (t-j)^{q-1} \varepsilon^{-\rho(t-j)} (\alpha_3 (x^1)^q(j) (x^2)^q(j)) \\ \xi_1(j+1) = \xi_1(j) + h \frac{1}{\Gamma(q)} (t-j)^{q-1} \varepsilon^{-\rho(t-j)} (k \alpha_1 ((x^2)^q(j) - x^2(j) \xi_2(j) + (\xi_2)^q(j))) \\ \xi_2(j+1) = \xi_2(j) + h \frac{1}{\Gamma(q)} (t-j)^{q-1} \varepsilon^{-\rho(t-j)} (k \alpha_2 ((x^1)^q(j) - x^1(j) \xi_1(j) + (\xi_1)^q(j))) \\ \xi_3(j+1) = \xi_3(j) + h \frac{1}{\Gamma(q)} (t-j)^{q-1} \varepsilon^{-\rho(t-j)} (\alpha_2 \alpha_3 (- (x^1)^q(j) x^2(j) \xi_2(j) + \\ + x^1(j) (x^2)^q(j) \xi_1(j) + x^1(j) (\xi_2)^q(j) - x^2(j) (\xi_1)^q(j))) \\ x^i(0) = \xi_i(0) = x_e^i + \varepsilon, \quad i = \overline{1,3}. \end{array} \right. \quad (4.10)$$

Using [5, 9], we have that the numerical algorithm given by (4.10) is convergent.

Example 4.1 Let us we present the numerical simulation of solutions of q -fractional Lagrange system with one control on a fractional Leibniz algebroid (3.9). Then, in (4.10) we take: $\alpha_1 = \alpha_2 = \alpha_3 = 1, k = 1$ and $q = 0.5$. For to integrate this q -fractional model, we consider $h = 0.01, \varepsilon = 0.01, N = 500, t = 502$ and the initial conditions $x^1(0) = \xi_1(0) = \varepsilon, x^2(0) = \xi_2(0) = \varepsilon, x^3(0) = \xi_3(0) = \varepsilon + 0.5$.

Using the software Maple, the orbits $(n, x^i(n)), i = \overline{1,3}$ and $(n, \xi_i(n)), i = \overline{1,3}$ of the fractional model (3.9) are represented in Figures 1-3 and Figures 4-6.

In the coordinate systems $Ox^1x^2x^3$ and $O\xi_1\xi_2\xi_3$, the orbits $(x^1(n), x^2(n), x^3(n))$ and $(\xi_1(n), \xi_2(n), \xi_3(n))$ for the solutions of equations (3.9) are represented in the Figures 7-8.




Figure 1. $(n, x^1(n))$




Figure 2. $(n, x^2(n))$




Figure 3. $(n, x^3(n))$




Figure 4. $(n, \xi_1(n))$




Figure 5. $(n, \xi_2(n))$




Figure 6. $(n, \xi_3(n))$




Figure 7. $(x^1(n), x^2(n), x^3(n))$ for $q = 0.5$




Figure 8. $(\xi_1(n), \xi_2(n), \xi_3(n))$ for $q = 0.5$. \square

Conclusions. This paper investigates a new family of fractional differential systems, called the fractional Euler top system with one control (2.4). The q -fractional Euler top system with one control on a fractional Leibniz algebroid (3.8) was constructed. The Adams-Bashforth-Moulton algorithm was applied for the numerical integration of the q -fractional Euler top system (3.8). Finally, the numerical simulation was given for the q -fractional Lagrange system with one control on a fractional Leibniz algebroid (3.9). \square

References

- [1] Agrawal O.P., Formulation of Euler-Lagrange equations for fractional variational problems. *J. Math. Anal. Appl.*, **272** (2002) 368–379.
- [2] Anastasiei M., Mechanical systems on Lie algebroids. *Algebras Groups Geometry*, **23**(2006), 235-245.
- [3] Băleanu D., Agrawal O.P., Fractional Hamilton formalism within Caputo’s derivative. *Czech. J. Physics*, **56**(2006), no. 10, 1087-1092.
- [4] Băleanu D., Guvenc Ziya B., Tenreiro J.A., *New Trends in Nanotechnology and Fractional Calculus Applications*. Springer, 2009.
- [5] El-Borai M.M., El-Said K., Mostafa O.L., Ahmed H.M., Volterra equations with fractional stochastic integrals. *Mathematical Problems in Engineering*, **5**(2004), 453-468.

- [6] Chiş O., Despi I., Opreş D., Fractional equations on algebroids and fractional algebroids. In volume: *New Trends in Nanotechnology and Fractional Calculus Applications*. Springer, 2009, 193-201.
- [7] Degeratu M., Ivan M., Linear connections on Lie algebroids. *Proceedings of the 5th Conference of Balkan Society of Geometers*, Aug. 29- Sept. 2, 2005. Mangalia Romania. Geometry Balkan Press, 2006, 44-53.
- [8] Diethelm K., *Fractional Differential Equations. Theory and Numerical Treatment*. TU Braunschweig, 2003.
- [9] Diethelm K., Ford N.J., Freed A.D., A predictor-corrector approach for the numerical solution of fractional differential equations. *Nonlinear Dynamics*, **29** (2002), no. 1-4, 3-22.
- [10] El-Nabulsi R.A., A fractional action-like variational approach of some classical quantum and geometrical dynamics. *Int. J. Appl. Math.*, **17**(2005), 299-317.
- [11] El-Nabulsi R.A., Fractional field theory from multi-dimensional fractional: variational problems. *Int. J. Geom. Meth. Mod. Phys.*, **5**(2008), no. 6, 863-892.
- [12] El-Nabulsi R.A., Fractional variational approach for dissipative mechanical systems. *Anal. Theory Appl.*, **30**(2014), no. 3, 249-259.
- [13] El-Nabulsi R.A., Complex Lie algebroids and Finsler manifold in time-dependent fractal dimension and their associated decomplexification. *Differential Geometry and its Applications*, **77**(2021), 101775.
- [14] Ezuck R.J., Quantum systems as Lie algebroids. *ArXiv:2105.01513v1[math-ph]*, 2021.
- [15] Ivan G., Geometrical and dynamical properties of general Euler top system. *Indian J. Pure Appl. Math.*, **44**(2013), no.1, 77-93. Doi:10.1007/s13226-013-0004-0.
- [16] Ivan G., On fractional differential equations of 3D Maxwell-Bloch type. *Int. J. Geom. Met. Mod. Phys.*, **11**(2014), no. 4, 1450028 (12 pages). Doi:10.1142/S02198878.14500285.
- [17] Ivan G., A general method for stability controllability in the theory of fractional-order differential systems. *ArXiv math.*, 1-10 (2022). Doi:10.48550/arxiv.2210.12751.
- [18] Ivan G., Ivan M., General Euler top system and its Lax representation. *Int. J. Geom. Meth. Mod. Phys.*, **8**(2011), no. 5, 937-944. Doi:10.1142/S0219887811003543.
- [19] Ivan G., Opreş D., Dynamical systems on Leibniz algebroids. *Differ. Geom.-Dyn. Syst.*, **8**(2006), 127-137.
- [20] Ivan G., Opreş D., Ivan M., Hybrid fractional differential systems associated to Toda lattice, Numerical simulation. *Proc. of the 12th Symposium of Math. and Its Appl.* "Politehnica" University of Timișoara, November 5-7, 2009, 227-234.
- [21] Ivan, G., Opreş, D., Ivan, M., Stochastic fractional equations associated to Euler top system. *Int. J. Geom. Met. Mod. Phys.*, **10**, no. 1, 1220018 (10 pages)(2013). Doi:10.1142/S02198878.12200186.
- [22] Ivan G., Ivan M., Opreş D., Fractional Euler-Lagrange and fractional Wong equations for Lie algebroids. *Proceedings of the 4-th International Colloquium Mathematics and Numerical Physics*, October 6-8, 2006, Bucharest, Romania, 73-80.
- [23] Ivan G., Ivan M., Opreş D., Fractional dynamical systems on fractional Leibniz algebroids. *An. Ştiinţ. Univ. Al. I. Cuza Iaşi, Mat. (N.S.)*, **53** (2007), Supl., 222-234. <https://arxiv.org/pdf/0709.2443.pdf>.

- [24] Ivan G., Ivan M., Pop C., Numerical integration and synchronization for the 3-dimensional metriplectic Volterra system. *Math. Probl. Eng.*, **2011**, Article ID 723629 (11 pages). Doi:10.1155/2011/723629.
- [25] Ivan M., Control chaos in the fractional Lorenz-Hamilton system. *Fractional Differ. Calc.*, **6**(2016), no. 1, 111-119. Doi:10.7153/fdc-06-07.
- [26] Ivan M., A fractional model for the single Stokes pulse from the nonlinear optics. *Journal of Applied Mathematics and Physics*, **10**(2022), 2856-2875. Doi:10.4236/jamp.2022.1010191.
- [27] Ivan M., Dynamics analysis of the fractional-order Lagrange system. *Int. J. Modern Eng. Research*, **12**(2022), no.8, 23-31.
- [28] Ivan M., Ivan G., On the fractional Euler top system with two parameters. *Int. J. Modern Eng. Research*, **8**(2018), no. 4, 10-22.
- [29] Ivan M., Ivan G., Opreș D., Fractional differential systems associated to Toda lattice. *BSG Proceed. 16. The Int. Conf. of Diff. Geom. and Dynamical Systems (DGDS-2008)*, Aug. 29 – Sept. 2, 2008, Mangalia, Romania, 80-90.
- [30] Ivan M., Ivan G., Opreș D., Fractional equations of the rigid body on the pseudo-orthogonal group $SO(2,1)$. *Int. J. Geom. Met. Mod. Phys.*, **6**(2009), no. 7, 1181-1192. Doi:10.1142/S0219887809004168.
- [31] Kilbas A.A., Srivastava H.M., Trujillo J.J., *Theory and Applications of Fractional Differential Equations*. North-Holland, Math. Studies, **204**, 2006.
- [32] Kulish V.V., Lage J.L., Application of fractional calculus to fluid mechanics. *J. Fluids Eng.*, **124**(2002), 803-806.
- [33] Martinez E., Geometric formulation of mechanics on Lie algebroids. In: *Proceedings of the VIII Fall Workshop on Geometry and Physics*, Medina del Campo, 1999, Publicaciones de la RSME, 2 (2001), 209-222.
- [34] Podlubny I., *Fractional Differential Equations*, Academic Press, New York, 1999.
- [35] Pop C., Aron A., Galea C., Ciobanu M., Ivan M., Some geometric aspects in theory of Lotka-Volterra system. *Proc. of the 11-th WSEAS Int. Conf. on Sust. in Sci. Eng.*, Timișoara, Romania, May (2009), 91-97.
- [36] Popescu L., Aspects of Lie algebroid geometry and hamiltonian formalism. *An. Științ. Univ. Al. I. Cuza Iași, Mat. (N.S.)*, **53** (2007), Supl., 297-308.
- [37] Takhtajan L., On foundation of the generalized Nambu mechanics. *Comm. Math. Phys.*, **160** (1994), no. 2, 295-315.
- [38] Tarasov V.E., Fractional variations for dynamical systems: Hamilton and Lagrange approaches. *J. Phys. A: Math. Gen.*, **39**(2006), no. 26, 8409-8425.
- [39] Yajima T., Yamasaki K., Geometry of surfaces with Caputo fractional derivatives and applications to incompressible two-dimensional flows. *J. Phys. A: Math. Theor.*, **45**(2012), no. 6, 15 pages.

Author's adress

West University of Timișoara. Seminarul de Geometrie și Topologie.
 Department of Mathematics. Timișoara, Romania.
 E-mail: gheorghe.ivan@e-uvv.ro