

Bounds on piercing and line-piercing numbers in families of convex sets in the plane

Shira Zerbib *

Abstract

A family of sets has the (p, q) property if among any p members of it some q intersect. It is shown that if a finite family of compact convex sets in \mathbb{R}^2 has the $(p + 1, 2)$ property then it is pierced by $\lfloor \frac{p}{2} \rfloor + 1$ lines. A colorful version of this result is proved as well. As a corollary, the following is proved: Let \mathcal{F} be a finite family of compact convex sets in the plane with no isolated sets, and let \mathcal{F}' be the family of its pairwise intersections. If \mathcal{F} has the $(p + 1, 2)$ property and \mathcal{F}' has the $(r + 1, 2)$ property, then \mathcal{F} is pierced by $(\lfloor \frac{r}{2} \rfloor^2 + \lfloor \frac{r}{2} \rfloor)p$ points when $r \geq 2$, and by p points otherwise. The proofs use the topological KKM theorem.

1 Introduction

For integers $p \geq q > 0$, a family of sets \mathcal{F} has the (p, q) *property* if among any p members of \mathcal{F} some q have a nonempty intersection. The matching number of \mathcal{F} is the largest size a subfamily of pairwise disjoint sets in \mathcal{F} . Note that \mathcal{F} has the $(p + 1, 2)$ property if and only if its matching number is at most p . We say that \mathcal{F} is *pierced by k points* if there exists a set $C \subseteq \bigcup \mathcal{F}$, such that $|C| = k$ and every $F \in \mathcal{F}$ contains a point from C . The *piercing number* of \mathcal{F} , denoted by $\tau(\mathcal{F})$, is the smallest k such that \mathcal{F} is pierced by k points.

The classical theorem of Helly [11] states that any family of compact convex sets in \mathbb{R}^d with the $(d + 1, d + 1)$ -property is pierced by one point.

*Department of Mathematics, Iowa State University, USA. Email: zerbib@iastate.edu. The author was supported by NSF grant DMS-195392.

Hadwiger and Debrunner [10] considered the more general problem of studying the piercing numbers of families of compact, convex sets in \mathbb{R}^d that satisfy the (p, q) -property. They proved that whenever $p \geq q \geq d + 1$ and $p(d - 1) < d(q - 1)$, any family of compact, convex sets in \mathbb{R}^d is pierced by $p - q + 1$ points. They also conjectured that for every $p \geq q \geq d + 1$ there exists a constant $c = c(p, q; d)$ such that any family of compact, convex sets in \mathbb{R}^d with the (p, q) property is pierced by c points. This conjecture was proved by Alon and Kleitman in 1992 [1].

It is known that such a constant c does not exist when $q \leq d$. For example, when $d = q = 2$, a family of pairwise intersecting line segments in general position in the plane has piercing number at least $|\mathcal{F}|/2$, which is not bounded by a constant. However, such a constant may exist if the family is further restricted. For example, Danzer [3] showed that a family of disks in the plane with the $(2, 2)$ property has piercing number at most 4; another example is a well-known conjecture of Wegner from 1965 [22], asserting that a family of axis-parallel rectangles in the plane with the $(p + 1, 2)$ property is pierced by $2p$ points.

The question of finding or improving bounds on the piercing numbers in general families of convex sets in \mathbb{R}^d with the (p, q) property for $q \geq d + 1$, or in restricted families when q may be smaller than $d + 1$, has received great attention over the years, see e.g., [6, 14, 16, 18] for general families, and [2, 9, 13, 17] for restricted families.

In this note we give a new bound for the piercing numbers in general families \mathcal{F} of compact convex sets in \mathbb{R}^2 with the $(p + 1, 2)$ property (namely, when $q = d = 2$), where the bound is given in terms of the matching number of the family $\mathcal{F}' = \{S \cap S' \mid S, S' \in \mathcal{F}\}$ of pairwise intersections in \mathcal{F} .

An *isolated set* is a set in \mathcal{F} that does not intersect any other set in \mathcal{F} .

Theorem 1.1. *Let p, r be positive integers. Let \mathcal{F} be a finite family of compact convex sets in \mathbb{R}^2 with no isolated sets, and let \mathcal{F}' be the family of its pairwise intersections. If \mathcal{F} has the $(p + 1, 2)$ property and \mathcal{F}' has the $(r + 1, 2)$ property then $\tau(\mathcal{F}) \leq (\lfloor \frac{r}{2} \rfloor^2 + \lfloor \frac{r}{2} \rfloor)p$ when $r \geq 2$, and $\tau(\mathcal{F}) \leq p$ otherwise.*

Note that the condition that \mathcal{F} has no isolated sets does not pose a significant restriction on the families \mathcal{F} we may consider in the theorem, as any isolated set would add exactly 1 to both the matching number and the piercing number of \mathcal{F} .

When $p = r = 1$, Theorem 1.1 implies Helly's theorem in the plane, as the condition that both \mathcal{F} and \mathcal{F}' have the $(2, 2)$ property is equivalent to the condition that \mathcal{F} has the $(3, 3)$ property, by Radon's theorem [20].

Moreover, if $p = 1$ and $r \geq 2$, then the conditions \mathcal{F} has the $(p + 1, 2)$ property and \mathcal{F}' has the $(r + 1, 2)$ property imply that \mathcal{F} has the $(t, 3)$ property for some $t \leq \sqrt{5r}$. Thus, by the Alon-Kleitman theorem, there exists some constant $c = c(t, 3; 2)$ such that $\tau(\mathcal{F}) \leq c$. The current best known bound on this constant, proved in [14], is $c(t, 3; 2) \leq O(t^4)$. Theorem 1.1 gives an improvement in the constants, under the stronger condition that \mathcal{F}' has the $(r + 1, 2)$ property.

Similarly, for any $p, r \geq 2$, the conditions \mathcal{F} has the $(p + 1, 2)$ property and \mathcal{F}' has the $(r + 1, 2)$ property imply that \mathcal{F} has the $(t, 4)$ property, for some $t \leq 2r + p + 1$. Indeed, let \mathcal{T} be a subfamily of \mathcal{F} of size $2r + p + 1$. At each step pick two sets that intersect and remove them, until less than $p + 1$ sets remain. We get $r + 1$ intersecting pairs, and by the $(r + 1, 2)$ property of \mathcal{F}' some two of them intersect. The bound proven in [14] then implies $\tau(\mathcal{F}) \leq O((2p + r)^3)$. Our bound provides an improvement in the constant, or when $p \gg r$ or $r \gg p$, under the stronger conditions of Theorem 1.1.

The proof of Theorem 1.1 follows from a bound on the line-piercing numbers in families of convex sets in the plane. We say that a family \mathcal{F} of sets in \mathbb{R}^d is *pierced by k lines*, if there exist k lines in \mathbb{R}^d whose union intersects every set in \mathcal{F} . The *line-piercing number* of \mathcal{F} is the smallest k such that \mathcal{F} is pierced by k lines. We say that \mathcal{F} has *property $T(r)$* if every r or fewer sets in \mathcal{F} are pierced by a line.

The problem of bounding the line-piercing numbers of families of convex sets in the plane with the $T(r)$ property has been studied since the 1940's. In 1969 Eckhoff [4] proved that if a family of compact convex sets has the $T(k)$ property, for $k \geq 4$, then it is pierced by two lines. In 1993 Eckhoff [5] proved that a family of compact convex sets with the $T(3)$ property is pierced by 4 lines, and conjectured that this bound can be improved to 3. This conjecture was proved in [19].

If \mathcal{F} is a finite family of compact convex sets in \mathbb{R}^2 with the $(p + 1, 2)$ property, then \mathcal{F} is pierced by p lines; this follows from projecting \mathcal{F} onto a line and using a theorem of Gallai (Theorem 1.6 below), asserting that the matching and piercing numbers in families of intervals in \mathbb{R} are equal. Here we obtain a tight bound on the line-piercing numbers of such families:

Theorem 1.2. *If \mathcal{F} is a finite family of compact convex sets in \mathbb{R}^2 with the $(p + 1, 2)$ property then it is pierced by $\lfloor \frac{p}{2} \rfloor + 1$ lines.*

The bound in Theorem 1.2 is tight by the following example. Let \mathcal{F} be the family of edges of a regular $(2p + 1)$ -gon. Then \mathcal{F} has the $(p + 1)$ -property, and since every line in \mathbb{R}^2 intersects at most 4 elements of \mathcal{F} , at least $\lceil \frac{2p+1}{4} \rceil = \lfloor \frac{p}{2} \rfloor + 1$ lines are needed to pierce \mathcal{F} .

We shall also prove the following colorful version of Theorem 1.2:

Theorem 1.3. *Suppose $\mathcal{F}_1, \dots, \mathcal{F}_{2k}$ are finite families of compact convex sets in \mathbb{R}^2 , such that for any choice of sets $F_i \in \mathcal{F}_i$, $i \in [2k]$, some two sets among F_1, \dots, F_{2k} intersect. Then there exists $i \in [2k]$ such that \mathcal{F}_i is pierced by k lines.*

The proof of Theorem 1.2 uses the topological KKM theorem, due to Knaster, Kuratowski and Mazurkiewicz [15]. Let

$$\Delta_k = \left\{ (x_1, \dots, x_{k+1}) \mid \sum_{i=1}^{k+1} x_i = 1, x_i \geq 0 \text{ for all } i \in [k+1] \right\}$$

be the standard k -dimensional simplex in \mathbb{R}^{k+1} . For a face σ of Δ_k , we write $i \in \sigma$ if σ contains the i -th vertex of Δ_k , namely the point (x_1, \dots, x_{k+1}) with $x_i = 1$ and $x_j = 0$ for $i \neq j$. Note that if $x_i > 0$ for some point $(x_1, \dots, x_{k+1}) \in \sigma$ then $i \in \sigma$.

Theorem 1.4 (The KKM Theorem). *If open subsets A_1, \dots, A_{k+1} of the k -dimensional simplex Δ_k satisfy $\sigma \subseteq \bigcup_{i \in \sigma} A_i$ for every face σ of Δ_k , then $\bigcap_{i=1}^{k+1} A_i \neq \emptyset$.*

A family (A_1, \dots, A_{k+1}) of subsets of Δ_k satisfying the conditions of Theorem 1.4 is called a *KKM cover* of Δ_k . Let S_n be the group of permutations of the elements in $[n] = \{1, 2, \dots, n\}$. The proof of Theorem 1.3 will use a colorful generalization of the KKM theorem, due to Gale [7].

Theorem 1.5 (The colorful KKM Theorem). *If $(A_1^i, \dots, A_{k+1}^i), i \in [k+1]$, are $k+1$ KKM covers of Δ_k , then there exists a permutation $\pi \in S_{k+1}$ such that $\bigcap_{i=1}^{k+1} A_i^{\pi(i)} \neq \emptyset$.*

For the proof of Theorem 1.1 we will also use the following theorem of Gallai [8], which is a special case of the theorem of Hadwiger and Debrunner mentioned above.

Theorem 1.6. *If \mathcal{F} a finite family of compact intervals in \mathbb{R} with the $(p + 1, 2)$ property, then $\tau(\mathcal{F}) \leq p$.*

Finally, we will use a theorem bounding the piercing numbers in families of d -intervals. A (*separated*) d -interval is a union of at most d compact (possibly empty) intervals, one on each of d copies of \mathbb{R} . The following was proved by Tardos [21] for $d = 2$ and by Kaiser [12] for all $d \geq 2$.

Theorem 1.7. *Let $d \geq 2$. If \mathcal{F} a finite family of compact d -intervals with the $(p + 1, 2)$ property, then $\tau(\mathcal{F}) \leq (d^2 - d)p$.*

2 Proofs of Theorems 1.2 and 1.3

Proof of Theorem 1.2. Since \mathcal{F} is finite and the sets in it are compact, we may assume that all the sets in \mathcal{F} are contained in the open unit disk D . Let $f(t)$ be a parameterization of $U = \partial D$ defined by $f(t) = (\cos(2\pi t), \sin(2\pi t))$.

Let $k = \lfloor \frac{p}{2} \rfloor$. A point $x = (x_1, \dots, x_{2k+2}) \in \Delta_{2k+1}$ corresponds to $2k + 2$ points on U , given by $f_i(x) = f(\sum_{j=0}^i x_j)$ for $0 \leq i \leq 2k + 1$, where $x_0 = 0$. For $i = 0, \dots, k$, let $\ell_i(x) = \ell_{i+k+1}(x)$ be the line segment connecting $f_i(x)$ and $f_{i+k+1}(x)$. Let $L(x) = \bigcup_{i=0}^k \ell_i(x)$. For $i \in [2k + 2]$, let $P_i(x)$ be the open region bounded by the lines $\ell_i(x)$ and $\ell_{i-1}(x)$ and the arc between $f_{i-1}(x)$ and $f_i(x)$. Further, for $i \in [2k + 2]$, define recursively the regions $T_i(x)$ as follows: $T_1(x) = P_1(x)$, and $T_i(x) = P_i(x) \setminus \bigcup_{j=1}^{i-1} T_j(x)$ for all $2 \leq i \leq 2k + 2$. Finally, let $R_i(x) = T_i(x) \setminus L(x)$. Note that since $L(x) \cup (\bigcup_{i=1}^{2k+2} P_i(x)) = D - U$, we have $L(x) \cup (\bigcup_{i=1}^{2k+2} R_i(x)) = D - U$. Moreover, the regions $R_1(x), \dots, R_{2k+2}(x)$ are pairwise disjoint and open (see Figure 1). Notice that $R_i(x) = \emptyset$ when $x_i = 0$, because in this case the arc between $f_{i-1}(x)$ and $f_i(x)$ is of length 0, and thus $P_i(x) = \emptyset$.

For every $i \in [2k + 2]$, define sets $A_i \subseteq \Delta_{2k+1}$ as follows: $x \in A_i$ if and only if there exists a set $S \in \mathcal{F}$ so that S is contained in $R_i(x)$.

Assume for contradiction that $\Delta_{2k+1} = \bigcup_{i=1}^{2k+2} A_i$. We claim that the KKM conditions hold for A_1, \dots, A_{2k+2} . First, the sets A_i are open because the sets in \mathcal{F} are closed. Moreover, if $x \in \sigma$ for some face σ of Δ_{2k+1} , then for any $i \notin \sigma$ we have $x_i = 0$ and thus $R_i(x) = \emptyset$, so $R_i(x)$ cannot contain a set of \mathcal{F} . Therefore $x \notin A_i$ when $i \notin \sigma$. Since $\Delta_{2k+1} = \bigcup_{i=1}^{2k+2} A_i$ by our assumption, it follows that $x \in A_i$ for some $i \in \sigma$, showing $\sigma \subseteq \bigcup_{i \in \sigma} A_i$. Hence, by the KKM theorem, there exists $x \in \bigcap_{i=1}^{2k+2} A_i$, and thus for every $i \in [2k + 2]$




Figure 1: The regions $R_i(x)$

there exists a set $F_i \in \mathcal{F}$ with $F_i \subset R_i(x)$. But then the sets F_1, \dots, F_{2k+2} are pairwise disjoint, and since $2k + 2 \geq p + 1$, this is a contradiction to the $(p + 1, 2)$ property of \mathcal{F} .

We conclude that there exists $x = (x_1, \dots, x_{2k+2}) \in \Delta_{2k+1} \setminus (\bigcup_{i=1}^{2k+2} A_i)$. Since $x \notin A_i$, there is no set in \mathcal{F} lying in $R_i(x)$, for all $i \in [2k + 2]$. But then $L(x) \cup (\bigcup_{i=1}^{2k+2} R_i(x)) = D - U$ and the fact that all the sets in \mathcal{F} lie in $D - U$ entail that every set in \mathcal{F} intersects one of the $k + 1$ lines $\ell_i(x)$, $0 \leq i \leq k$. This completes the proof of the theorem. \square

Proof of Theorem 1.3. The proof follows the lines of the previous proof, with some modifications. Again, we may assume that all the sets in $\bigcap_{j=1}^{2k} \mathcal{F}_j$ are contained in the open unit disk D . Let $f(t) = (\cos(2\pi t), \sin(2\pi t))$.

A point $x = (x_1, \dots, x_{2k}) \in \Delta_{2k-1}$ corresponds to $2k$ points on U , given by $f_i(x) = f(\sum_{j=0}^i x_j)$ for $0 \leq i \leq 2k - 1$, where $x_0 = 0$. For $i = 0, \dots, k - 1$, let $\ell_i(x) = \ell_{i+k}(x)$ be the line segment connecting $f_i(x)$ and $f_{i+k}(x)$. Let $L(x) = \bigcup_{i=0}^{k-1} \ell_i(x)$. As before, for $i \in [2k]$, let $P_i(x)$ be the open region bounded by the lines $\ell_i(x)$ and $\ell_{i-1}(x)$ and the arc between $f_{i-1}(x)$ and $f_i(x)$. Further, for $i \in [2k]$, define recursively the regions $T_i(x)$ as before: $T_1(x) = P_1(x)$, and $T_i(x) = P_i(x) \setminus \bigcup_{j=1}^{i-1} T_j(x)$ for all $2 \leq i \leq 2k$. Finally, let $R_i(x) = T_i(x) \setminus L(x)$. Note that $L(x) \cup (\bigcup_{i=1}^{2k} R_i(x)) = D - U$, the regions $R_1(x), \dots, R_{2k}(x)$ are pairwise disjoint and open, and $R_i(x) = \emptyset$ when $x_i = 0$.

For every $i, j \in [2k]$, define sets $A_i^j \subseteq \Delta_{2k-1}$ as follows: $x \in A_i^j$ if and only if there exists a set $S \in \mathcal{F}_j$ so that S is contained in $R_i(x)$.

Assume for contradiction that for every $j \in [2k]$ we have $\Delta_{2k-1} = \bigcup_{i=1}^{2k} A_i^j$. Then by the same argument as in the previous proof, for every $j \in [2k]$, the family (A_1^j, \dots, A_{2k}^j) is a KKM cover of Δ_{2k-1} .

Hence, by the colorful KKM theorem, there exists $\pi \in S_{2k}$ such that $\bigcap_{i=1}^{2k} A_i^{\pi(i)} \neq \emptyset$. Let $x \in \bigcap_{i=1}^{2k} A_i^{\pi(i)}$. Then for every $i \in [2k]$ there exists a set $F_i \in \mathcal{F}_{\pi(i)}$ satisfying $F_i \subset R_i(x)$. But then the sets F_1, \dots, F_{2k} are pairwise disjoint, violating the condition of the theorem.

We conclude that there exists $j \in [2k]$ and $x \in \Delta_{2k-1}$ such that $x = (x_1, \dots, x_{2k}) \in \Delta_{2k+1} \setminus (\bigcup_{i=1}^{2k} A_i^j)$. Like before, this means that \mathcal{F}_j is pierced by the k lines $\ell_i(x)$, $0 \leq i \leq k-1$. This completes the proof of the theorem. \square

3 Proof of Theorem 1.1

We apply Theorem 1.2 with the family \mathcal{F}' of pairwise intersections. Let $k = \lfloor \frac{r}{2} \rfloor$. By Theorem 1.2, there exist $k+1$ lines ℓ_0, \dots, ℓ_k in \mathbb{R}^2 piercing \mathcal{F}' . Write $L = \bigcup_{i=0}^k \ell_i$.

For $S \in \mathcal{F}$, let $\bar{S} = S \cap L$. Then $\bar{\mathcal{F}} = \{\bar{S} \mid S \in \mathcal{F}\}$ is a family of $(k+1)$ -intervals. We claim that $\bar{\mathcal{F}}$ has the $(p+1, 2)$ property. Indeed, suppose for contradiction that there exist pairwise disjoint sets $\bar{S}_1, \dots, \bar{S}_{p+1} \in \bar{\mathcal{F}}$. Since \mathcal{F} has the $(p+1, 2)$ property, there exist integers $1 \leq q < q' \leq p+1$ so that $S_q \cap S_{q'} \neq \emptyset$. But then $S_q \cap S_{q'} \in \mathcal{F}'$, which means that there exists some line ℓ_i , so that $S_j \cap S_k \cap \ell_i \neq \emptyset$. However, this contradicts $\bar{S}_q \cap \bar{S}_{q'} = \emptyset$.

If $k = 0$ then $r = 1$ and $\bar{\mathcal{F}}$ is a family of intervals. Therefore, by theorem 1.6, we have $\tau(\bar{\mathcal{F}}) \leq p$. If $k \geq 1$, then by theorem 1.7 we have $\tau(\bar{\mathcal{F}}) \leq ((k+1)^2 - (k+1))p = (\lfloor \frac{r}{2} \rfloor^2 + \lfloor \frac{r}{2} \rfloor)p$.

Finally, we claim that $\tau(\mathcal{F}) \leq \tau(\bar{\mathcal{F}})$, which will conclude the proof of the theorem. Indeed, let $S \in \mathcal{F}$. Since S is not isolated, there exists some $S' \in \mathcal{F}$ with $S \cap S' \neq \emptyset$, and moreover, there exists some $i \in \{0, \dots, k\}$ with $S \cap S' \cap \ell_i \neq \emptyset$. This implies that $\bar{S} \neq \emptyset$ for all $S \in \mathcal{F}$, entailing the claim. \square

Acknowledgment

The author is grateful to Joe Miller for his help with finding the example after Theorem 1.2. The author is also grateful to Daniel McGinnis for useful comments and for his help with generating the figures.

References

- [1] N. Alon and D. Kleitman, Piercing convex sets and the Hadwiger-Debrunner (p, q) -problem, *Adv. Math.* (1992), 103–112.
- [2] M. Chudnovsky, S. Spirkl and S. Zerbib, Piercing axis-parallel boxes. *Electronic Journal of Combinatorics*, Volume 25, Issue 1 (2018).
- [3] L. Danzer, Zur Lösung des Gallaischen Problems über Kreisscheiben in der Euklidischen Ebene, *Studia Sci. Math. Hungar.* 21 (1986), no. 1–2, 111–134.
- [4] J. Eckhoff, *Transversalenprobleme vom Gallaischen Typ*, Dissertation, Universität Göttingen, 1969.
- [5] J. Eckhoff, A Gallai-type transversal problem in the plane. *Discrete Comput. Geom.* 9 (1993), no. 2, 203–214.
- [6] J. Eckhoff, A survey of the Hadwiger-Debrunner (p, q) -problem, *Discrete Comput. Geom.* 347–377, *Algorithms Combin.*, 25, Springer, Berlin, (2003).
- [7] D. Gale. Equilibrium in a discrete exchange economy with money. *Internat. J. Game Theory*, 13(1):61–64, 1984.
- [8] T. Gallai, Über extreme Punktund Kantenmengen (in German), *Ann. Univ. Sci. Budapest, Eötvös Sect. Math.* 2 (1959), 133–138.
- [9] S. Gao and S. Zerbib, The $(2,2)$ and $(4,3)$ properties in families of fat sets in the plane. *SIAM Journal of Discrete Math*, 33 (2019), no. 3, 1326–1337.
- [10] H. Hadwiger and H. Debrunner, Über eine Variante zum Hellyschen Satz, *Arch. Math. (Basel)* 8 (1957), 309–313.
- [11] E. Helly, Über Mengen konvexer Körper mit gemeinschaftlichen Punkten, *Jahresber. Deutsch. Math. Verein.* 32 (1923), 175–176.
- [12] T. Kaiser, Transversals of d -intervals, *Discrete. Comput. Geom.* 18, 2, (1997), 195–203.

- [13] R. N. Karasev, Piercing families of convex sets with the d -intersection property in \mathbb{R}^d , *Discrete. Comput. Geom.* 39 (2008), no. 4, 766–777.
- [14] C. Keller, S. Smorodinsky and G. Tardos, On max-clique for intersection graphs of sets and the Hadwiger-Debrunner numbers, *Proceedings of the 28th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA 2017)*.
- [15] B. Knaster, C. Kuratowski, and S. Mazurkiewicz. Ein beweis des fixpunktsatzes für n -dimensionale simplexe. *Fund. Math.*, 14(1):132–137, 1929.
- [16] D. J. Kleitman, A. Gyárfás and G. Tóth, Convex sets in the plane with three of every four meeting, *Combinatorica* 21(2) (2001), 221–232.
- [17] J. Kynčl and M. Tancer, The maximum piercing number for some classes of convex sets with the $(4, 3)$ -property. *Electron. J. Combin.* 15 (2008), no. 1.
- [18] D. McGinnis, A Family of Convex Sets in the Plane Satisfying the $(4, 3)$ -Property can be Pierced by Nine Points. *Discrete. Comput. Geom.* 68, 860–880 (2022).
- [19] D. McGinnis and S. Zerbib, Line transversals in families of connected sets the plane. To appear in *SIAM Journal of Discrete Math.*
- [20] J. Radon, Mengen konvexer Körper, die einen gemeinsamen Punkt enthalten, *Mathematische Annalen*, 83 (1–2): 113–115 (1921).
- [21] G. Tardos, Transversals of 2-intervals, a topological approach, *Combinatorica* 15, 1 (1995), 123–134.
- [22] G. Wegner. Über eine kombinatorisch-geometrische Frage von Hadwiger und Debrunner. *Israel Journal of Mathematics*, 3(4):187–198, 1965.