

## SEPARATION OF HOMOGENEOUS CONTINUA

VESKO VALOV

ABSTRACT. We prove that any region  $\Gamma$  in a homogeneous  $n$ -dimensional and locally compact separable metric space cannot be irreducibly separated by a closed  $(n - 1)$ -dimensional subset  $C$  with the following property:  $C$  is acyclic in dimension  $n - 1$  and there is a point  $b \in C \cap \Gamma$  having a local base  $\mathcal{B}_C^b$  in  $C$  such that the boundary of each  $U \in \mathcal{B}_C^b$  is acyclic in dimension  $n - 2$ . The acyclicity means triviality of the corresponding Čech cohomology groups. This implies all known results concerning the separation of regions in homogeneous spaces.

## 1. INTRODUCTION

By a *space* we mean a locally compact separable metric space, and *maps* are continuous mappings. We also consider reduced in dimension zero Čech cohomology groups  $H^n(X; G)$  with coefficients from an Abelian group  $G$ . If  $G$  is the group of the integers  $\mathbb{Z}$ , we simply write  $H^n(X)$ . Recall that a space  $X$  is *separated by a set*  $C \subset X$  if  $C$  is closed in  $X$  and  $X \setminus C$  is the union of two disjoint open subsets  $G_1, G_2$  of  $X$ . When  $C$  is the intersection of the closures  $\overline{G_1} \cap \overline{G_2}$ , we say that  $C$  is an *irreducible separator*. A *partition* between two disjoint closed sets  $A, B$  in  $X$  is a closed set  $P$  such that  $X \setminus P$  is the union of two open disjoint sets  $U, V$  in  $X$  such that  $A \subset U$  and  $B \subset V$ . By  $\dim X$  we denote the covering dimension of  $X$ , and  $\dim_G X$  stands for the cohomological dimension of  $X$  with respect to a group  $G$ . Also, the boundary of a given set  $U \subset X$  in  $X$  is denoted by  $\text{bd}U$ ; if  $U \subset C \subset X$ , then  $\text{bd}_C U$  denotes the boundary of  $U$  in  $C$ .

One of the first results concerning the separation of homogenous metric spaces is the celebrated theorem of Krupski [12], [13] stating that every region in an  $n$ -dimensional homogeneous space cannot be separated by a subset of dimension  $\leq n - 2$ . Kallipoliti and Papasoglu [8] established that any locally connected, simply connected, homogeneous

---

2020 *Mathematics Subject Classification*. Primary 55M10; Secondary 54F45.

*Key words and phrases*. acyclic partitions, cohomological dimension, homogeneous spaces.

The author was partially by NSERC Grant 261914-19.

metric continuum cannot be separated by arcs (according to Krupski's theorem, mentioned above, the Kallipoliti-Papasoglu result is interesting for spaces of dimension two). van Mill and the author [17] proved that the Kallipoliti-Papasoglu theorem remains true without simply connectedness, but requiring strong local homogeneity instead of homogeneity. Recently, Karashev-Valov [9] settle the two-dimensional case by proving that any region of a homogeneous space cannot be separated by an arc.

In the present paper we establish the following theorem which captures all mentioned above results:

**Theorem 1.1.** *Let  $\Gamma$  be a region in a homogeneous space  $X$  with  $\dim X = n \geq 1$ . Then  $\Gamma$  cannot be irreducibly separated by any closed set  $C \subset X$  with the following property:*

- (i)  $\dim C \leq n - 1$  and  $H^{n-1}(C) = 0$ ;
- (ii) *There is a point  $b \in C \cap \Gamma$  having a local base  $\mathcal{B}_C^b$  in  $C$  such that  $H^{n-2}(\text{bd}_C U) = 0$  for every  $U \in \mathcal{B}_C^b$ .*

**Remark 1.2.** According to [20, Corollary 1.6], if  $X$  in Theorem 1.1 is a compactum with  $H^n(X) \neq 0$ , then  $X$  is not separated by any  $C$  satisfying condition (i).

Since  $H^{k+1}(Y) = 0$  for any  $k$ -dimensional space  $Y$ , we have the following fact: If  $\dim Y \leq n - 2$ , then  $H^{n-1}(Y) = 0$  and every  $x \in Y$  has a basis of neighborhoods  $U$  with  $H^{n-2}(\text{bd}U) = 0$ . Moreover, any  $(n - 2)$ -dimensional separator contains a closed subset which is an irreducible  $(n - 2)$ -dimensional separator. Therefore, Theorem 1.1 implies directly that any region in a homogeneous  $n$ -dimensional space cannot be separated by a subset of dimension  $\leq n - 2$ . Similar arguments show that if  $G$  is any countable Abelian group, then any homogeneous connected space of cohomological dimension  $\dim_G X \leq n$  cannot be separated by a closed subset of dimension  $\dim_G \leq n - 2$  (this fact was established by different methods in [10]).

If a region  $\Gamma$  in a two-dimensional homogeneous space is separated by an arc  $C$ , then there is a closed  $C' \subset C$  irreducibly separating  $\Gamma$ , see [17]. Then  $H^1(C') = 0$  and the point  $b = \max\{x : x \in C'\}$  satisfies condition (ii) from Theorem 1.1.

Theorem 1.1 follows from its cohomological version, Theorem 3.2, which is established in Section 3. Section 2 contains some definitions and preliminary results. In the final Section 4 we present an analogue of Mazurkiewicz' theorem [16] that any  $(n - 2)$ -dimensional subset of  $\mathbb{R}^n$  cannot cut  $\mathbb{R}^n$ .

## 2. DEFINITIONS AND PRELIMINARY RESULTS

Recall that for any nontrivial Abelian group  $G$  the Čech cohomology group  $H^n(X; G)$  is isomorphic to the group  $[X, K(G, n)]$  of pointed homotopy classes of maps from  $X$  to  $K(G, n)$ , where  $K(G, n)$  is a  $CW$ -complex of type  $(G, n)$ , see [6]. It is also well known that the circle group  $\mathbb{S}^1$  is a space of type  $(\mathbb{Z}, 1)$ . The cohomological dimension  $\dim_G(X)$  is the largest number  $n$  such that there exists a closed subset  $A \subset X$  with  $H^n(X, A; G) \neq 0$ . Equivalently, for a metric space  $X$  we have  $\dim_G X \leq n$  if and only if for any closed pair  $A \subset B$  in  $X$  the homomorphism  $j_{X,A}^n : H^n(B; G) \rightarrow H^n(A; G)$ , generated by the inclusion  $A \hookrightarrow B$ , is surjective, see [2]. This means that every map from  $A$  to  $K(G, n)$  can be extended over  $B$ . For every  $G$  we have  $\dim_G X \leq \dim_{\mathbb{Z}} X \leq \dim X$ , and  $\dim_{\mathbb{Z}} X = \dim X$  in case  $\dim X < \infty$  [15] (on the other hand, there is an infinite-dimensional compactum  $X$  with  $\dim_{\mathbb{Z}} X = 3$ , see [1]).

Suppose  $(K, A)$  is a pair of compact sets in a space  $X$  with  $A \subset K$ . We say that  $K$  is an  $n$ -cohomology membrane spanned on  $A$  for an element  $\gamma \in H^n(A; G)$  if  $\gamma$  is not extendable over  $K$ , but it is extendable over every proper closed subset of  $K$  containing  $A$ . Here,  $\gamma \in H^n(A; G)$  is *extendable over  $K$*  means that  $\gamma$  is contained in the image  $j_{K,A}^n(H^n(K; G))$ . Concerning extendability, we are using the following simple fact:

**Lemma 2.1.** *Let  $A, B$  be closed sets in  $X$  with  $X = A \cup B$ . Then  $\gamma \in H^n(A; G)$  is extendable over  $X$  if and only if  $j_{A,\Gamma}^n(\gamma)$  is extendable over  $B$ , where  $\Gamma = A \cap B$ .*

*Proof.* This follows from the Mayer-Vietoris exact sequence

$$H^n(X; G) \xrightarrow{\varphi^n} H^n(A; G) \oplus H^n(B; G) \xrightarrow{\psi^n} H^n(\Gamma; G),$$

where  $\varphi^n(\gamma) = (j_{X,A}^n(\gamma), j_{X,B}^n(\gamma))$  and  $\psi^n(\gamma_1, \gamma_2) = j_{A,\Gamma}^n(\gamma_1) - j_{B,\Gamma}^n(\gamma_2)$ . Indeed, suppose  $\gamma_\Gamma = j_{A,\Gamma}^n(\gamma)$  is extendable over  $B$ . So, there is  $\alpha \in H^n(B; G)$  with  $j_{B,\Gamma}^n(\alpha) = \gamma_\Gamma$ . Then,  $\psi^n(\gamma, \alpha) = 0$ , which implies the existence of  $\beta \in H^n(X; G)$  such that  $\varphi^n(\beta) = (\gamma, \alpha)$ . This yields  $j_{X,A}^n(\beta) = \gamma$ . Hence,  $\gamma$  is extendable over  $X$ .

To prove the other implication, suppose  $j_{X,A}^n(\beta) = \gamma$  for some  $\beta \in H^n(X; G)$ , and let  $\alpha = j_{X,B}^n(\beta)$ . Then  $\psi^n(\gamma, \alpha) = 0$ , which means that  $j_{B,\Gamma}^n(\alpha) = j_{A,\Gamma}^n(\gamma)$ . Therefore,  $j_{A,\Gamma}^n(\gamma)$  is extendable over  $B$ .  $\square$

We use below the following notation: Suppose  $A$  is partition in a space  $Z$  between two closed disjoint sets  $P, Q \subset Z$ . Then there are two

open disjoint subset  $W_P, W_Q$  of  $Z$  containing  $P$  and  $Q$ , respectively, such that  $Z \setminus A = W_P \cup W_Q$ . Then we denote  $\Lambda_P = W_P \cup A$  and  $\Lambda_Q = W_Q \cup A$ .

**Lemma 2.2.** *Let  $X$  be a homogeneous space with  $\dim_G X = n$  and  $x \in \Gamma$  be a fixed point, where  $\Gamma \subset X$  is open. Then there exist a compact set  $M \subset \Gamma$  with  $x \in M$  and a local basis  $\mathcal{B}_x$  in  $X$  with the following property: For every  $U, V \in \mathcal{B}_x$  with  $\bar{U} \subset V$  there exist a nontrivial  $\gamma_{U,V} \in H^{n-1}(M(U, V); G)$ , where  $M(U, V) = (\bar{V} \setminus U) \cap M$ , such that:*

- (i) *There is a compactum  $K_U \subset \bar{U}$  such that  $K_U$  is an  $(n-1)$ -cohomology membrane for  $\gamma_U = j_{M(U,V), M \cap \text{bd}\bar{U}}^{n-1}(\gamma_{U,V})$  spanned on  $M \cap \text{bd}\bar{U}$ ;*
- (ii)  *$\gamma_V = j_{M(U,V), M \cap \text{bd}\bar{V}}^{n-1}(\gamma_{U,V})$  is not extendable over  $M \cap \bar{V}$ ;*
- (iii) *If  $A$  is a partition in  $M(U, V)$  between  $M \cap \text{bd}\bar{U}$  and  $M \cap \text{bd}\bar{V}$ , then  $\gamma_A = j_{M(U,V), A}^{n-1}(\gamma_{U,V})$  is extendable over  $\Lambda_{\text{bd}\bar{U}} \cup B$  for any proper closed set  $B \subset K_U$ , but not extendable over  $\Lambda_{\text{bd}\bar{U}} \cup \bar{U}$ .*

*Proof.* Since  $X$  is homogeneous,  $\dim_G \Gamma = n$ . Hence,  $\Gamma$  is a countable union of compact sets and at least one of them should be of dimension  $\dim_G = n$  (otherwise, by the countable sum theorem for  $\dim_G$ , we would have  $\dim_G \Gamma \leq n-1$ ). Therefore, there exists a compactum  $Y \subset \Gamma$  with  $\dim_G Y = n$ . This implies the existence of a proper closed set  $F \subset Y$  and  $\gamma \in H^{n-1}(F; G)$  such that  $\gamma$  is not extendable over  $Y$ . Using the continuity of Čech cohomology [19], we can apply Zorn's lemma to conclude there exists a minimal compact set  $M \subset Y$  containing  $F$  such that  $\gamma$  is not extendable over  $M$ , but it is extendable over every proper closed subset of  $M$  containing  $F$ . Since  $X$  is homogeneous, we can assume that  $x \in M \setminus F$ . Now, let  $\mathcal{B}_x$  be the family of all open sets  $U$  in  $X$  such that:

$$U = \text{int}\bar{U} \text{ and } \bar{U} \cap F = \emptyset.$$

Suppose  $U, V \in \mathcal{B}_x$  with  $\bar{U} \subset V$ . Then  $M \setminus U$  is a proper closed subset of  $M$  containing  $F$ . Hence, there is  $\gamma' \in H^{n-1}(M \setminus U; G)$  with  $j_{M \setminus U, F}^{n-1}(\gamma') = \gamma$ . Let  $\gamma_{U,V} = j_{M \setminus U, M(U,V)}^{n-1}(\gamma')$ . So,  $\gamma_U = j_{M(U,V), M \cap \text{bd}\bar{U}}^{n-1}(\gamma_{U,V})$  is not extendable over  $M \cap \bar{U}$ , otherwise  $\gamma$  would be extendable over  $M$ , see Lemma 2.1. Let  $K_U$  be a minimal subset of  $\bar{U}$  containing  $M \cap \text{bd}\bar{U}$  such that  $\gamma_U$  is not extendable over  $K_U$ . Then  $\gamma_U$  is extendable over any proper closed subset of  $K_U$ . So,  $K_U$  is an  $(n-1)$ -cohomology membrane for  $\gamma_U$  spanned on  $M \cap \text{bd}\bar{U}$ . Similarly, we can see that  $\gamma_V = j_{M(U,V), M \cap \text{bd}\bar{V}}^{n-1}(\gamma_{U,V})$  is not extendable over  $M \cap \bar{V}$ .

To prove item (iii), suppose  $A$  is a partition in  $M(U, V)$  between  $M \cap \text{bd}\overline{U}$  and  $M \cap \text{bd}\overline{V}$ . Then  $\gamma_A$  is the restriction of  $\gamma(\Lambda_{\text{bd}\overline{U}})$  on  $A$ , where  $\gamma(\Lambda_{\text{bd}\overline{U}}) = j_{M(U,V), \Lambda_{\text{bd}\overline{U}}}^{n-1}(\gamma_{U,V})$ . So,  $\gamma(\Lambda_{\text{bd}\overline{U}})$  is an extension  $\gamma_A$  over the set  $\Lambda_{\text{bd}\overline{U}}$  and, since  $\gamma_U = j_{\Lambda_{\text{bd}\overline{U}}, M \cap \text{bd}\overline{U}}^{n-1}(\gamma(\Lambda_{\text{bd}\overline{U}}))$ ,  $\gamma_A$  is extendable over  $\Lambda_{\text{bd}\overline{U}} \cup B$  for any proper closed set  $B \subset K_U$ . On the other hand,  $\gamma_A$  is the restriction of  $\gamma(\Lambda_{\text{bd}\overline{V}}) = j_{M(U,V), \Lambda_{\text{bd}\overline{V}}}^{n-1}(\gamma_{U,V})$  on  $A$ . This implies that  $\gamma_A$  is not extendable over  $\Lambda_{\text{bd}\overline{U}} \cup \overline{U}$ , otherwise  $\gamma_V$  would be extendable over  $M \cap \overline{V}$ .  $\square$

**Proposition 2.3.** *Let  $A \subset K$  be a compact pair and  $\gamma$  be a nontrivial element of  $H^{n-1}(A; G)$ . Suppose there are closed subsets  $P_1, P_2$  of  $K$  satisfying the following conditions:*

- $P_1 \cup P_2 = K$  and  $P_1 \cap P_2 = C \neq \emptyset$ ;
- $\gamma$  is extendable over  $P_i \cup A$  for each  $i = 1, 2$ , but  $\gamma$  is not extendable over  $K$ .

Then  $H^{n-1}(C, C \cap A; G) \neq 0$ .

*Proof.* Consider the commutative diagram below whose rows are parts of Mayer-Vietoris exact sequences, while the columns are exact sequences for the corresponding couples:

$$\begin{array}{ccc}
 H^{n-1}(K; G) & \xrightarrow{\varphi_K^{n-1}} & H^{n-1}(P_1; G) \oplus H^{n-1}(P_2; G) \\
 \downarrow j_{K,A}^{n-1} & & \downarrow j_{P_1 \oplus P_2}^{n-1} \\
 H^{n-1}(A; G) & \xrightarrow{\varphi_A^{n-1}} & H^{n-1}(A \cap P_1; G) \oplus H^{n-1}(A \cap P_2; G) \\
 \downarrow \partial_{K,A} & & \downarrow \partial_{P_1 \oplus P_2} \\
 H^n(K, A; G) & \xrightarrow{\varphi_{K,A}^n} & H^n(P_1, P_1 \cap A; G) \oplus H^n(P_2, P_2 \cap A; G)
 \end{array}$$

Here, the maps  $j_{P_1 \oplus P_2}^{n-1}$  and  $\partial_{P_1 \oplus P_2}$  are defined by

$$j_{P_1, A \cap P_1}^{n-1} \oplus j_{P_2, A \cap P_2}^{n-1},$$

$$\partial_{P_1 \oplus P_2} = \partial_{P_1, P_1 \cap A} \oplus \partial_{P_2, P_2 \cap A}.$$

Recall also that  $\varphi_K^{n-1} = (j_{K, P_1}^{n-1}, j_{K, P_2}^{n-1})$ , the maps  $\varphi_A^{n-1}$ ,  $\varphi_{K,A}^n$  and  $\varphi_K^n$  are defined similarly.

Denote  $\alpha_i = j_{A, A \cap P_i}^{n-1}(\gamma)$ ,  $i = 1, 2$ . Since  $\gamma$  is extendable over  $A \cup P_i$ , there exists  $\gamma_i \in H^{n-1}(A \cup P_i)$  extending  $\gamma$ , i.e.  $j_{A \cup P_i, A}^{n-1}(\gamma_i) = \gamma$ . Let

$\beta_i = j_{A \cup P_i, P_i}^{n-1}(\gamma_i)$ . It follows from the Mayer-Vietoris exact sequence

$$H^{n-1}(A \cup P_i; G) \rightarrow H^{n-1}(A; G) \oplus H^{n-1}(P_i; G) \rightarrow H^{n-1}(A \cap P_i; G) \rightarrow \dots$$

that  $j_{P_i, A \cap P_i}^{n-1}(\beta_i) = \alpha_i$  for every  $i = 1, 2$ . This implies  $j_{P_1 \oplus P_2}^{n-1}((\beta_1, \beta_2)) = (\alpha_1, \alpha_2)$ . Since the second column is a part of an exact sequence, the last equality yields  $\partial_{P_1 \oplus P_2}(\varphi_A^{n-1}(\gamma)) = 0$ . Hence,  $\varphi_{K,A}^n(\partial_{K,A}(\gamma)) = 0$ . Note that  $\tilde{\gamma} = \partial_{K,A}(\gamma) \neq 0$  because the first column is exact and  $\gamma$  is not extendable over  $K$ .

Finally, since  $\varphi_{K,A}^n(\tilde{\gamma}) = 0$ , Proposition 2.3 follows from the Mayer-Vietoris exact sequence

$$H^{n-1}(C, C \cap A; G) \xrightarrow{\Delta} H^n(K, A; G) \xrightarrow{\varphi_{K,A}^n} H^n(P_1, P_1 \cap A; G) \oplus H^n(P_2, P_2 \cap A; G).$$

□

**Corollary 2.4.** *Let  $X$  be a homogeneous space with  $\dim_G X = n$  and  $C \subset X$  separate  $X$ . If there is an open set  $U$  such that  $C \subset U$  and  $\bar{U}$  is an  $(n-1)$ -cohomology membrane for some  $\gamma \in H^{n-1}(\text{bd}\bar{U}; G)$  spanned on  $\text{bd}\bar{U}$ , then  $H^{n-1}(C; G) \neq 0$ .*

*Proof.* Since  $C \cap \text{bd}\bar{U} = \emptyset$ , the proof follows from Proposition 2.3 and the exact sequence

$$H^{n-2}(C \cap \text{bd}\bar{U}; G) \longrightarrow H^{n-1}(C, C \cap \text{bd}\bar{U}; G) \longrightarrow H^{n-1}(C; G) \rightarrow \dots$$

□

Corollary 2.4 implies the well known fact [7] that  $H^{n-1}(C; G) \neq 0$  for any compact separator  $C$  of  $\mathbb{R}^n$ . Indeed, take any ball  $\mathbb{B}^n$  with  $C \subset \text{int}\mathbb{B}^n$ .

### 3. PROOF OF THEOREM 1.1

We need the following version of Effros' [5] theorem (see [12, Proposition 1.4]):

**Theorem 3.1.** *Let  $X$  be a homogeneous space with a metric  $\rho$ ,  $a \in X$  and  $\varepsilon > 0$ . Then there is  $\delta > 0$  such that for every  $x \in X$  with  $\rho(x, a) < \delta$  there exists a homeomorphism  $h : X \rightarrow X$  with  $h(a) = x$  and  $\rho(h(y), y) < \varepsilon$  for all  $y \in X$ .*

Theorem 1.1 is a particular case of the following theorem when  $G = \mathbb{Z}$ :

**Theorem 3.2.** *Let  $\Gamma$  be a region in a homogeneous space  $X$  with  $\dim_G X = n \geq 1$ , where  $G$  is a countable Abelian group. Then  $\Gamma$  cannot be irreducibly separated by any closed set  $C \subset X$  with the following property:*

- (i)  $\dim_G C \leq n - 1$  and  $H^{n-1}(C; G) = 0$ ;
- (ii) *There is a point  $b \in C \cap \Gamma$  having a local base  $\mathcal{B}_C^b$  in  $C$  such that  $H^{n-2}(\text{bd}_C U; G) = 0$  for every  $U \in \mathcal{B}_C^b$ .*

*Proof.* Let  $C \subset X$  be closed such that  $C \cap \Gamma$  irreducibly separates  $\Gamma$  and satisfies conditions (i) – (ii). Then  $\Gamma \setminus C = G_1 \cup G_2$  with  $C' = \overline{G_1} \cap \overline{G_2} \cap \Gamma \subset C$ , where  $G_1, G_2$  are disjoint open subsets of  $\Gamma$ . By Lemma 2.2, there exist a compactum  $M$  containing  $b$  and a local base  $\mathcal{B}_b$  satisfying the hypothesis of that lemma. Actually, it follows from the proof of Lemma 2.2 that there exists an open neighborhood  $W_0$  of  $b$  in  $\Gamma$  such that any open  $U \subset \Gamma$  with  $b \in U \subset \overline{U} \subset W_0$  belongs to  $\mathcal{B}_b$ . We assume that  $\overline{W_0} \subset \Gamma$  is compact and  $C' \setminus \overline{W_0} \neq \emptyset$ . Let  $U, V \in \mathcal{B}_b$  be any pair with  $\overline{U} \subset V$ . Following the notations from Lemma 2.2, note that the compact set  $M$  is contained in  $\Gamma$ , and observe that  $\dim_G(K_U \setminus \text{bd}\overline{U}) = n$  (otherwise  $\gamma_U$  would be extendable over  $K_U$ ).

*Claim 1.* We can suppose that  $b \in K_U \setminus \text{bd}\overline{U}$  and  $K_U \setminus \text{bd}\overline{U}$  meets both sets  $G_1$  and  $G_2$ .

Indeed, let  $\varepsilon$  be the distance between  $\overline{W_0}$  and  $X \setminus \Gamma$  and  $\delta > 0$  be a number from Theorem 3.1 corresponding to  $\varepsilon$  and the point  $b$ . Choose  $U$  to be so small that its diameter is less than  $\delta$ . Then there is a  $\delta$ -small homeomorphism  $h$  on  $X$  so that  $\overline{h(V)} \subset \Gamma$  and  $b \in h(K_U) \setminus \text{bd}h(\overline{U})$ . Hence, considering the sets  $h^{-1}(U), h^{-1}(V)$  and  $h^{-1}(K_U)$  instead of  $U, V$  and  $K_U$ , we can assume that  $b \in K_U \setminus \text{bd}\overline{U}$ . Since  $\dim_G C \leq n - 1$ ,  $K_U \setminus \text{bd}\overline{U}$  is not contained in  $C$ . So,  $K_U \setminus \text{bd}\overline{U}$  meets at least one  $G_i$ ,  $i = 1, 2$ . If  $K_U \setminus \text{bd}\overline{U}$  intersects only  $G_1$ , then Theorem 3.1 allows us to push  $K_U$  towards  $G_2$  by a small homeomorphism  $h : X \rightarrow X$  such that  $h(K_U \setminus \text{bd}\overline{U}) \cap G_i \neq \emptyset$ ,  $i = 1, 2$ , and  $h(U)$  still contains  $b$ . So, everywhere below we can assume that  $K_U \setminus \text{bd}\overline{U}$  meets both  $G_1$  and  $G_2$ , and  $b \in K_U \setminus \text{bd}\overline{U}$ .

We can take  $U$  so small that there is  $W^* \in \mathcal{B}_C^b$  such that  $\text{bd}_C W^* \subset V \setminus \overline{U}$ . Then the set  $F = M \cap \text{bd}_C W^*$  is a partition in  $C \cap M_{U,V}$  between  $C \cap M \cap \text{bd}\overline{U}$  and  $C \cap M \cap \text{bd}\overline{V}$ . Because  $G$  is countable and  $\dim_G C \leq n - 1$ ,  $C$  has a base  $\mathcal{B}_C$  such that  $\dim_G \text{bd}_C O \leq n - 2$  for every  $O \in \mathcal{B}_C$ , see [3] and [4]. So, we can suppose that  $\dim_G \text{bd}_C W^* \leq n - 2$ . This inequality together with  $H^{n-2}(\text{bd}_C W^*; G) = 0$  implies  $H^{n-2}(F; G) = 0$ . Now, consider the space  $M_{U,V}$  and its closed disjoint sets  $M \cap \text{bd}\overline{V}$ ,  $M \cap \text{bd}\overline{U}$ . By [18, Corollary 3.5], there exists a partition  $T$  in  $M_{U,V}$  between  $M \cap \text{bd}\overline{V}$  and  $M \cap \text{bd}\overline{U}$  such that  $T \cap C \subset F$ . Hence,  $A = T \cup F$  is a partition in  $M_{U,V}$  between  $M \cap \text{bd}\overline{V}$  and  $M \cap \text{bd}\overline{U}$  with  $A \cap C = F$ . Then, by Lemma 2.2,  $\gamma_A = j_{M(U,V),A}^{n-1}(\gamma_{U,V})$  is a nontrivial element which is extendable over  $\Lambda_{\text{bd}\overline{U}} \cup B$  for any proper closed set

$B \subset K_U$ , but not extendable over  $\Lambda_{\text{bd}\bar{U}} \cup \bar{U}$ . There are two possible cases, where  $K' = \Lambda_{\text{bd}\bar{U}} \cup K_U$ :

- (1)  $\gamma_A$  is not extendable over  $K'$ ;
- (2)  $\gamma_A$  is extendable over  $K'$ .

*Claim 2.* Case (1) is not possible.

Let  $C_V = C \cap \bar{V}$  and  $K = K' \cup C_V$ . Since  $\gamma_A$  is not extendable over  $K'$  it is also not extendable over  $K$ . We obviously have  $K = P_1 \cup P_2$  and  $C_V = P_1 \cap P_2$ , where  $P_i = C_V \cup (K \cap \bar{G}_i)$ ,  $i = 1, 2$ . Since each  $K_U \cap \bar{G}_i$  is a proper subset of  $K_U$ ,  $\gamma_A$  is extendable over each of the sets  $A \cup (K \cap \bar{G}_i)$ . Finally, since  $\dim_G C_V \leq n - 1$ ,  $\gamma_A$  is extendable over each of the sets  $A \cup P_i$ . Therefore, we can apply Proposition 2.3 to conclude that  $H^{n-1}(C_V, C_V \cap A) \neq 0$ . On the other hand we have the exact sequence

$$H^{n-2}(C_V \cap A; G) \longrightarrow H^{n-1}(C_V, C_V \cap A; G) \longrightarrow H^{n-1}(C_V; G).$$

To complete the proof of Claim 2, observe that  $C_V \cap A = C \cap A = F$ . Hence,  $H^{n-2}(C_V \cap A; G) = H^{n-2}(F; G) = 0$ . Since  $\dim_G C \leq n - 1$  with  $H^{n-1}(C; G) = 0$  and  $C_V$  is closed in  $C$ ,  $H^{n-1}(C_V; G) = 0$ . Hence,  $H^{n-1}(C_V, C_V \cap A; G) = 0$ , a contradiction.

*Claim 3.* Case (2) is not possible.

Since  $\gamma_A$  is not extendable over  $\Lambda_{\text{bd}\bar{U}} \cup \bar{U}$ , there is a minimal subset  $K_A$  of  $\Lambda_{\text{bd}\bar{U}} \cup \bar{U}$  containing  $K'$  such that  $K_A$  is an  $(n - 1)$ -cohomology membrane for  $\gamma_A$  spanned on  $A$ . Because  $K_A$  contains  $K_U$ ,  $K_A$  meets both  $G_1$  and  $G_2$ . Hence, we can repeat the proof of Claim 2 with  $K_U$  replaced by  $K_A$  and obtain again a contradiction with Proposition 2.3.  $\square$

**Remark 3.3.** (i) The proof of Theorem 3.2 works for arbitrary  $G$  if condition (ii) is replaced by the following one:  $H^{n-2}(\Phi \cap \text{bd}_C U; G) = 0$  for every  $U \in \mathcal{B}_C^b$  and every closed set  $\Phi \subset C$ .

(ii) Moreover, if  $C \cap \Gamma$  has a dense set  $C^*$  of points  $b$  satisfying condition (ii) from Theorem 3.2, then  $C$  does not separate  $\Gamma$ . Indeed, assuming that  $\Gamma \setminus C = G_1 \cup G_2$  with  $G_1, G_2$  disjoint open subsets of  $\Gamma$ , take arbitrary  $b \in \bar{G}_1 \cap \bar{G}_2$ . Following the proof of Theorem 3.2, choose neighborhoods  $U, V$  of  $b$  such that  $\bar{U} \subset V$ ,  $K_U \setminus \text{bd}\bar{U}$  meets both  $G_1$  and  $G_2$ , and  $b \in K_U \setminus \text{bd}\bar{U}$ . Then  $(V \setminus \bar{U}) \cap C$  is open in  $C$  and contains a point  $b' \in C^*$ . Next, we take a neighborhood  $W^*$  of  $b'$  in  $C$  such that  $\text{bd}_C \bar{W}^* \subset V \setminus \bar{U}$ ,  $H^{n-1}(\text{bd}_C \bar{W}^*; G) = 0$  and  $\dim_G \text{bd}_C \bar{W}^* \leq n - 2$ . Further, the proof goes as in Theorem 3.2.



## 4. CUTS

Mazurkiewicz [16] proved that any  $(n - 2)$ -dimensional set  $M \subset \mathbb{R}^n$  does not cut  $\mathbb{R}^n$  (i.e., for every two points from  $\mathbb{R}^n \setminus M$  there is a continuum in  $\mathbb{R}^n \setminus M$  joining both points). In this section we establish an analogue of this theorem for homogeneous spaces.

For any space  $X$  denote by  $\mathcal{H}_X(n - 1, G)$  all closed sets  $F \subset X$  with the following property:  $\dim_G F \leq n - 1$ ,  $H^{n-1}(F; G) = 0$  and  $F$  has a dense set of points  $x \in F$  each having a base  $\mathcal{B}_x$  in  $F$  such that  $H^{n-2}(\text{bd}\bar{U}; G) = 0$ ,  $U \in \mathcal{B}_x$ .

Theorem 4.1 below implies the case of [11, Theorem 4] when the class of all closed subsets  $F \subset X$  with  $\dim_G F \leq n - 2$  is considered.

**Theorem 4.1.** *Let  $\Gamma$  be a locally connected region in a homogeneous space  $X$  with  $\dim_G X = n \geq 1$ , where  $G$  is a countable Abelian group. Then  $\Gamma$  cannot be cut by any set  $M = \bigcup_{i=1}^{\infty} F_i$  such that all  $F_i$  are closed in  $\Gamma$  and  $F_i \in \mathcal{H}_\Gamma(n - 1, G)$ .*

*Proof.* Our proof follows the arguments from the proof of the mentioned above [9, Theorem 4]. Suppose there is an  $F_\sigma$ -subset  $\bigcup_{i=1}^{\infty} F_i$  of  $\Gamma$  cutting  $\Gamma$  with  $F_i \in \mathcal{H}_\Gamma(n - 1, G)$ ,  $i \geq 1$ . Then, by [9, Lemma 3], there exists  $j$  and a region  $\Gamma_1 \subset \Gamma$  such that  $F_j$  does not cut  $\Gamma_1$ . Since  $\Gamma_1$  is locally connected,  $F_j \cap \Gamma_1$  separates  $\Gamma_1$ , see [14, Theorem 1, page 238]. Consider now the set  $Q_j$  of the points  $x \in F_j$  having a base  $\mathcal{B}_x$  in  $F_j$  with  $H^{n-2}(\text{bd}\bar{U}; G) = 0$  for all  $U \in \mathcal{B}_x$ . Since  $Q_j$  is dense in  $F_j$ ,  $Q_j \cap \Gamma_1$  is dense in  $F_j \cap \Gamma_1$ . Then, according to Remark 3.3(ii),  $F_j \cap \Gamma_1$  cannot separate  $\Gamma_1$ , a contradiction.  $\square$

Let us note that Theorem 4.1 is not anymore true without the assumption each  $F_i$  to have a dense set of points  $x \in F_i$  having a base  $\mathcal{B}_x$  in  $F_i$  such that  $H^{n-2}(\text{bd}\bar{U}; G) = 0$ ,  $U \in \mathcal{B}_x$ . Indeed, let  $X = \mathbb{R}^2$  and  $M$  be any line in  $\mathbb{R}^2$ . Then  $M$  is a countable union of segments  $F_i$  and  $M$  cuts  $\mathbb{R}^2$ .

## REFERENCES

- [1] A. Dranishnikov, *On a problem of P.S. Alexandrov*, Math. USSR Sbornik **63** (1988), 412–426.
- [2] J. Dydak, *Cohomological dimension and metrizable spaces*, Trans. Amer. Math. Soc. **337** (1993), 219–234.
- [3] J. Dydak and A. Koyama, *Strong cohomological dimension*, Bull. Polish Acad. Sci. **56** (2008), 193–189.
- [4] J. Dydak and A. Koyama, *Cohomological dimension of locally connected compacta*, Topol. Appl. **113** (2001), 139–150.
- [5] E. Effros, *Transformation groups and  $C^*$ -algebras*, Ann. of Math. **81** (1965), 38–55.

- [6] P. Huber, *Homotopical cohomology and Čech cohomology*, Math. Annalen **144** (1961), 73–76.
- [7] W. Hurewicz and H. Wallman, *Dimension theory*, Princeton University Press, Princeton, 1948.
- [8] M. Kallipoliti and P. Papasoglu, *Simply connected homogeneous continua are not separated by arcs*, Topol. Appl. **154** (2007), 3039–3047.
- [9] A. Karashev and V. Valov, *Homogeneous spaces not separated by arcs*, arXiv:2302.06735v1 [math.GN].
- [10] A. Karashev, P. Krupski, V. Todorov and V. Valov, *Generalized Cantor manifolds and homogeneity*, Houston J. Math. **38** (2012), 583–609.
- [11] P. Krupski and V. Valov, *Mazurkiewicz manifolds and homogeneity*, Rocky Mountain J. Math. **41** (2011), 1933–1938.
- [12] P. Krupski, *Recent results on homogeneous curves and ANR's*, Topology Proc. **16** (1991), 109–118.
- [13] P. Krupski, *Homogeneity and Cantor manifolds*, Proc. Amer. Math. Soc. **109** (1990), 1135–1142.
- [14] K. Kuratowski, *Topology*, vol. II, Academic Press, New York, 1968.
- [15] V. Kuz'minov, *Homological dimension theory*, Russian Math. Surveys **23** (1968), 1–45.
- [16] S. Mazurkiewicz, *Sur le ensembles de dimension faible*, Fund. Math. **13** (1929), 209–217.
- [17] J. van Mill and V. Valov, *Homogeneous continua that are not separated by arcs*, Acta Math. Hung. **157**, (2019), 364–370.
- [18] J. van Mill, *The infinite-dimensional topology of function spaces*, North-Holland Publishing Co., Amsterdam, 2001.
- [19] E. Spanier, *Algebraic Topology*, McGraw-Hill Book Company, 1966.
- [20] V. Valov, *Homogeneous ANR spaces and Alexandroff manifolds*, Topol. Appl. **173** (2014), 227–233.

DEPARTMENT OF COMPUTER SCIENCE AND MATHEMATICS, NIPISSING UNIVERSITY, 100 COLLEGE DRIVE, P.O. BOX 5002, NORTH BAY, ON, P1B 8L7, CANADA

*Email address:* veskov@nipissingu.ca