

ON A BESSEL FUNCTION SERIES RELATED TO THE RIEMANN XI FUNCTION

ALEXANDER E. PATKOWSKI

ABSTRACT. We establish new Fourier integral evaluations involving the Riemann xi function related to a series involving Bessel function of the first kind. We show this infinite series involving the Bessel function of the first kind solves a boundary value problem for the cylindrical heat equation.

Keywords: Fourier Integrals; Riemann xi function; Heat equation.

2010 Mathematics Subject Classification 11M06, 33C15.

1. INTRODUCTION AND MAIN FORMULAS

The Riemann xi function is $\Xi(t) = \xi(\frac{1}{2} + it)$, where $\xi(s) := \frac{1}{2}s(s-1)\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})\zeta(s)$ and the Riemann zeta function is defined for $\Re(s) > 1$, to be the series $\sum_{n \geq 1} n^{-s}$ (see [7, 13] for background information). Evaluations of Fourier integrals for the Riemann xi function have a long history, dating back to the work of Koshliakov and Ramanujan [8, 12]. The classical application is found in Titchmarsh [13] to illustrate a proof of Hardy's theorem that infinitely many critical zeros of the Riemann zeta function lie on the line $\Re(s) = \frac{1}{2}$. Perhaps the most common example is [13]

$$(1.1) \quad \int_0^\infty \frac{\Xi(y)}{y^2 + \frac{1}{4}} \cos(xy) dy = \frac{\pi}{2} \left(e^{x/2} - 2e^{-x/2} \psi(e^{-2x}) \right),$$

which gives a nice connection to the theta function $\psi(y) = \sum_{n \geq 1} e^{-n^2 y}$. In fact, (1.1) implies the well known functional equation [9, pg.120, eq.(4.1.12)] by replacing x by $-x$.

Among many authors, we note [3, 4, 5, 8, 10, 12] have developed further evaluations of integrals of the same class as (1.1), many of which contain certain series involving Bessel functions. Infinite series identities involving Bessel functions have been extended by several authors [1, 2] to have relevance in a number theoretic context. The purpose herein is to note an example which appears to have a connection to a curious boundary value problem. As usual [9, pg.115] (or [6]), the Bessel series of

the first kind is given as

$$J_v(r) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n! \Gamma(n+v+1)} \left(\frac{r}{2}\right)^{2n+v}.$$

Theorem 1.1. For $x \in \mathbb{R}$,

$$\int_0^{\infty} \frac{\Xi^2(y)}{(y^2 + \frac{1}{4})^2} \left| {}_1F_1\left(\frac{1}{4} + \frac{iy}{2}; 1; -\frac{r^2}{4t}\right) \right|^2 \cos(xy) dy = e^{-x/2} \sqrt{t} \int_0^{\infty} H_1(ye^{-x}) H_1(y) dy,$$

where

$$H_1(y) := \sum_{n=1}^{\infty} e^{-tn^2 y^2} J_0(rny) - 2\sqrt{t\pi_1} F_1\left(\frac{1}{2}; 1; -\frac{r^2}{4t}\right) y^{-1}.$$

Next we offer a curious integral formula representation for $H_1(y)$, which can be seen as a generalized form of the functional equation for the Jacobi theta function [9, pg.120, eq.(4.1.12)]. This can be seen by setting $r = 0$, and evaluating the integral on the right side.

Theorem 1.2. For $x > 0$, $\Re(t) > 0$, and $|\arg(-\frac{r^2}{t})| < \frac{\pi}{2}$,

$$(1.2) \quad \sum_{n=1}^{\infty} e^{-tn^2 x^2} J_0(rnx) - 2\sqrt{t\pi_1} F_1\left(\frac{1}{2}; 1; -\frac{r^2}{4t}\right) x^{-1} \\ = e^{-\frac{r^2}{4t}} \int_1^{\infty} \frac{1}{\sqrt{t(yx)^2 - \pi^2}} \left(\sum_{n \geq 1} G_1(ny) - \frac{1}{y} \int_0^{\infty} G_1(y') dy' \right) dy,$$

where $G_1(z) = e^{-z^2} z J_0(2\sqrt{-\frac{r^2}{4t}} z)$.

We mention in passing that the asymptotic estimate [9, pg.130] (or [9, pg.209, eq.(5.4.7)]) of $J_0(r)$ for large r may be applied to see that

$$\sqrt{r} \sum_{n=1}^{\infty} J_0(nr) e^{-n^2 \kappa t} = \frac{1}{\sqrt{\pi}} \sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} (\cos(nr) + \sin(nr)) e^{-n^2 \kappa t} + O(r^{-1}),$$

as $r \rightarrow \infty$. While the series on the right side of this estimate is not quite the Jacobi theta function, it does indeed still provide a solution to the heat equation. For an expansion of the Fourier series on the right side when $t = 0$, see [9, pg.128, eq.(4.2.14)].

It is also possible to obtain a Riemann xi Fourier integral directly connected with $H_1(y)$.

Theorem 1.3. Set $F^+(s) = e^{xs} {}_1F_1(s/2; 1; -\frac{r^2}{4t}) + e^{x(1-s)} {}_1F_1((1-s)/2; 1; -\frac{r^2}{4t})$.
For $x \in \mathbb{R}$,

$$\int_0^\infty \frac{\Xi(y)}{(y^2 + \frac{1}{4})} F^+\left(\frac{1}{2} + iy\right) dy = \sum_{n=1}^\infty e^{-tn^2 e^{-2x}} J_0(rne^{-x}) {}_2F_1\left(\frac{1}{2}; 1; -\frac{r^2}{4t}\right) e^x.$$

2. REMARK ON AN ASSOCIATED BOUNDARY VALUE PROBLEM

In this section we detail a special boundary value problem associated with the series appearing in our Theorem 1.1. We consider the heat equation in cylindrical form

$$(2.1) \quad \kappa \nabla^2 u \equiv \kappa \left(\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} \right) = \frac{\partial u}{\partial t},$$

with the boundary conditions:

$$(i), u(0, t) = \sqrt{\frac{\pi}{t}} u(0, \frac{x^2}{t}) + \frac{1}{2} \sqrt{\frac{\pi}{t}} - \frac{1}{2}, \text{ for } t > 0,$$

$$(ii), u(r, 0) = -\frac{1}{2} + \frac{1}{r} + 2 \sum_{m=1}^n \frac{1}{\sqrt{r^2 - 4m^2 \pi^2}}, \text{ for } 2n\pi < r < 2(n+1)\pi.$$

The condition (i) is the functional equation for the theta function [9, pg.120, eq.(4.1.12)], and condition (ii) is equivalent to a series evaluation found in [6, pg.936]. The solution is given as our series in Theorem 1.1:

$$(2.2) \quad u(r, t) = \sum_{n=1}^\infty J_0(nr) e^{-n^2 \kappa t}.$$

This can be shown through the standard separation of variable method [11, pg.316].

Alternatively, one can show this through the known relations

$$\frac{\partial J_0(nr)}{\partial r} = -n J_1(nr),$$

$$\frac{\partial^2 J_0(nr)}{\partial r^2} = -n^2 \frac{\partial J_1(nr)}{\partial r} = \frac{n}{r} J_1(nr) - n^2 J_0(nr).$$

The growth condition $\lim_{r \rightarrow \infty} u(r, t) = 0$, may also be included in this problem, but $u(r', 0) = 0$ can not be satisfied for any $r' \in (0, \infty)$. This can be interpreted to mean that the initial temperature $u(r, 0)$ can never be 0, no matter the size of chosen radius r' . The general solution to (2.1) with $u(r', 0) = 0$, may be found in [11, pg.317, eq.(10)], and should be compared with (2.2).

3. PROOF OF MAIN RESULTS

In proving our Riemann xi function integrals we will regularly make use of Parseval's theorem for Mellin transforms [9, pg.83, eq.(3.1.11)]. Define

$$(3.1) \quad \int_0^\infty f(y)y^{s-1}dy =: \mathfrak{F}(s),$$

with inverse given by [9, pg.89],

$$(3.2) \quad f(y) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \mathfrak{F}(s)y^{-s} ds.$$

Lemma 3.1. *Suppose $\Re(s) = c$ is a real number confined to a region where $F(s)G(1-s)$ is analytic. Then,*

$$\int_0^\infty f(y)g(y)dy = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \mathfrak{F}(s)\mathfrak{G}(1-s)ds.$$

Proof of Theorem 1.1. By [6, pg.706, eq.6.631, #1] for $\Re(t) > 0$, $\Re(s+v) > 0$,

$$(3.3) \quad \int_0^\infty y^{s-1}e^{-ty^2} J_v(ry)dy = \frac{r^v \Gamma(\frac{v}{2} + \frac{s}{2})}{2t^{(s+v)/2} \Gamma(v+1)} {}_1F_1\left(\frac{v+s}{2}; v+1; -\frac{r^2}{4t}\right) \\ = \frac{\Gamma(\frac{v}{2} + \frac{s}{2})}{rt^{(s-1)/2} \Gamma(v+1)} e^{-r^2/8t} M_{(s-1)/2, v/2}\left(\frac{r^2}{4t}\right).$$

Setting $v = 0$, in (3.3), applying Mellin inversion, and inverting the desired sum gives us

$$(3.4) \quad \sum_{n=1}^\infty e^{-tn^2 y^2} J_0(rny) = \frac{1}{2\pi i} \int_{(p)} \frac{\Gamma(\frac{s}{2})}{2t^{s/2}} {}_1F_1\left(\frac{s}{2}; 1; -\frac{r^2}{4t}\right) \zeta(s) y^{-s} ds,$$

for $p > 1$, by absolute convergence. Computing the residue at the pole $s = 1$ gives us

$$(3.5) \quad H_1(y) := \sum_{n=1}^\infty e^{-tn^2 y^2} J_0(rny) - \frac{\sqrt{\pi}}{2t^{1/2}} {}_1F_1\left(\frac{1}{2}; 1; -\frac{r^2}{4t}\right) y^{-1} \\ = \frac{1}{2\pi i} \int_{(p')} \frac{\Gamma(\frac{s}{2})}{2t^{s/2}} {}_1F_1\left(\frac{s}{2}; 1; -\frac{r^2}{4t}\right) \zeta(s) y^{-s} ds,$$

for $0 < p' < 1$. Now applying Lemma 3.1 to (3.5) with $f(y) = H_1(ye^{-x})$, $g(y) = H_1(y)$, gives

$$(3.6) \quad \sqrt{t} \int_0^\infty H_1(ye^{-x}) H_1(y) dy = \frac{1}{2\pi i} \int_{(p')} \Gamma\left(\frac{s}{2}\right)^2 {}_1F_1\left(\frac{s}{2}; 1; -\frac{r^2}{4t}\right) {}_1F_1\left(\frac{1-s}{2}; 1; -\frac{r^2}{4t}\right) \zeta(s)^2 \pi^{\frac{1}{2}-s} e^{xs} ds,$$

where we have employed the functional equation for the Riemann xi function.

Putting $p' = \frac{1}{2}$, multiplying by $e^{-x/2}$, and making the change of variable $s = \frac{1}{2} + it$ now gives the result. \square

Proof of Theorem 1.2. Recalling (3.5), we have

$$\begin{aligned}
 H_1(x) &:= \sum_{n=1}^{\infty} e^{-tn^2x^2} J_0(rnx) - \frac{\sqrt{\pi}}{2t^{1/2}} {}_1F_1\left(\frac{1}{2}; 1; -\frac{r^2}{4t}\right) x^{-1} \\
 &= \frac{1}{2\pi i} \int_{(p')} \frac{\Gamma(\frac{s}{2})}{2t^{s/2}} {}_1F_1\left(\frac{s}{2}; 1; -\frac{r^2}{4t}\right) \zeta(s) x^{-s} ds \\
 &= \frac{1}{2\pi i} \int_{(1-p')} \frac{\Gamma(\frac{1-s}{2})}{2t^{(1-s)/2}} {}_1F_1\left(\frac{1-s}{2}; 1; -\frac{r^2}{4t}\right) \zeta(1-s) x^{s-1} ds \\
 &= \frac{1}{2\sqrt{\pi}i} \int_{(1-p')} \frac{\Gamma(\frac{s}{2})}{2t^{(1-s)/2}} {}_1F_1\left(\frac{1-s}{2}; 1; -\frac{r^2}{4t}\right) \zeta(s) \pi^{-s} x^{s-1} ds,
 \end{aligned} \tag{3.7}$$

for $0 < p' < 1$ by the functional equation for $\zeta(s)$ after making a change of variable s to $1-s$. From [9, pg.95, eq.(3.3.28)], $c > 0$,

$$\frac{1}{2\pi i} \int_{(c)} x^{-s} \frac{\Gamma(s)}{\Gamma(s+a+1)} ds = \frac{(1-x)^a}{\Gamma(a+1)}, \tag{3.8}$$

if $0 < x \leq 1$, and 0 if $x > 1$. From [6, pg.1023, eq.9.211,#3], we have that for $\Re(\alpha + v + 1) > 0$, $|\arg(x)| < \frac{\pi}{2}$,

$${}_1F_1(-v, \alpha + 1; x) = \frac{\Gamma(\alpha + 1)e^x x^{-\alpha/2}}{\Gamma(\alpha + v + 1)} \int_0^{\infty} e^{-y} y^{v+\alpha/2} J_{\alpha}(2\sqrt{xy}) dy, \tag{3.9}$$

Putting $v = (s-1)/2$, and $\alpha = 0$, in (3.9) gives for $\Re(s+1) > 0$, $|\arg(x)| < \frac{\pi}{2}$,

$${}_1F_1\left(\frac{1-s}{2}, 1; x\right) = \frac{e^x}{\Gamma(\frac{s+1}{2})} \int_0^{\infty} e^{-y} y^{(s-1)/2} J_0(2\sqrt{xy}) dy, \tag{3.10}$$

By the Müntz formula [13, pg.29, eq.(2.11.1)], if $F(x)$ satisfies (i) $F'(x)$ is bounded in any finite interval, (ii) continuous, and (iii) decays like $O(x^{-N})$, $N > 1$ as $x \rightarrow \infty$, then

$$\zeta(s) \int_0^{\infty} y^{s-1} F(y) dy = \int_0^{\infty} y^{s-1} \left(\sum_{n \geq 1} F(ny) - \frac{1}{y} \int_0^{\infty} F(y') dy' \right) dy, \tag{3.11}$$

for $0 < \Re(s) < 1$. Notice that the common region of holomorphy of (3.8) with s replaced by $1-s$, and (3.10) is $\{s : \Re(s) > -1\} \cap \{s \in \mathbb{C} : \Re(1-s) > 0\}$. Hence, putting $a = -\frac{1}{2}$ in (3.8) with $s \rightarrow s/2$, and applying (3.10) to Lemma 3.1 with

(3.11), tells us that (3.7) gives

$$\begin{aligned}
(3.12) \quad & \frac{1}{2\sqrt{\pi i}} \int_{(1-p')} \frac{\Gamma(\frac{s}{2})e^{-\frac{r^2}{4t}}}{2\Gamma(\frac{s+1}{2})t^{(1-s)/2}} \zeta(s) \left(\int_0^\infty e^{-z} z^{(s-1)/2} J_0(2\sqrt{-\frac{r^2}{4t}}z) dz \right) \pi^{-s} x^{s-1} ds \\
&= \frac{1}{2\sqrt{\pi i}} \int_{(1-p')} \frac{\Gamma(\frac{s}{2})e^{-\frac{r^2}{4t}}}{2\Gamma(\frac{s+1}{2})t^{(1-s)/2}} \left(\int_0^\infty z^{s-1} \left(\sum_{n \geq 1} G_1(nz) - \frac{1}{y} \int_0^\infty G_1(y') dy' \right) dz \right) \pi^{-s} x^{s-1} ds \\
&= e^{-\frac{r^2}{4t}} \int_1^\infty \frac{1}{\sqrt{t(yx)^2 - \pi^2}} \left(\sum_{n \geq 1} G_1(ny) - \frac{1}{y} \int_0^\infty G_1(y') dy' \right) dy,
\end{aligned}$$

where $G_1(z) = e^{-z^2} z J_0(2\sqrt{-\frac{r^2}{4t}}z)$. □

Proof of Theorem 1.3. This result follows from the second line in (3.7) with x replaced by e^{-x} upon noting $F^+(\frac{1}{2} + it) = F^+(\frac{1}{2} - it)$. □

REFERENCES

- [1] B.C. Berndt, S. Kim, A. Zaharescu, Weighted divisor sums and Bessel function series, II, *Adv. Math.* 229 (2012) 2055–2097.
- [2] B.C. Berndt, S. Kim, A. Zaharescu, Weighted divisor sums and Bessel function series, V, *J. of Approximation Theory* 197 (2015) 101–114.
- [3] A. Dixit. *Analogues of the general theta transformation formula*, *Proc. Roy. Soc. Edinburgh, Sect. A*, 143:371–399, 2013.
- [4] A. Dixit, B. C. Berndt and J. Sohn, *Character analogues of theorems of Ramanujan, Koshliakov and Guinand*, *Adv. Appl. Math.*, 46, Issues 1–4, (2011), 54–70 (Special issue in honor of Dennis Stanton).
- [5] A. Dixit, V. H. Moll *Self-reciprocal functions, powers of the Riemann zeta function and modular-type transformations*, *J. Number Theory* 147 (2015), 211–249.
- [6] I. S. Gradshteyn and I. M. Ryzhik. *Table of Integrals, Series, and Products*. Edited by A. Jeffrey and D. Zwillinger. Academic Press, New York, 7th edition, 2007.
- [7] H. Iwaniec and E. Kowalski, *Analytic number theory*, American Mathematical Society Colloquium Publications, vol. 53, American Mathematical Society, Providence, RI, 2004.
- [8] N. S. Koshlyakov, *On a transformation of definite integrals and its application to the theory of Riemann's function $\zeta(s)$* , *Comp. Rend. (Doklady) Acad. Sci. URSS* 15 (1937), 3–8.
- [9] R. B. Paris, D. Kaminski, *Asymptotics and Mellin–Barnes Integrals*. Cambridge University Press. (2001)
- [10] A. E. Patkowski, *Some Mellin transforms for the Riemann zeta function in the critical strip*, *International Journal of Number Theory*, Vol. 15, No. 01, pp. 153–156 (2019)

- [11] D. L. Powers, *Boundary Value Problems*, Harcourt Academic Press, 4th edition, 1999.
- [12] S. Ramanujan, *New expressions for Riemann's functions $\xi(s)$ and $\Xi(t)$* , *Quart. J. Math.*, 46: 253–260, 1915.
- [13] E. C. Titchmarsh, *The theory of the Riemann zeta function*, Oxford University Press, 2nd edition, 1986.

1390 Bumps River Rd.

Centerville, MA 02632

USA

E-mail: alexpatk@hotmail.com, alexepatkowski@gmail.com