

**PROPERTIES OF LOCAL ORTHONORMAL SYSTEMS**  
**PART I: UNCONDITIONALITY IN  $L^p$ ,  $1 < p < \infty$**

JACEK GULGOWSKI, ANNA KAMONT, AND MARKUS PASSENBRUNNER

**ABSTRACT.** Assume that we are given a filtration  $(\mathcal{F}_n)$  on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  of the form that each  $\mathcal{F}_n$  is generated by the partition of one atom of  $\mathcal{F}_{n-1}$  into two atoms of  $\mathcal{F}_n$  having positive measure. Additionally, assume that we are given a finite-dimensional linear space  $S$  of  $\mathcal{F}$ -measurable, bounded functions on  $\Omega$  so that on each atom  $A$  of any  $\sigma$ -algebra  $\mathcal{F}_n$ , all  $L^p$ -norms of functions in  $S$  are comparable independently of  $n$  or  $A$ . Denote by  $S_n$  the space of functions that are given locally, on atoms of  $\mathcal{F}_n$ , by functions in  $S$  and by  $P_n$  the orthoprojector (with respect to the inner product in  $L^2(\Omega)$ ) onto  $S_n$ .

Since  $S = \text{span}\{\mathbb{1}_\Omega\}$  satisfies the above assumption and  $P_n$  is then the conditional expectation  $\mathbb{E}_n$  with respect to  $\mathcal{F}_n$ , for such filtrations, martingales  $(\mathbb{E}_n f)$  are special cases of our setting.

We show in this article that certain convergence results that are known for martingales (or rather martingale differences) are also true in the general framework described above. More precisely, we show that the differences  $(P_n - P_{n-1})f$  converge unconditionally and are democratic in  $L^p$  for  $1 < p < \infty$ . This implies that those differences form a greedy basis in  $L^p$ -spaces for  $1 < p < \infty$ .

## 1. INTRODUCTION

Let  $(\Omega, \mathcal{F}, \mathbb{P}) = (\Omega, \mathcal{F}, |\cdot|)$  be a probability space and let  $(\mathcal{F}_n)_{n \geq 0}$  be a filtration. For  $1 < p < \infty$ , given a function  $f \in L^p$  and denoting by  $\mathbb{E}_n$  the conditional expectation of  $f$  with respect to the  $\sigma$ -algebra  $\mathcal{F}_n$ , the martingale differences  $(\mathbb{E}_n - \mathbb{E}_{n-1})f$  are orthogonal to each other and they converge unconditionally in  $L^p$  by D. Burkholder's inequality. We use [11] as our reference for the latter inequality and for other basic facts about martingales. Assume now that  $(\mathcal{F}_n)_{n \geq 0}$  is a *binary* filtration, meaning that

- (1)  $\mathcal{F}_0 = \{\emptyset, \Omega\}$ ,
- (2) for each  $n \geq 1$ ,  $\mathcal{F}_n$  is generated by  $\mathcal{F}_{n-1}$  and the subdivision of exactly one atom  $A_n$  of  $\mathcal{F}_{n-1}$  into two atoms  $A'_n, A''_n$  of  $\mathcal{F}_n$  satisfying  $|A''_n| \geq |A'_n| > 0$ .

For notational purpose, set  $\mathcal{F}_{-1} := \mathcal{F}_0 = \{\emptyset, \Omega\}$  and  $A_0 = \Omega$ . Moreover, denote the collection of all atoms in  $\mathcal{F}_n$  by  $\mathcal{A}_n$  and define  $\mathcal{A} = \cup_n \mathcal{A}_n$ . In this case, the range  $S_n$  of the conditional expectation  $\mathbb{E}_n$  consists of the piecewise constant functions. This precisely means that  $f \in S_n$  if and only if for all  $A \in \mathcal{A}_n$  there exists  $g \in \text{span}\{\mathbb{1}_\Omega\}$  with  $\mathbb{1}_A f = \mathbb{1}_A g$ . Here we denote by  $\mathbb{1}_A$  the characteristic function of the set  $A$  given by  $\mathbb{1}_A(x) = 1$  if  $x \in A$  and  $\mathbb{1}_A(x) = 0$  otherwise. Observe that clearly, for each atom  $A \in \mathcal{A}$ , we have

$$(1.1) \quad \{|\omega \in A : |f(\omega)| = \|f\|_A\} = |A|, \quad f \in \text{span}\{\mathbb{1}_\Omega\},$$

where by  $\|f\|_A$  we denote the  $L^\infty$ -norm of  $f$  on the set  $A$ .

---

*Date:* April 27, 2023.

*2020 Mathematics Subject Classification.* 42C05, 42C40, 46E30.

*Key words and phrases.* Local orthonormal systems, Unconditionality, Greedy bases, General measure spaces.

We now switch our viewpoint and assume that instead of  $\text{span}\{\mathbb{1}_\Omega\}$ , we are given an arbitrary vector space  $S \subseteq L^\infty$  of finite dimension  $k$ . Instead of (1.1), we assume for each  $A \in \mathcal{A}$  the inequality

$$(1.2) \quad |\{\omega \in A : |f(\omega)| \geq c_1 \|f\|_A\}| \geq c_2 |A|, \quad f \in S$$

for some constants  $c_1, c_2 \in (0, 1]$  independent of  $A$ . It can be checked that inequality (1.2) for fixed  $A$  is equivalent to the comparability of  $\|f\|_A$  and the mean value  $|A|^{-1} \int_A |f| d\mathbb{P}$  for all  $f \in S$ . For some explicit examples of spaces  $S$  and collections of atoms  $\mathcal{A}$  satisfying this inequality, we refer to Section 2.

Given any subspace  $V \subset L^1$  and any measurable set  $A \in \mathcal{F}$ , we let

$$V(A) = \{f \cdot \mathbb{1}_A : f \in V\}.$$

Let  $S_{-1} = \{0\}$  and, for each  $n \geq 0$ ,

$$S(\mathcal{F}_n) := S_n := \bigoplus_{A \in \mathcal{A}_n} S(A).$$

Thus,  $S(\mathcal{F}_n) = S_n$  consists of those functions that are piecewise (on each atom of  $\mathcal{F}_n$ ) contained in  $S$ . Letting  $P_n$  be the orthoprojector onto  $S_n$  (with respect to the inner product on the space  $L^2(\Omega, \mathcal{F}, \mathbb{P})$ ), we show in this article that the corresponding differences  $(P_n - P_{n-1})f$  also converge unconditionally in  $L^p$  for  $f \in L^p$ ,  $1 < p < \infty$ .

We even show a more general result. To describe it, observe first that the dimension of  $S_n/S_{n-1}$  generally is bigger than one. Therefore, we consider the following setting. For each  $n \geq 0$ , we assume that we are given a strictly increasing sequence of spaces  $S_{n-1} = V_{n,0} \subset V_{n,1} \subset \dots \subset V_{n,\ell_n} = S_n$ , such that  $\dim V_{n,j}/V_{n,j-1} = 1$  for each  $j = 1, \dots, \ell_n$ . Denote by  $P_{n,j}$  the orthoprojector onto  $V_{n,j}$ . We re-enumerate the spaces  $(V_{n,j})$  for  $n \geq 0$  and  $0 \leq j \leq \ell_n - 1$  so that we have a strictly increasing sequence of spaces  $(V_m)$ . Denote by  $R_m$  the orthoprojector onto  $V_m$ . For an integrable function  $f \in L^1(\Omega)$  and for an index  $m$ , set

$$f_m := R_m f, \quad df_m = f_m - f_{m-1}.$$

Then we show that those differences converge unconditionally in  $L^p$  for  $1 < p < \infty$  independently of the choice of spaces  $(V_{n,j})$ . More precisely,

**Theorem 1.1.** *Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space,  $(\mathcal{F}_n)$  a binary filtration and  $S$  a finite dimensional linear space of  $\mathcal{F}$ -measurable functions on  $\Omega$  satisfying (1.2) for every atom  $A \in \mathcal{A}$ .*

*Then, for every  $1 < p < \infty$  there exists a constant  $C_p$  such that for every  $f \in L^p$  and all positive integers  $n$ ,*

$$\left\| \sum_{m \leq n} \pm df_m \right\|_p \leq C_p \|f\|_p.$$

The result of Theorem 1.1 should be compared to corresponding results [4, 9] about the unconditionality (in  $L^p$  for  $1 < p < \infty$ ) of univariate orthogonal spline functions for arbitrary partitions. (In the piecewise linear case, those orthogonal spline functions are called general Franklin functions.) A necessary prerequisite for deriving unconditionality of those functions are sharp pointwise estimates for those orthogonal spline functions. Due to the fact that the considered splines admit smoothness conditions at the breakpoints, orthogonal spline functions are not localized, which makes them hard to estimate. On the other hand, note that discontinuous spline systems, i.e. such that each knot is of maximal admissible multiplicity, are examples of systems treated in this article.

Our setting in the present article considers differences  $df_m$  that have the same localization property than Haar systems. This allows us to prove general pointwise estimates for  $df_m$ , which is done in Section 4. As a consequence, we prove Theorem 1.1 under very mild assumptions on the domain of the involved functions. Here, we work with arbitrary probability spaces compared to some bounded interval, equipped with Lebesgue measure, that is needed for univariate orthogonal spline functions.

Denote by  $\psi_m^p$  a function with  $\|\psi_m^p\|_p = 1$ ,  $\psi_m^p \in V_m$  and  $\psi_m^p \perp V_{m-1}$ . Then, we additionally show that the functions  $(\psi_m^p)$  are democratic in  $L^p$  for  $1 < p < \infty$ , i.e. we show

**Theorem 1.2.** *For each  $1 < p < \infty$ , there exists a constant  $C_p$  depending on  $p$ ,  $c_1, c_2$  and the dimension of  $S$  so that*

$$C_p^{-1}(\text{card } \Lambda)^{1/p} \leq \left\| \sum_{m \in \Lambda} \psi_m^p \right\|_p \leq C_p(\text{card } \Lambda)^{1/p}$$

for every finite subset  $\Lambda$  of indices, where  $\text{card } \Lambda$  denotes the cardinality of  $\Lambda$ .

If we combine the results of Theorems 1.1 and 1.2, by a result of S. Konyagin and V. Temlyakov [7] (see also the book [13] by V. Temlyakov),  $(\psi_j^p)$  is also a greedy basis in its closed span  $U$  as a subset of  $L^p$ , meaning that the best  $m$ -term approximation of each function  $f = \sum_j c_j \psi_j^p \in U$  in terms of the functions  $(\psi_j^p)$  is achieved (up to a multiplicative constant) by greedy approximation, i.e. by taking just the terms in the expansion of  $f$  with the  $m$  highest absolute values of coefficients  $|c_j| = \|\langle f, \psi_j \rangle \psi_j\|_p$ .

The main motivation of the work presented in this article comes from the following question. It is known that the best  $n$ -term approximation of a function  $f \in L^p[0, 1]^d$ ,  $1 < p < \infty$  can be achieved by greedy approximation. On the other hand, P. Petrushev showed that the spaces corresponding to the best  $n$ -term approximation by linear combinations of characteristic functions of dyadic cubes and by linear combinations of Haar functions coincide, cf. [10, Theorems 3.3 and 5.3].

In [5], we ask the question whether this is also true in the more general situation given in the present article. Interesting special cases include general Haar systems or orthonormal piecewise polynomial systems corresponding to rectangular partitions, as discussed in Example 2.4. However, for this we need to know that the orthonormal systems under consideration are greedy basic sequences in  $L^p$ . This property is known (or can be easily deduced from the unconditionality) for general Haar systems (cf. [6]), or for orthonormal spline systems on the unit interval (cf. [4, 9]). The required result in general is provided by Theorems 1.1 and 1.2 above.

The organization of the article is as follows. Section 2 lists some explicit examples of spaces  $S$  and atoms  $\mathcal{A}$  satisfying inequality (1.2). In Section 3 we give a criterion under which  $\cup_n S_n$  is dense in  $L^p$ . In Section 4 we prove some crucial pointwise estimates for  $P_n f$  that are used subsequently. Sections 5 and 6 show Theorem 1.1 by deriving a decomposition of  $L^1$ -functions similar to Gundy's decomposition for martingales. In Section 7 we show Theorem 1.2. Section 8 treats the case of infinite measure spaces. In the last Section 9 we investigate the questions of unconditionality and democracy in the setting of non-binary filtrations, meaning that we consider splitting  $A_n \in \mathcal{A}_{n-1}$  into  $r \geq 2$  atoms  $A_{n,1}, \dots, A_{n,r}$  of  $\mathcal{F}_n$  instead of just two. In particular, we show that general results as Theorem 1.1 that show unconditionality independently of the choice of  $(V_{n,j})$ , are not possible in this case.

## 2. EXAMPLES

In this section we give a list of explicit examples of probability spaces  $(\Omega, \mathcal{F}, \mathbb{P})$ , finite-dimensional vector spaces  $S$  and binary filtrations  $(\mathcal{F}_n)$  satisfying inequality (1.2) for some constants  $c_1, c_2$ .

Before we discuss examples, we make the following simple observation:

**Lemma 2.1.** *Let  $(\Omega, \mathcal{F}, \mathbb{P})$  a probability space with  $|\cdot| = \mathbb{P}$  and  $(\mathcal{F}_n)$  be a binary filtration. Let  $S$  be chosen such that there exist two constants  $C, n$  so that for all  $A \in \mathcal{A}$  we have the inequality*

$$(2.1) \quad \|f\|_A \leq \left( \frac{C|A|}{|E|} \right)^n \|f\|_E, \quad f \in S,$$

for all  $\mathcal{F}$ -measurable subsets  $E \subset A$ .

Then, inequality (1.2) is satisfied for all  $A \in \mathcal{A}$  with the constants  $c_1 = (2C)^{-n}$  and  $c_2 = 1/2$ .

*Proof.* Let  $A \in \mathcal{A}$  be a fixed atom and let  $f \in S$  be arbitrary. We assume without restriction that  $\|f\|_A > 0$ . Then, we just need to apply assumption (2.1) to the  $\mathcal{F}$ -measurable set

$$E = \{\omega \in A : |f(\omega)| < \alpha \|f\|_A\},$$

with  $\alpha = (2C)^{-n}$ . Indeed, doing that implies

$$\|f\|_A \leq \left( \frac{C|A|}{|E|} \right)^n \|f\|_E \leq \alpha \left( \frac{C|A|}{|E|} \right)^n \|f\|_A.$$

Since  $\|f\|_A > 0$  we obtain  $|E|^n \leq \alpha(C|A|)^n$  and the specific value of  $\alpha$  yields  $|E| \leq |A|/2$ . Therefore,

$$|A \setminus E| = |\{\omega \in A : |f(\omega)| \geq \alpha \|f\|_A\}| \geq \frac{|A|}{2}.$$

Since  $A \in \mathcal{A}$  and  $f \in S$  were arbitrary, this shows (1.2) with the constants  $c_1 = \alpha = (2C)^{-n}$  and  $c_2 = 1/2$ .  $\square$

Thus, if we give examples satisfying (2.1), we in particular have (1.2) as well.

In the following two examples (Examples 2.2 and 2.3), let  $\Omega$  be a bounded subinterval of  $\mathbb{R}$ ,  $\mathcal{F}$  be the Borel  $\sigma$ -algebra,  $\lambda$  the (one-dimensional) Lebesgue measure and  $\mathbb{P} = |\cdot|$  the renormalized Lebesgue measure  $\lambda/\lambda(\Omega)$ .

**Example 2.2.** Let  $S$  be the space of polynomials of degree at most  $n$ . Then, the classical Remez inequality states that for every  $f \in S$  and every bounded interval  $A$  and every measurable subset  $E \subset A$  we have

$$\|f\|_A \leq \left( \frac{4|A|}{|E|} \right)^n \|f\|_E.$$

Thus, for every binary filtration  $(\mathcal{F}_n)$ , where all atoms  $A \in \mathcal{A}$  are intervals, we use Lemma 2.1 to deduce inequality (1.2) with the constants (e.g.)  $c_1 = 8^{-n}$  and  $c_2 = 1/2$ .

**Example 2.3.** Let  $(\lambda_k)_{k=1}^n$  be arbitrary complex numbers. Set

$$S := \text{span}\{\exp(\lambda_k \cdot) : k = 1, \dots, n\}.$$

F. Nazarov's extension [8] of P. Turan's lemma states that for every  $f \in S$ , every bounded interval  $A$  and every measurable subset  $E \subset A$  we have

$$(2.2) \quad \|f\|_A \leq \exp\left(\max_{1 \leq k \leq n} |\Re \lambda_k| \cdot \lambda(A)\right) \left(\frac{C|A|}{|E|}\right)^{n-1} \|f\|_E$$

for some constant  $C$ . Letting  $C_\Omega = \exp\left(\max_{1 \leq k \leq n} |\Re \lambda_k| \cdot \lambda(\Omega)/(n-1)\right)C$ , we have as a consequence

$$(2.3) \quad \|f\|_A \leq \left(\frac{C_\Omega|A|}{|E|}\right)^{n-1} \|f\|_E.$$

Inequality (2.3) shows assumption (2.1) of Lemma 2.1. Therefore, Lemma 2.1 implies inequality (1.2) with the constants  $c_1 = (2C_\Omega)^{-n}$  and  $c_2 = 1/2$ .

In particular, (1.2) is satisfied if  $S$  is the space of trigonometric polynomials up to some degree  $n$ .

Now, let  $\Omega \subset \mathbb{R}^d$  be a convex body (i.e. a compact convex subset of  $\mathbb{R}^d$  with nonempty interior),  $\mathcal{F}$  the Borel  $\sigma$ -algebra on  $\Omega$  and  $\mathbb{P} = |\cdot|$  be the  $d$ -dimensional Lebesgue measure renormalized so that  $|\Omega| = 1$ .

**Example 2.4.** If  $p(x) = \sum_{\alpha \in I} a_\alpha x^\alpha$  is a  $d$ -variate polynomial where  $I$  is a finite set containing  $d$ -dimensional multiindices, then the degree of  $p$  is defined as  $\max\{\sum_{i=1}^d \alpha_i : \alpha \in I\}$ . For a fixed non-negative integer  $n$ , let  $S$  be the space of polynomials of degree at most  $n$ .

Then, the multivariate Remez inequality (see [3, 1]) states that if  $A \subset \mathbb{R}^d$  is a convex body and  $E \subset A$  a measurable subset, for all  $f \in S$  we have

$$\|f\|_A \leq \left(\frac{4d|A|}{|E|}\right)^n \|f\|_E.$$

Thus, for every binary filtration  $(\mathcal{F}_n)$ , where all atoms  $A$  are convex sets whose closure  $\bar{A}$  has the same Lebesgue measure than  $A$ , we use Lemma 2.1 to deduce inequality (1.2) with the constants  $c_1 = (8d)^{-n}$  and  $c_2 = 1/2$ .

We remark here that similar to the multivariate Remez inequality as in Example 2.4, there are multivariate versions of (2.3), see for instance the result [2] by N. Fontes-Merz. Therefore, by Lemma 2.1, such spaces of functions also satisfy inequality (1.2) and are thus examples of the general framework discussed here.

**Example 2.5.** Let  $\gamma : [0, 1] \rightarrow \mathbb{C}$  be a curve of unit length that is two times continuously differentiable and let  $S$  be the space of polynomials of degree at most  $n$ . Let  $\mathbb{P}$  be the one-dimensional Lebesgue measure on the image of  $\gamma$ . Then, there exist constants  $c_1, c_2 \in (0, 1]$  that may depend on  $\gamma$  such that, for all intervals  $I \subset [0, 1]$  and denoting by  $\gamma_I$  the restriction of  $\gamma$  to the interval  $I$ , we have the inequality

$$\mathbb{P}(\{z \in \gamma_I : |f(z)| \geq c_1 \|f\|_{\gamma_I}\}) \geq c_2 |\gamma_I|, \quad f \in S.$$

This inequality is easy to see if  $\gamma$  is a straight line and if  $\gamma$  is not a straight line, we approximate  $\gamma$  by straight lines and use translation and dilation invariance of the class of polynomials of at most degree  $n$ .

**Remark 2.6.** If inequality (1.2) is satisfied for some vector space  $S$ , any subspace of  $S$  clearly satisfies (1.2) as well. Thus, in each of the above examples it is possible to replace the space  $S$  considered there by any linear subspace thereof.

3. DENSITY OF  $\cup_n S_n$  IN  $L^p$  SPACES

Here and in the following we assume that  $(\Omega, \mathcal{F}, \mathbb{P})$  is a probability space,  $(\mathcal{F}_n)$  is a binary filtration and  $S \subset L^\infty$  is a vector space satisfying inequality (1.2) for each atom  $A \in \mathcal{A}$ . We also use the notation introduced in Section 1.

Next, we give a condition in terms of the filtration  $(\mathcal{F}_n)$  under which  $\cup_n S_n$  is dense in  $L^p$ . Here we also have to assume something about the space  $S$ , but as we will see in the remark below, this condition is necessary.

**Theorem 3.1.** *Let  $0 < p < \infty$  and assume that the following two conditions are satisfied:*

- (1)  $|\bigcap_{f \in S} \{f = 0\}| = 0$ ,
- (2) for all  $\varepsilon > 0$  and every  $A \in \mathcal{F}$  there exists a positive integer  $N$  and a set  $B \in \mathcal{F}_N$  satisfying  $|A \Delta B| \leq \varepsilon$ .

Then,  $\cup_n S_n$  is dense in  $L^p(\Omega, \mathcal{F})$ .

**Remark 3.2.** (a) After the choice of a basis  $p_1, \dots, p_k$  of  $S$ , we have

$$\bigcap_{f \in S} \{f = 0\} = \bigcap_{i=1}^k \{p_i = 0\},$$

which in particular implies the measurability of the set  $\bigcap_{f \in S} \{f = 0\}$ .

(b) Condition (1) is necessary: for assume that condition (1) is not satisfied. Let  $D = \bigcap_{f \in S} \{f = 0\} \in \mathcal{F}$ , with  $|D| = c > 0$ . By definition of  $S_n$ , every function  $f \in \cup_n S_n$  vanishes identically on  $D$ , thus

$$c = \inf_{f \in \cup_n S_n} \|\mathbb{1}_D - f\|_{L^p(D)}^p \leq \inf_{f \in \cup_n S_n} \|\mathbb{1}_D - f\|_p^p.$$

This implies that  $\cup_n S_n$  is not dense in  $L^p(\Omega, \mathcal{F})$ .

(c) If the  $\sigma$ -algebra  $\mathcal{F}$  is generated by  $\cup_n \mathcal{F}_n$ , then condition (2) is satisfied.

*Proof of Theorem 3.1.* Choose a basis  $p_1, \dots, p_k$  of  $S$  and assume that (2) is satisfied. Observe that also, by (1), we have  $|\bigcap_{i=1}^k \{p_i = 0\}| = 0$ . Let  $A \in \mathcal{F}$ . We intend to approximate  $\mathbb{1}_A$  in  $L^p$  by functions in  $\cup_n S_n$ . Fix  $\varepsilon > 0$ . First, we split  $\mathbb{C} \setminus \{0\}$  into disjoint cubes  $(I_j)_{j \in \mathbb{Z} \setminus \{0\}}$  so that for each  $j$ , and every  $x, y \in I_j$ , we have that  $x/y \in B(1, \varepsilon) := \{z \in \mathbb{C} : |z - 1| < \varepsilon\}$ . For  $i = 1, \dots, k$  and  $j \in \mathbb{Z} \setminus \{0\}$ , define the sets

$$C_{i,j} = \{p_i \in I_j\}$$

and  $C_{i,0} = \{p_i = 0\}$ . For  $j = (j_1, \dots, j_k) \in \mathbb{Z}^k$ , define the disjoint sets

$$D_j := A \cap C_{1,j_1} \cap \dots \cap C_{k,j_k}.$$

Let  $W$  be a finite set of indices  $j \in \mathbb{Z}^k \setminus \{0\}$  so that  $|\cup_{j \in W} D_j| \geq (1 - \varepsilon)|A|$  and for each  $j \in W$  we have that  $|D_j| > 0$ . This is possible for every  $\varepsilon > 0$  since by assumption (1), we have  $|D_0| = 0$ . For each  $j \in W$ , choose a set  $B_j \in \mathcal{F}_{N_j}$  with

$$|B_j \Delta D_j| \leq \varepsilon_j,$$

where  $\varepsilon_j$  depending on  $\varepsilon, p, k$  will be chosen later. Since  $B_j \in \mathcal{F}_{N_j}$ , it can be written as a finite union of disjoint atoms  $(A_{j,\ell})_\ell$  from  $\mathcal{F}_{N_j}$ .

For  $q = 1 - c_2/2 \in (0, 1)$  with  $c_2$  being the constant in (1.2), split the indices  $\ell$  into

$$\Gamma_j := \{\ell : |A_{j,\ell} \cap D_j| \leq q|A_{j,\ell}|\}, \quad \Lambda_j := \{\ell : |A_{j,\ell} \cap D_j| > q|A_{j,\ell}|\}.$$

We have the following estimate

$$(3.1) \quad \varepsilon_j \geq |B_j \setminus D_j| = \sum_{\ell} |A_{j,\ell} \setminus D_j| \geq \sum_{\ell \in \Gamma_j} |A_{j,\ell} \setminus D_j| = \sum_{\ell \in \Gamma_j} |A_{j,\ell} \cap D_j^c| \geq (1-q) \sum_{\ell \in \Gamma_j} |A_{j,\ell}|.$$

For  $j \in W$ , let  $u_j := p_i$  with  $i \in \{1, \dots, k\}$  chosen so that  $j_i \neq 0$ . Let  $\omega_j \in D_j$  arbitrary. Then, set  $v_j := u_j/u_j(\omega_j)$  and  $E_j := \cup_{\ell \in \Lambda_j} A_{j,\ell}$ . Observe that (3.1) then implies

$$(3.2) \quad |E_j \Delta D_j| \leq |B_j \Delta D_j| + \left| \bigcup_{\ell \in \Gamma_j} A_{j,\ell} \right| \leq (1 + (1-q)^{-1}) \varepsilon_j.$$

Now, we define the approximation  $f$  of  $\mathbb{1}_A$  to be a suitable finite sum of the following form.

$$f = \sum_{j \in W} v_j \mathbb{1}_{E_j} \in \cup_n S_n.$$

The values of the function  $v_j$  on  $D_j$  are contained in the ball  $B(1, \varepsilon)$  by construction of the sets  $D_j$  and the intervals  $I_j$ .

Next we see that  $|v_j|$  is bounded on  $E_j$  by  $c_1^{-1}(1 + \varepsilon)$ . Assume the contrary, i.e., assume that  $\|v_j\|_{A_{j,\ell}} > (1 + \varepsilon)/c_1$  for some  $\ell \in \Lambda_j$ . This would imply using (1.2) that

$$(3.3) \quad c_2 |A_{j,\ell}| \leq |\{\omega \in A_{j,\ell} : |v_j| \geq c_1 \|v_j\|_{A_{j,\ell}}\}| \leq |\{\omega \in A_{j,\ell} : |v_j| > 1 + \varepsilon\}|.$$

On the other hand, since  $\ell \in \Lambda_j$ ,

$$(3.4) \quad q |A_{j,\ell}| \leq |D_j \cap A_{j,\ell}| \leq |\{\omega \in A_{j,\ell} : |v_j| \leq 1 + \varepsilon\}|.$$

Both (3.3) and (3.4) simultaneously are impossible by definition of  $q = 1 - c_2/2$  and thus we obtain  $\|v_j\|_{L^\infty(E_j)} \leq (1 + \varepsilon)/c_1$ .

Since we know the inequality  $|\cup_{j \in W} D_j| \geq (1 - \varepsilon)|A|$ , in order to estimate  $\|f - \mathbb{1}_A\|_p$ , it suffices to estimate the  $p$ -norm of the function  $\delta = f - \sum_{j \in W} \mathbb{1}_{D_j}$ . Define  $\delta_1 = \sum_{j \in W} (1 - v_j) \mathbb{1}_{D_j \cap E_j}$  and  $\delta_2 = \delta - \delta_1$ . Since the range of  $v_j$  on  $D_j$  is contained in  $B(1, \varepsilon)$  and the sets  $D_j \cap E_j$  are disjoint, we obtain  $\|\delta_1\|_p \leq \varepsilon$ . Moreover, we have the pointwise inequality  $|\delta_2| \leq \sum_{j \in W} (1 + \|v_j\|_{L^\infty(E_j)}) \mathbb{1}_{D_j \Delta E_j} \leq \sum_{j \in W} (1 + (1 + \varepsilon)c_1^{-1}) \mathbb{1}_{D_j \Delta E_j}$ . By (3.2), this implies in the case  $p \geq 1$

$$\|\delta_2\|_p \leq (1 + (1 + \varepsilon)c_1^{-1})(1 + (1 - q)^{-1})^{1/p} \sum_{j \in W} \varepsilon_j^{1/p},$$

and in the case  $p < 1$

$$\|\delta_2\|_p^p \leq (1 + (1 + \varepsilon)c_1^{-1})^p (1 + (1 - q)^{-1}) \sum_{j \in W} \varepsilon_j.$$

In both cases, we can choose  $(\varepsilon_j)$  depending on  $\varepsilon, p, k$  such that  $\|\delta_2\|_p \leq \varepsilon$ . Combining this with the estimate for  $\|\delta_1\|_p$ , we obtain that the characteristic function  $\mathbb{1}_A$  of each set  $A \in \mathcal{F}$  can be approximated arbitrarily closely by functions in  $\cup_n S_n$  in  $L^p$ , which implies the conclusion.  $\square$

#### 4. A POINTWISE BOUND FOR $P_{n,j}$

We continue to use the notation from Section 1. In particular,  $P_n$  denotes the orthoprojector onto  $S_n$ . Since  $S \subset L^\infty(\Omega)$ , the operators  $P_n$  can be extended to  $L^1(\Omega)$  and it is easy to see that for each  $n$  it satisfies

$$(4.1) \quad P_n(\mathbb{1}_B f) = \mathbb{1}_B P_n f, \quad B \in \mathcal{F}_n, f \in L^1.$$

Indeed, if  $g \in S_n$ , we have

$$\int (\mathbb{1}_B f - \mathbb{1}_B P_n f) \bar{g} \, d\mathbb{P} = \int (f - P_n f) \overline{\mathbb{1}_B g} \, d\mathbb{P},$$

which vanishes since  $\mathbb{1}_B g$ , together with  $g$ , is contained in  $S_n$ , which is the range of  $P_n$ . If  $n, j$  are chosen so that  $V_{n,j} = V_m$ , we have of course  $P_{n,j} = R_m$ . Define the index set  $I_n = \{m : S_{n-1} \subsetneq V_m \subseteq S_n\}$ .

**Lemma 4.1.** *Suppose that  $n \geq 1$  and let  $g \in S_n$  with  $g \perp S_{n-1}$ . Then  $g$  vanishes identically a.e. on the complement  $A_n^c$  of  $A_n$ .*

*Proof.* Let  $A = A_n^c$ . Then,  $\mathbb{1}_A g \in S_{n-1}$ , which implies that

$$0 = \langle g, \mathbb{1}_A g \rangle = \int_A |g|^2 \, d\mathbb{P},$$

which gives that  $g$  vanishes a.e. on  $A = A_n^c$ .  $\square$

**Remark 4.2.** Let  $n \geq 1$ . Since for  $m \in I_n$ , the function  $df_m$  is orthogonal to  $S_{n-1}$ , this implies that the support of  $df_m$  is a subset of  $A_n$  for  $m \in I_n$ .

Also this implies that for each  $B \in \mathcal{F}_{n-1}$  and each  $m \in I_n$  and every  $f \in L^1$ , we have  $R_m(\mathbb{1}_B f) = \mathbb{1}_B R_m f$ .

We denote by  $\mathbb{E}_n$  the conditional expectation with respect to  $\mathcal{F}_n$ . We use the symbol  $A(t) \lesssim B(t)$  in order to denote the fact that there exists a constant  $C$  depending only on  $c_1, c_2$  from (1.2) and  $k$  so that for all  $t$  we have the inequality  $A(t) \leq CB(t)$ , where  $t$  denotes all implicit or explicit dependencies that the objects  $A$  and  $B$  might have. Similarly, we use the notations  $A(t) \gtrsim B(t)$  and  $A(t) \simeq B(t)$ .

**Lemma 4.3.** *Let  $n \geq 1$  and let  $\psi \in S_n$  with  $\psi \perp S_{n-1}$  and  $\|\psi\|_2 = 1$ .*

*Then we have the estimates*

$$\|\psi\|_{A_n^c} = 0, \quad \|\psi\|_{A_n'} \lesssim |A_n'|^{-1/2}, \quad \|\psi\|_{A_n''} \lesssim \frac{|A_n'|^{1/2}}{|A_n|}.$$

*If  $|A_n''| \geq (1 - c_2/2)|A_n|$ , the latter estimate can be improved to*

$$(4.2) \quad \|\psi\|_{A_n''} \lesssim \sup_{u \in S} \frac{\|u\|_{A_n'} |A_n'|^{1/2}}{\|u\|_{A_n} |A_n|}.$$

*Proof.* We first show that  $\|\psi\|_{A_n'}$  and  $\|\psi\|_{A_n''}$  are both positive. Write  $\psi = \psi_1 \mathbb{1}_{A_n'} + \psi_2 \mathbb{1}_{A_n''}$  with  $\psi_1, \psi_2 \in S$  and assume that  $\psi_1 = 0$  a.s. on  $A_n'$ . Since  $\psi$  is orthogonal to  $S = S(\Omega)$  and  $\psi_2 \in S$ , we have

$$0 = \langle \psi, \psi_2 \rangle = \int_{A_n''} |\psi_2|^2 \, d\mathbb{P},$$

which implies  $\psi_2 = 0$  a.s. on  $A_n''$  and therefore,  $\psi = 0$  a.s. contradicting the assumption  $\|\psi\|_2 = 1$ . Thus we have  $\|\psi_1\|_{A_n'} > 0$ . By a similar argument, testing  $\psi$  with  $\psi_1$  instead of  $\psi_2$ , we obtain  $\|\psi_2\|_{A_n''} > 0$ .

The assertion about  $\|\psi\|_{A_n^c}$  is just Lemma 4.1. Observe that the  $L^2$ -normalization condition of  $\psi$  becomes

$$(4.3) \quad 1 = \int_{A_n} |\psi|^2 \, d\mathbb{P} = \int_{A_n''} |\psi|^2 \, d\mathbb{P} + \int_{A_n'} |\psi|^2 \, d\mathbb{P} \gtrsim \|\psi\|_{A_n''}^2 |A_n''| + \|\psi\|_{A_n'}^2 |A_n'|,$$

where in the last inequality we used (1.2) for the atoms  $A_n''$  and  $A_n'$ , respectively. In particular, this yields the estimates

$$(4.4) \quad \|\psi\|_{A_n'} \lesssim |A_n'|^{-1/2}, \quad \|\psi\|_{A_n''} \lesssim |A_n''|^{-1/2}.$$



Now we distinguish the following cases.

CASE I: Assume  $|A_n''| \leq (1 - c_2/2)|A_n|$ . In this case we have  $|A_n| \simeq |A_n''| \simeq |A_n'|$ . Therefore, the second estimate in (4.4) implies the assertion.

CASE II: Assume  $|A_n''| \geq (1 - c_2/2)|A_n|$ . By orthogonality of  $\psi$  to  $S$ , we have  $\langle \psi, \psi_2 \rangle = 0$ . Evaluating this condition yields

$$0 = \int_{A_n''} \psi_1 \overline{\psi_2} \, d\mathbb{P} + \int_{A_n''} |\psi_2|^2 \, d\mathbb{P},$$

which implies the inequality

$$(4.5) \quad |A_n''| \|\psi_2\|_{A_n''}^2 \lesssim \int_{A_n''} |\psi_2|^2 \, d\mathbb{P} = \left| \int_{A_n''} \psi_1 \overline{\psi_2} \, d\mathbb{P} \right| \leq \|\psi_1\|_{A_n'} \|\psi_2\|_{A_n'} |A_n'|.$$

Thus, since  $\|\psi_2\|_{A_n''} > 0$  by the first part of the proof, we divide by  $|A_n''| \|\psi_2\|_{A_n''}$  and use the estimate for  $\|\psi\|_{A_n'} = \|\psi_1\|_{A_n'}$  from (4.4) above to get

$$\|\psi\|_{A_n''} = \|\psi_2\|_{A_n''} \lesssim \frac{\|\psi_2\|_{A_n'} |A_n'|^{1/2}}{\|\psi_2\|_{A_n''} |A_n''|}.$$

Since  $|A_n''| \geq (1 - c_2/2)|A_n|$ , inequality (1.2) implies  $\|\psi_2\|_{A_n''} \simeq \|\psi_2\|_{A_n}$ . Thus, this inequality and the fact that  $\psi_2 \in S$  in particular implies (4.2).  $\square$

**Theorem 4.4.** *For every integer  $n \geq 0$ , every  $g \in L^1$  and every  $j \in \{0, \dots, \ell_n\}$ , we have the following pointwise inequality:*

$$|P_{n,j}g| \lesssim \mathbb{E}_{n-1}|g| + \mathbb{E}_n|g|.$$

*Proof.* Assume that  $n \geq 0$  and  $j \in \{0, \dots, \ell_n\}$ . Let  $A$  be an atom of  $\mathcal{F}_{n-1}$  and let  $g_1, \dots, g_M$  be an orthonormal basis of  $\{f\mathbb{1}_A : f \in \text{ran } P_{n,j}\}$  so that  $g_1, \dots, g_L$  is an orthonormal basis of  $\{f\mathbb{1}_A : f \in S_{n-1}\}$ . We know that  $L \leq M \leq 2k$ . Additionally, we know that  $L < M$  only if  $A = A_n$  and  $j > 0$ . Thus, in order to estimate  $P_{n,j}g = \sum_{i=1}^M \langle g, g_i \rangle g_i$  on  $A$ , it suffices to estimate the expression  $\langle g, g_i \rangle g_i$  on  $A$  for a fixed  $1 \leq i \leq M$ .

Firstly, assume that  $i \leq L$ . Then, since  $g_i$  is  $L^2$ -normalized and contained in  $S_{n-1}$  and  $A$  is an atom of  $\mathcal{F}_{n-1}$ , (1.2) implies

$$1 = \int_A |g_i|^2 \, d\mathbb{P} \geq d|A| \|g_i\|_A^2.$$

with  $d = c_1^2 c_2$ . Therefore we estimate on  $A$  as follows

$$(4.6) \quad |\langle g, g_i \rangle g_i| \leq \int_A |g| \, d\mathbb{P} \cdot \|g_i\|_A^2 \leq \frac{1}{d|A|} \int_A |g| \, d\mathbb{P} = d^{-1} \cdot \mathbb{E}_{n-1}|g|.$$

Secondly, assume that  $L < i \leq M$ , which implies  $A = A_n$ . We also assume that  $n \geq 1$  since if  $n = 0$ , the result follows in the same way as in (4.6). We have to estimate the expression  $\langle g, g_i \rangle g_i$  both on  $A_n''$  and on  $A_n'$ . Denote  $\ell := \|g_i\|_{A_n''}$  and  $s := \|g_i\|_{A_n'}$ . Thus, we can estimate  $|\langle g, g_i \rangle g_i|$  on  $A_n''$  using Lemma 4.3 as follows.

$$\begin{aligned} |\langle g, g_i \rangle g_i| &\leq \ell^2 \int_{A_n''} |g| \, d\mathbb{P} + \ell s \int_{A_n'} |g| \, d\mathbb{P} \\ &\lesssim \frac{|A_n'|}{|A_n''|^2} \int_{A_n''} |g| \, d\mathbb{P} + \frac{1}{|A_n|} \int_{A_n'} |g| \, d\mathbb{P} \lesssim \mathbb{E}_{n-1}|g| + \mathbb{E}_n|g| \end{aligned}$$

and similarly on  $A'_n$

$$\begin{aligned} |\langle g, g_i \rangle g_i| &\leq \ell s \int_{A''_n} |g| \, d\mathbb{P} + s^2 \int_{A'_n} |g| \, d\mathbb{P} \\ &\lesssim \frac{1}{|A_n|} \int_{A''_n} |g| \, d\mathbb{P} + \frac{1}{|A'_n|} \int_{A'_n} |g| \, d\mathbb{P} \lesssim \mathbb{E}_{n-1}|g| + \mathbb{E}_n|g|. \end{aligned}$$

This completes the proof of the theorem.  $\square$

A simple consequence of Theorem 4.4 is that the operators  $P_{n,j}$  and  $R_m$  are uniformly bounded on  $L^1$  and on  $L^\infty$  by a constant that depends only on  $c_1, c_2$  and  $k$ .

**Example 4.5.** Here we give an example to show that the assertion of Theorem 4.4 is not true in general if we split the set  $A_n$  into more than two subsets. We consider the case  $\Omega = [0, 1]$ , Lebesgue measure  $|\cdot|$ ,  $k = 1$  and  $S = \text{span}\{\mathbb{1}_\Omega\}$ . Let  $C_1 \cup C_2 \cup C_3$  be a partition of  $\Omega$  with the following properties:  $|C_1| = |C_2|$  and  $|C_3| = \varepsilon$  for some small parameter  $\varepsilon > 0$ . Define

$$h = \alpha_1 \mathbb{1}_{C_1} - \alpha_2 \mathbb{1}_{C_2} + \alpha_3 \mathbb{1}_{C_3}$$

where we choose the positive numbers  $\alpha_1, \alpha_2 \simeq 1$  and  $\alpha_3 \simeq |C_3|^{-1/2} = \varepsilon^{-1/2}$  such that  $\mathbb{E}h = 0$  and  $\mathbb{E}h^2 = 1$ . Let  $g$  be a function with  $g = 0$  on  $C_3$  that has the same sign than  $h$  on  $C_1 \cup C_2$ . Then, we evaluate for  $t \in C_3$

$$\langle g, h \rangle h(t) = \int_{C_1 \cup C_2} |g(x)| |h(x)| \, dx \cdot h(t) \gtrsim \varepsilon^{-1/2} \int_{C_1 \cup C_2} |g(x)| \, dx.$$

This implies that an inequality as in Theorem 4.4 is not possible as  $\varepsilon > 0$  was arbitrary. For more results in this direction, we refer to Section 9.

## 5. DECOMPOSITION OF $L^1$ -FUNCTIONS

Let  $T$  be a stopping time with respect to the filtration  $(\mathcal{F}_n)$ . Moreover, let  $f \in L^1$ . We set  $g_n := P_n f$  for  $n \geq -1$ . Then, we define  $g_T := (g_{T(x)})(x)$  if  $T(x) < \infty$  and we set  $g_T(x) = f(x)$  if  $T(x) = \infty$ .

**Proposition 5.1.** *Let  $T$  be a stopping time and  $f \in L^1$ . With the above notation, we have*

- (1)  $\mathbb{E}|g_T| \lesssim \mathbb{E}|f|$ .
- (2) If  $n \geq 0$  and  $j \in \{1, \dots, \ell_n\}$  we have

$$\mathbb{1}_{\{T \geq n\}}(P_{n,j} f) = \mathbb{1}_{\{T \geq n\}}(P_{n,j} g_T).$$

*Proof.* We have by (4.1) and the uniform  $L^1$ -boundedness of the operators  $P_{n,j}$

$$\begin{aligned} \mathbb{E}|g_T| &= \int_{\{T=\infty\}} |f| \, d\mathbb{P} + \sum_n \int_{\{T=n\}} |g_n| \, d\mathbb{P} \\ &\lesssim \int_{\{T=\infty\}} |f| \, d\mathbb{P} + \sum_n \int_{\{T=n\}} |f| \, d\mathbb{P} = \mathbb{E}|f|, \end{aligned}$$

which proves item (1).

Now we prove item (2): Since the set  $\{T \geq n\}$  is  $\mathcal{F}_{n-1}$ -measurable and by Remark 4.2, we see that

$$\mathbb{1}_{\{T \geq n\}}(P_{n,j} g_T) = P_{n,j}(\mathbb{1}_{\{T \geq n\}} g_T) = P_{n,j} \left( (\mathbb{1}_{\{T=\infty\}} + \sum_{\ell \geq n} \mathbb{1}_{\{T=\ell\}}) g_T \right)$$

$$= P_{n,j} \left( \mathbb{1}_{\{T=\infty\}} f + \sum_{\ell \geq n} \mathbb{1}_{\{T=\ell\}} P_\ell f \right)$$

with the series converging in  $L^1$ . By (4.1) and the continuity of  $P_{n,j}$  on  $L^1$ , we obtain

$$\begin{aligned} \mathbb{1}_{\{T \geq n\}}(P_{n,j} g_T) &= P_{n,j}(\mathbb{1}_{\{T=\infty\}} f) + \sum_{\ell \geq n} P_{n,j}(P_\ell(\mathbb{1}_{\{T=\ell\}} f)) \\ &= P_{n,j}(\mathbb{1}_{\{T=\infty\}} f) + \sum_{\ell \geq n} P_{n,j}(\mathbb{1}_{\{T=\ell\}} f), \end{aligned}$$

where the last equation is true since the range of the projector  $P_{n,j}$  is a subset of the range of the projector  $P_\ell$  for  $\ell \geq n$ . We thus obtain

$$\mathbb{1}_{\{T \geq n\}}(P_{n,j} g_T) = P_{n,j} \left( (\mathbb{1}_{\{T=\infty\}} + \sum_{\ell \geq n} \mathbb{1}_{\{T=\ell\}}) f \right) = P_{n,j}(\mathbb{1}_{\{T \geq n\}} f) = \mathbb{1}_{\{T \geq n\}}(P_{n,j} f),$$

since the set  $\{T \geq n\}$  is  $\mathcal{F}_{n-1}$  measurable.  $\square$

Utilizing the above results—in particular Theorem 4.4—we next show that an integrable function  $f$  admits a decomposition similar to Gundy's decomposition for martingales (see e.g. [11]).

**Theorem 5.2.** *Let  $f \in S_N$  for some positive integer  $N$  with  $\mathbb{E}|f| \leq 1$ . Then, for each  $\lambda > 0$  there exists a decomposition  $f = a + b + c$  with  $a, b, c \in L^1$  satisfying*

- (1)  $\|a\|_1 \lesssim 1$  and  $\mathbb{P}(\sup_m |da_m| \neq 0) \lesssim \lambda^{-1}$ .
- (2)  $\left\| \sum_m |db_m| \right\|_1 \lesssim 1$ .
- (3)  $\|c\|_1 \lesssim 1$  and  $\|c\|_\infty \lesssim \lambda$ .

*Proof.* Let  $\lambda > 0$  and let  $f \in S_N$  with  $\mathbb{E}|f| \leq 1$ . We recall the notation  $g_n = P_n f$  and the definition of the index set  $I_n = \{m : S_{n-1} \subsetneq V_m \subseteq S_n\}$ . Let  $c_3$  be the implicit constant from Theorem 4.4. Define the stopping time  $r$  with respect to  $(\mathcal{F}_n)$  to be

$$r = \inf\{n \geq 0 : \mathbb{E}_n |g_n| > \lambda/(4c_3) \text{ or } \max_{m \in I_n} \mathbb{E}_n |f_m| > \lambda\}.$$

Moreover, define  $v_{-1} = 0$  and

$$v_n = \sum_{m \in I_n} \mathbb{1}_{\{r=n\}} \mathbb{E}_n(|df_m|) = \sum_{m \in I_n} \mathbb{E}_n(\mathbb{1}_{\{r=n\}} |df_m|), \quad n \geq 0.$$

Let

$$s = \inf\{n \geq -1 : \sum_{\ell=-1}^n \mathbb{E}_\ell v_{\ell+1} + \sum_{\ell=0}^n v_\ell > \lambda\}.$$

Then,  $T := \min(r, s)$  is also a stopping time with respect to  $(\mathcal{F}_n)$ .

PART I: Define

$$a := f - g_T.$$

Then, Proposition 5.1 and the assumption  $\|f\|_1 \leq 1$  immediately implies  $\|a\|_1 \lesssim 1$ . If  $T(x) = \infty$  we have  $da_m(x) = 0$  for all  $m$  by item (2) of Proposition 5.1 and thus

$$\{\sup_m |da_m| \neq 0\} \subset \{T < \infty\} = \{r < \infty\} \cup \{s < \infty\}.$$

Therefore, by Theorem 4.4 and Doob's maximal inequality,

(5.1)

$$\mathbb{P}(r < \infty) \leq \mathbb{P}(\sup_{n \geq 0} \max_{m \in I_n} \mathbb{E}_n |f_m| > \lambda/(4c_3)) \leq \mathbb{P}(8c_3^2 \sup_{n \geq 0} \mathbb{E}_n |f| > \lambda) \leq 8c_3^2 \frac{\mathbb{E}|f|}{\lambda} \lesssim \lambda^{-1}.$$

Now we show that  $\mathbb{P}(s < \infty) \lesssim \lambda^{-1}$ . We begin by using Chebyshev's inequality:  
(5.2)

$$\mathbb{P}(s < \infty) = \mathbb{P}\left(\sum_{n=-1}^{\infty} (\mathbb{E}_n v_{n+1} + v_n) > \lambda\right) \leq \lambda^{-1} \sum_{n=-1}^{\infty} \mathbb{E}(\mathbb{E}_n(v_{n+1}) + v_n) \leq 2\lambda^{-1} \sum_{n=0}^{\infty} \mathbb{E}v_n.$$

In order to estimate the integral of  $v_n = \sum_{m \in I_n} \mathbb{E}_n(|df_m|) \mathbb{1}_{\{r=n\}}$ , we fix an index  $m \in I_n$  and write

$$(5.3) \quad \mathbb{E}(|df_m| \mathbb{1}_{\{r=n\}}) \leq \mathbb{E}(|f_m| \mathbb{1}_{\{r=n\}}) + \mathbb{E}(|f_{m-1}| \mathbb{1}_{\{r=n\}}).$$

Since for  $m \in I_n$ , the support of  $df_m$  is contained in  $A_n$  by Lemma 4.1, we know that  $\{r = n\} \subset A_n$ . If we assume that  $\{r = n\}$  is not empty, this implies by the  $\mathcal{F}_n$ -measurability of the set  $\{r = n\}$  that it can either be  $A_n''$ ,  $A_n'$  or  $A_n$  if  $n \geq 1$  and  $\{r = 0\}$  can only be  $\Omega$  if it is nonempty.

In order to continue the estimate in (5.3), we are going to show that  $\mathbb{E}(\mathbb{1}_{\{r=n\}}|f_m|) \lesssim \mathbb{E}(\mathbb{1}_{\{r=n\}}|g_n|)$  for  $m \in I_n$  or  $m$  such that  $f_m = g_{n-1}$ . To this end, we distinguish two possibilities for the sets  $\{r = n\}$ :

CASE 1:  $\{r = n\} = A_n$  with the atom  $A_n$  of  $\mathcal{F}_{n-1}$ : Since  $f_m = P_{n,j}g_n$  for some  $j$  and the operator  $P_{n,j}$  is uniformly bounded on  $L^1$ ,

$$(5.4) \quad \mathbb{E}_{n-1}(|f_m| \mathbb{1}_{\{r=n\}}) = \mathbb{1}_{\{r=n\}} \mathbb{E}_{n-1}|f_m| \lesssim \mathbb{1}_{\{r=n\}} \mathbb{E}_{n-1}|g_n| = \mathbb{E}_{n-1}(|g_n| \mathbb{1}_{\{r=n\}}).$$

CASE 2: Let  $A := \{r = n\} \in \{A_n', A_n''\}$ , which means that  $n \geq 1$  and  $B := (A_n \setminus A) \subset \{r > n\}$ . By Theorem 4.4,

$$(5.5) \quad \begin{aligned} \int_A |f_m| &\leq c_3 \int_A (\mathbb{E}_{n-1}|g_n| + \mathbb{E}_n|g_n|) = c_3 \left( \frac{|A|}{|A_n|} \int_{A_n} |g_n| + \int_A |g_n| \right) \\ &= c_3 \left( \frac{|A|}{|A| + |B|} \int_B |g_n| + \left(1 + \frac{|A|}{|A_n|}\right) \int_A |g_n| \right) \\ &\leq c_3 \left( \frac{|A|}{|A| + |B|} \int_B |g_n| + 2 \int_A |g_n| \right) \leq \frac{\lambda}{4} |A| + 2c_3 \int_A |g_n|, \end{aligned}$$

where in the last inequality, we used the inequality  $\int_B |g_n| \leq \lambda|B|/(4c_3)$ , which is implied by  $B \subset \{r > n\}$ . Since  $A = \{r = n\}$ , we know that either  $\mathbb{E}_n|g_n| > \lambda/(4c_3)$  on  $A$  or there exists  $m_1 \in I_n$  so that  $\mathbb{E}_n|f_{m_1}| > \lambda$  on  $A$ . In both cases, using (5.5) for  $m = m_1$  in the second case, we obtain the inequality

$$(5.6) \quad \int_A |g_n| \gtrsim \lambda|A|.$$

Using (5.6) we continue the calculation in (5.5) to deduce

$$\int_A |f_m| \lesssim \int_A |g_n|,$$

which implies

$$(5.7) \quad \mathbb{E}(\mathbb{1}_{\{r=n\}}|f_m|) = \mathbb{E}(\mathbb{1}_A|f_m|) \lesssim \mathbb{E}(\mathbb{1}_A|g_n|) = \mathbb{E}(\mathbb{1}_{\{r=n\}}|g_n|).$$

Combining cases 1 and 2, we continue the calculation in (5.3) to deduce

$$\mathbb{E}(|df_m| \mathbb{1}_{\{r=n\}}) \lesssim \mathbb{E}(|g_n| \mathbb{1}_{\{r=n\}}) \lesssim \mathbb{E}(|f| \mathbb{1}_{\{r=n\}}),$$

where the last inequality follows from  $P_n f = g_n$  and the fact that the operators  $P_n$  are uniformly bounded on  $L^1$ . Therefore, the definition of  $v_n$  implies

$$(5.8) \quad \sum_n \mathbb{E}v_n \lesssim \mathbb{E}|f| \leq 1.$$

Inserting this inequality in (5.2), we finally obtain

$$\mathbb{P}(s < \infty) \lesssim \lambda^{-1}.$$

PART II: Observe that

$$g_{T \wedge n} - g_{T \wedge (n-1)} = \mathbb{1}_{\{T \geq n\}}(g_n - g_{n-1}) = \mathbb{1}_{\{r \geq n\}} \mathbb{1}_{\{s \geq n\}}(g_n - g_{n-1}).$$

This gives

$$g_T = \sum_{n \geq 0} (g_{T \wedge n} - g_{T \wedge (n-1)}) = \sum_{n \geq 0} \sum_{m \in I_n} \mathbb{1}_{\{r \geq n\}} \mathbb{1}_{\{s \geq n\}} df_m.$$

Now, let  $n \geq 0$  and  $m \in I_n$  and define

$$\begin{aligned} \gamma_m &= \mathbb{1}_{\{r > n\}} \mathbb{1}_{\{s \geq n\}} df_m, \\ \delta_m &= \mathbb{1}_{\{r = n\}} \mathbb{1}_{\{s \geq n\}} df_m. \end{aligned}$$

Recall that by  $R_m$  we denote the orthoprojector onto the space  $V_m$  so that  $df_m = R_m f - R_{m-1} f$ . Then, observe that since  $\{r \geq n\}$  and  $\{s \geq n\}$  are  $\mathcal{F}_{n-1}$ -measurable, we have  $R_{m-1}(\gamma_m + \delta_m) = \mathbb{1}_{\{r \geq n\}} \mathbb{1}_{\{s \geq n\}} R_{m-1}(df_m) = 0$  by Remark 4.2. This means that  $\gamma_m + \delta_m$  is a difference sequence with respect to the operators  $(R_m)$ . Thus, the difference sequences

$$db_m = \delta_m - R_{m-1} \delta_m, \quad dc_m = \gamma_m + R_{m-1} \delta_m$$

define the functions  $b$  and  $c$ . Fix the parameter  $n$  and consider

$$\begin{aligned} \mathbb{E} \left[ \sum_{m \in I_n} |\delta_m| \right] &= \mathbb{E} \left[ \sum_{m \in I_n} \mathbb{1}_{\{r = n\}} \mathbb{1}_{\{s \geq n\}} |df_m| \right] \\ &\leq \mathbb{E} \left[ \sum_{m \in I_n} \mathbb{1}_{\{r = n\}} |df_m| \right] = \mathbb{E} v_n. \end{aligned}$$

Therefore, by estimate (5.8)

$$\sum_m \mathbb{E} |\delta_m| \lesssim \sum_n \mathbb{E} v_n \lesssim 1.$$

Since the operators  $R_m$  are uniformly bounded on  $L^1$ , this implies  $\mathbb{E} \sum_m |db_m| \lesssim 1$  and also,

$$\|c\|_1 = \left\| \sum_m dc_m \right\|_1 \leq \|g_T\|_1 + \left\| \sum_m |db_m| \right\|_1 \lesssim 1.$$

It remains to estimate the  $L^\infty$ -norm of  $c$ . We have

$$\sum_m \gamma_m = g_{(r-1) \wedge s}.$$

Let  $x \in \{r = n\}$  and for  $\ell < n$ , let  $B$  be the atom of  $\mathcal{F}_\ell$  that contains  $x$ . Since  $g_\ell \in S_\ell$ ,

$$\|g_\ell\|_B \lesssim (\mathbb{E}_\ell |g_\ell|)(x).$$

By the minimality of  $n$  in the definition of the stopping time  $r$ , this implies

$$\|g_\ell\|_B \lesssim \lambda.$$

This argument yields the inequality

$$\left\| \sum_m \gamma_m \right\|_\infty = \|g_{(r-1) \wedge s}\|_\infty \lesssim \lambda.$$

Now we estimate the second part of  $c$ . Since  $\{s \geq n\}$  is  $\mathcal{F}_{n-1}$ -measurable, by Remark 4.2,

$$\begin{aligned}
\left| \sum_m R_{m-1} \delta_m \right| &= \left| \sum_{n \geq 0} \sum_{m \in I_n} R_{m-1} (\mathbb{1}_{\{s \geq n\}} \mathbb{1}_{\{r=n\}}(df_m)) \right| \\
&= \left| \sum_{n \geq 0} \mathbb{1}_{\{s \geq n\}} \sum_{m \in I_n} R_{m-1} (\mathbb{1}_{\{r=n\}} df_m) \right| \\
&= \left| \sum_{n=0}^s \sum_{m \in I_n} R_{m-1} (\mathbb{1}_{\{r=n\}} df_m) \right| \\
&\leq \left| \sum_{n=0}^{s-1} \sum_{m \in I_n} R_{m-1} (\mathbb{1}_{\{r=n\}} df_m) \right| + \left| \sum_{n=s} \sum_{m \in I_n} R_{m-1} (\mathbb{1}_{\{r=n\}} df_m) \right| \\
&\leq c_3 \sum_{n=0}^{s-1} \sum_{m \in I_n} (\mathbb{E}_{n-1} + \mathbb{E}_n)(\mathbb{1}_{\{r=n\}} |df_m|) + \sum_{n=s} \sum_{m \in I_n} |R_{m-1} (\mathbb{1}_{\{r=n\}} df_m)|,
\end{aligned}$$

where the last inequality follows from Theorem 4.4. Looking at the definition of the stopping time  $s$ , we see that the first term is bounded by  $2c_3\lambda$ . For the second term we distinguish the cases where  $\{r = n\} = A_n$  is an atom of  $\mathcal{F}_{n-1}$  or  $\{r = n\}$  is only an atom of  $\mathcal{F}_n$ . Let  $m \in I_n$ . In the first case, we have by Remark 4.2 that

$$R_{m-1}(\mathbb{1}_{\{r=n\}} df_m) = \mathbb{1}_{\{r=n\}} R_{m-1}(df_m) = 0.$$

In the second case, we assume that  $A := \{r = n\} \in \{A'_n, A''_n\}$ , which yields that  $B := (A_n \setminus A) \subset \{r > n\}$ . Since

$$0 = R_{m-1}(\mathbb{1}_{A_n} df_m) = R_{m-1}(\mathbb{1}_A df_m) + R_{m-1}(\mathbb{1}_B df_m),$$

we get

$$|R_{m-1}(\mathbb{1}_{\{r=n\}} df_m)| = |R_{m-1}(\mathbb{1}_B df_m)|.$$

The fact that  $r > n$  on  $B$  now yields

$$\|\mathbb{1}_B df_m\|_\infty \leq \|\mathbb{1}_B f_m\|_\infty + \|\mathbb{1}_B f_{m-1}\|_\infty \lesssim \lambda$$

by definition of the stopping time  $r$  and inequality (1.2). Since the operator  $R_{m-1}$  is uniformly bounded on  $L^\infty$ , we obtain  $\|R_{m-1}(\mathbb{1}_B df_m)\|_\infty \lesssim \lambda$ . Inserting those estimates into the above, we get that

$$\left\| \sum_m R_{m-1} \delta_m \right\|_\infty \lesssim \lambda.$$

This finishes the proof of the theorem.  $\square$

## 6. UNCONDITIONALITY OF THE DIFFERENCES $df_m$

Once we know this Gundy-like decomposition, almost the same proof as in the martingale case yields the following result. We include it for completeness.

**Theorem 6.1.** *Let  $f \in L^1$  and let  $\tilde{f}_m = \sum_{j \leq m} \varepsilon_j df_j$  for some sequence of signs  $(\varepsilon_j)$ . Then*

$$(6.1) \quad \sup_{\lambda > 0} \lambda \mathbb{P}(\sup_m |\tilde{f}_m| > \lambda) \lesssim \mathbb{E}|f|.$$

*Proof.* It suffices to prove the result for functions in  $\cup_n S_n$ . Indeed, assuming the inequality for functions in  $\cup_n S_n$  and  $f \in L^1$ , we obtain

$$\mathbb{P}(\sup_m |\tilde{f}_m| > \lambda) = \lim_n \mathbb{P}(\sup_{m \leq n} |\tilde{f}_m| > \lambda) \lesssim \lim_n \inf \frac{\|f_n\|_1}{\lambda} \lesssim \frac{\|f\|_1}{\lambda},$$

where we used (6.1) for functions in  $S_n$  and the uniform boundedness of the operators  $(P_n)$  on  $L^1$ , respectively.

Therefore, we now prove the stated inequality for functions  $f \in \cup_n S_n$  using Theorem 5.2. Assume w.l.o.g. that  $\mathbb{E}|f| \leq 1$ . Then we decompose the function  $f$  by using Theorem 5.2 into  $f = a + b + c$ . Then, we see that  $\tilde{f}_m = \tilde{a}_m + \tilde{b}_m + \tilde{c}_m$  and we have

$$\mathbb{P}(\sup_m |\tilde{f}_m| > 3\lambda) \leq \mathbb{P}(\sup_m |\tilde{a}_m| > \lambda) + \mathbb{P}(\sup_m |\tilde{b}_m| > \lambda) + \mathbb{P}(\sup_m |\tilde{c}_m| > \lambda)$$

and we estimate each of those three parts separately.

- (1) If  $\sup_m |\tilde{a}_m| > \lambda$  this implies in particular that  $\sup_m |da_m| \neq 0$  and therefore

$$\mathbb{P}(\sup_m |\tilde{a}_m| > \lambda) \leq \mathbb{P}(\sup_m |da_m| \neq 0) \lesssim \lambda^{-1}$$

by item (1) of Theorem 5.2.

- (2) For the second term, we estimate

$$\sup_m |\tilde{b}_m| \leq \sum_m |db_m|,$$

which implies by Chebyshev's inequality

$$\mathbb{P}(\sup_m |\tilde{b}_m| > \lambda) \leq \lambda^{-1} \mathbb{E} \sum_m |db_m| \lesssim \lambda^{-1},$$

where the latter inequality is a consequence of item (2) in Theorem 5.2.

- (3) First we note that  $\tilde{c}_m = R_m \tilde{c}_N$ . Therefore, by Theorem 4.4 and Doob's inequality,

$$\|\sup_m |\tilde{c}_m|\|_2 \leq 2c_3 \|\tilde{c}_N\|_2.$$

Since the terms  $dc_j$  are orthogonal to each other for different indices  $j$ , we obtain

$$\|\tilde{c}_N\|_2 = \|c_N\|_2,$$

which yields by Chebyshev's inequality

$$\mathbb{P}(\sup_m |\tilde{c}_m| > \lambda) \leq \lambda^{-2} \mathbb{E} \sup_m |\tilde{c}_m|^2 \lesssim \lambda^{-2} \mathbb{E} |c_N|^2 \leq \lambda^{-2} \|c_N\|_\infty \mathbb{E} |c_N|.$$

Using now item (3) of Theorem 5.2 yields

$$\mathbb{P}(\sup_m |\tilde{c}_m| > \lambda) \lesssim \lambda^{-1}.$$

Combining those steps yields the assertion of the theorem.  $\square$

Now it is a simple task to prove the unconditionality in  $L^p$ ,  $1 < p < \infty$  for the differences  $(df_m)$ :

*Proof of Theorem 1.1.* Just use Theorem 6.1 and the Marcinkiewicz interpolation theorem.  $\square$

7. DEMOCRACY OF THE DIFFERENCES  $df_m$ 

We denote by  $\psi_{n,j}$  the unique (up to sign) function contained in  $V_{n,j}$  that is orthonormal to  $V_{n,j-1}$ . Recall the pointwise estimate from Lemma 4.3 which reads

$$(7.1) \quad |\psi_{n,j}| \lesssim \begin{cases} 0, & \text{on } A_n^c, \\ |A'_n|^{-1/2}, & \text{on } A'_n, \\ |A'_n|^{1/2}/|A''_n|, & \text{on } A''_n. \end{cases}$$

We use this estimate to perform the following calculation:

$$(7.2) \quad \begin{aligned} 1 = \|\psi_{n,j}\|_2^2 &= \int_{A'_n} |\psi_{n,j}|^2 d\mathbb{P} + \int_{A''_n} |\psi_{n,j}|^2 d\mathbb{P} \leq |A'_n| \|\psi_{n,j}\|_{A'_n}^2 + |A''_n| \|\psi_{n,j}\|_{A''_n}^2 \\ &\lesssim |A'_n| \|\psi_{n,j}\|_{A'_n}^2 + \frac{|A'_n|}{|A''_n|}, \end{aligned}$$

which yields that there exists a constant  $d > 0$  so that if  $|A'_n|/|A''_n| \leq d$ , we get the inequality

$$(7.3) \quad \|\psi_{n,j}\|_{A'_n} \simeq |A'_n|^{-1/2}.$$

For  $1 < p < \infty$ , Theorem 1.1 implies that

$$(7.4) \quad \left\| \sum_{n,j} a_{n,j} \psi_{n,j} \right\|_p \simeq \left\| \left( \sum_{n,j} |a_{n,j} \psi_{n,j}|^2 \right)^{1/2} \right\|_p$$

with constants that also depend on  $p$ .

Next, we want to check that we can also use some variant of the square function. Before we formulate this result we recall Stein's martingale inequality [12], which reads for  $1 < p < \infty$

$$(7.5) \quad \left\| \left( \sum_n |\mathbb{E}_n h_n|^2 \right)^{1/2} \right\|_p \lesssim \left\| \left( \sum_n |h_n|^2 \right)^{1/2} \right\|_p,$$

where  $h_n$  are arbitrary functions and the implied constant only depends on  $p$ .

**Lemma 7.1.** *For all  $1 < p < \infty$ , we have*

$$\left\| \left( \sum_{n,j} |a_{n,j} \psi_{n,j}|^2 \right)^{1/2} \right\|_p \simeq \left\| \left( \sum_{n,j} |a_{n,j}|^2 \frac{\mathbb{1}_{A'_n}}{|A'_n|} \right)^{1/2} \right\|_p,$$

where the implied constants depend on  $c_1, c_2, k$  and  $p$ .

*Proof.* Note that estimate (7.1) implies also that

$$|\psi_{n,j}| \lesssim \mathbb{E}_{n-1} \left( \frac{\mathbb{1}_{A'_n}}{|A'_n|^{1/2}} \right) \quad \text{on } A''_n.$$

Therefore, by Stein's inequality (7.5)

$$(7.6) \quad \left\| \left( \sum_{n,j} |a_{n,j} \psi_{n,j}|^2 \right)^{1/2} \right\|_p \lesssim \left\| \left( \sum_{n,j} |a_{n,j}|^2 \frac{\mathbb{1}_{A'_n}}{|A'_n|} \right)^{1/2} \right\|_p.$$

To get the converse inequality, first fix  $n$  and consider the two cases

CASE 1:  $|A'_n|/|A''_n| \leq d$ . Here we estimate for all  $j$ , using (7.3) and (1.2)

$$\frac{\mathbb{1}_{A'_n}}{|A'_n|^{1/2}} \lesssim \mathbb{E}_n |\psi_{n,j}|.$$



CASE 2:  $|A'_n|/|A''_n| > d$ . Here, by (7.2), we have

$$\|\psi_{n,j}\|_{A'_n} \gtrsim |A'_n|^{-1/2} \quad \text{or} \quad \|\psi_{n,j}\|_{A''_n} \gtrsim |A''_n|^{-1/2} \geq d^{-1/2}|A'_n|^{-1/2}.$$

Using again (1.2), this implies

$$\frac{\mathbb{1}_{A'_n}}{|A'_n|^{1/2}} \lesssim \mathbb{E}_{n-1}|\psi_{n,j}|.$$

Combining those two cases, we arrive at the inequality

$$\frac{\mathbb{1}_{A'_n}}{|A'_n|^{1/2}} \lesssim \mathbb{E}_{n-1}|\psi_{n,j}| + \mathbb{E}_n|\psi_{n,j}|.$$

Using Stein's inequality (7.5), we conclude

$$\left\| \left( \sum_{n,j} |a_{n,j}|^2 \frac{\mathbb{1}_{A'_n}}{|A'_n|} \right)^{1/2} \right\|_p \lesssim \left\| \left( \sum_{n,j} |a_{n,j}\psi_{n,j}|^2 \right)^{1/2} \right\|_p,$$

which, together with (7.6) completes the proof of the lemma.  $\square$

For  $1 < p < \infty$ , let  $\psi_{n,j}^p = \psi_{n,j}/\|\psi_{n,j}\|_p$  be the renormalized version of the functions  $\psi_{n,j}$ . Note that using the above lemma, we obtain

$$(7.7) \quad \|\psi_{n,j}\|_p \simeq |A'_n|^{1/p-1/2}.$$

Then, we prove the following theorem.

**Theorem 7.2.** *For  $1 < p < \infty$ , the functions  $(\psi_{n,j}^p)$  are democratic in  $L^p$ , i.e., we have for all finite sets  $\Lambda$  of indices  $(n, j)$  the equivalence*

$$\left\| \sum_{(n,j) \in \Lambda} \psi_{n,j}^p \right\|_p \simeq (\text{card } \Lambda)^{1/p},$$

where  $\text{card } \Lambda$  denotes the cardinality of  $\Lambda$ .

*Proof.* First, we remark that Lemma 7.1 and (7.7) imply

$$\left\| \left( \sum_{n,j} |a_{n,j}\psi_{n,j}^p|^2 \right)^{1/2} \right\|_p \simeq \left\| \left( \sum_{n,j} |a_{n,j}|^2 \frac{\mathbb{1}_{A'_n}}{|A'_n|^{2/p}} \right)^{1/2} \right\|_p.$$

Therefore, using also (7.4), we obtain

$$(7.8) \quad \left\| \sum_{(n,j) \in \Lambda} \psi_{n,j}^p \right\|_p^p \simeq \left\| \left( \sum_{(n,j) \in \Lambda} \frac{\mathbb{1}_{A'_n}}{|A'_n|^{2/p}} \right)^{1/2} \right\|_p^p = \int_{\Omega} \left( \sum_{(n,j) \in \Lambda} \frac{\mathbb{1}_{A'_n}}{|A'_n|^{2/p}} \right)^{p/2} d\mathbb{P}.$$

Note that for different indices  $n, m$  we have either  $A'_n \cap A'_m = \emptyset$  or one is strictly included in the other. Moreover, if  $A'_m \subsetneq A'_n$  we have geometric decay of measures  $|A'_m| \leq |A'_n|/2$ . Therefore, by a geometric series argument and the remark that for each  $n$ , there are at most  $k$  numbers  $j$  so that  $(n, j) \in \Lambda$ , we get pointwise

$$\left( \sum_{(n,j) \in \Lambda} \frac{\mathbb{1}_{A'_n}}{|A'_n|^{2/p}} \right)^{p/2} \simeq \left( \max_{(n,j) \in \Lambda} \frac{\mathbb{1}_{A'_n}}{|A'_n|^{2/p}} \right)^{p/2} = \max_{(n,j) \in \Lambda} \frac{\mathbb{1}_{A'_n}}{|A'_n|} \simeq \sum_{(n,j) \in \Lambda} \frac{\mathbb{1}_{A'_n}}{|A'_n|}.$$

Inserting this in (7.8), we obtain

$$\left\| \sum_{(n,j) \in \Lambda} \psi_{n,j}^p \right\|_p^p \simeq \int_{\Omega} \sum_{(n,j) \in \Lambda} \frac{\mathbb{1}_{A'_n}}{|A'_n|} d\mathbb{P} = \text{card } \Lambda,$$

which concludes the proof.  $\square$

If we use this theorem, together with Theorem 1.1 and Theorem 3.1 we get

**Corollary 7.3.** *For  $1 < p < \infty$ , the functions  $(\psi_{n,j}^p)$  are a greedy basis in the  $L^p$ -closure of  $\cup_n S_n$ .*

*If additionally  $|\bigcap_{f \in S} \{f = 0\}| = 0$  and  $\cup_n \mathcal{F}_n$  generates  $\mathcal{F}$ , the functions  $(\psi_{n,j}^p)$  form a greedy basis in  $L^p$ .*

## 8. INFINITE MEASURE SPACES

Let  $(\Omega, \mathcal{G}, \mathbf{m})$  be a measure space with  $\mathbf{m}(\Omega) = \infty$ . Let  $(\mathcal{G}_n)_{n \in \mathbb{Z}}$  be a filtration on  $\Omega$  such that for each  $n \in \mathbb{Z}$ , the  $\sigma$ -algebra  $\mathcal{G}_n$  is generated by  $\mathcal{G}_{n-1}$  and the subdivision of each atom  $A$  of  $\mathcal{G}_{n-1}$  into two atoms  $A', A''$  of  $\mathcal{G}_n$  having positive and finite measure.

Let  $\mathcal{A}_n$  be the collection of all atoms of  $\mathcal{G}_n$  and set  $\mathcal{A} := \cup_{n \in \mathbb{Z}} \mathcal{A}_n$ . Let  $S$  be a finite-dimensional space of  $\mathcal{G}$ -measurable functions on  $\Omega$  such that there exist constants  $c_1, c_2 \in (0, 1]$  so that for each  $f \in S$  and each  $A \in \mathcal{A}$  we have

$$|\{\omega \in A : |f(\omega)| \geq c_1 \|f\|_A\}| \geq c_2 |A|.$$

For each atom  $A \in \mathcal{A}$ , consider the spaces  $V_{A,m}$  satisfying

$$S(A) = V_{A,0} \subseteq \cdots \subseteq V_{A,\ell_A} = S(A') \oplus S(A'')$$

and  $\dim V_{A,m}/V_{A,m-1} = 1$  for each  $m = 1, \dots, \ell_A$ . Moreover, consider the functions  $\psi_{A,m} \in V_{A,m}$  satisfying  $\psi_{A,m} \perp V_{A,m-1}$  and  $\|\psi_{A,m}\|_2 = 1$ .

We now show that for each  $1 < p < \infty$ , the renormalized system  $\{\psi_{A,m}^p : A \in \mathcal{A}, 1 \leq m \leq \ell_A\}$ , given by  $\psi_{A,m}^p = \psi_{A,m}/\|\psi_{A,m}\|_p$ , is unconditional and democratic in  $L^p$ . We achieve this by using our results for probability spaces locally on atoms.

To this end, consider finitely many pairs  $\Lambda$  of indices  $(A, m)$  with  $A \in \mathcal{A}$  and  $1 \leq m \leq \ell_A$ . Choose  $n_0 \in \mathbb{Z}$  sufficiently small such that for each  $(A, m) \in \Lambda$ , there exists an atom  $B \in \mathcal{A}_{n_0}$  such that  $A \subseteq B$ . Let

$$\Gamma = \{B \in \mathcal{A}_{n_0} \mid \text{there exists } (A, m) \in \Lambda \text{ with } A \subseteq B\}.$$

For each  $B \in \Gamma$ , we consider the probability space  $(B, \mathcal{G} \cap B, \mathbf{m}(\cdot)/\mathbf{m}(B))$  and a binary filtration  $(\mathcal{F}_n^B)_{n \geq 0}$  on this probability space that contains of the  $\sigma$ -algebras  $(\mathcal{G}_n \cap B)_{n \geq n_0}$ . Moreover, we choose the spaces  $(V_{n,j})$  (as given in Section 1) corresponding to  $(\mathcal{F}_n^B)_{n \geq 0}$  such that

$$(8.1) \quad \{V_{n,j} : n \geq 0, 0 \leq j \leq \ell_n\} = \{V_{A,m} : A \in \mathcal{A}, A \subseteq B, 0 \leq m \leq \ell_A\}.$$

Define  $(\psi_{n,j})$  corresponding to the spaces  $(V_{n,j})$  as in the beginning of Section 7 and consider, for  $1 < p < \infty$ , the renormalized version  $\psi_{n,j}^p = \psi_{n,j}/\|\psi_{n,j}\|_p$ . Because of equality (8.1), the corresponding functions  $\psi_{n,j}$  and  $\psi_{A,m}$  also coincide and we deduce from the results in Sections 6 and 7 that for each fixed  $B \in \Gamma$ , the system

$$\{\psi_{A,m}^p : A \in \mathcal{A}, A \subseteq B, 1 \leq m \leq \ell_A\}$$

is unconditional and democratic in  $L^p$  with the same constants for each  $B \in \Gamma$ . Let  $f = \sum_{(A,m) \in \Lambda} a_{A,m} \psi_{A,m}^p$  for some coefficients  $(a_{A,m})$ . Then we estimate

$$\begin{aligned} \left\| \sum_{(A,m) \in \Lambda} \pm a_{A,m} \psi_{A,m}^p \right\|_p^p &= \left\| \sum_{B \in \Gamma} \sum_{(A,m) \in \Lambda: A \subseteq B} \pm a_{A,m} \psi_{A,m}^p \right\|_p^p \\ &= \sum_{B \in \Gamma} \left\| \sum_{(A,m) \in \Lambda: A \subseteq B} \pm a_{A,m} \psi_{A,m}^p \right\|_p^p, \end{aligned}$$

which is true since the sets  $B \in \Gamma$  are disjoint and the support of  $\psi_{A,m}^p$  is a subset of  $A$ . By the unconditionality of  $\{\psi_{A,m}^p : A \subseteq B, 1 \leq m \leq \ell_A\}$  for each  $B \in \Gamma$ , we obtain

$$\sum_{B \in \Gamma} \left\| \sum_{(A,m) \in \Lambda: A \subseteq B} \pm a_{A,m} \psi_{A,m}^p \right\|_p^p \simeq \sum_{B \in \Gamma} \|f \mathbb{1}_B\|_p^p.$$

The latter term is now equal to  $\|f\|_p^p$ , which shows the unconditionality of  $(\psi_{A,m}^p)$  in  $L^p$ . Similarly, by the uniform democracy of  $\{\psi_{A,m}^p : A \subseteq B, 1 \leq m \leq \ell_A\}$  for each  $B \in \Gamma$ ,

$$\begin{aligned} \left\| \sum_{(A,m) \in \Lambda} \psi_{A,m}^p \right\|_p^p &= \left\| \sum_{B \in \Gamma} \sum_{(A,m) \in \Lambda: A \subseteq B} \psi_{A,m}^p \right\|_p^p = \sum_{B \in \Gamma} \left\| \sum_{(A,m) \in \Lambda: A \subseteq B} \psi_{A,m}^p \right\|_p^p \\ &\simeq \sum_{B \in \Gamma} \text{card}\{(A,m) \in \Lambda : A \subseteq B\} = \text{card } \Lambda, \end{aligned}$$

which also shows the democracy of  $(\psi_{A,m}^p)$  in  $L^p$ .

## 9. NON-BINARY FILTRATIONS

In this last section, we expand on the setting when the filtration  $(\mathcal{F}_n)$  on the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  is not necessarily binary, meaning that an atom  $A_n$  of  $\mathcal{F}_{n-1}$  is divided into  $r = r(n) \geq 2$  atoms  $A_{n,1}, \dots, A_{n,r}$  of  $\mathcal{F}_n$ . We assume that for all  $n$ , the ordering of those atoms is such that  $|A_{n,1}| \leq \dots \leq |A_{n,r}|$  and that  $r(n)$  is bounded independently of  $n$ .

Define the intermediate  $\sigma$ -algebras  $\mathcal{F}_{n-1} = \mathcal{F}_{n,1} \subseteq \dots \subseteq \mathcal{F}_{n,r} = \mathcal{F}_n$  by the condition that for  $\mu = 1, \dots, r$  and restricted to  $A_n$ , the  $\sigma$ -algebra  $\mathcal{F}_{n,\mu}$  consists of the atoms

$$A_{n,1}, \dots, A_{n,\mu-1}, A_{n,\mu} \cup \dots \cup A_{n,r}.$$

Create a binary filtration  $(\mathcal{F}'_n)$  that consists precisely of the  $\sigma$ -algebras  $\mathcal{F}_{n,\mu}$  with  $n$  being a positive integer and  $1 \leq \mu \leq r(n)$ . Denote by  $\mathcal{A}, \mathcal{A}'$  the collection of atoms of the filtrations  $(\mathcal{F}_n)$  and  $(\mathcal{F}'_n)$  respectively.

We now show that if a space  $S$  satisfies (1.2) for the filtration  $(\mathcal{F}_n)$ , it also satisfies (1.2) for the binary filtration  $(\mathcal{F}'_n)$ . It is essential here that the refinement of  $\mathcal{F}_{n-1}$  to  $\mathcal{F}_n$  is done such that we take away the smallest remaining atom in each step.

**Proposition 9.1.** *Let  $S \subseteq L^\infty$  be a finite-dimensional vector space so that there exist constants  $c_1, c_2 \in (0, 1]$  such that for each atom  $A \in \mathcal{A}$  we have (1.2), i.e.,*

$$(9.1) \quad |\{\omega \in A : |f(\omega)| \geq c_1 \|f\|_A\}| \geq c_2 |A|, \quad f \in S.$$

*Then, there exists a constant  $c'_2 \in (0, 1]$ , depending only on  $c_2$  and  $\sup_n r(n)$ , such that for all  $A \in \mathcal{A}'$  we have*

$$(9.2) \quad |\{\omega \in A : |f(\omega)| \geq c_1 \|f\|_A\}| \geq c'_2 |A|, \quad f \in S.$$

*Proof.* Fix  $n$  and assume (9.1) for  $A \in \{A_n, A_{n,1}, \dots, A_{n,r}\}$ . Let  $d_1 = c_2$  and  $d_2 = d_1^2/(1 + d_1) = (1 - \alpha)d_1$  with  $\alpha = (1 + d_1)^{-1} < 1$ . We show that the set  $A' := A_{n,2} \cup \dots \cup A_{n,r}$  satisfies (9.2) with the constant  $c'_2 = d_2$ . To this end, we distinguish two cases:

CASE 1:  $|A_{n,1}| \leq \frac{d_1}{1+d_1} |A_n| = \alpha d_1 |A_n|$ : Here we estimate for  $f \in S$

$$\begin{aligned} |\{\omega \in A' : |f(\omega)| \geq c_1 \|f\|_{A'}\}| &\geq |\{\omega \in A' : |f(\omega)| \geq c_1 \|f\|_{A_n}\}| \\ &\geq (1 - \alpha) d_1 |A_n| \geq (1 - \alpha) d_1 |A'| = d_2 |A'|, \end{aligned}$$

where we used (9.1) for  $A = A_n$ .

CASE 2:  $|A_{n,1}| \geq \alpha d_1 |A_n|$ : Let  $\mu \in \{2, \dots, r\}$  be such that  $\|f\|_{A'} = \|f\|_{A_{n,\mu}}$ . Then we estimate

$$|\{\omega \in A' : |f(\omega)| \geq c_1 \|f\|_{A'}\}| \geq |\{\omega \in A_{n,\mu} : |f(\omega)| \geq c_1 \|f\|_{A_{n,\mu}}\}| \geq d_1 |A_{n,\mu}|,$$

where the last inequality follows from (9.1) with  $A = A_{n,\mu}$ . Continue with  $d_1 |A_{n,\mu}| \geq d_1 |A_{n,1}| \geq \alpha d_1^2 |A_n| \geq \alpha d_1^2 |A'| = d_2 |A'|$ .

Summarizing the two cases, we get inequality (9.2) for  $A = A'$  with the parameter  $c'_2 = d_2$ . Defining the numbers  $d_{j+1} = d_j^2 / (1 + d_j)$  recursively, we perform this argument inductively to obtain that all atoms  $A' \subseteq A_n$  in  $\mathcal{A}'$  satisfy inequality (9.2) with the parameter  $c'_2 = d_{r-1} \in (0, 1]$ .  $\square$

**Example 9.2.** Consider the space  $\Omega = [0, 1]$  and  $S = \text{span}\{p\}$  with  $p(t) = t$  for  $t \in [0, 1]$ . Divide  $[0, 1]$  into the partition of the intervals  $[0, \delta]$ ,  $(\delta, 1 - \varepsilon)$ ,  $[1 - \varepsilon, 1]$ . Then, Section 2 shows that (1.2) is satisfied in this case for the sets  $\{[0, 1], [0, \delta], (\delta, 1 - \varepsilon), [1 - \varepsilon, 1]\}$  with some uniform constants  $c_1, c_2 \in (0, 1]$  in  $\delta, \varepsilon$ .

On the other hand, it is also clear that for each choice of constants  $c'_1, c'_2 \in (0, 1]$  there are parameters  $\delta, \varepsilon > 0$  such that the converse of (1.2) is satisfied for the set  $K = [0, \delta] \cup [1 - \varepsilon, 1]$  and the function  $f = p_1$ , i.e., we have

$$|\{t \in K : p_1(t) \geq c'_1\}| < c'_2 |K| = c'_2 (\delta + \varepsilon).$$

Thus, if the splitting is not chosen in the way described above, for the intermediate atoms we not necessarily have (1.2) uniformly for all partitions.

Let  $S \subseteq L^\infty$  be a finite-dimensional vector space that satisfies (9.1) for all  $A \in \mathcal{A}$ . The above proposition shows that (9.2) is satisfied for all  $A \in \mathcal{A}'$  and some different constant  $c'_2$ . Since  $(\mathcal{F}'_n)$  is a binary filtration, we get from our previous results that the corresponding orthonormal system  $(\psi'_{n,j})$  (defined in the beginning of Section 7) is unconditional in  $L^p$  and its  $p$ -renormalization is democratic in  $L^p$  for  $1 < p < \infty$ . Therefore, for  $1 < p < \infty$  and the orthonormal system  $(\psi'_{n,j})$ , we have the estimate

$$(9.3) \quad \left\| \sum_{n,j} \pm \langle f, \psi'_{n,j} \rangle \psi'_{n,j} \right\|_p \leq C \|f\|_p, \quad f \in L^p$$

for some constant  $C$  independent of  $f$  and independent of the choice of signs. For each  $n$  and  $\mu = 1, \dots, r-1$ , let  $(\psi_{n,\mu,\ell})_\ell$  consist of those finitely many functions  $\psi'_{n,j}$  satisfying  $\psi'_{n,j} \in S(\mathcal{F}_{n,\mu+1})$  and  $\psi'_{n,j} \perp S(\mathcal{F}_{n,\mu})$ . Inequality (7.1) implies, for each  $\mu = 1, \dots, r-1$  and each  $\ell$ ,

$$(9.4) \quad |\psi_{n,\mu,\ell}| \lesssim \begin{cases} 0, & \text{on } (A_{n,\mu} \cup \dots \cup A_{n,r})^c, \\ |A_{n,\mu}|^{-1/2}, & \text{on } A_{n,\mu}, \\ |A_{n,\mu}|^{1/2} / |A_n|, & \text{on } A_{n,\mu+1} \cup \dots \cup A_{n,r}. \end{cases}$$

On the other hand, we choose, for each  $n$ , an  $L^2$ -normalized function  $\psi_{A_n}$  that is contained in  $S(\mathcal{F}_n)$  and orthogonal to  $S(\mathcal{F}_{n-1})$ , meaning that

$$(9.5) \quad \psi_{A_n} = \sum_{\mu=1}^{r-1} \sum_{\ell} a_{n,\mu,\ell} \psi_{n,\mu,\ell}$$

for some scalars  $(a_{n,\mu,\ell})$  with  $\sum_{\mu=1}^{r-1} \sum_{\ell} |a_{n,\mu,\ell}|^2 = 1$ . The number of terms in the inner sum over  $\ell$  is bounded from above by the dimension of the space  $S$ .

Having set up the notation, we now investigate the question whether the functions  $(\psi_{A_n})_n$  can also be unconditional/democratic in  $L^p$ ,  $1 < p < \infty$ . As we will see below,

this is not possible in general and this fact extends Example 4.5, where we showed that certain pointwise inequalities for  $\psi_{A_n}$  are not possible in general.

Since  $(\psi'_{n,j})$  is unconditional in  $L^p$  for  $1 < p < \infty$ , Lemma 7.1 implies

$$(9.6) \quad \begin{aligned} \|\psi_{A_n}\|_p &\simeq \left\| \left( \sum_{\mu=1}^{r-1} \sum_{\ell} |a_{n,\mu,\ell}|^2 \frac{\mathbb{1}_{A_{n,\mu}}}{|A_{n,\mu}|} \right)^{1/2} \right\|_p = \left( \sum_{\mu=1}^{r-1} |A_{n,\mu}|^{1-p/2} \left( \sum_{\ell} |a_{n,\mu,\ell}|^2 \right)^{p/2} \right)^{1/p} \\ &\simeq \sum_{\mu=1}^{r-1} |A_{n,\mu}|^{1/p-1/2} \sum_{\ell} |a_{n,\mu,\ell}|. \end{aligned}$$

In the analysis of the functions  $(\psi_{A_n})$ , the latter expressions are important, so we make the following definition.

**Definition 9.3.** Let  $1 \leq p \leq \infty$ . We say that the system  $(\psi_{A_n})$  satisfies condition  $(O_p)$ , if there exists a constant  $C$  such that for all  $n$  and all  $\mu, \nu = 1, \dots, r-1$ ,

$$\sum_{\ell} |a_{n,\mu,\ell}| \sum_m |a_{n,\nu,m}| |A_{n,\mu}|^{1/p-1/2} |A_{n,\nu}|^{1/2-1/p} \leq C,$$

We note that condition  $(O_2)$  is satisfied for any choice of functions  $(\psi_{A_n})$ . Moreover,  $(O_p)$  is satisfied if and only if  $(O_{p'})$  is satisfied with  $p' = p/(p-1)$ . Additionally, if  $(O_p)$  is satisfied for some  $p \in [1, 2)$ , then we also have  $(O_q)$  for all  $q \in (p, 2]$ .

Note that condition  $(O_p)$  is satisfied in particular if the measures of the atoms  $A_{n,1}, \dots, A_{n,r-1}$  are comparable with a constant independent of  $n$ . Moreover, if  $r = r(n) = 2$  for all  $n$ , condition  $(O_p)$  is also satisfied, since the measure  $|A_{n,r}|$  of the largest of the atoms  $A_{n,1}, \dots, A_{n,r}$  does not appear in  $(O_p)$ .

**Theorem 9.4.** Let  $1 < p < \infty$ . Then, the following assertions are true.

(1) If  $(\psi_{A_n})$  satisfies the estimate

$$(9.7) \quad \left\| \sum_n \pm \langle f, \psi_{A_n} \rangle \psi_{A_n} \right\|_p \leq C \|f\|_p, \quad f \in L^p,$$

for some constant  $C$  independent of  $f$  and the choice of signs, then condition  $(O_p)$  is satisfied.

(2) If condition  $(O_1)$  is satisfied, then  $(\psi_{A_n})$  satisfies estimate (9.7) for some constant  $C$  independent of  $f$  and the choice of signs.

(3) If condition  $(O_p)$  is satisfied, then the renormalized system  $(\psi_{A_n}^p)$  given by  $\psi_{A_n}^p = \psi_{A_n} / \|\psi_{A_n}\|_p$  is democratic in  $L^p$ .

Statement (2) in particular gives that  $(O_1)$  implies unconditionality of  $(\psi_{A_n})$  in  $L^p$ .

*Proof.* First consider statement (1). If  $(\psi_{A_n})$  satisfies (9.7), we have in particular that there exists a constant  $C$  such that for all  $n$  and all  $f \in L^p$

$$|\langle f, \psi_{A_n} \rangle| \cdot \|\psi_{A_n}\|_p \leq C \|f\|_p.$$

This condition is equivalent to the fact that for all  $n$

$$\|\psi_{A_n}\|_{p'} \cdot \|\psi_{A_n}\|_p \leq C.$$

with  $p' = p/(p-1)$  and the same constant  $C$ . Inserting (9.6) in this inequality implies condition  $(O_p)$ .

Next we prove statement (2). Observe that for fixed  $n$  and  $f \in L^p$  we have that

$$\langle f, \psi_{A_n} \rangle \psi_{A_n} = \sum_{\mu, \nu=1}^{r-1} \sum_{\ell, m} \overline{a_{n, \mu, \ell}} a_{n, \nu, m} \langle f, \psi_{n, \mu, \ell} \rangle \psi_{n, \nu, m}.$$

Using the square functions, the fact that  $(\psi'_{n, j}) = (\psi_{n, \mu, \ell})$  satisfies estimate (9.3) and Lemma 7.1, it is enough to show the following estimate for establishing (9.7):

$$(9.8) \quad \left\| \left( \sum_n \sum_{\nu=1}^{r-1} \sum_m \left| \sum_{\mu=1}^{r-1} \sum_{\ell} \pm \overline{a_{n, \mu, \ell}} a_{n, \nu, m} \langle f, \psi_{n, \mu, \ell} \rangle \right|^2 \frac{\mathbb{1}_{A_{n, \nu}}}{|A_{n, \nu}|} \right)^{1/2} \right\|_p \\ \lesssim \left\| \left( \sum_n \sum_{\mu=1}^{r-1} \sum_{\ell} |\langle f, \psi_{n, \mu, \ell} \rangle|^2 \frac{\mathbb{1}_{A_{n, \mu}}}{|A_{n, \mu}|} \right)^{1/2} \right\|_p.$$

To this end, we investigate, for fixed  $n$ , fixed  $\mu, \nu = 1, \dots, r-1$  and fixed  $\ell, m$  the expression

$$|a_{n, \mu, \ell} a_{n, \nu, m}|^2 \frac{\mathbb{1}_{A_{n, \nu}}}{|A_{n, \nu}|}$$

and consider the trivial  $\sigma$ -algebra  $\mathcal{G} = \mathcal{G}_{n, \mu, \nu} = \{\emptyset, A_{n, \mu} \cup A_{n, \nu}\}$  on  $A_{n, \mu} \cup A_{n, \nu}$ . We observe

$$\mathbb{E}_{\mathcal{G}} \mathbb{1}_{A_{n, \mu}} = \frac{|A_{n, \mu}|}{|A_{n, \mu}| + |A_{n, \nu}|} \mathbb{1}_{A_{n, \mu} \cup A_{n, \nu}}$$

and thus we write

$$\mathbb{1}_{A_{n, \mu} \cup A_{n, \nu}} = \mathbb{1}_{A_{n, \mu} \cup A_{n, \nu}}^2 = \frac{(|A_{n, \mu}| + |A_{n, \nu}|)^2}{|A_{n, \mu}|} \left( \mathbb{E}_{\mathcal{G}} \frac{\mathbb{1}_{A_{n, \mu}}}{|A_{n, \mu}|^{1/2}} \right)^2.$$

Then we estimate

$$|a_{n, \mu, \ell} a_{n, \nu, m}|^2 \frac{\mathbb{1}_{A_{n, \nu}}}{|A_{n, \nu}|} = |a_{n, \mu, \ell} a_{n, \nu, m}|^2 \left( \frac{\mathbb{1}_{A_{n, \nu}}}{|A_{n, \nu}|^{1/2}} \right)^2 \leq |a_{n, \mu, \ell} a_{n, \nu, m}|^2 \left( \frac{\mathbb{1}_{A_{n, \mu} \cup A_{n, \nu}}}{|A_{n, \nu}|^{1/2}} \right)^2 \\ = |a_{n, \mu, \ell} a_{n, \nu, m}|^2 \frac{(|A_{n, \mu}| + |A_{n, \nu}|)^2}{|A_{n, \mu}| |A_{n, \nu}|} \left( \mathbb{E}_{\mathcal{G}} \frac{\mathbb{1}_{A_{n, \mu}}}{|A_{n, \mu}|^{1/2}} \right)^2 \\ \leq 4C^2 \left( \mathbb{E}_{\mathcal{G}} \frac{\mathbb{1}_{A_{n, \mu}}}{|A_{n, \mu}|^{1/2}} \right)^2,$$

where the last inequality follows from  $(O_1)$ . Inserting this inequality in the left hand side of inequality (9.8) and using Stein's martingale inequality (7.5) for the  $\sigma$ -algebras  $\mathcal{G}_{n, \mu, \nu}$  gives the right hand side of (9.8). This finishes the proof of statement (2).

In order to prove statement (3), we first show that under condition  $(O_p)$ , we have the inequality

$$(9.9) \quad \sum_{\ell} |a_{n, \mu, \ell}| |A_{n, \mu}|^{1/p-1/2} \lesssim |A_{n, \mu_0}|^{1/p-1/2}$$

for all  $\mu = 1, \dots, r-1$ , where  $\mu_0 = \mu_0(n) \in \{1, \dots, r-1\}$  is chosen so that  $\sum_{\ell} |a_{n, \mu_0, \ell}| = \max_{\mu=1, \dots, r-1} \sum_{\ell} |a_{n, \mu, \ell}|$ . Without restriction, assume that  $p \leq 2$  so that  $1/p - 1/2 \geq 0$ . Observe that due to the assumption  $\sum_{\mu=1}^{r-1} \sum_{\ell} |a_{n, \mu, \ell}|^2 = 1$  we have  $\sum_{\ell} |a_{n, \mu_0, \ell}| \simeq 1$ . We distinguish two cases for the value of  $\mu$ .

CASE 1:  $\mu \leq \mu_0$ . In this case we have  $|A_{n, \mu}| \leq |A_{n, \mu_0}|$  and condition  $(O_p)$  implies

$$\sum_{\ell} |a_{n, \mu, \ell}| |A_{n, \mu}|^{1/p-1/2} \lesssim \frac{|A_{n, \mu}|^{2(1/p-1/2)}}{|A_{n, \mu_0}|^{1/p-1/2}} \leq |A_{n, \mu}|^{1/p-1/2} \leq |A_{n, \mu_0}|^{1/p-1/2}.$$

proving (9.9) for  $\mu \leq \mu_0$ .

CASE 2:  $\mu > \mu_0$ . In this case we have  $|A_{n,\mu_0}| \leq |A_{n,\mu}|$  and condition  $(O_p)$  implies

$$\sum_{\ell} |a_{n,\mu,\ell}| |A_{n,\mu}|^{1/p-1/2} \lesssim |A_{n,\mu_0}|^{1/p-1/2},$$

proving (9.9) also for  $\mu > \mu_0$ .

Therefore, by (9.6) and (9.9),

$$(9.10) \quad \|\psi_{A_n}\|_p \simeq \sum_{\mu=1}^{r-1} |A_{n,\mu}|^{1/p-1/2} \sum_{\ell} |a_{n,\mu,\ell}| \simeq |A_{n,\mu_0}|^{1/p-1/2}.$$

Now, we write

$$\psi_{A_n}^p = \sum_{\mu=1}^{r-1} \sum_{\ell} b_{n,\mu,\ell} \psi_{n,\mu,\ell}^p$$

with

$$b_{n,\mu,\ell} = a_{n,\mu,\ell} \frac{\|\psi_{n,\mu,\ell}\|_p}{\|\psi_{A_n}\|_p} \quad \text{and} \quad \psi_{n,\mu,\ell}^p = \frac{\psi_{n,\mu,\ell}}{\|\psi_{n,\mu,\ell}\|_p}.$$

Note that for all  $\mu = 1, \dots, r-1$ ,

$$\sum_{\ell} |b_{n,\mu,\ell}| = \sum_{\ell} |a_{n,\mu,\ell}| \frac{\|\psi_{n,\mu,\ell}\|_p}{\|\psi_{A_n}\|_p} \simeq \sum_{\ell} |a_{n,\mu,\ell}| \frac{|A_{n,\mu}|^{1/p-1/2}}{|A_{n,\mu_0}|^{1/p-1/2}}$$

by (9.10). Therefore, (9.9) implies  $\sum_{\ell} |b_{n,\mu,\ell}| \lesssim 1$  for all  $\mu = 1, \dots, r-1$  and  $\sum_{\ell} |b_{n,\mu_0,\ell}| \simeq 1$ . Let now  $\Lambda$  be a finite set of positive integers. Then, this bound on the numbers  $(b_{n,\mu,\ell})$  and the unconditionality and democracy of the functions  $(\psi_{n,\mu,\ell}^p)$  gives us

$$\left\| \sum_{n \in \Lambda} \psi_{A_n}^p \right\|_p = \left\| \sum_{n \in \Lambda} \sum_{\mu=1}^{r-1} \sum_{\ell} b_{n,\mu,\ell} \psi_{n,\mu,\ell}^p \right\|_p \gtrsim \left\| \sum_{n \in \Lambda} \sum_{\ell} |b_{n,\mu_0,\ell}| \psi_{n,\mu_0,\ell}^p \right\|_p \gtrsim (\text{card } \Lambda)^{1/p}.$$

Similarly, the converse inequality  $\|\sum_{n \in \Lambda} \psi_{A_n}^p\|_p \lesssim (\text{card } \Lambda)^{1/p}$  is a consequence of  $\sum_{\ell} |b_{n,\mu,\ell}| \lesssim 1$ , the unconditionality and democracy of the functions  $(\psi_{n,\mu,\ell}^p)$ , where we also use that the number  $r = r(n)$  is uniformly bounded in  $n$  and the number of terms in the sum over  $\ell$  is bounded by the dimension of the space  $S$ .  $\square$

**Proposition 9.5.** *Suppose that the functions  $\psi_{A_n} = \sum_{\mu=1}^{r-1} \sum_{\ell} a_{n,\mu,\ell} \psi_{n,\mu,\ell}$  satisfy the pointwise inequality*

$$|\psi_{A_n}| \lesssim \frac{|A_{n,1}|^{1/2}}{|A_{n,\kappa}|} \quad \text{on } A_{n,\kappa}$$

for each  $\kappa = 1, \dots, r$ .

Then, for each  $1 < p < \infty$ , the system  $(\psi_{A_n})$  satisfies (9.7) and the renormalized system  $(\psi_{A_n}^p)$  is democratic in  $L^p$ .

*Proof.* We show condition  $(O_1)$ , i.e., we show for each  $\mu, \nu = 1, \dots, r-1$  the inequality

$$(9.11) \quad \sum_{\ell} |a_{n,\mu,\ell}| \sum_m |a_{n,\nu,m}| |A_{n,\mu}|^{1/2} |A_{n,\nu}|^{-1/2} \leq C,$$

for some constant  $C$  independent of  $n$ . For each  $\mu = 1, \dots, r-1$  and each  $\ell$ , we observe that the pointwise estimates for  $\psi_{A_n}$  in the assumptions and the pointwise estimates

(9.4) for  $\psi_{n,\mu,\ell}$  imply

$$\begin{aligned} |a_{n,\mu,\ell}| &= |\langle \psi_{A_n}, \psi_{n,\mu,\ell} \rangle| \leq \sum_{\kappa=\mu}^r \int_{A_{n,\kappa}} |\psi_{A_n}| \cdot |\psi_{n,\mu,\ell}| \\ &\lesssim |A_{n,\mu}| \cdot \frac{|A_{n,1}|^{1/2}}{|A_{n,\mu}|} \cdot |A_{n,\mu}|^{-1/2} + \sum_{\kappa=\mu+1}^r |A_{n,\kappa}| \cdot \frac{|A_{n,1}|^{1/2}}{|A_{n,\kappa}|} \cdot \frac{|A_{n,\mu}|^{1/2}}{|A_n|} \\ &= \frac{|A_{n,1}|^{1/2}}{|A_{n,\mu}|^{1/2}} + \frac{(r-\mu) \cdot |A_{n,1}|^{1/2} |A_{n,\mu}|^{1/2}}{|A_n|} \lesssim \frac{|A_{n,1}|^{1/2}}{|A_{n,\mu}|^{1/2}}. \end{aligned}$$

Inserting this inequality in the left hand side of (9.11), we obtain (9.11) since  $|A_{n,1}| \leq |A_{n,\mu}|$  for all  $\mu = 1, \dots, r-1$ . Thus, Theorem 9.4 now implies that (9.7) is true for  $(\psi_{A_n})$  and  $(\psi_{A_n}^p)$  is democratic in  $L^p$ .  $\square$

We now use this result to give an explicit example of an orthonormal system of functions that does not fall in the category of functions of the form  $(\psi'_{n,j})$  as considered above, but is still unconditional and democratic in  $L^p$  for  $1 < p < \infty$ . The example we want to consider consists of local tensor products of functions. Assume that we are given probability spaces  $(\Omega_\delta, \mathcal{F}^\delta, \mathbb{P}^\delta)$  for  $\delta = 1, \dots, d$ . Assume that  $(\mathcal{F}_n)$  is a filtration on  $\Omega_1 \times \dots \times \Omega_d$  with  $\mathcal{F}_0 = \{\emptyset, \Omega_1 \times \dots \times \Omega_d\}$  and  $\mathcal{F}_n$  is generated by  $\mathcal{F}_{n-1}$  and the partition of one atom  $A_n = A_n^1 \times \dots \times A_n^d$  of  $\mathcal{F}_{n-1}$  into the atoms  $A_{n,m_1}^1 \times \dots \times A_{n,m_d}^d$  with  $m_\delta \in \{1, 2\}$  for each  $\delta = 1, \dots, d$  and the same ordering

$$|A_{n,1}^\delta| \leq |A_{n,2}^\delta|, \quad \delta = 1, \dots, d$$

as above in each direction  $\delta$ . Denote by  $\mathcal{A}_n$  the atoms of  $\mathcal{F}_n$  and  $\mathcal{A} = \cup_n \mathcal{A}_n$ . Let  $p_\delta : \Omega_1 \times \dots \times \Omega_d \rightarrow \Omega_\delta$  be the projection onto the  $\delta$ -th coordinate. Then, let  $\mathcal{G}^\delta := p_\delta(\mathcal{A})$  be the collection of subsets of  $\Omega_\delta$  that appear as the  $\delta$ -th coordinate projection of some atom  $A \in \mathcal{A}$ . Moreover, assume that we are given finite-dimensional subspaces  $S^\delta \subset L^\infty(\Omega_\delta)$  that satisfy (1.2) for each  $G_\delta \in \mathcal{G}^\delta$ , i.e.,

$$(9.12) \quad |\{\omega_\delta \in G_\delta : |f(\omega_\delta)| \geq c_1 \|f\|_{G_\delta}\}| \geq c_2 |G_\delta|, \quad f \in S^\delta.$$

Then we consider the tensor product space  $S := S^1 \otimes \dots \otimes S^d$  on  $\Omega_1 \times \dots \times \Omega_d$ . We now show that  $S$  satisfies (1.2) for some constants  $c'_1, c'_2 \in (0, 1]$  and for every  $G = G_1 \times \dots \times G_d$  with  $G_\delta \in \mathcal{G}^\delta$ , i.e.,

$$(9.13) \quad |\{\omega \in G : |f(\omega)| \geq c'_1 \|f\|_G\}| \geq c'_2 |G|, \quad f \in S.$$

It is enough to show that for each  $G$  of this form and each  $f \in S$ , the mean value of  $|f|$  on  $G$  is comparable to the  $L^\infty$ -norm of  $f$  on  $G$ . Let  $d \geq 2$  and  $f \in S$ . Let  $x \in G_1 \times \dots \times G_{d-1}$  and  $y \in G_d$  and let  $f_x(y) = f_y(x) = f(x, y)$ . The latter functions are measurable in the respective measure spaces. Then, using the notation  $G' = G_1 \times \dots \times G_{d-1}$ , we calculate by Fubini's theorem and the fact that on the coordinate projection  $G_d$ , mean values of positive functions in  $S^d$  are comparable to their  $L^\infty$ -norms by (9.12),

$$\begin{aligned} \frac{1}{|G|} \int_G |f(x, y)| \, d(x, y) &= \frac{1}{|G'|} \int_{G'} \frac{1}{|G_d|} \int_{G_d} |f(x, y)| \, dy \, dx \\ &\gtrsim \frac{1}{|G'|} \int_{G'} \|f_x\|_{G_d} \, dx \geq \left\| \frac{1}{|G'|} \int_{G'} |f_y| \, dx \right\|_{G_d}. \end{aligned}$$

By induction, this implies that the mean of  $|f|$  on  $G$  is comparable to the  $L^\infty$ -norm of  $f$  on  $G$ . Thus, we have established (9.13) for some constants  $c'_1, c'_2 \in (0, 1]$ .



Then we get, for each  $n$  and in each direction  $\delta = 1, \dots, d$  an orthonormal basis  $(\psi_{n,0,j_\delta}^\delta)_{j_\delta}$  of  $S^\delta(A_n^\delta)$  and an orthonormal basis  $(\psi_{n,1,j_\delta}^\delta)_{j_\delta}$  of the orthogonal complement of  $S^\delta(A_n^\delta)$  in  $S^\delta(A_{n,1}^\delta) \oplus S^\delta(A_{n,2}^\delta)$ . Then we consider the orthonormal system  $(\psi_{n,\mu,j})$  for indices  $\mu = (\mu_1, \dots, \mu_d)$  and  $j = (j_1, \dots, j_d)$  with  $\mu \neq 0$  consisting of the functions

$$(9.14) \quad \psi_{n,\mu,j} = \psi_{n,\mu_1,j_1}^1 \otimes \cdots \otimes \psi_{n,\mu_d,j_d}^d.$$

For fixed  $n$ , those functions are orthogonal to the space  $S^1(A_n^1) \otimes \cdots \otimes S^d(A_n^d)$ .

We use Lemma 4.3 to obtain the following estimate for every  $\delta = 1, \dots, d$  and every  $j_\delta$ ,

$$(9.15) \quad |\psi_{n,1,j_\delta}^\delta| \lesssim \begin{cases} 0, & \text{on } (A_n^\delta)^c, \\ |A_{n,1}^\delta|^{-1/2}, & \text{on } A_{n,1}^\delta, \\ |A_{n,1}^\delta|^{1/2}/|A_{n,2}^\delta|, & \text{on } A_{n,2}^\delta. \end{cases}$$

In the case  $\mu_\delta = 0$  we have the estimate

$$|\psi_{n,0,j_\delta}^\delta| \lesssim |A_n^\delta|^{-1/2} \quad \text{on } A_n^\delta,$$

which is a consequence of (9.12). Assume  $\mu_1 = \cdots = \mu_s = 0$  and  $\mu_{s+1} = \cdots = \mu_d = 1$  and  $m_{s+1} = \cdots = m_t = 1$ ,  $m_{t+1} = \cdots = m_d = 2$ . Then, we estimate on the set  $A_n^1 \times \cdots \times A_n^s \times A_{n,m_{s+1}}^{s+1} \times \cdots \times A_{n,m_d}^d$  as follows:

$$(9.16) \quad \begin{aligned} |\psi_{n,\mu,j}| &\lesssim (|A_n^1| \cdots |A_n^s| |A_{n,1}^{s+1}| \cdots |A_{n,1}^t|)^{-1/2} \left( \frac{|A_{n,1}^{t+1}|^{1/2}}{|A_{n,2}^{t+1}|} \cdots \frac{|A_{n,1}^d|^{1/2}}{|A_{n,2}^d|} \right) \\ &= \frac{|A_n^1 \times \cdots \times A_n^s \times A_{n,1}^{s+1} \times \cdots \times A_{n,1}^d|^{1/2}}{|A_n^1 \times \cdots \times A_n^s \times A_{n,1}^{s+1} \times \cdots \times A_{n,1}^t \times A_{n,2}^{t+1} \times \cdots \times A_{n,2}^d|}. \end{aligned}$$

For each fixed value of  $\mu \in \{0, 1\}^d \setminus \{0\}$  and  $j$  we now define a filtration on  $\Omega_1 \times \cdots \times \Omega_d$  so that we can use the pointwise estimate (9.16) to show the assumption of Proposition 9.5 for the sequence of functions  $(\psi_{n,\mu,j})_n$  defined in (9.14). Up to permutation of variables, we assume without restriction that  $\mu_1 = \cdots = \mu_s = 0$  and  $\mu_{s+1} = \cdots = \mu_d = 1$  for some  $s \in \{1, \dots, d\}$ . Then we introduce the intermediate  $\sigma$ -algebra  $\mathcal{F}_n^\mu$  satisfying  $\mathcal{F}_{n-1} \subseteq \mathcal{F}_n^\mu \subseteq \mathcal{F}_n$  by defining the atoms of  $\mathcal{F}_n^\mu$  to be the sets

$$A_n^1 \times \cdots \times A_n^s \times A_{n,m_{s+1}}^{s+1} \times \cdots \times A_{n,m_d}^d, \quad (m_{s+1}, \dots, m_d) \in \{1, 2\}^{d-s}.$$

Then, the function  $\psi_{n,\mu,j}$  is contained in the orthogonal complement of  $S(\mathcal{F}_{n-1})$  in  $S(\mathcal{F}_n^\mu)$ . Consider the filtration  $(\widetilde{\mathcal{F}}_n)$  consisting of the  $\sigma$ -algebras  $(\mathcal{F}_n)_n$  and  $(\mathcal{F}_n^\mu)_n$  and let  $\mathcal{A}$  be the collection of atoms of  $(\widetilde{\mathcal{F}}_n)$ . We showed above that (1.2) is satisfied for every atom  $A \in \mathcal{A}$  and every function  $f \in S$ . Observe also that estimate (9.16) implies the assumption of Proposition 9.5 for the filtration  $(\widetilde{\mathcal{F}}_n)$ . Therefore, we can use our theory, in particular Proposition 9.5, to deduce that for every  $f \in L^p$ , we have the estimate

$$\left\| \sum_n \pm \langle f, \psi_{n,\mu,j} \rangle \psi_{n,\mu,j} \right\|_p \leq C \|f\|_p,$$

where the constant  $C$  depends on  $p$  and on  $\mu, j$ , but not on  $f$ . Since there are only uniformly finitely many values of  $\mu, j$ , we deduce that the orthonormal system  $(\psi_{n,\mu,j})_{n,\mu,j}$  is unconditional in  $L^p$ .

Additionally, Proposition 9.5 implies that for each  $\mu, j$ , the renormalized system  $(\psi_{n,\mu,j}^p)_n$  is democratic in  $L^p$ . Choose a finite set  $\Lambda$  of values  $(n, \mu, j)$ . Since  $(\psi_{n,\mu,j}^p)_{n,\mu,j}$

is unconditional in  $L^p$ , we get

$$\left\| \sum_{(n,\mu,j) \in \Lambda} \psi_{n,\mu,j}^p \right\|_p \simeq \left\| \left( \sum_{(n,\mu,j) \in \Lambda} (\psi_{n,\mu,j}^p)^2 \right)^{1/2} \right\|_p \simeq \left\| \sum_{\mu,j} \left( \sum_{n:(n,\mu,j) \in \Lambda} (\psi_{n,\mu,j}^p)^2 \right)^{1/2} \right\|_p,$$

where the latter equivalence is a consequence of the fact that the number of indices  $(\mu, j)$  is uniformly bounded in  $n$ . Thus, there exist values  $(\mu_0, j_0)$  such that we have

$$\left\| \sum_{(n,\mu,j) \in \Lambda} \psi_{n,\mu,j}^p \right\|_p \simeq \left\| \left( \sum_{n:(n,\mu_0,j_0) \in \Lambda} (\psi_{n,\mu_0,j_0}^p)^2 \right)^{1/2} \right\|_p \simeq (\text{card } \Lambda)^{1/p},$$

implying the democracy of the system  $(\psi_{n,\mu,j}^p)_{n,\mu,j}$ .

**Acknowledgments.** M. Passenbrunner is supported by the Austrian Science Fund FWF, project P32342.

## REFERENCES

- [1] Ju. A. Brudnyĭ and M. I. Ganzburg. A certain extremal problem for polynomials in  $n$  variables. *Izv. Akad. Nauk SSSR Ser. Mat.*, 37:344–355, 1973.
- [2] N. Fontes-Merz. A multidimensional version of Turán’s lemma. *J. Approx. Theory*, 140(1):27–30, 2006.
- [3] M. I. Ganzburg. Polynomial inequalities on measurable sets and their applications. *Constructive Approximation*, 17(2):275, 2001.
- [4] G. G. Gevorkyan and A. Kamont. Unconditionality of general Franklin systems in  $L^p[0, 1]$ ,  $1 < p < \infty$ . *Studia Math.*, 164(2):161–204, 2004.
- [5] J. Gulgowski, A. Kamont, and M. Passenbrunner. Properties of local orthonormal systems, Part II: Geometric characterization of Bernstein inequalities. *arXiv:2304.05647*, 2023.
- [6] A. Kamont. General Haar systems and greedy approximation. *Studia Math.*, 145(2):165–184, 2001.
- [7] S. V. Konyagin and V. N. Temlyakov. A remark on greedy approximation in Banach spaces. *East J. Approx.*, 5(3):365–379, 1999.
- [8] F. L. Nazarov. Local estimates for exponential polynomials and their applications to inequalities of the uncertainty principle type. *Algebra i Analiz*, 5(4):3–66, 1993.
- [9] M. Passenbrunner. Unconditionality of orthogonal spline systems in  $L^p$ . *Studia Math.*, 222(1):51–86, 2014.
- [10] P. Petrushev. Multivariate  $n$ -term rational and piecewise polynomial approximation. *J. Approx. Theory*, 121(1):158–197, 2003.
- [11] G. Pisier. *Martingales in Banach spaces*, volume 155 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2016.
- [12] E. M. Stein. *Topics in harmonic analysis related to the Littlewood-Paley theory*. Annals of Mathematics Studies, No. 63. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1970.
- [13] V. Temlyakov. *Greedy approximation*, volume 20 of *Cambridge Monographs on Applied and Computational Mathematics*. Cambridge University Press, Cambridge, 2011.

FACULTY OF MATHEMATICS, PHYSICS AND INFORMATICS, UNIVERSITY OF GDAŃSK, UL. WITA STWOSZA 57, 80-308 GDAŃSK, POLAND

*Email address:* jacek.gulgowski@ug.edu.pl

INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, BRANCH IN GDAŃSK, UL. ABRAHAMAMA 18, 81-825 SOPOT, POLAND

*Email address:* Anna.Kamont@impan.pl

INSTITUTE OF ANALYSIS, JOHANNES KEPLER UNIVERSITY LINZ, AUSTRIA, 4040 LINZ, ALTENBERGER STRASSE 69

*Email address:* markus.passenbrunner@jku.at, markus.passenbrunner@gmail.com