

ULYANOV INEQUALITIES FOR THE MIXED MODULI OF SMOOTHNESS IN MIXED METRICS ¹

B.V. Simonov¹, A.A. Jumabayeva²

¹ *Volgograd State Technical University
Volgograd, 400131, Russia*

E-mail: simonov-b2002@yandex.ru

² *L.N.Gumilyov Eurasian National University
Satpayev 2, 010000 Astana, Kazakhstan*

E-mail: ainurjumabay@gmail.com

In this paper we prove Ulyanov-type inequalities between mixed moduli of smoothness of positive orders in different metrics.

Keywords: Mixed moduli of smoothness of positive order, Ulyanov type inequality.

1. Introduction

Let $L_{p_1 p_2}$, $1 \leq p_i \leq \infty$, $i = 1, 2$ be the set of measurable functions of two variables $f(x_1, x_2)$, 2π -periodic in each variable, for which $\|f\|_{p_1 p_2} = \|\{|f|\}_{p_1}\|_{p_2} < \infty$, where

$$\|f\|_{p_i} = \left(\int_0^{2\pi} |f|^{p_i} dx_i \right)^{\frac{1}{p_i}}, \text{ if } 1 \leq p_i < \infty,$$

$$\|f\|_{p_i} = \sup_{0 \leq x_i \leq 2\pi} |f|, \text{ if } p_i = \infty.$$

Let $L_{p_1 p_2}^0$ be the space of functions $f \in L_{p_1 p_2}$ such that $\int_0^{2\pi} f(x_1, x_2) dx_1 = 0$ for almost all x_2 and $\int_0^{2\pi} f(x_1, x_2) dx_2 = 0$ for almost all x_1 .

For the function $f \in L_{p_1 p_2}$, we define the fractional differences of positive order α_1 and α_2 with steps h_1 and h_2 respectively, by variables x_1 and x_2 as follows:

$$\Delta_{h_1}^{\alpha_1}(f) = \sum_{\nu_1=0}^{\infty} (-1)^{\nu_1} \binom{\alpha_1}{\nu_1} f(x_1 + (\alpha_1 - \nu_1)h_1, x_2),$$

$$\Delta_{h_2}^{\alpha_2}(f) = \sum_{\nu_2=0}^{\infty} (-1)^{\nu_2} \binom{\alpha_2}{\nu_2} f(x_1, x_2 + (\alpha_2 - \nu_2)h_2),$$

where $\binom{\alpha}{\nu} = 1$ for $\nu = 0$, $\binom{\alpha}{\nu} = \alpha$ for $\nu = 1$, $\binom{\alpha}{\nu} = \frac{\alpha(\alpha-1)\dots(\alpha-\nu+1)}{\nu!}$ for $\nu \geq 2$.

Denote (cf. [12]) by $\omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{p_1 p_2}$ the mixed modulus of smoothness of positive orders α_1 and α_2 , respectively, in the variables x_1 and x_2 of a function $f \in L_{p_1 p_2}$, that is,

$$\omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{p_1 p_2} = \sup_{|h_i| \leq \delta_i, i=1,2} \|\Delta_{h_1}^{\alpha_1}(\Delta_{h_2}^{\alpha_2}(f))\|_{p_1 p_2}.$$

Let $Y_{m_1, m_2}(f)_{p_1 p_2}$ be the best approximation by the angle of the function $f \in L_{p_1 p_2}$, i.e.

$$Y_{m_1, m_2}(f)_{p_1 p_2} = \inf_{T_{m_1, \infty}, T_{\infty, m_2}} \|f - T_{\infty, m_2} - T_{m_1, \infty}\|_{p_1 p_2},$$

¹This research was partially funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. AP09260223).

where $T_{m_1, \infty}(x_1, x_2)$ is a trigonometric polynomial of order at most m_1 ($m_1 \in \mathbb{N} \cup \{0\}$) with respect to x_1 and such that $T_{m_1, \infty} \in L_{p_1, p_2}$, $T_{\infty, m_2}(x_1, x_2)$ is a trigonometric polynomial of order at most m_2 ($m_2 \in \mathbb{N} \cup \{0\}$) with respect to x_2 and such that $T_{\infty, m_2} \in L_{p_1, p_2}$.

Let $f^{(\rho_1, \rho_2)}(x_1, x_2)$ ($\rho_1 \geq 0, \rho_2 \geq 0$) be a derivative in the sense of Weyl of the function $f(x_1, x_2) \in L_{p_1, p_2}^0$ of order ρ_1 with respect to x_1 and of order ρ_2 with respect to x_2 (see [1], p. 238).

For a given number q , we will use the notation q^* , which will be defined as follows: $q^* = q$ if $1 \leq q < \infty$ and $q^* = 1$ if $q = \infty$.

For non-negative functionals $F(f, \delta_1, \delta_2)$ and $G(f, \delta_1, \delta_2)$ we write that $F(f, \delta_1, \delta_2) \ll G(f, \delta_1, \delta_2)$, if there exists a positive constant C , independent of f, δ_1 and δ_2 such that $F(f, \delta_1, \delta_2) \leq CG(f, \delta_1, \delta_2)$. If $F(f, \delta_1, \delta_2) \ll G(f, \delta_1, \delta_2)$ and $G(f, \delta_1, \delta_2) \ll F(f, \delta_1, \delta_2)$, then we will write that $F(f, \delta_1, \delta_2) \asymp G(f, \delta_1, \delta_2)$.

Denote by $\omega_\alpha(f, t)_p^{(1)}$ the moduli of smoothness of a fractional order α , $\alpha > 0$, of the function $f \in L_p^{(1)}$, i.e., $\omega_\alpha(f, t)_p^{(1)} = \sup_{|h| \leq t} \|\Delta_h^\alpha f(x)\|_p^{(1)}$, where

$$\Delta_h^\alpha f(x) = \sum_{\nu=0}^{\infty} (-1)^\nu \binom{\alpha}{\nu} f(x + (\alpha - \nu)h).$$

The following (p, q) -inequality between moduli of smoothness in different metrics, nowadays called sharp Ulyanov type inequalities, is known (see [23]):

$$\omega_1(f, \delta)_q^{(1)} \ll \left(\int_0^\delta \left(t^{-\theta} \omega_1(f, t)_p^{(1)} \right)^{q^*} \frac{dt}{t} \right)^{\frac{1}{q^*}},$$

where $1 \leq p < q \leq \infty, \theta = \frac{1}{p} - \frac{1}{q}$. For the Lebesgue spaces, Ulyanov type inequalities in the one-dimensional case have been studied for the moduli of smoothness of any positive order by many authors, in particular, Ulyanov [29], V.I. Kolyada [9], W. Trebels [28], O. Domingues [3], B. Simonov [23], S. Tikhonov [24].

In the multidimensional case Ulyanov type inequalities were considered by many authors. Inequalities for the total moduli of smoothness of two variables were obtained in [3], [6], [10], [11], [23], and [24]. Ulyanov type inequalities were also proved for mixed moduli of smoothness in ([12]-[16]). The (p, q) inequalities for the moduli of smoothness of derivatives in terms of moduli of smoothness of the function itself were established in [7], [10], [27]. Moreover, such inequalities were handled in various spaces in [2], [4], and [5].

In this paper, we study Ulyanov-type inequalities for mixed moduli of smoothness. For the case when $p_1 = p_2 = p$, the following relation between the mixed moduli of smoothness is known ([12]-[16]). Let $f \in L_{pp}^0$, $\alpha_i > 0, \delta_i \in (0, 1)$ ($i = 1, 2$). Then

a) for $1 < p < q < \infty$

$$\omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{qq} \ll \left(\int_0^{\delta_2} \int_0^{\delta_1} \left(t_1^{-\frac{1}{p} + \frac{1}{q}} t_2^{-\frac{1}{p} + \frac{1}{q}} \omega_{\alpha_1 + \frac{1}{p} - \frac{1}{q}, \alpha_2 + \frac{1}{p} - \frac{1}{q}}(f, t_1, t_2)_{pp} \right)^q \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right)^{\frac{1}{q}},$$

b) for $1 = p < q = \infty$

$$\omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{\infty\infty} \ll \int_0^{\delta_2} \int_0^{\delta_1} (t_1 t_2)^{-1} \omega_{\alpha_1 + 1, \alpha_2 + 1}(f, t_1, t_2)_{11} \frac{dt_1}{t_1} \frac{dt_2}{t_2},$$

c) for $1 = p < q < \infty$

$$\begin{aligned} & \omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{qq} \ll \\ & \ll \left(\int_0^{\delta_2(\log_2 \frac{2}{\delta_2})^{\frac{1}{\alpha_2 q}}} \int_0^{\delta_1(\log_2 \frac{2}{\delta_1})^{\frac{1}{\alpha_1 q}}} ((t_1 t_2)^{-1 + \frac{1}{q}} \omega_{\alpha_1 + 1 - \frac{1}{q}, \alpha_2 + 1 - \frac{1}{q}}(f, t_1, t_2)_{11})^q \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right)^{\frac{1}{q}}, \end{aligned}$$

d) for $1 < p < q = \infty$

$$\omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{\infty\infty} \ll \int_0^{\delta_2(\log_2 \frac{2}{\delta_2})^{\frac{1}{\alpha_2 p'}}} \int_0^{\delta_1(\log_2 \frac{2}{\delta_1})^{\frac{1}{\alpha_1 p'}}} (t_1 t_2)^{-\frac{1}{p}} \omega_{\alpha_1 + \frac{1}{p}, \alpha_2 + \frac{1}{p}}(f, t_1, t_2)_{pp} \frac{dt_1}{t_1} \frac{dt_2}{t_2}.$$

Our main goal is to prove Ulyanov-type inequalities between mixed moduli of smoothness of positive orders, in particular, we strengthen the above inequalities. In this work, statements a) and b) are extended to the relation between mixed moduli of smoothness in the case when $p_1 \neq p_2$. Let us formulate the main results.

Theorem 1 *Let $f \in L_{p_1 p_2}^0$, where $1 < p_1 < q_1 < \infty$ or $1 = p_1 < q_1 = \infty$ and $1 < p_2 < q_2 < \infty$ or $1 = p_2 < q_2 = \infty$. Let for $\alpha_i > 0, \delta_i \in (0, 1), i = 1, 2$, we have*

$$\begin{aligned} & \omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{q_1 q_2} \ll \\ & \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}}. \end{aligned} \quad (1)$$

Remark 1 *Theorem 1 is optimal in the sense that there exists a function $f_0(x_1, x_2)$ such that for it in relation (1) the sign \ll can be replaced by the sign \asymp .*

Remark 2 *In relation (1), generally speaking, it is impossible to replace the sign \ll with the sign \asymp .*

Let us now estimate the mixed moduli of smoothness of the derivative of functions in terms of the mixed moduli of smoothness of the function itself.

Theorem 2 *Let $f \in L_{p_1 p_2}^0$, where $1 < p_1 < q_1 < \infty$ or $1 = p_1 < q_1 = \infty$ and $1 < p_2 < q_2 < \infty$ or $1 = p_2 < q_2 = \infty$. Let for $\alpha_i > 0, \rho_i \geq 0, \delta_i \in (0, 1), i = 1, 2$. we have*

$$\begin{aligned} & \omega_{\alpha_1, \alpha_2}(f^{(\rho_1, \rho_2)}, \delta_1, \delta_2)_{q_1 q_2} \ll \\ & \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\rho_1 - \frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\rho_2 - \frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \rho_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \rho_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

The paper is organized as follows: in the next section, we provide some basic notions and results that will be needed throughout the paper. In Sect. 3, we prove Theorems 1 and 2.

2. Some auxiliary results

In this section, we give some notations and lemmas which will be used in the proof of our main results. Let $L_p^{(1)}, 1 \leq p \leq \infty$ be the space of 2π -periodic measurable functions for which $\|f\|_p^{(1)} < \infty$,

where $\|f\|_p^{(1)} = \left(\int_0^{2\pi} |f|^p dx \right)^{\frac{1}{p}}$, when $1 \leq p < \infty$, and $\|f\|_p^{(1)} = \sup_{0 \leq x \leq 2\pi} |f|$, if $p = \infty$. Let $L_p^{0(1)}$ be set of functions $f \in L_p^{(1)}$ such that $\int_0^{2\pi} f(x) dx = 0$.

Let $\sigma(f)^{(1)}$ be the Fourier series of the functions $f(x)$ i.e.

$$\sigma(f)^{(1)} \equiv \sigma(f, x)^{(1)} \equiv \frac{a_0}{2} + \sum_{k=1}^{\infty} (a_k \cos kx + b_k \sin kx) \equiv \sum_{k=0}^{\infty} A_k(x),$$

where a_k and b_k are the Fourier coefficients of the function $f(x)$.

Let $\tilde{f}(x)$ be the conjugate function of $f(x)$ and $f^{(\alpha)}(x)$ be the Weyl derivative of the function $f(x)$ of order $\alpha > 0$.

$T_n(\{\lambda_n\}; x)$ is the trigonometric polynomial $T_n(x) = \sum_{\nu=0}^n (c_\nu \cos \nu x + d_\nu \sin \nu x)$ transformed by the sequence $\{\lambda_n\}$ ([8], [30]) i.e.

$$T_n(\{\lambda_n\}; x) = \sum_{\nu=0}^n \lambda_\nu (c_\nu \cos \nu x + d_\nu \sin \nu x).$$

Let $V_{(1)m}(f)$ denote the de la Vallée-Poussin sum of the Fourier series of a function $f(x)$,

$$V_{(1)m}(f) = \frac{1}{\pi} \int_0^{2\pi} f(x+t) V_m^{2m}(t) dt, \quad m = 0, 1, 2, \dots$$

The two-dimensional de la Vallée-Poussin sums of the Fourier series of a function $f(x_1, x_2)$ are defined as follows

$$V_{m_1, \infty}(f) = \frac{1}{\pi} \int_0^{2\pi} f(x_1 + t_1, x_2) V_{m_1}^{2m_1}(t_1) dt_1,$$

$$V_{\infty, m_2}(f) = \frac{1}{\pi} \int_0^{2\pi} f(x_1, x_2 + t_2) V_{m_2}^{2m_2}(t_2) dt_2,$$

$$V_{m_1, m_2}(f) = \frac{1}{\pi^2} \int_0^{2\pi} \int_0^{2\pi} f(x_1 + t_1, x_2 + t_2) V_{m_1}^{2m_1}(t_1) V_{m_2}^{2m_2}(t_2) dt_1 dt_2 \quad (m_i = 0, 1, 2, \dots, i = 1, 2),$$

where $V_0^0(t) = D_0(t)$, $V_n^{2n}(t) = \frac{D_n(t) + \dots + D_{2n-1}(t)}{n}$, $n = 1, 2, \dots$, $D_m(t) = \frac{\sin(m + \frac{1}{2})t}{2 \sin \frac{t}{2}}$, $m = 0, 1, 2, \dots$

We need the following lemmas.

Lemma 1 [17]. Let $f \in L_{p_1 p_2}^0$, $1 \leq p_i \leq \infty$, $\alpha_i > 0$, $n_i \in \mathbb{N}$, $i = 1, 2$. Then

$$\begin{aligned} \omega_{\alpha_1, \alpha_2} \left(f, \frac{1}{n_1}, \frac{1}{n_2} \right)_{p_1 p_2} &\asymp n_1^{-\alpha_1} n_2^{-\alpha_2} \left\| V_{n_1, n_2}^{(\alpha_1, \alpha_2)}(f) \right\|_{p_1 p_2} + n_1^{-\alpha_1} \left\| V_{n_1, \infty}^{(\alpha_1, 0)}(f - V_{\infty, n_2}(f)) \right\|_{p_1 p_2} \\ &+ n_2^{-\alpha_2} \left\| V_{\infty, n_2}^{(0, \alpha_2)}(f - V_{n_1, \infty}(f)) \right\|_{p_1 p_2} + \left\| f - V_{n_1, \infty}(f) - V_{\infty, n_2}(f) + V_{n_1, n_2}(f) \right\|_{p_1 p_2}, \\ \left\| f - V_{n_1-1, \infty}(f) - V_{\infty, n_2-1}(f) + V_{n_1-1, n_2-1}(f) \right\|_{p_1 p_2} &\ll Y_{n_1-1, n_2-1}(f)_{p_1 p_2}. \end{aligned}$$

Lemma 2 [16]. Let $a_n \geq 0, b_n \geq 0$, $\sum_{k=1}^n a_k = a_n \gamma_n$, $1 \leq p < \infty$. Then

$$\sum_{k=1}^{\infty} \left(\sum_{n=k}^{\infty} b_n \right)^p \ll \sum_{k=1}^{\infty} a_k (b_k \gamma_k)^p.$$

Lemma 3 [10, 12, 17]. Let $f \in L_{p_1 p_2}^0, g \in L_{p_1 p_2}^0, 1 \leq p_i \leq \infty, \beta_i > \alpha_i > 0, n_i \in N, i = 1, 2$. Then

- (1) $\omega_{\alpha_1, \alpha_2}(f, \delta_1, 0)_{p_1 p_2} = \omega_{\alpha_1, \alpha_2}(f, 0, \delta_2)_{p_1 p_2} = \omega_{\alpha_1, \alpha_2}(f, 0, 0)_{p_1 p_2} = 0$.
- (2) $\omega_{\alpha_1, \alpha_2}(f + g, \delta_1, \delta_2)_{p_1 p_2} \ll \omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{p_1 p_2} + \omega_{\alpha_1, \alpha_2}(g, \delta_1, \delta_2)_{p_1 p_2}$.
- (3) $\omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{p_1 p_2} \ll \omega_{\alpha_1, \alpha_2}(f, t_1, t_2)_{p_1 p_2}$, if $0 < \delta_i \leq t_i, i = 1, 2$.
- (4) $\frac{\omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{p_1 p_2}}{\delta_1^{\alpha_1} \delta_2^{\alpha_2}} \ll \frac{\omega_{\alpha_1, \alpha_2}(f, t_1, t_2)_{p_1 p_2}}{t_1^{\alpha_1} t_2^{\alpha_2}}$, if $0 < t_i \leq \delta_i \leq \pi, i = 1, 2$.
- (5) $\omega_{\alpha_1, \alpha_2}(f, \lambda_1 \delta_1, \lambda_2 \delta_2)_{p_1 p_2} \ll (\lambda_1 + 1)^{\alpha_1} (\lambda_2 + 1)^{\alpha_2} \omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{p_1 p_2}$, if $\lambda_i > 0, i = 1, 2$.
- (6) $Y_{n_1-1, n_2-1}(f)_{p_1 p_2} \ll \omega_{\alpha_1, \alpha_2}(f, \frac{1}{n_1}, \frac{1}{n_2})_{p_1 p_2} \ll$
 $\frac{1}{n_1^{\alpha_1}} \frac{1}{n_2^{\alpha_2}} \sum_{v_1=1}^{n_1+1} \sum_{v_2=1}^{n_2+1} v_1^{\alpha_1-1} v_2^{\alpha_2-1} Y_{v_1-1, v_2-1}(f)_{p_1 p_2}$.
- (7) $\omega_{\beta_1, \beta_2}(f, \delta_1, \delta_2)_{p_1 p_2} \ll \omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{p_1 p_2}$.
- (8) $\frac{\omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{p_1 p_2}}{\delta_1^{\alpha_1} \delta_2^{\alpha_2}} \ll \frac{\omega_{\beta_1, \beta_2}(f, \delta_1, \delta_2)_{p_1 p_2}}{\delta_1^{\beta_1} \delta_2^{\beta_2}}$, if $0 < \delta_i \leq \pi, i = 1, 2$.
- (9) $\delta_1^{-r_1} \omega_{\alpha_1+r_1, \alpha_2}(f, \delta_1, \delta_2)_{p_1 p_2} \ll \omega_{\alpha_1, \alpha_2}(f^{(r_1, 0)}, \delta_1, \delta_2)_{p_1 p_2} \ll$
 $\ll \int_0^{\delta_1} t_1^{-r_1} \omega_{\alpha_1+r_1, \alpha_2}(f, t_1, \delta_2)_{p_1 p_2} \frac{dt_1}{t_1}$, if $0 < \delta_1 < \pi$ and $f^{(r_1, 0)} \in L_{p_1 p_2}^0$,
 $\delta_2^{-r_2} \omega_{\alpha_1, \alpha_2+r_2}(f, \delta_1, \delta_2)_{p_1 p_2} \ll \omega_{\alpha_1, \alpha_2}(f^{(0, r_2)}, \delta_1, \delta_2)_{p_1 p_2} \ll$
 $\ll \int_0^{\delta_2} t_2^{-r_2} \omega_{\alpha_1, \alpha_2+r_2}(f, \delta_1, t_2)_{p_1 p_2} \frac{dt_2}{t_2}$, if $0 < \delta_2 < \pi$ and $f^{(0, r_2)} \in L_{p_1 p_2}^0$,
 $\delta_2^{-r_2} \delta_1^{-r_1} \omega_{\alpha_1+r_1, \alpha_2+r_2}(f, \delta_1, \delta_2)_{p_1 p_2} \ll \omega_{\alpha_1, \alpha_2}(f^{(r_1, r_2)}, \delta_1, \delta_2)_{p_1 p_2} \ll$
 $\ll \int_0^{\delta_2} \int_0^{\delta_1} t_2^{-r_2} t_1^{-r_1} \omega_{\alpha_1+r_1, \alpha_2+r_2}(f, t_1, t_2)_{p_1 p_2} \frac{dt_2}{t_2}$, if $0 < \delta_i < \pi (i = 1, 2)$ and $f^{(r_1, r_2)} \in L_{p_1 p_2}^0$.

Lemma 4 [18]. Let $f \in L_{p_1 p_2}^0, 1 \leq p_i < q_i \leq \infty, n_i = 0, 1, 2, \dots, i = 1, 2$. Then

$$Y_{2^{n_1-1}, 2^{n_2-1}}(f)_{q_1 q_2} \ll \left(\sum_{\nu_2=n_2}^{\infty} \left(\sum_{\nu_1=n_1}^{\infty} \left(2^{\nu_1 \left(\frac{1}{p_1} - \frac{1}{q_1} \right)} 2^{\nu_2 \left(\frac{1}{p_2} - \frac{1}{q_2} \right)} Y_{2^{\nu_1-1}, 2^{\nu_2-1}}(f)_{p_1 p_2} \right)^{q_1^*} \right)^{\frac{q_2^*}{q_1^*}} \right)^{\frac{1}{q_2^*}}.$$

Lemma 5 [19]. (a) Let the function $T_{\infty, 2^{n_2}} \in L_{p_1 p_2}, 1 \leq p_i \leq \infty (i = 1, 2)$ be a trigonometric polynomial of order 2^{n_2} in variable $x_2, 1 \leq p_2 < q_2 \leq \infty$. Then

$$\left\| \sum_{n_2=M_1}^{M_2} T_{\infty, 2^{n_2}} \right\|_{p_1 q_2} \leq C \left(\sum_{n_2=M_1}^{M_2} 2^{n_2 q_2^* \left(\frac{1}{p_2} - \frac{1}{q_2} \right)} \|T_{\infty, 2^{n_2}}\|_{p_1 p_2}^{q_2^*} \right)^{1/q_2^*},$$

where constant C does not depend on N_1, N_2 and $T_{2^{n_1}, \infty}$.

(b) Let the function $T_{2^{n_1}, \infty} \in L_{p_1 p_2}, 1 \leq p_i \leq \infty (i = 1, 2)$ be a trigonometric polynomial of order 2^{n_1} in variable $x_1, 1 \leq p_1 < q_1 \leq \infty$. Then

$$\left\| \sum_{n_1=N_1}^{N_2} T_{2^{n_1}, \infty} \right\|_{q_1 p_2} \leq C \left(\sum_{n_1=N_1}^{N_2} 2^{n_1 q_1^* \left(\frac{1}{p_1} - \frac{1}{q_1} \right)} \|T_{2^{n_1}, \infty}\|_{p_1 p_2}^{q_1^*} \right)^{1/q_1^*},$$

where constant C does not depend on N_1, N_2 and $T_{2^{n_1}, \infty}$.

Lemma 6 [20]. Let $f \in L_{p_1 p_2}^0, 1 \leq p_i \leq \infty, n_i = 0, 1, 2, \dots, i = 1, 2$. Then

- (a) $\|V_{\infty 2^{n_2}}^{(0, \alpha_2)}(f - V_{2^{n_1} \infty}(f))\|_{p_1 p_2} \ll \left\| \sum_{\nu_1=n_1}^{\infty} V_{\infty 2^{\nu_2}}^{(0, \alpha_2)}(V_{2^{\nu_1} \infty}(f) - V_{2^{\nu_1+1} \infty}(f)) \right\|_{p_1 p_2}$,
- (b) $\|V_{2^{n_1} \infty}^{(\alpha_1, 0)}(f - V_{\infty 2^{n_2}}(f))\|_{p_1 p_2} \ll \left\| \sum_{\nu_2=n_2}^{\infty} V_{2^{n_1} \infty}^{(\alpha_1, 0)}(V_{\infty 2^{\nu_2}}(f) - V_{\infty 2^{\nu_2+1}}(f)) \right\|_{p_1 p_2}$.

Lemma 7 ([30], p. 213). Let $f(x) \in L_p^{0(1)}$, $1 < p < q < \infty$, $\theta = \frac{1}{p} - \frac{1}{q}$, $\alpha \geq 0$. Then

$$\|f^{(\alpha)}\|_q^{(1)} \ll \|f^{(\alpha+\theta)}\|_p^{(1)}.$$

Lemma 8 [20]. (a) Let the function $T_{n_1, \infty}(x_1, x_2) \in L_{p_1 p_2}^0$, $1 = p_1 < q_1 = \infty$, $1 \leq p_2 \leq \infty$ be a trigonometric polynomial of order at most n_1 ($n_1 \in \mathbb{N}$), in the variable x_1 . Then

$$\|T_{n_1, \infty}\|_{\infty p_2} \leq C_1 \|T_{n_1, \infty}^{(1,0)}\|_{1, p_2},$$

where the constant C_1 does not depend on $T_{n_1, \infty}$.

(b) Let the function $T_{\infty, n_2}(x_1, x_2) \in L_{p_1 p_2}^0$, $1 = p_2 < q_2 = \infty$, $1 \leq p_1 \leq \infty$, be a trigonometric polynomial of order at most n_2 ($n_2 \in \mathbb{N}$), in the variable x_2 . Then

$$\|T_{\infty, n_2}\|_{p_1 \infty} \leq C_2 \|T_{\infty, n_2}^{(0,1)}\|_{p_1, 1},$$

where the constant C_2 does not depend on T_{∞, n_2} .

Lemma 9 ([21], p. 13). Let $\alpha > 0$, $f(x) \in L_1^{0(1)}$, $\sigma(f)^{(1)} = \sum_{k=1}^{\infty} A_k(x)$, where

$A_k(x) = a_k \cos kx + b_k \sin kx$. Let the function $\varphi(x)$ be such that $\sigma(\varphi)^{(1)} = \sum_{k=1}^{\infty} k^\alpha A_k(x)$. Then

$$\varphi(x) = \cos \frac{\pi\alpha}{2} f^{(\alpha)}(x) + \sin \frac{\pi\alpha}{2} \tilde{f}^{(\alpha)}(x).$$

Lemma 10 [22]. Let $0 < \alpha < 1$, $n \in \mathbb{N}$, $0 < |x| \leq \pi$. Then

$$\left| \sum_{\nu=1}^n \frac{\cos \nu x}{\nu^\alpha} \right| \leq C(\alpha) \cdot |x|^{\alpha-1},$$

$$\left| \sum_{\nu=1}^n \frac{\sin \nu x}{\nu^\alpha} \right| \leq C(\alpha) \cdot |x|^{\alpha-1},$$

where the positive constant $C(\alpha)$ does not depend on n and x , but depends only on α .

Lemma 11 ([1], p. 31). (Hardy-Littlewood inequality). Let $1 < p < q < \infty$, $\mu = 1 - \frac{1}{p} + \frac{1}{q}$, $f \in L_p^{(1)}(\mathbb{R}_1)$, $\mathbb{R}_1 = (-\infty, +\infty)$, that is $\|f\|_{L_p^{(1)}(\mathbb{R}_1)} = \left(\int_{-\infty}^{\infty} |f(x)|^p dx \right)^{\frac{1}{p}} < \infty$,

$$J(x) = \int_{\mathbb{R}_1} f(y) |y - x|^{-\mu} dy.$$

Then the inequality is true

$$\|J\|_{L_q^{(1)}(\mathbb{R}_1)} \leq K(p, q) \|f\|_{L_p^{(1)}(\mathbb{R}_1)},$$

where the constant $K(p, q)$ depends only on p and q .

Lemma 12 Let $1 < p_1 < q_1 < \infty$, $1 \leq p_2 \leq \infty$. Let the function $T_{n_1, \infty}(x_1, x_2) \in L_{p_1 p_2}^0$ be a trigonometric polynomial of order at most n_1 ($n_1 \in \mathbb{N}$), in x_1 . Then

$$\|T_{n_1, \infty}\|_{q_1 p_2} \leq C_1 \|T_{n_1, \infty}^{(\frac{1}{p_1} - \frac{1}{q_1}, 0)}\|_{p_1, p_2},$$

where the constant C_1 does not depend on $T_{n_1, \infty}$.

Proof. Let $1 \leq p_2 < \infty$. Let us estimate

$$\|T_{n_1, \infty}\|_{q_1 p_2} = \left(\int_0^{2\pi} \left(\left(\int_0^{2\pi} |T_{n_1, \infty}(x_1, x_2)|^{q_1} dx_1 \right)^{\frac{1}{q_1}} \right)^{p_2} dx_2 \right)^{\frac{1}{p_2}}.$$

For almost all fixed x_2^0 , the function $T_{n_1, \infty}(x_1, x_2^0) = P_{n_1}(x_1)$ is a function of one variable x_1 . Applying Lemma 7, we get

$$\left(\int_0^{2\pi} |P_{n_1}(x_1)|^{q_1} dx_1 \right)^{\frac{1}{q_1}} \ll \left(\int_0^{2\pi} |P_{n_1}^{(\frac{1}{p_1} - \frac{1}{q_1})}(x_1)|^{p_1} dx_1 \right)^{\frac{1}{p_1}}.$$

Then the following inequality is true

$$\begin{aligned} & \left(\int_0^{2\pi} \left(\left(\int_0^{2\pi} |T_{n_1, \infty}(x_1, x_2)|^{q_1} dx_1 \right)^{\frac{1}{q_1}} \right)^{p_2} dx_2 \right)^{\frac{1}{p_2}} \\ & \ll \left(\int_0^{2\pi} \left(\left(\int_0^{2\pi} |T_{n_1, \infty}^{(\frac{1}{p_1} - \frac{1}{q_1}, 0)}(x_1, x_2)|^{p_1} dx_1 \right)^{\frac{1}{p_1}} \right)^{p_2} dx_2 \right)^{\frac{1}{p_2}}, \end{aligned}$$

whence follows Lemma 12. Lemma 12 is verified similarly in the case when $p_2 = \infty$. Lemma 12 is proved.

Lemma 13 *Let $1 < p_2 < q_2 < \infty, 1 \leq p_1 \leq \infty$. Let the function $T_{\infty, n_2}(x_1, x_2) \in L_{p_1 p_2}^0$ be a trigonometric polynomial of order at most $n_2 (n_2 \in \mathbb{N})$, in x_2 . Then*

$$\|T_{\infty, n_2}\|_{p_1 q_2} \leq C_2 \|T_{\infty, n_2}^{(0, \frac{1}{p_2} - \frac{1}{q_2})}\|_{p_1, p_2},$$

where the constant C_2 does not depend on T_{∞, n_2} .

Proof. For almost all fixed x_1^0 , the function $P_{n_2}(x_2) = T_{\infty, n_2}(x_1^0, x_2)$ is a function of one variable x_2 . Then

$$P_{n_2}(x_2) = \sum_{\nu=1}^{n_2} (c_\nu \cos \nu x_2 + d_\nu \sin \nu x_2).$$

Let $\theta_2 = \frac{1}{p_2} - \frac{1}{q_2}$. Then, applying Lemma 9, we have

$$\begin{aligned} P_{n_2}(x_2) &= \sum_{\nu=1}^{n_2} (c_\nu \cos \nu x_2 + d_\nu \sin \nu x_2) = \sum_{\nu=1}^{n_2} \frac{1}{\nu^{\theta_2}} \nu^{\theta_2} (c_\nu \cos \nu x_2 + d_\nu \sin \nu x_2) \\ &= \sum_{\nu=1}^{n_2} \frac{1}{\nu^{\theta_2}} \left(\cos \frac{\theta_2 \pi}{2} (c_\nu \cos \nu x_2 + d_\nu \sin \nu x_2)^{(\theta_2)} + \sin \frac{\theta_2 \pi}{2} \widetilde{(c_\nu \cos \nu x_2 + d_\nu \sin \nu x_2)^{(\theta_2)}} \right) \\ &= \cos \frac{\theta_2 \pi}{2} (P_{n_2}(x_2) \left(\left\{ \frac{1}{m^{\theta_2}} \right\}; x_2 \right))^{(\theta_2)} + \sin \frac{\theta_2 \pi}{2} (\widetilde{P}_{n_2} \left(\left\{ \frac{1}{m^{\theta_2}} \right\}; x_2 \right))^{(\theta_2)} \\ &= \cos \frac{\theta_2 \pi}{2} \frac{1}{\pi} \int_{-\pi}^{\pi} (P_{n_2}(t_2))^{(\theta_2)} \sum_{\nu=1}^{n_2} \frac{1}{\nu^{\theta_2}} \cos \nu(t_2 - x_2) dt_2 \\ & \quad - \sin \frac{\theta_2 \pi}{2} \frac{1}{\pi} \int_{-\pi}^{\pi} (P_{n_2}(t_2))^{(\theta_2)} \sum_{\nu=1}^{n_2} \frac{1}{\nu^{\theta_2}} \sin \nu(t_2 - x_2) dt_2. \end{aligned}$$

Now, using Lemma 10, for almost all x_2 we obtain

$$\begin{aligned}
|P_{n_2}(x_2)| &\leq \frac{1}{\pi} \int_{-\pi}^{\pi} \left| \left(P_{n_2}(t_2) \right)^{(\theta_2)} \right| \left| \sum_{\nu=1}^{n_2+1} \frac{1}{\nu^{\theta_2}} \cos \nu(t_2 - x_2) \right| dt_2 \\
&\quad + \frac{1}{\pi} \int_{-\pi}^{\pi} \left| \left(P_{n_2}(t_2) \right)^{(\theta_2)} \right| \left| \sum_{\nu=1}^{n_2} \frac{1}{\nu^{\theta_2}} \sin \nu(t_2 - x_2) \right| dt_2 \\
&\leq C(\theta_2) \frac{1}{\pi} \int_{-\pi}^{\pi} \left| \left(P_{n_2}(t_2) \right)^{(\theta_2)} \right| |t_2 - x_2|^{\theta_2-1} dt_2 + C(\theta_2) \frac{1}{\pi} \int_{-\pi}^{\pi} \left| \left(P_{n_2}(t_2) \right)^{(\theta_2)} \right| |t_2 - x_2|^{\theta_2-1} dt_2 \\
&\leq C_1(\theta_2) \int_{-\pi}^{\pi} \left| \left(P_{n_2}(t_2) \right)^{(\theta_2)} \right| |t_2 - x_2|^{\theta_2-1} dt_2.
\end{aligned}$$

Thus, for almost all x_2 the following inequality holds

$$|P_{n_2}(x_2)| \leq C(p_2, q_2) \int_{-\pi}^{\pi} \left| P_{n_2}^{(\theta_2)}(t_2) \right| |t_2 - x_2|^{\theta_2-1} dt_2.$$

Therefore, for almost all x_1 and x_2 , the inequality

$$|T_{\infty, n_2}(x_1, x_2)| \leq C(p_2, q_2) \int_{-\pi}^{\pi} \left| T_{\infty, n_2}^{(0, \theta_2)}(x_1, t_2) \right| |t_2 - x_2|^{\theta_2-1} dt_2, \quad (2)$$

is true, where the constant $C(p_2, q_2)$ depends only on p_2 and q_2 .

If $p_1 < \infty$, then, using (2), we obtain

$$\begin{aligned}
\|T_{\infty, n_2}\|_{p_1 q_2} &= \left(\int_{-\pi}^{\pi} \left(\left\{ \int_{-\pi}^{\pi} \left| T_{\infty, n_2}(x_1, x_2) \right|^{p_1} dx_1 \right\}^{\frac{1}{p_1}} \right)^{q_2} dx_2 \right)^{\frac{1}{q_2}} \leq \\
&\leq C(p_2, q_2) \left(\int_{-\pi}^{\pi} \left(\left\{ \int_{-\pi}^{\pi} \left| \int_{-\pi}^{\pi} \left| T_{\infty, n_2}^{(0, \theta_2)}(x_1, t_2) \right| |t_2 - x_2|^{\theta_2-1} dt_2 \right|^{p_1} dx_1 \right\}^{\frac{1}{p_1}} \right)^{q_2} dx_2 \right)^{\frac{1}{q_2}}.
\end{aligned}$$

Using the Minkowski inequality, we have

$$\begin{aligned}
\|T_{\infty, n_2}\|_{p_1 q_2} &\leq \\
&\leq C(p_2, q_2) \left(\int_{-\pi}^{\pi} \left(\int_{-\pi}^{\pi} \left\{ \int_{-\pi}^{\pi} \left| T_{\infty, n_2}^{(0, \theta_2)}(x_1, x_2)(x_1, t_2) \right|^{p_1} dx_1 \right\}^{\frac{1}{p_1}} |t_2 - x_2|^{\theta_2-1} dt_2 \right)^{q_2} dx_2 \right)^{\frac{1}{q_2}}.
\end{aligned}$$

We introduce the function $F(t_2)$, which is equal to $\left\{ \int_{-\pi}^{\pi} \left| T_{\infty, n_2}^{(0, \theta_2)}(x_1, t_2) \right|^{p_1} dx_1 \right\}^{\frac{1}{p_1}}$, if $t_2 \in [-\pi, \pi]$

and

$$\|T_{\infty, n_2}\|_{p_1 q_2} \leq C(p_2, q_2) \left(\int_{\mathbb{R}_1} \left(\int_{\mathbb{R}_1} F(t_2) |t_2 - x_2|^{\theta_2-1} dt_2 \right)^{q_2} dx_2 \right)^{\frac{1}{q_2}}.$$

Therefore, from Lemma 11, we deduce that

$$\|T_{\infty, n_2}\|_{p_1 q_2} \leq C_1(p_2, q_2) \|F\|_{L_{p_2}^{(1)}(\mathbb{R}_1)} = C_1(p_2, q_2) \|T_{\infty, n_2}^{(0, \theta_2)}\|_{p_1 p_2}.$$

If $p_1 = \infty$, then, applying inequality (2), we get

$$\begin{aligned} \|T_{\infty, n_2}\|_{p_1 q_2} &= \left(\int_{-\pi}^{\pi} \left[\sup_{0 \leq x_1 \leq 2\pi} \text{vrai} |T_{\infty, n_2}(x_1, x_2)| \right]^{q_2} dx_2 \right)^{\frac{1}{q_2}} \\ &\ll \left(\int_{-\pi}^{\pi} \left[\sup_{0 \leq x_1 \leq 2\pi} \text{vrai} \left| \int_{-\pi}^{\pi} T_{\infty, n_2}^{(0, \theta_2)}(x_1, t_2) \left| t_2 - x_2 \right|^{\theta_2 - 1} dt_2 \right| \right]^{q_2} dx_2 \right)^{\frac{1}{q_2}} \\ &\ll \left(\int_{-\pi}^{\pi} \left[\int_{-\pi}^{\pi} \sup_{0 \leq x_1 \leq 2\pi} \text{vrai} \left| T_{\infty, n_2}^{(0, \theta_2)}(x_1, t_2) \left| t_2 - x_2 \right|^{\theta_2 - 1} dt_2 \right| \right]^{q_2} dx_2 \right)^{\frac{1}{q_2}}. \end{aligned}$$

Consider the function $F_1(t_2)$, which is equal to $\sup_{0 \leq x_1 \leq 2\pi} \text{vrai} |T_{\infty, n_2}^{(0, \theta_2)}(x_1, t_2)|$, if $t_2 \in [-\pi, \pi]$ and is zero otherwise. Then we obtain

$$\|T_{\infty, n_2}\|_{p_1 q_2} \leq C(p_2, q_2) \left(\int_{\mathbb{R}_1} \left(\int_{\mathbb{R}_1} F_1(t_2) \left| t_2 - x_2 \right|^{\theta_2 - 1} dt_2 \right)^{q_2} dx_2 \right)^{\frac{1}{q_2}}.$$

Applying Lemma 11, we conclude that

$$\|T_{\infty, n_2}\|_{p_1 q_2} \leq \|F_1\|_{L_{p_2}^{(1)}(\mathbb{R}_1)} \leq C_1(p_2, q_2) \|T_{\infty, n_2}^{(0, \theta_2)}\|_{p_1 p_2}.$$

Thus, we have shown that

$$\|T_{\infty, n_2}\|_{p_1 q_2} \leq C_1(p_2, q_2) \|T_{\infty, n_2}^{(0, \theta_2)}\|_{p_1 p_2}.$$

Lemma 13 is proved.

Lemma 14 ([21],[25]). (a) Let $1 < p < q < \infty$, $\delta \in (0, 1)$. Consider the function $g_1(x) = \sum_{\nu=0}^{\infty} \frac{(\nu+1)^\beta}{2^{\nu\alpha}} \cos 2^\nu x$, where $\beta > -\frac{1}{2}$, $\alpha > \frac{1}{p} - \frac{1}{q}$. Then

$$\omega_\alpha(g_1, \delta)_q^{(1)} \asymp \delta^\alpha \left(\log_2 \frac{2}{\delta} \right)^{\beta + \frac{1}{2}},$$

$$\left(\int_0^\delta \left[t^{-\theta} \omega_{\alpha + \frac{1}{p} - \frac{1}{q}}(g_1, t)_p^{(1)} \right]^q \frac{dt}{t} \right)^{\frac{1}{q}} \asymp \delta^{\alpha - \frac{1}{p} + \frac{1}{q}} \left(\log_2 \frac{2}{\delta} \right)^\beta.$$

(b) Let $1 = p < q = \infty$, $\delta \in (0, 1)$. Consider the function $g_2(x) = \sum_{\nu=0}^{\infty} \frac{(\nu+1)^\beta}{2^{\nu\alpha}} \cos(2^\nu x - \frac{\pi\alpha}{2})$, where $\beta > -1$, $\alpha > 1$. Then

$$\omega_\alpha(g_2, \delta)_\infty^{(1)} \asymp \delta^\alpha \left(\log_2 \frac{2}{\delta} \right)^{\beta + 1},$$

$$\left(\int_0^\delta t^{-1} \omega_{\alpha+1}(g_2, t)_1^{(1)} \frac{dt}{t} \right)^{\frac{1}{q}} \asymp \delta^{\alpha-1} \left(\log_2 \frac{2}{\delta} \right)^\beta.$$

3. Proof of main results

Proof of Theorems 1. For every $\delta_i \in (0, 1)$, take a non-negative integer n_i such that $\frac{1}{2^{n_i+1}} \leq \delta_i < \frac{1}{2^{n_i}}$, $i = 1, 2$. Using Lemma 3, we have

$$I = \omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{q_1 q_2} \ll \omega_{\alpha_1, \alpha_2}\left(f, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}}\right)_{q_1 q_2}.$$

Applying Lemma 1, we derive the following

$$\begin{aligned} I &\ll 2^{-n_1 \alpha_1 - n_2 \alpha_2} \|V_{2^{n_1}, 2^{n_2}}^{(\alpha_1, \alpha_2)}(f)\|_{q_1 q_2} + 2^{-n_1 \alpha_1} \|V_{2^{n_1}, \infty}^{(\alpha_1, 0)}(f - V_{\infty, 2^{n_2}}(f))\|_{q_1 q_2} \\ &+ 2^{-n_2 \alpha_2} \|V_{\infty, 2^{n_2}}^{(0, \alpha_2)}(f - V_{2^{n_1}, \infty}(f))\|_{q_1 q_2} + \|f - V_{2^{n_1}, \infty}(f) - V_{\infty, 2^{n_2}}(f) + V_{2^{n_1}, 2^{n_2}}(f)\|_{q_1 q_2} \\ &\equiv 2^{-n_1 \alpha_1 - n_2 \alpha_2} I_1 + 2^{-n_1 \alpha_1} I_2 + 2^{-n_2 \alpha_2} I_3 + I_4. \end{aligned}$$

By Lemmas 1, 4 using the mixed moduli of smoothness property, we obtain

$$\begin{aligned} I_4 &\ll \left(\sum_{\nu_2=n_2}^{\infty} \left(\sum_{\nu_1=n_1}^{\infty} \left(2^{\nu_1 \left(\frac{1}{p_1} - \frac{1}{q_1}\right)} 2^{\nu_2 \left(\frac{1}{p_2} - \frac{1}{q_2}\right)} Y_{2^{\nu_1-1}, 2^{\nu_2-1}}(f)_{p_1 p_2} \right)^{q_1^*} \right)^{\frac{q_2^*}{q_1^*}} \right)^{\frac{1}{q_2^*}} \\ &\ll \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

It follows from Lemma 6 (b) that

$$I_2 \ll \left\| \sum_{\nu_2=n_2}^{\infty} V_{2^{n_1} \infty}^{(\alpha_1, 0)}(V_{\infty 2^{\nu_2}}(f) - V_{\infty 2^{\nu_2+1}}(f)) \right\|_{q_1 q_2}.$$

Thus, by Lemma 5 (a), we deduce that

$$I_2 \ll \left(\sum_{\nu_2=n_2}^{\infty} \left(2^{\nu_2 \left(\frac{1}{p_2} - \frac{1}{q_2}\right)} \|V_{2^{n_1} \infty}^{(\alpha_1, 0)}(V_{\infty 2^{\nu_2}}(f) - V_{\infty 2^{\nu_2+1}}(f))\|_{q_1 p_2} \right)^{q_2^*} \right)^{\frac{1}{q_2^*}}.$$

Applying Lemma 12 in the case $1 < p_1 < q_1 < \infty$ or Lemma 8 (a) in the case $1 = p_1 < q_1 = \infty$, we have

$$I_2 \ll \left(\sum_{\nu_2=n_2}^{\infty} \left(2^{\nu_2 \left(\frac{1}{p_2} - \frac{1}{q_2}\right)} \|V_{2^{n_1} \infty}^{(\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, 0)}(V_{\infty 2^{\nu_2}}(f) - V_{\infty 2^{\nu_2+1}}(f))\|_{p_1 p_2} \right)^{q_2^*} \right)^{\frac{1}{q_2^*}}.$$

Using Lemma 1, we derive the following

$$\begin{aligned} 2^{-n_1 \alpha_1} I_2 &\ll \\ &\ll 2^{n_1 \left(\frac{1}{p_1} - \frac{1}{q_1}\right)} \left(\sum_{\nu_2=n_2}^{\infty} \left(2^{\nu_2 \left(\frac{1}{p_2} - \frac{1}{q_2}\right)} \times \right. \right. \\ &\quad \left. \left. \times \frac{1}{2^{n_1 \left(\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}\right)}} \|V_{2^{n_1} \infty}^{(\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, 0)}(V_{\infty 2^{\nu_2}}(f) - V_{\infty 2^{\nu_2+1}}(f))\|_{p_1 p_2} \right)^{q_2^*} \right)^{\frac{1}{q_2^*}} \\ &\ll 2^{n_1 \left(\frac{1}{p_1} - \frac{1}{q_1}\right)} \left(\sum_{\nu_2=n_2}^{\infty} \left(2^{\nu_2 \left(\frac{1}{p_2} - \frac{1}{q_2}\right)} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}\left(f, \frac{1}{2^{n_1}}, \frac{1}{2^{\nu_2}}\right)_{p_1 p_2} \right)^{q_2^*} \right)^{\frac{1}{q_2^*}} \\ &\ll \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

Now, we begin to estimate the $I_3 = \|V_{\infty, 2^{n_2}}^{(0, \alpha_2)}(f - V_{2^{n_1}, \infty}(f))\|_{q_1 q_2}$. Applying Lemma 6 (a), we have

$$I_3 \ll \left\| \sum_{\nu_1=n_1}^{\infty} V_{\infty, 2^{n_2}}^{(0, \alpha_2)}(V_{2^{\nu_1}, \infty}(f) - V_{2^{\nu_1+1}, \infty}(f)) \right\|_{q_1 q_2}.$$

Lemma 5 (b) yields that

$$I_3 \ll \left(\sum_{\nu_1=n_1}^{\infty} \left(2^{\nu_1(\frac{1}{p_1} - \frac{1}{q_1})} \|V_{\infty, 2^{n_2}}^{(0, \alpha_2)}(V_{2^{\nu_1}, \infty}(f) - V_{2^{\nu_1+1}, \infty}(f))\|_{p_1 q_2} \right)^{q_1^*} \right)^{\frac{1}{q_1^*}}.$$

Applying Lemma 13 in the case $1 < p_2 < q_2 < \infty$ or Lemma 8 (b) in the case $1 = p_2 < q_2 = \infty$, we get

$$I_3 \ll \left(\sum_{\nu_1=n_1}^{\infty} \left(2^{\nu_1(\frac{1}{p_1} - \frac{1}{q_1})} \|V_{\infty, 2^{n_2}}^{(0, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2})}(V_{2^{\nu_1}, \infty}(f) - V_{2^{\nu_1+1}, \infty}(f))\|_{p_1 p_2} \right)^{q_1^*} \right)^{\frac{1}{q_1^*}}.$$

Thus, using Lemma 1, we deduce

$$\begin{aligned} 2^{-n_2 \alpha_2} I_3 &\ll \\ &\ll 2^{n_2(\frac{1}{p_2} - \frac{1}{q_2})} \left(\sum_{\nu_1=n_1}^{\infty} \left(2^{\nu_1(\frac{1}{p_1} - \frac{1}{q_1})} \times \right. \right. \\ &\quad \times \left. \frac{1}{2^{n_2(\alpha_2 + \frac{1}{p_2} - \frac{1}{q_2})}} \|V_{\infty, 2^{n_2}}^{(0, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2})}(V_{2^{\nu_1}, \infty}(f) - V_{2^{\nu_1+1}, \infty}(f))\|_{p_1 p_2} \right)^{q_1^*} \right)^{\frac{1}{q_1^*}} \\ &\ll 2^{n_2(\frac{1}{p_2} - \frac{1}{q_2})} \left(\sum_{\nu_1=n_1}^{\infty} \left(2^{\nu_1(\frac{1}{p_1} - \frac{1}{q_1})} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f, \frac{1}{2^{\nu_1}}, \frac{1}{2^{n_2}})_{p_1 p_2} \right)^{q_1^*} \right)^{\frac{1}{q_1^*}} \\ &\ll \left(\int_{\frac{1}{2^{n_2+1}}}^{\frac{1}{2^{n_2}}} \left(t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \left(\sum_{\nu_1=n_1}^{\infty} \left(2^{\nu_1(\frac{1}{p_1} - \frac{1}{q_1})} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f, \frac{1}{2^{\nu_1}}, \frac{1}{2^{n_2+1}})_{p_1 p_2} \right)^{q_1^*} \right)^{\frac{1}{q_1^*}} \right)^{q_2^*} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}} \\ &\ll \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

Let us estimate $2^{-n_1 \alpha_1 - n_2 \alpha_2} I_1 = 2^{-n_1 \alpha_1 - n_2 \alpha_2} \|V_{2^{n_1}, 2^{n_2}}^{(\alpha_1, \alpha_2)}(f)\|_{q_1 q_2}$. Applying Lemma 12 in the case $1 < p_1 < q_1 < \infty$ or Lemma 8 (a) in the case $1 = p_1 < q_1 = \infty$, we obtain

$$2^{-n_1 \alpha_1 - n_2 \alpha_2} I_1 \ll 2^{-n_1 \alpha_1 - n_2 \alpha_2} \|V_{2^{n_1}, 2^{n_2}}^{(\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, 0)}(f)\|_{p_1 q_2}.$$

Using Lemma 13 in the case $1 < p_2 < q_2 < \infty$ or Lemma 8 (b) in the case $1 = p_2 < q_2 = \infty$, we finally obtain

$$2^{-n_1 \alpha_1 - n_2 \alpha_2} I_1 \ll 2^{-n_1 \alpha_1 - n_2 \alpha_2} \|V_{2^{n_1}, 2^{n_2}}^{(\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2})}(f)\|_{p_1 p_2}.$$

By Lemma 1 and using the mixed moduli of smoothness property, we obtain

$$\begin{aligned} &2^{-n_1 \alpha_1 - n_2 \alpha_2} I_1 \ll \\ &\ll \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

Combining the estimates for I_1, I_2, I_3 and I_4 , we finally get

$$\begin{aligned} & \omega_{\alpha_1, \alpha_2}(f, \delta_1, \delta_2)_{q_1 q_2} \ll \\ & \ll \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

Inequality (1) is proved.

Proof of Remark 1. Let us prove that Theorem 1 is optimal in the sense that there exists a function $f_0(x_1, x_2)$ such that for (1) the sign \ll can be replaced by the sign \asymp .

Consider the function $f_0(x_1, x_2) = \sin x_1 \sin x_2$. For $\delta_i \in (0, 1)$ there is a non-negative integer a number n_i such that $\frac{1}{2^{n_i+1}} \leq \delta_i < \frac{1}{2^{n_i}}, i = 1, 2$. Applying Lemmas 1 and 3, for any $r_i > 0, i = 1, 2$, and any $p_i \in [1, \infty]$ ($i = 1, 2$), we have

$$\omega_{r_1, r_2}(f_0, \delta_1, \delta_2)_{p_1, p_2} \asymp \omega_{r_1, r_2}\left(f_0, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}}\right)_{p_1, p_2} \asymp \frac{1}{2^{n_1 r_1 + n_2 r_2}} \asymp \delta_1^{r_1} \delta_2^{r_2}.$$

But then

$$\begin{aligned} & \omega_{\alpha_1, \alpha_2}(f_0, \delta_1, \delta_2)_{q_1, q_2} \asymp \delta_1^{\alpha_1} \delta_2^{\alpha_2}, \\ & \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}} \asymp \delta_1^{\alpha_1} \delta_2^{\alpha_2}. \end{aligned}$$

It follows from these estimates that for the function $f_0(x_1, x_2)$ in relation (1) the sign \ll can be replaced by the sign \asymp .

Another way to see the optimality of the estimate in Theorem 1 is to consider the series with monotone or general monotone coefficients, see [26].

Proof of Remark 2. Let us now show that, generally speaking, in inequality (1) the sign \ll cannot be replaced by the sign \asymp . Let $1 < p_2 < q_2 < \infty$. Consider the function $f_1(x_1, x_2) = \sin x_1 \cdot g(x_2)$, where $g(x_2) = \sum_{\nu=0}^{\infty} \frac{(\nu+1)^\beta}{2^{\nu \alpha_2}} \cos 2^\nu x_2$, $\beta > -\frac{1}{2}, \alpha_2 > \frac{1}{p_2} - \frac{1}{q_2}$. Since

$$\begin{aligned} \omega_{\alpha_1, \alpha_2}(f_1, \delta_1, \delta_2)_{q_1 q_2} &= \omega_{\alpha_1}(\sin x_1, \delta_1)_{q_1}^{(1)} \cdot \omega_{\alpha_2}(g, \delta_2)_{q_2}^{(1)} \asymp \delta_1^{\alpha_1} \cdot \omega_{\alpha_2}(g, \delta_2)_{q_2}^{(1)}, \\ \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f_1, t_1, t_2)_{p_1 p_2} &= \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}}(\sin x_1, t_1)_{p_1}^{(1)} \cdot \omega_{\alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(g, t_2)_{p_2}^{(1)} \\ &\asymp t_1^{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}} \cdot \omega_{\alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(g, t_2)_{p_2}^{(1)}, \end{aligned}$$

then, applying Lemma 14, item (a) and substituting into inequality (1), we obtain

$$\begin{aligned} \omega_{\alpha_1, \alpha_2}(f_1, \delta_1, \delta_2)_{q_1 q_2} &\asymp \delta_1^{\alpha_1} \cdot \delta_2^{\alpha_2} \left(\log_2 \frac{2}{\delta_2} \right)^{\beta + \frac{1}{2}}, \\ & \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f_1, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}} \\ & \asymp \delta_1^{\alpha_1} \cdot \delta_2^{\alpha_2 - \frac{1}{p_2} + \frac{1}{q_2}} \left(\log_2 \frac{2}{\delta_2} \right)^\beta. \end{aligned}$$

Thus, for the function $f_1(x_1, x_2)$, the right and left hand sides of relation (1) have different orders as functions of the variable δ_2 for a fixed δ_1 , which means that in relation (1) it is not possible replacing the sign \ll with the sign \asymp .

It can be verified similarly that in relation (1), generally speaking, the sign \ll cannot be replaced by the sign \asymp in the case $1 < p_1 < q_1 < \infty$.

Let $1 = p_2 < q_2 = \infty$. Consider the function $f_2(x_1, x_2) = \sin x_1 \cdot g(x_2)$, where $g(x_2) = \sum_{\nu=0}^{\infty} \frac{(\nu+1)^\beta}{2^{\nu\alpha_2}} \cos\left(2^\nu x_2 - \frac{\pi\alpha_2}{2}\right)$, $\beta > -1, \alpha_2 > 1$. Since

$$\begin{aligned} \omega_{\alpha_1, \alpha_2}(f_2, \delta_1, \delta_2)_{q_1 q_2} &= \omega_{\alpha_1}(\sin x_1, \delta_1)_q^{(1)} \cdot \omega_{\alpha_2}(g, \delta_2)_{q_2}^{(1)} \asymp \delta_1^{\alpha_1} \cdot \omega_{\alpha_2}(g, \delta_2)_{q_2}^{(1)}, \\ \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f_2, t_1, t_2)_{p_1 p_2} &= \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}}(\sin x_1, t_1)_{p_1}^{(1)} \cdot \omega_{\alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(g, t_2)_{p_2}^{(1)} \asymp \\ &\asymp t_1^{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}} \cdot \omega_{\alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(g, t_2)_{p_2}^{(1)}, \end{aligned}$$

then, applying Lemma 14 item (b) and substituting into inequality (1), we obtain

$$\begin{aligned} \omega_{\alpha_1, \alpha_2}(f_2, \delta_1, \delta_2)_{q_1 q_2} &\asymp \delta_1^{\alpha_1} \cdot \delta_2^{\alpha_2} \left(\log_2 \frac{2}{\delta_2} \right)^{\beta+1}, \\ \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f_2, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}} &\asymp \\ &\asymp \delta_1^{\alpha_1} \cdot \delta_2^{\alpha_2 - 1} \left(\log_2 \frac{2}{\delta_2} \right)^{\beta+1}. \end{aligned}$$

Therefore, for the function $f_2(x_1, x_2)$, the right and left parts of relation (1) have different orders as functions of the variable δ_2 for a fixed δ_1 , which means that in relation (1), it is impossible to replace the sign \ll with the sign \asymp .

In a similar way, it is verified that, in relation to (1), the sign \ll cannot be replaced by the sign \asymp in the case $1 = p_1 < q_1 = \infty$.

Thus the proof of Remark 2 is complete.

Proof of Theorem 2. Using Theorem 1, we have

$$\begin{aligned} \omega_{\alpha_1, \alpha_2}(f^{(\rho_1, \rho_2)}, \delta_1, \delta_2)_{q_1 q_2} &\ll \\ &\ll \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f^{(\rho_1, \rho_2)}, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

For each $\delta_i \in (0, 1)$ there is a non-negative integer n_i such that $\frac{1}{2^{n_i+1}} \leq \delta_i < \frac{1}{2^{n_i}}, i = 1, 2$. Then

$$\begin{aligned} \omega_{\alpha_1, \alpha_2}(f^{(\rho_1, \rho_2)}, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}})_{q_1 q_2} &\ll \\ &\ll \left(\int_0^{\frac{1}{2^{n_2}}} \left(\int_0^{\frac{1}{2^{n_1}}} \left(t_1^{-\frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f^{(\rho_1, \rho_2)}, t_1, t_2)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}} \\ &\ll \left(\sum_{\nu_2=n_2}^{\infty} \left(\sum_{\nu_1=n_1}^{\infty} \left(2^{\nu_1(\frac{1}{p_1} - \frac{1}{q_1})} 2^{\nu_2(\frac{1}{p_2} - \frac{1}{q_2})} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}}(f^{(\rho_1, \rho_2)}, \frac{1}{2^{\nu_1}}, \frac{1}{2^{\nu_2}})_{p_1 p_2} \right)^{q_1^*} \right)^{\frac{q_2^*}{q_1^*}} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

a). Let $\rho_1 > 0, \rho_2 = 0$. Using Lemma 3, we deduce

$$\begin{aligned} & \omega_{\alpha_1, \alpha_2} \left(f^{(\rho_1, \rho_2)}, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}} \right)_{q_1 q_2} \ll \\ & \left(\sum_{\nu_2=n_2}^{\infty} \left\{ \sum_{\nu_1=n_1}^{\infty} 2^{\nu_1 q_1^* \left(\frac{1}{p_1} - \frac{1}{q_1} \right)} \times \right. \right. \\ & \quad \left. \left. \times \left(\sum_{m_1=\nu_1}^{\infty} 2^{m_1 \rho_1} \omega_{\alpha_1 + \rho_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, \frac{1}{2^{m_1}}, \frac{1}{2^{\nu_2}} \right)_{p_1 p_2} \right)^{q_1^*} \right\}^{\frac{q_2^*}{q_1^*}} 2^{\nu_2 q_2^* \left(\frac{1}{p_2} - \frac{1}{q_2} \right)} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

Applying Lemma 2 to the inner sum, we obtain

$$\begin{aligned} & \omega_{\alpha_1, \alpha_2} \left(f^{(\rho_1, \rho_2)}, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}} \right)_{q_1 q_2} \ll \\ & \left(\sum_{\nu_2=n_2}^{\infty} \left\{ \sum_{\nu_1=n_1}^{\infty} 2^{\nu_1 \left(\rho_1 + \frac{1}{p_1} - \frac{1}{q_1} \right)} \omega_{\alpha_1 + \rho_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, \frac{1}{2^{\nu_1}}, \frac{1}{2^{\nu_2}} \right)_{p_1 p_2} \right\}^{q_1^*} \right)^{\frac{q_2^*}{q_1^*}} 2^{\nu_2 q_2^* \left(\frac{1}{p_2} - \frac{1}{q_2} \right)} \frac{1}{q_2^*} \ll \\ & \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\rho_1 - \frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\rho_2 - \frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \rho_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \rho_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, t_1, t_2 \right)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

b). Let $\rho_1 = 0, \rho_2 > 0$. By Lemma 3, we have

$$\begin{aligned} & \omega_{\alpha_1, \alpha_2} \left(f^{(\rho_1, \rho_2)}, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}} \right)_{q_1 q_2} \\ & \ll \left(\sum_{\nu_2=n_2}^{\infty} 2^{\nu_2 q_2^* \left(\frac{1}{p_2} - \frac{1}{q_2} \right)} \left\{ \left(\sum_{\nu_1=n_1}^{\infty} 2^{\nu_1 q_1^* \left(\frac{1}{p_1} - \frac{1}{q_1} \right)} \right. \right. \\ & \quad \left. \left. \left(\sum_{m_2=\nu_2}^{\infty} 2^{m_2 \rho_2} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \rho_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, \frac{1}{2^{\nu_1}}, \frac{1}{2^{m_2}} \right)_{p_1 p_2} \right)^{q_1^*} \right\}^{\frac{1}{q_1^*}} \right)^{\frac{q_2^*}{q_1^*}} \frac{1}{q_2^*}. \end{aligned}$$

Applying the Minkowski inequality to the inner sum, we get

$$\begin{aligned} & \omega_{\alpha_1, \alpha_2} \left(f^{(\rho_1, \rho_2)}, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}} \right)_{q_1 q_2} \\ & \ll \left(\sum_{\nu_2=n_2}^{\infty} 2^{\nu_2 q_2^* \left(\frac{1}{p_2} - \frac{1}{q_2} \right)} \left\{ \sum_{m_2=\nu_2}^{\infty} 2^{m_2 \rho_2} \right. \right. \\ & \quad \left. \left. \left(\sum_{\nu_1=n_1}^{\infty} \left(2^{\nu_1 \left(\frac{1}{p_1} - \frac{1}{q_1} \right)} \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \rho_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, \frac{1}{2^{\nu_1}}, \frac{1}{2^{m_2}} \right)_{p_1 p_2} \right)^{q_1^*} \right\}^{\frac{1}{q_1^*}} \right)^{\frac{q_2^*}{q_1^*}} \frac{1}{q_2^*}. \end{aligned}$$

Furthermore, it follows from Lemma 2 that

$$\begin{aligned} & \omega_{\alpha_1, \alpha_2} \left(f^{(\rho_1, \rho_2)}, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}} \right)_{q_1 q_2} \\ & \ll \left(\sum_{\nu_2=n_2}^{\infty} 2^{\nu_2 q_2^* \left(\rho_2 + \frac{1}{p_2} - \frac{1}{q_2} \right)} \left\{ \left(\sum_{\nu_1=n_1}^{\infty} \left(2^{\nu_1 \left(\frac{1}{p_1} - \frac{1}{q_1} \right)} \times \right. \right. \right. \\ & \quad \left. \left. \left. \times \omega_{\alpha_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \rho_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, \frac{1}{2^{\nu_1}}, \frac{1}{2^{\nu_2}} \right)_{p_1 p_2} \right)^{q_1^*} \right\}^{\frac{1}{q_1^*}} \right)^{\frac{q_2^*}{q_1^*}} \frac{1}{q_2^*} \ll \\ & \left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\rho_1 - \frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\rho_2 - \frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \rho_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \rho_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, t_1, t_2 \right)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

c). Let $\rho_1 > 0, \rho_2 > 0$. Using Lemma 3, we obtain

$$\begin{aligned} \omega_{\alpha_1, \alpha_2} \left(f^{(\rho_1, \rho_2)}, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}} \right)_{q_1 q_2} &\ll \left(\sum_{\nu_2=n_2}^{\infty} 2^{\nu_2 q_2^* \left(\frac{1}{p_2} - \frac{1}{q_2} \right)} \left\{ \left(\sum_{\nu_1=n_1}^{\infty} 2^{\nu_1 q_1^* \left(\frac{1}{p_1} - \frac{1}{q_1} \right)} \times \right. \right. \right. \\ &\times \left. \left. \left\langle \sum_{m_1=\nu_1}^{\infty} 2^{m_1 \rho_1} \sum_{m_2=\nu_2}^{\infty} 2^{m_2 \rho_2} \omega_{\alpha_1 + \rho_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \rho_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, \frac{1}{2^{m_1}}, \frac{1}{2^{m_2}} \right)_{p_1 p_2} \right\rangle^{q_1^*} \right)^{\frac{1}{q_1^*}} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

Applying Lemma 2 to the sum in the "angle" brackets, we get

$$\begin{aligned} \omega_{\alpha_1, \alpha_2} \left(f^{(\rho_1, \rho_2)}, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}} \right)_{q_1 q_2} &\ll \left(\sum_{\nu_2=n_2}^{\infty} 2^{\nu_2 q_2^* \left(\frac{1}{p_2} - \frac{1}{q_2} \right)} \left\{ \left(\sum_{\nu_1=n_1}^{\infty} 2^{\nu_1 q_1^* \left(\rho_1 + \frac{1}{p_1} - \frac{1}{q_1} \right)} \times \right. \right. \\ &\times \left. \left. \left\langle \sum_{m_2=\nu_2}^{\infty} 2^{m_2 \rho_2} \omega_{\alpha_1 + \rho_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \rho_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, \frac{1}{2^{\nu_1}}, \frac{1}{2^{m_2}} \right)_{p_1 p_2} \right\rangle^{q_1^*} \right)^{\frac{1}{q_1^*}} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

By application of Minkowski's inequality, we obtain

$$\begin{aligned} \omega_{\alpha_1, \alpha_2} \left(f^{(\rho_1, \rho_2)}, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}} \right)_{q_1 q_2} &\ll \left(\sum_{\nu_2=n_2}^{\infty} 2^{\nu_2 q_2^* \left(\frac{1}{p_2} - \frac{1}{q_2} \right)} \left\{ \sum_{m_2=\nu_2}^{\infty} 2^{m_2 \rho_2} \times \right. \right. \\ &\times \left. \left. \left(\sum_{\nu_1=n_1}^{\infty} \left\langle 2^{\nu_1 \left(\rho_1 + \frac{1}{p_1} - \frac{1}{q_1} \right)} \omega_{\alpha_1 + \rho_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \rho_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, \frac{1}{2^{\nu_1}}, \frac{1}{2^{m_2}} \right)_{p_1 p_2} \right\rangle^{q_1^*} \right)^{\frac{1}{q_1^*}} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

using Lemma 2 again, we deduce that

$$\begin{aligned} \omega_{\alpha_1, \alpha_2} \left(f^{(\rho_1, \rho_2)}, \frac{1}{2^{n_1}}, \frac{1}{2^{n_2}} \right)_{q_1 q_2} &\ll \left(\sum_{\nu_2=n_2}^{\infty} 2^{\nu_2 q_2^* \left(\rho_2 + \frac{1}{p_2} - \frac{1}{q_2} \right)} \times \right. \\ &\times \left\{ \left(\sum_{\nu_1=n_1}^{\infty} \left\langle 2^{\nu_1 \left(\rho_1 + \frac{1}{p_1} - \frac{1}{q_1} \right)} \omega_{\alpha_1 + \rho_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \rho_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, \frac{1}{2^{\nu_1}}, \frac{1}{2^{\nu_2}} \right)_{p_1 p_2} \right\rangle^{q_1^*} \right)^{\frac{1}{q_1^*}} \right\}^{\frac{1}{q_2^*}} \ll \\ &\left(\int_0^{\delta_2} \left(\int_0^{\delta_1} \left(t_1^{-\rho_1 - \frac{1}{p_1} + \frac{1}{q_1}} t_2^{-\rho_2 - \frac{1}{p_2} + \frac{1}{q_2}} \omega_{\alpha_1 + \rho_1 + \frac{1}{p_1} - \frac{1}{q_1}, \alpha_2 + \rho_2 + \frac{1}{p_2} - \frac{1}{q_2}} \left(f, t_1, t_2 \right)_{p_1 p_2} \right)^{q_1^*} \frac{dt_1}{t_1} \right)^{\frac{q_2^*}{q_1^*}} \frac{dt_2}{t_2} \right)^{\frac{1}{q_2^*}}. \end{aligned}$$

Theorem 2 is completely proved.

References

- [1] O.V. Besov, V.P. Ilyin, S.M. Nikolsky, *Integral representation of functions and the embedding theorem*, J. Wiley and Sons, New York, 1978; translated from Russian: Nauka, Moscow, 1975.
- [2] Z. Ditzian, S. Tikhonov, *Ul'yanov and Nikol'skii-type inequalities*, Journal of Approx. Theory 133(1) (2005), 100–133.
- [3] O. Domingues, S. Tikhonov, *Embedding of smooth function spaces extrapolations, and related inequalities*, ArXiv: 1909.12818v2 [math.FA] (2019), 1–71.
- [4] P. Glazyrina, S. Tikhonov, *Jacobi weights, fractional integration, and sharp Ulyanov inequalities*, J. Approx. Theory 195 (2015), 122–140.

- [5] A. Gogatishvili, B. Opic, S. Tikhonov, W. Trebels, *Ulyanov-type inequalities between Lorentz-Zygmund spaces*, J. Fourier Anal. Appl. 20(5) (2014), 1020–1049.
- [6] M.L. Gol'dman, *Embedding of constructive and structural Lipschitz spaces in symmetric spaces*, Tr. Mat. Inst. Steklova 173 (1986), 90–112 (in Russian); translated in: Proc Steklov Inst. Math. 173(4) (1987), 93–118.
- [7] A.A. Jumabayeva, *Sharp Ul'yanov inequalities for generalized Liouville-Weyl derivatives*, Anal. Math. 43(2) (2017), 279–302.
- [8] A.A. Jumabayeva, B.V. Simonov, *Transformation of Fourier series by means of general monotone sequences*, Mat. Zametki 107(5) (2020), 674–692.
- [9] V.I. Kolyada, *On the relations between the moduli of continuity in different metrics*, Trudy Mat. Institute of the Academy of Sciences of the USSR. 181 (1988), 117–136.
- [10] Yu. Kolomoitsev, S. Tikhonov, *Properties of moduli of smoothness in $L_p(\mathbb{R}^d)$* , J. Approx. Theory 257 (2020), Article 105423. Arxiv: 1907.12788, 1–29.
- [11] Yu. Kolomoitsev, S. Tikhonov, *Hardy-Littlewood and Ulyanov inequalities*, Mem Amer. Soc. 271(1325) (2021), Arxiv: 1711.08163.
- [12] M.K. Potapov, B.V. Simonov, S. Yu. Tikhonov, *Mixed moduli of smoothness in $L_p, 1 < p < \infty$: a survey*, Surveys in Approximation Theory, 8 (2013), 1–57.
- [13] M. K. Potapov, B. V. Simonov, S. Yu. Tikhonov, *Relations between mixed moduli of smoothness and embedding theorems for Nikol'skii classes*, Proc. Steklov Inst. Math., 269 (2010), 197–207
- [14] M. K. Potapov, B. V. Simonov, S. Yu. Tikhonov, *Analogues of Ulyanov Inequalities for Mixed Moduli of Smoothness*, Methods of Fourier Analysis and Approximation Theory, Applied and Numerical Harmonic Analysis, Springer International Publishing, Switzerland, 2016, 161–179.
- [15] M. K. Potapov, B. V. Simonov, *Refinement of the relations between mixed smoothness moduli in L_1 and L_q metrics*, Siberian Math. J. 62(4) (2021), 661–677.
- [16] M. K. Potapov, B. V. Simonov, *Refinement of Relations between Mixed Moduli of Smoothness in the Metrics of L_p and L_∞* , Math. Notes 110(3) (2021), 347–362.
- [17] M. K. Potapov, B. V. Simonov, *Properties of mixed moduli of smoothness of positive order in a mixed metric*, Vestn. Mosk. un-ta. Ser. Matem. mekh. 6 (2014), 31–40.
- [18] M. K. Potapov, *Embedding theorems in a mixed metric*, Tr. MIAN SSSR 156 (1980), 143–156.
- [19] M.K. Potapov, B.V. Simonov, *Inequalities of different metrics for trigonometric polynomials*, Izvestiya vuzov. Math. 1 (2019), 49–62.
- [20] M.K. Potapov, B.V. Simonov, *Estimates of partial moduli of smoothness in metrics of $L_{p_1\infty}$ and $L_{\infty p_2}$ through partial moduli of smoothness in metrics of $L_{p_1 p_2}$* , Moscow Univ. Math. Bull. 75 (2020), 3–17.
- [21] M. K. Potapov, B. Simonov, S. Tikhonov, *Fractional moduli of smoothness*, Maks-Press, Moscow, 2016.

- [22] M. K. Potapov, B.V. Simonov, *Strengthened Ul'yanov's inequalities for partial moduli of smoothness for functions from spaces with various metrics*, Vestnik Moskov. Univ. Ser. Mat. Mekh. 3 (2019), 26–38.
- [23] B. Simonov, S. Tikhonov, *Sharp Ul'yanov-type inequalities using fractional smoothness*, J. Approx. Theory, 162(9) (2010), 1654–1684.
- [24] S. Tikhonov, *Weak type inequalities for moduli of smoothness: the case of limit value parameters*, J. Fourier Anal. Appl. 16(4) (2010), 590–608.
- [25] S. Tikhonov, *Trigonometric series of Nikol'skii classes*, Acta Math. Hungar. 114(1-2) (2007), 61–78.
- [26] S. Tikhonov, *Trigonometric series with general monotone coefficients*, Jour. Math. Anal. Appl. 326(1) (2007), 721–735.
- [27] S. Tikhonov, W. Trebels, *Ulyanov inequalities and generalized Liouville derivatives*, Proc. Roy. Soc. Edinburgh Sect. A 141(1) (2011), 205–224.
- [28] W. Trebels, *Inequalities for moduli of smoothness versus embeddings of function spaces*, Arch. Math. 94 (2010), 155–164.
- [29] P.L. Ul'yanov, *The imbedding of certain function classes H_p^ω* , Izv. Akad. Nauk SSSR Ser. Mat. 32(3) (1968), 649–686.
- [30] A. Zygmund, *Trigonometric Series*, Vol. I. II Third edition. Cambridge 2002.