

NONLINEAR RESOLVENTS AND DECREASING LOEWNER CHAINS

IKKEI HOTTA, SEBASTIAN SCHLEISSINGER, AND TOSHIYUKI SUGAWA

Dedicated to the memory of Gabriela Kohr.

ABSTRACT. In this article we prove that nonlinear resolvents of infinitesimal generators on bounded and convex subdomains of \mathbb{C}^n are decreasing Loewner chains. Furthermore, we consider the problem of the existence of nonlinear resolvents on unbounded convex domains in \mathbb{C} . In the case of the upper half-plane, we obtain a complete solution by using that nonlinear resolvents of certain generators correspond to semigroups of probability measures with respect to free convolution.

CONTENTS

| | |
|---|----|
| 1. Introduction | 1 |
| 2. Semigroups and nonlinear resolvents | 3 |
| 3. Existence of nonlinear resolvents in unbounded domains | 5 |
| 3.1. Proof of Theorem 1.2 | 5 |
| 3.2. Parallel strips | 6 |
| 4. Decreasing Loewner chains | 7 |
| 5. Upper half-plane and Denjoy-Wolff point at infinity | 12 |
| 5.1. Additive convolution | 12 |
| 5.2. Multiplicative convolution | 14 |
| References | 15 |

1. INTRODUCTION

Let $D \subset \mathbb{C}^n$ be a complete (Kobayashi) hyperbolic domain. The (decreasing) Loewner partial differential equation on D has the form

$$\frac{\partial}{\partial t} f_t(z) = f'_t(z) \cdot G(t, z) \quad \text{for a.e. } t \geq 0, \quad f_0(z) = z \in D, \quad (1.1)$$

where f'_t denotes the Jacobian of f_t and $G(t, z)$ is a Herglotz vector field on D , i.e. $z \mapsto G(t, z)$ is an infinitesimal generator of a semigroup on D for almost every $t \geq 0$ and $(t, z) \mapsto G(t, z)$ satisfies certain regularity conditions. Equation (1.1) has a unique solution of univalent mappings $f_t : D \rightarrow D$, which form a decreasing Loewner chain, i.e. $f_t(D) \subset f_s(D)$ whenever $s \leq t$, f_0 is the identity, and $t \mapsto f_t$ is continuous. This result is well-known, but as the literature usually focuses on increasing Loewner chains, we also provide a proof (Theorem 4.4).

Now let us replace $G(t, z)$ by $G(t, f_t(z))$ to obtain the following modified equation:

$$\frac{\partial}{\partial t} f_t(z) = f'_t(z) \cdot G(t, f_t(z)) \quad \text{for a.e. } t \geq 0, \quad f_0(z) = z \in D. \quad (1.2)$$

At first sight, it seems unnatural to look at this equation, for, if G is a generator on D and $\varphi : D \rightarrow D$ a holomorphic self-mapping, then $G \circ \varphi$ need not be a generator. On the other hand, equation (1.2)

2020 *Mathematics Subject Classification.* Primary 37L05, Secondary 30C45, 46L54.

Key words and phrases. nonlinear resolvents, semigroups, decreasing Loewner chains, free convolution.

The first author was supported by JSPS KAKENHI Grant no.20K03632.

The second author was supported by the German Research Foundation (DFG), project no. 401281084.

The third author was supported by JSPS KAKENHI Grant Number JP17H02847.

is simpler than (1.1) in the following sense. While the inverse functions $g_t = f_t^{-1}$ of the solution to (1.1) satisfy Loewner's ordinary differential equation

$$\frac{\partial}{\partial t} g_t(z) = -G(t, g_t(z)),$$

the inverse functions g_t of the solution to (1.2), if it exists, satisfy

$$\frac{\partial}{\partial t} g_t(z) = -G(t, z), \quad \text{i.e.} \quad g_t(z) = z - \int_0^t G(s, z) ds.$$

Note that we can define g_t by this equation on the whole domain D . Special cases of (1.2) appear in different contexts: theory of continuous semigroups of holomorphic functions ([ESS20] and [GKH20]), limit of the radial/chordal multiple SLEs ([dMS16], [dMHS18], [HS20]) and semigroups of probability measures with respect to free convolution ([HS20]). In this article we explain these relations in a more general setting.

In the simplest case, $G(t, z)$ does not depend on t , i.e. $G(t, z) = G(z)$ for an infinitesimal generator G on D . The solution to (1.1) is then simply the semigroup associated to G . By approximating the general version of (1.1) by a piecewise constant Herglotz vector field, we can think of the solution as an infinite composition of semigroup mappings which are infinitesimally close to the identity.

Equation (1.2), on the other hand, is related to the nonlinear resolvents of G . Assume that D is bounded and convex. It is well-known that the nonlinear resolvent equation (at time t)

$$w = z - tG(z)$$

has a unique solution $f_t(z)$ for every $w \in D$ and every $t \geq 0$. Each f_t is a univalent self-map of D .

Theorem 1.1. *Let $D \subset \mathbb{C}^n$ be a domain which is bounded and convex.*

- (1) *Let G be an infinitesimal generator on D . Then there exists a unique solution f_t to*

$$\frac{\partial}{\partial t} f_t(z) = f_t'(z) \cdot G(f_t(z)) \quad \text{for all } t \geq 0, \quad f_0(z) = z \in D. \quad (1.3)$$

The mappings f_t are the nonlinear resolvents of G and they form a decreasing Loewner chain.

- (2) *More generally, let $G : [0, \infty) \times D \rightarrow \mathbb{C}^n$ be such that $z \mapsto G(t, z)$ is holomorphic for a.e. $t \geq 0$, $t \mapsto G(t, z)$ is locally integrable for all $z \in D$, and $H_t(z) := \int_0^t G(s, z) ds$ is an infinitesimal generator on D for every $t \geq 0$. Let J_t be the nonlinear resolvent of H_t at time 1. Then $(J_t)_{t \geq 0}$ is the unique solution of holomorphic self-mappings of D , locally absolutely continuous in t , of*

$$\frac{\partial}{\partial t} J_t(z) = J_t'(z) \cdot G(t, J_t(z)) \quad \text{for a.e. } t \geq 0, \quad J_0(z) = z \in D.$$

In contrast to equation (1.3), the solution in Theorem 1.1 (2) is not necessarily a decreasing Loewner chain, see Example 4.8.

Convexity of D is a natural assumption for the resolvent equation. On the upper half-plane \mathbb{H} – which is unbounded, convex, and still complete hyperbolic – the resolvent equation does not have solutions for all cases, see Example 2.4. The following theorem shows that this is rather an exceptional case.

Theorem 1.2. *Let D be an unbounded convex domain in \mathbb{C} with $D \neq \mathbb{C}$ and $G : D \rightarrow \mathbb{C}$ be an infinitesimal generator of a semigroup (F_t) on D . If the Denjoy-Wolff point σ of (F_t) is finite, then G admits nonlinear resolvents J_t on D for all $t \geq 0$.*

A definition of the Denjoy-Wolff point will be given in Section 2. If the Denjoy-Wolff point is ∞ , the question concerning the existence of nonlinear resolvents is more complicated. In Section 3.2 we give a sufficient condition for generators on a horizontal strip having Denjoy-Wolff point ∞ . For the case $D = \mathbb{H}$, we obtain a complete characterization.

Theorem 1.3. *Let $G : \mathbb{H} \rightarrow \mathbb{C}$ be an infinitesimal generator on \mathbb{H} .*

- (1) *Assume that $G(\mathbb{H}) \subset \mathbb{H} \cup \mathbb{R}$. (This is equivalent to either $G(z) \equiv 0$ or the associated semigroup has Denjoy-Wolff point ∞ .) Let $b = \lim_{y \rightarrow \infty} G(iy)/(iy)$. Then b is a non-negative real number and the resolvent equation for G can be solved if and only if $t \in [0, 1/b)$.*

If $b = 0$, then the resolvents f_t satisfy

$$\frac{1}{f_t(z)} = \int_{\mathbb{R}} \frac{1}{z - x} \mu_t(dx),$$

where $(\mu_t)_{t \geq 0}$ is a semigroup of probability measures on \mathbb{R} with respect to additive free convolution.

(2) In all other cases, the nonlinear resolvent of G exists for every $t \geq 0$.

The resolvents in Theorem 1.3 also satisfy (1.2). Note that Theorem 1.3 (2) follows directly from Theorem 1.2.

For the case of Theorem 1.3 (1) we also prove that the non-autonomous version has a solution.

Theorem 1.4. *Let $G(t, z)$ be a Herglotz vector field on \mathbb{H} such that, for a.e. $t \geq 0$, $z \mapsto G(t, z)$ maps \mathbb{H} into $\mathbb{H} \cup \mathbb{R}$ and $\lim_{y \rightarrow \infty} G(t, iy)/y = 0$. Then there exists a unique solution $(f_t)_{t \geq 0}$, locally absolutely continuous in t , to*

$$\frac{\partial}{\partial t} f_t(z) = f'_t(z) \cdot G(t, f_t(z)) \quad \text{for a.e. } t \geq 0, \quad f_0(z) = z \in \mathbb{H}.$$

The solution can also be written as $1/f_t(z) = \int_{\mathbb{R}} \frac{1}{z-x} \mu_t(dx)$ for a family of freely infinitely divisible probability measures on \mathbb{R} and $(f_t)_{t \geq 0}$ forms a decreasing Loewner chain on \mathbb{H} .

The paper is organized as follows.

- In Section 2 we briefly recall some facts on semigroups and nonlinear resolvents.
- The proof of Theorem 1.2 is provided in Section 3.
- In Section 4 we look at the general theory of decreasing Loewner chains and we prove Theorem 1.1.
- The proofs of Theorem 1.4 and part (1) of Theorem 1.3 are based on notions from free probability theory, which we discuss in Section 5.

2. SEMIGROUPS AND NONLINEAR RESOLVENTS

Let $D \subset \mathbb{C}^n$ be a domain. For a holomorphic map $f : D \rightarrow \mathbb{C}^n$ we denote by $f'(z)$ its (complex) Jacobian matrix at z . For $n = 1$, $f'(z)$ is the ordinary derivative of $f(z)$. We denote by $I : \mathbb{C}^n \rightarrow \mathbb{C}^n$ the identity mapping.

Let $(F_t)_{t \geq 0}$ be a semigroup of holomorphic functions on D , i.e. F_0 is the identity, $F_{t+s} = F_s \circ F_t$, and $t \mapsto F_t$ is continuous with respect to locally uniform convergence on D . In the sequel, we sometimes call it a *semigroup* for short. Then the locally uniform limit

$$\lim_{t \downarrow 0} \frac{F_t(z) - z}{t} =: G(z)$$

exists and the holomorphic function $G : D \rightarrow \mathbb{C}^n$ is called the *infinitesimal generator* (or, simply, the generator) of the semigroup, see [Aba92, Theorem 5]. (Sometimes, e.g. in [ESS20], the function $-G$ is called the infinitesimal generator.) We denote by $\mathcal{G}(D)$ the set of infinitesimal generators on D .

The Berkson-Porta formula [BP78] gives a form of infinitesimal generators on the unit disc \mathbb{D} .

Lemma 2.1. *Let $G \in \mathcal{G}(\mathbb{D})$. Then, there exist a point $\tau \in \overline{\mathbb{D}} := \mathbb{D} \cup \partial\mathbb{D}$ and a holomorphic function $p : \mathbb{D} \rightarrow \mathbb{C}$ with $\operatorname{Re}(p) \geq 0$ such that*

$$G(z) = (\tau - z)(1 - \bar{\tau}z)p(z). \tag{2.1}$$

Conversely, any function $G(z)$ of the form (2.1) is an infinitesimal generator on \mathbb{D} .

If $G(z) \not\equiv 0$, we call τ the *Denjoy-Wolff point* of the semigroup $(F_t)_{t \geq 0}$ generated by G . Unless $(F_t)_{t \geq 0}$ consists of elliptic automorphisms, we have $F_t(z) \rightarrow \tau$ as $t \rightarrow \infty$ locally uniformly on \mathbb{D} . Here, we recall that an automorphism of \mathbb{D} is called *elliptic* if it has an interior fixed point in \mathbb{D} . If $(F_t)_{t \geq 0}$ does consist of elliptic automorphisms, then its Denjoy-Wolff point is defined to be its unique common fixed point, i.e. $\tau \in \mathbb{D}$ and $F_t(\tau) = \tau$ for all $t \geq 0$. See [BCDM20] for details.

Let φ be a biholomorphic map of \mathbb{D} onto a simply connected domain D in the Riemann sphere $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$. It is known that φ extends to a continuous map $\overline{\mathbb{D}}$ to $\overline{D} \subset \widehat{\mathbb{C}}$ if and only if ∂D is locally connected (see e.g. [Pom92, Theorem 2.1]). In this case, we can generalize the notion of the Denjoy-Wolff point as follows. Let $(F_t)_{t \geq 0}$ be a semigroup on D . Suppose that the conjugated semigroup $\hat{F}_t = \varphi^{-1} \circ F_t \circ \varphi$ on \mathbb{D} has the Denjoy-Wolff point at $\tau \in \overline{\mathbb{D}}$. We may define $\sigma = \varphi(\tau) \in \overline{D}$ to be the Denjoy-Wolff point of the semigroup (F_t) on D . Indeed, $F_t \rightarrow \sigma$ locally uniformly on D as $t \rightarrow \infty$ unless (F_t) consists of automorphisms of D with a common interior fixed point at σ .

We are interested in general characterizations of infinitesimal generators. One of them is described in terms of nonlinear resolvents as an analogue of the Hille-Yosida theory for semigroups of linear operators on a Banach space.

Definition 2.2. Let D be a domain in \mathbb{C}^n and $G : D \rightarrow \mathbb{C}^n$ be a holomorphic map. Let $t \geq 0$. If the equation

$$w = z - t \cdot G(z) \quad (2.2)$$

has a unique solution z in D for each $w \in D$, the mapping $z = J_t(w) = J_t(w, G)$ is called the *nonlinear resolvent* (at “time” t) of G .

If J_t exists, it is a holomorphic mapping from D into itself.

The following characterization of infinitesimal generators is due to Reich and Shoikhet [RS97, Theorem 1.1 and Corollary 1.2].

Theorem 2.3 ([RS97]). *Let D be a bounded convex domain in \mathbb{C}^n . A holomorphic map $G : D \rightarrow \mathbb{C}^n$ admits nonlinear resolvents $z = J_t(w)$ in (2.2) for all $t \geq 0$ if and only if G is an infinitesimal generator of a semigroup (F_t) on D . Moreover, in this case,*

$$F_t = \lim_{n \rightarrow \infty} J_{\frac{t}{n}} \circ \cdots \circ J_{\frac{t}{n}} \text{ (} n\text{-times)} = \lim_{n \rightarrow \infty} J_{\frac{t}{n}}^{\circ n}$$

Example 2.4. Let

$$F_t(\zeta) = \frac{\zeta + \tanh(t/2)}{1 + \zeta \tanh(t/2)}, \quad \zeta \in \mathbb{D}, t \geq 0.$$

Then $(F_t)_{t \geq 0}$ is a semigroup of holomorphic mappings on \mathbb{D} whose Denjoy-Wolff point is 1. The generator H is given by $H(\zeta) = (1 - \zeta^2)/2$, and due to Theorem 2.3 the nonlinear resolvent always exists. On the other hand, consider the conjugate \hat{F}_t by $C(z) = (z - i)/(z + i) : \mathbb{H} \rightarrow \mathbb{D}$, namely $\hat{F}_t := C^{-1} \circ F_t \circ C$. Then

$$\hat{F}_t(z) = e^t z, \quad z \in \mathbb{H}, t \geq 0,$$

is a semigroup of holomorphic mappings on \mathbb{H} with generator $G(z) = z$. In this case the equation $w = z - tz$ has no solution in \mathbb{H} for $t \geq 1$, and the resolvent $J_t(w) = \frac{w}{1-t}$ exists only for $t \in [0, 1)$. We can consider the generator $G(z) = z$ also for $D = \mathbb{C}$, where J_t exists if and only if $t \neq 1$.

Lemma 2.5. *If D is bounded and convex, then $G \circ J_t \in \mathcal{G}(D)$ for all $t \geq 0$.*

Proof. This is clearly true for $t = 0$. So let $t > 0$. Then $(G \circ J_t)(w) = (J_t(w) - w)/t$. Now we use the fact that $f - I, f : D \rightarrow D$ holomorphic, is always an infinitesimal generator on bounded convex domains, see [RS96, Proposition 4.3], and the fact that $r \cdot G$ is an infinitesimal generator for every infinitesimal generator G and $r > 0$. \square

The following lemma can be checked easily.

Lemma 2.6. *Let $(F_t)_{t \geq 0}$ be a semigroup on a domain $D \subset \mathbb{C}^n$ with generator $G : D \rightarrow \mathbb{C}^n$ and $\varphi : \hat{D} \rightarrow D$ be a biholomorphic mapping. Then the maps $\hat{F}_t = \varphi^{-1} \circ F_t \circ \varphi$ form a semigroup on \hat{D} with generator $\hat{G}(z) = (\varphi'(z))^{-1} G(\varphi(z))$. Here, $\varphi'(z)$ is the Jacobian matrix of φ at $z \in \hat{D}$.*

Proof. Differentiation of both sides of $\varphi \circ \hat{F}_t = F_t \circ \varphi$ with respect to t gives us the formula

$$(\varphi' \circ \hat{F}_t) \frac{d\hat{F}_t}{dt} = \frac{dF_t}{dt} \circ \varphi.$$

Letting $t = 0$, we obtain the required formula. \square

We now use the Cayley transform $C : \mathbb{H} \rightarrow \mathbb{D}$, $C(z) = (z - i)/(z + i)$, to obtain a general form of generators on the upper half-plane \mathbb{H} . Let $G \in \mathcal{G}(\mathbb{D})$. Then G is expressed in the form (2.1). By the last lemma, the corresponding generator \hat{G} on \mathbb{H} has the form

$$\begin{aligned} \hat{G}(z) &= G(C(z))/C'(z) \\ &= G(C(z)) \frac{-i(z+i)^2}{2} \\ &= -(z+i)^2 (\tau - C(z))(1 - \bar{\tau}C(z)) \cdot \frac{i}{2} p(C(z)). \end{aligned}$$

If $\tau = 1$, this expression reduces to $G(z) = 2ip(C(z))$. If $\tau \neq 1$, then the first three factors form a polynomial of degree 2 with zeros $\sigma := C^{-1}(\tau) \in \mathbb{H} \cup \mathbb{R}$ and $\bar{\sigma}$. In this case we obtain $G(z) = (1 - \tau)(1 - \bar{\tau})(z - \sigma)(z - \bar{\sigma}) \cdot \frac{i}{2}p(C(z))$. Clearly, if $G(z) \neq 0$, then σ is the Denjoy-Wolff point of the semigroup \hat{F}_t . Note that $ip(C(z))$ is a *Pick function*, i.e. a holomorphic function from \mathbb{H} into $\mathbb{H} \cup \mathbb{R}$. We denote by \mathcal{N} the set of all Pick functions. We can summarize our calculation as follows (it is essentially contained in [BP78], where the right half-plane is considered).

Lemma 2.7. *Every $G \in \mathcal{G}(\mathbb{H})$ has the form*

$$G(z) = q(z) \quad \text{or} \quad G(z) = (z - \sigma)(z - \bar{\sigma})q(z),$$

with $q \in \mathcal{N}$, $\sigma \in \mathbb{H} \cup \mathbb{R}$. Conversely, all functions of these forms are infinitesimal generators on \mathbb{H} . The first case corresponds to all semigroups whose Denjoy-Wolff point is ∞ unless $G = 0$. In the second case, σ is the Denjoy-Wolff point unless $G = 0$.

Example 2.8. For the semigroup $F_t(z) = e^{-t}z$ on \mathbb{D} with generator $H(z) = -z$ we obtain the resolvents $J_t(w) = w/(1+t)$ for all $t \geq 0$. The conjugated resolvent on \mathbb{H} is given by $(C^{-1} \circ J_t \circ C)(z) = i[2z + t(z+i)]/[2i + t(z+i)]$. On the other hand, by the above argument, the generator of the conjugated semigroup \hat{F}_t on \mathbb{H} is calculated as $G(z) = \frac{i}{2}(z^2 + 1) = (z - i)(z + i) \cdot \frac{i}{2}$. Since one can observe that the equation $w = z - t \cdot G(z)$ does not lead to a Möbius map, the nonlinear resolvents of a conjugated semigroup are in general not equal to the conjugated resolvents.

3. EXISTENCE OF NONLINEAR RESOLVENTS IN UNBOUNDED DOMAINS

3.1. Proof of Theorem 1.2. We denote by $\mathbb{D}(a, r)$ the disk $|z - a| < r$ in the complex plane \mathbb{C} . For $\tau \in \mathbb{D}$ and $0 < \rho < 1$, we define

$$\Delta(\tau, \rho) = \{z \in \mathbb{D} : \left| \frac{z - \tau}{1 - \bar{\tau}z} \right| < \rho\} = \mathbb{D}\left(\frac{(1 - \rho^2)\tau}{1 - \rho^2|\tau|^2}, \frac{(1 - |\tau|^2)\rho}{1 - \rho^2|\tau|^2}\right),$$

which is the hyperbolic disc of center τ with radius $\frac{1}{2} \log \frac{1 + \rho}{1 - \rho}$. For $\tau \in \partial\mathbb{D}$ and $0 < R < +\infty$, we define

$$E(\tau, R) = \{z \in \mathbb{D} : \frac{|z - \tau|^2}{1 - |z|^2} < R\} = \mathbb{D}\left(\frac{\tau}{1 + R}, \frac{R}{1 + R}\right),$$

which is often called a horocycle of center τ . Note that

$$\bigcup_{0 < \rho < 1} \Delta(a, \rho) = \mathbb{D} \quad \text{and} \quad \bigcup_{0 < R < +\infty} E(\tau, R) = \mathbb{D}$$

for each $a \in \mathbb{D}$ and $\tau \in \partial\mathbb{D}$.

Let $f : \mathbb{D} \rightarrow \mathbb{D}$ be a non-identity holomorphic map. We define the *Denjoy-Wolff point* of f similar to the definition for semigroups. If f is not an elliptic automorphism, then the Denjoy-Wolff point is the unique point $\tau \in \overline{\mathbb{D}}$ such that $f^{\circ n}$, the n -th iterate of f , converges to τ locally uniformly on \mathbb{D} as $n \rightarrow \infty$. If f is an elliptic automorphism, we define the Denjoy-Wolff point of f as the unique fixed point $\tau \in \mathbb{D}$ of f .

If $(f_t)_{t \geq 0}$ is a semigroup on \mathbb{D} with Denjoy-Wolff point τ , then the Denjoy-Wolff point of each f_t is equal to τ as well provided f_t is not the identity.

Lemma 3.1. *Let $f : \mathbb{D} \rightarrow \mathbb{D}$ be a non-identity holomorphic map with Denjoy-Wolff point $\tau \in \overline{\mathbb{D}}$.*

- (i) *When $|\tau| < 1$, $f(\Delta(\tau, \rho)) \subset \Delta(\tau, \rho)$ for all $\rho \in (0, 1)$.*
- (ii) *When $|\tau| = 1$, $f(E(\tau, R)) \subset E(\tau, R)$ for all $R > 0$.*

The assertion (i) follows from the Schwarz-Pick lemma and the assertion (ii) is contained in the Denjoy-Wolff theorem (see [BCDM20, Theorem 1.8.4]).

For the proof of Theorem 1.2, we also need the following well-known result due to Study [Stu11] and Robertson [Rob36].

Lemma 3.2. *Let φ be a holomorphic function mapping \mathbb{D} univalently onto a convex domain. Then $\varphi(\Delta)$ is convex for every disk Δ contained in \mathbb{D} .*

Proof of Theorem 1.2. Let D be an unbounded convex domain in \mathbb{C} with $D \neq \mathbb{C}$ and $G : D \rightarrow \mathbb{C}$ be an infinitesimal generator of a semigroup (F_t) on D . Furthermore, we assume that the Denjoy-Wolff point σ of (F_t) is finite. We need to show that G admits nonlinear resolvents J_t on D for all $t \geq 0$. First, let $\varphi : \mathbb{D} \rightarrow D$ be a conformal mapping. As D is convex, it is a Jordan domain in the Riemann sphere unless D is a parallel strip, see [Koe87, Corollary 2]. Therefore, if D is a Jordan domain, φ

extends to a homeomorphism $\overline{\mathbb{D}} \rightarrow \overline{D}$ by a theorem of Carathéodory. Even when D is a parallel strip; i.e., $\varphi = A \log \frac{1+\psi}{1-\psi} + B$ for some disk automorphism ψ and constants A, B with $A \neq 0$, φ extends to a continuous map $\overline{\mathbb{D}} \rightarrow \overline{D}$.

We now consider the conjugated semigroup $\hat{F}_t = \varphi^{-1} \circ F_t \circ \varphi$ and denote by \hat{G} its generator. Then the Denjoy-Wolff point τ of (\hat{F}_t) is mapped to σ by φ .

Case 1: $\sigma \in D$: Then $\tau \in \mathbb{D}$. Let $D_\rho = \varphi(\Delta(\tau, \rho))$ for $0 < \rho < 1$. Note that D_ρ is a bounded convex domain by Lemma 3.2. Then Lemma 3.1 now implies that $F_t(D_\rho) \subset D_\rho$ for each ρ . (If F_t is the identity for some t , then trivially $F_t(D_\rho) \subset D_\rho$.) It enables us to regard (F_t) as a semigroup on the bounded convex domain D_ρ . By Theorem 2.3, we obtain nonlinear resolvents $z = J_t(w)$ on D_ρ for all $t \geq 0$; that is to say, the equation $w = z - t \cdot G(z)$ has a unique solution $z \in D_\rho$ for each $w \in D_\rho$. Since $D = \cup_{0 < \rho < 1} D_\rho$, we readily see that the equation $w = z - t \cdot G(z)$ has a unique solution $z \in D$ for each $w \in D$. Thus the assertion has been shown in this case.

Case 2: $\sigma \in \partial D$: In this case we have $\tau \in \partial \mathbb{D}$ so that we use $E(\tau, R)$ instead. Let $D_R = \varphi(E(\tau, R))$ for $R > 0$. Then Lemma 3.1 again implies that $F_t(D_R) \subset D_R$ for each R . Since $\varphi(\tau) = \sigma$ is finite by assumption, D_R is bounded convex by Lemma 3.2. We can now show the assertion in the same way as Case 1. \square

3.2. Parallel strips. As another interesting example, we consider parallel strips here. Let D be the standard one defined by $|\operatorname{Im} z| < \pi/2$. First we observe the semigroup $F_t(z) = z + t$, $z \in D, t \geq 0$. Then the generator is $G(z) = 1$ and the nonlinear resolvent exists for each t and is given by $J_t(w) = w + t$. It is the same as F_t incidentally. Note that the Denjoy-Wolff point is ∞ . So far, we have the following sufficient condition for existence of nonlinear resolvents on the parallel strips when the Denjoy-Wolff point is at infinity. Recall that Theorem 1.2 covers the case when the Denjoy-Wolff point is finite.

Theorem 3.3. *Let $D = \{z \in \mathbb{C} : |\operatorname{Im} z| < \pi/2\}$ and $G \in \mathcal{G}(D)$. Suppose that the Denjoy-Wolff point σ of the semigroup generated by G is ∞ . Then $G(z) = e^{-z}q(z)$ for a holomorphic function q with $\operatorname{Re} q \geq 0$. Moreover, the nonlinear resolvents of G exist for all $t \in [0, \frac{\pi}{2c})$, where*

$$c = \inf_{x \in \mathbb{R}} |\operatorname{Im} G(x)| = \inf_{x \in \mathbb{R}} e^{-x} |\operatorname{Im} q(x)|.$$

Proof. The function $\varphi(\zeta) = \log \frac{1+\zeta}{1-\zeta}$ maps \mathbb{D} onto D . We may assume that the point $\tau = 1$ corresponds to σ under the mapping φ . Then, by Lemmas 2.1 and 2.6, $\hat{G} = G \circ \varphi / \varphi'$ has the form $(1 - \zeta)^2 p(\zeta)$, where p is a holomorphic function on \mathbb{D} with $\operatorname{Re} p \geq 0$. Hence, for $z = \varphi(\zeta)$, we obtain

$$G(z) = \hat{G}(\zeta) \varphi'(\zeta) = 2 \frac{1 - \zeta}{1 + \zeta} p(\zeta) = e^{-z} q(z).$$

Here, we put $q(z) = 2p(\varphi^{-1}(z))$. For a while, we assume that q satisfies the additional conditions

- (i) $q(z)$ is holomorphic on an open set containing the closed parallel strip $|\operatorname{Im} z| \leq \pi/2$, and
- (ii) q satisfies the inequality $|q - 1| \leq k|q + 1|$ on D for a constant $0 < k < 1$.

Condition (ii) says that the image $q(D)$ is contained in a compact subset of the right half-plane $\operatorname{Re} w > 0$.

Under these conditions, we will show that the nonlinear resolvents of G exist for all $t \geq 0$. For a given point $w_0 \in D$ and a positive number t , we consider the function $w = f(z) = z - tG(z) - w_0$. We need to show that f has a unique zero in D . Note that $|\operatorname{Im} w_0| < \pi/2$. First we observe that for $x \in \mathbb{R}$, $f(x + i\pi/2) = x + i\pi/2 + ite^{-x}q(x + i\pi/2) - w$ so that $\operatorname{Im} f(z) = \pi/2 - \operatorname{Im} w_0 + te^{-x} \operatorname{Re} q(z) > 0$ on $\operatorname{Im} z = \pi/2$. Similarly, we see that $\operatorname{Im} f(z) < 0$ on $\operatorname{Im} z = -\pi/2$. When $x = \operatorname{Re} z \rightarrow +\infty$ in D , we have $\operatorname{Re} f(z) = x + O(1)$. In particular, $\operatorname{Re} f(z) > 0$ if $\operatorname{Re} z$ is large enough. When $x = \operatorname{Re} z \rightarrow -\infty$ in D , we observe

$$\arg(-f(z)) = \arg e^{-z} + \arg(q(z) - e^z(z - w_0)/t) = -\operatorname{Im} z + \arg(q(z) + o(1)).$$

Note that $|\arg q| \leq 2 \arctan k < \pi/2$ on D by assumption. We fix a number α so that $2 \arctan k < \alpha < \pi/2$. If $-x = -\operatorname{Re} z$ is positive and large enough, then $|\arg(-f(z))| < \pi/2 + \alpha < \pi$. Now we take the boundary of the rectangle $|\operatorname{Re} z| \leq T$ and $|\operatorname{Im} z| \leq \pi/2$ as a contour Γ for a large enough $T > 0$. Then we see that

$$\frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz = \frac{1}{2\pi} \int_{f(\Gamma)} d \arg w = 1$$

as required.

Next we consider the general $G(z) = e^{-z}q(z)$ and let $0 < t < \pi/(2c)$. We first assume that $\operatorname{Re} q > 0$. Take a number c' so that $c < c' \leq \pi/(2t)$. Then by the definition of c , we can find a real number x_0 so that $t|\operatorname{Im} G(x_0)| < tc' \leq \pi/2$. Thus $w_0 = x_0 - tG(x_0) \in D$. Let $p(\zeta) = q(\varphi(\zeta) + x_0)$. Then p is holomorphic and satisfies $\operatorname{Re} p > 0$ on \mathbb{D} . For $0 < \rho < 1$, we set $G_\rho(z) = e^{-z}q_\rho(z)$, where $q_\rho(z) = p(\rho\varphi^{-1}(z - x_0))$. Then G_ρ satisfies the conditions (i) and (ii) and $G_\rho(x_0) = G(x_0)$ so that we can find the nonlinear resolvent $g_\rho(w) = J_t(w, G_\rho)$ for each ρ . Note that $g_\rho : D \rightarrow D$ is holomorphic and satisfies $g_\rho(w_0) = x_0$. Since (g_ρ) is a normal family, there exists a local uniform limit $g = \lim g_{\rho_j} : D \rightarrow \overline{D}$ for a suitable sequence $\rho_j \rightarrow 1$. Since $g(w_0) = x_0 \in D$, we see that g maps D into itself. By definition of the nonlinear resolvents, the functions g_ρ satisfy the relation

$$w = g_\rho(w) - tG_\rho(g_\rho(w)), \quad w \in D.$$

We now take the limit through the sequence ρ_j to get the relation $w = g(w) - tG(g(w))$.

Next we consider the case $\operatorname{Re} q = 0$ at some point in D . Then $q = ia$ for a real constant a . In this case, we consider $q_\rho = \rho + ia$ for a positive number $\rho > 0$. Then the same argument works as $\rho \rightarrow 0$. Hence, we have obtained the nonlinear resolvent $g(w) = J_t(w, G)$. \square

4. DECREASING LOEWNER CHAINS

Definition 4.1. A family $(f_t)_{t \geq 0}$ of univalent mappings $f_t : D \rightarrow D$ is called a *decreasing Loewner chain* if f_0 is the identity, $f_t(D) \subset f_s(D)$ whenever $s \leq t$, and $t \mapsto f_t$ is continuous in the topology of locally uniform convergence on D .

Decreasing Loewner chains can be obtained by solving a certain Loewner PDE. First, we need the following definition, which was introduced in [BCDM09] for general complete (Kobayashi) hyperbolic manifolds. We will only consider complete hyperbolic subdomains of \mathbb{C}^n . For example, every bounded and convex domain is complete hyperbolic, see [BS09, Theorem 1.1].

Definition 4.2. Let $D \subset \mathbb{C}^n$ be a complete hyperbolic domain. A *Herglotz vector field* of order $d \in [1, +\infty]$ on D is a mapping $G : [0, \infty) \times D \rightarrow \mathbb{C}^n$ with the following properties:

- (1) The function $t \mapsto G(t, z)$ is measurable on $[0, \infty)$ for all $z \in D$.
- (2) The function $z \mapsto G(t, z)$ is an infinitesimal generator for almost every $t \in [0, \infty)$.
- (3) For any compact set $K \subset D$ and $T > 0$, there exists a function $C_{T,K} \in L^d([0, T], \mathbb{R}_0^+)$ such that $\|G(t, z)\| \leq C_{T,K}(t)$ for all $z \in K$ and for almost all $t \in [0, T]$.

Remark 4.3. In [BCDM09] the definition of a Herglotz vector field involves a condition on the function $G(\cdot, t)$ (for almost all t) using the Kobayashi metric of D instead of property (2). In [AB11, Theorem 1.1], however, it was shown that this condition is equivalent to condition (2).

Theorem 4.4. Let $G(t, z)$ be a Herglotz vector field on a complete hyperbolic domain. Then there exists a decreasing Loewner chain $(f_t)_{t \geq 0}$, locally absolutely continuous in t , satisfying the partial differential equation

$$\frac{\partial}{\partial t} f_t(z) = f_t'(z) \cdot G(t, z) \quad \text{for a.e. } t \geq 0, \quad f_0(z) = z \in D, \quad (4.1)$$

for all $z \in D$. Conversely, if $(J_t)_{t \geq 0}$ is a family of holomorphic functions $J_t : D \rightarrow D$, locally absolutely continuous in t , which satisfies (4.1), then $J_t = f_t$ for all $t \geq 0$.

Proof. We follow [CDMG14, Theorem 3.2], where decreasing Loewner chains are handled by studying evolution families (arising from increasing Loewner chains).

Fix $T > 0$ and consider the Loewner equation

$$\frac{\partial}{\partial t} g_{s,t}(z) = G(T - t, g_{s,t}(z)), \quad g_{s,s}(z) = z \in D, \quad 0 \leq s \leq t \leq T,$$

which has a unique solution of univalent mappings $g_{s,t} : D \rightarrow D$ with $g_{s,s}(z) \equiv z$, $g_{s,t} = g_{u,t} \circ g_{s,u}$ for all $0 \leq s \leq u \leq t \leq T$ and for any compact subset $K \subset D$ and for any $0 < S \leq T$ there exists a non-negative function $k_{K,S} \in L^d([0, S], \mathbb{R}_0^+)$ such that for all $0 \leq s \leq u \leq t \leq S$ and for all $z \in K$,

$$\|g_{s,u}(z) - g_{s,t}(z)\| \leq \int_u^t k_{K,S}(\tau) d\tau,$$

see [BCDM09, Def. 1.2, Prop. 3.1, Prop. 5.1]. ($g_{s,t}$ is (a part of) an evolution family of order d .) In particular $g_{s,T} \circ g_{0,s} = g_{0,T}$. Differentiation with respect to s gives

$$\frac{\partial}{\partial s} g_{s,T}(g_{0,s}(z)) + g'_{s,T}(g_{0,s}(z)) \cdot G(T-s, g_{0,s}(z)) = 0,$$

i.e.

$$\frac{\partial}{\partial s} g_{s,T}(w) = -g'_{s,T}(w) \cdot G(T-s, w)$$

for all $w \in g_{0,s}(D)$. However, the right side (and thus its integral w.r.t s) can be extended holomorphically to D and thus $\frac{\partial}{\partial s} g_{s,T}(w)$ satisfies the PDE for all $w \in D$.

Now define $f_t = g_{T-t,T}$, $0 \leq t \leq T$. We have $f_t = g_{T-t,T} = g_{T-s,T} \circ g_{T-t,T-s} = f_s \circ g_{T-t,T-s}$ whenever $s \leq t$. Hence $(f_t)_{0 \leq t \leq T}$ is (a part of) a decreasing Loewner chain. We have

$$\frac{\partial}{\partial t} f_t(z) = g'_{T-t,T}(z) \cdot G(t, z) = f'_t(z) \cdot G(t, z), \quad f_0(z) = z \in D, \quad 0 \leq t \leq T.$$

By choosing another $\hat{T} > T$, we obtain a family $(\hat{g}_{s,t})_{0 \leq s \leq t \leq \hat{T}}$ with $\hat{g}_{s,\hat{T}} = g_{s+T-\hat{T}, \hat{T}+T-\hat{T}} = g_{s+T-\hat{T}, T}$ for all $s \in [\hat{T}-T, \hat{T}]$. Hence $\hat{f}_t := \hat{g}_{\hat{T}-t, \hat{T}} = g_{T-t, T} = f_t$ for all $t \in [0, T]$.

As we can choose $T > 0$ arbitrarily large, we conclude that there exists a decreasing Loewner chain $(f_t)_{t \geq 0}$ satisfying (4.1).

Finally, let $(J_t)_{t \geq 0}$ be a family of holomorphic mappings, locally absolutely continuous in t , satisfying (4.1). Let $T > 0$ and define $g_{s,t}$ as above. We have

$$\frac{\partial}{\partial t} [(J_t(g_{0,T-t}))(z)] = J'_t(g_{0,T-t}(z)) \cdot G(t, g_{0,T-t}(z)) - J'_t(g_{0,T-t}(z)) \cdot G(T, g_{0,T-t}(z)) = 0$$

and thus $J_t(g_{0,T-t}(z)) = J_0(g_{0,T}(z)) = g_{0,T}(z)$ for all $z \in D$ and $0 \leq t \leq T$. This implies $J_t = g_{0,T} \circ g_{0,T-t}^{-1}$ on $g_{0,T-t}(D)$. As $g_{0,T-t}(D)$ is an open set, the identity theorem implies $J_t = f_t$ on D . \square

Next we prove Theorem 1.1. Part (1) generalizes [ESS20, Proposition 4.5], which considers the case of the unit disc and generators with Denjoy-Wolff point 0.

Theorem 4.5.

- (1) Let $D \subset \mathbb{C}^n$ be a bounded and convex domain and let $G \in \mathcal{G}(D)$ with nonlinear resolvents $J_t : D \rightarrow D$. Then $(J_t)_{t \geq 0}$ is a decreasing Loewner chain satisfying the Loewner partial differential equation

$$\frac{\partial}{\partial t} J_t(w) = J'_t(w) \cdot G(J_t(w)) \quad \text{for all } t \geq 0, \quad J_0(w) = w \in D. \quad (4.2)$$

The domains $J_t(D)$ contract to the zero set of G , i.e. $\bigcap_{t \geq 0} J_t(D) = G^{-1}(0)$.

- (2) Let $D \subset \mathbb{C}^n$ be a (possibly unbounded) convex domain and let $G \in \mathcal{G}(D)$. Furthermore, let $J_t : D \rightarrow D$ be a family of holomorphic functions satisfying (4.2). Then J_t are the nonlinear resolvents of G on D .
- (3) Let $D \subset \mathbb{C}^n$ be a bounded and convex domain and let $G : [0, \infty) \times D \rightarrow \mathbb{C}^n$ be such that $z \mapsto G(t, z)$ is holomorphic for a.e. $t \geq 0$, $t \mapsto G(t, z)$ is locally integrable for all $z \in D$, and $H_t(z) := \int_0^t G(s, z) ds$ is an infinitesimal generator on D for every $t \geq 0$. Let $J_t : D \rightarrow D$ be the nonlinear resolvent of H_t at time 1. Then $(J_t)_{t \geq 0}$ is the unique solution of holomorphic self-mappings of D , locally absolutely continuous in t , of

$$\frac{\partial}{\partial t} J_t(z) = J'_t(z) \cdot G(t, J_t(z)) \quad \text{for a.e. } t \geq 0, \quad J_0(z) = z \in D. \quad (4.3)$$

Proof.

- (1) Let J_t be the resolvents of G . From $w = J_t(w) - tG(J_t(w))$ we get by differentiation

$$J'_t(w) - tG'(J_t(w)) \cdot J'_t(w) = I,$$

i.e. $J'_t(w) = (I - tG'(J_t(w)))^{-1}$. Differentiation with respect to t yields

$$0 = (I - tG'(J_t(w))) \frac{\partial}{\partial t} J_t(w) - G(J_t(w)),$$

which gives us

$$\frac{\partial}{\partial t} J_t(w) = J'_t(w) \cdot G(J_t(w)).$$

Put $\varphi_t(z) = z - tG(z)$. As $(\varphi_t \circ J_t)(z) = z$, each J_t is clearly a univalent function. Moreover, the continuity of $t \mapsto \varphi_t$ implies that also $t \mapsto J_t$ is continuous.

Now let $z \in D$. The statement $z \in J_t(D)$ is equivalent to $\varphi_t(z) \in D$. Hence, as D is convex, we see that $z \in J_t(D)$ implies $z \in J_s(D)$ for all $s \in [0, t]$. We conclude that $J_t(D) \subset J_s(D)$ whenever $s \leq t$ and that $(J_t)_{t \geq 0}$ is a decreasing Loewner chain. Due to Lemma 2.5, we see that $(t, z) \mapsto G(J_t(z))$ is a Herglotz vector field and thus equation (4.2) is a Loewner partial differential equation of the form (4.1).

(We can also argue as follows: J_t solves (4.2) and $G \circ J_t$ is a Herglotz vector field. Hence, Theorem 4.4 implies that $(J_t)_{t \geq 0}$ is a decreasing Loewner chain.)

Let $z \in D$. If $G(z) = 0$, then $J_t(z) = z$ for all $t \geq 0$. If $G(z) \neq 0$, then there exists $T > 0$ such that $z - t \cdot G(z) \notin D$ for all $t \geq T$ as D is bounded. Hence, $z \notin J_t(D)$ for all $t \geq T$ and we conclude $\bigcap_{t \geq 0} J_t(D) = G^{-1}(0)$.

- (2) Now let $J_t : D \rightarrow D$ be a family of holomorphic functions satisfying (4.2), where D is a convex domain. (Note that now, we do not know yet whether $(t, z) \mapsto G(J_t(z))$ is a Herglotz vector field.) Consider the differential equation

$$\frac{\partial}{\partial t} \varphi_t(z) = -G(J_t(\varphi_t(z))), \quad \varphi_0(z) = z.$$

Fix $z \in D$. We can solve this equation at least for t small enough. A small computation shows that $\frac{d}{dt}[J_t(\varphi_t(z))] = 0$, i.e. $J_t(\varphi_t(z))$ does not depend on t and $J_t(\varphi_t(z)) = J_0(\varphi_0(z)) = z$. Hence $\frac{\partial}{\partial t} \varphi_t(z) = -G(z)$. We conclude that $t \mapsto \varphi_t(z)$ simply describes a straight line:

$$\varphi_t(z) = z - tG(z).$$

In particular, we can now define $\varphi_t(z)$ for all $z \in D$ and all $t \geq 0$ by $\varphi_t(z) = z - tG(z)$.

Let $D_t = \{z \in D \mid \varphi_t(z) \in D\}$. The convexity of D implies that $D_t \subset D_s$ whenever $s \leq t$. Thus, for all $z \in D_t$, we have $J_t(\varphi_t(z)) = z$. Applying φ_t gives $\varphi_t(J_t(w)) = w$ for all $w \in \varphi_t(D_t)$. As the left side extends holomorphically to D , we see that $\varphi_t(J_t(w)) = w$ for all $w \in D$. Applying J_t gives $J_t(\varphi_t(z)) = z$ for all $z \in J_t(D)$. We conclude that J_t is the nonlinear resolvent of G .

- (3) By definition, J_t is the inverse of $\varphi_t : z \mapsto z - \int_0^t G(s, z) ds$. Clearly, $t \mapsto \varphi_t(z)$ is locally absolutely continuous for any $z \in D$. Furthermore, as D is bounded, the set $\{J_\tau \mid \tau \geq 0\}$ is a normal family and thus $\{J'_\tau \mid \tau \geq 0\}$ is locally uniformly bounded.

Let $s, t \geq 0$ and $z \in D$. Put $w = J_t(z)$. If s is close enough to t , then $z'(s) = \varphi_s(w) \in D$. We have

$$J_t(z) - J_s(z) = J_t(z) - J_s(z'(s)) + J_s(z'(s)) - J_s(z) = J_s(z'(s)) - J_s(z).$$

From $z'(s) - z = \varphi_s(w) - \varphi_t(w)$, we see that $\tau \mapsto J_\tau(z)$ is locally absolutely continuous for any $z \in D$. J_t solves the equation $w = J_t(w) - \int_0^t G(s, J_t(w)) ds$. We get by differentiation

$$J'_t(w) - \int_0^t G'(s, J_t(w)) \cdot J'_t(w) ds = I,$$

i.e. $J'_t(w) = (I - \int_0^t G'(s, J_t(w)) ds)^{-1}$. Differentiation with respect to t yields

$$0 = (I - \int_0^t G'(s, J_t(w)) ds) \frac{\partial}{\partial t} J_t(w) - G(t, J_t(w)),$$

which gives us

$$\frac{\partial}{\partial t} J_t(w) = J'_t(w) \cdot G(t, J_t(w)).$$

Next we show uniqueness of the solution. Let $(f_t)_{t \geq 0}$ be another solution of holomorphic mappings $f_t : D \rightarrow D$, locally absolutely continuous in t , satisfying (4.3).

Choose some $z_0 \in D$. Then we find some open disc $B \subset D$ with center z_0 and some $\varepsilon > 0$ such that f_t is injective on B for all $t \in [0, \varepsilon]$. The inverse functions g_t satisfy

$$\frac{\partial}{\partial t} g_t(w) = -G(t, w)$$

for a.e. $t \in [0, \varepsilon]$ and all $w \in \bigcap_{t \in [0, \varepsilon]} f_t(B)$ (which is non-empty for ε small enough). This implies $g_t(w) = w - \int_0^t G(s, w) ds$, which shows $f_t = J_t$ on B for all $t \in [0, \varepsilon]$. The identity theorem implies $f_t = J_t$ on D for all $t \in [0, \varepsilon]$.

In this way, we also see that the set of all $t \geq 0$ with $f_t = J_t$ is open (in $[0, \infty)$). At the same time, it is also closed, and thus equal to $[0, \infty)$. \square

We can also prove a slight variation of the second part of the theorem.

Lemma 4.6. *Let $D \subset \mathbb{C}^n$ be bounded and convex and let $G : D \rightarrow \mathbb{C}^n$ be bounded and holomorphic. Let $(J_t)_{t \geq 0}$ be a family of holomorphic self-mappings of D such that*

$$\frac{\partial}{\partial t} J_t(w) = J_t'(w) \cdot G(J_t(w)) \quad \text{for all } t \geq 0, \quad J_0(w) = w \in D.$$

Then $G \in \mathcal{G}(D)$.

Proof. The proof of part (2) of Theorem 4.5 shows that the nonlinear resolvent equation of G can be solved for all $t \geq 0$. It follows from the boundedness of G and [RS97, Corollary 1.2] that $G \in \mathcal{G}(D)$. \square

Finally, we see how (4.2) behaves with a different initial value.

Lemma 4.7. *Let $G : D \rightarrow \mathbb{C}^n$ be a holomorphic function on a bounded and convex domain $D \subset \mathbb{C}^n$ and let $\varphi : D \rightarrow D$ be univalent such that $G \circ \varphi \in \mathcal{G}(D)$. Then there exists a unique solution to the initial value problem*

$$\frac{\partial}{\partial t} J_t(w) = J_t'(w) \cdot G(J_t(w)) \quad \text{for all } t \geq 0, \quad J_0 = \varphi. \quad (4.4)$$

We have $(J_t)_{t \geq 0} = (\varphi \circ K_t)_{t \geq 0}$, where $(K_t)_{t \geq 0}$ are the nonlinear resolvents of $G \circ \varphi$.

Proof. Let $(K_t)_{t \geq 0}$ be the nonlinear resolvents of the generator $G \circ \varphi$. They satisfy

$$\frac{\partial}{\partial t} K_t(z) = K_t'(z) \cdot G(\varphi(K_t(z))), \quad K_0(z) = z. \quad (4.5)$$

Now put $J_t = \varphi \circ K_t$. Then

$$\frac{\partial}{\partial t} J_t(z) = (\varphi(K_t(z)))' \cdot K_t'(z) \cdot G(\varphi(K_t(z))) = J_t'(z) \cdot G(J_t(z)), \quad J_0 = \varphi.$$

Next we show uniqueness of the solution. Let $(\hat{J}_t)_{t \geq 0}$ be another solution of (4.4). Choose a compact ball $B \subset D$. There exists $\varepsilon > 0$ such that for all $t < \varepsilon$ and for all $z \in B$ we have $\hat{J}_t(z) \in \varphi(D)$. Hence we can define $\hat{K}_t(z) = (\varphi^{-1} \circ \hat{J}_t)(z)$ for all such z and t . A small calculation shows that $t \mapsto \hat{K}_t(z)$ satisfies (4.5). But the solution to this equation is unique due to Theorem 4.5 and we conclude $\hat{K}_t(z) = K_t(z)$ for all $z \in B$ and $t < \varepsilon$. The identity theorem implies $\hat{J}_t(z) = J_t(z)$ for all $z \in D$ and all $t < \varepsilon$.

Let $T > 0$ the infimum of all t with $\hat{J}_t \neq J_t$ and assume $T < \infty$. Then $\hat{J}_T = J_T$ due to continuity and thus $\hat{J}_T(D) = (\varphi \circ K_T)(D) \subset \varphi(D)$. Now a similar argument shows that $\hat{J}_t = J_t$ for all $t \in [0, T + \varepsilon]$ for some $\varepsilon > 0$, a contradiction. \square

For the rest of this section, we look at the unit disc. First, we construct an example of a vector field $G(t, z)$ as in Theorem 4.5 (3) whose solution is not a decreasing Loewner chain.

Example 4.8. Let $D = \mathbb{D}$ and consider the two infinitesimal generators

$$G_1(z) = -z \frac{1+z}{1-z} \quad \text{and} \quad G_2(z) = -z \frac{1-z}{1+z}.$$

We define the Herglotz vector field $G(t, z)$ by $G(t, z) = G_1(z)$, $t \leq 1$, $G(t, z) = G_2(z)$, $t > 1$. As any combination $sG_1 + tG_2$, $s, t \geq 0$, is again a generator on \mathbb{D} , the conditions in Theorem 4.5 (3) are satisfied and we let $(J_t)_{t \geq 0}$ be the solution to (1.2).

For $t \leq 1$, J_t is simply the resolvent of G_1 . For $t = 1$, the solution of $w = z + tz(1+z)/(1-z)$ is given by $J_1(w) = w/(w+2)$.

For $t \geq 1$, we have

$$\frac{\partial}{\partial t} J_t(w) = J_t'(z) \cdot G_2(J_t(z)), \quad J_1(z) = z/(z+2).$$

Here, the conditions of Lemma 4.7 are not satisfied: Assume that $H := G_2 \circ J_1 \in \mathcal{G}(\mathbb{D})$.

We have $H(z) = -J_1(z)(1 - J_1(z))/(1 + J_1(z))$. As $H(0) = 0$ and H is not constant 0, the Berkson-Porta formula (2.1) implies $\operatorname{Re}(-H(z)/z) > 0$ on \mathbb{D} . However, for $z = e^{3i}$ we obtain $\operatorname{Re}(J_1(z)(1 - J_1(z))/(1 + J_1(z))/z) = \operatorname{Re}\left(\frac{1}{2+3e^{3i}+e^{6i}}\right) = -0.47113... < 0$.

The solution $(J_t)_{t \geq 0}$ can thus not be a decreasing Loewner chain. This can be seen as follows:

If it was a decreasing Loewner chain, we can define the mappings $(J_1^{-1} \circ J_t)_{t \geq 1}$. With the proof of Lemma 4.7, we see that these functions satisfy the conditions of Lemma 4.6 for the function $G_2 \circ J_1$. Hence, $G_2 \circ J_1$ would be a generator. (In order to apply Lemma 4.6, $G_2 \circ J_1$ has to be bounded. But we can consider the whole construction with $G_2(z)$ being replaced by the generator $G_2(Rz)$ for some $R \in (0, 1)$ and close enough to 1.)

Example 4.9. If we replace the generator G_2 in Example 4.8 by $G_2(z) = -z$, then $-G_2(J_1(z))/z = J_1(z)/z = 1/(z + 2)$ has positive real part in \mathbb{D} . By Lemma 4.7 and Theorem 4.5 (1), equation (1.2) has a unique solution, which is a decreasing Loewner chain. Hence we see that there do exist vector fields as in Theorem 4.5 (3) which are non-constant with respect to time such that the solution to (1.2) is a decreasing Loewner chain.

Remark 4.10. Numerical calculations suggest that every Herglotz vector field which jumps at some $T > 0$ from $G_1(z) = -z(1 + e^{i\alpha}z)/(1 - e^{i\alpha}z)$ to $G_2(z) = -z(1 + e^{i\beta}z)/(1 - e^{i\beta}z)$, $\alpha, \beta \in \mathbb{R}$, leads to the same situation as in Example 4.8, i.e. the solution is not a decreasing Loewner chain.

It seems to be less clear to decide whether the solution to (1.2) is a decreasing Loewner chain if $G(t, z)$ varies continuously with respect to t . For example, consider the ‘‘slit’’ Herglotz vector field

$$G(t, z) = -z \frac{\kappa(t) - z}{\kappa(t) + z}$$

for a continuous function $\kappa : [0, \infty) \rightarrow \partial\mathbb{D}$. There is quite some literature on the question as to which conditions on κ guarantee that (1.1) produces slit mappings, i.e., for all $t > 0$, the image domain of f_t should have the form $f_t(\mathbb{D}) = \mathbb{D} \setminus \gamma_t$ for a simple curve $\gamma_t \subset \overline{\mathbb{D}}$. (Roughly speaking, if κ is smooth enough, e.g. continuously differentiable, then f is a slit mapping. We refer to [ZZ18] and the references therein for such results.)

We might ask: for which κ does the solution to (1.2) form a decreasing Loewner chain?

In the case of the unit disc, we can obtain some further information about the Herglotz vector field $G \circ J_t$ appearing in (4.2).

Theorem 4.11. *Let $G(z) = (\tau - z)(1 - \bar{\tau}z)p(z) \in \mathcal{G}(\mathbb{D})$ with resolvents $(J_t)_{t \geq 0}$. Then the Herglotz vector field $G_t := G \circ J_t$ can be written as*

$$G_t(z) = (\tau - z)(1 - \bar{\tau}z)p_t(z) \quad \text{with} \quad p_t(z) = \frac{1}{t} \frac{z - J_t(z)}{(z - \tau)(1 - \bar{\tau}z)}, \quad \operatorname{Re} p_t \geq 0, \quad \text{for all } t > 0. \quad (4.6)$$

Proof. If $p(z) \equiv 0$, then $J_t(z) \equiv z$ and the statement trivially holds true. So assume that $p \not\equiv 0$. Then the Herglotz vector field $G_t = G \circ J_t$ can be written as $G_t(z) = \frac{z - J_t(z)}{t} = (z - \tau_t)(1 - \bar{\tau}_t z)p_t(z)$ for some functions $\tau_t : [0, \infty) \rightarrow \overline{\mathbb{D}}$, $p_t : [0, \infty) \rightarrow \{f : \mathbb{D} \rightarrow \mathbb{C} \mid f \text{ holomorphic, } \operatorname{Re} f \geq 0\}$.

Case 1: $\tau \in \mathbb{D}$. Then $G_t(\tau) = 0$ and thus $J_t(\tau) = \tau$, which shows $G_t(\tau) = 0$ and thus $\tau_t = \tau$ for all $t \geq 0$.

Case 2: $\tau \in \partial\mathbb{D}$. It is shown in [ESS20, Lemma 5.1] (plus the text following Lemma 5.1) that the Denjoy-Wolff point of J_t is equal to τ for each $t > 0$. For the sake of completeness, we add the necessary details.

By [CDMP06, Theorem 1] and [Sha93, The Grand Iteration Theorem, p. 78], we know that $G(\tau) = 0$ and $G'(\tau) \leq 0$ in the sense of angular limits. Thus, for $t > 0$, the function $\varphi_t(z) = z - tG(z)$ satisfies $\varphi_t(\tau) = \tau$ and $\varphi_t'(\tau) \geq 1$.

As G has no zeros in \mathbb{D} , φ_t , and thus also J_t , has no fixed points in \mathbb{D} . Hence, the Denjoy-Wolff point ρ of J_t lies on $\partial\mathbb{D}$. We have $J_t(\rho) = \rho$ and $J_t'(\rho) \in (0, 1]$. This implies $\varphi_t'(\rho) = \rho$ and $\varphi_t'(\rho) \geq 1$ and thus $G(\rho) = 0$ and $G'(\rho) \leq 0$. But then ρ is the Denjoy-Wolff point of the semigroup associated to G , i.e. $\rho = \tau$, again, by [CDMP06, Theorem 1] and [Sha93, The Grand Iteration Theorem, p. 78].

We conclude $J_t(\tau) = \tau$ and $J_t'(\tau) \in (0, 1]$. As $G_t(z) = (J_t(z) - z)/t$, we have $G_t(\tau) = 0$ and $G_t'(\tau) \leq 0$, again in the sense of angular limits. This implies that the Denjoy-Wolff point of the semigroup associated to G_t is equal to τ . We conclude $\tau_t = \tau$ for all $t \geq 0$. \square

5. UPPER HALF-PLANE AND DENJOY-WOLFF POINT AT INFINITY

In [HS20, Section 4 and Appendix A1], the authors explain that free semigroups of probability measures lead to certain decreasing Loewner chains. By comparing the differential equation of these Loewner chains to (1.2), we see that such semigroups are closely related to nonlinear resolvents of certain infinitesimal generators on the upper half-plane \mathbb{H} .

In the following we will use the Nevanlinna representation formula for Pick functions, which corresponds to the Herglotz representation for Carathéodory functions.

Lemma 5.1 (e.g. [Cau32, Theorem 1]). *If $q \in \mathcal{N}$, then there exist real constants $\alpha \geq 0$ and β and a finite non-negative Borel measure ρ on \mathbb{R} such that*

$$q(z) = \alpha z + \beta + \int_{\mathbb{R}} \frac{1+tz}{t-z} d\rho(t), \quad z \in \mathbb{H}. \quad (5.1)$$

The number α and β are calculated via $\alpha = \lim_{y \rightarrow \infty} f(iy)/(iy)$ and $\beta = \operatorname{Re} q(i)$, respectively. Conversely, the function expressed as above is a Pick function.

5.1. Additive convolution. Let μ be a probability measure on \mathbb{R} . The Cauchy transform (or Stieltjes transform) of a probability measure μ on \mathbb{R} is given by

$$G_{\mu}(z) := \int_{\mathbb{R}} \frac{1}{z-t} \mu(dt), \quad z \in \mathbb{H} = \{z \in \mathbb{C} \mid \operatorname{Im}(z) > 0\}.$$

The F -transform of μ is simply defined by $F_{\mu}(z) := 1/G_{\mu}(z)$ for $z \in \mathbb{H}$.

Remark 5.2. By the definition, the F -transform belongs to \mathcal{N} . Further it is known (e.g. [Maa92]) that it has the Nevanlinna representation

$$F_{\mu}(z) = z + \beta + \int_{\mathbb{R}} \frac{1+xz}{x-z} \rho(dx), \quad z \in \mathbb{H},$$

with $\beta \in \mathbb{R}$ and ρ is a finite non-negative Borel measure on \mathbb{R} . In view of Lemma 5.1, the above fact implies that F_{μ} satisfies $\lim_{y \rightarrow \infty} F_{\mu}(iy)/iy = 1$.

G_{μ} maps the upper half-plane \mathbb{H} into the lower half-plane. For some positive real constants $\alpha, \beta > 0$, G_{μ} is univalent in the set $\Gamma_{\alpha, \beta} = \{z \in \mathbb{H} \mid \operatorname{Re}(z) < \alpha \operatorname{Im}(z), |z| > \beta\}$ and we can define the compositional right inverse G_{μ}^{-1} there. We define $R_{\mu}(z) := G_{\mu}^{-1}(z) - \frac{1}{z}$, which is a holomorphic function that maps some subdomain of the lower half-plane into the lower half-plane or into \mathbb{R} ; see [BV93, Section 5].

Instead of R_{μ} , one might also regard the Voiculescu transform $\varphi_{\mu}(z)$ defined by $\varphi_{\mu}(z) := F_{\mu}^{-1}(z) - z = G_{\mu}^{-1}(1/z) - z = R_{\mu}(1/z)$. It maps some subdomain of \mathbb{H} into the lower half-plane or \mathbb{R} .

Example 5.3. The semicircle distribution $W(0, 1)$, with mean 0 and variance 1, is given by the density

$$\frac{1}{2\pi} \sqrt{4-x^2}, \quad x \in [-2, 2].$$

We have $G_{W(0,1)}(z) = \frac{z - \sqrt{z^2 - 4}}{2}$, where the branch of the square root is chosen such that $\sqrt{\cdot}$ maps $\mathbb{C}^2 \setminus [-4, \infty)$ into the upper half-plane. Solving the equation $\frac{z - \sqrt{z^2 - 4}}{4} = w$ gives $z = w + \frac{1}{w}$ and thus

$$R_{W(0,1)}(z) = z, \quad \varphi_{\mu}(z) = \frac{1}{z}.$$

Furthermore, $F_{W(0,1)}(z) = \frac{2}{z - \sqrt{z^2 - 4}}$, and as $F_{W(0,1)}$ extends continuously to $\mathbb{H} \cup \mathbb{R}$, $F_{W(0,1)}(\mathbb{H})$ is the unbounded complement of the curve $\{F_{W(0,1)}(x) \mid x \in (-2, 2)\}$ in \mathbb{H} . A small calculation yields $\operatorname{Re}(G_{W(0,1)}(x)) = \frac{x}{2}$ and $\operatorname{Im}(G_{W(0,1)}(x)) = \frac{-\sqrt{4-x^2}}{2}$ for $x \in [-2, 2]$. Thus, for $x \in (-2, 2)$, we have


$$F_{W(0,1)}(x) = \frac{1}{\frac{x}{2} + i\frac{-\sqrt{4-x^2}}{2}} = \frac{2}{x - i\sqrt{4-x^2}} = \frac{2x + 2i\sqrt{4-x^2}}{4} = \frac{x}{2} + i\sqrt{1 - \left(\frac{x}{2}\right)^2},$$

and we see that the curve is the semicircle $(\partial\mathbb{D}) \cap \mathbb{H}$.

For two probability measures μ and ν on \mathbb{R} , the *additive free convolution* \boxplus is defined by (see [BV93])

$$\varphi_{\mu \boxplus \nu}(z) = \varphi_{\mu}(z) + \varphi_{\nu}(z).$$

This convolution arises from the notion of *free independence* introduced by D. Voiculescu. Introductions to free probability theory can be found in [NS06], [Voi97].

FIGURE 1. F -transform of the semicircle distribution $W(0,1)$.

A probability measure μ on \mathbb{R} is called *freely infinitely divisible* if for every $n \in \mathbb{N}$ there exists a probability measure μ_n on \mathbb{R} such that $\mu = \mu_n \boxplus \cdots \boxplus \mu_n$ (n -fold convolution).

Theorem 5.4 (Theorem 5.10 in [BV93]). *For a probability measure μ on \mathbb{R} , the following statements are equivalent.*

- (1) μ is freely infinitely divisible.
- (2) $\mu = \mu_1$ for a \boxplus -semigroup $(\mu_t)_{t \geq 0}$ (i.e. $\mu_0 = \delta_0$, $\mu_{t+s} = \mu_t \boxplus \mu_s$ for all $s, t \geq 0$ and $t \mapsto \mu_t$ is continuous with respect to weak convergence).
- (3) $-\varphi_\mu$ extends to a Pick function, i.e. a holomorphic function from \mathbb{H} into $\mathbb{H} \cup \mathbb{R}$.
- (4) There exist $a \in \mathbb{R}$ and a finite non-negative measure ρ on \mathbb{R} such that

$$\varphi_\mu(z) = a + \int_{\mathbb{R}} \frac{1+zx}{z-x} \rho(dx), \quad z \in \mathbb{H}. \quad (5.2)$$

Conversely, given $a \in \mathbb{R}$ and a finite non-negative measure ρ on \mathbb{R} , there exists a unique \boxplus -infinitely divisible distribution μ which has the Voiculescu transform of the form (5.2).

Example 5.5. The Dirac measure $\mu = \delta_0$ is freely infinitely divisible as $\varphi_\mu(z) \equiv 0$. Example 5.3 shows that the semicircle distribution $W(0,1)$ is also freely infinitely divisible. $W(0,1)$ is the “normal distribution of free probability theory”, i.e. the limit distribution in the free central limit theorem, see [MS17, Chapter 2].

Next we prove Theorem 1.4.

Theorem 5.6. *Let $G(t, z)$ be a Herglotz vector field on \mathbb{H} such that, for a.e. $t \geq 0$, $z \mapsto G(t, z)$ maps \mathbb{H} into $\mathbb{H} \cup \mathbb{R}$ and $\lim_{y \rightarrow \infty} G(t, iy)/y = 0$. Then there exists a unique solution $(f_t)_{t \geq 0}$, locally absolutely continuous in t , to*

$$\frac{\partial}{\partial t} f_t(z) = f'_t(z) \cdot G(t, f_t(z)) \quad \text{for a.e. } t \geq 0, \quad f_0(z) = z \in \mathbb{H}. \quad (5.3)$$

The solution can also be written as $(f_t) = (F_{\mu_t})$ for a family of freely infinitely divisible probability measures $(\mu_t)_{t \geq 0}$ on \mathbb{R} and $(f_t)_{t \geq 0}$ is a decreasing Loewner chain.

If $G(t, z)$ does not depend on t , then $G(t, z) = -\varphi_{\mu_1}(z)$ and $(\mu_t)_{t \geq 0}$ is a free \boxplus -semigroup. Furthermore, the family $(f_t)_{t \geq 0}$ are the nonlinear resolvents of $-\varphi_{\mu_1}$ in this case.

Proof. Consider $H_t(z) := \int_0^t -G(s, z) ds$, $t \geq 0$. This function is of the form (5.2) and Theorem 5.4 implies that we find freely infinitely divisible probability measures $(\mu_t)_{t \geq 0}$ such that $\varphi_{\mu_t}(z) = H_t(z)$. Put $f_t = F_{\mu_t}$. Then $f_t(\varphi_{\mu_t}(z) + z) = z$ for all $z \in \mathbb{H}$. Differentiation yields

$$0 = \frac{\partial}{\partial t} f_t(\varphi_{\mu_t}(z) + z) + f'_t(\varphi_{\mu_t}(z) + z) \frac{\partial}{\partial t} \varphi_{\mu_t}(z) = \frac{\partial}{\partial t} f_t(\varphi_{\mu_t}(z) + z) - f'_t(\varphi_{\mu_t}(z) + z) G(t, z).$$

Put $w = z + \varphi_{\mu_t}(z) = f_t^{-1}(z)$. Then

$$\frac{\partial}{\partial t} f_t(w) = f'_t(w) G(t, f_t(w))$$

for all w in the image domain $f_t^{-1}(\mathbb{H})$, and thus, in particular, for all $w \in \mathbb{H}$. Clearly, f_0 is the identity as $\varphi_{\mu_0} = 0$. Due to Lemma 2.7, $G(t, f_t(z))$ is a Herglotz vector field and Theorem 4.4 implies that (f_t) is a decreasing Loewner chain. Uniqueness of the solution is shown as in the proof of Theorem 4.5 (3).

If $G(t, z)$ does not depend on t , then $G(t, z) = -\varphi_\nu$ for some probability measure ν and $\varphi_{\mu_t} = H_t(z) = t\varphi_\nu$. This shows that $\nu = \mu_1$ and that $(\mu_t)_{t \geq 0}$ is a free semigroup due to Theorem 5.4. Theorem 4.5 (2) now implies that the functions $(J_t)_{t \geq 0}$ are the resolvents of the generator $-\varphi_{\mu_1}$. \square

Remark 5.7. The fact that the F -transforms of a \boxplus -semigroup $(\mu_t)_{t \geq 0}$ form a decreasing Loewner chain is also proved in [Jek20, Proposition 5.15], even for the more general case of operator-valued free probability theory. Also in this higher dimensional case, the F -transforms, which are holomorphic functions on certain matricial upper half-planes, are the resolvents of -1 times the Voiculescu transform of μ_1 by essentially the same proof.

Remark 5.8. Let G be a generator of the form $-\varphi_{\mu_1}$. The nonlinear resolvents $(f_t)_{t \geq 0}$ of G form a decreasing Loewner chain satisfying the Loewner PDE with Herglotz vector field $G(t, z) = (G \circ f_t)(z)$. The semigroup $(g_t)_{t \geq 0}$ associated to G satisfies the Loewner PDE with Herglotz vector field $G(t, z) = G(z)$. While the mappings f_t are the F -transforms of a semigroup with respect to free convolution, the mappings g_t have a quite similar interpretation. For two probability measures μ and ν , the monotone convolution $\mu \triangleright \nu$ is defined by $F_{\mu \triangleright \nu}(z) = (F_\mu \circ F_\nu)(z)$, $z \in \mathbb{H}$.

Now there exists a family of probability measures $(\mu_t)_{t \geq 0}$ on \mathbb{R} such that $g_t = F_{\mu_t}$. As $F_{\mu_{s+t}} = F_{\mu_s} \circ F_{\mu_t}$, we see that this family is a semigroup with respect to monotone convolution, see [FHS20, Section 3], in particular [FHS20, Proposition 3.11].

A consequence of Theorem 5.6 is the following result, which finishes the proof of part (1) of Theorem 1.3.

Corollary 5.9. *Let $G : \mathbb{H} \rightarrow \mathbb{H} \cup \mathbb{R}$ be holomorphic with $b = \lim_{y \rightarrow \infty} G(iy)/(iy)$. Then the nonlinear resolvents J_t of G exist for all $t \in [0, 1/b)$. If $b \neq 0$, then the resolvent equation cannot be solved for $t \geq 1/b$.*

Proof. If $b = 0$, then G generates a free semigroup whose F -transforms yield the nonlinear resolvents of G for all $t \geq 0$ by Theorem 5.6.

Now assume that $b \neq 0$. Then (5.1) shows that $\hat{G}(z) = G(z) - bz$ also maps \mathbb{H} into $\mathbb{H} \cup \mathbb{R}$. The equation $w = z - tG(z)$ becomes $w = (1 - tb)z - t\hat{G}(z)$. For $t \geq 1/b$ and $z \in \mathbb{H}$, the right side belongs to $-\mathbb{H} \cup \mathbb{R}$. Thus it has no solution $z \in \mathbb{H}$ for any $w \in \mathbb{H}$. For $t < 1/b$, we can also write $w/(1 - tb) = z - t\hat{G}(z)/(1 - tb)$ and the previous case applied to the generator $\hat{G}(z)/(1 - tb)$ shows the existence of the nonlinear resolvent J_t of G . \square

5.2. Multiplicative convolution. For the sake of completeness we also discuss the case of multiplicative free convolution of probability measures on the unit circle \mathbb{T} . These measures again lead to nonlinear resolvents of certain generators, however, not on \mathbb{D} , as one might expect in the first place. Roughly speaking, the pullbacks of these measures to \mathbb{R} via $x \mapsto e^{ix}$ lead again to nonlinear resolvents of generators on \mathbb{H} .

Let μ be a probability measure on \mathbb{T} . The moment generating function of μ is a holomorphic function on \mathbb{D} defined by

$$\psi_\mu(z) := \int_{\mathbb{T}} \frac{xz}{1 - xz} \mu(dx) = \sum_{n=1}^{\infty} \left(\int_{\mathbb{T}} x^n \mu(dx) \right) z^n, \quad z \in \mathbb{D}.$$

The classical independence of random variables leads to the classical convolution, or Hadamard convolution, $\mu \star \nu$, with $\psi_{\mu \star \nu} = \sum_{n=1}^{\infty} \left(\int_{\mathbb{T}} x^n \mu(dx) \right) \left(\int_{\mathbb{T}} x^n \nu(dx) \right) z^n$. Other notions of independence from non-commutative probability theory lead to further convolutions. First, we need to define the η -transform of μ . Let

$$\eta_\mu(z) := \frac{\psi_\mu(z)}{1 + \psi_\mu(z)}, \quad z \in \mathbb{D}.$$

We denote by $\mathbf{P}(\mathbb{T})$ the set of all probability measures μ on \mathbb{T} . We let $\mathbf{P}_\times(\mathbb{T})$ be the set of all $\mu \in \mathbf{P}(\mathbb{T})$ with $\eta'_\mu(0) \neq 0$, i.e. the first moment of μ is $\neq 0$. Then we can invert η_μ in a neighbourhood of 0. Denote this locally defined function by η_μ^{-1} . The Σ -transform of μ is defined by

$$\Sigma_\mu(z) := \frac{1}{z} \eta_\mu^{-1}(z).$$

For two probability measures $\mu, \nu \in \mathbf{P}_\times(\mathbb{T})$, Voiculescu [Voi87] characterized *multiplicative free convolution* \boxtimes by

$$\Sigma_{\mu \boxtimes \nu}(z) = \Sigma_\mu(z) \Sigma_\nu(z) \tag{5.4}$$

in a neighbourhood of 0.

$\mu \in \mathbf{P}_\times(\mathbb{T})$ is called *freely infinitely divisible* if for every $n \in \mathbb{N}$ there exists $\mu_n \in \mathbf{P}_\times(\mathbb{T})$ such that $\mu = \mu_n \boxtimes \cdots \boxtimes \mu_n$ (n -fold convolution). Freely infinitely divisible distributions can be characterized in the following way.

Theorem 5.10 (See Lemma 6.6 in [BV92]). *Let $\mu \in \mathbf{P}_\times(\mathbb{T})$. Then the following three statements are equivalent.*

- (1) μ is freely infinitely divisible.
 - (2) There exists a \boxtimes -semigroup $(\mu_t)_{t \geq 0}$ (i.e. $\mu_0 = \delta_1$, $\mu_{t+s} = \mu_t \boxtimes \mu_s$ for all $s, t \geq 0$ and $t \mapsto \mu_t$ is continuous with respect to weak convergence) such that $\mu_1 = \mu$.
 - (3) There exists a holomorphic function $u_\mu: \mathbb{D} \rightarrow \mathbb{C}$ with $\operatorname{Re}(u_\mu) \geq 0$ such that $\Sigma_\mu(z) = \exp(u_\mu(z))$.
- Moreover, the analytic map u_μ in (3) can be characterized by the Herglotz representation

$$u_\mu(z) = -i\alpha + \int_{\mathbb{T}} \frac{1+z\zeta}{1-z\zeta} \rho(d\zeta), \quad (5.5)$$

where $\alpha \in \mathbb{R}$ and ρ is a finite non-negative Borel measure on \mathbb{T} .

Conversely, for any analytic map $u: \mathbb{D} \rightarrow \mathbb{C}$ with $\operatorname{Re}(u) \geq 0$, the function $\exp(u(z))$ is the Σ -transform of some freely infinitely divisible μ .

Remark 5.11. The multiplicative free convolution can also be considered for probability measures μ with zero mean. Such a measure is freely infinitely divisible if and only if μ is the uniform distribution on \mathbb{T} , i.e. $\psi_\mu(z) \equiv 0$, see [BV92, Lemma 6.1].

Theorem 5.12. *Let J_t be defined by $(i\frac{1+e^{iz}}{1-e^{iz}}) \circ J_t = iM_t(e^{iz})$, where $M_t(z) = \int_{\mathbb{T}} \frac{x+z}{x-z} \mu_t(dx)$ for a free \boxtimes -semigroup of probability measures μ_t on \mathbb{T} . Then J_t are the resolvents of the infinitesimal generator $G(z) = iu_{\mu_1}(-e^{iz})$ on \mathbb{H} .*

Proof. Let $\eta_t = \eta_{\mu_t}$, where $(\mu_t)_{t \geq 0}$ is a semigroup as in (2) of Theorem 5.10. Then by (3) of Theorem 5.10, we have

$$\eta_t^{-1}(z) = z \exp(tu_\mu(z)).$$

This yields the differential equation

$$\frac{\partial}{\partial t} \eta_t(z) = -zu_\mu(\eta_t(z)) \cdot \frac{\partial}{\partial z} \eta_t(z).$$

We put $M_t(z) = \frac{1+\eta_t(z)}{1-\eta_t(z)} = 1 + 2\psi_{\mu_t}$. Then we obtain the partial differential equation

$$\frac{\partial}{\partial t} M_t(z) = -zS_\mu(M_t(z)) \cdot \frac{\partial}{\partial z} M_t(z), \quad M_0(z) = \frac{1+z}{1-z},$$

with $S_\mu(z) = u_\mu(\frac{1-z}{1+z})$. Note that S_μ maps the right half-plane RH holomorphically into $RH \cup i\mathbb{R}$.

For $\operatorname{Im}(z) > 0$ define $V_t(z) = iM_t(e^{iz})$. Then V_t maps the upper half-plane into itself and we obtain

$$\frac{\partial V_t(z)}{\partial t} = iS(-iV_t(z)) \frac{\partial V_t(z)}{\partial z}, \quad V_0(z) = i\frac{1+e^{iz}}{1-e^{iz}}.$$

According to [HS20, Theorem 4.7], V_t is a decreasing subordination chain and we can write $V_t = V_0 \circ J_t$ with J_t satisfying the same equation (see the proof of [HS20, Theorem 2.2]), i.e.

$$\frac{\partial J_t(z)}{\partial t} = iS(-iV_t(z)) \frac{\partial J_t(z)}{\partial z} = iS(-i(V_0 \circ J_t)(z)) \frac{\partial J_t(z)}{\partial z} = G(J_t(z)) \frac{\partial J_t(z)}{\partial z}, \quad J_0(z) = z \in \mathbb{H},$$

with $G(z) = iS(-iV_0(z)) = iu_{\mu_1}(-e^{iz})$, which maps \mathbb{H} into $\mathbb{H} \cup \mathbb{R}$. Theorem 4.5 (2) finishes the proof. \square

Remark 5.13. As $iu_{\mu_1}(-e^{iz})$ maps \mathbb{H} into $\mathbb{H} \cup \mathbb{R}$ and $\lim_{y \rightarrow \infty} iu_{\mu_1}(-e^{-y})/(iy) = 0$, we see that the generators from Theorem 5.12 form a subset of the generators appearing in Theorem 1.3 (1).

REFERENCES

- [Aba92] M. Abate, *The Infinitesimal Generators of Semigroups of Holomorphic Maps*, Annali di Matematica pura ed applicata (IV), Vol. CLXI (1992), 167–180.
- [AB11] L. Arosio, F. Bracci, *Infinitesimal generators and the Loewner equation on complete hyperbolic manifolds*, Anal. Math. Phys. 1 (2011), no. 4, 337–350.
- [BV92] H. Bercovici, D. Voiculescu, *Lévy-Hinčîn type theorems for multiplicative and additive free convolution*, Pacific J. Math. 153 (1992), no. 2, 217–248.

- [BV93] ———, *Free convolution of measures with unbounded support*, Indiana Univ. Math. J. 42 (1993), no. 3, 733–773.
- [BP78] E. Berkson, H. Porta, *Semigroups of analytic functions and composition operators*, Michigan Math. J. 25 (1978), no. 1, 101–115.
- [BS09] F. Bracci, A. Saracco, *Hyperbolicity in unbounded convex domains*, Forum Math. 21 (2009), 815–825.
- [BCDM09] F. Bracci, M. D. Contreras, S. Díaz-Madrigal, *Evolution families and the Loewner equation. II. Complex hyperbolic manifolds*, Math. Ann. 344 (2009), no. 4, 947–962.
- [BCDM20] ———, *Continuous semigroups of holomorphic self-maps of the unit disc*, Springer Monographs in Mathematics. Springer, Cham, 2020. xxvii+566 pp.
- [CDM13] T. Casavecchia, S. Díaz-Madrigal, *A Non-Autonomous Version of the Denjoy-Wolff Theorem*, Complex Analysis and Operator Theory 7 (2013), 1457–1479.
- [Cau32] W. Cauer, *The Poisson integral for functions with positive real part*, Bull. Amer. Math. Soc. 38 (1932), 713–717.
- [CDMP06] M. D. Contreras, S. Díaz-Madrigal, C. Pommerenke, *On boundary critical points for semigroups of analytic functions*, Math. Scand. 98 (2006), 125–142.
- [CDMG10] M. D. Contreras, S. Díaz-Madrigal, P. Gumenyuk, *Loewner chains in the unit disk*, Rev. Mat. Iberoam. 26 (2010), no. 3, 975–1012.
- [CDMG14] ———, *Local duality in Loewner equations*, J. Nonlinear Convex Anal. 15 (2014), no. 2, 269–297.
- [dMHS18] A. del Monaco, I. Hotta, S. Schleißinger, *Tightness results for infinite-slit limits of the chordal Loewner equation*, Comput. Methods Funct. Theory 18 (2018), no. 1, 9–33.
- [dMS16] A. del Monaco, S. Schleißinger, *Multiple SLE and the complex Burgers equation*, Math. Nachr. 289 (2016), 2007–2018.
- [ESS20] M. Elin, D. Shoikhet, T. Sugawa, *Geometric properties of the nonlinear resolvent of holomorphic generators*, J. Math. Anal. Appl. 483 (2020).
- [FHS20] U. Franz, T. Hasebe, S. Schleißinger, *Monotone Increment Processes, Classical Markov Processes, and Loewner Chains*, Dissertationes Math. 552 (2020), 119 pp.
- [GKH20] I. Graham, H. Hamada, G. Kohr, *Loewner chains and nonlinear resolvents of the Carathéodory family on the unit ball in \mathbb{C}^n* . J. Math. Anal. Appl. 491 (2020), no. 1, 124289, 29 pp.
- [HS20] I. Hotta, S. Schleißinger, *Limits of radial multiple SLE and a Burgers-Loewner differential equation*, J. Theoret. Probab. 34 (2021), no. 2, 755–783.
- [Jek20] D. Jekel, *Operator-valued chordal Loewner chains and non-commutative probability*, J. Funct. Anal. 278 (2020), no. 10, 108452, 100 pp.
- [Koe87] W. Koepf, *Convex functions and the Nehari univalence criterion*, in: Laine, I., Sorvali, T., Rickman, S., Complex Analysis Joensuu 1987, Lecture Notes in Mathematics, vol 1351, Springer, Berlin, Heidelberg, 1987.
- [Maa92] H. Maassen, *Addition of freely independent random variables*, J. Funct. Anal. 106 (1992), no. 2, 409–438.
- [MS17] J. A. Mingo, R. Speicher, *Free Probability and Random Matrices*, Springer, Berlin, 2017.
- [NS06] A. Nica, R. Speicher, *Lectures on the Combinatorics of Free Probability*, London Mathematical Society, Lecture Note Series 335, Cambridge University Press, 2006.
- [Pom92] Ch. Pommerenke, *Boundary behaviour of conformal maps*, Grundlehren der Mathematischen Wissenschaften, vol. 299, Springer-Verlag, Berlin, 1992.
- [RS96] S. Reich, D. Shoikhet, *Generation theory for semigroups of holomorphic mappings in Banach spaces*, Abstr. Appl. Anal. 1 (1996), no. 1, 1–44.
- [RS97] S. Reich, D. Shoikhet, *Semigroups and generators on convex domains with the hyperbolic metric*, Atti della Accademia Nazionale dei Lincei. Classe di Scienze Fisiche, Matematiche e Naturali. Rendiconti Lincei. Matematica e Applicazioni, Serie 9, Vol. 8 (1997), no. 4, p. 231–250.
- [Rob36] M. S. Robertson, *On the theory of univalent functions*, Ann. Math. 37 (1936), no. 2, 374–408.
- [Sha93] J. H. Shapiro, *Composition Operators and Classical Function Theory*, Springer, 1993.
- [Stu11] E. Study, *Konforme Abbildung einfach-zusammenhängender Bereiche*, Volume 2 of Vorlesungen über ausgewählte Gegenstände der Geometrie, Leipzig, Berlin, B.G. Teubner, 1911–13.
- [Voi87] D. Voiculescu, *Multiplication of certain noncommuting random variables*, J. Operator Theory 18 (1987), 223–235.
- [Voi97] ———, *Free Probability Theory*, Fields Inst. Commun. 12, Amer. Math. Soc., 1997.
- [ZZ18] H. Zhang, M. Zinsmeister, *Local Analysis of Loewner Equation*, arXiv:1804.03410.