

# Difference-based covariance matrix estimate in time series nonparametric regression with applications to specification tests

Lujia Bai

Center for Statistical Science,  
Department of Industrial Engineering,  
Tsinghua University

Weichi Wu

Center for Statistical Science,  
Department of Industrial Engineering,  
Tsinghua University

## Abstract

Long-run covariance matrix estimation is the building block of time series inference problems. The corresponding difference-based estimator, which avoids detrending, has attracted considerable interest due to its robustness to both smooth and abrupt structural breaks and its competitive finite sample performance. However, existing methods mainly focus on estimators for the univariate process while their direct and multivariate extensions for most linear models are asymptotically biased. We propose a novel difference-based and debiased long-run covariance matrix estimator for functional linear models with time-varying regression coefficients, allowing time series non-stationarity, long-range dependence, state-heteroscedasticity and their mixtures. We apply the new estimator to i) the structural stability test, overcoming the notorious non-monotonic power phenomena caused by piecewise smooth alternatives for regression coefficients, and (ii) the nonparametric residual-based tests for long memory, improving the performance via the residual-free formula of the proposed estimator. The effectiveness of the proposed method is justified theoretically and demonstrated by superior performance in simulation studies, while its usefulness is elaborated by means of real data analysis.

**Keywords:** debias; difference-based; locally stationary; long-run variance; monotonic power; time-varying linear model;

## 1 Introduction

The long-run variance plays a central role in the statistical inference of time series linear models. Consider the following functional linear model (Zhou and Wu (2010)) for the time series observations  $(y_{i,n}, x_{i,n})$ ,  $1 \leq i \leq n$ ,

$$y_{i,n} = x_{i,n}^\top \beta_{i,n} + e_{i,n}, \quad \beta_{i,n} = \beta(i/n), \quad 1 \leq i \leq n, \quad (1.1)$$

---

<sup>1</sup>E-mail addresses: [blj20@mails.tsinghua.edu.cn](mailto:blj20@mails.tsinghua.edu.cn)(L.Bai), [wuweichi@mail.tsinghua.edu.cn](mailto:wuweichi@mail.tsinghua.edu.cn)(W.Wu)

where  $x_{i,n}$  is a  $p$ -dimensional covariate process whose first element is 1 and  $e_{i,n}$  is the error process, both of which can be non-stationary,  $(\beta_{i,n})$  are the time-varying regression coefficients and  $\beta(\cdot) : [0, 1] \rightarrow \mathbb{R}^p$  is the regression function. If  $\hat{\beta}_{i,n}$  is assumed to be constant or smoothly changing, it can be estimated by ordinary least squares or nonparametric methods (see [Chan and Zhang \(2010\)](#), [Li et al. \(2011\)](#), [Su et al. \(2019\)](#)) and the variation of such estimates is determined by the  $p$ -dimensional possibly time-varying *long-run covariance matrix* of  $(x_{i,n}e_{i,n})$ , see [\(2.1\)](#) for exact definition. Classic estimation of the long-run covariance matrix in linear models, see for instances [Newey and West \(1987\)](#) and [Andrews \(1991\)](#), requires consistent estimation of regression coefficients, which is difficult to achieve under structural changes. A prevalent approach to overcome the obstacle is the difference-based estimation, which is built on  $y_{i+m,n} - y_{i,n}$  where  $m$  is a diverging tuning parameter such that  $m = o(n)$  so that most  $y_{i+m,n} - y_{i,n}$  are approximately zero-mean under both smooth and abrupt structural changes. Most of the existing results assume stationarity, see for example, [Tecuatpetla-Gómez and Munk \(2017\)](#) and [Chan \(2022\)](#). Exceptions include [Dette and Wu \(2019\)](#), which allows non-stationary errors. To the best of the author's knowledge, the existing difference-based long-run variance estimators for time series data are designed for  $p = 1$ . When  $(e_{i,n})$  are *i.i.d.*, the difference-based estimators for variance has been studied by for example [Müller and Stadtmüller \(1987\)](#), [Hall et al. \(1990\)](#) and [Brown and Levine \(2007\)](#).

For  $p > 1$ , the long-run variance is referred as the long-run covariance matrix, whose estimation is much more involved, see [Jansson \(2002\)](#), [Hirukawa \(2021\)](#) and reviews therein. Most existing methods are plug-in methods that utilize estimated residuals which have been widely applied to the goodness of fit tests, tests of structural breaks ([Aue and Horváth \(2013\)](#), [Wu and Zhou \(2018\)](#), [Kao et al. \(2018\)](#)), detecting gradual changes ([Vogt and Dette \(2015\)](#)), simultaneous confidence bands for coefficient functions ([Zhou and Wu \(2010\)](#)), tests for long memory ([Beran et al. \(2013\)](#), [Bai and Wu \(2021\)](#)), etc. Since they crucially depend on the pre-estimation of regression coefficients, they are inconsistent under abrupt structural breaks, causing the notorious non-monotonic power when applied to structural stability tests. Moreover, most nonparametric specification tests involve both  $\hat{\beta}_{i,n}$  and the long-run covariance matrix, while  $\hat{\beta}_{i,n}$  is also used in the formula of the plug-in estimator for the long-run covariance matrix. Therefore, the plug-in estimate tends to sensitize those tests to the tuning parameters chosen for  $\hat{\beta}_{i,n}$ .

In this paper, we develop a new difference-based estimator of the time-varying long-run covariance matrix to overcome the aforementioned difficulties for functional linear models with non-stationary dependent errors. Our estimator has an extra debiased term compared to their univariate counterpart because the extension of the latter estimator to linear models with  $p > 1$  is asymptotically biased. We derive the consistency of the new estimator under time series nonstationarity allowing flexible predictor-error relationships in [Section 4](#). In particular, our difference-based estimator achieves faster convergence rates under certain moment conditions than that of the plug-in estimator in [Zhou and Wu \(2010\)](#). We then apply the new estimator to the structural stability tests and the long-memory tests, studying the theoretical and numerical performance in [Section 5](#) and [6](#), respectively. An empirical illustration using public health data can be found in [Section 7](#).

## 2 Preliminaries

In this paper, we consider a general form of non-stationary called local stationarity for the covariate and the error processes of (1.1), which has received substantial attention in the literature. We employ the definition of locally stationary processes based on Bernoulli shift processes, see Wu (2005) and Zhou and Wu (2010), while there are also many other formulations, see Dahlhaus (1997), Nason et al. (2000), and Dahlhaus et al. (2019) for a comprehensive review.

We start by introducing necessary notation that will be used in the rest of the paper. Define  $[a]$  as the largest integer smaller than  $a$ , and  $\lambda_{\min}(A)$  as the smallest eigenvalue of any symmetric squared matrix  $A$ . Let  $|\cdot|$  denote the absolute value for scalars and the Frobenius norm for vectors. Let  $\mathcal{F}_i = (\cdots, \eta_{i-1}, \eta_i)$ , where  $(\eta_i)_{i \in \mathbb{Z}}$  are *i.i.d.* random elements, and the couple process  $\mathcal{F}_{i, \{0\}} = (\cdots, \eta_{-1}, \eta'_0, \eta_1, \cdots, \eta_i)$ , where  $(\eta'_i)_{i \in \mathbb{Z}}$  is the *i.i.d.* copy of  $(\eta_i)_{i \in \mathbb{Z}}$ . We write  $f(\cdot) \in C^q[0, 1]$  if  $f(\cdot)$  has  $q$ -th order continuous derivative. Consider the data generating mechanism  $L(t, \mathcal{F}_i) \in \mathbb{R}^p$ , where  $L$  is a filter function such that  $L(t, \mathcal{F}_i)$  is well-defined. We say  $L(t, \mathcal{F}_0) \in \text{Lip}_s(I)$  if there exists a constant  $c > 0$  such that for any  $t_1, t_2 \in I, t_1 < t_2$ ,

$$\|L(t_1, \mathcal{F}_0) - L(t_2, \mathcal{F}_0)\|_s \leq c|t_1 - t_2|,$$

where  $\|X\|_s = (\mathbb{E}|X|^s)^{1/s}$ . To measure the dependence of time series, we adopt the physical dependence (see Wu (2005)). For  $L(t, \mathcal{F}_i)$  on interval  $I$ , the physical dependence in  $\mathcal{L}^r$  norm is defined by

$$\delta_r(L, k, I) = \sup_{t \in I} \|L(t, \mathcal{F}_k) - L(t, \mathcal{F}_{k, \{0\}})\|_r.$$

The following definition gives the definition of locally stationarity and short-range dependence.

**Definition 2.1.** *The process  $G(t, \mathcal{F}_i)$  is of  $r$ -order ( $r \geq 1$ ) short-range dependence on interval  $I$  if  $\sup_{t \in I} \|G(t, \mathcal{F}_0)\|_r < \infty$ ,  $\delta_r(H, k, I) = O(\chi^k)$ , and  $s$ -order ( $s \geq 2$ ) locally stationary on interval  $I$ , if  $G(t, \mathcal{F}_0) \in \text{Lip}_s(I)$ .*

In the functional linear model (1.1), we consider  $x_{i,n} = W(t_i, \mathcal{F}_i)$ ,  $e_{i,n} = H(t_i, \mathcal{F}_i)$ , where  $t_i = i/n$ ,  $W$  and  $H$  are measurable nonlinear filters mapping from  $[0, 1] \times \mathbb{R}^{\mathbb{Z}}$  to  $\mathbb{R}^p$  and  $(-\infty, 1] \times \mathbb{R}^{\mathbb{Z}}$  to  $\mathbb{R}$ , respectively. Permitting  $i$  to approach  $-\infty$  enables us to develop our proposed method under long memory, see Section 5.2 for detailed discussion. We further assume  $e_{i,n}$  has the following state-heteroscedastic form, i.e.,  $e_{i,n} = \tilde{H}(t_i, \mathcal{H}_i) \tilde{G}(t_i, \mathcal{G}_i)$  for  $-\infty < i \leq n$ , where the nonlinear filters  $\tilde{H}(t, \cdot)$  and  $\tilde{G}(t, \cdot)$  are  $\mathcal{H}_i$  and  $\mathcal{G}_i$  measurable functions for  $t \in (-\infty, 1]$ , and filtrations  $\mathcal{G}_i$  and  $\mathcal{H}_i$  are sub- $\sigma$ -fields of  $\mathcal{F}_i$ , independent of each other. Furthermore,  $\mathcal{H}_i \subset \sigma(x_{1,n}, \dots, x_{i,n})$  if  $1 \leq i \leq n$ , and for any  $t \in (-\infty, 1]$ ,  $\mathbb{E} \tilde{G}(t, \mathcal{G}_i) = 0$ . The above formulation admits the heteroscedastic errors considered in He and Zhu (2003) and Kulik and Wichelhaus (2012), where they assume  $\tilde{H}(t_i, \mathcal{H}_i)$  to have the form of  $s(x_{i,n})$  for some unknown smooth function  $s(\cdot)$ . Moreover,  $\tilde{G}(t_i, \mathcal{G}_i)$  allows the conditional heteroscedasticity as considered by assumption 1(b) of Cavaliere et al. (2017). Further, let  $U(t_i, \mathcal{F}_i) = W(t_i, \mathcal{F}_i)H(t_i, \mathcal{F}_i)$  such that  $x_{i,n}e_{i,n} = U(t_i, \mathcal{F}_i)$ .

The (time-varying) long-run covariance function for the functional linear model (1.1) is defined by

$$\Sigma(t) := \sum_{j=-\infty}^{\infty} \text{Cov} \{U(t, \mathcal{F}_0), U(t, \mathcal{F}_j)\}. \quad (2.1)$$

For stationary  $x_{i,n}$  and  $e_{i,n}$ ,  $\Sigma(t)$  will be time invariant. In this case for linear models with no structural changes, Zhou and Shao (2013) proposes a self-normalization method for statistical inference so that the estimation of the long-run covariance matrix can be avoided. However, their method relies crucially on strict stationarity. Therefore, for the statistical inference of non-stationary time series linear models, the accurate and robust estimation of  $\Sigma(t)$  is essential.

### 3 The debiased difference-based estimator

When  $p = 1$ , Dette and Wu (2019) proposes the following difference-based estimator  $\hat{\Sigma}(t)$  based on the difference of  $x_{i,n}y_{i,n}$  for the long run variance of  $e_{i,n}$ . Let  $Q_{k,m} = \sum_{i=k}^{k+m-1} x_{i,n}y_{i,n}$ , and for  $t \in [m/n, 1 - m/n]$ ,

$$\hat{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m\Delta_j\Delta_j^\top}{2} \omega(t, j), \quad \Delta_j = \frac{Q_{j-m+1,m} - Q_{j+1,m}}{m},$$

where for some bandwidth  $\tau_n$  and the kernel function  $K(\cdot)$  with support  $(-1, 1)$ ,

$$\omega(t, i) = K_{\tau_n}(t_i - t) / \sum_{i=1}^n K_{\tau_n}(t_i - t).$$

For  $t \in [0, m/n)$ ,  $\hat{\Sigma}(t) = \hat{\Sigma}(m/n)$  and for  $t \in (1 - m/n, 1]$ ,  $\hat{\Sigma}(t) = \hat{\Sigma}(1 - m/n)$ .

Unfortunately, in Appendix B.2.1, we find that the difference-based estimator  $\hat{\Sigma}(t)$  is asymptotically biased for  $p \geq 2$ , where we also show that  $\mathbb{E}(\hat{\Sigma}(t)) - \Sigma(t)$  can be uniformly approximated by the expectation of the following  $\Sigma^A(t)$ :

$$\Sigma^A(t) = \sum_{j=m}^{n-m} \frac{mA_{j,m}A_{j,m}^\top}{2} \omega(t, j), \quad \text{for } t \in [m/n, 1 - m/n],$$

where for  $j \geq m$ ,

$$A_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^j (x_{i,n}x_{i,n}^\top \beta(t_i) - x_{i+m,n}x_{i+m,n}^\top \beta(t_{i+m})),$$

and  $\Sigma^A(t) = \Sigma^A(m/n)$  for  $t \in [0, m/n)$ ,  $\Sigma^A(t) = \Sigma^A(1 - m/n)$  for  $t \in (1 - m/n, 1]$ . To utilize  $\Sigma^A(t)$ , we shall substitute  $\beta(\cdot)$  in  $A_{j,m}$  via an estimator without introducing additional tuning parameters. For this purpose, define  $\tilde{Y}_{i,m} = x_{i,n}y_{i,n} - x_{i+m,n}y_{i+m,n}$ ,  $\tilde{X}_{i,m} = x_{i,n}x_{i,n}^\top - x_{i+m,n}x_{i+m,n}^\top$ , and

$\tilde{E}_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^j (x_{i,n}e_{i,n} - x_{i+m,n}e_{i+m,n})$ . By the continuity of  $\beta(t)$ ,

$$\begin{aligned} \frac{1}{m} \sum_{i=j-m+1}^j \tilde{Y}_{i,m} &= A_{j,m} + \tilde{E}_{j,m} \approx \frac{1}{m} \sum_{i=j-m+1}^j \tilde{X}_{i,m} \beta(t_i) + \tilde{E}_{j,m} \\ &\approx \frac{1}{m} \beta(t_j) \sum_{i=j-m+1}^j \tilde{X}_{i,m} + \tilde{E}_{j,m}. \end{aligned} \quad (3.1)$$

Since  $\mathbb{E}\tilde{E}_{j,m} = 0$ , the random vector  $\sqrt{m}\tilde{E}_{j,m}$  is  $O_p(1)$ . Treating  $\sum_{i=j-m+1}^j \tilde{Y}_{i,m}/\sqrt{m}$  as the response variable,  $\sum_{i=j-m+1}^j \tilde{X}_{i,m}/\sqrt{m}$  as covariates and  $\sqrt{m}\tilde{E}_{j,m}$  as errors, (3.1) motivates us to approximate  $\beta(t)$  in  $A_{j,m}$  by

$$\check{\beta}(t) = \Omega^{-1}(t)\varpi(t), \quad (3.2)$$

where  $\Omega(t)$  and  $\varpi(t)$  are the smoothed versions of  $\hat{\Delta}_j/2 := \frac{1}{2m} \sum_{i=j-m+1}^j \tilde{X}_{i,m} \tilde{X}_{i,m}^\top$ , and  $\check{\Delta}_j/2 := \frac{1}{2m} \sum_{i=j-m+1}^j \tilde{X}_{i,m}^\top \tilde{Y}_{i,m}$ , i.e., for  $t \in [m/n, 1 - m/n]$ ,

$$\Omega(t) = \sum_{j=m}^{n-m} \hat{\Delta}_j \omega(t, j)/2, \quad \varpi(t) = \sum_{j=m}^{n-m} \check{\Delta}_j \omega(t, j)/2,$$

while for  $t \in [0, m/n)$ ,  $\Omega(t) = \Omega(m/n)$ ,  $\varpi(t) = \varpi(m/n)$  and for  $t \in (1 - m/n, 1]$ ,  $\Omega(t) = \Omega(1 - m/n)$ ,  $\varpi(t) = \varpi(1 - m/n)$ . The estimator  $\check{\beta}(t)$ , which is also based on difference series, is accurate except in the vicinity of abrupt changes. Fortunately, the effect of abrupt changes can be mitigated by the local averaging in the formula of  $A_{j,m}$ . Replacing  $\beta(t)$  by  $\check{\beta}(t)$  in  $A_{j,m}$  we obtain

$$\hat{A}_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^j (x_{i,n} x_{i,n}^\top \check{\beta}(t_i) - x_{i+m,n} x_{i+m,n}^\top \check{\beta}(t_{i+m})),$$

and the corresponding debiased difference-based estimator  $\hat{\Sigma}(t)$  for  $t \in [0, 1]$ :

$$\hat{\Sigma}(t) = \hat{\Sigma}(t) - \check{\Sigma}(t), \quad \text{where } \check{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m \hat{A}_{j,m} \hat{A}_{j,m}^\top}{2} \omega(t, j), \quad (3.3)$$

which is robust to structural breaks in regression coefficient functions due to differencing, and except for  $m$ ,  $\tau$  used for  $\hat{\Sigma}(t)$ , the correction  $\check{\Sigma}(t)$  does not involve additional tuning parameters. Thanks to its residual-free formula, we can obtain more accurate type I error rates for the nonparametric specification tests by using  $\hat{\Sigma}(t)$ , see Section 4.

## 4 Consistency under smooth structural changes

In this section, we discuss the uniform convergence of the debiased difference-based long-run covariance matrix estimator (3.3) for the functional linear model (1.1) under smooth structural changes and short-range dependence with locally stationary predictors and errors, which accommodates many null hypotheses of nonparametric specification tests. The convergence of  $\hat{\Sigma}(t)$  when both smooth and abrupt changes occur is deferred to Section 5.1, and the performance of  $\hat{\Sigma}(t)$  under long memory is postponed to Section 5.2.

**Assumption 4.1.** *Each coordinate of  $\beta(\cdot)$  lies in  $C^3[0, 1]$ .*

Assumption 4.1 imposes smooth structural change, i.e., the coefficient function  $\beta(\cdot)$  is smooth. In Section 5.1, we shall relax Assumption 4.1 to allow abrupt structural changes.

**Assumption 4.2.** *The value  $\lambda_{\min}(\Sigma(t))$  is bounded from 0 uniformly on  $[0, 1]$ , and each element of  $\Sigma(t) \in C^2[0, 1]$ .*

Assumption 4.2 guarantees the non-degeneracy of the long-run covariance matrix of the process of  $U(t, \mathcal{F}_i)$  and each component of  $\Sigma(t)$  is smooth, which are common in the analysis of functional linear models of locally stationary time series, see for example Zhou and Wu (2010). Let  $J(t, \mathcal{F}_0) = W(t, \mathcal{F}_0)W^\top(t, \mathcal{F}_0)$ . Define  $\mu_W(\cdot) = \mathbb{E}(W(t, \mathcal{F}_0))$ ,  $M(t) := \mathbb{E}(J(t, \mathcal{F}_0))$ ,  $t \in [0, 1]$ .

**Assumption 4.3.** *The functions  $M(\cdot) \in C^1[0, 1]$ ,  $\mu_W(\cdot) \in C^1[0, 1]$ , and  $\inf_{t \in [0, 1]} M(t) > 0$ . The covariate process  $W(t, \mathcal{F}_i)$  is of 4-order locally stationarity and  $16\kappa$ -order short-range dependence on  $[0, 1]$  for some  $\kappa \geq 1$ .*

Assumption 4.3 requires that  $M(t)$  is non-degenerate, implies that  $\sup_{t \in [0, 1]} \|J(t, \mathcal{F}_0)\|_{8\kappa} < \infty$ ,  $\delta_{8\kappa}(J, k) = O(\chi_1^k)$  for some  $\chi_1 \in (0, 1)$ , and ensures that each element of  $(x_{i,n}x_{i,n}^\top)$  is 2-order locally stationary.

The following assumption ensures the invertibility of  $\Omega(t)$  in (3.2).

**Assumption 4.4.** *The covariance function  $\text{Cov}(J(t, \mathcal{F}_0), J(t, \mathcal{F}_0)) \in C^2[0, 1]$  and its smallest eigenvalue is strictly positive on  $[0, 1]$ .*

**Assumption 4.5.** *The process  $H(\cdot, \mathcal{F}_i)$  satisfies  $16\kappa$ -order of short-range dependence and 4-order of locally stationarity on  $[0, 1]$ .*

Assumptions 4.3 and 4.5 ensure that  $U(t, \mathcal{F}_i)$  is of 2-order local stationarity and  $8\kappa$ -order of short-range dependence. The existence of  $16\kappa_{th}$  moments of covariates and errors are assumed for technical convenience, and it is satisfied by sub-exponential random variables. We conjecture that it can be relaxed by substantially more involved mathematical arguments, see the simulation study of Bai and Wu (2021). Assumption 4.4 is mild, and it excludes the scenario in which all of the time series covariates reduce to deterministic smooth trends; in this case, we recommend using the direct multivariate extension of the difference-based long-run covariance estimator of Dette and Wu (2019), namely  $\hat{\Sigma}(t)$ .

Let  $\mathcal{I} = [\gamma_n, 1 - \gamma_n] \subset (0, 1)$ , where  $\gamma_n = \tau_n + (m + 1)/n$ . The following theorem justifies the uniform consistency of  $\hat{\Sigma}(t)$  over  $\mathcal{I}$ .

**Theorem 4.1.** *Under Assumptions 4.1, 4.2, 4.3, 4.4 and 4.5 with constant  $\kappa \geq 1$ , suppose  $m = O(n^{1/3})$ ,  $m \rightarrow \infty$ ,  $\tau_n \rightarrow 0$ ,  $n\tau_n^3 \rightarrow \infty$ ,  $m/(n\tau_n^2) \rightarrow 0$ ,  $\tau_n^{2-1/\kappa} \sqrt{m} \rightarrow 0$ , we have*

$$\sup_{t \in \mathcal{I}} \left| \hat{\Sigma}(t) - \Sigma(t) \right| = O_{\mathbb{P}} \left( \frac{m^{1/2}}{n^{1/2} \tau_n^{1/2+1/\kappa}} + \frac{1}{m} + m^{1/2} \tau_n^{2-1/\kappa} \right).$$

The above equation and the bandwidth conditions for Theorem 4.1 ensure the uniform consistency of  $\hat{\Sigma}(t)$ . We compare our results with Zhou and Wu (2010) which proposes a plug-in estimator of long-run covariance matrix estimator using nonparametric residuals. According to Theorem 5 in Zhou and Wu (2010), their best approximation rate can be close to but not faster than  $n^{-1/4}$ . In contrast, our uniform rate in Theorem 4.1 is  $n^{-4/15+1/(5\kappa)}$  by taking  $m \asymp n^{4/15}$ ,  $\tau_n \asymp n^{-1/5}$ , which is better when  $\kappa$  is sufficiently large.

## 5 Applications

In this section, we elaborate two applications in which the long-run covariance matrix estimators are important, i.e., structural change detection and tests for long memory. We show that with the difference-based long-run variance-covariance estimate (3.3) both tests are asymptotically correct and powerful. Moreover, the application of the difference-based long-run variance-covariance estimate (3.3) can effectively and consistently yield higher power for all the tests considered compared to the residual-based testing procedures, see Section 6 for simulation evidence.

### 5.1 Structural change detection with monotonic power

We are interested in the structural changes in the stochastic linear regression model

$$y_{i,n} = x_{i,n}^{\top} \beta_{i,n} + e_{i,n}, \quad 1 \leq i \leq n,$$

where  $\{x_{i,n}\}_{i=1}^n$  is the  $p$ -dimensional covariate process and  $\{e_{i,n}\}_{i=1}^n$  is the error process. The test for structural changes of  $(\beta_{i,n})$  considers the hypothesis  $H_0 : \beta_{1,n} = \beta_{2,n} = \dots = \beta_{n,n}$ . Allowing general non-stationarity in the covariates and the errors, Wu and Zhou (2018) proposes to use the test statistic

$$T_n = \max_{1 \leq j \leq n} \left| \sum_{i=1}^j \hat{e}_{i,n} x_{i,n} / \sqrt{n} \right|, \quad \text{where } \hat{e}_{i,n} = y_{i,n} - x_{i,n}^{\top} \hat{\beta}_n,$$

where  $\hat{\beta}_n = \operatorname{argmin}_{\beta} \sum_{i=1}^n (y_{i,n} - x_{i,n}^{\top} \beta)^2$ . For the alternative hypothesis, we consider

$$y_{i,n} = x_{i,n}^{\top} \beta(i/n) + e_{i,n}, \quad \beta(t) = \beta + L_n g(t), \quad (5.1)$$

where  $\beta, g(t) \in \mathbb{R}^p$ ,  $g(0) = 0$ ,  $L_n$  is a positive real sequence,  $\beta(t)$  has potential abrupt changes, i.e.,  $\beta(t) = \sum_{j=0}^{q_n} b_j(t) \mathbb{I}_{\{a_j \leq t < a_{j+1}\}}$ ,  $0 = a_0 < a_1 < \dots < a_{q_n} < a_{q_n+1} = 1$ ,  $b_j(t) \in \mathcal{C}^3[a_j, a_{j+1}]$ ,  $\sup_{0 \leq j \leq q_n} |b_j(a_j) - b_j(a_j^-)| < \infty$ , and  $q_n$  is the number of abrupt changes. Wu and Zhou (2018) proposes a general bootstrap

statistics according to (5.2), which relies on the residual  $\hat{\varepsilon}_{i,n} = y_{i,n} - x_{i,n}^\top \hat{\beta}_n$ . Their test can be applied to piecewise locally stationary covariates and errors, and is unified for testing structural changes in general  $M$  estimation. They show their power approaches 1 under local alternatives  $L_n = n^{-1/2}$  or  $L_n = o(1)$ ,  $\sqrt{n}L_n \rightarrow \infty$ . However, if  $L_n$  does not vanish,  $\hat{\beta}_n$  is not consistent and the power of the test in Wu and Zhou (2018) in this case is not theoretically guaranteed. If we focus on the locally stationary process which is quite general and on least squares regression which is arguably the most widely applied M-estimator in practice, we could improve the power of the test for structural changes via our proposed long-run covariance matrix estimator  $\hat{\Sigma}(t)$  and an alternative bootstrap procedure to Wu and Zhou (2018). For this purpose, we define

$$F_r = \max_{m \leq i \leq n-m+1} |\Psi_{i,m}^{(r)} - \hat{\Lambda}(i/n) \hat{\Lambda}^{-1}(1) \Psi_{n-m+1,m}^{(r)}|,$$

where  $\Psi_{i,m}^{(r)} = n^{-1/2} \sum_{j=1}^i \hat{\Sigma}^{1/2}(t_j) R_j^{(r)}$ ,  $\hat{\Lambda}(i/n) = \sum_{j=1}^i x_{j,n} x_{j,n}^\top / n$ ,  $\{R_j^{(r)}\}_{j=1}^n$  are the *i.i.d.* standard norm random variables independent of data and are independently generated in the  $r$ th bootstrap iteration, and  $\hat{\Sigma}(t)$  is the proposed difference-based estimator of  $\Sigma(t)$ . Let  $B$  denote the number of bootstrap iterations. Let  $F_{(1)} \leq F_{(2)} \leq \dots \leq F_{(B)}$  be the order statistics of  $(F_r)$ . We reject the structural stability test at the significance level of  $\alpha$  if  $T_n$  is greater than the  $F_{[(1-\alpha)B]}$ . We proceed to relax Assumption 4.1 in Theorem 4.1 allowing for the possible presence of abrupt changes and discuss the property of the bootstrap procedure. Let  $\Lambda(t) = \int_0^t M(s) ds$ .

**Theorem 5.1.** *Assume  $q_n = o(\min\{n\tau_n^3/m, (n\tau_n)^2/m\})$  under the alternative hypothesis (5.1). Then, under the Assumptions 4.2, 4.3, 4.4 and 4.5 with constant  $\kappa \geq 1$ , and the bandwidth conditions  $m = O(n^{1/3})$ ,  $m \rightarrow \infty$ ,  $\tau_n \rightarrow 0$ ,  $n\tau_n^3 \rightarrow \infty$ ,  $m/(n\tau_n^2) \rightarrow 0$ ,  $\tau_n^{2-1/\kappa} \sqrt{m} \rightarrow 0$ , we have*

$$\sup_{t \in \mathcal{I}} |\hat{\Sigma}(t) - \Sigma(t)| = o_{\mathbb{P}}(1),$$

and as a result there exists a  $p$ -dimensional zero-mean Gaussian process  $Z(t)$  with covariance function  $\gamma(t, s) = \int_0^{\min(t,s)} \Sigma(r) dr$  such that

$$F_r \Rightarrow \sup_{t \in (0,1]} |G(t)| := \sup_{t \in (0,1]} |Z(t) - \Lambda(t) \Lambda^{-1}(1) Z(1)|.$$

Theorem 5.1 shows that our difference-based estimator  $\hat{\Sigma}(t)$  is consistent under the alternatives of both smooth and abrupt changes. Taking  $m \asymp n^{4/15}$ ,  $\tau_n \asymp n^{-1/5}$ , it follows that  $\min\{n\tau_n^3/m, (n\tau_n)^2/m\} \asymp n^{2/15}$  which allows the number of abrupt jumps diverges to infinity as  $n \rightarrow \infty$ . The following proposition gives the limiting distribution of the test statistic  $T_n$  under the null hypothesis and ensures the monotonic power of the bootstrap procedure under  $H_A : \beta(t) = \beta + L_n g(t)$ . Define

$$\Lambda(t) = \int_0^t M(s) ds, \quad \Lambda(s, g(\cdot)) = \int_0^s M(r) g(r) dr,$$

and  $F(t, g(\cdot)) = \Lambda(t, g(\cdot)) - \Lambda(t) \Lambda^{-1}(1) \Lambda(1, g(\cdot))$ .



**Proposition 5.1.** (i) Under the conditions of [Theorem 4.1](#) and the null hypothesis of no structural changes, we have

$$T_n \Rightarrow \sup_{t \in (0,1]} |G(t)|, \quad (5.2)$$

where  $G(t)$  is as defined in [Theorem 5.1](#).

(ii) Under the conditions of [Theorem 5.1](#) and the alternative hypothesis [\(5.1\)](#) with  $L_n = O(1)$ ,  $n^{1/2}L_n \rightarrow \infty$ , we have  $T_n \rightarrow \infty$  in probability at the rate  $\sqrt{n}L_n$ , and

$$|\mathbb{P}(T_n \geq \hat{q}_{1-\alpha}) - \mathbb{P}\left(\sup_{t \in [0,1]} |G(t) + n^{1/2}L_n F(t, g(\cdot))| \geq \hat{q}_{1-\alpha}\right)| = o(1),$$

where  $\hat{q}_{1-\alpha}$  is the bootstrap critical value of  $F_r$  at the significance level  $\alpha$ .

[Proposition 5.1](#) shows that  $T_n$  is of order  $\max(1, \sqrt{n}L_n)$  under null hypothesis. [Theorem 5.1](#) and [Proposition 5.1](#) guarantee the asymptotic correctness of the bootstrap procedure with the difference-based estimator and that its asymptotic power approaches 1 under the fixed alternative. Meanwhile, the bootstrap procedure with the difference-based estimator can detect the local alternatives at the parametric rate  $\sqrt{n}$  in the sense that if  $L_n = n^{-1/2}$ ,  $T_n \Rightarrow \sup_{t \in [0,1]} |G(t) + F(t, g(\cdot))|$ . Notably,  $F_r$  shares the same limiting distribution as the limiting distribution of  $T_n$  under the null hypothesis, while [Proposition B.1](#) of [Wu and Zhou \(2018\)](#) shows that with the ordinary least squares residuals, the magnitude of  $F_r$  using their bootstrap procedure is  $\sqrt{m} \max(L_n, \log^2 n / \sqrt{n}) \log n$  when  $\sqrt{m}L_n \rightarrow \infty$ ,  $L_n \rightarrow 0$ , implying power loss due to the divergence of  $F_r$ . Therefore, the bootstrap procedure equipped with the difference-based estimator will be more powerful than that with ordinary least squares residuals and overcome non-monotonic power caused by the inflation of the bootstrap statistics in [Wu and Zhou \(2018\)](#). In earlier work on the remedy of the non-monotonic power for the test of smooth structural changes, [Juhl and Xiao \(2009\)](#) proposes to estimate the long-run variance via plugging in the nonparametric residuals for  $p = 1$ . However, the improvement in the power of their approach does not carry over in the presence of abrupt structure changes.

## 5.2 Testing for long memory

In this section, we consider the functional linear model

$$y_{i,n} = x_{i,n}^\top \beta(t_i) + (1 - \mathcal{B})^d e_{i,n}, \quad i = 1, 2, \dots, n,$$

where  $\mathcal{B}$  is the lag operator,  $d \in [0, 1/2)$  is the long-memory parameter. When  $d = 0$ , the error process of the model is locally stationary and short-range dependent. We are interested in the following hypothesis testing problem

$$H_0 : d = 0 \quad \text{versus} \quad H_A : 0 < d < 1/2. \quad (5.3)$$

The rejection of  $H_0$  implies that the short-memory linear model is inadequate for the data and long-range dependence should be considered. [Bai and Wu \(2021\)](#) proposes to test  $H_0$  using the jackknife corrected nonparametric residuals. They obtain the local linear estimate of  $\beta(\cdot)$ , i.e.,

$$(\hat{\beta}_{b_n}(t), \hat{\beta}'_{b_n}(t)) = \arg \min_{\eta_0, \eta_1 \in \mathbb{R}^p} \sum_{i=1}^n \{y_{i,n} - x_{i,n}^\top \eta_0 - x_{i,n}^\top \eta_1 (t_i - t)\}^2 K_{b_n}(t_i - t), \quad (5.4)$$

where  $K(\cdot)$  is a kernel function with finite support  $(-1, 1)$ ,  $b_n$  is a bandwidth and  $K_{b_n}(\cdot) = K(\cdot/b_n)$ . Then, they consider the jackknife estimator  $\tilde{\beta}_{b_n}(t) = 2\hat{\beta}_{b_n/\sqrt{2}}(t) - \hat{\beta}_{b_n}(t)$  of which the asymptotic bias terms involving  $\beta''(\cdot)$  in the formula of  $\hat{\beta}_{b_n}$  and  $\hat{\beta}_{b_n/\sqrt{2}}$  are canceled. Let  $K^*(\cdot)$  denote the jackknife equivalent kernel  $2\sqrt{2}K(\sqrt{2}x) - K(x)$ . For the sake of simplicity, we write  $n'$  short for  $\lfloor nb_n \rfloor$ . Define the nonparametric residuals and their partial sum as  $\tilde{e}_{i,n} = y_{i,n} - x_{i,n}^\top \tilde{\beta}_{b_n}(t_i)$  and  $\tilde{S}_{r,n} = \sum_{i=n'+1}^r \tilde{e}_{i,n}$ ,  $r = n'+1, \dots, n-n'$ , respectively. The KPSS, R/S, V/S and K/S-type test statistics of [Bai and Wu \(2021\)](#) are

1. KPSS-type statistic  $K_n = \frac{1}{n(n-2n')} \sum_{r=n'+1}^{n-n'} (\tilde{S}_{r,n})^2$ .
2. R/S-type statistic  $Q_n = \max_{n'+1 \leq k \leq n-n'} \tilde{S}_{k,n} - \min_{n'+1 \leq k \leq n-n'} \tilde{S}_{k,n}$ .
3. V/S-type statistic  $M_n = \frac{1}{n(n-2n')} \left\{ \sum_{k=n'+1}^{n-n'} \tilde{S}_{k,n}^2 - \frac{1}{n-2n'} \left( \sum_{k=n'+1}^{n-n'} \tilde{S}_{k,n} \right)^2 \right\}$ .
4. K/S-type statistic  $G_n = \max_{n'+1 \leq k \leq n-n'} |\tilde{S}_{k,n}|$ .

[Bai and Wu \(2021\)](#) proposes to implement the above tests via the following bootstrap-assisted procedure. Define  $\hat{M}(t) = \frac{1}{n\eta_n} \sum_{i=1}^n x_{i,n} x_{i,n}^\top K_{\eta_n}(t_i - t^*)$ , where  $t^* = \max\{\eta_n, \min(t, 1 - \eta_n)\}$  for some bandwidth  $\eta_n \rightarrow 0$ ,  $n\eta_n^2 \rightarrow \infty$ . Let  $\hat{\Sigma}^*(\cdot)$  be any consistent long-run covariance matrix estimator satisfying the regularity condition 5.1 in their paper, and  $\hat{\sigma}_H^{*2}(t) = (\hat{\Sigma}^*(t))_{1,1}$ . Generate  $B$  *i.i.d.* copies of  $N(0, I_p)$  vectors  $V_i^{(r)} = (V_{i,1}^{(r)}, \dots, V_{i,p}^{(r)})^\top$ ,  $1 \leq r \leq B$ , and for each  $r$  calculate

$$\tilde{G}_k^{(r)} = - \sum_{j=1}^n \left( \frac{1}{nb_n} \sum_{i=n'+1}^k x_{i,n}^\top \hat{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \hat{\Sigma}^{*,1/2}(t_j) V_j^{(r)} + \sum_{i=n'+1}^k \hat{\sigma}_H^*(t_i) V_{i,1}^{(r)}$$

as well as the bootstrap statistics:  $\tilde{K}_n^{(r)}$ ,  $\tilde{RS}_n^{(r)}$ ,  $\tilde{VS}_n^{(r)}$  and  $\tilde{KS}_n^{(r)}$  which can be obtained by substituting  $\tilde{S}_{k,n}$  in the corresponding statistics by  $\tilde{G}_k^{(r)}$ . Let  $\tilde{K}_{n,(1)} \leq \tilde{K}_{n,(2)} \leq \dots \leq \tilde{K}_{n,(B)}$  be the ordered statistics of  $\tilde{K}_n^{(r)}$ ,  $r = 1, 2, \dots, B$ , and  $B^* = \max\{r : \tilde{K}_{n,(r)} \leq K_n\}$ . Then the  $p$ -value of the KPSS-type test is  $1 - B^*/B$ , and the  $p$ -values of R/S, V/S, and K/S-type tests can be obtained similarly. Given a nominal level  $\alpha$ , if the  $p$ -value is smaller than  $\alpha$ , we reject the null hypothesis of short memory.

In this paper we propose to set  $\hat{\Sigma}^*(t) = \hat{\Sigma}(t)$ . With the new difference-based long-run covariance matrix estimator, our testing procedure will be more robust than that using the plug-in estimator for  $\hat{\Sigma}^*(t)$ , since for the latter procedure, the parameter  $b_n$  in (5.4) for the estimation of the regression coefficients will additionally affect the estimate of the long-run covariance matrix through the nonparametric residuals  $\tilde{e}_{i,n}$  as well as the selection of  $m$  and  $\tau_n$ , as indicated by the discussion of Theorem 5 in [Zhou and Wu \(2010\)](#) that  $m$  and  $\tau_n$  should be chosen from an interval determined (implicitly) by  $b_n$ . Moreover, adopting  $\hat{\Sigma}(t)$  leads to more accurate type-I error control due to the faster convergence rate, see the discussion below

**Theorem 4.1.**

In the following, we show the validity of  $\hat{\Sigma}(t)$  via studying the asymptotic behavior of  $\hat{\Sigma}(t)$  under the fixed and local alternatives for the testing problem (5.3), which is essential for the consistency of the aforementioned bootstrap tests.

**Assumption 5.1.** *Assumption 4.5 holds over  $(-\infty, 1]$ ,  $H(t, \mathcal{F}_i)$  is of  $2\kappa$ -order locally stationarity on  $(-\infty, 1]$ , and its long-run variance function*

$$\sigma_H^2(t) := \sum_{k=-\infty}^{\infty} \text{Cov}(H(t, \mathcal{F}_0), H(t, \mathcal{F}_k)), \quad t \in (-\infty, 1],$$

satisfies that  $\inf_{t \in (-\infty, 1]} \sigma_H^2(t) > 0$ ,  $\sup_{t \in (-\infty, 1]} \sigma_H^2(t) < \infty$ , and  $\sigma_H^2(\cdot)$  is twice continuous differentiable on  $[0, 1]$ .

**Theorem 5.2.** *Under Assumptions 4.1, 4.3, 4.4, and 5.1, assuming  $m\tau_n^{3/2}/\log n \rightarrow \infty$ ,  $\tau_n \rightarrow 0$ ,  $n\tau_n^3 \rightarrow \infty$ ,  $m/(n\tau_n^2) \rightarrow 0$ ,  $\tau_n^{2-1/\kappa}\sqrt{m} \rightarrow 0$ ,  $m = O(n^{1/3})$ ,  $\kappa \geq \max\{\frac{4}{1/2-d}, \frac{2}{3d}, 4\}$ , it follows that under  $H_A$*

$$\sup_{t \in \mathcal{I}} \left| m^{-2d} \hat{\Sigma}(t) - \kappa_2(d) \sigma_H^2(t) \mu_W(t) \mu_W^\top(t) \right| = o_{\mathbb{P}}(1),$$

where  $\kappa_2(d) = \Gamma^{-2}(d+1) \int_0^\infty (t^d - (t-1)_+^d)(2t^d - (t-1)_+^d - (t+1)^d) dt$ .

Theorem 5.2 shows that the proposed difference-based estimator  $\hat{\Sigma}(t)$  in (3.3) inflates at the rate of  $m^{2d}$  under long-range dependence with parameter  $d > 0$ , while its limit (normalized by  $m^{2d}$ ) depends on the  $\mu_W(t)$  along with the long-run variance of  $e_{i,n}$  and the long-memory parameter  $d$ . The long-memory parameter  $d$  also affects the theoretical properties of the long-run variance-covariance estimate through the moment condition of  $\kappa$ . The exact convergence rate is displayed in Step 6 of the proof of Theorem 5.2 in supplement due to page limit. In the following Theorem 5.3, we investigate the performance of the estimator  $\hat{\Sigma}(t)$  under the local alternatives  $d_n = c/\log n$  for some constant  $c > 0$ . For this purpose, we define the long-run cross covariance vector between the locally stationary processes  $(U(t, \mathcal{F}_i))$  and  $(H(t, \mathcal{F}_j))$ .

**Definition 5.1.** *Define the long-run cross-covariance vector  $s_{UH}(t) \in \mathbb{R}^p$  by*

$$s_{UH}(t) = \sum_{j=-\infty}^{\infty} \text{Cov}(U(t, \mathcal{F}_0), H(t, \mathcal{F}_j)), \quad t \in [0, 1].$$

For given constants  $c > 0$  and  $\alpha_1 \in (0, 1)$ , define for  $0 \leq t \leq 1$ , the symmetric matrix

$$\check{\Sigma}(t) := \Sigma(t) + (e^{c\alpha_1} - 1)^2 \sigma_H^2(t) \mu_W(t) \mu_W^\top(t) + (e^{c\alpha_1} - 1)(s_{UH}(t) \mu_W^\top(t) + \mu_W(t) s_{UH}^\top(t)).$$

The following Assumption 5.2 guarantees that  $\check{\Sigma}(t)$  is smooth and non-degenerate.

**Assumption 5.2.**  *$\check{\Sigma}(\cdot) \in C^2[0, 1]$ , and  $\lambda_{\min}(\check{\Sigma}(t))$  is bounded above 0 on  $[0, 1]$ .*

Since  $\Sigma(t)$  and  $(e^{c\alpha} - 1)^2 \sigma_H^2(t) \mu_W(t) \mu_W^\top(t)$  are positive definite, by Weyl's inequality Assumption 5.2 is satisfied for sufficiently small positive  $c$ .

**Theorem 5.3.** *Let Assumptions 4.1, 4.3, 4.4, 5.1 and 5.2 be satisfied. If  $m\tau_n^{3/2} \rightarrow \infty$ ,  $\tau_n \rightarrow 0$ ,  $n\tau_n^3 \rightarrow \infty$ ,  $m/(n\tau_n^2) \rightarrow 0$ ,  $\tau_n^{2-1/\kappa} \sqrt{m} \rightarrow 0$ ,  $m = \lfloor n^{\alpha_1} \rfloor$ ,  $\alpha_1 \in (0, 1/3)$ , we have*

$$\sup_{t \in \mathcal{I}} \left| \hat{\Sigma}(t) - \check{\Sigma}(t) \right| = o_{\mathbb{P}}(1).$$

Theorem 5.2 and Theorem 5.3 lead to the desired limiting distribution of bootstrap statistics under the fixed and local alternatives achieving satisfactory power performance in finite samples, see Bai and Wu (2021) for theoretical justification and numerical evidence. In finite samples, we demonstrate that long memory tests with difference-based long-run covariance matrix estimates can achieve sizes closer to the nominal level and are more powerful than their counterparts using plug-in estimates, see Section 6.

## 6 Simulation

We investigate the finite sample power performance of the tests introduced in Section 5 and compare the results using difference-based estimator (3.3) with the results equipped with existing long-run variance estimators. Let  $(\varepsilon_l)_{l \in \mathbb{Z}}, (\zeta_l)_{l \in \mathbb{Z}}, (\eta_l)_{l \in \mathbb{Z}}$  be *i.i.d.*  $N(0, 1)$ ,  $\vartheta_i = (\eta_i + \varepsilon_i)/2$ , and consider the filtrations

$$\mathcal{F}_j = (\cdots, \zeta_{j-1}, \zeta_j), \quad \mathcal{G}_j = (\cdots, \varepsilon_{j-1}, \varepsilon_j), \quad \mathcal{H}_j = (\cdots, \varepsilon_j, \eta_j), \quad j = -\infty, \dots, n.$$

### 6.1 Testing for structural changes

We generate the locally stationary process  $x_{i,n,1}$  from  $G_1(t, \mathcal{H}_i) = \sum_{j=0}^{\infty} (0.5 - 0.5t)^j \vartheta_{i-j}$ , the locally stationary process  $x_{i,n,2}$  from  $G_2(t, \mathcal{H}_i) = \sum_{j=0}^{\infty} (0.25 + 0.5(t - 0.5)^2)^j \varepsilon_{i-j}$ , and the locally stationary process  $u_{i,n}$  from  $G(t, \mathcal{F}_i) = 0.65 \cos(2\pi t)G(t, \mathcal{F}_{i-1}) + \zeta_i$ . We consider the following heteroscedastic linear regression model:

$$y_{i,n} = 1 + m_{i,n} + x_{i,n,1} + x_{i,n,2} + e_{i,n}, \quad e_{i,n} = (1 + 0.1x_{i,n,1})u_{i,n},$$

where the function  $m_{i,n}$  includes the following scenarios:

CP1:  $m_{i,n} = 2\delta \sin(2\pi t_i)x_{i,n,1}1(0.5 \leq t_i \leq 1)$ .

CP2:  $m_{i,n} = \delta \sin(2\pi t_i)1(0 \leq t_i \leq 0.4) + \delta x_{i,n,1}1(0.7 \leq t_i \leq 1)/2$ .

CP4:  $m_{i,n} = 1.5\delta \sin(2\pi t)1(0 \leq t \leq 0.2 \text{ or } 0.4 \leq t \leq 0.6 \text{ or } 0.8 \leq t \leq 1)$ .

We conduct our simulation with sample size 300. As shown in Figure 6.1, when there are 4 change points, the block bootstrap tests based on ordinary least squares residuals (Wu and Zhou (2018)) suffer from low and non-monotonic power that can not approach 1. By contrast, the newly proposed difference-based long-run covariance matrix estimator enhances the simulated power significantly and addresses the non-monotonic power issue by taking the difference. To further illustrate the role of long-run covariance matrix estimators in bootstrap tests of structural breaks, we investigate the estimation accuracy of both




Figure 6.1: The power performance of gradient-based structural change point tests as  $\delta$  increases from 0 to 1 with sample size  $n = 300$  and simulation times 2000, under three scenarios CP1(blue), CP2(orange), CP4(red). Dotted curves denote simulated rejection rates of bootstrap tests using blocks of ordinary least squares residuals, and solid curves denote simulated rejection rates of those equipped with the difference-based long-run covariance matrix estimator. Left panel: nominal size 0.05; Right panel: nominal size 0.1.

long-run covariance matrix estimators and find that our proposed estimator halves the empirical mean square error in the presence of change points, see [Appendix A](#) of the supplement for extra simulation results.

## 6.2 Testing for long-range dependence

Consider the following heteroscedastic functional linear model,

$$y_{i,n} = \beta_1(t_i) + \beta_2(t_i)x_{i,n} + (1 - \mathcal{B})^d e_{i,n}, \quad i = 1, \dots, n, \quad (6.1)$$

where  $\mathcal{B}$  is the lag operator,  $\beta_1(t) = 4 \sin(\pi t)$ ,  $\beta_2(t) = 4 \exp\{-2(t - 0.5)^2\}$ ,  $x_{i,n} = W(t_i, \mathcal{F}_i)$ , and  $e_{j,n} = H(t_j, \mathcal{F}_j, \mathcal{G}_j)$ , where

$$H(t, \mathcal{F}_i, \mathcal{G}_i) = B(t, \mathcal{G}_i) (1 + W^2(t, \mathcal{F}_i))^{1/2},$$

where  $W(t, \mathcal{F}_i) = (0.1 + 0.1 \cos(2\pi t))W(t, \mathcal{F}_{i-1}) + 0.2\zeta_i + 0.7(t - 0.5)^2$ , and  $B(t, \mathcal{G}_i) = (0.3 - 0.4(t - 0.5)^2)B(t, \mathcal{G}_{i-1}) + 0.8\varepsilon_i$ . [Figure 6.2](#) compares the power performance of the KPSS, V/S, R/S, and K/S tests using difference-based and plug-in long-run covariance matrix estimators for model (6.1) of sample size 1500 with nominal level 0.05 (left) and 0.1 (right) with long memory parameter  $d$  increasing from 0 to 0.5. The difference-based long-run covariance matrix estimator leads to consistent improvement for the power of KPSS, V/S, R/S, and K/S tests against  $0 < d \leq 1/2$  in finite samples. Notably, equipped with the difference-based estimator the simulated power of K/S, R/S, and KPSS tests can reach 1 with the sample size 1500 as  $d$  increases to 0.5, while using the plug-in long-run covariance matrix estimator the power is much lower and stays far below 1, except for the V/S test.




Figure 6.2: Power performance of KPSS(orange), K/S(blue), R/S(red), and V/S(green) tests with sample size 1500. Dotted curves denote simulated rejection rates of tests using the plug-in method, while solid curves denote simulated rejection rates of tests using the difference-based method. Left panel: nominal size 0.05; Right panel: nominal size 0.1.

## 7 Data analysis

We apply our newly proposed long-run covariance matrix estimator (3.3) to the analysis of Hong Kong hospital data, including structural change detection and tests for long memory. The data set consists of daily hospital admissions in Hong Kong as well as daily measurements of pollutants between January 1, 1994, and December 31, 1995, whose sample size is 730. Consider the functional linear model for this data set, i.e.,

$$y_{i,n} = \beta_1(t_i) + \sum_{p=2}^4 \beta_p(t_i)x_{i,p,n} + \varepsilon_{i,n}, \quad i = 1, \dots, n,$$

where  $(y_{i,n})$  is the series of daily total number of hospital admissions of circulation and respiration and  $(x_{i,p,n})$ ,  $p = 2, 3, 4$ , are the series of daily levels (in micrograms per cubic meter) of  $\text{SO}_2$ ,  $\text{NO}_2$  and dust, respectively.

As illustrated in Section 5 of Wu and Zhou (2018), it is of practical concern to test whether  $\beta(\cdot) := (\beta_p(\cdot), 1 \leq p \leq 4)^\top$  is a constant vector. The test for structural changes equipped with the difference-based estimator yields  $p$ -value 0.0318, which rejects the null hypothesis of no structural change, while the test procedure proposed in Wu and Zhou (2018) based on ordinary least squares residuals yields  $p$ -value greater than 0.1. The different testing results can be attributed to the power loss of Wu and Zhou (2018) under structural change, as shown in Figure 6.1.

We then consider the test for the long memory of  $\varepsilon_{i,n}$ . Bai and Wu (2021) performs long-memory tests on each covariate process and concludes that they are short-range dependent. Therefore, we could apply the tests introduced in Section 5.2 to this data, and compare the  $p$ -values of the tests equipped with the difference-based estimator (3.3) and with the plug-in estimator of Zhou and Wu (2010), respectively. The bandwidth  $b_n$  in Section 5.2 are selected by the GCV method advocated by Zhou and Wu (2010) and Bai and Wu (2021). As in Table 7.1, the  $p$ -values of four types of tests for long memory based on

Table 7.1:  $p$ -values of tests for long memory.

	KPSS	RS	VS	KS		KPSS	RS	VS	KS
plug	0.353	0.276	0.126	0.411	diff	0.672	0.895	0.705	0.840

plug-in estimates are much smaller than those based on  $\hat{\Sigma}(t)$ . The smaller  $p$ -values might result from the inaccurate size performance associated with the plug-in estimator, which is evidenced by extra simulation results in [Table A.1](#) of the supplemental material showing that the methods with plug-in estimates tend to over-reject and result in smaller  $p$ -values. A further sensitivity check shows that when using  $1.2 \times \text{GCV}$  bandwidths, the R/S test with the plug-in estimator yields  $p$ -value 0.08 rejecting the null hypothesis at the significance level of 10%, while the  $p$ -values of tests equipped with the difference-based estimator remain large leading to the same decision of accepting the null for all four tests.

## 8 Conclusion

In this paper, we propose a debiased difference-based long-run covariance matrix estimator for the functional linear model under complex dynamics. We establish the uniform consistency of the proposed estimator and justify theoretically its application to structural change tests and detection of long memory. We conduct extensive simulation studies and achieve significant improvements in the finite samples. In particular, the new difference-based statistic is robust to both smooth and abrupt changes in the regression coefficients, largely eliminating the non-monotonic issue arising in the change point test. Finally, we apply our new estimators to the nonparametric analysis of Hong Kong circulatory and respiratory data, providing new uncertainty on structural changes and long memory. The optimal long-run variance for time series with stationary errors has been thoroughly discussed recently by [Chan \(2022\)](#). However, the approach therein is not applicable when non-stationarity is present. We leave the optimal estimation of the long-run covariance matrix under time series non-stationarity as rewarding future work. In addition, the generalization of our method beyond linear models will also be of great importance.

## 9 Acknowledgement

Weichi Wu is the corresponding author and is supported by NSFC 12271287.

## 10 Supplementary material

In the supplement, we present extra simulation studies and the proofs of results in this paper as well as auxiliary technical results.



# Supplement to “Difference-based covariance matrix estimate in time series nonparametric regression with applications to specification tests”

We organize the supplementary material as follows: We investigate the simulated sizes of tests for long memory and the magnitude of long-run covariance matrix estimates in the presence of smooth and abrupt changes in [Appendix A](#). [Appendix B](#) presents proofs of the results in the main paper. [Appendix C](#) provides auxiliary results which are used in the proofs.

## A Extra simulation

### A.1 Simulated rejected rates with sample size 750

The following [Table A.1](#) reports the simulated sizes of KPSS, R/S, V/S and K/S-type tests when  $n = 750$ .

	KPSS		R/S		V/S		K/S	
	5%	10%	5%	10%	5%	10%	5%	10%
plug-in	7.60	15.00	14.90	28.60	18.00	36.20	7.70	16.50
diff	4.80	8.90	3.70	6.20	3.60	5.90	4.50	9.40

Table A.1: Simulated rejection rates of KPSS, R/S, V/S and K/S-type tests for long memory when  $d = 0$ ,  $n = 750$ .

### A.2 Long-run covariance estimates with change points

To further illustrate the role of long-run variance estimators in bootstrap tests, we conduct 1000 times simulation, where we find the MSE of long-run variance estimators using ordinary least square residuals is 695.33, more than twice the magnitude of the MSE of that of difference-based method 313.72 when there are four change points in the time series structure (scenario CP4 with  $\delta = 1$ ). Moreover, two-sample t test also shows that under the alternative with  $\delta = 1$ , the difference-based method yields long-run variance estimate with much smaller MSE than the ordinary least square method with  $p$ -value smaller than 0.01%.

## B Proof

### B.1 Notation

We first introduce some notation that will be frequently used in the mathematical argument of this section. In the following proofs, we will omit the index  $n$  in  $e_{i,n}, x_{i,n}, y_{i,n}$  for simplicity. Define filtration  $\mathcal{F}_i = (\varepsilon_{-\infty}, \dots, \varepsilon_i)$  for *i.i.d.* random variables  $(\varepsilon_i)_{i \in \mathbb{Z}}$ . For a random vector  $(v_i)_{i=1}^n \in \mathcal{F}_s$ , let  $v_{i,\{s\}}$  denote the series replacing the  $\varepsilon_s$  with its *i.i.d.* copy. For a random matrix  $(A_i)_{i=1}^n \in \mathcal{F}_s$ , define  $A_{j,\{s\}}$  as the random matrix replacing  $\varepsilon_s$  in  $A_j$  with its *i.i.d.* copy. Recall that  $e_{i,n}^{(d)} = \sum_{j=0}^{\infty} \psi_j(d) e_{i-j,n}$ ,  $e_{i,n}^{(d_n)} = \sum_{j=0}^{\infty} \psi_j(d_n) e_{i-j,n}$ . For the sake of simplicity, we use  $\psi_j$  to represent  $\psi_j(d)$  when we discuss the fixed



alternatives and  $\psi_j(d_n)$  for the theory of the local alternatives. Recall  $t_i = i/n$ , and  $K^*(x)$  denotes the jackknife equivalent kernel  $2\sqrt{2}K(\sqrt{2}x) - K(x)$ . Let  $0 \times \infty = 0$ ,  $a_n \sim b_n$  denote  $\lim_{n \rightarrow \infty} a_n/b_n = 1$  for real sequences  $a_n$  and  $b_n$ . Let  $\mathcal{I} = [\gamma_n, 1 - \gamma_n] \subset (0, 1)$ ,  $\gamma_n = \tau_n + (m + 1)/n$ . Recall

$$A_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^j \{x_i x_i^\top \beta(t_i) - x_{i+m} x_{i+m}^\top \beta(t_{i+m})\}, \quad \Sigma^A(t) = \sum_{j=m}^{n-m} \frac{m A_{j,m} A_{j,m}^\top}{2} \omega(t, j),$$

$$\hat{A}_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^j \{x_i x_i^\top \check{\beta}(t_i) - x_{i+m} x_{i+m}^\top \check{\beta}(t_{i+m})\}, \quad \check{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m \hat{A}_{j,m} \hat{A}_{j,m}^\top}{2} \omega(t, j),$$

where  $\omega(t, i) = K_{\tau_n}(t_i - t) / \sum_{i=1}^n K_{\tau_n}(t_i - t)$ . Let

$$\dot{A}_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^j (x_i x_i^\top - x_{i+m} x_{i+m}^\top) (x_i x_i^\top \beta(t_i) - x_{i+m} x_{i+m}^\top \beta(t_{i+m})), \quad (\text{B.1})$$

and

$$\dot{\Delta}_j = \frac{1}{m} \sum_{i=j-m+1}^j (x_i x_i^\top - x_{i+m} x_{i+m}^\top) (x_i e_i - x_{i+m} e_{i+m}). \quad (\text{B.2})$$

We define below the counterparts of  $Q_{k,m}$ ,  $\Delta_j$  and  $\check{\Sigma}(\cdot)$  in [Section 3](#). Define for  $m \geq 2$ ,  $t \in [m/n, 1 - m/n]$ ,

$$\tilde{Q}_{k,m} = \sum_{i=k}^{k+m-1} x_i e_i, \quad \tilde{\Delta}_j = \frac{\tilde{Q}_{j-m+1,m} - \tilde{Q}_{j+1,m}}{m}, \quad \tilde{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m \tilde{\Delta}_j \tilde{\Delta}_j^\top}{2} \omega(t, j). \quad (\text{B.3})$$

## B.2 Proof of [Theorem 4.1](#)

The following lemma presents the consistency of bias correction for the difference-based estimator with time series covariates, which is crucial to establish the consistency result of [Theorem 4.1](#).

**Lemma B.1.** *Under the condition of [Theorem 4.1](#), we have*

$$\sup_{t \in \mathcal{I}} \left| \check{\Sigma}(t) - \Sigma^A(t) \right| = O_{\mathbb{P}} \left( \sqrt{m} \tau_n^{2-1/\kappa} + \sqrt{\frac{m}{n \tau_n^{1+2/\kappa}}} \right).$$

*Proof.* Let  $l_n$  be a sequence of real numbers so that  $l_n \rightarrow \infty$  arbitrarily slow. Define  $A_n = \{\sup_{t \in \mathcal{I}} |\Omega(t) - M^+(t)| \leq l_n ((m \tau_n)^{-1/2} + m/(n \tau_n^{1/2}) + \tau_n^{1/2})\}$ . By [\(B.9\)](#),  $\lim_{n \rightarrow \infty} \mathbb{P}(A_n) = 1$ . Since  $0 \leq \sum_{j=m}^{n-m} \omega(t, j) \leq 1$ ,

we have

$$\begin{aligned}
& \sup_{t \in \mathcal{I}} \left\| (\check{\Sigma}(t) - \Sigma^A(t)) \mathbf{1}(A_n) \right\|_{2\kappa} \\
& \leq \sup_{t \in \mathcal{I}} \sum_{j=m}^{n-m} \frac{m\omega(t, j)}{2} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} \left( \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} + 2\|A_{j,m}\|_{2\kappa} \right) \\
& \leq m \max_{m \leq j \leq n-m} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} \\
& \times \left( \max_{m \leq j \leq n-m} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} + 2 \max_{m \leq j \leq n-m} \|A_{j,m}\|_{2\kappa} \right). \tag{B.4}
\end{aligned}$$

First, we shall show that

$$\max_{m \leq j \leq n-m} \|A_{j,m}\|_{2\kappa} = O(m^{-1/2} + m/n) = O(m^{-1/2}). \tag{B.5}$$

Define  $B_{j,m} = \frac{1}{m} \sum_{i=j-m+1}^j x_i x_i^\top$ . Notice that

$$\|A_{j,m}\|_{2\kappa} \leq \sup_{m \leq i \leq n-m} \|\beta(t_i) - \beta(t_{i+m})\| \|B_{j,m}\|_{2\kappa} + \sup_{t \in [0,1]} |\beta(t)| \|B_{j,m} - B_{j+m,m}\|_{2\kappa} := A_1 + A_2.$$

Since  $\beta(t)$  is Lipschitz continuous, and under Assumption 4.3,  $\max_{m \leq j \leq n} \|B_{j,m}\|_{2\kappa}$  is bounded, we have

$$A_1 = O(m/n). \tag{B.6}$$

For the calculation of  $A_2$ , notice that

$$\begin{aligned}
\|B_{j,m} - B_{j+m,m}\|_{2\kappa} & \leq \|B_{j,m} - \mathbb{E}(B_{j,m})\|_{2\kappa} + \|B_{j+m,m} - \mathbb{E}(B_{j+m,m})\|_{2\kappa} \\
& \quad + \|\mathbb{E}(B_{j,m}) - M(t_j)\|_{2\kappa} + \|\mathbb{E}(B_{j+m,m}) - M(t_{j+m})\|_{2\kappa}.
\end{aligned}$$

Similar to Lemma 6 in Zhou and Wu (2010), using rectangular kernel with bandwidth  $m/n$ , under Assumption 4.3, we have

$$\sup_{m \leq j \leq n} \|B_{j,m} - \mathbb{E}(B_{j,m})\|_{2\kappa} = O(m^{-1/2}).$$

Since  $\mathbb{E}(B_{j,m}) = \frac{1}{m} \sum_{i=j-m+1}^j M(t_j)$  and  $M(t)$  is Lipschitz continuous,  $\|\mathbb{E}(B_{j,m}) - M(t_j)\|_{2\kappa} = O(m/n)$ . Finally, by the boundedness of  $\sup_{t \in [0,1]} |\beta(t)|$ , we have

$$A_2 = O(m^{-1/2} + m/n). \tag{B.7}$$

Therefore, by (B.6) and (B.7), we have shown (B.5).

Second, by triangle inequality, we have

$$\max_{m \leq j \leq n-m} \left\| (\hat{A}_{j,m} - A_{j,m}) 1(A_n) \right\|_{2\kappa} \leq 2 \max_{1 \leq i \leq n} \left\| x_i x_i^\top \right\|_{4\kappa} \|(\beta(t_i) - \check{\beta}(t_i)) 1(A_n)\|_{4\kappa}.$$

Since under Assumption 4.3,  $\max_{1 \leq i \leq n} \|x_i x_i^\top\|_{4\kappa} = O(1)$ , we shall show that

$$\sup_{t \in \mathcal{I}} \|(\beta(t) - \check{\beta}(t)) 1(A_n)\|_{4\kappa} = O(\tau_n^2). \quad (\text{B.8})$$

Let  $M^+(t) = \mathbb{E}\{\bar{J}(t, \mathcal{F}_0) \bar{J}^\top(t, \mathcal{F}_0)\}$ . Following similar arguments in Lemma 6 of [Zhou and Wu \(2010\)](#) and Theorem 5.2 of [Dette and Wu \(2019\)](#), under Assumptions 4.3 and 4.4, we have

$$\sup_{m \leq j \leq n} \|\hat{\Delta}_j/2 - M^+(t_j)\| = O(m^{-1/2} + m/n).$$

Then, it follows that

$$\sup_{t \in \mathcal{I}} \|\Omega(t) - M^+(t)\| = O(m^{-1/2} + m/n + \tau_n).$$

By the chaining argument in Propostion B.1 in Section B.2 in [Dette et al. \(2019\)](#), we have

$$\sup_{t \in \mathcal{I}} |\Omega(t) - M^+(t)| = O_{\mathbb{P}}((m\tau_n)^{-1/2} + m/(n\tau_n^{1/2}) + \tau_n^{1/2}). \quad (\text{B.9})$$

Note that  $\Omega(t)$  is invertible on  $A_n$ . Then, for a sufficiently large constant  $C$ , we have

$$\begin{aligned} \|(\check{\beta}(t) - \beta(t)) 1(A_n)\|_{4\kappa} &= \|\Omega^{-1}(t)(\varpi(t) - \Omega(t)\beta(t)) 1(A_n)\|_{4\kappa} \\ &\leq \|\rho(\Omega^{-1}(t))|\varpi(t) - \Omega(t)\beta(t)| 1(A_n)\|_{4\kappa} \leq C\|\varpi(t) - \Omega(t)\beta(t)\|_{4\kappa}. \end{aligned}$$

Then, it's sufficient to show that

$$\sup_{t \in \mathcal{I}} \|\varpi(t) - \Omega(t)\beta(t)\|_{4\kappa} = O(\tau_n^2).$$

Recall the definition of  $\hat{\Delta}_j$  and  $\hat{A}_{j,m}$  in (B.2) and (B.1) respectively. Observe that

$$\varpi(t) = \sum_{j=m}^{n-m} \frac{\omega(t, j)}{2} \hat{A}_{j,m} + \sum_{j=m}^{n-m} \frac{\omega(t, j)}{2} \hat{\Delta}_j := W_1 + W_2, \quad (\text{B.10})$$

where  $W_1, W_2$  are defined in the obvious way. Recall that  $\hat{\Delta}_j = \frac{1}{m} \sum_{i=j-m+1}^j (x_{i,n} x_{i,n}^\top - x_{i+m,n} x_{i+m,n}^\top)^2$ .

By triangle inequality, we have

$$\begin{aligned} \sup_{t \in \mathcal{I}} \|W_1 - \Omega(t)\beta(t)\|_{4\kappa} &\leq \sup_{t \in \mathcal{I}} \left\| W_1 - \sum_{j=m}^{n-m} \frac{\dot{\Delta}_j \omega(t, j)}{2} \beta(t_j) \right\|_{4\kappa} + \sup_{t \in \mathcal{I}} \left\| \sum_{j=m}^{n-m} \frac{\dot{\Delta}_j \omega(t, j)}{2} (\beta(t_j) - \beta(t)) \right\|_{4\kappa} \\ &:= W_{11} + W_{12}. \end{aligned}$$

Again, by triangle inequality, under Assumptions 4.1 and 4.3, we obtain

$$\begin{aligned} W_{11} &\leq \max_{m \leq j \leq n-m} \|\dot{A}_{j,m} - \dot{\Delta}_j \beta(t_j)\|_{4\kappa}, \\ &\leq \frac{1}{m} \max_{m \leq j \leq n-m} \left\{ \sum_{i=j-m+1}^{j+m} \|x_i x_i^\top - x_{i+m} x_{i+m}^\top\|_{8\kappa} \|x_i x_i^\top\|_{8\kappa} |\beta(t_i) - \beta(t_j)| \right\} \\ &= O(m/n). \end{aligned} \tag{B.11}$$

Under Assumption 4.3, we have  $\max_{m \leq j \leq n-m} \|\dot{\Delta}_j\|_{4\kappa} = O(1)$ . Then, by similar arguments in Lemmas 3 of Zhou and Wu (2010) and the continuity of  $M^+(t)$ , we obtain

$$\begin{aligned} W_{12} &= \sup_{t \in \mathcal{I}} \left\| \sum_{j=m}^{n-m} \frac{\dot{\Delta}_j \omega(t, j)}{2} (\beta'(t)(t_j - t) + O(\tau_n^2)) \right\|_{4\kappa} \\ &\leq \sup_{t \in \mathcal{I}} \left\| \sum_{j=m}^{n-m} (\dot{\Delta}_j/2 - M^+(t_j)) \omega(t, j) \beta'(t)(t_j - t) \right\|_{4\kappa} \\ &\quad + \sup_{t \in \mathcal{I}} \left\| \sum_{j=m}^{n-m} M^+(t_j) \omega(t, j) \beta'(t)(t_j - t) \right\|_{4\kappa} + O(\tau_n^2) \\ &= \sup_{t \in \mathcal{I}} \left\| \sum_{j=m}^{n-m} \omega(t, j) \beta'(t) M^+(t)(t_j - t) \right\|_{4\kappa} + O(\tau_n((n\tau_n)^{-1/2} + m/n) + \tau_n^2) \\ &= O(\tau_n^2). \end{aligned} \tag{B.12}$$

Therefore, combining (B.11) and (B.12), since  $n\tau_n^3 \rightarrow \infty$ ,  $m/(n\tau_n^2) \rightarrow 0$ , we have

$$\sup_{t \in \mathcal{I}} \|W_1 - \Omega(t)\beta(t)\|_{4\kappa} = O(\tau_n^2). \tag{B.13}$$

To proceed, we shall show that

$$\sup_{t \in \mathcal{I}} \|W_2\|_{4\kappa} = O((n\tau_n)^{-1/2} + \chi^m). \tag{B.14}$$

Under Assumption 4.3, we have

$$\left| \mathbb{E}(\dot{\Delta}_j) \right| = \frac{1}{m} \left| \sum_{i=j-m+1}^j \mathbb{E} \left( x_i x_i^\top x_{i+m} e_{i+m} \right) + \mathbb{E} \left( x_{i+m} x_{i+m}^\top x_i e_i \right) \right| = O(\chi^m). \quad (\text{B.15})$$

Then, by Burkholder's inequality, for a sufficiently large  $C$ , we have

$$\begin{aligned} \left\| \sum_{j=m}^{n-m} \frac{\omega(t, j)}{2} (\dot{\Delta}_j - \mathbb{E} \dot{\Delta}_j) \right\|_{4\kappa} &= \left\| \sum_{s=-m}^{\infty} \sum_{j=m}^{n-m} \frac{\omega(t, j)}{2} \mathcal{P}_{j-s} \dot{\Delta}_j \right\|_{4\kappa} \\ &\leq C \sum_{s=-m}^{\infty} \left\{ \sum_{j=m}^{n-m} \frac{\omega^2(t, j)}{4} \left\| \mathcal{P}_{j-s} \dot{\Delta}_j \right\|_{4\kappa}^2 \right\}^{1/2}. \end{aligned} \quad (\text{B.16})$$

Under Assumptions 4.3 and 4.5, using similar techniques in Lemma 3 of Zhou and Wu (2010), we have

$$\left\| \mathcal{P}_{j-s} \dot{\Delta}_j \right\|_{4\kappa} \leq \left\| \dot{\Delta}_j - \dot{\Delta}_{j, \{j-s\}} \right\|_{4\kappa} \leq \frac{1}{m} \sum_{i=j-m+1}^{j+m} (\delta_{8\kappa}(J, i-j+s) + \delta_{8\kappa}(U, i-j+s)). \quad (\text{B.17})$$

Then, (B.14) follows from (B.15), (B.16) and (B.17). Combining (B.13) and (B.14), since  $m = O(n^{1/3})$ , we have (B.8). Hence, by (B.4), under conditions  $m = O(n^{1/3})$ ,  $n\tau_n^3 \rightarrow \infty$ , we have

$$\sup_{t \in \mathcal{I}} \left\| (\check{\Sigma}(t) - \Sigma^A(t)) 1(A_n) \right\|_{\kappa} = O \left( \sqrt{m} \tau_n^2 + \sqrt{\frac{m}{n\tau_n}} \right).$$

Finally, the result follows from the chaining argument in Proposition B.1 in Section B.2 in Dette et al. (2019) and Proposition A.1 in Wu and Zhou (2018).  $\square$

### B.2.1 Proof of Theorem 4.1

Recall the definition of  $\tilde{\Sigma}(t)$  in (B.3). Observe that

$$\begin{aligned} \hat{\Sigma}(t) - \tilde{\Sigma}(t) &= \sum_{j=m}^{n-m} \frac{m \left( \Delta_j \Delta_j^\top - \tilde{\Delta}_j \tilde{\Delta}_j^\top \right)}{2} \omega(t, j) - \check{\Sigma}(t) \\ &= \Sigma^A(t) - \check{\Sigma}(t) + \frac{m}{2} \sum_{j=m}^{n-m} \left( A_{j,m} \tilde{\Delta}_j^\top + \tilde{\Delta}_j A_{j,m}^\top \right) \omega(t, j). \end{aligned}$$

Then, we have

$$\sup_{t \in \mathcal{I}} \left| \hat{\Sigma}(t) - \tilde{\Sigma}(t) \right| \leq \sup_{t \in \mathcal{I}} \left| \Sigma^A(t) - \check{\Sigma}(t) \right| + m \sup_{t \in \mathcal{I}} \left| \sum_{j=m}^{n-m} \omega(t, j) \tilde{\Delta}_j A_{j,m}^\top \right|. \quad (\text{B.18})$$

Take note that  $\Sigma^A(t)$  is the leading term of bias, so we introduce the correction. By [Lemma B.1](#), we have

$$\sup_{t \in \mathcal{I}} \left| \Sigma^A(t) - \check{\Sigma}(t) \right| = O_{\mathbb{P}} \left( \sqrt{m} \tau_n^{2-1/\kappa} + \sqrt{\frac{m}{n \tau_n^{1+2/\kappa}}} \right). \quad (\text{B.19})$$

To proceed, define

$$h_s(t) = \sum_{j=m}^{n-m} \omega(t, j) \mathcal{P}_{j-s} \{ \tilde{\Delta}_j A_{j,m}^\top \} := \sum_{j=m}^{n-m} h_{s,j}(t).$$

Under [Assumption 4.5](#), it's straightforward that  $\mathbb{E}(\tilde{\Delta}_j A_{j,m}^\top) = 0$ , for  $m \leq j \leq n - m$ . Then, we can write  $\sum_{j=m}^{n-m} \omega(t, j) \tilde{\Delta}_j A_{j,m}^\top$  as a summation of martingale differences, i.e.

$$\sum_{j=m}^{n-m} \omega(t, j) \tilde{\Delta}_j A_{j,m}^\top = \sum_{s=-m}^{\infty} h_s(t), \quad (\text{B.20})$$

Next, we shall show that  $\|h_{s,j}(t)\|_{\kappa} = O\{(1/n + m^{-3/2}) \min\{\chi^{s-m}, 1\}\}$ . Under [Assumptions 4.4](#) and [4.5](#), similar to (36) in [Lemma 3](#) in [Zhou and Wu \(2010\)](#), we have

$$\begin{aligned} \delta_{2\kappa}(\tilde{\Delta}(m), k) &:= \sup_{1 \leq j \leq n} \|\tilde{\Delta}_j - \tilde{\Delta}_{j,\{j-k\}}\|_{2\kappa} = O \left( \frac{1}{m} \sum_{i=-m+1}^m \delta_{2\kappa}(U, k+i) \right) \\ &= O(\min\{\chi^{k-m}, 1\}/m). \end{aligned} \quad (\text{B.21})$$

Using similar arguments in [\(B.21\)](#), by the boundedness of  $\beta(\cdot)$ , we have

$$\sup_{1 \leq j \leq n} \|A_{j,m} - A_{j,m,\{j-k\}}\|_{2\kappa} \leq \frac{M}{m} \sum_{i=-m+1}^m \delta_{4\kappa}(W, k+i) = O(\min\{\chi^{k-m}, 1\}/m), \quad (\text{B.22})$$

where  $M$  is a sufficiently large positive constant. Following similar arguments in [Theorem 1](#) of [Wu \(2007\)](#),  $\sup_j \|\tilde{\Delta}_j\|_{2\kappa} = O(m^{-1/2})$ . Similar to [\(B.5\)](#), we have uniformly for  $m \leq j \leq n - m$ ,

$$\|A_{j,m}\|_{2\kappa} = O(m/n + m^{-1/2}). \quad (\text{B.23})$$

Under [Assumption 4.2](#), from [\(B.21\)](#), [\(B.22\)](#) and [\(B.23\)](#), we obtain uniformly for  $m \leq j \leq n - m$ ,  $s \geq -m$ ,

$$\begin{aligned} \|h_{s,j}\|_{\kappa} / \omega(t, j) &= \|\mathcal{P}_{j-s} \{ \tilde{\Delta}_j A_{j,m}^\top \}\|_{\kappa} \\ &\leq \|A_{j,m,\{j-s\}} - A_{j,m}\|_{2\kappa} \|\tilde{\Delta}_j^\top\|_{2\kappa} + \|A_{j,m,\{j-s\}}\|_{2\kappa} \|\tilde{\Delta}_j^\top - \tilde{\Delta}_{j,\{j-s\}}^\top\|_{2\kappa} \\ &= O\{(1/n + m^{-3/2}) \min\{\chi^{s-m}, 1\}\}. \end{aligned}$$

Since  $h_{s,j}(t)$  are martingale differences with respect to  $j$ , we have for  $t \in [m/n, 1 - m/n]$ ,

$$\|h_s(t)\|_\kappa^2 = \sum_{j=m}^{n-m} \|h_{s,j}(t)\|_\kappa^2 = O \left[ (n\tau_n)^{-1} (1/n^2 + m^{-3}) \min\{\chi^{2s-2m}, 1\} \right]. \quad (\text{B.24})$$

By (B.20) and (B.24), we obtain

$$\left\| m \sum_{j=m}^{n-m} \omega(t, j) \tilde{\Delta}_j A_{j,m}^\top \right\|_\kappa \leq m \sum_{s=-m}^m \|h_s(t)\|_\kappa + m \sum_{s=m+1}^{\infty} \|h_s(t)\|_\kappa = O \left( \sqrt{\frac{m}{n\tau_n}} \right). \quad (\text{B.25})$$

By the chaining argument in Propostion B.1 in Section B.2 in Dette et al. (2019), we have

$$\sup_{t \in \mathcal{I}} \left| m \sum_{j=m}^{n-m} \omega(t, j) \tilde{\Delta}_j A_{j,m}^\top \right| = O_{\mathbb{P}} \left( \sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}} \right). \quad (\text{B.26})$$

Combining (B.18), (B.19) and (B.26), we obtain

$$\sup_{t \in \mathcal{I}} \left| \hat{\Sigma}(t) - \tilde{\Sigma}(t) \right| = O_{\mathbb{P}} \left( \sqrt{m\tau_n^{2-1/\kappa}} + \sqrt{\frac{m}{n\tau_n^{1+2\kappa}}} \right). \quad (\text{B.27})$$

By Lemma 3 in Zhou and Wu (2010), Proposition B.1 in Section B.2 in Dette et al. (2019), we have

$$\sup_{t \in \mathcal{I}} \left| \tilde{\Sigma}(t) - \mathbb{E}\tilde{\Sigma}(t) \right| = O_{\mathbb{P}} \left( \sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}} \right). \quad (\text{B.28})$$

Using similar techniques in Lemma 4 and Lemma 5 of Zhou and Wu (2010), which hold uniformly for  $t \in (0, 1)$ , (B.28) leads to

$$\sup_{t \in \mathcal{I}} \left| \tilde{\Sigma}(t) - \Sigma(t) \right| = O_{\mathbb{P}} \left( \sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}} + \frac{1}{m} + \sqrt{m\tau_n^{2-1/\kappa}} \right), \quad (\text{B.29})$$

With (B.27) and (B.29), the supreme bound is thus proved.  $\square$

### B.3 Proof of Theorem 5.1

Let  $\mathcal{C}_m = \{j - m \leq i \leq j + m, \beta'''(t_i) \text{ exists and continuous}\}$ . Similar to (B.4),

$$\begin{aligned}
& \sup_{t \in \mathcal{I}} \left\| (\check{\Sigma}(t) - \Sigma^A(t)) \mathbf{1}(A_n) \right\|_{\kappa} \\
& \leq \sup_{t \in \mathcal{I}} \sum_{j=m}^{n-m} \frac{m\omega(t, j)}{2} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} \left( \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} + 2\|A_{j,m}\|_{2\kappa} \right) \\
& \leq \sup_{t \in \mathcal{I}} \left( \sum_{j \in \mathcal{C}_m, m \leq j \leq n-m} + \sum_{j \notin \mathcal{C}_m, m \leq j \leq n-m} \right) \frac{m\omega(t, j)}{2} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} \\
& \quad \times \left( \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} + 2\|A_{j,m}\|_{2\kappa} \right). \tag{B.30}
\end{aligned}$$

For  $j \notin \mathcal{C}_m$ , we have  $\|A_{j,m}\|_{2\kappa} = O(1)$  and  $\left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} = O(1)$ . We shall show that,

$$\max_{j \in \mathcal{C}_m} \left\| (\hat{A}_{j,m} - A_{j,m}) \mathbf{1}(A_n) \right\|_{2\kappa} = O(\tau_n^2).$$

It suffices to show that

$$\sup_{j \in \mathcal{C}_m} \left\| (\beta(t_j) - \check{\beta}(t_j)) \mathbf{1}(A_n) \right\|_{4\kappa} = O(\tau_n^2). \tag{B.31}$$

Following Lemma B.1, it's sufficient to show that

$$\sup_{t \in \mathcal{I}} \left\| \varpi(t) - \Omega(t)\beta(t) \right\|_{4\kappa} = O(\tau_n^2).$$

Recall the definition of  $\hat{\Delta}_j$  and  $\hat{A}_{j,m}$  in (B.2) and (B.1) and we have from (B.10) that

$$\varpi(t) = \sum_{j=m}^{n-m} \frac{\omega(t, j)}{2} \hat{A}_{j,m} + \sum_{j=m}^{n-m} \frac{\omega(t, j)}{2} \hat{\Delta}_j := W_1 + W_2,$$



where  $W_1, W_2$  are defined in the obvious way. Following (B.11), (B.12) and (B.14), we have

$$\begin{aligned}
& \sup_{r \in \mathcal{C}_m} \|W_1 - \Omega(t_r)\beta(t_r)\| \\
& \leq \sum_{j=m}^{n-m} \frac{\omega(t_r, j)}{2m} \left\{ \sum_{i=j-m+1}^j \left\| x_i x_i^\top - x_{i+m} x_{i+m}^\top \right\|_{8\kappa} \right. \\
& \quad \times \left. \left\| x_i x_i^\top (\beta(t_i) - \beta(t_r)) - x_{i+m} x_{i+m}^\top (\beta(t_{i+m}) - \beta(t_r)) \right\|_{8\kappa} \right\} \\
& \leq \sup_{r \in \mathcal{C}_m} \sum_{j=m}^{n-m} \frac{\omega(t_r, j)}{2m} \left( \sum_{j \in \mathcal{C}_m, m \leq j \leq n-m} + \sum_{j \notin \mathcal{C}_m, m \leq j \leq n-m} \right) \left\{ \sum_{i=j-m+1}^j \left\| x_i x_i^\top - x_{i+m} x_{i+m}^\top \right\|_{8\kappa} \right. \\
& \quad \times \left. \left\| x_i x_i^\top (\beta(t_i) - \beta(t_r)) - x_{i+m} x_{i+m}^\top (\beta(t_{i+m}) - \beta(t_r)) \right\|_{8\kappa} \right\} \\
& = O\left(\frac{mq_n}{n\tau} + \frac{m}{n}\right), \tag{B.32}
\end{aligned}$$

as well as

$$\sup_{r \in \mathcal{C}_m} \|W_2\| = O((n\tau)^{-1/2} + \chi^m). \tag{B.33}$$

Since  $q_n = o(n\tau_n^3/m)$ , following similar arguments of Lemma B.1, combining (B.32) and (B.33), we have proved (B.31). Then, by (B.30) and the chaining argument of Proposition B.1 of Dette et al. (2019), we still have

$$\begin{aligned}
\sup_{t \in \mathcal{I}} \left| \check{\Sigma}(t) - \Sigma^A(t) \right| &= O_{\mathbb{P}} \left( \sqrt{m\tau_n^{2-1/\kappa}} + \sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}} + \frac{m^2 q_n}{n\tau^{1+1/\kappa}} \right) \\
&= O_{\mathbb{P}} \left( \sqrt{m\tau_n^{2-1/\kappa}} + \sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}} + \frac{mq_n}{n\tau^3} \right). \tag{B.34}
\end{aligned}$$

We proceed to investigate Theorem 4.1 in the presence of change points in  $\beta(t)$ . Recall that from (B.18),

$$\sup_{t \in \mathcal{I}} \left| \hat{\Sigma}(t) - \check{\Sigma}(t) \right| \leq \sup_{t \in \mathcal{I}} \left| \Sigma^A(t) - \check{\Sigma}(t) \right| + m \sup_{t \in \mathcal{I}} \left| \sum_{j=m}^{n-m} \omega(t, j) \tilde{\Delta}_j A_{j,m}^\top \right|. \tag{B.35}$$

Inspecting (B.24), we have

$$\begin{aligned}
\|h_s(t)\|_{\kappa}^2 &= \sum_{j \in \mathcal{C}_m, m \leq j \leq n-m} \|h_{s,j}(t)\|_{\kappa}^2 + \sum_{j \notin \mathcal{C}_m, m \leq j \leq n-m} \|h_{s,j}(t)\|_{\kappa}^2 \\
&= O\left( (n\tau_n)^{-1} (1/n^2 + m^{-3}) \min\{\chi^{2s-2m}, 1\} + \frac{q_n}{m(n\tau_n)^2} \min\{\chi^{2s-2m}, 1\} \right).
\end{aligned}$$

Similar to (B.25), we have

$$\begin{aligned} \left\| m \sum_{j=m}^{n-m} \omega(t, j) \tilde{\Delta}_j A_{j,m}^\top \right\|_\kappa &\leq m \sum_{s=-m}^m \|h_s(t)\|_\kappa + m \sum_{s=m+1}^{\infty} \|h_s(t)\|_\kappa \\ &= O\left(\sqrt{\frac{m}{n\tau_n}} + \sqrt{\frac{q_n m^3}{(n\tau_n)^2}}\right) = O\left(\sqrt{\frac{m}{n\tau_n}}\right). \end{aligned}$$

By the chaining argument in Proposition B.1 in Section B.2 in Dette et al. (2019), similar to (B.26), we have

$$\sup_{t \in \mathcal{I}} \left| m \sum_{j=m}^{n-m} \omega(t, j) \tilde{\Delta}_j A_{j,m}^\top \right| = O_{\mathbb{P}}\left(\sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}}\right). \quad (\text{B.36})$$

Finally, combining (B.34), (B.35), and (B.36), following similar argument of Theorem 4.1, we have shown the first part of the results. The second part follows from Yu et al. (2015) and Theorem 3.4 of Wu and Zhou (2018).  $\square$ .

## B.4 Proof of Proposition 5.1

Proposition 5.1 follows from similar but easier arguments of Theorem 3.1 and Theorem 3.2 of Wu and Zhou (2018).

## B.5 Proof of Theorem 5.2

For the clarity of presentation, write  $\hat{\Sigma}_d(t)$  for  $\hat{\Sigma}(t)$  defined in (3.3) under the fixed alternative. Define

$$\Sigma_d(t) = \kappa_2(d) \sigma_H^2(t) \mu_W(t) \mu_W^\top(t), \quad t \in [0, 1].$$

For the quantities under the fixed alternatives, let  $\tilde{Q}_{k,m}^{(d)} = \sum_{i=k}^{k+m-1} x_i e_i^{(d)}$ ,  $\tilde{\Delta}_j^{(d)} = \frac{\tilde{Q}_{j-m+1,m}^{(d)} - \tilde{Q}_{j+1,m}^{(d)}}{m}$ ,  $\tilde{\Sigma}_d(t) = \sum_{j=m}^{n-m} \frac{m \tilde{\Delta}_j^{(d)} \{\tilde{\Delta}_j^{(d)}\}^\top}{2} \omega(t, j)$ .  $\tilde{\Delta}_j^{(d)} = \frac{1}{m} \sum_{i=j-m+1}^j (x_i x_i^\top - x_{i+m} x_{i+m}^\top) (x_i e_i^{(d)} - x_{i+m} e_{i+m}^{(d)})$ . Let  $\hat{A}^{(d)}$ ,  $\varpi^{(d)}(\cdot)$ ,  $\check{\beta}^{(d)}(\cdot)$ ,  $\check{\Sigma}_d(\cdot)$  denote the counterparts of  $\hat{A}$ ,  $\varpi(\cdot)$ ,  $\check{\beta}(\cdot)$  and  $\check{\Sigma}(\cdot)$  in (3.3) under the fixed alternatives. Define  $H^{(d)}(t_i, \mathcal{F}_i) = \sum_{k=0}^{\infty} \psi_k H(t_{i-k}, \mathcal{F}_{i-k})$ ,  $U^{(d)}(t, \mathcal{F}_i) = W(t, \mathcal{F}_i) H^{(d)}(t, \mathcal{F}_i)$ . The following proposition is from Theorem 4.2 of Bai and Wu (2021).

**Proposition B.1.** *Under Assumptions 4.3 and 5.1, we have*

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r x_{i,n} e_{i,n}^{(d)} - \sum_{i=\lfloor nb_n \rfloor + 1}^r \mu_W(t_i) e_{i,n}^{(d)} \right| = O_{\mathbb{P}}(\sqrt{n}(\log n)^d).$$

**Corollary B.1.** *Under the conditions of Proposition B.1, assume  $l/\log n \rightarrow \infty$ ,  $l/n \rightarrow 0$ , we have*

$$\max_{1 \leq k \leq n-l+1} \left\| \sum_{i=k}^{k+l-1} (x_i - \mu_W(t_i)) e_i^{(d)} \right\| = O(\sqrt{l}(\log n)^d).$$

*Proof.* The corollary follows from a careful check of the proof of Theorem 4.2 of [Bai and Wu \(2021\)](#).  $\square$

### B.5.1 Asymptotic behavior of the bias correction term under the fixed alternatives

**Lemma B.2.** *Under the conditions of [Theorem 5.2](#), we have*

$$\sup_{t \in \mathcal{I}} \left| \check{\Sigma}_d(t) - \Sigma^A(t) \right| = O_{\mathbb{P}} \left( \sqrt{m}((n\tau_n)^{d-1/2} + \tau_n^2)\tau_n^{-1/\kappa} + m((n\tau_n)^{2d-1} + \tau_n^4)\tau_n^{-1/\kappa} \right) = o_{\mathbb{P}}(m^{2d}).$$

*Proof.* After a careful check of the proof of [Lemma B.1](#), the behavior of  $W_1$  is unchanged under the fixed alternatives and it's sufficient to show that  $W_2^{(d)}$ ,  $W_2$  under the fixed alternatives, s.t.

$$\sup_{t \in \mathcal{I}} \left\| W_2^{(d)} \right\|_{4\kappa} = O((n\tau_n)^{d-1/2}). \quad (\text{B.37})$$

Then the lemma will follow from the similar steps in [Lemma B.1](#). Similar to [\(B.17\)](#), under Assumptions [4.3](#) and [4.5](#), following similar arguments in [Lemma C.2](#), we have

$$\left\| \mathcal{P}_{j-s} \dot{\Delta}_j^{(d)} \right\|_{4\kappa} \leq \frac{1}{m} \sum_{i=j-m+1}^{j+m} \left( \delta_{8\kappa}(J, i-j+s) + \delta_{8\kappa}(U^{(d)}, i-j+s) \right) = O \left( \frac{1}{m} \sum_{i=-m+1}^m (i+s)^{d-1} \right).$$

Let  $N = N_n = \lfloor n\tau_n \rfloor$ . Note that under Assumption [4.5](#), we can write

$$W_2^{(d)} = \sum_{s=0}^N \sum_{j=m}^{n-m} \frac{\omega(t, j)}{2} \mathcal{P}_{j-s} \dot{\Delta}_j^{(d)} + \sum_{j=m}^{n-m} \frac{\omega(t, j)}{2} \sum_{s=N+1}^{\infty} \mathcal{P}_{j-s} \dot{\Delta}_j^{(d)} = W_{21}^{(d)} + W_{22}^{(d)}.$$

Since  $\mathcal{P}_{j-s} \dot{\Delta}_j^{(d)}$  are martingale differences with respect to  $j$ , for  $0 \leq s \leq N$ , we have

$$\left\| \sum_{j=m}^{n-m} \frac{\omega(t, j)}{2} \mathcal{P}_{j-s} \dot{\Delta}_j^{(d)} \right\|_{4\kappa}^2 = \sum_{j=m}^{n-m} \frac{\omega^2(t, j)}{4} \left\| \mathcal{P}_{j-s} \dot{\Delta}_j^{(d)} \right\|_{4\kappa}^2 = O \left( \frac{1}{n\tau_n m^2} \left( \sum_{i=-m+1}^m (i+s)^{d-1} \right)^2 \right). \quad (\text{B.38})$$

Therefore, by [\(B.38\)](#), we have

$$\left\| W_{21}^{(d)} \right\|_{4\kappa} = O(N^d / \sqrt{n\tau_n}) = O((n\tau_n)^{d-1/2}). \quad (\text{B.39})$$

Since  $\mathcal{P}_{j-s} \dot{\Delta}_j^{(d)}$  are martingale differences with respect to  $s$ , elementary calculations shows

$$\left\| \sum_{s=N+1}^{\infty} \mathcal{P}_{j-s} \dot{\Delta}_j^{(d)} \right\|_{4\kappa}^2 = O \left( m^{-2} \sum_{s=N+1}^{\infty} \left( \sum_{i=-m+1}^m (i+s)^{d-1} \right)^2 \right) = O(N^{2d-1}). \quad (\text{B.40})$$

Therefore, by [\(B.40\)](#) and triangle inequality we have

$$\left\| W_{22}^{(d)} \right\|_{4\kappa} = O(N^{d-1/2}) = O((n\tau_n)^{d-1/2}). \quad (\text{B.41})$$

Therefore, (B.37) follows from (B.39) and (B.41). □

### B.5.2 Proof of Theorem 5.2

Recall  $\tilde{Q}_{k,m}^{(d)} = \sum_{i=k}^{k+m-1} x_i e_i^{(d)}$ ,  $\tilde{\Delta}_j^{(d)} = \frac{\tilde{Q}_{j-m+1,m}^{(d)} - \tilde{Q}_{j+1,m}^{(d)}}{m}$ ,  $\tilde{\Sigma}_d(t) = \sum_{j=m}^{n-m} \frac{m \tilde{\Delta}_j^{(d)} \{\tilde{\Delta}_j^{(d)}\}^\top}{2} \omega(t, j)$ . We break the proof into 6 steps.

**Step 1:** We shall prove that under the bandwidth conditions  $\kappa \geq 4/(1/2 - d)$ ,  $m/(n\tau_n^2) \rightarrow 0$ ,  $\sqrt{m}\tau_n^{2-1/k} \rightarrow 0$ ,  $m = O(n^{1/3})$ ,  $m \rightarrow \infty$ ,  $n\tau_n^3 \rightarrow \infty$ ,

$$\sup_{t \in [0,1]} \left| \hat{\Sigma}_d(t) - \tilde{\Sigma}_d(t) \right| = O_{\mathbb{P}} \left( \sqrt{m} ((n\tau_n)^{d-1/2} + \tau_n^2) \tau_n^{-1/\kappa} + m ((n\tau_n)^{2d-1} + \tau_n^4) \tau_n^{-1/\kappa} \right) = o_{\mathbb{P}}(m^{2d}).$$

Recall that  $\Sigma^A(t) = \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} A_{j,m} A_{j,m}^\top$ , and similar to (B.18), we have

$$\sup_{t \in \mathcal{I}} \left| \hat{\Sigma}_d(t) - \tilde{\Sigma}_d(t) \right| \leq \sup_{t \in \mathcal{I}} \left| \check{\Sigma}_d(t) - \Sigma^A(t) \right| + \sup_{t \in \mathcal{I}} \left| \sum_{j=m}^{n-m} m\omega(t,j) \tilde{\Delta}_j^{(d)} A_{j,m}^\top \right|,$$

where the first term has been investigated in Lemma B.2.

Define  $h_{s,j}^{(d)}(t) = \mathcal{P}_{j-s}(\tilde{\Delta}_j^{(d)} A_{j,m}^\top)$ . Let  $N = N_n = (n\tau_n)$ . Observe that under Assumption 4.5,

$$\sum_{j=m}^{n-m} \omega(t,j) \tilde{\Delta}_j^{(d)} A_{j,m}^\top = \sum_{s=0}^N \sum_{j=m}^{n-m} \omega(t,j) h_{s,j}^{(d)} + \sum_{j=m}^{n-m} \omega(t,j) \sum_{s=N+1}^{\infty} h_{s,j}^{(d)} := S_1 + S_2,$$

where  $S_1$  and  $S_2$  are defined in the obvious way. To proceed, we first calculate  $\|h_{s,j}^{(d)}(t)\|_\kappa$ .

By Lemma B.4,  $\sup_j \|\tilde{\Delta}_j^{(d)}\|_{2\kappa} = O(m^{d-1/2})$ . Then, we have

$$\|h_{s,j}^{(d)}(t)\|_\kappa = O \left( m^{d-3/2} \min\{\chi^{s-m}, 1\} + (1/n + m^{-3/2}) \sum_{i=-m+1}^m \psi_{s+i} \right). \quad (\text{B.42})$$

Since  $h_{s,h}^{(d)}(t)$  are martingale differences with respect to  $j$ , since  $m/(n\tau_n) \rightarrow 0$ , we have for  $t \in \mathcal{I}$ ,

$$\|S_1\|_\kappa \leq \sum_{s=0}^N \left\| \sum_{j=m}^{n-m} \omega(t,j) h_{s,j}^{(d)} \right\|_\kappa = O \left( \sum_{s=0}^N \left( \sum_{j=m}^{n-m} \omega^2(t,j) \|h_{s,j}^{(d)}\|_\kappa^2 \right)^{1/2} \right) = O(m^{-1/2} (n\tau_n)^{d-1/2}). \quad (\text{B.43})$$

By (B.42) and triangle inequality, elementary calculation shows that

$$\|S_2\|_\kappa \leq \sum_{j=m}^{n-m} \omega(t,j) \left\| \sum_{s=N+1}^{\infty} h_{s,j}^{(d)} \right\|_\kappa = O \left\{ \left( \sum_{s=N+1}^{\infty} \max_{m \leq j \leq n-m} \|h_{s,j}^{(d)}\|_\kappa^2 \right)^{1/2} \right\} = O(m^{-1/2} (n\tau_n)^{d-1/2}). \quad (\text{B.44})$$

where the first big  $O$  follows from the fact that  $h_{s,j}^{(d)}$  are martingale differences with respect to  $s$ . Combining (B.43) and (B.44), by chaining argument in Proposition B.1 in Dette et al. (2019), we have

$$\sup_{t \in \mathcal{I}} \left| \sum_{j=m}^{n-m} m\omega(t, j) \tilde{\Delta}_j^{(d)} A_{j,m}^\top \right| = O_{\mathbb{P}} \left( \sqrt{\frac{m}{(n\tau_n)^{1-2d}}} \tau_n^{-1/\kappa} \right). \quad (\text{B.45})$$

Combining Lemma B.3 and (B.45), we have shown (B.65).

**Step 2:** Define  $\bar{Q}_{k,m}^{(d)} = \sum_{i=k}^{k+m-1} \mu_W(t_i) e_i^{(d)}$ ,  $1 \leq k \leq n-m+1$ ,  $\bar{\Delta}_j^{(d)} = \frac{\bar{Q}_{j-m+1,m}^{(d)} - \bar{Q}_{j+1,m}^{(d)}}{m}$ ,  $\bar{\Sigma}_d(t) = \sum_{j=m}^{n-m} \frac{m \bar{\Delta}_j^{(d)} \{\bar{\Delta}_j^{(d)}\}^\top}{2} \omega(t, j)$ . We shall show that under the bandwidth condition  $m\tau_n^{3/2}/\log n \rightarrow \infty$ ,  $\kappa \geq 2/(3d)$ ,

$$\sup_{t \in \mathcal{I}} \left| \tilde{\Sigma}_d(t) - \bar{\Sigma}_d(t) \right| = O_{\mathbb{P}}(m^d (\log n)^d \tau_n^{-1/\kappa}) = o_{\mathbb{P}}(m^{2d}). \quad (\text{B.46})$$

Following similar arguments in Corollary B.1, we have

$$\max_{1 \leq k \leq n-m+1} \left\| \bar{Q}_{k,m}^{(d)} - \tilde{Q}_{k,m}^{(d)} \right\|_{2\kappa} = O(\sqrt{m} (\log n)^d). \quad (\text{B.47})$$

Using (B.47), and the fact  $\sup_j \|\tilde{\Delta}_j^{(d)}\|_{2\kappa} = O(m^{d-1/2})$ ,  $\sup_j \|\bar{\Delta}_j^{(d)}\|_{2\kappa} = O(m^{d-1/2})$ , we have

$$\begin{aligned} \left\| \tilde{\Delta}_j^{(d)} \left\{ \tilde{\Delta}_j^{(d)} \right\}^\top - \bar{\Delta}_j^{(d)} \left\{ \bar{\Delta}_j^{(d)} \right\}^\top \right\|_{\kappa} &\leq \left\| \tilde{\Delta}_j^{(d)} - \bar{\Delta}_j^{(d)} \right\|_{2\kappa} \left\| \left\{ \bar{\Delta}_j^{(d)} \right\}^\top \right\|_{2\kappa} + \left\| \tilde{\Delta}_j^{(d)} \right\|_{2\kappa} \left\| \left\{ \tilde{\Delta}_j^{(d)} - \bar{\Delta}_j^{(d)} \right\}^\top \right\|_{2\kappa} \\ &= O(m^{d-1} (\log n)^d). \end{aligned}$$

Since  $m\tau_n^{3/2}/\log n \rightarrow \infty$ , (B.46) follows from triangle inequality and Proposition B.1 in Dette et al. (2019).

**Step 3:** Let  $\zeta_j = \sum_{i=j}^{\infty} \mathcal{P}_j e_i$ ,  $\zeta_j^\circ = \zeta_j(t_j) = \sum_{i=j}^{\infty} \mathcal{P}_j H(t_j, \mathcal{F}_i)$ . Define

$$\bar{Z}_{k,m} = \sum_{j=0}^L \psi_j \sum_{i=k}^{k+m-1} \mu_W(t_i) \zeta_{i-j}^\circ, \quad \Delta_j^{(d),\circ} = \frac{\bar{Z}_{j-m+1,m} - \bar{Z}_{j+1,m}}{m}, \quad \Sigma_d^\circ(t) = \sum_{j=m}^{n-m} \frac{m \Delta_j^{(d),\circ} \{\Delta_j^{(d),\circ}\}^\top}{2} \omega(t, j).$$

Let  $L = Mm^{1+\frac{1}{2d+1}} \tau_n^{1/2}$ , where  $M$  is a sufficiently large constant. We will show that

$$\sup_{t \in \mathcal{I}} |\bar{\Sigma}_d(t) - \Sigma_d^\circ(t)| = O_{\mathbb{P}} \left\{ m^{2d} \left( m^{-\frac{1/2-d}{2d+1}} \tau_n^{d/2-1/4-1/\kappa} \right) \right\} = o_{\mathbb{P}}(m^{2d}). \quad (\text{B.48})$$

Since  $m\tau_n \rightarrow \infty$ ,  $m^2 \tau_n^{1/2}/n = O(n^{-1/3} \tau_n^{1/2}) = o(1)$ , then  $L/m \rightarrow \infty$ ,  $L/m^2 \rightarrow 0$ ,  $m^{1+1/(2d)}/L \rightarrow \infty$ ,

$L(\log n)^2/n \rightarrow 0$ . Observe that

$$\begin{aligned}
& \|\bar{\Sigma}_d(t) - \Sigma_d^\circ(t)\|_\kappa \\
& \leq \sum_{j=m}^{n-m} \frac{m\omega(t, j)}{2} \left\| \bar{\Delta}_j^{(d)} \left\{ \bar{\Delta}_j^{(d)} \right\}^\top - \Delta_j^{(d), \circ} \left\{ \Delta_j^{(d), \circ} \right\}^\top \right\|_\kappa \\
& \leq m \max_{m \leq j \leq n-m} \left\{ \left\| \bar{\Delta}_j^{(d)} \right\|_{2\kappa} \left\| \left\{ \bar{\Delta}_j^{(d)} \right\}^\top - \left\{ \Delta_j^{(d), \circ} \right\}^\top \right\|_{2\kappa} + \left\| \bar{\Delta}_j^{(d)} - \Delta_j^{(d), \circ} \right\|_{2\kappa} \left\| \left\{ \Delta_j^{(d), \circ} \right\}^\top \right\|_{2\kappa} \right\}, \tag{B.49}
\end{aligned}$$

where

$$\left\| \bar{\Delta}_j^{(d)} - \Delta_j^{(d), \circ} \right\|_{2\kappa} \leq \frac{1}{m} \left( \left\| \bar{Q}_{j-m+1, m}^{(d)} - \bar{Z}_{j-m+1, m} \right\|_{2\kappa} + \left\| \bar{Q}_{j+1, m}^{(d)} - \bar{Z}_{j+1, m} \right\|_{2\kappa} \right).$$

Define

$$\bar{W}_{k, m} = \sum_{j=0}^L \psi_j \sum_{i=k}^{k+m-1} \mu_W(t_i) e_{i-j}, \quad 1 \leq k \leq n-m+1.$$

Then, we have for  $1 \leq k \leq n-m+1$  that

$$\begin{aligned}
\left\| \bar{Q}_{k-m+1, m}^{(d)} - \bar{Z}_{k-m+1, m} \right\|_{2\kappa} & \leq \left\| \bar{Q}_{k-m+1, m}^{(d)} - \bar{W}_{k-m+1, m} \right\|_{2\kappa} + \left\| \bar{W}_{k-m+1, m} - \bar{Z}_{k-m+1, m} \right\|_{2\kappa} \\
& =: C_{2\kappa, 1} + C_{2\kappa, 2}, \tag{B.50}
\end{aligned}$$

where  $C_{2\kappa, 1}$  and  $C_{2\kappa, 2}$  are defined in the obvious way.

Under conditions 4.5 and 4.3, by Burkholder's inequality, we have for  $k \geq m$ ,

$$\begin{aligned}
C_{2\kappa, 1} & = \left\| \sum_{j=L+1}^{\infty} \left\{ \sum_{i=0}^{\min\{m, j-L\}-1} \psi_{j-i} \mu_W \left( \frac{k-i}{n} \right) \right\} e_{k-j} \right\|_{2\kappa} \\
& \leq \sum_{t=0}^{\infty} \left[ \sum_{j=L+1}^{\infty} \left\{ \sum_{i=0}^{\min\{m, j-L\}-1} \psi_{j-i} \mu_W \left( \frac{k-i}{n} \right) \right\}^2 \left\| \mathcal{P}_{k-j-t} e_{k-j} \right\|_{2\kappa}^2 \right]^{1/2} \\
& \leq \left[ \sum_{j=L+1}^{\infty} \left\{ \sum_{i=0}^m \psi_{j-i} \mu_W \left( \frac{k-i}{n} \right) \right\}^2 \right]^{1/2} \sum_{t=0}^{\infty} \delta_{2\kappa}(H, t, (-\infty, 1]) \\
& = O(L^{d-1/2} m). \tag{B.51}
\end{aligned}$$

Then, we consider the upper bound of  $C_{2\kappa, 2}$ . Let  $p_{j, k, m} = \sum_{i=(j-L)_+}^{(m-1) \wedge j} \psi_{j-i} \mu_W \left( \frac{k-i}{n} \right)$ ,  $m \leq k \leq n-m$ . Then, for  $m \leq k \leq n-m$ , we can write

$$\bar{W}_{k-m+1, m} - \bar{Z}_{k-m+1, m} = \sum_{j=0}^{L+m-1} p_{j, k, m} (e_{k-j} - \zeta_{k-j}^\circ).$$

After a careful check on Lemma 2 in [Wu and Zhou \(2011\)](#), we have

$$\max_{0 \leq l \leq L+m-1} \left\| \sum_{j=0}^l (e_{k-j} - \zeta_{k-j}) \right\|_{2\kappa}^2 \leq M \sum_{i=1}^{L+m} \left( \sum_{j=i}^{\infty} \delta_{2\kappa}(H, j, (-\infty, 1]) \right)^2 = O(1), \quad (\text{B.52})$$

where  $M$  is a sufficiently large constant. Following the proof of Corollary 2 in [Wu and Zhou \(2011\)](#), under Assumption 5.1, we obtain

$$\|\zeta_i - \zeta_i^\circ\|_{2\kappa} = O((\log n)^2/n). \quad (\text{B.53})$$

Observe that

$$\sum_{l=0}^{L+m-1} |p_{l,k,m} - p_{l-1,k,m}| = O(L^d). \quad (\text{B.54})$$

Then, by the summation-by-parts formula, combining (B.52), (B.53) and (B.54), since  $L(\log n)^2/n \rightarrow 0$ , we have

$$\|\bar{W}_{k-m+1,m} - \bar{Z}_{k-m+1,m}\|_{2\kappa} = \left\| \sum_{j=0}^{L+m-1} p_{j,k,m} (e_{k-j} - \zeta_{k-j}^\circ) \right\|_{2\kappa} = O(L^d). \quad (\text{B.55})$$

Since  $\sup_j \|\bar{\Delta}_j^{(d)}\|_{2\kappa} = O(m^{d-1/2})$ ,  $\sup_j \|\Delta_j^{(d),\circ}\|_{2\kappa} = O(m^{d-1/2})$ , by (B.49), (B.50), (B.51) and (B.55), since  $L/m^2 \rightarrow 0$ , we obtain

$$\|\bar{\Sigma}_d(t) - \Sigma_d^\circ(t)\|_{\kappa} = O(L^{d-1/2} m^{d+1/2}).$$

Under the conditions  $m\tau_n^{3/2} \rightarrow \infty$  and  $\kappa \geq 4/(1/2 - d)$ , (B.48) then follows from Proposition B.1 in [Dette et al. \(2019\)](#).

**Step 4:** We shall show that under condition  $m\tau_n^{3/2} \rightarrow \infty$ ,

$$\sup_{t \in \mathcal{I}} |\Sigma_d^\circ(t) - \mathbb{E}\Sigma_d^\circ(t)| = O_{\mathbb{P}} \left\{ m^{2d} (m\tau_n^{3/2})^{-1/2} \right\} = o_{\mathbb{P}}(m^{2d}). \quad (\text{B.56})$$

Following similar arguments in the proof of Theorem 3.1 [Wu and Shao \(2006\)](#), for  $0 \leq k \leq \lfloor \frac{n-m}{2L} \rfloor$ , let  $D_{k,i} = \Delta_{2kL+i}^{(d),\circ} \{\Delta_{2kL+i}^{(d),\circ}\}^\top - \mathbb{E}(\Delta_{2kL+i}^{(d),\circ} \{\Delta_{2kL+i}^{(d),\circ}\}^\top | \mathcal{F}_{2kL+i-2L})$ ,  $i = 0, 1, 2, \dots, 2L-1$ , and  $E_h = \mathbb{E}(\Delta_h^{(d),\circ} \{\Delta_h^{(d),\circ}\}^\top | \mathcal{F}_{h-2L})$ ,  $\mathbb{E}(\Delta_h^{(d),\circ} \{\Delta_h^{(d),\circ}\}^\top)$ ,  $m \leq h \leq n - m$ . Let  $D_{k,i} = 0$ , if  $2kL + i < m$  or  $2kL + i > n - m$ . Then, we have

$$\Sigma_d^\circ(t) - \mathbb{E}\Sigma_d^\circ(t) = \sum_{h=m}^{n-m} \frac{m\omega(t, h)}{2} E_h + \sum_{i=0}^{2L-1} \sum_{k=0}^{\lfloor n/(2L) \rfloor} \frac{m\omega(t, 2kL + i)}{2} D_{k,i}. \quad (\text{B.57})$$

Recall that  $\Delta_h^{(d),\circ} = \frac{\bar{Z}_{h-m+1,m} - \bar{Z}_{h+1,m}}{m}$ , in which

$$\bar{Z}_{h,m} = \sum_{j=0}^L \psi_j \sum_{i=h}^{h+m-1} \mu_W(t_i) \zeta_{i-j}^\circ = \sum_{j=0}^{L+m-1} p_{j,h+m-1,m} \zeta_{h+m-1-j}^\circ,$$

where  $p_{j,h+m-1,m} = \sum_{i=(j-L)_+}^{(m-1)\wedge j} \psi_{j-i} \mu_W\left(\frac{h+m-1-i}{n}\right)$ ,  $\{\zeta_j^\circ\}$  are martingale differences.

Under the geometric measure contraction condition, for  $0 \leq j \leq L$ , we have

$$\|\mathbb{E}((\zeta_{r-j}^\circ)^2 | \mathcal{F}_{r-2L}) - \mathbb{E}(\zeta_{r-j}^\circ)^2\| = O(\chi^L). \quad (\text{B.58})$$

By Lemma C.1 and elementary calculation, we have

$$\sum_{j=0}^{L+m-1} |p_{j,s,m} p_{j,s,m}^\top| = O(m^{2d+1}), \quad \sum_{j=0}^{L-1} |p_{j,s,m} p_{j+m,s,m}^\top| = O(m^{2d+1}). \quad (\text{B.59})$$

Therefore, combining (B.58) and (B.59) we derive

$$\|E_h\| = O(m^{2d-1} \chi^L). \quad (\text{B.60})$$

By Burkholder's inequality, uniformly for all  $i$ ,

$$\begin{aligned} \left\| \sum_{k=1}^{\lfloor n/(2L) \rfloor} \omega(t, 2kL+i) D_{k,i} \right\|^2 &\leq C \sum_{k=1}^{\lfloor n/(2L) \rfloor} \omega^2(t, 2kL+i) \|D_{k,i}\|^2 \\ &\leq 2C \sum_{k \in \{r: |2rL+i-nt| \leq n\tau_n\}} (\|\Delta_{2kL+i}^{(d),\circ}\|_4 \|\{\Delta_{2kL+i}^{(d),\circ}\}^\top\|_4)^2 / (n\tau_n)^2 \\ &= O((n\tau_n)^{-1} L^{-1} m^{4d-2}), \end{aligned} \quad (\text{B.61})$$

where  $C$  is a sufficiently large constant. Therefore, since  $L/(n\tau_n) = m^{1+\frac{1}{2d+1}} / (n\tau_n^{1/2}) = O(1/(m\tau_n^{1/2}))$ , by (B.57), (B.60) and (B.61), and Proposition B.1 in Dette et al. (2019), we have shown (B.56).

**Step 5:** Recall that  $L = Mm^{1+\frac{1}{2d+1}} \tau_n^{1/2}$ ,  $m \rightarrow \infty$ ,  $m = O(n^{1/3})$ . It follows that  $m^{1+\frac{1}{d+1}}/L \rightarrow \infty$ ,  $L^2/(mn) = O(m^3\tau_n/n) = o(1)$ . We shall show that uniformly for  $s \in \mathcal{I}$ ,

$$m^{-2d} \mathbb{E} \Sigma_d^\circ(s) = \kappa_2(d) \mu_W(s) \mu_W^\top(s) \sigma_H^2(s) + O(f_n), \quad (\text{B.62})$$

where  $\kappa_2(d) = \Gamma^{-2}(d+1) \int_0^\infty (t^d - (t-1)_+^d) (2t^d - (t-1)_+^d - (t+1)^d) dt$ , and  $f_n = m^{-d} + \tau_n^2 + L^{d+1}/m^{d+2} + L^2/(mn) + (L/m^2)^{-\frac{1}{d-2}} = o(1)$ . Note that  $\bar{Z}_{k-m+1,m} = \sum_{j=0}^{L+m-1} p_{j,k,m} \zeta_{k-j}^\circ$ . Recall

$$p_{j,k,m} = \sum_{i=(j-L)_+}^{(m-1)\wedge j} \psi_{j-i} \mu_W\left(\frac{k-i}{n}\right) = \begin{cases} \sum_{i=0}^{j-1} \psi_{j-i} \mu_W\left(\frac{k-i}{n}\right) + \mu_W\left(\frac{k-j}{n}\right), & j \leq m-1 \\ \sum_{i=0}^{m-1} \psi_{j-i} \mu_W\left(\frac{k-i}{n}\right), & m \leq j \leq L \\ \sum_{i=j-L}^{m-1} \psi_{j-i} \mu_W\left(\frac{k-i}{n}\right) = O(mL^{d-1}), & j \geq L+1. \end{cases}$$



Then, approximate  $p_{j,k,m}$  by integrals. When  $j \leq m-1$ , by the continuity of  $\mu_W$  and [Lemma C.1](#),

$$m^{-d}\Gamma(d)p_{j,k,m} = d^{-1}\mu_W(k/n)(j/m)^d + O(m^{-d} + m/n).$$

When  $m \leq j \leq L$ ,

$$m^{-d}\Gamma(d)p_{j,k,m} = d^{-1}\mu_W(k/n)((j/m)^d - ((j+1)/m - 1)^d) + O(m/n + m^{-1}(j/m - 1)^{d-1}).$$

Since  $\{\zeta_i^\circ\}$  are martingale differences, and  $\sigma_H(t_j) = \|\sum_{i=j}^\infty \mathcal{P}_j H(t_j, \mathcal{F}_i)\| = \|\zeta_j^\circ\|$ , [\(B.62\)](#) then follows from elementary calculation.

**Step 6:** Let  $g_{\kappa,n} = m^{-d}(\log n)^d \tau_n^{-1/\kappa} + \sqrt{m}((n\tau_n)^{d-1/2} + \tau_n^2)\tau_n^{-1/\kappa} + m((n\tau_n)^{2d-1} + \tau_n^4)\tau_n^{-1/\kappa} + m^{-\frac{1/2-d}{2d+1}}\tau_n^{d/2-1/4-1/\kappa} + (m\tau_n^{3/2})^{-1/2} + f_n$ . Summarizing Step 1-5, we have

$$\sup_{t \in \mathcal{I}} \left| m^{-2d} \hat{\Sigma}_d(t) - \kappa_2(d) \sigma_H^2(t) \mu_W(t) \mu_W^\top(t) \right| = O_{\mathbb{P}}(g_{\kappa,n}) = o_{\mathbb{P}}(1).$$

## B.6 Proof of [Theorem 5.3](#)

We define the notation under the local alternatives by replacing  $d$  with  $d_n$ .

### B.6.1 Technical results for the proof of [Theorem 5.3](#)

[Lemma B.3](#) studies the asymptotic behavior of the bias correction term under the local alternatives. [Lemma B.4](#) investigates the physical dependence of  $U^{(d_n)}(t, \mathcal{F}_i)$  as well as the order of its partial sum process under the local alternatives.

**Lemma B.3.** *Under the conditions of [Theorem 5.3](#), we have*

$$\sup_{t \in \mathcal{I}} \left| \check{\Sigma}_{d_n}(t) - \Sigma^A(t) \right| = O_{\mathbb{P}} \left( \sqrt{m} \tau_n^{2-1/\kappa} + \sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}} \right) = o_{\mathbb{P}}(1).$$

*Proof of [Lemma B.3](#).* Letting  $d_n = c/\log n$ , the proof follows from similar steps in [Lemma B.2](#). □

**Lemma B.4.** *Under Assumptions [4.3](#) and [5.1](#),  $m \rightarrow \infty$ ,  $m = O(n)$ , we have*

$$\sup_{1 \leq k \leq n-m+1} \left\| \sum_{i=k}^{k+m-1} x_i e_i^{(d_n)} \right\|_4 = O(\sqrt{m}).$$

*Proof.* Define  $\tilde{Q}_{k,m}^{(d_n)} = \sum_{i=k}^{k+m-1} x_i e_i^{(d_n)}$ . By similar arguments in [Lemma C.2](#), we obtain

$$\delta_4(U^{(d_n)}, k) = O(\psi_k(d_n)), \quad k \geq 0, \tag{B.63}$$

and  $\delta_4(U^{(d_n)}, k) = 0$ , for  $k < 0$ . Then, uniformly for  $1 \leq k \leq n-m+1$ , by Burkholder's inequality we

have

$$\left\| \tilde{Q}_{k,m}^{(d_n)} \right\|_4^2 \leq B_4^2 \left\| \sum_{l=-\infty}^{\infty} \left| \mathcal{P}_l \tilde{Q}_{k,m}^{(d_n)} \right|^2 \right\| \leq B_4^2 \sum_{l=-\infty}^{k+m-1} \left\| \mathcal{P}_l \tilde{Q}_{k,m}^{(d_n)} \right\|_4^2 \leq B_4^2 \sum_{l=-\infty}^{k+m-1} \left( \sum_{i=k-l}^{k+m-1-l} \delta_4(U^{(d_n)}, i) \right)^2, \quad (\text{B.64})$$

where  $B_4$  is a constant. Therefore, combining (B.63) and (B.64), it follows from Lemma C.3 that

$$\max_{1 \leq k \leq n-m+1} \left\| \tilde{Q}_{k,m}^{(d_n)} \right\|_4^2 = O \left( \sum_{l=-\infty}^k \left( (k+m-l)^{d_n} - (k-l+1)^{d_n} \right)^2 + \sum_{l=k+1}^{k+m-1} (k+m-l)^{2d_n} \right) = O(m).$$

□

### B.6.2 Proof of Theorem 5.3

Recall that

$$\tilde{\Sigma}_{d_n}(t) = \sum_{j=m}^{n-m} \frac{m \tilde{\Delta}_j^{(d_n)} \{ \tilde{\Delta}_j^{(d_n)} \}^\top}{2} \omega(t, j),$$

where  $\tilde{\Delta}_j^{(d_n)} = \frac{\tilde{Q}_{j-m+1,m}^{(d_n)} - \tilde{Q}_{j+1,m}^{(d_n)}}{m}$ ,  $\tilde{Q}_{k,m}^{(d_n)} = \sum_{i=k}^{k+m-1} x_i e_i^{(d_n)}$ .

We break the proof in the following 8 steps.

**Step 1:** Following the proof of Theorem 5.2 by replacing  $d$  by  $d_n$ , since  $\kappa \geq 4$ , we have

$$\sup_{t \in \mathcal{I}} \left| \hat{\Sigma}_{d_n}(t) - \tilde{\Sigma}_{d_n}(t) \right| = O_{\mathbb{P}} \left( \sqrt{m} \tau_n^{2-1/\kappa} + \sqrt{\frac{m}{n \tau_n^{1+2/\kappa}}} \right) = o_{\mathbb{P}}(1). \quad (\text{B.65})$$

**Step 2:** Let  $L = m^2 \tau_n^{1/2}$ ,  $\check{e}_{i,L}^{(d_n)} = \sum_{j=0}^L \psi_j e_{i-j}$ . Define

$$\check{Q}_{k,m}^{(d_n)} =: \sum_{i=k}^{k+m-1} x_i \check{e}_{i,L}^{(d_n)}, \quad \check{\Delta}_j^{(d_n)} = \frac{\check{Q}_{j-m+1,m}^{(d_n)} - \check{Q}_{j+1,m}^{(d_n)}}{m}, \quad \check{\Sigma}_{d_n}(t) = \sum_{j=m}^{n-m} \frac{m \check{\Delta}_j^{(d_n)} \{ \check{\Delta}_j^{(d_n)} \}^\top}{2} \omega(t, j).$$

In this step, we shall show that under bandwidth condition  $m \tau_n^{3/2} \rightarrow \infty$ ,

$$\sup_{t \in \mathcal{I}} \left| \tilde{\Sigma}_{d_n}(t) - \check{\Sigma}_{d_n}(t) \right| = O_{\mathbb{P}}(m^{-1/2} \tau_n^{-3/4}) = o_{\mathbb{P}}(1). \quad (\text{B.66})$$

Observe that

$$e_i^{(d_n)} = \check{e}_{i,L}^{(d_n)} + \tilde{e}_{i,L}^{(d_n)}, \quad \text{where } \check{e}_{i,L}^{(d_n)} = \sum_{j=0}^L \psi_j e_{i-j}, \quad \tilde{e}_{i,L}^{(d_n)} = \sum_{j=L+1}^{\infty} \psi_j e_{i-j}.$$

By Proposition C.1, we have  $\|\tilde{e}_{i,L}^{(d_n)}\|_4^2 = O(\sum_{s=L+1}^{\infty} (s+1)^{2d_n-2}) = O(L^{-1})$ . Then, under Assumption 4.3,

we have uniformly for  $1 \leq k \leq n - m + 1$ ,

$$\left\| \check{Q}_{k,m}^{(d_n)} - \tilde{Q}_{k,m}^{(d_n)} \right\| \leq m \max_{1 \leq i \leq n} \left\| x_i \tilde{e}_{i,L}^{(d_n)} \right\| \leq m \max_{1 \leq i \leq n} \|x_i\|_4 \left\| \tilde{e}_{i,L}^{(d_n)} \right\|_4 = O(m/\sqrt{L}). \quad (\text{B.67})$$

By Lemma B.4 and (B.67), we have

$$\left\| \check{\Sigma}_{d_n}(t) - \tilde{\Sigma}_{d_n}(t) \right\| \leq m \max_{m \leq j \leq n-m} \left\| \check{\Delta}_j^{(d_n)} - \tilde{\Delta}_j^{(d_n)} \right\|_4 \left( \left\| \check{\Delta}_j^{(d_n)} - \tilde{\Delta}_j^{(d_n)} \right\|_4 + 2 \left\| \check{\Delta}_j^{(d_n)} \right\|_4 \right) = O\left(\sqrt{m/L}\right).$$

By Proposition B.1 in Dette et al. (2019), since  $m/(L\tau_n) = m^{-1}\tau_n^{-3/2} \rightarrow 0$ , (B.66) is proved.

**Step 3 :** Define  $\bar{Q}_{k,m}^{(d_n)} =: \sum_{i=k}^{k+m-1} (x_i e_i + \mu_W(t_i)(\tilde{e}_{i,L}^{(d_n)} - e_i))$ ,

$$\bar{\Delta}_j^{(d_n)} = \frac{\bar{Q}_{j-m+1,m}^{(d_n)} - \bar{Q}_{j+1,m}^{(d_n)}}{m}, \quad \bar{\Sigma}_{d_n}(t) = \sum_{j=m}^{n-m} \frac{m \bar{\Delta}_j^{(d_n)} \{\bar{\Delta}_j^{(d_n)}\}^\top}{2} \omega(t, j).$$

We shall show that

$$\sup_{t \in \mathcal{I}} |\check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t)| = O_{\mathbb{P}} \left( \sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}} + d_n \right) = o_{\mathbb{P}}(1).$$

Observe that

$$\check{Q}_{k,m}^{(d_n)} - \bar{Q}_{k,m}^{(d_n)} = \sum_{j=1}^L \sum_{i=k}^{k+m-1} (x_i - \mu_W(t_i)) \psi_j e_{i-j} = \sum_{j=1}^L \sum_{i=k}^{k+m-1} \psi_j \bar{x}_i e_{i-j},$$

where  $\bar{x}_i = x_i - \mu_W(t_i)$ . Let

$$\vartheta_{k,m} = \frac{1}{m} \sum_{j=1}^L \sum_{i=k-m+1}^k \psi_j (\bar{x}_i e_{i-j} - \bar{x}_{i+m} e_{i+m-j}).$$

Then, it follows that

$$\check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t) = \sum_{j=m}^{n-m} \frac{m\omega(t, j)}{2} (\check{\Delta}_j^{(d_n)} \{\check{\Delta}_j^{(d_n)}\}^\top - \bar{\Delta}_j^{(d_n)} \{\bar{\Delta}_j^{(d_n)}\}^\top), \quad (\text{B.68})$$

and

$$\check{\Delta}_j^{(d_n)} \{\check{\Delta}_j^{(d_n)}\}^\top - \bar{\Delta}_j^{(d_n)} \{\bar{\Delta}_j^{(d_n)}\}^\top = \vartheta_{k,m} \vartheta_{k,m}^\top + \vartheta_{k,m} \{\bar{\Delta}_j^{(d_n)}\}^\top + \bar{\Delta}_j^{(d_n)} \vartheta_{k,m}^\top. \quad (\text{B.69})$$

**Step 3.1** We first show that

$$\sup_{t \in \mathcal{I}} |\mathbb{E}(\check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t))| = O(d_n) = o(1). \quad (\text{B.70})$$

Observe that  $\{\bar{x}_i\}_{i=1}^n$ ,  $\{e_i\}_{i=-\infty}^n$  are two centered sequences. Under Assumptions 4.3 and 5.1, by

Lemma 7 in [Zhou \(2014\)](#) we have for  $l, j > 0$ ,

$$|\mathbb{E}(\bar{x}_i e_{i-j} \bar{x}_{i+k}^\top e_{i+k-l})| = O(\chi^{\rho^*}),$$

where following the lines in the proof of Theorem 2 in [Zhou \(2014\)](#), we have

$$\rho^* \geq \frac{1}{2} \min\{\max(|k|, |k-l+j|), \max(|k-l|, |k+j|)\} := \rho_{k,l,j}.$$

Then, we are able to bound the expectation of [\(B.69\)](#), for  $1 \leq p \leq L$ ,  $1 \leq q \leq L$ ,

$$\begin{aligned} & |\mathbb{E}(\vartheta_{k,m} \vartheta_{k,m}^\top)| \\ & \leq \frac{1}{m^2} \sum_{p,q=1}^L \psi_p \psi_q \left| \mathbb{E} \left[ \left\{ \sum_{i=k-m+1}^k (\bar{x}_i e_{i-p} - \bar{x}_{i+m} e_{i+m-p}) \right\} \left\{ \sum_{j=k-m+1}^k (\bar{x}_j e_{j-q} - \bar{x}_{j+m} e_{j+m-q}) \right\}^\top \right] \right| \\ & = O \left( \frac{1}{m^2} \sum_{p,q=1}^L \psi_p \psi_q \sum_{i,j=k-m+1}^{k+m} \chi^{\rho_{j-i,q,p}} \right). \end{aligned}$$

Consider  $q \leq p$ , since when  $q \geq 1$ ,  $\psi_q = O(d_n(1+q)^{d_n-1})$ , we have

$$\begin{aligned} & \frac{1}{m^2} \sum_{q=1}^L \sum_{p=q}^L \psi_p \psi_q \sum_{i,j=k-m+1}^{k+m} \chi^{\rho_{j-i,q,p}} \\ & = \frac{1}{m^2} \sum_{q=1}^L \sum_{p=q}^L \psi_p \psi_q \sum_{i=k-m+1}^{k+m} \left( \sum_{j>i+(q-p)/2} \chi^{(j-i-q+p)/2} + \sum_{j \leq i+(q-p)/2} \chi^{(i-j)/2} \right) \\ & = O \left( \sum_{q=1}^L \psi_q^2 / m \right) = O(d_n/m). \end{aligned}$$

Similarly,

$$\frac{1}{m^2} \sum_{q=1}^L \sum_{p=1}^{q-1} \psi_p \psi_q \sum_{i,j=k-m+1}^{k+m} \chi^{\rho_{j-i,q,p}} = O(d_n/m).$$

Then, we have

$$|\mathbb{E}(\vartheta_{k,m} \vartheta_{k,m}^\top)| = O(d_n/m). \tag{B.71}$$

Following similar arguments in [Lemma B.4](#),  $\|\bar{\Delta}_j^{(d_n)}\| = O(m^{-1/2})$ . Then, it follows that

$$|\mathbb{E}(\bar{\Delta}_j^{(d_n)} \vartheta_{k,m}^\top)| \leq \|\bar{\Delta}_j^{(d_n)}\| \|\vartheta_{k,m}^\top\| = O(d_n^{1/2}/m). \tag{B.72}$$

Therefore, by [\(B.69\)](#), [\(B.68\)](#), [\(B.71\)](#), and [\(B.72\)](#), we have [\(B.70\)](#).

**Step 3.2** We proceed to show that

$$\sup_{t \in \mathcal{I}} |\check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t) - \mathbb{E}(\check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t))| = O_{\mathbb{P}} \left( \sqrt{\frac{m}{nT_n^{1+2/\kappa}}} \right) = o_{\mathbb{P}}(1). \quad (\text{B.73})$$

Notice that  $\check{e}_{i,L}^{(d_n)} - e_i$  has summable physical dependence. Specifically,

$$\begin{aligned} \|\vartheta_{k,m} - \vartheta_{k,m,\{k-s\}}\|_{2\kappa} &= O \left\{ \frac{1}{m} \sum_{j=1}^L \psi_j \sum_{i=k-m+1}^{k+m} (\delta_{4\kappa}(W, i-k+s) + \delta_{4\kappa}(H, i-j-k+s)) \right\} \\ &= O \left( \frac{d_n}{m} \sum_{i=-m+1}^m \min \left\{ \chi^{i-L+s} L^{d_n-1}, (i+s)^{d_n-1} \right\} 1(i+s > 0) \right), \end{aligned}$$

where in the last equality, we use the fact  $\delta_{4\kappa}(H, k) = 0$ , if  $k \leq 0$  and  $\sum_{j=1}^L \psi_j \chi^{L-j} = O(\psi_L)$ . From (B.71), we have  $\sup_j \|\vartheta_{j,m}\|_{2\kappa} = O(m^{-1/2})$ .

For simplicity, write  $r_{i,s,n} = \frac{d_n}{m} \min \left\{ \chi^{i-L+s} L^{d_n-1}, (i+s)^{d_n-1} \right\} 1(i+s > 0)$ . Then, we obtain for  $m \leq j \leq n$ ,  $s \geq 0$ ,

$$\begin{aligned} \left\| \mathcal{P}_{j-s} \vartheta_{j,m} \vartheta_{j,m}^{\top} \right\|_{\kappa} &\leq \|\vartheta_{j,m}\|_{2\kappa} \|\vartheta_{j,m}^{\top} - \vartheta_{j,m,\{j-s\}}^{\top}\|_{2\kappa} + \|\vartheta_{j,m} - \vartheta_{j,m,\{j-s\}}\|_{2\kappa} \|\vartheta_{j,m,\{j-s\}}^{\top}\|_{2\kappa} \\ &= O \left( m^{-1/2} r_{i,s,n} \right). \end{aligned} \quad (\text{B.74})$$

Under [Assumption 4.4](#), similar to (36) in Lemma 3 in [Zhou and Wu \(2010\)](#), we have

$$\begin{aligned} \delta_{2\kappa}(\tilde{\Delta}(m), k) &:= \sup_{1 \leq j \leq n} \|\tilde{\Delta}_j - \tilde{\Delta}_{j,\{j-k\}}\|_{2\kappa} = O \left( \frac{1}{m} \sum_{i=-m+1}^m \delta_{2\kappa}(U, k+i) \right) \\ &= O(\min\{\chi^{k-m}, 1\}/m). \end{aligned}$$

Similar to (B.74), and by (B.21), we have

$$\begin{aligned} \left\| \bar{\Delta}_k^{(d_n)} - \bar{\Delta}_{k,\{k-s\}}^{(d_n)} \right\|_{2\kappa} &\leq \left\| \tilde{\Delta}_k - \tilde{\Delta}_{k,\{k-s\}} \right\|_{2\kappa} + \frac{1}{m} \sum_{i=k-m+1}^{k+m} \sum_{j=1}^L \psi_j \left\| \mu_W(t_i) (e_{i-j} - e_{i-j,\{k-s\}}) \right\|_{2\kappa} \\ &= O \left( \min\{\chi^{s-m}, 1\}/m + \sum_{i=-m+1}^m r_{i,s,n} \right). \end{aligned} \quad (\text{B.75})$$

Since  $\sup_j \|\bar{\Delta}_j^{(d_n)}\|_{\kappa} = O(m^{-1/2})$ , by (B.75), similar to (B.74), we obtain

$$\left\| \mathcal{P}_{j-s} \bar{\Delta}_j^{(d_n)} \vartheta_{j,m}^{\top} \right\|_{\kappa} = O \left( m^{-1/2} r_{i,s,n} + m^{-3/2} \min\{\chi^{s-m}, 1\} \right). \quad (\text{B.76})$$

By Burkholder's inequality, by (B.68) and (B.69), combining (B.74) and (B.76), we have for  $t \in \mathcal{I}$ ,

$$\begin{aligned} & \left\| \check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t) - \mathbb{E}(\check{\Sigma}_{d_n}(t) - \bar{\Sigma}_{d_n}(t)) \right\|_{\kappa} \\ &= O \left\{ \sum_{s=0}^{\infty} \left( \sum_{j=m}^{n-m} \omega^2(t, j) m^2 \left\| \mathcal{P}_{j-s} \vartheta_{j,m} \vartheta_{j,m}^{\top} + \mathcal{P}_{j-s} \bar{\Delta}_j^{(d_n)} \vartheta_{j,m}^{\top} + \mathcal{P}_{j-s} \bar{\Delta}_j^{(d_n)} \vartheta_{j,m}^{\top} \right\|_{\kappa}^2 \right)^{1/2} \right\} \\ &= O \left( \sqrt{\frac{m}{n\tau_n}} \right), \end{aligned}$$

where in the last equality, we consider  $0 < i + s < L$ , and  $i + s \geq L$  separately and use the fact  $\sum_{i=1}^L i^{-1} = O(\log L)$ . Then, (B.73) follows from the chaining argument in Proposition B.1 in Dette et al. (2019).

**Step 4: Decomposition** Recall that  $\tilde{Q}_{k,m} = \sum_{i=k}^{k+m-1} x_i e_i$ ,  $\tilde{\Delta}_j = \frac{\tilde{Q}_{j-m+1,m} - \tilde{Q}_{j+1,m}}{m}$ , and

$$\tilde{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m \tilde{\Delta}_j \tilde{\Delta}_j^{\top}}{2} \omega(t, j).$$

Define  $\check{\Delta}_j^{(d_n)} = \bar{\Delta}_j^{(d_n)} - \tilde{\Delta}_j = \frac{1}{m} \sum_{i=j-m+1}^j \mu_W(t_i) (\check{e}_{i,L}^{(d_n)} - e_i) - \mu_W(t_{i+m}) (\check{e}_{i+m,L}^{(d_n)} - e_{i+m})$ . Let

$$\tilde{s}_1(t) = \sum_{j=m}^{n-m} \frac{m \omega(t, j)}{2} \check{\Delta}_j^{(d_n)} \{\check{\Delta}_j^{(d_n)}\}^{\top}, \quad \tilde{s}_2(t) = \sum_{j=m}^{n-m} \frac{m \omega(t, j)}{2} \tilde{\Delta}_j \{\check{\Delta}_j^{(d_n)}\}^{\top}.$$

Observe that

$$\bar{\Sigma}_{d_n}(t) = \tilde{\Sigma}(t) + \tilde{s}_1(t) + \tilde{s}_2(t) + \tilde{s}_2^{\top}(t).$$

### Step 5: Martingale approximation

Let

$$z_j = \sum_{i=j}^{\infty} \mathcal{P}_j \{x_i e_i\}, \quad z_j^{\circ} = z_j(t_j) = \sum_{i=j}^{\infty} \mathcal{P}_j U(t_j, \mathcal{F}_i).$$

Recall that in Theorem 5.2,  $\zeta_j = \sum_{i=j}^{\infty} \mathcal{P}_j e_i$ ,  $\zeta_j^{\circ} = \zeta_j(t_j) = \sum_{i=j}^{\infty} \mathcal{P}_j H(t_j, \mathcal{F}_i)$ . Let  $z_{j,1}$  denote the first element in  $z_j$ . Then, it follows that  $z_{j,1} = \zeta_j$ ,  $z_{j,1}^{\circ} = \zeta_j^{\circ}$ .

Define  $\bar{Z}_{k,m}^{(d_n)} = \sum_{i=k}^{k+m-1} \left( z_i^{\circ} + \sum_{j=1}^L \psi_j \mu_W(t_i) \zeta_{i-j}^{\circ} \right)$ ,

$$\Delta_j^{(d_n), \circ} = \frac{\bar{Z}_{j-m+1,m}^{(d_n)} - \bar{Z}_{j+1,m}^{(d_n)}}{m}, \quad \Sigma_{d_n}^{\circ}(t) = \sum_{j=m}^{n-m} \frac{m \Delta_j^{(d_n), \circ} \{\Delta_j^{(d_n), \circ}\}^{\top}}{2} \omega(t, j).$$

Similarly to  $p_{j,k,m}$  defined in Step 3 of Theorem 5.2, we define  $\underline{p}_{j,k,m} = \sum_{i=(j-L)_+}^{(m \wedge j)-1} \psi_{j-i} \mu_W \left( \frac{k-i}{n} \right)$ . By

(B.55) and similar arguments in Theorem 1(ii) of Wu (2007), we have uniformly for  $1 \leq k \leq n - m + 1$ ,

$$\|\bar{Q}_{k,m}^{(d_n)} - \bar{Z}_{k,m}^{(d_n)}\|_4 \leq \left\| \sum_{j=1}^{L+m-1} p_{j,k+m-1,m} (e_{k+m-1-j} - \zeta_{k+m-1-j}^\circ) \right\|_4 + \left\| \sum_{i=k}^{k+m-1} x_i e_i - \sum_{i=k}^{k+m-1} z_i^\circ \right\|_4 = O(1).$$

Since  $\sup_j \|\bar{\Delta}_j^{(d_n)}\| = O(m^{-1/2})$ , by triangle inequality and Cauchy–Schwarz inequality, we have for  $t \in \mathcal{I}$ ,

$$\|\bar{\Sigma}_{d_n}(t) - \Sigma_{d_n}^\circ(t)\| \leq \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \left\| \bar{\Delta}_j^{(d_n)} \{\bar{\Delta}_j^{(d_n)}\}^\top - \Delta_j^{(d_n),\circ} \{\Delta_j^{(d_n),\circ}\}^\top \right\| = O(m^{-1/2}).$$

By chaining argument in Proposition B.1 in Dette et al. (2019), we have

$$\sup_{t \in \mathcal{I}} |\bar{\Sigma}_{d_n}(t) - \Sigma_{d_n}^\circ(t)| = O_{\mathbb{P}}((m\tau_n)^{-1/2}) = o_{\mathbb{P}}(1).$$

**Step 6** Observe that

$$\bar{Z}_{k,m}^{(d_n)} = \sum_{i=k}^{k+m-1} (z_i^\circ + \sum_{j=1}^L \psi_j \mu_W(t_i) \zeta_{i-j}^\circ) = \sum_{i=k}^{k+m-1} z_i^\circ + \sum_{j=1}^{L+m-1} p_{j,k+m-1,m} \zeta_{k+m-1-j}^\circ.$$

After a careful inspection of Step 4 of Theorem 5.2, we have

$$\sup_{t \in \mathcal{I}} |\Sigma_{d_n}^\circ(t) - \mathbb{E}\Sigma_{d_n}^\circ(t)| = O_{\mathbb{P}}\left(\sqrt{\frac{1}{m\tau_n^{3/2}}}\right) = o_{\mathbb{P}}(1),$$

**Step 7** Recall that  $\check{\Sigma}(t) = \Sigma(t) + (e^{c\alpha_1} - 1)^2 \sigma_H^2(t) \mu_W(t) \mu_W^\top(t) + (e^{c\alpha_1} - 1) s_{UH}(t) \mu_W^\top(t) + (e^{c\alpha_1} - 1) \mu_W(t) s_{UH}^\top(t)$ . We shall show that uniformly for  $t \in \mathcal{I}$ ,

$$\mathbb{E}\Sigma_{d_n}^\circ(t) = \check{\Sigma}(t) + O((\log n)^{-1}).$$

Let  $\check{\Delta}_k^{(d_n),\circ} = \frac{1}{m} \sum_{i=k-m+1}^k \sum_{j=1}^L \psi_j \left\{ \mu_W(t_i) \zeta_{i-j}^\circ - \mu_W(t_{i+m}) \zeta_{i+m-j}^\circ \right\}$ ,  $\Delta_k^\circ = \frac{1}{m} \sum_{i=k-m+1}^k (z_i^\circ - z_{i+m}^\circ)$ .

Define for  $t \in [0, 1]$ ,  $\check{\Sigma}^\circ(t) = \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \Delta_j^\circ \Delta_j^{\circ,\top}$ ,

$$\check{s}_1^\circ(t) = \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \check{\Delta}_j^{(d_n),\circ} \{\check{\Delta}_j^{(d_n),\circ}\}^\top, \text{ and } \check{s}_2^\circ(t) = \sum_{j=m}^{n-m} \frac{m\omega(t,j)}{2} \Delta_j^\circ \{\Delta_j^\circ\}^\top.$$

Then it follows that

$$\mathbb{E}\Sigma_{d_n}^\circ(t) = \mathbb{E}\check{\Sigma}^\circ(t) + \mathbb{E}\check{s}_1^\circ(t) + \mathbb{E}\check{s}_2^\circ(t) + \mathbb{E}\check{s}_2^{\circ,\top}(t).$$

Following similar arguments in Step 5 of [Theorem 5.2](#), by the continuity of  $\mu_W$ , we have when  $j \leq m$ ,

$$\underline{p}_{j,k,m} = (e^{c\alpha_1} - 1)\mu_W(k/n) + O(m/n + d_n). \quad (\text{B.77})$$

when  $j \geq m + 1$ , since  $L/n \rightarrow 0$ ,

$$\underline{p}_{j,k,m} = \mu_W(k/n)(j^{d_n} - (j - m + 1)^{d_n}) + O(m/n + d_n + m/L). \quad (\text{B.78})$$

Since  $z_i^\circ$  are martingale differences, and  $\sigma_H(t_j) = \|\sum_{i=j}^\infty \mathcal{P}_j H(t_j, \mathcal{F}_i)\| = \|\zeta_j^\circ\|$ ,

$$\begin{aligned} \mathbb{E}(\overline{Z}_{k-m+1,m}^{(d_n)} \overline{Z}_{k-m+1,m}^{(d_n),\top}) &= \sum_{j=1}^{L+m-1} \underline{p}_{j,k,m} \underline{p}_{j,k,m}^\top \mathbb{E}(\zeta_{k-j}^\circ)^2 + \sum_{i=k-m+1}^k \mathbb{E}(z_i^\circ z_i^{\circ,\top}) \\ &\quad + \sum_{j=1}^{m-1} \underline{p}_{j,k,m} \mathbb{E}(\zeta_{k-j}^\circ z_{k-j}^{\circ,\top}) + \sum_{j=1}^{m-1} \mathbb{E}(z_{k-j}^\circ \zeta_{k-j}^\circ) \underline{p}_{j,k,m}^\top \\ &:= Z_1 + Z_2 + Z_3 + Z_4. \end{aligned} \quad (\text{B.79})$$

By [Lemma C.3](#), we have

$$\begin{aligned} Z_1/m &= \frac{1}{m} \sum_{j=1}^{m-1} \underline{p}_{j,k,m} \underline{p}_{j,k,m}^\top \mathbb{E}(\zeta_{k-j}^\circ)^2 + \frac{1}{m} \sum_{j=m}^{L+m-1} \underline{p}_{j,k,m} \underline{p}_{j,k,m}^\top \mathbb{E}(\zeta_{k-j}^\circ)^2 \\ &= (e^{c\alpha_1} - 1)^2 \mu_W(k/n) \mu_W^\top(k/n) \sigma_H^2(k/n) + O(m/n + d_n + (\log m)^{-1}). \end{aligned}$$

Under [Assumption 4.2](#), we have

$$Z_2/m = \Sigma(t) + O(m/n).$$

Observe that

$$\mathbb{E}(z_j^\circ \zeta_j^\circ) = \sum_{k=-\infty}^{\infty} \text{Cov}(U(t_j, \mathcal{F}_0), H(t_j, \mathcal{F}_k)) = s_{UH}(t_j).$$

Under [Assumption 5.2](#), similar arguments in the calculation of  $Z_1$  and  $Z_2$  imply,

$$Z_3/m = (e^{c\alpha_1} - 1)\mu_W(k/n) s_{UH}^\top(k/n) + O(m/n + d_n),$$

and

$$Z_4/m = (e^{c\alpha_1} - 1) s_{UH}(k/n) \mu_W^\top(k/n) + O(m/n + d_n).$$



By Lemma C.3 (a), (B.77) and (B.78), similar techniques of (B.79) show that

$$\begin{aligned}\mathbb{E}(\overline{Z}_{k+1,m}^{(d_n)} \overline{Z}_{k-m+1,m}^{(d_n),\top}) &= \sum_{j=1}^{L-1} \underline{p}_{j+m,k+m,m} \underline{p}_{j,k,m}^\top \mathbb{E}(c_{k-j}^\circ)^2 + \sum_{j=1}^m \underline{p}_{j+m-1,k+m,m} \mathbb{E}(c_{k+1-j}^\circ z_{k+1-j}^{\circ,\top}) \\ &= O(m(\log m)^{-1}).\end{aligned}\tag{B.80}$$

Therefore, by (B.79) and (B.80), we have

$$\mathbb{E}\Sigma_{d_n}^\circ(t) = \sum_{j=m}^{n-m} \check{\Sigma}(t_j)\omega(t, j) + O(m/n + d_n + (\log m)^{-1}) = \check{\Sigma}(t) + O((\log n)^{-1}).$$

**Step 8** Summarizing Step 1 - Step 7, we have

$$\sup_{t \in \mathcal{I}} |\hat{\Sigma}_{d_n}(t) - \check{\Sigma}(t)| = O_{\mathbb{P}} \left( \sqrt{\frac{m}{n\tau_n^{1+2/\kappa}}} + \sqrt{m\tau_n^{2-1/\kappa}} + \sqrt{\frac{1}{m\tau_n^{3/2}}} + (\log n)^{-1} \right) = o_{\mathbb{P}}(1).$$

□

## C Auxiliary results

**Proposition C.1.** *Suppose  $Q_i = L(t_i, \mathcal{F}_i)$ ,  $t_i \in \mathcal{I}$ , for  $q \geq 1$ , we have*

$$\|\mathcal{P}_{i-l}Q_i\|_q \leq \delta_q(L, l, \mathcal{I}).$$

*Proof.* The proposition follows after a careful investigation of Theorem 1 in Wu (2005). □

**Proposition C.2.** *Under Assumption 5.1, we have uniformly for  $l \geq 0$ ,  $0 < d < 1/2$ ,*

$$\delta_p(H^{(d)}, l, (-\infty, 1]) = O\{(1+l)^{d-1}\}.$$

*Proof.* Under Assumption 5.1, by Lemma 3.2 of Kokoszka and Taqqu (1995) and Proposition C.1, we have

$$\delta_p(H^{(d)}, l, (-\infty, 1]) \leq \sum_{k=0}^l \psi_k(d) \delta_p(H, l-k, (-\infty, 1]) = O\{(1+l)^{d-1}\}.$$

□

**Lemma C.1.** *The following argument shows the properties of long memory coefficient  $\psi_j = \psi_j(d)$ .  $\psi_0 = 1$ , and for  $j \geq 1$ ,*

$$\psi_j = j^{d-1}l_d(j),$$

where  $l_d(j) = 1/\Gamma(d)(1 + O(1/j))$ .

*Proof.* By Stirling's formula,

$$\begin{aligned}\frac{\Gamma(j+d)}{\Gamma(j+1)} &= \frac{\sqrt{\frac{2\pi}{j+d}} \left(\frac{j+d}{e}\right)^{j+d} (1 + O(\frac{1}{j+d}))}{\sqrt{\frac{2\pi}{j+1}} \left(\frac{j+1}{e}\right)^{j+1} (1 + O(\frac{1}{j+1}))} = \frac{j^{d-1} (1 + d/j)^j (1 + d/j)^{d-1/2} (1 + O(1/j))}{e^{d-1} (1 + 1/j)^j (1 + 1/j)^{1/2} (1 + O(1/j))} \\ &= j^{d-1} + O(j^{d-2}).\end{aligned}$$

Since  $\ln \Gamma(z) \sim z \ln z - z + \frac{1}{2} \ln \frac{2\pi}{z} + \sum_{n=1}^{N-1} \frac{B_{2n}}{2n(2n-1)z^{2n-1}}$ , the constant in the big  $O$  of  $\Gamma(z) = \sqrt{\frac{2\pi}{z}} \left(\frac{z}{e}\right)^z (1 + O(\frac{1}{z}))$  is always  $B_2/2 = 1/12$ .  $\square$

**Lemma C.2.** *Assuming  $\sup_{t \in (-\infty, 1]} \|H(t, \mathcal{F}_0)\|_{2p} < \infty$ ,  $\delta_{2p}(H, k, (-\infty, 1]) = O(\chi^k)$ ,  $\delta_{2p}(W, k) = O(\chi^k)$ ,  $\chi \in (0, 1)$ ,  $\sup_{t \in [0, 1]} \|W(t, \mathcal{F}_0)\|_{2p} < \infty$ , we have*

$$\delta_p(U^{(d)}, k) = O(k^{d-1}).$$

*Proof.* Note that for  $j \leq i$ ,

$$\delta_p(U^{(d)}, i-j) \leq \|W(t_i, \mathcal{F}_i)\|_{2p} \delta_{2p}(H^{(d)}, i-j) + \|H^{(d)}(t_i, \mathcal{F}_{i-j}^*)\|_{2p} \delta_{2p}(W, i-j). \quad (\text{C.1})$$

By Burkholder's inequality and [Proposition C.2](#), we have

$$\|H^{(d)}(t_i, \mathcal{F}_i)\|_{2p}^2 \leq M \left\| \sum_{j \in \mathbb{Z}} \left( \mathcal{P}_j H^{(d)}(t_i, \mathcal{F}_i) \right)^2 \right\|_p \leq M \sum_{j \in \mathbb{Z}} \left\| \mathcal{P}_j H^{(d)}(t_i, \mathcal{F}_i) \right\|_{2p}^2 = O(1),$$

where  $M$  is a sufficiently large constant.

Then by [Proposition C.2](#) and [Equation \(C.1\)](#), we have proved the desired result.  $\square$

**Lemma C.3.** *Let  $\psi_j$  denote  $\psi_j(d_n)$ . (a) For  $h = o(n)$ ,  $h \rightarrow \infty$  we have*

$$\sum_{l=0}^{\infty} \left( \sum_{j=l}^{h+l} \psi_j \right)^2 \sim \sum_{l=0}^{\infty} ((h+l)^{d_n} - l^{d_n})^2 = O(h/(\log h)).$$

(b) If we further assume  $h = \lfloor n^{\alpha_1} \rfloor$ ,  $\alpha_1 \in (0, 1)$ , we have

$$h^{-1} \sum_{l=0}^{h-1} \left( \sum_{j=0}^l \psi_j \right)^2 \rightarrow e^{2c\alpha_1}.$$

**Remark C.1.** *These two results correspond to the conclusions in Lemma 2 of [Shao and Wu \(2007\)](#).*

*Proof.* Proof of (a). We first show

$$\sum_{l=0}^{\infty} ((h+l)^{d_n} - l^{d_n})^2 = O(h/(\log h)).$$

Let  $N_1 = \lfloor h^{1-\alpha_h} \rfloor$ ,  $N_2 = \lfloor h^{1+\alpha_h} \rfloor$ ,  $\alpha_h = (\log h)^{-1} \log \log h$ . Then,  $h/N_1 = O(h^{\alpha_h}) = O(\log h)$ .  $N_1^{d_n} = O(1)$ ,  $N_2^{d_n} = O(1)$ . By [Lemma C.1](#) and Taylor's expansion, we have

$$\begin{aligned} \sum_{l=0}^{\infty} ((h+l)^{d_n} - l^{d_n})^2 &= \sum_{l=0}^{N_1} ((h+l)^{d_n} - l^{d_n})^2 + \sum_{l=N_1+1}^{N_2} ((h+l)^{d_n} - l^{d_n})^2 + \sum_{l=N_2+1}^{\infty} ((h+l)^{d_n} - l^{d_n})^2 \\ &= O(h(\log h)^{-1}). \end{aligned}$$

Then by [Lemma C.1](#), we have

$$\sum_{l=0}^{\infty} \left( \sum_{j=l}^{h+l} \psi_j \right)^2 = \sum_{l=0}^{\infty} ((h+l)^{d_n} - l^{d_n})^2 + O(d_n h (\log h)^{-1}).$$

Proof of (b). By [Lemma C.1](#), we have

$$\sum_{l=0}^{h-1} \left( \sum_{j=0}^l \psi_j \right)^2 = 1 + \sum_{l=1}^{h-1} l^{2d_n} + O\left(d_n \sum_{l=1}^{h-1} l^{2d_n}\right), \quad (\text{C.2})$$

where for  $h = O(n)$ ,

$$\sum_{l=1}^h l^{2d_n} = e^{2c\alpha_1} \sum_{l=1}^h (l/h)^{2d_n} = e^{2c\alpha_1} h \int_0^1 t^{2d_n} dt + O(1) = e^{2c\alpha_1} h / (2d_n + 1) + O(1) = e^{2c\alpha_1} h + O(1). \quad (\text{C.3})$$

Combining [\(C.2\)](#) and [\(C.3\)](#), we have shown the desired result. □

## References

- Andrews, D. (1991). Heteroskedasticity and autocorrelation consistent covariant matrix estimation. *Econometrica*, 59(3):817–858.
- Aue, A. and Horváth, L. (2013). Structural breaks in time series. *Journal of Time Series Analysis*, 34(1):1–16.
- Bai, L. and Wu, W. (2021). Detecting long-range dependence for time-varying linear models.
- Beran, J., Feng, Y., Ghosh, S., and Kulik, R. (2013). *Long-Memory Processes*. Springer.
- Brown, L. D. and Levine, M. (2007). Variance estimation in nonparametric regression via the difference sequence method. *The Annals of Statistics*, 35(5):2219 – 2232.
- Cavaliere, G., Nielsen, M. Ø., and Taylor, A. R. (2017). Quasi-maximum likelihood estimation and

- bootstrap inference in fractional time series models with heteroskedasticity of unknown form. *Journal of Econometrics*, 198(1):165–188.
- Chan, K. W. (2022). Optimal difference-based variance estimators in time series: A general framework. *The Annals of Statistics*, 50(3):1376–1400.
- Chan, S.-C. and Zhang, Z. (2010). Local polynomial modeling and variable bandwidth selection for time-varying linear systems. *IEEE Transactions on Instrumentation and Measurement*, 60(3):1102–1117.
- Dahlhaus, R. (1997). Fitting time series models to nonstationary processes. *Ann. Statist.*, 25(1):1–37.
- Dahlhaus, R., Richter, S., and Wu, W. B. (2019). Towards a general theory for nonlinear locally stationary processes. *Bernoulli*, 25(2):1013 – 1044.
- Dette, H. and Wu, W. (2019). Detecting relevant changes in the mean of nonstationary processes—a mass excess approach. *The Annals of Statistics*, 47(6):3578–3608.
- Dette, H., Wu, W., and Zhou, Z. (2019). Change point analysis of correlation in non-stationary time series. *Statistica Sinica*, 29(2):611–643.
- Hall, P., Kay, J. W., and Titterton, D. M. (1990). Asymptotically optimal difference-based estimation of variance in nonparametric regression. *Biometrika*, 77(3):521–528.
- He, X. and Zhu, L.-X. (2003). A lack-of-fit test for quantile regression. *Journal of the American Statistical Association*, 98(464):1013–1022.
- Hirukawa, M. (2021). Robust covariance matrix estimation in time series: A review. *Econometrics and Statistics*.
- Jansson, M. (2002). Consistent covariance matrix estimation for linear processes. *Econometric Theory*, 18(6):1449–1459.
- Juhl, T. and Xiao, Z. (2009). Tests for changing mean with monotonic power. *Journal of Econometrics*, 148(1):14–24.
- Kao, C., Trapani, L., and Urga, G. (2018). Testing for instability in covariance structures. *Bernoulli*, 24(1):740 – 771.
- Kokoszka, P. S. and Taqqu, M. S. (1995). Fractional arima with stable innovations. *Stochastic Processes and their Applications*, 60(1):19 – 47.
- Kulik, R. and Wichelhaus, C. (2012). Conditional variance estimation in regression models with long memory. *Journal of Time Series Analysis*, 33(3):468–483.
- Li, D., Chen, J., and Gao, J. (2011). Non-parametric time-varying coefficient panel data models with fixed effects. *The Econometrics Journal*, 14(3):387–408.

- Müller, H.-G. and Stadtmüller, U. (1987). Estimation of heteroscedasticity in regression analysis. *The Annals of Statistics*, 15(2):610–625.
- Nason, G. P., Von Sachs, R., and Kroisandt, G. (2000). Wavelet processes and adaptive estimation of the evolutionary wavelet spectrum. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 62(2):271–292.
- Newey, W. K. and West, K. D. (1987). A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica*, 55(3):703–708.
- Shao, X. and Wu, W. B. (2007). Local asymptotic powers of nonparametric and semiparametric tests for fractional integration. *Stochastic processes and their Applications*, 117(2):251–261.
- Su, L., Wang, X., and Jin, S. (2019). Sieve estimation of time-varying panel data models with latent structures. *Journal of Business & Economic Statistics*, 37(2):334–349.
- Tecuapetla-Gómez, I. and Munk, A. (2017). Autocovariance estimation in regression with a discontinuous signal and m-dependent errors: A difference-based approach. *Scandinavian Journal of Statistics*, 44(2):346–368.
- Vogt, M. and Dette, H. (2015). Detecting gradual changes in locally stationary processes. *The Annals of Statistics*, 43(2):713–740.
- Wu, W. and Zhou, Z. (2018). Gradient-based structural change detection for nonstationary time series M-estimation. *The Annals of Statistics*, 46(3):1197 – 1224.
- Wu, W. B. (2005). Nonlinear system theory: Another look at dependence. *Proceedings of the National Academy of Sciences*, 102(40):14150–14154.
- Wu, W. B. (2007). Strong invariance principles for dependent random variables. *The Annals of Probability*, 35(6):2294–2320.
- Wu, W. B. and Shao, X. (2006). Invariance principles for fractionally integrated nonlinear processes. In *Recent developments in nonparametric inference and probability*, pages 20–30. Institute of Mathematical Statistics.
- Wu, W. B. and Zhou, Z. (2011). Gaussian approximations for non-stationary multiple time series. *Statistica Sinica*, 21(3):1397–1413.
- Yu, Y., Wang, T., and Samworth, R. J. (2015). A useful variant of the davis–kahan theorem for statisticians. *Biometrika*, 102(2):315–323.
- Zhou, Z. (2014). Inference of weighted  $V$ -statistics for nonstationary time series and its applications. *The Annals of Statistics*, 42(1):87 – 114.
- Zhou, Z. and Shao, X. (2013). Inference for linear models with dependent errors. *Journal of the Royal Statistical Society. Series B (Statistical Methodology)*, 75(2):323–343.

Zhou, Z. and Wu, W. B. (2010). Simultaneous inference of linear models with time varying coefficients. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 72(4):513–531.