

# SPECTRAL GAP OF THE SYMMETRIC INCLUSION PROCESS

SEONWOO KIM AND FEDERICO SAU

ABSTRACT. We consider the symmetric inclusion process on a general finite graph. Our main result establishes universal upper and lower bounds for the spectral gap of this interacting particle system in terms of the spectral gap of the random walk on the same graph. In the regime in which the gamma-like reversible measures of the particle systems are log-concave, our bounds match, yielding a version for the symmetric inclusion process of the celebrated Aldous' spectral gap conjecture originally formulated for the interchange process. Finally, by means of duality techniques, we draw analogous conclusions for an interacting diffusion-like unbounded conservative spin system known as Brownian energy process.

## 1. INTRODUCTION, MODELS, AND MAIN RESULTS

Establishing spectral gap inequalities is a central topic in the analysis of convergence to equilibrium for large scale stochastic dynamics. In the context of interacting particle systems (e.g., [Lig05]), a spectral gap estimate with a sharp dependence on the system size grants a good control on the local mixing properties of the dynamics. This yields a number of applications, for instance, in scaling limits and metastability (e.g., [KL99, GJ14, LMS23] and references therein).

In this paper, we focus on spectral gap inequalities for a specific model, namely the *symmetric inclusion process* (SIP). This process was first introduced in [GKR07] as a particle analogue of statistical mechanics' models of interacting diffusions, and corresponds to a spatial version of the mean-field Moran model with mutation from population genetics (e.g., [Cor23]). SIP is also related to SEP, i.e., the *symmetric exclusion process* (e.g., [CLR10]): SEP's hard-core exclusion interaction is replaced here by attraction between particles.

SIP has been shown to exhibit interesting qualitative behaviors, such as condensation and metastability (e.g., [GRV11, GRV13, BDG17, JCG20, Kim21, KS21]); representations in terms of Lie algebras and limit theorems out of equilibrium have also been rigorously investigated (e.g., [GKRV09, FGS22, DSPS21]). The present work represents a first step in the quantitative analysis of convergence to equilibrium for this interacting particle system, together with the same analysis for a related diffusion-like unbounded spin system called the Brownian energy process.

**1.1. SIP and its spectral gap.** Let  $G = (V, (c_{xy})_{x,y \in V})$  be a connected weighted finite graph, with  $|V| = n$  sites and symmetric non-negative edge-weights, i.e.,  $c_{xy} = c_{yx} \geq 0$  for all  $x, y \in V$ . For any positive site-weights  $\alpha = (\alpha_x)_{x \in V}$  and  $k \in \mathbb{N}$  total number of particles, we let  $\text{SIP}_k(G, \alpha)$  be defined as the Markov process  $(\eta(t))_{t \geq 0}$  on the configuration space with  $k$  particles

$$\Xi_k := \{ \eta \in \mathbb{N}_0^V : |\eta| = k \} , \quad |\eta| := \sum_{x \in V} \eta_x ,$$

---

2020 *Mathematics Subject Classification.* Primary 60K35; secondary 60J27, 05C50.

*Key words and phrases.* Interacting particle systems; unbounded conservative spin systems; spectral gap; symmetric inclusion process; Brownian energy process.

and with infinitesimal generator  $L_k$  given, for all functions  $f \in \mathbb{R}^{\Xi_k}$ , by

$$L_k f(\eta) = \sum_{x \in V} \eta_x \sum_{y \in V} c_{xy} (\alpha_y + \eta_y) (f(\eta - \delta_x + \delta_y) - f(\eta)) , \quad \eta \in \Xi_k . \quad (1.1)$$

In this formula,  $\eta - \delta_x + \delta_y$  denotes the configuration in  $\Xi_k$  obtained from  $\eta$  (with  $\eta_x \geq 1$ ) by moving a particle from site  $x$  to  $y$ . We observe that, if  $(\alpha_y + \eta_y)$  were to be replaced by  $(1 - \eta_y)$ ,  $L_k$  in (1.1) would describe a SEP-dynamics (e.g., [CLR10]); while, if  $c_{xy} \equiv 1/n$ , we would obtain the Moran model with parent-independent mutation (e.g., [Cor23]).

For each  $k \in \mathbb{N}$ , due to the connectedness of the graph  $G$ , the generator  $L_k$  describes an irreducible Markov chain with a unique invariant measure, which we call  $\mu_{\alpha,k}$ . As a simple detailed balance computation shows, the particle system is actually reversible with respect to  $\mu_{\alpha,k}$ , which reads as follows:

$$\mu_{\alpha,k}(\eta) = \frac{1}{Z_{\alpha,k}} \prod_{x \in V} \frac{\Gamma(\alpha_x + \eta_x)}{\Gamma(\alpha_x) \eta_x!} , \quad \eta \in \Xi_k , \quad (1.2)$$

with

$$Z_{\alpha,k} := \frac{\Gamma(|\alpha| + k)}{\Gamma(|\alpha|) k!} , \quad |\alpha| := \sum_{x \in V} \alpha_x ,$$

being the normalization constant. Here,  $\Gamma(\beta)$  is the usual gamma function, for which we recall  $\Gamma(\beta + \ell)/\Gamma(\beta) = \beta(\beta + 1) \cdots (\beta + \ell - 1)$  for  $\beta > 0$  and  $\ell \in \mathbb{N}$ .

By the ergodic theorem for finite-state Markov chains, the law of the  $k$ -particle system converges in the long-run to the invariant measure  $\mu_{\alpha,k}$ . Moreover, since the process is reversible, the generator  $L_k$  has real and non-positive eigenvalues

$$-\lambda_{k,n-1} \leq \dots \leq -\lambda_{k,1} < -\lambda_{k,0} = 0 ,$$

and all admit a variational characterization. Among these, the spectral gap — namely, the second smallest eigenvalue of  $-L_k$  — measures the exponential rate of such a convergence.

In what follows, we let

$$\text{gap}_k(G, \alpha) := \lambda_{k,1}$$

denote the spectral gap of  $\text{SIP}_k(G, \alpha)$ . Our main goal is to estimate  $\text{gap}_k(G, \alpha)$  in terms of the spectral gap of the corresponding random walk on the same graph, i.e., the Markov process, referred to as  $\text{RW}(G, \alpha)$ , on  $V$  with generator  $A_\alpha$  acting on functions  $f \in \mathbb{R}^V$  as

$$A_\alpha f(x) = \sum_{y \in V} c_{xy} \alpha_y (f(y) - f(x)) , \quad x \in V .$$

Since the rate to jump from site  $x$  to  $y$  equals  $c_{xy} \alpha_y$ , detailed balance shows that the reversible measure for  $\text{RW}(G, \alpha)$  is proportional to  $\alpha = (\alpha_x)_{x \in V}$ .

We observe that  $\text{SIP}_k(G, \alpha)$  with just one particle ( $k = 1$ ) corresponds to a single random walk (thus, non-interacting) evolving like  $\text{RW}(G, \alpha)$  on the sites of  $G$ ; the non-trivial inclusion interaction between particles occurs as soon as  $k \geq 2$ . Let

$$\text{gap}_{\text{SIP}}(G, \alpha) := \inf_{k \geq 2} \text{gap}_k(G, \alpha) \quad \text{and} \quad \text{gap}_{\text{RW}}(G, \alpha) := \text{gap}_1(G, \alpha)$$

denote the spectral gaps of the interacting particle system and of the random walk, respectively. We are now ready to state our main result.

**Theorem 1.1.** *For every  $G = (V, (c_{xy})_{x,y \in V})$  and  $\alpha = (\alpha_x)_{x \in V}$ ,*

$$(1 \wedge \alpha_{\min}) \text{gap}_{\text{RW}}(G, \alpha) \leq \text{gap}_{\text{SIP}}(G, \alpha) \leq \text{gap}_{\text{RW}}(G, \alpha) ,$$

where  $\alpha_{\min} := \min_{x \in V} \alpha_x$ .

These bounds can be used to efficiently estimate the spectral gap of SIP in concrete examples. Remarkably, we observe that the inequalities in Theorem 1.1 saturate to identities as soon as  $\alpha_{\min} \geq 1$  (which is equivalent to the log-concavity of  $\mu_{\alpha,k}$ ), yielding the following spectral gaps' identity:

**Corollary 1.2.** *For every  $G = (V, (c_{xy})_{x,y \in V})$  and  $\alpha = (\alpha_x)_{x \in V}$  such that  $\alpha_{\min} \geq 1$ ,*

$$\text{gap}_{\text{SIP}}(G, \alpha) = \text{gap}_{\text{RW}}(G, \alpha) . \quad (1.3)$$

The result in Corollary 1.2 may be interpreted as a SIP's version of the celebrated *Aldous' spectral gap conjecture*, originally formulated for the interchange process and SEP, and recently solved by Caputo *et al.* in [CLR10]. An identity of the type (1.3) may be viewed as an exact tensorization of the Poincaré inequality over the  $k$  components, a property which trivially holds when considering  $k$  independent particles. Such an identity is, in general, not expected to hold for truly interacting systems. This property — apart from situations in which the spectrum is fully explicit (e.g., [Shi77, Gri14, Cor23]) — has been established on *any graph* only for a few examples other than the interchange process and SEP:

- SEP in contact with equilibrium reservoirs in [Sal23];
- the binomial splitting process in [QS23] (see also [BC23]).

Corollary 1.2 proves that SIP with  $\alpha_{\min} \geq 1$  also satisfies this exact tensorization property. See also Section 1.3.2 for a more detailed discussion on Aldous' spectral gap conjecture and related work.

**1.2. BEP and its spectral gap.** The Brownian energy process (BEP) is an interacting system of continuous spins placed on the sites of a graph (e.g., [GKRV09]). This process falls into the larger class of *unbounded conservative spin systems*, and is intimately related to SIP. The spins (or, energies) evolve as diffusions. Moreover, the dynamics preserves the total amount of energy of the system, and is reversible with respect to measures associated to gamma distributions. Just like SIP and the Moran model are related, BEP on the complete graph corresponds to the multi-type Wright-Fisher diffusion with mutation.

Let us now describe the model more formally. Given a graph  $G$  and site-weights  $\alpha = (\alpha_x)_{x \in V}$ ,  $\text{BEP}(G, \alpha)$  is the diffusion process  $(\zeta(t))_{t \geq 0}$  on  $[0, \infty)^V$ , and whose infinitesimal evolution is described by the following generator:

$$\mathcal{L} = \frac{1}{2} \sum_{x,y \in V} c_{xy} \left\{ -(\alpha_y \zeta_x - \alpha_x \zeta_y) (\partial_{\zeta_x} - \partial_{\zeta_y}) + \zeta_x \zeta_y (\partial_{\zeta_x} - \partial_{\zeta_y})^2 \right\} . \quad (1.4)$$

The diffusion  $(\zeta(t))_{t \geq 0}$  admits  $\nu_\theta := \otimes_{x \in V} \text{Gamma}(\alpha_x, \theta)$ ,  $\theta > 0$ , as a one-parameter family of reversible product measures, fully supported on  $[0, \infty)^V$ . However, all features of the system are well-captured by the dynamics which only considers configurations with unit total energy: on the one side, applying the generator  $\mathcal{L}$  to the function  $\zeta \mapsto |\zeta| := \sum_{x \in V} \zeta_x$  shows that the dynamics conserves the total energy of the system; on the other side, a simple scaling argument demonstrates that the action of  $\mathcal{L}$  does not depend on  $|\zeta|$ . Therefore, all throughout, it suffices to consider  $\zeta(t)$  as evolving on  $\Delta_V$ , the simplex of probability measures on  $V$ , for which  $\pi := \nu_\theta(\cdot \mid \zeta \in \Delta_V)$ ,  $\theta > 0$ , is reversible, and given by

$$\pi(d\zeta) = \left( \frac{1}{B(\alpha)} \prod_{x \in V} \zeta_x^{\alpha_x - 1} \right) d\zeta , \quad \text{with } \zeta \in \Delta_V , B(\alpha) := \frac{\prod_{x \in V} \Gamma(\alpha_x)}{\Gamma(|\alpha|)} ,$$

where  $d\zeta$  denotes the uniform measure on  $\Delta_V$ . Note that  $\pi$  is independent of  $\theta > 0$ .

Quantifying the exponential rate of convergence to equilibrium goes through a spectral analysis of the generator  $\mathcal{L}$  on  $L^2(\Delta_V, \pi)$ . Since  $\mathcal{L}$  is self-adjoint on  $L^2(\Delta_V, \pi)$ , its spectrum is real. Moreover, as we will show in Section 4,  $-\mathcal{L}$  is non-negative and has a pure-point spectrum, with only one zero eigenvalue corresponding to the constant eigenfunction.

In our next result, we provide an analogue of Theorem 1.1 for  $\text{gap}_{\text{BEP}}(G, \alpha) > 0$ , the smallest non-zero eigenvalue of  $-\mathcal{L}$  on  $L^2(\Delta_V, \pi)$ .

**Theorem 1.3.** *For every  $G = (V, (c_{xy})_{x,y \in V})$  and  $\alpha = (\alpha_x)_{x \in V}$ ,*

$$(1 \wedge \alpha_{\min}) \text{gap}_{\text{RW}}(G, \alpha) \leq \text{gap}_{\text{BEP}}(G, \alpha) \leq \text{gap}_{\text{RW}}(G, \alpha) .$$

Hence, provided  $\alpha_{\min} \geq 1$ ,

$$\text{gap}_{\text{BEP}}(G, \alpha) = \text{gap}_{\text{RW}}(G, \alpha) . \tag{1.5}$$

### 1.3. Related work, proof strategy, and open problems.

1.3.1. *Functional inequalities and comparison techniques.* Functional inequalities play a major role in PDE and probability theory, and several approaches have been developed for this purpose. Comparison techniques (e.g., [SC97, MT06]) are among the most robust and well-established ones, and proved to be especially effective when estimating spectral gaps (or Poincaré constants, their inverses), log-Sobolev constants, and Nash inequalities.

In the more specific context of interacting particle systems and unbounded spin systems subjected to conservation laws, comparing Dirichlet forms is key in the so-called martingale method and its variants (e.g., [Qua92, LY93, DSC93, LSV96, JLQY99, LPY02, Cap03, Cap04, Cap08, Sas13] and references therein). In a nutshell, this strategy compares the system’s Dirichlet form on the geometry of interest (e.g., SEP on  $\mathbb{Z}^d$ -boxes of size  $\ell$ ), to that on more tractable geometries (e.g., the complete graph), and finally transfers the gained information back through a path counting argument. In most examples, this method captures the correct dependence on the size  $\ell$  of the system, but the universal prefactor is typically not optimal (e.g., it deteriorates with  $d$ , the dimension of the box).

1.3.2. *Aldous’ spectral gap conjecture and related examples.* Sharper identities like the one expressed in Aldous’ spectral gap conjecture [CLR10] holding true on general graphs have been verified only for a handful of models (as already discussed below Corollary 1.2), each of these examples requiring *ad hoc* proof arguments.

The proof in [CLR10] for the interchange process and SEP combines in a non-trivial way a nonlinear network reduction and a hard correlation inequality (which became known as *Octopus inequality*). The first ingredient allows an induction argument on  $n$ , the size of the graph, well-compatible with the particle-hole symmetry of SEP. Such a symmetry (or, alternatively, the fact that SEP may be obtained as a projection of the interchange process) is a property that seems to be lacking for SIP.

Negative dependence — a form of negativity of correlations of all orders — is nicely exploited in [Sal23] for the non-conservative reversible SEP. We remark that SIP is positive (rather than negative) dependent, see, e.g., [GRV10, FRS22].

The arguments in [QS23] for the binomial splitting process build on a  $L^2$ -contraction inequality established in [AL12] for the “dual” averaging process. BEP is one of the continuum duals of SIP (e.g., [GKRV09]); nevertheless, the analogue of such a contraction estimate is not known for BEP.

1.3.3. *Discussion on proof strategy.* Our approach combines in an elementary way two main ingredients:

- self-duality of the interacting particle system;
- comparison inequalities.

Self-duality/consistency of SIP (cf. (2.2) and (3.3) below) ensures a certain rigidity and structure of eigenvalues and eigenfunctions. This immediately yields the upper bound in Theorem 1.1 (Section 2), and this is what allows to effectively set off an induction argument on  $k$ , the total number of particles (rather than  $n$ , the size of the graph, as in [CLR10]), for the lower bound in Section 3. In closing the proof of the induction step from  $k - 1$  to  $k$ , we employ comparison inequalities. Inspired by the recent work [HS19] on zero-range dynamics, in Lemma 3.2 we rearrange SIP’s Dirichlet form so to reduce our task to an estimate of the spectral gap not of the whole system, but rather of the  $k$ th particle only, uniformly over the positions of the remaining  $k - 1$  particles. Finally, the min-max theorem for eigenvalues (e.g., [SC97, Theorem 1.2.10]) yields the desired single-particle spectral gap inequality (Lemma 3.3). For this step, we compare both Dirichlet forms and  $L^2$ -norms of such a particle to those of a non-interacting walk. We emphasize that, although the estimate that we get in Lemma 3.3 deteriorates by a factor  $k$ , the reduction applied in Lemma 3.2 returns back exactly the same factor  $k$ , thus, allows us to conclude the proof for SIP’s spectral gap estimates.

The analogous result for BEP is derived by remapping SIP into BEP via an intertwining relation, translating all spectral information from the particle to the diffusion system.

1.3.4. *Spectral gap identity and Gibbs samplers.* Focusing on interacting systems on more specific graphs, a spectral gap identity in the spirit of that in (1.3) has been proved on 1D geometries also for two models of continuous spins with Gibbs sampler dynamics in [CLL20, CLL22]. Especially the first one of these models studied in [CLL20] is relevant for the present work. Indeed, when considered on a general graph  $G = (V, (c_{xy})_{x,y \in V})$  and with general site-weights  $\alpha = (\alpha_x)_{x \in V}$ , it corresponds to the Gibbs sampler of  $\text{BEP}(G, \alpha)$ : instead of letting energies diffuse as in BEP, the energies are instantaneously set to their local beta-like equilibrium among randomly chosen edges.

While our Theorem 1.3 shows that, as soon as  $\alpha_{\min} \geq 1$ , (1.5) for BEP holds on *any graph*, rather surprisingly an analogous identity for its Gibbs sampler version fails on specific geometries. Indeed, the proof in [CLL20] crucially relies on the one-dimensional structure of the model, which grants a monotonicity property of the spectral gap’s eigenfunction (similar to that for birth-and-death chains). However, as discussed in [CLL20, Remark 4], the mean-field version of the same model provides a counterexample to such an identity.

It is well-known (e.g., [GKRV09, Section 6.3]) that, at least for symmetric systems, this procedure of “instantaneous thermalization” among edges does not alter qualitative properties of the system as, e.g., the form of the reversible measures and the richness of the duality relations. Nonetheless, it does affect dramatically the eigenstructure of the processes and other quantitative features determining convergence to equilibrium, as the comparison between the model in [CLL20] and our result on BEP illustrates. We emphasize that this example also shows that (self-)duality does not guarantee *per se* the validity of a spectral gap identity.

1.3.5. *Open problems.* Besides the problem of quantifying the sensitivity with respect to Gibbs-sampler perturbations of the model (as discussed in, e.g., [CLL20, Section 1.2]; see also the previous paragraph), settling the role of the threshold  $\alpha_{\min} = 1$  remains open.

More specifically, our results provide only partial answers in the regime

$$\alpha_{\min} \in (0, 1) .$$

This regime corresponds, roughly speaking, to the case in which particle/energy interaction becomes predominant over the mechanism of independent diffusion. Here, our results state that a spectral gap comparison is robust over the underlying geometry  $G = (V, (c_{xy})_{x,y \in V})$ , but do not say anything about the sharpness of the first-order dependence on the parameter  $\alpha_{\min} \in (0, 1)$ . We emphasize that such a threshold appears also in other related works, e.g., [CLL20, pp. 2453–4], as well as [BW09, LP22], and also there sharp results are not available.

When  $\alpha_{\min} \in (0, 1)$ , we observe that our proof techniques fail to give a lower bound for  $\text{gap}_{\text{SIP}}$  matching the upper bound. This can be checked already on simple geometries with  $n = 3$  or  $n = 4$  sites, in which the lower bound in Lemma 3.3 is essentially sharp, thus, cannot be further improved. However, even in those examples in which such estimates fail, a direct inspection with small values of  $k \in \mathbb{N}$  confirms the validity of (1.3).

Finally, we recall that on the complete graph ( $c_{xy} \equiv 1/n$ ) the spectrum of  $-L_k$  is fully explicit and given (without counting multiplicities) by

$$\frac{\ell}{n} (|\alpha| + \ell - 1) , \quad \ell = 0, 1, \dots, k .$$

Hence, the spectral gap identity in (1.3) holds true for all positive site-weights  $\alpha = (\alpha_x)_{x \in V}$  in this mean-field setting. We conclude by remarking that a non-trivial metastable picture of SIP emerges from the asymptotic regime  $\alpha_x \equiv \alpha \rightarrow 0$  only on the timescale  $\alpha^{-1}$  and for  $k$ -particle systems with  $\alpha \log k = o(1)$  ([GRV13, BDG17]). Since  $\text{gap}_{\text{RW}}(\alpha) \approx \alpha^{-1}$  as  $\alpha \rightarrow 0$ , this suggests — at least at the heuristic level — that  $\text{gap}_{\text{SIP}}(\alpha) \approx \text{gap}_{\text{RW}}(\alpha)$ . If this were true, the lower bound in Theorem 1.1 would not capture this.

**1.4. Structure of the paper.** The rest of the paper is organized as follows. The upper bound in Theorem 1.1 is proved in Section 2. The proof of the lower bound in Theorem 1.1 occupies the whole Section 3, which is further divided into four subsections. In Section 4, we present the proof of Theorem 1.3.

## 2. PROOF OF UPPER BOUND IN THEOREM 1.1

In the remainder of the article, since the graph  $G$  is fixed, we may drop the dependence on  $G$  for the objects. For example, we express

$$\text{gap}_{\text{SIP}}(G, \alpha) = \text{gap}_{\text{SIP}}(\alpha) \quad \text{and} \quad \text{gap}_{\text{RW}}(G, \alpha) = \text{gap}_{\text{RW}}(\alpha) .$$

We start with establishing the upper bound in Theorem 1.1. The simple idea underlying its proof is that, as in the case of SEP and other systems enjoying a suitable form of consistency/self-duality, observables of a few particles may be “lifted” to observables of many particles, yet yielding coherent statistics. In particular, eigenfunctions for  $\text{RW}(\alpha)$  “lift” to eigenfunctions for  $\text{SIP}_k(\alpha)$ ; this is rigorously demonstrated in Lemma 2.3 below.

For each  $k \in \mathbb{N}$ , the *annihilation operator*  $\mathbf{a}_k : \mathbb{R}^{\Xi_{k-1}} \rightarrow \mathbb{R}^{\Xi_k}$  is defined, for  $g \in \mathbb{R}^{\Xi_{k-1}}$  and  $\eta \in \Xi_k$ , as

$$\mathbf{a}_k g(\eta) := \sum_{x \in V} \eta_x g(\eta - \delta_x) ,$$

where  $\mathbb{R}^\emptyset$  is conventionally understood as the space of constants. Intuitively,  $\mathbf{a}_k g$  evaluates the value at  $\eta \in \Xi_k$  by summing up all the values of  $g$  evaluated at a  $(k-1)$ -particle configuration chosen inside  $\eta$  uniformly at random. This motivates also to say that  $\mathbf{a}_k$

corresponds to the operation of “removing a particle uniformly at random”. Moreover, it holds, for every  $k > \ell \in \mathbb{N}$ ,  $g \in \mathbb{R}^{\Xi_\ell}$ , and  $\eta \in \Xi_k$ , that

$$(\mathbf{a}_k \circ \cdots \circ \mathbf{a}_{\ell+1})g(\eta) = \sum_{\zeta \in \Xi_\ell} \left( \prod_{x \in V} \binom{\eta_x}{\zeta_x} \right) g(\zeta) . \quad (2.1)$$

Indeed, the left-hand side of (2.1) can be calculated by summing up the values of  $g$  evaluated at a  $\ell$ -particle configuration chosen from  $\eta$  uniformly, which is exactly the right-hand side of (2.1).

In this section, we use two important properties of  $\mathbf{a}_k$ ,  $k \in \mathbb{N}$ , which can be easily verified:

- the operator  $\mathbf{a}_k : \mathbb{R}^{\Xi_{k-1}} \rightarrow \mathbb{R}^{\Xi_k}$  is injective;
- it holds that

$$\mathbf{a}_k L_{k-1} = L_k \mathbf{a}_k . \quad (2.2)$$

Especially, (2.2) implies that removing a particle at random first and then running the system (right-hand side) is equivalent, in distribution, to running the system first and then removing a particle at random (left-hand side).

**Remark 2.1.** *The concept in  $\mathbf{a}_k : \mathbb{R}^{\Xi_{k-1}} \rightarrow \mathbb{R}^{\Xi_k}$  of removing a particle uniformly at random is essential in SIP. This is not the case in other related models such as the interchange process or the binomial splitting process. In these examples, the dynamics restricted to a subset of labeled particles is still Markovian and of the same type as the larger system. Thus, therein, one can fix a specific particle and then lift the remaining particle configuration. Such a property holds for SIP not for all subsets of particles, but only for subsets chosen uniformly at random.*

The identity (2.2) has the following consequence. Suppose that  $-L_{k-1}g = \lambda g$  holds for some  $\lambda \in \mathbb{R}$  and non-zero  $g \in \mathbb{R}^{\Xi_{k-1}}$ . Then,

$$-L_k(\mathbf{a}_k g) = \mathbf{a}_k(-L_{k-1}g) = \mathbf{a}_k(\lambda g) = \lambda \mathbf{a}_k g ,$$

so that the new function  $\mathbf{a}_k g \in \mathbb{R}^{\Xi_k}$ , which is non-zero since  $\mathbf{a}_k$  is injective, is an eigenfunction of  $-L_k$  subjected to the same eigenvalue  $\lambda$ . Thus, the operator  $\mathbf{a}_k$  lifts the eigenspace of  $-L_{k-1}$  to the eigenspace of  $-L_k$ . Then, inductively, the composition  $\mathbf{a}_k \circ \cdots \circ \mathbf{a}_{\ell+1} : \mathbb{R}^{\Xi_\ell} \rightarrow \mathbb{R}^{\Xi_k}$  lifts the eigenspace of the operator  $-L_\ell$  to the eigenspace of  $-L_k$  for all  $\ell < k \in \mathbb{N}$ . Since all the eigenvalues of  $-L_\ell$  are also eigenvalues of  $-L_k$ , it holds in particular that

$$\text{gap}_k(\alpha) \leq \text{gap}_\ell(\alpha) .$$

Considering the special case  $\ell = 1$  and taking the infimum over all  $k \geq 2$  in the left-hand side, we have verified the upper bound in Theorem 1.1:

**Theorem 2.2** (Upper bound). *For every  $\alpha = (\alpha_x)_{x \in V}$ ,  $\text{gap}_{\text{SIP}}(\alpha) \leq \text{gap}_{\text{RW}}(\alpha)$ .*

Before concluding this section, we record a lemma which presents the eigenfunction of  $-L_k$  lifted from the original eigenfunction of  $-A_\alpha$  subjected to the same eigenvalue.

**Lemma 2.3.** *Let  $\psi : V \rightarrow \mathbb{R}$  be an eigenfunction for  $-A_\alpha$  with eigenvalue  $\lambda \geq 0$ . Then, for every  $k \in \mathbb{N}$ , the function  $f_{\psi,k} \in \mathbb{R}^{\Xi_k}$  given by*

$$f_{\psi,k}(\eta) := \sum_{x \in V} \psi(x) \eta_x , \quad \eta \in \Xi_k ,$$

*is an eigenfunction for  $-L_k$  with the same eigenvalue  $\lambda \geq 0$ .*

*Proof.* Define  $g \in \mathbb{R}^{\Xi_1}$  as  $g(\delta_x) := \psi(x)$ . It is clear that  $g$  becomes an eigenfunction for  $-L_1$  with eigenvalue  $\lambda \geq 0$ . Substituting  $\ell = 1$  in (2.1), we obtain that

$$(\mathbf{a}_k \circ \cdots \circ \mathbf{a}_2)g(\eta) = \sum_{x \in V} \eta_x g(\delta_x) = \sum_{x \in V} \eta_x \psi(x) ,$$

so that we have

$$f_{\psi,k} = (\mathbf{a}_k \circ \cdots \circ \mathbf{a}_2)g . \quad (2.3)$$

Since  $g \neq 0$  (which follows from the fact that  $\psi$  is an eigenfunction) and the operators  $\mathbf{a}_2$  through  $\mathbf{a}_k$  are all injective,  $f_{\psi,k}$  is a non-zero function.

It remains to verify that  $-L_k f_{\psi,k} = \lambda f_{\psi,k}$  holds. This is an easy consequence of (2.3) and the intertwining relation (2.2):

$$-L_k f_{\psi,k} = -L_k(\mathbf{a}_k \circ \cdots \circ \mathbf{a}_2)g = (\mathbf{a}_k \circ \cdots \circ \mathbf{a}_2)(-L_1 g) = (\mathbf{a}_k \circ \cdots \circ \mathbf{a}_2)(\lambda g) = \lambda f_{\psi,k} .$$

Thus, we conclude the proof.  $\square$

### 3. PROOF OF LOWER BOUND IN THEOREM 1.1

In this section, we tackle the lower bound in Theorem 1.1; namely, we prove

$$(1 \wedge \alpha_{\min}) \text{gap}_{\text{RW}}(\alpha) \leq \text{gap}_{\text{SIP}}(\alpha) . \quad (3.1)$$

**3.1. Preliminaries.** In Section 2, we demonstrated that the eigenfunctions of the operator  $-L_{k-1}$  are lifted via  $\mathbf{a}_k$  to the eigenfunctions of the operator  $-L_k$ , thereby providing the eigenvalues of  $-L_{k-1}$  as also eigenvalues of  $-L_k$ . Hence, to find the remaining eigenvalues of  $-L_k$  that do not come from the lifting property of  $\mathbf{a}_k$ , since  $-L_k$  is self-adjoint, it suffices to investigate functions  $f \in L^2(\mu_{\alpha,k})$  (cf. (3.2)) that belong to the orthogonal complement of the image of the operator  $\mathbf{a}_k$ . Since  $\mathbf{a}_k$  injectively takes all the functions in  $\mathbb{R}^{\Xi_{k-1}}$  into  $\mathbb{R}^{\Xi_k}$ , such an  $f$  should satisfy a certain mean-zero condition subjected to each  $(k-1)$ -particle configuration:

$$\sum_{x \in V} (\xi_x + \alpha_x) f(\xi + \delta_x) = 0 , \quad \xi \in \Xi_{k-1} .$$

In this sense, it is natural to define, for  $k \in \mathbb{N}$ , the *creation operator*  $\mathbf{a}_{k-1}^\dagger : \mathbb{R}^{\Xi_k} \rightarrow \mathbb{R}^{\Xi_{k-1}}$  as follows: for all  $f \in \mathbb{R}^{\Xi_k}$  and  $\xi \in \Xi_{k-1}$ ,

$$\mathbf{a}_{k-1}^\dagger f(\xi) := \sum_{x \in V} (\xi_x + \alpha_x) f(\xi + \delta_x) ,$$

where, again,  $\mathbb{R}^\emptyset$  is considered as the space of constants.

It turns out that the two operators  $\mathbf{a}_k$  and  $\mathbf{a}_{k-1}^\dagger$  are indeed closely related to each other, as the following proposition shows. For two functions  $f, g \in \mathbb{R}^{\Xi_k}$ , we define the inner product  $\langle f | g \rangle_{\alpha,k}$  as

$$\langle f | g \rangle_{\alpha,k} := \sum_{\eta \in \Xi_k} \mu_{\alpha,k}(\eta) f(\eta) g(\eta) , \quad (3.2)$$

and let  $L^2(\mu_{\alpha,k})$  denote the corresponding  $L^2$ -space of functions on  $\Xi_k$ .

**Proposition 3.1.** *The following two properties are valid for all  $k \in \mathbb{N}$ :*

(a) (adjoint property) for all  $f \in \mathbb{R}^{\Xi_k}$  and  $g \in \mathbb{R}^{\Xi_{k-1}}$ ,

$$\langle \mathbf{a}_k g | f \rangle_{\alpha,k} = \frac{k}{|\alpha| + k - 1} \langle g | \mathbf{a}_{k-1}^\dagger f \rangle_{\alpha,k-1} ;$$

(b) (orthogonal decomposition)  $L^2(\mu_{\alpha,k}) = \text{Im } \mathbf{a}_k \oplus_\perp \text{Ker } \mathbf{a}_{k-1}^\dagger$ .



*Proof.* We fix  $f \in \mathbb{R}^{\Xi_k}$  and  $g \in \mathbb{R}^{\Xi_{k-1}}$ . Then,

$$\langle \mathbf{a}_k g | f \rangle_{\alpha, k} = \sum_{\eta \in \Xi_k} \mu_{\alpha, k}(\eta) \mathbf{a}_k g(\eta) f(\eta) = \sum_{x \in V} \sum_{\eta \in \Xi_k: \eta_x \geq 1} \mu_{\alpha, k}(\eta) \eta_x f(\eta) g(\eta - \delta_x).$$

Rearranging by substituting  $\xi := \eta - \delta_x$ , this becomes

$$\begin{aligned} & \sum_{x \in V} \sum_{\xi \in \Xi_{k-1}} \mu_{\alpha, k}(\xi + \delta_x) (\xi_x + 1) f(\xi + \delta_x) g(\xi) \\ &= \sum_{x \in V} \sum_{\xi \in \Xi_{k-1}} \frac{k(\alpha_x + \xi_x)}{|\alpha| + k - 1} \mu_{\alpha, k-1}(\xi) f(\xi + \delta_x) g(\xi), \end{aligned}$$

where in the equality we used (1.2). Thus, applying the definitions of  $\mathbf{a}_{k-1}^\dagger$  and  $\langle \cdot | \cdot \rangle_{\alpha, k-1}$ , the right-hand side equals

$$\sum_{\xi \in \Xi_{k-1}} \frac{k}{|\alpha| + k - 1} \mu_{\alpha, k-1}(\xi) \mathbf{a}_{k-1}^\dagger f(\xi) g(\xi) = \frac{k}{|\alpha| + k - 1} \langle g | \mathbf{a}_{k-1}^\dagger f \rangle_{\alpha, k-1},$$

which concludes the proof of part (a).

(b) Suppose that  $f \in \text{Im } \mathbf{a}_k$  and  $g \in \text{Ker } \mathbf{a}_{k-1}^\dagger$ . Then, by part (a), since  $f = \mathbf{a}_k h$  for some  $h \in \mathbb{R}^{\Xi_{k-1}}$ ,

$$\langle f | g \rangle_{\alpha, k} = \langle \mathbf{a}_k h | g \rangle_{\alpha, k} = \frac{k}{|\alpha| + k - 1} \langle h | \mathbf{a}_{k-1}^\dagger g \rangle_{\alpha, k-1} = 0,$$

where the last equality holds since  $g \in \text{Ker } \mathbf{a}_{k-1}^\dagger$ . This proves that  $\text{Im } \mathbf{a}_k$  and  $\text{Ker } \mathbf{a}_{k-1}^\dagger$  are orthogonal. Moreover,  $\dim \text{Im } \mathbf{a}_k = |\Xi_{k-1}|$  since  $\mathbf{a}_k$  is injective, and  $\dim \text{Ker } \mathbf{a}_{k-1}^\dagger \geq |\Xi_k| - |\Xi_{k-1}|$  by the dimension theorem. Thus, by orthogonality, we conclude that  $\dim \text{Ker } \mathbf{a}_{k-1}^\dagger = |\Xi_k| - |\Xi_{k-1}|$ , and that  $L^2(\mu_{\alpha, k}) = \text{Im } \mathbf{a}_k \oplus_\perp \text{Ker } \mathbf{a}_{k-1}^\dagger$ .  $\square$

A simple consequence of the previous proposition and (2.2) is that the following identity holds: for  $k \in \mathbb{N}$ ,

$$\mathbf{a}_{k-1}^\dagger L_k = L_{k-1} \mathbf{a}_{k-1}^\dagger. \quad (3.3)$$

Indeed, for all  $f \in \mathbb{R}^{\Xi_k}$  and  $g \in \mathbb{R}^{\Xi_{k-1}}$ , we calculate using part (a) of Proposition 3.1 as

$$\langle g | \mathbf{a}_{k-1}^\dagger L_k f \rangle_{\alpha, k-1} = \frac{|\alpha| + k - 1}{k} \langle \mathbf{a}_k g | L_k f \rangle_{\alpha, k} = \frac{|\alpha| + k - 1}{k} \langle L_k \mathbf{a}_k g | f \rangle_{\alpha, k},$$

where the second identity holds since  $L_k$  is self-adjoint on  $L^2(\mu_{\alpha, k})$ . Then, by (2.2) and again by part (a) of Proposition 3.1, this equals

$$\frac{|\alpha| + k - 1}{k} \langle \mathbf{a}_k L_{k-1} g | f \rangle_{\alpha, k} = \langle L_{k-1} g | \mathbf{a}_{k-1}^\dagger f \rangle_{\alpha, k-1} = \langle g | L_{k-1} \mathbf{a}_{k-1}^\dagger f \rangle_{\alpha, k-1},$$

where the last equality follows from the fact that  $L_{k-1}$  is self-adjoint on  $L^2(\mu_{\alpha, k-1})$ . Thus, we have proved that

$$\langle g | \mathbf{a}_{k-1}^\dagger L_k f \rangle_{\alpha, k-1} = \langle g | L_{k-1} \mathbf{a}_{k-1}^\dagger f \rangle_{\alpha, k-1}$$

holds for all  $g \in \mathbb{R}^{\Xi_{k-1}}$  and  $f \in \mathbb{R}^{\Xi_k}$ , which indeed implies (3.3).

According to part (b) of Proposition 3.1, we easily obtain the following fact:

$$-L_k f = \lambda f \quad \text{if and only if} \quad f \in \text{Im } \mathbf{a}_k \text{ or } f \in \text{Ker } \mathbf{a}_{k-1}^\dagger. \quad (3.4)$$

**3.2. Decomposition of Dirichlet forms.** For  $k \in \mathbb{N}$ , we define the *Dirichlet form*  $\mathcal{E}_{\alpha,k}(f)$  evaluated at  $f \in \mathbb{R}^{\Xi_k}$  as

$$\mathcal{E}_{\alpha,k}(f) := \langle f | -L_k f \rangle_{\alpha,k} .$$

Moreover, we let  $\mathcal{D}_\alpha(\phi)$  denote the Dirichlet form at  $\phi \in \mathbb{R}^V$  subjected to  $\text{RW}(\alpha)$ :

$$\mathcal{D}_\alpha(\phi) := \sum_{x \in V} \frac{\alpha_x}{|\alpha|} \phi(x) (-A_\alpha \phi)(x) = \frac{1}{|\alpha|} \sum_{x,y \in V} \alpha_x \alpha_y c_{xy} \phi(x) (\phi(x) - \phi(y)) . \quad (3.5)$$

Then, we have the following variational representation of the spectral gap (e.g., [SC97]):

$$\text{gap}_k(\alpha) = \inf_{f \in \mathbb{R}^{\Xi_k}: f \neq \text{const.}} \frac{\mathcal{E}_{\alpha,k}(f)}{\text{Var}_{\alpha,k}(f)} , \quad (3.6)$$

where  $\text{Var}_{\alpha,k}(f) := \langle f | f \rangle_{\alpha,k} - \langle f \rangle_{\alpha,k}^2$ , with  $\langle f \rangle_{\alpha,k} := \langle f | 1 \rangle_{\alpha,k}$ . Similarly, it holds that

$$\text{gap}_{\text{RW}}(\alpha) = \inf_{\phi \in \mathbb{R}^V: \phi \neq \text{const.}} \frac{\mathcal{D}_\alpha(\phi)}{\text{var}_\alpha(\phi)} , \quad (3.7)$$

where  $\text{var}_\alpha(\phi) := \langle \phi | \phi \rangle_{L^2(\alpha)} - \langle \phi \rangle_{L^2(\alpha)}^2$ , with  $\langle \phi \rangle_{L^2(\alpha)} := \langle \phi | 1 \rangle_{L^2(\alpha)}$ . Here,  $L^2(\alpha)$  denotes the  $L^2$  function space on  $V$  with respect to the probability measure  $(\alpha_x/|\alpha|)_{x \in V}$ .

Suppose that  $f \in \text{Ker } \mathbf{a}_{k-1}^\dagger$  for  $k \in \mathbb{N}$ . Then, since  $\mathbf{a}_k 1(\eta) = \sum_{x \in V} \eta_x = k$ , we calculate

$$\langle f | 1 \rangle_{\alpha,k} = \frac{1}{k} \langle f | \mathbf{a}_k 1 \rangle_{\alpha,k} = \frac{1}{|\alpha| + k - 1} \langle \mathbf{a}_{k-1}^\dagger f | 1 \rangle_{\alpha,k-1} = 0 ,$$

where the second equality holds by part (a) of Proposition 3.1. This implies that the expectation of  $f$  with respect to  $\mu_{\alpha,k}$  is zero, and thus

$$\text{Var}_{\alpha,k}(f) = \sum_{\eta \in \Xi_k} \mu_{\alpha,k}(\eta) f(\eta)^2 . \quad (3.8)$$

In this subsection, we prove the following lemma, which is partially motivated from [HS19]. The idea of proof is to decompose on  $\text{Ker } \mathbf{a}_{k-1}^\dagger$  the  $k$ -Dirichlet form  $\mathcal{E}_{\alpha,k}(\cdot)$  into lower-order Dirichlet forms  $\mathcal{D}_\beta(\cdot)$  for some suitably chosen  $\beta = \beta(\alpha)$ .

**Lemma 3.2.** *Suppose that  $f \in \text{Ker } \mathbf{a}_{k-1}^\dagger$ . Then, it holds that*

$$\mathcal{E}_{\alpha,k}(f) \geq k \left( \inf_{\xi \in \Xi_{k-1}} \text{gap}_{\text{RW}}(\alpha + \xi) \right) \text{Var}_{\alpha,k}(f) .$$

*Proof.* For  $k \in \mathbb{N}$  and  $f \in \text{Ker } \mathbf{a}_{k-1}^\dagger$ , we calculate  $\mathcal{E}_{\alpha,k}(f) = \langle f | -L_k f \rangle_{\alpha,k}$  as

$$\begin{aligned} & \sum_{\eta \in \Xi_k} \sum_{x,y \in V} \mu_{\alpha,k}(\eta) f(\eta) \eta_x c_{xy} (\alpha_y + \eta_y) (f(\eta) - f(\eta - \delta_x + \delta_y)) \\ &= \sum_{x \in V} \left( \sum_{\eta \in \Xi_k: \eta_x \geq 1} \mu_{\alpha,k}(\eta) \eta_x f(\eta) \sum_{y \in V} c_{xy} (\alpha_y + \eta_y) (f(\eta) - f(\eta - \delta_x + \delta_y)) \right) . \end{aligned}$$

Writing  $\eta - \delta_x =: \xi \in \Xi_{k-1}$  for each fixed  $x \in V$ , the right-hand side can be rewritten as

$$\sum_{x \in V} \sum_{\xi \in \Xi_{k-1}} \mu_{\alpha,k}(\xi + \delta_x) (\xi_x + 1) f(\xi + \delta_x) \sum_{y \in V} c_{xy} (\alpha_y + \xi_y) (f(\xi + \delta_x) - f(\xi + \delta_y)) .$$

By (1.2), it holds that

$$\mu_{\alpha,k}(\xi + \delta_x) (\xi_x + 1) = \frac{Z_{\alpha,k-1}}{Z_{\alpha,k}} \mu_{\alpha,k-1}(\xi) (\alpha_x + \xi_x) , \quad x \in V , \xi \in \Xi_{k-1} . \quad (3.9)$$

Thus, by using the shortcut  $f_\xi(x) := f(\xi + \delta_x)$  for  $x \in V$  and  $\xi \in \Xi_{k-1}$ , we get

$$\mathcal{E}_{\alpha,k}(f) = \frac{Z_{\alpha,k-1}}{Z_{\alpha,k}} \sum_{x \in V} \sum_{\xi \in \Xi_{k-1}} \mu_{\alpha,k-1}(\xi) f_\xi(x) \sum_{y \in V} c_{xy} (\alpha_x + \xi_x) (\alpha_y + \xi_y) (f_\xi(x) - f_\xi(y)) .$$

Renormalizing and rewriting, this is equal to (recall (3.5))

$$\begin{aligned} & (|\alpha| + k - 1) \frac{Z_{\alpha,k-1}}{Z_{\alpha,k}} \sum_{\xi \in \Xi_{k-1}} \mu_{\alpha,k-1}(\xi) \sum_{x,y \in V} f_\xi(x) \frac{c_{xy} (\alpha_x + \xi_x) (\alpha_y + \xi_y)}{|\alpha| + k - 1} (f_\xi(x) - f_\xi(y)) \\ &= (|\alpha| + k - 1) \frac{Z_{\alpha,k-1}}{Z_{\alpha,k}} \sum_{\xi \in \Xi_{k-1}} \mu_{\alpha,k-1}(\xi) \mathcal{D}_{\alpha+\xi}(f_\xi) . \end{aligned}$$

By (3.7) with  $\alpha + \xi$  in place of  $\alpha$ , the Dirichlet form  $\mathcal{D}_{\alpha+\xi}(f_\xi)$  in the right-hand side is bounded from below by

$$\text{gap}_{\text{RW}}(\alpha + \xi) \text{var}_{\alpha+\xi}(f_\xi) .$$

Thus, we have verified that

$$\mathcal{E}_{\alpha,k}(f) \geq (|\alpha| + k - 1) \frac{Z_{\alpha,k-1}}{Z_{\alpha,k}} \sum_{\xi \in \Xi_{k-1}} \mu_{\alpha,k-1}(\xi) \text{gap}_{\text{RW}}(\alpha + \xi) \text{var}_{\alpha+\xi}(f_\xi) . \quad (3.10)$$

Observe that since  $f \in \text{Ker } \mathbf{a}_{k-1}^\dagger$ , we have

$$\begin{aligned} \text{var}_{\alpha+\xi}(f_\xi) &= \sum_{x \in V} \frac{\alpha_x + \xi_x}{|\alpha| + k - 1} f_\xi(x)^2 - \left( \sum_{x \in V} \frac{\alpha_x + \xi_x}{|\alpha| + k - 1} f(\xi + \delta_x) \right)^2 \\ &= \sum_{x \in V} \frac{\alpha_x + \xi_x}{|\alpha| + k - 1} f_\xi(x)^2 . \end{aligned}$$

Thus, plugging this identity into (3.10) yields

$$\begin{aligned} \mathcal{E}_{\alpha,k}(f) &\geq \frac{Z_{\alpha,k-1}}{Z_{\alpha,k}} \sum_{\xi \in \Xi_{k-1}} \sum_{x \in V} \mu_{\alpha,k-1}(\xi) \text{gap}_{\text{RW}}(\alpha + \xi) (\alpha_x + \xi_x) f_\xi(x)^2 \\ &\geq \left( \inf_{\xi \in \Xi_{k-1}} \text{gap}_{\text{RW}}(\alpha + \xi) \right) \frac{Z_{\alpha,k-1}}{Z_{\alpha,k}} \sum_{x \in V} \sum_{\xi \in \Xi_{k-1}} \mu_{\alpha,k-1}(\xi) (\alpha_x + \xi_x) f(\xi + \delta_x)^2 . \end{aligned}$$

This last expression outside parenthesis can be rewritten as

$$\begin{aligned} & \frac{Z_{\alpha,k-1}}{Z_{\alpha,k}} \sum_{x \in V} \sum_{\xi \in \Xi_{k-1}} \mu_{\alpha,k-1}(\xi) (\alpha_x + \xi_x) f(\xi + \delta_x)^2 \\ &= \sum_{x \in V} \sum_{\xi \in \Xi_{k-1}} \mu_{\alpha,k}(\xi + \delta_x) (\xi_x + 1) f(\xi + \delta_x)^2 \\ &= \sum_{x \in V} \sum_{\eta \in \Xi_k: \eta_x \geq 1} \mu_{\alpha,k}(\eta) \eta_x f(\eta)^2 = k \sum_{\eta \in \Xi_k} \mu_{\alpha,k}(\eta) f(\eta)^2 = k \text{Var}_{\alpha,k}(f) , \end{aligned}$$

where the first identity holds by (3.9), the second one holds by substituting  $\eta := \xi + \delta_x$ , the third one holds by exchanging the order of summations and  $\eta \in \Xi_k$ , and the fourth one holds by (3.8). Therefore, we conclude the proof of the lemma.  $\square$

**3.3. Min-max theorem for eigenvalues.** Here, we apply the well-known min-max theorem for eigenvalues (e.g., [SC97, Theorem 1.2.10]) to obtain a lower bound for the term  $\inf_{\xi \in \Xi_{k-1}} \text{gap}_{\text{RW}}(\alpha + \xi)$  that appears in Lemma 3.2.

**Lemma 3.3.** *For every  $\alpha = (\alpha_x)_{x \in V}$  and  $k \in \mathbb{N}$ , it holds that*

$$\inf_{\xi \in \Xi_{k-1}} \text{gap}_{\text{RW}}(\alpha + \xi) \geq \frac{\alpha_{\min}}{\alpha_{\min} + k - 1} \text{gap}_{\text{RW}}(\alpha) .$$

*Proof.* Let us compare the Dirichlet forms and the  $L^2$ -norms associated to  $\text{RW}(\alpha)$  and  $\text{RW}(\alpha + \xi)$ . We claim that, for all  $\phi : V \rightarrow \mathbb{R}$  and  $\xi \in \Xi_{k-1}$ ,

$$\frac{|\alpha|}{|\alpha| + k - 1} \mathcal{D}_\alpha(\phi) \leq \mathcal{D}_{\alpha + \xi}(\phi) , \quad \frac{\alpha_{\min}(|\alpha| + k - 1)}{|\alpha|(\alpha_{\min} + k - 1)} \|\phi\|_{L^2(\alpha + \xi)}^2 \leq \|\phi\|_{L^2(\alpha)}^2 . \quad (3.11)$$

The first inequality of (3.11) is trivial, since

$$\begin{aligned} \frac{|\alpha|}{|\alpha| + k - 1} \mathcal{D}_\alpha(\phi) &= \frac{1}{2(|\alpha| + k - 1)} \sum_{x, y \in V} c_{xy} \alpha_x \alpha_y (\phi(x) - \phi(y))^2 \\ &\leq \frac{1}{2} \sum_{x, y \in V} c_{xy} \frac{\alpha_x + \xi_x}{|\alpha| + k - 1} (\alpha_y + \xi_y) (\phi(x) - \phi(y))^2 = \mathcal{D}_{\alpha + \xi}(\phi) . \end{aligned}$$

The second inequality of (3.11) is also immediate by observing that

$$\begin{aligned} \frac{\alpha_{\min}(|\alpha| + k - 1)}{|\alpha|(\alpha_{\min} + k - 1)} \sum_{x \in V} \frac{\alpha_x + \xi_x}{|\alpha| + k - 1} \phi(x)^2 &\leq \frac{1}{|\alpha|} \sum_{x \in V} \alpha_x \frac{\alpha_x + \xi_x}{\alpha_x + k - 1} \phi(x)^2 \\ &\leq \sum_{x \in V} \frac{\alpha_x}{|\alpha|} \phi(x)^2 , \end{aligned}$$

where for the first and second inequalities we used, respectively,

$$\frac{\alpha_{\min}}{\alpha_{\min} + k - 1} \leq \frac{\alpha_x}{\alpha_x + k - 1} \quad \text{and} \quad \xi_x \leq k - 1 , \quad x \in V .$$

By applying as, e.g., in [SC97, Theorem 1.2.11] the min-max theorem for eigenvalues and the comparison inequalities in (3.11), we get

$$\frac{\alpha_{\min}}{\alpha_{\min} + k - 1} \lambda_j^\alpha \leq \lambda_j^{\alpha + \xi} , \quad j = 0, 1, \dots, n - 1 ,$$

where  $0 = \lambda_0^\alpha < \lambda_1^\alpha \leq \dots \leq \lambda_{n-1}^\alpha$  are the eigenvalues of the generator  $-A_\alpha$  and  $0 = \lambda_0^{\alpha + \xi} < \lambda_1^{\alpha + \xi} \leq \dots \leq \lambda_{n-1}^{\alpha + \xi}$  are the eigenvalues of the generator  $-A_{\alpha + \xi}$ . In particular, for  $j = 1$ , we obtain the desired comparison inequality for the spectral gaps, which concludes the proof of the lemma.  $\square$

**3.4. Proof of lower bound in Theorem 1.1.** Finally, we present a formal proof of the lower bound in Theorem 1.1.

*Proof of lower bound in Theorem 1.1.* Recall from (3.1) that we aim to prove that

$$(1 \wedge \alpha_{\min}) \text{gap}_{\text{RW}}(\alpha) \leq \text{gap}_k(\alpha) , \quad k \in \mathbb{N} . \quad (3.12)$$

We proceed by an induction on  $k \in \mathbb{N}$ . First, (3.12) is obvious for  $k = 1$ . Next, suppose that (3.12) holds for  $k - 1$ , and we prove (3.12) for  $k \geq 2$ . By (3.6) and (3.4), we have

$$\begin{aligned} \text{gap}_k(\alpha) &= \inf_{f \in \mathbb{R}^{\Xi_k}: f \neq \text{const.}} \frac{\mathcal{E}_{\alpha, k}(f)}{\text{Var}_{\alpha, k}(f)} \\ &= \left( \inf_{f \in \text{Im } \mathfrak{a}_k: f \neq \text{const.}} \frac{\mathcal{E}_{\alpha, k}(f)}{\text{Var}_{\alpha, k}(f)} \right) \wedge \left( \inf_{f \in \text{Ker } \mathfrak{a}_{k-1}^\dagger: f \neq \text{const.}} \frac{\mathcal{E}_{\alpha, k}(f)}{\text{Var}_{\alpha, k}(f)} \right) . \end{aligned} \quad (3.13)$$

Since  $\mathfrak{a}_k : \mathbb{R}^{\Xi_{k-1}} \rightarrow \mathbb{R}^{\Xi_k}$  lifts all the eigenfunctions of  $-L_{k-1}$  to  $-L_k$ , it is readily verified that

$$\inf_{f \in \text{Im } \mathfrak{a}_k : f \neq \text{const.}} \frac{\mathcal{E}_{\alpha,k}(f)}{\text{Var}_{\alpha,k}(f)} = \text{gap}_{k-1}(\alpha) \geq (1 \wedge \alpha_{\min}) \text{gap}_{\text{RW}}(\alpha) , \quad (3.14)$$

where the inequality holds by the induction hypothesis. Moreover, by Lemmas 3.2 and 3.3, we have

$$\inf_{f \in \text{Ker } \mathfrak{a}_{k-1}^+ : f \neq \text{const.}} \frac{\mathcal{E}_{\alpha,k}(f)}{\text{Var}_{\alpha,k}(f)} \geq k \left( \inf_{\xi \in \Xi_{k-1}} \text{gap}_{\text{RW}}(\alpha + \xi) \right) \geq \frac{\alpha_{\min} k}{\alpha_{\min} + k - 1} \text{gap}_{\text{RW}}(\alpha) . \quad (3.15)$$

It is straightforward to check that

$$\frac{\alpha_{\min} k}{\alpha_{\min} + k - 1} \geq 1 \wedge \alpha_{\min} , \quad \text{for all } k \geq 2 . \quad (3.16)$$

Collecting (3.13), (3.14), (3.15), and (3.16), we conclude that

$$\text{gap}_k(\alpha) \geq (1 \wedge \alpha_{\min}) \text{gap}_{\text{RW}}(\alpha) ,$$

which proves (3.12) for  $k$ . Therefore, by induction on  $k$ , we conclude the proof of (3.12) and thus the proof of Theorem 1.1.  $\square$

#### 4. PROOF OF THEOREM 1.3

BEP and SIP are related via an intertwining relation (e.g., [RS18, Proposition 5.1]): for all  $k \in \mathbb{N}$  and  $f \in \mathbb{R}^{\Xi_k}$ , we have

$$\mathcal{L} \Lambda f = \Lambda L_k f , \quad \text{with } \Lambda f(\zeta) := \sum_{\eta \in \Xi_k} \left( \prod_{x \in V} \frac{\zeta_x^{\eta_x}}{\eta_x!} \right) f(\eta) . \quad (4.1)$$

Thanks to this connection and the assertion in Theorem 1.1, once we check that the spectrum of  $\mathcal{L}$  on  $L^2(\Delta_V, \pi)$  is pure point, the proof of Theorem 1.3 boils down to verifying the validity of the following two claims:

- $-\text{gap}_{\text{RW}}(\alpha)$  belongs to the spectrum of  $\mathcal{L}$  (Lemma 4.1);
- each eigenvalue of  $\mathcal{L}$  is also an eigenvalue of  $L_k$  given in (1.1), for some  $k \in \mathbb{N}$  (Lemma 4.2).

The fact that  $\mathcal{L}$  has a pure point spectrum may be shown as follows. The generator  $\mathcal{L}$  is self-adjoint on  $L^2(\Delta_V, \pi)$ , thus, its spectrum is real. Moreover, for every  $k \in \mathbb{N}$ , the generator in (1.4) is easily seen to leave invariant the subspace of all polynomials of degree at most  $k$  in the variables  $(\zeta_x)_{x \in V}$ . Each of these subspaces is finite-dimensional, ensuring a decomposition of  $\mathcal{L}$  in terms of finitely-many eigenvalue/eigenfunction pairs when restricted therein. By density of polynomials in  $L^2(\Delta_V, \pi)$ , this eigendecomposition, suitably orthonormalized, gives rise to an orthonormal basis of  $L^2(\Delta_V, \pi)$  consisting of eigenfunctions of  $\mathcal{L}$ . In conclusion, the generator  $\mathcal{L}$  on  $L^2(\Delta_V, \pi)$  admits a pure point real spectrum.

We verify that  $-\text{gap}_{\text{RW}}(\alpha)$  belongs to the spectrum of  $\mathcal{L}$  in the following lemma, whose proof is analogous to that of Lemma 2.3.

**Lemma 4.1.** *Let  $\psi : V \rightarrow \mathbb{R}$  be an eigenfunction for  $A_\alpha$  associated to  $-\text{gap}_{\text{RW}}(\alpha)$ . Then, the first-order polynomial  $f : \Delta_V \rightarrow \mathbb{R}$  in the variables  $(\zeta_x)_{x \in V}$  given by*

$$f_\psi(\zeta) := \sum_{x \in V} \psi(x) \zeta_x , \quad \zeta \in \Delta_V , \quad (4.2)$$

*is an eigenfunction for  $\mathcal{L}$  associated to the same eigenvalue.*

*Proof.* Recalling (4.1) and writing  $g(\delta_x) := \psi(x)$ , the function in (4.2) reads as

$$f_\psi = \Lambda g .$$

In view of this representation for  $f_\psi$ , the injectivity of  $\Lambda$ , and the intertwining relation (4.1), the desired claim follows as in the proof of Lemma 2.3.  $\square$

In the following lemma, we show that each eigenfunction  $f$  of  $\mathcal{L}$  corresponds to an eigenfunction of  $L_k$ , for some  $k \in \mathbb{N}$ , both associated to the same eigenvalue.

**Lemma 4.2.** *Let  $f \in L^2(\Delta_V, \pi)$  be a non-constant eigenfunction of  $\mathcal{L}$  associated to the eigenvalue  $-\lambda < 0$ . Then, there exist  $k \in \mathbb{N}$  and  $g \in \mathbb{R}^{\Xi^k}$  such that  $g$  is an eigenfunction of  $L_k$  associated to the eigenvalue  $-\lambda < 0$ .*

*Proof.* By the discussion at the beginning of this section, all eigenfunctions of  $\mathcal{L}$  are polynomials in the variables  $(\zeta_x)_{x \in V}$  of degree at most  $k$ , for some  $k \in \mathbb{N}$ . Hence, we may rewrite  $f$  as follows:

$$f(\zeta) = \sum_{\ell=0}^k \sum_{\eta \in \Xi_\ell} g_\ell(\eta) \prod_{x \in V} \frac{\zeta_x^{\eta_x}}{\eta_x!} = \sum_{\ell=0}^k \Lambda g_\ell(\zeta) , \quad (4.3)$$

for some  $k \in \mathbb{N}$  and functions  $g_\ell \in \mathbb{R}^{\Xi_\ell}$ ,  $\ell = 0, 1, \dots, k$ . In the formula above, we included the factor  $(\prod_{x \in V} \eta_x!)^{-1}$  just for convenience, so to recover an expression like that in (4.1). Now, by applying the generator  $\mathcal{L}$  to the function  $f$  in (4.3) and using (4.1), we get

$$\mathcal{L}f(\zeta) = \sum_{\ell=0}^k \sum_{\eta \in \Xi_\ell} L_\ell g_\ell(\eta) \prod_{x \in V} \frac{\zeta_x^{\eta_x}}{\eta_x!} = \sum_{\ell=0}^k \Lambda L_\ell g_\ell(\zeta) .$$

The assumption  $\mathcal{L}f + \lambda f = 0$  yields

$$\sum_{\ell=0}^k \sum_{\eta \in \Xi_\ell} \{L_\ell g_\ell(\eta) + \lambda g_\ell(\eta)\} \prod_{x \in V} \frac{\zeta_x^{\eta_x}}{\eta_x!} = 0 ,$$

which holds for all  $\zeta \in \Delta_V$  if and only if each term in the bracket equals zero. Finally, since at least one among the functions  $g_\ell$ ,  $\ell > 0$ , is non-zero, this ensures that  $-\lambda < 0$  is an eigenvalue for  $L_\ell$ .  $\square$

**Acknowledgement.** *SK was supported by NRF-2019-Fostering Core Leaders of the Future Basic Science Program/Global Ph.D. Fellowship Program and the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (No. 2017R1A5A1015626, 2022R1F1A106366811 and 2022R1A5A6000840). SK wishes to express his gratitude to Institute of Science and Technology Austria (where the project was initiated) and University of Trieste (where the project was completed) for the warm hospitality during his stays. FS thanks Pietro Caputo and Matteo Quattropani for fruitful discussions.*

## REFERENCES

- [AL12] David Aldous and Daniel Lanoue. A lecture on the averaging process. *Probab. Surv.*, 9:90–102, 2012.
- [BC23] Alexandre Bristiel and Pietro Caputo. Entropy inequalities for random walks and permutations. *Ann. Inst. Henri Poincaré Probab. Stat. (to appear)*, 2023.
- [BDG17] Alessandra Bianchi, Sander Dommers, and Cristian Giardinà. Metastability in the reversible inclusion process. *Electron. J. Probab.*, 22:no. 70, 34, 2017.
- [BW09] F. Barthe and P. Wolff. Remarks on non-interacting conservative spin systems: the case of gamma distributions. *Stochastic Process. Appl.*, 119(8):2711–2723, 2009.

- [Cap03] Pietro Caputo. Uniform Poincaré inequalities for unbounded conservative spin systems: the non-interacting case. *Stochastic Process. Appl.*, 106(2):223–244, 2003.
- [Cap04] Pietro Caputo. Spectral gap inequalities in product spaces with conservation laws. In *Stochastic analysis on large scale interacting systems*, volume 39 of *Adv. Stud. Pure Math.*, pages 53–88. Math. Soc. Japan, Tokyo, 2004.
- [Cap08] Pietro Caputo. On the spectral gap of the Kac walk and other binary collision processes. *ALEA Lat. Am. J. Probab. Math. Stat.*, 4:205–222, 2008.
- [CLL20] Pietro Caputo, Cyril Labbé, and Hubert Lacoïn. Mixing time of the adjacent walk on the simplex. *Ann. Probab.*, 48(5):2449–2493, 2020.
- [CLL22] Pietro Caputo, Cyril Labbé, and Hubert Lacoïn. Spectral gap and cutoff phenomenon for the Gibbs sampler of  $\nabla\varphi$  interfaces with convex potential. *Ann. Inst. Henri Poincaré Probab. Stat.*, 58(2):794–826, 2022.
- [CLR10] Pietro Caputo, Thomas M. Liggett, and Thomas Richthammer. Proof of Aldous’ spectral gap conjecture. *J. Amer. Math. Soc.*, 23(3):831–851, 2010.
- [Cor23] Josué Corujo. On the spectrum of a neutral multi-allelic Moran model. *ALEA Lat. Am. J. Probab. Math. Stat.*, 20:505–546, 2023.
- [DSC93] Persi Diaconis and Laurent Saloff-Coste. Comparison theorems for reversible Markov chains. *Ann. Appl. Probab.*, 3(3):696–730, 1993.
- [DSPS21] Lorenzo Dello Schiavo, Lorenzo Portinale, and Federico Sau. Scaling limits of random walks, harmonic profiles, and stationary non-equilibrium states in Lipschitz domains. *arXiv:2112.14196*, 2021.
- [FGS22] Chiara Franceschini, Patrícia Gonçalves, and Federico Sau. Symmetric inclusion process with slow boundary: hydrodynamics and hydrostatics. *Bernoulli*, 28(2):1340–1381, 2022.
- [FRS22] Simone Floreani, Frank Redig, and Federico Sau. Orthogonal polynomial duality of boundary driven particle systems and non-equilibrium correlations. *Ann. Inst. Henri Poincaré Probab. Stat.*, 58(1):220–247, 2022.
- [GJ14] Patrícia Gonçalves and Milton Jara. Nonlinear fluctuations of weakly asymmetric interacting particle systems. *Arch. Ration. Mech. Anal.*, 212(2):597–644, 2014.
- [GKR07] Cristian Giardinà, Jorge Kurchan, and Frank Redig. Duality and exact correlations for a model of heat conduction. *J. Math. Phys.*, 48(3):033301, 15, 2007.
- [GKRV09] Cristian Giardinà, Jorge Kurchan, Frank Redig, and Kiamars Vafayi. Duality and hidden symmetries in interacting particle systems. *J. Stat. Phys.*, 135(1):25–55, 2009.
- [Gri14] Robert C. Griffiths. The  $\Lambda$ -Fleming-Viot process and a connection with Wright-Fisher diffusion. *Adv. in Appl. Probab.*, 46(4):1009–1035, 2014.
- [GRV10] C. Giardinà, F. Redig, and K. Vafayi. Correlation inequalities for interacting particle systems with duality. *J. Stat. Phys.*, 141(2):242–263, 2010.
- [GRV11] Stefan Grosskinsky, Frank Redig, and Kiamars Vafayi. Condensation in the inclusion process and related models. *J. Stat. Phys.*, 142(5):952–974, 2011.
- [GRV13] Stefan Grosskinsky, Frank Redig, and Kiamars Vafayi. Dynamics of condensation in the symmetric inclusion process. *Electron. J. Probab.*, 18:no. 66, 23, 2013.
- [HS19] Jonathan Hermon and Justin Salez. A version of Aldous’ spectral-gap conjecture for the zero range process. *Ann. Appl. Probab.*, 29(4):2217–2229, 2019.
- [JCG20] Watthanan Jatuviriyapornchai, Paul Chleboun, and Stefan Grosskinsky. Structure of the condensed phase in the inclusion process. *J. Stat. Phys.*, 178(3):682–710, 2020.
- [JLQY99] E. Janvresse, C. Landim, J. Quastel, and H. T. Yau. Relaxation to equilibrium of conservative dynamics. I. Zero-range processes. *Ann. Probab.*, 27(1):325–360, 1999.
- [Kim21] Seonwoo Kim. Second time scale of the metastability of reversible inclusion processes. *Probab. Theory Related Fields*, 180(3-4):1135–1187, 2021.
- [KL99] Claude Kipnis and Claudio Landim. *Scaling limits of interacting particle systems*, volume 320 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1999.
- [KS21] Seonwoo Kim and Insuk Seo. Condensation and metastable behavior of non-reversible inclusion processes. *Comm. Math. Phys.*, 382(2):1343–1401, 2021.
- [Lig05] Thomas M. Liggett. *Interacting particle systems*. Classics in Mathematics. Springer-Verlag, Berlin, 2005. Reprint of the 1985 original.
- [LMS23] C. Landim, D. Marcondes, and I. Seo. Metastable behavior of weakly mixing Markov chains: the case of reversible, critical zero-range processes. *Ann. Probab.*, 51(1):157–227, 2023.
- [LP22] Cyril Labbé and Enguérand Petit. Hydrodynamic limit and cutoff for the biased adjacent walk on the simplex. *arXiv:2203.04008*, 2022.

- [LPY02] C. Landim, G. Panizo, and H. T. Yau. Spectral gap and logarithmic Sobolev inequality for unbounded conservative spin systems. *Ann. Inst. H. Poincaré Probab. Statist.*, 38(5):739–777, 2002.
- [LSV96] C. Landim, S. Sethuraman, and S. Varadhan. Spectral gap for zero-range dynamics. *Ann. Probab.*, 24(4):1871–1902, 1996.
- [LY93] Sheng Lin Lu and Horng-Tzer Yau. Spectral gap and logarithmic Sobolev inequality for Kawasaki and Glauber dynamics. *Comm. Math. Phys.*, 156(2):399–433, 1993.
- [MT06] Ravi Montenegro and Prasad Tetali. Mathematical aspects of mixing times in Markov chains. *Found. Trends Theor. Comput. Sci.*, 1(3):x+121, 2006.
- [QS23] Matteo Quattropani and Federico Sau. Mixing of the averaging process and its discrete dual on finite-dimensional geometries. *Ann. Appl. Probab.*, 33(2):936–971, 2023.
- [Qua92] Jeremy Quastel. Diffusion of color in the simple exclusion process. *Comm. Pure Appl. Math.*, 45(6):623–679, 1992.
- [RS18] Frank Redig and Federico Sau. Factorized duality, stationary product measures and generating functions. *J. Stat. Phys.*, 172(4):980–1008, 2018.
- [Sal23] Justin Salez. Universality of cutoff for exclusion with reservoirs. *Ann. Probab.*, 51(2):478–494, 2023.
- [Sas13] Makiko Sasada. On the spectral gap of the Kac walk and other binary collision processes on  $d$ -dimensional lattice. In *Symmetries, integrable systems and representations*, volume 40 of *Springer Proc. Math. Stat.*, pages 543–560. Springer, Heidelberg, 2013.
- [SC97] Laurent Saloff-Coste. Lectures on finite Markov chains. In *Lectures on probability theory and statistics (Saint-Flour, 1996)*, volume 1665 of *Lecture Notes in Math.*, pages 301–413. Springer, Berlin, 1997.
- [Shi77] Norio Shimakura. Équations différentielles provenant de la génétique des populations. *Tôhoku Math. J.*, 29(2):287–318, 1977.

SEONWOO KIM, DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, SEOUL, REPUBLIC OF KOREA  
*Email address:* `ksw61eta@snu.ac.kr`

FEDERICO SAU, DIPARTIMENTO DI MATEMATICA E GEOSCIENZE, UNIVERSITÀ DEGLI STUDI DI TRIESTE, TRIESTE, ITALY  
*Email address:* `federico.sau@units.it`