

STRICTIFICATION AND NON-STRICTIFICATION OF MONOIDAL CATEGORIES

JORGE BECERRA

ABSTRACT. It is a classical result that any monoidal category is monoidally equivalent to a strict one, called its strictification. In this paper, we prove that any monoidal category is also monoidally equivalent to a non-strict monoidal category, called its non-strictification. Furthermore, we show that these two constructions give the free (non-)strict monoidal category generated by a monoidal category. Moreover, we prove in detail that these two constructions are part of a pair of free-forgetful 2-adjunctions.

CONTENTS

1. Introduction	1
2. Monoidal categories	3
3. Strictification of monoidal categories	5
4. Non-strictification of monoidal categories	10
5. Higher categorical perspective	15
References	22

1. INTRODUCTION

Monoidal categories are ubiquitous in mathematics, having a far-reaching role in homotopy theory [Hov99, MMSS01, Sch99], theoretical computer science [Bar13, Pav13] or quantum topology [GHW22, KRT97, Tur16], among many other areas. Monoidal categories can be thought of as the categorification of the algebraic structure of monoid. Namely, the product operation $\cdot : M \times M \rightarrow M$ and the unit element $1 \in M$ of a monoid M get categorified to a bifunctor $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ and an object $\mathbf{1} \in \mathcal{C}$. The associativity axiom $(m_1 m_2) m_3 = m_1 (m_2 m_3)$ and the unit axiom $1 \cdot m = m = m \cdot 1$ in a monoid can be categorified to equalities of objects $(X \otimes Y) \otimes Z = X \otimes (Y \otimes Z)$ and $1 \otimes X = X = X \otimes 1$, but in category theory this is a too strong requirement to demand. Instead, it is much more natural to ask that the objects $(X \otimes Y) \otimes Z$ and $X \otimes (Y \otimes Z)$ are simply isomorphic, and the same applies to $1 \otimes X$, X and $X \otimes 1$. If the equalities hold, one generally talks about a *strict* monoidal category.

It is a folklore result that any monoidal category is monoidally equivalent (that is, equivalent through a functor that respects the monoidal structures) to a strict one [JS93]. Concretely, for any monoidal category \mathcal{C} , Kassel [Kas95] constructs a monoidally equivalent category \mathcal{C}^{str} , called its *strictification*. Similarly, one is

Date: 30 March 2023.

2020 *Mathematics Subject Classification.* 18M05.

sometimes also interested in obtaining a non-strict category out of a monoidal category [BN97, HM21]. Mimicking Kassel's construction and inspired by [HM21], we will show

Theorem 1.1 (Theorem 4.2). *Any monoidal category \mathcal{C} is monoidally equivalent to a non-strict one \mathcal{C}_q .*

By analogy, we call \mathcal{C}_q the *non-strictification* of \mathcal{C} . A natural question is to ask whether either of these constructions are functorial, that is, whether a monoidal functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between monoidal categories induces functors

$$F^{\text{str}} : \mathcal{C}^{\text{str}} \rightarrow \mathcal{D}^{\text{str}} \quad , \quad F_q : \mathcal{C}_q \rightarrow \mathcal{D}_q.$$

If so, one step further would be to ask whether a monoidal natural transformation $\alpha : F \Rightarrow G$ between monoidal functors induces monoidal natural transformations

$$\alpha^{\text{str}} : F^{\text{str}} \Rightarrow G^{\text{str}} \quad , \quad \alpha_q : F_q \Rightarrow G_q.$$

The answer to both questions is affirmative.

Proposition 1.2 (Proposition 5.1 and Proposition 5.3). *The strictification and non-strictification give rise to 2-functors*

$$\text{str} : \text{MonCat} \rightarrow \text{strMonCat} \quad , \quad q : \text{MonCat} \rightarrow \text{nonstrMonCat}.$$

Above we have written MonCat (resp. strMonCat , nonstrMonCat) for the 2-categories with 0-cells monoidal categories (resp. strict monoidal categories, non-strict monoidal categories), 1-cells strong monoidal functors (resp. strict monoidal functors, in both cases) and 2-cells monoidal natural transformations (in the three cases).

Note that there are canonical forgetful 2-functors

$$\text{strMonCat} \rightarrow \text{MonCat} \quad , \quad \text{nonstrMonCat} \rightarrow \text{MonCat}.$$

We will also show that the strictification and non-strictification provide left-adjoints to these 2-functors.

Theorem 1.3 (Theorem 5.2 and Theorem 5.4). *There is a pair of free-forgetful 2-adjunctions*

$$\begin{array}{ccccc} & & q & & \text{str} \\ & \curvearrowright & & \curvearrowleft & \\ \text{nonstrMonCat} & \perp & \text{MonCat} & \perp & \text{strMonCat} \\ & \curvearrowleft & & \curvearrowright & \end{array}$$

In particular, the previous theorem ensures that \mathcal{C}^{str} (resp. \mathcal{C}_q) is the free strict (resp. non-strict) monoidal category generated by a monoidal category \mathcal{C} .

Organisation of the paper. In Section 2, we recall the basic concepts in monoidal category theory. In Section 3, we recall Kassel's construction of the strictification of a monoidal category, where we slightly simplify his argument. We also prove the universal property of the strictification (both for monoidal functors and monoidal natural transformations) and conclude with a concrete realisation of this construction for categories whose collection of objects is given by the free magma on a set. In Section 4, in a completely analogous way to the previous one, we start constructing the non-strictification of a monoidal category, later we show its universal property (for monoidal functors and monoidal natural transformations) and conclude with a concrete realisation for categories whose collection of objects is given

by the free monoid on a set. In Section 5, we review 2-categories, 2-functors and 2-adjunctions from the point of view of enrichments. Later we prove Proposition 1.2 and Theorem 1.3 and conclude obtaining a pair of adjoint equivalences for the core truncations of the aforementioned 2-categories.

Acknowledgments. The author would like to thank Roland van der Veen for valuable comments on the manuscript.

2. MONOIDAL CATEGORIES

In this section, we review the notion of monoidal category and the construction of a strictification of a monoidal category. The basic definitions can be found, for instance, in [TV17].

2.1. Monoidal categories. Let \mathcal{C} be a category. A *monoidal structure* on \mathcal{C} is the data of

- (1) a functor

$$\otimes : \mathcal{C} \times \mathcal{C} \longrightarrow \mathcal{C},$$

called the *monoidal product*,

- (2) an object $\mathbf{1} \in \mathcal{C}$, called the *unit object*,
 (3) a natural isomorphism

$$a : \otimes \circ (\otimes \times \text{Id}_{\mathcal{C}}) \xrightarrow{\cong} \otimes \circ (\text{Id}_{\mathcal{C}} \times \otimes)$$

of functors $\mathcal{C} \times \mathcal{C} \times \mathcal{C} \longrightarrow \mathcal{C}$, called the *associativity constrain*,

- (4) two natural isomorphisms

$$\ell : \mathbf{1} \otimes \text{Id}_{\mathcal{C}} \xrightarrow{\cong} \text{Id}_{\mathcal{C}} \quad , \quad r : \text{Id}_{\mathcal{C}} \otimes \mathbf{1} \xrightarrow{\cong} \text{Id}_{\mathcal{C}}$$

of functors $\mathcal{C} \longrightarrow \mathcal{C}$, called the *left and right unitality constrains*, respectively,

with the property that the following two diagrams commute for all objects X, Y, Z, M in \mathcal{C} :

$$\begin{array}{ccc}
 & ((X \otimes Y) \otimes Z) \otimes M & \\
 a_{X,Y,Z} \otimes \text{Id}_M \swarrow & & \searrow a_{X \otimes Y, Z, M} \\
 (X \otimes (Y \otimes Z)) \otimes M & & (X \otimes Y) \otimes (Z \otimes M) \\
 a_{X,Y \otimes Z, M} \searrow & & \swarrow a_{X,Y,Z \otimes M} \\
 X \otimes ((Y \otimes Z) \otimes M) & \xrightarrow{\text{Id}_X \otimes a_{Y,Z,M}} & X \otimes (Y \otimes (Z \otimes M))
 \end{array}
 \quad (\text{Pentagon axiom})$$

$$\begin{array}{ccc}
 & (X \otimes \mathbf{1}) \otimes Y & \\
 a_{X,\mathbf{1},Y} \swarrow & & \searrow r_X \otimes \text{Id}_Y \\
 X \otimes (\mathbf{1} \otimes Y) & \xrightarrow{\text{Id}_X \otimes \ell_Y} & X \otimes Y
 \end{array}
 \quad (\text{Triangle axiom})$$

We call the tuple $(\mathcal{C}, \otimes, \mathbf{1}, a, \ell, r)$ a *monoidal category*.

A monoidal category is said to be *strict* if the associativity, left and right unit constrains are the identity natural transformations, so that

$$(X \otimes Y) \otimes Z = X \otimes (Y \otimes Z) \quad , \quad \mathbf{1} \otimes X = X \quad , \quad X \otimes \mathbf{1} = X$$

for any object X in \mathcal{C} . We say that \mathcal{C} is *non-strict* if it is not strict.

2.2. Monoidal functors. Let $(\mathcal{C}, \otimes, \mathbf{1})$ and $(\mathcal{D}, \boxtimes, \mathbf{1})$ be monoidal categories. A *monoidal* functor from \mathcal{C} to \mathcal{D} is the data of

- (1) a functor $F : \mathcal{C} \rightarrow \mathcal{D}$,
- (2) a natural transformation

$$\gamma^F = \gamma : \boxtimes \circ (F \times F) \Longrightarrow F \circ \otimes$$

of functors $\mathcal{C} \times \mathcal{C} \rightarrow \mathcal{D}$,

- (3) an arrow in \mathcal{D}

$$u : \mathbf{1} \rightarrow F(\mathbf{1}),$$

which are compatible with the associativity and left and right unit constrains in the following sense: for any objects X, Y, Z in \mathcal{C} , the following diagrams commute:

$$\begin{array}{ccc}
(F(X) \boxtimes F(Y)) \boxtimes F(Z) & \xrightarrow{a'_{FX, FY, FZ}} & F(X) \boxtimes (F(Y) \boxtimes F(Z)) \\
\gamma_{X,Y} \boxtimes \text{Id}_{FZ} \swarrow & & \searrow \text{Id}_{FX} \boxtimes \gamma_{Y,Z} \\
F(X \otimes Y) \boxtimes F(Z) & & F(X) \boxtimes F(Y \otimes Z) \quad (\text{Hexagon axiom}) \\
\gamma_{X \otimes Y, Z} \searrow & & \swarrow \gamma_{X, Y \otimes Z} \\
F((X \otimes Y) \otimes Z) & \xrightarrow{F(a_{X,Y,Z})} & F(X \otimes (Y \otimes Z))
\end{array}$$

$$\begin{array}{ccc}
\mathbf{1} \boxtimes F(X) & \xrightarrow{\ell'_{FX}} & F(X) & & F(X) \boxtimes \mathbf{1} & \xrightarrow{r'_{FX}} & F(X) \\
u \boxtimes \text{Id}_{FX} \downarrow & & \downarrow F(\ell_X) & & \text{Id}_{FX} \boxtimes u \downarrow & & \downarrow F(r_X) \quad (2.1) \\
F(\mathbf{1}) \boxtimes F(X) & \xrightarrow{\gamma_{\mathbf{1}, X}} & F(\mathbf{1} \otimes X) & & F(X) \boxtimes F(\mathbf{1}) & \xrightarrow{\gamma_{X, \mathbf{1}}} & F(X \otimes \mathbf{1})
\end{array}$$

where we have written (a, ℓ, r) for the constrains of \mathcal{C} and (a', ℓ', r') for the constrains of \mathcal{D} .

The composite

$$(\mathcal{C}, \otimes, \mathbf{1}) \xrightarrow{F} (\mathcal{D}, \boxtimes, \mathbf{1}) \xrightarrow{F'} (\mathcal{E}, \odot, \mathbf{1})$$

of two monoidal functors (F, γ, u) and (F', γ', u') is also monoidal with coherence constrains given by the composites

$$\mathbf{1} \xrightarrow{u'} F'(\mathbf{1}) \xrightarrow{F'(u)} (F'F)(\mathbf{1})$$

and

$$\odot \circ (F'F \times F'F) \xrightarrow{\gamma'(F \times F)} F' \circ \boxtimes \circ (F \times F) \xrightarrow{F'\gamma} F'F \circ \otimes.$$

A monoidal functor $F = (F, \gamma, u) : \mathcal{C} \rightarrow \mathcal{D}$ as above is *strong* if γ is a natural isomorphism of functors and u is an isomorphism in \mathcal{D} . We say that F is *strict* if

γ is the identity natural transformation and u is the identity arrow. If F is not strong or strict, it is usually called *lax* to distinguish it from these other two.

A *monoidal equivalence* between monoidal categories \mathcal{C} and \mathcal{D} is a strong monoidal functor $F : \mathcal{C} \rightarrow \mathcal{D}$ which is an equivalence of ordinary categories. If the functor F is strict, then we call it a *strict monoidal equivalence*.

2.3. Monoidal natural transformations. Let $F, G : \mathcal{C} \rightarrow \mathcal{D}$ be (lax, strong or strict) monoidal functors between monoidal categories $(\mathcal{C}, \otimes, \mathbf{1})$ and $(\mathcal{D}, \boxtimes, \mathbf{1})$, a natural transformation $\alpha : F \Rightarrow G$ is called *monoidal* if it is compatible with the monoidal constraints of F and G in the sense that the following two diagrams commute for all objects X, Y in \mathcal{C} :

$$\begin{array}{ccc}
 & \mathbf{1} & \\
 u \swarrow & & \searrow u' \\
 F(\mathbf{1}) & \xrightarrow{\alpha_{\mathbf{1}}} & G(\mathbf{1})
 \end{array}
 \quad , \quad
 \begin{array}{ccc}
 F(X) \boxtimes F(Y) & \xrightarrow{\gamma_{X,Y}} & F(X \otimes Y) \\
 \alpha_X \boxtimes \alpha_Y \downarrow & & \downarrow \alpha_{X \otimes Y} \\
 G(X) \boxtimes G(Y) & \xrightarrow{\gamma'_{X,Y}} & G(X \otimes Y)
 \end{array}$$

where $F = (F, \gamma, u)$ and $G = (G, \gamma', u')$. If in addition α is a natural isomorphism, we say that it is a *monoidal natural isomorphism*.

3. STRICTIFICATION OF MONOIDAL CATEGORIES

In this section we recall Kassel's construction of a strict monoidal category \mathcal{C}^{str} monoidally equivalent to a given monoidal category \mathcal{C} [Kas95, §XI.5]. We also show that this is the free strict monoidal category generated by \mathcal{C} .

3.1. Construction. Let $(\mathcal{C}, \otimes, \mathbf{1}, a, \ell, r)$ be a monoidal category. We define the category \mathcal{C}^{str} as follows: its objects are finite sequences $S = (X_1, \dots, X_n)$ of objects of \mathcal{C} , $n \geq 0$ (this includes the empty sequence \emptyset). If the *parenthesisation* of a sequence S is

$$Par(S) := (\dots(X_1 \otimes X_2) \otimes X_3) \otimes \dots \otimes X_n \in \mathcal{C}$$

for any sequence S of length $n > 0$ and $Par(\emptyset) := \mathbf{1}$, define

$$Hom_{\mathcal{C}^{\text{str}}}(S, S') := Hom_{\mathcal{C}}(Par(S), Par(S')), \quad (3.1)$$

that is, the datum of a map $f : S \rightarrow S'$ in \mathcal{C}^{str} is the same as the datum of a map $Par(f) : Par(S) \rightarrow Par(S')$ in \mathcal{C} . The composite law and identities are given by those of \mathcal{C} , so that parenthesisation gives rise to a functor $Par : \mathcal{C}^{\text{str}} \rightarrow \mathcal{C}$. Moreover, there is a canonical full embedding (i.e. fully faithful injective-on-objects functor) $i : \mathcal{C} \hookrightarrow \mathcal{C}^{\text{str}}$ given by $i(X) := (X)$, the length-one sequence whose only object is X .

Lemma 3.1. *The canonical embedding*

$$i : \mathcal{C} \hookrightarrow \mathcal{C}^{\text{str}}$$

is an equivalence of categories.

Proof. Let us see that Par can be taken as a quasi-inverse of i . It is clear that $Par \circ i = Id_{\mathcal{C}}$. Now, for $S \in \mathcal{C}^{\text{str}}$, let

$$\delta_S : S \rightarrow (Par(S))$$

be the unique arrow that corresponds to $Id_{Par(S)}$ under (3.1). These maps assemble into a natural isomorphism $\delta : Id_{\mathcal{C}^{\text{str}}} \Rightarrow i \circ Par$ since for an arrow $f : S \rightarrow S'$,

the equality $(i \circ \text{Par})(f) \circ \delta_S = \delta_{S'} \circ f$ (the naturality of δ) translates into $f = f$ under (3.1). \square

The category \mathcal{C}^{str} can be endowed with a strict monoidal structure, as follows: for non-empty sequences $S = (X_1, \dots, X_n)$ and $S' = (X_{n+1}, \dots, X_{n+m})$, set

$$S * S' := (X_1, \dots, X_{n+m}),$$

and we also put $S * \emptyset := S = \emptyset * S$, where S is possibly empty.

To upgrade the concadenation $*$ to a functor, we must define first a family of natural maps

$$\theta_{S,S'} : \text{Par}(S) \otimes \text{Par}(S') \longrightarrow \text{Par}(S * S') \quad (3.2)$$

inductively on the length of S' . Set $\theta_{\emptyset,S} := \ell_{\text{Par}(S)}$ and $\theta_{S,\emptyset} := r_{\text{Par}(S)}$. For $S' = (X)$ a sequence with one object, we set

$$\theta_{S,S'} := \text{Id}_{\text{Par}(S) \otimes X} : \text{Par}(S) \otimes X \longrightarrow \text{Par}(S) \otimes X = \text{Par}(S * (X)).$$

In general for $S' = \bar{S} * (X)$, then we define $\theta_{S,S'}$ as the composite

$$\begin{array}{ccc} \text{Par}(S) \otimes \text{Par}(S') & \xrightarrow{\theta_{S,S'}} & \text{Par}(S * S') \\ \parallel & & \parallel \\ \text{Par}(S) \otimes (\text{Par}(\bar{S}) \otimes X) & \xrightarrow{\theta_{S,S'}} & \text{Par}(S * \bar{S}) \otimes X \\ & \searrow \scriptstyle a_{\text{Par}(S), \text{Par}(\bar{S}), X}^{-1} \quad \theta_{S, \bar{S}} \otimes \text{Id}_X \nearrow & \\ & (\text{Par}(S) \otimes \text{Par}(\bar{S})) \otimes X & \end{array}$$

The naturality of θ then follows from the naturality of a, ℓ and r .

Now, given arrows $f : S_1 \longrightarrow S_2$ and $g : S'_1 \longrightarrow S'_2$ in \mathcal{C}^{str} , we define the arrow $f * g : S_1 * S'_1 \longrightarrow S_2 * S'_2$ as the composite given by the following dashed arrow:

$$\begin{array}{ccc} \text{Par}(S_1) \otimes \text{Par}(S'_1) & \xleftarrow{\theta_{S_1, S'_1}^{-1}} & \text{Par}(S_1 * S'_1) \\ f \otimes g \downarrow & & \downarrow \text{Par}(f * g) \\ \text{Par}(S_2) \otimes \text{Par}(S'_2) & \xrightarrow{\theta_{S_2, S'_2}} & \text{Par}(S_2 * S'_2) \end{array} \quad (3.3)$$

The functoriality of $*$ then follows from functoriality of \otimes by the above diagram. Since the concadenation $*$ is strictly associative and unital, it makes \mathcal{C}^{str} into a strict monoidal category, where the unit object is the empty sequence \emptyset .

Theorem 3.2 (Strictness, [ML63]). *Let \mathcal{C} be a monoidal category. The canonical embedding*

$$i : \mathcal{C} \xrightarrow{\cong} \mathcal{C}^{\text{str}}$$

is a monoidal equivalence of categories.

Proof. It is only left to exhibit i as a strong monoidal functor. Let $u : \emptyset \longrightarrow (\mathbf{1})$ be the unique map corresponding to $\text{Id}_{\mathbf{1}}$ under (3.1), that is, let $\text{Par}(u) := \text{Id}_{\mathbf{1}}$, and define

$$\eta_{X,Y} : (X) * (Y) = (X, Y) \longrightarrow (X \otimes Y)$$

as $\text{Par}(\eta_{X,Y}) := \text{Id}_{X \otimes Y}$. Clearly, η assembles into a natural transformation $\eta : * \circ (i \times i) \Longrightarrow i \circ \otimes$. The Hexagon axiom and the left and right squares (2.1) hold trivially, because their commutativity correspond under (3.1) to the equalities $a_{X,Y,Z} = a_{X,Y,Z}$, $\ell_X = \ell_X$ and $r_X = r_X$, respectively. \square

Remark 3.3. Our argument, that uses i instead of with Par , simplifies the one given by Kassel, as it avoids a lengthy computation to show that Par is strong monoidal [Kas95, XI.5.2], where the monoidal constraints for Par are given precisely by θ and Id_1 . From our perspective that follows from generalities of monoidal categories (e.g. [TV17, §1.4.9]).

Remark 3.4. Another way of exhibiting a monoidal equivalence between a monoidal category \mathcal{C} and a strict monoidal category is by constructing a monoidal equivalence between \mathcal{C} and the category of right \mathcal{C} -module endofunctors of \mathcal{C} , see [JS93].

Remark 3.5. Mac Lane Strictness Theorem 3.2 is equivalent to the celebrated Mac Lane Coherence Theorem, which states that in a monoidal category, any *formal* diagram made out of associativity, left and right unit constraints and identities commute. Here, the word “formal” means that no other isomorphism or equality of objects in the category may appear. An easy argument to derive this from the Strictness theorem can be found in [EGNO15].

3.2. Properties of strictification. Let us now discuss some properties of the previous construction. First, we will state the universal property of the strictification, which ensures that the strictification of a monoidal category \mathcal{C} is the free strict monoidal category generated by \mathcal{C} .

Theorem 3.6. *Let \mathcal{C} be a monoidal category and let \mathcal{D} be a strict monoidal category. Given a strong monoidal functor $F : \mathcal{C} \rightarrow \mathcal{D}$, there exists a unique strict monoidal functor $\widehat{F} : \mathcal{C}^{\text{str}} \rightarrow \mathcal{D}$ such that $\widehat{F} \circ i = F$,*

$$\begin{array}{ccc} & \mathcal{C}^{\text{str}} & \\ & \uparrow i & \searrow \widehat{F} \\ \mathcal{C} & \xrightarrow{F} & \mathcal{D} \end{array}$$

Proof. Let us write $(\mathcal{C}, \otimes, \mathbf{1})$ and $(\mathcal{D}, \boxtimes, \mathbb{1})$ for the monoidal structures. For a sequence $S = (X_1, \dots, X_n)$, since \widehat{F} (if it exists) is strict monoidal, we must have

$$\widehat{F}(S) = F(X_1) \boxtimes \dots \boxtimes F(X_n)$$

(regardless of parentheses as \mathcal{D} is strict) and $\widehat{F}(\emptyset) = \mathbb{1}$, hence this is the only possible definition for \widehat{F} . To see what \widehat{F} must be in arrows, consider the natural isomorphism $\delta : \text{Id}_{\mathcal{C}^{\text{str}}} \Rightarrow i \circ Par$ from the proof of Lemma 3.1. Given an arrow $f : S \rightarrow S'$, we have $(i \circ Par)(f) \circ \delta_S = \delta_{S'} \circ f$ by the naturality of δ and applying \widehat{F} to this we obtain a commutative diagram

$$\begin{array}{ccc} \widehat{F}(S) & \xrightarrow{\widehat{F}(f)} & \widehat{F}(S') \\ \widehat{F}(\delta_S) \downarrow & & \downarrow \widehat{F}(\delta_{S'}) \\ F(Par(S)) & \xrightarrow{F(Par(f))} & F(Par(S')) \end{array}$$

We will show now that for \widehat{F} as in the statement, the map $\widehat{F}(\delta_S)$ is fully determined by data of F , and so we will obtain a single possible definition for $\widehat{F}(f)$.

Let us now define a family of arrows

$$\beta_S^F = \beta_S : \widehat{F}(S) \rightarrow F(Par(S)).$$

Write $\gamma : \boxtimes \circ (F \times F) \xrightarrow{\cong} F \circ \boxtimes$ and $u : \mathbf{1} \rightarrow F(\mathbf{1})$ for the coherence data associated to the strong monoidal functor F . Inductively on the length of S , define $\beta_\emptyset := u$, $\beta_{(X)} := \text{Id}_{FX}$ and for $\bar{S} = S * (X)$ let $\beta_{\bar{S}}$ be the composite

$$\widehat{F}(\bar{S}) = \widehat{F}(S) \boxtimes F(X) \xrightarrow{\beta_S \boxtimes \text{Id}_{FX_n}} F(\text{Par}(S)) \boxtimes F(X) \xrightarrow{\gamma_{\text{Par}(S), X}} F(\text{Par}(\bar{S}))$$

Now we claim that for any such \widehat{F} we must have

$$\beta_S = \widehat{F}(\delta_S). \quad (3.4)$$

For we observe first that δ_S can be described inductively using η , the monoidal constraint of i , in a similar fashion as how β_S was defined. On the other hand, the equality of monoidal functors $\widehat{F} \circ i = F$ signifies at the level of the monoidal constraints that

$$\begin{array}{ccc} F(X) \boxtimes F(Y) & \xrightarrow{\gamma_{X,Y}} & F(X \otimes Y) \\ \parallel & & \parallel \\ \widehat{F}(X, Y) & \xrightarrow{\widehat{F}(\eta_{X,Y})} & \widehat{F}(i(X \otimes Y)) \end{array}$$

which by induction implies (3.4).

In conclusion, if \widehat{F} is as in the statement, then for any $f : S \rightarrow S'$ in \mathcal{C}^{str} we have a commutative diagram

$$\begin{array}{ccc} \widehat{F}(S) & \xrightarrow{\widehat{F}(f)} & \widehat{F}(S') \\ \beta_S \downarrow \cong & & \cong \downarrow \beta_{S'} \\ F(\text{Par}(S)) & \xrightarrow{F(\text{Par}(f))} & F(\text{Par}(S')) \end{array}$$

and so $\widehat{F}(f) = \beta_{S'}^{-1} \circ F(\text{Par}(f)) \circ \beta_S$ is the only possible definition for $\widehat{F}(f)$, and moreover we have $\widehat{F} \circ i = F$ as strong monoidal functors by construction. \square

A similar result also applies to monoidal natural transformations:

Proposition 3.7. *Let \mathcal{C} be a monoidal category and let \mathcal{D} be a strict monoidal category. Given a monoidal natural transformation $\alpha : F \Rightarrow G$ between strong monoidal functors $F, G : \mathcal{C} \rightarrow \mathcal{D}$, there exists a unique monoidal natural transformation $\widehat{\alpha} : \widehat{F} \Rightarrow \widehat{G}$ such that $\widehat{\alpha}i = \alpha$,*

$$\begin{array}{ccc} \mathcal{C}^{\text{str}} & & \mathcal{D} \\ \uparrow i & \searrow \widehat{F} & \\ \mathcal{C} & \xrightarrow{\widehat{G}} & \mathcal{D} \\ \downarrow F & \searrow \widehat{\alpha} & \\ \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ & \searrow \alpha & \\ & \mathcal{D} & \end{array}$$

Proof. The condition $\widehat{\alpha}i = \alpha$ means that for the length-one sequence (X) , we must have $\widehat{\alpha}_{(X)} = \alpha_X$. This implies that for an arbitrary sequence $S = (X_1, \dots, X_n)$ in \mathcal{C}^{str} ,

$$\widehat{\alpha}_S = \widehat{\alpha}_{(X_1)*\dots*(X_n)} = \widehat{\alpha}_{(X_1)} \boxtimes \dots \boxtimes \widehat{\alpha}_{(X_n)} = \alpha_{X_1} \boxtimes \dots \boxtimes \alpha_{X_n},$$

where in the second equality we have used that $\widehat{\alpha}$ is a monoidal natural transformation between strict monoidal functors. Besides, $\widehat{\alpha}_\emptyset = \text{Id}_{\mathbf{1}}$. Therefore, this is the

only possible definition. It is only left to check that $\widehat{\alpha}_S := \alpha_{X_1} \boxtimes \cdots \boxtimes \alpha_{X_n}$ is natural on S . Given an arrow $f : S \rightarrow S'$ in \mathcal{C}^{str} , let us contemplate the following cube:

$$\begin{array}{ccccc}
 & & \widehat{F}(S) & \xrightarrow{\widehat{F}(f)} & \widehat{F}(S') \\
 & \swarrow & \downarrow & \swarrow & \downarrow \\
 & & F(\text{Par}(S)) & \xrightarrow{F(\text{Par}(f))} & F(\text{Par}(S')) \\
 & \downarrow & \downarrow & \downarrow & \downarrow \\
 & & \widehat{G}(S) & \xrightarrow{\widehat{G}(f)} & \widehat{G}(S') \\
 & \swarrow & \downarrow & \swarrow & \downarrow \\
 & & G(\text{Par}(S)) & \xrightarrow{G(\text{Par}(f))} & G(\text{Par}(S'))
 \end{array}$$

$\alpha_{\text{Par}(S)}$ (left vertical arrow), β_S (top-left diagonal), $\beta_{S'}$ (top-right diagonal), $\widehat{\alpha}_S$ (middle-left vertical), $\widehat{\alpha}_{S'}$ (middle-right vertical), $\alpha_{\text{Par}(S')}$ (right vertical), $\bar{\beta}_S$ (bottom-left diagonal), $\bar{\beta}_{S'}$ (bottom-right diagonal).

We have put $\beta = \beta^F$ and $\bar{\beta} = \beta^G$. The top and bottom faces commute by the definition of \widehat{F} and \widehat{G} , the left and right faces commute because of the monoidality of α , and the front face also commutes by the naturality of α . Since $\beta_S, \beta_{S'}, \bar{\beta}_S$ and $\bar{\beta}_{S'}$ are isomorphisms, this implies the commutativity of the back face, which is precisely the naturality of $\widehat{\alpha}$. \square

3.3. A common realisation. Many examples “in nature” of non-strict monoidal categories, especially in quantum topology, have as the collection of objects the free (unital) magma $\text{Mag}(X)$ on some set X . We would like to make the previous construction more transparent in this case.

Suppose that \mathcal{C} is a monoidal category with $\text{ob}(\mathcal{C}) = \text{Mag}(X)$ and monoidal product given by the magma product. Let us now define a new category $\widetilde{\mathcal{C}}^{\text{str}}$. Its objects are given by the elements of the free monoid $\text{Mon}(X)$ on X . Given an object $w = x_1 \cdots x_n$ of $\widetilde{\mathcal{C}}^{\text{str}}$, its *sequencing* is the object of \mathcal{C}^{str}

$$\text{Seq}(w) := (x_1, \dots, x_n) \in \mathcal{C}^{\text{str}},$$

with $\text{Seq}(\emptyset) := \emptyset^1$. The set of arrows in $\widetilde{\mathcal{C}}^{\text{str}}$ is defined as

$$\text{Hom}_{\widetilde{\mathcal{C}}^{\text{str}}}(w, w') := \text{Hom}_{\mathcal{C}^{\text{str}}}(\text{Seq}(w), \text{Seq}(w')).$$

As before, we readily see that Seq extends to a fully faithful functor

$$\text{Seq} : \widetilde{\mathcal{C}}^{\text{str}} \rightarrow \mathcal{C}^{\text{str}}.$$

We define a monoidal product on $\widetilde{\mathcal{C}}^{\text{str}}$ as follows: on objects, it is simply given by the monoid product, $w \star w' := ww'$. Now observe that if $w = x_1 \cdots x_n$ and $w' = y_1 \cdots y_m$, then

$$\begin{aligned}
 \text{Seq}(w \star w') &= \text{Seq}(x_1 \cdots x_n y_1 \cdots y_m) = (x_1, \dots, x_n, y_1, \dots, y_m) \\
 &= (x_1, \dots, x_n) * (y_1, \dots, y_m) = \text{Seq}(w) * \text{Seq}(w').
 \end{aligned}$$

This observation allows us to define the monoidal product of arrows: given $f : w_1 \rightarrow w_2$ and $g : w'_1 \rightarrow w'_2$, then $f \star g : w_1 \star w_2 \rightarrow w'_1 \star w'_2$ is given by

$$\text{Seq}(w_1 \star w'_1) = \text{Seq}(w_1) * \text{Seq}(w'_1) \xrightarrow{f \star g} \text{Seq}(w_2) * \text{Seq}(w'_2) = \text{Seq}(w_2 \star w'_2).$$

Setting the empty word \emptyset as the unit, all this data determines a strict monoidal structure on $\widetilde{\mathcal{C}}^{\text{str}}$ such that Seq is naturally a strict monoidal functor.

¹Note that the first \emptyset refers to the empty word in $\text{Mon}(X)$ whereas the second \emptyset refers to the empty sequence in \mathcal{C}^{str} .

Let us summarise our findings:

Proposition 3.8. *Let \mathcal{C} be a monoidal category with $\text{ob}(\mathcal{C}) = \text{Mag}(X)$ and monoidal product given by the magma product. Then there is a strict monoidal equivalence*

$$\text{Seq} : \widetilde{\mathcal{C}^{\text{str}}} \xrightarrow{\cong} \mathcal{C}^{\text{str}}$$

between \mathcal{C}^{str} and a (strict) monoidal category $\widetilde{\mathcal{C}^{\text{str}}}$ such that $\text{ob}(\widetilde{\mathcal{C}^{\text{str}}}) = \text{Mon}(X)$ and whose monoidal product is given by the monoid product.

Proof. It is only left to check that Seq is essentially surjective. Let $S = (v_1, \dots, v_n)$ be an element of \mathcal{C}^{str} , where $v_i \in \text{Mag}(X)$. If $U : \text{Mag}(X) \rightarrow \text{Mon}(X)$ is the canonical map that forgets parentheses, then we will exhibit an isomorphism $\text{Seq}(U(v_1) \cdots U(v_n)) \cong S$, or in other words, an isomorphism

$$\text{Par}(\text{Seq}(U(v_1) \cdots U(v_n))) \cong \text{Par}(S)$$

in \mathcal{C} . For this, it suffices to show that if $v \in \text{Mag}(X)$ and $U(v) = x_1 \cdots x_p$, then there is an isomorphism

$$\rho_v : v \xrightarrow{\cong} (\cdots (x_1 x_2) x_3) \cdots x_p = \text{Par}(\text{Seq}(U(v)))$$

in \mathcal{C} .

Let $\gamma : \star \circ (\text{Par} \times \text{Par}) \implies \text{Par} \circ \star$ be the natural isomorphism (3.2), where we also denote by \star the monoidal product on \mathcal{C} . We will define this isomorphism inductively on the length $|v|$ of v : put $\rho_\emptyset := \text{Id}_\emptyset$ and $\rho_x := \text{Id}_x$. Suppose we have defined ρ for objects of \mathcal{C} of length less than n . Let $v \in \text{Mag}(X)$ such that $|v| = n > 1$, and let $v_1, v_2 \in \text{Mag}(X)$ the unique elements of positive length such that $v = v_1 v_2$. Then define ρ_v as the composite

$$\begin{aligned} v = v_1 \star v_2 &\xrightarrow{\rho_{v_1 \star v_2}} \text{Par}(\text{Seq}(U(v_1))) \star \text{Par}(\text{Seq}(U(v_2))) \\ &\xrightarrow{\gamma_{\text{Seq}(U(v_1)), \text{Seq}(U(v_2))}} \text{Par}(\text{Seq}(U(v_1)) \star \text{Seq}(U(v_2))) \\ &\xlongequal{\quad\quad\quad} \text{Par}(\text{Seq}(U(v))) \end{aligned}$$

Since γ is a natural isomorphism, ρ_v is an isomorphism as well. \square

4. NON-STRICTIFICATION OF MONOIDAL CATEGORIES

The goal of this section is to show, using an analogous construction to the one in the previous section, that any monoidal category is monoidally equivalent to a non-strict one. Our construction is inspired by [HM21]. Later we will similarly prove the universal property of this construction.

4.1. Main construction. Let $\text{Mag}(\bullet)$ be the free unital magma generated by the singleton \bullet . Recall that elements of $\text{Mag}(\bullet)$ are parenthesised sequences of bullets (included the empty sequence), e.g. $((\bullet\bullet)\bullet)(\bullet\bullet)$. Forgetting parenthesis gives a map $|\cdot| : \text{Mag}(\bullet) \rightarrow \mathbb{N}$ that counts the number of bullets.

Given a monoidal category $(\mathcal{C}, \otimes, \mathbf{1}, a, \ell, r)$, we define \mathcal{C}_q as follows: its objects are pairs (S, t) , where S is a finite sequence of objects of \mathcal{C} , $t \in \text{Mag}(\bullet)$ and $|t| = |S|$, where $|S|$ is the length of S . For such a pair (S, t) , define its *parenthesisation* $\text{Par}(S, t) \in \mathcal{C}$ as the object in \mathcal{C} given by inserting the i -th element of S in the i -th bullet of t and inserting tensor products between them, e.g.

$$\text{Par}((X_1, \dots, X_5), ((\bullet\bullet)\bullet)(\bullet\bullet)) = ((X_1 \otimes X_2) \otimes X_3) \otimes (X_4 \otimes X_5),$$

and $Par(\emptyset, \emptyset) := \mathbf{1}$. More precisely, $Par(S, w)$ is defined by the following inductive rules:

- (1) $Par(\emptyset, \emptyset) := \mathbf{1}$,
- (2) $Par((X), \bullet) := X$,
- (3) Given $(S, t) \in \mathcal{C}_q$ with $|t| > 1$, there exist unique $t_1, t_2 \in \text{Mag}(\bullet)$ with $|t_1|, |t_2| \geq 1$, $|t_1| + |t_2| = |t|$ such that $t = t_1 t_2$. If S_1, S_2 are the unique subsequences of S of lengths $|t_1|, |t_2|$ respectively such that $S = S_1 * S_2$, then define

$$Par(S, t) := Par(S_1, t_1) \otimes Par(S_2, t_2).$$

The set of morphisms in \mathcal{C}_q is given by

$$\text{Hom}_{\mathcal{C}_q}((S, t), (S', t')) := \text{Hom}_{\mathcal{C}}(Par(S, t), Par(S', t')) \quad (4.1)$$

with the composite and identities determined by those of \mathcal{C} . Once again, the datum of a map $f : (S, t) \rightarrow (S', t')$ in \mathcal{C}_q is the same as the datum of a map $Par(f) : Par(S, t) \rightarrow Par(S', t')$ in \mathcal{C} . This means that the parenthesisation gives rise to a functor $Par : \mathcal{C}_q \rightarrow \mathcal{C}$. On the other hand, note that there is a canonical full embedding $j : \mathcal{C} \hookrightarrow \mathcal{C}_q$ defined by $j(X) := ((X), \bullet)$.

Lemma 4.1. *The canonical embedding*

$$j : \mathcal{C} \hookrightarrow \mathcal{C}_q$$

is an equivalence of categories.

Proof. We will prove that Par is a quasi-inverse of j . On the one hand, we have $Par \circ j = \text{Id}_{\mathcal{C}}$. On the other hand, for $(S, t) \in \mathcal{C}_q$, define

$$\delta_{(S, t)} : (S, t) \rightarrow (Par(S, t), \bullet)$$

as $Par(\delta_{(S, t)}) := \text{Id}_{Par(S, t)}$. As in Lemma 3.1, these maps trivially assemble into a natural transformation $\delta : \text{Id}_{\mathcal{C}_q} \Rightarrow j \circ Par$, which gives the result. \square

Let us now endow \mathcal{C}_q with a canonical non-strict monoidal structure. Given objects $(S, t), (S', t')$ in \mathcal{C}_q , set

$$(S, t) * (S', t') := (S * S', tt').$$

We also agree that $(S, t) * (\emptyset, \emptyset) := (S, t) =: (\emptyset, \emptyset) * (S, t)$. We immediately see from the definition of Par that

$$Par((S, t) * (S', t')) = Par(S * S', tt') = Par(S, t) \otimes Par(S', t'). \quad (4.2)$$

We can now define $*$ on arrows directly: given $f : (S_1, t_1) \rightarrow (S_2, t_2)$ and $g : (S'_1, t'_1) \rightarrow (S'_2, t'_2)$, we define $f * g$ as $Par(f * g) := f \otimes g$. More precisely, the arrow $f * g : (S_1, t_1) * (S'_1, t'_1) \rightarrow (S_2, t_2) * (S'_2, t'_2)$ is determined by the composite

$$\begin{aligned} Par((S_1, t_1) * (S'_1, t'_1)) &= Par(S_1, t_1) \otimes Par(S'_1, t'_1) \\ &\xrightarrow{f \otimes g} Par(S_2, t_2) \otimes Par(S'_2, t'_2) \\ &= Par((S_2, t_2) * (S'_2, t'_2)) \end{aligned}$$

The functoriality of $*$ now follows directly from that of \otimes . The unit for this monoidal product is given by (\emptyset, \emptyset) . We define the left and right unitality constraints as

the identity natural isomorphisms, so that $*$ is strictly left and right unital. The associativity constrain a_q

$$\begin{array}{ccc} [(S, t) * (S', t')] * (S'', t'') & \xrightarrow{a_q} & (S, t) * [(S', t') * (S'', t'')] \\ \parallel & & \parallel \\ (S * S' * S'', (tt')t'') & & (S * S' * S'', t(t't'')) \end{array}$$

is defined as $Par(a_q) := a_{Par(S,t), Par(S',t'), Par(S'',t'')}$. All these elements make \mathcal{C}_q into a monoidal category: the Triangle axiom holds trivially and the Pentagon axiom follows from that for \mathcal{C} . Note that this is a non-strict monoidal category, even if \mathcal{C} is strict, since $(tt')t''$ and $t(t't'')$ are in general different elements in $\text{Mag}(\bullet)$.

Theorem 4.2. *The previous non-strict monoidal structure on \mathcal{C}_q makes*

$$j : \mathcal{C} \xrightarrow{\cong} \mathcal{C}_q$$

a monoidal equivalence of categories.

Proof. It suffices to exhibit j as a strong monoidal functor. Let $u : (\emptyset, \emptyset) \longrightarrow ((\mathbf{1}), \bullet)$ be given by $Par(u) := \text{Id}_{\mathbf{1}}$, and set

$$\eta_{X,Y} : ((X), \bullet) * ((Y), \bullet) = ((X, Y), \bullet\bullet) \longrightarrow ((X \otimes Y), \bullet)$$

to be determined by $Par(\eta_{X,Y}) := \text{Id}_{X \otimes Y}$. Trivially, η assembles into a natural transformation $\eta : * \circ (i \times i) \Longrightarrow i \circ \otimes$. As in Theorem 3.2, the commutativity of the hexagon axiom and the left and right squares (2.1) correspond under (3.1) to the equalities $a_{X,Y,Z} = a_{X,Y,Z}$, $\ell_X = \ell_X$ and $r_X = r_X$, respectively, so they trivially hold. \square

4.2. Properties of non-strictification. We now state the main properties of the previous construction, that will say that the non-strictification is the free non-strict monoidal category generated by a monoidal category, in a completely analogous manner as Theorem 3.6 and Proposition 3.7.

Theorem 4.3. *Let \mathcal{C} be a monoidal category and let \mathcal{D} be a non-strict monoidal category. Given a strong monoidal functor $F : \mathcal{C} \longrightarrow \mathcal{D}$, there exists a unique strict monoidal functor $\widehat{F} : \mathcal{C}_q \longrightarrow \mathcal{D}$ such that $\widehat{F} \circ j = F$,*

$$\begin{array}{ccc} \mathcal{C}_q & & \\ \uparrow j & \searrow \widehat{F} & \\ \mathcal{C} & \xrightarrow{F} & \mathcal{D} \end{array}$$

Proof. Let us put $(\mathcal{C}, \otimes, \mathbf{1})$ and $(\mathcal{D}, \boxtimes, \mathbf{1})$ for the monoidal structures. If such \widehat{F} exists, then we must have

$$\widehat{F}(\emptyset, \emptyset) = \mathbf{1} \quad , \quad \widehat{F}((X), \bullet) = F(X) \quad , \quad \widehat{F}((S_1, t_1) * (S_2, t_2)) = \widehat{F}(S_1, t_1) \boxtimes \widehat{F}(S_2, t_2)$$

so since every object (S, t) with $|S| > 1$ can be uniquely written as $(S, t) = (S_1, t_1) * (S_2, t_2)$ with $|S_i| > 0$, the previous equalities inductively determine the only possible definition of \widehat{F} on objects.

To see what \widehat{F} must be on arrows, we argue closely to Theorem 3.6: if $\delta : \text{Id}_{\mathcal{C}_q} \Longrightarrow j \circ Par$ is the natural isomorphism of Lemma 4.1, then applying \widehat{F} to the

naturality of δ yields the commutative diagram

$$\begin{array}{ccc} \widehat{F}(S, t) & \xrightarrow{\widehat{F}(f)} & \widehat{F}(S', t') \\ \widehat{F}(\delta_{(S, t)}) \downarrow & & \downarrow \widehat{F}(\delta_{(S', t')}) \\ F(\text{Par}(S, t)) & \xrightarrow{F(\text{Par}(f))} & F(\text{Par}(S', t')) \end{array}$$

for any arrow $f : (S, t) \rightarrow (S', t')$ in \mathcal{C}_q . To see that $\widehat{F}(\delta_{(S, t)})$ only depends on F , we define a family of maps

$$\beta_{(S, t)}^F = \beta_{(S, t)} : \widehat{F}(S, t) \rightarrow F(\text{Par}(S, t))$$

inductively on the length as follows: if γ and u are the coherence constraints of F , set $\beta_{(\emptyset, \emptyset)} := u$, $\beta_{((X), \bullet)} := \text{Id}_{FX}$ and for $(S, t) = (S_1, t_1) * (S_2, t_2)$, let $\beta_{(S, t)}$ be the composite

$$\begin{aligned} \widehat{F}(S, t) &= \widehat{F}(S_1, t_1) \boxtimes \widehat{F}(S_2, t_2) \xrightarrow{\beta_{(S_1, t_1)} \boxtimes \beta_{(S_2, t_2)}} F(\text{Par}(S_1, t_1)) \boxtimes F(\text{Par}(S_2, t_2)) \\ &\xrightarrow{\gamma_{\text{Par}(S_1, t_1), \text{Par}(S_2, t_2)}} F(\text{Par}(S_1, t_1) \otimes \text{Par}(S_2, t_2)) \\ &\xlongequal{\hspace{10em}} F(\text{Par}(S, t)) \end{aligned}$$

An argument identical to that given in Theorem 3.6 shows that

$$\beta_{(S, t)} = \widehat{F}(\delta_{(S, t)}),$$

hence $\widehat{F}(f) = \beta_{(S', t')}^{-1} \circ F(\text{Par}(f)) \circ \beta_{(S, t)}$ is the only possible definition for $\widehat{F}(f)$. Furthermore, we have $\widehat{F} \circ j = F$ as strong monoidal functors by construction. \square

Proposition 4.4. *Let \mathcal{C} be a monoidal category and let \mathcal{D} be a non-strict monoidal category. Given a monoidal natural transformation $\alpha : F \Rightarrow G$ between strong monoidal functors $F, G : \mathcal{C} \rightarrow \mathcal{D}$, there exists a unique monoidal natural transformation $\widehat{\alpha} : \widehat{F} \Rightarrow \widehat{G}$ such that $\widehat{\alpha}j = \alpha$,*

Proof. If such $\widehat{\alpha}$ exists, then it must satisfy

$$\widehat{\alpha}_{(\emptyset, \emptyset)} = \text{Id}_{\mathbf{1}} \quad , \quad \widehat{\alpha}_{((X), \bullet)} = \alpha_X \quad , \quad \widehat{\alpha}_{(S, t) * (S', t')} = \widehat{\alpha}_{(S, t)} \boxtimes \widehat{\alpha}_{(S', t')},$$

and as in the previous arguments these rules inductively determine the value $\widehat{\alpha}_{(S, t)}$ for any object (S, t) , so this is the only possible definition. The naturality of $\widehat{\alpha}$ on (S, t) follows as in Proposition 3.7 considering a similar cube and arguing that it must be commutative. \square

4.3. Common realisations. In nature one finds many examples of strict monoidal categories whose collection of objects is given by the free monoid $\text{Mon}(X)$ on some set X . As we did with the strictification, we would like to make the non-strictification more concrete for this particular case.

Suppose that \mathcal{C} is a monoidal category with $\text{ob}(\mathcal{C}) = \text{Mon}(X)$ for some set X and whose monoidal product is given by the monoid product. Define a new category $\widetilde{\mathcal{C}}_q$ as follows: its objects are given by the elements of the free magma $\text{Mag}(X)$ on X . Write $U : \text{Mag}(X) \rightarrow \text{Mon}(X)$ for the canonical map that forgets parentheses and $p : \text{Mag}(X) \rightarrow \text{Mag}(\bullet)$ for the magma map induced by the unique map of sets $X \rightarrow \bullet$. For $v \in \text{Mag}(X)$, define its *sequencing* as

$$\text{Seq}(v) := ((x_1, \dots, x_n), p(v)) \in \mathcal{C}_q$$

where $U(v) = x_1 \cdots x_n$, and set $\text{Seq}(\emptyset) := (\emptyset, \emptyset)$. The hom sets in $\widetilde{\mathcal{C}}_q$ are defined as

$$\text{Hom}_{\widetilde{\mathcal{C}}_q}(v, v') := \text{Hom}_{\mathcal{C}_q}(\text{Seq}(v), \text{Seq}(v')),$$

and the identities and composite law are defined as those in \mathcal{C}_q . As before, this gives rise to a fully faithful functor

$$\text{Seq} : \widetilde{\mathcal{C}}_q \rightarrow \mathcal{C}_q.$$

Let us now define a monoidal structure on $\widetilde{\mathcal{C}}_q$: on objects, it is given by the magma product, $v \star v' := vv'$. Now, if $v, v' \in \text{Mag}(X)$, $U(v) = x_1 \cdots x_n$ and $U(v') = y_1 \cdots y_m$, then note that

$$\begin{aligned} \text{Seq}(v \star v') &= ((x_1, \dots, x_n, y_1, \dots, y_m), p(vv')) \\ &= ((x_1, \dots, x_n) * (y_1, \dots, y_m), p(v)p(v')) = \text{Seq}(v) * \text{Seq}(v') \end{aligned}$$

This allows us to define the monoidal product $f \star g$ of morphisms $f : v_1 \rightarrow v_2$ and $g : v'_1 \rightarrow v'_2$ as the composite

$$\text{Seq}(v_1 \star v'_1) = \text{Seq}(v_1) * \text{Seq}(v'_1) \xrightarrow{f \star g} \text{Seq}(v_2) * \text{Seq}(v'_2) = \text{Seq}(v_2 \star v'_2).$$

Once more, setting the empty word \emptyset as the unit, all this data determines a non-strict monoidal structure on $\widetilde{\mathcal{C}}_q$ such that Seq is naturally a strict monoidal functor.

As before, this implies

Proposition 4.5. *Let \mathcal{C} be a monoidal category with $\text{ob}(\mathcal{C}) = \text{Mon}(X)$ and monoidal product given by the monoid product. Then there is a strict monoidal equivalence*

$$\text{Seq} : \widetilde{\mathcal{C}}_q \xrightarrow{\cong} \mathcal{C}_q$$

between \mathcal{C}_q and a (non-strict) monoidal category $\widetilde{\mathcal{C}}_q$ such that $\text{ob}(\widetilde{\mathcal{C}}_q) = \text{Mag}(X)$ and whose monoidal product is given by the magma product.

Proof. Again it is left to check that Seq is essentially surjective also in this case. Given an object $(S, t) \in \mathcal{C}_q$, with $S = (w_1, \dots, w_n)$ and $t \in \text{Mag}(\bullet)$, let $v \in \text{Mag}(X)$ be an arbitrary parenthesisation of the product $w_1 \cdots w_n \in \text{Mon}(X)$. Then since the monoidal product of \mathcal{C} on object is given by concatenation we have

$$\text{Par}(S, t) = w_1 \cdots w_n = \text{Par}(\text{Seq}(v)),$$

which exhibits an isomorphism $(S, w) \cong \text{Seq}(v)$ in \mathcal{C}_q . \square

The previous result recovers the notion of non-strictification given in [HM21].

5. HIGHER CATEGORICAL PERSPECTIVE

In this section, we will upgrade the strictification and non-strictification to the 2-categorical level.

5.1. Enriched categories. Let $(\mathcal{C}, \otimes, \mathbf{1}, a, \ell, r)$ be a monoidal category. A \mathcal{C} -enriched category \mathcal{V} (or *category enriched over \mathcal{C}*) is the data of

- (1) a family of objects A, B, C, \dots of \mathcal{V} ,
- (2) for every pair of objects $A, B \in \mathcal{V}$, a hom-object $\mathcal{V}(A, B) \in \mathcal{C}$,
- (3) for each $A \in \mathcal{V}$, an arrow $1_A : \mathbf{1} \rightarrow \mathcal{V}(A, A)$ in \mathcal{C} ,
- (4) for every triple $A, B, C \in \mathcal{V}$, an arrow in \mathcal{C}

$$\circ : \mathcal{V}(B, C) \otimes \mathcal{V}(A, B) \rightarrow \mathcal{V}(A, C)$$

satisfying the following coherence conditions: for every $A, B, C, D \in \mathcal{V}$, we have the following commutative diagrams in \mathcal{C} :

$$\begin{array}{ccc} (\mathcal{V}(C, D) \otimes \mathcal{V}(B, C)) \otimes \mathcal{V}(A, B) & \xrightarrow{a} & \mathcal{V}(C, D) \otimes (\mathcal{V}(B, C) \otimes \mathcal{V}(A, B)) \\ \circ \otimes \text{Id}_{\mathcal{V}(A, B)} \downarrow & & \downarrow \text{Id}_{\mathcal{V}(C, D)} \otimes \circ \\ \mathcal{V}(B, D) \otimes \mathcal{V}(A, B) & & \mathcal{V}(C, D) \otimes \mathcal{V}(A, C) \\ & \searrow \circ & \swarrow \circ \\ & \mathcal{V}(A, D) & \end{array}$$

$$\begin{array}{ccc} \mathcal{V}(A, B) \otimes \mathbf{1} & \xrightarrow{\text{Id} \otimes 1_A} & \mathcal{V}(A, B) \otimes \mathcal{V}(A, A) \\ & \searrow r & \downarrow \circ \\ & & \mathcal{V}(A, B) \end{array} \quad \begin{array}{ccc} \mathcal{V}(B, B) \otimes \mathcal{V}(A, B) & \xleftarrow{1_B \otimes \text{Id}} & \mathbf{1} \otimes \mathcal{V}(A, B) \\ & \downarrow \circ & \swarrow \ell \\ & \mathcal{V}(A, B) & \end{array}$$

Every \mathcal{C} -enriched category \mathcal{V} has an underlying ordinary category \mathcal{V}_0 whose objects are those of \mathcal{V} and whose arrows are given by

$$\text{Hom}_{\mathcal{V}_0}(A, B) := \text{Hom}_{\mathcal{C}}(\mathbf{1}, \mathcal{V}(A, B)),$$

see e.g. [Rie14, §3.4].

5.2. Enriched functors. Let $(\mathcal{C}, \otimes, \mathbf{1})$ be a monoidal category and let \mathcal{V}, \mathcal{W} be \mathcal{C} -enriched categories. A \mathcal{C} -enriched functor $F : \mathcal{V} \rightarrow \mathcal{W}$ is the data of

- (1) for every object $A \in \mathcal{V}$, an object $F(A) \in \mathcal{W}$,
- (2) for every pair of objects $A, B \in \mathcal{V}$, an arrow in \mathcal{C}

$$F_{A, B} : \mathcal{V}(A, B) \rightarrow \mathcal{W}(F(A), F(B))$$

such that for every triple $A, B, C \in \mathcal{V}$ the following two diagrams commute:

$$\begin{array}{ccc} \mathcal{V}(B, C) \otimes \mathcal{V}(A, B) & \xrightarrow{\circ} & \mathcal{V}(A, C) \\ F_{B, C} \otimes F_{A, B} \downarrow & & \downarrow F_{A, C} \\ \mathcal{W}(F(B), F(C)) \otimes \mathcal{W}(F(A), F(B)) & \xrightarrow{\circ} & \mathcal{W}(F(A), F(C)) \end{array} \quad (5.1)$$

$$\begin{array}{ccc} & \mathbf{1} & \\ 1_A \swarrow & & \searrow 1_{FA} \\ \mathcal{V}(A, A) & \xrightarrow{F_{A, A}} & \mathcal{W}(F(A), F(A)) \end{array} \quad (5.2)$$

We observe that \mathcal{C} -enriched categories and \mathcal{C} -enriched functors form an ordinary category $\mathcal{C}\text{-Cat}$.

5.3. Enriched adjunctions. Let $(\mathcal{C}, \otimes, \mathbf{1})$ be a monoidal category. A \mathcal{C} -enriched adjunction is a pair $F : \mathcal{V} \rightarrow \mathcal{W}$ and $G : \mathcal{W} \rightarrow \mathcal{V}$ of \mathcal{C} -enriched functors together with a family of isomorphisms in \mathcal{C}

$$\varphi_{V,W} : \mathcal{W}(FV, W) \xrightarrow{\cong} \mathcal{V}(V, GW)$$

for every $V \in \mathcal{V}$ and $W \in \mathcal{W}$ which are \mathcal{C} -natural, meaning that for every $V, V' \in \mathcal{V}$ and $W, W' \in \mathcal{W}$ the following two diagrams are commutative:

$$\begin{array}{ccc} \mathcal{W}(FV, W) \otimes \mathcal{V}(V', V) & \xrightarrow{\text{Id} \otimes F_{V',V}} & \mathcal{W}(FV, W) \otimes \mathcal{W}(FV', FV) \\ \varphi_{V,W} \otimes \text{Id} \downarrow & & \downarrow \circ \\ \mathcal{V}(V, GW) \otimes \mathcal{V}(V', V) & \xrightarrow{\circ} & \mathcal{V}(V', GW) \\ & \swarrow \circ & \nwarrow \varphi_{V',W} \\ & \mathcal{V}(V', GW) & \end{array}$$

$$\begin{array}{ccc} \mathcal{W}(W, W') \otimes \mathcal{W}(FV, W) & \xrightarrow{G_{W,W'} \otimes \varphi_{V,W}} & \mathcal{V}(GW, GW') \otimes \mathcal{V}(V, GW) \\ \downarrow \circ & & \downarrow \circ \\ \mathcal{W}(FV, W') & \xrightarrow{\varphi_{V,W'}} & \mathcal{V}(V, GW') \end{array}$$

It should be clear from the definitions that, if $(\mathbf{Set}, \times, \bullet)$ denotes the monoidal category of sets and set-theoretical maps with the cartesian product \times as monoidal product and the singleton \bullet as unit, then a \mathbf{Set} -enriched category is the same as a locally small category, a \mathbf{Set} -enriched functor the same as an ordinary functor and a \mathbf{Set} -enriched adjunction the same as an ordinary adjunction.

Let \mathbf{Cat} be the category whose objects are small² categories and whose morphisms are functors with the usual composition. This is a monoidal category with respect to the cartesian product as monoidal product and the one-object discrete category as unit. A \mathbf{Cat} -enriched category is called a *2-category*, a \mathbf{Cat} -enriched functor is called a *2-functor* and a \mathbf{Cat} -enriched adjunction is called a *2-adjunction*. We write $\mathbf{2-Cat} := \mathbf{Cat-Cat}$ for the category of 2-categories and 2-functors.

If \mathcal{V} is a 2-category, then the objects A, B, C, \dots are typically called *0-cells*, the objects of the hom-categories $\mathcal{V}(A, B)$ *1-cells*, and the arrows of these hom-categories $\mathcal{V}(A, B)$ *2-cells*.

Notice that in a 2-category, we have the notion 1-cells and 2-cells being isomorphisms, that we call them *1-isomorphisms* and *2-isomorphisms*. Besides, we also have the notion of equivalence of objects: we say that a 1-cell $f : A \rightarrow B$ is an *equivalence* if there exists another 1-cell $g : B \rightarrow A$ and two 2-isomorphisms $\alpha_1 : g \circ f \rightarrow \text{Id}_A$ and $\alpha_2 : f \circ g \rightarrow \text{Id}_B$, where Id_C of an object C is the 1-cell image of the functor 1_C .

²For set-theoretical reasons.

5.4. Strictification as a 2-functor. Let MonCat be the 2-category whose 0-cells are monoidal categories, whose 1-cells are strong monoidal functors and whose 2-cells are monoidal natural isomorphisms. The composition of 2-cells inside a hom-category is given by vertical composition of the natural isomorphisms, whereas the composition \circ from (4) in Section 5.1 for 2-cells is given by horizontal composition. Similarly, we let strMonCat (resp. nonstrMonCat) be the 2-categories of strict (resp. non-strict) monoidal categories as 0-cells, strict monoidal functors as 1-cells and monoidal natural transformations as 2-cells.

Our aim is to upgrade the strictification of monoidal categories that we constructed in Section 3.1 to a 2-functor

$$\text{str} : \text{MonCat} \longrightarrow \text{strMonCat},$$

as follows: given a monoidal category \mathcal{C} , let $\text{str}(\mathcal{C}) := \mathcal{C}^{\text{str}}$. Given a strong monoidal functor $F : \mathcal{C} \longrightarrow \mathcal{D}$ between monoidal categories, define $\text{str}(F) = F^{\text{str}}$ in this manner: on objects, for $S = (X_1, \dots, X_n)$ in \mathcal{C} , put

$$F^{\text{str}}(S) := (FX_1, \dots, FX_n)$$

and $F^{\text{str}}(\emptyset) := \emptyset$. Given an arrow $f : S \longrightarrow S'$ in \mathcal{C} , we define $F^{\text{str}}(f)$ as the following dashed arrow:

$$\begin{array}{ccc} \text{Par}(F^{\text{str}}(S)) & \xrightarrow{\text{Par}(F^{\text{str}}(f))} & \text{Par}(F^{\text{str}}(S')) \\ \beta_S \downarrow & & \downarrow \beta_{S'} \\ F(\text{Par}(S)) & \xrightarrow{F(\text{Par}f)} & F(\text{Par}(S')) \end{array} \quad (5.3)$$

where β was defined in the proof of Theorem 3.6. It is clear that F^{str} is a functor $F^{\text{str}} : \mathcal{C}^{\text{str}} \longrightarrow \mathcal{D}^{\text{str}}$, and furthermore strict monoidal as we have

$$F^{\text{str}}(S) * F^{\text{str}}(S') = F^{\text{str}}(S * S').$$

Now let $\alpha : F \Longrightarrow G$ be a monoidal natural transformation of functors $\mathcal{C} \longrightarrow \mathcal{D}$. Define $\text{str}(\alpha) = \alpha^{\text{str}}$ as

$$\alpha_S^{\text{str}} : F^{\text{str}}(S) \longrightarrow G^{\text{str}}(S)$$

determined by

$$\text{Par}(\alpha_S^{\text{str}}) := (\dots(\alpha_{X_1} \boxtimes \alpha_{X_2}) \boxtimes \dots) \boxtimes \alpha_{X_n} : \text{Par}(F^{\text{str}}(S)) \longrightarrow \text{Par}(G^{\text{str}}(S))$$

where $S = (X_1, \dots, X_n)$. To see that α^{str} is natural on S , let $f : S \longrightarrow S'$ be an arrow in \mathcal{C}^{str} and consider the following cube:

$$\begin{array}{ccccc} & & \text{Par}(F^{\text{str}}(S)) & \xrightarrow{\text{Par}(F^{\text{str}}(f))} & \text{Par}(F^{\text{str}}(S')) \\ & \swarrow \beta_S & \downarrow F(\text{Par}(f)) & & \swarrow \beta_{S'} \\ F(\text{Par}(S)) & \xrightarrow{\quad} & F(\text{Par}(S')) & & \\ \downarrow \alpha_{\text{Par}(S)} & & \downarrow \alpha_{\text{Par}(S')} & & \downarrow \text{Par}(\alpha_{S'}^{\text{str}}) \\ & \swarrow \beta_S & \text{Par}(G^{\text{str}}(S)) & \xrightarrow{\text{Par}(G^{\text{str}}(f))} & \text{Par}(G^{\text{str}}(S')) \\ & \downarrow \alpha_{\text{Par}(S)} & \downarrow \alpha_{\text{Par}(S')} & & \downarrow \alpha_{\text{Par}(S')} \\ G(\text{Par}(S)) & \xrightarrow{\quad} & G(\text{Par}(S')) & & \end{array} \quad (5.4)$$

We have put $\beta_S = \beta_S^F$ and $\bar{\beta}_S = \beta_S^G$. The top and bottom faces commute by (5.3), the left and right faces by the monoidality of α and the front face by the

naturality of α . Since the arrows $\beta_S, \beta_{S'}, \bar{\beta}_S$ and $\bar{\beta}_{S'}$ are isomorphisms, this implies the commutativity of the back face, which is precisely the naturality of α^{str} . On the other hand, it is readily seen that $\alpha_S^{\text{str}} = i(\alpha_{X_1}) * \cdots * i(\alpha_{X_n})$, which means that α^{str} is a monoidal natural transformation. It is clear from the definition that for the vertical composition we have $(\alpha_2 \circ \alpha_1)^{\text{str}} = \alpha_2^{\text{str}} \circ \alpha_1^{\text{str}}$ and that the identity natural transformation maps to the identity natural transformation, so that for any pair of monoidal categories \mathcal{C}, \mathcal{D} , strictification induces a functor

$$\text{str}_{\mathcal{C}, \mathcal{D}} : \text{MonCat}(\mathcal{C}, \mathcal{D}) \longrightarrow \text{strMonCat}(\mathcal{C}^{\text{str}}, \mathcal{D}^{\text{str}}).$$

Proposition 5.1. *Strictification defines a 2-functor*

$$\text{str} : \text{MonCat} \longrightarrow \text{strMonCat}.$$

Proof. It is only left to check the commutativity of the diagrams (5.1) and (5.2). The first diagram (5.1) amounts to the equality of functors

$$(F_2 \circ F_1)^{\text{str}} = F_2^{\text{str}} \circ F_1^{\text{str}} \quad (5.5)$$

for $F_1 : \mathcal{C} \longrightarrow \mathcal{D}$ and $F_2 : \mathcal{D} \longrightarrow \mathcal{E}$, and the equality of natural transformations

$$(\alpha_2 * \alpha_1)^{\text{str}} = \alpha_2^{\text{str}} * \alpha_1^{\text{str}}, \quad (5.6)$$

where if $\alpha_i : F_i \implies G_i$ then $\alpha_2 * \alpha_1 : F_2 F_1 \implies G_2 G_1$ denotes the horizontal composition. Now, (5.5) holds trivially on objects, and on arrows we consider the following commutative diagram:

$$\begin{array}{ccc} \text{Par}(F_2^{\text{str}} F_1^{\text{str}} S) & \xrightarrow{F_2^{\text{str}} F_1^{\text{str}}(f)} & \text{Par}(F_2^{\text{str}} F_1^{\text{str}} S') \\ \beta_{F_1^{\text{str}} S}^2 \downarrow & & \downarrow \beta_{F_1^{\text{str}} S'}^2 \\ F_2(\text{Par}(F_1^{\text{str}} S)) & \xrightarrow{F_2(F_1^{\text{str}}(f))} & F_2(\text{Par}(F_1^{\text{str}} S')) \\ F_2 \beta_S^1 \downarrow & & \downarrow F_2 \beta_{S'}^1 \\ F_2 F_1(\text{Par}(S)) & \xrightarrow{F_2 F_1(\text{Par}f)} & F_2 F_1(\text{Par}(S')) \end{array}$$

In this diagram, we have put $\beta^i = \beta^{F_i}$. The bottom square is the image under F_2 of the square (5.3) for F_1^{str} , and the top square is the square (5.3) for F_1^{str} and the arrow $F_1^{\text{str}}(f)$. Now the key point to note is that $F_2 \beta_S^1 \circ \beta_{F_1^{\text{str}} S}^2 = \beta_S^{F_2 F_1}$, which readily follows from the definition of the monoidal constrain of the composite of monoidal functors. Moreover, (5.6) follows from the following computation (where we have removed the paranthesisation of the strictification of a monoidal natural transformation for clarity), where we take $S = (X_1, \dots, X_n) \in \mathcal{C}^{\text{str}}$:

$$\begin{aligned} (\alpha_2 * \alpha_1)_{X_1} \odot \cdots \odot (\alpha_2 * \alpha_1)_{X_n} &= \\ &= [G_2(\alpha_{1, X_1}) \circ \alpha_{2, F_1 X_1}] \odot \cdots \odot [G_2(\alpha_{1, X_n}) \circ \alpha_{2, F_1 X_n}] \\ &= [G_2(\alpha_{1, X_1}) \odot \cdots \odot G_2(\alpha_{1, X_n})] \odot [\alpha_{2, F_1 X_1} \odot \cdots \odot \alpha_{2, F_1 X_n}] \\ &= G_2^{\text{str}}((\alpha_1^{\text{str}})_S) \circ (\alpha_2^{\text{str}})_{F_1^{\text{str}} S}. \end{aligned}$$

Finally, (5.2) amounts to the equalities $(\text{Id}_{\mathcal{C}})^{\text{str}} = \text{Id}_{\mathcal{C}^{\text{str}}}$ and $(\text{id}_{\text{Id}_{\mathcal{C}}})^{\text{str}} = \text{id}_{\text{Id}_{\mathcal{C}^{\text{str}}}}$, where by $\text{id}_{\text{Id}_{\mathcal{C}}}$ we denote the identity natural transformation of $\text{Id}_{\mathcal{C}}$. These hold trivially by the definitions. \square

There is a canonical forgetful 2-functor

$$U : \text{strMonCat} \longrightarrow \text{MonCat}$$

sending every strict monoidal category, strict monoidal functor and monoidal natural transformation to their underlying monoidal category, underlying strong monoidal functor and the same natural transformation.

The following result upgrades Theorem 3.6 to the 2-categorical level, stating that the strictification is part of a free-forgetful adjunction at the 2-categorical level:

Theorem 5.2. *There is a 2-adjunction*

$$\begin{array}{ccc} & \xrightarrow{\text{str}} & \\ \text{MonCat} & \perp & \text{strMonCat} \\ & \xleftarrow{U} & \end{array}$$

Proof. For every monoidal category \mathcal{C} and every strict monoidal category \mathcal{D} , Theorem 3.6 and Proposition 3.7 together give an isomorphism of categories

$$\varphi_{\mathcal{C}, \mathcal{D}} : \text{strMonCat}(\mathcal{C}^{\text{str}}, \mathcal{D}) \xrightarrow{\cong} \text{MonCat}(\mathcal{C}, U\mathcal{D}),$$

so it is only left to check that this isomorphism is **Cat**-natural on \mathcal{C} and \mathcal{D} .

Let $H : \mathcal{C}_2 \rightarrow \mathcal{C}_1$ be a strong monoidal functor between monoidal categories, and let $F : \mathcal{C}_1 \rightarrow \mathcal{D}$ be a strong monoidal functor. Now consider the following diagram of categories and functors:

$$\begin{array}{ccccc} \mathcal{C}_2^{\text{str}} & \xrightarrow{H^{\text{str}}} & \mathcal{C}_1^{\text{str}} & & \\ \uparrow i_2 & & \uparrow i_1 & \searrow \widehat{F} & \\ \mathcal{C}_2 & \xrightarrow{H} & \mathcal{C}_1 & \xrightarrow{F} & \mathcal{D} \end{array}$$

The leftmost square commutes by the definition of H^{str} , so the whole diagram commutes. This means that $F \circ H = \widehat{F} \circ H^{\text{str}} \circ i_2$, which is precisely the naturality on \mathcal{C} for functors. The naturality for natural transformations follows by a similar argument from the following: if $H, H' : \mathcal{C}_1 \rightarrow \mathcal{D}$ are strong monoidal functors and $\varepsilon : H \Rightarrow H'$ is a monoidal natural transformation, then we have the equality of natural transformations

$$\varepsilon^{\text{str}} i_2 = i_1 \varepsilon,$$

which readily follows from the definition of ε^{str} .

The **Cat**-naturality on \mathcal{D} is a tautology since U simply forgets the strict structure. \square

We believe that the previous result should be known to experts, but to the author's knowledge there is no literature about it.

5.5. Non-strictification as a 2-functor. Now let us present a similar construction for the non-strictification, and showing that it similarly gives rise to a 2-functor

$$q : \text{MonCat} \rightarrow \text{nonstrMonCat}$$

which will be shown to be left-adjoint to the canonical forgetful functor

$$U : \text{nonstrMonCat} \rightarrow \text{MonCat}.$$

For a monoidal category \mathcal{C} , set $q(\mathcal{C}) := \mathcal{C}_q$. If $F : \mathcal{C} \rightarrow \mathcal{D}$ is a strong monoidal functor between monoidal categories, we define $F_q : \mathcal{C}_q \rightarrow \mathcal{D}_q$ as follows: on objects, put

$$F_q((X_1, \dots, X_n), t) := ((FX_1, \dots, FX_n), t)$$

and $F_q(\emptyset, \emptyset) := (\emptyset, \emptyset)$. For an arrow $f : (S, t) \longrightarrow (S', t')$ in \mathcal{C}_q , we let $F_q(f)$ be determined by following dashed arrow:

$$\begin{array}{ccc} \text{Par}(F_q(S, t)) & \overset{\text{Par}(F_q(f))}{\dashrightarrow} & \text{Par}(F_q(S', t')) \\ \beta_{(S, t)} \downarrow & & \downarrow \beta_{(S', t')} \\ F(\text{Par}(S, t)) & \xrightarrow{F(\text{Par}f)} & F(\text{Par}(S', t')) \end{array} \quad (5.7)$$

where β is this time as defined in the proof of Theorem 4.3. It is readily seen that F_q is a strict monoidal functor, in particular note that

$$F_q(S, t) * F_q(S', t') = F_q((S, t) * (S', t')).$$

Let us now describe the non-strictification for 2-cells. Let $F, G : \mathcal{C} \longrightarrow \mathcal{D}$ be strong monoidal functors between monoidal categories and let $\alpha : F \Longrightarrow G$ be a monoidal natural transformation. Then $\alpha_q : F_q \Longrightarrow G_q$ is defined objectwise as

$$(\alpha_q)_{(S, t)} : F_q(S, t) \longrightarrow G_q(S, t)$$

being given by

$$\text{Par}((\alpha_q)_{(S, t)}) := \alpha_{X_1} \boxtimes \cdots \boxtimes \alpha_{X_n} : \text{Par}(F_q(S, t)) \longrightarrow \text{Par}(G_q(S, t)),$$

where $S = (X_1, \dots, X_n)$ and $\alpha_{X_1} \boxtimes \cdots \boxtimes \alpha_{X_n}$ is supposed to be parenthesised according to $t \in \text{Mag}(\bullet)$. That α_q is indeed a natural transformation follows after considering a commutative cube similar to the one displayed in (5.4) for the strictification. We also have

$$(\alpha_q)_{(S, t) * (S', t')} = (\alpha_q)_{(S, t)} * (\alpha_q)_{(S', t')}$$

by definition which exhibits α_q as a monoidal natural transformation and besides $(\alpha_2 \circ \alpha_1)_q = (\alpha_2)_q \circ (\alpha_1)_q$. The upshot of this discussion is that we obtain a functor

$$q_{\mathcal{C}, \mathcal{D}} : \text{MonCat}(\mathcal{C}, \mathcal{D}) \longrightarrow \text{nonstrMonCat}(\mathcal{C}_q, \mathcal{D}_q).$$

Proposition 5.3. *Non-strictification defines a 2-functor*

$$q : \text{MonCat} \longrightarrow \text{nonstrMonCat}.$$

Proof. The argument is entirely analogous to the one given in Proposition 5.1. \square

Theorem 5.4. *There is a 2-adjunction*

$$\begin{array}{ccc} & q & \\ \text{MonCat} & \overset{\curvearrowright}{\longrightarrow} & \text{nonstrMonCat} \\ & \perp & \\ & \underset{\curvearrowleft}{\longleftarrow} & \\ & U & \end{array}$$

Proof. If \mathcal{C} is a monoidal category and \mathcal{D} is a non-strict monoidal category, then Theorem 4.3 and Proposition 4.4 define an isomorphism of categories

$$\varphi_{\mathcal{C}, \mathcal{D}} : \text{nonstrMonCat}(\mathcal{C}_q, \mathcal{D}) \xrightarrow{\cong} \text{MonCat}(\mathcal{C}, U\mathcal{D}).$$

The proof that this isomorphism is **Cat**-natural on \mathcal{C} and \mathcal{D} is identical to that given in Theorem 5.2. \square

5.6. Relation between the strictification and non-strictification. We have seen that any monoidal category is monoidally equivalent to both a strict and a non-strict monoidal category. This means that, up to monoidal equivalence, there is no distinction between strict and non-strict categories. We would like to make this idea more precise.

Let \mathcal{V} be a 2-category. The *core truncation* of \mathcal{V} is the ordinary category $h\mathcal{V}$ whose objects are those of \mathcal{V} and whose arrows are 2-isomorphism classes of 1-cells [Ban20]. More precisely, given a (small) ordinary category \mathcal{C} , let $[\mathcal{C}]$ be the discrete category whose objects are isomorphism classes of objects of \mathcal{C} . Trivially this defines a functor $[-] : \mathbf{Cat} \rightarrow \mathbf{Cat}$. Then the set of arrows in $h\mathcal{V}$ is defined as

$$\mathrm{Hom}_{h\mathcal{V}}(A, B) := \mathrm{Hom}_{\mathbf{Cat}}(\mathbf{1}, [\mathcal{V}(A, B)]),$$

where here $\mathbf{1}$ denotes the one-object discrete category. The identities and the composite law are inherited from those of \mathcal{V} in the obvious way. Note that the core truncation satisfies the property that if $f \in \mathcal{V}(A, B)$ is an equivalence, then the arrow $A \rightarrow B$ corresponding to the 2-isomorphism class of f is an isomorphism.

If $F : \mathcal{V} \rightarrow \mathcal{W}$ is a 2-functor between 2-categories, then define a functor

$$hF : h\mathcal{V} \rightarrow h\mathcal{W}$$

as $(hF)(A) := F(A)$ on objects and

$$\mathrm{Hom}_{\mathbf{Cat}}(\mathbf{1}, [\mathcal{V}(A, B)]) \xrightarrow{[F_{A,B}]^*} \mathrm{Hom}_{\mathbf{Cat}}(\mathbf{1}, [\mathcal{W}(FA, FB)])$$

on arrows. This defines the *core truncation functor*

$$h : 2\text{-}\mathbf{Cat} \rightarrow \mathbf{Cat}.$$

Remark 5.5. For the model/homotopical category theory minded reader, we would like to place the core truncation construction into a more general context. Suppose that \mathcal{C} is a closed symmetric monoidal homotopical category [Rie14], and \mathcal{V} is a \mathcal{C} -enriched category. If $\mathrm{Ho}(\mathcal{C})$ denotes the homotopy category of \mathcal{C} , then \mathcal{V} gives rise to a $\mathrm{Ho}(\mathcal{C})$ -enriched category $\bar{h}\mathcal{V}$, where the hom-objects are given by the images of the hom-objects $\mathcal{V}(A, B)$ under the localisation functor $\mathcal{C} \rightarrow \mathrm{Ho}(\mathcal{C})$. These hom-objects $\bar{h}\mathcal{V}(A, B)$ are thought as “homotopy classes”. We can furthermore take the underlying ordinary category of $\bar{h}\mathcal{V}$, that we denote $h\mathcal{V} := (\bar{h}\mathcal{V})_0$. Note that we have

$$\mathrm{Hom}_{h\mathcal{V}}(A, B) = \mathrm{Hom}_{\mathrm{Ho}(\mathcal{C})}(\mathbf{1}, \bar{h}\mathcal{V}(A, B)) = \mathrm{Hom}_{\mathrm{Ho}(\mathcal{C})}(\mathbf{1}, \mathcal{V}(A, B)).$$

Having said that, take $\mathcal{C} = \mathbf{Cat}$, the category of small categories with its canonical symmetric monoidal model (hence homotopical) structure, where the weak equivalences are given by the equivalences of categories. The hom-sets in $\mathrm{Ho}(\mathbf{Cat})$ are given by natural isomorphism classes of functors. If \mathcal{V} is a 2-category, then

$$\mathrm{Hom}_{\mathrm{Ho}(\mathbf{Cat})}(\mathbf{1}, \mathcal{V}(A, B)) = \mathrm{Hom}_{\mathbf{Cat}}(\mathbf{1}, [\mathcal{V}(A, B)]),$$

because the datum of a natural isomorphism between functors $i_f, i_g : \mathbf{1} \rightarrow \mathcal{V}(A, B)$ is equivalent to the datum of a 2-isomorphism $\alpha : f \rightarrow g$. This makes our notation $h\mathcal{V}$ consistent, and realises the core truncation as an instance of a construction in homotopical category theory.

Lemma 5.6. *The core truncation functor $h : 2\text{-Cat} \rightarrow \mathbf{Cat}$ preserves adjunctions, that is, if*

$$\begin{array}{ccc} & F & \\ \mathcal{V} & \xrightarrow{\quad} & \mathcal{W} \\ & \perp & \\ & G & \end{array}$$

is a 2-adjunction, then

$$\begin{array}{ccc} & hF & \\ h\mathcal{V} & \xrightarrow{\quad} & h\mathcal{W} \\ & \perp & \\ & hG & \end{array}$$

is an ordinary adjunction.

Proof. This follows directly applying the functor

$$\mathrm{Hom}_{\mathbf{Cat}}(\mathbf{1}, [-]) : \mathbf{Cat} \rightarrow \mathbf{Set}$$

to each of the isomorphisms of categories

$$\varphi_{V,W} : \mathcal{W}(FV, W) \xrightarrow{\cong} \mathcal{V}(V, GW)$$

of the 2-adjunction and to each of the commutative diagrams expressing the \mathbf{Cat} -naturality. \square

Theorem 5.7. *The core truncations of the (non-)strictification 2-functors define a pair of adjoint equivalences (i.e., adjunctions and equivalences of categories)*

$$\begin{array}{ccccc} & & hq & & hstr \\ & \curvearrowright & & \curvearrowleft & \\ h\mathrm{nonstrMonCat} & \perp & h\mathrm{MonCat} & \perp & h\mathrm{strMonCat} \\ & \curvearrowleft & & \curvearrowright & \\ & & hU & & hU \end{array}$$

Proof. That both pairs are adjunctions follows from the previous lemma. To see that they are also equivalences of categories, recall that given an adjunction $F \dashv G$, we have that F is fully faithful if and only if the unit of the adjunction is a natural isomorphism (e.g. [Rie17, 4.5.13]). The unit η of the adjunction $hstr \dashv hU$ is objectwise precisely the 2-isomorphism class of the functor $i : \mathcal{C} \rightarrow \mathcal{C}^{\mathrm{str}}$, hence η must be a natural isomorphism in light of Theorem 3.2, so that $hstr$ is fully faithful. On the other hand, if \mathcal{D} is a strict monoidal category, then we have that $Par : \mathcal{D}^{\mathrm{str}} \rightarrow \mathcal{D}$ is a strict equivalence of categories, which means that \mathcal{D} and $\mathcal{D}^{\mathrm{str}} = hstr(\mathcal{D})$ are isomorphic in $h\mathrm{strMonCat}$, and therefore $hstr$ is essentially surjective. For the non-strictification, the argument is identical. \square

REFERENCES

- [Ban20] P. Banks. Extended TQFTs and algebraic geometry, 2020, [2011.02394](#).
- [Bar13] M. Bartha. The monoidal structure of Turing machines. *Math. Structures Comput. Sci.*, 23(2):204–246, 2013. [doi:10.1017/S0960129512000096](#).
- [BN97] D. Bar-Natan. Non-associative tangles. In *Geometric topology (Athens, GA, 1993)*, volume 2 of *AMS/IP Stud. Adv. Math.*, pages 139–183. Amer. Math. Soc., Providence, RI, 1997. [doi:10.1090/amsip/002.1/09](#).
- [EGNO15] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik. *Tensor categories*, volume 205 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015. [doi:10.1090/surv/205](#).

- [GHW22] E. Gorsky, M. Hogancamp, and P. Wedrich. Derived traces of Soergel categories. *Int. Math. Res. Not. IMRN*, (15):11304–11400, 2022. doi:[10.1093/imrn/rnab019](https://doi.org/10.1093/imrn/rnab019).
- [HM21] K. Habiro and G. Massuyeau. The Kontsevich integral for bottom tangles in handlebodies. *Quantum Topol.*, 12(4):593–703, 2021. doi:[10.4171/qt/155](https://doi.org/10.4171/qt/155).
- [Hov99] M. Hovey. *Model categories*, volume 63 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999.
- [JS93] A. Joyal and R. Street. Braided tensor categories. *Adv. Math.*, 102(1):20–78, 1993. doi:[10.1006/aima.1993.1055](https://doi.org/10.1006/aima.1993.1055).
- [Kas95] C. Kassel. *Quantum groups*, volume 155 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. doi:[10.1007/978-1-4612-0783-2](https://doi.org/10.1007/978-1-4612-0783-2).
- [KRT97] C. Kassel, M. Rosso, and V. Turaev. *Quantum groups and knot invariants*, volume 5 of *Panoramas et Synthèses [Panoramas and Syntheses]*. Société Mathématique de France, Paris, 1997.
- [ML63] S. Mac Lane. Natural associativity and commutativity. *Rice Univ. Stud.*, 49(4):28–46, 1963.
- [MMSS01] M. A. Mandell, J. P. May, S. Schwede, and B. Shipley. Model categories of diagram spectra. *Proc. London Math. Soc. (3)*, 82(2):441–512, 2001. doi:[10.1112/S0024611501012692](https://doi.org/10.1112/S0024611501012692).
- [Pav13] D. Pavlovic. Monoidal computer I: Basic computability by string diagrams. *Inform. and Comput.*, 226:94–116, 2013. doi:[10.1016/j.ic.2013.03.007](https://doi.org/10.1016/j.ic.2013.03.007).
- [Rie14] E. Riehl. *Categorical homotopy theory*, volume 24 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2014. doi:[10.1017/CBO9781107261457](https://doi.org/10.1017/CBO9781107261457).
- [Rie17] E. Riehl. *Category theory in context*. Courier Dover Publications, 2017.
- [Sch99] S. Schwede. Stable homotopical algebra and Γ -spaces. *Math. Proc. Cambridge Philos. Soc.*, 126(2):329–356, 1999. doi:[10.1017/S0305004198003272](https://doi.org/10.1017/S0305004198003272).
- [Tur16] V. G. Turaev. *Quantum invariants of knots and 3-manifolds*, volume 18 of *De Gruyter Studies in Mathematics*. De Gruyter, Berlin, 2016. doi:[10.1515/9783110435221](https://doi.org/10.1515/9783110435221). Third edition [of MR1292673].
- [TV17] V. Turaev and A. Virelizier. *Monoidal categories and topological field theory*, volume 322 of *Progress in Mathematics*. Birkhäuser/Springer, Cham, 2017. doi:[10.1007/978-3-319-49834-8](https://doi.org/10.1007/978-3-319-49834-8).

BERNOULLI INSTITUTE, UNIVERSITY OF GRONINGEN, NIJENBORGH 9, 9747 AG, GRONINGEN, THE NETHERLANDS

Email address: j.becerra@rug.nl

URL: <https://sites.google.com/view/becerra/>